

SOME PROPERTIES OF MODULAR RELATIONS

By S. Bochner

(Received April 10, 1950)

Introduction

A modular relation will be for us a relation resembling an identity

$$\sum_0^{\infty} a_m e^{-\lambda_m x} = x^{-\delta} \sum_0^{\infty} b_n \exp\left(-\frac{\mu_n}{x}\right) \quad (1)$$

in the half-line $0 < x < \infty$ or rather in the half-plane $\operatorname{Re}(x) > 0$; and beyond always assuming $\delta > 0, \lambda_m \geq 0, \mu_n \geq 0$ and sometimes also

$$0 < \lambda_1 < \lambda_2 < \lambda_3 < \cdots \rightarrow \infty \quad (2)$$

$$0 < \mu_1 < \mu_2 < \mu_3 < \cdots \rightarrow \infty \quad (3)$$

and introducing some convergence assumptions, we will impose on the constants $a_m, \lambda_m, b_n, \mu_n$ no explicit restrictions in addition to those implied in the validity of (1). If we write (1) as

$$\Phi(x) = x^{-\delta} \Psi\left(\frac{1}{x}\right) \quad (4)$$

where

$$\Phi(x) = \sum_0^{\infty} a_m e^{-\lambda_m x} \quad (5)$$

$$\Psi(x) = \sum_0^{\infty} b_n e^{-\mu_n x} \quad (6)$$

then the relation (4) as such is of no interest whatsoever. It can be always inverted to

$$\Psi(x) = x^{-\delta} \Phi\left(\frac{1}{x}\right), \quad (7)$$

and we may start with any analytic function $\Phi(x)$ in $\operatorname{Re}(x) > 0$ and define $\Psi(x)$ by (7), and vice-versa. Thus some structural requirements of the type (5), (6) must be added. In the classical set-up the functions (5), (6) are "periodic" that is we have

$$\lambda_m = \lambda^* m, \quad \mu_n = \mu^* n \quad (8)$$

but even if we start from a family of such periodic relations

$$\Phi_j(x) = x^{-\delta_j} \Psi_j \left(\frac{1}{x} \right)$$

then very general “almost periodic” relations will ensue if we form products

$$\prod_j \Phi_j(x) = \prod_j x^{-\delta_j} \Psi_j \left(\frac{1}{x} \right)$$

for any δ_j , and finite or uniformly convergent infinite sums

$$\sum_j c_j \Phi_j(x) = x^{-\delta} \sum_j c_j \Psi_j \left(\frac{1}{x} \right)$$

for equal δ_j , and combinations of such.

Another source of general relations are, for instance, multidimensional relations

$$\sum_m A_m e^{-\text{trace}(L_m X)} = (\det |X|)^{-\delta} \sum_n B_n e^{-\text{trace}(M_n X^{-1})} \quad (9)$$

in which A_m, B_n are scalar constants, each L_m and M_n is a real symmetric positive semi-definite matrix $\{\lambda_{pq}\}$ and $\{\mu_{pq}\}$ respectively, $p, q = 1, \dots, k$, and X is a real symmetric strictly positive definite matrix $\{x_{pq}\}$ of the same finite dimension k , see [13]. If we replace the matrix X by the product Xt in which t is a positive scalar, then (9) implies

$$\sum_m A_m \exp(-\text{trace}(L_m X)t) = t^{-\delta} (\det |X|)^{-\delta} \sum_n B_n \exp(-\text{trace}(M_n X^{-1})t^{-1}), \quad (10)$$

and this is of the form (1) if we put

$$a_m = A_m (\det |X|)^{-\delta}, \quad b_n = B_n, \quad \lambda_m = \text{trace}(L_m X),$$

$$\mu_n = \text{trace}(M_n X^{-1}), \quad x = t,$$

and even if in the original matrices L_m, M_n the coefficients are multiples of integers, then for a general matrix X the resulting exponents λ_m, μ_n will no longer so be.

In the classical cases of (8) it is a familiar fact, compare [10], that the relation (1) is equivalent with the functional equation

$$\Gamma(\delta - s) \psi(\delta - s) = \Gamma(s) \varphi(s) \quad (11)$$

where

$$\varphi(s) = \sum_1^\infty \frac{a_m}{\lambda_m^s} \quad (12)$$

$$\psi(s) = \sum_1^\infty \frac{b_n}{\mu_n^s} \quad (13)$$

and that the constant terms a_0, b_0 of (8) which do not appear in (12) and (13) are the residues of the function $\chi(s)$ which is the joint value of the two sides in (11), this function having simple poles at $s = 0$ and $s = \delta$ at most. Now, in Section I we will examine this equivalence from a broad approach and we will see that the singularities of $\chi(s)$ may contribute a rather large additional term to the modular relation, but that it can be absorbed into it structurally by introducing a relation

$$\int e^{-\lambda x} dR(\lambda) = x^{-\delta} \int \exp\left(-\frac{\mu}{x}\right) dS(\mu) \quad (14)$$

which is more comprehensive than (1). Also near the end of Section III we will have some supplementary remarks about functional equations other than (11).

For particular cases of (1) it has been known and it is formally true in general that relation (1) implies the summation formula

$$\sum_0^{\infty} a_m f(\lambda_m) = \sum_0^{\infty} b_n g(\mu_n), \quad (15)$$

or more generally

$$\sum_0^{\infty} a_m f(\lambda_m x) = x^{-\delta} \sum_0^{\infty} b_n g\left(\frac{\mu_n}{x}\right), \quad x > 0 \quad (16)$$

for a pair of functions $f(\lambda), g(\mu)$ which are connected by the Hankel transformations

$$g(\mu) = \int_0^{\infty} J_{\delta-1}(2(\lambda\mu)^{1/2}) \lambda^{\frac{1}{2}(\delta-1)} f(\lambda) d\lambda \quad (17)$$

$$f(\lambda) = \int_0^{\infty} J_{\delta-1}(2(\lambda\mu)^{1/2}) \mu^{\frac{1}{2}(\delta-1)} g(\mu) d\mu \quad (18)$$

of which each is a formal reciprocal to the other; with $J_{\nu}(x)$ being the ordinary Bessel function as usual.

As regards validity of (15), a primitive but important situation arises if it is required that both sides in (15) shall also be absolutely convergent,

$$\sum |a_m| |f(\lambda_m)| < \infty, \quad \sum |b_n| |g(\mu_n)| < \infty, \quad (19)$$

and for this situation we will draw up in Section II a kind of master criterion from which specific ones, old and new, may be obtained. We also note that formula (16), if applied to (9) by way of (10) reads

$$\sum_m A_m f(\text{trace}(L_m X)) = (\det |X|)^{-\delta} \sum_n B_n g(\text{trace}(M_n X^{-1})) \quad (20)$$

where $f(\lambda)$ and $g(\mu)$ are connected by the same relations as before and we are stating here the following consequence from Theorem 10 in Section II.

Theorem 1. *If a relation (9) is given then for a matrix X and a number $\gamma > 0$ we have relation*

$$\Gamma(\gamma + 1) \sum_m A_m (\text{trace}(L_m X^{-1}))^{-\gamma-1} = \sum_n B_n (\text{trace}(M_n X))^{\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2(\text{trace}(M_n X))^{1/2}) \quad (21)$$

provided we have

$$\sum_{\text{tr}(M_n X) \leq 1} |B_n| + \sum_{\text{tr}(M_n X) > 1} |B_n| \{\text{tr}(M_n X)\}^{-\frac{1}{2}(\delta+\gamma)-\frac{1}{4}} < \infty,$$

and also

$$\sum_{\text{tr}(L_m X^{-1}) \leq 1} |A_m| + \sum_{\text{tr}(L_m X^{-1}) > 1} |A_m| \{\text{tr}(L_m X^{-1})\}^{-\alpha} < \infty,$$

for some $\alpha > 0$.

Finally in Section IV we will have statements on Abel and Riesz summability of the modular series

$$\sum_{\lambda_m \geq 1} \lambda_m^l \exp(i(\lambda_m x)^{1/2}) \quad (22)$$

for arbitrary l , and on the validity of the formula

$$\Gamma(\delta + 1) \sum a_m \left(\frac{x}{\lambda_m} \right)^{\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2(\lambda_m x)^{1/2}) = \sum_{\mu_n \leq x} b_n (x - \mu_n)^\gamma \quad (23)$$

for arbitrary values of γ , positive or negative, but only for such values of the variable x as are different from all exponents μ_n occurring. Our statements as such will not be new, see for instance, Ferrar [8], but our line of reasoning will, and we think that, syllogistically, the proof to Theorem 11 is the main contribution of the present paper. We have nothing to contribute to the case of x being equal an exponent μ_n and for completeness' sake we note that as far as is known to us, the latest "delicate" result bearing on this situation is the one due to Wilton and Hardy [18].

Section I. Functional Equations

1.

We recall that we always assume

$$\delta > 0.$$

In the complex s -plane, $s = \sigma + it$, we take two Mellin integrals

$$\chi_1(s) = \int_0^\infty x^{s-1} \Phi(x) dx \quad (24)$$

$$\chi_2(s) = \int_0^\infty x^{s-1} \Psi(x) dx \quad (25)$$

and we assume that they are absolutely convergent in a right half-plane, that is

$$\int_0^\infty x^{\sigma-1} |\Phi(x)| dx < \infty, \quad \int_0^\infty x^{\sigma-1} |\Psi(x)| dx < \infty \quad (26)$$

for $\sigma \geq \sigma_0$ for some $\sigma_0 > 0$. We also assume that $\Phi(x)$ and $\Psi(x)$ are differentiable in $0 < x < \infty$, although this assumption could be circumvented altogether.

Definition 1. We will say that two functions $\chi_1(s), \chi_2(s)$, as just introduced, satisfy the functional equation

$$\chi_1(s) = \chi_2(\delta - s) \quad (27)$$

if the following situation prevails. There is given in the s -plane a domain D which is the entire exterior of a bounded closed set S , with D and S both symmetric with respect to the line

$$\sigma = \frac{\delta}{2}, \quad -\infty < t < \infty \quad (28)$$

and in D there is given a holomorphic function $\chi(s)$ which in a right half-plane coincides with $\chi_1(s)$ and in a left half-plane with $\chi_2(\delta - s)$, and furthermore we have

$$\lim_{|t| \rightarrow \infty} \chi(\sigma + it) = 0 \quad (29)$$

uniformly in every bounded interval

$$\sigma_1 \leq \sigma \leq \sigma_2, \quad -\infty < \sigma_1 < \sigma_2 < \infty. \quad (30)$$

Before proceeding we will immediately introduce another definition.

Definition 2. We will say that a function $P(x)$ is a *residual function* if it has the following properties: (i) It is defined and differentiable in $0 < x < \infty$, and for some $\gamma > 0$ we have $P(x) = O(x^{-\gamma})$ as $x \rightarrow 0$ and $= O(x^\gamma)$ as $x \rightarrow \infty$, so that we may introduce the functions

$$\chi_r(s) = \int_0^1 P(x) x^{s-1} dx \quad (31)$$

$$\chi_l(s) = - \int_1^\infty P(x) x^{s-1} dx \quad (32)$$

the first in a right half-plane and the second in a left half-plane. (ii) These two functions can be continued, the one into the other, in a domain D as introduced in Definition 1. (iii) We also have

$$\lim_{|t| \rightarrow \infty} \chi_r(\sigma + it) = 0 \quad (33)$$

uniformly in (30), where $\chi_r(s)$ simply denotes the joint value of the two functions (31), (32) in the domain D .

Lemma 1. Any integral

$$P(x) = \frac{1}{2\pi i} \int_C \chi(\zeta) x^{-\zeta} d\zeta \quad (34)$$

over a bounded curve or curves C , with $x^{-\zeta} = \exp(-\zeta \log x)$, is a residual function. So is also any finite sum

$$\sum_{\zeta} a_{\zeta} x^{-\zeta} \quad (35)$$

with

$$-a_0 + b_0 x^{-\delta} \quad (36)$$

occurring most frequently, and more generally any finite sum

$$\sum a_{p\zeta} (\log x)^p x^{-\zeta} \quad (37)$$

where $p = 0, 1, 2, \dots$, and ζ is complex.

Proof. If $\gamma > 0$ is such that C lies in the strip $-\gamma < \sigma < \gamma$ then property (i) of Definition 2 is fulfilled. Now, for $\sigma > \gamma$ we have

$$\begin{aligned} \int_0^1 P(x) x^{s-1} dx &= \frac{1}{2\pi i} \int_0^1 dx \cdot x^{s-1} \int_C d\zeta \cdot \chi(\zeta) x^{-\zeta} \\ &= \frac{1}{2\pi i} \int_C d\zeta \cdot \chi(\zeta) \int_0^1 x^{s-\zeta-1} dx = \frac{1}{2\pi i} \int_C \frac{\chi(\zeta) d\zeta}{s-\zeta} \end{aligned}$$

and for $\sigma < -\gamma$ we have

$$\begin{aligned} - \int_1^\infty P(x) x^{s-1} dx &= -\frac{1}{2\pi i} \int_1^\infty dx \cdot x^{s-1} \int_C d\zeta \cdot \chi(\zeta) x^{-\zeta} \\ &= -\frac{1}{2\pi i} \int_C d\zeta \cdot \chi(\zeta) \int_1^\infty x^{s-\zeta-1} dx = \frac{1}{2\pi i} \int_C \frac{\chi(\zeta) d\zeta}{s-\zeta} \end{aligned}$$

and from this (ii) and (iii) follow easily.

Also, for $P(x) = (\log x)^p x^{-\zeta}$ we obtain for (31) and (32) the same expression $-\Gamma(p+1)(\zeta-s)^{-p-1}$ and thus (37) is a residual function.

Theorem 2. *If two Mellin integrals (24), (25) satisfy the functional equation (27), then their integrands satisfy the (modular) relation*

$$\Phi(x) - x^{-\delta} \Psi\left(\frac{1}{x}\right) = P(x) \quad (38)$$

in which $P(x)$ is a residual function representable as an integral (34) in which C is a curve or curves in D encircling all of S .

Proof. If we choose a sufficiently large number $a > 0$, then by Mellin inversion of (24) we obtain

$$\Phi(x) = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{a-iT}^{a+iT} \chi_1(s) x^{-s} ds, \quad (39)$$

and due to (29) the right side is

$$\frac{1}{2\pi i} \int_C + \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{\delta-a-iT}^{\delta-a+iT} \chi(s) x^{-s} ds \quad (40)$$

provided C is located in the strip $\delta - a < \sigma < a$, and this will take place if a is sufficiently large. The first term in (40) is our residual function; and if in the second we write $\chi_2(\delta - s)$ for $\chi(s)$ and replace s by $\delta - s$, we obtain

$$x^{-\delta} \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{a-iT}^{a+iT} \chi_2(s) \left(\frac{1}{x}\right)^{-s} ds$$

which by Mellin inversion of (25) is $x^{-\delta} \Psi\left(\frac{1}{x}\right)$, as claimed.

Theorem 3. *Conversely, if we are given in $0 < x < \infty$ a (modular) relation (38) in which $P(x)$ is any residual function according to Definition 2, and if $\Phi(x)$ and $\Psi(x)$ are differentiable functions for which the restriction (26) holds so as to give rise to Mellin integrals (24), (25), then the latter satisfy the functional equation (27) in the sense of Definition 1.*

Remark. In particular, by Theorem 2, $P(x)$ can be represented by an integral (34) and this statement is a kind of converse to Lemma 1.

Proof. We have

$$\begin{aligned} \chi_1(s) &= \int_0^\infty x^{s-1} \Phi(x) dx \\ &= \int_0^1 x^{s-1} \Phi(x) dx + \int_1^\infty x^{s-1} \Phi(x) dx + \int_0^1 x^{s-1} P(x) dx + \int_1^\infty x^{s-1} P(x) dx \\ &= \chi_1^*(s) + \chi_r(s), \end{aligned}$$

where $\chi_r(s)$ is defined by (31) and

$$\chi_1^*(s) = \int_0^1 x^{s-1} \Phi(x) dx + \int_1^\infty x^{s-1} \Psi(x) dx. \quad (41)$$

Now, for $\chi_r(s)$ we have stipulated explicitly that it can be extended into a domain D and that (33) holds for it, and for $\chi_1^*(s)$ we are going to prove this. It follows easily from (26) that the two integrals comprising (41) are each absolutely uniformly convergent in every strip (30) so that (41) is actually an entire function. As for the validity of

$$\lim_{|t| \rightarrow \infty} \chi_1^*(\sigma + it) = 0 \quad (42)$$

uniformly in $\sigma_1 \leq \sigma \leq \sigma_2$ we note that this is nothing else but the Riemann-Lebesgue Lemma for the sum of Fourier integrals

$$\int_0^1 (e^{i(s \log x)} \Phi(x) + e^{i(\delta-s) \log x} \Psi(x)) d(\log x)$$

uniformly in the parameter σ and can be easily verified.

Next we note that relation (38) implies

$$\Psi(x) = x^{-\delta} \Phi\left(\frac{1}{x}\right) - x^{-\delta} P\left(\frac{1}{x}\right) \quad (43)$$

and therefore we have

$$\begin{aligned}
\chi_2(s) &= \int_0^\infty x^{s-1} \Psi(x) dx \\
&= \int_0^1 x^{s-1} \Psi(x) dx + \int_1^\infty x^{s-1} \Psi(x) dx \\
&= \int_0^1 x^{s-1-\delta} \Phi\left(\frac{1}{x}\right) dx - \int_0^1 x^{s-1-\delta} P\left(\frac{1}{x}\right) dx + \int_1^\infty x^{s-1} \Psi(x) dx \\
&= \int_1^\infty x^{\delta-s-1} \Phi(x) dx + \chi_l(\delta-s) + \int_1^\infty x^{s-1} \Psi(x) dx \\
&= \chi_2^*(\delta-s) + \chi_l(\delta-s),
\end{aligned}$$

and equation (27) follows now from the postulated equality of (31) and (32).

2.

We now take a Dirichlet series (12) with (2) and we assume that for a constant $\alpha > 0$ we have

$$\sum_1^\infty \frac{|a_m|}{\lambda_m^\alpha} < \infty. \quad (44)$$

If with the Gamma integral

$$\Gamma(s) = \int_0^\infty x^{s-1} e^{-x} dx$$

we form the product

$$\chi_1(s) = \Gamma(s) \varphi(s) \quad (45)$$

then we can write for it the Mellin integral (24) where $\Phi(x)$ is given by (5). Similarly if we introduce the Dirichlet series (13) with (3) and if for a constant $\beta > 0$ we have

$$\sum_1^\infty \frac{|b_n|}{\mu_n^\beta} < \infty \quad (46)$$

then for

$$\chi_2(s) = \Gamma(s) \psi(s) \quad (47)$$

we have the Mellin integral (25) where $\Psi(x)$ is (6), and our Theorems 2 and 3 express the following equivalence, subject to details.

Theorem 4. *A Hecke functional equation*

$$\Gamma(s) \varphi(s) = \Gamma(\delta-s) \psi(\delta-s) = \chi(s) \quad (48)$$

where

$$\varphi(s) = \sum_1^\infty \frac{a_m}{\lambda_m^s}, \quad \psi(s) = \sum_1^\infty \frac{b_n}{\mu_n^s} \quad (49)$$

implies the modular relation

$$\sum_1^\infty a_m e^{-\lambda_m x} - x^{-\delta} \sum_1^\infty b_n \exp\left(-\frac{\mu_n}{x}\right) = P(x) \quad (50)$$

in which $P(x)$ is a residual function and is given by an integral (34); and conversely any relation (50) with any residual function $P(x)$ leads back to an equation (48).

We may generalize this by introducing in (48) the LaplaceStieltjes integrals

$$\varphi(s) = \int_{+0}^\infty \lambda^{-s} dA(\lambda) \quad (51)$$

$$\psi(s) = \int_{+0}^\infty \mu^{-s} dB(\mu) \quad (52)$$

with

$$\int_{+0}^\infty \lambda^{-\alpha} |dA(\lambda)| < \infty, \quad \int_{+0}^\infty \mu^{-\beta} |dB(\mu)| < \infty \quad (53)$$

in which case (50) is to be replaced by

$$\int_{+0}^\infty e^{-\lambda x} dA(\lambda) - x^{-\delta} \int_{+0}^\infty \exp\left(-\frac{\mu}{x}\right) dB(\mu) = P(x) \quad (54)$$

and this includes the previous case if we put

$$A(\lambda) = \sum_{\lambda_m < \lambda} a_m, \quad B(\mu) = \sum_{\mu_n < \mu} b_n. \quad (55)$$

In forming the expressions (55) we may abandon the requirement that the exponents λ_m, μ_n shall be simple sequences (2), (3), and if we apply this to (10), recalling that (36) is a residual function, we obtain the following statement.

Theorem 5. *If there is given a multi-dimensional relation (9), and if corresponding to a matrix X there are numbers $\alpha > 0, \beta > 0$ such that*

$$\sum_{\text{tr}(L_m X)=0} |A_m| + \sum_{\text{tr}(L_m X)>0} (\text{tr}(L_m X))^{-\alpha} |A_m| < \infty \quad (56)$$

and

$$\sum_{\text{tr}(M_n X^{-1})=0} |B_n| + \sum_{\text{tr}(M_n X^{-1})>0} (\text{tr}(M_n X^{-1}))^{-\beta} |B_n| < \infty, \quad (57)$$

then we have the functional equation

$$\Gamma(s) \varphi(s; X) = (\det |X|)^{-\delta} \Gamma(\delta - s) \psi(\delta - s; X^{-1}) \quad (58)$$

where

$$\varphi(s; X) = \sum_{\text{tr}(L_m X)>0} A_m (\text{tr}(L_m X))^{-s} \quad (59)$$

and

$$\psi(s; X^{-1}) = \sum_{\text{tr}(M_n X^{-1})>0} B_n (\text{tr}(M_n X^{-1}))^{-s}. \quad (60)$$

3.

Usually the function $\chi(s)$ will have but a finite number of unramified singularities and the point set S will consist of these points only. In this case the point set S certainly does not contain a continuum linking the line $\sigma = 0$ with the line $\sigma = \delta$. If now we impose on S the non-linking property just stated (and nothing more specific) then it is possible to replace in (34) the system of curves C encircling S by a combinatorial sum $-C_1 + C_2$ in such a manner that C_1 is located in the right half-plane $\sigma > 0$ and C_2 in the left half-plane $\sigma < \delta$. We can then put

$$P(x) = -M(x) + x^{-\delta} N\left(\frac{1}{x}\right) \quad (61)$$

where

$$M(x) = \frac{1}{2\pi i} \int_{C_1} \chi(s) x^{-s} ds \quad (62)$$

and

$$N(x) = \frac{1}{2\pi i} \int_{C_2} \chi(s) x^{s-\delta} ds, \quad (63)$$

and by the use of the integrals

$$\Gamma(s) x^{-s} = \int_0^\infty e^{-\lambda x} \lambda^{s-1} d\lambda, \quad \Gamma(\delta - s) x^{s-\delta} = \int_0^\infty e^{-\mu x} \mu^{\delta-s-1} d\mu$$

we obtain

$$M(x) = \int_0^\infty e^{-\lambda x} u(\lambda) d\lambda \quad (64)$$

$$N(x) = \int_0^\infty e^{-\mu x} v(\mu) d\mu \quad (65)$$

where

$$u(\lambda) = \frac{1}{2\pi i} \int_{C_1} \varphi(s) \lambda^{s-1} ds, \quad v(\mu) = \frac{1}{2\pi i} \int_{C_2} \psi(\zeta) \mu^{\delta-\zeta-1} d\zeta. \quad (66)$$

If now we introduce the indefinite integrals

$$U(\lambda) = \int_0^\lambda u(\lambda) d\lambda, \quad V(\mu) = \int_0^\mu v(\mu) d\mu \quad (67)$$

and then put

$$R(\lambda) = A(\lambda) + U(\lambda), \quad S(\mu) = B(\mu) + V(\mu) \quad (68)$$

then the relation (54) assumes the “unified” appearance

$$\int_0^\infty e^{-\lambda x} dR(\lambda) = x^{-\delta} \int_0^\infty \exp\left(-\frac{\mu}{x}\right) dS(\mu) \quad (69)$$

in which the residual part $P(x)$ has been assimilated to the modular part proper.

This assimilation, however, is a very superficial one in that relation (69) as such is no longer equivalent to a functional equation

$$\Gamma(s) \tilde{\varphi}(s) = \Gamma(\delta - s) \tilde{\psi}(\delta - s) \quad (70)$$

with would-be expansions

$$\tilde{\varphi}(s) = \int_0^\infty \lambda^{-s} dR(\lambda), \quad \tilde{\psi}(s) = \int_0^\infty \mu^{-s} dS(\mu), \quad (71)$$

such expansions not converging. In fact if we take an equation (48) of the classical type in which $\chi(s)$ has two simple poles, at $s = 0$ and $s = \delta$ then $P(x)$ is of the form (36), and our decomposition (61) can be done in only one way with

$$M(x) = -b_0 x^{-\delta}, \quad N(x) = -a_0 x^{-\delta} \quad (72)$$

$$u(\lambda) = -b_0 \lambda^{\delta-1}, \quad v(\mu) = -a_0 \mu^{\delta-1}. \quad (73)$$

The would-be expansion $\tilde{\varphi}(s)$ would then be the sum of the previous expansion $\varphi(s)$, which exists, and of a would-be integral

$$\int_0^\infty \lambda^{-s} u(\lambda) d\lambda = -b_0 \int_0^\infty \lambda^{-s+\delta-1} d\lambda,$$

which latter however is undefinable.

We further notice that if there are singularities of $\chi(s)$ in the interior of the strip $0 < \sigma < \delta$, the decomposition (61) of $P(x)$ is no longer unique, and that the terms originating from the poles at $s = 0$ and $s = \delta$ are assigned wrong positions as it were. Thus if we start from a relation

$$\sum a_m e^{-\lambda_m x} - x^{-\delta} \sum b_n \exp(-\mu_n x^{-1}) = -a_0 + b_0 x^{-\delta} \quad (74)$$

then the decomposition (61) leads to

$$-b_0 x^{-\delta} + \sum a_m e^{-\lambda_m x} = x^{-\delta} (-a_0 x + \sum b_n \exp(\mu_n x^{-1})) \quad (75)$$

whereas, obviously, the “natural” decomposition would be

$$\sum a_m e^{-\lambda_m x} = x^{-\delta} \sum b_n \exp(\mu_n x^{-1}) \quad (76)$$

with $\lambda_0 = \mu_0 = 1$.

We are for the present terminating the general discussion of “relations” and “equations” and further supplementary remarks will be made in Section III.

Section II. Summation Formulas

4.

We now take any relation of the form (69), that is

$$\int_0^\infty e^{-\lambda x} dR(\lambda) = x^{-\delta} \int_0^\infty \exp\left(-\frac{\mu}{x}\right) dS(\mu), \quad (77)$$

no matter how originating, and we only postulate as follows. The function $R(\lambda)$ is of bounded variation in any finite interval $0 \leq \lambda \leq \lambda'$ and so is $S(\mu)$ in any finite interval $0 \leq \mu \leq \mu'$, and either function may have a saltus at the origin so as to include equation (76) with a_0 and b_0 not zero; and as before we assume that there are constants $\alpha > 0, \beta > 0$, which will be always denoted by these letters, such that

$$\int_1^\infty \lambda^{-\alpha} |dR(\lambda)| < \infty \quad (78)$$

$$\int_1^\infty \mu^{-\beta} |dS(\mu)| < \infty. \quad (79)$$

Now, such a relation (77) “generates” the “summation formula”

$$\int_0^\infty f(\lambda) dR(\lambda) = \int_0^\infty g(\mu) dS(\mu) \quad (80)$$

in which $f(\lambda)$ is an “arbitrary” function in $0 \leq \lambda < \infty$ and $g(\mu)$ is the Hankel transform

$$g(\mu) = \mu^{-\frac{1}{2}(\delta-1)} \int_0^\infty J_{\delta-1}(2(\lambda\mu)^{1/2}) \lambda^{\frac{1}{2}(\delta-1)} f(\lambda) d\lambda, \quad (81)$$

in which $J_\nu(x)$ is the ordinary Bessel function. The transformation (81) is self-inversive, so that, at least formally, it implies and is implied by the formula

$$f(\lambda) = \lambda^{-\frac{1}{2}(\delta-1)} \int_0^\infty J_{\delta-1}(2(\lambda\mu)^{1/2}) \mu^{\frac{1}{2}(\delta-1)} g(\mu) d\mu; \quad (82)$$

and we also note that on substituting

$$J_\nu(x) = \frac{x^\nu}{\Gamma(\nu+1)} \{1 + (\text{higher powers})\},$$

the transforms (81) and (82) obtain the limiting values

$$g(0) = \int_0^\infty \frac{\lambda^{\delta-1}}{\Gamma(\delta)} f(\lambda) d\lambda$$

$$f(0) = \int_0^\infty \frac{\mu^{\delta-1}}{\Gamma(\delta)} g(\mu) d\mu$$

and that such a limiting value is to be employed whenever $S(\mu)$ or $R(\lambda)$ has a discontinuity at the origin.

Furthermore if

$$\{f(\lambda), g(\mu)\}$$

are a pair of functions thus connected then for every $x > 0$ the pair

$$\left\{ f(\lambda x), \quad x^{-\delta} g\left(\frac{\mu}{x}\right) \right\}$$

will also so be, and thus we might just as well write (80) in the seemingly more general form

$$\int_0^\infty f(\lambda x) dR(\lambda) = x^{-\delta} \int_0^\infty g\left(\frac{\mu}{x}\right) dS(\mu) \quad (83)$$

of which it is technically the special case $x = 1$. If so written, the summation formula subsumes the original relation (77) as a particular case of itself in agreement with the known formula, see [17], p. 394,

$$e^{-\lambda} = \lambda^{-\frac{1}{2}(\delta-1)} \int_0^\infty J_{\delta-1}(2(\lambda\mu)^{1/2}) \mu^{\frac{1}{2}(\delta-1)} e^{-\mu} d\mu \quad (84)$$

and we are going to use this fact very directly towards obtaining a rigorous criterion for the validity of our summation formula; and the criterion will be such that in consequence of its hypothesis both sides in (80) will be convergent absolutely.

5.

Relation (77) is valid not only for real x in the half-line

$$0 < x < \infty$$

but for complex $x = \xi + i\eta$ in the half-plane

$$H : 0 < \xi < \infty, \quad -\infty < \eta < \infty$$

thus,

$$\int_0^\infty e^{-\lambda(\xi+i\eta)} dR(\lambda) = (\xi + i\eta)^{-\delta} \int_0^\infty \exp\left(-\frac{\mu}{\xi + i\eta}\right) dS(\mu)$$

and this implies formally the equality

$$\int_H \left(\int_0^\infty e^{-\lambda(\xi+i\eta)} dR(\lambda) \right) d\rho(\xi, \eta) = \int_H (\xi + i\eta)^{-\delta} \left(\int_0^\infty e^{-\mu(\xi+i\eta)^{-1}} dS(\mu) \right) d\rho(\xi, \eta)$$

for an “arbitrary” StieltjesRadon “distribution” $\rho(\xi, \eta)$ in H . If we invert integrations we obtain

$$\int_0^\infty f(\lambda) dR(\lambda) = \int_0^\infty \tilde{g}(\mu) dS(\mu)$$

where

$$f(\lambda) = \int_H e^{-\lambda(\xi+i\eta)} d\rho(\xi, \eta) \quad (85)$$

$$\tilde{g}(\mu) = \int_H (\xi + i\eta)^{-\delta} e^{-\mu(\xi+i\eta)^{-1}} d\rho(\xi, \eta), \quad (86)$$

and by Fubini’s theorem all this is rigorous, with all occurring integrals converging absolutely, provided we have

$$\int_H |d\rho(\xi, \eta)| < \infty \quad (87)$$

$$\int_H (\xi^2 + \eta^2)^{-\delta/2} |d\rho(\xi, \eta)| < \infty$$

$$\int_H e^{-\lambda\xi} |d\rho(\xi, \eta)| = O(\lambda^{-\alpha}), \quad \lambda \rightarrow \infty \quad (88)$$

$$\int_H (\xi^2 + \eta^2)^{-\delta/2} \exp\left(-\frac{\mu\xi}{\xi^2 + \eta^2}\right) |d\rho(\xi, \eta)| = O(\mu^{-\beta}), \quad \mu \rightarrow \infty. \quad (89)$$

However we do not yet know whether the function $\tilde{g}(\mu)$ is identical with the function $g(\mu)$ as related to $f(\lambda)$ by means of relation (81). But from (84) we deduce

$$(\xi + i\eta)^{-\delta} e^{-\mu(\xi+i\eta)^{-1}} = \mu^{-\frac{1}{2}(\delta-1)} \int_0^\infty J_{\delta-1}(2(\mu\lambda)^{1/2}) \lambda^{\frac{1}{2}(\delta-1)} e^{-\lambda(\xi+i\eta)} d\lambda \quad (90)$$

and if we substitute this in (86) and apply Fubini's theorem we obtain indeed (81), provided we have

$$\int_{\lambda=0}^\infty \mu^{-\frac{1}{2}(\delta-1)} \int_H |J_{\delta-1}(2(\lambda\mu)^{1/2})| \lambda^{\frac{1}{2}(\delta-1)} e^{-\lambda\xi} |d\rho(\xi, \eta)| d\lambda < \infty \quad (91)$$

for every $\mu \geq 0$. Now, for $\delta \geq \frac{1}{2}$ we have

$$|J_{\delta-1}(2x)| \leq Mx^{\delta-1}, \quad 0 < x < \infty \quad (92)$$

whereas for $0 < \delta < \frac{1}{2}$ we have

$$|J_{\delta-1}(2x)| \leq \begin{cases} Mx^{\delta-1}, & 0 < x \leq 1 \\ Mx^{-\frac{1}{2}}, & 1 < x < \infty \end{cases} \quad (93)$$

and therefore (91) is satisfied if we have

$$\int_H \xi^{-\delta} |d\rho(\xi, \eta)| < \infty, \quad \text{for } 0 < \delta < \infty, \quad (94)$$

and also

$$\int_H \xi^{-(\delta/2+1/4)} |d\rho(\xi, \eta)| < \infty \quad \text{for } 0 < \delta < \frac{1}{2}, \quad (95)$$

and this is certainly the case if for $0 < \delta < \infty$ assumptions (87) and (94) hold and for $0 < \delta < \frac{1}{2}$ also (95).

6.

If the set-function in H is concentrated entirely on the half-line $0 < \xi < \infty$, $\eta = 0$ we have a representation

$$f(\lambda) = \int_0^\infty e^{-\lambda\xi} d\rho(\xi), \quad (96)$$

and if the set function is also non-negative then on account of (87) we have

$$d\rho(\xi) \geq 0, \quad \int_0^\infty d\rho(\xi) < \infty \quad (97)$$

and our function $f(\lambda)$ is completely monotone in $0 \leq \lambda < \infty$. The two conditions (87) and (94) merge into the one

$$\int_0^\infty \xi^{-\delta} d\rho(\xi) < \infty, \quad 0 < \delta < \infty, \quad (98)$$

and the further condition (95) reads

$$\int_0^\infty \xi^{-(\delta/2+1/4)} d\rho(\xi) < \infty, \quad 0 < \delta < \frac{1}{2} \quad (99)$$

and it can also be seen that (98) and (99) are then equivalent with

$$\int_0^\infty \lambda^{\delta-1} f(\lambda) d\lambda < \infty, \quad 0 < \delta < \infty \quad (100)$$

$$\int_0^\infty \lambda^{\delta/2-3/4} f(\lambda) d\lambda < \infty, \quad 0 < \delta < \frac{1}{2}, \quad (101)$$

respectively. We furthermore note that (86) reads

$$\tilde{g}(\mu) = \int_0^\infty \xi^{-\delta} e^{-\mu\xi^{-1}} d\rho(\xi) = \int_0^\infty e^{-\mu\xi} \xi^\delta d\rho(1/\xi) \quad (102)$$

so that this function is also completely monotone in $0 \leq \mu < \infty$ and we may state the following theorem.

Theorem 6. *If $f(\lambda)$ is completely monotone in $0 \leq \lambda < \infty$ and (100) and (101) holds, then the transform (81) exists and is completely monotone in $0 \leq \mu < \infty$.*

If moreover we have

$$f(\lambda) = O(\lambda^{-\alpha}), \quad g(\mu) = O(\mu^{-\beta}) \quad (103)$$

for large λ and μ , then our summation formula (80), and also (83) holds with both sides converging absolutely.

If $R(\lambda)$ and $S(\mu)$ are monotonely non-decreasing as they are for instance in the case of relation (76) with $a_m \geq 0, b_n \geq 0$, then assumption (103) may be omitted and formula (80) holds whether the value of any of its two sides is finite or $+\infty$.

The last part of the theorem follows from the fact that for $dR(\lambda) \geq 0, dS(\mu) \geq 0, d\rho(\xi) \geq 0$ the order of integrations in the repeated integrals occurring in

$$\int_0^\infty \int_0^\infty e^{-\lambda\xi} dR(\lambda) d\rho(\xi) = \int_0^\infty \int_0^\infty \xi^{-\delta} e^{-\mu\xi^{-1}} dS(\mu) d\rho(\xi) \quad (104)$$

may be always interchanged.

7.

Another situation arises if the set function is concentrated on a line

$$\xi = \epsilon > 0, \quad -\infty < \eta < \infty \quad (105)$$

which is parallel to the η -axis, that is if we have a representation

$$f(\lambda) = \int_{-\infty}^{\infty} e^{-\lambda(\epsilon+i\eta)} d\sigma(\eta) \quad (106)$$

and the reader will verify the following statements.

Theorem 7. *If a function $f(\lambda)$ in $0 \leq \lambda < \infty$ can be continued into the entire line $-\infty < \lambda < \infty$ in such a manner that for some $\epsilon > 0$ the function $f_\epsilon(\lambda) = e^{\epsilon\lambda}f(\lambda)$ has a representation*

$$f_\epsilon(\lambda) = \int_{-\infty}^{\infty} e^{-i\lambda\eta} d\sigma(\eta) \quad (107)$$

with

$$\int_{-\infty}^{\infty} |d\sigma(\eta)| < \infty \quad (108)$$

then the integral (81) by which $g(\mu)$ is defined is absolutely convergent, and if moreover

$$\left(\int_{-\infty}^{-1} + \int_{+1}^{\infty} \right) \exp \left(-\frac{\mu\epsilon}{\epsilon^2 + \eta^2} \right) |d\sigma(\eta)| = O(\mu^{-\beta}), \quad \mu \rightarrow \infty, \quad (109)$$

then the summation formula (80) holds with both sides converging absolutely.

This leads to the following corollary

Theorem 8. *The summation formula holds if for some $\epsilon > 0$ we have a representation*

$$e^{\epsilon\lambda}f(\lambda) = \int_{-\infty}^{\infty} e^{-i\lambda\eta} \tau(\eta) d\eta \quad (110)$$

with

$$\int_{-\infty}^{\infty} |\tau(\eta)| d\eta < \infty \quad (111)$$

and

$$\left(\int_{-\infty}^{-1} + \int_{+1}^{\infty} \right) \eta^{4\beta-2\delta+1} |\tau(\eta)|^2 d\eta < \infty. \quad (112)$$

In fact, in order to verify (109) we note that we have

$$\begin{aligned} \left(\int_1^{\infty} \eta^{-\delta} \exp \left(-\frac{\mu\epsilon}{\epsilon^2 + \eta^2} \right) |\tau(\eta)| d\eta \right)^2 &\leq \int_1^{\infty} \eta^{-4\beta-1} \exp \left(\frac{2\mu\epsilon}{\epsilon^2 + \eta^2} \right) d\eta \\ &\cdot \int_1^{\infty} \eta^{4\beta-2\delta+1} |\tau(\eta)|^2 d\eta, \end{aligned}$$

and that the first factor on the right is $O(\mu^{-2\beta})$ as $\mu \rightarrow \infty$; and similarly for the interval $(-\infty, -1)$.

Since

$$\left(\int_1^{\infty} |\tau(\eta)| d\eta \right)^2 \leq \int_1^{\infty} \eta^{-2} d\eta \int_1^{\infty} \eta^2 |\tau(\eta)|^2 d\eta, \quad (113)$$

then by the theory of Plancherel transforms, see [5], the following conclusion from Theorem 8 will be drawn.

Theorem 9. *If p is an integer such that*

$$p \geq \max(1, 2\beta - \delta + \frac{1}{2}); \quad (114)$$

and if $f(\lambda)$ in $0 \leq \lambda < \infty$ has $p-1$ continuous derivatives, and if $f^{(p-1)}(\lambda)$ is the indefinite integral of a function $f^{(p)}(\lambda)$ which latter function is square integrable in every finite interval; and if the $p+1$ functions $f(\lambda), f'(\lambda), \dots, f^{(p)}(\lambda)$ are $O(e^{-2\epsilon\lambda})$ as $\lambda \rightarrow +\infty$ for some $\epsilon > 0$; then the summation formula holds with both sides converging absolutely.

Proof. We can extend the function $f(\lambda)$ into $-\infty < \lambda < 0$ so that it vanish in a left half-line $\lambda \leq \lambda' < 0$ and preserve all the differentiability properties stipulated for the entire line $-\infty < \lambda < \infty$. The function

$$f_\epsilon(\lambda) = e^{\epsilon\lambda} f(\lambda) \quad (115)$$

will then have p derivatives $f_\epsilon^{(q)}(\lambda), q = 1, \dots, p$, having similar properties in $-\infty < \lambda < \infty$, and the $p+1$ functions $f_\epsilon, f'_\epsilon, \dots, f_\epsilon^{(p)}$ are each of integrable square in $(-\infty, \infty)$ and for the Plancherel transform

$$\tau(\eta) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\eta\lambda} f_\epsilon(\lambda) d\lambda \quad (116)$$

we then have, see [5],

$$\int_{-\infty}^{\infty} \eta^{2p} |\tau(\eta)|^2 d\eta < \infty, \quad (117)$$

and this implies (111) and (112), q.e.d.

For instance the Riesz kernel

$$f(\lambda) = \begin{cases} (1-\lambda)^\gamma & 0 < \lambda < 1 \\ 0 & 1 \leq \lambda < \infty \end{cases} \quad (118)$$

falls under Theorem 9 for $\gamma > p - \frac{1}{2}$, and since

$$\mu^{-\frac{1}{2}(\delta-1)} \int_0^1 J_{\delta-1}(2(\mu\lambda)^{1/2}) \lambda^{\frac{1}{2}(\delta-1)} (1-\lambda)^\gamma d\lambda = \Gamma(\gamma+1) \mu^{-\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2\mu^{1/2}), \quad (119)$$

we obtain for $0 < x < \infty$ the formula

$$\frac{1}{\Gamma(\gamma+1)} \int_0^x (x-\lambda)^\gamma dR(\lambda) = \int_0^\infty \left(\frac{x}{\mu}\right)^{\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2(x\mu)^{1/2}) dS(\mu) \quad (120)$$

for $\gamma > p - \frac{1}{2}$ where $p \geq 1, p \geq 2\beta - \delta + \frac{1}{2}$.

Now, due to (99) the right side in (120) is absolutely convergent for

$$\gamma \geq 2\beta - \delta - \frac{1}{2} \quad (121)$$

which is a smaller value, and if we differentiate both sides in (120) with regard to x a suitable number of times we obtain the following conclusion.

Theorem 10. *Relation (120) is valid if $\gamma \geq 2\beta - \delta - \frac{1}{2}$ and $\gamma > 0$ simultaneously.*

Also the restriction $\gamma > 0$ may be dropped for values of x in the neighborhood of which $R(\lambda)$ is constant or only differentiable sufficiently often.

The topic of Theorem 10 will be continued in Section IV and in the next section we will first assemble some remarks on subjects we will not discuss in detail.

Section III. Supplementary Remarks

8.

For fixed $\epsilon > 0, \rho > 0$, the function

$$e^{-\epsilon\lambda} J_{\delta-1}(2(\lambda\rho)^{1/2}) \lambda^{-\frac{1}{2}(\delta-1)} \quad (122)$$

if viewed as an $f(\lambda)$ falls under Theorem 9 and this leads to the rigorous relation

$$\int_0^\infty e^{-\epsilon\lambda} \lambda^{-\frac{1}{2}(\delta-1)} J_{\delta-1}(2(\lambda\rho)^{1/2}) dR(\lambda) = \int_0^\infty \mu^{-\frac{1}{2}(\delta-1)} \frac{1}{\epsilon} \exp\left(-\frac{\rho+\mu}{\epsilon}\right) I_{\delta-1}\left(\frac{2}{\epsilon}(\rho\mu)^{1/2}\right) dS(\mu) \quad (123)$$

in which $I_\nu(x)$ is the Bessel function with imaginary argument, see [17], p. 396.

But now start with a function $g(\rho)$ in $0 \leq \rho < \infty$, multiply both sides of (123) with it, integrate with respect to ρ and interchange integrations. We then obtain formally

$$\int f(\lambda) dR(\lambda) = \int g_\epsilon(\mu) dS(\mu) \quad (124)$$

where $f(\lambda)$ is defined by (82) and $g_\epsilon(\mu)$ by

$$g_\epsilon(\mu) = \frac{1}{\epsilon} \int_0^\infty \exp\left(-\frac{\rho+\mu}{\epsilon}\right) I_{\delta-1}\left(\frac{2}{\epsilon}(\rho\mu)^{1/2}\right) g(\rho) d\rho. \quad (125)$$

Now, the comparison of (125) with (80) suggests that we ought to have the relation

$$\lim_{\epsilon \rightarrow 0} g_\epsilon(\mu) = g(\mu) \quad (126)$$

in the sense that for a general function $g(\mu)$ the associated function $g_\epsilon(\mu)$ as defined by (125) is a kind of approximation to it and that relation (126) holds literally subject to criteria for $g(\mu)$. Such is indeed the case, see for instance [4], and on another occasion we will show that it is possible to strengthen the criteria so as to obtain the further relation

$$\lim_{\epsilon \rightarrow 0} \int_0^\infty g_\epsilon(\mu) dS(\mu) = \int_0^\infty g(\mu) dS(\mu) \quad (127)$$

with the right side converging absolutely, say. Now, whenever this takes place, (124) leads to

$$\lim_{\epsilon \rightarrow 0} \int_0^\infty e^{-\epsilon\lambda} f(\lambda) dR(\lambda) = \int_0^\infty g(\mu) dS(\mu), \quad (128)$$

and this is obviously (80) itself with the understanding that the left side in (80) shall be evaluated by a Gauss limiting process as indicated.

9.

In the ordinary theory of Fourier series and integrals see [5], the Gauss summability factor is $e^{-\epsilon x^2}$, and $e^{-\epsilon x}$ is the abelian one, but in our present normalization these factors are $e^{-\epsilon\lambda}$ and

$e^{-\epsilon\lambda^{1/2}}$ respectively. This square-root has been introduced into our set-up by the manner in which it appears in the kernels of the transformations (81) and (82), and it must be taken into account when comparing our present summation formula with the classical summation formula of Poisson. If applied to relation (76) our formula (80) reads

$$\sum_0^\infty a_m f(\lambda_m) = \sum_0^\infty b_n g(\mu_n) \quad (129)$$

and in the case of the Jacobi relation

$$\sum_0^\infty r_k(n) e^{-\pi x n} = x^{-k/2} \sum_0^\infty r_k(n) \exp\left(-\frac{\pi n}{x}\right) \quad (130)$$

in which $r_k(n)$ is the number of representations $n = n_1^2 + \dots + n_k^2$, with $r_k(0) = 1$, (129) is

$$\sum_0^\infty r_k(n) f(\pi n) = \sum_0^\infty r_k(n) g(\pi n). \quad (131)$$

On the other hand if we take any function $F(x_1, \dots, x_k)$ with period 2π in each variable and put

$$G(y_1, \dots, y_k) = \int_{E_k} F(x_1, \dots, x_k) \exp(i(y_1 x_1 + \dots + y_k x_k)) dx_1 \dots dx_k, \quad (132)$$

then we obtain the Poisson summation formula

$$\sum_{n_1 \dots n_k} G(2\pi n_1, \dots, 2\pi n_k) = \sum_{m_1 \dots m_k} F(m_1, \dots, m_k). \quad (133)$$

If $F(x_1, \dots, x_k)$ is a “radial” function, see [2], p. 186,

$$F(x_1, \dots, x_k) = F((x_1^2 + \dots + x_k^2)^{1/2}) \quad (134)$$

then we similarly have

$$G(y_1, \dots, y_k) = G((y_1^2 + \dots + y_k^2)^{1/2}) \quad (135)$$

where, on putting $\delta = k/2$, the connection is

$$G(y) y^{\delta-1} = (2\pi)^\delta \int_0^\infty J_{\delta-1}(yx) x^\delta F(x) dx, \quad (136)$$

and (133) becomes

$$\sum_{m=0}^\infty r_k(m) G(2\pi m^{1/2}) = \sum_{n=0}^\infty r_k(n) F(n^{1/2}). \quad (137)$$

Since (136) implies

$$G(2(\pi y)^{1/2}) y^{\frac{1}{2}(\delta-1)} = \int_0^\infty J_{\delta-1}(2(yx)^{1/2}) x^{\frac{1}{2}(\delta-1)} F\left(\left(\frac{x}{\pi}\right)^{1/2}\right) dx, \quad (138)$$

we obtain (81) by putting

$$f(\lambda) = F\left(\left(\frac{\lambda}{\pi}\right)^{1/2}\right), \quad g(\mu) = G(2(\pi\mu)^{1/2})$$

and this is the origin of the square-root appearing subsequently.

We also note that for the same reason the summability factor $(x - \lambda)^\gamma$ in (120) does not correspond to the “ordinary” Fejér-Riesz factor as such but to the factor $(R^2 - \nu^2)^\gamma$ of “spherical summability”, see [3], which is also tellingly indicated by the presence of a Bessel function proper on the other side of (120).

Now, although spherical summability has usually the advantage of greater technical elegance over the ordinary one, yet in the theory of modular functions there is at least one very important occasion known to us on which spherical summability apparently fails where the ordinary one succeeds. In the proof of H. Hamburger, [9], for the unique determination of the Riemann Zeta function by its functional equation, use is made of a relation of the form (120) for $\gamma = 1$, but it is its “ordinary” counterpart, and we are unable to substitute in the reasoning formula (120) proper in its stead. Such a substitution would be of greatest importance, since it might open the way for obtaining uniqueness theorem for Zeta functions other than the very obvious ones.

Finally we note that the matrix trace in (20) and (21) is a quadratic form of the square roots of the eigenvalues of the matrix and that therefore these formulas collect the terms by “spheres” and not by “diagonals”, first impressions notwithstanding.

10.

We will now make some remarks on the functional equation

$$\Delta(\delta - s)\psi(\delta - s) = \Delta(s)\varphi(s) \quad (139)$$

in which $\varphi(s)$, $\psi(s)$ are Dirichlet series (49), but the factor $\Delta(s)$ is not the Gamma function itself, but some other function suitably circumscribed the most notable example of which is the Hecke factor

$$\Delta(s) = \Gamma\left(\frac{s}{2}\right)^{r_1} \Gamma(s)^{r_2} \quad (140)$$

where $r_2 + 2r_1$ is the degree of an algebraic number field. If $\Delta(s)$ is holomorphic in the right half plane $\sigma > 0$, and $\Delta(\sigma + it)$ belongs to $L_1(-\infty, \infty)$ uniformly in $0 < \sigma_1 \leq \sigma \leq \sigma_2$, then we can set up for $x > 0$ the Mellin transform

$$E(x) = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} \Delta(s)x^{-s} ds, \quad a > 0 \quad (141)$$

and we first of all have

$$E(x) = O(x^{-\epsilon}), \quad x \rightarrow 0, \quad (142)$$

for every $\epsilon > 0$. For (140), or more generally for

$$\Delta(s) = \prod_{j=1}^n \Gamma(p_j s + q_j) \quad (143)$$

where

$$1 \geq p_j > 0, \quad \operatorname{Re}(q_j) \geq 0 \quad (144)$$

we also have

$$E(x) = O(e^{-cx}), \quad x \rightarrow \infty \quad (145)$$

for some suitable $c > 0$; and in fact, we have

$$E(\xi + i\eta) \leq Me^{-c\xi}, \quad \xi \rightarrow \infty. \quad (146)$$

For a simple factor $\Delta(s) = \Gamma(ps + q)$ this is obviously so, since we then have

$$E(x) = \frac{1}{p} x^{q/p} e^{-x^{1/p}} \quad (147)$$

and for a product (143) this follows then by successively applying the composition rule

$$\frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} \Delta_1(s) \Delta_2(s) x^{-s} ds = \int_0^\infty E_1(u) E_2\left(\frac{x}{u}\right) \frac{du}{u} \quad (148)$$

Now, if we have (2) and (3), and also (145), we can again obtain from (139) the relation

$$\sum_1^\infty a_m E(\lambda_m x) = x^{-\delta} \sum_1^\infty b_n E\left(\frac{\mu_n}{x}\right) + P(x), \quad (149)$$

where $P(x)$ is a residual function as before, and the two series in (149) converge absolutely for $x > 0$, and under assumption (146) also for $\xi > 0$. We can again re-write (149) into the “unified” relation

$$\int_0^\infty E(\lambda x) dR(\lambda) = x^{-\delta} \int_0^\infty E\left(\frac{\mu}{x}\right) dS(\mu), \quad (150)$$

but the very simple relation (76) is usually no longer available, not even for the Hecke factor (140), since $E(x)$ usually does not have a (finite) limit for $x \rightarrow 0$.

On the other hand, (141) can be always inverted to

$$\Delta(s) = \int_0^\infty E(x) x^{s-1} dx \quad (151)$$

and consequently it is always possible to proceed from a relation (149) to an equation (139) very much under the same kind of conditions as before.

The relation (150) as such has apparently not been investigated systematically in the literature, if it has been previously introduced at all. However (150) implies the summation formula

$$\int_0^\infty f(\lambda) dR(\lambda) = \int_0^\infty g(\mu) dS(\mu) \quad (152)$$

with

$$g(\mu) = \mu^{-\frac{1}{2}(\delta-1)} \int_0^\infty K(\lambda\mu) f(\lambda) d\lambda, \quad (153)$$

and self-reciprocally, where

$$K(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Delta\left(\frac{\delta-1}{2} + s\right) \Delta\left(\frac{\delta+1}{2} - s\right) x^{-s} ds \quad (154)$$

and instances of the formula (152), in particular for $f(\lambda)$ being the Riesz kernel (118), do occur in considerable number. One of the most important papers in this direction, one which has not been superseded by other important papers dealing either with the summation formulas as such or with the nature of the kernels (154) employed therein, is the thesis of Walfisz [15] as supplemented by his later paper [16], and a further study of the topics introduced in [15] would be indicated.

Also, if we start from (150) then (152) can be fulfilled by putting

$$f(\lambda) = \int_0^\infty E(\lambda\xi) d\rho(\xi), \quad g(\mu) = \int_0^\infty \xi^{-\delta} E\left(\frac{\mu}{\xi}\right) d\rho(\xi) = \int_0^\infty E(\mu\xi) \xi^\delta d\rho\left(\frac{1}{\xi}\right) \quad (155)$$

and this suggests the very interesting problem of describing the class of functions $f(\lambda)$ which can be represented by an integral (155) with $E(x)$ corresponding to the Hecke factor (140) say. The companion problem of satisfying (152) by putting

$$f(\lambda) = \int_{-\infty}^\infty E(\lambda(\epsilon + i\eta)) \tau(\eta) d\eta, \quad g(\mu) = \int_{-\infty}^\infty (\epsilon + i\eta)^{-\delta} E\left(\frac{\mu}{\epsilon + i\eta}\right) \tau(\eta) d\eta$$

might also be worth studying, but this problem is less likely to lead to polished results, in our opinion.

Section IV. Modular Exponential Sums

11.

If we substitute the known identity, see [2], p. 57,

$$\exp(-2x\lambda^{1/2}) = \frac{x}{\pi^{1/2}} \int_0^\infty y^{-3/2} \exp\left(-\frac{x^2}{y}\right) e^{-\lambda y} dy \quad (156)$$

into

$$\int_0^\infty \exp(-2x\lambda^{1/2}) dR(\lambda) \quad (157)$$

and interchange integrations, and if our modular relation

$$\int_0^\infty e^{-\lambda y} dR(\lambda) = y^{-\delta} \int_0^\infty \exp\left(-\frac{\mu}{y}\right) dS(\mu) \quad (158)$$

holds, then (157) assumes the value

$$\frac{x}{\pi^{1/2}} \int_0^\infty dy \exp\left(-\frac{x^2}{y}\right) y^{-3/2-\delta} \int_0^\infty \exp\left(-\frac{\mu}{y}\right) dS(\mu)$$

$$= \frac{x}{\pi^{1/2}} \int_0^\infty dy \int_0^\infty y^{-3/2-\delta} \exp\left(-\frac{x^2+\mu}{y}\right) dS(\mu);$$

and thus (158) implies the relation

$$\int_0^\infty \exp(-2x\lambda^{1/2}) dR(\lambda) = \frac{\Gamma(\delta + \frac{1}{2})}{\pi^{1/2}} \int_0^\infty \frac{x dS(\mu)}{(x^2 + \mu)^{\delta + \frac{1}{2}}}, \quad (159)$$

and under assumptions (78), (79) this is rigorously valid for $\text{Re}(x) > 0$ provided

$$\beta \leq \delta + \frac{1}{2}. \quad (160)$$

On putting

$$F(x; \lambda) = \pi^{-1/2} x^{-1} \exp(-2x\lambda^{1/2}) \quad (161)$$

$$G(x; \mu) = \Gamma(\delta + \frac{1}{2}) (x^2 + \mu)^{-\delta - \frac{1}{2}} \quad (162)$$

(159) implies for $A > 0$ and any integer $p > 0$ the relation

$$\int_0^\infty \sum_{q=0}^p (-1)^q F((x^2 + qA)^{1/2}; \lambda) dR(\lambda) = \int_0^\infty \sum_{q=0}^p (-1)^q G(x^2 + qA; \mu) dS(\mu) \quad (163)$$

but for the latter we may state as follows.

Lemma 2. *Relation (163) holds in $0 < x < \infty$ for*

$$\beta \leq \delta + \frac{1}{2} + p \quad (164)$$

with both sides converging absolutely.

Proof. The absolute convergence follows from (78), (79). If we substitute (156) into the left side of (163) we obtain for the latter

$$\int_{\lambda=0}^\infty \left(\int_{y=0}^\infty y^{-3/2} e^{-x^2 y^{-1}} (1 - e^{-Ay^{-1}})^p e^{-\lambda y} dy \right) dR(\lambda),$$

and, on account of the positiveness of the integrand, if we substitute (158), this becomes

$$\int_{y=0}^\infty dy \left(\int_{\mu=0}^\infty y^{-\delta-3/2} \exp\left(-\frac{x^2+\mu}{y}\right) (1 - e^{-Ay^{-1}})^p dS(\mu) \right),$$

and, for (164), we may again interchange integrations which produce the right side in (163), q.e.d.

Theorem 11. *For $\mu^0 > 0$, if in an interval $|\mu - \mu^0| < \delta$ the function $S(\mu)$ is a constant, (or more generally is there analytic), then for*

$$t = \pm(\mu^0)^{1/2} \quad (165)$$

the integral

$$\int_1^\infty \lambda^l e^{-2i\lambda^{1/2}t} dR(\lambda) \quad (166)$$

exists for every $l \leq 0$ as the Abel limit

$$\lim_{\epsilon \rightarrow 0} \int_1^\infty e^{-\epsilon \lambda^{1/2}} \lambda^l e^{-2i\lambda^{1/2}t} dR(\lambda), \quad (167)$$

and for a neighborhood

$$|t^2 - \mu^0| < \delta - \eta \quad (168)$$

it exists uniformly in t .

Also, for the same values of t , and again uniformly, the Abel limit

$$\lim_{\epsilon \rightarrow 0} \int_0^\infty e^{-\epsilon \lambda^{1/2}} \left(\frac{t^2}{\lambda} \right)^{\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2t\lambda^{1/2}) dR(\lambda) \quad (169)$$

exists for $\gamma \geq 0$, and its value is

$$\frac{1}{\Gamma(\gamma+1)} \int_0^{t^2} (t^2 - \mu)^\gamma dS(\mu) \quad (170)$$

except for

$$\gamma = -1, -2, -3, \dots, \quad (171)$$

for which the limit has value 0.

Thus, in particular, in the case of a modular relation

$$\sum_0^\infty a_m e^{-\lambda_m x} = x^{-\delta} \sum_{n=0}^\infty b_n e^{-\mu_n x^{-1}} \quad (172)$$

with (2), (3), (44), (46), the series

$$\sum_1^\infty a_m \lambda_m^l e^{-2i\lambda_m^{1/2}t}, \quad l \leq 0 \quad (173)$$

$$\sum_0^\infty a_m \left(\frac{t^2}{\lambda_m} \right)^{\frac{1}{2}(\delta+\gamma)} J_{\delta+\gamma}(2t\lambda_m^{1/2}), \quad \gamma \geq 0 \quad (174)$$

are Abel summable for $t^2 > 0$, if $t^2 \neq \mu_1, \mu_2, \mu_3, \dots$, the second series having the value previously stated.

Proof. For $p = 0$, if we put $x = \epsilon + it$ into the right side of (159), then uniformly in (168) a limit exists for $\epsilon \rightarrow 0$, and this established the corresponding limit of (166) for $l = 0$. For $p > 0$, if we choose $A = \mu^0 + \delta$ then for $q > 0$ we have

$$\operatorname{Re}\{x^2 + qA\} \geq \operatorname{Re}\{x^2 + A\} = \epsilon^2 - t^2 + A, \quad (175)$$

and it follows that the right side of (163) has a limit as before so that the left side of (163) also has one. We now split the left side into the principal term

$$\int_0^\infty F(x; \lambda) dR(\lambda) \quad (176)$$

and the secondary term

$$\int_0^\infty \sum_{q=1}^p (-1)^q F(x^2 + qA; \lambda) dR(\lambda) \quad (177)$$

and we notice that (177) is absolutely convergent uniformly for small ϵ and t in (168). Therefore (176) itself must have a limit, as $\epsilon \rightarrow 0$, uniformly in (168) which proves our assertion about (166) for $l = 0$.

Our next step will be to discuss it for $l = k/2, k = 1, 2, 3, \dots$, by induction on k . If we differentiate both sides in (163), term by term, k times with respect to x , then the right side retains the previously stated limit behavior, and so does the secondary part (177). But the principal part (176) constitutes a term

$$(-2i)^k \int_0^\infty \lambda^{k/2} F(x; \lambda) dR(\lambda) \quad (178)$$

and similar terms involving $\lambda^{h/2} F(x; \lambda)$ for $h \leq k - 1$, which justifies the induction.

Finally, for a general real l , whether positive or negative, we put $l = k/2 - \alpha/2$ for some integer k and $\alpha > 0$, and on setting

$$\lambda = \xi^2, \quad d_\xi A(t, \xi) = e^{-2it\xi} dR(\xi^2)$$

we easily obtain the desired conclusion from the following lemma.

Lemma 3. *If*

$$\int_1^\infty e^{-x\xi} |d_\xi A(t, \xi)| \leq M(\epsilon) e^{-\epsilon}, \quad x \geq \epsilon, \quad (179)$$

uniformly in t , and if the function

$$L(t, x) = \int_1^\infty e^{-x\xi} d_\xi A(t, \xi) \quad (180)$$

has a limit $L(t)$ as $x \downarrow 0$ uniformly in t , then for $\rho > 0$ the function

$$L_\rho(t, x) = \int_1^\infty \xi^{-\rho} e^{-x\xi} d_\xi A(t, \xi) \quad (181)$$

has a limit of the same type.

Proof of Lemma 3. For fixed t and x we may substitute in (181)

$$\xi^{-\rho} = \frac{1}{\Gamma(\rho)} \int_0^\infty e^{-\xi\eta} \eta^{\rho-1} d\eta$$

and we obtain

$$L_\rho(t, x) = \frac{1}{\Gamma(\rho)} \int_0^\infty y^{\rho-1} L(t, x+y) dy.$$

If we split the last integral into $\int_0^1 y^{\rho-1} L(t, x+y) dy + \int_1^\infty$ then the first part converges, uniformly in t , as $x \downarrow 0$, in consequence of $L(t, x)$ so converging, whereas the second part

so converges in consequence of inequality (179); which finally proves our assertion regarding (166).

In turning to (169) we note that of the two assumptions (78), (79) until now only the second has been exploited fully, but that at present the first will come into its own. For

$$x = 2t\lambda^{1/2} \geq 1$$

and a fixed $\gamma \geq 0$ we introduce in (169) the asymptotic expansion

$$J_{\delta+\gamma}(x) = \sum_{\nu=0}^M \frac{a_{\nu}e^{ix} + b_{\nu}e^{-ix}}{x^{\nu+1/2}} + O\left(\frac{1}{x^{M+3/2}}\right), \quad (182)$$

and on the basis of (78) the contribution due to the error term is absolutely convergent for M sufficiently large. However the other terms of (182) lead to expressions of the form (167), and the uniform existence of (169) follows. As regards the value given for (169) we recall that for γ sufficiently large we have obtained it in Theorem 10, and for other values of γ it may be obtained by differentiation with respect to t , which completes the proof of Theorem 11.

12.

We are now going to quote an important theorem from Walfisz [16], his theorems A and A_1 , in a trivial extension from Dirichlet series to LaplaceStieltjes integrals.

Lemma 4. *If the integral*

$$f(\epsilon) = \int_1^{\infty} e^{-\epsilon\xi} dA(\xi) \quad (183)$$

is absolutely convergent for $\epsilon > 0$; if

$$\lim_{\epsilon \rightarrow 0} f(\epsilon) \text{ exists} \quad (184)$$

and if for $\tau > 0$

$$\int_1^{\infty} \xi^{-\tau} |dA(\xi)| < \infty \quad (185)$$

then the Riesz mean

$$A_{\tau}(x) = \frac{x^{-\tau}}{\Gamma(\tau+1)} \int_0^x (x-\xi)^{\tau} dA(\xi) \quad (186)$$

has a limit for $x \rightarrow \infty$.

If the function $A(\xi) = A(\xi, t)$ depends on a parameter t ; if (184) holds uniformly in t , and if the variation of $A(\xi, t)$ with regards to ξ is majorized by a function $B(\xi)$ for which

$$\int_1^{\infty} \xi^{-\tau} dB(\xi) < \infty \quad (187)$$

holds, then the limit of (186) exists uniformly in t .

Now, if we put

$$A(\xi, t) = \int_{\lambda=1}^{\xi^2} \lambda^l e^{-2it\lambda^{1/2}} dR(\lambda) \quad (188)$$

then we may put

$$B(\xi) = \int_{\lambda=1}^{\xi^2} \lambda^l |dR(\lambda)|,$$

and (78) implies (187) with

$$\tau = 2\alpha + 2l \quad (189)$$

and if we apply Hardy's equivalence of $(R; \lambda^{1/2})$ with $(R; \lambda)$ we obtain the following theorem.

Theorem 12. *If (78) holds, then for $l > -\alpha$ the integral (166) is summable $(R; \lambda; 2l + 2\alpha)$, that is the limit*

$$\lim_{x \rightarrow \infty} \{\Gamma(2l + 2\alpha + 1)\}^{-1} \int_1^x \left(1 - \frac{\lambda}{x}\right)^{2l+2\alpha} \lambda^l e^{-2it\lambda^{1/2}} dR(\lambda) \quad (190)$$

exists, and equals (167) uniformly in (168).

We note that for $l < -\alpha$ assumption (78) makes the integral (166) absolutely convergent so that Theorem 12 covers all other possibilities $l > -\alpha$, by securing for them a Riesz summability at least. But no intermediary interval of ordinary (non-absolute) convergence is provided for, and perhaps in general none is available, however for modular relations of the classical structure or nearly so the following theorem of AnandaRau [1] opens up a possibility.

Lemma 5. *If $R_0(\lambda)$ in $1 \leq \lambda < \infty$ is continuous from the right; if for $\sigma > 0$ we have*

$$\int_{\lambda}^{\lambda+\xi} |dR_0(\lambda)| \leq K \xi^{\sigma-1} \lambda \quad (191)$$

in

$$\lambda \leq \xi \leq H\lambda \quad (192)$$

and if for some $l \leq 0$ and some $c \geq 0$ the integral

$$\int_1^{\infty} \lambda^{l+c} dR_0(\lambda) \quad (193)$$

is summable $(R; \lambda; b)$ for some $b > 0$; then for $0 \leq \eta < b$ the integral

$$\int_1^{\infty} \lambda^l dR_0(\lambda) \quad (194)$$

is summable $(R; \lambda; \eta)$ for

$$\eta > \frac{(\sigma + l)b - c}{\sigma + l + c}. \quad (195)$$

In our case it implies the following theorem.

Theorem 13. *If for some $\sigma > 0$ we have*

$$\int_{\lambda}^{\lambda+\xi} |dR(\lambda)| \leq K \xi^{\sigma-1} \lambda, \quad 1 \leq \lambda < \infty \quad (196)$$

for (192), then the integral (166) is $(R; \lambda, \eta)$ if simultaneously $\eta \geq 0$ and $\eta > 2l + 2\sigma - 1$.

In particular, the integral

$$\int_1^\infty \frac{e^{-2it\lambda^{1/2}}}{\lambda^{\sigma-1/2+\epsilon}} dR(\lambda) \quad (197)$$

is convergent for $\epsilon > 0$.

Proof. For $l + c$ instead of l , our integral is summable $(R; \lambda; 2\sigma + 2l + 2c)$ by Theorem 12. Now, if we put $b = 2\sigma + 2l + 2c$ in (195) we obtain

$$\eta > \frac{(\sigma + l)(2\sigma + 2l + 2c) - c}{\sigma + l + c} \quad (198)$$

and on the basis of Lemma 5 we may claim that for fixed l our integral is summable $(R; \lambda; \eta)$ with $\eta \geq 0$, provided there exists some positive c for which (198) holds. Now, on letting $c \rightarrow \infty$, the right side in (198) converges to $2\sigma + 2l - 1$, and thus $\eta > 2\sigma + 2l - 1$ is a sufficient condition, which proves our theorem.

Remarks. Assumption (196) implies

$$\int_1^\infty \lambda^{-\sigma-\epsilon} |dR(\lambda)| < \infty$$

for $\epsilon > 0$, and this, with (78) known, it would have been reasonable to assume $\sigma \geq \alpha$ and not only $\sigma > 0$; for a similar reason the second half of the theorem is of interest only for $\sigma \leq \alpha + \frac{1}{2}$.

In the lattice point relation (130) our assumptions are fulfilled for $\alpha = k/2 + \epsilon$, $\sigma = \alpha$, so that Theorem 13 gives the result that the series

$$\sum_1^\infty \frac{r_k(n)}{n^{(k-1)/2+\epsilon}} e^{-it(\pi n)^{1/2}}$$

converges for $t \neq (\pi m)^{1/2}$, for $\epsilon > 0$, and this tallies with a best result known for this case, see [7] and [6].

Addendum

Remark on Hecke's Functional Equation for Zeta Functions in Algebraic Fields (Received June 8, 1950)

We note that the non-arithmetical part of Hecke's reasoning in his paper: *Eine neue Art von Zetafunktionen und ihre Beziehungen zur Verteilung der Primzahlen, Zweite Mitteilung*, Mathematische Zeitschrift 6 (1920), 1151, can be formalized and technically generalized in the following manner [compare also: Ch. H. Müntz, Hamburgische Abh. 3 (1924), 111, and L. J. Mordell, Proc. London Math. Soc. (2) 3 (1931), 501506].

Consider in k -space a real affine transformation

$$Z_\mu = a_{\mu 1} z_1 + a_{\mu 2} z_2 + \cdots + a_{\mu k} z_k, \quad \mu = 1, \dots, k,$$

with determinant 1,

$$|a_{\mu\nu}|_{\mu,\nu=1,\dots,k} = 1,$$

and for a fixed decomposition

$$k = p_1 + p_2 + \dots + p_{r+1}, \quad 1 \leq r+1 \leq k,$$

into any integers p_j , (not necessarily = 1 or 2 only),

$$1 \leq p_j \leq k, \quad j = 1, \dots, r+1,$$

we introduce the $r+1$ semi-definite forms

$$A_1(z) = Z_1^2 + \dots + Z_{p_1}^2, \quad A_2(z) = Z_{p_1+1}^2 + \dots + Z_{p_2}^2, \dots, A_{r+1}(z) = Z_{p_r+1}^2 + \dots + Z_{p_{r+1}}^2.$$

If we also introduce the inverse transformation

$$z_\nu = b_{\nu 1} Z_1 + \dots + b_{\nu k} Z_k,$$

$$\sum_j b_{\nu j} a_{j\mu} = \delta_{\nu\mu},$$

and the corresponding semi-definite forms

$$B_1(Z) = z_1^2 + \dots + z_{p_1}^2, \quad B_2(Z) = z_{p_1+1}^2 + \dots + z_{p_2}^2, \dots,$$

then the classical theta formula implies the relation

$$\sum_m \exp \left(-\pi \sum_{j=1}^{r+1} A_j(m) t_j \right) = \prod_{j=1}^{r+1} t_j^{-p_j/2} \sum_n \exp \left(-\pi \sum_{j=1}^{r+1} B_j(n) t_j^{-1} \right)$$

for variables $t_1 > 0, \dots, t_{r+1} > 0$, the summation extending over all lattice points $m = (m_1, \dots, m_k)$ and n respectively. This relation leads only to what corresponds to Zeta functions for the trivial character $\chi = 1$, and in order to include more general characters it is necessary to envisage more general relations, such as exist, of the form

$$\sum_m \varphi_m \exp \left(-\pi \sum_j A_j(m) t_j \right) = \prod_j t_j^{p_j s - q_j} \sum_n \psi_n \exp \left(-\pi \sum_j B_j(n) t_j^{-1} \right)$$

with suitable constant coefficients $\{\varphi_m\}$, $\{\psi_n\}$ and additional exponents q_j for which $\text{Re } q_j \geq 0$. For brevity we denote the left side by $\theta(t; A; \varphi)$ and the relation then reads

$$\theta(t; A; \varphi) = \prod_j t_j^{-\frac{1}{2} p_j - q_j} \theta \left(\frac{1}{t}; B; \psi \right). \quad (199)$$

We assume that for $m \neq (0, 0, \dots, 0)$ all $A_j(m) > 0$, and that the same holds for $B_j(n)$, and we introduce the series

$$\zeta(s; A; \varphi) = \sum_m'' \varphi_m \prod_{j=1}^{r+1} (\pi A_j(m))^{-\frac{1}{2}(p_j s + q_j)} \quad (200)$$

$$\theta''(t; A; \varphi) = \sum_m'' \varphi_m \exp \left(-\pi \sum_j A_j(m) t_j \right) \quad (201)$$

in which the summation extends only over a certain fixed subset of all lattice points (not including the origin) whose precise nature will be described later. From

$$\Gamma \left(\frac{p}{2}x + \frac{q}{2} \right) = \int_0^\infty e^{-t} t^{\frac{1}{2}(px+q)-1} dt$$

we then obtain

$$\Delta(s) \zeta(s; A; \varphi) = \int_0^\infty \cdots \int_0^\infty \theta''(t; A; \varphi) \prod_j t_j^{\frac{1}{2}(p_j s + q_j) - 1} dt_j \quad (202)$$

$$\Delta(s) \zeta(s; B; \psi) = \int_0^\infty \cdots \int_0^\infty \theta''(t; B; \psi) \prod_j t_j^{\frac{1}{2}(p_j s + q_j) - 1} dt_j \quad (203)$$

where

$$\Delta(s) = \prod_{j=1}^{r+1} \Gamma \left(\frac{p_j}{2}s + \frac{q_j}{2} \right),$$

whatever the subset of the lattice points may be.

We now substitute for the variables $t_1, \dots, t_{r+1}$ new variables $v; y_1, \dots, y_r$ by means of the transformation

$$t_j = \exp \left(v + \sum_{\rho=1}^r \alpha_{j\rho} y_\rho \right), \quad j = 1, \dots, r+1,$$

with suitable constants $\{\alpha_{j\rho}\}$ which will be subjected, axiomatically, to rather heavy assumptions expressive of the behavior of units in algebraic fields. We first of all assume that we have

$$\sum_{j=1}^{r+1} p_j \alpha_{j\rho} = 0, \quad \rho = 1, \dots, r. \quad (204)$$

and that the determinant

$$R = |1, \alpha_{j1}, \alpha_{j2}, \dots, \alpha_{jr}| \quad j = 1, \dots, r+1$$

is $\neq 0$, and in consequence of (204), if we put

$$q = \sum_{j=1}^{r+1} q_j, \quad \alpha_\rho = \sum_{j=1}^{r+1} q_j \alpha_{j\rho},$$

we obtain

$$\sum_{j=1}^{r+1} A_j t_j = e^v \sum_{j=1}^{r+1} A_j e^{\sum_\rho \alpha_{j\rho} y_\rho}$$

and

$$\prod_{j=1}^{r+1} t_j^{\frac{1}{2}(p_j s + q_j) - 1} dt_j = e^{v/2(ks+q)} dv \prod_{\rho=1}^r e^{\frac{1}{2}\alpha_\rho y_\rho} dy_\rho.$$

We now put

$$y_\rho = g_\rho + \eta_\rho, \quad \rho = 1, \dots, r$$

where

$$-\frac{1}{2} \leq \eta_\rho < \frac{1}{2}, \quad \rho = 1, \dots, r \quad (205)$$

and

$$g_\rho = 0, \pm 1, \pm 2, \pm 3, \dots,$$

and if we introduce the expression

$$\theta'(t; A; \varphi) = \sum_g \theta''(t; A_j e^{\sum \alpha_{j\rho} g_\rho}; \varphi e^{\frac{1}{2} \sum \alpha_\rho g_\rho})$$

with summation over all lattice point $(g_1, \dots, g_r)$ then the right side of (202) is

$$R \int_{-\infty}^{\infty} dv e^{v/2(ks+q)} \int_I \theta'(e^{v+\sum \alpha_{j\rho} \eta_\rho}; A; \varphi) e^{\frac{1}{2} \sum \alpha_\rho \eta_\rho} d\eta \quad (206)$$

where $d\eta = d\eta_1 \dots d\eta_r$ and I is the cube (205).

The decisive assumption now is that for a certain constant w we have

$$w\theta'(t; A; \varphi) = \theta(t; A; \varphi) - \varphi_0 \quad (207)$$

$$w\theta'(t; B; \psi) = \theta(t; B; \psi) - \psi_0 \quad (208)$$

and Hecke satisfies it by the following arithmetical data. The constant w is an integer, (it is the number of roots of unity in the field), and, except for a multiplicity w , the multitude of all vectors $\{A_j(m)\}$, $j = 1, \dots, r+1$, in conjunction with the coefficients $\{\varphi(m)\}$, as taken over all lattice points $\{m\}$ not identically zero, is the same as the multitude of quantities

$$\{A_j(m'') e^{\sum \alpha_{j\rho} g_\rho}\}, \quad \{\varphi(m'') e^{\frac{1}{2} \sum \alpha_\rho g_\rho}\}$$

as taken, independently, over all integers $\{g_\rho\}$ and the elements $\{m''\}$ of a certain subset of the set of all lattice points $\{m\}$, and similarly for B, ψ , the subset being precisely the one over which the summation in (200), (201) was meant to be taken.

Once relations (207), (208) are secured we obtain the functional equation

$$\Delta(s)\zeta(s; A; \varphi) = \Delta(1-s)\zeta(1-s; B; \psi)$$

by the familiar reasoning of decomposing in (206) the integration with respect to v into $(-\infty, 0)$ and $(0, \infty)$; substituting from (207); replacing in one of the half-line $\theta(t; A; \varphi)$ by $\theta(t^{-1}; B; \psi)$ from (199); and then again substituting from (208). See Hecke loc. cit., and also E. Landau, *Ueber Ideale und Primideale in Idealklassen*, Mathematische Zeitschrift 2 (1918), 52154.

PRINCETON UNIVERSITY

REFERENCES

1. K. Ananda-Rau, On the convergence and summability of Dirichlet series. Proc. London Math. Soc. (2) 34 (1932), 41440.
2. Bochner, S., Vorlesungen ueber Fouriersche Integrale. 1937, 1948.
3. Bochner, S., Spherical summability of multiple Fourier series. Trans. Amer. Math. Soc. 40 (1936), 175207; also Die Poissonsche Summationsformel in mehreren Veraenderlichen. Math. Annalen 106 (1932), 5663.
4. Bochner, S., and Chandrasekharan, K., On the localization property for multiple Fourier series. Ann. of Math., 49 (1948), 966978.
5. Bochner, S., and Chandrasekharan, K., Fourier transforms. 1949.
6. Bochner, S. and Chandrasekharan, K., Summation over lattice points in k -space. Quart. J. Math. Oxford Ser. 19 (1948), 238248.
7. A. L. Dixon and W. L. Ferrar, Some summations over the lattice points of a circle (I), (II). Quart. J. Math. Oxford Ser., 5 (1934), 4663; 172185.
8. M. L. Ferrar, Summation formulae and their relation to Dirichlet series I, II. Compositio Math., 1 (1934), 344360; 4 (1936), 396405.
9. H. Hamburger, Ueber die Riemannsche Funktionalgleichung der ζ -Funktion I, II, III; Math. Zeit. 10 (1921), 240254; 11 (1921), 224245; 13 (1922), 283311; also: Ueber einige Beziehungen die mit der Funktionalgleichung der Riemannschen ζ -Funktion aequivalent sind, Math. Ann. 85 (1922), 129140; also E. Hecke, Ueber die Loesungen der Riemannschen Funktionalgleichung, Math. Zeit. 16 (1923), 301307.
10. E. Hecke, Ueber die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung, Math. Ann. 112 (1936), 664699; also H. Rademacher, Trends in research, the analytic number theory. Bull. Amer. Math. Soc. 48 (1942), p. 393.
11. N. S. Koshliakov, Application of the theory of sumformulae to the investigation of a class of one-valued analytical functions in the theory of numbers. Messenger of Mathematics, 58 (1929), 123.
12. H. Maass, Ueber eine neue Art von nicht-analytischen automorphen Funktionen und die Bestimmung Dirichletscher Reihen durch Funktionalgleichungen. Math. Ann. 121 (1949), 141183; also, Automorphe Funktionen von mehreren Veraenderlichen und Dirichletsche Reihen. Abh. Math. Sem. Hansischen Univ. 16 (1949), 72100.
13. C. L. Siegel, Einfuehrung in die Theorie der Modulfunktionen n -ten Grades. Math. Ann. 116 (1939), 617657.
14. G. Voronoï, Sur une fonction transcendante et ses application a la sommation de quelques series. Ann. École Norm., (3) 21 (1904), 207267, 459533.
15. A. Walfisz, Ueber die Summatorischen Funktionen einiger Dirichletscher Reihen. Inaugural dissertation, Göttingen, 1922.

16. A. Walfisz, Ueber Summabilitätssätze von Marcel Riesz. Math. Ann. 93 (1924), 130148.
17. G. N. Watson, Bessel functions. 1922, 1944.
18. J. R. Wilton, A note on Ramanujan's arithmetical function $\tau(n)$. Proc. Cambridge Philos. Soc. 25 (1929), 1219; and G. H. Hardy, A further note on Ramanujan's arithmetical function $\tau(n)$. Ibid., 34 (1938), 309315.