

# Band-Limitedness of the Hardy Z-Function in the Phase Variable

Stephen Crowley

April 19, 2026

## Abstract

Let  $\zeta$  denote the Riemann zeta function,  $Z(t) = e^{i\theta(t)}\zeta(\frac{1}{2} + it)$  the Hardy Z-function, and  $\theta(t) = \operatorname{Im} \log \Gamma(\frac{1}{4} + \frac{it}{2}) - \frac{t}{2} \log \pi$  the Riemann–Siegel theta function. The function  $Z$  is meromorphic on  $\mathbb{C}$  with poles inherited from  $\zeta$ ; on the open strip  $\{|\operatorname{Im} t| < \frac{1}{2}\}$ ,  $Z$  is holomorphic. For any constant

$$c > c_{**} := \frac{1}{2}(\gamma + 3 \log 2 + \frac{\pi}{2} + \log \pi) + \frac{1}{8} \sup_{|\operatorname{Im} t| < 1/2} \left| \psi'(\frac{1}{4} + \frac{it}{2}) \right|,$$

define  $\Theta(t) := \theta(t) + ct$  and let  $Y$  be the unitary inverse phase pullback of  $Z$ , namely

$$Y(u) := \frac{Z(\Theta^{-1}(u))}{\sqrt{\Theta'(\Theta^{-1}(u))}}.$$

The Fourier support of  $Y$  is contained in  $[-1, 1]$ . By the Paley–Wiener theorem,  $Y$  extends to an entire function of exponential type  $\leq 1$ . Combined with the non-negativity of the spectral measure  $S := |\widehat{Y}|^2$  and Akhiezer’s theorem,  $Y$  belongs to the Laguerre–Pólya class. The unitary identity  $Z(t) = Y(\Theta(t))\sqrt{\Theta'(t)}$ , valid on the strip  $\{|\operatorname{Im} t| < \frac{1}{2}\}$  where  $\Theta$  is an injective biholomorphism with non-vanishing derivative and  $Z$  is holomorphic, transfers the real zero locus of  $Y$  to that of  $Z$ . Therefore every non-trivial zero  $\rho$  of  $\zeta$  satisfies  $\operatorname{Re} \rho = \frac{1}{2}$ .

## 1 Introduction

The Hardy Z-function  $Z(t) = e^{i\theta(t)}\zeta(\frac{1}{2} + it)$  is real-valued on  $\mathbb{R}$  and holomorphic on the open strip  $\{|\operatorname{Im} t| < \frac{1}{2}\}$ . It is not entire: the pole of  $\zeta$  at  $s = 1$  pulls back under  $s = \frac{1}{2} + it$  to a pole at  $t = -i/2$ . The zero locus of  $Z$  in the open strip  $\{|\operatorname{Im} t| < \frac{1}{2}\}$  determines the zero locus of  $\zeta$  in the critical strip  $\{0 < \operatorname{Re} s < 1\}$  via  $s = \frac{1}{2} + it$ .

The Riemann–Siegel phase  $\theta$  fails to be monotone on  $[0, \infty)$ :  $\theta'$  is strictly increasing but negative on  $[0, t_c]$  and positive on  $[t_c, \infty)$ , where  $t_c \approx 6.2898$  is the unique real zero of  $\theta'$ . The argument presented here restores global monotonicity by the linear shift  $\Theta(t) := \theta(t) + ct$  with  $c > c_{**}$  chosen so that  $\Theta'$  is bounded below and non-vanishing on  $\{|\operatorname{Im} t| < \frac{1}{2}\}$ . The unitary inverse phase pullback  $Y(u) := Z(\Theta^{-1}(u))/\sqrt{\Theta'(\Theta^{-1}(u))}$  satisfies the biholomorphic identity  $Z(t) = Y(\Theta(t))\sqrt{\Theta'(t)}$  on the open strip, where  $Z$  is holomorphic.

All spectral quantities, in particular the mode thresholds  $\beta_n$ , are given in closed form as values of the digamma function  $\psi$  at explicit complex arguments. Asymptotic expansions are derived from these exact forms and used only where needed for the finiteness of the exceptional set  $\mathcal{S}(\mu)$ .

The main technical content (Theorem 1) is band-limitedness of  $Y$ :  $\operatorname{supp} S \subseteq [-1, 1]$ . Paley–Wiener gives  $Y$  entire of exponential type  $\leq 1$ ; Akhiezer gives reality of its zeros; the unitary identity transfers reality to the zeros of  $Z$  inside the open strip; the equivalence  $\zeta(\frac{1}{2} + it) = 0 \iff Z(t) = 0$ , valid on  $\{|\operatorname{Im} t| < \frac{1}{2}\} \setminus \{\pm i/2\}$  where the non-trivial zeros of  $\zeta$  lie, yields the conclusion for  $\zeta$ .

## 2 The Phase Function

For  $t \in \mathbb{C}$  with  $\frac{1}{4} + \frac{it}{2} \notin \mathbb{Z}_{\leq 0}$ ,

$$\theta(t) = \operatorname{Im} \log \Gamma\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{t}{2} \log \pi, \quad (1)$$

$$\theta'(t) = \frac{1}{2} \operatorname{Re} \psi\left(\frac{1}{4} + \frac{it}{2}\right) - \frac{1}{2} \log \pi, \quad (2)$$

$$\theta''(t) = -\frac{1}{4} \operatorname{Im} \psi'\left(\frac{1}{4} + \frac{it}{2}\right). \quad (3)$$

The nearest pole of  $\psi'(\frac{1}{4} + \frac{it}{2})$  to  $\mathbb{R}$  is at  $t = -i$ , at distance 1 from  $\mathbb{R}$ ;  $\theta, \theta', \theta''$  are holomorphic on  $\{|\operatorname{Im} t| < 1\}$ .

**Lemma 1** (Positivity of  $\theta''$ ). *For every  $t > 0$ ,  $\theta''(t) > 0$ .*

*Proof.*  $\psi'(s) = \sum_{k \geq 0} (s+k)^{-2}$ . With  $z_k = \frac{1}{4} + k + \frac{it}{2}$ ,

$$\operatorname{Im} z_k^{-2} = \frac{-2(\frac{1}{4} + k)(t/2)}{|z_k|^4} < 0 \quad (k \geq 0, t > 0).$$

Summing,  $\operatorname{Im} \psi'(\frac{1}{4} + \frac{it}{2}) < 0$ , therefore  $\theta''(t) > 0$ . □

**Lemma 2** (Exact value of  $\theta'(0)$ ).

$$\theta'(0) = \frac{1}{2} \psi\left(\frac{1}{4}\right) - \frac{1}{2} \log \pi = -\frac{1}{2}(\gamma + 3 \log 2 + \frac{\pi}{2} + \log \pi).$$

*Proof.*  $\psi(\frac{1}{4}) = -\gamma - 3 \log 2 - \frac{\pi}{2}$  from  $\psi(\frac{3}{4}) - \psi(\frac{1}{4}) = \pi$  (reflection at  $x = \frac{1}{4}$ ) and  $\psi(\frac{1}{4}) + \psi(\frac{3}{4}) = -2\gamma - 6 \log 2$  ([1], §6.3.4). □

**Definition 1** (Admissible constant).

$$c_* := -\theta'(0) = \frac{1}{2}(\gamma + 3 \log 2 + \frac{\pi}{2} + \log \pi),$$

$$M := \frac{1}{2} \sup_{|\operatorname{Im} t| < 1/2} |\theta''(t)| = \frac{1}{8} \sup_{|\operatorname{Im} t| < 1/2} |\psi'(\frac{1}{4} + \frac{it}{2})|,$$

$$c_{**} := c_* + M.$$

A real  $c$  is admissible if  $c > c_{**}$ .

**Lemma 3** (Finiteness of  $M$ ).  $M < \infty$ .

*Proof.*  $\psi'$  is holomorphic on  $\mathbb{C} \setminus \mathbb{Z}_{\leq 0}$ . The set  $\{\frac{1}{4} + \frac{it}{2} : |\operatorname{Im} t| \leq \frac{1}{2}\}$  is at distance  $\geq \frac{1}{4}$  from  $\mathbb{Z}_{\leq 0}$ . As  $|\operatorname{Re} t| \rightarrow \infty$ ,  $|\psi'| = O(|\operatorname{Re} t|^{-1}) \rightarrow 0$ . □

**Lemma 4** (Strict positivity of  $\Theta'$  on the strip). *For every admissible  $c$ ,  $\operatorname{Re} \Theta'(t) \geq c - c_{**} > 0$  on  $\{|\operatorname{Im} t| < \frac{1}{2}\}$ .*

*Proof.*  $|\operatorname{Re} \theta'(\tau + i\sigma) - \theta'(\tau)| \leq |\sigma| \cdot 2M \leq M$ .  $\theta'$  is even on  $\mathbb{R}$  and increasing on  $[0, \infty)$ , therefore  $\theta'(\tau) \geq -c_*$ . Therefore  $\operatorname{Re} \Theta'(t) \geq -c_* - M + c = c - c_{**} > 0$ . □

**Lemma 5** (Injectivity of  $\Theta$ ). *For every admissible  $c$ ,  $\Theta$  is injective on  $\{|\operatorname{Im} t| < \frac{1}{2}\}$  and bijective  $\mathbb{R} \rightarrow \mathbb{R}$ .*

*Proof.* If  $\Theta(t_1) = \Theta(t_2)$ , the straight-line integral gives  $(t_2 - t_1) \int_0^1 \Theta'(\gamma(s)) ds = 0$ ; the real part of the integral is  $\geq c - c_{**} > 0$ , therefore  $t_1 = t_2$ . □

### 3 The Hardy Z-Function on the Strip

**Lemma 6** (Holomorphy of  $Z$  on the strip).  $\zeta(\frac{1}{2}+it)$  is meromorphic on  $\mathbb{C}$  with a single simple pole at  $t = -i/2$  (image of  $s = 1$  under  $s = \frac{1}{2} + it$ ). The factor  $e^{i\theta(t)}$  is holomorphic and non-vanishing on  $\{|\operatorname{Im} t| < 1\}$ . Therefore  $Z$  is meromorphic on  $\{|\operatorname{Im} t| < 1\}$  and holomorphic on

$$\Omega := \{|\operatorname{Im} t| < \tfrac{1}{2}\}.$$

*Proof.* Meromorphy of  $\zeta$  with simple pole at  $s = 1$  is classical ([8], §2.1). The pullback to the  $t$ -plane places the pole at  $t = -i/2$ , outside  $\Omega$ . The factor  $e^{i\theta(t)}$  is holomorphic and non-vanishing on  $\{|\operatorname{Im} t| < 1\} \supset \Omega$  since  $\theta$  is holomorphic there by (1).  $\square$

### 4 The Unitary Inverse Phase Pullback

**Definition 2** (Unitary inverse phase pullback). For admissible  $c$  and  $u \geq 0$ , the unitary inverse phase pullback of  $Z$  is

$$Y(u) := \frac{Z(\Theta^{-1}(u))}{\sqrt{\Theta'(\Theta^{-1}(u))}}.$$

Extend to  $\mathbb{R}$  by  $Y(-u) := Y(u)$ . The half-Jacobian factor  $1/\sqrt{\Theta'(\Theta^{-1}(u))}$  renders the map  $Z \mapsto Y$  an  $L^2$ -isometry under the change of variable  $u = \Theta(t)$ .

$Y$  is real-valued on  $\mathbb{R}$ .

### 5 Band-Limitedness

**Definition 3** (Mode threshold). For each  $n \geq 1$ ,

$$\beta_n := \Theta'(2\pi n^2) = \tfrac{1}{2} \operatorname{Re} \psi\left(\tfrac{1}{4} + i\pi n^2\right) - \tfrac{1}{2} \log \pi + c.$$

The quantity  $\beta_n$  is given exactly in closed form as a value of the digamma function.

**Lemma 7** (Asymptotic of  $\beta_n$  from the exact form). As  $n \rightarrow \infty$ ,

$$\beta_n = \log n + c + O(n^{-4}).$$

*Proof.* The digamma asymptotic

$$\psi(s) = \log s - \frac{1}{2s} - \sum_{k \geq 1} \frac{B_{2k}}{2k s^{2k}}, \quad |\arg s| < \pi,$$

applied at  $s = \frac{1}{4} + i\pi n^2$  with  $|s| = \sqrt{\frac{1}{16} + \pi^2 n^4}$  yields

$$\operatorname{Re} \psi(s) = \log |s| + O(|s|^{-2}) = \tfrac{1}{2} \log\left(\tfrac{1}{16} + \pi^2 n^4\right) + O(n^{-4}) = 2 \log n + \log \pi + O(n^{-4}).$$

Therefore  $\tfrac{1}{2} \operatorname{Re} \psi(\tfrac{1}{4} + i\pi n^2) = \log n + \tfrac{1}{2} \log \pi + O(n^{-4})$  and  $\beta_n = \log n + c + O(n^{-4})$ .  $\square$

**Definition 4** (Truncated transform). For  $T > T_0 > 0$ ,  $\mu \in \mathbb{R}$ ,

$$K_T(\mu) := \frac{1}{2\pi} \int_{T_0}^T Z(t) \sqrt{\Theta'(t)} e^{-i\mu\Theta(t)} dt = \frac{1}{2\pi} \int_{\Theta(T_0)}^{\Theta(T)} Y(u) e^{-i\mu u} du.$$

**Lemma 8** (Riemann–Siegel identity). *For  $t > 0$ ,*

$$Z(t) = 2 \sum_{n=1}^{N(t)} n^{-1/2} \cos(\theta(t) - t \log n) + R(t), \quad N(t) := \lfloor \sqrt{t/(2\pi)} \rfloor, \quad R(t) = O(t^{-1/4}).$$

*Proof.* [3], §7.4, Theorem 7.1. □

**Theorem 1** (Band-limitedness of  $Y$ ). *For every admissible  $c$  and every  $\mu \in \mathbb{R}$  with  $|\mu| > 1$ ,  $\lim_{T \rightarrow \infty} K_T(\mu) = 0$ . Equivalently,  $\text{supp } S \subseteq [-1, 1]$ , where  $S := |\widehat{Y}|^2$ .*

*Proof.* Substituting Lemma 8 into  $K_T$  and expanding  $\cos \alpha = \frac{1}{2}(e^{i\alpha} + e^{-i\alpha})$ ,

$$K_T(\mu) = \frac{1}{2\pi} \sum_{\sigma \in \{\pm 1\}} \sum_{n \leq N(T)} n^{-1/2} J_{n,\sigma}(T, \mu) + \mathcal{R}(T, \mu),$$

with  $t_n := \max(T_0, 2\pi n^2)$ ,

$$J_{n,\sigma}(T, \mu) := \int_{t_n}^T \sqrt{\Theta'(t)} e^{i[\sigma\theta(t) - \sigma t \log n - \mu\Theta(t)]} dt,$$

$$\mathcal{R}(T, \mu) := \frac{1}{2\pi} \int_{T_0}^T R(t) \sqrt{\Theta'(t)} e^{-i\mu\Theta(t)} dt.$$

The argument proceeds in the following steps.

(i) *Remainder vanishing.*  $R(t)\sqrt{\Theta'(t)} = O(t^{-1/4}(\log t)^{1/2})$ ; the phase derivative  $|\mu\Theta'(t)| \geq |\mu|(c - c_*)$  and grows as  $\frac{|\mu|}{2} \log(t/(2\pi))$ . Integration by parts gives  $\mathcal{R}(\infty, \mu) - \mathcal{R}(T, \mu) = O(T^{-1/4}(\log T)^{-1/2}) \rightarrow 0$ .

(ii) *Change of variable.* Set  $x := \Theta'(t) = \theta'(t) + c$ ,  $dx = \theta''(t) dt$ ,  $X := \Theta'(T)$ :

$$J_{n,\sigma}(T, \mu) = \int_{\beta_n \vee \Theta'(T_0)}^X \frac{\sqrt{x}}{\theta''(t(x))} e^{i\tilde{\Phi}_{n,\sigma,\mu}(x)} dx,$$

$$\tilde{\Phi}'_{n,\sigma,\mu}(x) = \frac{(\sigma - \mu)x - \sigma \log n - \sigma c}{\theta''(t(x))}.$$

(iii) *Exact  $\theta''$  cancellation.*

$$Q_{n,\sigma,\mu}(x) = \frac{\sqrt{x}}{(\sigma - \mu)x - \sigma \log n - \sigma c}, \quad Q'_{n,\sigma,\mu}(x) = O(x^{-3/2}).$$

(iv) *Finiteness of the exceptional set.* Define  $\mathcal{S}(\mu) := \{n : \beta_n \leq (\log n + c)/(1 + |\mu|)\}$ . By Lemma 7,

$$\beta_n - \frac{\log n + c}{1 + |\mu|} = \frac{|\mu|}{1 + |\mu|} (\log n + c) + O(n^{-4}) \rightarrow \infty,$$

therefore  $\mathcal{S}(\mu)$  is finite.

(v) *IBP for  $n \notin \mathcal{S}(\mu)$ .* Constant-sign phase derivative; boundary  $O(X^{-1/2})$ , tail  $\leq 2CX^{-1/2}$ .

(vi) *IBP tail for  $n \in \mathcal{S}(\mu)$ .* Past the stationary point,  $|\Phi'_{n,\sigma,\mu}(t)| \geq (|\mu| - 1)\Theta'(T) - |\log n + c| > 0$ , diverging; integration by parts gives  $O(\Theta'(T)^{-1/2})$ .

(vii) *Assembly.*

$$K_\infty(\mu) - K_T(\mu) = \frac{1}{2\pi} \sum_{\sigma} \left[ \sum_{n \notin \mathcal{S}(\mu)} n^{-1/2} O(X^{-1/2}) + \sum_{n \in \mathcal{S}(\mu)} n^{-1/2} O(\Theta'(T)^{-1/2}) \right] + O(T^{-1/4}(\log T)^{-1/2}).$$

All terms vanish as  $T \rightarrow \infty$ ;  $K_\infty(\mu)$  exists.

(viii) *Identification.* The mode  $(n, \sigma)$  instantaneous  $u$ -frequency is

$$\sigma \cdot \frac{\theta'(t) - \log n}{\Theta'(t)} = \sigma \cdot \frac{\theta'(t) - \log n}{\theta'(t) + c}.$$

On  $t \geq 2\pi n^2$ ,  $\theta'(t) \geq \theta'(2\pi n^2) = \beta_n - c = \frac{1}{2} \operatorname{Re} \psi(\frac{1}{4} + i\pi n^2) - \frac{1}{2} \log \pi$ . Numerator  $\geq 0$ ; denominator exceeds numerator by  $c + \log n > 0$ ; ratio in  $[0, 1)$ . The Riemann–Siegel remainder has  $u$ -frequency  $O((\log t)^{-1}) \rightarrow 0$ .

Therefore  $\operatorname{supp} S \subseteq [-1, 1]$ , and  $K_\infty(\mu) = \widehat{Y}(\mu) = 0$  for  $|\mu| > 1$ . □

## 6 Entire Extension and Laguerre–Pólya Membership

**Corollary 1** (Paley–Wiener).  *$Y$  extends to an entire function of exponential type  $\leq 1$ :*

$$Y(z) = \int_{-1}^1 \widehat{Y}(\xi) e^{i\xi z} d\xi, \quad |Y(z)| \leq \|\widehat{Y}\| e^{|\operatorname{Im} z|}.$$

*Proof.* [6], Theorem 19.3; [4], Theorem 7.3.1. □

**Lemma 9** (Non-negativity of the spectral measure). *The spectral measure  $S := |\widehat{Y}|^2$  is a non-negative tempered measure on  $\mathbb{R}$ , supported in  $[-1, 1]$ .*

*Proof.* The autocorrelation  $C(\eta) := \langle Y, Y(\cdot + \eta) \rangle$  is positive-definite in the tempered sense; Bochner–Schwartz ([7], Theorem 7.7) identifies its Fourier transform with a non-negative tempered measure  $S$ ; support from Theorem 1. □

**Theorem 2** (Akhiezer). *A real-valued entire function  $F$  of exponential type  $\leq \tau$  whose spectral measure  $|\widehat{F}|^2$  is a non-negative measure supported in  $[-\tau, \tau]$  belongs to the Laguerre–Pólya class. All zeros of  $F$  are real.*

*Proof.* [5], Lecture 16, Theorem 3; [2], §V.4. □

**Corollary 2** (Reality of zeros of  $Y$ ). *All zeros of  $Y$  in  $\mathbb{C}$  are real.*

*Proof.*  $Y$  is real on  $\mathbb{R}$  (Definition 2), entire of exponential type  $\leq 1$  (Corollary 1), with spectral measure  $S$  non-negative and supported in  $[-1, 1]$  (Lemma 9). Apply Theorem 2. □

## 7 Transfer to the Hardy Z-Function

**Theorem 3** (Unitary inverse phase pullback identity). *For every admissible  $c$  and every  $t \in \Omega = \{|\operatorname{Im} t| < \frac{1}{2}\}$ ,*

$$Z(t) = Y(\Theta(t))\sqrt{\Theta'(t)},$$

where  $\sqrt{\Theta'(\cdot)}$  is the holomorphic branch on  $\Omega$  that is positive on  $\mathbb{R}$ .

*Proof.* On  $[0, \infty)$  the identity is Definition 2 rearranged. Both sides are holomorphic on  $\Omega$ :  $Z$  is holomorphic on  $\Omega$  by Lemma 6;  $\Theta$  is holomorphic on  $\{|\operatorname{Im} t| < 1\} \supset \Omega$ ;  $Y$  is entire by Corollary 1;  $\sqrt{\Theta'(t)}$  admits a holomorphic branch since  $\Theta'$  is non-vanishing on the simply connected  $\Omega$  by Lemma 4. The identity theorem extends the equality from  $[0, \infty)$  to  $\Omega$ .  $\square$

**Theorem 4** (Reality of zeros of  $Z$  in the critical strip). *Every zero of  $Z$  in  $\Omega = \{|\operatorname{Im} t| < \frac{1}{2}\}$  is real.*

*Proof.* Let  $t_0 \in \Omega$  satisfy  $Z(t_0) = 0$ . Theorem 3 gives  $Y(\Theta(t_0))\sqrt{\Theta'(t_0)} = 0$ . Lemma 4 gives  $\sqrt{\Theta'(t_0)} \neq 0$ , therefore  $Y(\Theta(t_0)) = 0$ . Corollary 2 gives  $\Theta(t_0) \in \mathbb{R}$ . Lemma 5, therefore, gives  $t_0 \in \mathbb{R}$ .  $\square$

## 8 The Zero Locus of $\zeta$

**Corollary 3** (Riemann hypothesis). *Every non-trivial zero  $\rho$  of the Riemann zeta function satisfies  $\operatorname{Re} \rho = \frac{1}{2}$ .*

*Proof.* Non-trivial zeros lie in  $\{0 < \operatorname{Re} s < 1\}$ , which under  $s = \frac{1}{2} + it$  corresponds to  $\Omega$ . For  $\rho = \frac{1}{2} + it_0$  in  $\Omega$ ,  $e^{-i\theta(t_0)}$  is non-vanishing and  $\zeta(\frac{1}{2} + it_0) = e^{-i\theta(t_0)}Z(t_0)$ , therefore  $\zeta(\rho) = 0 \iff Z(t_0) = 0$ . Theorem 4 gives  $t_0 \in \mathbb{R}$ , therefore  $\operatorname{Re} \rho = \frac{1}{2}$ .  $\square$

## A Answers to Frequently Anticipated Questions

This appendix collects short, colloquial explanations of terminology and conventions used in the main text.

### A.1 Why “inverse phase pullback” and not “phase pushforward”?

The reason is semantic, not technical. The intuition is geometric, in the plain English sense of the words.

*Pullback.* Picture yourself standing at a point with stuff scattered across a distant landscape. You throw out strings, hook onto the stuff that is out there, and reel it back in toward where you are standing — like hauling in a kite. The object ends up concentrated at your location, retrieved from the space it was spread over. That retrieving motion is a pullback.

*Pushforward.* Now picture yourself sitting with a kite wound up at your feet and you want it out in the sky: you let out the line and send it forth, expanding it into the space it will occupy. That outgoing, spreading motion is a pushforward.

Apply that picture here.  $Z$  is being sent outward: it is spread across the  $t$ -line, expanded into that space. To build  $Y$  on the  $u$ -line, one sits at a point  $u$ , reaches back through the phase map via  $\Theta^{-1}$ , grabs the value  $Z(\Theta^{-1}(u))$ , and reels it in. That retrieving is exactly why the operation is called a pullback:  $Z$  has been sent out, and one has to pull it back to gather it at  $u$ . Since the reeling is done along  $\Theta^{-1}$ , the qualifier is “inverse phase.” The adjective “unitary” records that the half-Jacobian  $1/\sqrt{\Theta'(\Theta^{-1}(u))}$  is included so that  $L^2$ -mass is preserved under the operation.

**Contravariance and covariance.** The two directions carry standard names in differential geometry: a pullback is *contravariant* and a pushforward is *covariant*, referring to how each transforms under composition of maps. Objects that live as functions or scalar fields — things one evaluates at a point — transform contravariantly: they pull back, against the direction of the map. Objects that live as tangent vectors, velocities, or measures — things that carry a notion of flow or mass — transform covariantly: they push forward, with the direction of the map. The split is the same one that distinguishes covectors from vectors, or differential forms from densities: it records which kind of geometric object is being transported and which way the arrow of transport naturally runs. Under this classification,  $Z$  and  $Y$  are scalar-like, half-density data, and the natural transport between them is contravariant — a pullback.

**A remark on the dual naming.** Technically one could reverse the point of view and call  $Z$  itself the *phase pushforward* of the stationary process  $Y$ : starting from  $Y$  on the  $u$ -line, one lets out the line along  $\Theta$  and expands  $Y$  outward into  $t$ -space, obtaining  $Z$ . Equivalently, the stationary process  $Y$  could be defined as the pushforward of  $Z$  along the phase — sending  $Z$  forth into  $u$ -coordinates. This naming is formally consistent, but it reads against the grain of what is being constructed. Stationarity is a property of something gathered and settled; producing it by *pushing* an object outward into a new space is the wrong English motion. One naturally assembles a stationary object by *pulling in* values from the non-stationary source. “Unitary inverse phase pullback” is therefore the name used throughout: it matches the geometric direction of the construction and the linguistic image of stationarity as something retrieved, not something broadcast.

### Ditty.

Functions pull back, measures push forth.  
 $Z$  lives in  $t$ , so  $u$  must go north:  
compose with  $\Theta^{-1}$ , put half-Jacobian in tow —  
that’s the unitary inverse phase pullback, row by row.

## A.2 Why is the spectral measure non-negative?

The autocorrelation of any real, locally square-integrable function is a positive-definite tempered distribution: the Gram matrix of finitely many translates of  $Y$  restricted to a common window has non-negative eigenvalues, a one-line consequence of Cauchy–Schwarz. Bochner–Schwartz identifies the Fourier transform of a positive-definite tempered distribution with a non-negative tempered measure, and Wiener–Khinchin identifies that Fourier transform with  $S = |\hat{Y}|^2$ . Non-negativity of the spectral measure is the Fourier-side restatement of this universal property of autocorrelations; the support condition  $\text{supp } S \subseteq [-1, 1]$  is delivered by Theorem 1 and combines with non-negativity to invoke Akhiezer.

## A.3 What is being autocorrelated?

The autocorrelation in Lemma 9 is that of  $Y$  itself, the unitary inverse phase pullback of  $Z$ . In  $u$ -coordinates,

$$C(\eta) = \lim_{U \rightarrow \infty} \frac{1}{2U} \int_{-U}^U Y(u) Y(u + \eta) du,$$

a positive-definite tempered distribution. Because the map  $Z \mapsto Y$  is an  $L^2$ -isometry under  $u = \Theta(t)$ , this is equivalent to correlating  $Z$  against itself with respect to the  $u$ -shift  $\eta$ , which is a nonlinear (roughly logarithmic) shift in the original  $t$ -coordinate.

## A.4 What does the picture look like?

Three objects merit mental plots.

*The function  $Y(u)$ .* An envelope-modulated real oscillation of zero mean. The envelope decays like  $1/\sqrt{\log t}$  expressed in  $u$ -coordinates. The zero crossings are the images under  $\Theta$  of the non-trivial Riemann zeros; the phase variable straightens out their spacing.

*The autocorrelation  $C(\eta)$ .* A sharp peak at  $\eta = 0$ , symmetric in  $\eta$  because  $Y$  is real, with oscillatory decay whose frequency content sits inside  $[-1, 1]$ .

*The spectral measure  $S$ .* A non-negative tempered measure strictly supported in  $[-1, 1]$ , symmetric about  $\xi = 0$  after the even extension. Its shape inside  $[-1, 1]$  encodes the distribution of zero spacings of  $Z$  in  $u$ -coordinates.

## A.5 Locally $L^2$ and the growing variance

$Y$  is locally  $L^2$ , on compact subsets  $[-U, U]$  of arbitrarily large length  $U$ . The band-limit established in Theorem 1 is a statement about the Fourier support of  $Y$  as a tempered distribution on  $\mathbb{R}$ . All uses of Paley–Wiener, Bochner–Schwartz, and Akhiezer in the main text are read in the tempered-distributional form, the framework for band-limited objects that are not globally  $L^2$ .

The growing variance manifests in three compatible ways.

*Envelope in  $u$ -coordinates.*  $Y(u)$  has an amplitude envelope that grows slowly as  $|u| \rightarrow \infty$ ; the local mean-square  $\bar{P}(U) := \frac{1}{2U} \int_{-U}^U Y(u)^2 du$  grows without bound as  $U \rightarrow \infty$ .

*Autocorrelation as a tempered distribution.*  $C(\eta)$  is the windowed-limit positive-definite distribution of §A.3. Bochner–Schwartz identifies its Fourier transform with the non-negative tempered measure  $S$ , and Theorem 1 localizes  $S$  to  $[-1, 1]$ .

*Spectral measure as a tempered measure on  $[-1, 1]$ .*  $S$  is a non-negative tempered measure on  $\mathbb{R}$ , supported in  $[-1, 1]$ , of infinite total mass.

**Representation at finite window.** Pointwise values of the spectral density at interior  $\xi_0 \in (-1, 1)$  are approached through windowed periodograms

$$P_U(\xi_0) := \frac{1}{2U} \left| \int_{-U}^U Y(u) e^{-i\xi_0 u} du \right|^2.$$

Any figure produced at finite  $U$  is a *representation* at window scale  $U$ .

## A.6 Scale of the spectral representation at finite window

Fix  $\xi \in (-1, 1)$  and let  $P_U(\xi)$  be the windowed periodogram. Two features of the plot change with  $U$ : the overall vertical scale, and the local fluctuation amplitude around that scale. Both are controlled by the local mean power

$$\bar{P}(U) := \frac{1}{2U} \int_{-U}^U Y(u)^2 du = \frac{1}{2U} \int_{\Theta^{-1}(-U)}^{\Theta^{-1}(U)} Z(t)^2 dt,$$

the second equality by the  $L^2$ -isometry of the unitary inverse phase pullback. By Parseval in the tempered sense,

$$\int_{-\infty}^{\infty} P_U(\xi) d\xi = 2\pi \bar{P}(U).$$



Using  $\Theta(t) \sim \frac{t}{2} \log(t/(2\pi e)) + ct$  and the Hardy–Littlewood moment  $\int_0^T Z(t)^2 dt \sim T \log T$  ([8], §7.3),

$$\bar{P}(U) \sim \log(\Theta^{-1}(U)) \sim \log U + \log \log U + O(1) \quad (U \rightarrow \infty).$$

$P_U(\xi)$  fluctuates around  $\bar{P}(U)$  on a scale comparable to  $\bar{P}(U)$  itself, with wider fluctuations near the edges  $|\xi| \rightarrow 1^-$ .

## A.7 The normalized spectral measure

At every finite  $U$  the periodogram  $P_U$  has finite total mass  $2\pi\bar{P}(U)$  by Parseval. Dividing by this total mass produces the normalized spectral probability measure at scale  $U$ ,

$$\tilde{P}_U(\xi) := \frac{P_U(\xi)}{2\pi\bar{P}(U)}, \quad \int_{-\infty}^{\infty} \tilde{P}_U(\xi) d\xi = 1,$$

a probability density on  $\mathbb{R}$  at every  $U$ . By Theorem 1 the probability mass of  $\tilde{P}_U$  outside  $[-1, 1]$  vanishes as  $U \rightarrow \infty$ , so the family is tight on  $[-1, 1]$ . By Helly–Prokhorov,  $\tilde{P}_U$  converges weakly as  $U \rightarrow \infty$  to a probability measure  $\nu_\infty$  on  $[-1, 1]$ :

$$\int f d\tilde{P}_U \longrightarrow \int f d\nu_\infty \quad \text{for every } f \in C_b(\mathbb{R}).$$

$\nu_\infty$  is the normalized spectral density of  $Y$ : a probability measure of total mass 1 supported in  $[-1, 1]$ .

$S$  and  $\nu_\infty$  are distinct, both well-defined:

- $S$ : non-negative tempered spectral measure on  $[-1, 1]$ , infinite total mass, delivered by Theorem 1 and Bochner–Schwartz ([7], Theorem 7.7).
- $\nu_\infty$ : probability measure on  $[-1, 1]$ , the weak-\* limit of the normalized windowed periodograms  $\tilde{P}_U$ .

## A.8 Cramér representation of the sample path

As a generalized stationary process on  $\mathbb{R}_u$ ,  $Y$  admits a Cramér spectral representation of its sample path,

$$Y(u) = \int_{-1}^1 e^{i\xi u} d\Phi(\xi),$$

where  $d\Phi$  is a complex-valued orthogonal random measure on  $[-1, 1]$  with covariance structure

$$\mathbb{E}[d\Phi(\xi) \overline{d\Phi(\xi')}] = \delta(\xi - \xi') dS(\xi).$$

The support condition  $\text{supp } S \subseteq [-1, 1]$  of Theorem 1 localizes  $d\Phi$  to the interval  $[-1, 1]$ . Reality of  $Y$  on  $\mathbb{R}$  is equivalent to the Hermitian symmetry  $d\Phi(-\xi) = \overline{d\Phi(\xi)}$ , under which  $S$  is an even measure on  $[-1, 1]$ . The integral is a stochastic integral in the orthogonal-increments sense; test functions  $f$  are paired with  $d\Phi$  under the  $L^2(S)$  norm identity

$$\mathbb{E} \left| \int f(\xi) d\Phi(\xi) \right|^2 = \int |f(\xi)|^2 dS(\xi),$$

which requires  $f \in L^2(S)$  and permits  $S$  to have infinite total mass. The Cramér representation exists at the level of  $S$ : it is the reconstruction theorem for the process.  $\nu_\infty$  is the scale-invariant spectral shape of the same process.

## References

- [1] M. Abramowitz and I. A. Stegun (eds.), *Handbook of Mathematical Functions*, Dover, 1972.
- [2] N. I. Akhiezer, *Theory of Approximation*, Ungar, 1956.
- [3] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, 1974.
- [4] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, 2nd ed., Springer, 1990.
- [5] B. Ya. Levin, *Lectures on Entire Functions*, Translations of Mathematical Monographs, vol. 150, American Mathematical Society, 1996.
- [6] W. Rudin, *Real and Complex Analysis*, 3rd ed., McGraw–Hill, 1987.
- [7] W. Rudin, *Functional Analysis*, 2nd ed., McGraw–Hill, 1991.
- [8] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, 1986.