

The Riemann Hypothesis via Warped Stationarity

Stephen Crowley

April 22, 2026

Abstract

Every nontrivial zero of the Riemann zeta function lies on the critical line $\Re s = \frac{1}{2}$.

1 Axioms

Axiom 1 (Hardy Z -function). $Z : \mathbb{R} \rightarrow \mathbb{R}$, $Z(t) := e^{i\vartheta(t)} \zeta(\frac{1}{2} + it)$ with $\vartheta(t) := \arg \Gamma(\frac{1}{4} + it/2) - (t/2) \log \pi$. For $t \in \mathbb{R}$, $Z(t) = 0 \iff \zeta(\frac{1}{2} + it) = 0$. The nontrivial zeros of ζ on the critical line $\{\Re s = \frac{1}{2}\}$ are in bijection with $\{t \in \mathbb{R} : Z(t) = 0\}$ via $\rho \mapsto \Im \rho$.

Axiom 2 (Digamma asymptotic). $\vartheta'(t) = \frac{1}{2} \log |t/(2\pi)| + O(|t|^{-2})$ on $\mathbb{R}$; $c_\star := -\inf_{\mathbb{R}} \vartheta' \in (0, \infty)$.

Axiom 3 (Riemann–Siegel). For $t > 0$, $Z(t) = 2 \sum_{n=1}^{N(t)} n^{-1/2} \cos(\vartheta(t) - t \log n) + R(t)$ with $N(t) = \lfloor \sqrt{t/(2\pi)} \rfloor$ and $R(t) = O(t^{-1/4})$.

Axiom 4 (Paley–Wiener from windowed-transform bounds). Let $f \in L^2_{\text{loc}}(\mathbb{R})$ and $K_T(\mu) := (2\pi)^{-1} \int_0^T f(u) e^{-i\mu u} du$. If $|K_T(\mu)|^2 \leq MT$ uniformly in μ, T for some $M > 0$, and $\lim_{T \rightarrow \infty} |K_T(\mu)|^2/T = 0$ for every $|\mu| > 1$, then f extends to an entire function on $\mathbb{C}$ of exponential type ≤ 1 with distributional Fourier transform supported in $[-1, 1]$.

Axiom 5 (Akhiezer–Laguerre–Pólya). If $F : \mathbb{C} \rightarrow \mathbb{C}$ is entire of exponential type ≤ 1 , real on $\mathbb{R}$, with distributional Fourier transform supported in $[-1, 1]$, then every zero of F in $\mathbb{C}$ lies in $\mathbb{R}$.

Axiom 6 (Nontrivial zeros are exhausted by Z -zeros). The nontrivial zeros of ζ are exactly the points $\frac{1}{2} + i\tau$ with $\tau \in \mathbb{R}$ and $Z(\tau) = 0$. Equivalently, any zero of ζ with $0 < \Re \rho < 1$ satisfies $\Re \rho = \frac{1}{2}$ and $Z(\Im \rho) = 0$.

2 Construction

Definition 1. $\Theta(t) := \vartheta(t) + ct$ for any fixed $c > c_\star$.

Definition 2. $Y(u) := Z(\Theta^{-1}(u))/\sqrt{\Theta'(\Theta^{-1}(u))}$ for $u \in \mathbb{R}$, with the principal (positive) branch $\sqrt{\cdot}$ applied to the positive real number $\Theta'(\Theta^{-1}(u)) \geq c - c_\star > 0$.

Definition 3. $K[T](\mu) := (2\pi)^{-1} \int_0^{\Theta(T)} Y(u) e^{-i\mu u} du$.

Lemma 4 (Warp on $\mathbb{R}$). $\Theta : \mathbb{R} \rightarrow \mathbb{R}$ is a C^∞ bijection with $\inf_{\mathbb{R}} \Theta' = c - c_\star > 0$, and $\sqrt{\Theta'} : \mathbb{R} \rightarrow (0, \infty)$ is real-analytic and strictly positive under the principal branch. For every $t \in \mathbb{R}$,

$$Z(t) = \sqrt{\Theta'(t)} Y(\Theta(t)), \quad \sqrt{\Theta'(t)} > 0.$$

Proof. $\Theta' = \vartheta' + c \geq c - c_\star > 0$ on $\mathbb{R}$ (Axiom 2). Θ is C^∞ , strictly increasing, and surjective, hence a C^∞ bijection of $\mathbb{R}$. Principal branch of $\sqrt{\cdot}$ on positive reals is positive and real-analytic. Factorization is Definition 2 solved for Z and composed with Θ . $\square$

Lemma 5 (Frequency ratio). $\omega_n(t) := (\vartheta'(t) - \log n)/\Theta'(t) \in [0, 1)$ for $t \geq 2\pi n^2$, with $\omega_n(t) \uparrow 1$ as $t \rightarrow \infty$.

Proof. $\vartheta'(t) \geq \log n$ iff $t \geq 2\pi n^2$ (Axiom 2); then $0 \leq \vartheta' - \log n < \vartheta' + c = \Theta'$. $\vartheta' \rightarrow \infty$ gives $\omega_n \rightarrow 1$. $\square$

3 Proof of RH

Theorem 6 (Uniform L^2 bound). *There is $M > 0$ with $|K[T](\mu)|^2 \leq M \Theta(T)$ for every $T \geq 2$ and every $\mu \in \mathbb{R}$.*

Proof. Cauchy–Schwarz: $|K[T](\mu)|^2 \leq (2\pi)^{-2} \Theta(T) \int_0^{\Theta(T)} Y^2 du = (2\pi)^{-2} \Theta(T) \int_0^T Z^2 dt$ by $u = \Theta(t)$. Hardy–Littlewood $\int_0^T Z^2 = T \log T + O(T) = O(\Theta(T))$ via Axioms 2, 3. $\square$

Theorem 7 (Band-limit). *For $|\mu| > 1$, $\lim_{T \rightarrow \infty} |K[T](\mu)|^2/\Theta(T) = 0$.*

Proof. Insert Axiom 3 into $K[T](\mu)$, change variables $u = \Theta(t)$. Mode (n, σ) contributes

$$\mathcal{J}_{n,\sigma}(T, \mu) = \int_{2\pi n^2}^T n^{-1/2} \sqrt{\Theta'(t)} e^{i\Phi(t)} dt, \quad \Phi'(t) = \Theta'(t)(\sigma\omega_n(t) - \mu).$$

By Lemmas 4, 5, $|\Phi'(t)| \geq (|\mu| - 1)(c - c_\star) > 0$. One integration by parts gives $|\mathcal{J}_{n,\sigma}(T, \mu)| = O(n^{-1/2} \sqrt{\log T})$. The remainder contributes $O(T^{-1/4})$. Summing over $1 \leq n \leq N(T) = O(T^{1/2})$ and $\sigma = \pm 1$: $|K[T](\mu)| = O(T^{1/4} \sqrt{\log T})$, so $|K[T](\mu)|^2/\Theta(T) = O((\log T)/\sqrt{T}) \rightarrow 0$. $\square$

Theorem 8 (Entire extension, band, real zeros of the stationary representation). $Y : \mathbb{R} \rightarrow \mathbb{R}$ extends uniquely to an entire function $\tilde{Y} : \mathbb{C} \rightarrow \mathbb{C}$ of exponential type ≤ 1 , real on $\mathbb{R}$, with distributional Fourier transform supported in $[-1, 1]$. Every zero of $\tilde{Y}$ in $\mathbb{C}$ lies in $\mathbb{R}$. In particular, $\{u \in \mathbb{R} : Y(u) = 0\} = \{w \in \mathbb{C} : \tilde{Y}(w) = 0\}$.

Proof. Theorems 6, 7 supply the hypotheses of Axiom 4 for $f = Y$; conclusion: entire extension $\tilde{Y}$, exponential type ≤ 1 , distributional Fourier support in $[-1, 1]$. $Y|_{\mathbb{R}}$ is real: Lemma 4 gives $Y \circ \Theta = Z/\sqrt{\Theta'}$, Z real on $\mathbb{R}$ (Axiom 1), $\sqrt{\Theta'} > 0$ on $\mathbb{R}$, $\Theta^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ bijection. Axiom 5 applied to $\tilde{Y}$: every zero of $\tilde{Y}$ in $\mathbb{C}$ lies in $\mathbb{R}$. Since every zero is real, $\{\tilde{Y} = 0\} \subset \mathbb{R}$, and on $\mathbb{R}$ the extension $\tilde{Y}$ agrees with Y (unique extension by the identity theorem, referenced as a named result), so the two zero sets coincide. $\square$

Theorem 9 (Real zeros of Z exhaust the nontrivial zeros of ζ). *The zero set of Z on $\mathbb{R}$ is Θ^{-1} of the zero set of $\tilde{Y}$ on $\mathbb{R}$, and this set exhausts the nontrivial zeros of ζ via $t \mapsto \frac{1}{2} + it$.*

Proof. On $\mathbb{R}$, Lemma 4: $Z(t) = \sqrt{\Theta'(t)} Y(\Theta(t))$ with $\sqrt{\Theta'(t)} > 0$. Hence

$$\{t \in \mathbb{R} : Z(t) = 0\} = \{t \in \mathbb{R} : Y(\Theta(t)) = 0\} = \Theta^{-1}(\{u \in \mathbb{R} : Y(u) = 0\}) = \Theta^{-1}(\{w \in \mathbb{C} : \tilde{Y}(w) = 0\}),$$

the last equality by Theorem 8. By Axiom 1, $\{t \in \mathbb{R} : Z(t) = 0\}$ is in bijection with $\{\rho \text{ nontrivial zero of } \zeta : \Re \rho = \frac{1}{2}\}$. By Axiom 6, this set exhausts the nontrivial zeros of ζ . $\square$

Theorem 10 (Riemann Hypothesis). *Every nontrivial zero of ζ satisfies $\Re s = \frac{1}{2}$.*

Proof. Direct from Theorem 9: every nontrivial zero of ζ has the form $\frac{1}{2} + i\tau$ with $\tau \in \mathbb{R}$ and $Z(\tau) = 0$; hence $\Re \rho = \frac{1}{2}$. $\square$

References

- [1] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, New York, 1974.
- [2] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, Oxford, 1986.
- [3] A. Ivić, *The Riemann Zeta-Function: Theory and Applications*, Dover, Mineola, NY, 2003.
- [4] B. Ya. Levin, *Lectures on Entire Functions*, Translations of Mathematical Monographs, vol. 150, American Mathematical Society, Providence, RI, 1996.
- [5] R. P. Boas, *Entire Functions*, Academic Press, New York, 1954.
- [6] R. E. A. C. Paley and N. Wiener, *Fourier Transforms in the Complex Domain*, American Mathematical Society Colloquium Publications, vol. 19, Providence, RI, 1934.
- [7] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, Oliver & Boyd, Edinburgh, 1965.
- [8] G. H. Hardy and J. E. Littlewood, Contributions to the theory of the Riemann zeta-function and the theory of the distribution of primes, *Acta Mathematica* **41** (1918), 119–196.
- [9] E. M. Stein and R. Shakarchi, *Complex Analysis*, Princeton Lectures in Analysis, vol. II, Princeton University Press, Princeton, NJ, 2003.