A Detailed Analysis of Weak and Strong Convergence in Normed Spaces

by The Collective Unconscious

January 17, 2024

1 Introduction

In this article, we delve into the concepts of weak and strong convergence within normed spaces, illustrating the principles with examples. This exploration is aimed at demystifying the behavior of sequences under these two types of convergence, particularly focusing on how the norms of sequence elements behave in each case.

2 Concepts of Convergence

2.1 Strong Convergence

Strong convergence of a sequence $\{x_n\}$ to x in a normed space occurs when the norm $\|x_n - x\|$ approaches zero as $n \to \infty$. This type of convergence is intuitive as it involves the actual 'distance' between sequence elements and the limit point decreasing in terms of the space's norm.

2.2 Weak Convergence

Weak convergence is subtler. A sequence $\{x_n\}$ converges weakly to x if for every continuous linear functional f, the sequence of scalars $\{f(x_n)\}$ converges to $f(x)$. This means the convergence is determined by the sequence's interaction with linear functionals, not by its norm-distance to the limit.

3 Examples and Analysis

3.1 Oscillating Norms

Example 1. Consider the sequence in l^2 space, $x_n = (1, \frac{1}{2}, \ldots, \frac{1}{n}, 0, \ldots)$. While this sequence converges weakly to the zero vector, the norms exhibit oscillating behavior and do not converge to zero.



Figure 1. Oscillating norms of x_n in l^2 space.

3.2 Bounded Norms, Not Converging

Example 2. In $L^2([0,1])$, the sequence $f_n(x) = \sin(n\pi x)$ converges weakly to 0. However, the L^2 norms of f_n are constantly $\frac{1}{\sqrt{2}}$, not converging to the norm of the limit function (zero).



Figure 2. Constant L^2 norms of f_n in $L^2([0,1])$.

3.3 Unbounded Norms

Example 3. In l^2 space, consider $x_n = (n, 0, \dots)$. This sequence weakly converges to the zero vector, but the norms $\|x_n\| = n$ are unbounded, showing that weak convergence can occur even with unbounded norms.



Figure 3. Unbounded norms in l^2 space.

4 Discussion

These examples illustrate the diversity in behaviors that sequences can exhibit under weak convergence. Unlike strong convergence, where the norm of the difference between sequence elements and the limit decreases, weak convergence does not impose constraints on how the norms behave. This distinction is crucial in understanding infinite-dimensional spaces and the subtleties of functional analysis.

5 Conclusion

Our exploration into weak and strong convergence reveals the intricate nature of these concepts in normed spaces. Understanding these differences is vital for advanced mathematical studies and provides insight into the behavior of sequences in various contexts of functional analysis and related fields.