


This article was downloaded by: [Universite Laval]

On: 26 December 2014, At: 02:09

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Complex Variables, Theory and Application: An International Journal: An International Journal

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gcov19>

A quadratic extremal problem on the dirichlet space

Stephen D. Fisher ^a

^a Department of Mathematics , Northwestern University , Evanston, IL, 60208, USA

Published online: 29 May 2007.

To cite this article: Stephen D. Fisher (1995) A quadratic extremal problem on the dirichlet space , Complex Variables, Theory and Application: An International Journal: An International Journal, 26:4, 367-380

To link to this article: <http://dx.doi.org/10.1080/17476939508814798>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

A Quadratic Extremal Problem on the Dirichlet Space*

STEPHEN D. FISHER

Department of Mathematics, Northwestern University, Evanston, IL 60208, USA

It is shown that there is a unique solution F to the problem

$$\lambda = \sup \left\{ \operatorname{Re} \int_{\Delta} \bar{f} f' dA : \int_{\Delta} |f'|^2 dA \leq 1 \right\}.$$

The function F is entire with a number of special properties. The number λ is the reciprocal of the smallest zero of the 0th Bessel function of the first kind.

AMS No. 30D50, 30D45, 30D10, 30C15

Communicated: R. P. Gilbert

(Received October 2, 1993)

INTRODUCTION

The Dirichlet space, D , on the open unit disc Δ consists of all analytic functions f

$$f(z) = \sum_{k=1}^{\infty} a_k z^k, \quad |z| < 1, \quad f(0) = 0,$$

for which the quantity

$$\int_{\Delta} |f'(z)|^2 dA(z) = \sum_{k=1}^{\infty} k |a_k|^2 =: \|f\|_D^2 \quad (1)$$

is finite. In connection with a generalization of Harnack's inequality, Boris Korenblum [2] has asked how large the quantity

$$\lambda =: \sup_{f \in D} \left\{ \frac{\operatorname{Re}(\sum_1^{\infty} a_k \bar{a}_{k+1})}{\sum_1^{\infty} k |a_k|^2} \right\} \quad (2)$$

is and, if possible, to characterize all functions F which attain the value λ in (2). The expression in the numerator in (2) is not a linear function of f but rather quadratic; hence, the title of this paper.

It is simple to show that

$$\sum_1^{\infty} a_k \bar{a}_{k+1} = \int_{\Delta} \bar{f}(z) f'(z) dA(z) \quad (3)$$

*In memory of Ralph P. Boas, Jr. (1912–1992).

and therefore Korenblum's problem has this alternate form:

$$\lambda = \sup \left\{ \operatorname{Re} \left(\int_{\Delta} \bar{f} f' dA \right) : \|f\|_D \leq 1 \right\} \quad (4)$$

We show here that the extremal problem (2) or (4) has a unique solution F , up to multiplication by a constant; moreover, F is an entire function of exponential type with infinitely many zeros, all in the left half-plane, none of which lie in Δ or on the real axis, except for a first order zero at the origin. Moreover, the number λ is the reciprocal of the smallest positive zero of $J_0(x)$, the 0th Bessel function. Finally,

$$F(z) = C \sum_{n=1}^{\infty} J_n(1/\lambda) z^n$$

where J_n is the n th Bessel function and C is a certain constant.

The conclusions above are proved in Sections 1 and 2; Section 3 contains a number of results which generalize the extremal problem (2).

1. EXISTENCE AND UNIQUENESS

We begin by establishing simple bounds on λ .

PROPOSITION 1 $1/\sqrt{6} < \lambda \leq 1/2$.

Proof Since $2\operatorname{Re}(z\bar{w}) \leq |z|^2 + |w|^2$, we have

$$\begin{aligned} 2\operatorname{Re}(a_1\bar{a}_2 + a_2\bar{a}_3 + \cdots) &\leq |a_1|^2 + |a_2|^2 + |a_2|^2 + |a_3|^2 + \cdots \\ &= |a_1|^2 + 2|a_2|^2 + 2|a_3|^2 + \cdots \\ &\leq \sum k|a_k|^2 \end{aligned}$$

which implies that $\lambda \leq 1/2$. The lower bound is obtained by the specific choices

$$a_2 = \sqrt{\frac{2}{3}}a_1, \quad a_3 = \frac{1}{3}a_1, \quad a_4 = a_5 = \cdots = 0$$

which give

$$\lambda \geq \left\{ \frac{a_1a_2 + a_2a_3}{a_1^2 + 2a_2^2 + 3a_3^2} \right\} = \frac{\sqrt{\frac{2}{3}} + \frac{1}{3}\sqrt{\frac{2}{3}}}{1 + 2 \cdot \frac{2}{3} + 3 \cdot \frac{1}{9}} = \frac{1}{\sqrt{6}}. \quad \blacksquare$$

To prove the existence of a solution, we shall need the following Lemma.

LEMMA Given $\varepsilon > 0$ there is an R_0 , $0 < R_0 < 1$, such that

$$\int_R^1 \int_0^{2\pi} |f(re^{it})|^2 dt r dr < \varepsilon \|f\|_D^2 \quad (5)$$

whenever $R_0 \leq R < 1$ and $f(0) = 0$.

Proof Let $f(z) = \sum_1^\infty c_k z^k$. Then

$$\begin{aligned} \frac{1}{\pi} \int_R^1 \int_0^{2\pi} |f(re^{it})|^2 dt r dr &= \sum_1^\infty |c_k|^2 \frac{1 - R^{2k+2}}{k+1} \\ &= \sum_1^\infty (k|c_k|^2) \left(\frac{1 - R^{2k+2}}{k(k+1)} \right) \\ &\leq \|f\|_D^2 \left(\sum_1^\infty \frac{1 - R^{2k+2}}{k(k+1)} \right). \end{aligned}$$

Here we used the simple inequality

$$k|c_k|^2 \leq \|f\|_D^2, \quad k = 1, 2, \dots$$

The expression

$$\sum_1^\infty \frac{(1 - R^{2k+2})}{k(k+1)}$$

goes to zero monotonically as R increases to 1. We are done. ■

THEOREM 1 *A solution to (4) exists.*

Proof Let $\{f_k\}$ be a sequence with $f_k(0) = 0$,

$$\|f_k\|_D = 1 \quad \text{and} \quad \operatorname{Re} \left(\int_\Delta \bar{f}_k f'_k dA \right) \rightarrow \lambda.$$

We may assume that $\{f_k\}$ converges weakly in the Hilbert space D to a function F , $F(0) = 0$, $\|F\|_D \leq 1$. This implies that $f'_k \rightarrow F'$ uniformly on compact subsets of Δ and also that $f_k \rightarrow F$ uniformly on compact subsets of Δ . Thus,

$$\left| \int_\Delta \bar{f}_k f'_k dA - \int_\Delta \bar{F} F' dA \right| \leq \left| \int_\Delta (\bar{f}_k f'_k - \bar{F} F') dA \right| + \left| \int_\Delta (\bar{F} F' - \bar{F} F') dA \right|.$$

The second term goes to zero since $f_k \rightarrow F$ weakly in D . The first term is no larger than

$$\|f_k\|_D \|f_k - F\|_{L^2} = \|f_k - F\|_{L^2}.$$

The latter goes to zero as $k \rightarrow \infty$ since

$$\begin{aligned} \left(\int_\Delta |f_k - F|^2 dA \right)^{1/2} &\leq \left(\int_{|z| \leq R} |f_k - F|^2 dA \right)^{1/2} + \left(\int_{R < |z| < 1} |f_k - F|^2 dA \right)^{1/2} \\ &\leq \left(\int_{|z| \leq R} |f_k - F|^2 dA \right)^{1/2} + \left(\int_{R < |z| < 1} |f_k|^2 dA \right)^{1/2} \\ &\quad + \left(\int_{R < |z| < 1} |F|^2 dA \right)^{1/2}. \end{aligned}$$

Given $\varepsilon > 0$, we may employ the lemma to choose R so near 1 that the second and third terms on the right-hand side are each less than $\varepsilon/3$. For this R , the first term is less than $\varepsilon/3$ when k is large since $f_k \rightarrow F$ uniformly on compact subsets of Δ . ■

THEOREM 2 *A solution to (4) exists which has positive coefficients; every other solution is a unimodular constant multiple of this solution. Moreover, the coefficients of the solution satisfy the recursion relation*

$$2\lambda k a_k = a_{k+1} + a_{k-1}, \quad k = 1, 2, \dots; \quad a_0 = 0. \quad (6)$$

Proof Let

$$\sum_1^{\infty} a_k z^k$$

be a solution to (4) and consider

$$F(z) = \sum_1^{\infty} |a_k| z^k.$$

Then $\|F\|_D = 1$ and

$$\begin{aligned} \lambda &\geq \operatorname{Re} \left(\int_{\Delta} \bar{F} F' dA \right) = \sum_1^{\infty} |a_k| |a_{k+1}| \\ &\geq \left| \sum_1^{\infty} a_k \bar{a}_{k+1} \right| \geq \operatorname{Re} \left(\sum_1^{\infty} a_k \bar{a}_{k+1} \right) = \lambda. \end{aligned}$$

Hence, F is also a solution and F has non-negative coefficients. Henceforth we shall assume that

$$F(z) = \sum_1^{\infty} a_k z^k, \quad a_k \geq 0$$

is a solution. We shall show that any other solution

$$g(z) = \sum_1^{\infty} b_k z^k$$

is a multiple of F . Let ε be a (small) complex number and set

$$g_{\varepsilon}(z) = g(z) + \varepsilon z^k.$$

Then

$$\lambda \|g_{\varepsilon}\|_D^2 \geq \operatorname{Re} \left\{ \int_{\Delta} \bar{g}_{\varepsilon} g'_{\varepsilon} dA \right\}$$

which yields

$$\lambda (\operatorname{Re} 2k \bar{b}_k \varepsilon + k |\varepsilon|^2) \geq \operatorname{Re} (\varepsilon \bar{b}_{k-1} + \varepsilon \bar{b}_{k+1}).$$

Since this holds for every (small) complex ε , we find that

$$2\lambda k b_k = b_{k-1} + b_{k+1}, \quad k = 1, 2, \dots; \quad b_0 = 0. \quad (6')$$

Applying (6') repeatedly, we determine that b_{k+1} can be expressed in terms of λ and b_1 , namely:

$$b_{k+1} = b_1 P_k(\lambda), \quad k = 1, 2, \dots \quad (7)$$

where P_k is a polynomial of degree k which is an even function if k is even and an odd function if k is odd. The first three P_k are

$$P_1(x) = 2x; \quad P_2(x) = 8x^2 - 1; \quad P_3(x) = 8x(6x^2 - 1).$$

The formulas in (7) show that $b_1 \neq 0$ and hence

$$b_{k+1} = b_1 P_k(\lambda) = \frac{b_1}{a_1} a_1 P_k(\lambda) = \frac{b_1}{a_1} a_{k+1}$$

so that $g = (b_1/a_1)F$, as desired. ■

Remark We shall assume henceforth that F is normalized so that $F'(0) = 1$; that is,

$$F(z) = z + a_2 z^2 + a_3 z^3 + \dots \quad (8)$$

where a_2, a_3, \dots are positive. (That all the a_k are positive follows directly from (6).)

2. PROPERTIES OF THE SOLUTION

PROPOSITION 2 *The coefficients $\{a_k\}$ are monotonically decreasing and satisfy*

$$0 < a_k \leq \left(\frac{1}{\lambda}\right)^{k-1} \frac{1}{k!}, \quad k = 2, 3, \dots \quad (9)$$

Proof First we note that

$$a_2 = 2\lambda \leq 1$$

and then that

$$2a_2 \geq 2\lambda 2a_2 = 1 + a_3 \geq a_2 + a_3$$

so that $1 \geq a_2 \geq a_3$.

Suppose N is some integer greater than or equal to 3 and $a_{N+1} < a_N$. Then

$$2a_N < 2\lambda N a_N = a_{N-1} + a_{N+1} < a_{N-1} + a_N$$

and hence $a_N < a_{N-1}$; now repeat this argument to conclude that $a_{N-1} < a_{N-2}$, etc. Since $N a_N^2 \rightarrow 0$, there are arbitrarily large N with $a_{N+1} < a_N$. Hence, $\{a_k\}$ decreases monotonically.

Since $a_1 \geq a_2 \geq \dots > 0$, we then have

$$2\lambda k a_k = a_{k+1} + a_{k-1} < 2a_{k-1}$$

so that

$$a_k < \frac{1}{\lambda} \cdot \frac{1}{k} a_{k-1}, \quad k = 2, 3, \dots$$

which gives (9) after iteration. ■

COROLLARY *The function F is entire of exponential type.*

THEOREM 3 *λ is the reciprocal of the smallest positive zero of J_0 , the 0th Bessel function of the first kind. Moreover,*

$$F(z) = C \sum_{n=1}^{\infty} J_n(1/\lambda) z^n \quad (10)$$

where C is a constant and J_n is the n th Bessel function of the first kind.

Proof The relation (6) implies that F satisfies the ordinary differential equation

$$2\lambda z^2 F'(z) = (1 + z^2)F(z) - z. \quad (11)$$

Moreover, because F is entire, (11) holds throughout the complex plane. In (11), replace z by $-1/z$ to obtain

$$2\lambda \frac{1}{z^2} F' \left(\frac{-1}{z} \right) = \left(1 + \frac{1}{z^2} \right) F \left(\frac{-1}{z} \right) + \frac{1}{z}. \quad (12)$$

Define $G(z) = F(-1/z)$. Then (12) may be written as

$$2\lambda G'(z) = \left(1 + \frac{1}{z^2} \right) G(z) + \frac{1}{z}$$

and hence

$$2\lambda z^2 G'(z) = (1 + z^2)G(z) + z. \quad (13)$$

Add (11) and (13) to conclude that $h = F + G$ satisfies

$$2\lambda z^2 h'(z) = (1 + z^2)h(z).$$

This equation is easily solved to give

$$h(z) = C \exp \left(\frac{1}{2\lambda} \left(z - \frac{1}{z} \right) \right)$$

where

$$C = h(1) = F(1) + F(-1) = 2(a_2 + a_4 + a_6 + \dots) > 0.$$

That is

$$\begin{aligned} F(z) + F(-1/z) &= C \exp \left(\frac{1}{2\lambda} (z - 1/z) \right) \\ &= C \sum_{n=-\infty}^{\infty} J_n(1/\lambda) z^n. \end{aligned} \quad (14)$$

This gives (10). Note, as well, that the left-hand side of (14) has no constant term. Therefore,

$$J_0(1/\lambda) = 0. \quad (15)$$

The first two zeros of J_0 are

$$\alpha_0 = 2.404826\dots$$

$$\alpha_1 = 3.831706\dots;$$

see [4; page 502]. The reciprocal of α_0 lies between $1/\sqrt{6}$ and $1/2$ while that of α_1 does not. Hence,

$$\lambda = 1/\alpha_0 = 0.4158305\dots \quad (16)$$

By the way, the value of λ gives

$$a_2 \doteq 0.8316, \quad a_3 \doteq 0.3832, \quad a_4 \doteq 0.1247, \quad a_5 \doteq 0.0316$$

and

$$C \doteq 1.912. \quad \blacksquare$$

THEOREM 4 *F has infinitely many zeros, all of which are simple and, except for the zero at the origin, all lie in the left half-plane. None of the zeros lie on the real axis or in the closed unit disc. The zeros in the second quadrant are asymptotic to the curve*

$$Cy = e^{-x/2\lambda}.$$

Finally, if P is the canonical product of the zeros of F , then

$$F(z) = ze^{2\lambda z} P(z). \quad (17)$$

Proof The formula (14) shows that F must have at least one zero away from the origin and that F is entire of order 1. Let z_1, z_2, \dots be the zeros of F ; then cf. [1; Theorem 2.5.18],

$$\sum_m \frac{1}{|z_m|^2} < \infty$$

and P is given by

$$P(z) = \prod_m \left(1 - \frac{z}{z_m}\right) e^{z/z_m}.$$

At this point we do not know whether F has finitely many or infinitely many zeros. The Hadamard factorization theorem [1; page 22] implies that

$$F(z) = ze^{bz} P(z).$$

Hence,

$$\begin{aligned} \frac{zF'(z)}{F(z)} &= 1 + zb - z^2 \sum_m \frac{1}{z_m(z_m - z)} \\ &= 1 + zb - \sum_{j=0}^{\infty} z^{j+3} \sum_m \left(\frac{1}{z_m}\right)^{j+2}, \end{aligned}$$

the latter being valid for $|z|$ small. Comparing power series coefficients, we find that

$$b = a_2 = 2\lambda$$

and, incidentally,

$$\sum_m \left(\frac{1}{z_m} \right)^2 = -2(6\lambda^2 - 1). \quad (18)$$

To see that F has infinitely many zeros, multiply both sides of (14) by z and let $z = x \rightarrow -\infty$. If F had finitely many zeros, then

$$x^2 e^{2\lambda x} P(x) \rightarrow 0$$

and we would conclude that

$$\lim_{x \rightarrow -\infty} P(-1/x) = 0,$$

which is clearly incorrect since $P(0) = 1$.

The differential equation (11) satisfied by F shows that F and F' do not vanish simultaneously; that is, the zeros of F are simple.

To prove that F has no zeros in the right half-plane requires two steps. We first show that any zero in the right half-plane must have modulus no more than $(1+C)/C$; we then show that any zero of F with modulus 4λ or less must lie in the left half-plane. Since $(1+C)/C \doteq 1.52$ and $4\lambda \doteq 1.66$, we will be done.

From (14) (with $z = re^{i\theta}$)

$$\begin{aligned} \left| zF(z) - \frac{F(-1/z)}{(-1/z)} \right| &= rC \exp[(r - 1/r) \cos \theta] \\ &\geq rC \quad \text{if } r \geq 1 \text{ and } \cos \theta \geq 0. \end{aligned}$$

If $F(z) = 0$, then

$$\left| \frac{F(w)}{w} \right| \geq rC, \quad w = -1/z.$$

Since $r \geq 1$, we have $|w| \leq 1$ so

$$\begin{aligned} \left| \frac{F(w)}{w} \right| &= |1 + a_2 w + a_3 w^2 + \dots| \\ &\leq 1 + 2a_2 + 2a_4 + \dots = 1 + C. \end{aligned}$$

Hence, $|z| \leq (1+C)/C$, if z is a zero of F in the right half-plane.

As to the second step, we need the following result.

PROPOSITION 2 [3; page 129] *If A_0, A_1, \dots is a sequence of positive numbers satisfying*

$$\sum_0^\infty A_k < \infty \quad \text{and} \quad A_{k+1} - 2A_k + A_{k+1} \geq 0 \quad (\text{all } k)$$

then

$$\frac{1}{2}A_0 + \sum_{k=1}^{\infty} A_k \cos kx \geq 0 \quad \text{for all } x.$$

We consider

$$g(z) =: \frac{F(z) - z}{z^2} = a_2 + a_3z + a_4z^2 + \dots.$$

We shall show that (with $R = 4\lambda$) the numbers

$$A_0 = 2a_2, \quad A_k = a_{k+2}R^k, \quad k = 1, 2, \dots \quad (20)$$

satisfy the hypotheses of Proposition 2. Once this is proved, the Proposition implies that

$$\operatorname{Re}(g(\operatorname{Re}^{ix})) \geq 0, \quad 0 \leq x \leq 2\pi$$

and hence $\operatorname{Re}(g(z)) > 0$ if $|z| < R = 4\lambda$. From this we see that if $F(z) = 0$ and $|z| < 4\lambda$, then

$$0 < \operatorname{Re} g(z) = -\operatorname{Re}(1/z)$$

and hence z lies in the left half-plane. Thus, to finish we need to establish (20).

First:

$$A_0 - 2A_1 + A_2 = 2a_2 - 2a_3R + a_4R^2 = 12\lambda(1 - 8\lambda^2)^2 > 0.$$

Next, for $k \geq 4$

$$\begin{aligned} & a_{k+1}R^{k-1} - 2a_kR^{k-2} + a_{k-1}R^{k-3} \\ &= R^{k-3}\{a_{k+1}R^2 - 2a_kR + a_{k-1}\} \\ &= R^{k-3}\{(R^2 - 1)a_{k+1} + (2\lambda k - 2R)a_k\} \quad (\text{from (6)}) \\ &> 0 \quad \text{since } \lambda k \geq 4\lambda = R \quad \text{and } R > 1. \end{aligned}$$

To show that F has no zeros in $0 < |z| < 1$, we use Proposition 2 for the function

$$f(z) =: \frac{F(z)}{z} = 1 + a_2z + a_3z^2 + \dots.$$

Let $a_0 = 2$, $A_k = a_{k+1}$, $k = 1, 2, \dots$. Then

$$A_0 - 2A_1 + A_2 = 2 - 2a_2 + a_3 > 0 \quad \text{since } a_2 = 2\lambda < 1.$$

Further, for $k \geq 2$

$$\begin{aligned} A_{k+1} - 2A_k + A_{k-1} &= a_{k+2} - 2a_{k+1} + a_k \\ &= (2\lambda(k+1) - 2)a_{k+1} > 0. \end{aligned}$$

Hence,

$$\operatorname{Re}\left(\frac{F(z)}{z}\right) > 0 \quad \text{if } |z| < 1. \quad (21)$$

To see that $F(x) \neq 0$ if $-\infty < x < 0$, suppose to the contrary that

$$F(c) = 0 \quad \text{and} \quad F(x) < 0 \quad \text{and} \quad c < x < 0.$$

From (11), we find $2\lambda c^2 F'(c) = -c > 0$ and so $F(x) > 0$ if $c < x < c + \delta$. This contradiction establishes that $F(x) < 0$ if $-\infty < x < 0$.

The asymptotic behavior of $\{z_m\}$ follows from (14). Let $\{z_m\}$ be the zeros of F in the second quadrant. Then

$$|z_m| |F(-1/z_m)| = C |z_m| e^{1/2\lambda(x_m - \operatorname{Re}(1/z_m))}.$$

The left-hand side of this expression approaches 1 as $m \rightarrow \infty$. Since the right-hand side exceeds

$$C |z_m| e^{(1/2\lambda)x_m}$$

it follows that $x_m \rightarrow -\infty$ as $m \rightarrow \infty$. This, in turn, implies that

$$x_m e^{(1/2\lambda)x_m} \rightarrow 0 \quad \text{as} \quad m \rightarrow \infty.$$

Therefore,

$$\frac{1}{C} = \lim_{m \rightarrow \infty} y_m e^{(1/2\lambda)x_m},$$

which is the desired conclusion.

3. EXTENSIONS

In this section we take up two extensions of the extremal problem (2). The first is to replace the Hilbert space D with the more general Hilbert space \mathcal{H} determined by the condition

$$\sum_{k=0}^{\infty} \varepsilon_k |a_k|^2 < \infty \quad (22)$$

where $\varepsilon_0, \varepsilon_1, \dots$ are given positive numbers. Here we determine a necessary and sufficient condition that the corresponding functional

$$\phi(a_0, a_1, \dots) =: \frac{\operatorname{Re}(\sum_{k=0}^{\infty} a_k \bar{a}_{k+1})}{\sum_{k=0}^{\infty} \varepsilon_k |a_k|^2} \quad (23)$$

be bounded; that is, that $\phi(a_0, a_1, \dots) \leq A < \infty$ for all sequences $\{a_k\}$ satisfying (22). We also determine a sufficient condition that there is a solution in \mathcal{H} to the extremal problem

$$\sup\{\phi(a_0, a_1, \dots) : \{a_k\} \in \mathcal{H}\}. \quad (24)$$

This condition is general enough to provide an alternate proof of Theorem 1.

The other extension of (2) which we take up is to replace the expression $\sum a_k \bar{a}_{k+1}$ in (2) by $\sum a_k \bar{a}_{k+m}$ where m is some positive integer. Here we shall find that the extremal problem

$$\mu =: \sup_{f \in D} \frac{\operatorname{Re}(\sum_{k=1}^{\infty} a_k \bar{a}_{k+m})}{\sum_{k=1}^{\infty} k |a_k|^2} \quad (25)$$

is easily solved, given that we have already solved (24), and its solution is directly connected to that of (24).

We note first that if (24) has a solution, then there is a sequence $\{b_k\}$ with

$$\sum_{k=0}^{\infty} \varepsilon_k |b_k|^2 = 1$$

and

$$\phi(b_0, b_1, \dots) = \sup\{\phi(a_0, a_1, \dots) : \{a_k\} \in \mathcal{H}\}.$$

This, in turn, obviously implies that ϕ is bounded.

THEOREM 5

(i) ϕ is bounded if and only if

$$\liminf_{k \rightarrow \infty} (\varepsilon_k \varepsilon_{k+1}) > 0 \quad (26)$$

(ii) ϕ has a solution if

$$\lim_{k \rightarrow \infty} \varepsilon_k \varepsilon_{k+1} = +\infty. \quad (27)$$

Proof (i) Suppose that $\varepsilon_{k+1} \varepsilon_k \geq \delta^2$ for all k . Then

$$2\delta \operatorname{Re}(a_k \bar{a}_{k+1}) \leq 2\operatorname{Re} \sqrt{\varepsilon_k \varepsilon_{k+1}} a_k \bar{a}_{k+1} \leq \varepsilon_k |a_k|^2 + \varepsilon_{k+1} |a_{k+1}|^2$$

and hence

$$\phi(a_0, a_1, \dots) \leq 2 \frac{1}{2\delta} = \frac{1}{\delta}.$$

Conversely, if ϕ is bounded, say $\phi(a_0, a_1, \dots) \leq A$, and

$$f(z) = z^k + \sqrt{\varepsilon_k / \varepsilon_{k+1}} z^{k+1},$$

then

$$\sqrt{\varepsilon_k / \varepsilon_{k+1}} \leq A(\varepsilon_k + \varepsilon_{k+1}) = 2A\varepsilon_k$$

which gives (26).

(ii) Define

$$\mu = \sup \left\{ \phi(a_0, a_1, \dots) : \sum \varepsilon_k |a_k|^2 < \infty \right\}$$

and assume that

$$\lim \varepsilon_k \varepsilon_{k+1} = +\infty.$$

Let $\{a_k^{(n)}\}$ be a maximizing sequence:

$$\sum_k \varepsilon_k |a_k^{(n)}|^2 = 1 \quad \text{and} \quad \lim_n \left(\operatorname{Re} \sum_k a_k^{(n)} \bar{a}_{k+1}^{(n)} \right) = \mu.$$

We may assume with no loss that $\{a_k^{(n)}\} \rightarrow \{a_k\}$, as $n \rightarrow \infty$, weakly in the Hilbert space

$$\mathcal{H} = \left\{ (a_0, a_1, \dots) : \sum_k \varepsilon_k |a_k|^2 < \infty \right\}.$$

In particular, $a_k^{(n)} \rightarrow a_k$ as $n \rightarrow \infty$ for each k and $\sum_k \varepsilon_k |a_k|^2 \leq 1$. We then have

$$\begin{aligned} \left| \sum_k a_k \bar{a}_{k+1} - \sum_k a_k^{(n)} \bar{a}_{k+1}^{(n)} \right| &\leq \left| \sum_k (a_k - a_k^{(n)}) \bar{a}_{k+1} \right| + \left| \sum_k a_k^{(n)} (\bar{a}_{k+1} - \bar{a}_{k+1}^{(n)}) \right| \\ &= I + II. \end{aligned}$$

We shall assume (with no loss of generality) that $\varepsilon_k \varepsilon_{k+1} \geq 1$ for all k . Then

$$\begin{aligned} I &\leq \sum_{k=1}^{\infty} |a_k - a_k^{(n)}| |a_{k+1}| \\ &\leq \sum_1^N |a_k - a_k^{(n)}| |a_{k+1}| + \sum_{N+1}^{\infty} \sqrt{\varepsilon_k \varepsilon_{k+1}} |a_k - a_k^{(n)}| |a_{k+1}| \\ &\leq \sum_1^N |a_k - a_k^{(n)}| |a_{k+1}| + \left(\sum_{N+1}^{\infty} \varepsilon_k |a_k - a_k^{(n)}|^2 \right)^{1/2} \left(\sum_{N+1}^{\infty} \varepsilon_{k+1} |a_{k+1}|^2 \right)^{1/2} \\ &\leq \sum_1^N |a_k - a_k^{(n)}| |a_{k+1}| + 2 \left(\sum_{N+1}^{\infty} \varepsilon_{k+1} |a_{k+1}|^2 \right)^{1/2}. \end{aligned}$$

Both of these terms are small: the second for N large and the first for $n \geq n_0$, once N is fixed. Next,

$$\begin{aligned} II &\leq \sum_1^{\infty} |a_k^{(n)}| |a_{k+1} - a_{k+1}^{(n)}| \\ &\leq \left(\sum_1^{\infty} \varepsilon_k |a_k^{(n)}|^2 \right)^{1/2} \left(\sum_1^{\infty} \frac{1}{\varepsilon_k} |a_{k+1} - a_{k+1}^{(n)}|^2 \right)^{1/2} \\ &\leq \left(\sum_1^N \frac{1}{\varepsilon_k} |a_{k+1} - a_{k+1}^{(n)}|^2 \right)^{1/2} + \left(\sum_{N+1}^{\infty} \left(\frac{1}{\varepsilon_k \varepsilon_{k+1}} \right) \varepsilon_{k+1} |a_{k+1} - a_{k+1}^{(n)}|^2 \right)^{1/2}. \end{aligned}$$

The second term is less than

$$2 \max_{k \geq N+1} \left(\frac{1}{\varepsilon_k \varepsilon_{k+1}} \right)^{1/2} < \varepsilon \quad \text{if } N \text{ is large}$$

and the first term goes to zero as $n \rightarrow \infty$ for N fixed. ■

THEOREM 6 *The number μ defined in (25) is equal to number γ defined by*

$$\gamma =: \sup \left\{ \frac{\operatorname{Re} \left(\sum_0^\infty a_k \bar{a}_{k+1} \right)}{\sum_0^\infty (mk+1) |a_k|^2} \right\}. \quad (28)$$

Proof Let $\{a_0, a_1, \dots\}$ be any sequence for which

$$\sum_0^\infty (mk+1) |a_k|^2 < \infty$$

and define

$$b_j = \begin{cases} a_k & \text{if } j = km+1, \quad k = 0, 1, \dots \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\mu \geq \frac{\operatorname{Re} \left(\sum j |b_j| \bar{b}_{j+m} \right)}{\sum j |b_j|^2} = \frac{\operatorname{Re} \left(\sum_0^\infty a_k \bar{a}_{k+1} \right)}{\sum_0^\infty (mk+1) |a_k|^2}$$

so that $\mu \geq \gamma$.

To prove the reverse inequality, we note that

$$\begin{aligned} \frac{\operatorname{Re} \left(\sum_{j=1}^\infty j |b_j| \bar{b}_{j+m} \right)}{\sum_{j=1}^\infty j |b_j|^2} &= \frac{\sum_{r=1}^m \operatorname{Re} \left(\sum_{k=0}^\infty b_{km+r} \bar{b}_{(k+1)m+r} \right)}{\sum_{r=1}^m \sum_{k=0}^\infty (km+r) |b_{km+r}|^2} \\ &\leq \frac{\sum_{r=1}^m \operatorname{Re} \left(\sum_{k=0}^\infty b_{km+r} \bar{b}_{(k+1)m+r} \right)}{\sum_{r=1}^m \sum_{k=0}^\infty (km+1) |b_{km+r}|^2} \\ &\leq \gamma \end{aligned} \quad \blacksquare$$

EXAMPLE *The extremal problem (24) has no solution when $\varepsilon_k = 1$ for all k ; that is, when $\mathcal{H} = H^2$, the Hardy space.*

Let μ be the supremum in (24). To see that $\mu = 1$, note that the Schwarz inequality gives

$$\phi(a_0, a_1, \dots) \leq \frac{(\sum_0^\infty |a_k|^2)^{1/2} (\sum_1^\infty |a_k|^2)^{1/2}}{\sum_0^\infty |a_k|^2} \leq 1 \quad (29)$$

while the choice $a_k = 1, k = 0, \dots, N, a_k = 0$ for $k \geq N+1$, gives

$$\mu \geq \frac{N}{N+1}. \quad (30)$$

(29) and (30) give $\mu = 1$. If $F(z) = \sum_0^\infty a_k z^k$ were a solution to (24), then it would necessarily follow that

$$2\mu\varepsilon_k a_k = a_{k+1} + a_{k-1}, \quad k = 0, 1, \dots; \quad a_{-1} = 0.$$

That is

$$2a_k = a_{k+1} + a_{k-1}.$$

Hence

$$a_1 = 2a_0$$

$$a_2 = 2a_1 - a_0 = 3a_0$$

$$a_3 = 2a_2 - a_1 = 4a_0$$

etc.

which yields $a_k = (k + 1)a_0$ and so

$$F(z) = a_0 \sum_{k=0}^{\infty} (k + 1)z^k = a_0 \frac{1}{(1 - z)^2}.$$

But this F does not lie in H^2 (unless $a_0 = 0$).

References

- [1] R. P. Boas, Jr., *Entire Functions*, Academic Press, New York, 1954.
- [2] B. Korenblum, personal communication.
- [3] G. G. Lorentz, *Approximation of Functions*, Holt, Rinehart and Winston, New York, 1966.
- [4] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd ed., Cambridge University Press, 1966.