


1. for all $x_0 + y_0 \in W$, $x_0 \in U$, $y_0 \in V$

since U is open, $\exists \varepsilon > 0$ s.t. $B(x_0, \varepsilon) \subseteq U$

since $\forall p \in B(x_0 + y_0, \varepsilon)$ can be expressed as $p = x_1 + y_0$ where $x_1 \in B(x_0, \varepsilon) \subseteq U$, $y_0 \in V$

then $B(x_0 + y_0, \varepsilon) \subseteq W$

thus W is open.

2. for all $(x, y) \in \mathbb{R}$, set $F(x, y) = f(x)g(y)$

since $f, g: \mathbb{R} \rightarrow \mathbb{R}$ is continuous,

$\forall x \in \mathbb{R} (\forall \varepsilon > 0, \exists \delta_x > 0 \text{ s.t. } \forall |x' - x| < \delta_x, |f(x') - f(x)| < \varepsilon)$

$\forall y \in \mathbb{R} (\forall \varepsilon > 0, \exists \delta_y > 0, \text{ s.t. } \forall |y' - y| < \delta_y, |g(y') - g(y)| < \varepsilon)$

then

$\forall (x, y) \in \mathbb{R}^2 (\forall \varepsilon > 0 \exists \delta = \min(\delta_x, \delta_y) > 0, \text{ s.t. } \forall \|(x', y') - (x, y)\| < \delta,$

$$|F(x', y') - F(x, y)| = |f(x')g(y') - f(x)g(y)|$$

$$= |f(x')g(y') - f(x')g(y) + f(x')g(y) - f(x)g(y)|$$


$$\leq |f(x')| |g(y') - g(y)| + |g(y)| |f(x') - f(x)|$$

$$< \varepsilon (|f(x')| + |g(y)|) \quad \text{where } |f(x')|, |g(y)| \text{ are curb by a maximum since it's continuous.}$$

thus $F(x, y)$ is continuous, $(-\infty, 1)$ is open

thus $R = F^{-1}((-\infty, 1))$ is open.

$$3. Q = \underbrace{\{(x,y) \mid xy \geq 0\}}_A \cap \underbrace{\{(x,y) \mid |x|+|y| \leq 4\}}_B \cap \underbrace{\{(x,y) \mid x^2+y^2 \geq 1\}}_C$$



$$A^c = \{(x,y) \mid xy < 0\}$$

for $(x_0, y_0) \in A^c$

$$(x_0, y_0) \in B \quad [(x_0, y_0), r_0] \in A^c$$

where $r_0 = \min(x_0, y_0)$

$$B^c = \{(x,y) \mid |x|+|y| > 4\}$$

for $(x_1, y_1) \in B^c$

$$(x_1, y_1) \in B \quad [(x_1, y_1), r_1] \in B^c$$

where $r_1 = \min\left(\frac{|x_1+y_1+4|}{\sqrt{2}}, \frac{|x_1+y_1-4|}{\sqrt{2}}, \frac{|x_1-y_1+4|}{\sqrt{2}}, \frac{|x_1-y_1-4|}{\sqrt{2}}\right)$

$$C^c = \{(x,y) \mid x^2+y^2 < 1\} \text{ is a open ball}$$

thus A, B, C is close, so is $Q = A \cap B \cap C$

$$\partial Q \cap \{(x, 0) \mid x \in \mathbb{R}\} = [0, 4] \times \{0\}$$

4. S is not open. consider $(x,y) \in [0,1) \times \{0\}$, no neighbour is included.

S is closed. S is the union of countable many 2-cells.

S is not bounded. since as $x \rightarrow \infty$, y can $\rightarrow \infty$, there's no upper bound.

5. (a) $D^o = \mathbb{R}^d / \partial D$, suppose $t \in D$

then all $z \in \{t+x \mid x \in \mathbb{R}^d / \partial D\} = S$ can be expressed.

consider $z_0 \in \{t+x \mid x \in \partial D\} = \partial S$

since O is open, $\exists r > 0$, $B(t, r) \subseteq O$

$\{t+x \mid x \in \partial D\}$ is the boundary of $\{t+x \mid x \in \mathbb{R}^d / \partial D\}$

thus $\exists (a_n)_n \in S$, $\forall \varepsilon > 0$, $\exists N$, $\forall n \geq N$, $\|a_n - z_0\| < \varepsilon$

choose the smallest $n' \geq N$ for $\varepsilon = r$, set $a_{n'} = b \in S$

$$\|b - z_0\| < r \quad b = t + x_0 \quad z_0 \in \{x_0 + t' \mid t' \in B(t, r)\}$$

thus z_0 can be expressed as $z = x + y$

1b) use the outcome of (a) where set $A = O = D$

6. The statement equivalent to:

If $A \cap U = \emptyset$, then U is \emptyset or not open.

U is \emptyset is obvious.

U is not open:

If U is open & $A \cap U = \emptyset$, contradict to 5(a)

$$1. (a) \quad H_0: \theta = 1 \quad H_1: \theta = \theta, < 1$$

$$L(\theta_0; X) = \left(\prod_{i=1}^n 4x_i^3 \right) \cdot \exp\left(-\sum_{i=1}^n x_i^4\right)$$

$$L(\theta_1; X) = \left(\prod_{i=1}^n \frac{4x_i^3}{\theta_1} \right) \cdot \exp\left(-\frac{1}{\theta_1} \sum_{i=1}^n x_i^4\right)$$

$$C = \left\{ X \mid \frac{\exp\left[-\left(\frac{1}{\theta_1} - 1\right) \frac{\sum_{i=1}^n x_i^4}{\theta_1^n}\right]}{\theta_1^n} > k \right\}$$

$$\text{where } \mathbb{P}(X \in C \mid H_0) = \alpha$$

$$\mathbb{P}\left(\exp\left[-\left(\frac{1}{\theta_1} - 1\right) \frac{\chi_{2n}^2}{2}\right] > k \theta_1^n\right) = \mathbb{P}\left(\chi_{2n}^2 < \frac{2\theta_1}{\theta_1 - 1} \ln(k \theta_1^n)\right) = \alpha$$

$$C = \left\{ X \mid \sum_{i=1}^n x_i^4 \leq \frac{1}{2} \chi_{2n, \alpha}^2 \right\}$$

$$(b) \quad p\text{-value} = \mathbb{P}(\chi_{10}^2 \leq 3.589) = 0.042$$

$$(c) \quad \mathbb{P}(\chi_{10}^2 \leq 2k_1) = \frac{1}{2}\alpha \quad \mathbb{P}(\chi_{10}^2 \geq 2k_2) = \frac{1}{2}\alpha$$

$$k_1 = 1.623$$

$$k_2 = 10.24$$

$$(d) \quad 1.623 < \sum_{i=1}^5 x_i^4 = 1.794 < 10.24$$

thus retain. (b) is rejected since it's one-tailed.

$$2. (a) \quad l(\lambda; X) = -n\lambda + (\sum x_i) \ln \lambda - \sum \ln(x_i)!$$

$$\frac{\partial}{\partial \lambda} l(\lambda; X) = -n + \frac{1}{\lambda} \sum x_i$$

$$\hat{\lambda} = \arg \max_{\lambda} l = \frac{1}{n} \sum x_i = 5.05$$

$$X_i \sim \text{Poisson}(\lambda)$$

for large n

$$\hat{\lambda} = \frac{1}{n} \sum X_i \sim N\left(\lambda, \frac{\lambda}{n}\right)$$

$$\left| \frac{\hat{\lambda} - \lambda}{\sqrt{\frac{\lambda}{n}}} \right| < 1.96$$

$$95\% \text{ CI} : (4.11, 6.13)$$

$$(b) \quad \sum X_i = 101$$

$$\sigma = \sqrt{80} \quad \mu = 80$$

$$p = 2 \left[1 - \Phi \left(\frac{101 - 80}{\sqrt{80}} \right) \right] = 0.0188 \quad \text{refuse.}$$

$$(c) \quad S_w = 101 \quad S_s = 63$$

$$E_w = E_s = 81.5$$

$$\sum \frac{(O_k - E_k)^2}{E_k} = 9.33 \sim \chi^2_1$$

$$\text{use } \alpha = 0.01 \quad 9.33 > 3.84 \quad \text{unreasonable.}$$

$$(d) \quad \text{under } H_0: \lambda = \mu = \theta$$

$$l(\theta, \theta) = -2n\lambda_0 + (S_w + S_s) \ln \theta - \ln \prod x_i \prod y_i$$

$$\frac{\partial}{\partial \theta} l = -2n + (S_w + S_s)/\theta = 0$$

$$\hat{\theta} = 4.074$$

under $H_1: \lambda \neq \mu$

$$L[\lambda, \mu] = -n(\lambda + \mu) + S_w \ln \lambda + S_r \ln \mu - \ln \prod_{i=1}^n x_i \prod_{j=1}^n y_j$$

$$\begin{cases} \frac{\partial}{\partial \lambda} L = -n + S_w / \lambda = 0 \\ \frac{\partial}{\partial \mu} L = -n + S_r / \mu = 0 \end{cases} \Rightarrow \begin{cases} \hat{\lambda} = 5.05 \\ \hat{\mu} = 3.1 \end{cases}$$

$$2 [L(\hat{\lambda}, \hat{\mu}) - L(\hat{\theta}, \hat{\theta})] = 2 [S_w \ln$$