

---

---

---

---

---



对已知函数有以下性质

在  $x_1 < x_2 < x_3 \dots < x_n$

$$f(x_1) = a_{1,1}$$

$$f(x_n) = a_{1,n}$$

$$\vdots$$

$$f^{(b_1)}(x_1) = a_{b_1,1}$$

$$\vdots$$

$$f^{(b_n)}(x_n) = a_{b_n,n}$$

可构建  $F(x) = f(x) - \sum_{i=0}^k a_i x^i \quad (k = \sum_{i=1}^n b_i)$

e.g.

1.  $f(x) \in C[0,1] \quad f(x) \in D^r(0,1)$

$$f(0) = 1, \quad f(\frac{1}{2}) = 2, \quad f(1) = 2$$

证:  $\exists \xi \in (0,1) \text{ s.t. } f'(\xi) = -4$

Proof. 构造  $F(x)$

$$F(x) = \begin{vmatrix} f(x) & x^2 & x & 1 \\ f(0) & 0 & 0 & 1 \\ f(\frac{1}{2}) & \frac{1}{4} & \frac{1}{2} & 1 \\ f(1) & 1 & 1 & 1 \end{vmatrix}$$

$$F(0) = F(\frac{1}{2}) = F(1) = 0$$

罗尔

$$\exists x_1 \in (0, \frac{1}{2}), x_2 \in (\frac{1}{2}, 1) \text{ s.t. } f'(x_1) = f'(x_2) = 0$$

$$F''(x) = \begin{vmatrix} f''(x) & 2 & 0 & 0 \\ f''(0) & 0 & 0 & 1 \\ f''(\frac{1}{2}) & \frac{1}{2} & \frac{1}{2} & 1 \\ f''(1) & 1 & 1 & 1 \end{vmatrix} = -\frac{1}{4}f''(x) + 1$$

Q.E.D.

2.  $f(x) \in D^3[0,1]$ ,  $f(0) = -1$ ,  $f'(0) = 0$ ,  $f(1) = 0$   
 We  $\exists \xi \in (0,1)$  s.t.  $f(x) = -1 + x^2 + \frac{x^2(x-1)}{6} f'''(\xi)$

Proof.

$$F(x) = \begin{vmatrix} f(x) & x^3 & x^2 & x & 1 \\ f(0) & 0 & 0 & 0 & 1 \\ f'(0) & 0 & 0 & 1 & 0 \\ f(1) & 1 & 1 & 1 & 1 \\ f(t) & t^3 & t^2 & t & 1 \end{vmatrix}$$

$$F(0) = F(1) = F(t) = 0$$

$$\exists x_1, x_2, \quad F'(x_1) = F'(x_2) = 0$$

$$\exists y_1, y_2, \quad F''(y_1) = F''(y_2) = 0$$

$$\exists \xi, \quad F'''(\xi) = 0$$

$$F'''(x) = f'''(x) t^2(t-1) - 6(f(t) + 1 - t^2)$$

$$0 = F'''(\xi) = f'''(\xi) t^2(t-1) - 6(f(t) + 1 - t^2)$$

Q.E.D.