

Recitation Session 5 Solutions

1. *CLT and LLN.*

CLT says,

- Distribution of the total number of successes looks like a normal distribution centered at np , whose standard deviation is a constant multiple of \sqrt{n} .
- Distribution of the proportion of successes looks like a normal distribution centered at p , whose standard deviation is a constant multiple of $1/\sqrt{n}$.

LLN says,

- The proportion of successes is close to p .

2. *Random walk.* Let X_i be the i th step. $E(X_i) = (-1)\frac{1}{3} + (0)\frac{1}{3} + (+1)\frac{1}{3} = 0$, and $\text{Var}(X_i) = E(X^2) - E(X)^2 = (-1)^2\frac{1}{3} + (0)^2\frac{1}{3} + (+1)^2\frac{1}{3} = 2/3$ hence its standard deviation is 0.8165. The position at the 10,000th step is given by $S_{10,000} = \sum_{i=1}^{10,000} X_i$. By the normal approximation $S_{10,000}$ is approximately normal with mean zero and standard deviation 81.65. Then $P(S_{10,000} > 100) = P(S_{10,000}/81.65 > 100/81.65) = Q(100/81.65) \approx 11\%$.
3. *Biased die.* Since we count number of sixes let X_i be 1 if the i th roll is a six and 0 otherwise. Then $\sum X_i = 180.000$ is observed, so the estimator is $\hat{p} = 0.18$, $n = 10^6$, $\sigma = \sqrt{np(1-p)}$. Now plug them into equation 38 in the lecture notes. This interval depends on p , however we can slightly enlarge the interval to a more conservative value so the confidence is not broken but is independent of p . This is done by the bound $p(1-p) < 1/4$. We arrive at $(0.178, 0.182)$.
4. *Biased coin.* Note that $Q^{-1}(0.5) \approx 2.575$ so we have $\frac{\sqrt{p(1-p)}}{\sqrt{n}} 2.575 = 0.1$ gives $n = 25.75^2 p(1-p) < 25.75^2 (0.5)^2 = 165.77$ To decrease the length of the interval 10 times, increase the number of samples by 100 times: 16,577.
5. *Two surveys.* The larger sample will have a shorter interval. Since each interval scales with $1/\sqrt{n}$ the shorter interval will be $\sqrt{\frac{350}{1000}} = 0.59$ times the size of the longer one.