

section 2.6 , Linear Transformations pt. 2

→ Cross Product :

same as cross product of vectors

$$v \times w = \det \begin{bmatrix} \hat{i} & x_1 & x_2 \\ \hat{j} & y_1 & y_2 \\ \hat{k} & z_1 & z_2 \end{bmatrix}$$

$$\Rightarrow \|v \times w\|^2 = \|v\|^2 \|w\|^2 - (v \cdot w)^2 = \|v\|^2 \|w\|^2 \sin^2 \theta$$

as a consequence we learn Area of parallelogram spanned by v & w is...

$$\begin{aligned} \text{Area parallelogram} &= \|v \times w\| \\ &= \|v\| \|w\| \sin \theta \end{aligned}$$

e.g find area of parallelogram spanned by

$$\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \text{ \& } \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix}$$

$$\text{let } v = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \quad w = \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}$$

$$\text{Area } \square = \|v\| \|w\| \sin \theta$$

$$= \|v \times w\|$$

$$= \left\| \det \begin{pmatrix} \hat{i} & 1 & 2 \\ \hat{j} & -1 & 3 \\ \hat{k} & 2 & -1 \end{pmatrix} \right\|$$

$$\begin{aligned}
&= \left\| \hat{i} \det \begin{pmatrix} -1 & 3 \\ 2 & -1 \end{pmatrix} - \hat{j} \det \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix} + \hat{k} \det \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} \right\| \\
&= \left\| \hat{i}(1-6) - \hat{j}(-1-4) + \hat{k}(3+2) \right\| \\
&= \left\| -5\hat{i} + 5\hat{j} + 5\hat{k} \right\| \\
&= \left\| (-5, 5, 5) \right\| \\
&= \sqrt{(-5)^2 + 5^2 + 5^2} \\
&= \sqrt{75}
\end{aligned}$$

Recall :

for $n \times n$ matrix A , vector v is eigenvector if...

$$Av = \lambda v, \quad \lambda \text{ is eigenvalue}$$

let polynomial $C_A(x) = \det(xI_n - A)$ be the characteristic polynomial

$\Rightarrow \lambda$ is eigenvalue of A if it is a root of $C_A(\lambda) = 0$

$\Rightarrow \lambda$ -eigenvectors are non-trivial solutions to homogeneous system $(\lambda I_n - A)x = 0$
(basic eigenvectors)

we say A is diagonalizable if there exist invertible P & diagonal matrix D s.t.

$$A = PDP^{-1} \Rightarrow D = P^{-1}AP$$

$$D := \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} \longleftrightarrow P := (v_1 \ v_2 \ \dots \ v_n)$$

e.g.

check if $A = \begin{pmatrix} 2 & 4 \\ 0 & -5 \end{pmatrix}$ is diagonalizable.
find invert. P and diagonal D s.t. $A = PDP^{-1}$

$$\begin{aligned} C_A(x) &= \det(xI_2 - A) \\ &= \det\left(\begin{bmatrix} x & 0 \\ 0 & x \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 0 & -5 \end{bmatrix}\right) \\ &= \det\begin{bmatrix} x-2 & -4 \\ 0 & x+5 \end{bmatrix} \\ &= (x-2)(x+5) - 0 \\ &= x^2 + 5x - 2x - 10 \\ &= x^2 + 3x - 10 \\ &= (x+5)(x-2) \end{aligned}$$

$$\lambda_1 = -5, \lambda_2 = 2$$

$$(\lambda_1 I_n - A)x = 0$$

$$\left(\begin{bmatrix} -5 & 0 \\ 0 & -5 \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 0 & -5 \end{bmatrix}\right)x = 0$$

$$\begin{pmatrix} -7 & -4 \\ 0 & 0 \end{pmatrix} x = 0$$

$$\left(\begin{array}{cc|c} -7 & -4 & 0 \\ 0 & 0 & 0 \end{array}\right)$$

$$x_1 = 7$$

$$x_2 = 4$$

$$(\lambda_2 I_n - A)x = 0$$

$$\left(\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 0 & -5 \end{bmatrix}\right)x = 0$$

$$\begin{pmatrix} 0 & -4 \\ 0 & 7 \end{pmatrix} x = 0$$

$$\left(\begin{array}{cc|c} 0 & -4 & 0 \\ 0 & 7 & 0 \end{array}\right)$$

$$0$$

$$0$$

$$x_2 = 0$$

$$\Rightarrow v_1 = \begin{pmatrix} 7 \\ 4 \end{pmatrix}$$

$$\Rightarrow v_2 = x_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$P(2, 5)$$

$$D = \begin{bmatrix} 7 & 1 \\ 4 & 0 \end{bmatrix}$$

→ we already went over linear transformations & standard matrix

e.g. let $T\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x - 2y + 7z \\ 2x - 4y + z \end{pmatrix}$, compute $T(e_1)$

$T(e_2)$ $T(e_3)$. Find A_T , the standard matrix for T

$$T(e_1) = \dots = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$T(e_2) = \dots = \begin{pmatrix} -2 \\ -4 \end{pmatrix}$$

\Rightarrow

$$A_T = \begin{bmatrix} 1 & -2 & 7 \\ 2 & -4 & 1 \end{bmatrix}$$

$$T(e_3) = \dots = \begin{pmatrix} 7 \\ 1 \end{pmatrix}$$

Tutorial Questions:

* 1. visualize each transformation. —

$\begin{bmatrix} 0 \\ 0 \end{bmatrix}$ $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ are your base vertices

a) $T\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3x + y \\ x + 2y \end{pmatrix}$

$$\Rightarrow T\begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad T\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \end{pmatrix} \quad T\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad T\begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$



2. sketch the transformation $T\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3x - 2y \\ x + 4y \end{pmatrix}$ to unit square

find the resulting area of parallelogram

sketch w/ base vertices

A_T will be used to find area

$$A_T = \begin{pmatrix} 3 & -2 \\ 1 & 4 \end{pmatrix}$$