

## Notes on the function gsw\_CT\_second\_derivatives(SA,pt)

This function, **gsw\_CT\_second\_derivatives**(SA,pt), evaluates the second derivatives of  $\tilde{\Theta}(S_A, \theta)$ , namely

$$\tilde{\Theta}_{S_A S_A}, \quad \tilde{\Theta}_{\theta\theta} \quad \text{and} \quad \tilde{\Theta}_{S_A \theta}. \quad (1)$$

These second derivatives are found by mathematically differentiating Eqns. (A.12.3) of the TEOS-10 Manual, repeated here,

$$\Theta_{\theta}|_{S_A} = c_p(S_A, \theta, 0)/c_p^0, \quad \Theta_{S_A}|_{\theta} = [\mu(S_A, \theta, 0) - (T_0 + \theta)\mu_T(S_A, \theta, 0)]/c_p^0. \quad (A.12.3)$$

The outputs  $\tilde{\Theta}_{\theta\theta}$  and  $\tilde{\Theta}_{S_A \theta}$  of this function, **gsw\_CT\_second\_derivatives**(SA,pt), remain well-behaved as the input Absolute Salinity approaches zero, including at  $S_A = 0 \text{ g kg}^{-1}$ , but the output  $\tilde{\Theta}_{S_A S_A}$  has a singularity at  $S_A = 0 \text{ g kg}^{-1}$ , and a nan is returned at  $S_A = 0 \text{ g kg}^{-1}$ .

### References

IOC, SCOR and IAPSO, 2010: *The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties*. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56, UNESCO (English), 196 pp. Available from <http://www.TEOS-10.org> See Eqn. (A.12.3) of this TEOS-10 Manual.

Here follows appendix A.12 of the TEOS-10 Manual (IOC *et al.*, 2010).

### A.12 Differential relationships between $\eta$ , $\theta$ , $\Theta$ and $S_A$

Evaluating the fundamental thermodynamic relation in the forms (A.11.6) and (A.11.12) and using the four boxed equations in appendix A.11, we find the relations

$$\begin{aligned} (T_0 + t)d\eta + \mu(p)dS_A &= \frac{(T_0 + t)}{(T_0 + \theta)} c_p(0) d\theta + [\mu(p) - (T_0 + t)\mu_T(0)]dS_A \\ &= \frac{(T_0 + t)}{(T_0 + \theta)} c_p^0 d\Theta + \left[ \mu(p) - \frac{(T_0 + t)}{(T_0 + \theta)} \mu(0) \right] dS_A. \end{aligned} \quad (A.12.1)$$

The quantity  $\mu(p)dS_A$  is now subtracted from each of these three expressions and the whole equation is then multiplied by  $(T_0 + \theta)/(T_0 + t)$  obtaining

$$(T_0 + \theta)d\eta = c_p(0) d\theta - (T_0 + \theta)\mu_T(0) dS_A = c_p^0 d\Theta - \mu(0) dS_A. \quad (A.12.2)$$

From this follows all the following partial derivatives between  $\eta$ ,  $\theta$ ,  $\Theta$  and  $S_A$ ,

$$\Theta_{\theta}|_{S_A} = c_p(S_A, \theta, 0)/c_p^0, \quad \Theta_{S_A}|_{\theta} = [\mu(S_A, \theta, 0) - (T_0 + \theta)\mu_T(S_A, \theta, 0)]/c_p^0, \quad (A.12.3)$$

$$\Theta_{\eta}|_{S_A} = (T_0 + \theta)/c_p^0, \quad \Theta_{S_A}|_{\eta} = \mu(S_A, \theta, 0)/c_p^0, \quad (A.12.4)$$

$$\theta_{\eta}|_{S_A} = (T_0 + \theta)/c_p(S_A, \theta, 0), \quad \theta_{S_A}|_{\eta} = (T_0 + \theta)\mu_T(S_A, \theta, 0)/c_p(S_A, \theta, 0), \quad (A.12.5)$$

$$\theta_{\Theta}|_{S_A} = c_p^0/c_p(S_A, \theta, 0), \quad \theta_{S_A}|_{\Theta} = -[\mu(S_A, \theta, 0) - (T_0 + \theta)\mu_T(S_A, \theta, 0)]/c_p(S_A, \theta, 0), \quad (A.12.6)$$

$$\eta_{\theta}|_{S_A} = c_p(S_A, \theta, 0)/(T_0 + \theta), \quad \eta_{S_A}|_{\theta} = -\mu_T(S_A, \theta, 0), \quad (\text{A.12.7})$$

$$\eta_{\theta}|_{S_A} = c_p^0/(T_0 + \theta), \quad \eta_{S_A}|_{\theta} = -\mu(S_A, \theta, 0)/(T_0 + \theta). \quad (\text{A.12.8})$$

The three second order derivatives of  $\hat{\eta}(S_A, \Theta)$  are listed in Eqns. (P.14) and (P.15) of appendix P. The corresponding derivatives of  $\hat{\theta}(S_A, \Theta)$ , namely  $\hat{\theta}_{\Theta}$ ,  $\hat{\theta}_{S_A}$ ,  $\hat{\theta}_{\Theta\Theta}$ ,  $\hat{\theta}_{S_AS_A}$  and  $\hat{\theta}_{S_A\Theta}$  can also be derived using Eqn. (P.13), obtaining

$$\hat{\theta}_{\Theta} = \frac{1}{\tilde{\Theta}_{\theta}}, \quad \hat{\theta}_{S_A} = -\frac{\tilde{\Theta}_{S_A}}{\tilde{\Theta}_{\theta}}, \quad \hat{\theta}_{\Theta\Theta} = -\frac{\tilde{\Theta}_{\theta\theta}}{(\tilde{\Theta}_{\theta})^3}, \quad \hat{\theta}_{S_A\Theta} = -\frac{\tilde{\Theta}_{\theta S_A}}{(\tilde{\Theta}_{\theta})^2} + \frac{\tilde{\Theta}_{S_A} \tilde{\Theta}_{\theta\theta}}{(\tilde{\Theta}_{\theta})^3}, \quad (\text{A.12.9a,b,c,d})$$

$$\text{and} \quad \hat{\theta}_{S_AS_A} = -\frac{\tilde{\Theta}_{S_AS_A}}{\tilde{\Theta}_{\theta}} + 2 \frac{\tilde{\Theta}_{S_A}}{\tilde{\Theta}_{\theta}} \frac{\tilde{\Theta}_{\theta S_A}}{\tilde{\Theta}_{\theta}} - \left( \frac{\tilde{\Theta}_{S_A}}{\tilde{\Theta}_{\theta}} \right)^2 \frac{\tilde{\Theta}_{\theta\theta}}{\tilde{\Theta}_{\theta}}, \quad (\text{A.12.10})$$

in terms of the partial derivatives  $\tilde{\Theta}_{\theta}$ ,  $\tilde{\Theta}_{S_A}$ ,  $\tilde{\Theta}_{\theta\theta}$ ,  $\tilde{\Theta}_{\theta S_A}$  and  $\tilde{\Theta}_{S_AS_A}$  which can be obtained by differentiating the polynomial  $\tilde{\Theta}(S_A, \theta)$  from the TEOS-10 Gibbs function.