

CONJURA · OPEN PROBLEM

Randomness Extraction from Unpredictable Random-Oracle Sources

A Leftover-Hash-Lemma Conjecture, Public Seed

Status. Statement: human-written, formalized in Lean (see `lean/Statement.lean`).

Category. Information-Theoretic Cryptography.

Setting. Fix nonempty finite sets $\mathcal{K}$ (seeds), $\mathcal{D}$ (inputs), $\mathcal{R}$ (outputs), and write $K := |\mathcal{K}|$, $D := |\mathcal{D}|$, $R := |\mathcal{R}|$. Let $\text{Fun}(\mathcal{K} \times \mathcal{D}, \mathcal{R})$ be the set of all functions $\mathcal{K} \times \mathcal{D} \rightarrow \mathcal{R}$, let $\text{SD}(\cdot, \cdot)$ denote statistical distance and $U_{\mathcal{R}}$ the uniform distribution on $\mathcal{R}$. The source S , the predictor P and the distinguisher D of Figure 1 are computationally unbounded and receive the *entire* function table of H as an explicit input. The seed sd is sampled only after S has terminated, and is given to D : that is what makes this the *public-seed* game.

Game Pred_P^S $H \xleftarrow{\\$} \text{Fun}(\mathcal{K} \times \mathcal{D}, \mathcal{R}); \quad (x, z) \xleftarrow{\\$} S(H); \quad x' \xleftarrow{\\$} P(H, z)$ return $(x = x')$
Game Ext-pub_D^S (public seed) $H \xleftarrow{\\$} \text{Fun}(\mathcal{K} \times \mathcal{D}, \mathcal{R}); \quad (x, z) \xleftarrow{\\$} S(H); \quad sd \xleftarrow{\\$} \mathcal{K}$ $y_0 \leftarrow H(sd, x); \quad y_1 \xleftarrow{\\$} \mathcal{R}; \quad b \xleftarrow{\\$} \{0, 1\}$ $b' \xleftarrow{\\$} D(H, sd, y_b, z); \quad \textbf{return } (b = b')$

Figure 1: Prediction game (top) and the public-seed extraction game (bottom). The first argument of S , P and D is the *full function table* of H ; the distinguisher additionally receives the seed sd .

Definition 1. Set $\text{Adv}_{\mathcal{K}, \mathcal{D}, \mathcal{R}, S}^{\text{pred}}(P) := \Pr[\text{Pred}_P^S \Rightarrow 1]$ and

$$\text{Adv}_{\mathcal{K}, \mathcal{D}, \mathcal{R}}^{\text{ext-pub}}(S, D) := 2 \Pr[\text{Ext-pub}_D^S \Rightarrow 1] - 1.$$

A source S is ϵ -unpredictable if $\text{Adv}_{\mathcal{K}, \mathcal{D}, \mathcal{R}, S}^{\text{pred}}(P) \leq \epsilon$ for every unbounded predictor P .

Conjecture 1 (Public seed). *There is a universal constant $c > 0$, independent of $\mathcal{K}$, $\mathcal{D}$, $\mathcal{R}$ and ϵ , such that for all nonempty finite $\mathcal{K}, \mathcal{D}, \mathcal{R}$, all $\epsilon \in (0, 1]$, every ϵ -unpredictable source S and every unbounded distinguisher D ,*

$$\text{Adv}_{\mathcal{K}, \mathcal{D}, \mathcal{R}}^{\text{ext-pub}}(S, D) \leq \delta_{\text{pub}}(\epsilon, K, D, R) := c \cdot \left(\sqrt{\epsilon R} + \sqrt{\frac{\log_2 D}{K}} \right).$$

Remark 1 (What the leftover hash lemma gives). Condition on the table and the auxiliary output, $(H, Z) = (h, z)$, and write $p_x := \Pr[X = x \mid h, z]$. Because sd is drawn only after S has terminated it is uniform on $\mathcal{K}$ and independent of X given (h, z) , so the distinguisher's task is to tell $(sd, h(sd, X))$ from uniform on $\mathcal{K} \times \mathcal{R}$: a strong-extractor question about the *fixed* family $\{h(k, \cdot)\}_{k \in \mathcal{K}}$, to which the leftover hash lemma applies in its almost-universal form. Writing $\text{CP} := \sum_x p_x^2$ for the source's collision probability and $\bar{c} := \frac{1}{K} \sum_{k \in \mathcal{K}} \Pr[h(k, X) = h(k, X')]$ for the family's mean collision probability on that source, it bounds the conditional advantage by

$$\frac{1}{2} \sqrt{R\bar{c} - 1}, \quad R\bar{c} - 1 = R \cdot \text{CP} + \gamma, \quad \gamma := R(\bar{c} - \text{CP}) - 1,$$

with $\gamma \leq 0$ exactly when the family is universal on the source's support. Averaging over (H, Z) preserves the shape, $\sqrt{\cdot}$ being concave, and $\text{CP} \leq \max_x p_x$ turns the first summand into $\frac{1}{2} \sqrt{\epsilon R}$, which is δ_{pub} 's first term with $c = \frac{1}{2}$. On that summand the lemma is already at the conjectured shape.

On the second it is not, and the obstruction is structural rather than statistical. The family $\{h(k, \cdot)\}$ is not universal and cannot be made so — a universal family on D inputs needs a seed of $\Omega(\log D)$ bits, that is $K \geq D^{\Omega(1)}$, whereas this conjecture is of interest at $K \approx \log_2 D$ (Vadhan, *Pseudorandomness*, the remark following Theorem 6.18) — so γ must be taken over the supports S may pick *after* seeing H , and those can be chosen to make it large. Fix seeds $k_1, \dots, k_j$ and symbols $r_1, \dots, r_j$ and let S output a uniform element

of

$$T := \{x \in \mathcal{D} : H(k_i, x) = r_i \text{ for } i = 1, \dots, j\}, \quad |T| \approx D/R^j.$$

This is admissible: S reads the whole table, and sd is drawn only afterwards. On T the j chosen rows are *constant*, so each contributes its maximum to $\bar{c}$ and $\gamma \geq (R-1)j/K$. More generally, confining j rows to a symbol set of size R/s leaves $|T| \approx D/s^j$ and gives $\gamma \geq (s-1)j/K$; since $(s-1)/\log_2 s$ increases, the best choice is $s = \min\{R, D\epsilon\}$. Writing $L := \min\{\log_2 R, \log_2 D - \log_2(1/\epsilon)\}$ for the number of bits that can be pinned per row, the lemma therefore cannot yield better than

$$\frac{1}{2} \sqrt{\epsilon R + \frac{(\log_2 D - \log_2(1/\epsilon))(2^L - 1)}{LK}}.$$

The rate in K is the same square root δ_{pub} asks for; what the lemma loses is a *factor* of about $\sqrt{2^L/L}$, which is $L - \log_2 L$ extra bits of seed. For $D = 2^{256}$, $R = 2^{64}$ and a source of min-entropy 128, reaching $\delta = 2^{-32}$ needs $\log_2 K \approx 69.5$ under δ_{pub} and 127 by this route.

The loss is the Cauchy–Schwarz step $\|v\|_1 \leq \sqrt{KR} \|v\|_2$ inside the lemma: support selection concentrates the deviation v on j of the K rows rather than spreading it over all KR cells, and on such a v that step is off by $\sqrt{RK/j}$. For the source above the true advantage is $\approx j/K$ while the lemma returns $\approx \frac{1}{2} \sqrt{Rj/K}$. So the conjecture is not a corollary of the leftover hash lemma, and its content is the second summand: the price of letting the source choose its support after seeing the entire table. The solution note in this folder pays that price at the conjectured rate by never passing through a collision probability, bounding the L_1 distance directly instead. (What is stated here is a limitation — the lemma cannot do better than the displayed bound. Whether it does that well is not claimed.)

The secret-seed companion. This note states one conjecture, the one the accompanying page states. There is a companion statement in the same setting, differing only in that D is called as $D(H, y_b, z)$ and never receives sd ; it conjectures the bound $c\sqrt{(\epsilon R + \log_2 D)/K}$,

in which the seed's factor $1/K$ attaches to both summands. It is a genuinely different claim, not a restatement: the solution note in this folder proves the conjecture above and separately shows that transposing the secret-seed expression verbatim into the public-seed game gives a *false* statement. The two were originally posed together in a single write-up and were split so that each note states exactly the conjecture its page states.