





MSSC2025

Cost-effective Composite HF/DFT Methods for High Throughput Screening and Machine Learning Applications






Mattia Raimondo






September 8 to 12, 2025
Turin, Italy

Large & Complex systems








Large & Complex systems






Large-scale screenings:

- *Drug design*
- *Crystal Structure Prediction*
- *High-Throughput Screening*






Large & Complex systems



Large-scale screenings:

- Drug design
- Crystal Structure Prediction
- High-Throughput Screening





1) Are DFT calculation affordable even with *m*-GGA and hybrid functional?

2) How far can we go with small-scale computing resources?



Low cost, yet accurate methods are needed





S. Grimme

ab initio composite methods: idea



J. G. Brandenburg

Hierarchy of simplified electronic structure methods for good structures, reasonable energetic properties and non-covalent interactions.

Key ingredients:

- *Based on HF and DFT*
- *From minimal to small-to-medium basis sets (atom centered BF, GTO)*

$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$



S. Grimme

ab initio composite methods: idea




J. G. Brandenburg

Hierarchy of simplified electronic structure methods for good structures, reasonable energetic properties and non-covalent interactions.

Key ingredients:

- *Based on HF and DFT*
- *From minimal to small-to-medium basis sets (atom centered BF, GTO)*

$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$





S. Grimme

ab initio composite methods: idea





J. G. Brandenburg

Hierarchy of simplified electronic structure methods for good structures, reasonable energetic properties and non-covalent interactions.

Key ingredients:

- **Based on HF and DFT**
- **From minimal to small-to-medium basis sets (atom centered BF, GTO)**
- **London dispersion through D3(BJ) correction**
- **BSSE counterpoise by gCP correction**
- **BSIE correction at short-range for HF (SRB)**
- **Fitting of the parameters in the D3 and gCP corrections on s66x8 high-level dataset of molecular adducts (CCSD(T)/CBS)**



$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$


XC functional and basis set dominated *dispersion and BSSE dominated*




non-covalent

medium-range correlation

covalent



interatomic distances / Å





S. Grimme

ab initio composite methods: idea





J. G. Brandenburg

Hierarchy of simplified electronic structure methods for good structures, reasonable energetic properties and non-covalent interactions.

Key ingredients:

- *Based on HF and DFT*
- *From minimal to small-to-medium basis sets (atom centered BF, GTO)*
- *London dispersion through D3(BJ) correction*
- *BSSE counterpoise by gCP correction*
- *BSIE correction at short-range for HF (SRB)*
- *Fitting of the parameters in the D3 and gCP corrections on s66x8 high-level dataset of molecular adducts (CCSD(T)/CBS)*



$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$


XC functional and basis set dominated *dispersion and BSSE dominated*




non-covalent

medium-range correlation

covalent



interatomic distances / Å





S. Grimme


ab initio composite methods: idea



J. G. Brandenburg

$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$

	HF-3c	PBEh-3c	HSE-3c
<i>Hamiltonian</i>	Hartree-Fock	Global hybrid	Range separated
<i>Basis set</i>	MINIX	def2-mSVP	def2-mSVP
<i>Dispersion</i>	D3 (2-body)	D3 (2&3-body)	D3 (2&3-body)
<i>BSSE</i>	gCP	gCP (damped)	gCP (damped)
<i>BSIE</i>	SRB	-	-





S. Grimme


ab initio composite methods: idea



J. G. Brandenburg

$$E_{tot}^{3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP} + E^{SRB}$$

	HF-3c	PBEh-3c	HSE-3c
<i>Hamiltonian</i>	Hartree-Fock	Global hybrid	Range separated
<i>Basis set</i>	MINIX	def2-mSVP	def2-mSVP
<i>Dispersion</i>	D3 (2-body)	D3 (2&3-body)	D3 (2&3-body)
<i>BSSE</i>	gCP	gCP (damped)	gCP (damped)
<i>BSIE</i>	SRB	-	-




Accurate for:

- *Protein-ligand binding affinity*
- *Molecular crystals with short hydrogen contacts*
- *Light elements of the Periodic Table*

Not applicable to:

- *Covalent and inorganic solids*
- *Layered and porous materials*
- *Heavier elements*




Extending applicability to solid state: sol-3c


$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$

Methods (hybrids)

- Employ XC functionals devised for solids (PBEsol, HSEsol)
- Reduce the amount of HF exchange for a better description of electronic and optical properties



L. Donà





Extending applicability to solid state: sol-3c



L. Donà

CRYSTAL23

$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$


Methods (hybrids)

- Employ XC functionals devised for solids (PBEsol, HSEsol)
- Reduce the amount of HF exchange for a better description of electronic and optical properties



Basis set (HF & DFT)

- Apply a simple recipe to make molecular BS suitable for solids (H-Xe)
- MINIX: Decontract outermost shells and scale most diffuse functions
- def2-mSVP: Upshift (> 0.1 a.u.) and scale the most diffuse functions

molecular basis set



solid state basis set





Extending applicability to solid state: sol-3c



L. Donà

CRYSTAL23

$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$


Methods (hybrids)

- Employ XC functionals devised for solids (PBEsol, HSEsol)
- Reduce the amount of HF exchange for a better description of electronic and optical properties


Basis set (HF & DFT)

- Apply a simple recipe to make molecular BS suitable for solids (H-Xe)
- MINIX: Decontract outermost shells and scale most diffuse functions
- def2-mSVP: Upshift (> 0.1 a.u.) and scale the most diffuse functions

molecular basis set





solid state basis set



3c

- Refitting of most of the parameters in the D3 and gCP corrections on s66x8 high-level dataset (CCSD(T)/CBS)






r²SCAN-derived composite methods




$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$

- *Based on DFT (mGGA & hybrids):
r²SCAN/r²SCAN0*



- *From small-to-medium basis sets (atom centered BF, GTO):
soldef2-mSVP & POB-TZVP-rev2*
- *London dispersion through D3(BJ) correction*
- *BSSE counterpoise by gCP correction*
- *Refitted on NENCI-2021* dataset*




r²SCAN-derived composite methods





$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$


- Based on DFT (mGGA & hybrids):
r²SCAN/r²SCAN0



- From small-to-medium basis sets (atom centered BF, GTO):
soldef2-mSVP & POB-TZVP-rev2

- London dispersion through D3(BJ) correction
- BSSE counterpoise by gCP correction
- Refitted on NENCI-2021* dataset






r²SCAN-derived composite methods






$$E_{tot}^{sol3c} = E_{tot}^{DFT/basis} + E_{disp}^{D3} + E_{BSSE}^{gCP}$$

- Based on DFT (mGGA & hybrids):
r²SCAN/r²SCAN0



- From small-to-medium basis sets (atom centered BF, GTO):
soldef2-mSVP & POB-TZVP-rev2


- London dispersion through D3(BJ) correction
- BSSE counterpoise by gCP correction
- Refitted on NENCI-2021* dataset



sol-3c composite methods: validation

Have these methods the same accuracy of the original ones?


sol-3c composite methods: validation



Have these methods the same accuracy of the original ones?


- **23 molecular crystals:** low-T crystal structure (X-ray/Neutron), accurate sublimation enthalpy (298K), no phase transitions ($0 < T < 298\text{K}$)
- Comparison with back-corrected (ZPVE and thermal contribution from DFT) experimental ΔH (Range: $28 < \Delta H < 160 \text{ kJ/mol}$)

sol-3c composite methods: validation




Have these methods the same accuracy of the original ones?

- **23 molecular crystals:** low-T crystal structure (X-ray/Neutron), accurate sublimation enthalpy (298K), no phase transitions ($0 < T < 298\text{K}$)
- Comparison with back-corrected (ZPVE and thermal contribution from DFT) experimental ΔH (Range: $28 < \Delta H < 160 \text{ kJ/mol}$)




sol-3c composite methods: validation




Have these methods the same accuracy of the original ones?


- **23 molecular crystals:** low-T crystal structure (X-ray/Neutron), accurate sublimation enthalpy (298K), no phase transitions ($0 < T < 298\text{K}$)
- Comparison with back-corrected (ZPVE and thermal contribution from DFT) experimental ΔH (Range: $28 < \Delta H < 160 \text{ kJ/mol}$)

Molar Volume




Cohesive Energy







sol-3c composite methods: validation

Are these methods of general applicability?






sol-3c composite methods: validation




Are these methods of general applicability?

SS20: 20 simple solids – cubic lattice – covalent and ionic systems – Low-T data*

The presence of very diffuse Gaussian functions meant that the original 3c methods were not suitable for inorganic solids



sol-3c composite methods: validation




Are these methods of general applicability?


SS20: 20 simple solids – cubic lattice – covalent and ionic systems – Low-T data*

The presence of very diffuse Gaussian functions meant that the original 3c methods were not suitable for inorganic solids

Lattice Parameter




Band Gap



Large Scale Screening

Are these methods really cost effective?



System types:

- 0D to 1D, 2D and 3D;
- molecular crystals and inorganic solids;
- from dense to porous materials


System sizes

- Up to **1.5k atoms** (16 k AOs) on **40 cores**
- Up to **≈3k atoms** (37 k AOs) on **80 cores**




**How to Model
Metal-Organic Frameworks
with CRYSTAL**

Lorenzo Donà




High-Throughput Screening: Metal-Organic Frameworks



- Inorganic Cluster
- Organic Linker
- 3D framework
- Porous material
- Crystalline structure
- Framework Modifications
- High chemical versatility and modular nature

Our accurate yet cost-effective composite methods make DFT calculations feasible for large and complex systems such as MOFs, enabling:

- **High-Throughput Screening**
- the generation of **extensive datasets** for Machine Learning applications



ML-derived atomic charges

H. Paja



1. Prediction of atomic charges
2. From big data to small data

ML-derived atomic charges

H. Paja



1. Prediction of atomic charges
2. From big data to small data



- Experimentally derived MOFs
- 20 375 cleaned structures
- Structures with no more than 300 atoms in the primitive cell
- PBE-D3(BJ)/520eV

Structural descriptors from




Chemical descriptors from




**Random Forest
Algorithm**

1. Prediction of atomic charges
2. From big data to small data



mendeleev 

Random Forest Algorithm



ML-derived atomic charges

H. Paja



1. Prediction of atomic charges
2. From big data to small data



- Experimentally derived MOFs
- 20 375 cleaned structures
- Structures with no more than 300 atoms in the primitive cell
- PBE-D3(BJ)/520eV


Structural descriptors from




Chemical descriptors from



**Random Forest
Algorithm**



balanced composition
with equated metal
abundance



ML-derived atomic charges

H. Paja



1. Prediction of atomic charges
2. From big data to small data



- Experimentally derived MOFs
- 20 375 cleaned structures
- Structures with no more than 300 atoms in the primitive cell
- PBE-D3(BJ)/520eV


Structural descriptors from



Chemical descriptors from




**Random Forest
Algorithm**



balanced composition
with equated metal
abundance




randomly selected
spanning from 500 to
3000 MOFs



ML-derived atomic charges

H. Paja




- Chemically-driven selection: more balanced composition with equated metal abundance of 2%
- Although the size of the datasets increase, no relevant improvement on ML predicted charges are achieved.
- Despite this *rf_eq* represents a surprising improvement with a **MAE lower than 0.02 electrons**.

Label	5	10	15	EQ	20	30	QMOF
MOFs	500	1000	1500	1474	2000	3000	20309
DDEC	53 K	111 K	165 K	165 K	221 K	334 K	2 M

ML-derived atomic charges


H. Paja



- Chemically-driven selection: more balanced composition with equated metal abundance of 2%
- Although the size of the datasets increase, no relevant improvement on ML predicted charges are achieved.
- Despite this *rf_eq* represents a surprising improvement with a **MAE lower than 0.02 electrons**.

Label	5	10	15	EQ	20	30	QMOF
MOFs	500	1000	1500	1474	2000	3000	20309
DDEC	53 K	111 K	165 K	165 K	221 K	334 K	2 M

For the balanced dataset we proceed to the calculation of **Hirshfeld-I** atomic charges using CRYSTAL23 with **r²SCAN0pob-3c**, retrained the model, and achieved MAE of 0.06 electrons.




Adsorption Simulation

UFF force field

$$E_{total} = \sum_{bonds} K_r (r - r_{eq})^2 + \sum_{angles} K_\theta (\theta - \theta_{eq})^2 + \sum_{dihedrals} \frac{V_n}{2} [1 + \cos(n\phi - \gamma)] + \sum_{i < j} \left[\frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6} + \frac{q_i q_j}{\epsilon R_{ij}} \right]$$



The diagram illustrates the components of the UFF force field. It is divided into **Bonded** and **Non-bonded** terms. The **Bonded** section includes three diagrams: a bond length r between two atoms, an angle θ between three atoms, and a dihedral angle ϕ between four atoms. The **Non-bonded** section includes a diagram of a Lennard-Jones potential well with distance R_{ij} and a diagram of an electrostatic interaction between two charged spheres (one red with '+', one blue with '-') with distance R_{ij} . The electrostatic term in the equation is highlighted with a red box.



Adsorption Simulation


UFF force field

$$E_{\text{total}} = \sum_{\text{bonds}} K_r (r - r_{\text{eq}})^2 + \sum_{\text{angles}} K_\theta (\theta - \theta_{\text{eq}})^2 + \sum_{\text{dihedrals}} \frac{V_n}{2} [1 + \cos(n\phi - \gamma)] + \sum_{i < j} \left[\frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6} + \frac{q_i q_j}{\epsilon R_{ij}} \right]$$




- Machine learning models can accurately predict Hirshfeld charges based on structural descriptors.
- These predicted charges can then be used to reliably perform GCMC simulations, producing results that are comparable to reference data.
- This approach can be scaled up to include a larger number of MOFs, improving our understanding of, and ability to design, materials for specific applications.

r²SCAN-derived MLIP




- *r²SCAN0pob-3c* method for Geometry Optimisation

r²SCAN-derived MLIP




- *r²SCAN0pob-3c* method for Geometry Optimisation
- 16877 MOF configurations with hundreds atoms in the primitive unit cell

r²SCAN-derived MLIP



- *r²SCAN0pob-3c* method for Geometry Optimisation
- 16877 MOF configurations with hundreds atoms in the primitive unit cell
- The training set comprises more than **5 million entries**, encompassing atomic positions, forces, and stress tensors.

r²SCAN-derived MLIP



- *r²SCAN0pob-3c* method for Geometry Optimisation
- 16877 MOF configurations with hundreds atoms in the primitive unit cell
- The training set comprises more than **5 million entries**, encompassing atomic positions, forces, and stress tensors.
- expanding the chemical design space with f-block elements
- general purpose universal **ML potential for MOFs**
- robustly validated

2 MACE model:

- D8 (from scratch)
- D8_ft (finetuning of MP0)

...pluggable with ASE, LAMMPS, Nequip, PyTorch...



Validation I



MOF-LC15 (low Temperature X-ray diffraction measurements)

Comparison with MP0 and OMAT models

CPO-27-Mg

CPO-27-Ni

CPO-27-Zn

HKUST-1-Cu

HKUST-1-Zn

MIL-125-Ti

MIL-140-Zr

MIL-47-V

MIL-53-Al-LP

MIL-53-Al-NP


MIL-68-Ga

MOF-5-Zn

UIO-66-Zr

UIO-66-Zr-OH

UIO-67-Zr-OH





Validation I



MOF-LC15 (low Temperature X-ray diffraction measurements)

Comparison with MP0 and OMAT models

MOF-LC15

r^2 SCAN0pob-3c D8 D8_ft MP0 OMAT

EXP

1.50 3.18 3.67 5.92 3.72

r^2 SCAN0pob-3c

2.46 2.95 5.75 3.40

D8

3.43 7.85 4.91

D8_ft


7.89 5.56

MP-0

3.92

OMAT

MARE%







Validation II



882 Experimental MOF from *CCDC* ($T < 100\text{ K}$ and $R\text{-factors} < 5\%$)







Validation II



882 Experimental MOF from CCDC ($T < 100\text{ K}$ and $R\text{-factors} < 5\%$)







Validation II



882 Experimental MOF from *CCDC* ($T < 100\text{ K}$ and $R\text{-factors} < 5\%$)



Despite their **universal applicability**, both our MLIPs and state-of-the-art universals, such as MP-0 and OMAT, **fail to optimise some MOFs**.

Future Perspective:

1. Expand Training Data
2. Active Learning Integration
3. Transfer Learning & Fine-Tuning
4. Hyperparameter Optimization


Acknowledgements



H. Paja

L. Donà

B. Civalleri



J. G. Brandenburg



I. Bush



D3(BJ) and gCP corrections

D3(BJ): London dispersion

$$E_{disp} = E^{(2)} + E^{(3)}$$

2-body damped atom-pair wise term

$$E_{disp}^{(2)} = -\frac{1}{2} \sum_{AB} \frac{C_6^{AB}}{R_{AB}^6 + f(R_{AB}^0)^6} + s_8 \frac{C_8^{AB}}{R_{AB}^8 + f(R_{AB}^0)^8}$$

Becke-Johnson damping function

$$f(R_{AB}^0) = a_1 R_{AB}^0 + a_2$$

3-body damped atom-triple wise term

$$E^{(3)} = \sum_{ABC} f_{d,(3)}(\bar{R}_{ABC}) E^{ABC}$$

gCP: geometrical counterpoise

$$E_{gCP} = \sigma \sum_a^{atoms} E_a^{atom}$$

Mimicks the Boys-Bernardi counterpoise method with an atom-pair wise term

$$E_{BSSE}^{gCP} = \sigma \sum_A^{atoms} \sum_{A \neq B}^{atoms} E_A^{miss} \frac{\exp(-\alpha(R_{AB})^\beta)}{\sqrt{S_{AB} N_B^{virt}}}$$

E_A^{miss} is a gauge of the basis set incompleteness per each atomic species tuned through a decay function

$$f_{dec}(R_{AB,g}) = \frac{\exp(-\alpha(R_{AB,g})^\beta)}{\sqrt{S_{AB,g} N_B^{virt}}}$$