

Chapter 5

TECHNIQUES OF INTEGRATION

Contents

5.1. The Infefine Integral

5.2 The Substitution Rule

5.3. Integration by Parts

5.4 Improper Integrals

5.1. The Indefinite integral

Definition

Due to the relation given by the FTC between antiderivatives and integrals, the notation $\int f(x) dx$ is traditionally used for an antiderivative of f and is called an indefinite integral.

Thus, $\int f(x) dx = F(x)$ means $F'(x) = f(x)$.

You should distinguish carefully between definite and indefinite integrals. A definite integral $\int_a^b f(x) dx$ is a number.

An indefinite integral $\int f(x) dx$ is a **function** (or family of functions).

5.1. The Indefinite integral

Formular

$$1. \int cf(x) dx = c \int f(x) dx$$

$$2. \int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx$$

$$3. \int k dx = kx + C$$

$$4. \int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (n \neq -1)$$

$$5. \int \frac{1}{x} dx = \ln|x| + C$$

5.1. The Indefinite integral

Formular

$$6a. \int \sin x \, dx = -\cos x + C$$

$$6b. \int \cos x \, dx = \sin x + C$$

$$7a. \int \sec^2 x \, dx = \tan x + C$$

$$7b. \int \csc^2 x \, dx = -\cot x + C$$

$$8a. \int \sec x \tan x \, dx = \sec x + C$$

$$8b. \int \csc x \cot x \, dx = -\csc x + C$$

5.2. The Substitution Rule

Example: Evaluate $\int \frac{dx}{(3-5x)^2}$

Solution:

Let $u = 3 - 5x$. Then, $du = -5dx$, so $dx = -\frac{1}{5}du$

$$\begin{aligned} \int \frac{dx}{(3-5x)^2} &= \int \frac{1}{u^2} \left(\frac{-du}{5} \right) = -\frac{1}{5} \int \frac{1}{u^2} du \\ &= -\frac{1}{5} \int u^{-2} du = -\frac{1}{5} \cdot \frac{u^{-1}}{-1} + C = \frac{1}{5u} + C = \frac{1}{5(3-5x)} + C \end{aligned}$$

Method 2:
$$\int \frac{dx}{(3-5x)^2} = \int (3-5x)^{-2} dx = -\frac{1}{5} \cdot \frac{(3-5x)^{-1}}{-1} + C$$

5.2. The Substitution Rule

General: $\int f(ax + b)dx; u = ax + b \Rightarrow du = adx \Rightarrow dx = \frac{1}{a} du$

$$\int f(ax + b)dx = \int f(u) \times \frac{1}{a} du$$

$$= \frac{1}{a} f(u) du$$

$$= \frac{1}{a} F(u) + C$$

$$= \frac{1}{a} F(ax + b) + C$$

$$\int f(ax + b)dx = \frac{1}{a} F(ax + b) + C,$$

5.2. The Substitution Rule

Example: Use the formular $\int f(ax + b)dx = \frac{1}{a} F(ax + b) + C$,
 find:

$$1. \int (2x - 1)^{10} dx$$

$$2. \int \frac{1}{1-x} dx$$

$$3. \int \frac{1}{(1-x)^2} dx$$

$$4. \int \frac{1}{\sqrt{1-x}} dx$$

$$5. \int e^{3x+4} dx$$

$$6. \int (\sin 5x + \cos 7x) dx$$

5.2. The Substitution Rule

Theorem

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous on I , then

$$\int f(g(x))g'(x)dx = \int f(u)du$$

Example 1: Evaluate $I = \int 2x\sqrt{1+x^2} dx$.

Solution: $u = \sqrt{1+x^2}$

$$u^2 = 1 + x^2 \Rightarrow 2u du = 2x dx$$

$$\Rightarrow I = \int 2u^2 du = \frac{2}{3}u^3 + C = \frac{2}{3}\left(\sqrt{1+x^2}\right)^3 + C$$

5.2. The Substitution Rule

Example 2: Evaluate $\int x^3 \cos(x^4 + 2) dx$.

Solution:

$$u = x^4 + 2 \Rightarrow du = 4x^3 dx \Rightarrow x^3 dx = \frac{du}{4}$$

$$\begin{aligned} \int x^3 \cos(x^4 + 2) dx &= \int \cos u \cdot \frac{1}{4} du = \frac{1}{4} \int \cos u du \\ &= \frac{1}{4} \sin u + C \\ &= \frac{1}{4} \sin(x^4 + 2) + C \end{aligned}$$

5.2. The Substitution Rule

Sub.rule for def. integrals:

If g' is continuous on $[a, b]$ and f is continuous on the range

of $u = g(x)$, then
$$\int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du.$$

Let F be an antiderivative of f . Then, $F(g(x))$ is an antiderivative of $f(g(x))g'(x)$.

So,
$$\int_a^b f(g(x))g'(x)dx = F(g(x)) \Big|_a^b = F(g(b)) - F(g(a))$$


5.2. The Substitution Rule

Integs.of symm.functions:


Suppose f is continuous on $[-a, a]$.

a/ If f is even, $[f(-x) = f(x)]$, then $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$

b/ If f is odd, $[f(-x) = -f(x)]$, then $\int_{-a}^a f(x) dx = 0$



(a) f even, $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$



(b) f odd, $\int_{-a}^a f(x) dx = 0$

5.3. Integration by parts

Formula $I = \int f(x)g'(x)dx$

$$\text{Set } \begin{cases} u = f(x) \\ dv = g'(x)dx \end{cases} \Rightarrow \begin{cases} u' = f'(x) \\ v = g(x) \end{cases}$$

Thus, by the Substitution Rule, the formula for integration by parts becomes:

$$\int f(x)g'(x)dx = uv - \int vdu = f(x)g(x) - \int g(x)f'(x)dx$$

Note: $u = \ln x; x^k; e^x; \sin x; \cos x$

Using the FTC, we obtain:

$$\int_a^b f(x)g'(x)dx = [f(x)g(x)]_a^b - \int_a^b g(x)f'(x)dx$$

5.3. Integration by parts

Example 1: Find $\int x \sin x dx$

Solution:

$$\begin{cases} u = x \Rightarrow du = dx \\ dv = \sin x dx \Rightarrow v = -\cos x \end{cases}$$

$$\begin{aligned} I &= u.v - \int v du = -x \cos x - \int -\cos x dx \\ &= -x \cos x + \int \cos x dx = -x \cos x + \sin x + C \end{aligned}$$

5.3. Integration by parts

Example 2: Find $\int e^x \sin x dx$

Solution:

Step 1: Set
$$\begin{cases} u = e^x \Rightarrow du = e^x dx \\ dv = \sin x dx \Rightarrow v = -\cos x \end{cases}$$

$$\int e^x \sin x dx = -e^x \cos x + \int e^x \cos x dx$$

Step 2: Set
$$\begin{cases} u = e^x \Rightarrow du = e^x dx \\ dv = \cos x dx \Rightarrow v = \sin x \end{cases}$$

$$\int e^x \cos x dx = e^x \sin x - \int e^x \sin x dx$$

5.3. Integration by parts

Step 3: This can be regarded as an equation to be solved for the unknown integral.

$$\int e^x \sin x \, dx = -e^x \cos x + e^x \sin x - \int e^x \sin x \, dx$$

Adding to both sides $\int e^x \sin x \, dx$, we obtain:

$$2 \int e^x \sin x \, dx = -e^x \cos x + e^x \sin x$$

Dividing by 2 and adding the constant of integration, we get:

$$\int e^x \sin x \, dx = \frac{1}{2} e^x (\sin x - \cos x) + C$$

5.3. Integration by parts

Example 3: Suppose $f(x)$ is continuous and differentiable,

$f(1)=4$ and $\int_0^1 f(x)dx = 5$. Find $\int_0^1 xf'(x)dx$

a	4/5
b	5/4
c	1
d	None of the others
e	-1

5.3. Integration by parts

Example 4: Suppose $f(x)$ is continuous and differentiable,

$$f(1)=3, f(3)=1 \text{ and } \int_1^3 xf'(x)dx = 13$$

What is the average value of f on the interval $[1,3]$?


5.4. Improper Integrals

Improper Integrals of type 1

Definition: If $\int_a^t f(x) dx$ exists for every number $t \geq a$, then

$$\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx \text{ provided this limit exists}$$

(as a finite number).



5.4. Improper Integrals

Improper Integrals of type 1

Convergent and divergent:

The improper integrals $\int_a^{\infty} f(x) dx$ and $\int_{-\infty}^b f(x) dx$ are called:

- **Convergent** if the corresponding limit exists.
- **Divergent** if the limit does not exist.

5.4. Improper Integrals

If both $\int_a^\infty f(x) dx$ and $\int_{-\infty}^a f(x) dx$ are convergent, then

we define:
$$\int_{-\infty}^\infty f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^\infty f(x) dx$$

Here, any real number a can be used.

5.4. Improper Integrals

Example 1: Evaluate:

$$\text{a/ } I_1 = \int_1^{+\infty} \frac{dx}{x}$$

$$\text{b/ } I_2 = \int_1^{+\infty} \frac{dx}{x^2}$$

$$\text{c/ } I_3 = \int_1^{+\infty} \frac{dx}{\sqrt{x}}$$

5.4. Improper Integrals

Solution:

$$\text{a/ } I_1 = \int_1^{\infty} \frac{dx}{x}$$

$$\text{Step 1: } A = \int_1^b \frac{dx}{x} = \ln|x|_1^b = \ln b - \ln 1 = \ln b$$

$$\text{Step 2: } I_1 = \lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} \ln b = +\infty$$

5.4. Improper Integrals

Solution:

$$b/ I_2 = \int_1^{\infty} \frac{dx}{x^2}$$

$$\text{Step 1: } A = \int_1^b \frac{dx}{x^2} = -\frac{1}{x} \Big|_1^b = -\frac{1}{b} + 1$$

Step 2:

$$I_2 = \lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} \left(-\frac{1}{b} + 1 \right) = 1$$

5.4. Improper Integrals

Solution:

$$c/ I_3 = \int_1^{\infty} \frac{dx}{\sqrt{x}}$$

$$\text{Step 1: } A = \int_1^b \frac{dx}{\sqrt{x}} = 2\sqrt{x} \Big|_1^b = 2\sqrt{b} - 2$$

Step 2:

$$I_3 = \lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} (2\sqrt{b} - 2) = +\infty$$

5.4. Improper Integrals

Example 2: Evaluate:

$$\text{a/ } I_1 = \int_1^{+\infty} \frac{dx}{x^2 + 1}$$

$$\text{b/ } I_2 = \int_1^{+\infty} e^{-2x} dx$$

$$\text{c/ } I_3 = \int_e^{+\infty} \frac{1}{x(\ln x)^2} dx$$

5.4. Improper Integrals

Solution:

$$\text{a/ } I_1 = \int_1^{\infty} \frac{dx}{x^2 + 1}$$

$$\text{Step 1: } A = \int_1^b \frac{dx}{x^2 + 1} = \left(\arctan x \right)_1^b = \arctan b - \frac{\pi}{4}$$

Step 2:

$$I_1 = \lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} \left(\arctan b - \frac{\pi}{4} \right) = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$$

5.4. Improper Integrals

Solution:

$$\text{b/ } I_2 = \int_1^{+\infty} e^{-2x} dx$$

$$\text{Step 1: } A = \int_1^b e^{-2x} dx = \left(-\frac{1}{2} e^{-2x} \right)_1^b = -\frac{1}{2} e^{-2b} + \frac{1}{2} e^{-2}$$

$$\begin{aligned} \text{Step 2: } I_3 &= \lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} \left(-\frac{1}{2} e^{-2b} + \frac{1}{2} e^{-2} \right) \\ &= \lim_{b \rightarrow +\infty} \left(-\frac{1}{2e^{2b}} + \frac{1}{2} e^{-2} \right) = \frac{1}{2e^2} \end{aligned}$$

5.4. Improper Integrals

Solution:

$$\text{c/ } I_3 = \int_e^{\infty} \frac{1}{x(\ln x)^2} dx$$

$$\text{Step 1: } A = \int_e^b \frac{1}{x(\ln x)^2} dx \quad t = \ln x \Rightarrow dt = \frac{dx}{x}$$

$$A = \int_1^{\ln b} \frac{1}{t^2} dt = \left(-\frac{1}{t} \right)_1^{\ln b} = -\frac{1}{\ln b} + 1$$

5.4. Improper Integrals

Step 2: $\lim_{b \rightarrow +\infty} A = \lim_{b \rightarrow +\infty} \left(-\frac{1}{\ln b} + 1 \right) = 1$

So, $I_3 = \int_e^{\infty} \frac{1}{x(\ln x)^2} dx = 1$


5.4. Improper Integrals

Improper Integrals of type 2

Definition 1a: If f is continuous on $[a, b)$ and is discontinuous

at b , then
$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

this limit exists (as a finite number).




5.4. Improper Integrals

Improper Integrals of type 2

Definition 1b: If f is continuous on $(a, b]$ and is discontinuous

at a , then
$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

this limit exists (as a finite number).



5.4. Improper Integrals

Improper Integrals of type 2

The improper integral $\int_a^b f(x) dx$ is called:

- Convergent if the corresponding limit exists.
- Divergent if the limit does not exist.


5.4. Improper Integrals

Improper Integrals of type 2

If f has a discontinuity at c , where $a < c < b$, and

both $\int_a^c f(x) dx$ and $\int_c^b f(x) dx$ are convergent, then we

define:
$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$



5.4. Improper Integrals

Example: Evaluate:

$$\text{a/ } I_1 = \int_2^4 \frac{dx}{\sqrt{x-2}}$$

$$\text{b/ } I_2 = \int_1^2 \frac{dx}{x-1}$$

5.4. Improper Integrals

Solution:

$$\text{a/ } I_1 = \int_2^4 \frac{dx}{\sqrt{x-2}}$$

$$\text{Step 1: } A = \int_t^4 \frac{dx}{\sqrt{x-2}} = \left(2\sqrt{x-2} \right)_t^4 = 2\sqrt{2} - 2\sqrt{t-2}$$

$$I_1 = \lim_{t \rightarrow 2^+} A = \lim_{t \rightarrow 2^+} \left(2\sqrt{2} - 2\sqrt{t-2} \right) = 2\sqrt{2}$$


5.4. Improper Integrals

Comparison theorem

Suppose f and g are continuous functions with $f(x) \geq g(x) \geq 0$ for $x \geq a$.

a/ If $\int_a^{\infty} f(x) dx$ is convergent, then $\int_a^{\infty} g(x) dx$ is convergent.

b/ If $\int_a^{\infty} g(x) dx$ is divergent, then $\int_a^{\infty} f(x) dx$ is divergent.



5.4. Improper Integrals

Theorem:

$$\int_{a>0}^{\infty} \frac{1}{x^p} dx \text{ is:}$$

- Convergent if $p > 1$
- Divergent if $p \leq 1$

$$\int_a^{b<\infty} \frac{1}{(x-a)^p} dx \text{ is:}$$

Convergent if $p < 1$.

Divergent if $p \geq 1$

5.4. Improper Integrals

Example:

Use the Comparison Theorem to determine whether the integral is convergent or divergent.

a/ $\int_0^{\infty} e^{-x^2} dx$

b/ $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$

c/ $\int_0^1 \frac{2}{\sqrt{x^3}} dx$

d/ $\int_0^{\pi} \frac{\sin^2 x}{\sqrt{x}} dx$

5.4. Improper Integrals

Solution:

$$a/ \int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$$

• $\int_0^1 e^{-x^2} dx$ is just an ordinary definite integral.

• $x \geq 1$ we have $x^2 \geq x$ so $-x^2 \leq -x \Rightarrow e^{-x^2} \leq e^{-x}$

Therefore

$$\int_1^{\infty} e^{-x^2} dx \leq \int_1^{\infty} e^{-x} dx \text{ and } \int_1^{\infty} e^{-x} dx = -e^{-x} \Big|_1^{\infty} = \frac{1}{e}$$

5.4. Improper Integrals

Solution:

a/ we see that $\int_1^{\infty} e^{-x^2} dx$ is convergent.

In the Comparison Theorem, It follows that $\int_0^{\infty} e^{-x^2} dx$ is convergent.

5.4. Improper Integrals

Solution:

$$\text{b/ } \int_1^{\infty} \frac{1 + e^{-x}}{x} dx$$

We have $\frac{1 + e^{-x}}{x} > \frac{1}{x}$

• $\int_1^{\infty} \frac{1}{x} dx$ is divergent ($p = 1$)

The integral $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$ is divergent by the Comparison Theorem.

5.4. Improper Integrals

Solution:

$$c/ \int_0^1 \frac{2}{\sqrt{x^3}} dx = 2 \int_0^1 \frac{1}{\sqrt{x^3}} dx$$

We have $\frac{2}{\sqrt{x^3}} = \frac{1}{x^{3/2}}$

• $\int_0^1 \frac{1}{x^{3/2}} dx$ is divergent $\left(p = \frac{3}{2} > 1 \right)$

The integral $\int_0^1 \frac{2}{\sqrt{x^3}} dx$ is divergent by the Comparison Theorem.

5.4. Improper Integrals

Solution:

$$\text{d/} \int_0^{\pi} \frac{\sin^2 x}{\sqrt{x}} dx$$

We have $\frac{\sin^2 x}{\sqrt{x}} \leq \frac{1}{\sqrt{x}}$, for all $x \in (0, \pi)$

$$\bullet \int_0^{\pi} \frac{1}{\sqrt{x}} dx \text{ is convergent } \left(p = \frac{1}{2} \right)$$

The integral $\int_0^{\pi} \frac{\sin^2 x}{\sqrt{x}} dx$ is convergent by the Comparison Theorem.