

---

# Ordinary Differential Equations

— *EYNTKA* Series —

---

NATHANAEL CHWOJKO-SRAWLEY

September 6, 2026

---

---

# *Contents*

---

|                                                                |           |
|----------------------------------------------------------------|-----------|
| <b>Contents</b>                                                | <b>3</b>  |
| <b>1 First-Order Scalar Equations</b>                          | <b>9</b>  |
| 1.1 The General Initial Value Problem (IVP) . . . . .          | 9         |
| 1.2 Separable Equations . . . . .                              | 11        |
| 1.3 Linear Equations and Integrating Factors . . . . .         | 12        |
| 1.4 Exact Equations and Differential Forms . . . . .           | 13        |
| 1.5 Bernoulli Equations . . . . .                              | 14        |
| <b>2 Second-Order Linear Equations</b>                         | <b>17</b> |
| 2.1 Definitions and Theoretical Framework . . . . .            | 17        |
| 2.2 The Structure of Solutions . . . . .                       | 18        |
| 2.3 Homogeneous Equations with Constant Coefficients . . . . . | 20        |
| 2.4 Cauchy-Euler Equations . . . . .                           | 22        |
| 2.5 Inhomogeneous Equations . . . . .                          | 23        |
| <b>3 Systems of Linear Equations</b>                           | <b>25</b> |
| 3.1 Linear Systems and Geometry . . . . .                      | 25        |
| 3.2 The Matrix Exponential . . . . .                           | 26        |
| 3.3 Eigenvalues and Diagonalization . . . . .                  | 27        |
| 3.4 Defective Matrices and Jordan Forms . . . . .              | 28        |
| 3.5 Phase Portraits and Stability . . . . .                    | 29        |

|          |                                                                   |           |
|----------|-------------------------------------------------------------------|-----------|
| <b>4</b> | <b>The Fundamental Theorem of Ordinary Differential Equations</b> | <b>31</b> |
| 4.1      | Grönwall's Inequality . . . . .                                   | 31        |
| 4.2      | Picard-Lindelöf for Systems . . . . .                             | 33        |
| 4.3      | The Maximal Interval of Existence . . . . .                       | 36        |
| 4.4      | Continuous Dependence on Initial Data . . . . .                   | 39        |
| 4.5      | Differentiability of the Flow . . . . .                           | 41        |
| 4.6      | The Flow of an Autonomous System . . . . .                        | 44        |
| <b>5</b> | <b>Qualitative Theory and Stability</b>                           | <b>47</b> |
| 5.1      | Autonomous Systems and Phase Space . . . . .                      | 47        |
| 5.2      | Lyapunov's Direct Method . . . . .                                | 49        |
| 5.3      | Limit Cycles . . . . .                                            | 50        |
| <b>6</b> | <b>Boundary Value Problems and Sturm-Liouville Theory</b>         | <b>53</b> |
| 6.1      | Introduction to Boundary Value Problems . . . . .                 | 53        |
| 6.2      | The Sturm-Liouville Theorems . . . . .                            | 55        |
| 6.3      | The Fredholm Alternative . . . . .                                | 56        |
| 6.4      | Green's Functions . . . . .                                       | 56        |
| <b>7</b> | <b>Series Solutions and Special Functions</b>                     | <b>59</b> |
| 7.1      | Analytic Functions and Ordinary Points . . . . .                  | 59        |
| 7.2      | Regular Singular Points and the Method of Frobenius . . . . .     | 61        |

## Prerequisites

It is assumed that the reader is comfortable with the concepts introduced in a first course in analysis and linear algebra, importantly that smooth functions and vector-spaces are comfortable concepts.

---

# *Introduction Differential Equations*

---

To avoid super-script counting, the book shall work with smooth functions (or else the order of differentiability must be kept track). This shall not change the theory for ODE's and PDEs, especially since later it shall be shown that even working with non-smooth differentiable functions shall end up having the same theory (due to the advent of Distribution Theory <sup>1</sup>)

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ . When working smooth functions, we may take the derivative of the function. This gives the following function taking smooth functions and returning smooth functions:

$$f \mapsto f'$$

Due to the fundamental theorem of calculus, the inverse of this procedure is called *integration*, and gives a unique inverse up to a constant:

$$f \mapsto F + c \quad c \in \mathbb{R}$$

where  $F$  may be defined as  $F(x) = \int_t^x f(x)dx$  for appropriate  $t$ . Using this result, it is possible to find the smooth function that satisfies an arbitrary degree of differentiation, for example:

$$f^{(n)} = 0$$

which is a polynomial of degree  $n$ , or more generally

$$f^{(n)} = g \quad \text{equiv.} \quad f^{(n)} - g = 0$$

for some smooth functions  $g$  <sup>2</sup>.

Given this simple scenario, the question may arise if we may solve more complicated algebraic expression. For example, does there exist a solution to any of these:

$$f = f' \quad f' + f = 0 \quad f'' + f' + f = 0$$

---

<sup>1</sup>see chapter ref:HERE

<sup>2</sup>Note that the solution may not always come in an elementary form, that is expressible as a polynomial-combination and composition of  $x$ ,  $\log$ ,  $e^x$ , or a trigonometric identity

More precisely, we ask if any arbitrary algebraic expression on a smooth function  $f$  combined by using the 4 usual arithmetic operators ( $+$ ,  $-$ ,  $\times$ ,  $\text{div}$ ) along with differentiation yields a solution:

$$p(f) = 0$$

more commonly written as:

$$F(x, f(x), f'(x), \dots, f^{(n)}(x)) = 0$$

The study of the solution to such equations is called the study of *Ordinary Differential Equations* (ODEs). If we have a smooth function of the form  $f : \mathbb{R} \rightarrow \mathbb{R}^m$ , this is still the study of ODEs, where it is solving a *system of ODEs*. When working with multi-variable functions and partial derivatives, the study of the solution to such equations is called the study of *Partial Differential Equations* (PDEs). The study of ODEs can be thought of as a special case of the study of PDEs.

This book shall start with the study of ODEs in the first chapter, and proceed with the rest of the chapter with the study of PDEs. Only one chapter is dedicated to ODEs as they are a solved problem where most ODEs have a general solution that can be found algorithmically, while this is provably not the case for most PDEs.

From a more physical perspective, differential equations will give you how certain parameters relate to each, for PDEs it is usually over time. As a simple example, take:

$$\frac{\partial}{\partial t} = \frac{\partial}{\partial x} \quad \text{or} \quad \frac{\partial}{\partial t} - \frac{\partial}{\partial x} = 0 \quad (1)$$

What this equation is giving us is a restriction we require a function  $f(t, x)$  to obey, that is:

$$\frac{\partial f}{\partial t} - \frac{\partial f}{\partial x} = 0$$

For example  $f(x, t) = x + t$  would satisfy this equation. One key skill to learn when solving problems in PDEs is to be able to “read” such an equation. For example, equation (1) shows that the speed at which our “system” is evolving over time is identically propositional to the speed at which our “system” is moving. Sometimes, it is quite subtle what an equation may represent. For example:

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = 0$$

is very similar to equation (1), however it represents how a *wave* evolves over time, with behavior that is very different from equation (1).

The reason why finding solutions to PDEs is interesting to me is because the derivative can give lot's of local information about a system, and sometimes that's all the information we have. For example, it is much easier to describe the force of gravity as

$$F = ma$$

where  $a = dv/dt$ , and  $v = d(\text{displacement})/dt = dD/dt$ . We can use this equation to construct a gravitational field to how any object at any point will have a certain force on them represented by a vector pointing downwards times the mass of the object, but this “global solution” has a lot more information than is needed. More generally, it is often easier to come up with the “localizations”, since they are easier to define or control most of the time.

Also, after seeing how heat equations and diffusion equations are described I have a new motivation: I wonder if at some point these techniques of relating “local information” in describing how it’s dynamics evolve (in physics, it would be how it changes in time, in more general scenarios it can be a recursive function or a sequence) will at some point lend itself useful in solving more general problems. For example, let’s say we have a particularly complicated mathematical object. Let’s say we can solve it in some nice special cases, perhaps with some restricting assumptions. Can we then step-by-step evolve the problem (or model an evolution of the problem) to eliminate the restrictions? Can we localize in some way to simplify it, and glue that information together? These feel like modelling something like a PDE.

As a final motivator, every differential form has a “canonical representative” in that there always exists representative  $\omega$  such that  $\omega$  vanishes under the Laplacian:  $\Delta\omega = 0$  (functions, and more generally forms, that vanishes under the laplacian are called *harmonic*). This is the beginning of *Hodge Theory*, which is central in studying the cohomology groups of smooth manifolds. In the same vein as geometry but from another perspective, PDEs are also used to model *Ricci flow*, which is about the flow of a manifold given certain parameters as it changes across time.

## Ordinary Differential Equations

We focus on solving *Ordinary differential equations* with the hope that we may generalize to the non-linear case. For example:

$$5y'' + 1y' - 3y = 0$$

is a linear differential equation. Since it is equal to 0, it is called a *homogeneous* linear ODE. Since trigonometric functions have very simple derivative properties, they are also studied regularly. Though this might seem limiting, these in fact cover a satisfyingly large sample of scenario’s which we may encounter in physics. If your motivation is within pure mathematics, then having a standard repertoire to solve linear and trigonometric ODE’s allows us to use these as “local tools” to solve non-linear ODE’s or ODE’s approximated by a trigonometric sequence<sup>3</sup>. Here is small a list of types of equations and techniques used to solve ODE’s:

### 1. First order ODE:

- (a) Separable: if the equation can be written in the form:  $\frac{dy(x)}{dx} = \frac{m(x)}{n(y)}$ , or equivalently:

$$\frac{dy(x)}{dx} + \frac{m(x)}{n(y)} = 0$$

Then we can solve this ODE

- (b) Linear (Using an integral factor): If the equation can be written in the form:

$$\frac{dy(x)}{dx} + p(x)y(x) = g(x)$$

then we can solve this ODE

---

<sup>3</sup>If you know some more advanced analysis, think of functions being equal to a series of trigonometric functions or polynomials

## 2. Second order ODE:

- (a) If the ODE has constant coefficients, is linear, and homogeneous:

$$ay'' + by' + cy = 0$$

then we can solve this ode, with  $y(x) = Ae^{rx}$  for some  $r \in \mathbb{C}$  and  $A \in \mathbb{C}$ .

- (b) If the ODE is Equidimensional (Cauchy-Euler), linear, homogeneous:

$$ax^2y'' + bxy' + cy = 0$$

then the solution is of the form  $y(x) = Ax^p$  for some constant  $p \in \mathbb{C}$ .

There are many more that I've not listed here, I just wanted to give a list to give an idea of what was done.



---

# *First-Order Scalar Equations*

---

The introduction posed equations such as  $f = f'$  and  $f' + f = 0$  and asked whether solutions exist and how to find them. A scalar first-order equation  $y' = f(t, y)$  puts both questions in their sharpest form: it prescribes a slope at every point of the plane, so a solution is a curve whose tangent everywhere matches the prescribed direction. That reading makes existence plausible, since a curve can be grown from any starting point by following the prescribed slopes; Picard-Lindelöf turns the plausibility into a theorem, and a Lipschitz condition is the price.

Once solutions are known to exist and to be unique, the remaining question is which equations can be integrated in closed form. Separable, linear, and exact equations are the three classes that can, and the three techniques are one technique: each recognizes the equation as a derivative that has already been taken, and integrates it back.

## 1.1 The General Initial Value Problem (IVP)

We begin our study with the scalar first-order ordinary differential equation in the normal form:

$$y' = f(t, y) \tag{1.1}$$

where  $y(t)$  is the unknown function and  $f : U \rightarrow \mathbb{R}$  is a function defined on some open domain  $U \subseteq \mathbb{R}^2$ . Physically, if we view  $t$  as time and  $y$  as position,  $f$  prescribes the velocity of a particle based on its current time and location. Geometrically,  $f$  assigns a slope to every point in  $U$ , generating a *slope field* (or direction field). A solution  $\varphi(t)$  is a curve whose tangent at every point  $(t, \varphi(t))$  matches the slope prescribed by  $f$ .

**Definition 1.1.1: Initial Value Problem (IVP)**

An *Initial Value Problem* consists of an ODE combined with a specified value at a point  $t_0$ :

$$\begin{cases} y' = f(t, y) \\ y(t_0) = y_0 \end{cases}$$

A solution to the IVP on an interval  $I$  containing  $t_0$  is a differentiable function  $\varphi : I \rightarrow \mathbb{R}$  such that  $\varphi'(t) = f(t, \varphi(t))$  for all  $t \in I$  and  $\varphi(t_0) = y_0$ .

Before we attempt to solve these equations, we must address the fundamental questions of analysis: Does a solution exist? Is it unique?

**Theorem 1.1.2: Picard-Lindelöf Existence and Uniqueness**

Let  $D = \{(t, y) \in \mathbb{R}^2 \mid |t - t_0| \leq a, |y - y_0| \leq b\}$  be a rectangular region centered at the initial data. Suppose that:

1.  $f(t, y)$  is continuous on  $D$ .
2.  $f$  is *Lipschitz continuous* with respect to  $y$  on  $D$ . That is, there exists a constant  $L > 0$  such that:

$$|f(t, y_1) - f(t, y_2)| \leq L|y_1 - y_2| \quad \forall (t, y_1), (t, y_2) \in D$$

Then there exists a unique solution  $y(t)$  to the IVP defined on some interval  $|t - t_0| \leq h$  (where  $h \leq a$ ).

**Proof:**

We recast the initial value problem as a fixed-point problem. Integrating  $y' = f(t, y)$  from  $t_0$  to  $t$  and using  $y(t_0) = y_0$  shows that a continuous  $y$  solves the IVP on  $|t - t_0| \leq h$  if and only if it satisfies the integral equation

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

Solutions are therefore precisely the fixed points of the *Picard operator*  $(Ty)(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds$ .

*Step 1: a complete space on which  $T$  acts.* Since  $f$  is continuous on the compact rectangle  $D$ , it is bounded there, say  $|f| \leq M$ . Choose  $h > 0$  with

$$h \leq a, \quad Mh \leq b, \quad Lh < 1,$$

and let

$$X = \{y \in C([t_0 - h, t_0 + h]) \mid |y(t) - y_0| \leq b \text{ for all } t\},$$

with the sup metric  $d(y_1, y_2) = \sup_t |y_1(t) - y_2(t)|$ . Being a closed subset of the complete space  $C([t_0 - h, t_0 + h])$ , the space  $(X, d)$  is itself complete.

*Step 2:  $T$  maps  $X$  into itself.* For  $y \in X$  the function  $Ty$  is continuous, and

$$|(Ty)(t) - y_0| = \left| \int_{t_0}^t f(s, y(s)) ds \right| \leq M |t - t_0| \leq Mh \leq b,$$

so  $Ty \in X$ .

*Step 3:  $T$  is a contraction.* For  $y_1, y_2 \in X$  the integrand is controlled by the Lipschitz hypothesis,  $|f(s, y_1(s)) - f(s, y_2(s))| \leq L |y_1(s) - y_2(s)| \leq L d(y_1, y_2)$ ; integrating over the interval from  $t_0$  to  $t$ , of length  $|t - t_0| \leq h$ ,

$$|(Ty_1)(t) - (Ty_2)(t)| \leq \left| \int_{t_0}^t |f(s, y_1(s)) - f(s, y_2(s))| ds \right| \leq L |t - t_0| d(y_1, y_2) \leq Lh d(y_1, y_2).$$

Taking the supremum over  $t$  gives  $d(Ty_1, Ty_2) \leq (Lh) d(y_1, y_2)$ , and  $Lh < 1$ , so  $T$  is a contraction.

By the Banach contraction principle (*Metric and Function Spaces* [1]),  $T$  has a unique fixed point  $y \in X$ . This  $y$  is the unique solution of the integral equation, hence of the IVP; and since  $s \mapsto f(s, y(s))$  is continuous, the fundamental theorem of calculus shows  $y$  is continuously differentiable with  $y'(t) = f(t, y(t))$ .

Note that if  $\frac{\partial f}{\partial y}$  exists and is continuous, the Lipschitz condition is automatically satisfied locally.

## 1.2 Separable Equations

The simplest class of first-order equations are those where the dependence on  $t$  and  $y$  can be decoupled.

### Definition 1.2.1: Separable Equation

A first-order ODE is *separable* if it can be written in the form:

$$\frac{dy}{dt} = g(t)h(y)$$

or equivalently, if  $h(y) \neq 0$ :

$$\frac{1}{h(y)} dy = g(t) dt \tag{1.2}$$

The solution technique relies on the chain rule. Integrating (1.2) with respect to  $t$ :

$$\int \frac{1}{h(y(t))} \frac{dy}{dt} dt = \int g(t) dt \implies \int \frac{1}{h(y)} dy = \int g(t) dt$$

This yields an implicit relationship  $H(y) = G(t) + C$ .

### Example 1.2.2: Solving a Separable IVP

Solve the IVP:

$$y' = -2ty^2, \quad y(0) = 1$$

*Solution:* We separate variables (assuming  $y \neq 0$ ):

$$\frac{dy}{y^2} = -2t \, dt$$

Integrating both sides:

$$\int y^{-2} \, dy = \int -2t \, dt \implies -\frac{1}{y} = -t^2 + C \implies y(t) = \frac{1}{t^2 - C}$$

Using the initial condition  $y(0) = 1$ :

$$1 = \frac{1}{0 - C} \implies C = -1$$

Thus, the solution is  $y(t) = \frac{1}{t^2 + 1}$ . Note that this solution is defined for all  $t \in \mathbb{R}$ . Had the initial condition been  $y(0) = -1$ , we would have found  $y(t) = \frac{1}{t^2 - 1}$ , which blows up at  $t = \pm 1$ , illustrating that solutions to non-linear ODEs may exist only locally.

**Remark on Singular Solutions:** In the separation step, we divided by  $h(y)$ . The roots of  $h(y) = 0$  represent constant solutions (equilibrium solutions)  $y(t) = c$ . These must be checked separately to see if they satisfy the initial conditions or if solutions converge to them.

### 1.3 Linear Equations and Integrating Factors

Linear equations are the only class of ODEs for which a complete, closed-form general solution is always guaranteed (provided the coefficient functions are continuous).

#### Definition 1.3.1: First-Order Linear ODE

A first-order ODE is *linear* if it can be written as:

$$y' + p(t)y = g(t)$$

If  $g(t) \equiv 0$ , the equation is *homogeneous*; otherwise, it is *inhomogeneous*.

To solve this, we seek a function  $\mu(t)$ , called an *integrating factor*, such that multiplying the equation by  $\mu(t)$  transforms the left-hand side into a total derivative of a product.

$$\mu(t)y' + \mu(t)p(t)y = \mu(t)g(t)$$

We want the LHS to be  $\frac{d}{dt}[\mu(t)y] = \mu(t)y' + \mu'(t)y$ . Matching terms, we require:

$$\mu'(t) = \mu(t)p(t)$$

This is a separable equation for  $\mu$ :  $\frac{d\mu}{\mu} = p(t)dt$ , yielding  $\mu(t) = \exp\left(\int p(t) \, dt\right)$ .

**Theorem 1.3.2: General Solution of Linear Equations**

The general solution to  $y' + p(t)y = g(t)$  is given by:

$$y(t) = \frac{1}{\mu(t)} \left[ \int \mu(t)g(t) dt + C \right]$$

where  $\mu(t) = e^{\int p(t) dt}$ .

**Proof:**

Multiply by  $\mu(t)$ . Then  $\frac{d}{dt}[\mu(t)y] = \mu(t)g(t)$ . Integrating both sides yields  $\mu(t)y = \int \mu(t)g(t)dt + C$ . Dividing by  $\mu(t)$  gives the result.

**Example 1.3.3: Linear Equation with Discontinuous Forcing**

Consider  $y' + 2y = g(t)$  where  $y(0) = 0$  and  $g(t) = \chi_{[0,1]}$  (the characteristic function of  $[0, 1]$ ).

*Solution:* The integrating factor is  $\mu(t) = e^{\int 2dt} = e^{2t}$ .

$$\frac{d}{dt}[e^{2t}y] = e^{2t}g(t)$$

Integrating from 0 to  $t$ :

$$e^{2t}y(t) - e^0y(0) = \int_0^t e^{2s}g(s) ds$$

If  $0 \leq t \leq 1$ :

$$e^{2t}y(t) = \int_0^t e^{2s}(1) ds = \frac{1}{2}(e^{2t} - 1) \implies y(t) = \frac{1}{2}(1 - e^{-2t})$$

If  $t > 1$ :

$$e^{2t}y(t) = \int_0^1 e^{2s} ds + \int_1^t 0 ds = \frac{1}{2}(e^2 - 1) \implies y(t) = \frac{1}{2}(e^2 - 1)e^{-2t}$$

Notice that while  $g(t)$  is discontinuous, the solution  $y(t)$  is continuous, though  $y'(t)$  is discontinuous at  $t = 1$ . This “smoothing” property is characteristic of ODEs.

## 1.4 Exact Equations and Differential Forms

Here we take a slightly more sophisticated view. We can rewrite a first-order equation  $y' = f(x, y)$  symmetrically using differential forms:

$$M(x, y) dx + N(x, y) dy = 0$$

We ask: does there exist a scalar function (a potential)  $\psi(x, y)$  such that  $d\psi = Mdx + Ndy$ ? If so, the differential equation becomes  $d\psi = 0$ , implying the general solution is the level set  $\psi(x, y) = C$ .

**Definition 1.4.1: Exact Differential Equation**

The equation  $M(x, y)dx + N(x, y)dy = 0$  is *exact* in a region  $R$  if there exists a function  $\psi$  such that:

$$\frac{\partial \psi}{\partial x} = M \quad \text{and} \quad \frac{\partial \psi}{\partial y} = N$$

By equality of mixed partials ( $\psi_{xy} = \psi_{yx}$ ), a necessary condition for exactness is:

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \quad (1.3)$$

On a simply connected domain (like a rectangle or all of  $\mathbb{R}^2$ ), the Poincaré Lemma states this condition is also sufficient.

**Example 1.4.2: Solving an Exact Equation**

Solve  $(2xy + 3)dx + (x^2 - 1)dy = 0$ .

*Check Exactness:*

$$\partial_y(2xy + 3) = 2x \quad \partial_x(x^2 - 1) = 2x$$

The equation is exact. We seek  $\psi$  such that  $\psi_x = 2xy + 3$ . Integrating with respect to  $x$ :

$$\psi(x, y) = x^2y + 3x + h(y)$$

Now differentiate with respect to  $y$  and match  $N$ :

$$\psi_y = x^2 + h'(y) \stackrel{!}{=} x^2 - 1 \implies h'(y) = -1 \implies h(y) = -y$$

Thus  $\psi(x, y) = x^2y + 3x - y$ . The implicit solution is  $x^2y + 3x - y = C$ .

**Integrating Factors for Non-Exact Equations**

If  $Mdx + Ndy = 0$  is not exact, we may look for a function  $\mu(x, y)$  such that  $(\mu M)dx + (\mu N)dy = 0$  is exact. This requires:

$$(\mu M)_y = (\mu N)_x \implies \mu_y M + \mu M_y = \mu_x N + \mu N_x$$

This is a PDE for  $\mu$ , which is generally harder to solve than the original ODE! However, if we assume  $\mu$  depends only on  $x$  (i.e.,  $\mu_y = 0$ ), the condition simplifies to:

$$\frac{d\mu}{dx} N = \mu(M_y - N_x) \implies \frac{1}{\mu} \frac{d\mu}{dx} = \frac{M_y - N_x}{N}$$

If the RHS depends only on  $x$ , we can solve for  $\mu$ . A similar check can be done assuming  $\mu = \mu(y)$ .

**1.5 Bernoulli Equations**

We conclude this chapter with a classic non-linear equation that can be linearized. A *Bernoulli equation* has the form:

$$y' + p(t)y = g(t)y^n \quad n \in \mathbb{R}, n \neq 0, 1 \quad (1.4)$$

The substitution  $v = y^{1-n}$  linearizes this equation.

$$v' = (1-n)y^{-n}y'$$

Multiplying (1.4) by  $(1-n)y^{-n}$ :

$$(1-n)y^{-n}y' + (1-n)p(t)y^{1-n} = (1-n)g(t)$$

$$v' + (1-n)p(t)v = (1-n)g(t)$$

This is now a standard linear equation for  $v(t)$ , which can be solved using an integrating factor.





---

## *Second-Order Linear Equations*

---

Raising the order to two is where the subject becomes linear algebra. The solutions of a homogeneous linear equation are closed under addition and scalar multiplication, so they form a vector space, and dimension, independence, and basis all apply to it: the general solution stops being a formula and becomes a statement about a basis. The Wronskian is the determinant that detects independence of solutions, and Abel's identity computes it without knowing the solutions themselves.

For constant coefficients the basis can be written down, since exponentials reduce the equation to its characteristic polynomial and solving the equation reduces to factoring. Variation of parameters then handles an inhomogeneous forcing term by letting the coefficients in that basis expansion vary with  $t$ .

### 2.1 Definitions and Theoretical Framework

We begin by establishing the precise vocabulary used to classify these equations.

#### Definition 2.1.1: Second-Order Linear ODE

A second-order ordinary differential equation is called *linear* if it can be written in the form:

$$P(t)y'' + Q(t)y' + R(t)y = G(t) \tag{2.1}$$

where  $P, Q, R, G$  are continuous functions on an interval  $I$ , and  $P(t) \neq 0$  on  $I$ .

If we divide by  $P(t)$ , we obtain the *standard form*:

$$y'' + p(t)y' + q(t)y = g(t)$$

**Definition 2.1.2: Homogeneous vs. Inhomogeneous**

Referring to the standard form  $y'' + p(t)y' + q(t)y = g(t)$ :

- If  $g(t) \equiv 0$  for all  $t$ , the equation is called *homogeneous*.
- If  $g(t) \not\equiv 0$ , the equation is called *inhomogeneous* (or non-homogeneous).

The distinction is crucial: homogeneous equations describe the “natural” behavior of a system, while inhomogeneous equations describe the system’s response to an external “forcing” function  $g(t)$ . The following theorem guarantees that as long as the coefficient functions  $p(t)$  and  $q(t)$  are continuous, the solution extends across the entire interval of continuity.

**Theorem 2.1.3: Existence and Uniqueness for Linear Equations**

Consider the Initial Value Problem (IVP):

$$y'' + p(t)y' + q(t)y = g(t), \quad y(t_0) = y_0, \quad y'(t_0) = y'_0$$

If  $p(t)$ ,  $q(t)$ , and  $g(t)$  are continuous on an open interval  $I$  containing  $t_0$ , then there exists a unique solution  $y(t)$  defined on the entire interval  $I$ .

**Proof:**

This is a special case of the general Existence and Uniqueness theorem for linear systems (which we will cover in Chapter 3). We convert the second-order scalar equation into a first-order system by setting  $x_1 = y$  and  $x_2 = y'$ . This yields  $\vec{x}' = A(t)\vec{x} + \vec{g}(t)$ . Since the entries of  $A(t)$  (which are  $p(t)$  and  $q(t)$ ) are continuous on  $I$ , the Picard-Lindelöf theorem for systems guarantees a unique global solution on  $I$ .

**Example 2.1.4: Interval of Validity**

Determine the largest interval on which the IVP has a unique solution:

$$(t^2 - 4)y'' + 3ty' + (\cos t)y = e^t, \quad y(0) = 1, \quad y'(0) = 0$$

*Solution:* First, write the equation in standard form by dividing by  $(t^2 - 4)$ :

$$y'' + \frac{3t}{t^2 - 4}y' + \frac{\cos t}{t^2 - 4}y = \frac{e^t}{t^2 - 4}$$

The coefficients are discontinuous at  $t = \pm 2$ . The initial point is  $t_0 = 0$ . The largest interval containing 0 where all functions are continuous is  $(-2, 2)$ . Thus, the solution is guaranteed to exist uniquely on  $t \in (-2, 2)$ .

## 2.2 The Structure of Solutions

The set of solutions to a homogeneous linear ODE forms a vector space.

**Theorem 2.2.1: Principle of Superposition**

If  $y_1$  and  $y_2$  are solutions to the homogeneous equation  $L[y] = y'' + p(t)y' + q(t)y = 0$ , then any linear combination  $c_1y_1 + c_2y_2$  is also a solution.

**Proof:**

This follows from the linearity of the differential operator  $L$ .

$$L[c_1y_1 + c_2y_2] = c_1L[y_1] + c_2L[y_2] = c_1(0) + c_2(0) = 0$$

To describe *all* solutions, we need a “basis” for this solution space.

**Definition 2.2.2: Wronskian**

Let  $y_1, y_2$  be two differentiable functions. The *Wronskian* of  $y_1$  and  $y_2$  is the determinant:

$$W[y_1, y_2](t) = \det \begin{pmatrix} y_1(t) & y_2(t) \\ y_1'(t) & y_2'(t) \end{pmatrix} = y_1(t)y_2'(t) - y_2(t)y_1'(t)$$

The Wronskian is crucial because it allows us to test if solutions are linearly independent at a specific point, which is sufficient for linear independence on the entire interval.

**Theorem 2.2.3: Abel's Identity**

For the ODE  $y'' + p(t)y' + q(t)y = 0$ , the Wronskian  $W(t)$  satisfies the first-order differential equation  $W' = -p(t)W$ . Thus,

$$W(t) = Ce^{-\int p(t)dt}$$

where  $C$  is a constant determined by the initial conditions.

**Proof:**

Compute the derivative of the Wronskian definition:

$$\begin{aligned} W' &= (y_1y_2' - y_2y_1')' \\ &= (y_1'y_2' + y_1y_2'') - (y_2'y_1' + y_2y_1'') \\ &= y_1y_2'' - y_2y_1'' \end{aligned}$$

Since  $y_1$  and  $y_2$  satisfy the ODE  $y'' = -p(t)y' - q(t)y$ , substitute this into the expression for  $W'$ :

$$\begin{aligned} W' &= y_1[-p(t)y_2' - q(t)y_2] - y_2[-p(t)y_1' - q(t)y_1] \\ &= -p(t)(y_1y_2' - y_2y_1') - q(t)(y_1y_2 - y_2y_1) \\ &= -p(t)W - q(t)(0) \\ &= -p(t)W \end{aligned}$$

This is a separable first-order linear equation, the solution of which is  $W(t) = Ce^{-\int p(t)dt}$ .

**Theorem 2.2.4: General Solution of Homogeneous Equations**

Let  $y_1, y_2$  be two solutions to the homogeneous equation  $y'' + p(t)y' + q(t)y = 0$  on an interval  $I$ . If there exists a point  $t_0 \in I$  such that  $W[y_1, y_2](t_0) \neq 0$ , then  $\{y_1, y_2\}$  form a *Fundamental Set of Solutions*.

The general solution is:

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

**Proof:**

Let  $y(t)$  be any solution. We want to find  $c_1, c_2$  such that  $y(t) = c_1 y_1(t) + c_2 y_2(t)$ . Consider the initial conditions at  $t_0$ :  $y(t_0) = Y_0$  and  $y'(t_0) = Y'_0$ . We set up the linear system:

$$\begin{cases} c_1 y_1(t_0) + c_2 y_2(t_0) = Y_0 \\ c_1 y'_1(t_0) + c_2 y'_2(t_0) = Y'_0 \end{cases}$$

The determinant of the coefficient matrix is exactly  $W[y_1, y_2](t_0)$ . Since  $W(t_0) \neq 0$ , a unique pair  $(c_1, c_2)$  exists. By the Uniqueness Theorem, since  $c_1 y_1 + c_2 y_2$  satisfies the ODE and the same initial conditions as  $y(t)$ , they must be the same function.

**Example 2.2.5: Checking Linear Independence**

Show that  $y_1(t) = e^t$  and  $y_2(t) = e^{-t}$  form a fundamental set of solutions for  $y'' - y = 0$ .

*Solution:* Check if they solve the ODE:

$$y_1'' - y_1 = (e^t) - e^t = 0 \quad \checkmark \quad y_2'' - y_2 = (e^{-t}) - e^{-t} = 0 \quad \checkmark$$

Now compute the Wronskian:

$$W(t) = \det \begin{pmatrix} e^t & e^{-t} \\ e^t & -e^{-t} \end{pmatrix} = (e^t)(-e^{-t}) - (e^{-t})(e^t) = -1 - 1 = -2$$

Since  $W(t) = -2 \neq 0$ , the functions are linearly independent.

## 2.3 Homogeneous Equations with Constant Coefficients

We now turn to solving specific equations. The simplest case is when  $P, Q, R$  are constants.

$$ay'' + by' + cy = 0 \quad a, b, c \in \mathbb{R}, a \neq 0 \quad (2.2)$$

Motivated by the property that exponentials reproduce themselves under differentiation, we guess a solution of the form  $y = e^{rt}$ . Substituting this into (2.2):

$$a(r^2 e^{rt}) + b(r e^{rt}) + c(e^{rt}) = 0 \implies e^{rt}(ar^2 + br + c) = 0$$

Since  $e^{rt} \neq 0$ , we must solve the *Characteristic Equation*:

$$ar^2 + br + c = 0 \quad (2.3)$$

The roots are  $r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

### Theorem 2.3.1: Solutions to Constant Coefficient Equations

Depending on the discriminant  $\Delta = b^2 - 4ac$ , we have three cases for the general solution  $y(t)$ :

1. *Distinct Real Roots* ( $\Delta > 0$ ): Two roots  $r_1 \neq r_2$ .

$$y(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}$$

2. *Repeated Real Root* ( $\Delta = 0$ ): One root  $r = -b/2a$ .

$$y(t) = c_1 e^{rt} + c_2 t e^{rt}$$

3. *Complex Conjugate Roots* ( $\Delta < 0$ ): Roots  $r = \lambda \pm i\mu$ .

$$y(t) = e^{\lambda t} (c_1 \cos(\mu t) + c_2 \sin(\mu t))$$

#### Proof:

Case 1 follows directly from the characteristic equation yielding distinct  $r_1, r_2$ , and the Wronskian  $W[e^{r_1 t}, e^{r_2 t}] = (r_2 - r_1)e^{(r_1 + r_2)t} \neq 0$ .

Case 2 is proven via reduction of order. Let  $y_1 = e^{rt}$ . Guess  $y_2(t) = v(t)e^{rt}$ . Then:

$$y_2' = (v' + rv)e^{rt}, \quad y_2'' = (v'' + 2rv' + r^2v)e^{rt}$$

Substituting into the ODE  $ay'' + by' + cy = 0$ :

$$a(v'' + 2rv' + r^2v) + b(v' + rv) + cv = 0$$

Rearranging terms by derivatives of  $v$ :

$$av'' + (2ar + b)v' + (ar^2 + br + c)v = 0$$

Since  $r$  satisfies the characteristic equation, the coefficient of  $v$  is zero. Since  $r = -b/2a$  is a repeated root,  $2ar + b = 2a(-b/2a) + b = 0$ , so the coefficient of  $v'$  is also zero. We are left with  $av'' = 0 \implies v(t) = c_1 t + c_2$ . Choosing the simplest non-constant solution gives  $v(t) = t$ , so  $y_2 = te^{rt}$ .

Case 3 uses Euler's formula  $e^{(\lambda + i\mu)t} = e^{\lambda t}(\cos \mu t + i \sin \mu t)$ . Since the ODE is linear, the real and imaginary parts of a complex solution are also solutions.

### Example 2.3.2: Solving Constant Coefficient Problems

Find the general solution for:

1.  $y'' - 5y' + 6y = 0$
2.  $y'' + 4y' + 4y = 0$
3.  $y'' + 4y' + 13y = 0$

*Solutions:*

1. Char. Eq:  $r^2 - 5r + 6 = (r - 2)(r - 3) = 0$ . Roots:  $r_1 = 2, r_2 = 3$ .

$$y(t) = c_1 e^{2t} + c_2 e^{3t}$$

2. Char. Eq:  $r^2 + 4r + 4 = (r + 2)^2 = 0$ . Root:  $r = -2$  (repeated).

$$y(t) = c_1 e^{-2t} + c_2 t e^{-2t}$$

3. Char. Eq:  $r^2 + 4r + 13 = 0$ . Using quadratic formula:

$$r = \frac{-4 \pm \sqrt{16 - 52}}{2} = -2 \pm 3i$$

Here  $\lambda = -2, \mu = 3$ .

$$y(t) = e^{-2t}(c_1 \cos(3t) + c_2 \sin(3t))$$

## 2.4 Cauchy-Euler Equations

### Definition 2.4.1: Cauchy-Euler Equation

An *Equidimensional* or *Cauchy-Euler* equation is of the form:

$$at^2 y'' + bty' + cy = 0 \quad t > 0$$

We guess solutions of the form  $y = t^m$ . Substituting:

$$at^2(m(m-1)t^{m-2}) + bt(mt^{m-1}) + c(t^m) = 0$$

This simplifies to the auxiliary equation:

$$am(m-1) + bm + c = 0 \implies am^2 + (b-a)m + c = 0$$

### Theorem 2.4.2: Solutions to Cauchy-Euler Equations

Let  $m_1, m_2$  be the roots of  $am^2 + (b-a)m + c = 0$ .

1. *Distinct Real Roots:*  $y(t) = c_1 t^{m_1} + c_2 t^{m_2}$
2. *Repeated Real Root* ( $m_1 = m_2 = m$ ):  $y(t) = c_1 t^m + c_2 t^m \ln(t)$
3. *Complex Roots* ( $m = \lambda \pm i\mu$ ):

$$y(t) = t^\lambda [c_1 \cos(\mu \ln t) + c_2 \sin(\mu \ln t)]$$

**Proof:**

Substitute  $t = e^x$  (so  $x = \ln t$ ). Then use the chain rule:

$$\begin{aligned}\frac{dy}{dt} &= \frac{dy}{dx} \frac{dx}{dt} = \frac{1}{t} \frac{dy}{dx} \\ \frac{d^2y}{dt^2} &= \frac{d}{dt} \left( \frac{1}{t} \frac{dy}{dx} \right) = -\frac{1}{t^2} \frac{dy}{dx} + \frac{1}{t} \frac{d^2y}{dx^2} \frac{1}{t} = \frac{1}{t^2} \left( \frac{d^2y}{dx^2} - \frac{dy}{dx} \right)\end{aligned}$$

Substituting these into the original ODE yields a constant coefficient equation in  $x$ :

$$a \left( \frac{d^2y}{dx^2} - \frac{dy}{dx} \right) + b \frac{dy}{dx} + cy = 0 \implies a \frac{d^2y}{dx^2} + (b-a) \frac{dy}{dx} + cy = 0$$

The auxiliary equation for  $t$  is exactly the characteristic equation for  $x$ . The solutions in terms of  $x$  (e.g.,  $e^{m_1 x}$ ) become solutions in terms of  $t$  (e.g.,  $(e^x)^{m_1} = t^{m_1}$ ).

**Example 2.4.3: Cauchy-Euler Application**

Solve  $t^2 y'' - 3ty' + 3y = 0$  for  $t > 0$ .

*Solution:* Auxiliary equation:  $1m(m-1) - 3m + 3 = 0 \implies m^2 - 4m + 3 = 0$ . Factors to  $(m-1)(m-3) = 0$ . Roots are 1, 3.

$$y(t) = c_1 t + c_2 t^3$$

## 2.5 Inhomogeneous Equations

To solve  $y'' + p(t)y' + q(t)y = g(t)$ , we rely on the following structural theorem.

### Theorem 2.5.1: General Solution of Inhomogeneous Equations

The general solution to the inhomogeneous equation is:

$$y(t) = y_h(t) + y_p(t)$$

where  $y_h(t)$  is the general homogeneous solution and  $y_p(t)$  is *any* particular solution.

**Proof:**

Let  $L[y] = y'' + p(t)y' + q(t)y$ . Let  $Y$  be any solution such that  $L[Y] = g$ . Let  $y_p$  be a particular solution such that  $L[y_p] = g$ . Then  $L[Y - y_p] = L[Y] - L[y_p] = g - g = 0$ . Thus  $Y - y_p$  solves the homogeneous equation, so  $Y - y_p = y_h$ . Hence  $Y = y_h + y_p$ .

### Definition 2.5.2: Variation of Parameters

Given a fundamental set  $\{y_1, y_2\}$  for the homogeneous equation, we seek a particular solution of the form:

$$y_p(t) = u_1(t)y_1(t) + u_2(t)y_2(t)$$

**Theorem 2.5.3: Variation of Parameters Formula**

The particular solution is given by:

$$y_p(t) = -y_1(t) \int \frac{y_2(t)g(t)}{W(t)} dt + y_2(t) \int \frac{y_1(t)g(t)}{W(t)} dt$$

where  $W(t) = W[y_1, y_2](t)$ .

**Proof:**

We differentiate our guess  $y_p = u_1 y_1 + u_2 y_2$ :

$$y_p' = u_1' y_1 + u_1 y_1' + u_2' y_2 + u_2 y_2'$$

To simplify, we impose the condition:  $u_1' y_1 + u_2' y_2 = 0$ . Then  $y_p' = u_1 y_1' + u_2 y_2'$ . Differentiating again:

$$y_p'' = u_1' y_1' + u_1 y_1'' + u_2' y_2' + u_2 y_2''$$

Substitute  $y_p, y_p', y_p''$  into the ODE. Since  $y_1, y_2$  are homogeneous solutions, terms involving  $u_1(\dots)$  and  $u_2(\dots)$  vanish. We are left with:

$$u_1' y_1' + u_2' y_2' = g(t)$$

We now have a system of two linear equations for the unknowns  $u_1', u_2'$ :

$$\begin{cases} u_1' y_1 + u_2' y_2 = 0 \\ u_1' y_1' + u_2' y_2' = g(t) \end{cases}$$

Using Cramer's Rule, the determinant is the Wronskian  $W$ . Thus:

$$u_1' = \frac{\det \begin{pmatrix} 0 & y_2 \\ g(t) & y_2' \end{pmatrix}}{W} = \frac{-y_2 g(t)}{W}, \quad u_2' = \frac{\det \begin{pmatrix} y_1 & 0 \\ y_1' & g(t) \end{pmatrix}}{W} = \frac{y_1 g(t)}{W}$$

Integrating yields the result.

**Example 2.5.4: Variation of Parameters**

Solve  $y'' + y = \sec t$ .

*Solution:* Homogeneous solution:  $y_1 = \cos t, y_2 = \sin t$ .  $W = 1$ .

$$u_1(t) = - \int \sin t \sec t dt = - \int \tan t dt = - \ln |\sec t|$$

$$u_2(t) = \int \cos t \sec t dt = \int 1 dt = t$$

Particular solution:  $y_p = (-\ln |\sec t|) \cos t + (t) \sin t$ .



---

# *Systems of Linear Equations*

---

A scalar equation tracks one quantity, and most models track several at once: interacting species, coupled oscillators, the concentrations in a reaction. What couples them is that each rate depends on the others, so no one of the quantities satisfies an equation of its own. Collecting the unknowns into a vector  $\mathbf{x}(t)$  restores a single equation,  $\mathbf{x}' = A\mathbf{x}$ , now with matrix-valued coefficients.

The scalar solution  $y' = ay$ ,  $y = e^{at}y_0$ , then generalizes without change of form: the matrix exponential  $e^{At}$  solves the system for every  $A$ . The content of the chapter is computing  $e^{At}$ , which is where the eigenstructure of  $A$  enters, and where a defective  $A$  stops being a technicality and forces the Jordan form.

## 3.1 Linear Systems and Geometry

We define a first-order system of dimension  $n$  using vector notation. Let  $\mathbf{x}(t) \in \mathbb{R}^n$ .

### Definition 3.1.1: Linear System

A first-order *linear system* is an equation of the form:

$$\mathbf{x}'(t) = A(t)\mathbf{x}(t) + \mathbf{f}(t)$$

where  $A(t)$  is an  $n \times n$  matrix of continuous functions, and  $\mathbf{f}(t)$  is a vector of continuous forcing functions. If  $\mathbf{f}(t) \equiv \mathbf{0}$ , the system is *homogeneous*.

Any  $n$ -th order scalar linear ODE  $y^{(n)} + p_{n-1}y^{(n-1)} + \cdots + p_0y = 0$  can be converted into a first-order system of dimension  $n$  by introducing variables  $x_1 = y, x_2 = y', \dots, x_n = y^{(n-1)}$ .

**Theorem 3.1.2: Existence and Uniqueness for Systems**

If  $A(t)$  and  $\mathbf{f}(t)$  are continuous on an interval  $I$  containing  $t_0$ , then for any initial vector  $\mathbf{x}_0 \in \mathbb{R}^n$ , the IVP:

$$\mathbf{x}' = A(t)\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

has a unique solution  $\mathbf{x}(t)$  defined on the entire interval  $I$ .

**Proof:**

The right-hand side  $F(t, \mathbf{x}) = A(t)\mathbf{x} + \mathbf{f}(t)$  is continuous, and on any compact sub-interval of  $I$  the continuous matrix entries are bounded, say  $\|A(t)\| \leq M$  there, so  $F$  is Lipschitz continuous in  $\mathbf{x}$ :

$$\|F(t, \mathbf{x}) - F(t, \mathbf{y})\| = \|A(t)(\mathbf{x} - \mathbf{y})\| \leq \|A(t)\|\|\mathbf{x} - \mathbf{y}\| \leq M\|\mathbf{x} - \mathbf{y}\|$$

The Picard-Lindelöf theorem for systems (theorem 4.2.3, proved in chapter 4 by the contraction argument of theorem 1.1.2 and using nothing from this chapter) gives a unique local solution through  $(t_0, \mathbf{x}_0)$ , and since  $\|F(t, \mathbf{x})\| \leq \|\mathbf{f}(t)\| + \|A(t)\|\|\mathbf{x}\|$  grows at most linearly in  $\|\mathbf{x}\|$  with continuous coefficients, corollary 4.3.6 extends the solution uniquely to all of  $I$ . (If  $I$  has an endpoint, extend  $A$  and  $\mathbf{f}$  continuously past it by freezing their endpoint values, apply the open-interval statement, and restrict.)

**Example 3.1.3: Converting Scalar to System**

Convert the damped harmonic oscillator  $y'' + 2y' + 2y = 0$  into a system.

*Solution:* Let  $x_1 = y$  and  $x_2 = y'$ . Then  $x_1' = x_2$ . From the ODE,  $x_2' = y'' = -2y' - 2y = -2x_2 - 2x_1$ . The system is:

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ -2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

## 3.2 The Matrix Exponential

For the scalar case  $x' = ax$ , the solution is  $x(t) = x_0 e^{at}$ . We generalize this to systems  $\mathbf{x}' = A\mathbf{x}$  (where  $A$  is a constant matrix) by defining the exponential of a matrix.

**Definition 3.2.1: Matrix Exponential**

For an  $n \times n$  matrix  $A$ , the *matrix exponential* is defined by the convergent power series:

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!} = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots$$

This series converges absolutely for any matrix  $A$  (under any matrix norm), making  $e^A$  well-defined. Crucially, if  $AB = BA$ , then  $e^{A+B} = e^A e^B$ . However, if matrices do not commute, this identity fails.

**Theorem 3.2.2: Fundamental Matrix Solution**

The matrix function  $\Phi(t) = e^{At}$  is the unique solution to the matrix IVP:

$$\Phi'(t) = A\Phi(t), \quad \Phi(0) = I$$

Consequently, the solution to the vector IVP  $\mathbf{x}' = A\mathbf{x}, \mathbf{x}(0) = \mathbf{x}_0$  is given by:

$$\mathbf{x}(t) = e^{At}\mathbf{x}_0$$

**Proof:**

We differentiate the power series term by term (justified by uniform convergence):

$$\begin{aligned} \frac{d}{dt}e^{At} &= \frac{d}{dt} \left( I + At + \frac{A^2t^2}{2!} + \frac{A^3t^3}{3!} + \dots \right) \\ &= 0 + A + \frac{A^2(2t)}{2!} + \frac{A^3(3t^2)}{3!} + \dots \\ &= A + A^2t + \frac{A^3t^2}{2!} + \dots \\ &= A \left( I + At + \frac{A^2t^2}{2!} + \dots \right) = Ae^{At} \end{aligned}$$

At  $t = 0$ , the series becomes  $I + 0 + \dots = I$ . By the Uniqueness Theorem, this is the unique solution.

**Example 3.2.3: Computing Matrix Exponential (Diagonal Case)**

Compute  $e^{At}$  for  $A = \begin{pmatrix} 2 & 0 \\ 0 & -3 \end{pmatrix}$ .

*Solution:* Since  $A$  is diagonal,  $A^k = \begin{pmatrix} 2^k & 0 \\ 0 & (-3)^k \end{pmatrix}$ . Substituting into the series:

$$e^{At} = \sum \frac{t^k}{k!} \begin{pmatrix} 2^k & 0 \\ 0 & (-3)^k \end{pmatrix} = \begin{pmatrix} \sum \frac{(2t)^k}{k!} & 0 \\ 0 & \sum \frac{(-3t)^k}{k!} \end{pmatrix} = \begin{pmatrix} e^{2t} & 0 \\ 0 & e^{-3t} \end{pmatrix}$$

### 3.3 Eigenvalues and Diagonalization

In practice, we do not compute infinite sums. We use eigenvalues to diagonalize the matrix.

**Theorem 3.3.1: Diagonalizable Case**

If  $A$  has  $n$  linearly independent eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  with eigenvalues  $\lambda_1, \dots, \lambda_n$ , then  $A$  is diagonalizable:  $A = PDP^{-1}$ , where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$  and  $P = [\mathbf{v}_1 \dots \mathbf{v}_n]$ .

The solution is:

$$e^{At} = Pe^{Dt}P^{-1}$$

Explicitly, the general solution is:

$$\mathbf{x}(t) = c_1 e^{\lambda_1 t} \mathbf{v}_1 + \dots + c_n e^{\lambda_n t} \mathbf{v}_n$$

**Proof:**

Since  $A = PDP^{-1}$ , we have  $A^k = (PDP^{-1})(PDP^{-1}) \dots = PD^k P^{-1}$ . Thus:

$$e^{At} = \sum \frac{t^k}{k!} P D^k P^{-1} = P \left( \sum \frac{(Dt)^k}{k!} \right) P^{-1} = P e^{Dt} P^{-1}$$

$e^{Dt}$  is simply the diagonal matrix with entries  $e^{\lambda_i t}$ .

**Example 3.3.2: Real Distinct Eigenvalues**

Solve  $\mathbf{x}' = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \mathbf{x}$ .

*Solution:* Find eigenvalues:  $\det(A - \lambda I) = (1 - \lambda)^2 - 4 = \lambda^2 - 2\lambda - 3 = (\lambda - 3)(\lambda + 1)$ . Roots:  $\lambda_1 = 3, \lambda_2 = -1$ . Eigenvectors: For  $\lambda_1 = 3$ :  $\begin{pmatrix} -2 & 1 \\ 4 & -2 \end{pmatrix} \mathbf{v} = 0 \implies \mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ . For  $\lambda_2 = -1$ :  $\begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix} \mathbf{v} = 0 \implies \mathbf{v}_2 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$ . General Solution:

$$\mathbf{x}(t) = c_1 e^{3t} \begin{pmatrix} 1 \\ 2 \end{pmatrix} + c_2 e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix}$$

## 3.4 Defective Matrices and Jordan Forms

What if  $A$  does not have  $n$  independent eigenvectors? This occurs when eigenvalues are repeated and the geometric multiplicity is less than the algebraic multiplicity.

We can decompose any matrix  $A$  as  $A = \lambda I + N$ , where  $N$  is nilpotent (meaning  $N^k = 0$  for some  $k$ ). Since  $\lambda I$  and  $N$  commute,  $e^{At} = e^{\lambda I t} e^{Nt} = e^{\lambda t} e^{Nt}$ . Because  $N$  is nilpotent, the series for  $e^{Nt}$  *truncates* to a finite polynomial!

**Definition 3.4.1: Generalized Eigenvector**

A vector  $\mathbf{v}$  is a *generalized eigenvector* of rank  $k$  corresponding to eigenvalue  $\lambda$  if:

$$(A - \lambda I)^k \mathbf{v} = \mathbf{0} \quad \text{but} \quad (A - \lambda I)^{k-1} \mathbf{v} \neq \mathbf{0}$$

**Theorem 3.4.2: Solutions for Defective Matrices**

If an eigenvalue  $\lambda$  has algebraic multiplicity 2 but only 1 eigenvector  $\mathbf{v}$ , we find a second generalized eigenvector  $\mathbf{w}$  such that:

$$(A - \lambda I)\mathbf{w} = \mathbf{v}$$

The two linearly independent solutions are:

$$\mathbf{x}_1(t) = e^{\lambda t} \mathbf{v}$$

$$\mathbf{x}_2(t) = e^{\lambda t}(\mathbf{w} + t(A - \lambda I)\mathbf{w}) = e^{\lambda t}(\mathbf{w} + t\mathbf{v})$$

**Proof:**

We compute  $e^{At}\mathbf{w}$ . Since  $A\mathbf{w} = \lambda\mathbf{w} + \mathbf{v}$ , we recall the decomposition  $A = \lambda I + (A - \lambda I)$ .

$$e^{At}\mathbf{w} = e^{\lambda t}e^{(A-\lambda I)t}\mathbf{w} = e^{\lambda t} \left( I + t(A - \lambda I) + \frac{t^2}{2}(A - \lambda I)^2 + \dots \right) \mathbf{w}$$

Since  $\mathbf{w}$  is a generalized eigenvector of rank 2,  $(A - \lambda I)^2\mathbf{w} = (A - \lambda I)\mathbf{v} = 0$ . The series terminates after the second term:

$$e^{At}\mathbf{w} = e^{\lambda t}(\mathbf{w} + t\mathbf{v})$$

This proves  $\mathbf{x}_2$  is a solution.

**Example 3.4.3: Repeated Eigenvalues**

Solve  $\mathbf{x}' = \begin{pmatrix} 1 & -1 \\ 1 & 3 \end{pmatrix} \mathbf{x}$ .

*Solution:* Char. Poly:  $(\lambda - 1)(\lambda - 3) + 1 = \lambda^2 - 4\lambda + 4 = (\lambda - 2)^2$ .  $\lambda = 2$ . Eigenvector:  $(A - 2I)\mathbf{v} = \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix} \mathbf{v} = 0 \implies \mathbf{v} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ . Only one eigenvector. We need a generalized eigenvector  $\mathbf{w}$ :

$$(A - 2I)\mathbf{w} = \mathbf{v} \implies \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

$-w_1 - w_2 = 1$ . Let  $w_1 = 0$ , then  $w_2 = -1$ .  $\mathbf{w} = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$ . General Solution:

$$\mathbf{x}(t) = c_1 e^{2t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + c_2 e^{2t} \left( \begin{pmatrix} 0 \\ -1 \end{pmatrix} + t \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right)$$

### 3.5 Phase Portraits and Stability

For  $2 \times 2$  systems, we can classify the qualitative behavior of solutions based on the trace  $\tau = \text{tr}(A)$  and determinant  $\Delta = \det(A)$ . The characteristic equation is  $\lambda^2 - \tau\lambda + \Delta = 0$ .

- *Saddle Point* ( $\Delta < 0$ ): Real eigenvalues with opposite signs. Unstable.
- *Node* ( $\Delta > 0, \tau^2 - 4\Delta > 0$ ): Real eigenvalues with same sign.
  - Stable (Sink) if  $\tau < 0$ .
  - Unstable (Source) if  $\tau > 0$ .
- *Spiral* ( $\Delta > 0, \tau^2 - 4\Delta < 0$ ): Complex eigenvalues  $\alpha \pm i\beta$ .
  - Stable Spiral if  $\tau < 0$  ( $\alpha < 0$ ).
  - Unstable Spiral if  $\tau > 0$  ( $\alpha > 0$ ).
  - Center if  $\tau = 0$  (Purely imaginary eigenvalues).

**Example 3.5.1: Classifying Stability**

Classify the origin for the system with  $A = \begin{pmatrix} 0 & 1 \\ -5 & -2 \end{pmatrix}$ .

*Solution:* Trace  $\tau = -2$ . Determinant  $\Delta = 5$ . Discriminant:  $\tau^2 - 4\Delta = 4 - 20 = -16 < 0$ . Complex eigenvalues. Since  $\tau = -2 < 0$ , the real part is negative. Classification: *Stable Spiral*.

---

# *The Fundamental Theorem of Ordinary Differential Equations*

---

Theorem 1.1.2 settled existence and uniqueness for a single scalar equation on a rectangle, and the proof of theorem 3.1.2 rests on the systems form of the same theorem, which it quotes from this chapter. This chapter proves that form and then answers the three questions the local statement leaves open: how far each solution extends, what happens at a finite endpoint of its interval of existence, and how the solution moves when the initial condition does.

The answers assemble into what is rightly called the fundamental theorem of the subject. Every solution extends to a unique maximal open interval, and at a finite endpoint the solution must leave every compact subset of the domain, so the only obstructions to global existence are approaching the boundary of the domain and blowing up in norm. Nearby initial conditions produce solutions that diverge at most exponentially fast, and the dependence on the initial condition is as differentiable as the equation allows: a  $C^k$  right-hand side produces a  $C^k$  flow. This last statement is the deepest in the chapter, and the qualitative theory of chapter 5 rests on it, since the phase portrait is a picture of the flow. One inequality, Grönwall's, does most of the analytic work, and the chapter opens with it.

## 4.1 Grönwall's Inequality

The bounds available for a solution of  $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$  come from the integral form  $\mathbf{x}(t) = \mathbf{x}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{x}(s)) ds$ , so any estimate extracted from that identity contains the unknown  $\|\mathbf{x}(s)\|$  under the integral sign. *Grönwall's inequality* unwinds bounds of exactly this self-referential shape: a function dominated by a constant  $a$  plus an integral of itself against a weight  $b$  is dominated by  $a$  times the exponential of the integrated weight. A preliminary lemma records the triangle inequality for vector-valued integrals, which converts the vector integral form into the scalar bounds that Grönwall's

inequality applies to.

**Lemma 4.1.1: Triangle Inequality for Vector Integrals**

Let  $\mathbf{g} : [c, d] \rightarrow \mathbb{R}^n$  be continuous, and let  $\int_c^d \mathbf{g}(s) ds$  denote the entrywise integral. Then

$$\left\| \int_c^d \mathbf{g}(s) ds \right\| \leq \int_c^d \|\mathbf{g}(s)\| ds$$

**Proof:**

Set  $\mathbf{v} = \int_c^d \mathbf{g}(s) ds$ . If  $\mathbf{v} = \mathbf{0}$  the claim is trivial. Otherwise, by bilinearity of the inner product and the Cauchy-Schwarz inequality,

$$\|\mathbf{v}\|^2 = \left\langle \mathbf{v}, \int_c^d \mathbf{g}(s) ds \right\rangle = \int_c^d \langle \mathbf{v}, \mathbf{g}(s) \rangle ds \leq \int_c^d \|\mathbf{v}\| \|\mathbf{g}(s)\| ds = \|\mathbf{v}\| \int_c^d \|\mathbf{g}(s)\| ds$$

and dividing by  $\|\mathbf{v}\| > 0$  gives the claim.

The same bound holds for continuous matrix-valued integrands in the *operator norm*  $\|B\| = \max_{\|\mathbf{v}\|=1} \|B\mathbf{v}\|$ , which is the matrix norm used throughout this chapter: for every unit vector  $\mathbf{v}$ ,  $\|(\int B(s) ds)\mathbf{v}\| = \|\int B(s)\mathbf{v} ds\| \leq \int \|B(s)\mathbf{v}\| ds \leq \int \|B(s)\| ds$ . The operator norm is also submultiplicative,  $\|BC\| \leq \|B\| \|C\|$ , directly from the definition.

**Theorem 4.1.2: Grönwall's Inequality**

Let  $I$  be an interval,  $t_0 \in I$ , and let  $u, b : I \rightarrow [0, \infty)$  be continuous. If there is a constant  $a \geq 0$  with

$$u(t) \leq a + \left| \int_{t_0}^t b(s) u(s) ds \right| \quad \text{for all } t \in I$$

then

$$u(t) \leq a \exp \left( \left| \int_{t_0}^t b(s) ds \right| \right) \quad \text{for all } t \in I$$

**Proof:**

First take  $t \geq t_0$ . Define

$$v(t) = a + \int_{t_0}^t b(s) u(s) ds$$

so that  $v(t_0) = a$  and, by hypothesis,  $u \leq v$  on  $I \cap [t_0, \infty)$ , the integral being non-negative there. Since the integrand is continuous,  $v$  is continuously differentiable with  $v'(t) = b(t)u(t)$ , and  $b \geq 0$  together with  $u \leq v$  gives  $v'(t) \leq b(t)v(t)$ . Setting  $B(t) = \int_{t_0}^t b(s) ds$ ,

$$\frac{d}{dt} [v(t)e^{-B(t)}] = (v'(t) - b(t)v(t))e^{-B(t)} \leq 0$$

so  $v(t)e^{-B(t)} \leq v(t_0)e^{-B(t_0)} = a$ , and therefore  $u(t) \leq v(t) \leq ae^{B(t)}$ , which is the claim for  $t \geq t_0$ .



Now take  $t \leq t_0$ , where the hypothesis reads  $u(t) \leq a + \int_t^{t_0} b(s)u(s) ds$ . Define  $w(t) = a + \int_t^{t_0} b(s)u(s) ds$  and  $\tilde{B}(t) = \int_t^{t_0} b(s) ds$ , so that  $u \leq w$ ,  $w(t_0) = a$ , and  $w'(t) = -b(t)u(t) \geq -b(t)w(t)$ . Since  $\tilde{B}'(t) = -b(t)$ ,

$$\frac{d}{dt} [w(t)e^{-\tilde{B}(t)}] = (w'(t) + b(t)w(t))e^{-\tilde{B}(t)} \geq 0$$

so  $t \mapsto w(t)e^{-\tilde{B}(t)}$  is non-decreasing on  $I \cap (-\infty, t_0]$ , whence  $w(t)e^{-\tilde{B}(t)} \leq w(t_0)e^0 = a$  and  $u(t) \leq w(t) \leq ae^{\tilde{B}(t)}$ . In both cases the exponent equals  $|\int_{t_0}^t b(s) ds|$ , completing the proof.

Two special cases recur below. With  $a = 0$  the conclusion is  $u \equiv 0$ , and this is how uniqueness of solutions will be proved. With  $b$  equal to a constant  $L$ , the conclusion is  $u(t) \leq ae^{L|t-t_0|}$ , and this is the exponential-divergence bound behind continuous dependence on initial data.

## 4.2 Picard-Lindelöf for Systems

The scalar proof of theorem 1.1.2 was a fixed-point argument in a space of continuous functions, and it generalizes to systems with no change of strategy: the absolute value becomes a norm, and the Picard operator is unchanged. Two things nevertheless improve in the passage. First, the smallness condition  $Lh < 1$  that the scalar proof imposed on the interval disappears: iterating the Picard operator  $m$  times improves its contraction ratio to  $(Lh)^m/m!$ , which falls below 1 for large  $m$  regardless of  $Lh$ , so the length of the interval of existence is governed by the geometry of the rectangle alone. Second, uniqueness is upgraded from uniqueness within a chosen function space to uniqueness among all solutions. Throughout,  $\bar{B}(\mathbf{x}_0, b)$  and  $B(\mathbf{x}_0, b)$  denote the closed and open balls of radius  $b$  about  $\mathbf{x}_0$  in  $\mathbb{R}^n$ .

### Definition 4.2.1: Lipschitz and Locally Lipschitz

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  and  $\mathbf{f} : D \rightarrow \mathbb{R}^n$ . Then  $\mathbf{f}$  is *Lipschitz in  $\mathbf{x}$*  on  $D$  with constant  $L \geq 0$  if

$$\|\mathbf{f}(t, \mathbf{x}) - \mathbf{f}(t, \mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\| \quad \text{for all } (t, \mathbf{x}), (t, \mathbf{y}) \in D$$

If  $D$  is open,  $\mathbf{f}$  is *locally Lipschitz in  $\mathbf{x}$*  if every point of  $D$  has a neighborhood contained in  $D$  on which  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  with some constant.

Similarly, a function  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  on an open set  $U \subseteq \mathbb{R}^n$  is *locally Lipschitz* if every point of  $U$  has a neighborhood on which  $\|\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\|$  for some constant  $L$ . In that case  $(t, \mathbf{x}) \mapsto \mathbf{f}(\mathbf{x})$  is continuous (a Lipschitz function is continuous) and locally Lipschitz in  $\mathbf{x}$  on  $\mathbb{R} \times U$ .

### Lemma 4.2.2: $C^1$ Implies Locally Lipschitz

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open and let  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  have continuous partial derivatives in the  $\mathbf{x}$ -variables, with  $D_{\mathbf{x}}\mathbf{f} = (\partial f_i / \partial x_j)_{i,j}$  the corresponding  $n \times n$  matrix. If  $Q = [c, d] \times \bar{B}(\mathbf{x}_1, \rho) \subseteq D$ , then  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  on  $Q$  with constant  $L = \max_Q \|D_{\mathbf{x}}\mathbf{f}\|$ . In particular, every  $C^1$  function on an open set is locally Lipschitz in  $\mathbf{x}$ , since every point of an open  $D$  has such a product neighborhood  $Q$ .

**Proof:**

Fix  $(t, \mathbf{x}), (t, \mathbf{y}) \in Q$ . The segment  $\sigma \mapsto \mathbf{y} + \sigma(\mathbf{x} - \mathbf{y})$ ,  $\sigma \in [0, 1]$ , stays in  $\bar{B}(\mathbf{x}_1, \rho)$  because the ball is convex, so  $g(\sigma) = \mathbf{f}(t, \mathbf{y} + \sigma(\mathbf{x} - \mathbf{y}))$  is defined, and by the chain rule it is  $C^1$  with  $g'(\sigma) = D_{\mathbf{x}}\mathbf{f}(t, \mathbf{y} + \sigma(\mathbf{x} - \mathbf{y}))(\mathbf{x} - \mathbf{y})$ . The fundamental theorem of calculus, applied entrywise, gives

$$\mathbf{f}(t, \mathbf{x}) - \mathbf{f}(t, \mathbf{y}) = \int_0^1 D_{\mathbf{x}}\mathbf{f}(t, \mathbf{y} + \sigma(\mathbf{x} - \mathbf{y}))(\mathbf{x} - \mathbf{y}) d\sigma \quad (4.1)$$

and lemma 4.1.1 bounds the norm of the right side by  $\int_0^1 \|D_{\mathbf{x}}\mathbf{f}\| \|\mathbf{x} - \mathbf{y}\| d\sigma \leq L\|\mathbf{x} - \mathbf{y}\|$ .

**Theorem 4.2.3: Picard-Lindelöf for Systems**

Let  $R = \{(t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^n \mid |t - t_0| \leq a, \|\mathbf{x} - \mathbf{x}_0\| \leq b\}$  and suppose that:

1.  $\mathbf{f} : R \rightarrow \mathbb{R}^n$  is continuous, with  $\|\mathbf{f}\| \leq M$  on  $R$  for some constant  $M > 0$ .
2.  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  on  $R$  with constant  $L$ .

Set  $h = \min(a, b/M)$ . Then the initial value problem

$$\mathbf{x}' = \mathbf{f}(t, \mathbf{x}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

has exactly one solution on  $[t_0 - h, t_0 + h]$ , and it satisfies  $\|\mathbf{x}(t) - \mathbf{x}_0\| \leq b$  for all  $|t - t_0| \leq h$ .

**Proof:**

As in the scalar case, we recast the problem as a fixed-point problem: a continuous  $\mathbf{x}$  on  $[t_0 - h, t_0 + h]$  with values in  $\bar{B}(\mathbf{x}_0, b)$  solves the IVP if and only if it satisfies

$$\mathbf{x}(t) = \mathbf{x}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{x}(s)) ds$$

Indeed, a solution integrates to this identity, and conversely a continuous solution of the identity has a continuous integrand  $s \mapsto \mathbf{f}(s, \mathbf{x}(s))$ , so the fundamental theorem of calculus makes it continuously differentiable with  $\mathbf{x}'(t) = \mathbf{f}(t, \mathbf{x}(t))$  and  $\mathbf{x}(t_0) = \mathbf{x}_0$ . Solutions with values in the closed ball are therefore precisely the fixed points of the *Picard operator*

$$(T\mathbf{x})(t) = \mathbf{x}_0 + \int_{t_0}^t \mathbf{f}(s, \mathbf{x}(s)) ds$$

*Step 1: a complete space on which  $T$  acts.* Let

$$X = \{\mathbf{x} \in C([t_0 - h, t_0 + h], \mathbb{R}^n) \mid \|\mathbf{x}(t) - \mathbf{x}_0\| \leq b \text{ for all } t\}$$

with the sup metric  $d(\mathbf{x}, \mathbf{y}) = \sup_t \|\mathbf{x}(t) - \mathbf{y}(t)\|$ . The space  $C([t_0 - h, t_0 + h], \mathbb{R}^n)$  is complete, since a uniformly Cauchy sequence of continuous  $\mathbb{R}^n$ -valued functions converges uniformly to a continuous function, and  $X$  is a closed subset (a uniform limit of functions with values in the closed ball again has values in the closed ball), so  $(X, d)$  is complete. It is nonempty, containing the constant function  $\mathbf{x}_0$ .

*Step 2:  $T$  maps  $X$  into itself.* For  $\mathbf{x} \in X$  the function  $T\mathbf{x}$  is continuous, and lemma 4.1.1 gives

$$\|(T\mathbf{x})(t) - \mathbf{x}_0\| = \left\| \int_{t_0}^t \mathbf{f}(s, \mathbf{x}(s)) ds \right\| \leq M|t - t_0| \leq Mh \leq b$$

*Step 3: some iterate of  $T$  is a contraction.* We claim that for all  $m \geq 0$ , all  $\mathbf{x}, \mathbf{y} \in X$ , and all  $|t - t_0| \leq h$ ,

$$\|(T^m \mathbf{x})(t) - (T^m \mathbf{y})(t)\| \leq \frac{L^m |t - t_0|^m}{m!} d(\mathbf{x}, \mathbf{y})$$

For  $m = 0$  this is the definition of  $d$ . Assuming it for  $m$ , the Lipschitz hypothesis and lemma 4.1.1 give, for  $t \geq t_0$  (the case  $t \leq t_0$  is identical),

$$\begin{aligned} \|(T^{m+1} \mathbf{x})(t) - (T^{m+1} \mathbf{y})(t)\| &\leq \int_{t_0}^t \|\mathbf{f}(s, (T^m \mathbf{x})(s)) - \mathbf{f}(s, (T^m \mathbf{y})(s))\| ds \\ &\leq \int_{t_0}^t L \frac{L^m (s - t_0)^m}{m!} d(\mathbf{x}, \mathbf{y}) ds = \frac{L^{m+1} (t - t_0)^{m+1}}{(m+1)!} d(\mathbf{x}, \mathbf{y}) \end{aligned}$$

Taking suprema,  $d(T^m \mathbf{x}, T^m \mathbf{y}) \leq \frac{(Lh)^m}{m!} d(\mathbf{x}, \mathbf{y})$ . Since  $(Lh)^m/m!$  is the general term of the convergent series for  $e^{Lh}$ , it tends to 0, so there is an  $m$  with  $(Lh)^m/m! < 1$ , and for this  $m$  the iterate  $T^m : X \rightarrow X$  is a contraction.

*Step 4:  $T$  has exactly one fixed point in  $X$ .* By the Banach contraction principle ([1, Banach Contraction Principle]),  $T^m$  has a unique fixed point  $\mathbf{x}^* \in X$ . Then  $T^m(T\mathbf{x}^*) = T(T^m \mathbf{x}^*) = T\mathbf{x}^*$ , so  $T\mathbf{x}^*$  is also a fixed point of  $T^m$ , and uniqueness forces  $T\mathbf{x}^* = \mathbf{x}^*$ . Conversely, every fixed point of  $T$  is a fixed point of  $T^m$ , so  $\mathbf{x}^*$  is the unique fixed point of  $T$  in  $X$ . By the equivalence above,  $\mathbf{x}^*$  is the unique solution of the IVP among continuous functions with values in  $\bar{B}(\mathbf{x}_0, b)$ .

*Step 5: every solution stays in the ball.* Let  $\mathbf{y}$  be any solution of the IVP on  $[t_0 - h, t_0 + h]$ . It suffices to show  $\|\mathbf{y}(t) - \mathbf{x}_0\| \leq b$  for all  $t$ , for then  $\mathbf{y} \in X$  and Step 4 gives  $\mathbf{y} = \mathbf{x}^*$ . Consider  $t \geq t_0$  (the other side is symmetric) and let

$$t^* = \sup \{ \tau \in [t_0, t_0 + h] \mid \|\mathbf{y}(s) - \mathbf{x}_0\| \leq b \text{ for all } s \in [t_0, \tau] \}$$

The set is nonempty ( $\tau = t_0$  belongs) and, by continuity,  $\|\mathbf{y}(s) - \mathbf{x}_0\| \leq b$  on all of  $[t_0, t^*]$ , so the graph of  $\mathbf{y}$  over  $[t_0, t^*]$  lies in  $R$  and the integral identity applies there:  $\|\mathbf{y}(t) - \mathbf{x}_0\| \leq M(t - t_0)$  for  $t \in [t_0, t^*]$ . If  $t^* < t_0 + h$ , then  $\|\mathbf{y}(t^*) - \mathbf{x}_0\| \leq M(t^* - t_0) < Mh \leq b$ , and continuity keeps  $\|\mathbf{y} - \mathbf{x}_0\| \leq b$  on  $[t_0, t^* + \eta]$  for some  $\eta > 0$ , contradicting the definition of  $t^*$ . Hence  $t^* = t_0 + h$ , completing the proof.

#### Lemma 4.2.4: Global Uniqueness

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open and  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  continuous and locally Lipschitz in  $\mathbf{x}$ . If  $\mathbf{x} : J_1 \rightarrow \mathbb{R}^n$  and  $\mathbf{y} : J_2 \rightarrow \mathbb{R}^n$  are solutions of  $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$  on intervals  $J_1, J_2$  with graphs in  $D$ , and  $\mathbf{x}(t_1) = \mathbf{y}(t_1)$  for some  $t_1 \in J_1 \cap J_2$ , then  $\mathbf{x} = \mathbf{y}$  on all of  $J_1 \cap J_2$ .

**Proof:**

The set  $J = J_1 \cap J_2$  is an interval containing  $t_1$ , and  $A = \{t \in J \mid \mathbf{x}(t) = \mathbf{y}(t)\}$  is nonempty and closed in  $J$  by continuity. We show  $A$  is also open in  $J$ : since an interval is connected,  $A = J$  then follows.

Let  $t_2 \in A$ . Local Lipschitz continuity provides a neighborhood of  $(t_2, \mathbf{x}(t_2))$  in  $D$  on which  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  with some constant  $L_2$ , and by continuity of  $\mathbf{x}$  and  $\mathbf{y}$  there is  $\eta > 0$  such that the graphs of both solutions over  $J' = J \cap (t_2 - \eta, t_2 + \eta)$  lie in that neighborhood. For  $t \in J'$ , integrating both equations from  $t_2$  and subtracting,

$$\mathbf{x}(t) - \mathbf{y}(t) = \int_{t_2}^t (\mathbf{f}(s, \mathbf{x}(s)) - \mathbf{f}(s, \mathbf{y}(s))) ds$$

so  $u(t) = \|\mathbf{x}(t) - \mathbf{y}(t)\|$  satisfies  $u(t) \leq \int_{t_2}^t L_2 u(s) ds$  by lemma 4.1.1. This is the hypothesis of theorem 4.1.2 with  $a = 0$ , so  $u \equiv 0$  on  $J'$ , that is,  $J' \subseteq A$ .

**Corollary 4.2.5: Existence and Uniqueness for Autonomous Systems**

Let  $U \subseteq \mathbb{R}^n$  be open and let  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  be locally Lipschitz, which holds in particular whenever  $\mathbf{f}$  is  $C^1$  or smooth (lemma 4.2.2). Then for every  $\mathbf{x}_0 \in U$  there is an  $\varepsilon > 0$  such that the autonomous initial value problem

$$\mathbf{x}' = \mathbf{f}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0$$

has a solution on  $(-\varepsilon, \varepsilon)$ , and any two solutions agree on the intersection of their intervals of definition.

**Proof:**

Choose  $b > 0$  such that  $\bar{B}(\mathbf{x}_0, b) \subseteq U$  and  $\mathbf{f}$  is Lipschitz with some constant  $L$  on  $\bar{B}(\mathbf{x}_0, b)$ , shrinking the neighborhood from the definition of locally Lipschitz to a closed ball. A bound  $\|\mathbf{f}\| \leq M$  with  $M > 0$  exists on the ball, since a Lipschitz function is continuous and the ball is compact. The function  $F(t, \mathbf{x}) = \mathbf{f}(\mathbf{x})$  on  $R = [-1, 1] \times \bar{B}(\mathbf{x}_0, b)$  is continuous, bounded by  $M$ , and Lipschitz in  $\mathbf{x}$  with constant  $L$ , so theorem 4.2.3 with  $a = 1$  gives a solution on  $[-\varepsilon, \varepsilon]$  for  $\varepsilon = \min(1, b/M)$ . Agreement of any two solutions is lemma 4.2.4 applied to  $F$  on  $D = \mathbb{R} \times U$ , which is continuous and locally Lipschitz in  $\mathbf{x}$  by definition 4.2.1.

### 4.3 The Maximal Interval of Existence

Local solutions glue. If one solution of an initial value problem is defined on  $J_1$  and another on  $J_2$ , they agree on  $J_1 \cap J_2$  by lemma 4.2.4, so together they define a solution on  $J_1 \cup J_2$ . Pushing this to its limit produces a single solution on the largest possible interval, and global existence becomes the question of what that interval is. The escape theorem below describes the only way it can fail to be everything: at a finite endpoint the solution must leave every compact subset of the domain, so it either approaches the boundary of the domain or blows up in norm.

**Theorem 4.3.1: The Maximal Interval of Existence**

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open,  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  continuous and locally Lipschitz in  $\mathbf{x}$ , and  $(t_0, \mathbf{x}_0) \in D$ . Among all solutions of

$$\mathbf{x}' = \mathbf{f}(t, \mathbf{x}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

there is exactly one, the *maximal solution*, of which every other solution is a restriction. Its domain  $J = (\alpha, \beta)$ , the *maximal interval of existence*, is an open interval containing  $t_0$ , with  $\alpha \in [-\infty, t_0)$  and  $\beta \in (t_0, +\infty]$ .

**Proof:**

At least one solution exists:  $(t_0, \mathbf{x}_0)$  has a neighborhood in  $D$  on which  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$ , this neighborhood contains a rectangle  $[t_0 - a, t_0 + a] \times \bar{B}(\mathbf{x}_0, b)$ , and theorem 4.2.3 applies to it.

Let  $J$  be the union of the domains of all solutions of the IVP, an interval containing  $t_0$  since each domain is one. For  $t \in J$  define  $\mathbf{x}(t) = \mathbf{y}(t)$ , where  $\mathbf{y}$  is any solution defined at  $t$ : by lemma 4.2.4 any two such solutions agree at  $t$ , so  $\mathbf{x}$  is well defined, and it restricts to  $\mathbf{y}$  on the domain of every solution  $\mathbf{y}$ .

Next,  $J$  is open and  $\mathbf{x}$  is a solution on  $J$ . Let  $t \in J$  and let  $\mathbf{y}$  be a solution of the IVP defined at  $t$ . The point  $(t, \mathbf{y}(t))$  lies in  $D$ , so the first paragraph, applied at this point, gives a solution  $\mathbf{z}$  of the equation with  $\mathbf{z}(t) = \mathbf{y}(t)$  on some  $[t - \eta, t + \eta]$ . By lemma 4.2.4,  $\mathbf{y}$  and  $\mathbf{z}$  agree on their common domain, so the function equal to  $\mathbf{y}$  on the domain of  $\mathbf{y}$  and to  $\mathbf{z}$  on  $[t - \eta, t + \eta]$  is well defined, and it is a solution of the IVP: near every point of its domain it coincides with  $\mathbf{y}$  or with  $\mathbf{z}$ , and it passes through  $(t_0, \mathbf{x}_0)$  because  $\mathbf{y}$  does. Its domain contains  $(t - \eta, t + \eta)$ , so  $J$  contains a neighborhood of  $t$ , and  $\mathbf{x}$ , agreeing with this solution near  $t$ , is differentiable at  $t$  with  $\mathbf{x}'(t) = \mathbf{f}(t, \mathbf{x}(t))$ .

Finally,  $\mathbf{x}$  extends every solution by construction, and a second solution with the same property would have domain  $J$  and agree with  $\mathbf{x}$  there by lemma 4.2.4, completing the proof.

**Lemma 4.3.2: Uniform Existence Time on Compact Sets**

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open and  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  continuous and locally Lipschitz in  $\mathbf{x}$ . For every compact  $K \subseteq D$  there is a  $\delta > 0$  such that, for every  $(t_1, \mathbf{x}_1) \in K$ , the initial value problem with data  $\mathbf{x}(t_1) = \mathbf{x}_1$  has a solution on  $[t_1 - \delta, t_1 + \delta]$ .

**Proof:**

Every point  $p = (s, \mathbf{y}) \in K$  has, inside a neighborhood on which  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$ , a rectangle

$$R_p = [s - 2a_p, s + 2a_p] \times \bar{B}(\mathbf{y}, 2b_p) \subseteq D$$

with Lipschitz constant  $L_p$ , and  $\mathbf{f}$  is bounded on the compact  $R_p$ , say  $\|\mathbf{f}\| \leq M_p$  with  $M_p > 0$ . The open sets  $V_p = (s - a_p, s + a_p) \times B(\mathbf{y}, b_p)$  cover  $K$ , so finitely many  $V_{p_1}, \dots, V_{p_N}$  do. Set

$$\delta = \min_{1 \leq i \leq N} \min \left( a_{p_i}, \frac{b_{p_i}}{M_{p_i}} \right)$$

Given  $(t_1, \mathbf{x}_1) \in K$ , choose  $i$  with  $(t_1, \mathbf{x}_1) \in V_{p_i}$ , and abbreviate  $a_i, b_i, M_i, L_i$  for the constants of  $p_i$ . Then

$$[t_1 - a_i, t_1 + a_i] \times \bar{B}(\mathbf{x}_1, b_i) \subseteq R_{p_i} \subseteq D$$

since  $|t - t_1| \leq a_i$  and  $|t_1 - s| < a_i$  force  $|t - s| < 2a_i$ , and likewise in the spatial factor. On this rectangle  $\mathbf{f}$  is continuous, bounded by  $M_i$ , and Lipschitz in  $\mathbf{x}$  with constant  $L_i$ , so theorem 4.2.3 yields a solution through  $(t_1, \mathbf{x}_1)$  on  $[t_1 - h_i, t_1 + h_i]$  with  $h_i = \min(a_i, b_i/M_i) \geq \delta$ .

### Theorem 4.3.3: Escape from Compact Sets

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open,  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  continuous and locally Lipschitz in  $\mathbf{x}$ , and let  $\mathbf{x} : (\alpha, \beta) \rightarrow \mathbb{R}^n$  be a maximal solution. If  $\beta < \infty$ , then for every compact  $K \subseteq D$  there is a  $\tau < \beta$  such that

$$(t, \mathbf{x}(t)) \notin K \quad \text{for all } t \in (\tau, \beta)$$

Symmetrically, if  $\alpha > -\infty$ , the graph of the solution leaves every compact subset of  $D$  as  $t \rightarrow \alpha^+$ .

#### Proof:

Suppose instead that some compact  $K \subseteq D$  contains graph points at times arbitrarily close to  $\beta$ : there is a sequence  $t_k \rightarrow \beta^-$  with  $(t_k, \mathbf{x}(t_k)) \in K$  for all  $k$ . Let  $\delta$  be the uniform existence time of lemma 4.3.2 for  $K$ , and fix  $k$  with  $\beta - t_k < \delta$ . The lemma gives a solution  $\mathbf{z}$  on  $[t_k - \delta, t_k + \delta]$  with  $\mathbf{z}(t_k) = \mathbf{x}(t_k)$ . Since  $\mathbf{x}$  and  $\mathbf{z}$  agree at  $t_k$ , lemma 4.2.4 makes them agree on their common domain, so

$$\mathbf{w}(t) = \begin{cases} \mathbf{x}(t) & t \in (\alpha, \beta) \\ \mathbf{z}(t) & t \in [t_k, t_k + \delta] \end{cases}$$

is well defined on  $(\alpha, t_k + \delta)$ , the two branches agreeing on  $[t_k, \beta)$ . Near every point of  $(\alpha, t_k + \delta)$  the function  $\mathbf{w}$  coincides with  $\mathbf{x}$  or with  $\mathbf{z}$ , so  $\mathbf{w}$  is a solution through  $(t_0, \mathbf{x}_0)$  whose domain properly contains  $(\alpha, \beta)$ , because  $t_k + \delta > \beta$ . This contradicts theorem 4.3.1, which makes every solution a restriction of  $\mathbf{x}$ .

For the statement at  $\alpha$ , apply the case already proved to  $\mathbf{y}(t) = \mathbf{x}(-t)$ , the maximal solution of  $\mathbf{y}' = \mathbf{g}(t, \mathbf{y})$  for  $\mathbf{g}(t, \mathbf{y}) = -\mathbf{f}(-t, \mathbf{y})$  on the reflected domain  $\{(t, \mathbf{y}) \mid (-t, \mathbf{y}) \in D\}$ : the function  $\mathbf{g}$  is continuous and locally Lipschitz in  $\mathbf{y}$ , the upper endpoint of the maximal interval of  $\mathbf{y}$  is  $-\alpha < \infty$ , and compact subsets of  $D$  reflect to compact subsets of the reflected domain.

#### Example 4.3.4: Blow-up in Finite Time

The scalar problem  $x' = x^2$ ,  $x(0) = 1$  has right-hand side  $f(t, x) = x^2$ , which is  $C^1$  on  $D = \mathbb{R} \times \mathbb{R}$ , and separation of variables (section 1.2) gives the maximal solution

$$x(t) = \frac{1}{1-t} \quad \text{on } (\alpha, \beta) = (-\infty, 1)$$

The right endpoint is finite even though the domain is all of  $\mathbb{R} \times \mathbb{R}$ , and theorem 4.3.3 says what must happen there. When  $D = \mathbb{R} \times \mathbb{R}^n$ , every set  $[t_0, \beta] \times \bar{B}(\mathbf{0}, R)$  is compact, so a maximal solution with  $\beta < \infty$  must leave each of them, forcing  $\|\mathbf{x}(t)\| \rightarrow \infty$  as  $t \rightarrow \beta^-$ : on a domain without boundary, escape from compact sets is blow-up in norm. Indeed  $x(t) = 1/(1-t) \rightarrow \infty$  as  $t \rightarrow 1^-$ .

### Corollary 4.3.5: Escape for Autonomous Systems

Let  $U \subseteq \mathbb{R}^n$  be open,  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  locally Lipschitz, and let  $\mathbf{x} : (\alpha, \beta) \rightarrow U$  be a maximal solution of  $\mathbf{x}' = \mathbf{f}(\mathbf{x})$ . If  $\beta < \infty$ , then for every compact  $K \subseteq U$  there is a  $\tau < \beta$  with  $\mathbf{x}(t) \notin K$  for all  $t \in (\tau, \beta)$ . In particular, a maximal solution that remains in a compact subset of  $U$  on  $[t_1, \beta)$  for some  $t_1$  has  $\beta = \infty$ . Symmetric statements hold at  $\alpha$ .

#### Proof:

View the system on  $D = \mathbb{R} \times U$  (definition 4.2.1). If there were  $t_k \rightarrow \beta^-$  with  $\mathbf{x}(t_k) \in K$ , then for any fixed  $t_1 \in (\alpha, \beta)$  the points  $(t_k, \mathbf{x}(t_k))$  with  $t_k \geq t_1$  would lie in  $[t_1, \beta] \times K$ , a compact subset of  $D$  because  $\beta < \infty$ , contradicting theorem 4.3.3. The second statement is the contrapositive, and the statements at  $\alpha$  follow by the reflection used in theorem 4.3.3.

### Corollary 4.3.6: Global Existence Under Linear Growth

Let  $I \subseteq \mathbb{R}$  be an open interval and let  $\mathbf{f} : I \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  be continuous and locally Lipschitz in  $\mathbf{x}$ . Suppose there are continuous functions  $c_1, c_2 : I \rightarrow [0, \infty)$  with

$$\|\mathbf{f}(t, \mathbf{x})\| \leq c_1(t) + c_2(t)\|\mathbf{x}\| \quad \text{for all } (t, \mathbf{x}) \in I \times \mathbb{R}^n$$

Then every maximal solution of  $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$  is defined on all of  $I$ .

#### Proof:

Let  $\mathbf{x}$  be a maximal solution on  $(\alpha, \beta)$  and suppose  $\beta < \sup I$ , so in particular  $\beta < \infty$ . Fix  $t_0 \in (\alpha, \beta)$ . The interval  $[t_0, \beta]$  is a compact subset of  $I$ , so  $c_1 \leq C_1$  and  $c_2 \leq C_2$  on it for some constants. For  $t \in [t_0, \beta)$ , the integral form and lemma 4.1.1 give

$$\|\mathbf{x}(t)\| \leq \|\mathbf{x}(t_0)\| + \int_{t_0}^t (C_1 + C_2\|\mathbf{x}(s)\|) ds \leq (\|\mathbf{x}(t_0)\| + C_1(\beta - t_0)) + \int_{t_0}^t C_2\|\mathbf{x}(s)\| ds$$

and theorem 4.1.2 on the interval  $[t_0, \beta)$  yields

$$\|\mathbf{x}(t)\| \leq (\|\mathbf{x}(t_0)\| + C_1(\beta - t_0)) e^{C_2(\beta - t_0)} = \rho$$

The graph of  $\mathbf{x}$  over  $[t_0, \beta)$  therefore stays inside  $[t_0, \beta] \times \bar{B}(\mathbf{0}, \rho)$ , a compact subset of  $I \times \mathbb{R}^n$ , contradicting theorem 4.3.3. Hence  $\beta = \sup I$ , and  $\alpha = \inf I$  follows symmetrically.

## 4.4 Continuous Dependence on Initial Data

Uniqueness attaches one solution to each initial condition, and the next question is the stability of that assignment: whether solutions launched from nearby initial values remain nearby. Grönwall's inequality gives the sharp answer, exponential divergence at worst, and applying the answer uniformly over a neighborhood produces the local flow, the map sending time and initial value jointly to the solution's position.

**Theorem 4.4.1: Continuous Dependence on Initial Data**

Let  $S \subseteq \mathbb{R} \times \mathbb{R}^n$  and let  $\mathbf{f} : S \rightarrow \mathbb{R}^n$  be continuous and Lipschitz in  $\mathbf{x}$  on  $S$  with constant  $L$ . If  $\mathbf{x}, \mathbf{y} : J \rightarrow \mathbb{R}^n$  are solutions of  $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$  on an interval  $J \ni t_0$  whose graphs lie in  $S$ , then

$$\|\mathbf{x}(t) - \mathbf{y}(t)\| \leq e^{L|t-t_0|} \|\mathbf{x}(t_0) - \mathbf{y}(t_0)\| \quad \text{for all } t \in J$$

**Proof:**

Integrating both equations from  $t_0$  and subtracting, then applying lemma 4.1.1 and the Lipschitz bound, the function  $u(t) = \|\mathbf{x}(t) - \mathbf{y}(t)\|$  satisfies

$$u(t) \leq u(t_0) + \left| \int_{t_0}^t L u(s) ds \right| \quad \text{for all } t \in J$$

and theorem 4.1.2 with  $a = u(t_0)$  and  $b \equiv L$  gives  $u(t) \leq u(t_0) e^{L|t-t_0|}$ .

**Theorem 4.4.2: The Local Flow**

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open,  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  continuous and locally Lipschitz in  $\mathbf{x}$ , and  $(t_0, \mathbf{x}_0) \in D$ . Choose a rectangle

$$R_0 = [t_0 - a, t_0 + a] \times \bar{B}(\mathbf{x}_0, b) \subseteq D$$

on which  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  with constant  $L$  (the Lipschitz neighborhood of  $(t_0, \mathbf{x}_0)$  contains such a rectangle), and a bound  $\|\mathbf{f}\| \leq M$  on  $R_0$  with  $M > 0$ . Set

$$r = \frac{b}{2}, \quad \delta = \min \left( a, \frac{b}{2M} \right), \quad I_\delta = [t_0 - \delta, t_0 + \delta]$$

Then for every  $\mathbf{y} \in \bar{B}(\mathbf{x}_0, r)$  the initial value problem with data  $\mathbf{x}(t_0) = \mathbf{y}$  has exactly one solution  $\varphi(\cdot, \mathbf{y})$  on  $I_\delta$ , its values lie in  $\bar{B}(\mathbf{x}_0, b)$ , and it is the restriction to  $I_\delta$  of the maximal solution through  $(t_0, \mathbf{y})$ . The resulting map  $\varphi : I_\delta \times \bar{B}(\mathbf{x}_0, r) \rightarrow \mathbb{R}^n$ , the *local flow* at  $(t_0, \mathbf{x}_0)$ , satisfies

$$\|\varphi(t, \mathbf{y}) - \varphi(s, \mathbf{z})\| \leq M|t - s| + e^{L\delta} \|\mathbf{y} - \mathbf{z}\|$$

for all  $t, s \in I_\delta$  and  $\mathbf{y}, \mathbf{z} \in \bar{B}(\mathbf{x}_0, r)$ . In particular  $\varphi$  is continuous.

**Proof:**

For  $\mathbf{y} \in \bar{B}(\mathbf{x}_0, r)$  the rectangle  $R_{\mathbf{y}} = [t_0 - a, t_0 + a] \times \bar{B}(\mathbf{y}, b/2)$  is contained in  $R_0$ , since  $\|\mathbf{z} - \mathbf{y}\| \leq b/2$  and  $\|\mathbf{y} - \mathbf{x}_0\| \leq b/2$  force  $\|\mathbf{z} - \mathbf{x}_0\| \leq b$ . On  $R_{\mathbf{y}}$  the function  $\mathbf{f}$  is continuous, bounded by  $M$ , and Lipschitz in  $\mathbf{x}$  with constant  $L$ , so theorem 4.2.3, with  $b/2$  in place of  $b$ , gives exactly one solution  $\varphi(\cdot, \mathbf{y})$  of the problem with data  $\mathbf{x}(t_0) = \mathbf{y}$  on the interval  $|t - t_0| \leq \min(a, b/2M) = \delta$ , with values in  $\bar{B}(\mathbf{y}, b/2) \subseteq \bar{B}(\mathbf{x}_0, b)$ . It is the restriction of the maximal solution by theorem 4.3.1.

For the estimate, split the difference into a time step and a space step:

$$\|\varphi(t, \mathbf{y}) - \varphi(s, \mathbf{z})\| \leq \|\varphi(t, \mathbf{z}) - \varphi(s, \mathbf{z})\| + \|\varphi(t, \mathbf{y}) - \varphi(t, \mathbf{z})\|$$



For the first term,  $\|\varphi(t, \mathbf{z}) - \varphi(s, \mathbf{z})\| = \left\| \int_s^t \mathbf{f}(\tau, \varphi(\tau, \mathbf{z})) d\tau \right\| \leq M|t - s|$  by lemma 4.1.1. For the second, the graphs of  $\varphi(\cdot, \mathbf{y})$  and  $\varphi(\cdot, \mathbf{z})$  over  $I_\delta$  lie in  $I_\delta \times \bar{B}(\mathbf{x}_0, b) \subseteq R_0$ , where  $\mathbf{f}$  is Lipschitz in  $\mathbf{x}$  with constant  $L$ , so theorem 4.4.1 bounds it by  $e^{L|t-t_0|}\|\mathbf{y} - \mathbf{z}\| \leq e^{L\delta}\|\mathbf{y} - \mathbf{z}\|$ .

## 4.5 Differentiability of the Flow

Continuity of the local flow is settled, and differentiability in the initial condition is the harder question. To see what the derivative would have to be, suppose for a moment that  $\varphi$  is continuously differentiable and differentiate the defining conditions

$$\partial_t \varphi(t, \mathbf{y}) = \mathbf{f}(t, \varphi(t, \mathbf{y})), \quad \varphi(t_0, \mathbf{y}) = \mathbf{y}$$

with respect to  $\mathbf{y}$ , exchanging the two differentiations: the matrix  $J(t) = D_{\mathbf{y}}\varphi(t, \mathbf{y})$  would have to satisfy

$$J'(t) = D_{\mathbf{x}}\mathbf{f}(t, \varphi(t, \mathbf{y})) J(t), \quad J(t_0) = I \quad (4.2)$$

where  $I$  is the identity matrix. This linear system is the *variational equation* of  $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$  along the solution  $\varphi(\cdot, \mathbf{y})$ : it transports an infinitesimal displacement of the initial value along the flow. Nothing in this computation is justified yet, since the derivative being differentiated is the one whose existence is in question. The proof below therefore runs in the opposite direction. The variational equation is a linear system, so it has a unique solution  $J$  by the theory already in hand, and a Grönwall estimate on the difference quotient shows that this  $J$  is the derivative. That estimate is the hardest point of the chapter.

### Theorem 4.5.1: Differentiability in the Initial Condition

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open, let  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  be  $C^1$ , and let  $(t_0, \mathbf{x}_0) \in D$ . Construct the local flow  $\varphi : I_\delta \times \bar{B}(\mathbf{x}_0, r) \rightarrow \mathbb{R}^n$  of theorem 4.4.2 on a rectangle  $R_0 = [t_0 - a, t_0 + a] \times \bar{B}(\mathbf{x}_0, b) \subseteq D$ , taking as Lipschitz constant  $L = \max_{R_0} \|D_{\mathbf{x}}\mathbf{f}\|$ , as lemma 4.2.2 permits. Write  $B^\circ = (t_0 - \delta, t_0 + \delta) \times B(\mathbf{x}_0, r)$  for the open box. Then:

1. For every  $\mathbf{y} \in B(\mathbf{x}_0, r)$  the variational equation (4.2) along  $\varphi(\cdot, \mathbf{y})$  has a unique solution  $J(\cdot; \mathbf{y})$  on  $(t_0 - \delta, t_0 + \delta)$ .
2. For every  $(t, \mathbf{y}) \in B^\circ$  the derivative of  $\varphi(t, \cdot)$  at  $\mathbf{y}$  exists and equals  $J(t; \mathbf{y})$ .
3.  $\varphi$  is  $C^1$  on  $B^\circ$ , with  $\partial_t \varphi(t, \mathbf{y}) = \mathbf{f}(t, \varphi(t, \mathbf{y}))$  and  $D_{\mathbf{y}}\varphi(t, \mathbf{y}) = J(t; \mathbf{y})$ .

#### Proof:

*Step 0: the candidate derivative exists.* For  $\mathbf{y} \in B(\mathbf{x}_0, r)$  define the matrix-valued coefficient

$$A(t; \mathbf{y}) = D_{\mathbf{x}}\mathbf{f}(t, \varphi(t, \mathbf{y}))$$

The map  $(t, \mathbf{y}) \mapsto A(t; \mathbf{y})$  is continuous on  $I_\delta \times \bar{B}(\mathbf{x}_0, r)$ , being a composition of the continuous maps  $\varphi$  (theorem 4.4.2) and  $D_{\mathbf{x}}\mathbf{f}$ , and  $\|A(t; \mathbf{y})\| \leq L$  because  $(t, \varphi(t, \mathbf{y})) \in R_0$  always. Identify the space of  $n \times n$  matrices with  $\mathbb{R}^{n^2}$ . For fixed  $\mathbf{y}$ , the right-hand side  $(t, J) \mapsto A(t; \mathbf{y})J$  of (4.2) is continuous on  $(t_0 - \delta, t_0 + \delta) \times \mathbb{R}^{n^2}$  and Lipschitz in  $J$  with constant  $L$ : column by column,  $\|A(t; \mathbf{y})(J - K)\mathbf{e}_i\| \leq L\|(J - K)\mathbf{e}_i\|$ , and squaring and summing over columns bounds

the Euclidean norm of  $A(J - K)$  in  $\mathbb{R}^{n^2}$  by  $L$  times that of  $J - K$ . The same bound shows the right-hand side grows at most linearly in  $J$ . Theorem 4.3.1 and corollary 4.3.6 therefore give a unique solution  $J(\cdot; \mathbf{y})$  of (4.2) on all of  $(t_0 - \delta, t_0 + \delta)$ , proving (1). Integrating (4.2) and applying the matrix form of lemma 4.1.1,

$$\|J(t; \mathbf{y})\| \leq 1 + \left| \int_{t_0}^t L \|J(s; \mathbf{y})\| ds \right|$$

so theorem 4.1.2 gives the uniform bound  $\|J(t; \mathbf{y})\| \leq e^{L\delta}$ .

*Step 1: the Taylor remainder of  $\mathbf{f}$ , with a uniform modulus.* Since  $D_{\mathbf{x}}\mathbf{f}$  is continuous on the compact set  $R_0$ , it is uniformly continuous there. Define, for  $\rho \geq 0$ ,

$$\omega(\rho) = \sup \{ \|D_{\mathbf{x}}\mathbf{f}(s, \mathbf{z}) - D_{\mathbf{x}}\mathbf{f}(s, \mathbf{z}')\| \mid (s, \mathbf{z}), (s, \mathbf{z}') \in R_0, \|\mathbf{z} - \mathbf{z}'\| \leq \rho \}$$

Then  $\omega$  is non-decreasing and  $\omega(\rho) \rightarrow 0$  as  $\rho \rightarrow 0^+$ . For  $(s, \mathbf{p}), (s, \mathbf{q}) \in R_0$ , subtracting  $D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p})(\mathbf{q} - \mathbf{p})$  from the segment identity (4.1) gives

$$\mathbf{f}(s, \mathbf{q}) - \mathbf{f}(s, \mathbf{p}) - D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p})(\mathbf{q} - \mathbf{p}) = \int_0^1 \left[ D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p} + \sigma(\mathbf{q} - \mathbf{p})) - D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p}) \right] (\mathbf{q} - \mathbf{p}) d\sigma$$

Every point  $\mathbf{p} + \sigma(\mathbf{q} - \mathbf{p})$  lies in  $\bar{B}(\mathbf{x}_0, b)$  by convexity of the ball, at distance  $\sigma\|\mathbf{q} - \mathbf{p}\| \leq \|\mathbf{q} - \mathbf{p}\|$  from  $\mathbf{p}$ , so lemma 4.1.1 bounds the remainder:

$$\|\mathbf{f}(s, \mathbf{q}) - \mathbf{f}(s, \mathbf{p}) - D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p})(\mathbf{q} - \mathbf{p})\| \leq \omega(\|\mathbf{q} - \mathbf{p}\|) \|\mathbf{q} - \mathbf{p}\| \quad (4.3)$$

*Step 2: the difference quotient.* Fix  $\mathbf{y} \in B(\mathbf{x}_0, r)$  and let  $\mathbf{v} \in \mathbb{R}^n$  satisfy  $0 < \|\mathbf{v}\| \leq r - \|\mathbf{y} - \mathbf{x}_0\|$ , so that  $\mathbf{y} + \mathbf{v} \in \bar{B}(\mathbf{x}_0, r)$  and  $\varphi(\cdot, \mathbf{y} + \mathbf{v})$  is defined. Abbreviate  $\mathbf{p}(t) = \varphi(t, \mathbf{y})$  and  $\mathbf{q}(t) = \varphi(t, \mathbf{y} + \mathbf{v})$ , both with values in  $\bar{B}(\mathbf{x}_0, b)$ , and set

$$\boldsymbol{\theta}(t) = \mathbf{q}(t) - \mathbf{p}(t) - J(t; \mathbf{y}) \mathbf{v}$$

The claim to be proved is that  $\|\boldsymbol{\theta}(t)\| = o(\|\mathbf{v}\|)$  uniformly in  $t$ . Integrating the differential equations satisfied by  $\mathbf{q}$ ,  $\mathbf{p}$ , and  $J(\cdot; \mathbf{y})\mathbf{v}$  from  $t_0$ , and using  $\mathbf{q}(t_0) - \mathbf{p}(t_0) = \mathbf{v} = J(t_0; \mathbf{y})\mathbf{v}$ , so that  $\boldsymbol{\theta}(t_0) = \mathbf{0}$ ,

$$\boldsymbol{\theta}(t) = \int_{t_0}^t \left[ \mathbf{f}(s, \mathbf{q}(s)) - \mathbf{f}(s, \mathbf{p}(s)) - A(s; \mathbf{y}) J(s; \mathbf{y}) \mathbf{v} \right] ds$$

Insert and remove the middle term  $A(s; \mathbf{y})(\mathbf{q}(s) - \mathbf{p}(s))$ : since  $A(s; \mathbf{y}) = D_{\mathbf{x}}\mathbf{f}(s, \mathbf{p}(s))$ , the integrand becomes the Taylor remainder of (4.3) at  $\mathbf{p}(s)$  plus  $A(s; \mathbf{y})\boldsymbol{\theta}(s)$ . By theorem 4.4.1, exactly as in the proof of theorem 4.4.2,  $\|\mathbf{q}(s) - \mathbf{p}(s)\| \leq e^{L\delta}\|\mathbf{v}\|$ , so (4.3) and the monotonicity of  $\omega$  bound the remainder in norm by  $\omega(e^{L\delta}\|\mathbf{v}\|) e^{L\delta}\|\mathbf{v}\|$ . Therefore, by lemma 4.1.1 and  $\|A\| \leq L$ ,

$$\|\boldsymbol{\theta}(t)\| \leq \delta e^{L\delta} \omega(e^{L\delta}\|\mathbf{v}\|) \|\mathbf{v}\| + \left| \int_{t_0}^t L \|\boldsymbol{\theta}(s)\| ds \right|$$

and theorem 4.1.2 gives, for all  $t \in (t_0 - \delta, t_0 + \delta)$ ,

$$\|\boldsymbol{\theta}(t)\| \leq \delta e^{2L\delta} \omega(e^{L\delta}\|\mathbf{v}\|) \|\mathbf{v}\|$$

As  $\mathbf{v} \rightarrow \mathbf{0}$  the factor  $\omega(e^{L\delta}\|\mathbf{v}\|)$  tends to 0, so

$$\varphi(t, \mathbf{y} + \mathbf{v}) - \varphi(t, \mathbf{y}) - J(t; \mathbf{y}) \mathbf{v} = o(\|\mathbf{v}\|) \quad \text{uniformly in } t$$

By the definition of the total derivative,  $D_{\mathbf{y}}\varphi(t, \mathbf{y})$  exists and equals  $J(t; \mathbf{y})$ , proving (2).

*Step 3: continuity of  $(t, \mathbf{y}) \mapsto J(t; \mathbf{y})$ .* For  $\mathbf{y}, \mathbf{z} \in B(\mathbf{x}_0, r)$  set  $\Delta(t) = J(t; \mathbf{y}) - J(t; \mathbf{z})$ , so that  $\Delta(t_0) = 0$  and, integrating the two variational equations and inserting  $A(s; \mathbf{y})J(s; \mathbf{z})$ ,

$$\Delta(t) = \int_{t_0}^t \left[ A(s; \mathbf{y}) \Delta(s) + (A(s; \mathbf{y}) - A(s; \mathbf{z})) J(s; \mathbf{z}) \right] ds$$

Both  $(s, \varphi(s, \mathbf{y}))$  and  $(s, \varphi(s, \mathbf{z}))$  lie in  $R_0$ , and  $\|\varphi(s, \mathbf{y}) - \varphi(s, \mathbf{z})\| \leq e^{L\delta} \|\mathbf{y} - \mathbf{z}\|$  by theorem 4.4.2, so  $\|A(s; \mathbf{y}) - A(s; \mathbf{z})\| \leq \omega(e^{L\delta} \|\mathbf{y} - \mathbf{z}\|)$ . Combining with the bound  $\|J(s; \mathbf{z})\| \leq e^{L\delta}$  of Step 0 and applying theorem 4.1.2 as in Step 2,

$$\|J(t; \mathbf{y}) - J(t; \mathbf{z})\| \leq \delta e^{2L\delta} \omega(e^{L\delta} \|\mathbf{y} - \mathbf{z}\|) \quad \text{for all } t$$

which tends to 0 as  $\mathbf{z} \rightarrow \mathbf{y}$ , uniformly in  $t$ . In the time direction,  $\|J(t; \mathbf{y}) - J(s; \mathbf{y})\| \leq \int_s^t \|A\| \|J\| d\tau \leq Le^{L\delta} |t - s|$ . The two bounds together give joint continuity of  $J$  on  $B^\circ$ .

*Step 4:  $\varphi$  is  $C^1$ .* Fix  $(t, \mathbf{y}) \in B^\circ$ . The claim is that  $\varphi$  is differentiable at  $(t, \mathbf{y})$  with

$$\varphi(t + \tau, \mathbf{y} + \mathbf{v}) - \varphi(t, \mathbf{y}) = \mathbf{f}(t, \varphi(t, \mathbf{y})) \tau + J(t; \mathbf{y}) \mathbf{v} + o(|\tau| + \|\mathbf{v}\|)$$

Split the increment as

$$\left[ \varphi(t + \tau, \mathbf{y} + \mathbf{v}) - \varphi(t, \mathbf{y} + \mathbf{v}) - \mathbf{f}(t, \varphi(t, \mathbf{y})) \tau \right] + \left[ \varphi(t, \mathbf{y} + \mathbf{v}) - \varphi(t, \mathbf{y}) - J(t; \mathbf{y}) \mathbf{v} \right]$$

The second bracket is  $o(\|\mathbf{v}\|)$  by Step 2. The first equals

$$\int_t^{t+\tau} \left[ \mathbf{f}(s, \varphi(s, \mathbf{y} + \mathbf{v})) - \mathbf{f}(t, \varphi(t, \mathbf{y})) \right] ds$$

whose norm is at most  $|\tau|$  times the supremum of the integrand's norm over  $|s - t| \leq |\tau|$ . The map  $(s, \mathbf{z}) \mapsto \mathbf{f}(s, \varphi(s, \mathbf{z}))$  is continuous at  $(t, \mathbf{y})$ , so that supremum tends to 0 as  $(\tau, \mathbf{v}) \rightarrow (0, \mathbf{0})$ , making the first bracket  $o(|\tau|)$ . Hence  $\varphi$  is differentiable at every point of  $B^\circ$ , and its derivative, with blocks  $\mathbf{f}(t, \varphi(t, \mathbf{y}))$  and  $J(t; \mathbf{y})$ , is continuous on  $B^\circ$  by theorem 4.4.2 and Step 3. This proves (3), completing the proof.

### Theorem 4.5.2: $C^k$ Dependence on Initial Data

Let  $D \subseteq \mathbb{R} \times \mathbb{R}^n$  be open, let  $k \geq 1$  be finite, and let  $\mathbf{f} : D \rightarrow \mathbb{R}^n$  be  $C^k$ . Then every  $(t_1, \mathbf{x}_1) \in D$  admits  $\delta, r > 0$  such that the flow map  $\varphi(t, \mathbf{y})$ , the value at time  $t$  of the maximal solution with data  $\mathbf{x}(t_1) = \mathbf{y}$ , is defined and  $C^k$  on the open box  $(t_1 - \delta, t_1 + \delta) \times B(\mathbf{x}_1, r)$ , with  $D_{\mathbf{y}}\varphi$  solving the variational equation (4.2).

#### Proof:

The proof is by induction on  $k$ , with the statement quantified over all dimensions  $n$  and all  $C^k$  right-hand sides. The case  $k = 1$  is theorem 4.5.1. Let  $k \geq 2$  and assume the statement for  $k - 1$ .

Consider the *extended system* for the pair  $(\mathbf{y}, J) \in \mathbb{R}^n \times \mathbb{R}^{n \times n}$ :

$$\mathbf{y}' = \mathbf{f}(t, \mathbf{y}), \quad J' = D_{\mathbf{x}}\mathbf{f}(t, \mathbf{y}) J$$

on the open set  $\tilde{D} = \{(t, \mathbf{y}, J) \mid (t, \mathbf{y}) \in D, J \in \mathbb{R}^{n \times n}\} \subseteq \mathbb{R} \times \mathbb{R}^{n+n^2}$ . Its right-hand side  $F(t, \mathbf{y}, J) = (\mathbf{f}(t, \mathbf{y}), D_{\mathbf{x}}\mathbf{f}(t, \mathbf{y})J)$  is  $C^{k-1}$ : the entries of  $\mathbf{f}$  are  $C^k$ , the entries of  $D_{\mathbf{x}}\mathbf{f}$  are  $C^{k-1}$ , and the second block is a sum of products of entries of  $D_{\mathbf{x}}\mathbf{f}$  with entries of  $J$ . Since  $k-1 \geq 1$ , the function  $F$  is in particular  $C^1$ , so lemma 4.2.2 and lemma 4.2.4 apply to the extended system.

By the inductive hypothesis applied to the extended system at the point  $(t_1, (\mathbf{x}_1, I))$ , with  $I$  the identity matrix, there are  $\tilde{\delta}, \tilde{r} > 0$  such that its flow map  $\Psi$ , sending  $(t, (\mathbf{y}_0, J_0))$  to the value at time  $t$  of the maximal solution with data  $(\mathbf{y}_0, J_0)$  at  $t_1$ , is defined and  $C^{k-1}$  on  $(t_1 - \tilde{\delta}, t_1 + \tilde{\delta}) \times B((\mathbf{x}_1, I), \tilde{r})$ . Shrink the box of theorem 4.5.1 at  $(t_1, \mathbf{x}_1)$  to  $\delta \leq \tilde{\delta}$  and  $r \leq \tilde{r}$ , so that both conclusions hold on  $B^\circ = (t_1 - \delta, t_1 + \delta) \times B(\mathbf{x}_1, r)$ .

Fix  $\mathbf{y}_0 \in B(\mathbf{x}_1, r)$  and consider  $\Psi(t, (\mathbf{y}_0, I))$ , defined for  $|t - t_1| < \delta$  because  $\|(\mathbf{y}_0, I) - (\mathbf{x}_1, I)\| = \|\mathbf{y}_0 - \mathbf{x}_1\| < r \leq \tilde{r}$ . The extended system decouples. Its first component solves  $\mathbf{y}' = \mathbf{f}(t, \mathbf{y})$  with  $\mathbf{y}(t_1) = \mathbf{y}_0$ , hence equals  $\varphi(t, \mathbf{y}_0)$  by lemma 4.2.4. Its second component then solves  $J' = D_{\mathbf{x}}\mathbf{f}(t, \varphi(t, \mathbf{y}_0))J$  with  $J(t_1) = I$ , which is the variational equation along  $\varphi(\cdot, \mathbf{y}_0)$ , hence equals  $J(t; \mathbf{y}_0) = D_{\mathbf{y}}\varphi(t, \mathbf{y}_0)$  by parts (1) and (2) of theorem 4.5.1. Therefore, on  $B^\circ$ ,

$$D_{\mathbf{y}}\varphi(t, \mathbf{y}_0) = (J\text{-component of } \Psi(t, (\mathbf{y}_0, I))), \quad \partial_t \varphi(t, \mathbf{y}_0) = \mathbf{f}(t, \varphi(t, \mathbf{y}_0))$$

The map  $(t, \mathbf{y}_0) \mapsto (t, (\mathbf{y}_0, I))$  is affine, hence smooth, so the first expression is  $C^{k-1}$  on  $B^\circ$  as a composition of  $C^{k-1}$  maps. For the second,  $\varphi$  itself is the first component of the same composition, hence  $C^{k-1}$ , and  $\mathbf{f}$  is  $C^k$ , so the chain rule makes  $\mathbf{f}(t, \varphi(t, \mathbf{y}_0))$  a  $C^{k-1}$  function of  $(t, \mathbf{y}_0)$ . Thus every first-order partial derivative of  $\varphi$  (namely  $\partial_t \varphi$  and the columns of  $D_{\mathbf{y}}\varphi$ ) is  $C^{k-1}$  on  $B^\circ$ , so every partial derivative of  $\varphi$  of order at most  $k$  exists and is continuous there. That is the definition of  $\varphi \in C^k(B^\circ)$ , closing the induction.

For a smooth right-hand side, the theorem gives, for each finite  $k$ , a possibly smaller box on which the local flow is  $C^k$ . Theorem 4.6.2 removes the shrinkage for autonomous systems: near every point of the full domain of the maximal flow, the flow is  $C^k$  for every finite  $k$ , so it is smooth on that domain.

## 4.6 The Flow of an Autonomous System

For an autonomous system the theory takes its final form. The right-hand side  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  has no time dependence, so time-translates of solutions are solutions and every initial time reduces to  $t = 0$ . Writing  $J(\mathbf{y}) = (\alpha(\mathbf{y}), \beta(\mathbf{y}))$  for the maximal interval of the solution with value  $\mathbf{y}$  at time 0, the *flow*  $\varphi(t, \mathbf{y})$  is the value at time  $t$  of that maximal solution, defined on the *flow domain*

$$\Omega = \{(t, \mathbf{y}) \in \mathbb{R} \times U \mid t \in J(\mathbf{y})\}$$

The theorem closing this section assembles the entire chapter: the flow domain is open, the flow is as regular as the field, flowing for time  $t + s$  is flowing for time  $s$  and then for time  $t$ , and a trajectory with a finite maximal time must leave every compact subset of  $U$ .

**Lemma 4.6.1: The Group Law**

Let  $U \subseteq \mathbb{R}^n$  be open,  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  locally Lipschitz, and  $\mathbf{y} \in U$ . If  $s \in J(\mathbf{y})$ , then  $J(\varphi(s, \mathbf{y})) = J(\mathbf{y}) - s$ , the translate  $\{t - s \mid t \in J(\mathbf{y})\}$ , and

$$\varphi(t, \varphi(s, \mathbf{y})) = \varphi(t + s, \mathbf{y}) \quad \text{for all } t \in J(\mathbf{y}) - s$$

**Proof:**

The system fits the general theory on  $D = \mathbb{R} \times U$  (definition 4.2.1), so through every point of  $U$  there is a maximal solution with open interval (theorem 4.3.1), and  $\varphi$  and  $J(\cdot)$  are defined. Set  $\mathbf{z} = \varphi(s, \mathbf{y})$  and define  $\gamma(t) = \varphi(t + s, \mathbf{y})$  for  $t \in J(\mathbf{y}) - s$ , an open interval containing 0. Because the right-hand side has no  $t$ -dependence,  $\gamma'(t) = \mathbf{f}(\varphi(t + s, \mathbf{y})) = \mathbf{f}(\gamma(t))$ , and  $\gamma(0) = \mathbf{z}$ , so  $\gamma$  is a solution through  $\mathbf{z}$ . Maximality (theorem 4.3.1) gives

$$J(\mathbf{z}) \supseteq J(\mathbf{y}) - s \quad \text{and} \quad \varphi(t, \mathbf{z}) = \varphi(t + s, \mathbf{y}) \quad \text{for } t \in J(\mathbf{y}) - s$$

In particular  $-s \in J(\mathbf{z})$  and  $\varphi(-s, \mathbf{z}) = \varphi(0, \mathbf{y}) = \mathbf{y}$ . Applying the inclusion just proved with  $(\mathbf{z}, -s)$  in place of  $(\mathbf{y}, s)$  gives  $J(\mathbf{y}) \supseteq J(\mathbf{z}) + s$ , so the inclusion is an equality.

**Theorem 4.6.2: The Fundamental Theorem for Autonomous Systems**

Let  $U \subseteq \mathbb{R}^n$  be open and let  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  be  $C^k$  with  $1 \leq k \leq \infty$ . With  $\varphi$ ,  $J(\cdot)$ , and  $\Omega$  as above:

1. Every  $\mathbf{y} \in U$  has an open maximal interval  $J(\mathbf{y}) = (\alpha(\mathbf{y}), \beta(\mathbf{y}))$  containing 0, and every solution of  $\mathbf{x}' = \mathbf{f}(\mathbf{x})$  with value  $\mathbf{y}$  at time 0 is a restriction of  $\varphi(\cdot, \mathbf{y})$ .
2. If  $s \in J(\mathbf{y})$ , then  $J(\varphi(s, \mathbf{y})) = J(\mathbf{y}) - s$  and  $\varphi(t, \varphi(s, \mathbf{y})) = \varphi(t + s, \mathbf{y})$  for  $t \in J(\mathbf{y}) - s$ .
3. If  $\beta(\mathbf{y}) < \infty$ , then for every compact  $K \subseteq U$  there is a  $\tau < \beta(\mathbf{y})$  with  $\varphi(t, \mathbf{y}) \notin K$  for all  $\tau < t < \beta(\mathbf{y})$ . The statement at  $\alpha(\mathbf{y})$  is symmetric.
4.  $\Omega$  is open in  $\mathbb{R} \times \mathbb{R}^n$  and  $\varphi : \Omega \rightarrow U$  is  $C^k$ .

**Proof:**

A  $C^k$  field with  $k \geq 1$  is  $C^1$ , hence locally Lipschitz (lemma 4.2.2), so (1) is theorem 4.3.1 on  $D = \mathbb{R} \times U$ , (2) is lemma 4.6.1, and (3) is corollary 4.3.5. It remains to prove (4). Since  $C^\infty$  means  $C^k$  for every finite  $k$ , and openness and the  $C^k$  property are local, it suffices to fix a finite  $k$  for which  $\mathbf{f}$  is  $C^k$  and a point  $(T, \mathbf{y}^*) \in \Omega$ , and to produce an open neighborhood of  $(T, \mathbf{y}^*)$  contained in  $\Omega$  on which  $\varphi$  is  $C^k$ .

First, the local statement. For  $\mathbf{z} \in U$ , theorem 4.5.2 applied to the autonomous system at  $(0, \mathbf{z})$  gives  $\delta_{\mathbf{z}}, r_{\mathbf{z}} > 0$  such that  $(t, \mathbf{w}) \mapsto \varphi(t, \mathbf{w})$  is defined and  $C^k$  on  $(-\delta_{\mathbf{z}}, \delta_{\mathbf{z}}) \times B(\mathbf{z}, r_{\mathbf{z}})$ , the flow map there being built from the same maximal solutions with data at time 0. In particular these boxes lie in  $\Omega$ , and the case  $T = 0$  is done. Take  $T > 0$ . (For  $T < 0$ , apply the result to the field  $-\mathbf{f}$ , whose flow is  $(t, \mathbf{y}) \mapsto \varphi(-t, \mathbf{y})$  with flow domain the reflection of  $\Omega$  in  $t$ , and which is  $C^k$  exactly when  $\mathbf{f}$  is.)

The trajectory segment  $K_0 = \{\varphi(s, \mathbf{y}^*) \mid 0 \leq s \leq T\}$  is a compact subset of  $U$ , being a continuous image of  $[0, T]$ . Cover  $K_0$  by the balls  $B(\mathbf{z}, r_{\mathbf{z}})$ ,  $\mathbf{z} \in K_0$ , extract a finite subcover  $B(\mathbf{z}_1, r_1), \dots, B(\mathbf{z}_m, r_m)$ , and set  $\delta = \min_j \delta_{\mathbf{z}_j}$ . Choose an integer  $N$  with  $T/N < \delta$  and set  $s_i = iT/N$  for  $i = 0, \dots, N$ . For each  $i < N$  fix an index  $j(i)$  with  $\varphi(s_i, \mathbf{y}^*) \in B(\mathbf{z}_{j(i)}, r_{j(i)})$ , and let

$$h_i : B(\mathbf{z}_{j(i)}, r_{j(i)}) \rightarrow U, \quad h_i(\mathbf{w}) = \varphi(T/N, \mathbf{w})$$

a  $C^k$  map, since  $T/N < \delta \leq \delta_{\mathbf{z}_{j(i)}}$  places it inside the box at  $\mathbf{z}_{j(i)}$ .

Now define open sets  $W_i \subseteq U$  and maps  $G_i : W_i \rightarrow U$  by  $W_0 = U$ ,  $G_0 = \text{id}$ , and

$$W_{i+1} = \{\mathbf{w} \in W_i \mid G_i(\mathbf{w}) \in B(\mathbf{z}_{j(i)}, r_{j(i)})\}, \quad G_{i+1} = h_i \circ G_i$$

By induction on  $i$ , each  $W_i$  is open and contains  $\mathbf{y}^*$ , each  $G_i$  is  $C^k$  on  $W_i$ , and for every  $\mathbf{w} \in W_i$  the time  $s_i$  lies in  $J(\mathbf{w})$  with  $G_i(\mathbf{w}) = \varphi(s_i, \mathbf{w})$ . Indeed, for  $i = 0$  all three claims are trivial. Assuming them for  $i$ : the set  $W_{i+1}$  is open as the preimage of an open ball under the continuous  $G_i$ , and contains  $\mathbf{y}^*$  because  $G_i(\mathbf{y}^*) = \varphi(s_i, \mathbf{y}^*) \in B(\mathbf{z}_{j(i)}, r_{j(i)})$  by the choice of  $j(i)$ . For  $\mathbf{w} \in W_{i+1}$ , the point  $\varphi(s_i, \mathbf{w}) = G_i(\mathbf{w})$  lies in  $B(\mathbf{z}_{j(i)}, r_{j(i)})$ , so the local box gives  $J(\varphi(s_i, \mathbf{w})) \supseteq (-\delta_{\mathbf{z}_{j(i)}}, \delta_{\mathbf{z}_{j(i)}}) \ni T/N$ , and the group law (2) turns this into  $s_{i+1} \in J(\mathbf{w})$  and

$$\varphi(s_{i+1}, \mathbf{w}) = \varphi(T/N, \varphi(s_i, \mathbf{w})) = h_i(G_i(\mathbf{w})) = G_{i+1}(\mathbf{w})$$

Finally  $G_{i+1}$  is  $C^k$  on  $W_{i+1}$  as a composition of  $C^k$  maps.

Take the open set  $V = \{(t, \mathbf{w}) \mid \mathbf{w} \in W_N, |t - s_{N-1}| < \delta_{\mathbf{z}_{j(N-1)}}\}$ . For  $(t, \mathbf{w}) \in V$ , the point  $\varphi(s_{N-1}, \mathbf{w}) = G_{N-1}(\mathbf{w})$  lies in  $B(\mathbf{z}_{j(N-1)}, r_{j(N-1)})$  by the definition of  $W_N$ , so by the group law  $t \in J(\mathbf{w})$  and

$$\varphi(t, \mathbf{w}) = \varphi(t - s_{N-1}, G_{N-1}(\mathbf{w}))$$

The right-hand side is a  $C^k$  function of  $(t, \mathbf{w})$ : the map  $(t, \mathbf{w}) \mapsto (t - s_{N-1}, G_{N-1}(\mathbf{w}))$  is  $C^k$  into the box at  $\mathbf{z}_{j(N-1)}$ , on which  $\varphi$  is  $C^k$ . And  $V$  contains  $(T, \mathbf{y}^*)$ , since  $\mathbf{y}^* \in W_N$  and  $|T - s_{N-1}| = T/N < \delta \leq \delta_{\mathbf{z}_{j(N-1)}}$ . Hence  $V$  is an open neighborhood of  $(T, \mathbf{y}^*)$  inside  $\Omega$  on which  $\varphi$  is  $C^k$ , completing the proof.

For a smooth field the flow is therefore smooth on all of  $\Omega$ , and chapter 5 draws on the consequences throughout: trajectories of an autonomous system never cross, each time- $t$  map  $\mathbf{y} \mapsto \varphi(t, \mathbf{y})$  is as regular as the field, and a trajectory confined to a compact subset of  $U$  exists for all forward time.

# 5

---

## *Qualitative Theory and Stability*

---

Most differential equations, and nearly all non-linear ones, cannot be solved in closed form. Giving up on explicit formulae loses less than it appears to, because the questions a model is usually asked are not questions about formulae: whether the system settles to an equilibrium, oscillates, or runs away, and whether a small error in the initial state stays small. All three are questions about the shape of the solution curves.

The phase plane displays that shape directly, plotting solutions as trajectories rather than graphs. Lyapunov's definitions make stability precise without producing a single solution. And near an equilibrium, linearization replaces a non-linear system by the constant-coefficient system of chapter 3, whose behaviour is already known; the theorems of this chapter say when that replacement is legitimate.

### 5.1 Autonomous Systems and Phase Space

We focus on *autonomous* systems, where the governing laws do not change with time.

#### **Definition 5.1.1: Autonomous System**

A system of ODEs is *autonomous* if it can be written as:

$$\mathbf{x}' = \mathbf{f}(\mathbf{x})$$

where  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  is a  $C^1$  vector field on an open set  $U \subseteq \mathbb{R}^n$ . The independent variable  $t$  does not appear explicitly on the right-hand side.

**Definition 5.1.2: Critical Point**

A point  $\mathbf{x}_0$  is a *critical point* (or equilibrium point) if  $\mathbf{f}(\mathbf{x}_0) = \mathbf{0}$ .

If the system starts at a critical point ( $\mathbf{x}(0) = \mathbf{x}_0$ ), the unique solution is the constant function  $\mathbf{x}(t) \equiv \mathbf{x}_0$  for all  $t$ .

The set of all solution curves (trajectories) in the domain  $U$  forms the *Phase Portrait*. For autonomous systems, trajectories never cross (by Uniqueness).

Intuitively, an equilibrium is “stable” if small disturbances do not grow over time. We formalize this using  $\epsilon - \delta$  language.

**Definition 5.1.3: Types of Stability**

Let  $\mathbf{x}_0$  be a critical point.

1. *Stable (in the sense of Lyapunov)*: For every  $\epsilon > 0$ , there exists a  $\delta > 0$  such that if  $\|\mathbf{x}(0) - \mathbf{x}_0\| < \delta$ , then  $\|\mathbf{x}(t) - \mathbf{x}_0\| < \epsilon$  for all  $t \geq 0$ .
2. *Asymptotically Stable*: The point is stable, and there exists a  $\delta_0 > 0$  such that if  $\|\mathbf{x}(0) - \mathbf{x}_0\| < \delta_0$ , then  $\lim_{t \rightarrow \infty} \mathbf{x}(t) = \mathbf{x}_0$ .
3. *Unstable*: The point is not stable.

Stability means the solution stays close. Asymptotic stability means it stays close *and* eventually converges to the equilibrium.

For a non-linear system  $\mathbf{x}' = \mathbf{f}(\mathbf{x})$ , the behavior near a critical point  $\mathbf{x}_0$  is often well-approximated by the linear term of its Taylor series.

$$\mathbf{f}(\mathbf{x}) \approx \mathbf{f}(\mathbf{x}_0) + D\mathbf{f}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0) = J(\mathbf{x} - \mathbf{x}_0)$$

where  $J = D\mathbf{f}(\mathbf{x}_0)$  is the *Jacobian matrix* evaluated at the critical point.

**Theorem 5.1.4: Principle of Linearized Stability**

Let  $\mathbf{x}' = \mathbf{f}(\mathbf{x})$  have a critical point at  $\mathbf{0}$  (w.l.o.g.). Let  $J = D\mathbf{f}(\mathbf{0})$ .

1. If all eigenvalues of  $J$  have strictly negative real parts ( $\text{Re}(\lambda_i) < 0$ ), then  $\mathbf{0}$  is *asymptotically stable*.
2. If at least one eigenvalue has a strictly positive real part ( $\text{Re}(\lambda_i) > 0$ ), then  $\mathbf{0}$  is *unstable*.
3. If some eigenvalues have zero real part (and none positive), the linearization is inconclusive (the center case).

**Proof:**

We sketch the proof for (1). Since all eigenvalues have negative real parts, there exists a specific inner product  $\langle \cdot, \cdot \rangle_*$  and norm  $\|\cdot\|_*$  such that  $\langle J\mathbf{x}, \mathbf{x} \rangle_* \leq -c\|\mathbf{x}\|_*^2$  for some  $c > 0$ . Consider the squared norm  $V(t) = \|\mathbf{x}(t)\|_*^2$ . Its derivative along trajectories is:

$$\frac{d}{dt}\|\mathbf{x}\|_*^2 = 2\langle \mathbf{x}', \mathbf{x} \rangle_* = 2\langle J\mathbf{x} + \mathbf{g}(\mathbf{x}), \mathbf{x} \rangle_*$$



where  $\mathbf{g}(\mathbf{x})$  represents higher-order terms ( $\|\mathbf{g}(\mathbf{x})\|/\|\mathbf{x}\| \rightarrow 0$ ).

$$\frac{dV}{dt} \leq -2c\|\mathbf{x}\|_*^2 + 2\|\mathbf{g}(\mathbf{x})\|_*\|\mathbf{x}\|_*$$

Near  $\mathbf{0}$ , the quadratic decay dominated the higher-order terms, proving  $V(t)$  decreases, so  $\mathbf{x}(t) \rightarrow \mathbf{0}$ .

### Example 5.1.5: Linearization Analysis

Analyze the stability of the critical points for the pendulum equation with damping:

$$\theta'' + \gamma\theta' + \sin \theta = 0, \quad \gamma > 0$$

*Solution:* Convert to a system:  $x = \theta, y = \theta'$ .

$$\begin{cases} x' = y \\ y' = -\sin x - \gamma y \end{cases}$$

Critical points occur where  $y = 0$  and  $\sin x = 0$ , so  $x = n\pi$ . The Jacobian is:

$$J(x, y) = \begin{pmatrix} 0 & 1 \\ -\cos x & -\gamma \end{pmatrix}$$

At  $(0, 0)$  (downward position):  $J = \begin{pmatrix} 0 & 1 \\ -1 & -\gamma \end{pmatrix}$ . Char. Poly:  $\lambda^2 + \gamma\lambda + 1 = 0$ . Roots  $\lambda = \frac{-\gamma \pm \sqrt{\gamma^2 - 4}}{2}$ . Real parts are negative. *Asymptotically Stable*.

At  $(\pi, 0)$  (upward position):  $J = \begin{pmatrix} 0 & 1 \\ 1 & -\gamma \end{pmatrix}$ . Char. Poly:  $\lambda^2 + \gamma\lambda - 1 = 0$ . Roots  $\lambda = \frac{-\gamma \pm \sqrt{\gamma^2 + 4}}{2}$ . One root is positive. *Unstable Saddle*.

## 5.2 Lyapunov's Direct Method

Linearization fails when eigenvalues have zero real parts. Lyapunov's Direct Method (or Second Method) allows us to determine stability without solving the equation, by generalizing the concept of energy.

### Definition 5.2.1: Lyapunov Function

Let  $\mathbf{x}^*$  be a critical point. A scalar function  $V : U \rightarrow \mathbb{R}$  is a *Lyapunov Function* if:

1.  $V(\mathbf{x}^*) = 0$  and  $V(\mathbf{x}) > 0$  for  $\mathbf{x} \neq \mathbf{x}^*$  (Positive Definite).
2.  $\dot{V}(\mathbf{x}) = \nabla V \cdot \mathbf{f}(\mathbf{x}) \leq 0$  for all  $\mathbf{x} \in U \setminus \{\mathbf{x}^*\}$  (Orbital Derivative is Negative Semi-Definite).

If the inequality in (2) is strict ( $\dot{V} < 0$ ), it is a *Strict Lyapunov Function*.

**Theorem 5.2.2: Lyapunov Stability Theorem**

Let  $\mathbf{x}^*$  be a critical point.

1. If there exists a Lyapunov Function  $V$  in a neighborhood of  $\mathbf{x}^*$ , then  $\mathbf{x}^*$  is *Stable*.
2. If there exists a Strict Lyapunov Function  $V$ , then  $\mathbf{x}^*$  is *Asymptotically Stable*.

**Proof:**

We prove (1). Let  $\epsilon > 0$  be small enough that the ball  $B_\epsilon(\mathbf{x}^*)$  is in the domain of  $V$ . Let  $m = \min_{\|\mathbf{x}\|=\epsilon} V(\mathbf{x})$ . Since  $V$  is continuous and positive definite,  $m > 0$ . Define the set  $U = \{\mathbf{x} | V(\mathbf{x}) < m\}$ . Since  $V(\mathbf{x}^*) = 0$  and  $V$  is continuous, there exists a  $\delta > 0$  such that  $B_\delta(\mathbf{x}^*) \subset U$ .

Now, let a trajectory start at  $\mathbf{x}_0$  with  $\|\mathbf{x}_0 - \mathbf{x}^*\| < \delta$ . Then  $V(\mathbf{x}_0) < m$ . Since  $\dot{V} \leq 0$ ,  $V(\mathbf{x}(t))$  is non-increasing. Thus,  $V(\mathbf{x}(t)) \leq V(\mathbf{x}_0) < m$  for all  $t \geq 0$ . Since the boundary of  $B_\epsilon$  has  $V \geq m$ , the trajectory can never reach the boundary  $\|\mathbf{x}\| = \epsilon$ . Therefore,  $\|\mathbf{x}(t) - \mathbf{x}^*\| < \epsilon$  for all  $t$ . This is the definition of stability.

**Example 5.2.3: Lyapunov Function**

Show that the origin is stable for the system:

$$x' = -x - xy^2, \quad y' = -y - yx^2$$

*Solution:* Try  $V(x, y) = x^2 + y^2$ . This is positive definite. Compute  $\dot{V}$ :

$$\begin{aligned} \dot{V} &= \frac{\partial V}{\partial x} x' + \frac{\partial V}{\partial y} y' \\ &= 2x(-x - xy^2) + 2y(-y - yx^2) \\ &= -2x^2 - 2x^2y^2 - 2y^2 - 2y^2x^2 \\ &= -2(x^2 + y^2) - 4x^2y^2 \end{aligned}$$

Clearly  $\dot{V} < 0$  for all  $(x, y) \neq (0, 0)$ . Thus,  $V$  is a strict Lyapunov function, and the origin is asymptotically stable.

## 5.3 Limit Cycles

In non-linear systems, solutions can approach a closed periodic orbit called a *limit cycle*.

**Theorem 5.3.1: Poincaré-Bendixson Theorem**

Suppose  $R$  is a closed, bounded region in the plane ( $\mathbb{R}^2$ ) that contains no critical points. If a trajectory  $\mathbf{x}(t)$  is confined in  $R$  for all  $t \geq 0$ , then  $\mathbf{x}(t)$  must either be a periodic orbit or approach a periodic orbit as  $t \rightarrow \infty$ .

**Proof:**

(Sketch) The proof relies on the Jordan Curve Theorem (unique to  $\mathbb{R}^2$ ). Since trajectories cannot cross, a bounded trajectory that doesn't terminate at a fixed point must eventually wind onto itself, forming a limit cycle.

**Example 5.3.2: Existence of Limit Cycle**

Consider  $r' = r(1 - r^2), \theta' = 1$  (in polar coordinates).

- If  $r < 1$ ,  $r' > 0$  (radius increases).
- If  $r > 1$ ,  $r' < 0$  (radius decreases).

Consider the annulus  $R = \{1/2 \leq r \leq 2\}$ . Trajectories enter  $R$  on both boundaries and never leave. Since there are no critical points (as  $\theta' = 1 \neq 0$ ), any solution starting in  $R$  must approach the limit cycle at  $r = 1$ .



# 6

---

## *Boundary Value Problems and Sturm-Liouville Theory*

---

Every problem so far has specified its conditions at a single point  $t_0$ . Imposing them instead at the two ends of an interval changes the theory rather than the technique. A linear initial value problem always has exactly one solution; a boundary value problem may have none, exactly one, or infinitely many, and which of the three occurs depends on the parameter in the equation only through the spectrum of the differential operator.

Solvability therefore becomes an eigenvalue question, and Sturm-Liouville theory is the eigenvalue theory that answers it. It is diagonalization transplanted from matrices to differential operators: a self-adjoint operator, real eigenvalues, and an orthogonal basis of eigenfunctions in which a solution may be expanded. Green's functions complete the picture by inverting the operator outright on the inhomogeneous problem.

### 6.1 Introduction to Boundary Value Problems

Consider the second-order linear equation on an interval  $[a, b]$ :

$$L[y] = y'' + p(x)y' + q(x)y = g(x), \quad a < x < b$$

subject to boundary conditions at the endpoints:

$$B_1[y] = \alpha_1 y(a) + \alpha_2 y'(a) = \gamma_1, \quad B_2[y] = \beta_1 y(b) + \beta_2 y'(b) = \gamma_2$$

**Definition 6.1.1: Homogeneous BVP**

A BVP is called *homogeneous* if both the differential equation ( $g(x) \equiv 0$ ) and the boundary conditions ( $\gamma_1 = \gamma_2 = 0$ ) are homogeneous.

The trivial solution  $y(x) \equiv 0$  is always a solution to a homogeneous BVP. The central question of this chapter is: *Under what conditions does a non-trivial solution exist?*

**Example 6.1.2: Eigenvalues of a BVP**

Consider the problem:

$$y'' + \lambda y = 0, \quad y(0) = 0, \quad y(\pi) = 0$$

Depending on the parameter  $\lambda$ , the nature of the solution changes:

1. If  $\lambda < 0$  (let  $\lambda = -\mu^2$ ):  $y(x) = c_1 \cosh \mu x + c_2 \sinh \mu x$ . The boundary conditions force  $c_1 = c_2 = 0$ .
2. If  $\lambda = 0$ :  $y(x) = c_1 x + c_2$ . Boundary conditions force  $c_1 = c_2 = 0$ .
3. If  $\lambda > 0$  (let  $\lambda = \mu^2$ ):  $y(x) = c_1 \cos \mu x + c_2 \sin \mu x$ .  $y(0) = 0 \implies c_1 = 0$ .  $y(\pi) = c_2 \sin(\mu\pi) = 0$ . For a non-trivial solution, we require  $\sin(\mu\pi) = 0$ , which implies  $\mu = n$  for  $n \in \mathbb{Z}^+$ .

Thus, non-trivial solutions (eigenfunctions)  $y_n(x) = \sin(nx)$  exist only for the discrete spectrum  $\lambda_n = n^2$ .

To understand the existence of eigenvalues generally, we must define the “transpose” of a differential operator.

**Definition 6.1.3: Formal Adjoint**

For the operator  $L[y] = p_2(x)y'' + p_1(x)y' + p_0(x)y$ , the *formal adjoint*  $L^*$  is defined by:

$$L^*[u] = \frac{d^2}{dx^2}(p_2 u) - \frac{d}{dx}(p_1 u) + p_0 u$$

The operator  $L$  is *formally self-adjoint* if  $L = L^*$ .

A straightforward calculation shows that  $L$  is self-adjoint if and only if  $p_1(x) = p_2'(x)$ . This motivates the *Sturm-Liouville form*.

**Definition 6.1.4: Sturm-Liouville Form**

A second-order differential operator  $L$  is in *Sturm-Liouville form* if:

$$L[y] = -\frac{d}{dx} \left[ p(x) \frac{dy}{dx} \right] + q(x)y$$

where  $p(x) > 0$ . This operator is formally self-adjoint.

*Note:* Any linear operator  $P(x)y'' + Q(x)y' + R(x)y$  can be multiplied by the integrating factor

$\mu(x) = \frac{1}{P(x)} e^{\int Q/P dx}$  to be converted into Sturm-Liouville form.

### Theorem 6.1.5: Lagrange's Identity

For a self-adjoint operator  $L$  in Sturm-Liouville form:

$$uL[v] - vL[u] = \frac{d}{dx} [p(x)(uv' - u'v)]$$

Integrating over  $[a, b]$  yields Green's Formula:

$$\langle L[u], v \rangle - \langle u, L[v] \rangle = [p(x)(uv' - u'v)]_a^b$$

If  $u$  and  $v$  satisfy *separated boundary conditions* (e.g.,  $\alpha_1 u(a) + \alpha_2 u'(a) = 0$ ), the boundary term vanishes, and we obtain the crucial symmetry property:  $\langle L[u], v \rangle = \langle u, L[v] \rangle$ .

## 6.2 The Sturm-Liouville Theorems

We now consider the eigenvalue problem:

$$L[y] = \lambda w(x)y$$

subject to homogeneous separated boundary conditions.

### Theorem 6.2.1: Sturm-Liouville Theorem

For a regular Sturm-Liouville problem:

1. All eigenvalues  $\lambda$  are real.
2. Eigenfunctions  $\varphi_n, \varphi_m$  corresponding to distinct eigenvalues are *orthogonal* with respect to the weight  $w(x)$ :

$$\int_a^b \varphi_n(x) \varphi_m(x) w(x) dx = 0 \quad \text{if } n \neq m$$

3. The eigenvalues form an infinite sequence  $\lambda_1 < \lambda_2 < \dots$  with  $\lim_{n \rightarrow \infty} \lambda_n = \infty$ .
4. The eigenfunctions form a *complete basis* for the space of piecewise continuous functions  $L_w^2[a, b]$ .

#### Proof:

*Real Eigenvalues:* Let  $L[u] = \lambda wu$ . Take the complex conjugate  $L[\bar{u}] = \bar{\lambda} w\bar{u}$ . Using Green's Formula (with boundary terms vanishing):

$$(\lambda - \bar{\lambda}) \langle wu, \bar{u} \rangle = \langle L[u], \bar{u} \rangle - \langle u, L[\bar{u}] \rangle = 0$$

Since  $w(x) > 0$ ,  $\langle wu, \bar{u} \rangle = \int w|u|^2 > 0$ , implying  $\lambda = \bar{\lambda}$ .

*Orthogonality:* Let  $L[\varphi_n] = \lambda_n w \varphi_n$  and  $L[\varphi_m] = \lambda_m w \varphi_m$ .

$$(\lambda_n - \lambda_m) \langle w \varphi_n, \varphi_m \rangle = \langle L[\varphi_n], \varphi_m \rangle - \langle \varphi_n, L[\varphi_m] \rangle = 0$$

If  $\lambda_n \neq \lambda_m$ , the inner product must vanish.

### Theorem 6.2.2: Eigenfunction Expansion

Any piecewise smooth function  $f(x)$  on  $[a, b]$  can be represented as:

$$f(x) \sim \sum_{n=1}^{\infty} c_n \varphi_n(x)$$

where the Fourier coefficients are given by the projection:

$$c_n = \frac{\langle f, \varphi_n \rangle_w}{\langle \varphi_n, \varphi_n \rangle_w} = \frac{\int_a^b f(x) \varphi_n(x) w(x) dx}{\int_a^b \varphi_n(x)^2 w(x) dx}$$

## 6.3 The Fredholm Alternative

We now address the inhomogeneous equation  $L[y] - \lambda w(x)y = f(x)$ . The solvability depends on whether  $\lambda$  is an eigenvalue.

### Theorem 6.3.1: Fredholm Alternative

Consider the inhomogeneous BVP  $L[y] - \sigma w(x)y = f(x)$ .

1. **Non-Resonant Case:** If  $\sigma$  is *not* an eigenvalue of the homogeneous problem, there exists a unique solution for any  $f(x)$ .
2. **Resonant Case:** If  $\sigma = \lambda_k$  is an eigenvalue with eigenfunction  $\varphi_k$ , a solution exists *if and only if*  $f$  is orthogonal to the eigenfunction:

$$\int_a^b f(x) \varphi_k(x) w(x) dx = 0$$

If this condition holds, there are infinitely many solutions (of the form  $y_p + c\varphi_k$ ).

## 6.4 Green's Functions

In the non-resonant case, the unique solution can be represented by an integral kernel called the Green's function.



**Definition 6.4.1: Green's Function**

The *Green's Function*  $G(x, \xi)$  for the operator  $L$  is the solution to the distributional equation:

$$L[G(x, \xi)] = \delta(x - \xi)$$

subject to the homogeneous boundary conditions.

To construct  $G(x, \xi)$  explicitly, we solve the homogeneous equation  $L[y] = 0$  on the intervals  $[a, \xi)$  and  $(\xi, b]$ . Let  $y_1(x)$  satisfy the boundary condition at  $a$ , and  $y_2(x)$  satisfy the condition at  $b$ . We assume:

$$G(x, \xi) = \begin{cases} c_1(\xi)y_1(x) & a \leq x < \xi \\ c_2(\xi)y_2(x) & \xi < x \leq b \end{cases}$$

We require two conditions at  $x = \xi$ :

1. *Continuity*:  $c_1y_1(\xi) = c_2y_2(\xi)$ .
2. *Jump in Derivative*: Integrating  $-\frac{d}{dx}[py'] = \delta$  from  $\xi - \epsilon$  to  $\xi + \epsilon$  implies a jump of  $-1/p(\xi)$ :

$$\frac{\partial G}{\partial x}(\xi^+, \xi) - \frac{\partial G}{\partial x}(\xi^-, \xi) = -\frac{1}{p(\xi)}$$

Solving this system for  $c_1, c_2$  using the Wronskian  $W(\xi) = y_1y_2' - y_2y_1'$  yields the symmetric form:

**Theorem 6.4.2: Structure of the Green's Function**

The Green's function is given by:

$$G(x, \xi) = \frac{1}{p(\xi)W(\xi)} \begin{cases} y_1(x)y_2(\xi) & a \leq x \leq \xi \\ y_1(\xi)y_2(x) & \xi \leq x \leq b \end{cases}$$

Note that  $p(\xi)W(\xi)$  is a constant. The solution to  $L[y] = f(x)$  is then:

$$y(x) = \int_a^b G(x, \xi) f(\xi) d\xi$$



---

## *Series Solutions and Special Functions*

---

The linear equations solved so far have had constant coefficients, which is what allowed exponentials to reduce them to polynomials. The equations of mathematical physics do not cooperate. Separating the heat, wave, and Schrödinger equations in cylindrical or spherical coordinates produces ordinary equations whose coefficients vary with the independent variable, and these almost never have solutions expressible in  $e^x$ ,  $\sin x$ , and the rest of the elementary vocabulary.

The response is to widen what counts as a solution. A power series determined term by term from the equation defines a function whether or not that function has an elementary name, and the Frobenius method extends the construction across the singular points where the leading coefficient vanishes, which is exactly where the physical problems place them. The functions this produces, Bessel and Legendre chief among them, are the standard vocabulary for those problems.

### **7.1 Analytic Functions and Ordinary Points**

We consider the homogeneous linear second-order equation:

$$P(x)y'' + Q(x)y' + R(x)y = 0 \tag{7.1}$$

Dividing by  $P(x)$ , we obtain the standard form  $y'' + p(x)y' + q(x)y = 0$ . The behavior of solutions depends entirely on the behavior of  $p(x)$  and  $q(x)$  at a given point  $x_0$ .

**Definition 7.1.1: Ordinary and Singular Points**

Let  $p(x) = Q(x)/P(x)$  and  $q(x) = R(x)/P(x)$ . A point  $x_0$  is called an *ordinary point* if both  $p(x)$  and  $q(x)$  are analytic at  $x_0$  (i.e., they have convergent Taylor series expansions in a neighborhood of  $x_0$ ).

If either function is not analytic at  $x_0$ , the point is called a *singular point*.

Since polynomials are analytic everywhere, if  $P, Q, R$  are polynomials, singular points only occur at the roots of  $P(x) = 0$ .

**Theorem 7.1.2: Existence of Power Series Solutions**

If  $x_0$  is an ordinary point of the differential equation, then there exist two linearly independent solutions of the form:

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$$

The radius of convergence of this series is at least as large as the distance from  $x_0$  to the nearest singular point (in the complex plane).

**Proof:**

(Sketch) Since  $p(x)$  and  $q(x)$  are analytic, they can be represented by power series. We assume  $y(x) = \sum a_n (x - x_0)^n$ , differentiate term-by-term, and substitute into the ODE. Equating coefficients of like powers of  $(x - x_0)$  yields a recurrence relation determining  $a_n$  for  $n \geq 2$  in terms of the arbitrary constants  $a_0, a_1$ . The convergence is guaranteed by the theory of analytic functions in complex analysis.

**Example 7.1.3: Legendre's Equation**

Consider Legendre's equation, which arises in spherical coordinates:

$$(1 - x^2)y'' - 2xy' + \alpha(\alpha + 1)y = 0$$

Here  $P(x) = 1 - x^2$ . The singular points are  $x = \pm 1$ . The point  $x_0 = 0$  is an ordinary point.

*Solution:* Let  $y = \sum a_n x^n$ .

$$y' = \sum n a_n x^{n-1}, \quad y'' = \sum n(n-1) a_n x^{n-2}$$

Substituting into the ODE:

$$(1 - x^2) \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 2x \sum_{n=1}^{\infty} n a_n x^{n-1} + \alpha(\alpha + 1) \sum_{n=0}^{\infty} a_n x^n = 0$$

Shifting indices to align powers of  $x^k$ :

$$\sum_{k=0}^{\infty} [(k+2)(k+1)a_{k+2} - (k(k-1) + 2k - \alpha(\alpha + 1))a_k] x^k = 0$$

The recurrence relation is:

$$a_{k+2} = -\frac{(\alpha - k)(\alpha + k + 1)}{(k + 2)(k + 1)}a_k$$

Notice that if  $\alpha = n$  is a non-negative integer, the numerator vanishes when  $k = n$ , causing the series to terminate. These terminating solutions are the *Legendre Polynomials*  $P_n(x)$ .

## 7.2 Regular Singular Points and the Method of Frobenius

Many physical potentials (like the Coulomb potential  $1/r$ ) introduce singularities at the origin. Standard power series fail here. We need a more robust method.

### Definition 7.2.1: Regular Singular Point

A singular point  $x_0$  is called a *regular singular point* if the singularities are no worse than a pole of order 1 for  $p(x)$  and order 2 for  $q(x)$ . Specifically, if:

$$\lim_{x \rightarrow x_0} (x - x_0)p(x) \quad \text{and} \quad \lim_{x \rightarrow x_0} (x - x_0)^2 q(x)$$

are both finite (analytic), then  $x_0$  is regular.

For such points, we assume a solution of the form:

$$y(x) = (x - x_0)^r \sum_{n=0}^{\infty} a_n (x - x_0)^n = \sum_{n=0}^{\infty} a_n (x - x_0)^{n+r}, \quad a_0 \neq 0$$

This is called the *Frobenius Method*. The exponent  $r$  is determined by the “Indicial Equation.”

### Theorem 7.2.2: Frobenius Theorem

Let  $x = 0$  be a regular singular point. Let  $p_0 = \lim_{x \rightarrow 0} xp(x)$  and  $q_0 = \lim_{x \rightarrow 0} x^2 q(x)$ . The *Indicial Equation* is:

$$r(r - 1) + p_0 r + q_0 = 0$$

Let  $r_1, r_2$  be the roots with  $\text{Re}(r_1) \geq \text{Re}(r_2)$ .

1. There always exists one solution  $y_1(x) = |x|^{r_1} \sum a_n x^n$ .
2. If  $r_1 - r_2$  is not an integer, the second solution is  $y_2(x) = |x|^{r_2} \sum b_n x^n$ .
3. If  $r_1 - r_2$  is an integer (or zero), the second solution may involve a logarithm:

$$y_2(x) = C y_1(x) \ln |x| + |x|^{r_2} \sum b_n x^n$$

The most important application of the Frobenius method is solving Bessel’s equation, which arises when separating variables in cylindrical coordinates (Laplacian in 2D or 3D).

**Example 7.2.3: Bessel's Equation of Order  $\nu$** 

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0, \quad \nu \geq 0$$

Here  $p(x) = 1/x$  and  $q(x) = (x^2 - \nu^2)/x^2$ .  $x = 0$  is a regular singular point.

*Indicial Equation:* Multiply by  $x^2$  to identify the leading terms:  $x^2 y'' + xy' - \nu^2 y \approx 0$  near  $x = 0$ . This is Cauchy-Euler! The characteristic equation is  $r(r-1) + r - \nu^2 = 0 \implies r^2 - \nu^2 = 0 \implies r = \pm \nu$ .

*Solution Construction:* Using  $r_1 = \nu$ , the Frobenius method yields the *Bessel Function of the First Kind*, denoted  $J_\nu(x)$ :

$$J_\nu(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n! \Gamma(n + \nu + 1)} \left(\frac{x}{2}\right)^{2n+\nu}$$

For the second solution, since  $r_1 - r_2 = 2\nu$  might be an integer, we encounter the logarithmic case. The second linearly independent solution is the *Bessel Function of the Second Kind*,  $Y_\nu(x)$ , which blows up at  $x = 0$ .

## Sturm-Liouville Properties of Special Functions

Why do we care about these complicated series? Because they form orthogonal bases, just like sines and cosines.

Recall Bessel's equation:  $(xy')' + (x - \frac{\nu^2}{x})y = 0$ . This is almost Sturm-Liouville form with  $p(x) = x$ ,  $q(x) = -\nu^2/x$ ,  $\lambda = 1$ ,  $w(x) = x$ . Consider the BVP on  $[0, R]$  with  $y(R) = 0$ . The solutions are  $J_\nu(\lambda_k x)$  where  $\lambda_k$  are roots of  $J_\nu(\lambda R) = 0$ .

**Theorem 7.2.4: Orthogonality of Bessel Functions**

Let  $\{\lambda_k\}$  be the positive zeros of  $J_\nu(x)$ . Then the functions  $\varphi_k(x) = J_\nu(\lambda_k x/R)$  are orthogonal on  $[0, R]$  with respect to the weight  $w(x) = x$ :

$$\int_0^R J_\nu\left(\frac{\lambda_k x}{R}\right) J_\nu\left(\frac{\lambda_m x}{R}\right) x dx = 0 \quad \text{if } k \neq m$$

This orthogonality allows us to expand any function  $f(r)$  on a disk into a Fourier-Bessel series:

$$f(r) = \sum_{k=1}^{\infty} c_k J_\nu\left(\frac{\lambda_k r}{R}\right)$$

This is the fundamental mechanism for solving heat flow on a circular plate or vibrations of a drum head.

---

# *Bibliography*

---

- [1] Nathanael Chwojko-Srawley. *Everything You Need To Know About Metric and Function Spaces*. Bootstrapped 2026-07-29 to metric-function-spaces/ from NateElemAnalysis Part II. Series home for metric completeness, the Banach contraction principle, Stone-Weierstrass, Arzela-Ascoli and Baire category; see DECISIONS 2026-07-29. 2026.