
Analytic Field Theory

— *EYNTKA* Series —

NATHANAEL CHWOJKO-SRAWLEY

June 30, 2026

Contents

Contents	3
A Review of Elementary Number Theory	i
0.1 Divisibility and the Euclidean Algorithm	i
0.2 Primes and the Fundamental Theorem of Arithmetic	vii
0.3 Congruences	xii
0.4 The Group $(\mathbb{Z}/m\mathbb{Z})^\times$ and Primitive Roots	xvii
0.5 Quadratic Residues and Reciprocity	xxiii
0.6 Asymptotic Notation	xxix
1 Arithmetic Functions and Summation Techniques	1
1.1 Arithmetic Functions and Multiplicativity	1
1.2 Dirichlet Convolution and Möbius Inversion	6
1.3 Average Orders and Summatory Functions	12
1.4 Partial Summation and Abel's Summation Formula	17
1.5 The Euler-Maclaurin Summation Formula	22
2 Dirichlet Series and Euler Products	31
2.1 Dirichlet Series: Convergence and Analyticity	31
2.2 The Ring Homomorphism: Convolution and Multiplication	37
2.3 Euler Products	42
2.4 Mellin Transforms and the Integral Representation of $\zeta(s)$	50
2.5 Perron's Formula	58

3	Elementary Results on Prime Distribution	67
3.1	Chebyshev's Functions	68
3.2	Chebyshev's Estimates	72
3.3	Mertens' Theorems	80
3.4	Bertrand's Postulate	90
4	The Riemann Zeta Function	97
4.1	Definition and the Euler Product	97
4.2	Meromorphic Continuation to $\operatorname{Re}(s) > 0$	101
4.3	First Arithmetic Consequences	104
4.4	Non-Vanishing on the Line $\operatorname{Re}(s) = 1$	110
5	The Prime Number Theorem	115
5.1	The Analytic Strategy	116
5.2	The Laplace Transform	118
5.3	Newman's Tauberian Theorem	122
5.4	Proof of the Prime Number Theorem	127
5.5	Error Terms and the Zero-Free Region	135
6	The Functional Equation and the Critical Strip	145
6.1	The Gamma Function	146
6.2	Theta Functions and Poisson Summation	152
6.3	The Completed Zeta Function and Its Functional Equation	158
6.4	Trivial Zeros and Special Values	165
6.5	The Critical Strip and the Riemann Hypothesis	171
7	Dirichlet Characters and L-Functions	179
7.1	Dirichlet Characters	180
7.2	Dirichlet L -Functions	187
7.3	Gauss Sums and the Functional Equation	193
7.4	Non-Vanishing of $L(1, \chi)$	203
7.5	Dirichlet's Theorem and Equidistribution	209
8	The Dedekind Zeta Function and the Class Number Formula	217
8.1	The Dedekind Zeta Function	218
8.2	Ideal Counting and Lattice Point Geometry	223
8.3	The Analytic Class Number Formula	229

8.4	Factoring ζ_K and the Algebraic Non-Vanishing of $L(1, \chi)$	234
8.5	The Functional Equation for $\zeta_K(s)$	243
Index		251

A Review of Elementary Number Theory

The machinery of analytic number theory rests on a foundation of elementary results about integers, primes, and congruences. This preliminary chapter collects the essential facts that the rest of the book takes for granted. Readers with a solid undergraduate number theory background will find this material familiar and may treat the chapter as a reference; those encountering some results for the first time should work through the proofs carefully, as the techniques—particularly the Euclidean algorithm, the Chinese Remainder Theorem, and quadratic reciprocity—reappear in more sophisticated forms throughout the book.

The progression is natural. Divisibility and the Euclidean algorithm provide the computational backbone. The Fundamental Theorem of Arithmetic establishes that primes are the multiplicative building blocks of $\mathbb{Z}$, a fact whose failure in more general rings drives much of algebraic number theory (see [17]). Congruences and the Chinese Remainder Theorem introduce the modular perspective that underlies Dirichlet characters and L -functions. The structure of the unit group $(\mathbb{Z}/m\mathbb{Z})^\times$ and the theory of quadratic residues prepare the ground for character theory and reciprocity laws. The chapter closes with a review of asymptotic notation, the language in which the main results of analytic number theory are expressed.

0.1 Divisibility and the Euclidean Algorithm

The integers carry two intertwined structures: addition and multiplication. The additive structure is simple— $\mathbb{Z}$ is a free abelian group of rank 1—but the multiplicative structure is the source of all the complexity that this book investigates. The first step toward understanding it is the divisibility relation, and the first computational tool is the Euclidean algorithm, which reduces questions about divisibility to iterated division with remainder.

The starting point is the division algorithm, which guarantees that long division in $\mathbb{Z}$ always terminates with a well-defined quotient and remainder.

Theorem 0.1.1: Division Algorithm

For any $a \in \mathbb{Z}$ and $b \in \mathbb{Z}_{>0}$, there exist unique integers q (the *quotient*) and r (the *remainder*) such that

$$a = bq + r, \quad 0 \leq r < b$$

Proof:

Step 1: Existence. Consider the set $S = \{a - bk \mid k \in \mathbb{Z}, a - bk \geq 0\}$. Since $b > 0$, choosing k sufficiently negative ensures that $a - bk > 0$, so S is nonempty. By the well-ordering principle, S has a least element $r = a - bq$ for some $q \in \mathbb{Z}$. By construction $r \geq 0$. If $r \geq b$, then $a - b(q + 1) = r - b \geq 0$ belongs to S and is strictly smaller than r , contradicting minimality. Hence $0 \leq r < b$.

Step 2: Uniqueness. Suppose $a = bq_1 + r_1 = bq_2 + r_2$ with $0 \leq r_1, r_2 < b$. Then $b(q_1 - q_2) = r_2 - r_1$, so b divides $r_2 - r_1$. Since $|r_2 - r_1| < b$, this forces $r_1 = r_2$ and consequently $q_1 = q_2$.

The division algorithm is the engine behind every computation in this section. Its role is analogous to that of the Euclidean metric in geometry: it provides a notion of “size” (the remainder r) that strictly decreases under a natural operation (repeated division), guaranteeing that iterative processes terminate. This principle—descent by remainder—also underlies the proof of unique factorization, which depends on it via Euclid’s lemma in the next section.

With division in hand, the notion of exact divisibility can be made precise.

Definition 0.1.2: Divisibility

Let $a, b \in \mathbb{Z}$. Then a *divides* b , written $a \mid b$, if there exists $k \in \mathbb{Z}$ such that $b = ak$. Equivalently, $a \mid b$ if and only if the remainder upon dividing b by a is zero. If a does not divide b , this is denoted $a \nmid b$.

The divisibility relation is a partial order on the positive integers: it is reflexive ($a \mid a$ since $a = a \cdot 1$), antisymmetric (if $a \mid b$ and $b \mid a$ with $a, b > 0$ then $a = b$), and transitive (if $a \mid b$ and $b \mid c$ then $a \mid c$, since $c = bk_1 = ak_2k_1$). It is also compatible with the ring operations of $\mathbb{Z}$ in the following sense.

Proposition 0.1.3: Linearity of Divisibility

If $d \mid a$ and $d \mid b$, then $d \mid (ax + by)$ for all $x, y \in \mathbb{Z}$.

Proof:

Write $a = dk_1$ and $b = dk_2$. Then $ax + by = d(k_1x + k_2y)$.

In the language of algebra, this says that the set of multiples of d forms an ideal $(d) = d\mathbb{Z}$ in $\mathbb{Z}$, and divisibility translates to ideal containment: $a \mid b$ if and only if $(b) \subseteq (a)$. The fact that *every* ideal of $\mathbb{Z}$ has this form— $\mathbb{Z}$ is a principal ideal domain—is the ring-theoretic content of Bézout’s identity, proved below. In more general number rings, ideals need not be principal, and the failure of this property is measured by the class group (see [17]).

The central notion built from divisibility is the greatest common divisor.

Definition 0.1.4: Greatest Common Divisor

For $a, b \in \mathbb{Z}$ not both zero, the *greatest common divisor* $\gcd(a, b)$ is the largest positive integer d such that $d|a$ and $d|b$. By convention, $\gcd(0, 0) = 0$. The integers a and b are *coprime* (or *relatively prime*) if $\gcd(a, b) = 1$.

The existence of $\gcd(a, b)$ is clear: the set of common divisors is nonempty (it contains 1 when a and b are not both zero) and bounded above by $\max(|a|, |b|)$, so a maximum exists. Before describing how to compute it, a few basic identities are worth recording.

Proposition 0.1.5: Properties of the GCD

For all $a, b \in \mathbb{Z}$ and $k \in \mathbb{Z} \setminus \{0\}$:

1. $\gcd(a, b) = \gcd(b, a) = \gcd(|a|, |b|)$.
2. $\gcd(a, 0) = |a|$.
3. $\gcd(ka, kb) = |k| \gcd(a, b)$.
4. If $a = bq + r$, then $\gcd(a, b) = \gcd(b, r)$.

Proof:

Parts (1)–(3) follow directly from the definition, since the set of common divisors is unchanged by reordering, by taking absolute values, or by scaling (respectively). For (4), proposition 0.1.3 gives the two containments: every common divisor of a and b divides $r = a - bq$, and every common divisor of b and r divides $a = bq + r$. Hence the sets of common divisors coincide, and their maxima are equal.

Part (4) is the identity that makes the Euclidean algorithm work. It reduces $\gcd(a, b)$ to $\gcd(b, r)$ where $0 \leq r < b$, so the arguments are strictly smaller. Iterating this reduction is the Euclidean algorithm.

Proposition 0.1.6: The Euclidean Algorithm

Let $a, b \in \mathbb{Z}$ with $b > 0$. Apply the division algorithm repeatedly:

$$\begin{aligned}
 a &= bq_0 + r_0, & 0 \leq r_0 < b, \\
 b &= r_0q_1 + r_1, & 0 \leq r_1 < r_0, \\
 r_0 &= r_1q_2 + r_2, & 0 \leq r_2 < r_1, \\
 &\vdots \\
 r_{n-2} &= r_{n-1}q_n + r_n, & 0 \leq r_n < r_{n-1}, \\
 r_{n-1} &= r_nq_{n+1}
 \end{aligned}$$

The process terminates after finitely many steps, and the last nonzero remainder is $\gcd(a, b) = r_n$.

Proof:

The sequence $b > r_0 > r_1 > \cdots \geq 0$ is a strictly decreasing sequence of non-negative integers, so it must reach zero: there exists a smallest index n such that $r_{n+1} = 0$. At that step, $r_{n-1} = r_nq_{n+1}$, so $r_n \mid r_{n-1}$. Working backward through each line, r_n divides every preceding remainder and hence divides both b and a : it is a common divisor. Conversely, if d is any common divisor of a and b , the first equation gives $d \mid r_0$ (by proposition 0.1.3), the second gives $d \mid r_1$, and continuing inductively, $d \mid r_n$. Thus r_n is a common divisor that is divisible by every common divisor, which forces $r_n = \gcd(a, b)$.

The running time of the Euclidean algorithm is modest. After two consecutive division steps, the remainder drops by at least a factor of two: $r_{k+2} < r_k/2$. This gives an upper bound of $2\lfloor \log_2 b \rfloor + 1$ steps, making the algorithm logarithmic in the size of its input. A finer analysis shows that the worst-case inputs are consecutive Fibonacci numbers, where each quotient q_i equals 1 and the remainders decrease as slowly as possible.

Example 0.1.7: Computing a GCD

Take $a = 252$ and $b = 105$. The Euclidean algorithm proceeds:

$$\begin{aligned}
 252 &= 105 \cdot 2 + 42, \\
 105 &= 42 \cdot 2 + 21, \\
 42 &= 21 \cdot 2 + 0
 \end{aligned}$$

The last nonzero remainder is 21, so $\gcd(252, 105) = 21$. The algorithm terminates in three steps, reflecting the fact that $252/105$ has a short continued fraction expansion.

The Euclidean algorithm does more than compute the GCD—it simultaneously produces an explicit $\mathbb{Z}$ -linear combination. Tracing back through the successive divisions recovers the coefficients in what is arguably the single most useful identity in elementary number theory.

Theorem 0.1.8: Bézout's Identity

For any $a, b \in \mathbb{Z}$ not both zero, there exist $x, y \in \mathbb{Z}$ such that

$$ax + by = \gcd(a, b)$$

In particular, a and b are coprime if and only if 1 is a $\mathbb{Z}$ -linear combination of a and b .

This is a statement about the ideal structure of $\mathbb{Z}$: the ideal generated by a and b ,

$$(a, b) = \{ax + by \mid x, y \in \mathbb{Z}\},$$

is principal, generated by $\gcd(a, b)$. Every ideal of $\mathbb{Z}$ is principal—the ring $\mathbb{Z}$ is a PID. In rings of integers of number fields, this property can fail: in $\mathbb{Z}[\sqrt{-5}]$, for instance, the ideal $(2, 1 + \sqrt{-5})$ is not principal. The deviation from principality is measured by the ideal class group, one of the central objects studied in [17] and a key player in the analytic class number formula of chapter 8.

Proof:

Define d to be the smallest positive element of $I = \{ax + by \mid x, y \in \mathbb{Z}\}$. This set contains positive elements (for instance, $a^2 + b^2$ if a and b are not both zero), so d exists by the well-ordering principle.

Step 1: d divides a and b . Apply the division algorithm to write $a = dq + r$ with $0 \leq r < d$. Since $d = ax_0 + by_0$ for some $x_0, y_0 \in \mathbb{Z}$, the remainder satisfies

$$r = a - dq = a(1 - qx_0) + b(-qy_0) \in I$$

If $r > 0$, then r is a positive element of I smaller than d , contradicting the choice of d . Hence $r = 0$, so $d \mid a$. The identical argument with b in place of a gives $d \mid b$.

Step 2: $d = \gcd(a, b)$. Since $d \mid a$ and $d \mid b$, the integer d is a common divisor. If c is any common divisor of a and b , then c divides every element of I (by proposition 0.1.3), and in particular $c \mid d$. Hence d is the greatest common divisor.

The proof is clean but non-constructive: it asserts existence via the well-ordering principle without producing x and y explicitly. To find them, one runs the Euclidean algorithm forward and then substitutes backward through the chain of divisions (see example 0.1.10 below).

A direct and frequently used consequence of Bézout's identity is the following cancellation property for coprime integers. It is the mechanism underlying unique factorization in the next section.

Corollary 0.1.9: Euclid's Lemma for Coprime Integers

If $a \mid bc$ and $\gcd(a, b) = 1$, then $a \mid c$.

Proof:

By theorem 0.1.8, there exist $x, y \in \mathbb{Z}$ with $ax + by = 1$. Multiplying through by c gives $acx + bcy = c$. Since $a \mid acx$ (trivially) and $a \mid bcy$ (by hypothesis), proposition 0.1.3 gives $a \mid c$.

The proof illustrates the standard technique: invoke Bézout to produce a linear combination equal to 1, then multiply by the desired target. This argument generalizes verbatim to any principal ideal domain, but fails in rings where Bézout's identity does not hold—such as $\mathbb{Z}[\sqrt{-5}]$, where $2 \mid (1 + \sqrt{-5})(1 - \sqrt{-5}) = 6$ but 2 divides neither factor.

Example 0.1.10: Back-Substitution for Bézout Coefficients

Continuing example 0.1.7, the Euclidean algorithm gave:

$$252 = 105 \cdot 2 + 42,$$

$$105 = 42 \cdot 2 + 21$$

From the second equation: $21 = 105 - 42 \cdot 2$. Substituting $42 = 252 - 105 \cdot 2$ from the first:

$$21 = 105 - (252 - 105 \cdot 2) \cdot 2 = 105 \cdot 5 + 252 \cdot (-2)$$

This gives $252 \cdot (-2) + 105 \cdot 5 = 21 = \gcd(252, 105)$, so the Bézout coefficients are $x = -2$ and $y = 5$. These are not unique: the general solution to $252x + 105y = 21$ is $(x, y) = (-2 + 5t, 5 - 12t)$ for $t \in \mathbb{Z}$, as the next result makes precise.

Bézout's identity settles the question of when a linear equation over $\mathbb{Z}$ has integer solutions.

Proposition 0.1.11: Linear Diophantine Equations

Let $a, b, c \in \mathbb{Z}$ with a and b not both zero, and set $d = \gcd(a, b)$. The equation

$$ax + by = c$$

has a solution $(x, y) \in \mathbb{Z}^2$ if and only if $d \mid c$. When solutions exist, they form the family

$$(x, y) = \left(x_0 + \frac{b}{d}t, y_0 - \frac{a}{d}t \right), \quad t \in \mathbb{Z},$$

where (x_0, y_0) is any particular solution.

Proof:

If (x, y) is a solution, then $d \mid a$ and $d \mid b$ give $d \mid (ax + by) = c$, establishing necessity. Conversely, if $d \mid c$, then theorem 0.1.8 provides $x'_0, y'_0 \in \mathbb{Z}$ with $ax'_0 + by'_0 = d$, and multiplying through by c/d gives the particular solution $(x_0, y_0) = ((c/d)x'_0, (c/d)y'_0)$.

For the general solution, note that (x, y) satisfies $ax + by = c$ if and only if $a(x - x_0) = -b(y - y_0)$. Dividing by d yields

$$\frac{a}{d}(x - x_0) = -\frac{b}{d}(y - y_0)$$

Since $\gcd(a/d, b/d) = 1$, corollary 0.1.9 gives $(b/d) \mid (x - x_0)$. Writing $x - x_0 = (b/d)t$ and substituting gives $y - y_0 = -(a/d)t$, as claimed.

The essential mechanism is the reduction to the coprime case: dividing by $d = \gcd(a, b)$ produces the coprimality condition $\gcd(a/d, b/d) = 1$, which then forces the structure of the solution set via

Bézout. This pattern—divide out the GCD, argue in the coprime case, and then scale back—recurs throughout number theory, from the analysis of congruences in the next section to the study of Dirichlet characters in chapter 7.

Example 0.1.12: Solving a Linear Diophantine Equation

Consider $14x + 35y = 63$. Here $\gcd(14, 35) = 7$ and $7 \mid 63$, so solutions exist. Dividing through by 7 reduces the problem to $2x + 5y = 9$. By inspection, $(x_0, y_0) = (2, 1)$ satisfies $2 \cdot 2 + 5 \cdot 1 = 9$. The general solution to the original equation is

$$(x, y) = (2 + 5t, 1 - 2t), \quad t \in \mathbb{Z},$$

producing the solutions $\dots, (-3, 3), (2, 1), (7, -1), (12, -3), \dots$. For instance, the unique solution with $0 \leq x < 5$ is $(x, y) = (2, 1)$.

0.2 Primes and the Fundamental Theorem of Arithmetic

The Euclidean algorithm and Bézout’s identity treat all integers uniformly, without distinguishing between those that are “composed” and those that are not. This section introduces the multiplicative building blocks of $\mathbb{Z}$ —the prime numbers—and proves that every positive integer factors uniquely into primes. Unique factorization is the structural fact that makes the Euler product formula work (chapter 4): the identity $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ is, at its core, a restatement of the Fundamental Theorem of Arithmetic in the language of Dirichlet series.

Definition 0.2.1: Prime and Irreducible Elements

An integer $p > 1$ is *irreducible* if its only positive divisors are 1 and p . An integer $p > 1$ is *prime* if whenever $p \mid ab$, either $p \mid a$ or $p \mid b$.

In $\mathbb{Z}$, these two notions coincide—a fact that depends on Euclid’s lemma and therefore on the Euclidean algorithm. The distinction matters in more general rings. In $\mathbb{Z}[\sqrt{-5}]$, for instance, 2 is irreducible (it cannot be written as a product of two non-units) but not prime: $2 \mid (1 + \sqrt{-5})(1 - \sqrt{-5}) = 6$, yet 2 divides neither factor. The failure of “irreducible implies prime” is precisely the obstruction to unique factorization, and its resolution in number rings via the passage to ideals is a central theme of [17].

Proposition 0.2.2: In $\mathbb{Z}$, Prime and Irreducible Coincide

An integer $p > 1$ is prime if and only if it is irreducible.

Proof:

Prime implies irreducible. Suppose p is prime and $p = ab$ with $a, b > 0$. Then $p \mid ab$, so $p \mid a$ or $p \mid b$. If $p \mid a$, then since $a \leq p$ (as $b \geq 1$), this forces $a = p$ and $b = 1$. Similarly if $p \mid b$. Hence p is irreducible.

Irreducible implies prime. Suppose p is irreducible and $p \mid ab$. If $p \nmid a$, then the only positive common divisor of p and a is 1 (since p is irreducible and its only positive divisors are 1 and p ,

and $p \nmid a$ rules out p). Hence $\gcd(p, a) = 1$, and corollary 0.1.9 gives $p \mid b$.

The key step—the implication “irreducible implies prime”—relies entirely on corollary 0.1.9, which in turn depends on Bézout’s identity. This chain of dependencies is the mechanism by which the division algorithm guarantees unique factorization in $\mathbb{Z}$. In rings without a division algorithm (or more precisely, without the property that irreducibles are prime), unique factorization can fail.

The first result about the collection of all primes is among the oldest in mathematics.

Proposition 0.2.3: Infinitude of Primes

There are infinitely many primes.

Proof:

Suppose for contradiction that $p_1, p_2, \dots, p_n$ is a complete list of all primes. Consider the integer $N = p_1 p_2 \cdots p_n + 1$. Since $N > 1$, it has a prime divisor p . For each i , the prime p_i divides $p_1 \cdots p_n$ but does not divide N (since dividing N by p_i leaves remainder 1). Hence p is not among $p_1, \dots, p_n$, contradicting the assumption that the list was complete.

Euclid’s argument establishes the infinitude of primes but says nothing about how they are distributed among the integers. The quantitative study of this distribution—how many primes are there up to x , and how are they spaced?—is the central question of analytic number theory. The Prime Number Theorem (chapter 5) gives the asymptotic answer $\pi(x) \sim x/\log x$, but reaching it requires the full power of complex analysis applied to $\zeta(s)$.

The Fundamental Theorem of Arithmetic asserts that every integer greater than 1 can be written as a product of primes, and that this product is essentially unique.

Theorem 0.2.4: Fundamental Theorem of Arithmetic

Every integer $n > 1$ has a factorization

$$n = p_1^{a_1} p_2^{a_2} \cdots p_k^{a_k}$$

where $p_1 < p_2 < \cdots < p_k$ are primes and $a_1, \dots, a_k \geq 1$. This factorization is unique: if $n = q_1^{b_1} \cdots q_\ell^{b_\ell}$ is another such factorization with $q_1 < \cdots < q_\ell$, then $k = \ell$, $p_i = q_i$, and $a_i = b_i$ for all i .

Proof:

Step 1: Existence. Proceed by strong induction on n . If $n = 2$, then n is prime and $n = 2$ is itself a prime factorization. For $n > 2$: if n is prime, the factorization $n = n^1$ works. If n is not prime, then $n = ab$ for some $1 < a, b < n$, and by the inductive hypothesis both a and b have prime factorizations. Concatenating and reordering gives a prime factorization of n .

Step 2: Uniqueness. Suppose $p_1^{a_1} \cdots p_k^{a_k} = q_1^{b_1} \cdots q_\ell^{b_\ell}$. Then p_1 divides the right-hand side, so $p_1 \mid q_j^{b_j}$ for some j (by proposition 0.2.2, applied repeatedly). Since q_j is prime and $p_1 > 1$, this forces $p_1 = q_j$. As both lists are sorted in increasing order: if $p_1 < q_1$, then $p_1 \mid q_1^{b_1} \cdots q_\ell^{b_\ell}$ but $p_1 \neq q_j$ for any j (since $p_1 < q_1 \leq q_j$), a contradiction. Similarly $q_1 < p_1$ is impossible. Hence

$$p_1 = q_1.$$

Dividing both sides by $p_1^{\min(a_1, b_1)}$ and applying the same argument inductively shows that the two factorizations agree term by term. In particular, $k = \ell$, $p_i = q_i$, and $a_i = b_i$ for all i .

The existence of a prime factorization uses only the well-ordering of $\mathbb{N}$ (via strong induction). The uniqueness, on the other hand, is the substantive content: it depends on the fact that primes in $\mathbb{Z}$ satisfy the cancellation property of proposition 0.2.2, which traces back through Euclid's lemma to the division algorithm. Removing any link in this chain allows uniqueness to fail, as the example $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$ in $\mathbb{Z}[\sqrt{-5}]$ demonstrates.

Example 0.2.5: Prime Factorizations

1. $360 = 2^3 \cdot 3^2 \cdot 5$. This is the unique way to write 360 as a product of prime powers (up to reordering).
2. $2310 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11$. This is a *primorial*: the product of the first five primes. Primorials are squarefree integers (every prime appears with exponent exactly 1) and appear naturally in sieve arguments (??).
3. $10! = 2^8 \cdot 3^4 \cdot 5^2 \cdot 7$. The exponents can be computed directly from the Fundamental Theorem, but Legendre's formula (proved at the end of this section) provides a much faster method.

The Fundamental Theorem is the reason why the study of multiplicative properties of $\mathbb{Z}$ reduces to the study of primes. Every question about divisibility, common factors, or the multiplicative structure of an integer can be answered by examining its prime factorization. This reductive principle has an analytic counterpart: the Euler product identity $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ converts a sum over all positive integers into a product over primes, and it is through this product that the analytic properties of $\zeta(s)$ yield information about the distribution of primes.

With unique factorization established, every positive integer n is completely described by its *prime signature*: the function $p \mapsto v_p(n)$ giving the exponent of each prime in the factorization of n .

Definition 0.2.6: The p -adic Valuation

For a prime p and a nonzero integer n , the *p -adic valuation* of n , denoted $v_p(n)$, is the largest integer $e \geq 0$ such that $p^e \mid n$. By convention, $v_p(0) = +\infty$. The Fundamental Theorem of Arithmetic gives

$$|n| = \prod_p p^{v_p(n)}$$

where the product is over all primes and is finite (since $v_p(n) = 0$ for all but finitely many p).

The p -adic valuation converts multiplicative questions about $\mathbb{Z}$ into additive questions about $\mathbb{Z}_{\geq 0}$: if $n = ab$ then $v_p(n) = v_p(a) + v_p(b)$ for every prime p . This decomposition—analyzing a global multiplicative problem one prime at a time—is the “local-to-global” philosophy that pervades number theory, from the Euler product for $\zeta(s)$ to the adelic methods of class field theory.

The valuation extends naturally to $\mathbb{Q}^\times$ by setting $v_p(a/b) = v_p(a) - v_p(b)$, giving a group homomorphism $v_p : \mathbb{Q}^\times \rightarrow \mathbb{Z}$. Every nonzero rational number r can then be written uniquely as $r = \pm \prod_p p^{v_p(r)}$, where the product is finite. The associated *p -adic absolute value* $|r|_p = p^{-v_p(r)}$

defines a metric on $\mathbb{Q}$ whose completion is the field $\mathbb{Q}_p$ of p -adic numbers (see [15]). The interplay between all these completions—one for each prime and one for the ordinary absolute value—is a recurring theme beginning with the completed zeta function in chapter 6.

Proposition 0.2.7: Divisibility via Valuations

For nonzero integers a and b :

1. $a|b$ if and only if $v_p(a) \leq v_p(b)$ for every prime p .
2. $\gcd(a, b) = \prod_p p^{\min(v_p(a), v_p(b))}$.
3. $\text{lcm}(a, b) = \prod_p p^{\max(v_p(a), v_p(b))}$.
4. $\gcd(a, b) \cdot \text{lcm}(a, b) = |ab|$.

Proof:

For (1): $a|b$ means $b = ak$ for some $k \in \mathbb{Z}$, which translates to $v_p(b) = v_p(a) + v_p(k) \geq v_p(a)$ for all p . The converse follows by writing $k = b/a$ and checking that $v_p(k) = v_p(b) - v_p(a) \geq 0$ for all p , so k is an integer. Parts (2) and (3) follow from (1): the greatest common divisor is the largest integer dividing both a and b , which means taking the minimum exponent at each prime; the least common multiple is the smallest integer divisible by both, which means taking the maximum. Part (4) follows from the identity $\min(x, y) + \max(x, y) = x + y$.

The valuation-based characterization of divisibility is the “right” perspective for many computations. For example, it makes the following formula for gcd immediate.

Example 0.2.8: GCD and LCM via Valuations

1. Take $a = 252 = 2^2 \cdot 3^2 \cdot 7$ and $b = 105 = 3 \cdot 5 \cdot 7$. Reading off:

$$\gcd(252, 105) = 2^0 \cdot 3^1 \cdot 5^0 \cdot 7^1 = 21, \quad \text{lcm}(252, 105) = 2^2 \cdot 3^2 \cdot 5 \cdot 7 = 1260$$

As a check: $\gcd \cdot \text{lcm} = 21 \cdot 1260 = 26460 = 252 \cdot 105 = |ab|$.

2. For $a = 360 = 2^3 \cdot 3^2 \cdot 5$ and $b = 750 = 2 \cdot 3 \cdot 5^3$:

$$\gcd(360, 750) = 2^1 \cdot 3^1 \cdot 5^1 = 30, \quad \text{lcm}(360, 750) = 2^3 \cdot 3^2 \cdot 5^3 = 9000$$

The Fundamental Theorem also provides a clean way to express many arithmetic functions. The number of positive divisors of $n = \prod p_i^{a_i}$ is $\prod (a_i + 1)$, since each divisor is obtained by independently choosing an exponent between 0 and a_i at each prime. The sum of divisors is $\prod (1 + p_i + p_i^2 + \cdots + p_i^{a_i}) = \prod \frac{p_i^{a_i+1} - 1}{p_i - 1}$. These formulas are simple consequences of unique factorization combined with the Chinese Remainder Theorem, and they generalize to the full family of multiplicative arithmetic functions studied in chapter 1.

Example 0.2.9: Divisor Count and Divisor Sum

1. The integer $n = 360 = 2^3 \cdot 3^2 \cdot 5$ has $(3+1)(2+1)(1+1) = 24$ positive divisors.
2. The sum of divisors of 360 is

$$\sigma(360) = \frac{2^4 - 1}{2 - 1} \cdot \frac{3^3 - 1}{3 - 1} \cdot \frac{5^2 - 1}{5 - 1} = 15 \cdot 13 \cdot 6 = 1170$$

3. For a prime p : the number of divisors of p^k is $k+1$, and $\sigma(p^k) = \frac{p^{k+1}-1}{p-1}$. In particular, $\sigma(p) = p+1$.

A useful bookkeeping device that appears throughout the book is the *canonical factorization notation*. For $n = \prod_{i=1}^k p_i^{a_i}$, a sum or product “over the prime factorization of n ” means over the pairs (p_i, a_i) . The notation $p^a \parallel n$ (read “ p^a exactly divides n ”) means $v_p(n) = a$, i.e., $p^a \mid n$ but $p^{a+1} \nmid n$.

Definition 0.2.10: Exact Divisibility

For a prime p and a positive integer n , the notation $p^a \parallel n$ means $v_p(n) = a$, that is, $p^a \mid n$ and $p^{a+1} \nmid n$.

To close this section, a formula that illustrates how the Fundamental Theorem interacts with factorials. It will be used in the proof of Chebyshev’s bounds in chapter 3.

Proposition 0.2.11: Legendre’s Formula

For any prime p and positive integer n :

$$v_p(n!) = \sum_{k=1}^{\infty} \left\lfloor \frac{n}{p^k} \right\rfloor$$

The sum is finite, as $\lfloor n/p^k \rfloor = 0$ for $p^k > n$.

Proof:

The exponent $v_p(n!)$ counts the total number of factors of p appearing in the product $1 \cdot 2 \cdots n$. Among the integers $1, 2, \dots, n$, there are exactly $\lfloor n/p \rfloor$ multiples of p , each contributing at least one factor. Of these, $\lfloor n/p^2 \rfloor$ are multiples of p^2 , each contributing a second factor. Continuing, $\lfloor n/p^k \rfloor$ multiples of p^k each contribute a k -th factor. Summing over all k counts each factor of p exactly once, giving the formula.

Example 0.2.12: Applying Legendre’s Formula

The exponent of 2 in $20!$ is

$$v_2(20!) = \left\lfloor \frac{20}{2} \right\rfloor + \left\lfloor \frac{20}{4} \right\rfloor + \left\lfloor \frac{20}{8} \right\rfloor + \left\lfloor \frac{20}{16} \right\rfloor = 10 + 5 + 2 + 1 = 18$$

Similarly, $v_3(20!) = 6 + 2 = 8$ and $v_5(20!) = 4$. In particular, $20!$ is divisible by $10^4 = 2^4 \cdot 5^4$ but not by 10^5 , so $20! = 2432902008176640000$ ends in exactly four trailing zeros.

An immediate consequence of Legendre’s formula is a clean expression for the p -adic valuation of binomial coefficients, which plays a role in Chebyshev’s estimates (chapter 3).

Corollary 0.2.13: Valuation of Binomial Coefficients

For a prime p and integers $0 \leq m \leq n$:

$$v_p \binom{n}{m} = \sum_{k=1}^{\infty} \left(\left\lfloor \frac{n}{p^k} \right\rfloor - \left\lfloor \frac{m}{p^k} \right\rfloor - \left\lfloor \frac{n-m}{p^k} \right\rfloor \right)$$

Each summand is either 0 or 1, and it equals 1 precisely when the addition $m + (n - m)$ produces a carry at the p^k -place in base p .

Proof:

Since $\binom{n}{m} = n!/(m!(n-m)!)$, the formula follows from Legendre’s formula and the additivity of v_p :

$$v_p \binom{n}{m} = v_p(n!) - v_p(m!) - v_p((n-m)!) = \sum_{k \geq 1} \left(\left\lfloor \frac{n}{p^k} \right\rfloor - \left\lfloor \frac{m}{p^k} \right\rfloor - \left\lfloor \frac{n-m}{p^k} \right\rfloor \right)$$

For the “carry” interpretation: let $a = m$ and $b = n - m$, so $a + b = n$. The identity $\lfloor (a+b)/p^k \rfloor - \lfloor a/p^k \rfloor - \lfloor b/p^k \rfloor$ equals 1 when the addition of the base- p digits of a and b produces a carry into the p^k -position, and 0 otherwise. This is Kummer’s theorem.

Example 0.2.14: Central Binomial Coefficient

Consider $\binom{10}{5} = 252$. In base 2: $5 = (101)_2$ and $5 + 5 = 10 = (1010)_2$. Adding $(101)_2 + (101)_2$, carries occur at positions 2^0 and 2^2 , so $v_2 \binom{10}{5} = 2$. Indeed, $252 = 2^2 \cdot 3^2 \cdot 7$. Similarly, in base 3: $5 = (12)_3$ and $5 + 5 = 10 = (101)_3$; adding $(12)_3 + (12)_3$ produces carries at 3^0 and 3^1 , giving $v_3 \binom{10}{5} = 2$.

0.3 Congruences

The previous sections studied the integers through their multiplicative structure: divisibility, primes, and factorization. Congruences introduce a complementary perspective by collapsing $\mathbb{Z}$ into finite quotient rings $\mathbb{Z}/m\mathbb{Z}$. This modular viewpoint is the natural setting for Dirichlet characters (chapter 7), and the Chinese Remainder Theorem—the structural result of this section—is the tool that decomposes these characters into local components, one for each prime power dividing the modulus.

Definition 0.3.1: Congruence

Let $m \in \mathbb{Z}_{>0}$. Two integers a and b are *congruent modulo m* , written $a \equiv b \pmod{m}$, if $m \mid (a - b)$. Equivalently, a and b leave the same remainder upon division by m .

Congruence modulo m is an equivalence relation on $\mathbb{Z}$, partitioning it into m equivalence classes called *residue classes*. The set of residue classes forms a ring under the inherited addition and multiplication:

Definition 0.3.2: The Ring $\mathbb{Z}/m\mathbb{Z}$

The *residue class ring modulo m* is the quotient ring

$$\mathbb{Z}/m\mathbb{Z} = \{0, 1, 2, \dots, m-1\}$$

with addition and multiplication performed modulo m . The canonical projection $\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ sends each integer to its residue class.

The ring $\mathbb{Z}/m\mathbb{Z}$ is the simplest example of a quotient ring, and its structure—which residue classes are units, which are zero divisors, and how the ring decomposes—is controlled entirely by the prime factorization of m . The key structural result is the Chinese Remainder Theorem, which says that if m factors into coprime parts, then $\mathbb{Z}/m\mathbb{Z}$ splits as a product of smaller rings.

Before stating it, the following criterion for invertibility modulo m is needed.

Proposition 0.3.3: Units of $\mathbb{Z}/m\mathbb{Z}$

An element $a \in \mathbb{Z}/m\mathbb{Z}$ is a unit (i.e., has a multiplicative inverse modulo m) if and only if $\gcd(a, m) = 1$. When this holds, the inverse can be computed via the Euclidean algorithm: if $ax + my = 1$ (from theorem 0.1.8), then $a^{-1} \equiv x \pmod{m}$.

Proof:

If $\gcd(a, m) = 1$, then theorem 0.1.8 gives $ax + my = 1$ for some $x, y \in \mathbb{Z}$, which reads $ax \equiv 1 \pmod{m}$, so x is the inverse of a . Conversely, if $ax \equiv 1 \pmod{m}$ for some x , then $ax = 1 + km$ for some k , giving $ax - km = 1$. By proposition 0.1.3, any common divisor of a and m divides 1, so $\gcd(a, m) = 1$.

The group of units $(\mathbb{Z}/m\mathbb{Z})^\times = \{a \in \mathbb{Z}/m\mathbb{Z} \mid \gcd(a, m) = 1\}$ has order $\varphi(m)$, where φ is the Euler totient function. This group is the domain on which Dirichlet characters live, and its structure will be analyzed in detail in the next section.

An element $a \in \mathbb{Z}/m\mathbb{Z}$ with $a \neq 0$ and $\gcd(a, m) > 1$ is a *zero divisor*: writing $d = \gcd(a, m)$ with $1 < d < m$, the product $a \cdot (m/d) \equiv 0 \pmod{m}$ is zero even though neither factor is. Thus $\mathbb{Z}/m\mathbb{Z}$ is an integral domain if and only if m is prime, in which case every nonzero element is a unit and $\mathbb{Z}/m\mathbb{Z} = \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p$ is a field with p elements. This is the finite field that appears throughout the book, from character sums (??) to the local factors of L -functions.

Example 0.3.4: Units and Zero Divisors in $\mathbb{Z}/12\mathbb{Z}$

The elements of $\mathbb{Z}/12\mathbb{Z}$ partition as follows:

1. *Units* ($\gcd(a, 12) = 1$): $\{1, 5, 7, 11\}$. Each has a multiplicative inverse: $5 \cdot 5 = 25 \equiv 1$, $7 \cdot 7 = 49 \equiv 1$, $11 \cdot 11 = 121 \equiv 1$. So $5^{-1} = 5$, $7^{-1} = 7$, $11^{-1} = 11$ in $\mathbb{Z}/12\mathbb{Z}$.
2. *Zero divisors* ($1 < \gcd(a, 12) < 12$, $a \neq 0$): $\{2, 3, 4, 6, 8, 9, 10\}$. For instance, $3 \cdot 4 = 12 \equiv 0$, $6 \cdot 2 = 12 \equiv 0$.
3. *The zero element*: $\{0\}$.

Definition 0.3.5: Euler Totient Function

For $m \in \mathbb{Z}_{>0}$, the *Euler totient function* $\varphi(m)$ is the number of integers a with $1 \leq a \leq m$ and $\gcd(a, m) = 1$. Equivalently, $\varphi(m) = |(\mathbb{Z}/m\mathbb{Z})^\times|$.

Example 0.3.6: Small Values of φ

1. $\varphi(1) = 1$, since $\gcd(1, 1) = 1$.
2. $\varphi(p) = p - 1$ for any prime p , since every integer from 1 to $p - 1$ is coprime to p .
3. $\varphi(12) = |\{1, 5, 7, 11\}| = 4$, as these are the residues coprime to 12.
4. $\varphi(p^k) = p^k - p^{k-1} = p^{k-1}(p - 1)$ for a prime power: among $1, \dots, p^k$, the multiples of p are $p, 2p, \dots, p^{k-1} \cdot p$, and there are p^{k-1} of them.

The multiplicativity of φ is a consequence of the Chinese Remainder Theorem, stated next.

Theorem 0.3.7: Chinese Remainder Theorem

Let $m_1, m_2, \dots, m_k$ be pairwise coprime positive integers, and set $M = m_1 m_2 \cdots m_k$. Then the map

$$\mathbb{Z}/M\mathbb{Z} \xrightarrow{\sim} \mathbb{Z}/m_1\mathbb{Z} \times \mathbb{Z}/m_2\mathbb{Z} \times \cdots \times \mathbb{Z}/m_k\mathbb{Z}, \quad a \mapsto (a \bmod m_1, a \bmod m_2, \dots, a \bmod m_k)$$

is a ring isomorphism.

The CRT says that specifying a residue class modulo M is the same as independently specifying a residue class modulo each m_i . This is the arithmetic analogue of the decomposition of a product of topological spaces into its factors, and it is the mechanism that allows global questions about $\mathbb{Z}/M\mathbb{Z}$ to be reduced to local questions about each $\mathbb{Z}/m_i\mathbb{Z}$.

Proof:

Step 1: The map is a ring homomorphism. The reduction map $a \mapsto (a \bmod m_1, \dots, a \bmod m_k)$ preserves addition and multiplication, since these operations in each factor are performed modulo m_i .

Step 2: Injectivity. If $a \equiv 0 \pmod{m_i}$ for all i , then $m_i \mid a$ for each i . Since the m_i are pairwise coprime, $M = m_1 \cdots m_k$ divides a (apply corollary 0.1.9 inductively: $m_1 \mid a$ and $\gcd(m_1, m_2) = 1$ together with $m_2 \mid a$ give $m_1 m_2 \mid a$; continuing gives $M \mid a$). Hence the kernel is trivial.

Step 3: Surjectivity. Both sides have the same cardinality $M = m_1 \cdots m_k$, so injectivity implies surjectivity.

Alternatively, surjectivity can be shown constructively. For each i , define $M_i = M/m_i$ and note that $\gcd(M_i, m_i) = 1$ (since the m_j for $j \neq i$ are all coprime to m_i). By theorem 0.1.8, there

exists y_i with $M_i y_i \equiv 1 \pmod{m_i}$. Given a target $(a_1, \dots, a_k)$, the integer

$$a = \sum_{i=1}^k a_i M_i y_i$$

satisfies $a \equiv a_i \pmod{m_i}$ for each i , since $M_j y_j \equiv 0 \pmod{m_i}$ when $j \neq i$ (as $m_i \mid M_j$) and $M_i y_i \equiv 1 \pmod{m_i}$.

The constructive proof is useful in practice: it gives an explicit formula for “gluing” local data into a global solution. The abstract proof via cardinality counting is cleaner but less informative.

Example 0.3.8: Solving a System of Congruences

Solve the system $x \equiv 2 \pmod{3}$, $x \equiv 3 \pmod{5}$, $x \equiv 2 \pmod{7}$. Here $M = 105$, $M_1 = 35$, $M_2 = 21$, $M_3 = 15$. The inverses are: $35y_1 \equiv 1 \pmod{3}$, giving $y_1 = 2$ (since $35 \cdot 2 = 70 \equiv 1$); $21y_2 \equiv 1 \pmod{5}$, giving $y_2 = 1$ (since $21 \equiv 1$); $15y_3 \equiv 1 \pmod{7}$, giving $y_3 = 1$ (since $15 \equiv 1$). Then

$$x = 2 \cdot 35 \cdot 2 + 3 \cdot 21 \cdot 1 + 2 \cdot 15 \cdot 1 = 140 + 63 + 30 = 233 \equiv 23 \pmod{105}$$

One verifies: $23 = 7 \cdot 3 + 2 \equiv 2 \pmod{3}$, $23 = 4 \cdot 5 + 3 \equiv 3 \pmod{5}$, $23 = 3 \cdot 7 + 2 \equiv 2 \pmod{7}$.

An immediate corollary of the CRT is the multiplicativity of the Euler totient.

Corollary 0.3.9: Multiplicativity of φ

If $\gcd(m, n) = 1$, then $\varphi(mn) = \varphi(m)\varphi(n)$.

Proof:

By theorem 0.3.7, $(\mathbb{Z}/mn\mathbb{Z})^\times \cong (\mathbb{Z}/m\mathbb{Z})^\times \times (\mathbb{Z}/n\mathbb{Z})^\times$, since the isomorphism $\mathbb{Z}/mn\mathbb{Z} \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$ restricts to an isomorphism on unit groups (an element is a unit in the product if and only if each component is a unit). Taking cardinalities: $\varphi(mn) = \varphi(m)\varphi(n)$.

Combined with the prime power formula $\varphi(p^k) = p^{k-1}(p-1)$ from example 0.3.6, this gives a complete formula for φ .

Proposition 0.3.10: Formula for the Euler Totient

For any $n = p_1^{a_1} \cdots p_k^{a_k} > 1$:

$$\varphi(n) = n \prod_{p \mid n} \left(1 - \frac{1}{p}\right) = \prod_{i=1}^k p_i^{a_i-1} (p_i - 1)$$

Proof:

By corollary 0.3.9 and induction, $\varphi(n) = \prod_{i=1}^k \varphi(p_i^{a_i}) = \prod_{i=1}^k p_i^{a_i-1} (p_i - 1)$. Factoring out $p_i^{a_i}$ from each term gives the product formula $\varphi(n) = n \prod_{p \mid n} (1 - 1/p)$.

Example 0.3.11: Computing φ

$$\varphi(360) = \varphi(2^3 \cdot 3^2 \cdot 5) = 360 \cdot (1 - \frac{1}{2})(1 - \frac{1}{3})(1 - \frac{1}{5}) = 360 \cdot \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{4}{5} = 96.$$

The section closes with the two classical results that connect the Euler totient to modular exponentiation. These are the finite-group-theoretic facts that underlie the convergence arguments for Dirichlet series twisted by characters.

Theorem 0.3.12: Euler's Theorem

If $\gcd(a, m) = 1$, then $a^{\varphi(m)} \equiv 1 \pmod{m}$.

Proof:

The residue class of a is an element of the group $(\mathbb{Z}/m\mathbb{Z})^\times$, which has order $\varphi(m)$. By Lagrange's theorem (see [13]), the order of every element divides the order of the group. Hence $a^{\varphi(m)} \equiv 1 \pmod{m}$.

Euler's theorem is a statement about the group $(\mathbb{Z}/m\mathbb{Z})^\times$: every element has finite order dividing $\varphi(m)$. The special case $m = p$ prime gives the result traditionally attributed to Fermat.

Corollary 0.3.13: Fermat's Little Theorem

If p is prime and $p \nmid a$, then $a^{p-1} \equiv 1 \pmod{p}$. Equivalently, $a^p \equiv a \pmod{p}$ for all $a \in \mathbb{Z}$.

Proof:

Since $\varphi(p) = p - 1$, the first statement is a special case of theorem 0.3.12. For the second: if $p \mid a$ then $a^p \equiv 0 \equiv a \pmod{p}$; if $p \nmid a$ then $a^p = a \cdot a^{p-1} \equiv a \cdot 1 = a \pmod{p}$.

Another classical characterization of primes via modular arithmetic is Wilson's theorem.

Proposition 0.3.14: Wilson's Theorem

An integer $n \geq 2$ is prime if and only if $(n-1)! \equiv -1 \pmod{n}$.

Proof:

If n is composite, say $n = ab$ with $1 < a \leq b < n$, then a appears as a factor in $(n-1)!$. If $a \neq b$, both appear, and $n = ab$ divides $(n-1)!$, so $(n-1)! \equiv 0 \not\equiv -1 \pmod{n}$. If $a = b$, then $n = a^2 \geq 4$ (since $a \geq 2$), and both a and $2a$ appear in $\{1, \dots, n-1\}$ when $n > 4$; the case $n = 4$ is verified directly: $3! = 6 \equiv 2 \not\equiv -1 \pmod{4}$.

If $n = p$ is prime, then $\mathbb{F}_p^\times = \{1, 2, \dots, p-1\}$ is a group of order $p-1$. Each element a pairs with its inverse a^{-1} . The elements satisfying $a = a^{-1}$, i.e., $a^2 \equiv 1 \pmod{p}$, are exactly $a \equiv \pm 1 \pmod{p}$ (since $p \mid (a-1)(a+1)$ forces $p \mid (a-1)$ or $p \mid (a+1)$). Hence in the product $1 \cdot 2 \cdots (p-1)$, all elements except 1 and $p-1$ cancel in pairs, leaving $(p-1)! \equiv 1 \cdot (p-1) = p-1 \equiv -1 \pmod{p}$.

Wilson's theorem gives an exact criterion for primality, but it is computationally impractical since computing $(n-1)!$ modulo n requires $\Theta(n)$ multiplications. Its theoretical importance lies in providing explicit generators of ideals: the identity $(p-1)! + 1 \equiv 0 \pmod{p}$ is used in the proof of quadratic reciprocity (section 0.5).

Fermat's little theorem has a useful contrapositive: if $a^{p-1} \not\equiv 1 \pmod{p}$, then p is not prime. This is the basis of primality testing algorithms. The converse fails: there exist composite numbers n (called *Carmichael numbers*) such that $a^{n-1} \equiv 1 \pmod{n}$ for all a coprime to n . The smallest is $n = 561 = 3 \cdot 11 \cdot 17$, which satisfies this property because $(p-1) \mid 560$ for each prime factor $p \in \{3, 11, 17\}$ (since $560 = 2 \cdot 280 = 10 \cdot 56 = 16 \cdot 35$), and the CRT then guarantees $a^{560} \equiv 1 \pmod{561}$ for all a with $\gcd(a, 561) = 1$. This criterion—known as *Korselt's criterion*—states that n is a Carmichael number if and only if n is squarefree and $(p-1) \mid (n-1)$ for every prime p dividing n .

Example 0.3.15: Modular Exponentiation via Euler's Theorem

To compute $7^{222} \bmod 10$, note that $\varphi(10) = 4$ and $\gcd(7, 10) = 1$, so $7^4 \equiv 1 \pmod{10}$ by theorem 0.3.12. Since $222 = 4 \cdot 55 + 2$:

$$7^{222} = (7^4)^{55} \cdot 7^2 \equiv 1^{55} \cdot 49 \equiv 9 \pmod{10}$$

The last digit of 7^{222} is therefore 9. More generally, the last digit of 7^n cycles with period 4: 7, 9, 3, 1, 7, 9, 3, 1, ... —a consequence of 7 having order exactly 4 in $(\mathbb{Z}/10\mathbb{Z})^\times$.

A complementary identity that counts divisors weighted by the totient is the following.

Proposition 0.3.16: Totient Summation Identity

For every positive integer n :

$$\sum_{d \mid n} \varphi(d) = n$$

Proof:

Partition the set $\{1, 2, \dots, n\}$ according to $\gcd(k, n)$. For each divisor d of n , the integers $k \in \{1, \dots, n\}$ with $\gcd(k, n) = d$ are exactly those of the form $k = d\ell$ where $1 \leq \ell \leq n/d$ and $\gcd(\ell, n/d) = 1$. There are $\varphi(n/d)$ such integers. Summing over all divisors d of n gives $n = \sum_{d \mid n} \varphi(n/d)$. Since $d \mapsto n/d$ is a bijection on the divisors of n , this equals $\sum_{d \mid n} \varphi(d)$.

This identity will reappear in chapter 1 as the Dirichlet convolution identity $\varphi * \mathbf{1} = \text{id}$, where $\mathbf{1}$ is the constant function 1 and id is the identity function $n \mapsto n$. Möbius inversion then recovers φ from this relation.

0.4 The Group $(\mathbb{Z}/m\mathbb{Z})^\times$ and Primitive Roots

Euler's theorem (theorem 0.3.12) guarantees that every element of $(\mathbb{Z}/m\mathbb{Z})^\times$ has finite multiplicative order dividing $\varphi(m)$. A natural question is whether this bound is achieved: does $(\mathbb{Z}/m\mathbb{Z})^\times$ always contain an element of order exactly $\varphi(m)$? Equivalently, is $(\mathbb{Z}/m\mathbb{Z})^\times$ cyclic? The answer is no in general— $(\mathbb{Z}/8\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ has exponent 2, so no element has order $\varphi(8) = 4$ —but a precise

characterization exists. Understanding the group structure of $(\mathbb{Z}/m\mathbb{Z})^\times$ is essential for the theory of Dirichlet characters in chapter 7, since the characters of a finite abelian group are determined by its invariant factor decomposition.

Definition 0.4.1: Multiplicative Order

For $a \in (\mathbb{Z}/m\mathbb{Z})^\times$, the *multiplicative order* of a modulo m , denoted $\text{ord}_m(a)$, is the smallest positive integer k such that $a^k \equiv 1 \pmod{m}$.

By theorem 0.3.12, $\text{ord}_m(a)$ exists and divides $\varphi(m)$. The order is the period of the cyclic subgroup $\langle a \rangle = \{1, a, a^2, \dots, a^{k-1}\}$ inside $(\mathbb{Z}/m\mathbb{Z})^\times$, and the standard theory of cyclic groups (see [13]) gives the following.

Proposition 0.4.2: Basic Properties of Multiplicative Order

Let $a \in (\mathbb{Z}/m\mathbb{Z})^\times$ with $\text{ord}_m(a) = k$. Then:

1. $a^n \equiv 1 \pmod{m}$ if and only if $k \mid n$.
2. $\text{ord}_m(a^j) = k / \gcd(j, k)$.
3. The number of elements of order k in $(\mathbb{Z}/m\mathbb{Z})^\times$ is either 0 or $\varphi(k)$.

Proof:

For (1): if $k \mid n$ then $a^n = (a^k)^{n/k} \equiv 1$. Conversely, write $n = kq + r$ with $0 \leq r < k$; then $a^r = a^n \cdot (a^k)^{-q} \equiv 1$, and minimality of k forces $r = 0$.

For (2): let $d = \gcd(j, k)$. Then $(a^j)^{k/d} = (a^k)^{j/d} \equiv 1$, so $\text{ord}_m(a^j)$ divides k/d . If $(a^j)^n \equiv 1$ then $k \mid jn$ by (1), hence $(k/d) \mid (j/d)n$. Since $\gcd(j/d, k/d) = 1$, corollary 0.1.9 gives $(k/d) \mid n$. Hence $\text{ord}_m(a^j) = k/d$.

For (3): if some element a has order k , then $\langle a \rangle$ contains exactly one element of each order dividing k , and by (2) the number of generators of $\langle a \rangle$ (i.e., elements of order k in $\langle a \rangle$) is $\varphi(k)$. To see that no element outside $\langle a \rangle$ has order k : the polynomial $x^k - 1$ has at most k roots in $\mathbb{Z}/p\mathbb{Z}$ (since $\mathbb{Z}/p\mathbb{Z}$ is a field), and the k elements of $\langle a \rangle$ already exhaust them. For composite m , the argument requires the structure theorem below.

Part (3) uses a critical property of fields: a polynomial of degree k over a field has at most k roots. This fails over $\mathbb{Z}/m\mathbb{Z}$ when m is composite—for example, $x^2 - 1 \equiv 0 \pmod{8}$ has four solutions $\{1, 3, 5, 7\}$ —and this failure is precisely why the counting argument in the existence theorem below works only for prime moduli.

Definition 0.4.3: Primitive Root

A *primitive root* modulo m is an element $g \in (\mathbb{Z}/m\mathbb{Z})^\times$ of order $\varphi(m)$, i.e., a generator of $(\mathbb{Z}/m\mathbb{Z})^\times$. Equivalently, $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic if and only if a primitive root exists.

When a primitive root g exists, every unit $a \in (\mathbb{Z}/m\mathbb{Z})^\times$ can be written as $a \equiv g^k \pmod{m}$ for a

unique $k \in \{0, 1, \dots, \varphi(m) - 1\}$. This integer k is the *discrete logarithm* of a to base g , and the map $a \mapsto k$ is an isomorphism $(\mathbb{Z}/m\mathbb{Z})^\times \xrightarrow{\sim} \mathbb{Z}/\varphi(m)\mathbb{Z}$ that converts multiplication to addition.

Example 0.4.4: The Discrete Logarithm Table Modulo 13

The group $(\mathbb{Z}/13\mathbb{Z})^\times$ has order 12. Testing $g = 2$: the successive powers are

$$\begin{aligned} 2^1 &= 2, & 2^2 &= 4, & 2^3 &= 8, & 2^4 &= 3, & 2^5 &= 6, & 2^6 &= 12, \\ 2^7 &= 11, & 2^8 &= 9, & 2^9 &= 5, & 2^{10} &= 10, & 2^{11} &= 7, & 2^{12} &= 1, \end{aligned}$$

confirming that 2 is a primitive root. The discrete logarithm $\log_2(a) \bmod 12$ converts multiplication in $(\mathbb{Z}/13\mathbb{Z})^\times$ to addition in $\mathbb{Z}/12\mathbb{Z}$. For instance, $5 \cdot 8 = 40 \equiv 1 \pmod{13}$, which corresponds to $\log_2(5) + \log_2(8) = 9 + 3 = 12 \equiv 0 \pmod{12}$. This conversion is the mechanism behind index calculus methods for computing in $(\mathbb{Z}/p\mathbb{Z})^\times$.

The following theorem characterizes exactly when primitive roots exist.

Theorem 0.4.5: Existence of Primitive Roots

The group $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic (i.e., a primitive root modulo m exists) if and only if $m \in \{1, 2, 4, p^k, 2p^k\}$ where p is an odd prime and $k \geq 1$.

The proof has two parts: showing cyclicity for odd prime powers (the hard direction), and showing non-cyclicity for all other moduli.

Proof:

The cases $m = 1$ and $m = 2$ are trivial ($(\mathbb{Z}/1\mathbb{Z})^\times$ and $(\mathbb{Z}/2\mathbb{Z})^\times$ are both trivial). The case $m = 4$ is checked directly: $(\mathbb{Z}/4\mathbb{Z})^\times = \{1, 3\}$ is cyclic of order 2, with primitive root 3.

Step 1: $(\mathbb{Z}/p\mathbb{Z})^\times$ is cyclic for p an odd prime. The group $(\mathbb{Z}/p\mathbb{Z})^\times$ has order $p-1$. For each divisor d of $p-1$, let $\psi(d)$ denote the number of elements of order d . Every element has some order dividing $p-1$, so $\sum_{d|(p-1)} \psi(d) = p-1$. By proposition 0.4.2(3), $\psi(d) \in \{0, \varphi(d)\}$ for each d . Since $\sum_{d|(p-1)} \varphi(d) = p-1$ (by proposition 0.3.16), and $\psi(d) \leq \varphi(d)$ for all d , the only possibility is $\psi(d) = \varphi(d)$ for every d . In particular, $\psi(p-1) = \varphi(p-1) \geq 1$, so an element of order $p-1$ exists.

Step 2: $(\mathbb{Z}/p^k\mathbb{Z})^\times$ is cyclic for p odd and $k \geq 2$. Let g be a primitive root modulo p . The group $(\mathbb{Z}/p^k\mathbb{Z})^\times$ has order $\varphi(p^k) = p^{k-1}(p-1)$. The reduction map $(\mathbb{Z}/p^k\mathbb{Z})^\times \rightarrow (\mathbb{Z}/p\mathbb{Z})^\times$ is surjective with kernel of order p^{k-1} , so $\text{ord}_{p^k}(g) = p^j(p-1)$ for some $0 \leq j \leq k-1$. It suffices to show $j = k-1$.

The key observation is: if $h \equiv 1 + tp \pmod{p^2}$ with $p \nmid t$, then $\text{ord}_{p^k}(h) = p^{k-1}$. To see this, use induction. The binomial theorem gives

$$h^p = (1 + tp)^p = 1 + tp^2 + \binom{p}{2} t^2 p^2 + \dots \equiv 1 + t' p^2 \pmod{p^3}$$

where $t' \equiv t \pmod{p}$ (since $p \mid \binom{p}{2}$ for $p \geq 3$), so $p \nmid t'$. Inductively, $h^{p^{j-1}} \equiv 1 + t_j p^j \pmod{p^{j+1}}$ with $p \nmid t_j$ for each j . In particular, $h^{p^{k-2}} \not\equiv 1 \pmod{p^k}$ while $h^{p^{k-1}} \equiv 1 \pmod{p^k}$, giving

$$\text{ord}_{p^k}(h) = p^{k-1}.$$

Applying this to $h = g^{p-1}$: if $g^{p-1} \not\equiv 1 \pmod{p^2}$, write $g^{p-1} = 1 + tp$ with $p \nmid t$, and the claim gives $\text{ord}_{p^k}(g^{p-1}) = p^{k-1}$. Then $\text{ord}_{p^k}(g) = p^{k-1}(p-1) = \varphi(p^k)$, so g is a primitive root modulo p^k . If instead $g^{p-1} \equiv 1 \pmod{p^2}$, replace g by $g' = g + p$ (which reduces to g modulo p , hence is still a primitive root modulo p). The binomial expansion gives $(g')^{p-1} \equiv g^{p-1} + (p-1)g^{p-2}p \equiv 1 - g^{p-2}p \pmod{p^2}$. Since $p \nmid g^{p-2}$, this shows $(g')^{p-1} \not\equiv 1 \pmod{p^2}$, so the argument applies to g' .

Step 3: $(\mathbb{Z}/2p^k\mathbb{Z})^\times$ is cyclic. By the Chinese Remainder Theorem (theorem 0.3.7), $(\mathbb{Z}/2p^k\mathbb{Z})^\times \cong (\mathbb{Z}/2\mathbb{Z})^\times \times (\mathbb{Z}/p^k\mathbb{Z})^\times \cong \{1\} \times (\mathbb{Z}/p^k\mathbb{Z})^\times$, which is cyclic.

Step 4: Non-cyclicity for other moduli. The remaining cases are handled by proposition 0.4.8 and proposition 0.4.9 below: $(\mathbb{Z}/2^k\mathbb{Z})^\times$ is not cyclic for $k \geq 3$, and the CRT decomposition for m with two or more distinct odd prime factors always produces a non-cyclic product (since each factor $(\mathbb{Z}/p_i^{a_i}\mathbb{Z})^\times$ has even order).

The proof that $(\mathbb{Z}/p\mathbb{Z})^\times$ is cyclic uses a counting argument that avoids constructing a primitive root explicitly. Finding a primitive root modulo a given prime is a computational problem with no known polynomial-time deterministic algorithm, though efficient probabilistic methods exist.

Example 0.4.6: Primitive Roots for Small Primes

1. Modulo 7: $(\mathbb{Z}/7\mathbb{Z})^\times$ has order 6. Testing $g = 3$: the powers $3^1 = 3, 3^2 = 2, 3^3 = 6, 3^4 = 4, 3^5 = 5, 3^6 = 1$ exhaust all of $(\mathbb{Z}/7\mathbb{Z})^\times$, so 3 is a primitive root. The other primitive roots are the elements of order 6: by proposition 0.4.2(2), these are 3^j with $\gcd(j, 6) = 1$, giving $3^1 = 3$ and $3^5 = 5$. Hence $(\mathbb{Z}/7\mathbb{Z})^\times$ has $\varphi(6) = 2$ primitive roots.
2. Modulo 11: $\varphi(11) = 10$. Testing $g = 2$: $2^1 = 2, 2^2 = 4, 2^3 = 8, 2^4 = 5, 2^5 = 10, 2^6 = 9, 2^7 = 7, 2^8 = 3, 2^9 = 6, 2^{10} = 1$. So 2 is a primitive root modulo 11. There are $\varphi(10) = 4$ primitive roots: $2, 2^3 = 8, 2^7 = 7, 2^9 = 6$.

A fixed integer $a > 1$ that is not a perfect square is a primitive root modulo infinitely many primes—this is a conjecture of Artin (1927), still open in general. Hooley proved in 1967 that the conjecture follows from the Generalized Riemann Hypothesis. For instance, 2 is conjectured to be a primitive root modulo a positive proportion (approximately 37.4%) of all primes. The unconditional result closest to Artin's conjecture, due to Gupta-Murty and Heath-Brown, shows that at least one of 2, 3, or 5 is a primitive root modulo infinitely many primes.

Primitive roots also arise in a familiar context: the decimal expansion of $1/p$ for a prime $p \neq 2, 5$ has period exactly $\text{ord}_p(10)$. When 10 is a primitive root modulo p , the period is $p-1$, the maximum possible. For example, $1/7 = 0.\overline{142857}$ has period 6 = $\varphi(7)$ because $\text{ord}_7(10) = 6$ (i.e., 10 is a primitive root modulo 7), while $1/13 = 0.\overline{076923}$ also has maximal period 12 = $\varphi(13)$. On the other hand, $1/11 = 0.\overline{09}$ has period 2 because $\text{ord}_{11}(10) = 2$.

Example 0.4.7: Lifting a Primitive Root from p to p^2

Take $p = 7$ and $g = 3$, a primitive root modulo 7. Is 3 a primitive root modulo 49? The group $(\mathbb{Z}/49\mathbb{Z})^\times$ has order $\varphi(49) = 42 = 6 \cdot 7$, so the question reduces to checking whether $3^6 \equiv 1 \pmod{49}$. Computing: $3^6 = 729 = 14 \cdot 49 + 43$, so $3^6 \equiv 43 \equiv 1 + 42 \equiv 1 + 6 \cdot 7 \pmod{49}$. Since $3^6 \not\equiv 1 \pmod{49}$ and the coefficient 6 satisfies $7 \nmid 6$, the lifting argument from Step 2 applies: 3 is a primitive root modulo 49, and modulo 7^k for all $k \geq 1$.

On the other hand, take $p = 29$ and $g = 14$, a primitive root modulo 29. Computing $14^{28} \pmod{29^2}$: since $14^{28} \equiv 1 \pmod{29}$ (Fermat), write $14^{28} = 1 + 29t$. If $29 \nmid t$, then 14 does not lift, and one must replace it by $14 + 29 = 43$. In practice, such “exceptional” primitive roots are rare; for most primes p and most primitive roots g modulo p , the condition $g^{p-1} \not\equiv 1 \pmod{p^2}$ holds.

For moduli that do *not* admit primitive roots, the structure of $(\mathbb{Z}/m\mathbb{Z})^\times$ is determined by the following two results.

Proposition 0.4.8: Structure of $(\mathbb{Z}/2^k\mathbb{Z})^\times$

For $k \geq 3$: $(\mathbb{Z}/2^k\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2^{k-2}\mathbb{Z}$, generated by -1 and 5.

Proof:

The group $(\mathbb{Z}/2^k\mathbb{Z})^\times$ has order $\varphi(2^k) = 2^{k-1}$. The element -1 has order 2, and 5 has order 2^{k-2} : to verify the latter, write $5 = 1 + 4$ and use induction. If $5^{2^{j-1}} \equiv 1 + c_j \cdot 2^{j+1} \pmod{2^{j+2}}$ with c_j odd, then

$$5^{2^j} = (5^{2^{j-1}})^2 \equiv 1 + c_j 2^{j+2} + c_j^2 2^{2j+2} \equiv 1 + c_j 2^{j+2} \pmod{2^{j+3}}$$

with c_j still odd (since only the linear term contributes modulo 2^{j+3} when $2j+2 \geq j+3$, i.e., $j \geq 1$). The base case $j = 0$: $5 = 1 + 1 \cdot 2^2$ with $c_0 = 1$. Inductively, $5^{2^{k-3}} \equiv 1 + c_{k-3} \cdot 2^{k-1} \not\equiv 1 \pmod{2^k}$, while $5^{2^{k-2}} \equiv 1 \pmod{2^k}$. Hence $\text{ord}_{2^k}(5) = 2^{k-2}$.

Since $5^j \equiv 1 \pmod{4}$ for all j (as $5 \equiv 1 \pmod{4}$) while $-1 \equiv 3 \pmod{4}$, no power of 5 equals -1 . Hence $\langle -1 \rangle \cap \langle 5 \rangle = \{1\}$, and the product $\langle -1 \rangle \times \langle 5 \rangle$ has order $2 \cdot 2^{k-2} = 2^{k-1} = |(\mathbb{Z}/2^k\mathbb{Z})^\times|$, giving the isomorphism.

Proposition 0.4.9: Structure of $(\mathbb{Z}/m\mathbb{Z})^\times$ for General m

For $m = 2^a p_1^{a_1} \cdots p_r^{a_r}$ with $p_1, \dots, p_r$ distinct odd primes:

$$(\mathbb{Z}/m\mathbb{Z})^\times \cong (\mathbb{Z}/2^a\mathbb{Z})^\times \times \prod_{i=1}^r (\mathbb{Z}/p_i^{a_i}\mathbb{Z})^\times$$

Each odd prime-power factor is cyclic of order $p_i^{a_i-1}(p_i - 1)$. A primitive root modulo m exists if and only if the product is cyclic, which happens exactly when $m \in \{1, 2, 4, p^k, 2p^k\}$.

Proof:

The decomposition is the CRT applied to unit groups, as in corollary 0.3.9. A direct product of cyclic groups is cyclic if and only if their orders are pairwise coprime (see [13]). If m has two distinct odd prime factors p_1, p_2 , then $\varphi(p_1^{a_1})$ and $\varphi(p_2^{a_2})$ are both even, so the product is not cyclic. Similarly, if $m = 2^a p^k$ with $a \geq 2$, then both $\varphi(2^a)$ and $\varphi(p^k)$ are even, ruling out cyclicity. The surviving cases are $m \in \{1, 2, 4, p^k, 2p^k\}$.

Example 0.4.10: Non-Cyclic Unit Groups

1. $(\mathbb{Z}/8\mathbb{Z})^\times = \{1, 3, 5, 7\}$. Every element satisfies $a^2 \equiv 1 \pmod{8}$, so the group has exponent 2 and $(\mathbb{Z}/8\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, the Klein four-group. The generators of the two $\mathbb{Z}/2\mathbb{Z}$ factors are $-1 = 7$ and 5 : indeed $\{(\pm 1)^a \cdot 5^b : a \in \{0, 1\}, b \in \{0, 1\}\} = \{1, 7, 5, 3\}$ gives all four elements. No primitive root exists.
2. $(\mathbb{Z}/15\mathbb{Z})^\times \cong (\mathbb{Z}/3\mathbb{Z})^\times \times (\mathbb{Z}/5\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$. This group has order 8 but exponent $\text{lcm}(2, 4) = 4$, so it is not cyclic. The maximum order 4 is achieved by elements like 2 (since $2^1 = 2, 2^2 = 4, 2^3 = 8, 2^4 = 16 \equiv 1 \pmod{15}$).
3. $(\mathbb{Z}/24\mathbb{Z})^\times \cong (\mathbb{Z}/8\mathbb{Z})^\times \times (\mathbb{Z}/3\mathbb{Z})^\times \cong (\mathbb{Z}/2\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z} \cong (\mathbb{Z}/2\mathbb{Z})^3$. This has order 8 and exponent 2: every element is its own inverse. There are $2^3 - 1 = 7$ elements of order 2, confirming that the group is as far from cyclic as possible.

The *exponent* of $(\mathbb{Z}/m\mathbb{Z})^\times$ —the least common multiple of the orders of all its elements—is captured by the Carmichael function.

Definition 0.4.11: Carmichael Function

The *Carmichael function* $\lambda(m)$ is the exponent of $(\mathbb{Z}/m\mathbb{Z})^\times$: the smallest positive integer e such that $a^e \equiv 1 \pmod{m}$ for all a with $\gcd(a, m) = 1$.

By the structure theorem, $\lambda(m) = \text{lcm}(\lambda(p_1^{a_1}), \dots, \lambda(p_r^{a_r}), \lambda(2^a))$, where $\lambda(p^a) = \varphi(p^a) = p^{a-1}(p-1)$ for odd primes, $\lambda(2) = 1$, $\lambda(4) = 2$, and $\lambda(2^a) = 2^{a-2}$ for $a \geq 3$. When $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic, $\lambda(m) = \varphi(m)$; otherwise $\lambda(m)$ is a proper divisor of $\varphi(m)$.

The Carmichael function refines Euler's theorem: $a^{\lambda(m)} \equiv 1 \pmod{m}$ for all a coprime to m , and $\lambda(m)$ is the *smallest* universal exponent with this property. When $\lambda(m) < \varphi(m)$, Euler's theorem with the exponent $\varphi(m)$ is valid but not sharp.

Example 0.4.12: Carmichael Function Computations

1. $\lambda(7) = \varphi(7) = 6$, since $(\mathbb{Z}/7\mathbb{Z})^\times$ is cyclic. The primitive root 3 achieves this order.
2. $\lambda(8) = 2$, since every element of $(\mathbb{Z}/8\mathbb{Z})^\times$ squares to 1, while $\varphi(8) = 4$. Euler's theorem gives $a^4 \equiv 1 \pmod{8}$, but the Carmichael function shows $a^2 \equiv 1$ already suffices.
3. $\lambda(15) = \text{lcm}(\lambda(3), \lambda(5)) = \text{lcm}(2, 4) = 4$, while $\varphi(15) = 8$. The element 2 has order exactly 4.
4. $\lambda(100) = \text{lcm}(\lambda(4), \lambda(25)) = \text{lcm}(2, 20) = 20$, while $\varphi(100) = 40$. Every element coprime to 100 satisfies $a^{20} \equiv 1 \pmod{100}$.

The structure theorem has a direct consequence for character theory. The dual group $(\widehat{\mathbb{Z}/m\mathbb{Z}})^\times$ of group homomorphisms $(\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ is (non-canonically) isomorphic to $(\mathbb{Z}/m\mathbb{Z})^\times$ itself (see [13]). When $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic, every character is determined by the image of a single generator; when it is not, specifying a character requires choosing the image of each invariant factor independently. The Dirichlet characters of modulus m (chapter 7) are the extensions of these group characters to all of $\mathbb{Z}$, and the invariant factor decomposition determines how many characters of each order exist.

Example 0.4.13: Orders and Characters Modulo 12

The group $(\mathbb{Z}/12\mathbb{Z})^\times = \{1, 5, 7, 11\} \cong (\mathbb{Z}/4\mathbb{Z})^\times \times (\mathbb{Z}/3\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is the Klein four-group. Every non-identity element has order 2:

$$5^2 = 25 \equiv 1, \quad 7^2 = 49 \equiv 1, \quad 11^2 = 121 \equiv 1 \pmod{12}$$

The dual group $(\widehat{\mathbb{Z}/12\mathbb{Z}})^\times \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ has four characters $\chi_0, \chi_1, \chi_2, \chi_3$, each taking values in $\{\pm 1\}$. The trivial character sends every unit to 1. The three nontrivial characters correspond to the three subgroups of index 2 in $(\mathbb{Z}/12\mathbb{Z})^\times$:

$$\chi_1(n) = \begin{cases} 1 & n \equiv 1, 5 \\ -1 & n \equiv 7, 11 \end{cases}, \quad \chi_2(n) = \begin{cases} 1 & n \equiv 1, 7 \\ -1 & n \equiv 5, 11 \end{cases}, \quad \chi_3(n) = \begin{cases} 1 & n \equiv 1, 11 \\ -1 & n \equiv 5, 7 \end{cases}$$

modulo 12. Each nontrivial character has order 2 in the dual group. These Dirichlet characters of modulus 12 control the distribution of primes in residue classes modulo 12, as developed in chapter 7.

This example illustrates a general principle: the number of Dirichlet characters of modulus m equals $\varphi(m) = |(\mathbb{Z}/m\mathbb{Z})^\times|$, and the order structure of the character group mirrors that of $(\mathbb{Z}/m\mathbb{Z})^\times$. When the group is cyclic of order n , there is a character of every order dividing n ; when it decomposes as a product $\mathbb{Z}/d_1\mathbb{Z} \times \cdots \times \mathbb{Z}/d_r\mathbb{Z}$, the characters decompose correspondingly.

0.5 Quadratic Residues and Reciprocity

The previous section determined when $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic. This section addresses a finer structural question: which elements of $(\mathbb{Z}/p\mathbb{Z})^\times$ are perfect squares? The answer is governed by the Legendre symbol, and the relationship between squareness modulo different primes is the content of the law of quadratic reciprocity—one of the central results of elementary number theory. Quadratic reciprocity has far-reaching generalizations: Artin reciprocity in class field theory ([9]) subsumes it as a special case, and the Langlands program seeks non-abelian analogues. In this book, quadratic characters reappear in the study of Dirichlet L -functions (chapter 7) and the factorization of Dedekind zeta functions of quadratic fields (chapter 8).

Definition 0.5.1: Quadratic Residue

Let p be an odd prime. An integer a with $p \nmid a$ is a *quadratic residue* modulo p if there exists $x \in \mathbb{Z}$ with $x^2 \equiv a \pmod{p}$. Otherwise, a is a *quadratic non-residue* modulo p .

Since the squaring map $x \mapsto x^2$ on $(\mathbb{Z}/p\mathbb{Z})^\times$ sends x and $-x$ to the same value, and $x \neq -x$ in $(\mathbb{Z}/p\mathbb{Z})^\times$ for p odd (as $2x \neq 0$), the image of squaring has exactly $(p-1)/2$ elements. Hence exactly

half the nonzero residues modulo p are quadratic residues, and half are non-residues.

Example 0.5.2: Quadratic Residues Modulo Small Primes

1. Modulo 7: the squares are $1^2 = 1$, $2^2 = 4$, $3^2 = 2$, so the quadratic residues are $\{1, 2, 4\}$ and the non-residues are $\{3, 5, 6\}$.
2. Modulo 13: the squares are $1^2 = 1$, $2^2 = 4$, $3^2 = 9$, $4^2 = 3$, $5^2 = 12$, $6^2 = 10$, giving quadratic residues $\{1, 3, 4, 9, 10, 12\}$ and non-residues $\{2, 5, 6, 7, 8, 11\}$.

The Legendre symbol packages the quadratic residue information into a multiplicative function.

Definition 0.5.3: Legendre Symbol

For an odd prime p and an integer a , the *Legendre symbol* is

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & \text{if } a \text{ is a quadratic residue modulo } p, \\ -1 & \text{if } a \text{ is a quadratic non-residue modulo } p, \\ 0 & \text{if } p|a. \end{cases}$$

The Legendre symbol $\left(\frac{\cdot}{p}\right) : \mathbb{Z} \rightarrow \{-1, 0, 1\}$ is a Dirichlet character of modulus p (as defined in chapter 7): it is totally multiplicative and p -periodic. This is a consequence of the following criterion.

Proposition 0.5.4: Euler's Criterion

For an odd prime p and a with $p \nmid a$:

$$\left(\frac{a}{p}\right) \equiv a^{(p-1)/2} \pmod{p}$$

Proof:

The group $(\mathbb{Z}/p\mathbb{Z})^\times$ is cyclic of order $p-1$ (theorem 0.4.5). Let g be a primitive root and write $a \equiv g^k \pmod{p}$. Then $a^{(p-1)/2} \equiv g^{k(p-1)/2} \pmod{p}$. Since g has order $p-1$, the value $g^{k(p-1)/2}$ is 1 if $(p-1) \mid k(p-1)/2$, i.e., if k is even, and $g^{(p-1)/2} = -1$ if k is odd (as $g^{(p-1)/2}$ is a root of $x^2 - 1$ other than 1, hence equals -1).

Now, $a = g^k$ is a quadratic residue if and only if $a = (g^{k/2})^2$ for some integer $k/2$, i.e., if and only if k is even. Hence $\left(\frac{a}{p}\right) = 1$ when k is even and -1 when k is odd, matching $a^{(p-1)/2} \pmod{p}$.

Euler's criterion immediately implies the total multiplicativity of the Legendre symbol: $\left(\frac{ab}{p}\right) = (ab)^{(p-1)/2} \equiv a^{(p-1)/2} \cdot b^{(p-1)/2} \equiv \left(\frac{a}{p}\right) \left(\frac{b}{p}\right) \pmod{p}$. Since both sides take values in $\{-1, 1\}$ and agree modulo $p > 2$, they are equal as integers. This multiplicativity reduces the computation of $\left(\frac{a}{p}\right)$ for general a to the case of prime arguments.

The Legendre symbol $\left(\frac{\cdot}{p}\right)$ is the unique nontrivial quadratic character of $(\mathbb{Z}/p\mathbb{Z})^\times$: it is the group homomorphism $(\mathbb{Z}/p\mathbb{Z})^\times \rightarrow \{\pm 1\}$ whose kernel is the subgroup of quadratic residues, an index-2 subgroup. In the language of chapter 7, it is the unique primitive Dirichlet character of modulus p and order 2. The Dirichlet L -function $L(s, \left(\frac{\cdot}{p}\right))$ controls the distribution of quadratic residues and non-residues among the primes, and its non-vanishing at $s = 1$ is one of the key inputs for Dirichlet's theorem on primes in arithmetic progressions.

The values of $\left(\frac{-1}{p}\right)$ and $\left(\frac{2}{p}\right)$ can be determined directly from Euler's criterion.

Proposition 0.5.5: First Supplement: -1 as a Quadratic Residue

For an odd prime p :

$$\left(\frac{-1}{p}\right) = (-1)^{(p-1)/2} = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4}, \\ -1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Proof:

By Euler's criterion, $\left(\frac{-1}{p}\right) \equiv (-1)^{(p-1)/2} \pmod{p}$. Since $(p-1)/2$ is even when $p \equiv 1 \pmod{4}$ and odd when $p \equiv 3 \pmod{4}$, the result follows.

The first supplement has a clean interpretation: -1 is a square modulo p precisely when $(\mathbb{Z}/p\mathbb{Z})^\times$ has an element of order 4, which happens if and only if $4 \mid (p-1)$.

Proposition 0.5.6: Second Supplement: 2 as a Quadratic Residue

For an odd prime p :

$$\left(\frac{2}{p}\right) = (-1)^{(p^2-1)/8} = \begin{cases} 1 & \text{if } p \equiv \pm 1 \pmod{8}, \\ -1 & \text{if } p \equiv \pm 3 \pmod{8}. \end{cases}$$

Proof:

The exponent $(p^2-1)/8 = (p-1)(p+1)/8$. When $p \equiv 1 \pmod{8}$, both $(p-1)/2$ and $(p+1)/2$ are even, so the product is divisible by 4 and $(p^2-1)/8$ is even. When $p \equiv 3 \pmod{8}$, then $(p-1)/2 = 1 + 4k$ for some k , and $(p+1)/4 = 1 + 2k$ is odd; working through the cases gives $(p^2-1)/8$ odd. The remaining cases $p \equiv 5, 7 \pmod{8}$ follow by symmetry. Alternatively, this can be deduced from Gauss's lemma (lemma 0.5.7 below) applied to $a = 2$.

Both supplements can also be proved using *Gauss sums*: the quantity $\tau(\chi) = \sum_{n=1}^{p-1} \left(\frac{n}{p}\right) e^{2\pi i n/p}$, which satisfies $\tau(\chi)^2 = (-1)^{(p-1)/2} p$. Gauss sums are the Fourier-analytic tool that connects quadratic residues to exponential sums, and they play a central role in the functional equations of Dirichlet L -functions (chapter 7).

The central tool for proving quadratic reciprocity is the following counting lemma.

Lemma 0.5.7: Gauss's Lemma

Let p be an odd prime and a an integer with $p \nmid a$. For each $k \in \{1, 2, \dots, (p-1)/2\}$, let r_k be the representative of $ka \pmod p$ in $\{-(p-1)/2, \dots, -1, 1, \dots, (p-1)/2\}$, and let n be the number of indices k for which $r_k < 0$. Then

$$\left(\frac{a}{p}\right) = (-1)^n$$

Proof:

The values $|r_1|, |r_2|, \dots, |r_{(p-1)/2}|$ are a permutation of $\{1, 2, \dots, (p-1)/2\}$: they are $(p-1)/2$ distinct elements of this set (distinct because $r_i = \pm r_j$ with $i \neq j$ would give $a(i \pm j) \equiv 0 \pmod p$, contradicting $p \nmid a$ and $0 < i \pm j < p$). Taking the product:

$$\prod_{k=1}^{(p-1)/2} r_k = (-1)^n \prod_{k=1}^{(p-1)/2} |r_k| = (-1)^n \prod_{k=1}^{(p-1)/2} k = (-1)^n \left(\frac{p-1}{2}\right)!$$

On the other hand, $r_k \equiv ka \pmod p$, so $\prod r_k \equiv a^{(p-1)/2} \prod_{k=1}^{(p-1)/2} k \pmod p$. Comparing with the line above and cancelling $((p-1)/2)!$ (which is nonzero modulo p) gives $a^{(p-1)/2} \equiv (-1)^n \pmod p$. The result follows from Euler's criterion.

Gauss's lemma converts the algebraic question “is a a square modulo p ?” into the combinatorial question “how many of the reductions $a, 2a, \dots, \frac{p-1}{2}a$ modulo p land in the upper half $\{(p+1)/2, \dots, p-1\}$?” This geometric reformulation is the key to Eisenstein's proof of quadratic reciprocity.

Example 0.5.8: Gauss's Lemma in Action

Take $a = 5$ and $p = 11$. The set $\{ka \pmod{11} : k = 1, \dots, 5\}$ is $\{5, 10, 4, 9, 3\}$. Representing these in $\{-5, \dots, -1, 1, \dots, 5\}$: $5 \mapsto 5, 10 \mapsto -1, 4 \mapsto 4, 9 \mapsto -2, 3 \mapsto 3$. The negative values are $\{-1, -2\}$, so $n = 2$ and $\left(\frac{5}{11}\right) = (-1)^2 = 1$. Indeed, $4^2 = 16 \equiv 5 \pmod{11}$, confirming that 5 is a quadratic residue.

Theorem 0.5.9: Quadratic Reciprocity

Let p and q be distinct odd primes. Then

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}$$

Equivalently: $\left(\frac{p}{q}\right) = \left(\frac{q}{p}\right)$ unless $p \equiv q \equiv 3 \pmod{4}$, in which case $\left(\frac{p}{q}\right) = -\left(\frac{q}{p}\right)$.

Quadratic reciprocity relates the solvability of $x^2 \equiv p \pmod q$ to that of $x^2 \equiv q \pmod p$ —two problems that, a priori, have nothing to do with each other. The sign $(-1)^{(p-1)(q-1)/4}$ is $+1$ unless both primes are congruent to 3 modulo 4, in which case the Legendre symbols “flip.” This single sign change is the entire content of the reciprocity law.

Proof:

By Gauss's lemma, $\left(\frac{q}{p}\right) = (-1)^{n_p}$ where $n_p = \#\{k \in \{1, \dots, (p-1)/2\} \mid kq \bmod p > p/2\}$. Since $kq \bmod p = kq - \lfloor kq/p \rfloor p$, the condition $kq \bmod p > p/2$ is equivalent to $kq - \lfloor kq/p \rfloor p > p/2$, which rearranges to $\lfloor kq/p \rfloor < kq/p - 1/2$. Counting more carefully:

$$n_p = \sum_{k=1}^{(p-1)/2} \left\lfloor \frac{kq}{p} \right\rfloor - 2 \sum_{k=1}^{(p-1)/2} \left\lfloor \frac{kq}{2p} \right\rfloor$$

Eisenstein's approach avoids this bookkeeping by counting lattice points. Consider the rectangle $R = (0, p/2) \times (0, q/2)$ in $\mathbb{R}^2$. The number of lattice points $(x, y) \in \mathbb{Z}^2$ with $0 < x < p/2$ and $0 < y < qx/p$ is $\sum_{x=1}^{(p-1)/2} \lfloor qx/p \rfloor$. Similarly, the number with $0 < y < q/2$ and $0 < x < py/q$ is $\sum_{y=1}^{(q-1)/2} \lfloor py/q \rfloor$. Since p and q are distinct odd primes, no lattice point (x, y) with $0 < x < p/2$ and $0 < y < q/2$ lies on the diagonal $qx = py$ (as this would require $p \mid x$ or $q \mid y$, contradicting the bounds). Hence every lattice point in $R \cap \mathbb{Z}^2$ lies strictly above or strictly below the diagonal, giving:

$$\sum_{x=1}^{(p-1)/2} \left\lfloor \frac{qx}{p} \right\rfloor + \sum_{y=1}^{(q-1)/2} \left\lfloor \frac{py}{q} \right\rfloor = \frac{p-1}{2} \cdot \frac{q-1}{2}$$

The left-hand side counts all lattice points in the open rectangle R : those below the diagonal contribute the first sum, and those above contribute the second. The total number of such points is $\frac{p-1}{2} \cdot \frac{q-1}{2}$.

By Gauss's lemma, one can verify that $(-1)^{n_p} = (-1)^{\sum_{x=1}^{(p-1)/2} \lfloor qx/p \rfloor}$ (since the parity of n_p matches that of this sum). Hence:

$$\left(\frac{q}{p}\right) \left(\frac{p}{q}\right) = (-1)^{n_p + n_q} = (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}$$

as claimed.

The lattice-point argument is due to Eisenstein and is among the most geometric of the many known proofs of quadratic reciprocity (over 240 proofs are known). The rectangle R and its diagonal will reappear—in a much more general form—in the lattice-point counting arguments used to prove the analytic class number formula in chapter 8.

Example 0.5.10: Lattice Point Verification

Take $p = 5$ and $q = 7$. The rectangle $R = (0, 5/2) \times (0, 7/2)$ contains lattice points with $1 \leq x \leq 2$ and $1 \leq y \leq 3$. The diagonal is $y = (7/5)x$. Below the diagonal: $(1, 1)$ (since $1 < 7/5$); above the diagonal: $(1, 2)$, $(1, 3)$, $(2, 2)$, $(2, 3)$ (since $2 > 14/5$), and $(2, 1)$ lies below (since $1 < 14/5$). Counting: $\sum_{x=1}^2 \lfloor 7x/5 \rfloor = 1 + 2 = 3$ below, $\sum_{y=1}^3 \lfloor 5y/7 \rfloor = 0 + 1 + 2 = 3$ above, total $6 = 2 \cdot 3 = \frac{p-1}{2} \cdot \frac{q-1}{2}$. Since 6 is even, $\left(\frac{5}{7}\right) \left(\frac{7}{5}\right) = 1$. One checks: $\left(\frac{5}{7}\right) = 5^3 = 125 \equiv 6 \equiv -1 \pmod{7}$ and $\left(\frac{7}{5}\right) = 7^2 = 49 \equiv 4 \equiv -1 \pmod{5}$, confirming the product is 1.

Example 0.5.11: Using Quadratic Reciprocity

1. Is 21 a quadratic residue modulo 71? By multiplicativity, $\left(\frac{21}{71}\right) = \left(\frac{3}{71}\right) \left(\frac{7}{71}\right)$. For $\left(\frac{3}{71}\right)$: since $71 \equiv 3 \pmod{4}$ and $3 \equiv 3 \pmod{4}$, reciprocity gives $\left(\frac{3}{71}\right) = -\left(\frac{71}{3}\right) = -\left(\frac{2}{3}\right) = -(-1) = 1$. For $\left(\frac{7}{71}\right)$: since $71 \equiv 3 \pmod{4}$ and $7 \equiv 3 \pmod{4}$, reciprocity gives $\left(\frac{7}{71}\right) = -\left(\frac{71}{7}\right) = -\left(\frac{1}{7}\right) = -1$. Hence $\left(\frac{21}{71}\right) = 1 \cdot (-1) = -1$: the integer 21 is a non-residue modulo 71.
2. Is 5 a quadratic residue modulo 23? Since $5 \equiv 1 \pmod{4}$, reciprocity gives $\left(\frac{5}{23}\right) = \left(\frac{23}{5}\right) = \left(\frac{3}{5}\right)$. Since $5 \equiv 1 \pmod{4}$, applying reciprocity again: $\left(\frac{3}{5}\right) = \left(\frac{5}{3}\right) = \left(\frac{2}{3}\right) = -1$ (by the second supplement, since $3 \equiv 3 \pmod{8}$). Hence 5 is a non-residue modulo 23.

The Legendre symbol is defined only for prime moduli. The Jacobi symbol extends it to odd composite moduli and is the standard tool for efficient computation.

Definition 0.5.12: Jacobi Symbol

For an odd positive integer $n = p_1^{a_1} \cdots p_k^{a_k}$ (not necessarily prime) and any integer a , the *Jacobi symbol* is

$$\left(\frac{a}{n}\right) = \prod_{i=1}^k \left(\frac{a}{p_i}\right)^{a_i}$$

The Jacobi symbol inherits the multiplicativity properties of the Legendre symbol: $\left(\frac{ab}{n}\right) = \left(\frac{a}{n}\right) \left(\frac{b}{n}\right)$ and $\left(\frac{a}{mn}\right) = \left(\frac{a}{m}\right) \left(\frac{a}{n}\right)$. The reciprocity law, first supplement, and second supplement all extend to Jacobi symbols with identical formulas:

$$\left(\frac{m}{n}\right) \left(\frac{n}{m}\right) = (-1)^{\frac{m-1}{2} \cdot \frac{n-1}{2}}, \quad \left(\frac{-1}{n}\right) = (-1)^{(n-1)/2}, \quad \left(\frac{2}{n}\right) = (-1)^{(n^2-1)/8}$$

for odd coprime $m, n > 0$. These follow from the corresponding Legendre symbol identities by multiplying over the prime factorizations.

A caution: $\left(\frac{a}{n}\right) = 1$ does *not* imply that a is a quadratic residue modulo n when n is composite. For example, $\left(\frac{2}{15}\right) = \left(\frac{2}{3}\right) \left(\frac{2}{5}\right) = (-1)(-1) = 1$, but $x^2 \equiv 2 \pmod{15}$ has no solution (since $x^2 \equiv 2 \pmod{3}$ has no solution). The Jacobi symbol detects quadratic residuosity only modulo each prime factor individually.

The computational value of the Jacobi symbol lies in its reciprocity law: evaluating $\left(\frac{a}{n}\right)$ reduces to repeated application of reciprocity and the supplements, without factoring n . The resulting algorithm runs in $O(\log^2 n)$ time, analogous to the Euclidean algorithm for GCD.

Example 0.5.13: Computing a Jacobi Symbol

Evaluate $\left(\frac{1001}{9907}\right)$. Neither number need be prime. Since $9907 \equiv 3 \pmod{4}$ and $1001 \equiv 1 \pmod{4}$, reciprocity gives $\left(\frac{1001}{9907}\right) = \left(\frac{9907}{1001}\right)$. Now $9907 = 9 \cdot 1001 + 898$, so $\left(\frac{9907}{1001}\right) = \left(\frac{898}{1001}\right) = \left(\frac{2}{1001}\right) \left(\frac{449}{1001}\right)$. Since $1001 \equiv 1 \pmod{8}$, the second supplement gives $\left(\frac{2}{1001}\right) = 1$. For $\left(\frac{449}{1001}\right)$: since $449 \equiv 1 \pmod{4}$, reciprocity gives $\left(\frac{449}{1001}\right) = \left(\frac{1001}{449}\right) = \left(\frac{103}{449}\right)$ (as $1001 = 2 \cdot 449 + 103$). Since $103 \equiv 3 \pmod{4}$ and $449 \equiv 1 \pmod{4}$, reciprocity gives $\left(\frac{103}{449}\right) = \left(\frac{449}{103}\right) = \left(\frac{37}{103}\right)$. Since $37 \equiv 1 \pmod{4}$: $\left(\frac{37}{103}\right) = \left(\frac{103}{37}\right) = \left(\frac{29}{37}\right)$. Since $29 \equiv 1 \pmod{4}$: $\left(\frac{29}{37}\right) = \left(\frac{37}{29}\right) = \left(\frac{8}{29}\right) = \left(\frac{2}{29}\right)^3 = (-1)^3 = -1$ (since $29 \equiv 5 \pmod{8}$). Hence $\left(\frac{1001}{9907}\right) = -1$.

Quadratic reciprocity is the first in a hierarchy of reciprocity laws. *Cubic reciprocity* governs when a is a perfect cube modulo p and is naturally formulated over the Eisenstein integers $\mathbb{Z}[\zeta_3]$; *quartic reciprocity* governs fourth powers and lives over the Gaussian integers $\mathbb{Z}[i]$. These higher reciprocity laws were developed by Gauss, Jacobi, and Eisenstein in the early 19th century and are part of the motivation for studying number rings beyond $\mathbb{Z}$. The ultimate generalization—Artin reciprocity—unifies all abelian reciprocity laws into a single statement about the Galois group $\text{Gal}(K^{\text{ab}}/K)$ of the maximal abelian extension of a number field K , and is the subject of [9]. The quadratic case treated here, while elementary, already exhibits the key phenomenon: a local condition (squareness modulo p) is controlled by global arithmetic data (the residue classes of p and q modulo 4).

0.6 Asymptotic Notation

The results of analytic number theory are expressed in the language of asymptotics: statements about how functions behave as their argument grows without bound. The Prime Number Theorem, for instance, asserts that $\pi(x) \sim x/\log x$ —a precise claim about the ratio $\pi(x)/(x/\log x)$ tending to 1, not about the difference $\pi(x) - x/\log x$. This section collects the standard asymptotic notations and their basic properties, establishing the conventions used throughout the book.

Definition 0.6.1: Big- O and Little- o

Let f and g be functions defined on a neighborhood of ∞ (or of some limit point), with g eventually positive.

1. $f(x) = O(g(x))$ as $x \rightarrow \infty$ means there exist constants $C > 0$ and x_0 such that $|f(x)| \leq Cg(x)$ for all $x \geq x_0$. The notation $f \ll g$ is synonymous.
2. $f(x) = o(g(x))$ as $x \rightarrow \infty$ means $\lim_{x \rightarrow \infty} f(x)/g(x) = 0$, i.e., $|f(x)| \leq \varepsilon g(x)$ for any $\varepsilon > 0$ and all sufficiently large x .

The big- O notation asserts a bound up to an unspecified multiplicative constant; the little- o notation asserts that the bound can be made arbitrarily small. Every $o(g)$ is also $O(g)$, but not conversely: $x = O(x)$ but $x \neq o(x)$.

A common source of confusion is that the “=” in $f = O(g)$ is not symmetric: it means “ f belongs to the class of functions bounded by Cg ,” not that two specific quantities are equal. The notation is best read as “ f is at most of order g .” When multiple big- O terms appear in an equation, each is understood to represent a (possibly different) function satisfying the stated bound. For instance, $O(x) + O(x^2) = O(x^2)$ is a valid statement: any function bounded by Cx plus any function bounded by $C'x^2$ is bounded by $C''x^2$ for $C'' = C + C'$.

Definition 0.6.2: Vinogradov Notation

The notation $f \ll g$ means $f = O(g)$, and $f \gg g$ means $g = O(f)$. If both $f \ll g$ and $f \gg g$ hold, this is written $f \asymp g$ (read “ f is of the same order as g ”).

The Vinogradov notation $\ll$ is standard in analytic number theory and avoids some of the awkwardness of the big- O symbol in chains of inequalities. For example, $\sum_{n \leq x} d(n) \ll x \log x$ is cleaner than $\sum_{n \leq x} d(n) = O(x \log x)$ when embedded in a longer computation. Both notations are used freely in

this book; the choice in any given instance is dictated by readability.

A critical convention: the implied constant in $f \ll g$ or $f = O(g)$ may depend on parameters that are considered fixed. When the dependence matters, it is indicated by a subscript: $f \ll_\varepsilon g$ means the implied constant depends on ε , and $f = O_k(g)$ means it depends on k . Failing to track such dependencies is a common source of error in analytic arguments, particularly when a parameter later varies (e.g., when $\varepsilon \rightarrow 0$ or $k \rightarrow \infty$).

Definition 0.6.3: Asymptotic Equivalence

Two functions f and g , both eventually nonzero, are *asymptotically equivalent*, written $f(x) \sim g(x)$, if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$$

Equivalently, $f \sim g$ if and only if $f = g + o(g)$, i.e., $f(x) = g(x)(1 + o(1))$.

Asymptotic equivalence is the relation used to state the main theorems of analytic number theory. The Prime Number Theorem says $\pi(x) \sim x/\log x$; Mertens' theorem says $\prod_{p \leq x} (1 - 1/p) \sim e^{-\gamma}/\log x$. These are statements about the *leading-order* behavior of the function, ignoring lower-order corrections. A stronger statement specifies the error: $\pi(x) = \text{Li}(x) + O(x \exp(-c\sqrt{\log x}))$ gives both the leading term and a bound on the remainder.

The relation $\sim$ is an equivalence relation on the set of eventually positive functions: it is reflexive ($f \sim f$), symmetric ($f \sim g$ implies $g \sim f$), and transitive ($f \sim g$ and $g \sim h$ imply $f \sim h$). It is also compatible with multiplication ($f_1 \sim g_1$ and $f_2 \sim g_2$ imply $f_1 f_2 \sim g_1 g_2$) but *not* with addition: $f_1 \sim g_1$ and $f_2 \sim g_2$ do not imply $f_1 + f_2 \sim g_1 + g_2$, since cancellation can occur. For instance, $(x+1) \sim x$ and $(-x) \sim (-x)$, but $(x+1) + (-x) = 1 \not\sim 0 = x + (-x)$.

Two further notations occasionally appear in the literature.

Definition 0.6.4: Ω and Θ Notation

1. $f(x) = \Omega(g(x))$ means $f(x) \neq o(g(x))$, i.e., $|f(x)| \geq c g(x)$ for some $c > 0$ and infinitely many $x \rightarrow \infty$. This is the negation of $f = o(g)$: the function f is “at least of order g ” infinitely often.
2. $f(x) = \Theta(g(x))$ means $f \asymp g$, i.e., both $f = O(g)$ and $f = \Omega(g)$. This gives a tight two-sided bound: $c_1 g(x) \leq |f(x)| \leq c_2 g(x)$ for constants $0 < c_1 \leq c_2$ and all sufficiently large x .

The Ω notation is used in number theory to express *lower bounds* on oscillation. For example, the statement $\pi(x) - \text{Li}(x) = \Omega(x^{1/2} \log \log \log x / \log x)$ (a theorem of Littlewood) asserts that the error in the Prime Number Theorem is not just $O(x^{1/2})$ —it is at least as large as $x^{1/2} \log \log \log x / \log x$ infinitely often.

The following rules govern the manipulation of asymptotic expressions. They are used without comment throughout the book.

Proposition 0.6.5: Arithmetic of Asymptotic Notation

Let f, f_1, f_2, g, g_1, g_2 be functions with g, g_1, g_2 eventually positive. Then:

1. *Transitivity*: if $f = O(g)$ and $g = O(h)$, then $f = O(h)$.
2. *Sum rule*: $O(g_1) + O(g_2) = O(\max(g_1, g_2))$, and $O(g) + O(g) = O(g)$.
3. *Product rule*: $O(g_1) \cdot O(g_2) = O(g_1 g_2)$.
4. *Scalar rule*: for any constant $c \neq 0$, $c \cdot O(g) = O(g)$.
5. *Absorption*: if $g_1 = O(g_2)$, then $O(g_1) + O(g_2) = O(g_2)$.
6. *Little-o absorbs*: $o(g) + O(g) = O(g)$, and $o(g) \cdot O(h) = o(gh)$.

The same rules hold with $\ll$ in place of O , and analogous rules hold for o and $\sim$.

Proof:

All follow from the definition. For (1): $|f| \leq C_1 g$ and $g \leq C_2 h$ give $|f| \leq C_1 C_2 h$. For (2): if $|f_1| \leq C_1 g_1$ and $|f_2| \leq C_2 g_2$, then $|f_1 + f_2| \leq C_1 g_1 + C_2 g_2 \leq (C_1 + C_2) \max(g_1, g_2)$. The remaining parts are similar.

Asymptotic notation interacts well with summation and integration.

Proposition 0.6.6: Summation and Integration of O -Terms

1. If $f(n) = O(g(n))$ for all n , and g is non-negative, then $\sum_{n=1}^N f(n) = O\left(\sum_{n=1}^N g(n)\right)$.
2. If $f(t) = O(g(t))$ for $t \geq t_0$, and g is non-negative and integrable, then $\int_{t_0}^x f(t) dt = O\left(\int_{t_0}^x g(t) dt\right)$.

Proof:

For (1): $|f(n)| \leq Cg(n)$ implies $|\sum f(n)| \leq \sum |f(n)| \leq C \sum g(n)$. Part (2) is identical with integrals replacing sums.

However, O -terms cannot be freely moved through nonlinear operations. For instance, $e^{O(1)}$ is *not* $O(1)$ —it denotes a function bounded between e^{-C} and e^C for some constant C , which is $\Theta(1)$. And $O(\log x)$ inside an exponent, as in $x^{O(1)} = e^{O(\log x)}$, denotes a function bounded by a power of x but with unspecified exponent.

The following table of growth rates, ordered from slowest to fastest, is a useful reference for comparing asymptotic estimates. Each function in the list is o of the next:

$$1 \ll \log \log x \ll \log x \ll (\log x)^A \ll x^\varepsilon \ll x^a \ll x^b \ll e^{x^c} \ll e^x \ll x! \ll x^x$$

where $0 < \varepsilon < a < b < 1 \leq c$ and $A > 0$. The gaps between consecutive entries in this hierarchy reflect qualitative differences in growth behavior. For example, the gap between $(\log x)^A$ and x^ε is

the reason that “logarithmic savings” in analytic number theory (gaining a factor of $(\log x)^4$ over a trivial bound) are fundamentally different from “power savings” (gaining x^ε). The Prime Number Theorem provides logarithmic savings over the trivial bound $\pi(x) \leq x$; the Riemann Hypothesis would upgrade this to a power saving.

Example 0.6.7: Manipulating Asymptotic Expressions

Suppose one has established that $f(x) = x^2 + 3x \log x + O(x)$ and wishes to estimate $f(x)/x^2$. Using the rules:

$$\frac{f(x)}{x^2} = 1 + \frac{3 \log x}{x} + O\left(\frac{1}{x}\right) = 1 + O\left(\frac{\log x}{x}\right)$$

since $\frac{3 \log x}{x} = O\left(\frac{\log x}{x}\right)$ and $\frac{1}{x} = O\left(\frac{\log x}{x}\right)$ (because $1 \leq \log x$ for $x \geq e$). Hence $f(x) \sim x^2$. The intermediate expression $1 + \frac{3 \log x}{x} + O(1/x)$ is more precise: it gives the leading term 1 and the next correction $3 \log x/x$, which tends to 0 but more slowly than $1/x$.

A common pitfall in asymptotic arguments is the hidden dependence of implied constants on parameters. Consider the estimate $\sum_{n \leq x} n^{-s} = O_s(1)$ for fixed $\operatorname{Re}(s) > 1$. This is valid for each fixed s , but the implied constant blows up as $s \rightarrow 1^+$ (since the sum diverges at $s = 1$). Any argument that takes $s \rightarrow 1^+$ *after* using this estimate produces an incorrect bound. The safest practice is to make all parameter dependencies explicit at the point of first use, as in $\sum_{n \leq x} n^{-\sigma} \leq \frac{x^{1-\sigma}}{1-\sigma} + \zeta(\sigma)$ for $\sigma > 1$, which exhibits the blowup at $\sigma = 1$ explicitly through the factor $1/(1-\sigma)$.

Example 0.6.8: Standard Asymptotic Estimates

The following estimates appear repeatedly throughout the book.

1. *Harmonic series:* $\sum_{n=1}^N \frac{1}{n} = \log N + \gamma + O(1/N)$, where $\gamma = 0.5772\dots$ is the Euler-Mascheroni constant. The error $O(1/N)$ can be refined to an asymptotic expansion in powers of $1/N$ using the Euler-Maclaurin formula (chapter 1).
2. *Geometric series:* for $|r| < 1$, $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$, and the partial sum satisfies $\sum_{n=0}^N r^n = \frac{1}{1-r} + O(r^N)$.
3. *Logarithmic growth:* $\log x = o(x^\varepsilon)$ for every $\varepsilon > 0$. Equivalently, $\log x \ll_\varepsilon x^\varepsilon$. This slow growth of the logarithm is the reason that the Prime Number Theorem $\pi(x) \sim x/\log x$ says primes are “barely” infinite: they thin out, but only logarithmically.
4. *Stirling:* $\log(n!) = n \log n - n + \frac{1}{2} \log(2\pi n) + O(1/n)$. In cruder form, $n! \sim \sqrt{2\pi n} (n/e)^n$. This is the asymptotic tool for estimating binomial coefficients and factorials.
5. *Partial summation prototype:* $\sum_{n \leq x} \frac{1}{n^s} = \frac{x^{1-s}}{1-s} + \zeta(s) + O(x^{-\operatorname{Re}(s)})$ for $\operatorname{Re}(s) > 0$, $s \neq 1$. This is proved in chapter 2 using partial summation.

The interaction between O , o , and $\sim$ deserves emphasis, as it governs the precision of every asymptotic statement in the book. Saying $f = O(g)$ gives an upper bound but no information about the actual size of f : the function f could be much smaller than g , or even identically zero. Saying $f \sim g$ pins down the leading behavior precisely, but says nothing about the error $f - g$ beyond it being $o(g)$. Giving an explicit error term, as in $f = g + O(h)$ with $h = o(g)$, is the strongest type of asymptotic statement and is the form in which the main theorems (PNT, class number formula, Chebotarev) are most useful.

Example 0.6.9: Distinguishing Asymptotic Statements

Consider the three statements about $\pi(x)$:

1. $\pi(x) = O(x/\log x)$: there is a constant C such that $\pi(x) \leq Cx/\log x$ for large x . This is a consequence of Chebyshev's estimates (chapter 3) and is proved without complex analysis.
2. $\pi(x) \sim x/\log x$: the ratio $\pi(x)/(x/\log x) \rightarrow 1$. This is the Prime Number Theorem (chapter 5), and its proof requires the non-vanishing of $\zeta(s)$ on $\text{Re}(s) = 1$.
3. $\pi(x) = \text{Li}(x) + O(x \exp(-c\sqrt{\log x}))$: the error between $\pi(x)$ and $\text{Li}(x)$ is bounded by $x \exp(-c\sqrt{\log x})$, which is $o(x/(\log x)^n)$ for every n but *not* $O(x^{1-\varepsilon})$ for any $\varepsilon > 0$. This sharper form requires a quantitative zero-free region for $\zeta(s)$.

Each successive statement is strictly more informative than the previous one, and each requires strictly more analytic input to prove. This hierarchy—from order of magnitude, to leading term, to explicit error—recurs throughout the book.

The distinction between $\sim$ and explicit error terms has practical consequences. The two approximations $\pi(x) \sim x/\log x$ and $\pi(x) \sim \text{Li}(x) = \int_2^x dt/\log t$ are both valid forms of the Prime Number Theorem, since $\text{Li}(x) \sim x/\log x$. However, $\text{Li}(x)$ is a significantly better numerical approximation: at $x = 10^9$, the prime counting function satisfies $\pi(10^9) = 50,847,534$, while $10^9/\log(10^9) \approx 48,254,942$ (an error of about 5%) and $\text{Li}(10^9) \approx 50,849,235$ (an error of about 0.003%). The asymptotic relation $\sim$ treats both approximations as equally valid, but the explicit error terms distinguish them.

Example 0.6.10: Growth Rate Comparisons

1. $n^{100} = o(1.01^n)$: any fixed polynomial is eventually dominated by any exponential with base greater than 1. This follows from $\lim_{n \rightarrow \infty} n^{100}/1.01^n = 0$, proved by taking logarithms: $100 \log n - n \log 1.01 \rightarrow -\infty$.
2. $(\log n)^{1000} = o(n^{0.001})$: any fixed power of $\log n$ is eventually dominated by any positive power of n . At $n = 10^6$, however, $(\log n)^{1000} \approx (13.8)^{1000}$ vastly exceeds $n^{0.001} \approx 1.014$. The crossover point is astronomically large, illustrating that asymptotic statements say nothing about the rate of convergence.
3. $n! \gg (n/e)^n$: Stirling's approximation gives $n! \sim \sqrt{2\pi n} (n/e)^n$, so $n! \asymp n^{1/2} (n/e)^n$. The factor $\sqrt{2\pi n}$ is asymptotically negligible compared to $(n/e)^n$ but is needed for precise estimates of binomial coefficients.

Example 0.6.11: Summing with Asymptotic Notation

Estimate $\sum_{n=1}^N \sqrt{n}$. Since $\sqrt{n}$ is monotonically increasing, comparison with the integral gives

$$\int_1^N \sqrt{t} dt \leq \sum_{n=1}^N \sqrt{n} \leq \sqrt{N} + \int_1^N \sqrt{t} dt$$

Computing $\int_1^N \sqrt{t} dt = \frac{2}{3}N^{3/2} - \frac{2}{3}$, this yields

$$\sum_{n=1}^N \sqrt{n} = \frac{2}{3}N^{3/2} + O(N^{1/2})$$

The Euler-Maclaurin formula (chapter 1) refines this to $\sum_{n=1}^N \sqrt{n} = \frac{2}{3}N^{3/2} + \frac{1}{2}N^{1/2} + \frac{1}{24}N^{-1/2} + O(N^{-3/2})$, but the leading term $\frac{2}{3}N^{3/2}$ is often sufficient.

Finally, a convention on logarithms: throughout this book, $\log$ denotes the natural logarithm $\ln = \log_e$. This is standard in analytic number theory (and in most of mathematics outside of computer science). Iterated logarithms are written $\log \log x = \log(\log x)$, and occasionally $\log_k x$ denotes the k -fold iterated logarithm.

Example 0.6.12: Asymptotic Expansion of a Sum

Estimate $S(N) = \sum_{n=1}^N \frac{\log n}{n}$. Using the integral comparison $\sum_{n=1}^N f(n) = \int_1^N f(t) dt + O(f(N))$ for monotone f (a rough form of Abel summation, proved in chapter 1):

$$S(N) = \int_1^N \frac{\log t}{t} dt + O\left(\frac{\log N}{N}\right) = \frac{(\log N)^2}{2} + O\left(\frac{\log N}{N}\right)$$

This gives $S(N) \sim \frac{1}{2}(\log N)^2$. For $N = 10^6$: the exact value is $S(10^6) \approx 95.38$, while $\frac{1}{2}(\log 10^6)^2 = \frac{1}{2}(6 \log 10)^2 \approx 95.40$ —close agreement, as expected from the small error term $O(\log N/N) \approx 10^{-5}$.

The notations introduced in this section form the grammar in which the results of this book are stated. The reader should be comfortable manipulating O , o , $\sim$, and $\ll$ fluently before proceeding, as every chapter from this point forward uses them without further comment.

Exercise 0.6.1

1. Use the Euclidean algorithm to compute $\gcd(1001, 385)$, and find integers x, y such that $1001x + 385y = \gcd(1001, 385)$.
2. Find all integer solutions (x, y) to $91x + 65y = 13$.
3. Let p be a prime and n a positive integer. Prove that

$$v_p(n!) = \frac{n - s_p(n)}{p - 1}$$

where $s_p(n)$ denotes the sum of the digits of n in base p .

4. Find the smallest positive integer x satisfying $x \equiv 3 \pmod{7}$, $x \equiv 5 \pmod{11}$, and $x \equiv 2 \pmod{13}$.
5.
 1. Find the order of 2 modulo 31.
 2. Determine whether 2 is a primitive root modulo 31.
 3. List all primitive roots modulo 13.
6. Compute $\lambda(360)$, where λ is the Carmichael function. Find an element of $(\mathbb{Z}/360\mathbb{Z})^\times$ achieving this order.

7. Evaluate the following Legendre and Jacobi symbols:

1. $\left(\frac{-3}{17}\right)$
2. $\left(\frac{11}{41}\right)$
3. $\left(\frac{1009}{2027}\right)$ (both are prime)

8. Using Wilson's theorem, prove that if p is a prime with $p \equiv 1 \pmod{4}$, then $x = \left(\frac{p-1}{2}\right)!$ satisfies $x^2 \equiv -1 \pmod{p}$.

9. Let p be an odd prime. Prove that the product of all quadratic residues modulo p in $\{1, 2, \dots, p-1\}$ is congruent to $(-1)^{(p+1)/2} \pmod{p}$.

10. 1. Show that $\sum_{n=1}^N \frac{1}{n^2} = \frac{\pi^2}{6} + O\left(\frac{1}{N}\right)$.
 2. Show that $\sum_{p \leq N} \frac{1}{p^2} = O(1)$, where the sum is over primes.
 3. Prove that $\prod_{n=1}^N \left(1 + \frac{1}{n}\right) \sim \frac{N}{e^\gamma} e^{1/N} \dots$ is false, and find the correct asymptotic: $\prod_{n=1}^N \left(1 + \frac{1}{n}\right) = N + 1$.

11. Prove that for any fixed $m \geq 1$:

$$\frac{\#\{n \leq x : \gcd(n, m) = 1\}}{x} = \frac{\varphi(m)}{m} + O\left(\frac{1}{x}\right)$$

Interpret this as saying that the “probability” that a random integer is coprime to m is $\varphi(m)/m = \prod_{p|m} (1 - 1/p)$.

12. Let n be an odd positive integer with $n > 1$. Prove that $\left(\frac{-1}{n}\right) = (-1)^{(n-1)/2}$ for the Jacobi symbol, without assuming n is prime.

Arithmetic Functions and Summation Techniques

The preliminary chapter studied individual integers: their divisibility, their prime factorizations, their residue classes. Analytic number theory asks a different kind of question. Rather than examining a single integer n , it attaches a numerical invariant $f(n)$ to every positive integer and studies how these values behave in aggregate—how $\sum_{n \leq x} f(n)$ grows as $x \rightarrow \infty$, what the “average” value of f is, and how the values fluctuate around that average. This shift from the particular to the statistical is the defining move of the subject.

This chapter builds the toolkit for that program. The first two sections are algebraic: they introduce the arithmetic functions that encode number-theoretic data and the Dirichlet convolution that organizes them into a ring. The last three sections are analytic: they develop the summation techniques—the hyperbola method, Abel’s formula, and the Euler-Maclaurin formula—that convert discrete sums into continuous integrals amenable to the methods of analysis. Every later chapter in this book depends on this material.

1.1 Arithmetic Functions and Multiplicativity

The starting point is to formalize the idea of a “number-theoretic quantity attached to each positive integer.” The divisor count $d(n)$, the Euler totient $\varphi(n)$, and the von Mangoldt function $\Lambda(n)$ are all examples of the same type of object: a function from the positive integers to the complex numbers. The structure that makes these functions tractable is *multiplicativity*—the property that $f(mn) = f(m)f(n)$ whenever $\gcd(m, n) = 1$ —which reduces the study of f on all positive integers to its behavior at prime powers.

Definition 1.1.1: Arithmetic Function

An *arithmetic function* is a function $f : \mathbb{Z}_{>0} \rightarrow \mathbb{C}$.

No regularity conditions are imposed: an arithmetic function is simply a sequence of complex numbers indexed by the positive integers. The interest lies in the additional structure that specific arithmetic functions possess.

Definition 1.1.2: Multiplicative, Totally Multiplicative, and Additive Functions

An arithmetic function f is:

1. *Multiplicative* if $f(1) = 1$ and $f(mn) = f(m)f(n)$ whenever $\gcd(m, n) = 1$.
2. *Totally multiplicative* if $f(1) = 1$ and $f(mn) = f(m)f(n)$ for all $m, n \geq 1$.
3. *Additive* if $f(mn) = f(m) + f(n)$ whenever $\gcd(m, n) = 1$.
4. *Totally additive* if $f(mn) = f(m) + f(n)$ for all $m, n \geq 1$.

The condition $f(1) = 1$ for multiplicative functions is a normalization: setting $m = n = 1$ in $f(mn) = f(m)f(n)$ gives $f(1) = f(1)^2$, so $f(1) \in \{0, 1\}$, and the convention excludes the zero function. The logarithm function $\log : \mathbb{Z}_{>0} \rightarrow \mathbb{R}$ is totally additive.

The fundamental observation is that multiplicativity reduces global information to local information at each prime.

Proposition 1.1.3: Determination by Prime Powers

A multiplicative function f is completely determined by its values at prime powers $\{f(p^k) : p \text{ prime}, k \geq 1\}$. Specifically, for $n = p_1^{a_1} \cdots p_r^{a_r}$:

$$f(n) = \prod_{i=1}^r f(p_i^{a_i})$$

If f is totally multiplicative, then $f(p^k) = f(p)^k$, so f is determined by its values at primes.

Proof:

Since $\gcd(p_i^{a_i}, p_j^{a_j}) = 1$ for $i \neq j$, repeated application of the multiplicativity condition gives $f(n) = f(p_1^{a_1}) \cdots f(p_r^{a_r})$. For totally multiplicative f : $f(p^k) = f(p \cdot p^{k-1}) = f(p)f(p^{k-1})$, and induction gives $f(p^k) = f(p)^k$.

This reduction to prime powers is the arithmetic analogue of the Chinese Remainder Theorem (theorem 0.3.7), which decomposes $\mathbb{Z}/n\mathbb{Z}$ into a product of rings $\mathbb{Z}/p_i^{a_i}\mathbb{Z}$. In both cases, the global object splits into independent local pieces, one for each prime. This “local-to-global” decomposition is the structural principle that, in the language of Dirichlet series (chapter 2), becomes the Euler product.

The rest of this section introduces the arithmetic functions that appear most frequently throughout the book. Each is defined, its values at prime powers are recorded, and a small table of values is

given for orientation.

Definition 1.1.4: The Möbius Function

The *Möbius function* $\mu : \mathbb{Z}_{>0} \rightarrow \{-1, 0, 1\}$ is defined by

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1, \\ (-1)^k & \text{if } n = p_1 p_2 \cdots p_k \text{ is a product of } k \text{ distinct primes,} \\ 0 & \text{if } p^2 | n \text{ for some prime } p. \end{cases}$$

Equivalently, μ is the multiplicative function with $\mu(p) = -1$ and $\mu(p^k) = 0$ for $k \geq 2$.

The Möbius function detects squarefreeness: $\mu(n) \neq 0$ if and only if n is squarefree. Its deeper role as the identity element for Möbius inversion, or equivalently as the Dirichlet inverse of the constant function **1**, is the subject of the next section. The alternative sign is for the inclusion-exclusion principle: for a finite set with k elements, the inclusion-exclusion formula alternates signs $(-1)^k$, and μ does the same over the prime factors of n .

Definition 1.1.5: Divisor Functions

For $k \in \mathbb{C}$, the k -th *divisor function* is

$$\sigma_k(n) = \sum_{d|n} d^k$$

The two most important special cases are the *number-of-divisors function* $d(n) = \sigma_0(n) = \sum_{d|n} 1$ and the *sum-of-divisors function* $\sigma(n) = \sigma_1(n) = \sum_{d|n} d$.

Each σ_k is multiplicative (but not totally multiplicative for $k \neq 0$). At a prime power:

$$\sigma_k(p^a) = 1 + p^k + p^{2k} + \cdots + p^{ak} = \frac{p^{(a+1)k} - 1}{p^k - 1}$$

for $k \neq 0$, and $d(p^a) = a + 1$. The multiplicativity of σ_k follows from the fact that the divisors of mn (with $\gcd(m, n) = 1$) are exactly the products $d_1 d_2$ where $d_1 | m$ and $d_2 | n$, so the sum factors.

Definition 1.1.6: The Liouville Function

The *Liouville function* $\lambda : \mathbb{Z}_{>0} \rightarrow \{-1, 1\}$ is defined by $\lambda(n) = (-1)^{\Omega(n)}$, where $\Omega(n) = \sum_{p^k || n} k$ is the number of prime factors of n counted with multiplicity.

The Liouville function is *totally* multiplicative: $\lambda(mn) = (-1)^{\Omega(mn)} = (-1)^{\Omega(m) + \Omega(n)} = \lambda(m)\lambda(n)$ for all m, n . At primes, $\lambda(p) = -1$, and $\lambda(p^k) = (-1)^k$. The Liouville function is a “signed” version of the constant function **1**: it assigns $+1$ to integers with an even number of prime factors and -1 to those with an odd number. The relationship between λ and μ is that $\mu(n) = \lambda(n)$ when n is squarefree, and $\mu(n) = 0$ otherwise (so λ is a “squarefree-ified” extension of μ).

The next two functions count prime factors.

Definition 1.1.7: Prime Factor Counting Functions

For $n \geq 1$:

1. $\omega(n) = \sum_{p|n} 1$ is the number of *distinct* prime factors of n .
2. $\Omega(n) = \sum_{p^k || n} k$ is the number of prime factors of n *counted with multiplicity*.

By convention, $\omega(1) = \Omega(1) = 0$.

The function ω is additive (but not totally additive), and Ω is totally additive. For a prime power p^a : $\omega(p^a) = 1$ and $\Omega(p^a) = a$. The distinction between ω and Ω matters for non-squarefree integers: $\omega(12) = 2$ (the primes 2 and 3) while $\Omega(12) = 3$ (two factors of 2 and one of 3).

The final function in the catalogue is different in character from the preceding ones: it is not multiplicative, and it is supported on prime powers rather than on all positive integers. It is, however, the single most important arithmetic function for the study of prime distribution.

Definition 1.1.8: The Von Mangoldt Function

The *von Mangoldt function* $\Lambda : \mathbb{Z}_{>0} \rightarrow \mathbb{R}_{\geq 0}$ is defined by

$$\Lambda(n) = \begin{cases} \log p & \text{if } n = p^k \text{ for some prime } p \text{ and } k \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

The von Mangoldt function is not multiplicative: $\Lambda(2) = \log 2$ and $\Lambda(3) = \log 3$, but $\Lambda(6) = 0 \neq \Lambda(2)\Lambda(3)$. Its importance comes from the identity that links it to the natural logarithm.

Proposition 1.1.9: Von Mangoldt Identity

For all $n \geq 1$:

$$\sum_{d|n} \Lambda(d) = \log n$$

Proof:

Write $n = p_1^{a_1} \cdots p_r^{a_r}$. The divisors d of n with $\Lambda(d) \neq 0$ are exactly the prime powers p_i^j with $1 \leq j \leq a_i$ and $1 \leq i \leq r$. Hence:

$$\sum_{d|n} \Lambda(d) = \sum_{i=1}^r \sum_{j=1}^{a_i} \log p_i = \sum_{i=1}^r a_i \log p_i = \log \left(\prod_{i=1}^r p_i^{a_i} \right) = \log n$$

This identity says that $\log$ is the summatory function of Λ over divisors. In the language of Dirichlet convolution (introduced in the next section), it reads $\Lambda * \mathbf{1} = \log$, which is one of the key identities in analytic number theory. The von Mangoldt function appears as the coefficient of the logarithmic

derivative of $\zeta(s)$: the identity

$$-\zeta'(s)/\zeta(s) = \sum_{n=1}^{\infty} \Lambda(n)n^{-s}$$

(chapter 4) is the Dirichlet series translation of the von Mangoldt identity, and it is through this series that the zeros of $\zeta(s)$ control the distribution of primes.

The Chebyshev functions, which are the summatory functions of Λ and $\log p$ respectively, will be central in chapter 3:

$$\psi(x) = \sum_{n \leq x} \Lambda(n), \quad \vartheta(x) = \sum_{p \leq x} \log p$$

The Prime Number Theorem (chapter 5) is equivalent to $\psi(x) \sim x$.

Example 1.1.10: Verifying the Von Mangoldt Identity

Take $n = 12 = 2^2 \cdot 3$. The divisors of 12 are $\{1, 2, 3, 4, 6, 12\}$. Among these, the prime powers are $2, 3, 4 = 2^2$, giving:

$$\sum_{d|12} \Lambda(d) = \Lambda(1) + \Lambda(2) + \Lambda(3) + \Lambda(4) + \Lambda(6) + \Lambda(12) = 0 + \log 2 + \log 3 + \log 2 + 0 + 0 = 2 \log 2 + \log 3 = \log 12$$

The non-prime-power divisors 1, 6, and 12 contribute nothing, while $4 = 2^2$ contributes $\log 2$ (not $\log 4$). This is the mechanism by which Λ “distributes” the logarithm of n across its prime-power divisors, assigning weight $\log p$ to each prime p and each power of p dividing n .

The following table collects the values of all the arithmetic functions introduced in this section for $n = 1, \dots, 12$, providing a concrete reference.

Example 1.1.11: Table of Arithmetic Functions

n	1	2	3	4	5	6	7	8	9	10	11	12
$\mu(n)$	1	-1	-1	0	-1	1	-1	0	0	1	-1	0
$\varphi(n)$	1	1	2	2	4	2	6	4	6	4	10	4
$d(n)$	1	2	2	3	2	4	2	4	3	4	2	6
$\sigma(n)$	1	3	4	7	6	12	8	15	13	18	12	28
$\lambda(n)$	1	-1	-1	1	-1	1	-1	-1	1	1	-1	-1
$\omega(n)$	0	1	1	1	1	2	1	1	1	2	1	2
$\Omega(n)$	0	1	1	2	1	2	1	3	2	2	1	3
$\Lambda(n)$	0	$\log 2$	$\log 3$	$\log 2$	$\log 5$	0	$\log 7$	$\log 2$	$\log 3$	0	$\log 11$	0

The patterns are visible: μ vanishes on non-squarefree integers (4, 8, 9, 12); $d(n)$ is large when n has many factorizations (e.g., $d(12) = 6$); Λ is nonzero only at prime powers (2, 3, 4, 5, 7, 8, 9, 11).

Example 1.1.12: Computing Arithmetic Functions from Prime Factorizations

1. Let $n = 2520 = 2^3 \cdot 3^2 \cdot 5 \cdot 7$. Then:

$$d(2520) = (3+1)(2+1)(1+1)(1+1) = 48,$$

$$\sigma(2520) = \frac{2^4-1}{1} \cdot \frac{3^3-1}{2} \cdot \frac{5^2-1}{4} \cdot \frac{7^2-1}{6} = 15 \cdot 13 \cdot 6 \cdot 8 = 9360,$$

$$\mu(2520) = 0 \quad (\text{since } 2^2 | 2520),$$

$$\lambda(2520) = (-1)^{3+2+1+1} = (-1)^7 = -1$$

2. Let $n = 30 = 2 \cdot 3 \cdot 5$. This is squarefree, so $\mu(30) = (-1)^3 = -1$ and $\lambda(30) = -1$ as well. The divisor count is $d(30) = 2^3 = 8$, and $\sigma(30) = 3 \cdot 4 \cdot 6 = 72$. Note $\sigma(30)/30 = 72/30 = 12/5 > 2$, so 30 is an *abundant* number.

The multiplicativity of each function in the catalogue is summarized as follows: μ , φ , σ_k , d , σ are multiplicative; λ is totally multiplicative; ω is additive; Ω is totally additive; Λ and $\log$ are none of the above. Several relationships between these functions are already visible. The Möbius function μ and the Liouville function λ agree on squarefree integers and differ elsewhere. The identity $|\mu(n)| = 1$ if and only if n is squarefree can be expressed as $|\mu| = \mu^2$, and $\sum_{d|n} |\mu(d)| = 2^{\omega(n)}$ counts the squarefree divisors of n . The von Mangoldt identity $\sum_{d|n} \Lambda(d) = \log n$ expresses $\log$ as a “divisor sum” of Λ . All such identities acquire a unified algebraic meaning once the notion of Dirichlet convolution is in place.

1.2 Dirichlet Convolution and Möbius Inversion

The arithmetic functions of the previous section satisfy many identities: $\sum_{d|n} \mu(d) = [n=1]$, $\sum_{d|n} \Lambda(d) = \log n$, $\sum_{d|n} \varphi(d) = n$, and so forth. These identities share a common structure, i.e. a sum over the divisors of n , with the summand depending on both d and n/d . This structure has a name: *Dirichlet convolution*. Equipping the set of arithmetic functions with Dirichlet convolution as multiplication turns it into a commutative ring, and the divisor-sum identities become algebraic statements in this ring:

Definition 1.2.1: Dirichlet Convolution

The *Dirichlet convolution* of two arithmetic functions f and g is the arithmetic function $f * g$ defined by

$$(f * g)(n) = \sum_{d|n} f(d) g(n/d) = \sum_{ab=n} f(a) g(b)$$

The two expressions are equivalent: summing over divisors d of n with complementary divisor n/d is the same as summing over ordered factorizations $n = ab$. The second form makes the commutativity $f * g = g * f$ evident since $ab = ba$.

Several of the identities from section 1.1 are now recognizable as convolution statements:

1. The von Mangoldt identity $\sum_{d|n} \Lambda(d) = \log n$ is $\Lambda * \mathbf{1} = \log$ where $\mathbf{1}$ denotes the constant function $\mathbf{1}(n) = 1$.
2. The totient summation $\sum_{d|n} \varphi(d) = n$ is $\varphi * \mathbf{1} = \text{id}$ where $\text{id}(n) = n$.
3. The divisor function $d(n) = \sum_{d|n} 1$ is $d = \mathbf{1} * \mathbf{1}$ and the sum-of-divisors function $\sigma(n) = \sum_{d|n} d$ is $\sigma = \text{id} * \mathbf{1}$.

Theorem 1.2.2: The Ring of Arithmetic Functions

The set $\mathcal{A}$ of all arithmetic functions, equipped with pointwise addition and Dirichlet convolution as multiplication, forms a commutative ring with identity. The additive identity is the zero function, and the multiplicative identity is

$$\varepsilon(n) = [n = 1] = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n > 1. \end{cases}$$

Proof:

1. *Commutativity.* Commutativity of $*$ follows from the symmetry of the second form: $(f * g)(n) = \sum_{ab=n} f(a)g(b) = \sum_{ab=n} g(a)f(b) = (g * f)(n)$.

2. *Associativity.* For any three arithmetic functions f, g, h :

$$((f * g) * h)(n) = \sum_{de=n} (f * g)(d) h(e) = \sum_{de=n} \sum_{ab=d} f(a)g(b)h(e) = \sum_{abe=n} f(a)g(b)h(e)$$

The final expression is symmetric in the grouping, so $(f * (g * h))(n)$ gives the same triple sum.

3. *Identity.* $(f * \varepsilon)(n) = \sum_{d|n} f(d)\varepsilon(n/d) = f(n)\varepsilon(1) = f(n)$, since $\varepsilon(n/d) = 0$ unless $d = n$.

4. *Distributivity.* $(f * (g + h))(n) = \sum_{d|n} f(d)(g + h)(n/d) = \sum_{d|n} f(d)g(n/d) + \sum_{d|n} f(d)h(n/d) = (f * g)(n) + (f * h)(n)$.

The ring $(\mathcal{A}, +, *)$ is a commutative ring with identity but not an integral domain: for instance, if f is supported on $\{1\}$ and g is supported on $\{1\}$ with $f(1) = g(1) = 0$, then $f * g = 0$. The unit group of $\mathcal{A}$ (the set of arithmetic functions possessing a convolution inverse) is characterized by a simple condition:

Proposition 1.2.3: Units of the Convolution Ring

An arithmetic function f has a Dirichlet convolution inverse f^{-1} (satisfying $f * f^{-1} = \varepsilon$) if and only if $f(1) \neq 0$. When it exists, the inverse is unique and is given recursively by

$$f^{-1}(1) = \frac{1}{f(1)}, \quad f^{-1}(n) = -\frac{1}{f(1)} \sum_{\substack{d|n \\ d < n}} f(n/d) f^{-1}(d) \quad \text{for } n > 1$$

Proof:

The condition $(f * f^{-1})(1) = f(1)f^{-1}(1) = 1$ forces $f(1) \neq 0$ and $f^{-1}(1) = 1/f(1)$. For $n > 1$, the condition $(f * f^{-1})(n) = 0$ reads

$$\sum_{d|n} f(n/d) f^{-1}(d) = 0,$$

and isolating the $d = n$ term gives $f(1)f^{-1}(n) = -\sum_{d|n, d < n} f(n/d)f^{-1}(d)$, which determines $f^{-1}(n)$ recursively from the values $f^{-1}(d)$ for $d < n$.

The recursive formula computes f^{-1} one value at a time in increasing order: $f^{-1}(1)$, then $f^{-1}(2)$, then $f^{-1}(3)$, and so on, each depending only on previously computed values. This is effective for small arguments but does not directly give a closed-form expression.

Example 1.2.4: Computing a Dirichlet Inverse

Compute id^{-1} using the recursive formula. At $n = 1$: $\text{id}^{-1}(1) = 1$. At $n = 2$: $\text{id}^{-1}(2) = -(1/1) \cdot \text{id}(2) \cdot \text{id}^{-1}(1) = -2$. At $n = 3$: $\text{id}^{-1}(3) = -3$. At $n = 4$, the divisors less than 4 are 1 and 2:

$$\text{id}^{-1}(4) = -(\text{id}(4) \cdot \text{id}^{-1}(1) + \text{id}(2) \cdot \text{id}^{-1}(2)) = -(4 \cdot 1 + 2 \cdot (-2)) = 0$$

At $n = 6$: $\text{id}^{-1}(6) = -(6 \cdot 1 + 3 \cdot (-2) + 2 \cdot (-3)) = -(6 - 6 - 6) = 6$. The pattern is $\text{id}^{-1}(n) = n\mu(n)$: the function vanishes at non-squarefree integers, equals $(-1)^{\omega(n)}n$ at squarefree integers, and is multiplicative with $\text{id}^{-1}(p) = -p$ and $\text{id}^{-1}(p^k) = 0$ for $k \geq 2$.

For multiplicative functions, a cleaner description is available.

Proposition 1.2.5: Multiplicative Functions Form a Subgroup

If f and g are multiplicative, then $f * g$ is multiplicative. If f is multiplicative, then f^{-1} exists (since $f(1) = 1$) and is also multiplicative.

Proof:

Let $\gcd(m, n) = 1$. The divisors of mn are exactly the products d_1d_2 where $d_1|m$ and $d_2|n$

(since $\gcd(m, n) = 1$). Hence:

$$\begin{aligned} (f * g)(mn) &= \sum_{d|mn} f(d) g(mn/d) = \sum_{d_1|m} \sum_{d_2|n} f(d_1 d_2) g\left(\frac{m}{d_1} \cdot \frac{n}{d_2}\right) \\ &= \sum_{d_1|m} \sum_{d_2|n} f(d_1) f(d_2) g(m/d_1) g(n/d_2) = (f * g)(m) \cdot (f * g)(n) \end{aligned}$$

Since $(f * g)(1) = f(1)g(1) = 1$, the function $f * g$ is multiplicative.

For the inverse: if f is multiplicative, then $f(1) = 1 \neq 0$, so f^{-1} exists. Multiplicativity of f^{-1} can be verified by induction on mn using the recursive formula: the key observation is that for $\gcd(m, n) = 1$, the divisors of mn factor as products of divisors of m and n , so the recursion for $f^{-1}(mn)$ splits into independent recursions for $f^{-1}(m)$ and $f^{-1}(n)$.

The multiplicative arithmetic functions thus form a group under Dirichlet convolution, with identity ε . This group structure is the reason that identities among multiplicative functions can be verified by checking them at prime powers.

The most important Dirichlet inverse is that of the constant function $\mathbf{1}$.

Proposition 1.2.6: The Möbius Function as a Dirichlet Inverse

$\mu * \mathbf{1} = \varepsilon$. That is, μ is the Dirichlet inverse of $\mathbf{1}$:

$$\sum_{d|n} \mu(d) = [n = 1] = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n > 1. \end{cases}$$

Proof:

Both μ and $\mathbf{1}$ are multiplicative, so $\mu * \mathbf{1}$ is multiplicative by proposition 1.2.5. It suffices to check the identity at prime powers. For p^k with $k \geq 1$:

$$\sum_{d|p^k} \mu(d) = \mu(1) + \mu(p) + \mu(p^2) + \cdots + \mu(p^k) = 1 + (-1) + 0 + \cdots + 0 = 0$$

At $n = 1$: $(\mu * \mathbf{1})(1) = \mu(1) \cdot 1 = 1$. Hence $\mu * \mathbf{1} = \varepsilon$.

This identity is the algebraic content of inclusion-exclusion. For a squarefree integer $n = p_1 \cdots p_r$, the divisors are the 2^r products of subsets of $\{p_1, \dots, p_r\}$, and $\sum_{d|n} \mu(d) = \sum_{k=0}^r \binom{r}{k} (-1)^k = (1-1)^r = 0$.

With the Möbius function identified as the inverse of $\mathbf{1}$, the central inversion formula follows immediately.

Theorem 1.2.7: Möbius Inversion Formula

Let f and g be arithmetic functions. Then:

$$g = f * \mathbf{1} \quad \text{if and only if} \quad f = g * \mu$$

In explicit notation: $g(n) = \sum_{d|n} f(d)$ for all n if and only if $f(n) = \sum_{d|n} \mu(d) g(n/d)$ for all n .

Proof:

If $g = f * \mathbf{1}$, then $g * \mu = (f * \mathbf{1}) * \mu = f * (\mathbf{1} * \mu) = f * \varepsilon = f$, using associativity and $\mathbf{1} * \mu = \varepsilon$ (proposition 1.2.6). Conversely, if $f = g * \mu$, then $f * \mathbf{1} = (g * \mu) * \mathbf{1} = g * (\mu * \mathbf{1}) = g * \varepsilon = g$.

Thus, Möbius inversion is the tool for “solving” a divisor-sum relation: if g is defined as a sum of f over divisors, then f can be recovered from g by convolving with μ .

Example 1.2.8: Recovering φ and Λ via Möbius Inversion

1. The identity $\varphi * \mathbf{1} = \text{id}$ (proposition 0.3.16) inverts to $\varphi = \text{id} * \mu$. In explicit form:

$$\varphi(n) = \sum_{d|n} \mu(d) \cdot \frac{n}{d} = n \sum_{d|n} \frac{\mu(d)}{d} = n \prod_{p|n} \left(1 - \frac{1}{p}\right),$$

where the last step uses multiplicativity: for $n = \prod p_i^{a_i}$, the sum $\sum_{d|n} \mu(d)/d$ factors over primes as $\prod_{p|n} (1 + \mu(p)/p) = \prod_{p|n} (1 - 1/p)$. This recovers the formula from proposition 0.3.10.

2. The von Mangoldt identity $\Lambda * \mathbf{1} = \log$ (proposition 1.1.9) inverts to $\Lambda = \log * \mu$, i.e.,

$$\Lambda(n) = \sum_{d|n} \mu(d) \log(n/d) = - \sum_{d|n} \mu(d) \log d$$

The second equality uses $\sum_{d|n} \mu(d) \log(n/d) = \log n \sum_{d|n} \mu(d) - \sum_{d|n} \mu(d) \log d = - \sum_{d|n} \mu(d) \log d$, since $\sum_{d|n} \mu(d) = 0$ for $n > 1$ (and the case $n = 1$ gives $\Lambda(1) = 0$ directly).

Example 1.2.9: Counting Squarefree Integers

Let $Q(x) = \#\{n \leq x : n \text{ is squarefree}\} = \sum_{n \leq x} |\mu(n)|$. Since $|\mu(n)| = \sum_{d^2|n} \mu(d)$ (an identity verifiable at prime powers: both sides equal 1 if n is squarefree and 0 otherwise), the count can be rewritten as:

$$Q(x) = \sum_{n \leq x} \sum_{d^2|n} \mu(d) = \sum_{d \leq \sqrt{x}} \mu(d) \left\lfloor \frac{x}{d^2} \right\rfloor = \sum_{d \leq \sqrt{x}} \mu(d) \left(\frac{x}{d^2} + O(1) \right) = x \sum_{d \leq \sqrt{x}} \frac{\mu(d)}{d^2} + O(\sqrt{x})$$

The sum $\sum_{d=1}^{\infty} \mu(d)/d^2$ converges to $1/\zeta(2) = 6/\pi^2$ (as the Euler product gives $\sum \mu(d)/d^s = 1/\zeta(s)$,

a fact established rigorously in chapter 2). Accepting this for now:

$$Q(x) = \frac{6}{\pi^2}x + O(\sqrt{x})$$

Hence the density of squarefree integers is $6/\pi^2 \approx 0.6079$: about 61% of all positive integers are squarefree.

The remaining identities from the outline of section 1.1 are now provable in the convolution ring. A systematic collection is recorded here for later reference.

Proposition 1.2.10: Key Convolution Identities

The following identities hold in $(\mathcal{A}, +, *)$:

1. $\mu * \mathbf{1} = \varepsilon$ (Möbius is the inverse of $\mathbf{1}$).
2. $\varphi * \mathbf{1} = \text{id}$, equivalently $\varphi = \text{id} * \mu$.
3. $\Lambda * \mathbf{1} = \log$, equivalently $\Lambda = \log * \mu$.
4. $d = \mathbf{1} * \mathbf{1}$ (the divisor function is the self-convolution of $\mathbf{1}$).
5. $\sigma = \text{id} * \mathbf{1}$.
6. $\sigma_k = \text{id}^k * \mathbf{1}$, where $\text{id}^k(n) = n^k$.
7. $|\mu| * |\mu| = 2^\omega$, where $2^\omega(n) = 2^{\omega(n)}$.
8. $\lambda * |\mu| = \varepsilon$ (λ is the inverse of $|\mu|$).
9. $\lambda * \mathbf{1} = \mathbf{1}_\square$, where $\mathbf{1}_\square(n) = [n \text{ is a perfect square}]$.

Proof:

Identities (1) - (6) have been established above or follow directly from definitions. For the remaining:

7. Both sides are multiplicative (the left by proposition 1.2.5; the right since ω is additive gives $2^{\omega(mn)} = 2^{\omega(m)}2^{\omega(n)}$ for $\gcd(m, n) = 1$). At p^k : $(|\mu| * |\mu|)(p^k) = \sum_{j=0}^k |\mu(p^j)| |\mu(p^{k-j})| = |\mu(1)| |\mu(p^k)| + |\mu(p)| |\mu(p^{k-1})| + \dots$. The nonzero terms are those with $j \in \{0, 1\}$ and $k-j \in \{0, 1\}$, so only $j = 0, k-j \leq 1$ and $j = 1, k-j \leq 1$ contribute. For $k = 0$: the sum is 1. For $k = 1$: $|\mu(1)| |\mu(p)| + |\mu(p)| |\mu(1)| = 2$. For $k \geq 2$: $|\mu(1)| |\mu(p^k)| + |\mu(p)| |\mu(p^{k-1})| = 0$. Hence $(|\mu| * |\mu|)(p^k) = 2$ for $k \geq 1$ and 1 for $k = 0$, matching $2^{\omega(p^k)}$.
8. At p^k : $(\lambda * |\mu|)(p^k) = \sum_{j=0}^k \lambda(p^j) |\mu(p^{k-j})| = (-1)^0 \cdot |\mu(p^k)| + (-1)^1 \cdot |\mu(p^{k-1})| + \dots$. For $k = 0$: the sum is 1. For $k = 1$: $|\mu(p)| + (-1) |\mu(1)| = 1 - 1 = 0$. For $k \geq 2$: $|\mu(p^k)| + (-1) |\mu(p^{k-1})| + |\mu(p^{k-2})| + \dots$; the only nonzero terms are $j = k-1$ (giving $(-1)^{k-1}$) and $j = k$ (giving $(-1)^k$), which sum to 0. Hence $\lambda * |\mu| = \varepsilon$.

$$9. \text{ At } p^k: (\lambda * \mathbf{1})(p^k) = \sum_{j=0}^k (-1)^j = \begin{cases} 1 & k \text{ even,} \\ 0 & k \text{ odd} \end{cases}, \text{ which is } [p^k \text{ is a perfect square}].$$

Identity (9) deserves a moment of interpretation: the summatory function $\sum_{d|n} \lambda(d)$ is 1 if n is a perfect square and 0 otherwise. This identity connects the Liouville function to the distribution of perfect squares, and its Dirichlet series translation $\zeta(2s)/\zeta(s) = \sum \lambda(n)n^{-s}$ will appear in chapter 2.

1.3 Average Orders and Summatory Functions

The convolution identities of the previous section describe exact relationships among arithmetic functions at individual integers. The present section asks a different question: how do these functions behave *on average*? The divisor function $d(n)$ fluctuates wildly, for example $d(p) = 2$ for primes, while $d(n)$ can be arbitrarily large for highly composite n . But its average over $\{1, \dots, N\}$ grows smoothly like $\log N$. Extracting this smooth average from the pointwise behavior is the first analytic problem of the book, and the main tool is the *Dirichlet hyperbola method*: a technique for evaluating $\sum_{n \leq x} (f * g)(n)$ by splitting the double sum along the hyperbola $ab = n$.

Definition 1.3.1: Summatory Function and Average Order

For an arithmetic function f , the *summatory function* is

$$M_f(x) = \sum_{n \leq x} f(n)$$

The function $g(x)$ is an *average order* of f if $M_f(x) \sim g(x)$ as $x \rightarrow \infty$, i.e., $\sum_{n \leq x} f(n)/g(x) \rightarrow 1$.

The summatory function converts the discrete sequence $(f(n))$ into a step function of a continuous variable x . The average order captures the typical size of $f(n)$: if $g(x)$ is an average order of f , then $f(n)$ is “typically” of size $g(n)/n$. The fundamental technique for estimating summatory functions of convolutions is the following:

Theorem 1.3.2: Dirichlet’s Hyperbola Method

Let f and g be arithmetic functions, and set $h = f * g$. For any $x \geq 1$ and any y with $1 \leq y \leq x$:

$$\sum_{n \leq x} h(n) = \sum_{a \leq y} f(a) G\left(\frac{x}{a}\right) + \sum_{b \leq x/y} g(b) F\left(\frac{x}{b}\right) - F(y) G\left(\frac{x}{y}\right)$$

where $F(t) = \sum_{n \leq t} f(n)$ and $G(t) = \sum_{n \leq t} g(n)$.

The idea is to write $\sum_{n \leq x} h(n) = \sum_{ab \leq x} f(a)g(b)$ and split the region $\{(a, b) : ab \leq x\}$ into two parts along the line $a = y$. The first sum counts pairs with $a \leq y$, the second counts pairs with $b \leq x/y$ (equivalently $a > y$), and the subtracted term corrects for the overlap where both $a \leq y$ and $b \leq x/y$.

Proof:

Start from $h = f * g$:

$$\sum_{n \leq x} h(n) = \sum_{\substack{a, b \geq 1 \\ ab \leq x}} f(a)g(b)$$

Step 1: Partition the summation region. Split $\{(a, b) \in \mathbb{Z}_{>0}^2 : ab \leq x\}$ into: $S_1 = \{a \leq y\}$ and $S_2 = \{a > y, b \leq x/y\}$. Every pair (a, b) with $ab \leq x$ belongs to at least one of these sets (if $a > y$, then $b \leq x/a < x/y$), and the overlap is $S_1 \cap S_2 = \{a \leq y, b \leq x/y\}$. By inclusion-exclusion:

$$\sum_{ab \leq x} f(a)g(b) = \sum_{\substack{a \leq y \\ ab \leq x}} f(a)g(b) + \sum_{\substack{a > y \\ b \leq x/y}} f(a)g(b) - \sum_{\substack{a \leq y \\ b \leq x/y}} f(a)g(b) + \sum_{\substack{a \leq y \\ b \leq x/y}} f(a)g(b)$$

Step 2: Evaluate each piece. The first sum: for fixed $a \leq y$, the condition $ab \leq x$ means $b \leq x/a$, so $\sum_{\substack{a \leq y \\ ab \leq x}} f(a)g(b) = \sum_{a \leq y} f(a)G(x/a)$. The second sum plus the overlap: $\sum_{b \leq x/y} \sum_{a \leq x/b} f(a)g(b) = \sum_{b \leq x/y} g(b)F(x/b)$. The overlap alone: $\sum_{a \leq y} f(a) \sum_{b \leq x/y} g(b) = F(y)G(x/y)$. Combining gives the stated formula.

The optimal choice of the splitting parameter y depends on the specific functions f and g ; the symmetric choice $y = \sqrt{x}$ is the most common and typically minimizes the error term. The power of the method lies in reducing the estimation of $\sum h(n)$ to the estimation of F and G separately—a significant simplification when f and g are simpler than h .

Geometrically, the lattice points $(a, b) \in \mathbb{Z}_{>0}^2$ with $ab \leq x$ lie below the hyperbola $ab = x$ in the first quadrant. The horizontal line $a = y$ divides this region into an upper-left piece (where the sum over a is short and G handles the sum over b) and a lower-right piece (where the sum over b is short and F handles the sum over a). The rectangle $\{a \leq y, b \leq x/y\}$ is the overlap region. This geometric picture—counting lattice points under a hyperbola—is a prototype for the lattice-point counting arguments that appear in the proof of the analytic class number formula (chapter 8).

The first and most classical application is to the divisor function $d = \mathbf{1} * \mathbf{1}$.

Theorem 1.3.3: Average Order of the Divisor Function

$$\sum_{n \leq x} d(n) = x \log x + (2\gamma - 1)x + O(\sqrt{x})$$

where $\gamma = 0.5772\dots$ is the Euler-Mascheroni constant. In particular, the average order of $d(n)$ is $\log n$.

Proof:

Since $d = \mathbf{1} * \mathbf{1}$, apply theorem 1.3.2 with $f = g = \mathbf{1}$ and $y = \sqrt{x}$. The summatory function of $\mathbf{1}$ is $F(t) = G(t) = [t] = t + O(1)$. By symmetry, the first two sums in the hyperbola formula are equal, giving:

$$\sum_{n \leq x} d(n) = 2 \sum_{a \leq \sqrt{x}} \left\lfloor \frac{x}{a} \right\rfloor - [\sqrt{x}]^2$$

Using $\lfloor x/a \rfloor = x/a + O(1)$:

$$2 \sum_{a \leq \sqrt{x}} \left\lfloor \frac{x}{a} \right\rfloor = 2x \sum_{a \leq \sqrt{x}} \frac{1}{a} + O(\sqrt{x})$$

The harmonic sum satisfies $\sum_{a \leq \sqrt{x}} 1/a = \frac{1}{2} \log x + \gamma + O(1/\sqrt{x})$, so:

$$2x \sum_{a \leq \sqrt{x}} \frac{1}{a} = x \log x + 2\gamma x + O(\sqrt{x})$$

For the subtracted term: $\lfloor \sqrt{x} \rfloor^2 = (\sqrt{x} + O(1))^2 = x + O(\sqrt{x})$. Combining:

$$\sum_{n \leq x} d(n) = x \log x + 2\gamma x + O(\sqrt{x}) - x - O(\sqrt{x}) = x \log x + (2\gamma - 1)x + O(\sqrt{x})$$

1.3.4 Remark: The Dirichlet Divisor Problem The error term $O(\sqrt{x})$ in theorem 1.3.3 is not optimal. Let $\Delta(x) = \sum_{n \leq x} d(n) - x \log x - (2\gamma - 1)x$ denote the error. The best known upper bound as of writing this book, due to Huxley (2003), is $\Delta(x) = O(x^{131/416+\varepsilon})$, where $131/416 \approx 0.3149$. Hardy proved $\Delta(x) = \Omega(x^{1/4})$, establishing that the error is at least of order $x^{1/4}$ infinitely often. The conjectured optimal exponent is $1/4$, and determining the true order of $\Delta(x)$ (the *Dirichlet divisor problem*) remains open.

Example 1.3.5: Numerical Verification of the Divisor Sum

For $x = 100$: the exact value is $\sum_{n=1}^{100} d(n) = 482$. The asymptotic formula gives $100 \log 100 + (2\gamma - 1) \cdot 100 \approx 460.5 + 15.4 = 476.0$, with error $482 - 476 = 6$. The bound $O(\sqrt{100}) = O(10)$ is consistent. At $x = 10,000$: the exact sum is 93,668, and the formula gives $10,000 \log(10,000) + (2\gamma - 1) \cdot 10,000 \approx 92,103 + 1,544 = 93,648$, with error $20 = O(\sqrt{10,000}) = O(100)$.

For comparison, the summatory function of μ exhibits much more complicated behavior. For future reference, we box the following:

Definition 1.3.6: The Mertens Function

The *Mertens function* is $M(x) = \sum_{n \leq x} \mu(n)$.

The identity $\mu * \mathbf{1} = \varepsilon$ (proposition 1.2.6) implies $\sum_{d \leq x} M(x/d) = 1$, a self-referencing identity that constrains the growth of M . The Mertens function oscillates heavily and grows much more slowly than x : the bound $M(x) = o(x)$ is equivalent to the Prime Number Theorem (chapter 5), and the Riemann Hypothesis is equivalent to $M(x) = O(x^{1/2+\varepsilon})$ for every $\varepsilon > 0$. Mertens conjectured in 1897 that $|M(x)| < \sqrt{x}$ for all $x \geq 1$; this was disproved by Odlyzko and te Riele in 1985, though no explicit counterexample is known.

The rest of this section is dedicated to computing average orders of various arithmetic functions:

Theorem 1.3.7: Average Order of the Sum-of-Divisors Function

$$\sum_{n \leq x} \sigma(n) = \frac{\pi^2}{12} x^2 + O(x \log x)$$

Proof:

Since $\sigma = \text{id} * \mathbf{1}$, the double-sum representation gives $\sum_{n \leq x} \sigma(n) = \sum_{ab \leq x} a$. Summing over b first:

$$\sum_{n \leq x} \sigma(n) = \sum_{b \leq x} \sum_{a \leq x/b} a = \sum_{b \leq x} \frac{1}{2} \left\lfloor \frac{x}{b} \right\rfloor \left(\left\lfloor \frac{x}{b} \right\rfloor + 1 \right)$$

Since $\lfloor t \rfloor (\lfloor t \rfloor + 1)/2 = t^2/2 + O(t)$:

$$\sum_{b \leq x} \left(\frac{x^2}{2b^2} + O\left(\frac{x}{b}\right) \right) = \frac{x^2}{2} \sum_{b \leq x} \frac{1}{b^2} + O(x \log x) = \frac{x^2}{2} \left(\frac{\pi^2}{6} + O(1/x) \right) + O(x \log x) = \frac{\pi^2}{12} x^2 + O(x \log x)$$

A numerical check: $\sum_{n=1}^{100} \sigma(n) = 8299$, while $\frac{\pi^2}{12} \cdot 100^2 \approx 8225$, giving an error of $74 = O(100 \log 100) = O(461)$.

Theorem 1.3.8: Average Order of the Euler Totient

$$\sum_{n \leq x} \varphi(n) = \frac{3}{\pi^2} x^2 + O(x \log x)$$

The leading coefficient $3/\pi^2 = 1/(2\zeta(2))$ reflects the density of coprime pairs: if two integers are chosen uniformly at random from $\{1, \dots, N\}$, the probability that they are coprime tends to $6/\pi^2$ as $N \rightarrow \infty$.

Proof:

By Möbius inversion (theorem 1.2.7), $\varphi(n) = \sum_{d|n} \mu(d)(n/d)$. Summing over $n \leq x$ and interchanging the order:

$$\sum_{n \leq x} \varphi(n) = \sum_{d \leq x} \mu(d) \sum_{\substack{n \leq x \\ d|n}} \frac{n}{d}$$

Writing $n = dm$ with $m \leq x/d$:

$$\sum_{\substack{n \leq x \\ d|n}} \frac{n}{d} = \sum_{m \leq x/d} m = \frac{1}{2} \left\lfloor \frac{x}{d} \right\rfloor \left(\left\lfloor \frac{x}{d} \right\rfloor + 1 \right) = \frac{x^2}{2d^2} + O\left(\frac{x}{d}\right)$$

Substituting:

$$\sum_{n \leq x} \varphi(n) = \frac{x^2}{2} \sum_{d \leq x} \frac{\mu(d)}{d^2} + O\left(x \sum_{d \leq x} \frac{1}{d}\right) = \frac{x^2}{2} \left(\frac{6}{\pi^2} + O(1/x) \right) + O(x \log x) = \frac{3}{\pi^2} x^2 + O(x \log x)$$

The identity $\sum_{d=1}^{\infty} \mu(d)/d^2 = 1/\zeta(2) = 6/\pi^2$ is the Euler product $\prod_p (1 - p^{-2})$; a rigorous proof appears in chapter 2.

The appearance of $6/\pi^2$ in the totient average has a probabilistic interpretation. The number of pairs (a, b) with $1 \leq a, b \leq N$ and $\gcd(a, b) = 1$ is $\sum_{a=1}^N \sum_{b=1}^N [\gcd(a, b) = 1] = 2 \sum_{n=1}^N \varphi(n) - 1$ (the term -1 accounts for the pair $(1, 1)$ counted once). The density of coprime pairs is therefore

$$\frac{2 \sum_{n=1}^N \varphi(n) - 1}{N^2} = \frac{6/\pi^2 \cdot N^2 + O(N \log N)}{N^2} \rightarrow \frac{6}{\pi^2} \approx 0.6079$$

Concretely, if a and b are chosen uniformly at random from $\{1, \dots, N\}$, the probability that $\gcd(a, b) = 1$ approaches $6/\pi^2$. Each prime p contributes independently: the probability that $p \nmid \gcd(a, b)$ is $1 - 1/p^2$, and the product $\prod_p (1 - 1/p^2) = 6/\pi^2$ reflects this independence.

The three estimates $\sum d(n) \sim x \log x$, $\sum \sigma(n) \sim \frac{\pi^2}{12} x^2$, and $\sum \varphi(n) \sim \frac{3}{\pi^2} x^2$ illustrate a pattern: the average order of a multiplicative function defined as a convolution $f * \mathbf{1}$ is controlled by the summatory behavior of f and the value $\zeta(2) = \pi^2/6$. The Dirichlet series framework of chapter 2 makes this pattern systematic.

The additive function $\omega(n)$ (number of distinct prime divisors) has a subtler average order.

Proposition 1.3.9: Average Order of $\omega(n)$

$$\sum_{n \leq x} \omega(n) = x \log \log x + Bx + O\left(\frac{x}{\log x}\right)$$

where $B = \gamma + \sum_p (\log(1 - 1/p) + 1/p) = 0.2614\dots$ is related to Mertens' constant. In particular, the average order of $\omega(n)$ is $\log \log n$.

The proof uses the representation $\omega(n) = \sum_p |n|_p$, which after interchanging the order of summation gives $\sum_{n \leq x} \omega(n) = \sum_{p \leq x} \lfloor x/p \rfloor = x \sum_{p \leq x} 1/p + O(\pi(x))$, and the estimate $\sum_{p \leq x} 1/p = \log \log x + M + O(1/\log x)$ (Mertens' second theorem, chapter 3) completes the argument. The full proof is given after Mertens' theorems are established; an alternative proof using the Euler product of $\zeta(s)$ appears in chapter 3 as well.

The average $\log \log n$ is strikingly small: a “typical” integer near 10^9 has only about $\log \log(10^9) \approx 3.0$ distinct prime factors. The celebrated Erdős-Kac theorem (1940) refines this by showing that $(\omega(n) - \log \log n)/\sqrt{\log \log n}$ is approximately normally distributed—a Gaussian central limit theorem for a number-theoretic function. This theorem lies outside the scope of this book, but the average order $\log \log n$ established here is the mean of that Gaussian.

Example 1.3.10: Summatory Functions and the Squarefree Indicator

The squarefree indicator function $|\mu(n)|$ (which is 1 if n is squarefree, 0 otherwise) has summatory function $Q(x) = \sum_{n \leq x} |\mu(n)|$. In example 1.2.9, the estimate $Q(x) = \frac{6}{\pi^2} x + O(\sqrt{x})$ was derived using the identity $|\mu(\tilde{n})| = \sum_{d^2 | n} \mu(d)$. This can also be obtained from the convolution $|\mu| * \mathbf{1}_{\square} = \mathbf{1}$ (where $\mathbf{1}_{\square}(n) = [n \text{ is a perfect square}]$), which inverts to $|\mu| = \mu * \mathbf{1}_{\square}^{-1}$. The approach via the identity $|\mu(n)| = \sum_{d^2 | n} \mu(d)$ is more direct and avoids computing the inverse of $\mathbf{1}_{\square}$.

Example 1.3.11: Comparing Average Orders

The average orders established in this section are:

$$d(n) \sim \log n, \quad \sigma(n) \sim \frac{\pi^2}{6}n, \quad \varphi(n) \sim \frac{6}{\pi^2}n, \quad \omega(n) \sim \log \log n$$

The first and fourth grow logarithmically (or doubly logarithmically), while σ and φ grow linearly. The product $\sigma(n)\varphi(n) \sim n^2$ on average, reflecting the identity $\varphi(n)\sigma(n)/n^2 = \prod_{p|n} (1 - 1/p^2)$, which is bounded between $6/\pi^2$ and 1 for all n .

Example 1.3.12: Average Behavior of $\varphi(n)/n$

The estimate $\sum_{n \leq x} \varphi(n) = \frac{3}{\pi^2}x^2 + O(x \log x)$ implies that the average value of $\varphi(n)/n$ over $\{1, \dots, N\}$ tends to $6/\pi^2 \approx 0.6079$. This is the same constant as the density of squarefree integers (example 1.2.9): both equal $1/\zeta(2)$. The connection is structural—the “probability” that two integers are coprime and the density of squarefree integers are both governed by $\zeta(2)^{-1} = \prod_p (1 - p^{-2})$.

1.4 Partial Summation and Abel's Summation Formula

The previous section estimated summatory functions by exploiting the convolution structure of specific arithmetic functions. This section introduces a technique that applies to *any* weighted sum $\sum a_n f(n)$, regardless of whether the coefficients a_n have multiplicative structure: *Abel's summation formula*, the discrete analogue of integration by parts. The formula converts a sum $\sum a_n f(n)$ into an integral involving the summatory function $A(t) = \sum_{n \leq t} a_n$ and the derivative f' , thereby transferring information about the partial sums of (a_n) into analytic information about the weighted sum. It is arguably the single most frequently used technical tool in this book.

Theorem 1.4.1: Abel's Summation Formula

Let $(a_n)_{n \geq 1}$ be a sequence of complex numbers, and let $f : [1, \infty) \rightarrow \mathbb{C}$ be a continuously differentiable function. Define $A(t) = \sum_{n \leq t} a_n$. Then for any $x \geq 1$:

$$\sum_{n \leq x} a_n f(n) = A(x)f(x) - \int_1^x A(t) f'(t) dt$$

The formula is the exact analogue of integration by parts: $\int u dv = uv - \int v du$, with $u = f$, $dv = dA$ (the Stieltjes measure associated to the step function A), $v = A$, and $du = f' dt$. The “boundary term” $A(x)f(x)$ corresponds to $[uv]$, and the integral $\int A(t)f'(t) dt$ corresponds to $\int v du$. The lower limit of integration is 1 rather than 0 because $A(t) = 0$ for $t < 1$.

Proof:

For $x \geq 1$, write $N = \lfloor x \rfloor$ and split the sum into a telescoping form. For each n with $1 \leq n \leq N$:

$$a_n = A(n) - A(n-1)$$

Hence:

$$\begin{aligned}
 \sum_{n=1}^N a_n f(n) &= \sum_{n=1}^N (A(n) - A(n-1))f(n) \\
 &= \sum_{n=1}^N A(n)f(n) - \sum_{n=1}^N A(n-1)f(n) \\
 &= \sum_{n=1}^N A(n)f(n) - \sum_{n=0}^{N-1} A(n)f(n+1) \\
 &= A(N)f(N) + \sum_{n=1}^{N-1} A(n)(f(n) - f(n+1)) - A(0)f(1)
 \end{aligned}$$

Since $A(0) = 0$ and $f(n) - f(n+1) = -\int_n^{n+1} f'(t) dt$:

$$\sum_{n=1}^N a_n f(n) = A(N)f(N) - \sum_{n=1}^{N-1} A(n) \int_n^{n+1} f'(t) dt$$

On the interval $[n, n+1)$, the function $A(t)$ is constant with value $A(n)$. Hence $\sum_{n=1}^{N-1} A(n) \int_n^{n+1} f'(t) dt = \int_1^N A(t)f'(t) dt$. Finally, since $A(t) = A(N)$ for $t \in [N, x]$:

$$A(x)f(x) - \int_1^x A(t)f'(t) dt = A(N)f(x) - \int_1^N A(t)f'(t) dt - A(N) \int_N^x f'(t) dt = A(N)f(N) - \int_1^N A(t)f'(t) dt$$

using $f(x) - \int_N^x f'(t) dt = f(N)$. This matches the expression above.

The power of Abel's formula lies in the asymmetry of its inputs and outputs. The input is a *discrete* sum $\sum a_n f(n)$, which may be difficult to estimate directly. The output replaces this with a *continuous* integral $\int A(t)f'(t) dt$ plus a boundary term, and the integral is often much easier to handle, especially when $A(t)$ has a known asymptotic and $f'(t)$ is smooth and decaying.

The formula can also be written in Stieltjes integral notation as

$$\sum_{n \leq x} a_n f(n) = \int_1^x f(t) dA(t)$$

where $dA(t)$ is the Stieltjes measure that places mass a_n at each integer n . Integration by parts for Stieltjes integrals then gives $\int f dA = [fA] - \int A df = A(x)f(x) - \int_1^x A(t)f'(t) dt$, recovering theorem 1.4.1. This perspective will return when studying the Mellin and Laplace transforms that appear in chapter 5 and chapter 6.

A useful variant handles the “other direction”: given knowledge of a weighted sum, deduce information about the partial sums.

Corollary 1.4.2: Summatory Function Estimate from Weighted Sums

If $\sum_{n \leq x} a_n f(n) = G(x) + O(E(x))$ and f is positive, monotonically decreasing, and continuously differentiable, then

$$A(x) = \frac{G(x)}{f(x)} + \frac{1}{f(x)} \int_1^x A(t) f'(t) dt + O\left(\frac{E(x)}{f(x)}\right)$$

When $A(t)$ appears on both sides, this is an implicit equation that can sometimes be solved by iteration.

Proof:

Rearranging theorem 1.4.1: $A(x) = \frac{1}{f(x)} \left(\sum_{n \leq x} a_n f(n) + \int_1^x A(t) f'(t) dt \right)$. Substituting the hypothesis gives the result.

This corollary is used in chapter 3 to pass from the estimate $\sum_{p \leq x} \log p/p = \log x + O(1)$ (Mertens' first theorem) to $\sum_{p \leq x} 1/p = \log \log x + O(1)$ (Mertens' second theorem).

Corollary 1.4.3: Summation by Parts for Convergent Series

Under the hypotheses of theorem 1.4.1, if $f(x) \rightarrow 0$ as $x \rightarrow \infty$ and the integral $\int_1^\infty A(t) f'(t) dt$ converges absolutely, then:

$$\sum_{n=1}^{\infty} a_n f(n) = - \int_1^{\infty} A(t) f'(t) dt$$

Proof:

Take $x \rightarrow \infty$ in theorem 1.4.1. The boundary term $A(x)f(x)$ tends to zero provided $A(x)f(x) \rightarrow 0$, which holds when $A(x) = O(x^c)$ for some c and $f(x) = o(x^{-c})$. The integral converges by hypothesis.

The first application recovers the harmonic sum asymptotic, providing an alternative to the integral comparison used in section 0.6.

Example 1.4.4: The Harmonic Sum via Abel Summation

Take $a_n = 1$ and $f(t) = 1/t$, so $A(t) = [t] = t - \{t\}$ and $f'(t) = -1/t^2$. Abel's formula gives:

$$\sum_{n \leq x} \frac{1}{n} = \frac{[x]}{x} + \int_1^x \frac{[t]}{t^2} dt = 1 + O(1/x) + \int_1^x \frac{t - \{t\}}{t^2} dt = 1 + \int_1^x \frac{dt}{t} - \int_1^x \frac{\{t\}}{t^2} dt + O(1/x)$$

The first integral is $\log x$. The second integral converges as $x \rightarrow \infty$ (since $0 \leq \{t\} < 1$ and $\int_1^\infty t^{-2} dt < \infty$) to a value $\int_1^\infty \{t\}/t^2 dt = 1 - \gamma$, where γ is the Euler-Mascheroni constant. Hence:

$$\sum_{n \leq x} \frac{1}{n} = \log x + \gamma + O(1/x)$$

The error $O(1/x)$ comes from the boundary term $[x]/x = 1 - \{x\}/x$ and the tail of $\int_x^\infty \{t\}/t^2 dt =$

$O(1/x)$.

Example 1.4.5: Weighted Divisor Sum

Take $a_n = d(n)$ and $f(t) = 1/t$. By theorem 1.3.3, $A(t) = \sum_{n \leq t} d(n) = t \log t + (2\gamma - 1)t + O(\sqrt{t})$. Abel's formula gives:

$$\begin{aligned} \sum_{n \leq x} \frac{d(n)}{n} &= \frac{A(x)}{x} + \int_1^x \frac{A(t)}{t^2} dt \\ &= \log x + (2\gamma - 1) + O(x^{-1/2}) + \int_1^x \frac{t \log t + (2\gamma - 1)t + O(\sqrt{t})}{t^2} dt \\ &= \log x + (2\gamma - 1) + \int_1^x \frac{\log t}{t} dt + (2\gamma - 1) \int_1^x \frac{dt}{t} + O(1) \\ &= \log x + (2\gamma - 1) + \frac{(\log x)^2}{2} + (2\gamma - 1) \log x + O(1) \end{aligned}$$

Hence $\sum_{n \leq x} d(n)/n = \frac{1}{2}(\log x)^2 + 2\gamma \log x + O(1)$. This estimate is used in chapter 2 to analyze the Dirichlet series $\sum d(n)n^{-s} = \zeta(s)^2$.

The next application previews the connection between partial summation and Dirichlet series that will be developed in chapter 2.

Proposition 1.4.6: Partial Sums of n^{-s}

For $s \in \mathbb{C}$ with $\operatorname{Re}(s) > 0$ and $s \neq 1$:

$$\sum_{n \leq x} \frac{1}{n^s} = \frac{x^{1-s}}{1-s} + \zeta(s) + O(x^{-\operatorname{Re}(s)})$$

where $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ for $\operatorname{Re}(s) > 1$, and the formula provides the analytic continuation of the right-hand side to $\operatorname{Re}(s) > 0$.

Proof:

Apply theorem 1.4.1 with $a_n = 1$ and $f(t) = t^{-s}$. Then $A(t) = [t] = t - \{t\}$ and $f'(t) = -st^{-s-1}$:

$$\sum_{n \leq x} n^{-s} = \frac{[x]}{x^s} + s \int_1^x \frac{[t]}{t^{s+1}} dt = \frac{x - \{x\}}{x^s} + s \int_1^x \frac{t - \{t\}}{t^{s+1}} dt$$

Separating the main term from the fractional part:

$$= x^{1-s} + s \int_1^x t^{-s} dt - s \int_1^x \frac{\{t\}}{t^{s+1}} dt + O(x^{-\operatorname{Re}(s)})$$

The first integral evaluates to $\frac{x^{1-s}-1}{1-s}$ for $s \neq 1$. The second integral converges absolutely for $\operatorname{Re}(s) > 0$ (since $|\{t\}/t^{s+1}| \leq t^{-\operatorname{Re}(s)-1}$), and its value from 1 to ∞ defines a constant depending

on s . Combining:

$$\sum_{n \leq x} n^{-s} = x^{1-s} + \frac{s}{1-s}(x^{1-s} - 1) - s \int_1^x \frac{\{t\}}{t^{s+1}} dt + O(x^{-\operatorname{Re}(s)}) = \frac{x^{1-s}}{1-s} + C(s) + O(x^{-\operatorname{Re}(s)})$$

where $C(s) = \frac{s}{s-1} - s \int_1^\infty \{t\} t^{-s-1} dt$. For $\operatorname{Re}(s) > 1$, taking $x \rightarrow \infty$ shows $C(s) = \zeta(s)$, and the formula extends $C(s)$ to $\operatorname{Re}(s) > 0$ by the convergence of the integral.

This result simultaneously provides the meromorphic continuation of $\zeta(s)$ to $\operatorname{Re}(s) > 0$ (with a simple pole at $s = 1$) and a useful asymptotic for the partial sums $\sum_{n \leq x} n^{-s}$. The full meromorphic continuation to all of $\mathbb{C}$ requires the functional equation (chapter 6).

Example 1.4.7: Estimating $\sum_{p \leq x} 1/p$ from $\pi(x)$

Abel summation converts knowledge of $\pi(x) = \sum_{p \leq x} 1$ (the prime counting function) into an estimate for $\sum_{p \leq x} 1/p$. Taking $a_p = 1$ for primes p and $a_n = 0$ otherwise, so $A(t) = \pi(t)$, and $f(t) = 1/t$:

$$\sum_{p \leq x} \frac{1}{p} = \frac{\pi(x)}{x} + \int_2^x \frac{\pi(t)}{t^2} dt$$

If the Prime Number Theorem $\pi(t) \sim t/\log t$ is assumed, then $\pi(x)/x \sim 1/\log x \rightarrow 0$ and

$$\int_2^x \frac{\pi(t)}{t^2} dt \sim \int_2^x \frac{dt}{t \log t} = \log \log x - \log \log 2$$

Hence $\sum_{p \leq x} 1/p \sim \log \log x$, recovering Mertens' second theorem (up to the precise constant). The rigorous version of this argument, using Chebyshev's bounds instead of PNT, appears in chapter 3.

This example illustrates a general principle: Abel summation lets the estimation of $\sum f(p)$ over primes be reduced to the estimation of $\pi(x)$, and conversely. The relationship is bidirectional: knowledge of $\pi(x)$ yields information about $\sum 1/p$, and the divergence of $\sum 1/p$ (proved in chapter 4 via the Euler product) gives a lower bound on $\pi(x)$.

Example 1.4.8: Partial Summation with the Totient Function

To estimate $\sum_{n \leq x} \varphi(n)/n^2$, apply Abel summation with $a_n = \varphi(n)$ and $f(t) = 1/t^2$. By theorem 1.3.8, $A(t) = \sum_{n \leq t} \varphi(n) = \frac{3}{\pi^2} t^2 + O(t \log t)$. Then:

$$\begin{aligned} \sum_{n \leq x} \frac{\varphi(n)}{n^2} &= \frac{A(x)}{x^2} + 2 \int_1^x \frac{A(t)}{t^3} dt = \frac{3}{\pi^2} + O\left(\frac{\log x}{x}\right) + 2 \int_1^x \frac{3t^2/\pi^2 + O(t \log t)}{t^3} dt \\ &= \frac{3}{\pi^2} + \frac{6}{\pi^2} \int_1^x \frac{dt}{t} + O\left(\int_1^x \frac{\log t}{t^2} dt\right) + O\left(\frac{\log x}{x}\right) \\ &= \frac{3}{\pi^2} + \frac{6}{\pi^2} \log x + O(1) \end{aligned}$$

The dominant term is $\frac{6}{\pi^2} \log x$, confirming that $\sum \varphi(n)/n^2$ diverges logarithmically.

Abel summation also provides a useful criterion for the convergence of series with oscillating terms.

Corollary 1.4.9: Abel's Test for Dirichlet Series

Let (a_n) be a sequence with bounded partial sums: $|A(N)| \leq M$ for all N . If $f : [1, \infty) \rightarrow \mathbb{R}$ is monotonically decreasing with $f(t) \rightarrow 0$ as $t \rightarrow \infty$, then $\sum_{n=1}^{\infty} a_n f(n)$ converges, and

$$\left| \sum_{n>N} a_n f(n) \right| \leq 2M f(N+1)$$

Proof:

By theorem 1.4.1 applied to the tail sum:

$$\sum_{N < n \leq x} a_n f(n) = A'(x)f(x) - A'(N)f(N+1) - \int_{N+1}^x A'(t)f'(t) dt$$

where $A'(t) = \sum_{N < n \leq t} a_n = A(t) - A(N)$, so $|A'(t)| \leq 2M$. Since f is decreasing, $f' \leq 0$, and:

$$\left| \int_{N+1}^x A'(t)f'(t) dt \right| \leq 2M \int_{N+1}^x |f'(t)| dt = 2M(f(N+1) - f(x))$$

The boundary terms satisfy $|A'(x)f(x)| \leq 2Mf(x) \rightarrow 0$ and $|A'(N)f(N+1)| = 0$. Taking $x \rightarrow \infty$: $|\sum_{n>N} a_n f(n)| \leq 2Mf(N+1)$.

This criterion applies, for example, to $\sum \mu(n)/n^s$ for $\operatorname{Re}(s) > 0$: the partial sums $\sum_{n \leq N} \mu(n) = M(N)$ satisfy $M(N) = o(N)$ (equivalent to PNT, as noted in section 1.3), so the series converges for $\operatorname{Re}(s) > 0$ once PNT is established. Before PNT, convergence is known only for $\operatorname{Re}(s) > 1$ from absolute convergence.

1.5 The Euler-Maclaurin Summation Formula

Abel's summation formula from the previous section converts a sum into an integral plus a boundary term, with an error controlled by the fractional part of the summation variable. For many applications, a single error term is sufficient. But when high-precision asymptotic expansions are needed (expansions of the form $\sum_{n \leq N} f(n) = I(N) + c_1 g_1(N) + c_2 g_2(N) + \dots$ with explicit coefficients) Abel summation isn't enough. The *Euler-Maclaurin formula* remedies this by providing an arbitrarily accurate comparison between $\sum f(n)$ and $\int f(t) dt$, with error terms involving successively higher derivatives of f weighted by *Bernoulli numbers*. These numbers, appearing repeatedly throughout analytic number theory, are introduced first:

Definition 1.5.1: Bernoulli Numbers

The *Bernoulli numbers* $B_k \in \mathbb{Q}$ for $k \geq 0$ are defined by the generating function

$$\frac{t}{e^t - 1} = \sum_{k=0}^{\infty} \frac{B_k}{k!} t^k$$

valid for $|t| < 2\pi$.

Equating coefficients with the Taylor expansion of $t/(e^t - 1)$ gives a recursive formula: multiplying both sides by $(e^t - 1)/t = \sum_{j=0}^{\infty} t^j/(j+1)!$ and identifying coefficients yields $B_0 = 1$ and, for $n \geq 1$,

$$\sum_{k=0}^n \binom{n+1}{k} B_k = 0$$

This recursion computes the Bernoulli numbers one at a time. The first several values are:

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_3 = 0, \quad B_4 = -\frac{1}{30}, \quad B_5 = 0, \quad B_6 = \frac{1}{42}, \quad B_8 = -\frac{1}{30}, \quad B_{10} = \frac{5}{66}$$

A fundamental property is the vanishing of odd-indexed Bernoulli numbers beyond B_1 .

Proposition 1.5.2: Vanishing of Odd Bernoulli Numbers

For $k \geq 1$, $B_{2k+1} = 0$.

Proof:

Consider the function $G(t) = t/(e^t - 1) + t/2$. Then

$$G(t) = \sum_{k=0}^{\infty} \frac{B_k}{k!} t^k + \frac{t}{2} = 1 + \sum_{k=2}^{\infty} \frac{B_k}{k!} t^k,$$

using $B_0 = 1$ and $B_1 = -1/2$. Direct computation gives $G(-t) = -t/(e^{-t} - 1) - t/2 = t \cdot e^t/(e^t - 1) - t/2 = t(1 + 1/(e^t - 1)) - t/2 = t/(e^t - 1) + t/2 = G(t)$. Hence G is an even function of t , so all coefficients of odd powers of t in $G(t)$ vanish. This forces $B_{2k+1} = 0$ for $k \geq 1$.

The Bernoulli numbers grow rapidly in absolute value: $|B_{2k}| \sim 2(2k)!/(2\pi)^{2k}$ as $k \rightarrow \infty$. This growth means that the Euler-Maclaurin expansion is typically an *asymptotic* series rather than a convergent one—truncating at a well-chosen index gives an excellent approximation, but including too many terms causes divergence.

Definition 1.5.3: Bernoulli Polynomials

The *Bernoulli polynomials* $B_k(x)$ are defined by

$$B_k(x) = \sum_{j=0}^k \binom{k}{j} B_j x^{k-j},$$

or equivalently by the generating function $\frac{te^{xt}}{e^t-1} = \sum_{k=0}^{\infty} \frac{B_k(x)}{k!} t^k$ for $|t| < 2\pi$.

The Bernoulli polynomials satisfy several identities that make them the right objects for Euler-Maclaurin. The first few are $B_0(x) = 1$, $B_1(x) = x - 1/2$, $B_2(x) = x^2 - x + 1/6$, $B_3(x) = x^3 - 3x^2/2 + x/2$, and $B_4(x) = x^4 - 2x^3 + x^2 - 1/30$. Note $B_k(0) = B_k$ for all k .

Proposition 1.5.4: Properties of Bernoulli Polynomials

For $k \geq 1$:

1. $B'_k(x) = k B_{k-1}(x)$.
2. $\int_0^1 B_k(x) dx = 0$.
3. $B_k(x+1) - B_k(x) = kx^{k-1}$.
4. $B_k(1-x) = (-1)^k B_k(x)$.

Proof:

For (1): differentiate the generating function with respect to x :

$$\frac{t^2 e^{xt}}{e^t - 1} = \sum_{k=0}^{\infty} \frac{B'_k(x)}{k!} t^k$$

The left side equals $t \cdot \frac{te^{xt}}{e^t-1} = \sum_{k=0}^{\infty} \frac{B_k(x)}{k!} t^{k+1} = \sum_{k=1}^{\infty} \frac{k B_{k-1}(x)}{k!} t^k$. Matching coefficients gives $B'_k(x) = k B_{k-1}(x)$ for $k \geq 1$.

For (2): integrate the generating function from $x = 0$ to $x = 1$:

$$\int_0^1 \frac{te^{xt}}{e^t-1} dx = \frac{e^t-1}{e^t-1} = 1$$

Equating this with $\sum_{k=0}^{\infty} \frac{1}{k!} \int_0^1 B_k(x) dx \cdot t^k$ forces the $k = 0$ coefficient to be 1 and all higher coefficients to vanish: $\int_0^1 B_k(x) dx = 0$ for $k \geq 1$.

For (3): the generating function satisfies $\frac{te^{(x+1)t}}{e^t-1} - \frac{te^{xt}}{e^t-1} = te^{xt}$. Matching coefficients of t^k : $\frac{B_k(x+1) - B_k(x)}{k!} = \frac{x^{k-1}}{(k-1)!}$ for $k \geq 1$, giving $B_k(x+1) - B_k(x) = kx^{k-1}$.

For (4): the generating function satisfies $\frac{te^{(1-x)t}}{e^t-1} = \frac{te^{-xt}e^t}{e^t-1}$. Using the identity $\frac{e^t}{e^t-1} = \frac{-1}{e^{-t}-1} \cdot (-1)$,

this equals $\frac{-te^{-xt}}{e^{-t}-1}$, which is the generating function of $B_k(x)$ evaluated at $-t$. Matching coefficients of t^k on both sides: $B_k(1-x)/k! = (-1)^k B_k(x)/k!$, giving $B_k(1-x) = (-1)^k B_k(x)$.

Property (3) is the key identity for Euler-Maclaurin: it expresses the power kx^{k-1} as the discrete difference $B_k(x+1) - B_k(x)$, which allows telescoping sums to be converted into integrals of Bernoulli polynomials. Property (4) gives the reflection symmetry: combined with (1), it shows that $B_{2k+1}(1/2) = 0$ for $k \geq 0$, generalizing the vanishing of odd Bernoulli numbers.

Definition 1.5.5: Periodic Bernoulli Functions

For $k \geq 0$, the *periodic Bernoulli function* $\tilde{B}_k : \mathbb{R} \rightarrow \mathbb{R}$ is

$$\tilde{B}_k(x) = B_k(\{x\})$$

where $\{x\} = x - \lfloor x \rfloor$ is the fractional part of x . The function $\tilde{B}_k$ is periodic with period 1.

For $k = 0$, $\tilde{B}_0(x) = 1$ is constant. For $k = 1$, $\tilde{B}_1(x) = \{x\} - 1/2$, which is the sawtooth function (discontinuous at integers). For $k \geq 2$, $\tilde{B}_k$ is continuous everywhere because $B_k(0) = B_k(1) = B_k$ for $k \geq 2$ (from property (3): $B_k(1) - B_k(0) = k \cdot 0^{k-1} = 0$).

With these tools in place, the Euler-Maclaurin formula follows from repeated integration by parts.

Theorem 1.5.6: Euler-Maclaurin Summation Formula

Let $f \in C^{2p}[a, b]$ with $a, b \in \mathbb{Z}$ and $p \geq 1$. Then

$$\sum_{n=a}^b f(n) = \int_a^b f(t) dt + \frac{f(a) + f(b)}{2} + \sum_{k=1}^p \frac{B_{2k}}{(2k)!} (f^{(2k-1)}(b) - f^{(2k-1)}(a)) + R_p$$

where the remainder term is

$$R_p = - \int_a^b \frac{\tilde{B}_{2p}(t)}{(2p)!} f^{(2p)}(t) dt$$

Proof:

Step 1: The base case. On each unit interval $[n, n+1]$, use $\tilde{B}_1(t) = t - n - 1/2$ for $t \in [n, n+1]$ and note that $\tilde{B}'_1(t) = 1$ on this interval. Then:

$$\int_n^{n+1} f(t) dt = \int_n^{n+1} f(t) \cdot 1 dt = \int_n^{n+1} f(t) d\tilde{B}_1(t)$$

Integration by parts gives

$$\int_n^{n+1} f(t) d\tilde{B}_1(t) = [f(t)\tilde{B}_1(t)]_n^{n+1} - \int_n^{n+1} \tilde{B}_1(t) f'(t) dt$$

Since $\tilde{B}_1(n^+) = -1/2$ and $\tilde{B}_1((n+1)^-) = 1/2$:

$$\int_n^{n+1} f(t) dt = \frac{f(n) + f(n+1)}{2} - \int_n^{n+1} \tilde{B}_1(t) f'(t) dt$$

Summing over $n = a, a + 1, \dots, b - 1$:

$$\int_a^b f(t) dt = \sum_{n=a}^{b-1} \frac{f(n) + f(n+1)}{2} - \int_a^b \tilde{B}_1(t) f'(t) dt = \sum_{n=a}^b f(n) - \frac{f(a) + f(b)}{2} - \int_a^b \tilde{B}_1(t) f'(t) dt$$

Rearranging:

$$\sum_{n=a}^b f(n) = \int_a^b f(t) dt + \frac{f(a) + f(b)}{2} + \int_a^b \tilde{B}_1(t) f'(t) dt$$

This is the formula for $p = 0$ (no Bernoulli correction terms), with remainder $\int \tilde{B}_1 f'$.

Step 2: Iteration. For the remainder integral, apply integration by parts using $\tilde{B}'_{k+1}(t) = (k+1)\tilde{B}_k(t)$ for $t \notin \mathbb{Z}$ (from proposition 1.5.4(1)). Then

$$\int_a^b \tilde{B}_k(t) f^{(k)}(t) dt = \left[\frac{\tilde{B}_{k+1}(t)}{k+1} f^{(k)}(t) \right]_a^b - \int_a^b \frac{\tilde{B}_{k+1}(t)}{k+1} f^{(k+1)}(t) dt$$

At the endpoints $a, b \in \mathbb{Z}$: $\tilde{B}_{k+1}(a) = \tilde{B}_{k+1}(b) = B_{k+1}(0) = B_{k+1}$. Hence

$$\int_a^b \tilde{B}_k(t) f^{(k)}(t) dt = \frac{B_{k+1}}{k+1} (f^{(k)}(b) - f^{(k)}(a)) - \int_a^b \frac{\tilde{B}_{k+1}(t)}{k+1} f^{(k+1)}(t) dt$$

Dividing by $k!$: $\int_a^b \frac{\tilde{B}_k(t)}{k!} f^{(k)}(t) dt = \frac{B_{k+1}}{(k+1)!} (f^{(k)}(b) - f^{(k)}(a)) - \int_a^b \frac{\tilde{B}_{k+1}(t)}{(k+1)!} f^{(k+1)}(t) dt$.

Step 3: Apply iteratively. Starting from $\int \tilde{B}_1 f'$ and applying the above recursion $2p - 1$ times, the odd-indexed Bernoulli numbers $B_3, B_5, \dots$ vanish, contributing nothing to the boundary terms. Only the terms with even index $2k$ survive, giving the stated formula with remainder $R_p = - \int_a^b \frac{\tilde{B}_{2p}(t)}{(2p)!} f^{(2p)}(t) dt$.

The power of the Euler-Maclaurin formula is that it gives an exact identity for any p , not just an asymptotic approximation. The practical question is: how should p be chosen? For smooth decaying f , the remainder R_p is typically small for moderate p but grows rapidly as $p \rightarrow \infty$ (reflecting the rapid growth of $|B_{2p}|$). The optimal p depends on the specific f and is generally the value that minimizes $|R_p|$.

The first major application is to Stirling's formula.

Theorem 1.5.7: Stirling's Formula

For $x > 0$ as $x \rightarrow \infty$:

$$\log \Gamma(x+1) = x \log x - x + \frac{1}{2} \log(2\pi x) + \sum_{k=1}^{p-1} \frac{B_{2k}}{2k(2k-1)} \cdot \frac{1}{x^{2k-1}} + O_p(x^{-2p+1})$$

In particular, the leading-order form is $\log(n!) = n \log n - n + \frac{1}{2} \log(2\pi n) + O(1/n)$, or equivalently $n! \sim \sqrt{2\pi n}(n/e)^n$.

Proof:

Apply theorem 1.5.6 to $f(t) = \log t$ on $[1, N]$ with N a positive integer. The derivatives are $f'(t) = 1/t$, $f''(t) = -1/t^2$, and generally $f^{(k)}(t) = (-1)^{k-1}(k-1)!/t^k$. The integral is $\int_1^N \log t \, dt = N \log N - N + 1$. The boundary terms and Bernoulli corrections give:

$$\sum_{n=1}^N \log n = N \log N - N + 1 + \frac{\log N}{2} + \sum_{k=1}^p \frac{B_{2k}}{(2k)!} (f^{(2k-1)}(N) - f^{(2k-1)}(1)) + R_p$$

With $f^{(2k-1)}(t) = (2k-2)!/t^{2k-1}$:

$$\frac{B_{2k}}{(2k)!} \cdot (2k-2)! \left(\frac{1}{N^{2k-1}} - 1 \right) = \frac{B_{2k}}{2k(2k-1)} \left(\frac{1}{N^{2k-1}} - 1 \right)$$

The sum $\log(N!) = \sum_{n=1}^N \log n$ on the left, and letting $N \rightarrow \infty$ with the correction terms at infinity vanishing, identifies the constant as $\frac{1}{2} \log(2\pi)$. The details of the constant determination use Wallis's product or the duplication formula for Γ ; the result is as stated.

Example 1.5.8: Numerical Stirling

For $n = 10$: the exact value is $10! = 3,628,800$. The leading Stirling approximation $\sqrt{20\pi}(10/e)^{10} \approx 3,598,696$ gives relative error about 0.83%. Including the $B_2/(2x) = 1/(12x)$ correction gives $\sqrt{20\pi}(10/e)^{10} \exp(1/120) \approx 3,628,810$, matching the exact value to 5 significant figures.

A second application gives an asymptotic expansion for the partial sums of n^{-s} refined beyond proposition 1.4.6.

Proposition 1.5.9: Asymptotic Expansion of $\sum n^{-s}$

For $s \in \mathbb{C}$ with $\operatorname{Re}(s) > 0$ and $s \neq 1$, and any integer $p \geq 1$:

$$\sum_{n=1}^N n^{-s} = \frac{N^{1-s}}{1-s} + \zeta(s) + \frac{N^{-s}}{2} + \sum_{k=1}^{p-1} \frac{B_{2k}}{(2k)!} s(s+1) \cdots (s+2k-2) \cdot N^{-s-2k+1} + O_{p,s}(N^{-\operatorname{Re}(s)-2p+1})$$

Proof:

Apply theorem 1.5.6 to $f(t) = t^{-s}$ on $[1, N]$. The integral is $\int_1^N t^{-s} dt = \frac{N^{1-s} - 1}{1-s}$. Derivatives: $f^{(k)}(t) = (-1)^k s(s+1) \cdots (s+k-1) t^{-s-k}$. The boundary terms $(f(1) + f(N))/2 = (1 + N^{-s})/2$ and the Bernoulli corrections combine to give the stated expansion. The constant $\zeta(s)$ emerges from the value at $N = 1$ when $\operatorname{Re}(s) > 1$, and the expansion extends the right-hand side to $\operatorname{Re}(s) > 1 - 2p$ as an analytic function of s .

1.5.10 Foreshadowing: Bernoulli Numbers and Special Values of $\zeta(s)$ The proposition above suggests the following identity. Setting $s = -n$ (a negative integer) formally in the expansion and taking $N = 1$, the boundary and Bernoulli terms combine to yield

$$\zeta(-n) = -\frac{B_{n+1}}{n+1} \quad \text{for } n \geq 1$$

This is not rigorous here: the meromorphic continuation of $\zeta(s)$ to all of $\mathbb{C}$ has not yet been established, and the formal manipulation of the expansion at $s = -n$ requires justification. Both issues are resolved in chapter 6, where the functional equation of $\zeta(s)$ provides the continuation and confirms the identity. In particular: $\zeta(0) = -1/2$, $\zeta(-1) = -1/12$, $\zeta(-3) = 1/120$, and $\zeta(-2k) = 0$ for $k \geq 1$ (the trivial zeros). The famous “identity” $1 + 2 + 3 + \cdots = -1/12$ is a shorthand for $\zeta(-1) = -1/12$.

Example 1.5.11: High-Precision Computation of the Harmonic Sum

To compute $H_{100} = \sum_{n=1}^{100} 1/n$ to high precision, apply theorem 1.5.6 with $f(t) = 1/t$ on $[1, 100]$ and $p = 3$. With $f^{(2k-1)}(t) = -(2k-1)!/t^{2k}$, the Bernoulli term for index $2k$ contributes

$$\frac{B_{2k}}{(2k)!} (f^{(2k-1)}(100) - f^{(2k-1)}(1)) = \frac{B_{2k}}{2k} \left(1 - \frac{1}{100^{2k}} \right)$$

The formula gives:

$$H_{100} = \log 100 + \frac{1 + 1/100}{2} + \frac{1/6}{2} \left(1 - \frac{1}{100^2} \right) + \frac{-1/30}{4} \left(1 - \frac{1}{100^4} \right) + \frac{1/42}{6} \left(1 - \frac{1}{100^6} \right) + R_3$$

Numerically: $\log 100 \approx 4.60517$, boundary = 0.505, $k = 1$ term ≈ 0.08333 , $k = 2$ term ≈ -0.00833 , $k = 3$ term ≈ 0.00397 . The sum is approximately 5.18914, matching the exact value $H_{100} \approx 5.18738$ to within 0.002. The discrepancy is the remainder R_3 , which is of order $|B_6|/(6 \cdot 100^0) \approx 0.004$ (the worst contribution being at the lower endpoint $t = 1$, where the derivatives of $1/t$ are largest).

Exercise 1.5.1

1. For each of the following arithmetic functions, determine whether it is multiplicative, totally multiplicative, additive, totally additive, or none of these. Justify each answer.
 1. $f(n) = n^2$
 2. $f(n) = \log n$
 3. $f(n) = \sigma(n)/n$

$$4. f(n) = \sigma_2(n) = \sum_{d|n} d^2$$

2.
 1. Compute $\Lambda(n)$ for $n = 1, 2, \dots, 12$.
 2. Verify the von Mangoldt identity $\sum_{d|n} \Lambda(d) = \log n$ for $n = 12$ and $n = 18$.
 3. Show that $\Lambda(n) \leq \log n$ for all $n \geq 1$, with equality if and only if n is prime.
3. Prove that σ_k is multiplicative for every $k \in \mathbb{C}$, and derive the formula

$$\sigma_k(p^a) = \frac{p^{(a+1)k} - 1}{p^k - 1}$$

for $k \neq 0$ and any prime p . State the corresponding formula for $k = 0$.

4. Prove that the Liouville function λ is totally multiplicative, and show that $\lambda(n)\mu(n) = \mu(n)^2$ for all $n \geq 1$. Deduce that λ and μ agree on squarefree integers.
5. Compute the Dirichlet convolution $(f * g)(12)$ for:
 1. $f = g = \mathbf{1}$.
 2. $f = \mu, g = \text{id}$.
 3. $f = \lambda, g = \mathbf{1}$.

Compare each result with the value predicted by the convolution identities in proposition 1.2.10.

6. Using the recursive formula for Dirichlet inverses, compute $\sigma^{-1}(n)$ for $n = 1, 2, 3, 4, 6$.
7. Prove the *multiplicative* version of Möbius inversion: if $F(n) = \prod_{d|n} f(d)$ for all $n \geq 1$, then $f(n) = \prod_{d|n} F(d)^{\mu(n/d)}$.
8. Prove the identity $d * \varphi = \sigma$. [Hint: use $d = \mathbf{1} * \mathbf{1}$ and $\varphi = \text{id} * \mu$.]
9. Let f be a multiplicative function. Show that f^{-1} (the Dirichlet inverse of f) is also multiplicative, and prove that for any prime p ,

$$f^{-1}(p^a) = \sum_{k=1}^a (-1)^k \sum_{\substack{a_1 + \dots + a_k = a \\ a_i \geq 1}} f(p^{a_1}) \cdots f(p^{a_k})$$

10. Using the hyperbola method, provide an independent derivation of the estimate $\sum_{n \leq x} d(n) = x \log x + O(x)$. Then verify numerically for $x = 20$: compute the exact sum $\sum_{n=1}^{20} d(n)$ and compare with $20 \log 20 + (2\gamma - 1) \cdot 20$.
11. Let $Q(x) = \sum_{n \leq x} |\mu(n)|$ count the squarefree integers up to x . Using the hyperbola method applied to the identity $|\mu(n)| = \sum_{d^2|n} \mu(d)$, give a self-contained proof that $Q(x) = 6x/\pi^2 + O(\sqrt{x})$.
12. Prove that

$$\sum_{\substack{n \leq x \\ n \text{ squarefree}}} \frac{1}{n} = \frac{6}{\pi^2} \log x + C + O(x^{-1/2})$$

for a constant C . [Hint: use Abel summation with $a_n = |\mu(n)|$ and $f(t) = 1/t$, then apply exercise 1.5.1 (11).]

13. Prove the trivial bound $|M(x)| = |\sum_{n \leq x} \mu(n)| \leq x$ for all $x \geq 1$. Then, using the identity $\sum_{d \leq x} M(x/d) = 1$, derive the recursion-based estimate $M(x) = 1 - \sum_{2 \leq d \leq x} M(x/d)$ and explain why this does not immediately yield $M(x) = o(x)$.
14. Using Abel summation and the estimate $\sum_{n \leq x} \sigma(n) = \frac{\pi^2}{12}x^2 + O(x \log x)$, prove that

$$\sum_{n \leq x} \frac{\sigma(n)}{n} = \frac{\pi^2}{6}x + O((\log x)^2)$$

15. Using Abel summation and $\sum_{n \leq x} \varphi(n) = \frac{3}{\pi^2}x^2 + O(x \log x)$, show that

$$\sum_{n \leq x} \frac{\varphi(n)}{n^2} = \frac{6}{\pi^2} \log x + C + O\left(\frac{\log x}{x}\right)$$

for a constant C . (An explicit formula for C is not required.)

16. Let a_n satisfy $|A(N)| = |\sum_{n \leq N} a_n| \leq M$ for all N . Use Abel summation to prove that $\sum_{n=1}^{\infty} a_n/n^s$ converges for every $s \in \mathbb{C}$ with $\operatorname{Re}(s) > 0$, and the convergence is uniform on compact subsets of $\{\operatorname{Re}(s) > 0\}$.
17. Using Abel summation with $a_n = 1$ and $f(t) = (\log t)/t$, prove that

$$\sum_{n \leq x} \frac{\log n}{n} = \frac{(\log x)^2}{2} + C_1 + O\left(\frac{\log x}{x}\right)$$

for a constant C_1 . (An explicit value for C_1 is not required.)

18. Using the recursion $\sum_{k=0}^n \binom{n+1}{k} B_k = 0$ for $n \geq 1$, compute B_6 , B_8 , and B_{10} directly from the values $B_0 = 1$, $B_1 = -1/2$, $B_2 = 1/6$, $B_4 = -1/30$, and the vanishing of odd-indexed B_k for $k \geq 3$.
19.
 1. Compute the Bernoulli polynomials $B_2(x)$, $B_3(x)$, and $B_4(x)$ from the formula $B_k(x) = \sum_{j=0}^k \binom{k}{j} B_j x^{k-j}$.
 2. Verify $B_k(1) = B_k(0)$ for $k \in \{2, 3, 4\}$.
 3. Use the telescoping identity $B_3(x+1) - B_3(x) = 3x^2$ to derive the closed form $\sum_{j=0}^{N-1} j^2 = N(N-1)(2N-1)/6$.
20. Apply the Euler-Maclaurin formula to $f(t) = 1/t^2$ on $[1, N]$ with $p = 2$ (two Bernoulli correction terms). Extract the asymptotic expansion for $\sum_{n=1}^N 1/n^2$ and identify the N -independent constant as $\zeta(2) = \pi^2/6$.
21. The functional equation of $\zeta(s)$ implies $\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k} B_{2k}}{2(2k)!}$ for $k \geq 1$ (this will be proved in chapter 6). Verify this identity for $k = 1$ (giving $\zeta(2) = \pi^2/6$) and $k = 2$ (giving $\zeta(4) = \pi^4/90$). What does the formula give for $k = 3$?

Dirichlet Series and Euler Products

Chapter 1 organized the arithmetic functions into a ring under Dirichlet convolution and extracted asymptotic information about their summatory functions through summation techniques. The present chapter introduces the analytic objects that encode the same information in a fundamentally different way: *Dirichlet series*. The Dirichlet series $D(f, s) = \sum_{n=1}^{\infty} f(n)/n^s$ associates to each arithmetic function f an analytic function of a complex variable s , and this association converts Dirichlet convolution into pointwise multiplication, multiplicativity into an Euler product over primes, and the summatory function into a contour integral. The poles of the Dirichlet series control the main terms in asymptotic estimates; its zeros govern the oscillatory corrections; and its analytic continuation beyond the region of absolute convergence is the mechanism by which the distribution of primes becomes accessible to the methods of complex analysis.

This chapter builds the dictionary between the algebraic side (arithmetic functions, convolution, multiplicativity) and the analytic side (Dirichlet series, products, Mellin transforms). Sections 2.1-2.3 establish the algebraic-analytic correspondence: convergence, the multiplication theorem, and Euler products. Section 2.4 develops the Mellin transform as a self-contained tool, and Section 2.5 proves Perron's formula, which inverts the Dirichlet series back into information about partial sums. Together, these tools are the analytic engine of the book.

2.1 Dirichlet Series: Convergence and Analyticity

An arithmetic function f is a sequence of complex numbers; its Dirichlet series is the analytic function obtained by summing $f(n)/n^s$ over all positive integers. The first question is: for which values of the complex variable s does the series converge? The answer has a clean geometric structure: there is a

critical vertical line in the complex plane, to the right of which the series converges and to the left of which it diverges. This *abscissa of convergence* is the Dirichlet series analogue of the radius of convergence for a power series, and it is determined by the growth rate of the partial sums of the coefficients.

Definition 2.1.1: Dirichlet Series

A *Dirichlet series* is a series of the form

$$D(f, s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s}$$

where $f : \mathbb{Z}_{>0} \rightarrow \mathbb{C}$ is an arithmetic function and $s \in \mathbb{C}$.

The variable s is conventionally written as $s = \sigma + it$ with $\sigma = \operatorname{Re}(s)$ and $t = \operatorname{Im}(s)$. Since $n^s = n^\sigma \cdot n^{it} = n^\sigma e^{it \log n}$, the factor $n^{-\sigma}$ controls convergence while n^{-it} contributes oscillation without affecting absolute convergence. This separation between the real and imaginary parts of s underlies the half-plane structure of the convergence region.

The fundamental convergence theorem is the following.

Theorem 2.1.2: Half-Plane of Convergence

Let $D(f, s) = \sum f(n)/n^s$ be a Dirichlet series. If $D(f, s_0)$ converges for some $s_0 \in \mathbb{C}$, then $D(f, s)$ converges for all s with $\operatorname{Re}(s) > \operatorname{Re}(s_0)$, and the convergence is uniform on compact subsets of this half-plane.

Proof:

Write $\sigma_0 = \operatorname{Re}(s_0)$ and fix s with $\sigma = \operatorname{Re}(s) > \sigma_0$. Set $a_n = f(n)/n^{s_0}$ and $g(t) = t^{s_0-s}$, so that $f(n)/n^s = a_n \cdot n^{s_0-s}$. By hypothesis, $A(N) = \sum_{n \leq N} a_n$ is bounded (as the partial sums of a convergent series). The function $g(t) = t^{s_0-s}$ has $|g(t)| = t^{\sigma_0-\sigma} \rightarrow 0$ as $t \rightarrow \infty$ (since $\sigma_0 - \sigma < 0$), and $g'(t) = (s_0 - s)t^{s_0-s-1}$.

By Abel summation (theorem 1.4.1):

$$\sum_{n=1}^N \frac{f(n)}{n^s} = \sum_{n=1}^N a_n \cdot n^{s_0-s} = A(N) \cdot N^{s_0-s} - \int_1^N A(t)(s_0 - s)t^{s_0-s-1} dt$$

As $N \rightarrow \infty$: $|A(N) \cdot N^{s_0-s}| \leq M \cdot N^{\sigma_0-\sigma} \rightarrow 0$, where $M = \sup_N |A(N)|$. The integral converges absolutely:

$$\int_1^\infty |A(t)| |s_0 - s| t^{\sigma_0-\sigma-1} dt \leq M |s_0 - s| \int_1^\infty t^{\sigma_0-\sigma-1} dt = \frac{M |s_0 - s|}{\sigma - \sigma_0} < \infty$$

Hence $D(f, s)$ converges.

For uniform convergence on a compact set $K \subset \{\operatorname{Re}(s) > \sigma_0\}$: let $\sigma_1 = \min_{s \in K} \operatorname{Re}(s) > \sigma_0$. The same bound gives $|\sum_{n > N} f(n)/n^s| \leq M |s_0 - s| N^{\sigma_0-\sigma_1} / (\sigma_1 - \sigma_0) + M N^{\sigma_0-\sigma_1}$, and $|s_0 - s|$ is bounded on K , so the tail tends to 0 uniformly.

This theorem shows that the region of convergence of a Dirichlet series is always a right half-plane (or the entire complex plane, or the empty set). The boundary of this half-plane defines the abscissa of convergence.

Definition 2.1.3: Abscissae of Convergence and Absolute Convergence

For a Dirichlet series $D(f, s)$:

1. The *abscissa of convergence* σ_c is the infimum of the set of $\sigma \in \mathbb{R}$ such that $D(f, \sigma + it)$ converges for all t . Equivalently, $D(f, s)$ converges for $\operatorname{Re}(s) > \sigma_c$ and diverges for $\operatorname{Re}(s) < \sigma_c$.
2. The *abscissa of absolute convergence* σ_a is the infimum of the set of σ such that $\sum |f(n)|/n^\sigma < \infty$.

By convention, $\sigma_c = -\infty$ if the series converges everywhere, $\sigma_c = +\infty$ if it converges nowhere, and similarly for σ_a .

By definition, $\sigma_c \leq \sigma_a$ (absolute convergence implies convergence). The gap between σ_c and σ_a reflects the amount of cancellation in the coefficients.

Proposition 2.1.4: Gap Between Abscissae

For any Dirichlet series: $\sigma_c \leq \sigma_a \leq \sigma_c + 1$. Both inequalities can be equalities.

Proof:

The inequality $\sigma_c \leq \sigma_a$ is immediate. For the upper bound: if $D(f, s_0)$ converges at s_0 with $\operatorname{Re}(s_0) = \sigma_0$, then $f(n)/n^{s_0} \rightarrow 0$, so $|f(n)| \leq Cn^{\sigma_0}$ for some constant C . Hence $\sum |f(n)|/n^\sigma \leq C \sum n^{\sigma_0 - \sigma}$, which converges for $\sigma > \sigma_0 + 1$. Taking the infimum over convergent s_0 gives $\sigma_a \leq \sigma_c + 1$.

Equality $\sigma_c = \sigma_a$ holds for $\zeta(s) = \sum 1/n^s$ (both equal 1, since the coefficients are non-negative). Equality $\sigma_a = \sigma_c + 1$ holds for the alternating series $\sum (-1)^{n+1}/n^s = (1 - 2^{1-s})\zeta(s)$: here $\sigma_c = 0$ (by Abel's test, since the partial sums of $(-1)^{n+1}$ are bounded), while $\sigma_a = 1$.

The gap between σ_c and σ_a measures the cancellation among the coefficients $f(n)$. For non-negative coefficients, there is no cancellation, and $\sigma_c = \sigma_a$. For coefficients with heavy sign alternation (like $(-1)^{n+1}$), the gap can reach its maximum value of 1. The intermediate case $0 < \sigma_a - \sigma_c < 1$ occurs for the Dirichlet series $\sum \mu(n)/n^s$: the cancellation in the Möbius function is sufficient to push σ_c below $\sigma_a = 1$, but determining σ_c precisely is equivalent to determining the supremum of the real parts of the zeros of $\zeta(s)$.

More generally, whenever the coefficients $f(n)$ are non-negative, the abscissa of convergence equals the abscissa of absolute convergence, and the Dirichlet series $D(f, s)$ has a singularity (non-removable) at the point $s = \sigma_a$ on the real axis. This is the Landau-Vivanti theorem for Dirichlet series.

Proposition 2.1.5: Landau's Theorem on Non-Negative Coefficients

If $f(n) \geq 0$ for all n , then $\sigma_c = \sigma_a$, and $D(f, s)$ has a singularity at $s = \sigma_a$.

Proof:

Since $f(n) \geq 0$: $|f(n)|/n^\sigma = f(n)/n^\sigma$, so $\sigma_c = \sigma_a$. For the singularity: suppose for contradiction that $D(f, s)$ extends analytically to a neighborhood of σ_a . Then the Taylor expansion of $D(f, s)$ at some point $s_1 > \sigma_a$ would converge in a disk centered at s_1 reaching past σ_a , giving

$$D(f, s) = \sum_{k=0}^{\infty} \frac{D^{(k)}(f, s_1)}{k!} (s - s_1)^k = \sum_{k=0}^{\infty} \frac{(s_1 - s)^k}{k!} \sum_{n=1}^{\infty} \frac{f(n)(\log n)^k}{n^{s_1}}$$

for $|s - s_1| < s_1 - \sigma_a + \varepsilon$. Since $f(n) \geq 0$, the double sum has non-negative terms, and Tonelli's theorem allows swapping the summations. At a real point $s_2 < \sigma_a$ inside the disk:

$$D(f, s_2) = \sum_{n=1}^{\infty} \frac{f(n)}{n^{s_1}} \sum_{k=0}^{\infty} \frac{(s_1 - s_2)^k (\log n)^k}{k!} = \sum_{n=1}^{\infty} \frac{f(n)}{n^{s_1}} n^{s_1 - s_2} = \sum_{n=1}^{\infty} \frac{f(n)}{n^{s_2}},$$

so the Dirichlet series converges at $s_2 < \sigma_a$, contradicting the definition of σ_a .

For $\zeta(s) = \sum 1/n^s$, this theorem guarantees a singularity at $s = 1$, consistent with the pole of $\zeta(s)$ at $s = 1$ established by the integral representation of section 1.4. For $\sum d(n)/n^s = \zeta(s)^2$, the theorem gives a singularity at $s = 1$ (a double pole). The theorem does not determine the *nature* of the singularity (pole, branch point, essential singularity), only its existence.

The abscissae can be computed from the coefficients via the following formulas.

Proposition 2.1.6: Formulas for the Abscissae

Let $A(x) = \sum_{n \leq x} f(n)$ and $A^*(x) = \sum_{n \leq x} |f(n)|$. Then:

$$\sigma_c = \limsup_{x \rightarrow \infty} \frac{\log |A(x)|}{\log x}, \quad \sigma_a = \limsup_{x \rightarrow \infty} \frac{\log A^*(x)}{\log x}$$

(With the convention that $\sigma_c = -\infty$ if $A(x)$ is eventually zero, etc.)

Proof:

For σ_a : if $\sigma > \limsup_{x \rightarrow \infty} \frac{\log A^*(x)}{\log x}$, then $A^*(x) \leq x^{\sigma - \varepsilon}$ for large x and some $\varepsilon > 0$. Abel summation applied to $\sum |f(n)|/n^\sigma$ with partial sums $A^*(x)$ gives convergence, since the integral $\int_1^\infty A^*(t)/t^{\sigma+1} dt \leq \int_1^\infty t^{-1-\varepsilon} dt < \infty$. Conversely, if $\sigma < \limsup \log A^*(x)/\log x$, then $A^*(x) \geq x^{\sigma + \varepsilon}$ for infinitely many x , and the series diverges. The proof for σ_c is analogous, replacing $A^*(x)$ with $A(x)$ and using Abel summation for conditional convergence.

The behavior at the boundary $\operatorname{Re}(s) = \sigma_c$ is delicate. The series may converge at every point of this line, at no point, or at some points but not others. For $\zeta(s)$, the boundary is $\operatorname{Re}(s) = 1$, and the series diverges at $s = 1$ but converges conditionally at every other point $1 + it$ with $t \neq 0$ (by Dirichlet's test, since the partial sums of n^{-it} are bounded for $t \neq 0$ and $1/n$ decreases to 0). For

the eta function $\eta(s)$, the boundary is $\operatorname{Re}(s) = 0$, and the series diverges at $s = 0$ by oscillation but converges at $s = it$ for $t \neq 0$.

A final remark on the strip $\sigma_c < \operatorname{Re}(s) < \sigma_a$ (when it is nonempty). In this region, the Dirichlet series converges but not absolutely. The convergence is conditional, and the series does not converge uniformly on all of $\{\operatorname{Re}(s) > \sigma_c\}$ (only on compact subsets). This strip is analogous to the annulus of conditional convergence for Laurent series. The arithmetic content of this strip is often significant: for $\sum \mu(n)/n^s$, the identity $1/\zeta(s) = \sum \mu(n)/n^s$ extends into the strip $\sigma_c < \operatorname{Re}(s) \leq 1$, and the non-vanishing of $\zeta(s)$ on the line $\operatorname{Re}(s) = 1$ is what makes the extension to $\sigma_c < 1$ possible.

A Dirichlet series is not just convergent in its half-plane—it is analytic there, and can be differentiated term by term.

Theorem 2.1.7: Analyticity and Term-by-Term Differentiation

A Dirichlet series $D(f, s) = \sum f(n)/n^s$ is an analytic function on $\{\operatorname{Re}(s) > \sigma_c\}$, and for all $k \geq 1$:

$$D^{(k)}(f, s) = \sum_{n=1}^{\infty} \frac{f(n)(-\log n)^k}{n^s}$$

in the same half-plane. This series also has abscissa of convergence σ_c .

Proof:

Fix a compact set $K \subset \{\operatorname{Re}(s) > \sigma_c\}$ with $\sigma_1 = \min_{s \in K} \operatorname{Re}(s) > \sigma_c$. Choose σ_0 with $\sigma_c < \sigma_0 < \sigma_1$. The partial sums $A(N) = \sum_{n \leq N} f(n)/n^{\sigma_0}$ are bounded for $\operatorname{Re}(s_0) = \sigma_0$, and by the proof of theorem 2.1.2, $D(f, s)$ converges uniformly on K . Since each term $f(n)/n^s$ is an entire function of s , and the series converges uniformly on compact subsets, the sum is analytic on $\{\operatorname{Re}(s) > \sigma_c\}$ (by Weierstrass's theorem on uniformly convergent series of analytic functions, [16]).

For differentiation: $\frac{d}{ds}(f(n)/n^s) = -f(n) \log n / n^s$. The series $\sum f(n) \log n / n^s$ has the same abscissa of convergence as $\sum f(n)/n^s$: the factor $\log n$ grows slower than any power n^ε , so for $\operatorname{Re}(s) > \sigma_c + \varepsilon$, the additional factor $\log n$ does not affect absolute convergence, and letting $\varepsilon \rightarrow 0$ shows the abscissa is unchanged. Uniform convergence on compacta allows exchanging the derivative and the sum.

The analyticity theorem has an important consequence: a Dirichlet series is determined by its coefficients.

Corollary 2.1.8: Uniqueness of Coefficients

If $D(f, s) = D(g, s)$ for all s in some half-plane $\{\operatorname{Re}(s) > \sigma_0\}$, then $f(n) = g(n)$ for all n .

Proof:

Set $h = f - g$, so $D(h, s) = 0$ for $\operatorname{Re}(s) > \sigma_0$. Restricting to real $s = \sigma > \sigma_0$: $D(h, \sigma) = \sum_{n=1}^{\infty} h(n)/n^\sigma = 0$. As $\sigma \rightarrow +\infty$, the terms $h(n)/n^\sigma$ decay at different rates (faster for larger n), and the dominant term is $h(1)$. Formally: dividing by $1^{-\sigma} = 1$ and taking $\sigma \rightarrow \infty$, the terms with $n \geq 2$ vanish, giving $h(1) = 0$. Now $0 = \sum_{n \geq 2} h(n)/n^\sigma$; multiplying by 2^σ and taking

$\sigma \rightarrow \infty$ shows $h(2) = 0$. Inductively, $h(n) = 0$ for all n .

This uniqueness theorem is the Dirichlet series analogue of the fact that a power series is determined by its coefficients. The proof exploits a feature specific to Dirichlet series: the “basis functions” n^{-s} decay at *different rates* as $\sigma \rightarrow \infty$, allowing the coefficients to be peeled off one at a time. For power series $\sum a_n z^n$, the analogous statement requires evaluating derivatives at $z = 0$; for Dirichlet series, the multiplicative structure of the basis $\{n^{-s}\}$ gives a simpler argument.

Example 2.1.9: Abscissae of Standard Dirichlet Series

1. $\zeta(s) = \sum 1/n^s$: since the coefficients are non-negative, proposition 2.1.5 gives $\sigma_c = \sigma_a$. The harmonic series $\sum 1/n$ diverges, so $\sigma_a \geq 1$. For $\sigma > 1$: $\sum 1/n^\sigma \leq 1 + \int_1^\infty t^{-\sigma} dt = \sigma/(\sigma - 1) < \infty$. Hence $\sigma_c = \sigma_a = 1$.
2. $\sum d(n)/n^s$: the divisor function satisfies $d(n) \geq 1$ for all n and $d(n) \leq C_\varepsilon n^\varepsilon$ for any $\varepsilon > 0$ (a proof appears in chapter 4), so $\sigma_a = 1$. The identity $\sum d(n)/n^s = \zeta(s)^2$ (proved in the next section) shows the series has a double pole at $s = 1$.
3. $\sum \Lambda(n)/n^s$: since $0 \leq \Lambda(n) \leq \log n \leq n^\varepsilon$ for any $\varepsilon > 0$, the abscissa $\sigma_a = 1$. By proposition 2.1.5, $\sigma_c = 1$ as well, and there is a singularity at $s = 1$. The identity $-\zeta'(s)/\zeta(s) = \sum \Lambda(n)/n^s$ (§2.2) identifies this singularity as a simple pole with residue 1.
4. $\sum \mu(n)/n^s$: since $|\mu(n)| \leq 1$, $\sigma_a = 1$. The exact value of σ_c is a major open problem. The bound $\sigma_c \leq 1$ follows from the Prime Number Theorem ($M(x) = o(x)$ implies convergence for $\operatorname{Re}(s) > 1$ by Abel summation). The Riemann Hypothesis is equivalent to $\sigma_c = 1/2$; the best unconditional result, $\sigma_c \leq 1$, does not pin down the exact value.

The term-by-term differentiation identity $D'(f, s) = D(f \cdot (-\log), s)$ has an important interpretation: differentiating a Dirichlet series with respect to s multiplies the coefficients by $-\log n$. This connects differentiation to the von Mangoldt function, since $-\zeta'(s) = \sum (\log n)/n^s$ is the Dirichlet series of $\log$, and dividing by $\zeta(s)$ gives $-\zeta'(s)/\zeta(s) = D(\Lambda, s)$ (the logarithmic derivative encodes the von Mangoldt function). More generally, for any Dirichlet series $D(f, s)$ with f multiplicative and $D(f, s) \neq 0$, the logarithmic derivative $-D'(f, s)/D(f, s)$ is the Dirichlet series of an arithmetic function whose values at prime powers are determined by those of f . This mechanism is developed systematically in the next section.

Example 2.1.10: A Dirichlet Series with $\sigma_c < \sigma_a$

The Dirichlet eta function $\eta(s) = \sum_{n=1}^\infty (-1)^{n+1}/n^s$ has $\sigma_a = 1$ and $\sigma_c = 0$. The partial sums $A(N) = \sum_{n=1}^N (-1)^{n+1}$ alternate between 1 and 0, so $|A(N)| \leq 1$ for all N , and Abel’s test (corollary 1.4.9) gives convergence for $\operatorname{Re}(s) > 0$. At $s = 0$: $\sum (-1)^{n+1} = 1 - 1 + 1 - \dots$ diverges by oscillation, confirming $\sigma_c = 0$.

The identity $\eta(s) = (1 - 2^{1-s})\zeta(s)$ is verified by writing $\eta(s) = \sum 1/n^s - 2 \sum 1/(2n)^s = \zeta(s) - 2^{1-s}\zeta(s)$. The factor $(1 - 2^{1-s})$ vanishes at $s = 1 + 2\pi i k / \log 2$ for $k \in \mathbb{Z}$, cancelling the pole of $\zeta(s)$ at $s = 1$ when $k = 0$. In particular, $\eta(1) = 1 - 1/2 + 1/3 - 1/4 + \dots = \log 2$, and this value can also be obtained as $\lim_{s \rightarrow 1} (1 - 2^{1-s})\zeta(s) = \log 2$ (using $\zeta(s) \sim 1/(s - 1)$ and $(1 - 2^{1-s}) \sim (s - 1) \log 2$ near $s = 1$).

The convergence theory of Dirichlet series should be compared with that of power series. A power

series $\sum a_n z^n$ converges in a *disk* $|z| < R$, determined by a single parameter R (the radius of convergence). A Dirichlet series $\sum a_n n^{-s}$ converges in a *half-plane* $\operatorname{Re}(s) > \sigma_c$, also determined by a single parameter. The disk is replaced by a half-plane because $|n^{-s}| = n^{-\sigma}$ depends only on $\operatorname{Re}(s)$, not on $\operatorname{Im}(s)$: changing the imaginary part of s rotates each term n^{-it} on the unit circle without affecting convergence. The analogy extends further: the Hadamard formula $1/R = \limsup |a_n|^{1/n}$ for the radius of convergence corresponds to the formula $\sigma_a = \limsup (\log A^*(x)) / \log x$ for the abscissa of absolute convergence.

Example 2.1.11: Dirichlet Series That Converge Everywhere or Nowhere

1. The Dirichlet series $D(s) = \sum_{n=1}^N a_n/n^s$ (a finite sum) is an entire function of s , with $\sigma_c = \sigma_a = -\infty$.
2. The series $\sum n!/n^s$ has $\sigma_c = \sigma_a = +\infty$: the terms $n!/n^s$ grow faster than any power of n , so the series diverges for all s .
3. The series $\sum 1/(n^s \log^2 n)$ (starting from $n = 2$) converges absolutely for $\operatorname{Re}(s) \geq 1$ (since $\sum 1/(n \log^2 n) < \infty$). Hence $\sigma_a \leq 1$. Since the coefficients are positive, $\sigma_c = \sigma_a$; and $\sum 1/(n^\sigma \log^2 n)$ diverges for $\sigma < 1$ (by comparison with $\sum 1/n$), so $\sigma_a = 1$ exactly. This series converges on the *closed* half-plane $\operatorname{Re}(s) \geq 1$, unlike $\zeta(s)$ which diverges at $s = 1$.

2.2 The Ring Homomorphism: Convolution and Multiplication

The previous section developed Dirichlet series as analytic functions on a half-plane. The next structural fact is that the operation of *Dirichlet convolution* on arithmetic functions, introduced in section 1.2, corresponds to ordinary *multiplication* of the associated Dirichlet series. This single identity is the algebraic engine of analytic number theory: every standard convolution identity from chapter 1 translates into an identity among Dirichlet series, and the convolution ring $(\mathcal{A}, +, *)$ realizes concretely as a ring of analytic functions on a half-plane. The isomorphism is also the route by which the von Mangoldt function Λ enters the analytic theory through the logarithmic derivative $-\zeta'(s)/\zeta(s)$, the identity that drives the analytic proof of the Prime Number Theorem in chapter 5. The crux is the following.

Theorem 2.2.1: Dirichlet Multiplication

Let f, g be arithmetic functions, and let $\sigma^* = \max(\sigma_a(f), \sigma_a(g))$. For $\operatorname{Re}(s) > \sigma^*$, the series $D(f, s)$, $D(g, s)$, and $D(f * g, s)$ all converge absolutely, and

$$D(f * g, s) = D(f, s) \cdot D(g, s)$$

Proof:

Fix s with $\sigma = \operatorname{Re}(s) > \sigma^*$. By the definition of σ^* , both $\sum |f(m)|/m^\sigma$ and $\sum |g(k)|/k^\sigma$ converge.

Step 1: Form the absolutely convergent product series. The product

$$\left(\sum_{m=1}^{\infty} \frac{f(m)}{m^s} \right) \left(\sum_{k=1}^{\infty} \frac{g(k)}{k^s} \right) = \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{f(m)g(k)}{(mk)^s}$$

converges absolutely, since the absolute value of the double sum is bounded by the product of the absolute series.

Step 2: Group terms by $n = mk$. For each $n \geq 1$, the pairs $(m, k) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0}$ with $mk = n$ are in bijection with the divisors d of n via $(m, k) = (d, n/d)$. Hence

$$\sum_{m,k} \frac{f(m)g(k)}{(mk)^s} = \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{d|n} f(d)g(n/d) = \sum_{n=1}^{\infty} \frac{(f * g)(n)}{n^s}$$

by the definition of Dirichlet convolution. The interchange of summation order is justified by the absolute convergence established in Step 1 (Tonelli's theorem applied to the counting measure on $\mathbb{Z}_{>0} \times \mathbb{Z}_{>0}$).

*Step 3: Absolute convergence of $D(f * g, s)$.* The same argument with $|f|$ and $|g|$ in place of f and g gives

$$\sum_{n=1}^{\infty} \frac{|(f * g)(n)|}{n^\sigma} \leq \sum_{n=1}^{\infty} \frac{(|f| * |g|)(n)}{n^\sigma} = \left(\sum_{m=1}^{\infty} \frac{|f(m)|}{m^\sigma} \right) \left(\sum_{k=1}^{\infty} \frac{|g(k)|}{k^\sigma} \right) < \infty$$

completing the proof.

The proof says something more than the bare identity. The same rearrangement, applied to $|f|$ and $|g|$, also bounds the abscissa of absolute convergence of the convolution: $\sigma_a(f * g) \leq \max(\sigma_a(f), \sigma_a(g))$. Equality need not hold (cancellation in $f * g$ can shift the abscissa downward), but the inequality places $D(f * g, s)$ on a half-plane no smaller than the common half-plane of $D(f, s)$ and $D(g, s)$.

The theorem converts every convolution identity from section 1.2 into a Dirichlet series identity. The corollary collects the most-used cases.

Corollary 2.2.2: Standard Identities from Convolution

For $\operatorname{Re}(s)$ in the appropriate half-plane:

1. $\zeta(s)^2 = \sum_{n=1}^{\infty} \frac{d(n)}{n^s}$ for $\operatorname{Re}(s) > 1$, from $d = \mathbf{1} * \mathbf{1}$.
2. $\zeta(s) \zeta(s-k) = \sum_{n=1}^{\infty} \frac{\sigma_k(n)}{n^s}$ for $\operatorname{Re}(s) > \max(1, k+1)$, from $\sigma_k = \operatorname{id}_k * \mathbf{1}$, where $\operatorname{id}_k(n) = n^k$.
3. $\frac{1}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s}$ for $\operatorname{Re}(s) > 1$, from $\mu * \mathbf{1} = \varepsilon$.
4. $\sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s} = \frac{\zeta(s-1)}{\zeta(s)}$ for $\operatorname{Re}(s) > 2$, from $\varphi * \mathbf{1} = \operatorname{id}$.
5. $\frac{\zeta(2s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s}$ for $\operatorname{Re}(s) > 1$, from $\lambda * |\mu| = \varepsilon$ together with $\sum |\mu(n)|/n^s = \zeta(s)/\zeta(2s)$.

Proof:

Each identity follows from theorem 2.2.1 together with the elementary computations $D(\varepsilon, s) = 1$, $D(\mathbf{1}, s) = \zeta(s)$, and $D(\operatorname{id}_k, s) = \sum n^k/n^s = \zeta(s-k)$ on $\operatorname{Re}(s) > k+1$.

Identity (1): $D(d, s) = D(\mathbf{1} * \mathbf{1}, s) = \zeta(s) \zeta(s) = \zeta(s)^2$.

Identity (2): $D(\sigma_k, s) = D(\operatorname{id}_k * \mathbf{1}, s) = D(\operatorname{id}_k, s) D(\mathbf{1}, s) = \zeta(s-k) \zeta(s)$.

Identity (3): $D(\varepsilon, s) = 1 = D(\mu * \mathbf{1}, s) = D(\mu, s) \zeta(s)$, so $D(\mu, s) = 1/\zeta(s)$.

Identity (4): $D(\operatorname{id}, s) = \zeta(s-1) = D(\varphi * \mathbf{1}, s) = D(\varphi, s) \zeta(s)$, so $D(\varphi, s) = \zeta(s-1)/\zeta(s)$.

Identity (5): The squarefree indicator $|\mu|$ satisfies $|\mu| * \mathbf{1}_{\square} = \mathbf{1}$, where $\mathbf{1}_{\square}(n) = 1$ if n is a square and 0 otherwise. Hence $D(|\mu|, s) D(\mathbf{1}_{\square}, s) = \zeta(s)$. Since $D(\mathbf{1}_{\square}, s) = \sum_{m \geq 1} 1/(m^2)^s = \zeta(2s)$, this gives $D(|\mu|, s) = \zeta(s)/\zeta(2s)$. The Liouville identity $\lambda * |\mu| = \varepsilon$ (proposition 1.2.10) then gives $D(\lambda, s) \cdot \zeta(s)/\zeta(2s) = 1$, i.e., $D(\lambda, s) = \zeta(2s)/\zeta(s)$, as claimed.

These identities form the fundamental dictionary at the level of $\zeta(s)$. Identity (1) makes the divisor function $d(n)$ accessible to analytic methods through $\zeta(s)^2$, whose double pole at $s = 1$ is the analytic source of the leading $x \log x$ term in the asymptotic $\sum_{n \leq x} d(n) \sim x \log x$ from section 1.3. Identity (2) generalizes this to all σ_k . Identity (3) inverts the role of the Möbius function at the analytic level: the reciprocal of the zeta function is the Dirichlet series of μ , linking the analytic behavior of $\zeta(s)$ to summatory questions about $\mu(n)$ and ultimately to the Prime Number Theorem and the Riemann Hypothesis. Identity (4) gives the Dirichlet series of the Euler totient as a ratio of zeta functions; its

2.2. The Ring Homomorphism: Convolution and Multiplication Dirichlet Series and Euler Products

residue at $s = 2$ is $1/\zeta(2) = 6/\pi^2$, the asymptotic density of coprime pairs (section 1.3). Identity (5) is the Liouville series, whose ratio form $\zeta(2s)/\zeta(s)$ encodes the multiplicative structure of squarefree integers analytically.

The first identity admits a direct verification that makes the convolution-multiplication mechanism transparent.

Example 2.2.3: Verifying $\zeta(s)^2 = \sum d(n)/n^s$ by Squaring

Squaring the series for $\zeta(s)$ on $\operatorname{Re}(s) > 1$:

$$\zeta(s)^2 = \left(\sum_{m=1}^{\infty} \frac{1}{m^s} \right) \left(\sum_{k=1}^{\infty} \frac{1}{k^s} \right) = \sum_{m,k \geq 1} \frac{1}{(mk)^s} = \sum_{n=1}^{\infty} \frac{c(n)}{n^s}$$

where $c(n) = \#\{(m, k) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0} : mk = n\}$. The pairs (m, k) with product n are in bijection with the divisors d of n via $(m, k) = (d, n/d)$, so $c(n) = d(n)$ exactly.

The first few coefficients confirm the count: $d(1) = 1$, $d(2) = d(3) = d(5) = d(7) = 2$, $d(4) = d(9) = 3$, $d(6) = d(8) = d(10) = 4$, $d(12) = 6$. Hence

$$\zeta(s)^2 = 1 + \frac{2}{2^s} + \frac{2}{3^s} + \frac{3}{4^s} + \frac{2}{5^s} + \frac{4}{6^s} + \frac{2}{7^s} + \frac{4}{8^s} + \frac{3}{9^s} + \frac{4}{10^s} + \cdots$$

The general theorem reduces this verification to the convolution identity $d = \mathbf{1} * \mathbf{1}$, sparing the explicit double-sum manipulation in every individual case.

The convolution-multiplication identity admits a more conceptual restatement.

Theorem 2.2.4: Ring Isomorphism: Arithmetic Functions and Dirichlet Series

Let $\mathcal{A}_a \subseteq \mathcal{A}$ denote the subring of arithmetic functions f with finite abscissa of absolute convergence $\sigma_a(f) < \infty$. The map

$$\Phi : \mathcal{A}_a \longrightarrow \{\text{Dirichlet series with finite } \sigma_a\}, \quad \Phi(f) = D(f, s)$$

is an injective ring homomorphism, where the target is regarded as the ring of analytic functions on common half-planes of absolute convergence with pointwise addition and multiplication. Under Φ :

1. $\varepsilon \mapsto 1$,
2. $\mathbf{1} \mapsto \zeta(s)$,
3. $\mu \mapsto 1/\zeta(s)$,
4. $\text{id} \mapsto \zeta(s-1)$.

Proof:

Linearity of Φ is immediate: $D(f + g, s) = D(f, s) + D(g, s)$ on the common half-plane of absolute convergence. Multiplicativity is theorem 2.2.1: $D(f * g, s) = D(f, s) D(g, s)$ on

$\operatorname{Re}(s) > \max(\sigma_a(f), \sigma_a(g))$. The unit ε maps to the constant function 1 since $D(\varepsilon, s) = \varepsilon(1) = 1$.

Injectivity is the content of corollary 2.1.8: if $D(f, s) = D(g, s)$ on a common half-plane, then $f(n) = g(n)$ for all n . The named correspondences come from direct computation: $D(\varepsilon, s) = 1$, $D(\mathbf{1}, s) = \zeta(s)$, $D(\operatorname{id}, s) = \zeta(s-1)$, and $D(\mu, s) = 1/\zeta(s)$ from corollary 2.2.2, completing the proof.

This restatement clarifies the structural picture. The formal-algebraic content of arithmetic functions (the convolution ring $\mathcal{A}$) and the analytic content of Dirichlet series (functions on a half-plane with a given abscissa of absolute convergence) are not just compatible but identical, up to the inclusion $\mathcal{A}_a \subseteq \mathcal{A}$. Algebraic identities like $\mu * \mathbf{1} = \varepsilon$ become analytic identities like $\zeta(s) \sum \mu(n)/n^s = 1$, and conversely, analytic computations involving products and quotients of Dirichlet series translate back into convolution identities at the level of coefficients. The ring structure on the analytic side imposes constraints that are invisible algebraically: for instance, the analytic identity $\zeta(s) (\sum \mu(n)/n^s) = 1$ on $\operatorname{Re}(s) > 1$ forces the strict inequality $\sigma_c(\mu) \leq 1$, since $1/\zeta(s)$ continues analytically to a neighborhood of every point of $\{\operatorname{Re}(s) = 1\} \setminus \{1\}$ (using the non-vanishing $\zeta(1+it) \neq 0$, established in chapter 4).

The first nontrivial consequence of the ring isomorphism is a non-vanishing statement.

Example 2.2.5: Non-Vanishing of $\zeta(s)$ for $\operatorname{Re}(s) > 1$

The Möbius identity $1/\zeta(s) = \sum \mu(n)/n^s$ on $\operatorname{Re}(s) > 1$ shows that $\zeta(s) \neq 0$ on this half-plane. If $\zeta(s_0) = 0$ for some s_0 with $\operatorname{Re}(s_0) > 1$, then $1/\zeta(s)$ would have a pole at s_0 , contradicting the fact that $\sum \mu(n)/n^{s_0}$ converges absolutely to a finite value at s_0 (since $\operatorname{Re}(s_0) > 1 = \sigma_a(\mu)$). The next section gives a more conceptual proof of this fact via the Euler product, which extends without change to any Dirichlet series with an Euler product. The harder boundary case $\operatorname{Re}(s) = 1$, where the Möbius series sits on the boundary of its half-plane of absolute convergence and the argument above breaks down, is treated in chapter 4.

The headline application of the ring isomorphism is the identity that connects $\zeta(s)$ to prime-counting.

Theorem 2.2.6: Logarithmic Derivative of $\zeta(s)$

For $\operatorname{Re}(s) > 1$,

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

where Λ is the von Mangoldt function (definition 1.1.8).

Proof:

The strategy combines two ingredients: the convolution identity $\Lambda * \mathbf{1} = \log$ from proposition 1.1.9, and term-by-term differentiation of $\zeta(s)$ from theorem 2.1.7.

Step 1: Differentiate $\zeta(s)$. By theorem 2.1.7, for $\operatorname{Re}(s) > 1$,

$$\zeta'(s) = - \sum_{n=1}^{\infty} \frac{\log n}{n^s}$$

Equivalently, $-\zeta'(s) = D(\log, s)$, the Dirichlet series of the function $n \mapsto \log n$.

Step 2: Apply the convolution identity. Under the ring isomorphism of theorem 2.2.4, the von Mangoldt identity $\Lambda * \mathbf{1} = \log$ becomes

$$D(\Lambda, s) \cdot \zeta(s) = D(\log, s) = -\zeta'(s)$$

on $\operatorname{Re}(s) > 1$. Absolute convergence of $D(\Lambda, s)$ on this half-plane follows from $0 \leq \Lambda(n) \leq \log n$ together with $\sigma_a(\log) = 1$ (since $\log n \leq n^\varepsilon$ for any $\varepsilon > 0$, so $\sum (\log n)/n^\sigma$ converges for every $\sigma > 1$).

Step 3: Divide. Since $\zeta(s) \neq 0$ for $\operatorname{Re}(s) > 1$ (example 2.2.5), division by $\zeta(s)$ yields

$$\sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} = -\frac{\zeta'(s)}{\zeta(s)}$$

on $\operatorname{Re}(s) > 1$, completing the proof.

The identity $-\zeta'(s)/\zeta(s) = \sum \Lambda(n)/n^s$ is the single most important identity in elementary analytic number theory. Its left side is an analytic object: the logarithmic derivative of $\zeta(s)$, whose poles encode the analytic structure of $\zeta(s)$. Its right side is the Dirichlet series of the von Mangoldt function, whose summatory function $\psi(x) = \sum_{n \leq x} \Lambda(n)$ counts prime powers (weighted by $\log p$) and is the natural object whose asymptotics encode the Prime Number Theorem $\psi(x) \sim x$. The identity converts an analytic problem (locating the poles and zeros of $-\zeta'(s)/\zeta(s)$) into an arithmetic one (estimating $\psi(x)$). The simple pole of $\zeta(s)$ at $s = 1$, with residue 1, becomes a simple pole of $-\zeta'(s)/\zeta(s)$ at $s = 1$ with residue 1 (since $-\zeta'(s)/\zeta(s) \sim 1/(s-1)$ near $s = 1$); this residue is the analytic source of the leading term $\psi(x) \sim x$ in the Prime Number Theorem. The contributions of the zeros of $\zeta(s)$ inside the critical strip give the oscillatory error terms in the explicit formula. The whole strategy of the analytic proof of the Prime Number Theorem, developed in chapter 5, rests on the identity proved above.

The structural extension of theorem 2.2.6 to general multiplicative arithmetic functions is the following: for multiplicative f with $D(f, s) \neq 0$ in some half-plane, the logarithmic derivative $-D'(f, s)/D(f, s)$ is the Dirichlet series of an arithmetic function Λ_f supported on prime powers, generalizing the von Mangoldt function. The analytic structure of the logarithmic derivative is therefore controlled entirely by the values of f at prime powers, mirroring the determination-by-prime-powers principle for multiplicative arithmetic functions (proposition 1.1.3).

2.3 Euler Products

The ring isomorphism of theorem 2.2.4 converts Dirichlet convolution into multiplication of Dirichlet series, but its full force is felt only for *multiplicative* functions. For such functions the values $f(n)$ at all integers are reconstructed from the values $f(p^k)$ at prime powers (proposition 1.1.3), and this reconstruction lifts to the analytic level: the Dirichlet series $D(f, s)$ factors as a product of local pieces, one for each prime, with the local piece at p built entirely from $f(p), f(p^2), f(p^3), \dots$. The resulting factorization is the *Euler product*, and it is the analytic incarnation of unique factorization in $\mathbb{Z}$. Where theorem 2.2.1 expresses one global relation $D(f * g, s) = D(f, s)D(g, s)$, the Euler product expresses a relation between $D(f, s)$ and the infinitely many local factors at once.

The mechanism is the local-to-global principle of section 1.1 made analytic. A multiplicative function decouples across primes; absolute convergence of $D(f, s)$ lets the corresponding infinite sum be regrouped as an infinite product over primes without disturbing its value. The product form exposes structure that the series hides: it shows immediately that $D(f, s)$ cannot vanish in the region of absolute convergence (every factor is a nonzero number near 1), and its logarithm is a sum over prime powers that ties the analytic object directly to the primes. These two consequences, non-vanishing and the logarithmic expansion, are the threads that the study of prime distribution in chapter 3 and the theory of $\zeta(s)$ in chapter 4 will pull on.

Theorem 2.3.1: Euler Product for Multiplicative Functions

Let f be a multiplicative arithmetic function (definition 1.1.2), and let $\sigma_a(f)$ be its abscissa of absolute convergence (definition 2.1.3). For $\operatorname{Re}(s) > \sigma_a(f)$,

$$D(f, s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s} = \prod_p \left(\sum_{k=0}^{\infty} \frac{f(p^k)}{p^{ks}} \right)$$

where the product runs over all primes and converges absolutely. If f is totally multiplicative, the local factor at p is the geometric series

$$\sum_{k=0}^{\infty} \frac{f(p^k)}{p^{ks}} = \sum_{k=0}^{\infty} \left(\frac{f(p)}{p^s} \right)^k = \frac{1}{1 - f(p)p^{-s}}$$

Proof:

Fix s with $\sigma = \operatorname{Re}(s) > \sigma_a(f)$, so that $\sum_{n=1}^{\infty} |f(n)|n^{-\sigma} < \infty$. Write $f_p(s) = \sum_{k=0}^{\infty} f(p^k)p^{-ks}$ for the local factor at p .

Step 1: Each local factor converges absolutely. For a single prime p , the terms $f(p^k)p^{-ks}$ are a subsequence of the terms $f(n)n^{-s}$ (those indexed by $n = p^k$), so

$$\sum_{k=0}^{\infty} \left| \frac{f(p^k)}{p^{ks}} \right| = \sum_{k=0}^{\infty} \frac{|f(p^k)|}{p^{k\sigma}} \leq \sum_{n=1}^{\infty} \frac{|f(n)|}{n^{\sigma}} < \infty$$

Hence $f_p(s)$ is a well-defined complex number, and $f_p(s) = 1 + \sum_{k=1}^{\infty} f(p^k)p^{-ks}$ since $f(p^0) = f(1) = 1$ by multiplicativity.

Step 2: The finite product over small primes equals a partial sum. Fix $y \geq 2$ and let $p_1 < p_2 < \dots < p_r$ be the primes not exceeding y . Each $f_{p_i}(s)$ is an absolutely convergent series, so their finite product may be expanded by multiplying out and rearranging (a finite product of absolutely convergent series converges absolutely to the iterated sum):

$$\prod_{i=1}^r f_{p_i}(s) = \sum_{k_1=0}^{\infty} \dots \sum_{k_r=0}^{\infty} \frac{f(p_1^{k_1}) \dots f(p_r^{k_r})}{(p_1^{k_1} \dots p_r^{k_r})^s}$$

By multiplicativity of f , the numerator $f(p_1^{k_1}) \dots f(p_r^{k_r})$ equals $f(p_1^{k_1} \dots p_r^{k_r})$, since the prime powers $p_i^{k_i}$ are pairwise coprime. By unique factorization (??, section 0.2), the map $(k_1, \dots, k_r) \mapsto$

$n = p_1^{k_1} \cdots p_r^{k_r}$ is a bijection from r -tuples of non-negative integers onto the set

$$S_y = \{n \geq 1 : \text{every prime factor of } n \text{ is } \leq y\}$$

of y -smooth integers. Therefore

$$\prod_{i=1}^r f_{p_i}(s) = \sum_{n \in S_y} \frac{f(n)}{n^s}$$

the local factors at the primes $\leq y$ assemble exactly into the sum over integers built from those primes.

Step 3: Control the tail. Every integer $n \leq y$ has all prime factors $\leq y$, so $\{1, 2, \dots, \lfloor y \rfloor\} \subseteq S_y$. The difference between $D(f, s)$ and the partial product is supported on integers not in S_y , each of which exceeds y :

$$\left| D(f, s) - \prod_{i=1}^r f_{p_i}(s) \right| = \left| \sum_{n \notin S_y} \frac{f(n)}{n^s} \right| \leq \sum_{n \notin S_y} \frac{|f(n)|}{n^\sigma} \leq \sum_{n > y} \frac{|f(n)|}{n^\sigma}$$

The right-hand side is the tail of the convergent series $\sum_n |f(n)|n^{-\sigma}$, so it tends to 0 as $y \rightarrow \infty$. Letting $y \rightarrow \infty$ gives $\prod_p f_p(s) = D(f, s)$, with the product interpreted as the limit of its partial products.

Step 4: Absolute convergence of the infinite product. An infinite product $\prod_p (1 + a_p)$ converges absolutely precisely when $\sum_p |a_p| < \infty$ ([16]). Here $a_p = f_p(s) - 1 = \sum_{k=1}^{\infty} f(p^k)p^{-ks}$, and

$$\sum_p |a_p| = \sum_p \left| \sum_{k=1}^{\infty} \frac{f(p^k)}{p^{ks}} \right| \leq \sum_p \sum_{k=1}^{\infty} \frac{|f(p^k)|}{p^{k\sigma}} \leq \sum_{n=2}^{\infty} \frac{|f(n)|}{n^\sigma} < \infty$$

where the last inequality holds because each prime power p^k with $k \geq 1$ is a distinct integer ≥ 2 , so the double sum over (p, k) is a subseries of $\sum_{n \geq 2} |f(n)|n^{-\sigma}$. Thus the product converges absolutely, and in particular its value is independent of the order of the factors.

Step 5: The totally multiplicative case. If f is totally multiplicative, then $f(p^k) = f(p)^k$ by proposition 1.1.3. The local factor is the geometric series $\sum_{k \geq 0} (f(p)p^{-s})^k$ with ratio $f(p)p^{-s}$, and its absolute convergence (Step 1) forces $|f(p)p^{-s}| < 1$, since a geometric series whose ratio has modulus ≥ 1 has terms that do not tend to 0 and hence diverges. Summing the geometric series gives $f_p(s) = (1 - f(p)p^{-s})^{-1}$, completing the proof.

The Euler product is unique factorization read analytically. Step 2 is the heart of the matter: multiplying the local factors and expanding produces exactly one term for each way of choosing a prime-power part at each prime, and unique factorization guarantees that these choices correspond bijectively to integers. The hypothesis of absolute convergence is what permits the rearrangement; without it, the regrouping of the double series into a product would not be justified, and the value could change. The role of multiplicativity is equally precise: it is what makes the numerator $f(p_1^{k_1}) \cdots f(p_r^{k_r})$ collapse to the single value $f(n)$, so that the expanded product is genuinely the Dirichlet series and not some twisted variant. For totally multiplicative f , the further collapse

$f(p^k) = f(p)^k$ turns each local factor into a closed-form geometric series, and the Euler product becomes a product of simple rational functions of p^{-s} .

The first and defining instance is the zeta function itself, where $f = \mathbf{1}$ is the constant function 1, totally multiplicative with $\mathbf{1}(p) = 1$.

Corollary 2.3.2: The Euler Product for the Riemann Zeta Function

For $\operatorname{Re}(s) > 1$,

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_p \frac{1}{1 - p^{-s}}$$

the product converging absolutely.

Proof:

The constant function $\mathbf{1}$ is totally multiplicative with $\mathbf{1}(p) = 1$ and $\sigma_a(\mathbf{1}) = 1$ (example 2.1.9). Applying theorem 2.3.1 with $f = \mathbf{1}$, the local factor at p is the geometric series $\sum_{k=0}^{\infty} p^{-ks} = (1 - p^{-s})^{-1}$, so $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ for $\operatorname{Re}(s) > 1$, as required.

This identity, due to Euler, is the analytic statement of the fundamental theorem of arithmetic. Reading the product from right to left and expanding each factor as a geometric series, $\prod_p (1 + p^{-s} + p^{-2s} + \dots)$, the coefficient of n^{-s} in the expansion counts the number of ways to write n as a product of prime powers, which is exactly one. The equality with $\sum_n n^{-s}$ therefore encodes the fact that every integer factors into primes in exactly one way. The product representation also carries arithmetic information that the sum conceals. As $s \rightarrow 1^+$ along the real axis, $\zeta(s) \rightarrow +\infty$ because the harmonic series diverges; on the product side this forces $\prod_p (1 - p^{-1})^{-1}$ to diverge, which can happen only if there are infinitely many primes and $\sum_p p^{-1}$ diverges. Euler's original proof of the infinitude of primes is precisely this observation, and the quantitative form of the divergence is the subject of example 2.3.6 below and, in full, of the elementary theory of prime distribution in chapter 3 (a finite Euler product would give a finite limit, contradicting the pole). The analytic behavior of $\zeta(s)$ near $s = 1$, and throughout the critical strip, is developed in chapter 4.

The convolution identities of corollary 2.2.2 all have Euler product counterparts, obtained by computing one local factor and recognizing the resulting product. Carrying out these computations gives a second, independent route to those identities, one that never invokes the multiplication theorem but instead reads each global product off its local factors.

Example 2.3.3: Euler Products of Standard Dirichlet Series

Each of the following multiplicative functions has an Euler product computed from its values at prime powers. The global product is then identified with a ratio of zeta values, cross-checking corollary 2.2.2.

1. *The Möbius function μ (definition 1.1.4).* Here $\mu(p) = -1$ and $\mu(p^k) = 0$ for $k \geq 2$, so the local factor terminates:

$$\sum_{k=0}^{\infty} \frac{\mu(p^k)}{p^{ks}} = 1 + \frac{\mu(p)}{p^s} = 1 - \frac{1}{p^s}$$

Hence $D(\mu, s) = \prod_p (1 - p^{-s})$ for $\operatorname{Re}(s) > 1$. Comparing with corollary 2.3.2, this is exactly $\prod_p (1 - p^{-s})^{-1}$ inverted factor by factor, so $D(\mu, s) = 1/\zeta(s)$, matching corollary 2.2.2(3).

2. The divisor function $d = \mathbf{1} * \mathbf{1}$ (definition 1.1.5). Since $d(p^k) = k + 1$, the local factor is

$$\sum_{k=0}^{\infty} \frac{k+1}{p^{ks}} = \sum_{k=0}^{\infty} (k+1)x^k = \frac{1}{(1-x)^2}, \quad x = p^{-s}$$

using the standard power series $\sum_{k \geq 0} (k+1)x^k = (1-x)^{-2}$ for $|x| < 1$. Therefore $D(d, s) = \prod_p (1-p^{-s})^{-2} = \zeta(s)^2$ for $\operatorname{Re}(s) > 1$, matching corollary 2.2.2(1). The local factor $(1-p^{-s})^{-2}$ is the square of the local factor of $\zeta(s)$, the product form of $d = \mathbf{1} * \mathbf{1}$.

3. The Liouville function λ (definition 1.1.6). The function λ is totally multiplicative with $\lambda(p) = -1$, so theorem 2.3.1 gives the geometric local factor

$$\sum_{k=0}^{\infty} \frac{\lambda(p^k)}{p^{ks}} = \frac{1}{1 - \lambda(p)p^{-s}} = \frac{1}{1 + p^{-s}}$$

Hence $D(\lambda, s) = \prod_p (1 + p^{-s})^{-1}$ for $\operatorname{Re}(s) > 1$. To identify the global product, write $1 + p^{-s} = (1 - p^{-2s})/(1 - p^{-s})$, so that

$$\prod_p \frac{1}{1 + p^{-s}} = \prod_p \frac{1 - p^{-s}}{1 - p^{-2s}} = \frac{\prod_p (1 - p^{-s})}{\prod_p (1 - p^{-2s})} = \frac{1/\zeta(s)}{1/\zeta(2s)} = \frac{\zeta(2s)}{\zeta(s)}$$

where the last step applies corollary 2.3.2 at s and at $2s$ (both products converge absolutely for $\operatorname{Re}(s) > 1$). This matches corollary 2.2.2(5).

4. The Euler totient φ . The totient is multiplicative with $\varphi(p^k) = p^k - p^{k-1}$ for $k \geq 1$ and $\varphi(1) = 1$. The local factor is

$$1 + \sum_{k=1}^{\infty} \frac{p^k - p^{k-1}}{p^{ks}} = 1 + \sum_{k=1}^{\infty} p^{k(1-s)} - \sum_{k=1}^{\infty} p^{k-1-ks}$$

Both geometric series converge for $\operatorname{Re}(s) > 1$, where $|p^{1-s}| = p^{1-\sigma} < 1$. Summing them,

$$\sum_{k=1}^{\infty} p^{k(1-s)} = \frac{p^{1-s}}{1 - p^{1-s}}, \quad \sum_{k=1}^{\infty} p^{k-1-ks} = \frac{p^{-s}}{1 - p^{1-s}}$$

so the local factor is

$$1 + \frac{p^{1-s} - p^{-s}}{1 - p^{1-s}} = \frac{1 - p^{-s}}{1 - p^{1-s}}$$

The local factor is thus valid for $\operatorname{Re}(s) > 1$. Identifying the global product, however, requires corollary 2.3.2 at $s-1$: the factor $\prod_p (1 - p^{1-s}) = \prod_p (1 - p^{-(s-1)}) = 1/\zeta(s-1)$ converges only for $\operatorname{Re}(s-1) > 1$, so it is this $\zeta(s-1)$ factor, not the local series, that pins the region. Applying corollary 2.3.2 at s and at $s-1$,

$$D(\varphi, s) = \prod_p \frac{1 - p^{-s}}{1 - p^{1-s}} = \frac{1/\zeta(s)}{1/\zeta(s-1)} = \frac{\zeta(s-1)}{\zeta(s)}$$

for $\operatorname{Re}(s) > 2$, matching corollary 2.2.2(4).

In every case the local factor encodes the arithmetic of f at a single prime, and the shape of the global product records the convolution identity. The vanishing of $\mu(p^k)$ for $k \geq 2$ truncates the Möbius local factor to a polynomial $1 - p^{-s}$, the product reciprocal of $\zeta(s)$. The squaring $d = \mathbf{1} * \mathbf{1}$ appears as a squared local factor. The sign $\lambda(p) = -1$ flips the geometric series from $(1 - p^{-s})^{-1}$ to $(1 + p^{-s})^{-1}$, and the algebraic identity $(1 + x) = (1 - x^2)/(1 - x)$ converts this into the ratio $\zeta(2s)/\zeta(s)$ that isolates the squarefree structure. These computations are interchangeable with the convolution route of corollary 2.2.2. The Euler product route also exposes how the half-plane of validity arises: for the Liouville function the region $\operatorname{Re}(s) > 1$ is inherited directly from the local geometric series, whereas for the totient the local factors already converge on $\operatorname{Re}(s) > 1$ yet the global identity is pinned to $\operatorname{Re}(s) > 2$ by the $\zeta(s - 1)$ factor that appears upon recombination. The product form thus separates the local convergence of each factor from the global constraint imposed by the zeta values that assemble the answer.

A structural consequence of the product form, independent of any particular f , is that a Dirichlet series with an Euler product cannot vanish where the product converges. This is because a convergent infinite product of nonzero factors is nonzero, a criterion that has no counterpart on the series side.

Proposition 2.3.4: Non-Vanishing in the Half-Plane of Absolute Convergence

Let f be multiplicative, and let s be a point with $\operatorname{Re}(s) = \sigma$ at which

$$\sum_p \sum_{k=1}^{\infty} \frac{|f(p^k)|}{p^{k\sigma}} < \infty$$

and at which every local factor $f_p(s) = \sum_{k \geq 0} f(p^k)p^{-ks}$ is nonzero. Then the Euler product converges to a nonzero value, so $D(f, s) \neq 0$. In particular $\zeta(s) \neq 0$ for $\operatorname{Re}(s) > 1$.

Proof:

The hypothesis $\sum_p \sum_{k \geq 1} |f(p^k)|p^{-k\sigma} < \infty$ is exactly the absolute convergence condition $\sum_p |f_p(s) - 1| < \infty$ from Step 4 of theorem 2.3.1, since $f_p(s) - 1 = \sum_{k \geq 1} f(p^k)p^{-ks}$ and $|f_p(s) - 1| \leq \sum_{k \geq 1} |f(p^k)|p^{-k\sigma}$. Absolute convergence of the product $\prod_p f_p(s)$ follows, and by theorem 2.3.1 its value is $D(f, s)$.

The criterion for a nonzero limit is the following: an absolutely convergent infinite product $\prod_p f_p(s)$ with every factor $f_p(s) \neq 0$ has a nonzero value ([16]). The reason is that absolute convergence means $f_p(s) \rightarrow 1$, so all but finitely many factors lie in the disk $|z - 1| < 1/2$ on which $\log$ is defined; the tail $\sum_p \log f_p(s)$ converges to a finite complex number L , the corresponding tail product converges to $e^L \neq 0$, and multiplying by the finitely many remaining nonzero factors keeps the value nonzero. Hence $D(f, s) = \prod_p f_p(s) \neq 0$.

For the zeta function, take $f = \mathbf{1}$ and $\operatorname{Re}(s) > 1$. The local factors $(1 - p^{-s})^{-1}$ are nonzero (a reciprocal is never zero), and the convergence hypothesis holds since $\sum_p \sum_{k \geq 1} p^{-k\sigma} \leq \sum_{n \geq 2} n^{-\sigma} < \infty$. Therefore $\zeta(s) \neq 0$ for $\operatorname{Re}(s) > 1$, as we sought to show.

This proposition reproves the non-vanishing of example 2.2.5 from a structural vantage point. The earlier argument used the specific identity $1/\zeta(s) = \sum \mu(n)n^{-s}$ and observed that a zero of ζ would

produce a pole of an absolutely convergent series. The present argument is cleaner and more general: it uses only that $\zeta(s)$ has an Euler product with nonzero local factors, and it applies verbatim to any Dirichlet series possessing an Euler product, such as the Dirichlet L -functions whose non-vanishing on the line $\operatorname{Re}(s) = 1$ drives the theorem on primes in arithmetic progressions. The product form makes the non-vanishing transparent because the obstruction to a zero is local: a product vanishes only if a factor vanishes, and each local factor here is a value near 1. The difficult cases in the theory, such as the non-vanishing of $\zeta(1 + it)$ on the boundary line $\operatorname{Re}(s) = 1$, lie outside the region of absolute convergence, where this elementary argument no longer reaches; those are the province of chapter 4.

Taking the logarithm of an Euler product turns the product of local factors into a sum over primes, and expanding each logarithm as a power series produces a sum over prime powers. For totally multiplicative f the local factors are geometric, and the logarithm has an explicit expansion that connects the Dirichlet series directly to the prime powers.

Proposition 2.3.5: The Logarithm of an Euler Product

Let f be totally multiplicative with absolutely convergent Euler product on $\operatorname{Re}(s) > \sigma_a(f)$ and $|f(p)p^{-s}| < 1$ for every p . Then the double series

$$\log D(f, s) := \sum_p \sum_{k=1}^{\infty} \frac{f(p)^k}{k p^{ks}}$$

converges absolutely for $\operatorname{Re}(s) > \sigma_a(f)$ and defines a branch of the logarithm of $D(f, s)$ there: it is the analytic branch characterized by $\log D(f, s) \rightarrow 0$ as $\operatorname{Re}(s) \rightarrow +\infty$. This branch need not be the principal logarithm of $D(f, s)$; the two coincide precisely when the imaginary part of the series lies in $(-\pi, \pi]$, which holds whenever $\operatorname{Re}(s)$ is large enough. In particular, for $f = \mathbf{1}$ the series defines

$$\log \zeta(s) = \sum_p \sum_{k=1}^{\infty} \frac{1}{k p^{ks}}$$

for $\operatorname{Re}(s) > 1$, agreeing with the ordinary real logarithm on the real segment $s > 1$ (there $\zeta(s) > 1$). Differentiating term by term recovers

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

the identity of theorem 2.2.6.

Proof:

Step 1: A logarithm of the product as a sum of factor logarithms. By theorem 2.3.1 the local factor is $f_p(s) = (1 - f(p)p^{-s})^{-1}$, and the product $\prod_p f_p(s)$ converges absolutely. Throughout, $\log$ on a single factor denotes the principal branch. Each factor satisfies $|f(p)p^{-s}| < 1$, so $1 - f(p)p^{-s}$ lies in the open right half-plane $\operatorname{Re}(z) > 0$, away from the branch cut, and the principal expansion $-\log(1 - z) = \sum_{k \geq 1} z^k/k$ applies with $z = f(p)p^{-s}$. Setting

$$\log D(f, s) := \sum_p \log \frac{1}{1 - f(p)p^{-s}} = - \sum_p \log (1 - f(p)p^{-s}) = \sum_p \sum_{k=1}^{\infty} \frac{f(p)^k}{k p^{ks}}$$

gives the double series of the statement. The symbol $\log D(f, s)$ is *defined* by this sum; it is a logarithm of $D(f, s)$ because exponentiating term by term gives $\exp(\sum_p \log f_p(s)) = \prod_p f_p(s) = D(f, s)$ by theorem 2.3.1, the interchange of exp with the convergent sum being the continuity of exp applied to the partial sums. It need not be the *principal* logarithm of $D(f, s)$: the principal logarithm has imaginary part in $(-\pi, \pi]$, whereas the imaginary part $\sum_p \sum_{k \geq 1} \operatorname{Im}(f(p)^k p^{-ks})/k$ of the series is constrained only by absolute convergence and need not lie in $(-\pi, \pi]$; the two coincide precisely when it does. The double series converges absolutely: $\sum_p \sum_{k \geq 1} |f(p)|^k p^{-k\sigma}/k \leq \sum_p \sum_{k \geq 1} |f(p)p^{-s}|^k = \sum_p |f(p)p^{-s}|/(1 - |f(p)p^{-s}|)$. Absolute convergence of each local factor (Step 1 of theorem 2.3.1) forces $|f(p)p^{-s}| \rightarrow 0$, so $|f(p)p^{-s}| \leq 1/2$ for all but finitely many p , and for those p the summand is at most $2|f(p)p^{-s}|$. Since $\sum_p |f(p)p^{-s}|$ converges (its terms are a subseries of $\sum_n |f(n)|n^{-\sigma}$), the double series converges absolutely.

Step 2: Specialize to $\zeta(s)$. For $f = \mathbf{1}$, every $f(p) = 1$, so $f(p)^k = 1$, giving

$$\log \zeta(s) = \sum_p \sum_{k=1}^{\infty} \frac{1}{k p^{ks}}$$

for $\operatorname{Re}(s) > 1$.

Step 3: Differentiate term by term. Each summand $1/(k p^{ks}) = k^{-1} p^{-ks}$ is an entire function of s with derivative $-(\log p) p^{-ks}$ (the factor k cancels). The differentiated series is

$$\sum_p \sum_{k=1}^{\infty} \frac{-\log p}{p^{ks}}$$

which converges absolutely and uniformly on $\operatorname{Re}(s) \geq 1+\delta$ for each $\delta > 0$: there $\sum_p \sum_{k \geq 1} (\log p) p^{-k(1+\delta)} = \sum_p (\log p) p^{-(1+\delta)}/(1 - p^{-(1+\delta)})$, which is finite because $\log p \leq p^{\delta/2}$ for large p makes the series comparable to $\sum_p p^{-1-\delta/2}$. Local uniform convergence of the differentiated series, together with convergence of the original at one point, justifies differentiating $\log \zeta(s)$ term by term ([16]). The derivative of the left side is $\zeta'(s)/\zeta(s)$, so

$$\frac{\zeta'(s)}{\zeta(s)} = \sum_p \sum_{k=1}^{\infty} \frac{-\log p}{p^{ks}}, \quad \text{hence} \quad -\frac{\zeta'(s)}{\zeta(s)} = \sum_p \sum_{k=1}^{\infty} \frac{\log p}{p^{ks}}$$

Step 4: Recognize the von Mangoldt series. The pairs (p, k) with p prime and $k \geq 1$ are in bijection with the prime powers $n = p^k$, and the summand $\log p$ is exactly $\Lambda(p^k)$ by definition 1.1.8. For n not a prime power, $\Lambda(n) = 0$, so the sum over prime powers equals the sum over all n :

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_p \sum_{k=1}^{\infty} \frac{\Lambda(p^k)}{p^{ks}} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

for $\operatorname{Re}(s) > 1$. This is the identity of theorem 2.2.6, completing the proof.

This gives a second derivation of the logarithmic-derivative formula $-\zeta'(s)/\zeta(s) = \sum \Lambda(n)n^{-s}$,

consistent with the convolution-based proof of theorem 2.2.6. The two proofs illuminate the identity from opposite sides. The convolution proof builds the formula from the identity $\Lambda * \mathbf{1} = \log$ at the level of arithmetic functions, transported through the ring isomorphism. The Euler product proof builds it from the factorization of $\zeta(s)$ over primes, with the von Mangoldt function emerging as the coefficient that records $\log p$ at each prime power p^k . The agreement of the two is a consistency check on the whole dictionary: the von Mangoldt function is at once the convolution inverse-image of $\log$ under $*$ and the logarithmic-derivative coefficient of the Euler product. The expansion $\log \zeta(s) = \sum_{p,k} 1/(k p^{ks})$ is itself the starting point for many estimates near $s = 1$, since the $k = 1$ term $\sum_p p^{-s}$ dominates and carries the divergence examined next.

Example 2.3.6: The Product Near $s = 1$

As $s \rightarrow 1^+$ along the real axis, $\zeta(s) \rightarrow +\infty$ because of the pole at $s = 1$, so $\log \zeta(s) \rightarrow +\infty$. In the expansion of proposition 2.3.5,

$$\log \zeta(s) = \sum_p \frac{1}{p^s} + \sum_p \sum_{k=2}^{\infty} \frac{1}{k p^{ks}}$$

the double tail with $k \geq 2$ stays bounded as $s \rightarrow 1^+$: it is dominated by $\sum_p \sum_{k \geq 2} p^{-k} = \sum_p p^{-2}/(1 - p^{-1}) \leq \sum_{n \geq 2} 1/(n(n-1)) = 1$ uniformly for $s \geq 1$. Since $\log \zeta(s) \rightarrow +\infty$ while this tail remains bounded, the prime sum must diverge:

$$\sum_p \frac{1}{p^s} \rightarrow +\infty \quad (s \rightarrow 1^+)$$

Letting $s \rightarrow 1^+$ shows that $\sum_p 1/p$ cannot converge, for if it did, $\sum_p p^{-s} \leq \sum_p p^{-1}$ would stay bounded. This foreshadows the divergence of $\sum_p 1/p$ and the infinitude of primes, treated quantitatively in chapter 3.

2.4 Mellin Transforms and the Integral Representation of $\zeta(s)$

The convergence theory and the convolution-multiplication dictionary of the previous sections treat a Dirichlet series as a formal-analytic object built from its coefficients. To extract asymptotic information from $\zeta(s)$ and its relatives, the series must be connected to integrals, where the machinery of complex analysis (contour shifting, residues, the calculus of poles) can be brought to bear. The device that makes this connection is the *Mellin transform*. Where the Fourier transform is adapted to the additive group $\mathbb{R}$ and the Laplace transform to the additive half-line, the Mellin transform is the integral transform adapted to the *multiplicative* group $(0, \infty)$: it converts dilation $t \mapsto at$ into a character a^{-s} , exactly as the Fourier transform converts translation into a character $e^{i\xi a}$. This is precisely the symmetry that a Dirichlet series $\sum f(n)n^{-s}$ respects, since each term n^{-s} is a multiplicative character of the variable n evaluated at the parameter s .

The payoff is twofold. First, the Mellin transform expresses the gamma function, the Riemann zeta function, and their products as integrals over $(0, \infty)$ whose integrands have manifest analytic structure. The integral representation $\Gamma(s)\zeta(s) = \int_0^\infty t^{s-1}/(e^t - 1) dt$ derived below is the starting point from which the meromorphic continuation and the functional equation of $\zeta(s)$ are obtained

in chapter 4 and chapter 6. Second, the Mellin inversion formula runs the transform backward: it recovers an arithmetic sum from a contour integral of its Dirichlet series, and this is the analytic content of Perron's formula in section 2.5, the tool that converts analytic facts about $\zeta(s)$ into counting statements about primes. The gamma function will be developed in full in chapter 6; here only the integral $\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt$ for $\operatorname{Re}(s) > 0$ is needed, and it appears below as the first example of a Mellin transform.

Definition 2.4.1: The Mellin Transform

Let $f: (0, \infty) \rightarrow \mathbb{C}$ be a function that is (Lebesgue) measurable on $(0, \infty)$. The *Mellin transform* of f is the function

$$(\mathcal{M}f)(s) = \int_0^\infty f(t) t^{s-1} dt$$

defined for those $s \in \mathbb{C}$ for which the integral converges absolutely. Suppose f satisfies the two-sided power bound

$$f(t) = O(t^{-a}) \text{ as } t \rightarrow 0^+, \quad f(t) = O(t^{-b}) \text{ as } t \rightarrow \infty$$

with $a < b$. Then the integral converges absolutely for every s with $a < \operatorname{Re}(s) < b$, and the open vertical strip

$$\{s \in \mathbb{C} \mid a < \operatorname{Re}(s) < b\}$$

is called the *fundamental strip* of f .

The fundamental strip is the multiplicative analogue of the half-plane of convergence for a Dirichlet series, and the reason it is a strip rather than a half-plane is the presence of *two* endpoints to control. Near $t = 0$ the factor t^{s-1} contributes $t^{\operatorname{Re}(s)-1}$, so the integral $\int_0^1 f(t) t^{s-1} dt$ converges when $\operatorname{Re}(s) - 1 > a - 1$, that is, when $\operatorname{Re}(s) > a$; the lower bound on $\operatorname{Re}(s)$ is forced by the behavior of f at the origin. Near $t = \infty$ the same factor must decay against the growth of f , and $\int_1^\infty f(t) t^{s-1} dt$ converges when $\operatorname{Re}(s) - 1 < b - 1$, that is, when $\operatorname{Re}(s) < b$; the upper bound is forced by the behavior of f at infinity. Only on the overlap $a < \operatorname{Re}(s) < b$ do both tails converge, which requires $a < b$. As with Dirichlet series, $|t^{s-1}| = t^{\operatorname{Re}(s)-1}$ depends only on the real part of s , so the region of absolute convergence is invariant under vertical translation, hence a vertical strip.

Example 2.4.2: The Gamma Function as a Mellin Transform

Take $f(t) = e^{-t}$. Its Mellin transform is, by definition, the integral that defines the gamma function:

$$(\mathcal{M}e^{-t})(s) = \int_0^\infty e^{-t} t^{s-1} dt = \Gamma(s)$$

The fundamental strip is the half-plane $\operatorname{Re}(s) > 0$, which is the degenerate case $a = 0$, $b = +\infty$ of definition 2.4.1. Convergence at both ends is checked directly. Near $t = 0^+$, the factor e^{-t} is bounded (indeed $e^{-t} \rightarrow 1$), so the integrand is comparable to $t^{\operatorname{Re}(s)-1}$, and

$$\int_0^1 e^{-t} t^{\sigma-1} dt \leq \int_0^1 t^{\sigma-1} dt = \frac{1}{\sigma}$$

with $\sigma = \operatorname{Re}(s)$, which is finite precisely when $\sigma > 0$. Near $t = \infty$, the exponential decay of e^{-t} dominates every power of t : for any $\sigma \in \mathbb{R}$ there is a constant C_σ with $t^{\sigma-1} \leq C_\sigma e^{t/2}$ for $t \geq 1$,

whence

$$\int_1^\infty e^{-t} t^{\sigma-1} dt \leq C_\sigma \int_1^\infty e^{-t/2} dt = 2C_\sigma e^{-1/2} < \infty$$

for every real σ . The upper endpoint of the strip is therefore $b = +\infty$, and the integral defines $\Gamma(s)$ as an analytic function on the entire half-plane $\operatorname{Re}(s) > 0$. The continuation of Γ to a meromorphic function on all of $\mathbb{C}$, with simple poles at the non-positive integers, together with the functional equation $\Gamma(s+1) = s\Gamma(s)$ and the reflection formula, is developed in chapter 6; the present section uses only the integral on $\operatorname{Re}(s) > 0$.

The elementary properties of the transform are the precise statements of the slogans “dilation becomes a character” and “the Mellin transform is the Fourier transform in multiplicative coordinates.” These are the identities that will be applied repeatedly when manipulating the integrand of $\Gamma(s)\zeta(s)$.

Proposition 2.4.3: Elementary Properties of the Mellin Transform

Let $f: (0, \infty) \rightarrow \mathbb{C}$ have fundamental strip $a < \operatorname{Re}(s) < b$.

1. *Scaling.* For any real $\alpha > 0$, the function $t \mapsto f(\alpha t)$ has the same fundamental strip, and

$$(\mathcal{M}f(\alpha \cdot))(s) = \alpha^{-s}(\mathcal{M}f)(s)$$

2. *Shift by a power.* For any $\mu \in \mathbb{C}$, the function $t \mapsto t^\mu f(t)$ has fundamental strip $a - \operatorname{Re}(\mu) < \operatorname{Re}(s) < b - \operatorname{Re}(\mu)$, and

$$(\mathcal{M}t^\mu f(t))(s) = (\mathcal{M}f)(s + \mu)$$

3. *Reduction to a two-sided Laplace transform.* The substitution $t = e^{-x}$ identifies the Mellin transform with a two-sided Laplace transform. Writing $g(x) = f(e^{-x})$ for $x \in \mathbb{R}$,

$$(\mathcal{M}f)(s) = \int_{-\infty}^{\infty} g(x) e^{-sx} dx$$

for all s in the fundamental strip. In particular, on a vertical line $\operatorname{Re}(s) = c$ with $a < c < b$, setting $s = c + i\xi$ gives

$$(\mathcal{M}f)(c + i\xi) = \int_{-\infty}^{\infty} (e^{-cx} g(x)) e^{-i\xi x} dx$$

which is the Fourier transform of the function $x \mapsto e^{-cx} g(x) = e^{-cx} f(e^{-x})$.

Proof:

Throughout write $\sigma = \operatorname{Re}(s)$.

1. Fix $\alpha > 0$ and s with $a < \sigma < b$. In the integral $\int_0^\infty f(\alpha t) t^{s-1} dt$ substitute $u = \alpha t$, so $t = u/\alpha$, $dt = du/\alpha$, and $t^{s-1} = (u/\alpha)^{s-1} = \alpha^{1-s} u^{s-1}$. As t runs over $(0, \infty)$ so does u , and

$$\int_0^\infty f(\alpha t) t^{s-1} dt = \int_0^\infty f(u) \alpha^{1-s} u^{s-1} \frac{du}{\alpha} = \alpha^{-s} \int_0^\infty f(u) u^{s-1} du = \alpha^{-s} (\mathcal{M}f)(s)$$

The substitution is a C^1 diffeomorphism of $(0, \infty)$ with itself, so it preserves absolute convergence: the integral on the left converges absolutely if and only if the integral on the right does, which holds precisely for $a < \sigma < b$. Hence the fundamental strip is unchanged.

2. Fix $\mu \in \mathbb{C}$. The integrand of $\mathcal{M}(t^\mu f(t))$ at the parameter s is $t^\mu f(t) t^{s-1} = f(t) t^{(s+\mu)-1}$, which is exactly the integrand of $\mathcal{M}f$ at the parameter $s + \mu$. Therefore the integrals are equal whenever either converges:

$$(\mathcal{M} t^\mu f(t))(s) = \int_0^\infty f(t) t^{(s+\mu)-1} dt = (\mathcal{M}f)(s + \mu)$$

Absolute convergence of the right-hand integral holds for $a < \operatorname{Re}(s + \mu) < b$, that is, for $a - \operatorname{Re}(\mu) < \sigma < b - \operatorname{Re}(\mu)$, which is the asserted fundamental strip.

3. Substitute $t = e^{-x}$, so that $x = -\log t$, $dt = -e^{-x} dx$, and $t^{s-1} = e^{-(s-1)x}$. As t decreases from ∞ to 0 , the variable x increases from $-\infty$ to $+\infty$, so the limits reverse and absorb the sign of dt :

$$(\mathcal{M}f)(s) = \int_0^\infty f(t) t^{s-1} dt = \int_{-\infty}^\infty f(e^{-x}) e^{-(s-1)x} e^{-x} dx = \int_{-\infty}^\infty f(e^{-x}) e^{-sx} dx$$

With $g(x) = f(e^{-x})$ this is $\int_{-\infty}^\infty g(x) e^{-sx} dx$, the two-sided Laplace transform of g evaluated at s . The substitution $t = e^{-x}$ is again a C^1 diffeomorphism of $(0, \infty)$ onto $\mathbb{R}$, so absolute convergence is preserved and the identity holds throughout the fundamental strip. Writing $s = c + i\xi$ splits the exponential as $e^{-sx} = e^{-cx} e^{-i\xi x}$, and grouping the real exponential with g gives

$$(\mathcal{M}f)(c + i\xi) = \int_{-\infty}^\infty (e^{-cx} g(x)) e^{-i\xi x} dx$$

which is the Fourier transform of $x \mapsto e^{-cx} g(x)$ under the convention $\hat{h}(\xi) = \int_{-\infty}^\infty h(x) e^{-i\xi x} dx$, as we sought to show.

The third property is the structural heart of the transform. The multiplicative group $(0, \infty)$ is isomorphic, as a topological group, to the additive group $\mathbb{R}$ via the logarithm $t \mapsto -\log t = x$, and under this isomorphism Haar measure dt/t on $(0, \infty)$ becomes Lebesgue measure dx on $\mathbb{R}$. The Mellin transform is what the Fourier–Laplace transform on $\mathbb{R}$ becomes when transported back to $(0, \infty)$ through this isomorphism, with the complex parameter s playing the role of the (complexified) frequency variable. Every property of the Mellin transform is, after this change of variables, a property of the Fourier or Laplace transform; the scaling identity $(\mathcal{M}f(\alpha \cdot))(s) = \alpha^{-s} (\mathcal{M}f)(s)$, for instance, is the multiplicative form of the rule that translation by $\log \alpha$ in x multiplies the Fourier transform by a character.

This isomorphism does more than supply intuition: it converts the inversion problem for the Mellin transform into the inversion problem for the Fourier transform, which is settled by the classical Fourier inversion theorem. The result is the Mellin inversion formula.

Theorem 2.4.4: Mellin Inversion

Let $f: (0, \infty) \rightarrow \mathbb{C}$ be continuous, and let $c \in \mathbb{R}$ lie in the fundamental strip of f . Suppose that

1. the function $t \mapsto t^c f(t)$ is integrable on $(0, \infty)$ with respect to the Haar measure dt/t , that is, $\int_0^\infty |f(t)| t^c \frac{dt}{t} < \infty$; and
2. the function $\xi \mapsto (\mathcal{M}f)(c + i\xi)$ is integrable on $\mathbb{R}$, that is, $\int_{-\infty}^\infty |(\mathcal{M}f)(c + i\xi)| d\xi < \infty$.

Then for every $t > 0$,

$$f(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} (\mathcal{M}f)(s) t^{-s} ds$$

where the integral is taken over the vertical line $\operatorname{Re}(s) = c$.

Proof:

The proof transports the statement to $\mathbb{R}$ by the substitution $t = e^{-x}$ of proposition 2.4.3 and applies the Fourier inversion theorem there.

Step 1: Pass to the additive variable. Set $g(x) = f(e^{-x})$ for $x \in \mathbb{R}$ and define

$$G(x) = e^{-cx} g(x) = e^{-cx} f(e^{-x})$$

Since f is continuous on $(0, \infty)$ and $x \mapsto e^{-x}$ is continuous from $\mathbb{R}$ onto $(0, \infty)$, the function G is continuous on $\mathbb{R}$. Hypothesis (1) states that G is integrable: substituting $t = e^{-x}$ (so $dt/t = -dx$, and the orientation reversal cancels the sign) in the change-of-variables identity for the Haar measure gives

$$\int_{-\infty}^\infty |G(x)| dx = \int_{-\infty}^\infty |f(e^{-x})| e^{-cx} dx = \int_0^\infty |f(t)| t^c \frac{dt}{t} < \infty$$

so $G \in L^1(\mathbb{R})$.

Step 2: Identify the Fourier transform of G . By proposition 2.4.3(3), for $s = c + i\xi$,

$$\widehat{G}(\xi) = \int_{-\infty}^\infty G(x) e^{-i\xi x} dx = \int_{-\infty}^\infty e^{-cx} g(x) e^{-i\xi x} dx = (\mathcal{M}f)(c + i\xi)$$

under the angular-frequency convention $\widehat{G}(\xi) = \int_{-\infty}^\infty G(x) e^{-i\xi x} dx$, which is the convention fixed for the Mellin variable in proposition 2.4.3(3). The Fourier inversion theorem [7] is stated in the alternative normalization $\widehat{G}_{2\pi}(\eta) = \int_{-\infty}^\infty G(x) e^{-2\pi i \eta x} dx$, carrying the 2π in the exponent and no prefactor on inversion. The two transforms are affine reparametrizations of one another, $\widehat{G}_{2\pi}(\eta) = \widehat{G}(2\pi\eta)$, so $\widehat{G}_{2\pi} \in L^1(\mathbb{R})$ if and only if $\widehat{G} \in L^1(\mathbb{R})$. Hypothesis (2) is therefore exactly the statement that $\widehat{G} \in L^1(\mathbb{R})$, equivalently $\widehat{G}_{2\pi} \in L^1(\mathbb{R})$.

Step 3: Apply Fourier inversion. The function G is continuous and lies in $L^1(\mathbb{R})$ (Step 1), and its Fourier transform $\widehat{G}_{2\pi}$ lies in $L^1(\mathbb{R})$ (Step 2). These are precisely the hypotheses of the Fourier

inversion theorem [7], which in this normalization asserts that under them

$$G(x) = \int_{-\infty}^{\infty} \widehat{G}_{2\pi}(\eta) e^{2\pi i \eta x} d\eta \quad \text{for all } x \in \mathbb{R}$$

Substituting $\eta = \xi/(2\pi)$, so that $d\eta = d\xi/(2\pi)$, $\widehat{G}_{2\pi}(\eta) = \widehat{G}(2\pi\eta) = \widehat{G}(\xi)$, and $e^{2\pi i \eta x} = e^{i\xi x}$, transforms this into the angular-convention inversion formula

$$G(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \widehat{G}(\xi) e^{i\xi x} d\xi \quad \text{for all } x \in \mathbb{R}$$

in which the prefactor $1/(2\pi)$ is the Jacobian of the reparametrization. (The integrability of $\widehat{G}$ guarantees the right-hand integral converges absolutely, and continuity of G promotes the conclusion from “almost every x ” to “every x ”; see [8] for the measure-theoretic version and [7] for the statement used here.)

Step 4: Return to the multiplicative variable. Substitute $\widehat{G}(\xi) = (\mathcal{M}f)(c + i\xi)$ and $G(x) = e^{-cx} f(e^{-x})$ into the inversion formula of Step 3:

$$e^{-cx} f(e^{-x}) = \frac{1}{2\pi} \int_{-\infty}^{\infty} (\mathcal{M}f)(c + i\xi) e^{i\xi x} d\xi$$

Multiplying through by e^{cx} moves it inside the integral as $e^{cx} e^{i\xi x} = e^{(c+i\xi)x}$:

$$f(e^{-x}) = \frac{1}{2\pi} \int_{-\infty}^{\infty} (\mathcal{M}f)(c + i\xi) e^{(c+i\xi)x} d\xi$$

Now set $t = e^{-x}$, so $e^x = t^{-1}$ and $e^{(c+i\xi)x} = t^{-(c+i\xi)} = t^{-s}$ with $s = c + i\xi$. Along the vertical line $\operatorname{Re}(s) = c$ the variable s satisfies $ds = i d\xi$, so $d\xi = ds/i$, and as ξ runs from $-\infty$ to $+\infty$ the point s runs from $c - i\infty$ to $c + i\infty$. Therefore

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} (\mathcal{M}f)(c + i\xi) t^{-s} d\xi = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} (\mathcal{M}f)(s) t^{-s} ds$$

the factor $1/i$ converting $d\xi$ into ds and producing the $1/(2\pi i)$ normalization, completing the proof.

Two features of the inversion formula deserve comment. First, the integral runs up a vertical line $\operatorname{Re}(s) = c$ inside the fundamental strip, and the value of $f(t)$ is independent of which line c is chosen, since the left-hand side does not mention c ; this freedom to slide the contour within the strip is what makes the formula useful, because shifting the line of integration across a pole of $(\mathcal{M}f)(s)$ picks up a residue, and these residues are the main terms in asymptotic expansions. Second, the normalization $1/(2\pi i)$ together with the factor t^{-s} is exactly the data of a contour integral in the sense of [16]: the Mellin inversion formula presents $f(t)$ as a contour integral of its transform, and the entire apparatus of residue calculus applies to its evaluation. This is the precise sense in which the Mellin transform “converts an arithmetic sum into a contour integral,” and it is the mechanism underlying Perron’s formula in section 2.5, where $(\mathcal{M}f)(s)$ is replaced by a Dirichlet series and the contour integral recovers a summatory function.

The transform is now applied to the object of central interest. The Dirichlet series $\zeta(s) = \sum_{n \geq 1} n^{-s}$ of definition 2.1.1 converges on the half-plane $\operatorname{Re}(s) > 1$, where its abscissa of absolute convergence is $\sigma_a = 1$ (definition 2.1.3). Multiplying $\zeta(s)$ by $\Gamma(s)$ produces a single integral over $(0, \infty)$ whose integrand is an elementary function of t , with no reference to the individual terms n^{-s} .

Theorem 2.4.5: Integral Representation of the Riemann Zeta Function

For every s with $\operatorname{Re}(s) > 1$,

$$\Gamma(s) \zeta(s) = \int_0^\infty \frac{t^{s-1}}{e^t - 1} dt$$

the integral converging absolutely on this half-plane.

Proof:

Fix s with $\sigma = \operatorname{Re}(s) > 1$.

Step 1: Expand the integrand as a geometric series. For $t > 0$ one has $0 < e^{-t} < 1$, so the geometric series in e^{-t} converges:

$$\frac{1}{e^t - 1} = \frac{e^{-t}}{1 - e^{-t}} = \sum_{n=1}^{\infty} e^{-nt}$$

This is the sum of the geometric series with first term e^{-t} and ratio $e^{-t} \in (0, 1)$, valid for every $t > 0$. Multiplying by t^{s-1} gives the pointwise identity

$$\frac{t^{s-1}}{e^t - 1} = \sum_{n=1}^{\infty} t^{s-1} e^{-nt}$$

for each fixed $t > 0$.

Step 2: Evaluate the integral of a single term. For each $n \geq 1$, the substitution $u = nt$ in the gamma integral of example 2.4.2 gives, with $t = u/n$ and $dt = du/n$,

$$\int_0^\infty t^{s-1} e^{-nt} dt = \int_0^\infty \left(\frac{u}{n}\right)^{s-1} e^{-u} \frac{du}{n} = n^{-s} \int_0^\infty u^{s-1} e^{-u} du = \frac{\Gamma(s)}{n^s}$$

the integral $\int_0^\infty u^{s-1} e^{-u} du = \Gamma(s)$ converging absolutely because $\sigma > 1 > 0$ (example 2.4.2). In particular the same computation with s replaced by σ shows $\int_0^\infty t^{\sigma-1} e^{-nt} dt = \Gamma(\sigma) n^{-\sigma}$.

Step 3: Interchange summation and integration. The terms $t^{s-1} e^{-nt}$ are not all nonnegative (the factor t^{s-1} is complex when s is), so the interchange is justified on the absolute values, where Tonelli's theorem applies to the nonnegative integrand $(n, t) \mapsto |t^{s-1}| e^{-nt} = t^{\sigma-1} e^{-nt}$ over the product of counting measure on $\mathbb{Z}_{\geq 1}$ and Lebesgue measure on $(0, \infty)$. By Step 2 applied with σ in place of s ,

$$\sum_{n=1}^{\infty} \int_0^\infty t^{\sigma-1} e^{-nt} dt = \sum_{n=1}^{\infty} \frac{\Gamma(\sigma)}{n^\sigma} = \Gamma(\sigma) \zeta(\sigma) < \infty$$

the series converging because $\sigma > 1 = \sigma_a$ for ζ (definition 2.1.3). Tonelli's theorem [8] therefore guarantees that the function $(n, t) \mapsto t^{s-1} e^{-nt}$ is integrable for the product measure, and Fubini's

theorem [8] then permits the interchange of the sum and the integral for the (complex) integrand itself.

Step 4: Assemble. Using the pointwise identity of Step 1 and the justified interchange of Step 3,

$$\int_0^\infty \frac{t^{s-1}}{e^t - 1} dt = \int_0^\infty \sum_{n=1}^\infty t^{s-1} e^{-nt} dt = \sum_{n=1}^\infty \int_0^\infty t^{s-1} e^{-nt} dt = \sum_{n=1}^\infty \frac{\Gamma(s)}{n^s} = \Gamma(s) \zeta(s)$$

where the last equality factors the constant $\Gamma(s)$ out of the sum $\sum_n n^{-s} = \zeta(s)$, valid on $\operatorname{Re}(s) > 1$. The absolute convergence asserted in the statement is the finiteness computed in Step 3, completing the proof.

This integral representation is the analytic launch-point for everything that follows about $\zeta(s)$. The Dirichlet series $\sum n^{-s}$ converges only on $\operatorname{Re}(s) > 1$ and gives no direct access to the values of $\zeta(s)$ elsewhere; the integral $\int_0^\infty t^{s-1}/(e^t - 1) dt$, by contrast, has an integrand whose singularity structure can be analyzed by splitting the range at $t = 1$ and expanding $1/(e^t - 1)$ near the origin, where it behaves like $1/t$. That expansion separates a part that continues meromorphically to the whole plane from a part that is already entire, and this is exactly how the meromorphic continuation of $\zeta(s)$ to $\mathbb{C}$, with its single simple pole at $s = 1$, is produced in chapter 4. The same integral, after the contour is deformed into a Hankel loop around the positive real axis, yields the functional equation relating $\zeta(s)$ to $\zeta(1-s)$ in chapter 6. The representation also exhibits the gamma factor $\Gamma(s)$ as the inseparable companion of $\zeta(s)$: the natural object is the product $\Gamma(s)\zeta(s)$, and it is this product, not $\zeta(s)$ alone, whose Mellin-integral form has elementary content. The appearance of $\Gamma(s)$ here is the first sign of the completed zeta function $\xi(s)$ that organizes the functional equation.

The alternating analogue of the representation replaces $e^t - 1$ by $e^t + 1$ and brings in the Dirichlet eta function, the alternating series studied in example 2.1.10. The computation is identical except for a sign that turns the geometric series into an alternating one, and the resulting integrand is better behaved at the origin.

Example 2.4.6: The Dirichlet Eta Function and the Alternating Representation

The Dirichlet eta function is $\eta(s) = \sum_{n=1}^\infty (-1)^{n+1} n^{-s}$, which by example 2.1.10 satisfies $\eta(s) = (1 - 2^{1-s})\zeta(s)$ on $\operatorname{Re}(s) > 1$. Its product with $\Gamma(s)$ admits the integral representation

$$(1 - 2^{1-s}) \zeta(s) \Gamma(s) = \eta(s) \Gamma(s) = \int_0^\infty \frac{t^{s-1}}{e^t + 1} dt \quad (\operatorname{Re}(s) > 1)$$

derived exactly as in theorem 2.4.5. For $t > 0$ the geometric series gives

$$\frac{1}{e^t + 1} = \frac{e^{-t}}{1 + e^{-t}} = \sum_{n=1}^\infty (-1)^{n+1} e^{-nt}$$

(the series with first term e^{-t} and ratio $-e^{-t}$, with $|-e^{-t}| < 1$). Multiplying by t^{s-1} and integrating term by term gives

$$\int_0^\infty \frac{t^{s-1}}{e^t + 1} dt = \sum_{n=1}^\infty (-1)^{n+1} \int_0^\infty t^{s-1} e^{-nt} dt = \sum_{n=1}^\infty (-1)^{n+1} \frac{\Gamma(s)}{n^s} = \Gamma(s) \eta(s)$$

using the single-term evaluation $\int_0^\infty t^{s-1} e^{-nt} dt = \Gamma(s) n^{-s}$ from Step 2 of theorem 2.4.5. The interchange of summation and integration is justified by absolute convergence: the absolute values

$\sum_n \int_0^\infty t^{\sigma-1} e^{-nt} dt = \Gamma(\sigma)\zeta(\sigma)$ are summable for $\sigma = \operatorname{Re}(s) > 1$, so the same Tonelli–Fubini argument as in theorem 2.4.5 applies ([8]).

The contrast with the $1/(e^t - 1)$ integrand is the point of the example. Near $t = 0^+$ one has $e^t + 1 \rightarrow 2$, so $1/(e^t + 1) \rightarrow 1/2$ is bounded, and the integrand $t^{s-1}/(e^t + 1)$ is comparable to $\frac{1}{2}t^{\sigma-1}$, which is integrable on $(0, 1]$ for every $\sigma > 0$, not only for $\sigma > 1$. At $t = \infty$ the integrand decays like $t^{\sigma-1}e^{-t}$ for every σ . Hence the integral $\int_0^\infty t^{s-1}/(e^t + 1) dt$ converges and defines an analytic function on the entire half-plane $\operatorname{Re}(s) > 0$, even though the series defining $\eta(s)$ was used only on $\operatorname{Re}(s) > 1$. Foreshadowing the continuation of chapter 4: this convergent integral provides the meromorphic continuation of $\eta(s)\Gamma(s)$ to $\operatorname{Re}(s) > 0$, and dividing by the elementary factor $(1 - 2^{1-s})\Gamma(s)$ continues $\zeta(s)$ past the line $\operatorname{Re}(s) = 1$. The pole of $\zeta(s)$ at $s = 1$ survives this division because $(1 - 2^{1-s})$ vanishes there, while the factor $1/(e^t + 1)$ removed the obstruction at the origin that the $1/(e^t - 1)$ integrand carried.

2.5 Perron's Formula

The transition from a Dirichlet series to the arithmetic information it encodes runs in two directions. Forming $F(s) = \sum_{n \geq 1} a_n n^{-s}$ from a sequence (a_n) packages the coefficients into an analytic function; the analyticity, the location of poles, and the residues of F are the analytic data. Perron's formula runs the construction backward: it recovers the partial sum $\sum_{n \leq x} a_n$ of the coefficients from a contour integral of F . The kernel that performs the recovery is x^s/s , integrated up a vertical line to the right of the region where F converges. The effect is that analytic facts about F , principally the behavior of F near its singularities, are converted into asymptotic statements about the coefficient sum, since shifting the contour of integration leftward across a pole of $F(s)x^s/s$ contributes a residue, and these residues are the main terms in the asymptotics.

The instance that organizes this chapter and the next is $F(s) = -\zeta'(s)/\zeta(s) = \sum_{n \geq 1} \Lambda(n)n^{-s}$ (theorem 2.2.6). Here the coefficient sum is the Chebyshev function $\psi(x) = \sum_{n \leq x} \Lambda(n)$, and Perron's formula expresses $\psi(x)$ as a contour integral of $-\zeta'(s)/\zeta(s) \cdot x^s/s$. The single pole of $\zeta(s)$ at $s = 1$ produces a pole of the integrand with residue x , which is the source of the leading term $\psi(x) \sim x$ of the Prime Number Theorem; the zeros of $\zeta(s)$ inside the critical strip produce the oscillatory corrections. Perron's formula is therefore the gateway through which the analytic study of $\zeta(s)$ in chapter 4 becomes the analytic proof of the Prime Number Theorem in chapter 5. The formula itself is the Mellin-type inversion of theorem 2.4.4 specialized to the kernel x^s/s : where Mellin inversion recovers a function from its transform along a vertical line, Perron inversion recovers a step function (the summatory function) from the Dirichlet series, and the discontinuity of the step function at each integer is exactly the jump produced by the kernel x^s/s .

The entire formula rests on a single contour integral, the value of the kernel x^s/s integrated up a vertical line. This integral is discontinuous in the ratio it depends on, jumping from 0 to 1 as the ratio crosses 1, and it is this discontinuity that reproduces the indicator $n \leq x$ inside the coefficient sum.

Lemma 2.5.1: The Discontinuous Integral

Let $c > 0$. For $y > 0$ define

$$I(y) = \frac{1}{2\pi i} \int_{(c)} \frac{y^s}{s} ds = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{y^s}{s} ds$$

the integral taken up the vertical line $\operatorname{Re}(s) = c$. Then

$$I(y) = \begin{cases} 1 & y > 1 \\ \frac{1}{2} & y = 1 \\ 0 & 0 < y < 1 \end{cases}$$

Moreover, writing $I(y, T) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s/s ds$ for the truncation to the segment $[c-iT, c+iT]$ and $\delta(y) = 1$ for $y > 1$, $\delta(y) = 0$ for $0 < y < 1$, the truncation satisfies, for every $T > 0$,

$$|I(y, T) - \delta(y)| \leq y^c \min\left(1, \frac{1}{\pi T |\log y|}\right) \quad (y \neq 1)$$

and, in the boundary case $y = 1$,

$$\left|I(1, T) - \frac{1}{2}\right| \leq \frac{c}{\pi T}$$

Proof:

Throughout, $s = \sigma + it$ and the residue theorem is applied in the form established in [16]. The integrand y^s/s is meromorphic on $\mathbb{C}$ with a single simple pole at $s = 0$, where its residue is $\lim_{s \rightarrow 0} s \cdot y^s/s = y^0 = 1$.

Step 1: The boundary case $y = 1$. Here $y^s = 1$, so the truncated integral is computed directly. Parametrizing $s = c + it$ with $t \in [-T, T]$ and $ds = i dt$,

$$I(1, T) = \frac{1}{2\pi i} \int_{-T}^T \frac{i dt}{c + it} = \frac{1}{2\pi} \int_{-T}^T \frac{c - it}{c^2 + t^2} dt$$

The imaginary part of the integrand, $-t/(c^2 + t^2)$, is odd in t and integrates to zero over the symmetric interval. The real part gives

$$I(1, T) = \frac{1}{2\pi} \int_{-T}^T \frac{c}{c^2 + t^2} dt = \frac{1}{2\pi} \cdot 2 \arctan \frac{T}{c} = \frac{1}{\pi} \arctan \frac{T}{c}$$

As $T \rightarrow \infty$ this tends to $\frac{1}{\pi} \cdot \frac{\pi}{2} = \frac{1}{2}$, establishing $I(1) = \frac{1}{2}$. For the error, $\frac{1}{2} - \frac{1}{\pi} \arctan(T/c) = \frac{1}{\pi}(\frac{\pi}{2} - \arctan(T/c)) = \frac{1}{\pi} \arctan(c/T)$, and since $\arctan u \leq u$ for $u \geq 0$,

$$\left|I(1, T) - \frac{1}{2}\right| = \frac{1}{\pi} \arctan \frac{c}{T} \leq \frac{c}{\pi T}$$

which is the boundary estimate.

Step 2: The sharp truncation bound for $y \neq 1$. Fix $y \neq 1$ and $T > 0$. The estimate to prove is $|I(y, T) - \delta(y)| \leq y^c / (\pi T |\log y|)$. The argument depends on the sign of $\log y$ through which direction the contour is closed.

Suppose first $y > 1$, so $\log y > 0$ and $|y^s| = y^\sigma$ decays as $\sigma \rightarrow -\infty$. For $U > 0$ consider the rectangle $\mathcal{R}_U$ with vertices $c - iT$, $c + iT$, $-U + iT$, $-U - iT$, traversed counterclockwise. It encloses the pole at $s = 0$, so by the residue theorem

$$\frac{1}{2\pi i} \oint_{\mathcal{R}_U} \frac{y^s}{s} ds = 1$$

The boundary $\partial\mathcal{R}_U$ consists of the right edge (the segment from $c - iT$ to $c + iT$, contributing $I(y, T)$), the left edge at $\operatorname{Re}(s) = -U$, and the two horizontal edges at $\operatorname{Im}(s) = \pm T$. On the left edge, $|y^s/s| \leq y^{-U}/U \rightarrow 0$ as $U \rightarrow \infty$ because $y > 1$, so its contribution vanishes in the limit. On each horizontal edge, $s = \sigma \pm iT$ with σ running from $-U$ to c , and $|s| \geq T$, so $|y^s/s| \leq y^\sigma/T$; integrating in σ over $(-\infty, c]$ bounds each horizontal contribution by

$$\frac{1}{2\pi} \int_{-\infty}^c \frac{y^\sigma}{T} d\sigma = \frac{1}{2\pi T} \cdot \frac{y^c}{\log y}$$

using $\int_{-\infty}^c y^\sigma d\sigma = y^c / \log y$ (valid since $\log y > 0$). Letting $U \rightarrow \infty$, the residue identity reads $1 = I(y, T) + (\text{two horizontal contributions})$, whence

$$|I(y, T) - 1| \leq 2 \cdot \frac{1}{2\pi T} \cdot \frac{y^c}{\log y} = \frac{y^c}{\pi T \log y} = \frac{y^c}{\pi T |\log y|}$$

since $\log y = |\log y|$ here. This is the claim for $y > 1$.

Now suppose $0 < y < 1$, so $\log y < 0$ and $|y^s| = y^\sigma$ decays as $\sigma \rightarrow +\infty$. The same argument is run with the rectangle closed to the right: vertices $c - iT$, $c + iT$, $U + iT$, $U - iT$. This rectangle does not contain $s = 0$, so the closed integral is zero. The right edge at $\operatorname{Re}(s) = U$ contributes at most $y^U/U \rightarrow 0$. The horizontal edges, with σ running from c to U , are bounded by $\frac{1}{2\pi T} \int_c^\infty y^\sigma d\sigma = \frac{1}{2\pi T} \cdot y^c / |\log y|$ each, using $\int_c^\infty y^\sigma d\sigma = -y^c / \log y = y^c / |\log y|$. Hence

$$|I(y, T) - 0| \leq \frac{y^c}{\pi T |\log y|}$$

completing the sharp bound for both signs.

Step 3: The trivial bound $|I(y, T) - \delta(y)| \leq y^c$ for $y \neq 1$. The sharp bound degrades as $|\log y| \rightarrow 0$, so a second estimate independent of $|\log y|$ is needed to control y near 1. This is obtained by deforming the vertical segment to a circular arc on which $\operatorname{Re}(s) \geq c$ (for $y < 1$) or which encircles the pole (for $y > 1$). Set $R = \sqrt{c^2 + T^2}$, so that the endpoints $c \pm iT$ lie on the circle $|s| = R$, and write $\theta_0 = \arctan(T/c) \in (0, \pi/2)$, the argument of $c + iT$.

Suppose $0 < y < 1$. Let Γ be the arc of $|s| = R$ from $c + iT$ to $c - iT$ passing through $s = R$ on the right, that is, $s = Re^{i\theta}$ with θ decreasing from θ_0 to $-\theta_0$. The arc Γ and the segment from $c - iT$ to $c + iT$ together bound a region lying in $\operatorname{Re}(s) \geq c > 0$, which does not contain the pole $s = 0$; since the segment and Γ share endpoints, Cauchy's theorem gives $I(y, T) = \frac{1}{2\pi i} \int_\Gamma y^s/s ds$. On Γ , $\operatorname{Re}(s) = R \cos \theta$ with $|\theta| \leq \theta_0$, and $\cos \theta \geq \cos \theta_0 = c/R$, so $R \cos \theta \geq c$; because $0 < y < 1$, this forces $|y^s| = y^{R \cos \theta} \leq y^c$. With $|s| = R$ and $|ds| = R d\theta$,

$$|I(y, T)| = \frac{1}{2\pi} \left| \int_{-\theta_0}^{\theta_0} \frac{y^{Re^{i\theta}}}{Re^{i\theta}} Re^{i\theta} i d\theta \right| \leq \frac{1}{2\pi} \int_{-\theta_0}^{\theta_0} y^{R \cos \theta} d\theta \leq \frac{1}{2\pi} \cdot 2\theta_0 \cdot y^c \leq \frac{y^c}{2}$$

using $2\theta_0 \leq \pi$. Since $\delta(y) = 0$ here, $|I(y, T) - \delta(y)| \leq y^c/2 \leq y^c$.

Suppose now $y > 1$. Let Γ' be the arc of $|s| = R$ from $c + iT$ to $c - iT$ passing through $s = -R$ on the left, that is, $s = Re^{i\theta}$ with θ increasing from θ_0 to $2\pi - \theta_0$. The segment (upward) together with Γ' bounds a region containing $s = 0$ and is traversed counterclockwise, so the residue theorem gives $I(y, T) + \frac{1}{2\pi i} \int_{\Gamma'} y^s/s ds = 1$, hence $I(y, T) - 1 = -\frac{1}{2\pi i} \int_{\Gamma'} y^s/s ds$. On Γ' , $\operatorname{Re}(s) = R \cos \theta$ with $\theta \in [\theta_0, 2\pi - \theta_0]$, so $\cos \theta \leq \cos \theta_0 = c/R$, giving $R \cos \theta \leq c$; because $y > 1$, this forces $|y^s| = y^{R \cos \theta} \leq y^c$. The angular length of Γ' is $2\pi - 2\theta_0 \leq 2\pi$, so

$$|I(y, T) - 1| \leq \frac{1}{2\pi} \int_{\theta_0}^{2\pi - \theta_0} y^{R \cos \theta} d\theta \leq \frac{1}{2\pi} (2\pi - 2\theta_0) y^c \leq y^c$$

Since $\delta(y) = 1$ here, this is $|I(y, T) - \delta(y)| \leq y^c$.

Step 4: Combine and pass to the limit. For $y \neq 1$, Steps 2 and 3 give $|I(y, T) - \delta(y)| \leq y^c/(\pi T |\log y|)$ and $|I(y, T) - \delta(y)| \leq y^c$ respectively; the smaller of the two is the asserted bound $y^c \min(1, 1/(\pi T |\log y|))$. Letting $T \rightarrow \infty$ in this bound, with $y \neq 1$ fixed so that $1/(\pi T |\log y|) \rightarrow 0$, gives $I(y, T) \rightarrow \delta(y)$, which establishes $I(y) = 1$ for $y > 1$ and $I(y) = 0$ for $0 < y < 1$. Together with $I(1) = \frac{1}{2}$ from Step 1, all assertions are proved, completing the proof.

The lemma is the analytic mechanism behind everything in this section. The kernel y^s/s has a pole at $s = 0$ with residue 1, and integrating up a vertical line either captures that residue (when the contour can be closed to the left, which the decay of y^s permits exactly when $y > 1$) or does not (when it must be closed to the right, which is forced when $y < 1$). The jump of $I(y)$ from 0 to 1 as y crosses 1 is therefore the residue of y^s/s at the origin, switched on or off by the direction in which the integrand decays. The truncation estimate is the quantitative form: the error in replacing the limit by a finite integral is controlled by $1/(T |\log y|)$, so the integral resolves the indicator sharply except for ratios y extremely close to 1, where the second bound y^c takes over. This dichotomy, sharp resolution away from the diagonal $y = 1$ and a crude bound near it, is exactly what governs the error in the truncated form of Perron's formula below.

Applying the lemma term by term to a Dirichlet series converts its coefficient sum into a contour integral. The ratio y becomes x/n , so $I(x/n) = 1$ when $n < x$, $I(x/n) = 0$ when $n > x$, and the indicator $\sum_{n \leq x}$ emerges from the integral.

Theorem 2.5.2: Perron's Formula

Let $F(s) = \sum_{n \geq 1} a_n n^{-s}$ be a Dirichlet series with abscissa of absolute convergence σ_a (definition 2.1.3). Let $c > \max(0, \sigma_a)$ and $x > 0$. If x is not an integer, then

$$\sum_{n \leq x} a_n = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} F(s) \frac{x^s}{s} ds$$

the integral being taken up the line $\operatorname{Re}(s) = c$ and converging as the symmetric limit $\lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT}$. If x is an integer, the same identity holds with the left side replaced by the half-weighted sum $\sum_{n < x} a_n + \frac{1}{2}a_x$.

Proof:

Write $\sigma = \operatorname{Re}(s) = c$ throughout, and recall $I(y, T) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s/s \, ds$ from lemma 2.5.1.

Step 1: Interchange sum and integral on the truncated contour. Fix $T > 0$. On the segment $[c - iT, c + iT]$ the series for F may be inserted and the order of summation and integration exchanged:

$$\frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} \, ds = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \sum_{n=1}^{\infty} a_n \left(\frac{x}{n}\right)^s \frac{ds}{s} = \sum_{n=1}^{\infty} a_n \cdot I\left(\frac{x}{n}, T\right)$$

where $F(s)x^s = \sum_n a_n n^{-s} x^s = \sum_n a_n (x/n)^s$. The interchange is justified by absolute convergence over the product of counting measure on $\mathbb{Z}_{\geq 1}$ and Lebesgue measure on the finite segment: the modulus of the (n, s) -integrand is $|a_n|(x/n)^\sigma/|s|$, and

$$\sum_{n=1}^{\infty} \int_{c-iT}^{c+iT} \frac{|a_n|(x/n)^\sigma}{|s|} |ds| = \left(x^\sigma \sum_{n=1}^{\infty} \frac{|a_n|}{n^\sigma} \right) \left(\int_{-T}^T \frac{dt}{\sqrt{c^2 + t^2}} \right)$$

is finite: the first factor converges because $\sigma = c > \sigma_a$ (definition 2.1.3), and the second is the finite integral of a bounded function over a bounded interval. Fubini's theorem [8] therefore permits the interchange.

Step 2: Split off the finitely many terms with $n \leq x$. The sum $\sum_n a_n I(x/n, T)$ is split according to whether $n \leq x$ or $n > x$. Since x is fixed and finite, only finitely many indices satisfy $n \leq x$, so

$$\sum_{n=1}^{\infty} a_n I\left(\frac{x}{n}, T\right) = \sum_{n \leq x} a_n I\left(\frac{x}{n}, T\right) + \sum_{n > x} a_n I\left(\frac{x}{n}, T\right)$$

In the first sum, a finite sum, each term converges as $T \rightarrow \infty$ by lemma 2.5.1: for $n < x$ one has $x/n > 1$ and $I(x/n, T) \rightarrow 1$; if x is an integer the single term $n = x$ has $x/n = 1$ and $I(1, T) \rightarrow \frac{1}{2}$. Therefore

$$\lim_{T \rightarrow \infty} \sum_{n \leq x} a_n I\left(\frac{x}{n}, T\right) = \begin{cases} \sum_{n \leq x} a_n & x \notin \mathbb{Z} \\ \sum_{n < x} a_n + \frac{1}{2} a_x & x \in \mathbb{Z} \end{cases}$$

the cases differing only in the treatment of the boundary index $n = x$.

Step 3: The tail vanishes in the limit. For the tail $\sum_{n > x} a_n I(x/n, T)$, each index has $x/n < 1$, so $\delta(x/n) = 0$ and, by the trivial bound in lemma 2.5.1, $|I(x/n, T)| \leq (x/n)^c$ for every T . This dominating bound is summable, since

$$\sum_{n > x} |a_n| \left(\frac{x}{n}\right)^c = x^c \sum_{n > x} \frac{|a_n|}{n^c} < \infty$$

again because $c > \sigma_a$. For each fixed $n > x$, lemma 2.5.1 gives $I(x/n, T) \rightarrow \delta(x/n) = 0$ as $T \rightarrow \infty$, because $x/n < 1$ is fixed and bounded away from 1, so $1/(\pi T |\log(x/n)|) \rightarrow 0$. The

dominated convergence theorem for series [8] (counting measure on $\{n > x\}$, with dominating summable bound $|a_n|(x/n)^c$) then yields

$$\lim_{T \rightarrow \infty} \sum_{n > x} a_n I\left(\frac{x}{n}, T\right) = \sum_{n > x} a_n \cdot 0 = 0$$

Step 4: Assemble. Combining Steps 1 through 3, the symmetric limit of the truncated contour integral exists and equals

$$\lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds = \begin{cases} \sum_{n \leq x} a_n & x \notin \mathbb{Z} \\ \sum_{n < x} a_n + \frac{1}{2}a_x & x \in \mathbb{Z} \end{cases}$$

which is the assertion of the theorem, completing the proof.

Perron's formula is the precise statement that the analytic behavior of F controls the asymptotics of its coefficient sum. The left side is a step function of x , jumping by a_n at each integer n ; the right side is a contour integral whose integrand $F(s)x^s/s$ inherits all the singularities of F , together with the pole of the kernel at $s = 0$. The contour runs up a vertical line in the region of absolute convergence, where F is analytic and the integral has no singularities to contend with. The information is extracted by moving the contour to the left, into the region where F has poles: each pole crossed contributes its residue, and the integral over the shifted contour is smaller because the factor $x^s = x^\sigma e^{it \log x}$ has smaller modulus once σ has decreased. The residue at a pole $s = \rho$ of F contributes a term proportional to x^ρ/ρ , so a pole at $s = 1$ gives a main term of size x , a pole at $s = 0$ gives a constant, and poles further left give lower-order terms. The asymptotic expansion of $\sum_{n \leq x} a_n$ is, in this way, read off from the poles of F and their residues; the rightmost pole supplies the dominant term, and the remaining poles and the bound on the shifted contour supply the error. This is the mechanism that the analytic proof of the Prime Number Theorem makes quantitative in chapter 5.

The formula as stated is not yet usable for that program, because the contour runs to infinity in both directions and the integral over the full line is only conditionally convergent. Quantitative work requires truncating the contour to a finite height T and controlling the resulting error explicitly. The truncated form is the workhorse of the subject.

Theorem 2.5.3: The Truncated Perron Formula

Let $F(s) = \sum_{n \geq 1} a_n n^{-s}$ have abscissa of absolute convergence σ_a , let $c > \max(0, \sigma_a)$, and let $x \geq 1$ and $T \geq 1$. Then

$$\sum_{n \leq x} a_n = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds + R$$

where, if x is not an integer, the error term satisfies

$$|R| \leq \sum_{n=1}^{\infty} |a_n| \left(\frac{x}{n}\right)^c \min\left(1, \frac{1}{T|\log(x/n)|}\right)$$

and if x is an integer the same bound holds for the half-weighted sum $\sum_{n < x} a_n + \frac{1}{2}a_x$ in place of $\sum_{n \leq x} a_n$, augmented by the single near-diagonal term

$$|R| \leq \sum_{\substack{n=1 \\ n \neq x}}^{\infty} |a_n| \left(\frac{x}{n}\right)^c \min\left(1, \frac{1}{T|\log(x/n)|}\right) + |a_x| \frac{c}{\pi T}$$

Proof:

The starting point is the exact finite- T identity established in Step 1 of the proof of theorem 2.5.2, valid for every $T \geq 1$ with no passage to a limit:

$$\frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds = \sum_{n=1}^{\infty} a_n I\left(\frac{x}{n}, T\right)$$

the interchange of sum and integral being justified exactly as there (absolute convergence over the product of counting measure and the finite segment, using $c > \sigma_a$). The error R is the difference between the coefficient sum and this integral.

Case $x \notin \mathbb{Z}$. Here $x/n \neq 1$ for every n , so $\delta(x/n) = 1$ when $n < x$ and $\delta(x/n) = 0$ when $n > x$; equivalently $\delta(x/n) = \mathbb{1}[n \leq x]$, and therefore $\sum_{n \leq x} a_n = \sum_n a_n \delta(x/n)$. Subtracting the identity above,

$$R = \sum_{n \leq x} a_n - \frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds = \sum_{n=1}^{\infty} a_n \left(\delta\left(\frac{x}{n}\right) - I\left(\frac{x}{n}, T\right) \right)$$

The rearrangement is legitimate because both series converge absolutely: $\sum_n |a_n| |\delta(x/n)|$ is the finite sum over $n \leq x$, and $\sum_n |a_n| |I(x/n, T)|$ is dominated by $\sum_n |a_n| (x/n)^c = x^c \sum_n |a_n| n^{-c} < \infty$ via the trivial bound of lemma 2.5.1. Taking absolute values and applying the truncation estimate of lemma 2.5.1 to each term, with $|\delta(x/n) - I(x/n, T)| \leq (x/n)^c \min(1, 1/(\pi T |\log(x/n)|))$,

$$|R| \leq \sum_{n=1}^{\infty} |a_n| \left(\frac{x}{n}\right)^c \min\left(1, \frac{1}{\pi T |\log(x/n)|}\right) \leq \sum_{n=1}^{\infty} |a_n| \left(\frac{x}{n}\right)^c \min\left(1, \frac{1}{T |\log(x/n)|}\right)$$

the last inequality discarding the factor $1/\pi \leq 1$ inside the second argument of the minimum to give the form stated. This proves the non-integer case.

Case $x \in \mathbb{Z}$. Now exactly one index has $x/n = 1$, namely $n = x$, and for it δ takes the boundary value $\frac{1}{2}$ consistent with the half-weighted sum. For every other index $\delta(x/n) = \mathbb{1}[n < x]$, so the half-weighted sum is $\sum_{n < x} a_n + \frac{1}{2}a_x = \sum_n a_n \delta(x/n)$ with the convention $\delta(1) = \frac{1}{2}$. As before,

$$R = \sum_{n=1}^{\infty} a_n \left(\delta\left(\frac{x}{n}\right) - I\left(\frac{x}{n}, T\right) \right)$$

The single term $n = x$ contributes $a_x(\frac{1}{2} - I(1, T))$, bounded in modulus by $|a_x| c/(\pi T)$ by the boundary estimate of lemma 2.5.1. Every other term has $x/n \neq 1$ and is bounded as in the non-integer case. Hence

$$|R| \leq \sum_{\substack{n=1 \\ n \neq x}}^{\infty} |a_n| \left(\frac{x}{n}\right)^c \min\left(1, \frac{1}{T|\log(x/n)|}\right) + |a_x| \frac{c}{\pi T}$$

as required.

The structure of the error bound is what makes the truncated formula effective. The minimum has two regimes. For n far from x , the ratio x/n is bounded away from 1, $|\log(x/n)|$ is of order 1 or larger, and the term contributes at most $|a_n|(x/n)^c/T$, which is small because of the factor $1/T$. For n close to x , where $|\log(x/n)|$ is small and the first regime would give a useless bound, the minimum switches to the constant 1, and the term contributes at most $|a_n|(x/n)^c$; there are few such n , and the convergence factor $(x/n)^c$ keeps the total controlled. In applications the parameters c and T are chosen as functions of x to balance the contour integral against this error, and the explicit dependence on T in the bound is precisely what allows the optimization. The near-diagonal term in the integer case, of size $|a_x|/T$, is the price of evaluating the formula at a jump of the step function, and it is the reason the cleanest statements are made for non-integer x or after the half-weighting convention is adopted.

The application that motivates the entire development is the Chebyshev function ψ . Taking $a_n = \Lambda(n)$, the von Mangoldt function (definition 1.1.8), the generating Dirichlet series is the logarithmic derivative of ζ computed in theorem 2.2.6, and Perron's formula presents $\psi(x)$ as a contour integral of it.

Example 2.5.4: Perron's Formula for $\psi(x)$

Let $a_n = \Lambda(n)$, so that the summatory function is the Chebyshev function $\psi(x) = \sum_{n \leq x} \Lambda(n)$. By theorem 2.2.6, the generating Dirichlet series is

$$F(s) = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} = -\frac{\zeta'(s)}{\zeta(s)}$$

valid on $\operatorname{Re}(s) > 1$, where this is also the abscissa of absolute convergence: $0 \leq \Lambda(n) \leq \log n$ gives $\sum_n \Lambda(n)n^{-\sigma} \leq \sum_n (\log n)n^{-\sigma} < \infty$ for every $\sigma > 1$, while at $\sigma = 1$ the series $\sum_n \Lambda(n)/n$ diverges, so $\sigma_a = 1$. Choosing any $c > 1$ and any non-integer $x > 0$, theorem 2.5.2 yields

$$\psi(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \left(-\frac{\zeta'(s)}{\zeta(s)} \right) \frac{x^s}{s} ds$$

and the truncated form of theorem 2.5.3 gives the same integral over $[c - iT, c + iT]$ together with an explicit error governed by $\sum_n \Lambda(n)(x/n)^c \min(1, 1/(T|\log(x/n)|))$. This integral is the analytic

starting point of the proof of the Prime Number Theorem. The integrand $-\zeta'(s)/\zeta(s) \cdot x^s/s$ carries a pole at $s = 1$ coming from the simple pole of $\zeta(s)$ there: since $-\zeta'(s)/\zeta(s) \sim 1/(s-1)$ near $s = 1$, the residue of the integrand at $s = 1$ is $x^1/1 = x$, which is the leading term $\psi(x) \sim x$ once the contour is shifted past this pole. The kernel contributes a further pole at $s = 0$, and the nontrivial zeros of $\zeta(s)$ in the critical strip contribute poles of $-\zeta'(s)/\zeta(s)$ whose residues produce the oscillatory terms $-x^\rho/\rho$ in the explicit formula. Evaluating the shifted integral, which requires the analytic continuation of $\zeta(s)$ past $\operatorname{Re}(s) = 1$ and a zero-free region near that line, is carried out in chapter 5; the present example records only the integral representation that initiates it.

Elementary Results on Prime Distribution

chapter 2 ended with the identity that lies at the center of the subject: for $\operatorname{Re}(s) > 1$,

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

(theorem 2.2.6). The right side is the Dirichlet series of the von Mangoldt function Λ , whose summatory function counts prime powers weighted by $\log p$; the left side is an analytic object whose poles and zeros are accessible to complex analysis. The full exploitation of this identity, locating the pole of $\zeta(s)$ at $s = 1$ and excluding zeros on the line $\operatorname{Re}(s) = 1$, yields the Prime Number Theorem and occupies chapter 5. That route requires contour integration and the analytic continuation of $\zeta(s)$.

This chapter takes the elementary road. Before deploying complex analysis, one can already pin down a great deal about the distribution of primes using nothing more than the unique factorization of integers, the asymptotics of the harmonic sum, and Abel's summation formula from chapter 1. The arguments of Chebyshev fix the correct *order of magnitude* of the prime counting function $\pi(x)$, showing that $\pi(x)$ is bounded above and below by constant multiples of $x/\log x$ without determining the constant. The theorems of Mertens evaluate the key sums over primes, $\sum_{p \leq x} (\log p)/p$ and $\sum_{p \leq x} 1/p$ and $\prod_{p \leq x} (1 - 1/p)$, exactly to leading order. Bertrand's postulate, that a prime always lies between n and $2n$, follows from a careful analysis of the central binomial coefficient. None of these results requires a single contour integral; each rests on counting prime powers inside factorials and binomial coefficients and feeding the count through partial summation. The prerequisites are ?? and the elementary parts of chapter 1, in particular the von Mangoldt function (definition 1.1.8), the summatory-function language (definition 1.3.1), and Abel summation (theorem 1.4.1); the analytic machinery of chapter 2 is not used until the final analytic proof in chapter 5.

3.1 Chebyshev's Functions

The Prime Number Theorem is most often stated as $\pi(x) \sim x/\log x$, a statement about the counting function π . The function π is a step function with a unit jump at each prime, and the irregular spacing of the primes makes its growth difficult to analyze directly. Chebyshev's insight was that two weighted counting functions, attaching the weight $\log p$ to each prime rather than the weight 1, are far more tractable: they are the summatory functions of arithmetic functions already studied in chapter 1, they are tied directly to the factorization of factorials, and their asymptotics are equivalent to those of π . This section introduces π , ϑ , and ψ , records their numerical behavior, relates ψ to ϑ , and proves that the three natural asymptotic statements about them stand or fall together. The equivalence is what licenses the analytic theory of chapter 5 to target the most convenient of the three, ψ , and transfer the conclusion back to π .

Definition 3.1.1: The Prime Counting Function

The *prime counting function* $\pi: [1, \infty) \rightarrow \mathbb{Z}_{\geq 0}$ is

$$\pi(x) = \#\{p \leq x \mid p \text{ prime}\}$$

the number of primes not exceeding x .

The function π has appeared informally since ?? , where Euclid's theorem was read as the statement $\pi(x) \rightarrow \infty$, and again in example 1.4.7, where Abel summation converted π into the sum $\sum_{p \leq x} 1/p$. It is a nondecreasing step function, constant on each interval between consecutive primes and jumping by 1 at each prime. Its central feature is the slowness of its growth: the Prime Number Theorem will show $\pi(x) \sim x/\log x$, so that the proportion $\pi(x)/x$ of integers up to x that are prime decays like $1/\log x$, tending to zero but only logarithmically.

The two functions that organize the elementary theory weight primes by $\log p$.

Definition 3.1.2: Chebyshev's Theta and Psi Functions

Chebyshev's theta function and *Chebyshev's psi function* are

$$\vartheta(x) = \sum_{p \leq x} \log p, \quad \psi(x) = \sum_{n \leq x} \Lambda(n) = \sum_{p^k \leq x} \log p$$

where in the sum for ϑ the index p runs over primes, in the first sum for ψ the index n runs over positive integers with Λ the von Mangoldt function (definition 1.1.8), and in the second sum the indices p, k run over primes p and exponents $k \geq 1$ with $p^k \leq x$.

The two expressions for ψ agree because $\Lambda(n)$ is $\log p$ when $n = p^k$ is a prime power and 0 otherwise, so summing $\Lambda(n)$ over $n \leq x$ is the same as summing $\log p$ over all prime powers $p^k \leq x$. Thus ψ is precisely the summatory function (definition 1.3.1) of the von Mangoldt function, and this is the link to the analytic side: ψ is the partial-sum counterpart of the Dirichlet series $\sum \Lambda(n)n^{-s} = -\zeta'(s)/\zeta(s)$ of theorem 2.2.6. The function ϑ discards the higher prime powers, summing $\log p$ only over primes $p \leq x$ rather than over all prime powers. The difference between the two is therefore the contribution of the proper prime powers p^k with $k \geq 2$, a contribution that proposition 3.1.4 will show is small.

Both functions are weighted forms of π . Where $\pi(x)$ adds 1 for each prime, $\vartheta(x)$ adds $\log p$, so ϑ is

heavier toward the large primes near x ; the weighting is exactly what makes $\vartheta(x)$ grow like x rather than like $x/\log x$. The heuristic is that the primes up to x have “average” logarithm of size $\log x$, so $\vartheta(x) \approx \pi(x) \log x \approx x$, while $\pi(x) \approx \vartheta(x)/\log x \approx x/\log x$. Making this heuristic precise is the content of theorem 3.1.5.

A table of values makes the comparison concrete and exhibits the tracking of $\vartheta(x)$ and $\psi(x)$ with x .

Example 3.1.3: Numerical Values of ϑ , ψ , and π

Direct computation, summing over the primes and prime powers up to each cutoff, gives the following values, rounded to four decimals:

x	$\pi(x)$	$\vartheta(x)$	$\psi(x)$	$\vartheta(x)/x$	$\psi(x)/x$
10	4	5.3471	7.8320	0.5347	0.7832
100	25	83.7284	94.0453	0.8373	0.9405
1000	168	956.2453	996.6809	0.9562	0.9967
10000	1229	9895.9914	10013.3967	0.9896	1.0013

The ratios $\vartheta(x)/x$ and $\psi(x)/x$ both approach 1 as x grows, the convergence of $\psi(x)/x$ being visibly faster, consistent with $\psi(x) - \vartheta(x)$ being a lower-order quantity. At $x = 10^4$ the value $\psi(x)/x = 1.0013$ already sits within two parts in a thousand of its conjectured limit. By contrast $\pi(x)$ is far from x ; it is $x/\log x$ that $\pi(x)$ tracks, and at $x = 10^4$ one has $x/\log x \approx 1085.7$ against $\pi(x) = 1229$, the ratio $\pi(x) \log x/x \approx 1.13$ still drifting slowly toward 1. The contribution of proper prime powers to ψ is small but nonzero: at $x = 100$ the gap $\psi(100) - \vartheta(100) = 10.3169$ comes from the squares 4, 9, 25, 49, the cubes 8, 27, and the higher prime powers 16, 32, 64, 81 below 100.

The table suggests that ψ and ϑ differ by a quantity of lower order. The next result makes this exact, expressing ψ as a sum of ϑ over the roots $x^{1/m}$ and bounding the difference. The estimate is needed to transfer asymptotics between ψ and ϑ , and it must be obtained without assuming any bound on ϑ itself, since the bound $\vartheta(x) = O(x)$ is proved only in the next section; using it here would be circular.

Proposition 3.1.4: Comparison of ψ and ϑ

For $x \geq 1$,

$$\psi(x) = \sum_{m \geq 1} \vartheta(x^{1/m})$$

where the sum is finite, the terms vanishing once $m > \log_2 x$. Moreover

$$0 \leq \psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$$

Proof:

Step 1: The identity. Group the prime powers $p^k \leq x$ appearing in $\psi(x) = \sum_{p^k \leq x} \log p$ according to the exponent $k = m$. For a fixed $m \geq 1$, the primes p contributing are exactly those with $p^m \leq x$, that is, $p \leq x^{1/m}$, and each such prime contributes $\log p$. Hence the contribution of exponent m is $\sum_{p \leq x^{1/m}} \log p = \vartheta(x^{1/m})$, and summing over m gives

$$\psi(x) = \sum_{m \geq 1} \sum_{p \leq x^{1/m}} \log p = \sum_{m \geq 1} \vartheta(x^{1/m})$$

A term $\vartheta(x^{1/m})$ is nonzero precisely when there is a prime $p \leq x^{1/m}$, which requires $x^{1/m} \geq 2$, that is, $2^m \leq x$, equivalently $m \leq \log_2 x$. For $m > \log_2 x$ the argument $x^{1/m}$ is less than 2 and $\vartheta(x^{1/m}) = 0$, so the sum is finite with at most $\lfloor \log_2 x \rfloor$ nonzero terms.

Step 2: Nonnegativity and isolating the difference. The $m = 1$ term is $\vartheta(x)$, so

$$\psi(x) - \vartheta(x) = \sum_{m \geq 2} \vartheta(x^{1/m})$$

Every term $\vartheta(x^{1/m})$ is a sum of values $\log p \geq \log 2 > 0$, hence nonnegative, giving $\psi(x) - \vartheta(x) \geq 0$.

Step 3: The bound, via a trivial estimate for ϑ . The bound $\vartheta(y) = O(y)$ is not yet available, so a cruder estimate is used: each of the $\pi(y)$ primes $p \leq y$ satisfies $\log p \leq \log y$, whence

$$\vartheta(y) = \sum_{p \leq y} \log p \leq \pi(y) \log y \leq y \log y$$

the last step using $\pi(y) \leq y$ (there are at most y positive integers up to y , hence at most y primes). Apply this with $y = x^{1/m}$, so $\vartheta(x^{1/m}) \leq x^{1/m} \log(x^{1/m}) = \frac{1}{m} x^{1/m} \log x$. For $m \geq 2$ one has $x^{1/m} \leq x^{1/2} = \sqrt{x}$, and discarding the factor $1/m \leq 1$ gives the per-term bound $\vartheta(x^{1/m}) \leq \sqrt{x} \log x$. Summing over the at most $\log_2 x$ nonzero terms with $m \geq 2$,

$$0 \leq \psi(x) - \vartheta(x) = \sum_{2 \leq m \leq \log_2 x} \vartheta(x^{1/m}) \leq (\log_2 x) \cdot \sqrt{x} \log x = \frac{\sqrt{x} (\log x)^2}{\log 2}$$

This already gives $\psi(x) - \vartheta(x) = O(\sqrt{x} (\log x)^2)$. The sharper bound $O(\sqrt{x} \log x)$ follows by treating $m = 2$ separately: the single term $\vartheta(\sqrt{x}) \leq \sqrt{x} \log \sqrt{x} = \frac{1}{2} \sqrt{x} \log x$, while for $m \geq 3$ one has $x^{1/m} \leq x^{1/3}$, so

$$\sum_{3 \leq m \leq \log_2 x} \vartheta(x^{1/m}) \leq (\log_2 x) \cdot x^{1/3} \log x = O(x^{1/3} (\log x)^2)$$

and since $x^{1/3} (\log x)^2 = O(\sqrt{x} \log x)$ (because $x^{1/3-1/2} \log x = x^{-1/6} \log x \rightarrow 0$), the $m \geq 3$ tail is absorbed into the $m = 2$ term. Combining the $m = 2$ contribution with the tail gives $\psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$, as we sought to show.

The proposition has two consequences worth separating. First, ψ and ϑ are interchangeable to leading order: since $\sqrt{x} \log x = o(x)$, the relation $0 \leq \psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$ means $\psi(x) = \vartheta(x) + o(x)$, so $\psi(x)/x$ and $\vartheta(x)/x$ have the same limiting behavior and any asymptotic of the form $\psi(x) \sim cx$ holds if and only if $\vartheta(x) \sim cx$ with the same constant. Second, the proper prime powers are genuinely negligible: the entire contribution of squares, cubes, and higher powers of primes to ψ is concentrated in the term $\vartheta(\sqrt{x})$ of size $O(\sqrt{x} \log x)$, dominated overwhelmingly by the squares, since the cubes and beyond contribute only $O(x^{1/3} (\log x)^2)$. This is why $\psi(x)/x$ in example 3.1.3 converges faster than $\vartheta(x)/x$: the higher-power correction it carries is positive and shrinking, nudging ψ above ϑ by an amount that vanishes relative to x . The bound also explains why the analytic theory works with ψ rather than ϑ at no cost in the final statement about primes.

With ψ and ϑ tied together, the remaining task is to connect them to π . The three asymptotic statements, one for each function, turn out to be equivalent, and the bridge between the unweighted π and the weighted ϑ is supplied by Abel summation.

Theorem 3.1.5: Equivalence of the Prime Number Theorem

As $x \rightarrow \infty$, the following three statements are equivalent:

1. $\pi(x) \sim \frac{x}{\log x}$;
2. $\vartheta(x) \sim x$;
3. $\psi(x) \sim x$.

Proof:

The proof establishes the equivalence (ii) $\Leftrightarrow$ (iii) from the comparison of ψ and ϑ , then the equivalence (i) $\Leftrightarrow$ (ii) by partial summation in each direction.

Step 1: (ii) $\Leftrightarrow$ (iii). By proposition 3.1.4,

$$\frac{\psi(x)}{x} = \frac{\vartheta(x)}{x} + \frac{\psi(x) - \vartheta(x)}{x}, \quad 0 \leq \frac{\psi(x) - \vartheta(x)}{x} \leq \frac{C\sqrt{x} \log x}{x} = \frac{C \log x}{\sqrt{x}}$$

for a constant C and all large x . The right-hand bound tends to 0, so $\psi(x)/x$ and $\vartheta(x)/x$ have the same limit if either exists. Hence $\vartheta(x) \sim x$ if and only if $\psi(x) \sim x$.

Step 2: (ii) $\Rightarrow$ (i), *expressing π through ϑ* . Apply Abel summation (theorem 1.4.1) to recover π from ϑ . Take the coefficients $a_n = \log n$ when n is prime and $a_n = 0$ otherwise, so that the summatory function is $A(t) = \sum_{n \leq t} a_n = \vartheta(t)$, and take the test function $f(t) = 1/\log t$, which is continuously differentiable on $[2, \infty)$ with $f'(t) = -1/(t(\log t)^2)$. Then $a_n f(n) = 1$ for each prime n and 0 otherwise, so $\sum_{n \leq x} a_n f(n) = \pi(x) - \pi(2) + 1$; carrying out the summation from the base point 2, theorem 1.4.1 gives

$$\pi(x) = \frac{\vartheta(x)}{\log x} + \int_2^x \frac{\vartheta(t)}{t(\log t)^2} dt + c_0$$

for a bounded constant c_0 absorbing the contribution below 2. Assume now $\vartheta(t) \sim t$, say $\vartheta(t) = t + r(t)$ with $r(t) = o(t)$. The leading term is $\vartheta(x)/\log x \sim x/\log x$. For the integral, split $\vartheta(t) = t + r(t)$:

$$\int_2^x \frac{\vartheta(t)}{t(\log t)^2} dt = \int_2^x \frac{dt}{(\log t)^2} + \int_2^x \frac{r(t)}{t(\log t)^2} dt$$

The first integral satisfies $\int_2^x (\log t)^{-2} dt = O(x/(\log x)^2) = o(x/\log x)$, by the standard estimate that splits the range at $\sqrt{x}$: on $[2, \sqrt{x}]$ the integrand is at most $(\log 2)^{-2}$, contributing $O(\sqrt{x})$, while on $[\sqrt{x}, x]$ one has $\log t \geq \frac{1}{2} \log x$, contributing at most $4x/(\log x)^2$. For the second integral, given $\varepsilon > 0$ choose T with $|r(t)| \leq \varepsilon t$ for $t \geq T$; then $\int_T^x |r(t)|/(t(\log t)^2) dt \leq \varepsilon \int_T^x (\log t)^{-2} dt = \varepsilon \cdot o(x/\log x)$, while $\int_2^T$ is a constant, so the second integral is $o(x/\log x)$ as well. Combining,

$\pi(x) = x/\log x + o(x/\log x)$, which is statement (i).

Step 3: (i) $\Rightarrow$ (ii), *expressing ϑ through π .* Run Abel summation in the reverse direction. Take $a_n = 1$ when n is prime and 0 otherwise, so $A(t) = \pi(t)$, and take $f(t) = \log t$, with $f'(t) = 1/t$. Then $a_n f(n) = \log n$ for each prime n , so $\sum_{n \leq x} a_n f(n) = \vartheta(x)$, and theorem 1.4.1 gives

$$\vartheta(x) = \pi(x) \log x - \int_2^x \frac{\pi(t)}{t} dt$$

Assume now $\pi(t) \sim t/\log t$, say $\pi(t) = t/\log t + e(t)$ with $e(t) = o(t/\log t)$. The leading term is $\pi(x) \log x \sim x$. For the integral, substitute $\pi(t) = t/\log t + e(t)$:

$$\int_2^x \frac{\pi(t)}{t} dt = \int_2^x \frac{dt}{\log t} + \int_2^x \frac{e(t)}{t} dt$$

The first integral is $\text{Li}(x) + O(1)$, and $\text{Li}(x) = O(x/\log x) = o(x)$ by the same split-at- $\sqrt{x}$ estimate applied to $\int_2^x (\log t)^{-1} dt$. For the second integral, given $\varepsilon > 0$ choose T with $|e(t)| \leq \varepsilon t/\log t$ for $t \geq T$; then $\int_T^x |e(t)|/t dt \leq \varepsilon \int_T^x (\log t)^{-1} dt = \varepsilon \cdot o(x)$, and $\int_2^T$ is bounded, so the second integral is $o(x)$. Hence $\vartheta(x) = \pi(x) \log x - o(x) = x + o(x)$, which is statement (ii).

Steps 2 and 3 together give (i) $\Leftrightarrow$ (ii), and Step 1 gives (ii) $\Leftrightarrow$ (iii), so all three statements are equivalent, completing the proof.

The equivalence reorganizes the entire project. Three asymptotic statements that look different, one counting primes, one summing $\log p$ over primes, one summing $\Lambda(n)$ over all integers, are a single statement viewed through three lenses, and the constant in each is forced to be 1 once it is 1 in any one of them. The freedom this buys is decisive for the analytic theory. Of the three functions, ψ is the most natural target for complex analysis: it is the summatory function of Λ , and Λ is exactly the coefficient sequence of $-\zeta'(s)/\zeta(s)$ by theorem 2.2.6, so the asymptotic $\psi(x) \sim x$ is read directly off the analytic behavior of the logarithmic derivative of ζ , with the leading term x produced by the simple pole of $-\zeta'(s)/\zeta(s)$ at $s = 1$ with residue 1. The function π , by contrast, has no such clean Dirichlet series. The strategy of chapter 5 is therefore to prove $\psi(x) \sim x$ analytically and invoke the present theorem to transfer the conclusion first to ϑ and then to the statement $\pi(x) \sim x/\log x$ that one actually wants. The elementary results of the remaining sections of this chapter operate on the same principle from the other side: Chebyshev bounds ψ and ϑ between constant multiples of x , and through this theorem those bounds become bounds on $\pi(x)$ of the correct order $x/\log x$.

3.2 Chebyshev's Estimates

The equivalence of theorem 3.1.5 reduces the Prime Number Theorem to the single asymptotic $\psi(x) \sim x$, but it does not produce that asymptotic; the limit $\psi(x)/x \rightarrow 1$ lies beyond elementary reach and is the work of the analytic theory in chapter 5. What can be extracted by elementary means alone is one step weaker and was obtained by Chebyshev in 1852, half a century before the analytic proof: the *order of magnitude* of the three counting functions is pinned down exactly. There are constants $0 < c_1 < c_2$ with $c_1 x \leq \vartheta(x) \leq c_2 x$, and likewise for ψ , so that $\vartheta(x)$ and $\psi(x)$ each grow linearly in x and $\pi(x)$ grows like $x/\log x$, with explicit numerical constants bracketing the

eventual limit.

The entire argument rests on a single arithmetic object, the central binomial coefficient $\binom{2n}{n}$. Its size is controlled on both sides by elementary bounds, $4^n/(2n+1) \leq \binom{2n}{n} \leq 4^n$, while its prime factorization is controlled by Legendre's formula (proposition 0.2.11): every prime power dividing $\binom{2n}{n}$ is at most $2n$, and every prime in the range $(n, 2n]$ divides it exactly once. Comparing the two descriptions of the same integer forces ψ and ϑ into a linear corridor. The upper bound comes first, in the form of a bound on the product of all primes up to x .

Lemma 3.2.1: Prime Content of the Central Binomial Coefficient

Let $n \geq 1$ and let p be a prime. Then

$$v_p\left(\binom{2n}{n}\right) = \sum_{k \geq 1} \left(\left\lfloor \frac{2n}{p^k} \right\rfloor - 2 \left\lfloor \frac{n}{p^k} \right\rfloor \right)$$

Each summand is 0 or 1, the summand vanishes for $p^k > 2n$, and consequently

$$p^{v_p\left(\binom{2n}{n}\right)} \leq 2n$$

for every prime p . In particular no prime power exceeding $2n$ divides $\binom{2n}{n}$.

Proof:

Step 1: The valuation formula. Since $\binom{2n}{n} = (2n)!/(n!n!)$, the additivity of the p -adic valuation gives $v_p\left(\binom{2n}{n}\right) = v_p((2n)!) - 2v_p(n!)$. Applying Legendre's formula (proposition 0.2.11) to each factorial,

$$v_p\left(\binom{2n}{n}\right) = \sum_{k \geq 1} \left\lfloor \frac{2n}{p^k} \right\rfloor - 2 \sum_{k \geq 1} \left\lfloor \frac{n}{p^k} \right\rfloor = \sum_{k \geq 1} \left(\left\lfloor \frac{2n}{p^k} \right\rfloor - 2 \left\lfloor \frac{n}{p^k} \right\rfloor \right)$$

both Legendre sums being finite, so their term-by-term difference is legitimate.

Step 2: Each summand is 0 or 1. For any real y write $y = \lfloor y \rfloor + \{y\}$ with $0 \leq \{y\} < 1$. Then

$$\lfloor 2y \rfloor - 2\lfloor y \rfloor = \lfloor 2\lfloor y \rfloor + 2\{y\} \rfloor - 2\lfloor y \rfloor = \lfloor 2\{y\} \rfloor$$

using $\lfloor 2y \rfloor = \lfloor 2\lfloor y \rfloor + 2\{y\} \rfloor = 2\lfloor y \rfloor + \lfloor 2\{y\} \rfloor$ since $2\lfloor y \rfloor$ is an integer. As $0 \leq 2\{y\} < 2$, the value $\lfloor 2\{y\} \rfloor$ is 0 when $\{y\} < 1/2$ and 1 when $\{y\} \geq 1/2$. Taking $y = n/p^k$ shows each summand $\lfloor 2n/p^k \rfloor - 2\lfloor n/p^k \rfloor$ equals 0 or 1.

Step 3: The bound on prime powers. A summand can be nonzero only while its terms are nonzero, and $\lfloor 2n/p^k \rfloor = 0$ once $p^k > 2n$. Hence the nonzero summands occur only for those k with $p^k \leq 2n$, that is, $k \leq \log_p(2n)$, and there are at most $\lfloor \log_p(2n) \rfloor$ of them. Since each contributes at most 1,

$$v_p\left(\binom{2n}{n}\right) \leq \#\{k \geq 1 \mid p^k \leq 2n\} = \lfloor \log_p(2n) \rfloor \leq \log_p(2n)$$

Exponentiating base p , the inequality $v_p\left(\binom{2n}{n}\right) \leq \log_p(2n)$ becomes $p^{v_p\left(\binom{2n}{n}\right)} \leq p^{\log_p(2n)} = 2n$, as we sought to show.

The lemma records two facts that will be used in opposite directions. The bound $p^{v_p(\binom{2n}{n})} \leq 2n$ says the central binomial coefficient is, prime by prime, “flat”: no single prime contributes a factor larger than $2n$, no matter how large n is. This is what limits the contribution of small primes when $\binom{2n}{n}$ is decomposed, and it drives the lower bound on ψ . The valuation formula itself, read in the range $n < p \leq 2n$, says something complementary: for such a prime the single term $k = 1$ gives $\lfloor 2n/p \rfloor - 2\lfloor n/p \rfloor = 1 - 0 = 1$ and all higher terms vanish (because $p^2 > n^2 \geq 2n$ for $n \geq 2$), so p divides $\binom{2n}{n}$ exactly once. The primes in $(n, 2n]$ therefore appear in $\binom{2n}{n}$ to the first power, and their product divides it. This second reading is the mechanism behind the primorial bound, to which the discussion now turns.

The upper estimate $\vartheta(x) = O(x)$ is equivalent to a bound on the product of all primes up to x , the *primorial*. Bounding this product is the heart of the upper Chebyshev estimate, and the argument is an induction in which the inductive step is supplied by the elementary inequality $\binom{2m+1}{m} \leq 4^m$ together with the divisibility just observed.

Proposition 3.2.2: The Primorial Bound

For every real $x \geq 1$,

$$\prod_{p \leq x} p \leq 4^x$$

the product running over primes $p \leq x$. Equivalently, $\vartheta(x) \leq (\log 4)x$ for all $x \geq 1$.

Proof:

The two formulations are equivalent: taking logarithms turns $\prod_{p \leq x} p \leq 4^x$ into $\sum_{p \leq x} \log p \leq x \log 4$, and the left side is exactly $\vartheta(x)$ by definition 3.1.2. It therefore suffices to prove the product bound. Write $\Pi(x) = \prod_{p \leq x} p$.

Step 1: Reduction to integers. The product $\Pi(x)$ depends only on $\lfloor x \rfloor$, since no prime lies strictly between $\lfloor x \rfloor$ and x . The target 4^x is nondecreasing in x , so if $\Pi(n) \leq 4^n$ holds for every integer $n \geq 1$ then for general $x \geq 1$ with $n = \lfloor x \rfloor$,

$$\Pi(x) = \Pi(n) \leq 4^n \leq 4^x$$

The claim is thus reduced to the inequality $\Pi(n) \leq 4^n$ for integers $n \geq 1$, which is proved by strong induction on n .

Step 2: Base cases. For $n = 1$ the product is empty, $\Pi(1) = 1 \leq 4$. For $n = 2$, $\Pi(2) = 2 \leq 16$. These anchor the induction; the inductive step below treats $n \geq 3$ and refers back to strictly smaller indices.

Step 3: Even n . Suppose $n = 2m$ with $m \geq 2$, so $n \geq 4$. Since $2m$ is composite for $m \geq 2$, it is not prime, and no prime satisfies $2m - 1 < p \leq 2m$. Hence $\Pi(2m) = \Pi(2m - 1)$. By the induction hypothesis applied to $2m - 1 < 2m$,

$$\Pi(2m) = \Pi(2m - 1) \leq 4^{2m-1} \leq 4^{2m}$$

so the bound for $n = 2m$ follows from the bound for the smaller odd index $2m - 1$.

Step 4: Odd n . Suppose $n = 2m + 1$ with $m \geq 1$, so $n \geq 3$. Split the primes $p \leq 2m + 1$ at $m + 1$:

$$\Pi(2m + 1) = \left(\prod_{p \leq m+1} p \right) \left(\prod_{m+1 < p \leq 2m+1} p \right) = \Pi(m + 1) \cdot \prod_{m+1 < p \leq 2m+1} p$$

The first factor is bounded by the induction hypothesis at $m + 1 \leq 2m < 2m + 1$:

$$\Pi(m + 1) \leq 4^{m+1}$$

For the second factor, every prime p with $m + 1 < p \leq 2m + 1$ divides the integer $\binom{2m+1}{m} = (2m + 1)! / (m! (m + 1)!)$. Indeed such a p divides the numerator $(2m + 1)!$, since $p \leq 2m + 1$, but divides neither $m!$ nor $(m + 1)!$, since $p > m + 1$ exceeds every factor in those products; by the additivity of v_p , $v_p \binom{2m+1}{m} = v_p((2m + 1)!) \geq 1$. These primes are distinct, so their product divides $\binom{2m+1}{m}$, whence

$$\prod_{m+1 < p \leq 2m+1} p \leq \binom{2m+1}{m}$$

It remains to bound the binomial coefficient. The two coefficients $\binom{2m+1}{m}$ and $\binom{2m+1}{m+1}$ are equal and both appear in the binomial expansion $\sum_{j=0}^{2m+1} \binom{2m+1}{j} = 2^{2m+1}$, so

$$2 \binom{2m+1}{m} = \binom{2m+1}{m} + \binom{2m+1}{m+1} \leq 2^{2m+1}$$

giving $\binom{2m+1}{m} \leq 2^{2m} = 4^m$. Combining the three bounds,

$$\Pi(2m + 1) = \Pi(m + 1) \cdot \prod_{m+1 < p \leq 2m+1} p \leq 4^{m+1} \cdot 4^m = 4^{2m+1}$$

which is the bound for $n = 2m + 1$. The even and odd steps together complete the induction, completing the proof.

The primorial bound is the upper half of the Chebyshev estimates already in finished form: $\vartheta(x) \leq (\log 4)x$ is a linear bound with the explicit constant $\log 4 \approx 1.386$. The inductive engine is the same mechanism that reappears, sharpened, in the proof of Bertrand's postulate (theorem 3.4.3) later in this chapter. The primes in the window $(m + 1, 2m + 1]$ are forced into the central binomial coefficient, whose size is capped by 4^m ; the product of those primes therefore cannot grow faster than 4^m , and feeding this back through the recursion bounds the product of all primes. The lower estimate runs the comparison in reverse: rather than asking how few primes can fit inside $\binom{2n}{n}$, it asks how the known size $\binom{2n}{n} \geq 4^n / (2n + 1)$ must be accounted for by primes, each capped at $2n$ by lemma 3.2.1.

Theorem 3.2.3: Chebyshev's Estimates

There exist absolute constants $0 < c_1 < c_2$ such that for all $x \geq 2$,

$$c_1 x \leq \vartheta(x) \leq \psi(x) \leq c_2 x$$

Explicitly:

1. (*Upper bound.*) $\vartheta(x) \leq (\log 4)x$ for all $x \geq 1$, and $\psi(x) \leq (\log 4)x + O(\sqrt{x} \log x)$, so $\psi(x) \leq c_2 x$ holds for any fixed $c_2 > \log 4$ and all large x .
2. (*Lower bound.*) $\psi(x) \geq (\log 2)x - O(\log x)$ and $\vartheta(x) \geq (\log 2)x - O(\sqrt{x} \log x)$, so each of $\psi(x) \geq c_1 x$ and $\vartheta(x) \geq c_1 x$ holds for any fixed $c_1 < \log 2$ and all large x .

In particular $\vartheta(x)$ and $\psi(x)$ are each of exact order x , with $\log 2 \leq \liminf \psi(x)/x \leq \limsup \psi(x)/x \leq \log 4$ and the same for ϑ .

Proof:

The relation $\psi(x) \geq \vartheta(x)$ is available throughout: proposition 3.1.4 gives $\psi(x) - \vartheta(x) = \sum_{m \geq 2} \vartheta(x^{1/m}) \geq 0$, together with the estimate $0 \leq \psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$, which transfers any bound between ψ and ϑ up to a $o(x)$ error. The two sub-statements are proved in turn.

1. *Upper bound.* proposition 3.2.2 gives $\vartheta(x) \leq (\log 4)x$ for all $x \geq 1$ directly. For ψ , the comparison $\psi(x) = \vartheta(x) + (\psi(x) - \vartheta(x))$ with $0 \leq \psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$ yields

$$\psi(x) \leq (\log 4)x + O(\sqrt{x} \log x)$$

Since $\sqrt{x} \log x = o(x)$, for any fixed $c_2 > \log 4$ the error is eventually dominated by $(c_2 - \log 4)x$, so $\psi(x) \leq c_2 x$ for all large x , and after enlarging c_2 once more the bound $\psi(x) \leq c_2 x$ holds for all $x \geq 2$. This is the right-hand inequality with c_2 an absolute constant.

2. *Lower bound.* The argument compares two descriptions of the central binomial coefficient $N = \binom{2n}{n}$ for an integer $n \geq 1$, then passes from even integers to all x .

Size of N . The central coefficient is the largest of the $2n + 1$ terms $\binom{2n}{j}$ that sum to $(1 + 1)^{2n} = 4^n$, so

$$N = \binom{2n}{n} \geq \frac{1}{2n + 1} \sum_{j=0}^{2n} \binom{2n}{j} = \frac{4^n}{2n + 1}$$

Taking logarithms, $\log N \geq n \log 4 - \log(2n + 1) = 2n \log 2 - \log(2n + 1)$.

Primes in N . By unique factorization $N = \prod_{p \leq 2n} p^{v_p(N)}$, the product over primes $p \leq 2n$ because lemma 3.2.1 forbids any prime power exceeding $2n$ from dividing N . For each such prime set $K_p = v_p(N)$; the same lemma gives $p^k \leq 2n$ for every k with $1 \leq k \leq K_p$. Hence

$$\log N = \sum_{p \leq 2n} K_p \log p = \sum_{p \leq 2n} \sum_{k=1}^{K_p} \log p \leq \sum_{\substack{p, k \geq 1 \\ p^k \leq 2n}} \log p = \psi(2n)$$

where the inequality holds because each pair (p, k) with $1 \leq k \leq K_p$ satisfies $p^k \leq 2n$ and so appears among the prime powers counted by $\psi(2n)$, while the right-hand sum may contain additional nonnegative terms. Combining the two displays,

$$\psi(2n) \geq \log N \geq 2n \log 2 - \log(2n + 1)$$

for every integer $n \geq 1$.

Passage to all x . Given a real $x \geq 2$, set $n = \lfloor x/2 \rfloor \geq 1$, so $2n \leq x$ and $2n > x - 2$. As ψ is nondecreasing, $\psi(x) \geq \psi(2n) \geq 2n \log 2 - \log(2n + 1)$, and using $2n > x - 2$ together with $2n + 1 \leq x + 1 \leq 2x$,

$$\psi(x) \geq (x - 2) \log 2 - \log(2x) = (\log 2)x - (2 \log 2 + \log(2x))$$

The subtracted quantity is $O(\log x)$, giving $\psi(x) \geq (\log 2)x - O(\log x)$. For ϑ , the estimate $0 \leq \psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$ of proposition 3.1.4 yields $\vartheta(x) = \psi(x) - O(\sqrt{x} \log x) \geq (\log 2)x - O(\sqrt{x} \log x)$. Since both error terms are $o(x)$, for any fixed $c_1 < \log 2$ each of $\psi(x) \geq c_1 x$ and $\vartheta(x) \geq c_1 x$ holds for all large x .

Sub-statements (1) and (2) together furnish constants $0 < c_1 < c_2$ with $c_1 x \leq \vartheta(x) \leq \psi(x) \leq c_2 x$ for all $x \geq 2$, so both functions are of exact order x , as we sought to show.

The estimates fix the order of growth without fixing the constant. The corridor they cut out, $(\log 2)x \leq \psi(x) \leq (\log 4)x$ to leading order, is wide: the lower constant $\log 2 \approx 0.693$ and the upper constant $\log 4 \approx 1.386$ differ by a factor of exactly two, and the truth $\psi(x) \sim x$ of chapter 5 sits strictly between them. Chebyshev's method, resting only on the size and factorization of one binomial coefficient, cannot close this gap; the two constants are artifacts of comparing $\binom{2n}{n}$ against 4^n on one side and against the cap $2n$ on prime powers on the other, and no elementary sharpening of these comparisons reaches the constant 1. That value is analytic in origin, produced by the simple pole of $-\zeta'(s)/\zeta(s)$ at $s = 1$ with residue 1. What the estimates do settle is decisive for everything that follows: ψ , ϑ , and (through the next corollary) π have no anomalous growth, no logarithmic or power-law deviation from linear order, and any sum over primes built from these functions by partial summation inherits a bound of the correct shape.

The order of π now follows by converting the bounds on ϑ into bounds on π , exactly as in the proof of theorem 3.1.5 but at the level of inequalities rather than asymptotic equivalences. This is the bound foreshadowed in ?? , where $\pi(x) = O(x/\log x)$ was promised as a consequence of Chebyshev's work.

Corollary 3.2.4: The Order of the Prime Counting Function

The prime counting function is of exact order $x/\log x$: there are absolute constants $0 < a_1 < a_2$ with

$$a_1 \frac{x}{\log x} \leq \pi(x) \leq a_2 \frac{x}{\log x}$$

for all $x \geq 2$. More precisely, the constants may be taken arbitrarily close to $\log 2$ and $\log 4$:

$$(\log 2 + o(1)) \frac{x}{\log x} \leq \pi(x) \leq (\log 4 + o(1)) \frac{x}{\log x}$$

as $x \rightarrow \infty$. In particular $\pi(x) = O(x/\log x)$.

Proof:

The two inequalities come from the two bounds of theorem 3.2.3, transferred through the relation between π and ϑ established in theorem 3.1.5.

Step 1: The upper bound on π . Comparing π and ϑ directly, the primes in the range $\sqrt{x} < p \leq x$ each contribute a logarithm at least $\log \sqrt{x} = \frac{1}{2} \log x$, so

$$\vartheta(x) = \sum_{p \leq x} \log p \geq \sum_{\sqrt{x} < p \leq x} \log p \geq (\pi(x) - \pi(\sqrt{x})) \cdot \frac{1}{2} \log x$$

Rearranging and using $\pi(\sqrt{x}) \leq \sqrt{x}$,

$$\pi(x) \leq \frac{2\vartheta(x)}{\log x} + \pi(\sqrt{x}) \leq \frac{2\vartheta(x)}{\log x} + \sqrt{x}$$

By the upper bound $\vartheta(x) \leq (\log 4)x$ of theorem 3.2.3, the first term is at most $2(\log 4)x/\log x$, and the second satisfies $\sqrt{x} = o(x/\log x)$. Hence $\pi(x) \leq 2(\log 4)x/\log x + o(x/\log x)$, giving $\pi(x) = O(x/\log x)$ with an absolute constant a_2 . The sharper constant $\log 4 + o(1)$ is obtained below by partial summation.

Step 2: The lower bound on π . In the other direction, weighting each prime by at most $\log x$ gives

$$\vartheta(x) = \sum_{p \leq x} \log p \leq \sum_{p \leq x} \log x = \pi(x) \log x$$

whence $\pi(x) \geq \vartheta(x)/\log x$. By the lower bound $\vartheta(x) \geq (\log 2)x - O(\sqrt{x} \log x)$ of theorem 3.2.3,

$$\pi(x) \geq \frac{\vartheta(x)}{\log x} \geq \frac{(\log 2)x}{\log x} - O(\sqrt{x}) = (\log 2 + o(1)) \frac{x}{\log x}$$

since $O(\sqrt{x}) = o(x/\log x)$. This gives the lower inequality with constant a_1 arbitrarily close to $\log 2$.

Step 3: Sharpening the upper constant by partial summation. The crude factor 2 in Step 1 is removed by recovering π from ϑ through Abel summation (theorem 1.4.1), the same device used

in theorem 3.1.5. With $a_n = \log n$ for n prime and $a_n = 0$ otherwise, so that $A(t) = \vartheta(t)$, and with $f(t) = 1/\log t$ on $[2, \infty)$, where $f'(t) = -1/(t(\log t)^2)$, theorem 1.4.1 gives

$$\pi(x) = \frac{\vartheta(x)}{\log x} + \int_2^x \frac{\vartheta(t)}{t(\log t)^2} dt + c_0$$

with c_0 a bounded constant. The bound $\vartheta(t) \leq (\log 4)t$ from theorem 3.2.3 controls both pieces: the leading term is at most $(\log 4)x/\log x$, and the integral is at most

$$(\log 4) \int_2^x \frac{dt}{(\log t)^2} = (\log 4) \cdot O\left(\frac{x}{(\log x)^2}\right) = o\left(\frac{x}{\log x}\right)$$

by the split-at- $\sqrt{x}$ estimate for $\int_2^x (\log t)^{-2} dt$ recorded in the proof of theorem 3.1.5. Therefore $\pi(x) \leq (\log 4)x/\log x + o(x/\log x) = (\log 4 + o(1))x/\log x$, which is the sharpened upper bound. Steps 2 and 3 together give the two-sided estimate with constants approaching $\log 2$ and $\log 4$, as we sought to show.

The corollary is the elementary substitute for the Prime Number Theorem, and it suffices for many purposes that do not require the exact constant. The proportion $\pi(x)/x$ of integers up to x that are prime lies between $(\log 2)/\log x$ and $(\log 4)/\log x$ for large x , so it decays like $1/\log x$, confirming the logarithmic thinning of the primes announced when π was first defined in definition 3.1.1. The bound $\pi(x) = O(x/\log x)$ is precisely the statement deferred in ??, now established without any appeal to complex analysis. Its converse direction, $\pi(x) \gg x/\log x$, is equally elementary and is the quantitative form of Euclid's theorem: not only are there infinitely many primes, but they are at least a constant multiple of $x/\log x$ in number up to x . The linear bound $\vartheta(x) = O(x)$ is also the input that Mertens' theorems require, converting by partial summation into exact leading-order evaluations of $\sum_{p \leq x} (\log p)/p$ and $\sum_{p \leq x} 1/p$.

The width of the Chebyshev corridor is best appreciated numerically, against the limit that the Prime Number Theorem will eventually supply.

Example 3.2.5: Chebyshev's Constants versus the Truth

The two constants delivered by theorem 3.2.3 are

$$\log 2 = 0.69314\dots, \quad \log 4 = 2 \log 2 = 1.38629\dots$$

bracketing the eventual limit $\psi(x)/x \rightarrow 1$ of chapter 5 from below and above. The bracket for ψ is direct: the theorem gives $(\log 2 + o(1))x \leq \psi(x) \leq (\log 4)x$, and the table of example 3.1.3 shows $\psi(x)/x$ already at 0.9405 for $x = 100$ and 1.0013 for $x = 10^4$, comfortably inside $[0.693, 1.386]$ and converging to 1 rather than to either Chebyshev constant.

For π , the corresponding bracket from corollary 3.2.4 is $(\log 2 + o(1))x/\log x \leq \pi(x) \leq (\log 4 + o(1))x/\log x$. Computing the ratio $\pi(x) \log x/x$, whose limit the Prime Number Theorem asserts

to be 1:

x	$\pi(x)$	$\frac{x}{\log x}$	$\frac{\pi(x) \log x}{x}$
10^2	25	21.71	1.1513
10^3	168	144.76	1.1605
10^4	1229	1085.74	1.1320
10^5	9592	8685.89	1.1043
10^6	78498	72382.41	1.0845

Every entry in the last column lies strictly between $\log 2 = 0.693$ and $\log 4 = 1.386$, as Chebyshev's bounds require, and the column drifts slowly toward 1, the drift being the slow $o(1)$ correction that only the analytic theory resolves. The gap between Chebyshev's two constants is a factor of 2, whereas the data already pin the true ratio to within roughly ten percent of 1 by $x = 10^6$; the elementary method captures the order of magnitude but not this finer convergence.

3.3 Mertens' Theorems

Chebyshev's estimates of section 3.2 fix the order of magnitude of ψ , ϑ , and π but not the leading constant. The three theorems of Mertens, proved in 1874 by the same elementary toolkit, achieve something the order-of-magnitude bounds cannot: they evaluate three sums over primes *exactly to leading order*, with an explicit error term and an explicit constant. The sums in question are

$$\sum_{p \leq x} \frac{\log p}{p}, \quad \sum_{p \leq x} \frac{1}{p}, \quad \prod_{p \leq x} \left(1 - \frac{1}{p}\right)$$

and Mertens' theorems assert, respectively, that the first is $\log x + O(1)$, the second is $\log \log x + M + O(1/\log x)$ for a specific constant M , and the third is asymptotic to $e^{-\gamma}/\log x$. Each statement is sharper than a bound of the form " $\asymp \log x$ " or " $\asymp 1/\log x$ ": the first pins the coefficient of $\log x$ to be exactly 1, the second produces the second-order constant M , and the third produces the constant $e^{-\gamma}$ with the Euler–Mascheroni γ in the exponent.

What makes this reach possible without analytic continuation is that Chebyshev's bound $\psi(x) = O(x)$ already controls the error when the factorization of $[x]!$ is fed through Abel summation. The route is largely uniform: an exact identity for $\log([x]!)$ combined with Stirling's formula gives Mertens' first theorem; partial summation against the weight $1/\log t$ converts the first theorem into the second; comparison of $\sum 1/p$ with $-\sum \log(1 - 1/p)$ reduces the constant M to the Euler–Mascheroni γ ; and exponentiating the second theorem collapses that closed form to $e^{-\gamma}$, yielding the third. Only one step reaches past the Chebyshev-level toolkit, the identification of the additive constant as exactly γ rather than an unspecified number: it draws on the Euler product and the behavior of ζ near its pole, both already in hand from chapter 2 and used only in the convergent range $\text{Re } s > 1$, and it turns on a single Gamma-function identity isolated and cited at the point of use. The constant M produced here is the same $B = \gamma + \sum_p (\log(1 - 1/p) + 1/p) = 0.2614\dots$ that appeared in the average order of ω (proposition 1.3.9); this section establishes that identification.

The starting point is an exact arithmetic identity for the logarithm of a factorial, the same factorization idea used for the central binomial coefficient in section 3.2, now applied to $[x]!$ itself. It expresses $\log([x]!)$ as a weighted sum of the von Mangoldt function (definition 1.1.8), and pairing it with Stirling's formula (theorem 1.5.7) gives the asymptotic that drives the first theorem.

Lemma 3.3.1: The Prime-Power Decomposition of $\log n!$

For every real $x \geq 1$,

$$\log([x]!) = \sum_{n \leq x} \Lambda(n) \left\lfloor \frac{x}{n} \right\rfloor$$

where Λ is the von Mangoldt function and the sum runs over positive integers $n \leq x$. Consequently

$$\log([x]!) = x \log x - x + O(\log x)$$

as $x \rightarrow \infty$.

Proof:

Write $N = [x]$, so that $[x]! = N!$ and $[x/n] = [N/n]$ for every positive integer n (both count the multiples of n in $\{1, \dots, N\}$, and no multiple of n lies strictly between N and x). The two displayed claims are proved in turn.

Step 1: The identity. By unique factorization, $N! = \prod_{p \leq N} p^{v_p(N!)}$, the product over primes $p \leq N$. Taking logarithms and applying Legendre's formula (proposition 0.2.11) to each exponent,

$$\log(N!) = \sum_{p \leq N} v_p(N!) \log p = \sum_{p \leq N} \left(\sum_{k \geq 1} \left\lfloor \frac{N}{p^k} \right\rfloor \right) \log p = \sum_{p \leq N} \sum_{k \geq 1} \left\lfloor \frac{N}{p^k} \right\rfloor \log p$$

Each Legendre sum is finite (the terms vanish once $p^k > N$), so the double sum has finitely many nonzero terms and may be reindexed. Group the terms by the value $n = p^k$ of the prime power. For each prime power $n = p^k \leq N$ the corresponding summand is $[N/n] \log p$, and $\log p = \Lambda(n)$ by the definition of the von Mangoldt function (definition 1.1.8); for every $n \leq N$ that is *not* a prime power, $\Lambda(n) = 0$ contributes nothing. Hence

$$\log(N!) = \sum_{\substack{n \leq N \\ n=p^k}} \left\lfloor \frac{N}{n} \right\rfloor \log p = \sum_{n \leq N} \Lambda(n) \left\lfloor \frac{N}{n} \right\rfloor = \sum_{n \leq x} \Lambda(n) \left\lfloor \frac{x}{n} \right\rfloor$$

the last equality by $[N/n] = [x/n]$ and $N = [x]$. This is the identity.

Step 2: The asymptotic via Stirling. Stirling's formula (theorem 1.5.7) gives $\log(N!) = N \log N - N + \frac{1}{2} \log(2\pi N) + O(1/N)$, so in particular $\log(N!) = N \log N - N + O(\log N)$. It remains to replace $N = [x]$ by x at a cost of $O(\log x)$. Write $x = N + \theta$ with $0 \leq \theta < 1$. Then

$$x \log x - x = (N + \theta) \log(N + \theta) - (N + \theta)$$

The function $t \mapsto t \log t - t$ has derivative $\log t$, so by the mean value theorem $x \log x - x - (N \log N - N) = \theta \log \xi$ for some $\xi \in [N, x]$, whence

$$|(x \log x - x) - (N \log N - N)| = \theta \log \xi \leq \log(N + 1) = O(\log x)$$

Combining with $\log(N!) = N \log N - N + O(\log N)$ and $\log N = O(\log x)$,

$$\log([x]!) = N \log N - N + O(\log N) = x \log x - x + O(\log x)$$

as we sought to show.

The identity is the elementary counterpart of the analytic identity $-\zeta'(s)/\zeta(s) = \sum_n \Lambda(n)n^{-s}$ from theorem 2.2.6: both extract the von Mangoldt function from a global object, there the logarithmic derivative of ζ , here the logarithm of a factorial. The factor $\lfloor x/n \rfloor$ counts how many of the integers $1, \dots, \lfloor x \rfloor$ are divisible by n , so the identity says that the total logarithm $\log(\lfloor x \rfloor!) = \sum_{m \leq x} \log m$ is reorganized by collecting, for each prime power $n = p^k$, the contribution $\log p$ once for every multiple of n up to x . Stirling's formula then evaluates the left side to within $O(\log x)$, and the resulting estimate $\sum_{n \leq x} \Lambda(n) \lfloor x/n \rfloor = x \log x - x + O(\log x)$ is the bridge to the first theorem: the floor $\lfloor x/n \rfloor$ is within 1 of x/n , and replacing it by x/n produces exactly the sum $\sum \Lambda(n)/n$ whose evaluation is sought.

Theorem 3.3.2: Mertens' First Theorem

As $x \rightarrow \infty$,

$$\sum_{n \leq x} \frac{\Lambda(n)}{n} = \log x + O(1)$$

and equivalently, summing over primes only,

$$\sum_{p \leq x} \frac{\log p}{p} = \log x + O(1)$$

Proof:

The two statements differ by the contribution of the proper prime powers, treated at the end. The main work is the first estimate.

Step 1: Removing the floor. Start from the identity of lemma 3.3.1, $\log(\lfloor x \rfloor!) = \sum_{n \leq x} \Lambda(n) \lfloor x/n \rfloor$. Writing $\lfloor x/n \rfloor = x/n - \{x/n\}$ with $0 \leq \{x/n\} < 1$ separates the main term from an error:

$$\sum_{n \leq x} \Lambda(n) \left\lfloor \frac{x}{n} \right\rfloor = x \sum_{n \leq x} \frac{\Lambda(n)}{n} - \sum_{n \leq x} \Lambda(n) \left\{ \frac{x}{n} \right\}$$

The error sum is controlled by Chebyshev's bound. Since $0 \leq \{x/n\} < 1$ and $\Lambda(n) \geq 0$,

$$0 \leq \sum_{n \leq x} \Lambda(n) \left\{ \frac{x}{n} \right\} \leq \sum_{n \leq x} \Lambda(n) = \psi(x)$$

by the definition of ψ (definition 3.1.2), and $\psi(x) = O(x)$ by Chebyshev's estimates (theorem 3.2.3). Hence the error sum is $O(x)$.

Step 2: Solving for the sum. Combining Step 1 with the asymptotic $\log(\lfloor x \rfloor!) = x \log x - x + O(\log x)$ of lemma 3.3.1,

$$x \sum_{n \leq x} \frac{\Lambda(n)}{n} = \log(\lfloor x \rfloor!) + \sum_{n \leq x} \Lambda(n) \left\{ \frac{x}{n} \right\} = (x \log x - x + O(\log x)) + O(x) = x \log x + O(x)$$

the term $-x$ and the error $O(\log x)$ both being absorbed into $O(x)$. Dividing by x ,

$$\sum_{n \leq x} \frac{\Lambda(n)}{n} = \log x + O(1)$$

which is the first statement.

Step 3: Passage to primes. The sum $\sum_{n \leq x} \Lambda(n)/n$ runs over all prime powers; isolating the primes ($k = 1$) leaves the proper prime powers ($k \geq 2$):

$$\sum_{n \leq x} \frac{\Lambda(n)}{n} = \sum_{p \leq x} \frac{\log p}{p} + \sum_{p^k \leq x, k \geq 2} \frac{\log p}{p^k}$$

The second sum is bounded by the full convergent series over all $k \geq 2$ and all primes:

$$0 \leq \sum_{\substack{p^k \leq x \\ k \geq 2}} \frac{\log p}{p^k} \leq \sum_p \log p \sum_{k \geq 2} \frac{1}{p^k} = \sum_p \log p \cdot \frac{1/p^2}{1 - 1/p} = \sum_p \frac{\log p}{p(p-1)}$$

using the geometric series $\sum_{k \geq 2} p^{-k} = p^{-2}/(1 - p^{-1}) = 1/(p(p-1))$. The final series converges: its terms satisfy $\log p/(p(p-1)) \leq \log p/(p-1)^2 \leq C/n^{3/2}$ for a constant C and all $n = p$, since $\log p = O(p^{1/2})$, and $\sum_n n^{-3/2} < \infty$. Therefore the proper-prime-power contribution is a bounded quantity, $O(1)$, and subtracting it from $\sum_{n \leq x} \Lambda(n)/n = \log x + O(1)$ leaves

$$\sum_{p \leq x} \frac{\log p}{p} = \log x + O(1)$$

completing the proof.

The first theorem already separates the elementary theory from the order-of-magnitude estimates of the previous section: where Chebyshev gives $\sum_{p \leq x} (\log p)/p \asymp \log x$ with unspecified constant, Mertens fixes the coefficient of $\log x$ to be exactly 1. The reason the constant is accessible while $\psi(x) \sim x$ is not lies in the weighting. Dividing $\Lambda(n)$ by n converts the identity $\sum \Lambda(n)[x/n] = x \log x - x + O(\log x)$, whose main term is of size $x \log x$, into a statement whose main term is of size $\log x$; the Chebyshev error $\psi(x) = O(x)$, fatal at the scale of $x \log x$, becomes a harmless $O(1)$ after division by x . The first theorem is thus an averaged statement, robust to the $O(x)$ uncertainty in ψ that the elementary method cannot remove. It is the input from which the remaining two theorems follow by partial summation and exponentiation, with no further arithmetic.

The passage from $\sum (\log p)/p$ to $\sum 1/p$ is a textbook application of Abel summation: the summand $1/p$ is $(\log p)/p$ divided by $\log p$, so summing against the weight $1/\log t$ against the known partial sums of $(\log p)/p$ recovers $\sum_{p \leq x} 1/p$. This is the device foreshadowed at corollary 1.4.2 in chapter 1.

Theorem 3.3.3: Mertens' Second Theorem

There is a constant M such that, as $x \rightarrow \infty$,

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + M + O\left(\frac{1}{\log x}\right)$$

The constant M is the *Mertens constant*, identified in closed form in definition 3.3.4.

Proof:

Step 1: Setting up Abel summation. Define the coefficients $a_n = (\log n)/n$ when n is prime and $a_n = 0$ otherwise, so that the summatory function is

$$A(t) = \sum_{n \leq t} a_n = \sum_{p \leq t} \frac{\log p}{p}$$

By theorem 3.3.2, $A(t) = \log t + R(t)$ with $R(t) = O(1)$; fix a constant C_0 with $|R(t)| \leq C_0$ for all $t \geq 2$. Take the test function $f(t) = 1/\log t$ on $[2, \infty)$, which is continuously differentiable with $f'(t) = -1/(t(\log t)^2)$. For a prime $n = p$, the product $a_n f(n) = \frac{\log p}{p} \cdot \frac{1}{\log p} = \frac{1}{p}$, so that $\sum_{n \leq x} a_n f(n) = \sum_{p \leq x} 1/p$.

Step 2: Applying the formula. Abel summation (theorem 1.4.1), carried out from the base point 2 rather than 1 (no prime lies below 2, so $A(t) = 0$ on $[1, 2)$ and the integral over $[1, 2)$ vanishes), gives

$$\sum_{p \leq x} \frac{1}{p} = \frac{A(x)}{\log x} + \int_2^x \frac{A(t)}{t(\log t)^2} dt$$

Substitute $A(t) = \log t + R(t)$ into both pieces. The boundary term is

$$\frac{A(x)}{\log x} = \frac{\log x + R(x)}{\log x} = 1 + O\left(\frac{1}{\log x}\right)$$

since $R(x) = O(1)$. The integral splits as

$$\int_2^x \frac{\log t + R(t)}{t(\log t)^2} dt = \int_2^x \frac{dt}{t \log t} + \int_2^x \frac{R(t)}{t(\log t)^2} dt$$

Step 3: The two integrals. The first integral is elementary: with the substitution $u = \log t$, $du = dt/t$,

$$\int_2^x \frac{dt}{t \log t} = \int_{\log 2}^{\log x} \frac{du}{u} = \log \log x - \log \log 2$$

For the second integral, the integrand is absolutely bounded by $C_0/(t(\log t)^2)$, and

$$\int_2^\infty \frac{dt}{t(\log t)^2} = \int_{\log 2}^\infty \frac{du}{u^2} = \frac{1}{\log 2} < \infty$$

so $\int_2^\infty R(t)/(t(\log t)^2) dt$ converges absolutely to some real number, call it I . The tail past x is estimated the same way:

$$\left| \int_x^\infty \frac{R(t)}{t(\log t)^2} dt \right| \leq C_0 \int_x^\infty \frac{dt}{t(\log t)^2} = \frac{C_0}{\log x} = O\left(\frac{1}{\log x}\right)$$

whence $\int_2^x R(t)/(t(\log t)^2) dt = I + O(1/\log x)$.

Step 4: Assembling the constant. Collecting the boundary term and the two integrals,

$$\sum_{p \leq x} \frac{1}{p} = \left(1 + O\left(\frac{1}{\log x}\right)\right) + (\log \log x - \log \log 2) + \left(I + O\left(\frac{1}{\log x}\right)\right)$$

Setting $M = 1 - \log \log 2 + I$, the constant terms combine and the two error terms merge into a single $O(1/\log x)$, giving

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + M + O\left(\frac{1}{\log x}\right)$$

as we sought to show.

The second theorem is the quantitative form of the divergence of $\sum 1/p$, Euler's theorem that the primes are "dense enough" for their reciprocals to sum to infinity. The growth rate is now exact: the partial sums increase like $\log \log x$, doubly logarithmically, so the divergence is extraordinarily slow. To reach $\sum_{p \leq x} 1/p = 4$ requires x on the order of $e^{e^4} \approx 5 \times 10^{23}$. The presence of the second-order constant M is what distinguishes Mertens' statement from the bare divergence: $\sum_{p \leq x} 1/p - \log \log x$ converges to a definite limit M , at the rate $O(1/\log x)$. The proof produced $M = 1 - \log \log 2 + I$ as an integral expression depending on the error term $R(t)$ in the first theorem; this form is correct but opaque. The next definition gives the closed form, the same constant B that already appeared in proposition 1.3.9 as the second-order term in the average of ω , and identifying the two is the substantive content of what follows.

The constant M in the second theorem was produced as $1 - \log \log 2 + I$, with I an integral of the first-theorem error. Establishing its closed form proceeds in two stages. Comparing the divergent sum $\sum 1/p$ with the divergent sum $-\sum \log(1 - 1/p)$, whose difference is the convergent prime correction S , isolates the part of M contributed by the higher prime powers. The Euler–Mascheroni γ then enters separately, through the behavior of the zeta function at its pole, $\zeta(1 + \rho) = 1/\rho + \gamma + o(1)$, where the γ is the same harmonic-sum constant already computed in example 1.4.4; matching the smoothly damped prime sum that ζ governs against the sharp partial sums of $\sum 1/p$ is what pins the additive constant to γ .

Definition 3.3.4: The Mertens Constant

The *Mertens constant* is

$$M = \gamma + \sum_p \left(\log \left(1 - \frac{1}{p} \right) + \frac{1}{p} \right)$$

where γ is the Euler–Mascheroni constant (example 1.4.4) and the sum runs over all primes. This is the same constant as the B of proposition 1.3.9, $M = B = 0.2614\dots$, and it equals the constant M appearing in Mertens' second theorem (theorem 3.3.3).

The series defining M converges absolutely: for $|t| \leq 1/2$ the Taylor expansion $\log(1 - t) = -t - t^2/2 - t^3/3 - \dots$ gives $\log(1 - t) + t = -\sum_{j \geq 2} t^j/j$, so $|\log(1 - 1/p) + 1/p| \leq \sum_{j \geq 2} p^{-j}/j \leq \frac{1}{2} \sum_{j \geq 2} p^{-j} = \frac{1}{2p(p-1)} \leq 1/p^2$ for every prime $p \geq 2$, and $\sum_p p^{-2}$ converges. The claim requiring proof is that this closed-form constant is the same number M delivered by the proof of theorem 3.3.3.

Proof:

Let $M_2 = \lim_{x \rightarrow \infty} (\sum_{p \leq x} 1/p - \log \log x)$ denote the constant produced in theorem 3.3.3 (the limit exists because the error there is $O(1/\log x) \rightarrow 0$), and let $M = \gamma + \sum_p (\log(1 - 1/p) + 1/p)$ denote the closed form above. The two are shown equal.

Step 1: Logarithm of the truncated Euler product. For a fixed $x \geq 2$, take logarithms of the finite product over primes $p \leq x$ and expand each factor by the Taylor series for $\log(1 - 1/p)$:

$$\sum_{p \leq x} \log\left(1 - \frac{1}{p}\right) = -\sum_{p \leq x} \frac{1}{p} - \sum_{p \leq x} \sum_{j \geq 2} \frac{1}{j p^j} = -\sum_{p \leq x} \frac{1}{p} - S(x)$$

where $S(x) = \sum_{p \leq x} \sum_{j \geq 2} \frac{1}{j p^j} = -\sum_{p \leq x} (\log(1 - 1/p) + 1/p)$. As established above, the double series $\sum_p \sum_{j \geq 2} (j p^j)^{-1}$ converges absolutely, so $S(x) \rightarrow S := -\sum_p (\log(1 - 1/p) + 1/p)$ as $x \rightarrow \infty$, with tail

$$0 \leq S - S(x) = \sum_{p > x} \sum_{j \geq 2} \frac{1}{j p^j} \leq \sum_{p > x} \frac{1}{p^2} \leq \sum_{n > x} \frac{1}{n^2} = O\left(\frac{1}{x}\right)$$

Rearranging the displayed identity,

$$\sum_{p \leq x} \frac{1}{p} = -\sum_{p \leq x} \log\left(1 - \frac{1}{p}\right) - S(x)$$

Step 2: The regularized prime sum and the entry of γ . Write $P(x) = \prod_{p \leq x} (1 - 1/p)^{-1}$, so that $\log P(x) = -\sum_{p \leq x} \log(1 - 1/p)$ and the identity of Step 1 becomes

$$\sum_{p \leq x} \frac{1}{p} = \log P(x) - S(x) \tag{3.1}$$

The number M_2 to be evaluated is $\lim_{x \rightarrow \infty} (\sum_{p \leq x} 1/p - \log \log x)$, and (3.1) together with $S(x) \rightarrow S$ rewrites it as $M_2 = \lim_{x \rightarrow \infty} (\log P(x) - \log \log x) - S$. The whole task is therefore to show that the additive constant in $\log P(x) = \log \log x + (\text{const}) + o(1)$ is exactly the Euler–Mascheroni γ . This constant is not produced by any termwise manipulation of the product; it is the precise discrepancy between two regularizations of the divergent sum $\sum_p 1/p$, the sharp cutoff $\sum_{p \leq x} 1/p$ and the analytic cutoff $F(\rho) := \sum_p p^{-1-\rho}$ as $\rho \rightarrow 0^+$. The strategy is to evaluate the analytic cutoff in closed form, where γ appears in-chapter, and then compare the two cutoffs.

The analytic regularization is evaluated first. Taking logarithms of the Euler product $\zeta(1 + \rho) = \prod_p (1 - p^{-1-\rho})^{-1}$ (valid for $\rho > 0$, where both sides converge absolutely) and expanding each factor exactly as in Step 1,

$$\log \zeta(1 + \rho) = \sum_p \sum_{j \geq 1} \frac{1}{j p^{j(1+\rho)}} = F(\rho) + S(\rho), \quad S(\rho) := \sum_p \sum_{j \geq 2} \frac{1}{j p^{j(1+\rho)}}$$

Since $0 \leq S(\rho) \leq S(0) = S$ for $\rho \geq 0$ and each term increases to its $\rho = 0$ value as $\rho \rightarrow 0^+$, monotone convergence gives $S(\rho) \rightarrow S$. The constant γ now enters through $\zeta(1 + \rho)$ as $\rho \rightarrow 0^+$,

evaluated entirely within the convergent range $\text{Re } s > 1$, with no appeal to continuation past the line. By the closed form of proposition 1.4.6, valid for $\text{Re } s > 1$, $\zeta(s) = \frac{1}{s-1} + 1 - s \int_1^\infty \{t\} t^{-s-1} dt$, and at $s = 1 + \rho$ this reads

$$\zeta(1 + \rho) = \frac{1}{\rho} + 1 - (1 + \rho) \int_1^\infty \frac{\{t\}}{t^{2+\rho}} dt = \frac{1}{\rho} + \gamma + o(1) \quad (\rho \rightarrow 0^+)$$

the limit by dominated convergence ($\{t\} t^{-2-\rho} \leq t^{-2}$ for $\rho \geq 0$) and the value $\int_1^\infty \{t\} t^{-2} dt = 1 - \gamma$, which is exactly the evaluation of the Euler–Mascheroni constant in example 1.4.4. Hence $\log \zeta(1 + \rho) = \log(1/\rho) + \log(1 + \gamma\rho + o(\rho)) = \log(1/\rho) + o(1)$, and subtracting $S(\rho) \rightarrow S$,

$$F(\rho) = \sum_p \frac{1}{p^{1+\rho}} = \log \frac{1}{\rho} - S + o(1) \quad (\rho \rightarrow 0^+) \quad (3.2)$$

This is the analytic regularization in closed form, with γ already discharged into the relation $M_2 = \gamma - S$ once the two regularizations are matched.

Step 3: Matching the two regularizations. It remains to compare the sharp cutoff with the analytic one at the calibration $\rho = 1/\log x$. The comparison is

$$\sum_{p \leq x} \frac{1}{p} - F\left(\frac{1}{\log x}\right) = \gamma + o(1) \quad (x \rightarrow \infty) \quad (3.3)$$

This single comparison is the one analytic input not delivered by the tools of this chapter, and it is isolated here as such. Its content is that replacing the sharp truncation $1_{p \leq x}$ by the smooth damping $p^{-1/\log x}$ shifts the (divergent) prime sum by exactly γ in the limit. The shift is computed by Mertens' own method: the partial sums of theorem 3.3.2 are fed through partial summation against the two cutoff weights, the difference is expressed as an integral over a smooth integrand by the Euler–Maclaurin formula (theorem 1.5.6), and the limit $\rho \rightarrow 0^+$ is taken. The constant that survives is the value of the integral $\int_0^\infty \left(\frac{1}{e^t-1} - \frac{1}{te^t}\right) dt = \gamma$, equivalently $\Gamma'(1) = -\gamma$, an identity for the Gamma function rather than for ζ . The full computation, with explicit error bounds and the evaluation of this integral, is carried out in [10, §I.1.6] and in [2, Theorem 429]; the historically faithful version of Mertens' 1874 argument, with the same integral producing γ , is reproduced in detail in [12]. The mechanical tools that (3.3) rests on, the Euler product, partial summation, and Euler–Maclaurin, are all in hand; only the Gamma-function constant lies outside the chapter, and it is a fact about the Gamma function, not a disguised appeal to the analytic theory of ζ developed in chapter 2.

Granting (3.3), substitute (3.2) at $\rho = 1/\log x$, where $\log(1/\rho) = \log \log x$, to obtain

$$\sum_{p \leq x} \frac{1}{p} = F\left(\frac{1}{\log x}\right) + \gamma + o(1) = (\log \log x - S + o(1)) + \gamma + o(1) = \log \log x + (\gamma - S) + o(1)$$

Hence $\lim_{x \rightarrow \infty} (\sum_{p \leq x} 1/p - \log \log x) = \gamma - S$. By theorem 3.3.3 the same limit equals M_2 , and limits are unique, so

$$M_2 = \gamma - S = \gamma + \sum_p \left(\log \left(1 - \frac{1}{p} \right) + \frac{1}{p} \right) = M$$

the middle equality by $S = -\sum_p (\log(1 - 1/p) + 1/p)$ from Step 1. This is the closed form claimed for the Mertens constant. The equality $M = B$ with the constant of proposition 1.3.9 is immediate, since B was defined there by the identical expression $\gamma + \sum_p (\log(1 - 1/p) + 1/p)$, completing the proof.

The closed form makes the Mertens constant computable and ties together three appearances of the same number: the second-order term in $\sum_{p \leq x} 1/p$, the second-order term in the average of ω from proposition 1.3.9, and (through theorem 3.3.5 below) the constant governing the product $\prod(1 - 1/p)$. The expression $M = \gamma + \sum_p (\log(1 - 1/p) + 1/p)$ exhibits M as the Euler–Mascheroni constant corrected by a rapidly convergent prime sum: each correction term $\log(1 - 1/p) + 1/p$ is $-1/(2p^2) + O(p^{-3})$, so the sum is dominated by its first few terms and $M = 0.2614972128\dots$ is computed to high precision from a short truncation. The dependence on γ is not incidental. The constant γ entered through the value $\zeta(1 + \rho) = 1/\rho + \gamma + o(1)$ of the zeta function near its pole, which is itself the harmonic-sum constant of example 1.4.4 read off from the analytic continuation in proposition 1.4.6; the prime sum $\sum 1/p$ inherits γ as the precise gap between its sharp partial sums and the smoothly damped sum $\sum_p p^{-1-\rho}$, the two regularizations of the same divergent series whose difference (3.3) pins to γ . The convergent correction $-S$ is a separate, smaller adjustment: it measures only the gap between $\sum 1/p$ and $\log \prod(1 - 1/p)^{-1}$ caused by the higher prime powers, not the appearance of γ itself.

The product form is the most frequently cited of the three theorems, appearing throughout sieve theory and the theory of arithmetic functions as the density of integers free of small prime factors. Taking the logarithm of the product converts it into the sum $\sum_{p \leq x} \log(1 - 1/p)$, which the second theorem and the closed form of M evaluate; the constant $e^{-\gamma}$ emerges because the convergent prime correction in M cancels precisely against the convergent correction in $\sum \log(1 - 1/p)$, leaving only $-\gamma$.

Theorem 3.3.5: Mertens' Third Theorem

As $x \rightarrow \infty$,

$$\prod_{p \leq x} \left(1 - \frac{1}{p}\right) \sim \frac{e^{-\gamma}}{\log x}$$

where γ is the Euler–Mascheroni constant. Equivalently,

$$\prod_{p \leq x} \left(1 - \frac{1}{p}\right)^{-1} = e^{\gamma} \log x (1 + o(1))$$

Proof:

Step 1: Taking logarithms. Let $L(x) = \log \prod_{p \leq x} (1 - 1/p) = \sum_{p \leq x} \log(1 - 1/p)$. The claim $\prod_{p \leq x} (1 - 1/p) \sim e^{-\gamma} / \log x$ is equivalent, after taking logarithms, to

$$L(x) = -\gamma - \log \log x + o(1)$$

since $\log(e^{-\gamma} / \log x) = -\gamma - \log \log x$ and the asymptotic $\sim$ between positive quantities is equivalent to the difference of their logarithms tending to 0.

Step 2: Splitting off the convergent correction. For each prime p write $\log(1 - 1/p) = -1/p +$

$(\log(1 - 1/p) + 1/p)$, and sum over $p \leq x$:

$$L(x) = - \sum_{p \leq x} \frac{1}{p} + \sum_{p \leq x} \left(\log \left(1 - \frac{1}{p} \right) + \frac{1}{p} \right)$$

The second sum converges as $x \rightarrow \infty$: by the bound established after definition 3.3.4, $|\log(1 - 1/p) + 1/p| \leq 1/p^2$, so the series $\sum_p (\log(1 - 1/p) + 1/p)$ converges absolutely, with value $M - \gamma$ by the definition of the Mertens constant (definition 3.3.4). Writing the partial sum as the full sum minus a tail,

$$\sum_{p \leq x} \left(\log \left(1 - \frac{1}{p} \right) + \frac{1}{p} \right) = (M - \gamma) - \sum_{p > x} \left(\log \left(1 - \frac{1}{p} \right) + \frac{1}{p} \right) = (M - \gamma) + o(1)$$

since the tail of a convergent series tends to 0 (explicitly, $|\sum_{p > x} (\dots)| \leq \sum_{n > x} n^{-2} = O(1/x)$).

Step 3: Applying the second theorem. By Mertens' second theorem (theorem 3.3.3) with the closed form M of definition 3.3.4,

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + M + O\left(\frac{1}{\log x}\right)$$

Substituting both evaluations into the expression for $L(x)$ from Step 2,

$$L(x) = - \left(\log \log x + M + O\left(\frac{1}{\log x}\right) \right) + ((M - \gamma) + o(1))$$

The constant M cancels: $-M + (M - \gamma) = -\gamma$. The error terms $O(1/\log x)$ and $o(1)$ combine into $o(1)$. Hence

$$L(x) = -\log \log x - \gamma + o(1)$$

which is exactly the equivalent form from Step 1.

Step 4: Exponentiating. Exponentiating $L(x) = -\gamma - \log \log x + o(1)$,

$$\prod_{p \leq x} \left(1 - \frac{1}{p} \right) = e^{L(x)} = e^{-\gamma - \log \log x + o(1)} = \frac{e^{-\gamma}}{\log x} \cdot e^{o(1)} = \frac{e^{-\gamma}}{\log x} (1 + o(1))$$

using $e^{o(1)} = 1 + o(1)$, which is the first stated form. Taking reciprocals gives $\prod_{p \leq x} (1 - 1/p)^{-1} = e^{\gamma} \log x (1 + o(1))$, the second form, completing the proof.

The cancellation in Step 3 is the mechanism that produces $e^{-\gamma}$, and it shows precisely why the Mertens constant had to be defined with the γ term: the constant M in $\sum 1/p$ and the convergent sum $\sum (\log(1 - 1/p) + 1/p) = M - \gamma$ in $\log \prod (1 - 1/p)$ share the prime-sum part, which cancels, while the γ that distinguishes them survives as the exponent. The third theorem quantifies a competition between two effects in the product $\prod_{p \leq x} (1 - 1/p)$: each factor $1 - 1/p$ is slightly less than 1, so the product decreases, but the factors approach 1 as p grows, so the decrease slows. The outcome $\sim e^{-\gamma}/\log x$ says the product tends to 0, but only as slowly as $1/\log x$, with the precise rate fixed by $e^{-\gamma} \approx 0.5615$. This product is the proportion of integers surviving the sieve that removes every

multiple of every prime up to x ; its value $e^{-\gamma}/\log x$, rather than the naive $1/\log x$ one might guess from the heuristic that “a $1/\log x$ fraction of integers near x are prime,” is the source of the constant $2e^{-\gamma}$ in the Mertens correction to sieve estimates and the reason the twin-prime constant and similar products carry factors of $e^{-\gamma}$.

A numerical check confirms that the two nontrivial constants, M in the second theorem and $e^{-\gamma}$ in the third, are the ones the data demand, and exhibits the slow convergence that the error terms predict.

Example 3.3.6: Numerical Check of Mertens' Theorems

The constants entering the second and third theorems are

$$M = 0.26150\dots, \quad \gamma = 0.57722\dots, \quad e^{-\gamma} = 0.56146\dots$$

For the second theorem, the table compares the partial sum $\sum_{p \leq x} 1/p$ against its predicted value $\log \log x + M$, recording the discrepancy:

x	$\sum_{p \leq x} 1/p$	$\log \log x + M$	difference
10^2	1.80282	1.78868	+0.01414
10^4	2.48306	2.48182	+0.00124
10^6	2.88733	2.88729	+0.00004

The difference $\sum_{p \leq x} 1/p - (\log \log x + M)$ shrinks steadily, from about 1.4×10^{-2} at $x = 10^2$ to 4×10^{-5} at $x = 10^6$, consistent with the $O(1/\log x)$ error term of theorem 3.3.3: the ratio of successive differences tracks the ratio of successive values of $1/\log x$, the predicted decay. For the third theorem, the table compares $\prod_{p \leq x} (1 - 1/p) \cdot \log x$ against the predicted limit $e^{-\gamma} = 0.56146$:

x	$\prod_{p \leq x} (1 - 1/p)$	$(\prod_{p \leq x} (1 - 1/p)) \log x$	$e^{-\gamma}$
10^2	0.12032	0.55408	0.56146
10^4	0.06088	0.56077	0.56146
10^6	0.04064	0.56144	0.56146

The product $(\prod_{p \leq x} (1 - 1/p)) \log x$ approaches $e^{-\gamma}$ from below, sitting within two parts in 10^5 of $e^{-\gamma}$ at $x = 10^6$. Both tables confirm the predicted constants, M in the second theorem and $e^{-\gamma}$ in the third, and both exhibit the slow convergence characteristic of error terms that decay only as a power of $1/\log x$ rather than a power of $1/x$.

3.4 Bertrand's Postulate

Chebyshev's estimates of section 3.2 pin the global count of primes up to x to the correct order $x/\log x$, but a global count says nothing about where the primes sit. A statement like $\pi(2n) - \pi(n) \approx n/\log n$ would assert, on average, many primes between n and $2n$, yet an average leaves open the possibility of long prime-free stretches compensated by dense clusters elsewhere. Bertrand's postulate is the first genuine *localization* result: for every integer $n \geq 1$ there is at least one prime in the interval $(n, 2n]$, with no exceptions and no averaging. Conjectured by Bertrand in 1845 on numerical evidence up to several million and proved by Chebyshev in 1852, it guarantees a prime in an interval whose length is comparable to its left endpoint, a far shorter window than the global count alone delivers.

The proof is a refinement of the binomial method already used for Chebyshev's bounds. The central binomial coefficient $\binom{2n}{n}$ is large, of size roughly 4^n , and section 3.2 controlled its prime factorization through Legendre's formula. Sharpening that control shows that if no prime fell in $(n, 2n]$, then $\binom{2n}{n}$ would be forced to assemble its entire size out of small primes alone, each contributing only a bounded factor; the product of those small contributions is too small to account for 4^n once n is large. The contradiction proves the postulate for all large n , and the finitely many remaining cases are settled by exhibiting an explicit chain of primes. The argument here is the one found by Erdős in 1932, which replaced Chebyshev's analytic estimates of the Gamma function by the elementary bounds of lemma 3.4.1.

Lemma 3.4.1: Bounds on the Central Binomial Coefficient

For every integer $n \geq 1$,

$$\frac{4^n}{2n+1} \leq \binom{2n}{n} \leq 4^n$$

Proof:

Both bounds compare $\binom{2n}{n}$ with the full binomial sum $\sum_{j=0}^{2n} \binom{2n}{j} = (1+1)^{2n} = 4^n$.

Step 1: The upper bound. The coefficient $\binom{2n}{n}$ is one of the $2n+1$ nonnegative terms $\binom{2n}{0}, \binom{2n}{1}, \dots, \binom{2n}{2n}$ whose sum is 4^n . A single term of a sum of nonnegative reals cannot exceed the whole sum, so

$$\binom{2n}{n} \leq \sum_{j=0}^{2n} \binom{2n}{j} = 4^n$$

which is the upper bound.

Step 2: The lower bound. The same $2n+1$ terms have $\binom{2n}{n}$ as their largest. Indeed the ratio of consecutive terms is

$$\frac{\binom{2n}{j+1}}{\binom{2n}{j}} = \frac{2n-j}{j+1}$$

which exceeds 1 exactly when $2n-j > j+1$, that is, $j < (2n-1)/2 = n - \frac{1}{2}$, and is less than 1 when $j > n - \frac{1}{2}$. Thus the terms strictly increase up to $j = n$ and strictly decrease after it, so $\binom{2n}{n}$ is the unique maximum. Being the largest of $2n+1$ terms, it is at least their average:

$$\binom{2n}{n} \geq \frac{1}{2n+1} \sum_{j=0}^{2n} \binom{2n}{j} = \frac{4^n}{2n+1}$$

which is the lower bound, completing the proof.

The two bounds bracket $\binom{2n}{n}$ between $4^n/(2n+1)$ and 4^n , a window narrow on the logarithmic scale: $\log \binom{2n}{n} = n \log 4 + O(\log n)$, so to leading order $\binom{2n}{n}$ behaves like $4^n = 2^{2n}$. The lower bound is the operative one for Bertrand's postulate, since the argument proceeds by showing that a hypothetical absence of primes in $(n, 2n]$ makes $\binom{2n}{n}$ too small, and "too small" is measured against $4^n/(2n+1)$. The factor $2n+1$ in the denominator is wasteful (Stirling's formula theorem 1.5.7 gives the sharper

$\binom{2n}{n} \sim 4^n / \sqrt{\pi n}$, but the crude bound suffices and keeps the proof elementary. The next lemma supplies the matching upper control on $\binom{2n}{n}$, not through its size but through its prime factorization.

The mechanism of Bertrand's postulate is a restriction on *which* primes can divide $\binom{2n}{n}$, and to what power. Three facts are needed. The first removes a whole band of primes just below n from the factorization; the second caps the exponent of every prime past $\sqrt{2n}$ at one; the third, already proved in section 3.2, caps every prime power at $2n$. Together they bound $\binom{2n}{n}$ from above by a product whose factors are small primes raised to controlled powers.

Lemma 3.4.2: Primes in the Central Binomial Coefficient

Let $n \geq 3$ be an integer and let p be a prime.

- (i) If $\frac{2n}{3} < p \leq n$, then $p \nmid \binom{2n}{n}$.
- (ii) If $\sqrt{2n} < p \leq 2n$, then $v_p\left(\binom{2n}{n}\right) \leq 1$, so p divides $\binom{2n}{n}$ to at most the first power.
- (iii) For every prime p , the prime power $p^{v_p\left(\binom{2n}{n}\right)}$ satisfies $p^{v_p\left(\binom{2n}{n}\right)} \leq 2n$.

Proof:

Throughout, the valuation formula of lemma 3.2.1 is used in the form

$$v_p\left(\binom{2n}{n}\right) = \sum_{k \geq 1} \left(\left\lfloor \frac{2n}{p^k} \right\rfloor - 2 \left\lfloor \frac{n}{p^k} \right\rfloor \right)$$

in which every summand is 0 or 1 and the summand vanishes once $p^k > 2n$.

- (i) *The band $(2n/3, n]$.* Assume $\frac{2n}{3} < p \leq n$. The hypotheses on p and $n \geq 3$ force $p \geq 3$: from $p > 2n/3 \geq 2$ one gets $p \geq 3$. The aim is to show every summand in the valuation formula vanishes. Consider the terms by exponent k .

For $k = 1$: from $p \leq n$ one has $n/p \geq 1$, and from $p > 2n/3$ one has $n/p < 3/2$, so $1 \leq n/p < 3/2$ and $\lfloor n/p \rfloor = 1$. Likewise $2 \leq 2n/p < 3$, so $\lfloor 2n/p \rfloor = 2$. The $k = 1$ summand is $\lfloor 2n/p \rfloor - 2\lfloor n/p \rfloor = 2 - 2 \cdot 1 = 0$.

For $k \geq 2$: since $p > 2n/3$ and $p \geq 3$,

$$p^2 > \frac{2n}{3} \cdot p \geq \frac{2n}{3} \cdot 3 = 2n$$

so $p^k \geq p^2 > 2n$ for all $k \geq 2$, and every such summand vanishes by lemma 3.2.1.

All summands being zero, $v_p\left(\binom{2n}{n}\right) = 0$, that is, $p \nmid \binom{2n}{n}$.

- (ii) *Primes past $\sqrt{2n}$.* Assume $\sqrt{2n} < p \leq 2n$. Then $p^2 > 2n$, so for every $k \geq 2$ the term $\lfloor 2n/p^k \rfloor = 0$ and the corresponding summand vanishes. Only the $k = 1$ summand can survive, and it is 0 or 1 by lemma 3.2.1. Hence $v_p\left(\binom{2n}{n}\right) \leq 1$.
- (iii) *The cap on prime powers.* This is exactly the bound $p^{v_p\left(\binom{2n}{n}\right)} \leq 2n$ proved in lemma 3.2.1, valid for every prime p and every $n \geq 1$, recorded here for use alongside (i) and (ii).

The three statements together control the factorization as required.

The three statements act on disjoint parts of the prime range. Statement (iii) is universal but coarse: it bounds each prime's contribution by $2n$ regardless of the prime. Statement (ii) improves this for primes past $\sqrt{2n}$, where the exponent drops to at most one, so the contribution is just p rather than a power. Statement (i) is the new ingredient, absent from the Chebyshev bounds of section 3.2: it asserts that the primes in the band $(2n/3, n]$ are entirely missing from $\binom{2n}{n}$. The reason is arithmetic cancellation. Such a prime p appears once in the numerator $(2n)!$ as the factor p and once more as $2p$ (since $2p \leq 2n$ when $p \leq n$), contributing two to $v_p((2n)!)$; in the denominator $n!n!$ it appears once in each factorial as p (since $p \leq n$), contributing $2 \cdot 1 = 2$ to $2v_p(n!)$. The two cancel exactly. The constraint $p > 2n/3$ ensures that $3p > 2n$, so $3p$ is not among the factors of $(2n)!$ and no third copy disturbs the balance. This band is precisely the range just below n where one might have hoped to find the divisors of $\binom{2n}{n}$; their absence is what forces the coefficient to draw on smaller primes and ultimately produces the contradiction.

With the size of $\binom{2n}{n}$ bounded below and its factorization bounded above, the postulate follows by comparing the two. The comparison yields a contradiction for all sufficiently large n ; the threshold is explicit, and the finitely many smaller values are checked by an explicit list of primes.

Theorem 3.4.3: Bertrand's Postulate

For every integer $n \geq 1$ there exists a prime p with $n < p \leq 2n$.

Proof:

The argument has two parts: a proof for all $n \geq 512$ by contradiction, and a direct verification of $1 \leq n \leq 511$ by an explicit chain of primes.

Step 1: Assume no prime lies in $(n, 2n]$ for some $n \geq 512$. Under this assumption the primes dividing $\binom{2n}{n}$ are confined to small values. By lemma 3.2.1 every prime dividing $\binom{2n}{n}$ is at most $2n$. The assumption removes the primes in $(n, 2n]$, and part (i) of lemma 3.4.2 removes the primes in $(2n/3, n]$ (here $n \geq 512 \geq 3$). What remains is

$$\binom{2n}{n} = \prod_{p \leq 2n/3} p^{v_p(\binom{2n}{n})}$$

the product over primes $p \leq 2n/3$ only.

Step 2: Bound the product from above. Split the surviving primes at $\sqrt{2n}$.

For primes $p \leq \sqrt{2n}$, part (iii) of lemma 3.4.2 gives $p^{v_p(\binom{2n}{n})} \leq 2n$. The number of primes $p \leq \sqrt{2n}$ is at most $\sqrt{2n}$ (there are at most $\sqrt{2n}$ positive integers up to $\sqrt{2n}$), so their combined contribution is

$$\prod_{p \leq \sqrt{2n}} p^{v_p(\binom{2n}{n})} \leq (2n)^{\sqrt{2n}}$$

For primes $\sqrt{2n} < p \leq 2n/3$, part (ii) of lemma 3.4.2 gives $v_p(\binom{2n}{n}) \leq 1$, so each contributes at most p itself, and their product is at most the product of all primes up to $2n/3$. The primorial bound proposition 3.2.2 controls this:

$$\prod_{\sqrt{2n} < p \leq 2n/3} p^{v_p(\binom{2n}{n})} \leq \prod_{p \leq 2n/3} p \leq 4^{2n/3}$$

Multiplying the two ranges,

$$\binom{2n}{n} \leq (2n)^{\sqrt{2n}} \cdot 4^{2n/3}$$

is the upper bound forced by the assumption.

Step 3: Contradiction with the lower bound. The lower bound of lemma 3.4.1 gives $\binom{2n}{n} \geq 4^n/(2n+1)$. Combined with Step 2,

$$\frac{4^n}{2n+1} \leq (2n)^{\sqrt{2n}} \cdot 4^{2n/3}$$

Dividing by $4^{2n/3}$ and noting $4^n/4^{2n/3} = 4^{n/3}$,

$$4^{n/3} \leq (2n+1)(2n)^{\sqrt{2n}}$$

Take logarithms. With $2n+1 \leq (2n)^2$ for $n \geq 1$ (since $2n+1 \leq 4n^2$ once $n \geq 1$), the right side satisfies $\log((2n+1)(2n)^{\sqrt{2n}}) \leq (\sqrt{2n}+2)\log(2n)$, so

$$\frac{n}{3} \log 4 \leq (\sqrt{2n}+2)\log(2n)$$

The left side grows linearly in n while the right side grows like $\sqrt{n} \log n$, so the inequality fails for all large n ; it remains to locate the threshold. Substitute $m = \sqrt{2n}$, so $n = m^2/2$ and the inequality reads

$$\frac{m^2}{6} \log 4 \leq (m+2) \cdot 2 \log m$$

that is, $m \log 4 \leq 12(1+2/m) \log m$. For $m \geq 32$ (which corresponds to $n \geq 512$) the factor $1+2/m \leq 1+1/16 = 17/16$, so the right side is at most $12 \cdot \frac{17}{16} \log m = \frac{51}{4} \log m$, and the inequality would force

$$m \log 4 \leq \frac{51}{4} \log m, \quad \text{i.e.} \quad \frac{m}{\log m} \leq \frac{51}{4 \log 4} = \frac{51}{4 \cdot 1.3862 \dots} = 9.197 \dots$$

But $m/\log m$ is increasing for $m \geq 3$, and at $m = 32$ it already equals $32/\log 32 = 32/3.4657 \dots = 9.233 \dots > 9.197 \dots$. Hence the inequality fails for $m = 32$, and being increasing, $m/\log m$ exceeds the bound for all $m \geq 32$. This contradicts the displayed inequality for every $n \geq 512$. Therefore the assumption is untenable: a prime exists in $(n, 2n]$ for every $n \geq 512$.

Step 4: The range $1 \leq n \leq 511$. Consider the increasing chain of primes

$$2, 3, 5, 7, 13, 23, 43, 83, 163, 317, 631$$

Each prime in the list is less than twice its predecessor: $3 < 4$, $5 < 6$, $7 < 10$, $13 < 14$, $23 < 26$, $43 < 46$, $83 < 86$, $163 < 166$, $317 < 326$, $631 < 634$. Write the chain as $q_0 = 2 < q_1 = 3 < \dots < q_{10} = 631$, so that $q_i < 2q_{i-1}$ for each $i \geq 1$. Given an integer n with $1 \leq n \leq 511$, choose the smallest chain prime q_i with $q_i > n$; such an index exists because $q_{10} = 631 > 511 \geq n$. Minimality gives $q_{i-1} \leq n$ when $i \geq 1$, and when $n = 1$ the prime $q_0 = 2$ already satisfies $1 < 2 \leq 2 = 2n$. For $i \geq 1$,

$$q_i < 2q_{i-1} \leq 2n$$

so $n < q_i \leq 2n$, exhibiting a prime in $(n, 2n]$. This covers every n in $[1, 511]$.

Steps 1 through 3 establish the postulate for $n \geq 512$ and Step 4 for $1 \leq n \leq 511$, so a prime lies in $(n, 2n]$ for every integer $n \geq 1$, as we sought to show.

Bertrand's postulate sharpens the picture of the primes from a global count to a guarantee on every dyadic scale. Where Chebyshev's bounds (theorem 3.2.3) assert that the primes up to x number a constant multiple of $x/\log x$, leaving open how they bunch, the postulate forbids any gap from doubling: between n and $2n$ there is always a prime, so the ratio of consecutive primes is bounded. The two results draw on the same object, the central binomial coefficient, and differ only in how finely its factorization is dissected. Chebyshev compared the size $4^n/(2n+1) \leq \binom{2n}{n} \leq 4^n$ against the universal cap $p^{v_p} \leq 2n$ on prime powers; Bertrand adds the observation that the primes in the band $(2n/3, n]$ contribute nothing, which concentrates the entire size of $\binom{2n}{n}$ onto primes below $2n/3$ and onto the at-most-once primes between $\sqrt{2n}$ and $2n/3$, controlled by the primorial bound $\prod_{p \leq 2n/3} p \leq 4^{2n/3}$ of proposition 3.2.2. The exponent $2n/3$ against the size 4^n leaves a deficit of $4^{n/3}$ that only primes in $(n, 2n]$ could supply, and that is exactly the interval whose nonemptiness the theorem asserts. The threshold $n \geq 512$ is not optimal; the inequality of Step 3 already fails for $n \geq 468$, and tightening the prime-power bookkeeping lowers it further, but a clean power of two suffices and the explicit chain disposes of the remainder. In the language of Chebyshev's functions (definition 3.1.2), the postulate is the statement that $\vartheta(2n) - \vartheta(n) > 0$ for all $n \geq 1$, a positivity that the order-of-magnitude estimate $\vartheta(x) \asymp x$ suggests but does not by itself prove.

The postulate yields immediate bounds on the growth of the prime sequence, both on the gap between consecutive primes and on the size of the n -th prime.

Corollary 3.4.4: Consequences of Bertrand's Postulate

Let $p_1 = 2 < p_2 = 3 < p_3 = 5 < \dots$ enumerate the primes in increasing order. Then for every $n \geq 1$:

- (i) the consecutive primes satisfy $p_{n+1} < 2p_n$;
- (ii) the n -th prime satisfies $p_n \leq 2^n$, with equality only at $n = 1$.

Proof:

The two statements are proved in turn, the second by induction using the first.

- (i) Apply theorem 3.4.3 with n replaced by p_n : there is a prime q with $p_n < q \leq 2p_n$. Since $q > p_n$, the prime q is one of $p_{n+1}, p_{n+2}, \dots$, so $p_{n+1} \leq q$ as p_{n+1} is the least prime exceeding p_n . The strict inequality $q \leq 2p_n$ combined with $2p_n$ being even and hence not prime (as $p_n \geq 2$ gives $2p_n \geq 4 > 2$) yields $q < 2p_n$, whence $p_{n+1} \leq q < 2p_n$.
- (ii) Induct on n . The base case $n = 1$ is the equality $p_1 = 2 = 2^1$, which gives $p_1 \leq 2^1$. Assume $p_n \leq 2^n$ for some $n \geq 1$. By part (i), $p_{n+1} < 2p_n \leq 2 \cdot 2^n = 2^{n+1}$, so $p_{n+1} \leq 2^{n+1}$, which is the claim for $n + 1$. By induction $p_n \leq 2^n$ for all $n \geq 1$. Equality occurs only at $n = 1$: for $n \geq 2$ the inductive step delivers the strict $p_{n+1} < 2^{n+1}$, so $p_n < 2^n$ for every $n \geq 2$.

Both bounds follow, completing the proof.

The bound $p_{n+1} < 2p_n$ records that the primes never thin out faster than geometrically: each prime is followed by another before the value doubles. The bound $p_n \leq 2^n$ (strict for $n \geq 2$) is a crude but unconditional handle on the size of the n -th prime, obtained with no analytic input. It is far from the truth, since the Prime Number Theorem (chapter 5, foreshadowed) gives $p_n \sim n \log n$, a function of much slower growth than 2^n ; the gap measures how much is lost by iterating a doubling bound

rather than tracking the actual density of primes. The value of $p_n \leq 2^n$ is its independence from any analytic input: it follows from Bertrand's postulate alone, which follows in turn from the size and factorization of a single binomial coefficient.

The Riemann Zeta Function

The two preceding chapters approached the zeta function from opposite directions. chapter 2 built $\zeta(s) = \sum_{n \geq 1} n^{-s}$ as a Dirichlet series, established its Euler product $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ as the analytic form of unique factorization (corollary 2.3.2), and produced the integral representation $\Gamma(s)\zeta(s) = \int_0^\infty t^{s-1}/(e^t - 1) dt$ (theorem 2.4.5), all on the half-plane $\operatorname{Re}(s) > 1$ where the defining series converges absolutely. chapter 3 took the elementary road, fixing the order of magnitude of $\pi(x)$ and evaluating the sums $\sum_{p \leq x} (\log p)/p$ and $\sum_{p \leq x} 1/p$ to leading order by counting prime powers, without any appeal to complex analysis. What both chapters left untouched is the behavior of $\zeta(s)$ as a function of a *complex* variable outside the half-plane of convergence. The series $\sum n^{-s}$ diverges at $s = 1$ and gives no value there, yet $\zeta(s)$ extends to a meromorphic function on a larger region, and it is the analytic structure of that extension, not the series, that controls the primes.

This chapter studies $\zeta(s)$ as a holomorphic function. The central facts are three. First, $\zeta(s)$ continues meromorphically to the half-plane $\operatorname{Re}(s) > 0$, holomorphic everywhere except for a single simple pole at $s = 1$ with residue 1; the continuation is already implicit in the partial-sum formula of chapter 1 and requires no new machinery. Second, the logarithmic derivative $-\zeta'(s)/\zeta(s) = \sum_n \Lambda(n)n^{-s}$ inherits a simple pole at $s = 1$ with residue 1, and this pole is the analytic source of the leading term $\psi(x) \sim x$ of the Prime Number Theorem. Third, $\zeta(s)$ does not vanish on the line $\operatorname{Re}(s) = 1$, a fact that lies just outside the reach of the elementary Euler-product argument of chapter 2 and demands a new idea. These three results are the analytic foundation on which the proof of the Prime Number Theorem rests in chapter 5; the deeper continuation to all of $\mathbb{C}$, together with the functional equation relating $\zeta(s)$ to $\zeta(1-s)$, is taken up in chapter 6.

4.1 Definition and the Euler Product

The starting point is the zeta function on its native half-plane $\operatorname{Re}(s) > 1$, where the series, the product, and the integral of chapter 2 all converge and agree. This section consolidates what is

already known there, recording the elementary analytic properties of $\zeta(s)$ as a holomorphic function of s , restating the Euler product and the non-vanishing it forces, and computing a few special values. The Euler product and the integral representation are not re-derived; they are imported from chapter 2 and cited. The one formula carried forward in full is the partial-sum identity of chapter 1, which already contains the continuation to $\operatorname{Re}(s) > 0$ and which the next section turns into the continuation theorem.

The defining series and the meaning of $\zeta(s)$ on $\operatorname{Re}(s) > 1$ were fixed in chapter 2; recall that $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ is the Dirichlet series of the constant function **1** (definition 2.1.1), with abscissa of absolute convergence $\sigma_a = 1$. The first task is to assemble the elementary consequences of this description into a single statement about ζ as a function.

Proposition 4.1.1: Analytic Properties of ζ on $\operatorname{Re}(s) > 1$

The Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ has the following properties on the half-plane $\operatorname{Re}(s) > 1$.

1. It is holomorphic, and its derivative is obtained by term-by-term differentiation:

$$\zeta'(s) = - \sum_{n=1}^{\infty} \frac{\log n}{n^s}$$

2. For every s with $\operatorname{Re}(s) = \sigma > 1$,

$$|\zeta(s) - 1| \leq \sum_{n=2}^{\infty} \frac{1}{n^{\sigma}}$$

and the right-hand side tends to 0 as $\sigma \rightarrow +\infty$; hence $\zeta(s) \rightarrow 1$ as $\operatorname{Re}(s) \rightarrow +\infty$.

3. On the real axis, $\zeta(\sigma)$ is real and strictly decreasing for $\sigma > 1$, with $\zeta(\sigma) > 1$ throughout, $\zeta(\sigma) \rightarrow 1$ as $\sigma \rightarrow +\infty$, and $\zeta(\sigma) \rightarrow +\infty$ as $\sigma \rightarrow 1^+$.

Proof:

The three parts are proved in turn.

1. The function ζ is the Dirichlet series $D(\mathbf{1}, s)$, whose abscissa of convergence is $\sigma_c = 1$ (example 2.1.9). By theorem 2.1.7, a Dirichlet series is holomorphic on the half-plane to the right of its abscissa of convergence, and there it may be differentiated term by term, with $D'(f, s) = \sum_n f(n)(-\log n)n^{-s}$. Taking $f = \mathbf{1}$ gives holomorphy on $\operatorname{Re}(s) > 1$ and

$$\zeta'(s) = \sum_{n=1}^{\infty} \frac{-\log n}{n^s} = - \sum_{n=1}^{\infty} \frac{\log n}{n^s}$$

the $n = 1$ term vanishing since $\log 1 = 0$.

2. Separating the $n = 1$ term, $\zeta(s) - 1 = \sum_{n \geq 2} n^{-s}$. Since $|n^{-s}| = n^{-\sigma}$ for $\operatorname{Re}(s) = \sigma$, the triangle inequality gives

$$|\zeta(s) - 1| = \left| \sum_{n=2}^{\infty} \frac{1}{n^s} \right| \leq \sum_{n=2}^{\infty} \frac{1}{n^{\sigma}}$$

which depends only on σ . The tail $\sum_{n \geq 2} n^{-\sigma}$ is bounded by the integral comparison $\sum_{n \geq 2} n^{-\sigma} \leq \int_1^\infty t^{-\sigma} dt = 1/(\sigma - 1)$, valid for $\sigma > 1$, and $1/(\sigma - 1) \rightarrow 0$ as $\sigma \rightarrow +\infty$. Hence $|\zeta(s) - 1| \rightarrow 0$, that is, $\zeta(s) \rightarrow 1$, uniformly in $\text{Im}(s)$ as $\text{Re}(s) \rightarrow +\infty$.

3. For real $s = \sigma > 1$, every term $n^{-\sigma}$ is a positive real number, so $\zeta(\sigma) = \sum_n n^{-\sigma}$ is real and exceeds its first term 1, giving $\zeta(\sigma) > 1$. For $1 < \sigma_1 < \sigma_2$ and each $n \geq 2$, the function $\sigma \mapsto n^{-\sigma} = e^{-\sigma \log n}$ is strictly decreasing because $\log n > 0$, so $n^{-\sigma_2} < n^{-\sigma_1}$; summing over $n \geq 2$ (the $n = 1$ term being constant) yields $\zeta(\sigma_2) < \zeta(\sigma_1)$, so ζ is strictly decreasing on $(1, \infty)$. The limit $\zeta(\sigma) \rightarrow 1$ as $\sigma \rightarrow +\infty$ is the real-axis case of part (2). For the limit at 1^+ , fix any N ; for $1 < \sigma$ the partial sum gives $\zeta(\sigma) \geq \sum_{n=1}^N n^{-\sigma}$, and letting $\sigma \rightarrow 1^+$ this finite sum tends to $\sum_{n=1}^N 1/n$, so $\liminf_{\sigma \rightarrow 1^+} \zeta(\sigma) \geq \sum_{n=1}^N 1/n$. Since N is arbitrary and the harmonic series $\sum_n 1/n$ diverges (example 1.4.4), the right-hand side is unbounded, forcing $\zeta(\sigma) \rightarrow +\infty$ as $\sigma \rightarrow 1^+$, as we sought to show.

These properties already record the qualitative shape of ζ along the real axis: a function descending strictly from $+\infty$ at the right edge of the pole to the horizontal asymptote $\zeta = 1$ at infinity. The blow-up at $\sigma = 1^+$ is the real-axis trace of the simple pole that the next section locates precisely, and it is inseparable from the divergence of the harmonic series, the most elementary statement about the density of the integers. Part (2) is the analytic counterpart of the fact that, far to the right, the $n = 1$ term dominates the series and every higher term is negligible; this is the same decay-at-different-rates phenomenon that made the coefficients of a Dirichlet series recoverable one at a time in corollary 2.1.8, and it will reappear as the normalization $\log \zeta(s) \rightarrow 0$ that pins the branch of the logarithm used in the non-vanishing argument of section 4.4. The holomorphy in part (1) is what entitles one to speak of the zeros and poles of ζ at all, and term-by-term differentiation makes $\zeta'(s)$ a Dirichlet series in its own right, the series of $-\log$, whose ratio with ζ produces the von Mangoldt series that carries the arithmetic.

The product form of ζ , established in chapter 2, is the single feature that ties the function to the primes; it is recalled here together with the non-vanishing it forces on the half-plane of absolute convergence.

Corollary 4.1.2: The Euler Product and Non-Vanishing on $\text{Re}(s) > 1$

For $\text{Re}(s) > 1$, the zeta function factors as an absolutely convergent product over the primes,

$$\zeta(s) = \prod_p \frac{1}{1 - p^{-s}}$$

and $\zeta(s) \neq 0$ on this half-plane.

Proof:

The factorization is corollary 2.3.2, the Euler product for ζ , obtained by applying the Euler product for multiplicative functions to the totally multiplicative function 1. The non-vanishing is proposition 2.3.4: an absolutely convergent infinite product whose factors are all nonzero has a nonzero value, and the local factors $(1 - p^{-s})^{-1}$ are reciprocals, hence never zero. Therefore $\zeta(s) = \prod_p (1 - p^{-s})^{-1} \neq 0$ for $\text{Re}(s) > 1$, as required.

The Euler product is the reason ζ encodes the primes at all: it expresses the global function as an

assembly of one local factor per prime, so that every analytic statement about ζ on $\operatorname{Re}(s) > 1$ is simultaneously a statement about the primes, and conversely. The non-vanishing recalled here is the easy half of the story. It holds throughout the region of absolute convergence precisely because the obstruction to a zero is local, a single vanishing factor, and no factor vanishes; this is the structural argument of proposition 2.3.4, sharper than the earlier proof through the Möbius series in example 2.2.5. The non-vanishing that matters for the distribution of primes, on the boundary line $\operatorname{Re}(s) = 1$, lies outside this region and is inaccessible to the product argument; it is the subject of section 4.4.

A few numerical values anchor the qualitative picture and connect ζ to constants from elsewhere in mathematics.

Example 4.1.3: Special Values of ζ

The two best-known special values are the even-argument evaluations

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}, \quad \zeta(4) = \sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90}$$

The first is the Basel problem, solved by Euler; both are proved in [16] (the standard derivations use the Fourier expansion of a periodic function or the product formula for $\sin$, neither of which is developed here). Numerically $\zeta(2) \approx 1.6449$ and $\zeta(4) \approx 1.0823$. The monotone decrease of proposition 4.1.1(3) toward the asymptote $\zeta = 1$ is visible in a short table of real values:

s	1.5	2	3	4	6	10
$\zeta(s)$	2.6124	1.6449	1.2021	1.0823	1.0173	1.0010

Each entry exceeds the next, and the values approach 1 as s grows, the contribution beyond the $n = 1$ term shrinking like 2^{-s} . The value $\zeta(3) = 1.2021\dots$, the constant appearing at $s = 3$, is *Apéry's constant*; Apéry proved it irrational in 1979. The even values $\zeta(2k)$ are all rational multiples of π^{2k} , a pattern that the functional equation of chapter 6 explains in one stroke, whereas the odd values $\zeta(3), \zeta(5), \dots$ have no such closed form and remain arithmetically obscure.

The contrast between the even and odd values is the first sign that ζ carries arithmetic information no elementary manipulation of the series reveals. The even values fall out of the functional equation because that equation relates $\zeta(2k)$ to $\zeta(1 - 2k)$, a value the equation forces to be rational; the odd values are paired by the same equation with $\zeta(-2k)$, which the functional equation makes vanish, so no comparable evaluation results. This asymmetry is what makes $\zeta(3)$ and its odd siblings difficult, and it foreshadows the role of the functional equation as the organizing principle of the next two chapters.

The single tool from chapter 1 that the continuation rests on is the partial-sum formula for $\sum_{n \leq x} n^{-s}$. For $\operatorname{Re}(s) > 0$ with $s \neq 1$, proposition 1.4.6 gives

$$\sum_{n \leq x} \frac{1}{n^s} = \frac{x^{1-s}}{1-s} + \zeta(s) + O(x^{-\operatorname{Re}(s)})$$

where the constant $\zeta(s)$ appearing here is, for $\operatorname{Re}(s) > 1$, the value of the defining series, and is given on the wider half-plane $\operatorname{Re}(s) > 0$ by the convergent integral

$$\zeta(s) = \frac{s}{s-1} - s \int_1^{\infty} \frac{\{t\}}{t^{s+1}} dt$$

with $\{t\}$ the fractional part of t . The integral converges absolutely for $\operatorname{Re}(s) > 0$ because $|\{t\}t^{-s-1}| \leq t^{-\operatorname{Re}(s)-1}$ and $\int_1^\infty t^{-\operatorname{Re}(s)-1} dt < \infty$ there. The right-hand side of this formula is defined and finite for every s with $\operatorname{Re}(s) > 0$ except $s = 1$, where the term $s/(s-1)$ has a simple pole; it therefore furnishes a value for $\zeta(s)$ on all of $\operatorname{Re}(s) > 0$, agreeing with the series where the series converges. This formula is the engine of the continuation, and the next section establishes that the function it defines is holomorphic on $\operatorname{Re}(s) > 0 \setminus \{1\}$ with a simple pole of residue 1 at $s = 1$, thereby extending ζ past the line $\operatorname{Re}(s) = 1$ to which the series confined it.

4.2 Meromorphic Continuation to $\operatorname{Re}(s) > 0$

The Dirichlet series $\sum_{n \geq 1} n^{-s}$ of definition 2.1.1 converges only on the half-plane $\operatorname{Re}(s) > 1$, and there it defines $\zeta(s)$ as a holomorphic function (proposition 4.1.1). Outside that half-plane the series is silent: it diverges, and the symbol $\zeta(s)$ has, as yet, no meaning. The whole analytic theory of the primes depends on giving it one. The Abel-summation analysis of proposition 1.4.6 already does so. Comparing the partial sums $\sum_{n \leq x} n^{-s}$ against the integral $\int_1^x t^{-s} dt$ produced the closed form

$$\zeta(s) = \frac{s}{s-1} - s \int_1^\infty \frac{\{t\}}{t^{s+1}} dt$$

in which the right-hand side makes sense on the larger half-plane $\operatorname{Re}(s) > 0$, because the integral converges there. The series-defined ζ on $\operatorname{Re}(s) > 1$ and this integral expression agree where both are defined, so the latter extends the former.

This section records that continuation as a theorem and reads off its local structure. The two elementary pieces on the right behave very differently near the line $\operatorname{Re}(s) = 1$: the rational term $s/(s-1)$ carries a single pole at $s = 1$, while the integral term is holomorphic across the whole half-plane and contributes nothing singular. Isolating the pole and computing the first two coefficients of the Laurent expansion of ζ at $s = 1$ is the work that remains, and it is the structural input that the proof of the Prime Number Theorem will draw on.

Theorem 4.2.1: Meromorphic Continuation of ζ to $\operatorname{Re}(s) > 0$

The function $\zeta(s)$, defined for $\operatorname{Re}(s) > 1$ by the series $\sum_{n=1}^\infty n^{-s}$, extends to a meromorphic function on the half-plane $\operatorname{Re}(s) > 0$ whose only singularity there is a simple pole at $s = 1$ with residue 1. Explicitly, for all s with $\operatorname{Re}(s) > 0$ and $s \neq 1$,

$$\zeta(s) = \frac{s}{s-1} - s \int_1^\infty \frac{\{t\}}{t^{s+1}} dt \quad (4.1)$$

where $\{t\} = t - [t]$ denotes the fractional part.

Proof:

Write $I(s) = \int_1^\infty \{t\} t^{-s-1} dt$ for the integral on the right of (4.1), so that the asserted identity reads $\zeta(s) = s/(s-1) - sI(s)$. The argument has three stages: the integral defines a holomorphic function on $\operatorname{Re}(s) > 0$; the right-hand side of (4.1) is therefore meromorphic on $\operatorname{Re}(s) > 0$ with a single simple pole at $s = 1$; and it agrees with the series on $\operatorname{Re}(s) > 1$, so it is the continuation. The residue is computed at the end.

Step 1: The integral $I(s)$ is holomorphic on $\operatorname{Re}(s) > 0$. The integrand $g(s, t) = \{t\} t^{-s-1}$ is, for each fixed $t \geq 1$, an entire function of s , and for each fixed s it is continuous in t . Since $0 \leq \{t\} < 1$, it is dominated by

$$|g(s, t)| = \{t\} t^{-\operatorname{Re}(s)-1} \leq t^{-\operatorname{Re}(s)-1}$$

Fix a compact set $K \subset \{\operatorname{Re}(s) > 0\}$ and let $\sigma_0 = \min_{s \in K} \operatorname{Re}(s) > 0$. Then $|g(s, t)| \leq t^{-\sigma_0-1}$ for all $s \in K$ and all $t \geq 1$, and $\int_1^\infty t^{-\sigma_0-1} dt = 1/\sigma_0 < \infty$. The dominating function $t^{-\sigma_0-1}$ is independent of s and integrable on $[1, \infty)$, so the integral defining $I(s)$ converges absolutely and uniformly for $s \in K$. A locally uniform limit of holomorphic functions is holomorphic, and a parameter integral with holomorphic integrand and a locally uniform integrable bound is holomorphic in the parameter, with differentiation permissible under the integral sign; this is the standard theorem on holomorphy of integrals depending on a parameter ([16]). Applying it on each such K , and noting that the half-plane $\operatorname{Re}(s) > 0$ is exhausted by such compact sets, shows that I is holomorphic on $\operatorname{Re}(s) > 0$.

Step 2: The right-hand side is meromorphic with a simple pole at $s = 1$. On $\operatorname{Re}(s) > 0$ the term $sI(s)$ is holomorphic, being the product of the entire function s with the holomorphic function $I(s)$ of Step 1. The term $s/(s-1)$ is a rational function whose only singularity is a simple pole at $s = 1$: writing $s/(s-1) = 1 + 1/(s-1)$ exhibits the principal part $1/(s-1)$ explicitly. The sum

$$R(s) := \frac{s}{s-1} - sI(s)$$

is therefore meromorphic on $\operatorname{Re}(s) > 0$, holomorphic away from $s = 1$, and has at $s = 1$ a simple pole, since the holomorphic term $sI(s)$ cannot cancel the simple pole of $s/(s-1)$.

Step 3: R agrees with ζ on $\operatorname{Re}(s) > 1$ and is the continuation. By proposition 1.4.6, for every s with $\operatorname{Re}(s) > 1$ the constant term in the asymptotic expansion of $\sum_{n \leq x} n^{-s}$ is exactly $R(s)$, and that constant equals $\lim_{x \rightarrow \infty} \sum_{n \leq x} n^{-s} = \sum_{n=1}^\infty n^{-s} = \zeta(s)$. Hence $R(s) = \zeta(s)$ for all s with $\operatorname{Re}(s) > 1$. The function ζ is holomorphic on the connected open set $\operatorname{Re}(s) > 1$, and R is meromorphic on the connected open set $\operatorname{Re}(s) > 0$ with its only pole at $s = 1$; the two agree on the nonempty open subset $\operatorname{Re}(s) > 1$, which has accumulation points in $\operatorname{Re}(s) > 0$. By the identity theorem ([16]), a meromorphic function on a connected open set is determined by its values on any subset with an accumulation point, so R is the unique meromorphic extension of ζ to $\operatorname{Re}(s) > 0$. This establishes (4.1) and shows that the only singularity of the continued ζ on $\operatorname{Re}(s) > 0$ is the simple pole of R at $s = 1$.

Step 4: The residue at $s = 1$ is 1. For a simple pole the residue is the limit of $(s-1)$ times the function. Using (4.1) and the holomorphy of $sI(s)$ at $s = 1$,

$$\lim_{s \rightarrow 1} (s-1)\zeta(s) = \lim_{s \rightarrow 1} \left[(s-1) \cdot \frac{s}{s-1} - (s-1)sI(s) \right] = \lim_{s \rightarrow 1} [s - (s-1)sI(s)] = 1$$

the second term vanishing in the limit because $sI(s)$ is finite at $s = 1$. Thus the pole is simple with residue 1, completing the proof.

The continuation is what lets the analytic machinery reach the line $\operatorname{Re}(s) = 1$ and the point $s = 1$, where the series itself gives nothing. Every later argument that extracts arithmetic from $\zeta(s)$ near

this line, the non-vanishing on $\operatorname{Re}(s) = 1$, the behavior of the logarithmic derivative, and ultimately the asymptotic $\psi(x) \sim x$, presupposes that ζ is defined and analytic there. The single simple pole at $s = 1$ is the one feature of this extended function that survives into the Prime Number Theorem: when Perron's formula expresses the Chebyshev sum $\psi(x)$ as a contour integral of $-\zeta'(s)/\zeta(s) \cdot x^s/s$, the pole of ζ at $s = 1$ becomes a pole of the integrand with residue x , and that residue is the main term $\psi(x) \sim x$ (chapter 5). That the residue of ζ at the pole is exactly 1, rather than some other constant, is what makes the main term x rather than a multiple of it. The pole is, in this precise sense, the analytic source of the leading order of the distribution of primes.

The continuation obtained here reaches only the half-plane $\operatorname{Re}(s) > 0$, governed by the convergence of the integral in (4.1), which fails once $\operatorname{Re}(s) \leq 0$. Extending ζ to all of $\mathbb{C}$ requires a different mechanism, the functional equation relating $\zeta(s)$ to $\zeta(1-s)$, developed in chapter 6. An alternative route to the same half-plane runs through the Dirichlet eta function: the integral representation of example 2.4.6 expresses $(1 - 2^{1-s})\Gamma(s)\zeta(s)$ by an integral that already converges on $\operatorname{Re}(s) > 0$, and dividing by the elementary factor $(1 - 2^{1-s})\Gamma(s)$ continues ζ across the line $\operatorname{Re}(s) = 1$ as well. The Abel-summation route taken here is the more elementary of the two, requiring no special functions, and it has the advantage of displaying the pole and its residue directly in the formula.

The pole at $s = 1$ is significant enough to isolate as a corollary in the form most often invoked, the value of the limit defining the residue.

Corollary 4.2.2: The Simple Pole at $s = 1$

The continued function $\zeta(s)$ has a simple pole at $s = 1$, and

$$\lim_{s \rightarrow 1} (s - 1) \zeta(s) = 1$$

Proof:

This is the statement of the pole and residue in theorem 4.2.1: the residue of a simple pole at $s = 1$ is by definition $\lim_{s \rightarrow 1} (s - 1)\zeta(s)$, and theorem 4.2.1 computes this limit to be 1, as required.

Knowing that the pole is simple with residue 1 fixes the leading term $1/(s - 1)$ of the Laurent expansion of ζ at $s = 1$. The constant term is the next datum, and it is not arbitrary: it is the Euler–Mascheroni constant γ , the same number that measures the gap between the harmonic sum $\sum_{n \leq x} 1/n$ and $\log x$ (example 1.4.4). The coincidence is not accidental. Both quantities arise from comparing a sum of n^{-s} against its integral, the harmonic sum at $s = 1$ and the zeta values as $s \rightarrow 1$, and the constant γ is the common discrepancy. The formula (4.1) makes the connection precise, because the integral $I(s)$ that appears in it specializes at $s = 1$ to the very integral whose value computes γ .

Proposition 4.2.3: The Laurent Expansion at $s = 1$

Near $s = 1$ the continued zeta function has the Laurent expansion

$$\zeta(s) = \frac{1}{s - 1} + \gamma + O(s - 1)$$

where γ is the Euler–Mascheroni constant.

Proof:

Start from the continuation formula (4.1) of theorem 4.2.1, valid for $\operatorname{Re}(s) > 0$ with $s \neq 1$, and split off the principal part of the rational term by writing $s/(s-1) = 1 + 1/(s-1)$. With $I(s) = \int_1^\infty \{t\} t^{-s-1} dt$ as before, this gives

$$\zeta(s) = \frac{1}{s-1} + 1 - sI(s) \quad (4.2)$$

The leading term $1/(s-1)$ is the principal part already identified in corollary 4.2.2; the remaining part $1 - sI(s)$ is holomorphic at $s = 1$ by Step 1 of the proof of theorem 4.2.1, so its value at $s = 1$ is the constant term of the Laurent expansion and its Taylor expansion supplies the higher-order terms. Evaluating at $s = 1$ requires the single number $I(1)$.

The function I is continuous at $s = 1$, again by the holomorphy of Step 1, so $I(1) = \int_1^\infty \{t\} t^{-2} dt$, the integral obtained by setting $s = 1$ in the integrand. This is exactly the integral whose value was computed in example 1.4.4 in the course of evaluating the Euler–Mascheroni constant, namely

$$\int_1^\infty \frac{\{t\}}{t^2} dt = 1 - \gamma$$

Substituting $s = 1$ and $I(1) = 1 - \gamma$ into the holomorphic part $1 - sI(s)$ of (4.2) gives the constant term

$$1 - 1 \cdot I(1) = 1 - (1 - \gamma) = \gamma$$

Since $1 - sI(s)$ is holomorphic at $s = 1$, it equals $\gamma + O(s-1)$ near $s = 1$, and combining with the principal part yields $\zeta(s) = 1/(s-1) + \gamma + O(s-1)$, as we sought to show.

The constant term γ ties the analytic behavior of ζ at its pole to the elementary asymptotics of the harmonic series, and this is the form in which the pole enters several refined estimates: the closed-form evaluation of the Mertens constant in theorem 3.3.3, for instance, isolates γ precisely as the value $\zeta(1+\rho) = 1/\rho + \gamma + o(1)$ read off from this expansion as $\rho \rightarrow 0^+$. The higher coefficients of the Laurent expansion are the *Stieltjes constants* γ_k , defined by $\zeta(s) = 1/(s-1) + \sum_{k \geq 0} \frac{(-1)^k}{k!} \gamma_k (s-1)^k$ with $\gamma_0 = \gamma$; they admit closed forms as regularized limits of $\sum_{n \leq x} (\log n)^k / n$ but do not enter the leading-order theory of the primes.

4.3 First Arithmetic Consequences

The continuation theorem of the previous section gives $\zeta(s)$ a single simple pole at $s = 1$ with residue 1 (theorem 4.2.1 and corollary 4.2.2), and the Laurent expansion $\zeta(s) = 1/(s-1) + \gamma + O(s-1)$ records its leading behavior there (proposition 4.2.3). This is a statement about ζ as an analytic function; on its own it says nothing about primes. The bridge to arithmetic is the Euler product, which on $\operatorname{Re}(s) > 1$ ties ζ to the primes one factor at a time, and the two operations that read the primes off the product, the logarithmic derivative $-\zeta'/\zeta$ and the logarithm $\log \zeta$. Applying these operations to a function with a known pole transfers that pole into analytic statements about prime-power sums.

This section carries out three such transfers. The logarithmic derivative $-\zeta'/\zeta$ acquires its own simple pole at $s = 1$, again of residue 1, and its Dirichlet series is the von Mangoldt series $\sum_n \Lambda(n) n^{-s}$

(theorem 2.2.6); this is the analytic object Perron's formula integrates to produce $\psi(x)$. Stripping the higher prime powers from $-\zeta'/\zeta$ isolates the prime series $\Phi(s) = \sum_p (\log p)p^{-s}$, which inherits the same pole. And the logarithm $\log \zeta$, expanded over prime powers, yields an analytic reproof that $\sum_p 1/p$ diverges, with the precise rate $\sum_p p^{-s} = \log(1/(s-1)) + O(1)$ as $s \rightarrow 1^+$. Each of these is a meromorphic fact about a half-plane; converting any of them into an asymptotic for $\psi(x)$ requires control of these functions on the boundary line $\operatorname{Re}(s) = 1$, supplied in the next section and exploited in chapter 5.

The first transfer is the most direct. The logarithmic derivative of a meromorphic function turns a pole or zero into a simple pole whose residue counts its order, and ζ has exactly one pole near $s = 1$ and (by the continuation) no zero there. The order of that pole is 1, so $-\zeta'/\zeta$ has a simple pole of residue 1.

Proposition 4.3.1: The Pole of the Logarithmic Derivative at $s = 1$

On the half-plane $\operatorname{Re}(s) > 1$,

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

where Λ is the von Mangoldt function (definition 1.1.8). Moreover $-\zeta'/\zeta$ continues to a meromorphic function on a neighborhood of $s = 1$, holomorphic there apart from a simple pole at $s = 1$, and that pole has residue 1.

Proof:

The Dirichlet series identity on $\operatorname{Re}(s) > 1$ is theorem 2.2.6: the von Mangoldt series $\sum_n \Lambda(n)n^{-s}$ converges absolutely on $\operatorname{Re}(s) > 1$ and equals $-\zeta'(s)/\zeta(s)$ there. It remains to analyze the behavior near $s = 1$.

By the continuation theorem (theorem 4.2.1) and the Laurent expansion of proposition 4.2.3, on a punctured neighborhood of $s = 1$ the zeta function has the form

$$\zeta(s) = \frac{1}{s-1} + \gamma + O(s-1)$$

Factor out the principal part: writing

$$\zeta(s) = \frac{1}{s-1} (1 + \gamma(s-1) + O((s-1)^2)) = \frac{1}{s-1} g(s)$$

defines $g(s) = 1 + \gamma(s-1) + O((s-1)^2)$, which is holomorphic at $s = 1$ with $g(1) = 1 \neq 0$. Since g is holomorphic and nonzero at $s = 1$, it is nonzero on some neighborhood U of $s = 1$, and on U the function $\zeta(s) = g(s)/(s-1)$ is meromorphic with its only pole the simple pole at $s = 1$ contributed by the factor $1/(s-1)$; in particular ζ has no zero on U .

On $U \setminus \{1\}$ the logarithmic derivative is computed from the factorization. Taking logarithmic derivatives turns the product $\zeta = (s-1)^{-1}g$ into a sum,

$$\frac{\zeta'(s)}{\zeta(s)} = \frac{d}{ds} \log((s-1)^{-1}) + \frac{g'(s)}{g(s)} = -\frac{1}{s-1} + \frac{g'(s)}{g(s)}$$

the first term being the logarithmic derivative of $(s-1)^{-1}$. Because g is holomorphic and nonvanishing on U , the quotient g'/g is holomorphic on U ; denote it $h(s)$, so that h is holomorphic at $s=1$. Negating,

$$-\frac{\zeta'(s)}{\zeta(s)} = \frac{1}{s-1} - h(s)$$

on $U \setminus \{1\}$. The right-hand side is meromorphic on U , holomorphic away from $s=1$, and has at $s=1$ a simple pole whose principal part is $1/(s-1)$, since the holomorphic term $-h(s)$ contributes nothing to the principal part. The coefficient of $(s-1)^{-1}$ is 1, so the residue is

$$\operatorname{Res}_{s=1} \left(-\frac{\zeta'(s)}{\zeta(s)} \right) = 1$$

This identifies the continuation of $-\zeta'/\zeta$ near $s=1$ as a simple pole of residue 1, as we sought to show.

The pole of $-\zeta'/\zeta$ at $s=1$ is the single analytic fact on which the leading order of the Prime Number Theorem rests. Perron's formula applied to the coefficients $a_n = \Lambda(n)$ expresses the Chebyshev function as the contour integral

$$\psi(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \left(-\frac{\zeta'(s)}{\zeta(s)} \right) \frac{x^s}{s} ds$$

for any $c > 1$ (theorem 2.5.2 and example 2.5.4). Shifting the contour to the left past the line $\operatorname{Re}(s) = 1$ picks up the residue of the integrand at the pole $s=1$; since $-\zeta'/\zeta$ has residue 1 there, the integrand $-\zeta'(s)/\zeta(s) \cdot x^s/s$ has residue $x^1/1 = x$, and this residue is exactly the main term $\psi(x) \sim x$. The agreement of the two residue-1 computations, the pole of ζ and the pole of $-\zeta'/\zeta$, is what fixes the main term as x with coefficient 1 rather than some other multiple. What the present proposition does not supply is the legitimacy of the contour shift: moving the line of integration from $\operatorname{Re}(s) = c > 1$ to a line $\operatorname{Re}(s) < 1$ requires that $-\zeta'/\zeta$ be holomorphic on the line $\operatorname{Re}(s) = 1$ itself, away from $s=1$, which in turn requires ζ to have no zeros there. That non-vanishing is the missing input, established in section 4.4, and the quantitative contour estimate that turns it into $\psi(x) \sim x$ is the work of chapter 5.

The von Mangoldt series $\sum_n \Lambda(n)n^{-s}$ runs over all prime powers p^k , weighting each by $\log p$. The contribution of the genuine primes, the terms with $k=1$, is a series over primes alone, and isolating it produces an object closer to the prime-counting function than $-\zeta'/\zeta$ is.

Definition 4.3.2: The Prime Series $\Phi(s)$

The *prime series* is

$$\Phi(s) = \sum_p \frac{\log p}{p^s}$$

the sum running over all primes p , defined for $\operatorname{Re}(s) > 1$.

The series converges absolutely on $\operatorname{Re}(s) > 1$: its terms are a subseries of the von Mangoldt series $\sum_n \Lambda(n)n^{-s}$, which converges absolutely there (theorem 2.2.6), so Φ is holomorphic on $\operatorname{Re}(s) > 1$. The function Φ is the Dirichlet generating function of the coefficients $(\log p)$ supported on primes, and its summatory function is the Chebyshev theta function $\vartheta(x) = \sum_{p \leq x} \log p$ of definition 3.1.2, just as $-\zeta'/\zeta$ generates $\psi(x) = \sum_{n \leq x} \Lambda(n)$. The two Chebyshev functions differ only by the prime-power

terms $\psi(x) - \vartheta(x) = \sum_{k \geq 2} \sum_{p^k \leq x} \log p$, and theorem 3.1.5 shows this difference to be of lower order, so $\vartheta(x) \sim x$ and $\psi(x) \sim x$ stand or fall together. The same redundancy appears analytically: Φ and $-\zeta'/\zeta$ differ by a function holomorphic well past the line $\operatorname{Re}(s) = 1$, so they carry the same singular data near $s = 1$.

Example 4.3.3: The Prime Series at $s = 2$

At $s = 2$ the series $\Phi(2) = \sum_p (\log p)p^{-2}$ converges, and its first terms are

$$\frac{\log 2}{4} + \frac{\log 3}{9} + \frac{\log 5}{25} + \frac{\log 7}{49} + \frac{\log 11}{121} + \dots$$

Numerically the first five terms contribute $0.1733 + 0.1221 + 0.0644 + 0.0397 + 0.0198 = 0.4193$, and the full sum is $\Phi(2) = 0.4931\dots$. The companion logarithmic-derivative value is $-\zeta'(2)/\zeta(2) = 0.5700\dots$, the full von Mangoldt sum $\sum_n \Lambda(n)n^{-2}$; the difference $-\zeta'(2)/\zeta(2) - \Phi(2) = 0.0769\dots$ collects the prime-power terms $\sum_p \sum_{k \geq 2} (\log p)p^{-2k}$, the largest of which is $(\log 2)(4^{-2} + 4^{-3} + \dots) = (\log 2)/12 = 0.0578$. The prime-power correction is small even at $s = 2$ and shrinks as $\operatorname{Re}(s)$ grows, the first sign of the general bound that makes Φ and $-\zeta'/\zeta$ interchangeable near the line $\operatorname{Re}(s) = 1$.

The relation between Φ and $-\zeta'/\zeta$ is exact and explicit: their difference is the sum of the prime-power terms with exponent $k \geq 2$, and that sum converges on a half-plane reaching well to the left of $\operatorname{Re}(s) = 1$. The difference is therefore holomorphic where it matters, and the pole of $-\zeta'/\zeta$ at $s = 1$ passes to Φ unchanged.

Proposition 4.3.4: Φ and the Logarithmic Derivative

For $\operatorname{Re}(s) > 1$,

$$-\frac{\zeta'(s)}{\zeta(s)} = \Phi(s) + R(s), \quad R(s) = \sum_p \sum_{k=2}^{\infty} \frac{\log p}{p^{ks}}$$

The remainder $R(s)$ converges absolutely on $\operatorname{Re}(s) > 1/2$ and is holomorphic there. Consequently $\Phi(s) - 1/(s-1)$ extends holomorphically to a neighborhood of $s = 1$: the prime series Φ has a simple pole at $s = 1$ with residue 1.

Proof:

Step 1: The decomposition. By theorem 2.2.6 the logarithmic derivative is the von Mangoldt series, which by definition 1.1.8 sums $\log p$ over all prime powers p^k with $k \geq 1$:

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s} = \sum_p \sum_{k=1}^{\infty} \frac{\log p}{p^{ks}}$$

on $\operatorname{Re}(s) > 1$, where the rearrangement of the absolutely convergent series into a double sum over (p, k) is the bijection between prime powers and pairs (p, k) . Splitting off the $k = 1$ terms, which form $\Phi(s)$ by definition 4.3.2, leaves

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_p \frac{\log p}{p^s} + \sum_p \sum_{k=2}^{\infty} \frac{\log p}{p^{ks}} = \Phi(s) + R(s)$$

which is the asserted identity.

Step 2: R is holomorphic on $\operatorname{Re}(s) > 1/2$. Fix a compact set $K \subset \{\operatorname{Re}(s) > 1/2\}$ and let $\sigma_0 = \min_{s \in K} \operatorname{Re}(s) > 1/2$. For $s \in K$ the modulus of the (p, k) -term is $(\log p)p^{-k\sigma_0}$ or smaller, and summing the geometric inner series over $k \geq 2$,

$$\sum_{k=2}^{\infty} \frac{\log p}{p^{k\sigma_0}} = \frac{\log p}{p^{2\sigma_0}} \cdot \frac{1}{1 - p^{-\sigma_0}} \leq \frac{\log p}{p^{2\sigma_0}} \cdot \frac{1}{1 - 2^{-\sigma_0}}$$

using $p \geq 2$ so that $1 - p^{-\sigma_0} \geq 1 - 2^{-\sigma_0} > 0$. The constant $(1 - 2^{-\sigma_0})^{-1}$ is finite and independent of p , so it suffices that $\sum_p (\log p)p^{-2\sigma_0}$ converges. Since $2\sigma_0 > 1$, choose $\varepsilon > 0$ with $2\sigma_0 - \varepsilon > 1$; then $\log p \leq C_\varepsilon p^\varepsilon$ for a constant C_ε and all p , whence

$$\sum_p \frac{\log p}{p^{2\sigma_0}} \leq C_\varepsilon \sum_p \frac{1}{p^{2\sigma_0 - \varepsilon}} \leq C_\varepsilon \sum_{n=1}^{\infty} \frac{1}{n^{2\sigma_0 - \varepsilon}} < \infty$$

the last series converging because its exponent exceeds 1. The bound is independent of $s \in K$, so the double series for R converges absolutely and uniformly on K . Each summand $(\log p)p^{-ks}$ is entire in s , and a locally uniform limit of holomorphic functions is holomorphic ([16]); exhausting $\{\operatorname{Re}(s) > 1/2\}$ by compact sets K shows R is holomorphic on $\operatorname{Re}(s) > 1/2$.

Step 3: The pole of Φ . On $\operatorname{Re}(s) > 1$ the identity of Step 1 rearranges to $\Phi(s) = -\zeta'(s)/\zeta(s) - R(s)$. By proposition 4.3.1 the function $-\zeta'/\zeta$ continues to a neighborhood U of $s = 1$ as a meromorphic function whose only singularity is a simple pole of residue 1, with $-\zeta'(s)/\zeta(s) = 1/(s-1) - h(s)$ for some h holomorphic on U . By Step 2, R is holomorphic on the neighborhood $U \cap \{\operatorname{Re}(s) > 1/2\}$ of $s = 1$. Therefore on this neighborhood

$$\Phi(s) - \frac{1}{s-1} = \left(-\frac{\zeta'(s)}{\zeta(s)} - \frac{1}{s-1} \right) - R(s) = -h(s) - R(s)$$

is a difference of two functions holomorphic at $s = 1$, hence holomorphic at $s = 1$. Thus $\Phi(s) = 1/(s-1) + (\text{holomorphic near } s = 1)$, so Φ has a simple pole at $s = 1$ whose principal part is $1/(s-1)$, that is, a simple pole of residue 1, completing the proof.

The prime series Φ is the analytic object that most directly governs the primes themselves, as opposed to the prime powers, and its pole at $s = 1$ is the analytic form of the statement that the primes have density $1/\log x$ in the sense made precise by the Prime Number Theorem. Perron's formula applied to Φ , with coefficients $a_n = \log p$ for $n = p$ prime and $a_n = 0$ otherwise, produces the contour integral for the Chebyshev theta function $\vartheta(x) = \sum_{p \leq x} \log p$, and the residue of $\Phi(s)x^s/s$ at the pole $s = 1$ is again x , giving the main term $\vartheta(x) \sim x$. The decomposition $-\zeta'/\zeta = \Phi + R$ with R holomorphic past the line $\operatorname{Re}(s) = 1$ is the analytic counterpart of the elementary estimate $\psi(x) - \vartheta(x) = O(\sqrt{x} \log x)$: in both pictures the prime powers of exponent $k \geq 2$ contribute a lower-order, harmless correction, and the substance lies entirely with the primes. The non-vanishing of ζ on $\operatorname{Re}(s) = 1$ enters here exactly as it does for $-\zeta'/\zeta$, since Φ differs from it only by the holomorphic R ; once that non-vanishing is in hand, Φ and $-\zeta'/\zeta$ may be used interchangeably in the contour argument of chapter 5.

The third transfer works with $\log \zeta$ rather than its derivative, and it reproves, by analytic means, the oldest quantitative fact about the primes. The expansion of $\log \zeta$ over prime powers (proposition 2.3.5) has the prime sum $\sum_p p^{-s}$ as its leading piece, and the pole of ζ forces $\log \zeta$ to blow up as $s \rightarrow 1^+$. Since the higher prime-power terms stay bounded, the blow-up must come from the prime sum, and

the Laurent expansion of ζ makes the rate explicit.

Proposition 4.3.5: Analytic Divergence of the Prime Sum

As $s \rightarrow 1^+$ along the real axis,

$$\sum_p \frac{1}{p^s} = \log \frac{1}{s-1} + O(1)$$

In particular $\sum_p 1/p$ diverges.

Proof:

Step 1: Separate the prime sum from the prime powers in $\log \zeta$. For real $s > 1$, proposition 2.3.5 gives the absolutely convergent expansion

$$\log \zeta(s) = \sum_p \sum_{k=1}^{\infty} \frac{1}{k p^{ks}} = \sum_p \frac{1}{p^s} + \sum_p \sum_{k=2}^{\infty} \frac{1}{k p^{ks}}$$

where the logarithm is the real logarithm of $\zeta(s) > 1$, and the second sum splits off the prime-power terms with $k \geq 2$. Write $E(s)$ for that second sum.

Step 2: The prime-power tail is bounded. For $s \geq 1$ the bound established in example 2.3.6 applies: dropping the factor $1/k \leq 1$ and summing the geometric inner series,

$$0 \leq E(s) = \sum_p \sum_{k=2}^{\infty} \frac{1}{k p^{ks}} \leq \sum_p \sum_{k=2}^{\infty} \frac{1}{p^{ks}} \leq \sum_p \sum_{k=2}^{\infty} \frac{1}{p^k} = \sum_p \frac{p^{-2}}{1 - p^{-1}}$$

the middle inequality using $p^{ks} \geq p^k$ for $s \geq 1$. The final sum is bounded by a convergent series independent of s : since $p^{-2}/(1 - p^{-1}) = 1/(p^2 - p) = 1/(p(p-1))$, comparison with all integers gives

$$\sum_p \frac{1}{p(p-1)} \leq \sum_{n=2}^{\infty} \frac{1}{n(n-1)} = \sum_{n=2}^{\infty} \left(\frac{1}{n-1} - \frac{1}{n} \right) = 1$$

the sum telescoping. Hence $0 \leq E(s) \leq 1$ for all $s \geq 1$, so $E(s) = O(1)$ as $s \rightarrow 1^+$.

Step 3: The size of $\log \zeta(s)$ near $s = 1$. By the Laurent expansion of proposition 4.2.3, near $s = 1$,

$$\zeta(s) = \frac{1}{s-1} + \gamma + O(s-1) = \frac{1}{s-1} (1 + \gamma(s-1) + O((s-1)^2))$$

For real s slightly greater than 1 both sides are positive, so taking the real logarithm,

$$\log \zeta(s) = \log \frac{1}{s-1} + \log(1 + \gamma(s-1) + O((s-1)^2))$$

As $s \rightarrow 1^+$ the argument of the second logarithm tends to 1, so that term tends to $\log 1 = 0$ and is in particular $O(1)$. Therefore

$$\log \zeta(s) = \log \frac{1}{s-1} + O(1) \quad (s \rightarrow 1^+)$$

Step 4: Combine. Subtracting the bounded tail of Step 2 from the expansion of Step 1 and inserting Step 3,

$$\sum_p \frac{1}{p^s} = \log \zeta(s) - E(s) = \left(\log \frac{1}{s-1} + O(1) \right) - O(1) = \log \frac{1}{s-1} + O(1)$$

as $s \rightarrow 1^+$. The right-hand side tends to $+\infty$ as $s \rightarrow 1^+$, since $\log(1/(s-1)) \rightarrow +\infty$. If the series $\sum_p 1/p$ converged, then for every $s > 1$ the comparison $\sum_p p^{-s} \leq \sum_p p^{-1}$ would bound $\sum_p p^{-s}$ by a finite constant uniformly in s , contradicting the divergence to $+\infty$ just established. Hence $\sum_p 1/p$ diverges, completing the proof.

The divergence of $\sum_p 1/p$ is Euler's theorem, and the rate $\sum_p p^{-s} \sim \log(1/(s-1))$ is its analytic sharpening: the prime sum grows like $\log(1/(s-1))$ as s approaches the pole, the same logarithmic rate by which $\log \zeta(s)$ itself blows up. This is the Dirichlet-series shadow of the Mertens estimate $\sum_{p \leq x} 1/p = \log \log x + M + O(1/\log x)$ of theorem 3.3.3. The two are linked by the heuristic $s-1 \leftrightarrow 1/\log x$ that runs throughout the subject: setting $s = 1 + 1/\log x$ turns $\log(1/(s-1))$ into $\log \log x$, and the analytic statement becomes the leading term of the Mertens statement. The agreement is not a coincidence but a reflection of the same arithmetic seen through two summation methods, the sharp cutoff $\sum_{p \leq x}$ and the smooth analytic weighting $\sum_p p^{-s}$; the constant M and the error $O(1/\log x)$ are the finer data that the smooth version compresses into its $O(1)$. The Mertens theorem, proved by elementary partial summation in chapter 3, is the stronger statement, with an explicit constant and error term that the present proof does not reach. What the analytic proof offers in exchange is method: it derives the divergence from the single fact that ζ has a pole at $s = 1$, and the same argument applied to a Dirichlet L -function $L(s, \chi)$, whose non-vanishing at $s = 1$ replaces the pole of ζ , yields the divergence of $\sum_{p \equiv a} 1/p$ over primes in an arithmetic progression, the analytic core of Dirichlet's theorem in chapter 7.

All three results of this section are statements about half-planes and about behavior as $s \rightarrow 1^+$ along the real axis. The pole of $-\zeta'/\zeta$, the pole of Φ , and the rate of divergence of $\sum_p p^{-s}$ each isolate the singular structure of an analytic function at the point $s = 1$. None of them yet delivers an asymptotic for $\psi(x)$, $\vartheta(x)$, or $\pi(x)$, because the passage from the singularity to the counting function runs through Perron's formula (theorem 2.5.2 and example 2.5.4), and that passage demands a contour shift across the line $\operatorname{Re}(s) = 1$. The shift is legitimate only where the integrand is holomorphic, and the integrand $-\zeta'(s)/\zeta(s) \cdot x^s/s$ is holomorphic on the line $\operatorname{Re}(s) = 1$ precisely when ζ has no zeros there. Establishing that $\zeta(1+it) \neq 0$ for all real t , the one piece of analytic input that lies beyond the elementary Euler-product arguments, is the subject of the next section, after which chapter 5 converts the pole at $s = 1$ into the asymptotic $\psi(x) \sim x$.

4.4 Non-Vanishing on the Line $\operatorname{Re}(s) = 1$

The preceding sections fixed the analytic structure of $\zeta(s)$ on the half-plane $\operatorname{Re}(s) > 0$: a single simple pole at $s = 1$ with residue 1 (theorem 4.2.1), and the logarithmic derivative $-\zeta'(s)/\zeta(s) = \sum_n \Lambda(n)n^{-s}$ carrying the arithmetic. What remains is the one analytic fact that the elementary Euler-product argument of chapter 2 could not reach. The non-vanishing of $\zeta(s)$ throughout $\operatorname{Re}(s) > 1$ followed from the factorization $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$, in which no local factor vanishes (corollary 4.1.2); but

that argument lives inside the region of absolute convergence and says nothing on the boundary line $\operatorname{Re}(s) = 1$, where the product no longer converges. The decisive analytic fact behind the Prime Number Theorem is that $\zeta(s)$ has no zeros on this line either: $\zeta(1 + it) \neq 0$ for every real t .

This section proves it by the positivity argument of Mertens and de la Vallée Poussin, organized around the elementary trigonometric inequality $3 + 4 \cos \theta + \cos 2\theta \geq 0$. The mechanism is the logarithmic expansion $\log \zeta(s) = \sum_{p,k} 1/(k p^{ks})$ of proposition 2.3.5, whose real part on a vertical line is a sum of cosines $\cos(kt \log p)$ weighted by positive coefficients. Feeding the three real parts at heights 0, t , and $2t$ into the inequality produces a single positive combination, and that positivity is incompatible with a zero of ζ on the line: a zero at $1 + it_0$ would force the combination to negative infinity as s approaches the line, a contradiction. The argument is short once the inequality is in hand.

Lemma 4.4.1: The Inequality $3 + 4 \cos \theta + \cos 2\theta \geq 0$

For every real number θ ,

$$3 + 4 \cos \theta + \cos 2\theta = 2(1 + \cos \theta)^2 \geq 0$$

Proof:

The double-angle identity $\cos 2\theta = 2 \cos^2 \theta - 1$ gives

$$3 + 4 \cos \theta + \cos 2\theta = 3 + 4 \cos \theta + (2 \cos^2 \theta - 1) = 2 \cos^2 \theta + 4 \cos \theta + 2$$

The right-hand side is $2(\cos^2 \theta + 2 \cos \theta + 1) = 2(\cos \theta + 1)^2$. A real square is nonnegative and the factor 2 is positive, so $2(1 + \cos \theta)^2 \geq 0$ for every real θ , as we sought to show.

The choice of coefficients 3, 4, 1 is not arbitrary. The argument below combines $\log \zeta$ at three heights, weighting the height 0 contribution by 3, the height t contribution by 4, and the height $2t$ contribution by 1, and the combination is useful only if the resulting trigonometric sum is nonnegative for every angle. Any nonnegative combination $a_0 + a_1 \cos \theta + a_2 \cos 2\theta$ with $a_0 = a_2$ doubled appropriately would serve in principle, but (3, 4, 1) is the lightest integer choice that both stays nonnegative and gives the height 0 term enough weight to overpower a single zero on the line; the equality case $\theta = \pi$, where the expression vanishes, shows the coefficients cannot be lowered while preserving positivity. The same inequality, applied to Dirichlet L -functions, drives the non-vanishing $L(1, \chi) \neq 0$ behind primes in arithmetic progressions (chapter 7).

The positivity now transfers from angles to the zeta function. The link is the expansion of $\log \zeta$ along a vertical line, whose real part is precisely a weighted sum of cosines.

Theorem 4.4.2: Non-Vanishing of ζ on $\operatorname{Re}(s) = 1$

The Riemann zeta function does not vanish on the line $\operatorname{Re}(s) = 1$ away from the pole: $\zeta(1 + it) \neq 0$ for every real $t \neq 0$.

Proof:

Throughout, write $s = \sigma + it$ with $\sigma > 1$ real, so that s lies in the half-plane of absolute convergence where the Euler product and its logarithm are valid. The argument has two stages: a positivity estimate valid for all $\sigma > 1$, and a contradiction derived from a hypothetical zero on the line.

Step 1: The fundamental inequality $|\zeta(\sigma)|^3 |\zeta(\sigma + it)|^4 |\zeta(\sigma + 2it)| \geq 1$ *for* $\sigma > 1$. By proposition 2.3.5, for $\operatorname{Re}(s) > 1$ the logarithm of ζ is given by the absolutely convergent double series

$$\log \zeta(s) = \sum_p \sum_{k=1}^{\infty} \frac{1}{k p^{ks}}$$

For $s = \sigma + it$ the summand is $k^{-1} p^{-k\sigma} p^{-ikt} = k^{-1} p^{-k\sigma} e^{-ikt \log p}$, whose real part is $k^{-1} p^{-k\sigma} \cos(kt \log p)$ since $\operatorname{Re}(e^{-i\alpha}) = \cos \alpha$. Absolute convergence permits taking the real part term by term, so

$$\operatorname{Re} \log \zeta(\sigma + it) = \sum_p \sum_{k=1}^{\infty} \frac{\cos(kt \log p)}{k p^{k\sigma}} \quad (4.3)$$

Now form the combination of (4.3) at the three heights 0, t , and $2t$ with weights 3, 4, 1. Each term carries the common positive coefficient $c_{p,k} = (k p^{k\sigma})^{-1}$, and writing $\theta_{p,k} = kt \log p$ for the angle at height t , the angle at height 0 is 0 and the angle at height $2t$ is $2\theta_{p,k}$. Hence

$$3\operatorname{Re} \log \zeta(\sigma) + 4\operatorname{Re} \log \zeta(\sigma + it) + \operatorname{Re} \log \zeta(\sigma + 2it) = \sum_p \sum_{k=1}^{\infty} c_{p,k} (3 + 4 \cos \theta_{p,k} + \cos 2\theta_{p,k})$$

the rearrangement of the three absolutely convergent series into one being permissible. By lemma 4.4.1 each bracket is nonnegative, and each coefficient $c_{p,k}$ is positive, so every term of the combined series is nonnegative and therefore

$$3\operatorname{Re} \log \zeta(\sigma) + 4\operatorname{Re} \log \zeta(\sigma + it) + \operatorname{Re} \log \zeta(\sigma + 2it) \geq 0 \quad (4.4)$$

For any nonzero complex number w and any branch of its logarithm, $\operatorname{Re} \log w = \log |w|$. Each ζ value here is nonzero, lying in the half-plane $\operatorname{Re}(s) > 1$ (corollary 4.1.2), so (4.4) reads $3 \log |\zeta(\sigma)| + 4 \log |\zeta(\sigma + it)| + \log |\zeta(\sigma + 2it)| \geq 0$. Exponentiating, and using that $\exp$ is increasing,

$$|\zeta(\sigma)|^3 |\zeta(\sigma + it)|^4 |\zeta(\sigma + 2it)| \geq 1 \quad (\sigma > 1) \quad (4.5)$$

This inequality holds for every real t and every $\sigma > 1$, and it is the only consequence of positivity that the argument needs.

Step 2: A zero on the line contradicts (4.5). Suppose, for contradiction, that $\zeta(1 + it_0) = 0$ for some real $t_0 \neq 0$. Apply (4.5) with $t = t_0$ and let $\sigma \rightarrow 1^+$ along the real axis; the three factors are estimated separately. The estimates compare each factor with a power of $\sigma - 1$, and the exponents are arranged so that the product tends to 0, contradicting the lower bound 1.

The first factor blows up at the controlled rate of the pole. By corollary 4.2.2 the continued ζ has a simple pole at $s = 1$ with $\lim_{\sigma \rightarrow 1^+} (\sigma - 1)\zeta(\sigma) = 1$, so $(\sigma - 1)\zeta(\sigma) \rightarrow 1$ and in particular $(\sigma - 1)|\zeta(\sigma)| \rightarrow 1$. Thus there is a constant $A > 0$ and a threshold $\sigma_1 \in (1, 2]$ such that

$$|\zeta(\sigma)| \leq \frac{A}{\sigma - 1} \quad (1 < \sigma \leq \sigma_1) \quad (4.6)$$

indeed any $A > 1$ works for σ near 1. Cubing, $|\zeta(\sigma)|^3 \leq A^3(\sigma - 1)^{-3}$.

The second factor vanishes at the controlled rate of the assumed zero. The point $1 + it_0$ is not the pole, since $t_0 \neq 0$, so by theorem 4.2.1 the function ζ is holomorphic in a neighborhood of $1 + it_0$. By hypothesis it vanishes there, hence vanishes to some finite order $m \geq 1$: there is a function h holomorphic near $1 + it_0$ with $h(1 + it_0) \neq 0$ and $\zeta(s) = (s - (1 + it_0))^m h(s)$ on that neighborhood (the local factorization of a holomorphic function at a zero, [16]). Restricting to $s = \sigma + it_0$ with $\sigma > 1$, the displacement from the zero is $s - (1 + it_0) = \sigma - 1$, a positive real number, so

$$\zeta(\sigma + it_0) = (\sigma - 1)^m h(\sigma + it_0)$$

As $\sigma \rightarrow 1^+$ the factor $h(\sigma + it_0) \rightarrow h(1 + it_0)$, a finite value, so $|h(\sigma + it_0)|$ is bounded by some constant $B > 0$ for σ in an interval $(1, \sigma_2]$ with $\sigma_2 \in (1, \sigma_1]$. Since $m \geq 1$ and $0 < \sigma - 1 < 1$ on this interval (shrinking σ_2 below 2 if needed), $(\sigma - 1)^m \leq (\sigma - 1)$, and therefore

$$|\zeta(\sigma + it_0)| = (\sigma - 1)^m |h(\sigma + it_0)| \leq B(\sigma - 1) \quad (1 < \sigma \leq \sigma_2) \quad (4.7)$$

Raising to the fourth power, $|\zeta(\sigma + it_0)|^4 \leq B^4(\sigma - 1)^4$.

The third factor stays bounded. The point $1 + 2it_0$ is also distinct from the pole, since $2t_0 \neq 0$, so by theorem 4.2.1 the function ζ is holomorphic at $1 + 2it_0$, hence continuous there. Consequently $\zeta(\sigma + 2it_0) \rightarrow \zeta(1 + 2it_0)$ as $\sigma \rightarrow 1^+$, a finite limit, and there is a constant $C > 0$ and a threshold $\sigma_3 \in (1, \sigma_2]$ with

$$|\zeta(\sigma + 2it_0)| \leq C \quad (1 < \sigma \leq \sigma_3) \quad (4.8)$$

No claim about the order of ζ at $1 + 2it_0$ is needed; mere boundedness suffices, and that is guaranteed by continuity whether or not ζ happens to vanish there.

Combining the three estimates (4.6), (4.7), and (4.8), valid simultaneously for $1 < \sigma \leq \sigma_3$, gives

$$|\zeta(\sigma)|^3 |\zeta(\sigma + it_0)|^4 |\zeta(\sigma + 2it_0)| \leq \frac{A^3}{(\sigma - 1)^3} \cdot B^4(\sigma - 1)^4 \cdot C = A^3 B^4 C (\sigma - 1)$$

The right-hand side tends to 0 as $\sigma \rightarrow 1^+$. Yet by (4.5) the left-hand side is at least 1 for every $\sigma > 1$. Letting $\sigma \rightarrow 1^+$ forces $1 \leq \lim_{\sigma \rightarrow 1^+} A^3 B^4 C (\sigma - 1) = 0$, an impossibility. The assumption $\zeta(1 + it_0) = 0$ is therefore untenable, and $\zeta(1 + it_0) \neq 0$ for every real $t_0 \neq 0$, completing the proof.

The arithmetic of the exponents is the heart of the matter. The pole contributes a singularity of order one, so the cube $|\zeta(\sigma)|^3$ grows like $(\sigma - 1)^{-3}$; a zero of order $m \geq 1$ makes the fourth power $|\zeta(\sigma + it_0)|^4$ vanish at least like $(\sigma - 1)^4$; and the bounded third factor contributes nothing. The net exponent $-3 + 4 = +1$ is positive, so the product vanishes, and this is exactly the contradiction. Had the weights been chosen differently, the balance could fail: weights $(1, 2, 1)$, for instance, would give net exponent $-1 + 2 = +1$ as well, but the corresponding trigonometric polynomial $1 + 2\cos\theta + \cos 2\theta = 2\cos^2\theta + 2\cos\theta$ is *not* nonnegative for all θ , so Step 1 would fail. The pair of requirements, nonnegativity of the cosine polynomial and a strictly positive net exponent, is what singles out $(3, 4, 1)$, and the equality $3 + 4\cos\theta + \cos 2\theta = 2(1 + \cos\theta)^2$ is the smallest integer solution meeting both. A single zero on the line cannot survive this squeeze.

This non-vanishing is the analytic linchpin of the Prime Number Theorem. The proof of $\psi(x) \sim x$ in chapter 5 begins from the Perron representation $\psi(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} (-\zeta'(s)/\zeta(s)) x^s/s ds$ of example 2.5.4, taken on a vertical line $\operatorname{Re}(s) = c$ with $c > 1$, and proceeds by pushing the contour

leftward onto the line $\operatorname{Re}(s) = 1$. Each zero of ζ on or near that line would be a pole of the integrand $-\zeta'(s)/\zeta(s)$, an obstruction to the contour shift contributing an extra term to $\psi(x)$ of size comparable to the main term. theorem 4.4.2 is precisely the statement that there is no such obstruction on the line itself: the only pole of the integrand encountered on $\operatorname{Re}(s) = 1$ is the one at $s = 1$ coming from the pole of ζ (proposition 4.3.1), whose residue x is the main term $\psi(x) \sim x$. The non-vanishing is what licenses the contour to rest on $\operatorname{Re}(s) = 1$ with no pole obstruction except $s = 1$, and it is in this sense that the asymptotic distribution of primes is encoded in a single line of the complex plane.

4.4.3 Historical Note: Hadamard and de la Vallée Poussin The Prime Number Theorem was conjectured by Gauss and Legendre near the close of the eighteenth century and resisted proof for nearly a hundred years. It was settled independently in 1896 by Jacques Hadamard and Charles-Jean de la Vallée Poussin, each building on Riemann's 1859 program of studying $\pi(x)$ through the complex-analytic properties of $\zeta(s)$. The technical core of both proofs is the non-vanishing of $\zeta(s)$ on the line $\operatorname{Re}(s) = 1$; the positivity argument through the inequality $3 + 4 \cos \theta + \cos 2\theta \geq 0$ presented here is the streamlined form later extracted by Mertens and refined by de la Vallée Poussin, and is the version that has become standard. Qualitative non-vanishing on the line suffices to prove $\psi(x) \sim x$, but not to control the error term. De la Vallée Poussin went further, converting the same positivity into a *quantitative* zero-free region of the form $\sigma > 1 - c/\log |t|$ to the left of the line, which yields an explicit error bound in the Prime Number Theorem; these zero-free regions, and the error terms they produce, are developed in chapter 5.

The non-vanishing is recorded once more in the form in which the proof of the Prime Number Theorem invokes it, as a statement about the logarithmic derivative.

Corollary 4.4.4: Regularity of the Logarithmic Derivative on $\operatorname{Re}(s) = 1$

The logarithmic derivative $-\zeta'(s)/\zeta(s)$ is holomorphic on the line $\operatorname{Re}(s) = 1$ except for a simple pole at $s = 1$. Equivalently, $\zeta(s)$ has no zeros on $\operatorname{Re}(s) = 1$.

Proof:

By theorem 4.2.1 the function ζ is holomorphic on a neighborhood of the line $\operatorname{Re}(s) = 1$ except at $s = 1$, where it has a simple pole. The logarithmic derivative $-\zeta'/\zeta$ of a meromorphic function is holomorphic at every point that is neither a zero nor a pole of ζ ; at a zero of order m it has a simple pole with residue $-m$, and at a pole of order n it has a simple pole with residue n (the standard local computation for $-\zeta'/\zeta$, [16]). On the line $\operatorname{Re}(s) = 1$ the only pole of ζ is the simple pole at $s = 1$, which contributes a simple pole of $-\zeta'/\zeta$ at $s = 1$ with residue 1 (proposition 4.3.1). It remains to rule out zeros of ζ on the line, the other possible source of poles of $-\zeta'/\zeta$. The point $s = 1$ is the pole, not a zero. For every other point $1 + it$ with $t \neq 0$, theorem 4.4.2 gives $\zeta(1 + it) \neq 0$. Hence ζ has no zeros on $\operatorname{Re}(s) = 1$, so $-\zeta'/\zeta$ has no poles there beyond the simple pole at $s = 1$, and is holomorphic on the rest of the line, as required.

The Prime Number Theorem

The four preceding chapters assembled, piece by piece, every ingredient the proof of the Prime Number Theorem requires. chapter 3 reduced the statement $\pi(x) \sim x/\log x$ to the assertion $\psi(x) \sim x$ about the Chebyshev function, and to the equivalent assertion $\vartheta(x) \sim x$ about its prime-only counterpart, through the equivalence of theorem 3.1.5; the same chapter pinned the order of magnitude of these functions, proving $\vartheta(x) \asymp x$ by Chebyshev's elementary count of prime powers (theorem 3.2.3). chapter 4 then supplied the analytic facts that the elementary theory could not reach: the zeta function continues meromorphically across the line $\operatorname{Re}(s) = 1$ with a single simple pole at $s = 1$ of residue 1 (theorem 4.2.1), its logarithmic derivative $-\zeta'(s)/\zeta(s)$ inherits that pole (proposition 4.3.1), and, the one fact lying just beyond the Euler-product arguments, $\zeta(s)$ has no zeros on the line $\operatorname{Re}(s) = 1$ (theorem 4.4.2). The goal of this chapter is to combine these into the asymptotic $\psi(x) \sim x$, equivalently $\vartheta(x) \sim x$ and $\pi(x) \sim x/\log x$, which is the Prime Number Theorem.

The route is the analytic proof of Hadamard and de la Vallée Poussin from 1896, in the streamlined form due to D. J. Newman in 1980 [6] and presented by Zagier [18], which trades the heavy contour estimates of the original arguments for a single Tauberian theorem. The strategy is to package the primes into the Dirichlet series $\Phi(s) = \sum_p (\log p)p^{-s}$, to convert the non-vanishing of ζ on $\operatorname{Re}(s) = 1$ into a holomorphic extension of Φ to the closed half-plane $\operatorname{Re}(s) \geq 1$ except for one pole, and to feed that extension into a Tauberian theorem that forces the integral $\int_1^\infty (\vartheta(x) - x)/x^2 dx$ to converge; the convergence of this integral, together with the monotonicity of ϑ , yields $\vartheta(x) \sim x$. This first section lays out the plan in full and states the central analytic claim that the next three sections establish. The Laplace-transform machinery and Newman's theorem occupy section 5.2 and section 5.3, the proof itself section 5.4; the final section, section 5.5, sharpens qualitative non-vanishing into a quantitative zero-free region and reads off an error term, the first quantitative refinement of the asymptotic.

5.1 The Analytic Strategy

The Prime Number Theorem is a single asymptotic statement, but its proof draws on two streams of input developed in separate chapters, and the first task is to see precisely how they join. The elementary side, from chapter 3, contributes two facts: that the three counting functions ψ , ϑ , and π rise or fall together, so that proving the asymptotic for any one of them proves it for all three; and that $\vartheta(x)$ is already known to be of the correct order x , bounded above and below by constant multiples of it. These reduce the entire theorem to the one limit $\vartheta(x)/x \rightarrow 1$. The analytic side, from chapter 4, contributes the behavior of $\zeta(s)$ and of the prime series $\Phi(s)$ near and on the line $\operatorname{Re}(s) = 1$. This section records the reduction, then states the plan by which the analytic inputs produce the missing limit.

The reduction itself is a restatement of results already proved, collected here so that the rest of the chapter may speak only of ϑ .

Proposition 5.1.1: Reduction of the Prime Number Theorem to $\vartheta(x) \sim x$

The following three asymptotic statements are equivalent:

$$\pi(x) \sim \frac{x}{\log x}, \quad \vartheta(x) \sim x, \quad \psi(x) \sim x$$

Moreover $\vartheta(x) = O(x)$, and the ratio $\vartheta(x)/x$ is bounded. Consequently the Prime Number Theorem in any of its three forms is equivalent to the single statement $\vartheta(x) \sim x$, and to prove it it suffices to prove that $\vartheta(x)/x \rightarrow 1$.

Proof:

The equivalence of the three asymptotics is theorem 3.1.5, where $\pi(x) \sim x/\log x$, $\vartheta(x) \sim x$, and $\psi(x) \sim x$ are shown to imply one another by Abel summation transferring between the prime-counting function π (definition 3.1.1) and the Chebyshev functions ϑ and ψ (definition 3.1.2). The bound $\vartheta(x) = O(x)$ is the upper half of Chebyshev's estimates (theorem 3.2.3), which produces constants $0 < c_1 < c_2$ with $c_1x \leq \vartheta(x) \leq \psi(x) \leq c_2x$ for all large x ; the right-hand inequality gives $\vartheta(x) \leq c_2x$, so $\vartheta(x) = O(x)$ and $\vartheta(x)/x \leq c_2$ is bounded above, while $\vartheta(x)/x \geq c_1$ bounds it below. Since the three statements are equivalent and $\vartheta(x) \sim x$ is by definition the assertion $\vartheta(x)/x \rightarrow 1$, establishing that single limit establishes all three, completing the proof.

The proposition is the hinge of the chapter. It says that the entire content of the Prime Number Theorem is concentrated in the behavior of one function, $\vartheta(x) = \sum_{p \leq x} \log p$, and that the work already done has narrowed the target to a single limit: $\vartheta(x)/x$ is known to lie between two positive constants, and what remains is to show that it converges, and that its limit is exactly 1. The Chebyshev bounds do half the job, confining the ratio to a bounded interval; the analytic theory must do the other half, collapsing that interval to the point 1. The boundedness recorded here is not incidental to the proof that follows. It is the hypothesis that makes the Tauberian step legitimate: a Tauberian theorem extracts an asymptotic for $\vartheta(x)$ from the analytic behavior of its generating function only under a one-sided growth condition, and $\vartheta(x) = O(x)$ is exactly that condition.

With the target fixed as $\vartheta(x) \sim x$, the plan of the proof can be stated in three steps. Each step is the subject of one of the next three sections; the purpose here is to show how they fit together and where each draws on the analytic structure of chapter 4.

5.1.2 Key Ideas: The Plan of the Proof The proof of $\vartheta(x) \sim x$ proceeds in three steps.

1. *The prime series extends to the line.* The prime series $\Phi(s) = \sum_p (\log p) p^{-s}$ of definition 4.3.2, holomorphic on $\operatorname{Re}(s) > 1$, has a simple pole at $s = 1$ of residue 1 and extends holomorphically to the closed half-plane $\operatorname{Re}(s) \geq 1$ with no singularity other than that pole. This is the analytic heart of the argument, and it is where the non-vanishing of ζ enters: by proposition 4.3.4 the difference $-\zeta'(s)/\zeta(s) - \Phi(s)$ is holomorphic on $\operatorname{Re}(s) > 1/2$, so Φ and $-\zeta'/\zeta$ have the same singularities up to that line, and by corollary 4.4.4 the function $-\zeta'/\zeta$ is holomorphic on all of $\operatorname{Re}(s) = 1$ except for the simple pole at $s = 1$, precisely because ζ has no zeros on the line (theorem 4.4.2).
2. *An integral converges.* The extension of Φ to the line forces the convergence of the integral

$$\int_1^\infty \frac{\vartheta(x) - x}{x^2} dx$$

The passage from step (1) to this convergence runs through a Laplace transform: Φ is recast as the Laplace transform of a function built from ϑ (the subject of section 5.2), and the holomorphic extension of that transform to the boundary line is exactly the hypothesis of the Tauberian theorem of Newman established in section 5.3, whose conclusion is the convergence of the integral. This step is carried out in section 5.4.

3. *Convergence forces the asymptotic.* The convergence of $\int_1^\infty (\vartheta(x) - x)/x^2 dx$, combined with the fact that ϑ is nondecreasing, forces $\vartheta(x) \sim x$. A monotone function whose normalized deviation from x has a convergent integral cannot stay bounded away from x : were $\vartheta(x) \geq \lambda x$ for some $\lambda > 1$ on a sequence of points, or $\vartheta(x) \leq \lambda x$ for some $\lambda < 1$, monotonicity would make the integrand bounded below by a positive constant over intervals of fixed logarithmic length, contradicting convergence. This final step, also in section 5.4, yields $\vartheta(x)/x \rightarrow 1$ and hence, by proposition 5.1.1, the Prime Number Theorem.

The three steps trace a path from the complex plane to the real line: a holomorphic extension across $\operatorname{Re}(s) = 1$ in step (1), a statement about a real integral in step (2), and a real-variable asymptotic in step (3). The transitions between them are the substance of the next three sections, and each transition is where a piece of analysis does work that no elementary argument could. What the roadmap makes visible, and what is worth dwelling on before the details, is why the non-vanishing of ζ on the line $\operatorname{Re}(s) = 1$ is the one input on which everything turns.

The decisive point is step (1), and within it the holomorphic extension of Φ to the closed half-plane. The series $\Phi(s) = \sum_p (\log p) p^{-s}$ converges only on $\operatorname{Re}(s) > 1$; by itself it gives no value on the boundary, and the convergence of the integral in step (2) is a statement about the behavior of ϑ that is sensitive to exactly that boundary. The bridge is the identity $-\zeta'(s)/\zeta(s) = \Phi(s) + R(s)$ of proposition 4.3.4, in which R is holomorphic well past the line, on $\operatorname{Re}(s) > 1/2$. This identity transfers the question about Φ on the line into a question about $-\zeta'/\zeta$ on the line, and the singularities of $-\zeta'/\zeta$ are dictated entirely by the zeros and the pole of ζ . The pole of ζ at $s = 1$ produces the one permitted singularity, the simple pole of Φ of residue 1, which becomes the main term in the asymptotic. Every zero of ζ on the line would produce an additional pole of $-\zeta'/\zeta$, hence of Φ , and would destroy the holomorphic extension that step (1) asserts. A single zero at a point $1 + it_0$ with $t_0 \neq 0$ would be a pole of Φ off the real axis, the Laplace transform of step (2) would fail to extend holomorphically to that point of the boundary, the hypothesis of Newman's theorem would not hold,

and the integral would have no reason to converge. The non-vanishing $\zeta(1+it) \neq 0$ for all real $t \neq 0$, proved in theorem 4.4.2, is precisely the statement that no such obstruction exists: the only singularity of Φ on the closed half-plane $\operatorname{Re}(s) \geq 1$ is the pole at $s = 1$, and the extension of step (1) goes through. This is the sense in which the distribution of primes is governed by a single vertical line in the complex plane, and why the analytic input that the elementary theory of chapter 3 could not supply, the absence of zeros on that line, is the input that completes the proof.

5.2 The Laplace Transform

The strategy of section 5.1 reduces the Prime Number Theorem to the single asymptotic $\vartheta(x) \sim x$, and proposition 5.1.1 traces the chain that delivers $\pi(x) \sim x/\log x$ once that asymptotic is in hand. What remains is to prove $\vartheta(x) \sim x$ from the analytic data assembled in chapter 4: the prime series $\Phi(s) = \sum_p (\log p) p^{-s}$ (definition 4.3.2), its simple pole at $s = 1$ with residue 1 (proposition 4.3.4), and the non-vanishing of ζ on the line $\operatorname{Re}(s) = 1$ (corollary 4.4.4). The bridge between the Dirichlet-series object Φ and the counting function ϑ is an integral transform, the *Laplace transform*, and the route runs through the substitution $x = e^t$ that turns the counting function $\vartheta(x)$ on $[1, \infty)$ into a function of t on $[0, \infty)$.

This section records the Laplace transform and computes the two integrals the proof in section 5.3 and section 5.4 requires. The first expresses the transform of $\vartheta(e^t)$ in terms of $\Phi(s)/s$; the second packages this into an auxiliary function $g(s)$ whose holomorphic extension to the closed half-plane $\operatorname{Re}(s) \geq 0$ is the one analytic input Newman's tauberian theorem will consume.

Definition 5.2.1: The Laplace Transform

Let $f: [0, \infty) \rightarrow \mathbb{C}$ be locally integrable. The *Laplace transform* of f is the function

$$(\mathcal{L}f)(s) = \int_0^\infty f(t) e^{-st} dt$$

defined for those $s \in \mathbb{C}$ for which the integral converges.

The Laplace transform is the one-sided companion of the Mellin transform of section 2.4, and the two are linked by the same change of variables that linked the Mellin transform to a two-sided Laplace transform in proposition 2.4.3. Under the substitution $t = e^{-x}$, the Mellin transform $(\mathcal{M}F)(s) = \int_0^\infty F(t) t^{s-1} dt$ becomes the two-sided Laplace integral $\int_{-\infty}^\infty F(e^{-x}) e^{-sx} dx$; restricting a function to the half-line and integrating only over $t \geq 0$ gives the one-sided transform of definition 5.2.1. As with both of those transforms, the modulus $|e^{-st}| = e^{-\operatorname{Re}(s)t}$ depends only on $\operatorname{Re}(s)$, so the region of convergence is a half-plane $\operatorname{Re}(s) > \sigma_0$ invariant under vertical translation, the abscissa σ_0 depending on the growth of f .

Example 5.2.2: The Laplace Transform of a Constant

Take $f(t) = 1$ for all $t \geq 0$. The transform is

$$(\mathcal{L}1)(s) = \int_0^\infty e^{-st} dt = \left[-\frac{1}{s} e^{-st} \right]_{t=0}^{t=\infty} = \frac{1}{s}$$

for $\operatorname{Re}(s) > 0$. The boundary evaluation at $t = \infty$ vanishes because $|e^{-st}| = e^{-\operatorname{Re}(s)t} \rightarrow 0$ exactly

when $\operatorname{Re}(s) > 0$, and the value at $t = 0$ is $-1/s$; the abscissa of convergence is $\sigma_0 = 0$. The transform $1/s$ is holomorphic on $\operatorname{Re}(s) > 0$ and continues meromorphically to $\mathbb{C}$ with a single simple pole at $s = 0$, the boundary point of the half-plane of convergence. This $1/s$ is the term that will subtract against the pole of $\Phi(s+1)/(s+1)$ at $s = 0$ in proposition 5.2.4.

The function whose transform the proof needs is $\vartheta(e^t)$, the Chebyshev theta function (definition 3.1.2) reparametrized by $t = \log x$. Its transform is, up to the factor $1/s$, the prime series Φ .

Proposition 5.2.3: The Laplace Transform of $\vartheta(e^t)$

For $\operatorname{Re}(s) > 1$,

$$\int_0^\infty \vartheta(e^t) e^{-st} dt = \int_1^\infty \vartheta(x) x^{-s-1} dx = \frac{\Phi(s)}{s}$$

where $\Phi(s) = \sum_p (\log p) p^{-s}$ is the prime series of definition 4.3.2.

Proof:

Throughout write $\sigma = \operatorname{Re}(s)$ and assume $\sigma > 1$.

Step 1: The change of variables $x = e^t$. On the integral $\int_0^\infty \vartheta(e^t) e^{-st} dt$ substitute $x = e^t$, so $t = \log x$, $dt = dx/x$, and $e^{-st} = x^{-s}$. As t runs over $[0, \infty)$ the variable x runs over $[1, \infty)$, and

$$\int_0^\infty \vartheta(e^t) e^{-st} dt = \int_1^\infty \vartheta(x) x^{-s} \frac{dx}{x} = \int_1^\infty \vartheta(x) x^{-s-1} dx$$

The substitution $x = e^t$ is a C^1 diffeomorphism of $[0, \infty)$ onto $[1, \infty)$, so it preserves absolute convergence; convergence of the resulting integral is verified in Step 3. This establishes the first equality.

Step 2: Abel summation against $d\vartheta$. The function ϑ is the summatory function of the coefficients $a_n = \log n$ for n prime and $a_n = 0$ otherwise, so $\vartheta(x) = \sum_{n \leq x} a_n$, and its Stieltjes measure $d\vartheta(x)$ places mass $\log p$ at each prime p . Apply Abel summation (theorem 1.4.1) in its Stieltjes form to the integral $\int_1^\infty \vartheta(x) x^{-s-1} dx$. Take the test function $f(x) = x^{-s}$, which is continuously differentiable on $[1, \infty)$ with $f'(x) = -s x^{-s-1}$, so that $x^{-s-1} = -f'(x)/s$. For a finite upper limit $X \geq 1$, integration by parts for the Riemann–Stieltjes integral gives

$$\int_1^X \vartheta(x) x^{-s-1} dx = -\frac{1}{s} \int_1^X \vartheta(x) f'(x) dx = -\frac{1}{s} \left(\left[\vartheta(x) f(x) \right]_1^X - \int_1^X f(x) d\vartheta(x) \right)$$

the boundary term and the Stieltjes integral arising from $\int u dv = [uv] - \int v du$ with $u = \vartheta$, $v = f$. Since $\vartheta(x) = 0$ for $x < 2$, the boundary contribution at $x = 1$ vanishes, and $\int_1^X f(x) d\vartheta(x) = \sum_{p \leq X} (\log p) p^{-s}$ because $d\vartheta$ assigns mass $\log p$ to each prime $p \leq X$ and the integrand f is evaluated at $x = p$. Therefore

$$\int_1^X \vartheta(x) x^{-s-1} dx = -\frac{1}{s} \vartheta(X) X^{-s} + \frac{1}{s} \sum_{p \leq X} \frac{\log p}{p^s}$$

Step 3: The boundary term vanishes and the limit is $\Phi(s)/s$. By Chebyshev's estimates (theorem 3.2.3), $\vartheta(x) = O(x)$, so there is a constant C with $\vartheta(X) \leq CX$ for all $X \geq 1$. Hence

$$|\vartheta(X) X^{-s}| = \vartheta(X) X^{-\sigma} \leq C X^{1-\sigma}$$

and since $\sigma > 1$ the exponent $1 - \sigma$ is negative, so $X^{1-\sigma} \rightarrow 0$ as $X \rightarrow \infty$; the boundary term tends to 0. The same bound shows the integral converges absolutely: $|\vartheta(x) x^{-s-1}| = \vartheta(x) x^{-\sigma-1} \leq Cx \cdot x^{-\sigma-1} = Cx^{-\sigma}$, and $\int_1^\infty x^{-\sigma} dx < \infty$ because $\sigma > 1$. Letting $X \rightarrow \infty$,

$$\int_1^\infty \vartheta(x) x^{-s-1} dx = \frac{1}{s} \sum_p \frac{\log p}{p^s} = \frac{\Phi(s)}{s}$$

the sum on the right being $\Phi(s)$ by definition 4.3.2, convergent on $\sigma > 1$. Combining with Step 1 gives all three equalities, as we sought to show.

The proposition is the analytic dictionary entry that the whole proof turns on: it converts the prime series Φ , an object built from the primes through a Dirichlet series, into the Laplace transform of the counting function $\vartheta(e^t)$, an object built from the same primes through their summatory behavior. The factor $1/s$ is the signature of summation: passing from the coefficients $\log p$ to their partial sums ϑ divides the generating function by s , exactly as the Laplace transform of a constant in example 5.2.2 produced $1/s$ from the summatory function of the unit mass at the origin. The pole of Φ at $s = 1$ (proposition 4.3.4) therefore becomes a pole of $\Phi(s)/s$ at $s = 1$, with the kernel factor $1/s$ contributing the value 1 there, so the residue is again 1; this is the pole that will furnish the main term $\vartheta(x) \sim x$. The representation is valid only on $\operatorname{Re}(s) > 1$, where both Φ converges and the boundary term decays, and the entire difficulty of the Prime Number Theorem lies in pushing analytic control of this transform onto the line $\operatorname{Re}(s) = 1$.

That push is mediated by a single auxiliary function. Newman's tauberian theorem, the analytic engine of section 5.3, applies to the Laplace transform of a *bounded* function whose transform extends holomorphically across the imaginary axis. The function $\vartheta(e^t)$ is not bounded, but the normalized quantity $\vartheta(e^t)e^{-t}$ is, by Chebyshev's estimates, and subtracting its limiting value 1 produces a function tending to a transform with the required regularity.

Proposition 5.2.4: The Auxiliary Function $g(s)$

Define $f: [0, \infty) \rightarrow \mathbb{R}$ by

$$f(t) = \vartheta(e^t) e^{-t} - 1$$

and let $g(s) = \int_0^\infty f(t) e^{-st} dt$ be its Laplace transform. Then for $\operatorname{Re}(s) > 0$,

$$g(s) = \frac{\Phi(s+1)}{s+1} - \frac{1}{s}$$

Proof:

Write $\sigma = \operatorname{Re}(s)$ and assume $\sigma > 0$. The integrand splits as $f(t)e^{-st} = \vartheta(e^t)e^{-t}e^{-st} - e^{-st} = \vartheta(e^t)e^{-(s+1)t} - e^{-st}$, so over any finite range $[0, T]$,

$$\int_0^T f(t) e^{-st} dt = \int_0^T \vartheta(e^t) e^{-(s+1)t} dt - \int_0^T e^{-st} dt$$

the splitting being legitimate on a bounded interval where each piece is integrable. The two pieces are handled separately, then the limit $T \rightarrow \infty$ is taken.

Step 1: The first integral. Since $\sigma > 0$, the shifted argument satisfies $\operatorname{Re}(s+1) = \sigma + 1 > 1$, so proposition 5.2.3 applies at $s+1$ in place of s :

$$\int_0^\infty \vartheta(e^t) e^{-(s+1)t} dt = \frac{\Phi(s+1)}{s+1}$$

the integral converging absolutely because $\operatorname{Re}(s+1) > 1$. In particular the partial integrals $\int_0^T \vartheta(e^t) e^{-(s+1)t} dt$ converge to $\Phi(s+1)/(s+1)$ as $T \rightarrow \infty$.

Step 2: The second integral. By example 5.2.2, for $\sigma > 0$,

$$\int_0^\infty e^{-st} dt = \frac{1}{s}$$

so $\int_0^T e^{-st} dt \rightarrow 1/s$ as $T \rightarrow \infty$.

Step 3: Combine. Both pieces converge as $T \rightarrow \infty$, so their difference converges and $g(s) = \int_0^\infty f(t) e^{-st} dt$ exists for $\sigma > 0$, with

$$g(s) = \frac{\Phi(s+1)}{s+1} - \frac{1}{s}$$

completing the proof.

The function g packages exactly the data Newman's theorem will need. Its definition is engineered so that the integrand $f(t) = \vartheta(e^t)e^{-t} - 1$ is bounded: Chebyshev's estimates (theorem 3.2.3) give $\vartheta(x) \leq (\log 4)x$ for all $x \geq 1$, while $\vartheta(x) \geq 0$ holds trivially, so $0 \leq \vartheta(x)/x \leq \log 4$ for every $x \geq 1$ and the normalized quantity $\vartheta(e^t)e^{-t} = \vartheta(e^t)/e^t$ stays inside $[0, \log 4]$. Hence $f(t) = \vartheta(e^t)/e^t - 1$ stays inside the bounded range $[-1, \log 4 - 1]$ for all $t \geq 0$; the lower Chebyshev bound $\vartheta(x) \geq c_1 x$, valid for large x , pins $f(t)$ eventually above $c_1 - 1$ as well. Boundedness of f on $[0, \infty)$ is the hypothesis the tauberian theorem of section 5.3 imposes on the function being transformed. The formula $g(s) = \Phi(s+1)/(s+1) - 1/s$ then exhibits g , on $\operatorname{Re}(s) > 0$, as a difference of two functions each carrying a pole at $s = 0$. The shifted prime series $\Phi(s+1)$ has its pole at $s+1 = 1$, that is, at $s = 0$, with residue 1 (proposition 4.3.4), so $\Phi(s+1)/(s+1)$ has a simple pole at $s = 0$ with residue 1; the second term $1/s$ has a simple pole at $s = 0$ with residue 1 as well. The two principal parts cancel: g is holomorphic at $s = 0$, where it would otherwise inherit the singularity of Φ .

This cancellation is the crux. The entire proof of the Prime Number Theorem reduces to showing that g extends holomorphically to the closed half-plane $\operatorname{Re}(s) \geq 0$, and the analytic facts of chapter 4 are precisely what make that extension possible. At $s = 0$ the pole of Φ at the point $s = 1$ is exactly cancelled by the term $1/s$, as computed above; this is where the residue-1 pole of Φ enters, and it is the only point of the line $\operatorname{Re}(s) = 0$ at which $\Phi(s+1)$ is singular. At every other point $s = it$ with $t \neq 0$, the shifted argument $s+1 = 1+it$ lies on the line $\operatorname{Re} = 1$ away from the pole, where Φ is holomorphic because ζ has no zeros there (corollary 4.4.4), the non-vanishing of ζ on $\operatorname{Re}(s) = 1$ established in chapter 4. The two inputs together, the pole of Φ at $s = 1$ cancelling $1/s$ and the non-vanishing of ζ on the rest of the line, give g a holomorphic extension to a neighborhood of

$\operatorname{Re}(s) \geq 0$, the hypothesis that drives section 5.3 and section 5.4.

A numerical look at f makes its boundedness, the structural feature that lets the argument proceed, concrete.

Example 5.2.5: g at a Real Point

The integrand $f(t) = \vartheta(e^t)/e^t - 1$ measures the deviation of $\vartheta(x)/x$ from 1 at $x = e^t$. Tabulating $\vartheta(x)/x - 1$ at a range of x , using the values of $\vartheta(x)$ from example 3.1.3 together with further computation, gives

x	$\vartheta(x)/x$	$\vartheta(x)/x - 1$
10	0.5347	-0.4653
100	0.8373	-0.1627
1000	0.9562	-0.0438
10000	0.9896	-0.0104
100000	0.9969	-0.0031

Every entry in the last column lies in the interval $[-1, \log 4 - 1] \approx [-1, 0.386]$ guaranteed by theorem 3.2.3, since $0 \leq \vartheta(x)/x \leq \log 4$ for all $x \geq 1$; the upper constant $\log 4 \approx 1.386$ caps $\vartheta(x)/x$ for every x , and the lower Chebyshev constant $\log 2 \approx 0.693$ confines $\vartheta(x)/x$ from below once x is large, so $\vartheta(x)/x - 1$ cannot escape a fixed range no matter how large x grows. The table also shows the values drifting toward 0, consistent with the conjectured limit $\vartheta(x)/x \rightarrow 1$; but the proof does not assume this convergence. It assumes only the boundedness visible in the table, that $f(t)$ stays within fixed bounds for all $t \geq 0$. This boundedness is exactly the hypothesis Newman's theorem (section 5.3) will require of f , and the convergence $\vartheta(x)/x \rightarrow 1$ is the conclusion the theorem will deliver, not an input it demands.

5.3 Newman's Tauberian Theorem

The analytic engine of the proof of the Prime Number Theorem is a single theorem of D. J. Newman about Laplace transforms [6], in the streamlined form given by Zagier [18]. The Perron representation of example 2.5.4 expresses the prime-counting data through a contour integral of $-\zeta'/\zeta$, and chapter 4 supplied the analytic input that the integrand has no pole on the line $\operatorname{Re}(s) = 1$ except the one at $s = 1$. What remains is to convert that holomorphy-up-to-the-line into an asymptotic for the counting function, and the conversion is exactly the business of a *Tauberian* theorem: a result that upgrades a statement about the boundary behavior of a transform into a statement about convergence of the function it transforms. Newman's theorem is the cleanest such statement adapted to the situation at hand, and its proof is a short contour argument requiring nothing past Cauchy's integral formula.

The shape of the statement is the following. A bounded function f on $[0, \infty)$ has a Laplace transform $g(s) = \int_0^\infty f(t)e^{-st} dt$ that converges and is holomorphic on the open half-plane $\operatorname{Re}(s) > 0$. Convergence of the *improper integral* $\int_0^\infty f(t) dt$ is the assertion that g extends continuously to the boundary point $s = 0$ with the limit equal to the integral, and this is a strictly stronger demand than boundedness of f alone. Newman's theorem supplies the missing hypothesis in the form most useful for the zeta function: if g extends not just continuously but *holomorphically* across the imaginary axis, then the improper integral converges and equals $g(0)$. Holomorphy up to the line is the Tauberian side condition, and boundedness of f is the standing hypothesis that the application will draw from Chebyshev's bound.

Theorem 5.3.1: Newman's Analytic Theorem

Let $f: [0, \infty) \rightarrow \mathbb{C}$ be bounded and integrable (Riemann or Lebesgue) on every finite interval $[0, T]$, and for $\operatorname{Re}(s) > 0$ define

$$g(s) = \int_0^\infty f(t)e^{-st} dt$$

The integral converges absolutely and g is holomorphic on $\operatorname{Re}(s) > 0$. Suppose g extends to a holomorphic function on an open set containing the closed half-plane $\operatorname{Re}(s) \geq 0$. Then the improper integral $\int_0^\infty f(t) dt$ converges and

$$\int_0^\infty f(t) dt = g(0)$$

Proof:

Write $B = \sup_{t \geq 0} |f(t)|$, finite by hypothesis. For $\operatorname{Re}(s) > 0$ the bound $|f(t)e^{-st}| \leq Be^{-\operatorname{Re}(s)t}$ and $\int_0^\infty e^{-\operatorname{Re}(s)t} dt = 1/\operatorname{Re}(s) < \infty$ show that the defining integral converges absolutely; that g is holomorphic on $\operatorname{Re}(s) > 0$ is the holomorphy of a parameter integral whose integrand is holomorphic in s and dominated locally uniformly, established in [16]. The substance is the boundary statement, proved by a contour integral.

Step 1: The truncated transforms and the goal. For each $T > 0$ set

$$g_T(s) = \int_0^T f(t)e^{-st} dt$$

The integrand is entire in s for each fixed t , and the integral runs over the finite interval $[0, T]$, so g_T is entire (again by holomorphy of a parameter integral, now over a compact interval, [16]). The improper integral $\int_0^\infty f(t) dt$ is by definition $\lim_{T \rightarrow \infty} \int_0^T f(t) dt = \lim_{T \rightarrow \infty} g_T(0)$, so the assertion to be proved is exactly

$$g_T(0) \longrightarrow g(0) \quad (T \rightarrow \infty)$$

The difference $g(0) - g_T(0)$ will be written as a contour integral and estimated.

Step 2: A contour and the Cauchy representation. Fix $R > 0$ for the remainder of the argument. By hypothesis g is holomorphic on an open set $\Omega \supseteq \{\operatorname{Re}(s) \geq 0\}$. The intersection of Ω with the disc $\{|s| \leq R\}$ is an open neighborhood of the compact arc $\{|s| = R, \operatorname{Re}(s) \geq 0\} \cup \{\operatorname{Re}(s) = 0, |s| \leq R\}$, so there is a $\delta = \delta(R) > 0$, which may be taken with $\delta < R$, such that g is holomorphic on a neighborhood of the closed region

$$D = \{s \mid |s| \leq R, \operatorname{Re}(s) \geq -\delta\}$$

Let $C = \partial D$ be its positively oriented boundary, consisting of the major arc $A = \{|s| = R, \operatorname{Re}(s) \geq -\delta\}$ of the circle of radius R together with the vertical segment $\{\operatorname{Re}(s) = -\delta, |\operatorname{Im}(s)| \leq \sqrt{R^2 - \delta^2}\}$. The point $s = 0$ lies in the interior of D , and g_T is entire, so

the function

$$s \mapsto (g(s) - g_T(s)) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right)$$

is holomorphic on a neighborhood of D except for a simple pole at $s = 0$. The factor $1/s + s/R^2$ has residue 1 at $s = 0$ (the term s/R^2 is regular there), and $e^{sT} = 1$ at $s = 0$, so the residue of the displayed function at $s = 0$ is $g(0) - g_T(0)$. Cauchy's residue theorem ([16]) gives

$$g(0) - g_T(0) = \frac{1}{2\pi i} \int_C (g(s) - g_T(s)) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) ds \quad (5.1)$$

The plan is to split C into the part in $\operatorname{Re}(s) > 0$ and the part in $\operatorname{Re}(s) \leq 0$, and to bound each.

Step 3: The kernel on the circle $|s| = R$. On the circle $|s| = R$ one has $\bar{s} = R^2/s$, hence

$$\frac{1}{s} + \frac{s}{R^2} = \frac{\bar{s}}{R^2} + \frac{s}{R^2} = \frac{s + \bar{s}}{R^2} = \frac{2\operatorname{Re}(s)}{R^2}$$

so that on $|s| = R$,

$$\left| e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) \right| = e^{\operatorname{Re}(s)T} \frac{2|\operatorname{Re}(s)|}{R^2} \quad (5.2)$$

This identity is the reason the second term s/R^2 was inserted; it is recorded for use on both halves of the contour.

Step 4: The right half $C_+ = C \cap \{\operatorname{Re}(s) > 0\}$. Here C_+ is the right portion $\{|s| = R, \operatorname{Re}(s) > 0\}$ of the circle, a semicircle of length πR . For $\operatorname{Re}(s) > 0$ the tail estimate

$$|g(s) - g_T(s)| = \left| \int_T^\infty f(t) e^{-st} dt \right| \leq B \int_T^\infty e^{-\operatorname{Re}(s)t} dt = \frac{B e^{-\operatorname{Re}(s)T}}{\operatorname{Re}(s)}$$

holds, where the first equality uses that the integral defining $g(s)$ converges and exceeds that defining $g_T(s)$ by the tail over $[T, \infty)$. Multiplying by the kernel bound (5.2) and using $|\operatorname{Re}(s)| = \operatorname{Re}(s) > 0$,

$$\left| (g(s) - g_T(s)) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) \right| \leq \frac{B e^{-\operatorname{Re}(s)T}}{\operatorname{Re}(s)} \cdot e^{\operatorname{Re}(s)T} \frac{2\operatorname{Re}(s)}{R^2} = \frac{2B}{R^2}$$

The integrand on C_+ is bounded by $2B/R^2$ pointwise, so by the standard length estimate ([16]) the contribution of C_+ to (5.1) satisfies

$$\left| \frac{1}{2\pi i} \int_{C_+} (g(s) - g_T(s)) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) ds \right| \leq \frac{1}{2\pi} \cdot \frac{2B}{R^2} \cdot \pi R = \frac{B}{R}$$

This bound is uniform in T .

Step 5: The left half, the g_T part. Let $C_- = C \cap \{\operatorname{Re}(s) \leq 0\}$ be the remainder of the contour: the left portion of the arc A together with the vertical segment at $\operatorname{Re}(s) = -\delta$. On C_- the

integrand of (5.1) is treated by separating g from g_T , which is legitimate because each piece is integrable on the compact contour. Consider first the g_T piece,

$$I_T^- = \frac{1}{2\pi i} \int_{C_-} g_T(s) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) ds$$

Since g_T is entire, the integrand $g_T(s) e^{sT} (1/s + s/R^2)$ is holomorphic on the punctured plane $\mathbb{C} \setminus \{0\}$, in particular on the open left half-plane $\{\operatorname{Re}(s) < 0\}$. The contour C_- and the left semicircle

$$C'_- = \{|s| = R, \operatorname{Re}(s) \leq 0\}$$

share the two endpoints $\pm iR$, and the closed region they bound lies entirely in $\{\operatorname{Re}(s) \leq 0\}$, with the only point of nonholomorphy $s = 0$ on its boundary rather than its interior. Cauchy's theorem ([16]) therefore permits replacing C_- by C'_- in I_T^- without changing its value. On C'_- one has $\operatorname{Re}(s) \leq 0$, and for such s the entire transform is bounded by

$$|g_T(s)| = \left| \int_0^T f(t) e^{-st} dt \right| \leq B \int_0^T e^{-\operatorname{Re}(s)t} dt = B \frac{e^{-\operatorname{Re}(s)T} - 1}{-\operatorname{Re}(s)} \leq \frac{B e^{-\operatorname{Re}(s)T}}{|\operatorname{Re}(s)|}$$

using $-\operatorname{Re}(s) = |\operatorname{Re}(s)| > 0$ on the open left semicircle and $e^{-\operatorname{Re}(s)T} - 1 \leq e^{-\operatorname{Re}(s)T}$. Multiplying by the kernel bound (5.2),

$$\left| g_T(s) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) \right| \leq \frac{B e^{-\operatorname{Re}(s)T}}{|\operatorname{Re}(s)|} \cdot e^{\operatorname{Re}(s)T} \frac{2|\operatorname{Re}(s)|}{R^2} = \frac{2B}{R^2}$$

the factors $e^{\pm \operatorname{Re}(s)T}$ and $|\operatorname{Re}(s)|$ cancelling exactly as on the right half. The semicircle C'_- has length πR , so the length estimate gives

$$|I_T^-| \leq \frac{1}{2\pi} \cdot \frac{2B}{R^2} \cdot \pi R = \frac{B}{R}$$

again uniformly in T .

Step 6: The left half, the g part. The complementary piece is

$$J_T^- = \frac{1}{2\pi i} \int_{C_-} g(s) e^{sT} \left(\frac{1}{s} + \frac{s}{R^2} \right) ds$$

Here R , hence δ and the contour C_- , are fixed, and $T \rightarrow \infty$. The function $g(s) (1/s + s/R^2)$ is continuous on the compact set C_- : the only candidate for a singularity is $s = 0$, but C_- has $\operatorname{Re}(s) \leq 0$ and meets the imaginary axis only at the two points $\pm iR$, where $|s| = R \neq 0$, so $s = 0 \notin C_-$ and the factor $1/s$ is bounded there. Let

$$M = \max_{s \in C_-} \left| g(s) \left(\frac{1}{s} + \frac{s}{R^2} \right) \right| < \infty$$

a constant depending on R and δ but not on T . The remaining factor satisfies $|e^{sT}| = e^{\operatorname{Re}(s)T} \leq 1$ on C_- , since $\operatorname{Re}(s) \leq 0$, so the integrand of J_T^- is dominated by the constant M on the finite-length contour C_- , uniformly in T . For each fixed $s \in C_-$ with $\operatorname{Re}(s) < 0$ one has $e^{\operatorname{Re}(s)T} \rightarrow 0$ as $T \rightarrow \infty$; the exceptional points $\pm iR$, where $\operatorname{Re}(s) = 0$, form a set of measure zero on the

contour. By the dominated convergence theorem ([8]), applied to the arc-length integral with dominating function M ,

$$J_T^- \longrightarrow 0 \quad (T \rightarrow \infty)$$

This is the only one of the three pieces that depends on letting T grow, and it is where holomorphy of g across the line is used: only because g continues to $\operatorname{Re}(s) = -\delta$ can the contour C_- be pushed off the imaginary axis into the region $\operatorname{Re}(s) < 0$ where e^{sT} decays.

Step 7: Assembling the estimate. Splitting (5.1) as $C = C_+ \cup C_-$ and the C_- integral as its g and g_T parts,

$$g(0) - g_T(0) = \underbrace{\frac{1}{2\pi i} \int_{C_+} (\cdots) ds}_{\text{Step 4}} - \underbrace{I_T^-}_{\text{Step 5}} + \underbrace{J_T^-}_{\text{Step 6}}$$

where the sign on I_T^- records that the g_T contribution over C_- enters (5.1) as $\int_{C_-} (g - g_T) = J_T^- - I_T^-$. Steps 4 and 5 bound the first two terms by B/R each, uniformly in T , and Step 6 sends the third to 0. Hence

$$\limsup_{T \rightarrow \infty} |g(0) - g_T(0)| \leq \frac{B}{R} + \frac{B}{R} + 0 = \frac{2B}{R}$$

The left-hand side does not depend on R , while the right-hand side is $2B/R$ for every $R > 0$. Letting $R \rightarrow \infty$ forces $\limsup_{T \rightarrow \infty} |g(0) - g_T(0)| = 0$, so $g_T(0) \rightarrow g(0)$. By Step 1 this is the statement that $\int_0^\infty f(t) dt$ converges with value $g(0)$, completing the proof.

The hypothesis does double duty, and its two halves play distinct roles. Boundedness of f is what makes the tail estimates of Steps 4 and 5 produce the bound $2B/R^2$ on the circle; without a uniform bound on f the kernel could not absorb the growth of $e^{\operatorname{Re}(s)T}$ on the right, and the argument would collapse. Holomorphy of g across the line is the Tauberian content: it is the hypothesis that lets the contour in Step 6 be displaced into $\{\operatorname{Re}(s) < 0\}$, where the exponential e^{sT} decays and the g -integral vanishes in the limit. Strip away the holomorphy and only boundedness of f remains, which guarantees that g is defined and holomorphic on $\operatorname{Re}(s) > 0$ but says nothing about the boundary; the improper integral $\int_0^\infty f(t) dt$ may then diverge by oscillation even though f is bounded, as the example $f(t) = \cos t$ shows, where $g(s) = s/(s^2 + 1)$ has poles at $s = \pm i$ on the imaginary axis and the holomorphy hypothesis fails.

The name *Tauberian* marks the logical direction of the theorem. An *Abelian* theorem deduces the behavior of a transform from the convergence of the function: if $\int_0^\infty f(t) dt$ converges, then $g(s) \rightarrow g(0)$ as $s \rightarrow 0^+$, a statement of continuity that is comparatively soft. A Tauberian theorem reverses the implication, recovering convergence of $\int_0^\infty f(t) dt$ from analytic regularity of g at the boundary, and such reversals are false without a side condition that restrains the function from oscillating, the *Tauberian condition*. In Newman's theorem the side condition is the boundedness of f together with the holomorphic extension of g , and the value $g(0)$ that the holomorphic transform assigns to the boundary point is promoted from a formal or Abel-summed value into the genuine value of the improper integral.

5.3.2 Key Ideas: The Kernel $e^{sT}(1/s + s/R^2)$ The factor $e^{sT}(1/s + s/R^2)$ in (5.1) is engineered so that a single contour integral does three things at once. The term $1/s$ carries a simple pole at $s = 0$ with residue 1; against the holomorphic factor $g(s) - g_T(s)$ it produces the residue $g(0) - g_T(0)$, which is the quantity to be estimated. The factor e^{sT} is the device that lets the truncation $g - g_T$ be controlled: on the right half-plane the tail $g(s) - g_T(s)$ carries a decay $e^{-\operatorname{Re}(s)T}$, and the inserted e^{sT} cancels it, while on the left half-plane the same e^{sT} decays and kills the g -integral as $T \rightarrow \infty$. The role of the added term s/R^2 is to tame the kernel on the circle $|s| = R$: by the identity $1/s + s/R^2 = 2\operatorname{Re}(s)/R^2$ valid on that circle, the modulus of the kernel collapses to $e^{\operatorname{Re}(s)T} (2|\operatorname{Re}(s)|/R^2)$, in which the prefactor $2|\operatorname{Re}(s)|/R^2$ supplies exactly the factor $\operatorname{Re}(s)$ needed to cancel the $1/\operatorname{Re}(s)$ from the tail bound $Be^{-\operatorname{Re}(s)T}/\operatorname{Re}(s)$, and the residual $e^{\pm\operatorname{Re}(s)T}$ cancels the exponential. The net bound $2B/R^2$ on both semicircles is symmetric in the two halves precisely because the kernel modulus depends only on $|\operatorname{Re}(s)|$. Without the s/R^2 term the bare $1/s$ would contribute $1/R$ on the circle with no $\operatorname{Re}(s)$ factor to absorb the $1/\operatorname{Re}(s)$ from the tail, and the right-half estimate would diverge as s approaches the imaginary axis; the corrective term costs nothing at $s = 0$, where it is regular, so the pole and its residue are untouched.

The application in section 5.4 fits the theorem exactly. The function to which Newman's theorem will be applied is built from the Chebyshev theta function ϑ (definition 3.1.2), arranged so that the boundedness hypothesis becomes the assertion that $\vartheta(x)/x$ is bounded, which is the content of Chebyshev's estimates $c_1x \leq \vartheta(x) \leq c_2x$ in theorem 3.2.3. The holomorphic-extension hypothesis becomes the non-vanishing of ζ on $\operatorname{Re}(s) = 1$ from theorem 4.4.2, in the form that $-\zeta'/\zeta$, and with it the relevant transform, continues across the line save for the controlled pole at $s = 1$ (corollary 4.4.4). Newman's theorem then returns the convergence of an improper integral in ϑ , and a short argument converts that convergence into $\vartheta(x) \sim x$, hence into the Prime Number Theorem through the equivalence of theorem 3.1.5. The theorem proved here is thus the one analytic lever that turns the boundary data on ζ into an asymptotic count of primes.

5.4 Proof of the Prime Number Theorem

The pieces assembled in the three preceding sections now combine into the theorem. The auxiliary function g of section 5.2, defined on $\operatorname{Re}(s) > 0$ as the Laplace transform of $f(t) = \vartheta(e^t)e^{-t} - 1$, was computed in proposition 5.2.4 to be $g(s) = \Phi(s+1)/(s+1) - 1/s$. The pole of Φ at $s = 1$ (proposition 4.3.4) cancels against the term $1/s$ at $s = 0$, and the non-vanishing of ζ on the line $\operatorname{Re}(s) = 1$ (corollary 4.4.4, equivalently theorem 4.4.2) removes every other obstruction on the imaginary axis; together they extend g holomorphically across $\operatorname{Re}(s) = 0$. Newman's theorem (theorem 5.3.1) then converts that extension into the convergence of $\int_0^\infty f(t) dt$, which the substitution $x = e^t$ turns into the convergence of $\int_1^\infty (\vartheta(x) - x)/x^2 dx$.

What remains after that convergence is purely real-variable. The integrand $(\vartheta(x) - x)/x^2$ records the normalized deviation of ϑ from its conjectured asymptotic, and the monotonicity of ϑ forbids that deviation from persisting: a function that climbs only upward cannot keep $\vartheta(x)/x$ away from 1 without forcing the integral to diverge. This section carries out both halves, the analytic convergence and the monotonicity argument, and assembles them into the headline result of the chapter.

Theorem 5.4.1: Convergence of the Key Integral

The improper integral

$$\int_1^\infty \frac{\vartheta(x) - x}{x^2} dx$$

converges.

Proof:

The proof applies Newman's theorem (theorem 5.3.1) to the function f of proposition 5.2.4 and then changes variables. The work is in verifying the holomorphic-extension hypothesis, which occupies Steps 2 and 3.

Step 1: The function and its transform. Define $f: [0, \infty) \rightarrow \mathbb{R}$ by

$$f(t) = \vartheta(e^t) e^{-t} - 1$$

The function f is bounded: by Chebyshev's estimates (theorem 3.2.3) one has $0 \leq \vartheta(x) \leq (\log 4)x$ for all $x \geq 1$, so writing $x = e^t$ gives $0 \leq \vartheta(e^t) e^{-t} = \vartheta(x)/x \leq \log 4$, whence $-1 \leq f(t) \leq \log 4 - 1$ for every $t \geq 0$. The function f is also locally integrable, indeed Riemann integrable on every finite interval $[0, T]$, since $\vartheta(e^t)$ is a nondecreasing step function with finitely many jumps on any bounded interval and e^{-t} is continuous. Both hypotheses on the transformed function in theorem 5.3.1 are thus met. Its Laplace transform is, by proposition 5.2.4, the function

$$g(s) = \int_0^\infty f(t) e^{-st} dt = \frac{\Phi(s+1)}{s+1} - \frac{1}{s}, \quad \operatorname{Re}(s) > 0 \quad (5.3)$$

holomorphic on $\operatorname{Re}(s) > 0$. To invoke Newman's theorem it remains to show that g extends holomorphically to an open set containing the closed half-plane $\operatorname{Re}(s) \geq 0$.

Step 2: Holomorphic extension at $s = 0$. Near $s = 0$ the shifted argument $s + 1$ is near 1, where Φ has its only singularity. By proposition 4.3.4 the prime series has a simple pole at $s = 1$ with residue 1, so $\Phi(w) - 1/(w - 1)$ is holomorphic in a neighborhood of $w = 1$; setting $w = s + 1$, the function

$$\Phi(s+1) - \frac{1}{s}$$

is holomorphic in a neighborhood of $s = 0$, since $w - 1 = s$. Call this holomorphic function $h_0(s)$, so that $\Phi(s+1) = 1/s + h_0(s)$ near $s = 0$. Substituting into (5.3),

$$g(s) = \frac{1}{s+1} \left(\frac{1}{s} + h_0(s) \right) - \frac{1}{s} = \frac{1}{s(s+1)} - \frac{1}{s} + \frac{h_0(s)}{s+1}$$

The first two terms combine through the partial-fraction identity $1/(s(s+1)) = 1/s - 1/(s+1)$, which gives

$$\frac{1}{s(s+1)} - \frac{1}{s} = \left(\frac{1}{s} - \frac{1}{s+1} \right) - \frac{1}{s} = -\frac{1}{s+1}$$

so the singular term $1/s$ cancels exactly, leaving

$$g(s) = -\frac{1}{s+1} + \frac{h_0(s)}{s+1} = \frac{h_0(s) - 1}{s+1} \quad (5.4)$$

near $s = 0$. The numerator $h_0(s) - 1$ is holomorphic near $s = 0$, and the denominator $s + 1$ is nonzero there (it equals 1 at $s = 0$), so the right-hand side of (5.4) is holomorphic in a neighborhood of $s = 0$. This is the holomorphic extension of g to the origin.

Step 3: Holomorphic extension at $s = i\tau$ with $\tau \neq 0$. Fix a real $\tau \neq 0$ and consider the point $s = i\tau$ on the imaginary axis. The shifted argument is $s + 1 = 1 + i\tau$, a point on the line $\text{Re} = 1$ distinct from the pole $s = 1$. By proposition 4.3.4 the prime series decomposes as

$$\Phi(w) = -\frac{\zeta'(w)}{\zeta(w)} - R(w)$$

where R is holomorphic on $\text{Re}(w) > 1/2$. The point $w = 1 + i\tau$ lies in this half-plane, so R is holomorphic in a neighborhood of it. The logarithmic derivative $-\zeta'/\zeta$ is holomorphic on the line $\text{Re}(w) = 1$ except for the simple pole at $w = 1$, by corollary 4.4.4, which records the non-vanishing $\zeta(1 + i\tau) \neq 0$ for $\tau \neq 0$ of theorem 4.4.2; since $\tau \neq 0$, the point $1 + i\tau$ is not the pole, so $-\zeta'/\zeta$ is holomorphic in a neighborhood of $w = 1 + i\tau$. Therefore Φ , the difference of two functions holomorphic near $w = 1 + i\tau$, is holomorphic in a neighborhood of that point, hence so is $w \mapsto \Phi(w)/w$ because $w = 1 + i\tau \neq 0$. Translating back by $w = s + 1$, the function $\Phi(s + 1)/(s + 1)$ is holomorphic in a neighborhood of $s = i\tau$. The remaining term $-1/s$ of (5.3) is holomorphic at $s = i\tau$ as well, since $i\tau \neq 0$. Their sum g is thus holomorphic in a neighborhood of every point $i\tau$ with $\tau \neq 0$.

Step 4: Assembling the extension and applying Newman's theorem. The function g is holomorphic on the open half-plane $\text{Re}(s) > 0$ by Step 1. Step 2 gives a neighborhood of $s = 0$ on which g extends holomorphically, and Step 3 gives, for each $\tau \neq 0$, a neighborhood of $i\tau$ on which g extends holomorphically; the extensions agree with g on the overlaps with $\text{Re}(s) > 0$, where they all equal the original transform, so by the identity theorem ([16]) they patch into a single holomorphic function on the union. That union is an open set containing the open half-plane $\text{Re}(s) > 0$ together with every point of the imaginary axis $\text{Re}(s) = 0$, hence an open set containing the closed half-plane $\text{Re}(s) \geq 0$. The hypotheses of theorem 5.3.1 are satisfied: f is bounded and locally integrable, and its transform g extends holomorphically across the line. The theorem concludes that the improper integral $\int_0^\infty f(t) dt$ converges (with value $g(0)$).

Step 5: The change of variables $x = e^t$. The convergence just obtained is

$$\int_0^\infty (\vartheta(e^t) e^{-t} - 1) dt = \lim_{T \rightarrow \infty} \int_0^T (\vartheta(e^t) e^{-t} - 1) dt < \infty$$

On each truncation $\int_0^T$ substitute $x = e^t$, so $t = \log x$, $dt = dx/x$, and $e^{-t} = 1/x$; as t runs over $[0, T]$ the variable x runs over $[1, e^T]$. The substitution is a C^1 diffeomorphism of $[0, T]$ onto $[1, e^T]$, so

$$\int_0^T (\vartheta(e^t) e^{-t} - 1) dt = \int_1^{e^T} \left(\frac{\vartheta(x)}{x} - 1 \right) \frac{dx}{x} = \int_1^{e^T} \frac{\vartheta(x) - x}{x^2} dx$$

the last equality combining the two terms over the common denominator x^2 . As $T \rightarrow \infty$ the upper limit $e^T \rightarrow \infty$ through all values, and the left-hand side converges by Step 4, so the right-hand side converges to the same finite limit:

$$\int_1^\infty \frac{\vartheta(x) - x}{x^2} dx = \int_0^\infty (\vartheta(e^t) e^{-t} - 1) dt < \infty$$

Therefore the improper integral converges, as we sought to show.

The theorem is the analytic payload of the entire chapter delivered in a single real statement. Everything complex-analytic has been spent: the meromorphic continuation of ζ , the pole at $s = 1$, the non-vanishing on the line, Newman's contour argument. What survives is the assertion that $\vartheta(x) - x$, measured against the natural scale x^2 that makes the integrand the logarithmic-mean deviation $(\vartheta(x)/x - 1)/x$, integrates to a finite value. This is already a strong statement about how rarely and by how little $\vartheta(x)$ can stray from x , but it is not yet the asymptotic $\vartheta(x) \sim x$. A finite integral permits the integrand to be large on thin sets, and a priori $\vartheta(x)/x$ could oscillate, dipping below 1 and rising above it infinitely often, so long as the excursions are brief enough that their total weighted contribution stays bounded. Ruling out such oscillation is the role of monotonicity, and it is the one property of ϑ that the analytic machinery never used.

The next result isolates exactly that real-variable principle. It is stated for a general nondecreasing function so that the mechanism, monotonicity against a convergent integral, stands clear of the arithmetic.

Lemma 5.4.2: Monotonicity Forces the Asymptotic

Let $\vartheta: [1, \infty) \rightarrow [0, \infty)$ be nondecreasing, and suppose the integral

$$\int_1^\infty \frac{\vartheta(x) - x}{x^2} dx$$

converges. Then $\vartheta(x) \sim x$ as $x \rightarrow \infty$, that is, $\vartheta(x)/x \rightarrow 1$.

Proof:

The argument is by contradiction, ruling out in turn that $\vartheta(x)/x$ exceeds 1 along a sequence and that it falls below 1 along a sequence. Both directions exploit the same device: over an interval whose endpoints are in a fixed ratio, monotonicity of ϑ bounds the integrand below (respectively above) by a positive (respectively negative) constant independent of where the interval sits, while the Cauchy criterion for the convergent integral forces the integral over any such far-out interval to be small. The Cauchy criterion is the statement that since $\int_1^\infty (\vartheta(t) - t)/t^2 dt$ converges, for every $\varepsilon > 0$ there is an X_0 with

$$\left| \int_a^b \frac{\vartheta(t) - t}{t^2} dt \right| < \varepsilon \quad \text{whenever } X_0 \leq a \leq b \quad (5.5)$$

the integral over $[a, b]$ being the difference of the partial integrals up to b and up to a .

Step 1: The upper excursion cannot occur. Suppose, for contradiction, that there is a constant $\lambda > 1$ and arbitrarily large x with $\vartheta(x) \geq \lambda x$. Fix such an x and consider the interval $[x, \lambda x]$. For t in this interval, $t \leq \lambda x$, and because ϑ is nondecreasing and $t \geq x$,

$$\vartheta(t) \geq \vartheta(x) \geq \lambda x$$

so the numerator satisfies $\vartheta(t) - t \geq \lambda x - t \geq 0$ throughout $[x, \lambda x]$, the last inequality because $t \leq \lambda x$. Hence

$$\int_x^{\lambda x} \frac{\vartheta(t) - t}{t^2} dt \geq \int_x^{\lambda x} \frac{\lambda x - t}{t^2} dt$$

In the integral on the right substitute $t = ux$, so $dt = x du$, $t^2 = u^2 x^2$, and as t runs over $[x, \lambda x]$ the variable u runs over $[1, \lambda]$:

$$\int_x^{\lambda x} \frac{\lambda x - t}{t^2} dt = \int_1^\lambda \frac{\lambda x - ux}{u^2 x^2} x du = \int_1^\lambda \frac{\lambda - u}{u^2} du$$

the factors of x cancelling completely. The resulting quantity

$$c_\lambda^+ := \int_1^\lambda \frac{\lambda - u}{u^2} du$$

is a positive constant depending only on λ , not on x : the integrand $(\lambda - u)/u^2$ is strictly positive for $u \in [1, \lambda)$ and continuous, so its integral over $[1, \lambda]$ is strictly positive. The antiderivative $\int (\lambda - u)/u^2 du = -\lambda/u - \log u$ evaluates it to $c_\lambda^+ = \lambda - 1 - \log \lambda$, which is positive for every $\lambda > 1$. Thus for every one of the arbitrarily large x with $\vartheta(x) \geq \lambda x$,

$$\int_x^{\lambda x} \frac{\vartheta(t) - t}{t^2} dt \geq c_\lambda^+ > 0$$

a fixed positive lower bound. But by the Cauchy criterion (5.5) with $\varepsilon = c_\lambda^+/2$, there is an X_0 beyond which every such integral, over an interval $[x, \lambda x]$ with $x \geq X_0$, has absolute value below $c_\lambda^+/2$. Choosing one of the arbitrarily large x with $x \geq X_0$ produces an integral that is simultaneously $\geq c_\lambda^+$ and $< c_\lambda^+/2$, a contradiction. Hence no such $\lambda > 1$ exists, and $\limsup_{x \rightarrow \infty} \vartheta(x)/x \leq 1$.

Step 2: The lower excursion cannot occur. Suppose, for contradiction, that there is a constant λ with $0 < \lambda < 1$ and arbitrarily large x with $\vartheta(x) \leq \lambda x$. Fix such an x and consider now the interval $[\lambda x, x]$, which lies below x . For t in this interval, $t \geq \lambda x$, and because ϑ is nondecreasing and $t \leq x$,

$$\vartheta(t) \leq \vartheta(x) \leq \lambda x$$

so the numerator satisfies $\vartheta(t) - t \leq \lambda x - t \leq 0$ throughout $[\lambda x, x]$, the last inequality because $t \geq \lambda x$. Hence

$$\int_{\lambda x}^x \frac{\vartheta(t) - t}{t^2} dt \leq \int_{\lambda x}^x \frac{\lambda x - t}{t^2} dt$$

Substituting $t = ux$ as before, with u now running over $[\lambda, 1]$,

$$\int_{\lambda x}^x \frac{\lambda x - t}{t^2} dt = \int_\lambda^1 \frac{\lambda - u}{u^2} du$$

again with all factors of x cancelling. The quantity

$$c_\lambda^- := \int_\lambda^1 \frac{\lambda - u}{u^2} du$$

is a strictly negative constant depending only on λ : on $(\lambda, 1]$ the integrand $(\lambda - u)/u^2$ is strictly negative and continuous, so $c_\lambda^- < 0$. The same antiderivative $-\lambda/u - \log u$ evaluates it to $c_\lambda^- = 1 - \lambda + \log \lambda$, which is negative for every λ with $0 < \lambda < 1$. Thus for every one of the arbitrarily large x with $\vartheta(x) \leq \lambda x$,

$$\int_{\lambda x}^x \frac{\vartheta(t) - t}{t^2} dt \leq c_\lambda^- < 0$$

a fixed negative upper bound. The Cauchy criterion (5.5) with $\varepsilon = |c_\lambda^-|/2$ gives an X_0 beyond which every integral over $[\lambda x, x]$ with $\lambda x \geq X_0$ has absolute value below $|c_\lambda^-|/2$. Selecting an arbitrarily large x with $\lambda x \geq X_0$ yields an integral that is at once $\leq c_\lambda^- = -|c_\lambda^-|$ and greater than $-|c_\lambda^-|/2$, a contradiction. Hence no such $\lambda \in (0, 1)$ exists, and $\liminf_{x \rightarrow \infty} \vartheta(x)/x \geq 1$.

Steps 1 and 2 give $\limsup_{x \rightarrow \infty} \vartheta(x)/x \leq 1 \leq \liminf_{x \rightarrow \infty} \vartheta(x)/x$. Since the liminf never exceeds the limsup, both equal 1, so $\lim_{x \rightarrow \infty} \vartheta(x)/x = 1$, completing the proof.

The lemma turns a statement about an average into a statement about a limit, and monotonicity is the entire reason the conversion is legitimate. The integrand $(\vartheta(t) - t)/t^2$ has a convergent integral, which controls only the cumulative deviation; the integrand could in principle be large at isolated places. Monotonicity forbids isolation. If ϑ overshoots λx at a single point x , it stays at least λx for the entire interval $[x, \lambda x]$, because it cannot come back down, and the overshoot is therefore spread across a window of multiplicative width λ , contributing a fixed positive amount to the integral. The same window appears in both directions, and its scale invariance, the fact that $c_\lambda^\pm$ depends on λ but not on x , is what lets a single excursion at arbitrarily large x contradict the smallness guaranteed far out by the Cauchy criterion. The choice of weight x^2 in the denominator is exactly what produces this invariance: it makes the substitution $t = ux$ scale the integral to a form free of x , so a deviation at height x costs the same regardless of x . Replace ϑ by a function that is not monotone, and the argument collapses, since a brief spike could be undone immediately and the window of forced deviation would vanish.

With both halves in place, the proof of the Prime Number Theorem is immediate. The function ϑ supplied by arithmetic is nondecreasing, and theorem 5.4.1 furnishes exactly the convergent integral the lemma requires.

Theorem 5.4.3: The Prime Number Theorem

As $x \rightarrow \infty$,

$$\vartheta(x) \sim x, \quad \psi(x) \sim x, \quad \pi(x) \sim \frac{x}{\log x}$$

where ϑ and ψ are the Chebyshev functions (definition 3.1.2) and π is the prime counting function (definition 3.1.1).

Proof:

The Chebyshev theta function $\vartheta(x) = \sum_{p \leq x} \log p$ is nondecreasing in x , since increasing x can only admit further primes into the sum, each contributing a nonnegative term $\log p$. By theorem 5.4.1 the integral $\int_1^\infty (\vartheta(x) - x)/x^2 dx$ converges. The two hypotheses of lemma 5.4.2, monotonicity of ϑ and convergence of this integral, are therefore met, and the lemma yields

$$\vartheta(x) \sim x$$

By the equivalence of the three asymptotic forms (theorem 3.1.5), $\vartheta(x) \sim x$ holds if and only if $\psi(x) \sim x$ and if and only if $\pi(x) \sim x/\log x$. Hence all three hold simultaneously, completing the proof.

This is the theorem the chapter was built to prove, and its history measures the distance the argument has travelled. The asymptotic $\pi(x) \sim x/\log x$ was conjectured by Gauss and Legendre

from numerical tables near the end of the eighteenth century, refined by Gauss into the sharper guess $\pi(x) \sim \text{Li}(x)$, and proved only in 1896, independently by Jacques Hadamard and Charles-Jean de la Vallée Poussin. The century of delay separates the two kinds of information the statement contains. The *order* of $\pi(x)$, that it grows like $x/\log x$ up to bounded factors, is elementary: Chebyshev obtained it in 1852 by comparing the central binomial coefficient against its prime factorization, and that argument, reproduced in theorem 3.2.3, places $\vartheta(x)/x$ between $\log 2$ and $\log 4$ using nothing past unique factorization and Legendre's formula. What the elementary method cannot reach is the *constant*. Chebyshev's corridor has width a factor of two, and no sharpening of the binomial comparison closes it to the single value 1. The constant is analytic in origin. It is the residue of the pole of ζ at $s = 1$, transmitted through the pole of $-\zeta'/\zeta$ and of Φ at $s = 1$, each of residue exactly 1 (proposition 4.3.4), and it reaches the counting function as the value $g(0)$ that Newman's theorem assigns at the boundary. The non-vanishing of ζ on the line $\text{Re}(s) = 1$ is what licenses that boundary value to exist, and it is precisely the input that the elementary theory of chapter 3 could not produce. The Prime Number Theorem is, in this sense, the meeting of two centuries of method: the order from Chebyshev's combinatorics, the constant from Riemann's complex analysis of ζ .

5.4.4 Historical Note: Riemann's Memoir and the Elementary Proof The path proved here is the one Riemann charted in his 1859 memoir, which proposed studying $\pi(x)$ through the analytic continuation of $\zeta(s)$ and the location of its zeros, though Riemann himself proved no asymptotic. Hadamard and de la Vallée Poussin completed the program in 1896 with the non-vanishing of ζ on $\text{Re}(s) = 1$ as the technical core. For half a century afterward it was believed that the analytic input was unavoidable, that no proof could avoid the complex zeros of ζ . This was refuted in 1949 by Atle Selberg and Paul Erdős, who gave an *elementary* proof, in the technical sense of avoiding complex analysis, built on Selberg's symmetry formula $\sum_{p \leq x} (\log p)^2 + \sum_{pq \leq x} (\log p)(\log q) = 2x \log x + O(x)$. The elementary proof is longer and less transparent than the analytic one and yields a weaker error term; the analytic method through zero-free regions, taken up in section 5.5, remains the route to quantitative refinements. Newman's 1980 contour argument [6], in Zagier's exposition [18], is the streamlining of the analytic proof followed in this chapter.

The theorem is best felt against the numbers that first suggested it, where the slow approach of $\pi(x)$ to $x/\log x$ and the closer fit of the logarithmic integral are both visible.

Example 5.4.5: The Prime Number Theorem Numerically

Write $\text{Li}(x) = \int_2^x dt/\log t$ for the logarithmic integral, the approximation to $\pi(x)$ that emerged inside the proof of theorem 3.1.5. Comparing $\pi(x)$ against the two approximations $x/\log x$ and $\text{Li}(x)$ at three widely spaced values of x gives

x	$\pi(x)$	$\frac{x}{\log x}$	$\frac{\pi(x)}{x/\log x}$	$\text{Li}(x)$	$\frac{\pi(x)}{\text{Li}(x)}$
10^3	168	144.8	1.1605	176.6	0.9515
10^6	78498	72382.4	1.0845	78626.5	0.9984
10^9	50847534	48254942.4	1.0537	50849233.9	0.99997

The fourth column, $\pi(x)/(x/\log x)$, is the ratio whose limit the theorem asserts to be 1. It does tend to 1, but slowly: even at $x = 10^9$ it is still 1.05, off by five percent, and the convergence is governed by the next-order term, which is of relative size $1/\log x$, equal to $1/\log 10^9 \approx 0.048$ at $x = 10^9$, matching the observed gap. The logarithmic integral $\text{Li}(x)$ is the better approximation

by a wide margin: the last column $\pi(x)/\text{Li}(x)$ reaches 1 far faster, already agreeing to four decimal places at $x = 10^9$. The reason is that $\text{Li}(x) = x/\log x + x/(\log x)^2 + \dots$ captures the secondary term $x/(\log x)^2$ that $x/\log x$ omits, and it is exactly this secondary term that accounts for the persistent excess of $\pi(x)$ over $x/\log x$ seen in the fourth column. Both approximations are asymptotically correct, $\pi(x)/(x/\log x) \rightarrow 1$ and $\pi(x)/\text{Li}(x) \rightarrow 1$, since $\text{Li}(x) \sim x/\log x$; the theorem is the statement that the ratios converge, and the table is the statement that for $x/\log x$ the convergence is unhurried. A consequence read directly from $\pi(x) \sim x/\log x$, made precise in corollary 5.4.6 below, is that the n -th prime p_n satisfies $p_n \sim n \log n$: at $n = 10^4$ the prime is $p_n = 104729$ against $n \log n \approx 92103$, the ratio 1.137 drifting toward 1 at the same logarithmic rate.

The asymptotic for p_n used at the close of the example is the most frequently quoted reformulation of the theorem, and it follows by inverting $\pi(x) \sim x/\log x$. Two further consequences are worth recording in the same breath: the precise sense in which primes thin out, and the inversion itself.

Corollary 5.4.6: Consequences

Let p_n denote the n -th prime. Then as $n \rightarrow \infty$,

$$p_n \sim n \log n$$

Moreover the density of primes near x is asymptotically $1/\log x$: the probability that a randomly chosen integer in $[1, x]$ is prime, namely $\pi(x)/x$, satisfies $\pi(x)/x \sim 1/\log x$.

Proof:

The density statement is a restatement of the theorem: dividing $\pi(x) \sim x/\log x$ from theorem 5.4.3 by x gives $\pi(x)/x \sim 1/\log x$, which is the asserted density. The substance is the asymptotic for p_n , proved by inverting the relation $\pi(x) \sim x/\log x$.

Step 1: A logarithmic comparison. Set $x = p_n$ in the theorem. By definition $\pi(p_n) = n$, the count of primes up to the n -th prime, so theorem 5.4.3 gives

$$n = \pi(p_n) \sim \frac{p_n}{\log p_n}, \quad \text{that is} \quad \frac{n \log p_n}{p_n} \rightarrow 1 \quad (5.6)$$

Taking logarithms of $n \sim p_n/\log p_n$, and using that $\log$ of a quantity tending to 1 tends to 0,

$$\log n = \log p_n - \log \log p_n + o(1)$$

Since $p_n \rightarrow \infty$, both $\log p_n \rightarrow \infty$ and $\log \log p_n \rightarrow \infty$, but the second grows far slower; dividing the last display by $\log p_n$,

$$\frac{\log n}{\log p_n} = 1 - \frac{\log \log p_n}{\log p_n} + \frac{o(1)}{\log p_n} \rightarrow 1$$

because $\log \log p_n / \log p_n \rightarrow 0$. Hence $\log n \sim \log p_n$.

Step 2: Inverting. Rewrite (5.6) as $p_n \sim n \log p_n$. Substituting the equivalence $\log p_n \sim \log n$ from Step 1,

$$p_n \sim n \log p_n \sim n \log n$$

where the replacement of $\log p_n$ by $\log n$ is legitimate because the two are asymptotically equal and n is a common factor: $p_n/(n \log n) = (p_n/(n \log p_n)) \cdot (\log p_n/\log n) \rightarrow 1 \cdot 1 = 1$. Therefore $p_n \sim n \log n$, as we sought to show.

The corollary translates the theorem into the language of the primes as a sequence. The statement $p_n \sim n \log n$ says that the n -th prime sits near $n \log n$, so the primes, though individually unpredictable, are on average spaced $\log n \approx \log p_n$ apart at height p_n : the gap between consecutive primes near x is, on average, about $\log x$. This is the same fact as the density $1/\log x$ viewed from the dual side, counting the spacing between primes rather than their frequency, and the two are reciprocal as one expects. The average gap $\log x$ grows without bound, so the primes do thin out, but only logarithmically, far more slowly than the squares or the powers of 2; between x and $2x$ there are still about $x/\log x$ primes for every large x . The estimate $p_n \sim n \log n$ is the crudest term of a longer expansion, $p_n = n \log n + n \log \log n - n + o(n)$, whose secondary terms come from the same logarithmic-integral correction that made $\text{Li}(x)$ outperform $x/\log x$ in example 5.4.5; the leading term alone is the direct content of the Prime Number Theorem.

5.5 Error Terms and the Zero-Free Region

The proof of the Prime Number Theorem completed in section 5.4 rests on a single qualitative fact: $\zeta(s)$ has no zeros on the line $\text{Re}(s) = 1$ (theorem 4.4.2). That fact is exactly enough to force $\psi(x) \sim x$, equivalently $\vartheta(x) \sim x$ and $\pi(x) \sim x/\log x$ (theorem 5.4.3), but it is silent on the rate of convergence. Non-vanishing on the line says the ratio $\psi(x)/x$ tends to 1; it does not say how fast, and the Tauberian route of Newman's theorem, which trades all quantitative information for a clean limit, gives no error term at all. To measure the speed one needs a quantitative strengthening: not just that ζ avoids the line, but that it avoids an open region to the *left* of the line, a region whose width near a given height controls how far the contour of integration can be pushed past $\text{Re}(s) = 1$ and hence how small the resulting error can be made.

This section proves the de la Vallée Poussin zero-free region, the first such quantitative result, in full: there is a constant $c > 0$ with $\zeta(s) \neq 0$ throughout $\sigma \geq 1 - c/\log(|t| + 2)$. The argument is the quantitative form of the positivity inequality $3 + 4\cos\theta + \cos 2\theta \geq 0$ that proved qualitative non-vanishing in chapter 4, sharpened by a growth bound for ζ near the line. The width of this region then dictates an error term, $\psi(x) = x + O(x \exp(-c\sqrt{\log x}))$, whose derivation by contour optimization is stated with a precise reference. The section closes with the Riemann–von Mangoldt explicit formula, the Riemann Hypothesis, and the modern Korobov–Vinogradov region, each placing the elementary error term in the larger landscape that the functional equation of the next chapter opens up.

The proof of the zero-free region needs one analytic input beyond the structural facts of chapter 4: a bound on the size of ζ and its derivative just to the left of $\text{Re}(s) = 1$. The continuation formula of proposition 1.4.6 supplies it directly.

Lemma 5.5.1: A Growth Bound for ζ near the Line

There is a constant $A > 0$ such that

$$|\zeta(s)| \leq A \log |t| \quad \text{and} \quad |\zeta'(s)| \leq A (\log |t|)^2$$

uniformly for all $s = \sigma + it$ with $\sigma \geq 1 - \frac{1}{\log |t|}$ and $|t| \geq 5$.

Proof:

By the reflection $\zeta(\bar{s}) = \overline{\zeta(s)}$, which holds because ζ is real on the real axis and analytic, both moduli are symmetric under $t \mapsto -t$, so it suffices to treat $t \geq 5$. Write $\sigma = \operatorname{Re}(s)$ throughout. The threshold $t \geq 5$ gives $\log t \geq \log 5 > \frac{3}{2}$, which keeps every point used below inside the half-plane $\operatorname{Re}(s) > 0$ where the continuation formula of proposition 1.4.6 is valid: the lower edge of the region satisfies $1 - 1/\log t \geq 1 - 1/\log 5 > 0$, and the Cauchy circle of Step 3 stays in $\operatorname{Re}(w) > 0$ as well.

Step 1: A truncated continuation formula. proposition 1.4.6 gives, for $\operatorname{Re}(s) > 0$ and $s \neq 1$ and every real $x \geq 1$,

$$\sum_{n \leq x} \frac{1}{n^s} = \frac{x^{1-s}}{1-s} + \zeta(s) + O(x^{-\sigma})$$

with an absolute implied constant. Solving for $\zeta(s)$,

$$\zeta(s) = \sum_{n \leq x} \frac{1}{n^s} - \frac{x^{1-s}}{1-s} + O(x^{-\sigma}) \quad (5.7)$$

The estimate of the three pieces will be carried out at the choice $x = t$, the truncation height matched to the imaginary part.

Step 2: The bound for $|\zeta(s)|$. The argument is carried out uniformly in a width parameter: for each fixed $\kappa \geq 1$ there is a constant $A_0(\kappa)$ with $|\zeta(s)| \leq A_0(\kappa) \log t$ for all $s = \sigma + it$ with $\sigma \geq 1 - \kappa/\log t$, $\sigma > 0$, and $t \geq 2$. The positivity constraint $\sigma > 0$ is what makes the continuation formula available; in the stated region of the lemma ($\kappa = 1$, $t \geq 5$) it holds automatically since $1 - 1/\log t > 0$ there, while in the $\kappa = 3$ application of Step 3 it is checked directly at each point. Fix $\kappa \geq 1$ and such an s , and take $x = t$ in (5.7); the continuation formula applies since $\operatorname{Re}(s) = \sigma > 0$ and $s \neq 1$. First note that for these σ ,

$$t^{1-\sigma} = \exp((1-\sigma) \log t) \leq \exp\left(\frac{\kappa}{\log t} \cdot \log t\right) = e^\kappa$$

since $1 - \sigma \leq \kappa/\log t$; this bound on $t^{1-\sigma}$ uses only the region constraint, not the sign of σ . The three pieces are estimated in turn.

For the Dirichlet sum, write $|n^{-s}| = n^{-\sigma} = n^{-1} n^{1-\sigma}$ and bound the factor $n^{1-\sigma}$ for $1 \leq n \leq x$ by $\max(x^{1-\sigma}, 1)$. For $\sigma \leq 1$ the exponent $1 - \sigma$ is nonnegative, so $n^{1-\sigma} \leq x^{1-\sigma}$; for $\sigma > 1$ it is negative, so $n^{1-\sigma} \leq 1$. In both cases $n^{1-\sigma} \leq \max(x^{1-\sigma}, 1) \leq e^\kappa$, using $x^{1-\sigma} = t^{1-\sigma} \leq e^\kappa$ and

$1 \leq e^\kappa$. Hence $n^{-\sigma} \leq n^{-1}e^\kappa$ for $n \leq x$, and

$$\left| \sum_{n \leq x} \frac{1}{n^s} \right| \leq e^\kappa \sum_{n \leq x} \frac{1}{n} \leq e^\kappa (1 + \log x) = e^\kappa (1 + \log t)$$

the harmonic sum bounded by $1 + \log x$ through comparison with $\int_1^x du/u$.

For the rational term, $|x^{1-s}/(1-s)| = x^{1-\sigma}/|1-s|$, and $|1-s| = |(1-\sigma) - it| \geq |t| = t \geq 2$, so

$$\left| \frac{x^{1-s}}{1-s} \right| = \frac{t^{1-\sigma}}{|1-s|} \leq \frac{e^\kappa}{t} \leq \frac{e^\kappa}{2}$$

For the error term, $x^{-\sigma} = t^{-\sigma} = t^{-1}t^{1-\sigma} \leq t^{-1}e^\kappa \leq e^\kappa/2$. Summing the three contributions,

$$|\zeta(s)| \leq e^\kappa(1 + \log t) + \frac{e^\kappa}{2} + O\left(\frac{e^\kappa}{2}\right) \leq A_0(\kappa) \log t$$

for a suitable constant $A_0(\kappa)$ and all $t \geq 2$, since $\log t$ dominates the constants once $t \geq 2$ and $1 + \log t \leq (1 + 1/\log 2) \log t$ there. This is the asserted bound on $|\zeta|$, valid on $\sigma \geq 1 - \kappa/\log t$ with $\sigma > 0$.

Step 3: The bound for $|\zeta'(s)|$ by a Cauchy estimate. Fix $s = \sigma + it$ with $\sigma \geq 1 - 1/\log t$ and $t \geq 5$, and let Γ be the positively oriented circle of radius $r = 1/(2 \log t)$ centered at s . Since $t \geq 5$ gives $\log t \geq \log 5$, the radius satisfies $r \leq 1/(2 \log 5) < 1$. Each point $w = u + iv$ on Γ satisfies

$$u \geq \sigma - r \geq 1 - \frac{1}{\log t} - \frac{1}{2 \log t} = 1 - \frac{3}{2 \log t}, \quad |v - t| \leq r < 1$$

so $|v| \geq t - 1 \geq 4$ and $|v| \leq t + 1 \leq 2t$, the last because $1 \leq t$. The lower bound on u is positive: $\log t \geq \log 5 > \frac{3}{2}$ gives $1 - 3/(2 \log t) > 0$, so $\operatorname{Re}(w) = u > 0$ and w lies in the half-plane where ζ is defined and the bound of Step 2 is available. From $|v| \leq 2t$ and $t \geq 5$ one gets $\log |v| \leq \log(2t) = \log 2 + \log t \leq 2 \log t$, hence $1/\log t \leq 2/\log |v|$, and therefore

$$u \geq 1 - \frac{3}{2 \log t} \geq 1 - \frac{3}{\log |v|}$$

so w lies in the region of Step 2 with $\kappa = 3$, imaginary part v satisfying $|v| \geq 4 \geq 2$, and $\operatorname{Re}(w) = u > 0$. Step 2 with $\kappa = 3$ therefore applies to every $w \in \Gamma$, giving $|\zeta(w)| \leq A_0(3) \log |v| \leq 2A_0(3) \log t$, the last step using $\log |v| \leq 2 \log t$. Thus $|\zeta(w)| \leq A_1 \log t$ for an absolute constant $A_1 = 2A_0(3)$ and every $w \in \Gamma$. The Cauchy integral formula for the derivative ([16]) gives

$$\zeta'(s) = \frac{1}{2\pi i} \int_{\Gamma} \frac{\zeta(w)}{(w-s)^2} dw$$

the contour Γ lying in $\operatorname{Re}(w) > 0$, $w \neq 1$, where ζ is holomorphic. On Γ one has $|w - s| = r$, so $|(w - s)^{-2}| = r^{-2}$, and the length of Γ is $2\pi r$; the standard length estimate yields

$$|\zeta'(s)| \leq \frac{1}{2\pi} \cdot \frac{\max_{w \in \Gamma} |\zeta(w)|}{r^2} \cdot 2\pi r = \frac{1}{r} \max_{w \in \Gamma} |\zeta(w)| \leq (2 \log t) \cdot A_1 \log t = 2A_1 (\log t)^2$$

Taking $A = \max(A_0(1), 2A_1)$ gives both bounds simultaneously on the stated region, completing the proof.

The bound is crude in its dependence on σ , holding uniformly across the thin strip $1 - 1/\log |t| \leq \sigma$ with no improvement as σ increases past 1, but its dependence on t is what matters: ζ grows no faster than a power of $\log |t|$ near the line. This polynomial-in-log growth is the quantitative substitute for the boundedness that holds deeper inside the half-plane $\operatorname{Re}(s) > 1$, and it is the only size information the zero-free argument consumes. The choice $x = |t|$ in the proof is the balance point: a larger truncation would shrink the error term $O(x^{-\sigma})$ but inflate the Dirichlet sum, and a smaller one would do the reverse, so matching the truncation height to the imaginary part is what produces the single factor of $\log |t|$ rather than a power of $|t|$. The derivative bound costs one further factor of $\log |t|$, the price of the radius $r \asymp 1/\log |t|$ in the Cauchy estimate, the largest scale of radius on which the growth bound for ζ still applies.

Example 5.5.2: The Growth Bound at $s = 1 + it$

On the line itself, $\sigma = 1$, the lemma reads $|\zeta(1 + it)| = O(\log |t|)$ and $|\zeta'(1 + it)| = O((\log |t|)^2)$. The first is the familiar statement that ζ grows slowly along the line $\operatorname{Re}(s) = 1$. At the concrete height $t = 100$ the bound gives $|\zeta(1 + 100i)| \leq A \log 100 = A \cdot 4.605 \dots$, while the true value is $|\zeta(1 + 100i)| \approx 1.63$, comfortably inside the bound for any reasonable constant; the gap reflects the crudeness of the constant A , not of the $\log |t|$ shape. The logarithmic growth is genuine, not an artifact: the partial sums $\sum_{n \leq |t|} n^{-1-it}$ that dominate (5.7) have modulus of order $\log |t|$ for typical t , since the phases $n^{-it} = e^{-it \log n}$ fail to produce full cancellation in the harmonic-weighted sum. The same computation with the second derivative, a Cauchy estimate on a circle of radius $\asymp 1/\log |t|$ applied to the bound for ζ' , would give $|\zeta''(1 + it)| = O((\log |t|)^3)$, each derivative costing one additional logarithm.

The growth bound is the missing quantitative ingredient. The qualitative non-vanishing theorem of chapter 4 ruled out zeros *on* the line by showing that a zero there would drive the positive combination $|\zeta(\sigma)|^3 |\zeta(\sigma + it)|^4 |\zeta(\sigma + 2it)|$ below 1 as $\sigma \rightarrow 1^+$. The quantitative version measures how far a zero must sit from the line for the same combination to remain admissible, and the distance it extracts is exactly $c/\log(|t| + 2)$.

Theorem 5.5.3: The de la Vallée Poussin Zero-Free Region

There is a constant $c > 0$ such that $\zeta(s) \neq 0$ throughout the region

$$\sigma \geq 1 - \frac{c}{\log(|t| + 2)}$$

In particular ζ has no zeros in an open neighborhood of the line $\operatorname{Re}(s) = 1$ whose width at height t shrinks like $1/\log |t|$.

Proof:

The half-plane $\operatorname{Re}(s) > 1$ contains no zeros, by the Euler product (corollary 4.1.2), and a finite portion of the line is handled by continuity, so it suffices to find $c > 0$ such that no zero $\rho = \beta + i\gamma$ has

$$\beta > 1 - \frac{c}{\log(|\gamma| + 2)}$$

for $|\gamma|$ large. By the symmetry $\zeta(\bar{s}) = \overline{\zeta(s)}$, zeros come in conjugate pairs, so it is enough to treat $\gamma \geq \gamma_0$ for a fixed $\gamma_0 \geq 7$, the range $0 \leq \gamma < \gamma_0$ being a compact piece of the closed half-plane $\operatorname{Re}(s) \geq 1$ on which ζ is nonzero (no zeros on $\operatorname{Re}(s) = 1$ by theorem 4.4.2, none in $\operatorname{Re}(s) > 1$)

and hence bounded away from zero on a strip $\sigma \geq 1 - c'$ by continuity and compactness. Fix a zero $\rho = \beta + i\gamma$ with $\gamma \geq \gamma_0$ and write $L = \log(\gamma + 2)$.

Step 1: The positivity inequality. For real $\sigma > 1$ the von Mangoldt series of proposition 4.3.1 converges absolutely, and taking real parts of $-\zeta'(s)/\zeta(s) = \sum_n \Lambda(n)n^{-s}$ at $s = \sigma + i\tau$ gives, since $\Lambda(n) \geq 0$ and $|n^{-i\tau}| = 1$ with $\operatorname{Re}(n^{-i\tau}) = \cos(\tau \log n)$,

$$\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\tau)\right) = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^{\sigma}} \cos(\tau \log n)$$

Form the combination with weights 3, 4, 1 at the heights $\tau = 0, \gamma, 2\gamma$:

$$3\left(-\frac{\zeta'}{\zeta}(\sigma)\right) + 4\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\gamma)\right) + \operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + 2i\gamma)\right) = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^{\sigma}} (3 + 4\cos\theta_n + \cos 2\theta_n)$$

with $\theta_n = \gamma \log n$. By lemma 4.4.1 every bracket $3 + 4\cos\theta_n + \cos 2\theta_n = 2(1 + \cos\theta_n)^2 \geq 0$, and the coefficients $\Lambda(n)n^{-\sigma}$ are nonnegative, so the entire sum is nonnegative:

$$3\left(-\frac{\zeta'}{\zeta}(\sigma)\right) + 4\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\gamma)\right) + \operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + 2i\gamma)\right) \geq 0 \quad (5.8)$$

This is the quantitative engine; the three terms are now bounded above individually, the first from the pole and the latter two from the growth bound, and the inequality is read backward to constrain β .

Step 2: The pole term. By proposition 4.3.1 the function $-\zeta'/\zeta$ has near $s = 1$ the form $-\zeta'(s)/\zeta(s) = 1/(s-1) - h(s)$ with h holomorphic in a fixed neighborhood of $s = 1$, hence bounded on the segment $1 < \sigma \leq 2$. Therefore there is a constant A_2 with

$$-\frac{\zeta'}{\zeta}(\sigma) \leq \frac{1}{\sigma-1} + A_2 \quad (1 < \sigma \leq 2) \quad (5.9)$$

the inequality being equality up to the bounded term $-h(\sigma)$, which contributes at most a constant.

Step 3: The off-line terms from the growth bound. The two remaining terms are controlled by a standard bound on the logarithmic derivative of a function of moderate growth, applied to ζ . The bound states that there is a constant $A_3 > 0$ such that for $1 < \sigma \leq 2$ and any height τ with $|\tau| \geq \gamma_0$,

$$\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\tau)\right) \leq A_3 \log(|\tau| + 2) - \sum_{\rho'} \operatorname{Re} \frac{1}{\sigma + i\tau - \rho'} \quad (5.10)$$

where the sum runs over the zeros ρ' of ζ with $|\operatorname{Im}\rho' - \tau| \leq 1$, and every term of that sum is nonnegative because $\operatorname{Re}(\sigma + i\tau - \rho') = \sigma - \beta' > 1 - 1 = 0$ for a zero $\rho' = \beta' + i\gamma'$ (which has $\beta' \leq 1$). The inequality (5.10) is the partial-fraction estimate for $-\zeta'/\zeta$ obtained from the Borel–Carathéodory theorem and Jensen’s formula applied to ζ on the disc of radius slightly larger than 1 centered at $2 + i\tau$: Borel–Carathéodory converts the upper bound $|\zeta(s)| \leq A \log |t|$ of lemma 5.5.1 on the boundary circle into a bound on ζ'/ζ in the interior, and Jensen’s formula identifies the zeros ρ' as the only obstruction, contributing the displayed negative partial-fraction

terms; the derivation is standard and is carried out in full in [1, Ch. 12] and [4, Ch. 6], using the complex-analytic tools of [16] (the Borel–Carathéodory theorem and Jensen’s formula), with $\log(|\tau| + 2)$ entering as the logarithm of the growth bound’s M . Two applications of (5.10) are needed.

At height $\tau = 2\gamma$, drop the entire nonnegative sum to obtain the clean upper bound

$$\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + 2i\gamma)\right) \leq A_3 \log(2\gamma + 2) \leq A_4 L \quad (5.11)$$

for a constant A_4 , using $\log(2\gamma + 2) \leq 2\log(\gamma + 2) = 2L$ for $\gamma \geq 1$.

At height $\tau = \gamma$, the chosen zero $\rho = \beta + i\gamma$ has $\operatorname{Im}\rho = \gamma = \tau$, so $|\operatorname{Im}\rho - \tau| = 0 \leq 1$ and ρ appears in the sum of (5.10). Keep only its term and drop the rest, all nonnegative:

$$\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\gamma)\right) \leq A_3 \log(\gamma + 2) - \operatorname{Re}\frac{1}{\sigma + i\gamma - \rho} = A_3 L - \operatorname{Re}\frac{1}{\sigma - \beta}$$

since $\sigma + i\gamma - \rho = \sigma + i\gamma - (\beta + i\gamma) = \sigma - \beta$ is real and positive. Thus

$$\operatorname{Re}\left(-\frac{\zeta'}{\zeta}(\sigma + i\gamma)\right) \leq A_3 L - \frac{1}{\sigma - \beta} \quad (5.12)$$

the term $-1/(\sigma - \beta)$ recording the proximity of $\sigma + i\gamma$ to the zero.

Step 4: Combining the bounds. Substitute (5.9), (5.12), and (5.11) into the positivity inequality (5.8):

$$0 \leq 3\left(\frac{1}{\sigma - 1} + A_2\right) + 4\left(A_3 L - \frac{1}{\sigma - \beta}\right) + A_4 L$$

Collecting the logarithmic terms into a single constant times L , there is a constant $A_5 = 3A_2/L + 4A_3 + A_4 \leq A_6$ (bounded since $L \geq \log(\gamma_0 + 2) \geq 1$) with

$$\frac{4}{\sigma - \beta} \leq \frac{3}{\sigma - 1} + A_6 L \quad (5.13)$$

valid for every $\sigma \in (1, 2]$. This single inequality, true for all such σ , will be made to contradict β being too close to 1 by an appropriate choice of σ .

Step 5: Optimizing σ . Choose

$$\sigma = 1 + \frac{\delta}{L}, \quad \delta > 0 \text{ a constant to be fixed,}$$

which lies in $(1, 2]$ once $\delta \leq L$, true for large γ since $L \rightarrow \infty$. Then $\sigma - 1 = \delta/L$, so $3/(\sigma - 1) = 3L/\delta$, and (5.13) becomes

$$\frac{4}{\sigma - \beta} \leq \frac{3L}{\delta} + A_6 L = \left(\frac{3}{\delta} + A_6\right) L$$

Inverting,

$$\sigma - \beta \geq \frac{4}{(3/\delta + A_6)L} = \frac{4\delta}{(3 + A_6\delta)L}$$

Now $\beta = \sigma - (\sigma - \beta) \leq \sigma - \frac{4\delta}{(3 + A_6\delta)L} = 1 + \frac{\delta}{L} - \frac{4\delta}{(3 + A_6\delta)L}$, that is,

$$\beta \leq 1 - \frac{1}{L} \left(\frac{4\delta}{3 + A_6\delta} - \delta \right) = 1 - \frac{\delta}{L} \cdot \frac{4 - (3 + A_6\delta)}{3 + A_6\delta} = 1 - \frac{\delta}{L} \cdot \frac{1 - A_6\delta}{3 + A_6\delta}$$

The bracketed factor $(1 - A_6\delta)/(3 + A_6\delta)$ is positive precisely when $\delta < 1/A_6$. Fix any such δ , for instance $\delta = 1/(2A_6)$, giving the positive constant

$$c = \frac{\delta(1 - A_6\delta)}{3 + A_6\delta} = \frac{1}{2A_6} \cdot \frac{1/2}{3 + 1/2} = \frac{1}{14A_6} > 0$$

With this δ ,

$$\beta \leq 1 - \frac{c}{L} = 1 - \frac{c}{\log(\gamma + 2)}$$

for every zero $\rho = \beta + i\gamma$ with $\gamma \geq \gamma_0$. Shrinking c if necessary to also cover the compact range $0 \leq \gamma < \gamma_0$, where a fixed zero-free strip $\sigma \geq 1 - c'$ exists by the non-vanishing on the line, the bound $\beta \leq 1 - c/\log(|\gamma| + 2)$ holds for all zeros, which is the assertion that $\zeta(s) \neq 0$ for $\sigma > 1 - c/\log(|t| + 2)$, completing the proof.

The proof is the qualitative non-vanishing argument run with a ruler. In chapter 4 the inequality (5.8) was used only to exclude $\beta = 1$: a zero on the line would make $-4/(\sigma - 1)$ blow up faster than $+3/(\sigma - 1)$ as $\sigma \rightarrow 1^+$, breaking nonnegativity. Here the same competition between the $3/(\sigma - 1)$ from the pole and the $4/(\sigma - \beta)$ from the zero is resolved quantitatively, and the factor $4 > 3$ is what does the work: because the zero is weighted by 4 and the pole by 3, the zero cannot approach the pole's line faster than a definite rate without the weighted sum turning negative. The growth bound enters only through the constant A_3L in (5.10), a term of size $\log \gamma$; it sets the scale $1/L = 1/\log(\gamma + 2)$ of the forbidden region, since balancing the constant against the pole's $1/(\sigma - 1)$ forces the optimal offset $\sigma - 1 \asymp 1/L$, and the zero is then trapped at distance $\asymp 1/L$ from the line. The shape $c/\log(|t| + 2)$ is thus the direct fingerprint of the logarithmic growth of ζ : a slower-growing ζ would widen the region, and the conjectural bound $|\zeta(1 + it)| = O((\log |t|)^\varepsilon)$ would not, by this argument alone, change the log in the denominator, because the method's loss is in the constant, not the shape.

The width of the zero-free region translates directly into an error term for $\psi(x)$. The mechanism is the truncated Perron formula: the contour can be pushed left to the boundary of the region, and the further left it goes, the smaller the factor x^σ on the shifted line, hence the smaller the error.

Theorem 5.5.4: The Prime Number Theorem with Error Term

There is a constant $c > 0$ such that

$$\psi(x) = x + O\left(x \exp(-c\sqrt{\log x})\right)$$

and correspondingly, with $\text{Li}(x) = \int_2^x dt/\log t$,

$$\pi(x) = \text{Li}(x) + O\left(x \exp(-c'\sqrt{\log x})\right)$$

for a constant $c' > 0$.

The full proof is a contour-optimization computation: the truncation height and the depth of the contour shift must be chosen as coupled functions of x , and tracking the resulting error through the curved boundary of the zero-free region is several pages of estimation with no conceptual turning point beyond the choices themselves. The argument is given in full in [1, Ch. 18] and [4, Ch. 6]; only the structure and the origin of the term $\exp(-c\sqrt{\log x})$ are recorded here.

Proof Key ideas:

Apply the truncated Perron formula (theorem 2.5.3) to the von Mangoldt coefficients $a_n = \Lambda(n)$, whose Dirichlet series is $-\zeta'/\zeta$ (example 2.5.4), at a height T to be chosen:

$$\psi(x) = \frac{1}{2\pi i} \int_{c_0-iT}^{c_0+iT} \left(-\frac{\zeta'}{\zeta}(s)\right) \frac{x^s}{s} ds + R(x, T)$$

with $c_0 = 1 + 1/\log x$ and an explicit error $R(x, T)$ from the truncation. *Step 1: shift the contour.* Move the segment of integration left to a contour hugging the boundary $\sigma = 1 - c/\log(|t| + 2)$ of the zero-free region of theorem 5.5.3, closing it off with horizontal segments at $\pm T$. The shift is legitimate because $-\zeta'/\zeta$ is holomorphic in the region swept, save for the single pole at $s = 1$; the residue of $(-\zeta'/\zeta)(s) x^s/s$ there is x , by the residue-1 pole of proposition 4.3.1, and this residue is the main term. *Step 2: bound the shifted integrand.* On and near the boundary contour the size of $-\zeta'/\zeta$ is controlled by the growth bound lemma 5.5.1, which gives $-\zeta'/\zeta = O((\log |t|)^2)$ there, while the factor x^s contributes $x^\sigma = x \cdot x^{-c/\log(|t|+2)} = x \exp(-c \log x / \log(|t|+2))$, exponentially smaller than x by an amount that worsens as $|t|$ grows. *Step 3: optimize T .* The truncation error $R(x, T)$ decreases as T grows, while the contribution of the shifted contour increases with T through the $\log(|t| + 2)$ in the exponent; balancing the two competing errors fixes the optimal height. Setting $\log T \asymp \sqrt{\log x}$ equalizes them, because the dominant saving $\exp(-c \log x / \log T)$ on the contour and the dominant loss from the truncation both become $\exp(-c''\sqrt{\log x})$ at that height. The common value $\exp(-c''\sqrt{\log x})$, times x , is the error term. *Step 4: pass to $\pi(x)$.* The estimate for ψ transfers to ϑ by removing the prime-power contribution (of size $O(\sqrt{x} \log x)$, absorbed into the error), and to $\pi(x)$ by partial summation against $1/\log t$ (theorem 1.4.1), which converts the main term x into $\text{Li}(x) = \int_2^x dt/\log t$ and preserves an error of the same exponential shape with a possibly smaller constant c' . The full execution of these four steps is carried out in [1, Ch. 18] and [4, Ch. 6], completing the proof.

The error term $x \exp(-c\sqrt{\log x})$ is smaller than $x/(\log x)^N$ for every fixed N , since $\exp(-c\sqrt{\log x})$ decays faster than any negative power of $\log x$, yet it is far larger than the power saving $x^{1-\delta}$ that a zero-free region of *fixed* width δ to the left of the line would yield. The square root in the exponent is

the exact arithmetic of the optimization: the contour saving is $\exp(-c \log x / \log T)$ and the truncation loss grows with $\log T$, so the two balance when $(\log x) / \log T \asymp \log T$, that is, $\log T \asymp \sqrt{\log x}$, making each error $\exp(-c\sqrt{\log x})$. A wider zero-free region, one whose boundary recedes from the line more slowly in $|t|$, would permit a deeper contour shift at each height and hence a larger exponent; the shape of the region and the shape of the error term are two readings of the same data. The constant c here is not the same as the constant in the zero-free region, but it is determined by it, and any improvement in the width of the region propagates to an improvement in the error.

The error term and the zero-free region are two faces of a single object, made precise by the explicit formula, which expresses the deviation $\psi(x) - x$ as a sum over the zeros of ζ .

5.5.5 Note: The Explicit Formula The deviation of ψ from its main term is given exactly, not just asymptotically, by the *Riemann-von Mangoldt explicit formula*: for $x > 1$ not a prime power,

$$\psi_0(x) = x - \sum_{\rho} \frac{x^{\rho}}{\rho} - \log(2\pi) - \frac{1}{2} \log\left(1 - \frac{1}{x^2}\right)$$

where $\psi_0(x) = \frac{1}{2}(\psi(x^-) + \psi(x^+))$ is the normalized Chebyshev function and the sum runs over the nontrivial zeros ρ of ζ , taken in conjugate pairs and in order of increasing $|\operatorname{Im}\rho|$. The main term x is the residue at the pole $s = 1$; each zero $\rho = \beta + i\gamma$ contributes an oscillation $-x^{\rho}/\rho$ of size $x^{\beta}/|\rho|$; the constant $-\log(2\pi)$ is the value at $s = 0$ from the kernel; and the final term collects the trivial zeros at $s = -2, -4, \dots$. Its proof lies beyond the present chapter: it requires the meromorphic continuation of ζ to all of $\mathbb{C}$ through the functional equation (chapter 6) and the Hadamard product factorization of the entire function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$, which expresses ξ as a product over its zeros and thereby converts the contour integral of $-\zeta'/\zeta$ into the sum over ρ . The full derivation is given in [1, Ch. 17]; see also [4]. The formula foreshadows the role of the functional equation in chapter 6: every term but the first is governed by the symmetry $s \leftrightarrow 1-s$ and the zeros it organizes, and the error term of theorem 5.5.4 is precisely a bound on the zero-sum $\sum_{\rho} x^{\rho}/\rho$ obtained by knowing that all $\beta < 1$ stay a definite distance below 1.

The explicit formula reframes every statement about the error term as a statement about the location of the zeros. The size of $\sum_{\rho} x^{\rho}/\rho$ is governed by $\sup_{\rho} \beta = \sup_{\rho} \operatorname{Re}(\rho)$: the larger the real parts of the zeros, the larger the oscillations x^{β} , and the worse the error. The de la Vallée Poussin region bounds β below 1 by a margin that narrows with height, and the resulting error $x \exp(-c\sqrt{\log x})$ is what that margin yields. The optimal possible control corresponds to the zeros lying as far left as they can, on the line $\operatorname{Re}(s) = 1/2$, which is the content of the Riemann Hypothesis.

Conjecture 5.5.6: The Riemann Hypothesis

Every nontrivial zero ρ of the Riemann zeta function satisfies $\operatorname{Re}(\rho) = \frac{1}{2}$.

The hypothesis is equivalent to the sharpest possible error term in the Prime Number Theorem. If every nontrivial zero has $\beta = \frac{1}{2}$, then each term x^{ρ}/ρ in the explicit formula has size $x^{1/2}/|\rho|$, and summing over zeros with the known density produces the bound below; conversely, that bound forces the zeros onto the line, so the equivalence runs both ways.

Proposition 5.5.7: Riemann Hypothesis and the Error Term

The Riemann Hypothesis holds if and only if

$$\psi(x) = x + O(\sqrt{x} (\log x)^2)$$

and this error term is essentially optimal: the deviation $\psi(x) - x$ is not $o(\sqrt{x})$, so the exponent $\frac{1}{2}$ cannot be lowered.

The equivalence is proved in [4, Ch. 13]; the forward direction reads the bound off the explicit formula under $\beta \equiv \frac{1}{2}$, and the reverse recovers $\sup \beta \leq \frac{1}{2}$ from the bound by a contour argument, while the optimality is a consequence of the oscillation forced by the infinitely many zeros on the line. Unconditionally, the best known zero-free region is wider than de la Vallée Poussin's but still far from the half-plane $\operatorname{Re}(s) > \frac{1}{2}$ that the Riemann Hypothesis asserts: the *Korobov–Vinogradov* region states that $\zeta(s) \neq 0$ for

$$\sigma \geq 1 - \frac{c}{(\log |t|)^{2/3} (\log \log |t|)^{1/3}}$$

established by Vinogradov's method for estimating exponential sums and proved in full in [11, Ch. 6]. By the same contour optimization that produced theorem 5.5.4 from de la Vallée Poussin's region, this wider region yields the best unconditional error term $\psi(x) = x + O(x \exp(-c(\log x)^{3/5} (\log \log x)^{-1/5}))$.

The progression from de la Vallée Poussin to Korobov–Vinogradov to the Riemann Hypothesis is a progression in the width of a zero-free region and, by the explicit formula, in the size of an error term. De la Vallée Poussin's region has width $\asymp 1/\log |t|$ and gives the error $x \exp(-c\sqrt{\log x})$; Korobov–Vinogradov widens the region near the real axis by a power of $\log \log |t|$ and improves the exponent from $\frac{1}{2}$ to $\frac{3}{5}$; the Riemann Hypothesis would push the region all the way to the open half-plane $\operatorname{Re}(s) > \frac{1}{2}$, the widest a zero-free region can be, since zeros are known to lie on $\operatorname{Re}(s) = \frac{1}{2}$ and so the region cannot extend past it, and would replace the exponential saving by the square-root power $x^{1/2}$. Each step trades a better understanding of where the zeros are not for a sharper count of the primes, and the line $\operatorname{Re}(s) = \frac{1}{2}$ marks the limit of what any zero-free region can achieve. The functional equation of the next chapter is what brings that line into view, by exhibiting the symmetry $s \leftrightarrow 1 - s$ under which the critical line $\operatorname{Re}(s) = \frac{1}{2}$ is the axis, and by supplying the continuation and the product factorization that the explicit formula and the Riemann Hypothesis both presuppose.

6

The Functional Equation and the Critical Strip

The previous two chapters extended $\zeta(s)$ only as far as the half-plane $\operatorname{Re}(s) > 0$ and drew the Prime Number Theorem out of its behavior near the line $\operatorname{Re}(s) = 1$: the simple pole at $s = 1$ (theorem 4.2.1) supplied the main term $\psi(x) \sim x$, and the non-vanishing on the line $\operatorname{Re}(s) = 1$ (theorem 4.4.2) supplied the convergence that turned that main term into theorem 5.4.3. None of it touched the region $\operatorname{Re}(s) \leq 0$, where the defining series diverges and the integral of (4.1) fails to converge. This chapter continues $\zeta(s)$ to the entire complex plane and uncovers the symmetry that governs the continuation: the *functional equation*, a single identity relating $\zeta(s)$ to $\zeta(1-s)$. The mechanism is a pairing of two transcendental functions, the gamma function $\Gamma(s)$ and the Jacobi theta function $\theta(t)$, the first supplying the archimedean factor that completes ζ into a symmetric object and the second supplying, through its own transformation law under $t \mapsto 1/t$, the reflection $s \leftrightarrow 1-s$ itself.

The consequences are immediate once the symmetry is in hand. The poles of Γ at the non-positive integers, transferred through the completed zeta function of section 6.3, force ζ to vanish at the negative even integers: these are the *trivial zeros*. The same machinery discharges the foreshadowed evaluations of example 4.1.3, expressing every even value $\zeta(2k)$ as a rational multiple of π^{2k} and giving in particular the value $\zeta(2) = \pi^2/6$ promised there. What the symmetry leaves undetermined is the location of the remaining zeros, the nontrivial ones, all of which lie in the *critical strip* $0 \leq \operatorname{Re}(s) \leq 1$; the assertion that they lie on the central line $\operatorname{Re}(s) = 1/2$ is the Riemann Hypothesis (?? 5.5.6), and the functional equation is what makes that line the natural object of the conjecture. The development begins with the gamma function, the analytic prerequisite on which the completed zeta function, and hence the entire chapter, rests.

6.1 The Gamma Function

The gamma function appeared in chapter 2 as the Mellin transform of e^{-t} (example 2.4.2) and as the inseparable companion of $\zeta(s)$ in the integral representation $\Gamma(s)\zeta(s) = \int_0^\infty t^{s-1}/(e^t - 1) dt$ of theorem 2.4.5. There it was used only on the half-plane $\operatorname{Re}(s) > 0$ where its defining integral converges. The functional equation requires Γ on the whole plane: the completed zeta function $\xi(s)$ of section 6.3 is built from the factor $\Gamma(s/2)$, and the symmetry $s \leftrightarrow 1 - s$ that ξ enjoys can only be read off once Γ is understood as a meromorphic function with its full pole structure. This section develops that structure in full: the meromorphic continuation of Γ to $\mathbb{C}$, its poles and their residues, its absence of zeros, and the two identities (reflection and duplication) that, together with the asymptotic estimate of Stirling, control its size and analytic behavior throughout the plane.

The starting point is the integral that defined Γ in chapter 2, recorded here as the definition for this chapter.

Definition 6.1.1: The Gamma Function

The *gamma function* is defined for $\operatorname{Re}(s) > 0$ by the integral

$$\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt$$

the Mellin transform of e^{-t} evaluated at s (example 2.4.2). The integral converges absolutely on $\operatorname{Re}(s) > 0$ and defines a holomorphic function there.

The convergence and holomorphy were established in example 2.4.2: near $t = 0^+$ the integrand is comparable to $t^{\operatorname{Re}(s)-1}$, integrable precisely when $\operatorname{Re}(s) > 0$, and near $t = \infty$ the factor e^{-t} forces convergence for every s . Holomorphy follows from the standard theorem on integrals depending holomorphically on a parameter, applied on compact subsets of the half-plane ([16]). The half-plane $\operatorname{Re}(s) > 0$ is the natural domain of the integral, but it is not the natural domain of the function: Γ extends past it, and the extension is forced by a single functional relation.

The relation in question is the integration-by-parts identity $\Gamma(s+1) = s\Gamma(s)$, the analytic shadow of the factorial recursion $n! = n \cdot (n-1)!$. Read forward it is a curiosity; read backward it is a continuation device, for it expresses $\Gamma(s)$ in terms of $\Gamma(s+1)$, a value with a larger real part, and so pushes the domain of definition leftward one unit at a time.

Theorem 6.1.2: Meromorphic Continuation of Γ

The gamma function satisfies the functional relation

$$\Gamma(s+1) = s\Gamma(s) \tag{6.1}$$

for all s with $\operatorname{Re}(s) > 0$. Through this relation Γ extends to a meromorphic function on all of $\mathbb{C}$, holomorphic everywhere except for simple poles at the non-positive integers $s = 0, -1, -2, \dots$, with

$$\operatorname{Res}_{s=-n}\Gamma(s) = \frac{(-1)^n}{n!} \quad (n = 0, 1, 2, \dots)$$

The continued function has no zeros: $\Gamma(s) \neq 0$ for every $s \in \mathbb{C}$.

Proof:

The proof has three stages: the functional relation on the half-plane of convergence; the continuation it forces, with the location and order of the poles; and the residues, followed by the absence of zeros.

Step 1: The functional relation. Fix s with $\operatorname{Re}(s) > 0$. Integration by parts in $\Gamma(s+1) = \int_0^\infty e^{-t} t^s dt$, with $u = t^s$ and $dv = e^{-t} dt$, gives $du = s t^{s-1} dt$ and $v = -e^{-t}$, so that

$$\Gamma(s+1) = \left[-e^{-t} t^s \right]_0^\infty + s \int_0^\infty e^{-t} t^{s-1} dt$$

The boundary term vanishes at both ends: as $t \rightarrow \infty$ the factor e^{-t} dominates the power t^s , and as $t \rightarrow 0^+$ the power $t^s \rightarrow 0$ because $\operatorname{Re}(s) > 0$. The remaining integral is $s \Gamma(s)$, establishing (6.1).

Step 2: The continuation and the poles. Solving (6.1) for $\Gamma(s)$ gives $\Gamma(s) = \Gamma(s+1)/s$. Iterating the relation N times, valid as an identity between meromorphic functions wherever both sides are defined, expresses

$$\Gamma(s) = \frac{\Gamma(s+N)}{s(s+1)(s+2)\cdots(s+N-1)} \quad (6.2)$$

Fix an integer $N \geq 1$ and consider (6.2) on the half-plane $\operatorname{Re}(s) > -N$. The numerator $\Gamma(s+N)$ is holomorphic there, because $\operatorname{Re}(s+N) > 0$ puts its argument in the domain of definition 6.1.1. The denominator is a polynomial in s vanishing exactly at $s = 0, -1, -2, \dots, -(N-1)$, each a simple zero. The right-hand side of (6.2) is therefore a meromorphic function on $\operatorname{Re}(s) > -N$, holomorphic except for simple poles at $s = 0, -1, \dots, -(N-1)$, and on the overlap $\operatorname{Re}(s) > 0$ it agrees with the integral $\Gamma(s)$ by (6.1). Two meromorphic functions on the connected open set $\operatorname{Re}(s) > -N$ agreeing on the nonempty open subset $\operatorname{Re}(s) > 0$ are equal, by the identity theorem ([16]); hence (6.2) furnishes the meromorphic continuation of Γ to $\operatorname{Re}(s) > -N$. The poles introduced for one value of N persist for every larger value, since enlarging N multiplies numerator and denominator by the additional factors $\Gamma(s+N+1)/\Gamma(s+N) = s+N$ in a way consistent with (6.1); letting $N \rightarrow \infty$ assembles a single meromorphic function on all of $\mathbb{C}$, holomorphic except for simple poles at every non-positive integer.

Step 3: The residues and the absence of zeros. Fix $n \geq 0$ and compute the residue at $s = -n$ from (6.2) with any $N > n$. Near $s = -n$ the only vanishing factor in the denominator is $(s+n)$, so the residue is the limit

$$\operatorname{Res}_{s=-n} \Gamma(s) = \lim_{s \rightarrow -n} (s+n) \Gamma(s) = \frac{\Gamma(-n+N)}{(-n)(-n+1)\cdots(-1) \cdot (1)(2)\cdots(N-1-n)}$$

where in the denominator the factor $(s+n)$ has been removed and the remaining factors evaluated at $s = -n$. The product of the negative factors $(-n)(-n+1)\cdots(-1)$ has n terms and equals $(-1)^n n!$; the product of the positive factors $(1)(2)\cdots(N-1-n)$ equals $(N-1-n)!$; and the numerator is $\Gamma(N-n) = (N-n-1)!$ since $N-n$ is a positive integer (example 6.1.5). The positive-factor product cancels the numerator, leaving

$$\operatorname{Res}_{s=-n} \Gamma(s) = \frac{(N-n-1)!}{(-1)^n n! (N-1-n)!} = \frac{(-1)^n}{n!}$$

as claimed. For the absence of zeros, suppose $\Gamma(s_0) = 0$ for some $s_0 \in \mathbb{C}$; necessarily s_0 is not a pole. Reflection (theorem 6.1.3) gives $\Gamma(s_0)\Gamma(1-s_0) = \pi/\sin(\pi s_0)$, and the right-hand side is finite and nonzero unless s_0 is an integer. If s_0 is not an integer, the product $\Gamma(s_0)\Gamma(1-s_0)$ is nonzero, so $\Gamma(s_0) \neq 0$, a contradiction. If s_0 is a positive integer, $\Gamma(s_0) = (s_0-1)! \neq 0$ (example 6.1.5); and s_0 cannot be a non-positive integer, those being the poles. Hence Γ has no zeros, completing the proof.

The pole structure recorded here is the feature that the rest of the chapter turns into arithmetic. The completed zeta function of section 6.3 carries the factor $\Gamma(s/2)$, whose argument $s/2$ runs through the poles $0, -1, -2, \dots$ of Γ exactly when s runs through $0, -2, -4, \dots$. At each such s the factor $\Gamma(s/2)$ becomes infinite; since the completed function is finite there (away from $s = 0$), the companion factor $\zeta(s)$ must vanish to compensate. The poles of Γ at the non-positive integers are, through this transfer, the source of the trivial zeros of ζ at the negative even integers $s = -2, -4, -6, \dots$, made precise in corollary 6.4.1. That Γ has no zeros is equally significant: it guarantees that the gamma factor introduces no *spurious* zeros into the completed function, so that every zero of $\xi(s)$ traces back to a zero of $\zeta(s)$ and not to an accident of the archimedean factor.

The recursion (6.1) fixes the values of Γ at integers and half-integers, but it says nothing about the relation between $\Gamma(s)$ and $\Gamma(1-s)$, the pairing that the functional equation will need. That relation is the reflection formula, and it is most cleanly obtained through the Beta integral, the two-variable companion of Γ .

Theorem 6.1.3: The Reflection Formula

For every $s \in \mathbb{C}$ that is not an integer,

$$\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)} \quad (6.3)$$

Proof:

The argument proves (6.3) first on the strip $0 < \operatorname{Re}(s) < 1$ and then extends it to all non-integer s by analytic continuation. The strip is reached through the Beta function, whose relation to Γ is established as the first step.

Step 1: The Beta integral and its evaluation by Γ . For complex x, y with $\operatorname{Re}(x) > 0$ and $\operatorname{Re}(y) > 0$ define the *Beta integral*

$$B(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt$$

which converges absolutely there: near $t = 0$ the integrand is comparable to $t^{\operatorname{Re}(x)-1}$ and near $t = 1$ to $(1-t)^{\operatorname{Re}(y)-1}$, both integrable. The identity $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x+y)$ is obtained from the product of the two gamma integrals. By definition 6.1.1,

$$\Gamma(x)\Gamma(y) = \int_0^\infty \int_0^\infty e^{-u-v} u^{x-1} v^{y-1} du dv$$

the product of two absolutely convergent integrals written as a double integral over the first quadrant, the interchange into a single double integral justified by Tonelli's theorem applied to the absolute values $e^{-u-v} u^{\operatorname{Re}(x)-1} v^{\operatorname{Re}(y)-1}$, whose double integral is the finite product $\Gamma(\operatorname{Re} x)\Gamma(\operatorname{Re} y)$ ([8]). Change variables by $u = r\tau$, $v = r(1-\tau)$ with $r \in (0, \infty)$ and $\tau \in (0, 1)$; this maps the

strip $\{(r, \tau)\}$ bijectively onto the first quadrant, with $u + v = r$ and Jacobian

$$\frac{\partial(u, v)}{\partial(r, \tau)} = \det \begin{pmatrix} \tau & r \\ 1 - \tau & -r \end{pmatrix} = -r\tau - r(1 - \tau) = -r$$

so $du dv = r dr d\tau$. Substituting,

$$\Gamma(x)\Gamma(y) = \int_0^1 \int_0^\infty e^{-r} (r\tau)^{x-1} (r(1-\tau))^{y-1} r dr d\tau = \int_0^\infty e^{-r} r^{x+y-1} dr \int_0^1 \tau^{x-1} (1-\tau)^{y-1} d\tau$$

the separation of the double integral into a product again justified by Tonelli on the absolute values. The first factor is $\Gamma(x+y)$ and the second is $B(x, y)$, giving

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad (6.4)$$

for $\operatorname{Re}(x), \operatorname{Re}(y) > 0$.

Step 2: Reduction of the reflection product to a single integral. Take $y = 1 - x$ with $0 < \operatorname{Re}(x) < 1$, so both $\operatorname{Re}(x) > 0$ and $\operatorname{Re}(1 - x) > 0$ hold and (6.4) applies. Since $x + y = 1$ and $\Gamma(1) = 1$ (example 6.1.5), (6.4) reads

$$\Gamma(x)\Gamma(1-x) = B(x, 1-x) = \int_0^1 t^{x-1} (1-t)^{-x} dt$$

The substitution $u = t/(1-t)$, equivalently $t = u/(1+u)$, carries $t \in (0, 1)$ to $u \in (0, \infty)$ with $1-t = 1/(1+u)$ and $dt = du/(1+u)^2$. The integrand transforms as

$$t^{x-1} (1-t)^{-x} dt = \left(\frac{u}{1+u} \right)^{x-1} (1+u)^x \cdot \frac{du}{(1+u)^2} = u^{x-1} (1+u)^{-(x-1)+x-2} du = \frac{u^{x-1}}{1+u} du$$

the powers of $(1+u)$ combining to the single exponent $-(x-1)+x-2 = -1$, so that

$$\Gamma(x)\Gamma(1-x) = \int_0^\infty \frac{u^{x-1}}{1+u} du \quad (6.5)$$

an integral converging for $0 < \operatorname{Re}(x) < 1$, since the integrand is comparable to $u^{\operatorname{Re}(x)-1}$ near $u = 0$ and to $u^{\operatorname{Re}(x)-2}$ near $u = \infty$.

Step 3: Evaluation of the integral and continuation. The integral on the right of (6.5) evaluates to $\pi/\sin(\pi x)$ for $0 < \operatorname{Re}(x) < 1$. This is a contour integral computed by the residue theorem: integrating $z^{x-1}/(1+z)$ around a keyhole contour about the branch cut along the positive real axis picks up the residue at the simple pole $z = -1$, and the two sides of the cut combine to give $(1 - e^{2\pi i(x-1)}) \int_0^\infty u^{x-1}/(1+u) du = 2\pi i e^{\pi i(x-1)}$, which rearranges to $\pi/\sin(\pi x)$. The full keyhole computation is carried out in [16, Residue theorem chapter]; the result is

$$\int_0^\infty \frac{u^{x-1}}{1+u} du = \frac{\pi}{\sin(\pi x)} \quad (0 < \operatorname{Re}(x) < 1)$$

Combining this with (6.5) gives (6.3) on the strip $0 < \operatorname{Re}(s) < 1$. Both sides of (6.3) are meromorphic on all of $\mathbb{C}$: the left side $\Gamma(s)\Gamma(1-s)$ by theorem 6.1.2, the right side $\pi/\sin(\pi s)$ as the ratio of an entire function and the entire function $\sin(\pi s)$. They agree on the strip $0 < \operatorname{Re}(s) < 1$, a set with accumulation points, so by the identity theorem ([16]) they agree wherever both are defined, that is, for every non-integer s . This establishes (6.3), as we sought to show.

The reflection formula is the algebraic heart of the symmetry $s \leftrightarrow 1 - s$. It pairs the value of Γ at s with its value at $1 - s$ through the elementary factor $\pi/\sin(\pi s)$, and this pairing is exactly the form the gamma factors take in the functional equation of theorem 6.3.3: the completed function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ inherits its reflection symmetry from the interplay of this formula with the theta transformation of section 6.2. The formula also re-derives the pole structure of theorem 6.1.2 from a different vantage. The right-hand side $\pi/\sin(\pi s)$ has simple poles at every integer; on the negative side $s = -n$ ($n \geq 0$), the factor $\Gamma(1-s) = \Gamma(1+n) = n!$ is finite and nonzero, so the pole must come from $\Gamma(s)$, confirming that Γ has a simple pole there. On the positive side $s = n \geq 1$, the factor $\Gamma(s) = (n-1)!$ is finite, so the pole of $\pi/\sin(\pi s)$ is absorbed by $\Gamma(1-s)$, which indeed has a pole at $1-n \leq 0$. The reflection formula thus encodes the entire pole pattern of Γ in the zeros of $\sin(\pi s)$. Combined with the residues of theorem 6.1.2, it is the tool that, in section 6.4, converts the poles of $\Gamma(s/2)$ into the trivial zeros of ζ and the values of Γ at half-integers into the special values $\zeta(2k)$.

A second multiplicative identity relates Γ at s and $s + 1/2$ to Γ at $2s$. Where reflection pairs a point with its mirror image across $\operatorname{Re}(s) = 1/2$, the duplication formula interpolates between the lattice of integers and the lattice of half-integers, and it is the gamma-function input to the relation between $\zeta(s)$ and the completed function built from $\Gamma(s/2)$.

Theorem 6.1.4: Legendre Duplication and Stirling

The gamma function satisfies the *Legendre duplication formula*

$$\Gamma(s)\Gamma\left(s + \frac{1}{2}\right) = 2^{1-2s}\sqrt{\pi}\Gamma(2s) \quad (6.6)$$

for all $s \in \mathbb{C}$ at which both sides are defined. Moreover Γ obeys the asymptotic estimate of Stirling: for $|\arg s| \leq \pi - \delta$ with $\delta > 0$ fixed, as $|s| \rightarrow \infty$,

$$\log \Gamma(s) = \left(s - \frac{1}{2}\right) \log s - s + \frac{1}{2} \log(2\pi) + O\left(\frac{1}{|s|}\right) \quad (6.7)$$

with the principal branch of the logarithm, the implied constant depending only on δ .

Proof:

The duplication formula is proved in full through the Beta integral; the complex Stirling estimate is treated by key ideas with a precise reference, for the reason stated below.

Step 1: The duplication formula. Assume first $\operatorname{Re}(s) > 0$, so that all Beta integrals below converge; the identity then extends to all admissible s by analytic continuation exactly as in theorem 6.1.3. Apply (6.4) with $x = y = s$:

$$B(s, s) = \frac{\Gamma(s)^2}{\Gamma(2s)} = \int_0^1 (t(1-t))^{s-1} dt$$

Substitute $t = (1+u)/2$, so $dt = du/2$ and $t(1-t) = (1-u^2)/4$, carrying $t \in (0, 1)$ to $u \in (-1, 1)$:

$$B(s, s) = \int_{-1}^1 \left(\frac{1-u^2}{4}\right)^{s-1} \frac{du}{2} = 2^{1-2s} \int_{-1}^1 (1-u^2)^{s-1} du$$

where $\frac{1}{2} \cdot 4^{1-s} = 2^{-1}2^{2-2s} = 2^{1-2s}$. The integrand is even in u , so the integral over $(-1, 1)$ is twice

the integral over $(0, 1)$. In that integral substitute $v = u^2$, $dv = 2u du$, hence $du = \frac{1}{2}v^{-1/2} dv$:

$$2 \int_0^1 (1 - u^2)^{s-1} du = 2 \int_0^1 (1 - v)^{s-1} \cdot \frac{1}{2}v^{-1/2} dv = \int_0^1 v^{-1/2}(1 - v)^{s-1} dv = B\left(\frac{1}{2}, s\right)$$

the last integral being the Beta integral with parameters $\frac{1}{2}$ and s . Assembling the two displays,

$$\frac{\Gamma(s)^2}{\Gamma(2s)} = 2^{1-2s} B\left(\frac{1}{2}, s\right) = 2^{1-2s} \frac{\Gamma(\frac{1}{2}) \Gamma(s)}{\Gamma(s + \frac{1}{2})}$$

using (6.4) once more for $B(\frac{1}{2}, s)$. Now $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ (example 6.1.5). Multiplying both sides by $\Gamma(s + \frac{1}{2})/\Gamma(s)$ clears the common factor $\Gamma(s)$ and yields

$$\frac{\Gamma(s) \Gamma(s + \frac{1}{2})}{\Gamma(2s)} = 2^{1-2s} \sqrt{\pi}$$

which is (6.6). Both sides of (6.6) are meromorphic on $\mathbb{C}$, so the identity, proved on $\operatorname{Re}(s) > 0$, holds wherever both sides are defined by the identity theorem ([16]).

Step 2: Stirling's estimate. The real-variable form $\log \Gamma(x+1) = x \log x - x + \frac{1}{2} \log(2\pi x) + O(1/x)$ as $x \rightarrow \infty$ is theorem 1.5.7; since $\log \Gamma(x+1) = \log x + \log \Gamma(x)$ and $\frac{1}{2} \log(2\pi x) = \frac{1}{2} \log(2\pi) + \frac{1}{2} \log x$, subtracting $\log x$ recovers (6.7) for real $s = x \rightarrow +\infty$, with $(x - \frac{1}{2}) \log x = x \log x - \frac{1}{2} \log x$. The extension to the complex sector $|\arg s| \leq \pi - \delta$ requires more than the Euler–Maclaurin computation that produced the real case: it requires controlling the error term uniformly as $\arg s$ ranges over the sector, which is most efficiently done either from the Binet integral representation of $\log \Gamma$ or by combining the real-axis asymptotic with the functional relation (6.1) and a Phragmén–Lindelöf argument to propagate the bound off the real axis. The full complex argument is several pages of estimation with no new conceptual content beyond the real case, and is carried out in [16, Stirling/asymptotics chapter]; see also [11, Ch. 4] for the form with explicit error terms in the sector. The estimate (6.7) is as stated, completing the proof.

Stirling's estimate is the quantitative control that the analytic theory of ζ ultimately rests on. The completed function $\xi(s)$ of section 6.3 is entire of order one, and that order is a direct consequence of (6.7): the gamma factor $\Gamma(s/2)$ grows like $\exp(\frac{1}{2}|s| \log |s|)$ in vertical strips, and it is this growth rate, balanced against the polynomial size of ζ in the strip, that determines the density of the zeros and underlies the explicit formula of box 5.5.5. The duplication formula, for its part, is the algebraic identity that allows the factor $\Gamma(s/2)$ appearing in $\xi(s)$ to be related to the factor $\Gamma(s)$ appearing in the integral representation $\Gamma(s)\zeta(s) = \int_0^\infty t^{s-1}/(e^t - 1) dt$ of theorem 2.4.5; the two normalizations differ by exactly the elementary factor that (6.6) supplies. Together, reflection and duplication exhaust the multiplicative identities of Γ that the functional equation needs.

The abstract development is best anchored by computing Γ at the values where it is most often used: the integers, where it reproduces the factorial, and the half-integers, where it produces the powers of $\sqrt{\pi}$ that appear throughout the theory.

Example 6.1.5: Special Values of Γ

1. *Integer values.* Direct evaluation of definition 6.1.1 at $s = 1$ gives

$$\Gamma(1) = \int_0^\infty e^{-t} dt = 1$$

The recursion (6.1) then gives $\Gamma(2) = 1 \cdot \Gamma(1) = 1$, $\Gamma(3) = 2 \cdot \Gamma(2) = 2$, and by induction $\Gamma(n) = (n-1)!$ for every positive integer n : if $\Gamma(n) = (n-1)!$ then $\Gamma(n+1) = n \Gamma(n) = n \cdot (n-1)! = n!$. Thus Γ interpolates the factorial, with $\Gamma(n+1) = n!$.

2. *The value at $\frac{1}{2}$.* Setting $s = \frac{1}{2}$ in the reflection formula (6.3) gives $\Gamma(\frac{1}{2})^2 = \Gamma(\frac{1}{2})\Gamma(1 - \frac{1}{2}) = \pi / \sin(\pi/2) = \pi$. Since $\Gamma(\frac{1}{2}) = \int_0^\infty e^{-t} t^{-1/2} dt > 0$ is a positive real number, taking the positive square root gives

$$\Gamma(\tfrac{1}{2}) = \sqrt{\pi}$$

The same value follows from the Gaussian integral: the substitution $t = w^2$ in $\Gamma(\frac{1}{2})$ gives $\Gamma(\frac{1}{2}) = \int_0^\infty e^{-w^2} \cdot w^{-1} \cdot 2w dw = 2 \int_0^\infty e^{-w^2} dw = \int_{-\infty}^\infty e^{-w^2} dw = \sqrt{\pi}$ ([7]).

3. *Half-integer values.* The recursion (6.1) propagates the value at $\frac{1}{2}$ to all half-integers. From $\Gamma(s+1) = s \Gamma(s)$ with $s = \frac{1}{2}, \frac{3}{2}, \dots$,

$$\Gamma(\tfrac{3}{2}) = \tfrac{1}{2}\sqrt{\pi}, \quad \Gamma(\tfrac{5}{2}) = \tfrac{3}{2} \cdot \tfrac{1}{2}\sqrt{\pi} = \tfrac{3}{4}\sqrt{\pi}$$

and in general, for $n \geq 0$,

$$\Gamma(\tfrac{1}{2} + n) = \frac{(2n)!}{4^n n!} \sqrt{\pi}$$

The closed form is verified by induction: it holds at $n = 0$, and the step from n to $n+1$ multiplies by $\frac{1}{2} + n = (2n+1)/2$, while the right side is multiplied by $\frac{(2n+2)!}{4^{n+1}(n+1)!} / \frac{(2n)!}{4^n n!} = \frac{(2n+1)(2n+2)}{4(n+1)} = \frac{2n+1}{2}$, the two factors agreeing.

4. *Stirling at $s = 10$.* The estimate (6.7) at the integer $s = 10$, where $\log \Gamma(10) = \log 9! = \log(362,880) \approx 12.8018$, gives the approximation

$$(10 - \tfrac{1}{2}) \log 10 - 10 + \tfrac{1}{2} \log(2\pi) \approx 9.5 \cdot 2.302585 - 10 + 0.918939 \approx 12.7935$$

matching $\log \Gamma(10)$ to three significant figures. The discrepancy is $12.8018 - 12.7935 \approx 0.0083$, in agreement with the predicted error $O(1/|s|) = O(1/10)$: the first correction term of the full Stirling expansion (theorem 1.5.7) is $B_2/(2 \cdot 1 \cdot 10) = 1/(12 \cdot 10) \approx 0.00833$, and adding it brings the approximation to 12.8018, recovering $\log \Gamma(10)$ to the displayed precision.

6.2 Theta Functions and Poisson Summation

The functional equation of the zeta function relates $\zeta(s)$ to $\zeta(1-s)$, exchanging the half-plane $\operatorname{Re}(s) > 1$ where the series converges with its mirror image $\operatorname{Re}(s) < 0$. The source of this symmetry is not visible in the Dirichlet series itself; it lies one level down, in a symmetry of the Gaussian. The function $x \mapsto e^{-\pi x^2}$ is its own Fourier transform, and rescaling the variable converts that self-duality

into the relation $t \leftrightarrow 1/t$ for the Gaussian $e^{-\pi x^2 t}$. Summing over the integer lattice turns this analytic fact into an identity for the Jacobi theta function, the generating function $\theta(t) = \sum_{n \in \mathbb{Z}} e^{-\pi n^2 t}$, by way of the Poisson summation formula, which equates a sum over $\mathbb{Z}$ to a sum of Fourier-transform values over $\mathbb{Z}$.

This section develops the three ingredients in order: Poisson summation, the Fourier transform of a Gaussian, and the theta function with its transformation law $\theta(1/t) = \sqrt{t} \theta(t)$. The theta functional equation is the geometric heart of the matter; the functional equation of ζ , obtained in section 6.3 by feeding θ into a Mellin transform, is its arithmetic shadow. The $t \leftrightarrow 1/t$ symmetry established here becomes, after the change of variables $t \mapsto t^{-1}$ inside the Mellin integral, the $s \leftrightarrow 1-s$ symmetry of the completed zeta function.

The first ingredient is Poisson summation. It is the statement that periodizing a function and reading off its Fourier coefficients recovers the values of the Fourier transform at integer points, so that the total mass of a function over the integer lattice equals the total mass of its transform over the dual lattice. The hypotheses must rule out the two ways such an identity can fail: the sum $\sum_n f(n)$ must converge, and the periodization must have a convergent Fourier series that represents it pointwise. Schwartz decay, or any decay forcing both f and $\hat{f}$ to fall off faster than $|x|^{-1-\delta}$, secures both.

Theorem 6.2.1: Poisson Summation

Let $f: \mathbb{R} \rightarrow \mathbb{C}$ be a function for which there exist constants $C > 0$ and $\delta > 0$ with

$$|f(x)| \leq \frac{C}{(1+|x|)^{1+\delta}}, \quad |\hat{f}(\xi)| \leq \frac{C}{(1+|\xi|)^{1+\delta}}$$

for all $x, \xi \in \mathbb{R}$, where the Fourier transform is normalized by

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx$$

(every Schwartz function satisfies these bounds, as does $f(x) = e^{-\pi x^2 t}$ for fixed $t > 0$). Then both series below converge absolutely and

$$\sum_{n \in \mathbb{Z}} f(n) = \sum_{n \in \mathbb{Z}} \hat{f}(n) \quad (6.8)$$

Proof:

The argument periodizes f , expands the periodization in a Fourier series, and evaluates at the origin. Each step is justified by the stated decay.

Step 1: The periodization is a continuous 1-periodic function. Define

$$F(x) = \sum_{n \in \mathbb{Z}} f(x+n)$$

For x in the compact interval $[0, 1]$ and $|n| \geq 2$ one has $|x+n| \geq |n| - 1 \geq |n|/2$, so $|f(x+n)| \leq C(1+|n|/2)^{-1-\delta}$, a bound independent of x and summable over n because $\sum_n (1+|n|/2)^{-1-\delta} < \infty$ for $\delta > 0$. By the Weierstrass M -test the series defining F converges absolutely and uniformly on $[0, 1]$, hence its sum F is continuous there; the same bound on every translate $[k, k+1]$ gives continuity on all of $\mathbb{R}$. Replacing x by $x+1$ permutes the summands,

so $F(x+1) = F(x)$, and F is 1-periodic.

Step 2: The Fourier coefficients of F are the values $\widehat{f}(m)$. The m -th Fourier coefficient of the continuous periodic function F is

$$c_m = \int_0^1 F(x) e^{-2\pi i m x} dx = \int_0^1 \sum_{n \in \mathbb{Z}} f(x+n) e^{-2\pi i m x} dx$$

The uniform convergence from Step 1 permits interchanging the sum and the integral over the bounded interval $[0, 1]$. Substituting $y = x + n$ in the n -th term, and using $e^{-2\pi i m x} = e^{-2\pi i m(y-n)} = e^{-2\pi i m y}$ since $e^{2\pi i m n} = 1$ for integers m, n , gives

$$c_m = \sum_{n \in \mathbb{Z}} \int_0^1 f(x+n) e^{-2\pi i m x} dx = \sum_{n \in \mathbb{Z}} \int_n^{n+1} f(y) e^{-2\pi i m y} dy = \int_{-\infty}^{\infty} f(y) e^{-2\pi i m y} dy$$

the consecutive intervals $[n, n+1]$ tiling $\mathbb{R}$, with the reassembly of the integral justified by absolute convergence of $\int_{\mathbb{R}} |f| < \infty$, itself a consequence of the decay hypothesis. The last integral is $\widehat{f}(m)$.

Step 3: The Fourier series represents F pointwise at $x = 0$. By Step 2 the Fourier coefficients satisfy $|c_m| = |\widehat{f}(m)| \leq C(1+|m|)^{-1-\delta}$, so $\sum_m |c_m| < \infty$ and the Fourier series $\sum_m c_m e^{2\pi i m x}$ converges absolutely and uniformly. A continuous periodic function whose Fourier coefficients are absolutely summable equals the sum of its Fourier series at every point ([7]); the uniform limit of the partial sums is a continuous function with the same Fourier coefficients as F , and a continuous periodic function is determined by its Fourier coefficients. Evaluating the resulting identity $F(x) = \sum_m c_m e^{2\pi i m x}$ at $x = 0$ yields

$$\sum_{n \in \mathbb{Z}} f(n) = F(0) = \sum_{m \in \mathbb{Z}} c_m = \sum_{m \in \mathbb{Z}} \widehat{f}(m)$$

which is (6.8), completing the proof.

Poisson summation is a self-duality of the integer lattice: the lattice $\mathbb{Z}$ is its own dual under the pairing $(x, \xi) \mapsto e^{2\pi i x \xi}$, and the formula says that a function and its Fourier transform deposit the same total mass on $\mathbb{Z}$. The two sides see the function at complementary scales. The left side, $\sum_n f(n)$, samples f at integer arguments and is dominated by the behavior of f near the origin when f is concentrated there; the right side, $\sum_n \widehat{f}(n)$, samples the transform, which is concentrated near the origin precisely when f is spread out. This exchange of “concentrated near 0” for “spread out” is the mechanism that will convert small- t behavior of the theta function into large- t behavior, and it is the analytic content of the functional equation. The decay hypotheses are exactly what is needed to make every interchange in the proof legitimate; for the single function this section applies the formula to, the Gaussian, the hypotheses hold with room to spare, since a Gaussian and its transform decay faster than any power.

The function Poisson summation will be applied to is the Gaussian $e^{-\pi x^2 t}$, and the right side of (6.8) requires its Fourier transform. The defining property of the Gaussian among all functions is that it is an eigenfunction of the Fourier transform: the standard Gaussian $e^{-\pi x^2}$ is fixed, and rescaling the width sends the transform to the reciprocal width, with a compensating factor of $t^{-1/2}$.

Lemma 6.2.2: The Fourier Transform of a Gaussian

For each $t > 0$, the Fourier transform of the Gaussian $g_t(x) = e^{-\pi x^2 t}$, in the normalization $\widehat{g}_t(\xi) = \int_{\mathbb{R}} g_t(x) e^{-2\pi i x \xi} dx$, is

$$\widehat{g}_t(\xi) = t^{-1/2} e^{-\pi \xi^2 / t} \quad (6.9)$$

In particular $g_1 = e^{-\pi x^2}$ is its own Fourier transform.

Proof:

The standard Gaussian is treated first, and the general case follows by a change of variables.

Step 1: The transform of $g_1(x) = e^{-\pi x^2}$ is g_1 . Set $\varphi(\xi) = \widehat{g}_1(\xi) = \int_{\mathbb{R}} e^{-\pi x^2} e^{-2\pi i x \xi} dx$. The integrand and its ξ -derivative $-2\pi i x e^{-\pi x^2} e^{-2\pi i x \xi}$ are dominated in absolute value by the integrable functions $e^{-\pi x^2}$ and $2\pi |x| e^{-\pi x^2}$, uniformly for ξ in any bounded set, so differentiation under the integral sign is justified ([8]) and

$$\varphi'(\xi) = \int_{\mathbb{R}} e^{-\pi x^2} (-2\pi i x) e^{-2\pi i x \xi} dx$$

The factor $-2\pi x e^{-\pi x^2}$ is the derivative $\frac{d}{dx} e^{-\pi x^2}$, so integration by parts with $u = e^{-2\pi i x \xi}$ and $dv = i \left(\frac{d}{dx} e^{-\pi x^2} \right) dx$ gives, the boundary term vanishing because $e^{-\pi x^2} \rightarrow 0$ as $x \rightarrow \pm\infty$,

$$\varphi'(\xi) = i \left[e^{-2\pi i x \xi} e^{-\pi x^2} \right]_{-\infty}^{\infty} - i \int_{\mathbb{R}} e^{-\pi x^2} (-2\pi i \xi) e^{-2\pi i x \xi} dx = -2\pi \xi \varphi(\xi)$$

This first-order linear differential equation has solution $\varphi(\xi) = \varphi(0) e^{-\pi \xi^2}$. The initial value is the Gaussian integral $\varphi(0) = \int_{\mathbb{R}} e^{-\pi x^2} dx = 1$, the normalization built into the factor π in the exponent ([8]). Hence $\varphi(\xi) = e^{-\pi \xi^2} = g_1(\xi)$.

Step 2: Rescale to obtain g_t . Fix $t > 0$ and write $g_t(x) = g_1(\sqrt{t}x)$. Substituting $u = \sqrt{t}x$, so $du = \sqrt{t}dx$, in the defining integral gives

$$\widehat{g}_t(\xi) = \int_{\mathbb{R}} g_1(\sqrt{t}x) e^{-2\pi i x \xi} dx = \frac{1}{\sqrt{t}} \int_{\mathbb{R}} g_1(u) e^{-2\pi i u (\xi/\sqrt{t})} du = \frac{1}{\sqrt{t}} \widehat{g}_1\left(\frac{\xi}{\sqrt{t}}\right)$$

By Step 1, $\widehat{g}_1(\xi/\sqrt{t}) = e^{-\pi (\xi/\sqrt{t})^2} = e^{-\pi \xi^2 / t}$, so $\widehat{g}_t(\xi) = t^{-1/2} e^{-\pi \xi^2 / t}$, which is (6.9). Setting $t = 1$ recovers $\widehat{g}_1 = g_1$, as we sought to show.

The identity (6.9) packages the entire dependence on t into the reciprocal $1/t$ in the exponent and the prefactor $t^{-1/2}$. A narrow Gaussian (large t , sharply peaked at the origin) transforms to a wide one (small $1/t$, spread out), and conversely; the Fourier transform interchanges the two regimes, with the prefactor $t^{-1/2}$ enforcing the correct normalization of total mass. This is the precise sense in which the Gaussian realizes the uncertainty principle with equality, and it is the single input that makes the theta function transform cleanly: summing (6.9) over the lattice will send t to $1/t$ and produce the factor $\sqrt{t}$ that appears in the theta functional equation.

The theta function is the lattice sum of the Gaussian. Collecting the values $e^{-\pi n^2 t}$ over all integers n produces a function of t that encodes the Gaussian at every integer point at once, and it is the

object on which Poisson summation acts.

Definition 6.2.3: The Jacobi Theta Function

The *Jacobi theta function* is the series

$$\theta(t) = \sum_{n \in \mathbb{Z}} e^{-\pi n^2 t}, \quad t > 0$$

Separating the term $n = 0$ from the pair $\pm n$ for $n \geq 1$, which contribute equally, gives the equivalent form

$$\theta(t) = 1 + 2 \sum_{n=1}^{\infty} e^{-\pi n^2 t} \quad (6.10)$$

For every $t > 0$ the series converges absolutely and rapidly: the n -th term is $e^{-\pi n^2 t}$, and for $n \geq 1$ this is bounded by $e^{-\pi n t} \cdot e^{-\pi(n^2-n)t} \leq (e^{-\pi t})^n$, so the tail is dominated by a convergent geometric series with ratio $e^{-\pi t} < 1$. The convergence is locally uniform on $t > 0$, since on any interval $t \geq t_0 > 0$ the same geometric bound holds with ratio $e^{-\pi t_0} < 1$; hence θ is continuous, indeed real-analytic, on $(0, \infty)$. As $t \rightarrow +\infty$ every term with $n \neq 0$ tends to 0, and the geometric bound shows the tail $\sum_{n \geq 1} e^{-\pi n^2 t} \leq e^{-\pi t} / (1 - e^{-\pi t})$ vanishes, so

$$\theta(t) = 1 + 2 \sum_{n=1}^{\infty} e^{-\pi n^2 t} \longrightarrow 1 \quad (t \rightarrow +\infty)$$

the $n = 0$ term surviving alone. The behavior as $t \rightarrow 0^+$ is the opposite extreme: every term tends to 1, the series loses its geometric decay, and $\theta(t)$ diverges. Quantifying that divergence is exactly what the functional equation does.

The functional equation is the transformation law of θ under $t \mapsto 1/t$. It is the lattice-sum form of the Gaussian self-duality of lemma 6.2.2, obtained by applying Poisson summation to the Gaussian and reading the prefactor $t^{-1/2}$ as the source of the $\sqrt{t}$.

Theorem 6.2.4: The Functional Equation of the Theta Function

For all $t > 0$,

$$\theta\left(\frac{1}{t}\right) = \sqrt{t} \theta(t) \quad (6.11)$$

Proof:

Fix $t > 0$ and apply Poisson summation (theorem 6.2.1) to the Gaussian $g_t(x) = e^{-\pi x^2 t}$. The function g_t satisfies the decay hypotheses of theorem 6.2.1: it decays faster than any power of $|x|$, and by lemma 6.2.2 its transform $\widehat{g}_t(\xi) = t^{-1/2} e^{-\pi \xi^2 / t}$ does as well, so both required bounds hold (with any $\delta > 0$, for a suitable constant C depending on t). The formula (6.8) therefore applies and gives

$$\sum_{n \in \mathbb{Z}} g_t(n) = \sum_{n \in \mathbb{Z}} \widehat{g}_t(n)$$

The left side is $\sum_{n \in \mathbb{Z}} e^{-\pi n^2 t} = \theta(t)$ by the definition of θ . For the right side, lemma 6.2.2 gives

$\widehat{g}_t(n) = t^{-1/2} e^{-\pi n^2/t}$, so

$$\sum_{n \in \mathbb{Z}} \widehat{g}_t(n) = t^{-1/2} \sum_{n \in \mathbb{Z}} e^{-\pi n^2/t} = t^{-1/2} \theta\left(\frac{1}{t}\right)$$

the last equality being the definition of θ evaluated at $1/t$. Equating the two sides yields

$$\theta(t) = t^{-1/2} \theta\left(\frac{1}{t}\right) \quad (6.12)$$

Multiplying both sides of (6.12) by $t^{1/2} = \sqrt{t}$ isolates $\theta(1/t)$ on one side and gives $\theta(1/t) = \sqrt{t} \theta(t)$, which is (6.11), completing the proof.

The transformation law (6.11) is the symmetry that the functional equation of ζ inherits. It relates the value of θ at t to its value at the reciprocal $1/t$, exchanging the regime t large, where $\theta(t) \rightarrow 1$, with the regime t small, where $\theta(t)$ blows up. The point $t = 1$ is the fixed point of the involution $t \mapsto 1/t$, and (6.11) reads $\theta(1) = \sqrt{1} \theta(1)$ there, a tautology that locates the axis of symmetry. This is the same structural picture that will govern ζ : an involution with a distinguished fixed point, relating the function on one side to its values on the other. The connection is made precise in section 6.3 through the Mellin transform of chapter 2. Integrating $\theta(t) - 1$ against $t^{s/2}$ produces, up to elementary factors, the completed zeta function $\xi(s)$; the change of variables $t \mapsto 1/t$ inside that integral turns the theta symmetry $t \leftrightarrow 1/t$ into the reflection $s \leftrightarrow 1 - s$, because $t^{s/2} dt/t$ becomes $t^{(1-s)/2} dt/t$ under $t \mapsto 1/t$ once the factor $\sqrt{t}$ from (6.11) is absorbed. The quadratic exponent n^2 in $e^{-\pi n^2/t}$ is what forces the Mellin pairing to be against the half-power $t^{s/2}$ rather than t^s , and this halving is what centers the zeta symmetry at $s = \frac{1}{2}$ rather than at an integer, the analytic origin of the critical line $\text{Re}(s) = \frac{1}{2}$. In this sense the entire functional-equation apparatus of the next section is a translation of the single identity (6.11) into the language of Dirichlet series.

The asymmetry between the two limiting regimes of θ is what the functional equation quantifies, and a numerical table makes both the asymmetry and the symmetry $t \leftrightarrow 1/t$ concrete before they are put to analytic use.

Example 6.2.5: Decay and the Modular Transformation

The functional equation (6.11) converts the elementary limit $\theta(s) \rightarrow 1$ as $s \rightarrow +\infty$ into the precise blow-up rate of θ at the origin. Setting $u = 1/t$, so that $t \rightarrow 0^+$ corresponds to $u \rightarrow +\infty$, the relation $\theta(1/t) = \sqrt{t} \theta(t)$ reads $\theta(u) = \sqrt{1/u} \theta(1/u)$, hence

$$\theta\left(\frac{1}{u}\right) = \sqrt{u} \theta(u) = \sqrt{u} (1 + o(1)) \quad (u \rightarrow +\infty)$$

since $\theta(u) \rightarrow 1$. Writing $t = 1/u$, this is the asymptotic

$$\theta(t) \sim t^{-1/2} \quad (t \rightarrow 0^+) \quad (6.13)$$

The two ends of the range thus behave oppositely: $\theta(t) - 1$ decays to 0 faster than any power of t as $t \rightarrow +\infty$ (geometrically, by the bound of definition 6.2.3), while $\theta(t)$ itself diverges like $t^{-1/2}$ as $t \rightarrow 0^+$. This asymmetry, rapid decay at one end against a $t^{-1/2}$ singularity at the other, is precisely what splits the Mellin integral defining $\xi(s)$ into a convergent piece and a pair of elementary polar terms in section 6.3.

A short table of values, computed by summing the rapidly convergent series (6.10) to ten significant figures, exhibits both the symmetry and the divergence. Each pair $(t, 1/t)$ is related by (6.11):

t	$1/4$	$1/2$	1	2	4
$\theta(t)$	2.000013949	1.419495488	1.086434811	1.003734885	1.000006975

The growth as t decreases toward 0 is the $t^{-1/2}$ trend of (6.13): at $t = 1/4$ the value $\theta(1/4) = 2.0000\dots$ is already close to $t^{-1/2} = 2$, and at $t = 4$ the value $\theta(4) = 1.0000\dots$ is close to its limit 1. The functional equation is verified directly against these entries. For the pair $t = 4$ and $1/t = 1/4$,

$$\sqrt{4} \theta(4) = 2 \times 1.000006975 = 2.000013950, \quad \theta\left(\frac{1}{4}\right) = 2.000013949$$

agreeing to the recorded precision, so $\theta(1/4) / (\sqrt{4} \theta(4)) = 1.0000000000$. The same check on the pair $t = 2$, $1/t = 1/2$ gives $\sqrt{2} \theta(2) = 1.41421356 \times 1.003734885 = 1.419495487$ against $\theta(1/2) = 1.419495488$, again a ratio of 1 to ten digits. The fixed point $t = 1$ of the involution $t \mapsto 1/t$ is self-paired, and the entry $\theta(1) = 1.086434811$ satisfies $\sqrt{1} \theta(1) = \theta(1)$ trivially; this value equals $\pi^{1/4} / \Gamma(3/4)$ in closed form, the one argument at which θ admits an evaluation in terms of the gamma function.

6.3 The Completed Zeta Function and Its Functional Equation

The two preceding sections assembled the ingredients: the gamma function with its full pole structure (section 6.1), and the Jacobi theta function with its transformation law $\theta(1/t) = \sqrt{t} \theta(t)$ under the involution $t \mapsto 1/t$ (section 6.2). Multiplying $\zeta(s)$ by the archimedean gamma factor $\pi^{-s/2} \Gamma(s/2)$ produces a single function, the completed zeta function, that is symmetric under $s \leftrightarrow 1 - s$. This symmetry is Riemann's functional equation. The mechanism that produces it is the Mellin transform of chapter 2: writing the completed function as the Mellin transform of $\theta(t) - 1$ converts the theta symmetry $t \leftrightarrow 1/t$, after the change of variables $t \mapsto 1/t$ inside the integral, into the reflection $s \leftrightarrow 1 - s$.

The functional equation does two things at once. It continues $\zeta(s)$ from the half-plane $\operatorname{Re}(s) > 0$ reached in chapter 4 to the entire complex plane, and it pins the continuation down by a reflection symmetry whose axis is the line $\operatorname{Re}(s) = 1/2$. Every later structural fact about ζ , the trivial zeros of section 6.4, the special values $\zeta(2k)$ promised in example 4.1.3, and the centering of the Riemann Hypothesis on the critical line, is read off from this one identity.

The object that carries the symmetry is ζ multiplied by the gamma factor at the real place.

Definition 6.3.1: The Completed Zeta Function

The *local gamma factor at the real place* is

$$\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right)$$

with Γ the gamma function of definition 6.1.1. The *completed zeta function* is the product

$$\Lambda(s) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \Gamma_{\mathbb{R}}(s) \zeta(s)$$

defined initially on the half-plane $\operatorname{Re}(s) > 1$, where both factors are holomorphic and $\zeta(s)$ is given by its Dirichlet series.

The notation $\Gamma_{\mathbb{R}}$ marks the factor as attached to the real place of $\mathbb{Q}$, the place corresponding to the ordinary absolute value, in contrast to the p -adic factors $(1 - p^{-s})^{-1}$ of the Euler product attached to the finite places (corollary 4.1.2). The completed function $\Lambda(s)$ multiplies the Euler product over the finite places by the one missing factor over the infinite place, and it is this complete product, not ζ alone, that enjoys the clean symmetry. The closely related *Riemann xi function*

$$\xi(s) = \frac{1}{2} s(s-1) \Lambda(s) = \frac{1}{2} s(s-1) \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s)$$

clears the two poles of Λ (located in theorem 6.3.3) by the elementary factor $\frac{1}{2}s(s-1)$, producing the entire function whose Hadamard product over its zeros drives the explicit formula of box 5.5.5; the factor $s(s-1)$ is itself invariant under $s \mapsto 1-s$, so ξ inherits the symmetry of Λ unchanged.

Example 6.3.2: The Completed Factor at $s = 2$

At $s = 2$ the gamma factor is $\Gamma_{\mathbb{R}}(2) = \pi^{-1} \Gamma(1) = \pi^{-1}$, using $\Gamma(1) = 1$ from example 6.1.5. Since $\zeta(2) = \pi^2/6$ (example 4.1.3), the completed value is

$$\Lambda(2) = \pi^{-1} \cdot \frac{\pi^2}{6} = \frac{\pi}{6}$$

The symmetry proved below forces $\Lambda(-1) = \Lambda(2) = \pi/6$ as well. This is a nontrivial constraint: the reflected point $s = -1$ lies outside the half-plane of definition 6.3.1, where neither $\zeta(-1)$ nor $\Lambda(-1)$ has yet been assigned a value, and the functional equation is precisely what defines them. Reading the constraint backward, $\Lambda(-1) = \Gamma_{\mathbb{R}}(-1) \zeta(-1) = \pi^{1/2} \Gamma(-\frac{1}{2}) \zeta(-1)$; with $\Gamma(-\frac{1}{2}) = -2\sqrt{\pi}$ from the recursion (6.1) and $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ (example 6.1.5), this reads $\pi^{1/2} \cdot (-2\sqrt{\pi}) \cdot \zeta(-1) = -2\pi \zeta(-1)$, and equating to $\pi/6$ gives $\zeta(-1) = -1/12$. The functional equation thus turns the elementary datum $\zeta(2) = \pi^2/6$ into the value $\zeta(-1) = -1/12$; this is the computation carried out systematically in section 6.4.

The motivating idea behind the gamma factor is the integral representation of example 2.4.2. The gamma function is the Mellin transform of e^{-t} , and rescaling its argument produces, for each fixed n , the Mellin transform of the Gaussian $e^{-\pi n^2 t}$. Summing over n assembles the gamma factor times $\zeta(s)$ as the Mellin transform of the theta series, and the theta functional equation then delivers the symmetry. This is Riemann's argument, carried out in full below.

Theorem 6.3.3: The Functional Equation of the Riemann Zeta Function

The completed zeta function extends to a meromorphic function on all of $\mathbb{C}$, holomorphic everywhere except for simple poles at $s = 0$ and $s = 1$, and satisfies the functional equation

$$\Lambda(s) = \Lambda(1 - s) \quad (6.14)$$

for all $s \in \mathbb{C}$. Equivalently,

$$\pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \pi^{-(1-s)/2} \Gamma\left(\frac{1-s}{2}\right) \zeta(1-s)$$

Proof:

The argument has four stages: the Mellin representation of Λ on $\operatorname{Re}(s) > 1$; the splitting of the Mellin integral at $t = 1$; the application of the theta functional equation to the integral over $(0, 1)$; and the reading-off of the symmetry, the continuation, and the poles from the resulting formula.

Step 1: The Mellin representation of Λ . Fix s with $\operatorname{Re}(s) > 1$. The starting point is the integral identity

$$\pi^{-s/2} \Gamma\left(\frac{s}{2}\right) n^{-s} = \int_0^\infty e^{-\pi n^2 t} t^{s/2-1} dt \quad (6.15)$$

valid for each integer $n \geq 1$. To see it, begin from the definition $\Gamma(s/2) = \int_0^\infty e^{-u} u^{s/2-1} du$ of definition 6.1.1, which converges since $\operatorname{Re}(s/2) > 0$, and substitute $u = \pi n^2 t$, so $du = \pi n^2 dt$ and $u^{s/2-1} = (\pi n^2)^{s/2-1} t^{s/2-1}$. This gives

$$\Gamma\left(\frac{s}{2}\right) = \int_0^\infty e^{-\pi n^2 t} (\pi n^2)^{s/2-1} t^{s/2-1} \cdot \pi n^2 dt = (\pi n^2)^{s/2} \int_0^\infty e^{-\pi n^2 t} t^{s/2-1} dt$$

and dividing by $(\pi n^2)^{s/2} = \pi^{s/2} n^s$ yields (6.15). Summing (6.15) over $n \geq 1$ and using $\sum_{n \geq 1} n^{-s} = \zeta(s)$ gives, on the left, $\pi^{-s/2} \Gamma(s/2) \zeta(s) = \Lambda(s)$. On the right the sum of the integrals is the integral of the sum:

$$\Lambda(s) = \sum_{n=1}^\infty \int_0^\infty e^{-\pi n^2 t} t^{s/2-1} dt = \int_0^\infty \left(\sum_{n=1}^\infty e^{-\pi n^2 t} \right) t^{s/2-1} dt = \int_0^\infty \psi(t) t^{s/2-1} dt \quad (6.16)$$

where $\psi(t) = \sum_{n \geq 1} e^{-\pi n^2 t} = \frac{1}{2}(\theta(t) - 1)$ by (6.10). The interchange of sum and integral is justified by Tonelli's theorem ([8]) applied to the nonnegative integrand $e^{-\pi n^2 t} t^{\operatorname{Re}(s)/2-1}$: the iterated integral $\sum_n \int_0^\infty e^{-\pi n^2 t} t^{\operatorname{Re}(s)/2-1} dt = \pi^{-\operatorname{Re}(s)/2} \Gamma(\operatorname{Re}(s)/2) \sum_n n^{-\operatorname{Re}(s)}$ is finite because $\operatorname{Re}(s) > 1$ makes the last series converge, so Tonelli permits the interchange for the absolute values, and the integrand here is already nonnegative. This establishes the Mellin representation (6.16) of Λ on $\operatorname{Re}(s) > 1$.

Step 2: Splitting at $t = 1$. Break the integral (6.16) at $t = 1$:

$$\Lambda(s) = \int_0^1 \psi(t) t^{s/2-1} dt + \int_1^\infty \psi(t) t^{s/2-1} dt \quad (6.17)$$

The second integral converges for every $s \in \mathbb{C}$ and defines an entire function. On $[1, \infty)$ the bound $\psi(t) \leq e^{-\pi t}/(1 - e^{-\pi t})$ established in section 6.2 shows $\psi(t)$ decays faster than any power of t , so for every fixed s the integrand $\psi(t)t^{s/2-1}$ is dominated by $\psi(t)t^{\operatorname{Re}(s)/2-1}$, integrable on $[1, \infty)$; holomorphy in s follows from differentiation under the integral sign, justified on compact subsets by the same domination ([16]). The first integral, over $(0, 1)$, is where the theta functional equation enters, because $\psi(t)$ is not small as $t \rightarrow 0^+$.

Step 3: The theta functional equation on $(0, 1)$. The functional equation $\theta(1/t) = \sqrt{t}\theta(t)$ of theorem 6.2.4, written in terms of $\psi = \frac{1}{2}(\theta - 1)$, relates $\psi(t)$ to $\psi(1/t)$. From $\theta(t) = 2\psi(t) + 1$ and $\theta(1/t) = 2\psi(1/t) + 1$, the relation $\theta(1/t) = \sqrt{t}\theta(t)$ reads $2\psi(1/t) + 1 = \sqrt{t}(2\psi(t) + 1)$, and solving for $\psi(t)$ gives

$$\psi(t) = \frac{1}{\sqrt{t}}\psi\left(\frac{1}{t}\right) + \frac{1}{2\sqrt{t}} - \frac{1}{2} \quad (6.18)$$

Now substitute $t \mapsto 1/t$ in the first integral of (6.17). With $t = 1/r$, $dt = -r^{-2}dr$, the range $t \in (0, 1)$ becomes $r \in (1, \infty)$, and $t^{s/2-1} = r^{1-s/2}$, so

$$\int_0^1 \psi(t)t^{s/2-1}dt = \int_1^\infty \psi\left(\frac{1}{r}\right)r^{1-s/2}r^{-2}dr = \int_1^\infty \psi\left(\frac{1}{r}\right)r^{-s/2-1}dr$$

By (6.18) with t replaced by $1/r$, namely $\psi(1/r) = \sqrt{r}\psi(r) + \frac{1}{2}\sqrt{r} - \frac{1}{2}$, the integrand becomes

$$\psi\left(\frac{1}{r}\right)r^{-s/2-1} = \left(\sqrt{r}\psi(r) + \frac{\sqrt{r}}{2} - \frac{1}{2}\right)r^{-s/2-1} = \psi(r)r^{-(s-1)/2-1} + \frac{1}{2}r^{-(s+1)/2} - \frac{1}{2}r^{-s/2-1}$$

the first power following from $\frac{1}{2} - s/2 - 1 = -(s-1)/2 - 1$. The two elementary terms integrate explicitly over $[1, \infty)$, convergently when $\operatorname{Re}(s) > 1$:

$$\frac{1}{2} \int_1^\infty r^{-(s+1)/2}dr = \frac{1}{2} \cdot \frac{1}{(s+1)/2-1} = \frac{1}{s-1}, \quad \frac{1}{2} \int_1^\infty r^{-s/2-1}dr = \frac{1}{2} \cdot \frac{1}{s/2} = \frac{1}{s}$$

Therefore

$$\int_0^1 \psi(t)t^{s/2-1}dt = \int_1^\infty \psi(r)r^{(1-s)/2-1}dr + \frac{1}{s-1} - \frac{1}{s} \quad (6.19)$$

where the surviving integral is written with the exponent $-(s-1)/2-1 = (1-s)/2-1$.

Step 4: Symmetry, continuation, and poles. Substituting (6.19) into (6.17) gives

$$\Lambda(s) = \frac{1}{s-1} - \frac{1}{s} + \int_1^\infty \psi(t)\left(t^{s/2-1} + t^{(1-s)/2-1}\right)dt \quad (6.20)$$

derived for $\operatorname{Re}(s) > 1$, where every step was valid. The right-hand side, however, makes sense for every $s \in \mathbb{C}$. The integral $\int_1^\infty \psi(t)(t^{s/2-1} + t^{(1-s)/2-1})dt$ converges for all s and is entire, by the rapid decay of ψ on $[1, \infty)$ exactly as in Step 2, applied now to each of the two powers; the remaining terms $1/(s-1) - 1/s$ are meromorphic on $\mathbb{C}$ with simple poles at $s = 1$ and $s = 0$. Hence the right-hand side of (6.20) is a meromorphic function on $\mathbb{C}$, holomorphic except for simple poles at $s = 0$ and $s = 1$, agreeing with $\Lambda(s)$ on $\operatorname{Re}(s) > 1$; by the identity theorem ([16]) it is the meromorphic continuation of Λ to all of $\mathbb{C}$. The residues are $\operatorname{Res}_{s=1}\Lambda = 1$ and $\operatorname{Res}_{s=0}\Lambda = -1$, the coefficients of $1/(s-1)$ and $1/s$.

The functional equation is now read directly off (6.20). The substitution $s \mapsto 1 - s$ fixes the right-hand side: it interchanges the two powers $t^{s/2-1}$ and $t^{(1-s)/2-1}$ inside the integral, leaving the integral unchanged, and it sends

$$\frac{1}{s-1} - \frac{1}{s} \mapsto \frac{1}{(1-s)-1} - \frac{1}{1-s} = \frac{1}{-s} - \frac{1}{1-s} = -\frac{1}{s} + \frac{1}{s-1}$$

which is the same pair of terms. Both the integral and the rational part are therefore invariant under $s \mapsto 1 - s$, so $\Lambda(s) = \Lambda(1 - s)$ for all s , which is (6.14). The equivalent form follows by writing out $\Lambda(s) = \pi^{-s/2}\Gamma(s/2)\zeta(s)$ and $\Lambda(1 - s) = \pi^{-(1-s)/2}\Gamma((1-s)/2)\zeta(1 - s)$ from definition 6.3.1, completing the proof.

The symmetry expressed by (6.14) has its axis at the line $\operatorname{Re}(s) = 1/2$, the fixed-point set of the involution $s \mapsto 1 - s$. The completed function takes equal values at any two points reflected across this line, and the line itself is where a point coincides with its mirror image. This is the analytic origin of the central role of $\operatorname{Re}(s) = 1/2$: it is not imposed on the theory but forced by the functional equation, descending from the fixed point $t = 1$ of the theta involution $t \mapsto 1/t$ through the half-power $t^{s/2}$ of the Mellin pairing in (6.16). The quadratic exponent n^2 in $e^{-\pi n^2 t}$ is what makes that pairing a pairing against $t^{s/2}$ rather than t^s , and the resulting halving is what centers the symmetry at $s = 1/2$ rather than at an integer; the Riemann Hypothesis (?? 5.5.6) places every nontrivial zero on this axis, and (6.14) is what singles the axis out.

The two poles of Λ sit symmetrically about the axis, at $s = 1$ and $s = 0$, and the symmetry exchanges them. The pole at $s = 1$ is the pole of ζ itself (corollary 4.2.2): the gamma factor $\Gamma_{\mathbb{R}}(1) = \pi^{-1/2}\Gamma(1/2) = 1$ is finite and nonzero there, so Λ inherits the simple pole of ζ with the same residue, $\operatorname{Res}_{s=1}\Lambda = \operatorname{Res}_{s=1}\zeta = 1$, consistent with Step 4. The pole at $s = 0$, by contrast, is not a pole of ζ , which is holomorphic at the origin; it is the pole of the gamma factor, since $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$ has a simple pole at $s = 0$ inherited from the pole of Γ at 0 (theorem 6.1.2). Under $s \leftrightarrow 1 - s$ the analytic pole of ζ at $s = 1$ is matched with the archimedean pole of the gamma factor at $s = 0$: the two poles are reflections of one another, one arithmetic and one transcendental, and the functional equation is what identifies them.

The gamma factor $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$ is the factor at the real place of $\mathbb{Q}$, and its appearance here is the first instance of a general pattern. A completed L -function is the product of the Dirichlet-series Euler factors over the finite places with an archimedean factor over the infinite place, built from $\Gamma_{\mathbb{R}}$ or its complex analogue, and it is the completed product, never the bare Dirichlet series, that satisfies a clean functional equation relating s to $1 - s$ (or to $k - s$ in the appropriate normalization). The completed Dirichlet L -functions, taken up in a later chapter of this book, carry exactly such a factor, chosen according to the parity of the character, and the completed Dedekind zeta functions, also treated later, carry one factor $\Gamma_{\mathbb{R}}$ for each real place and one factor $\Gamma_{\mathbb{C}}$ for each complex place of the number field. The single factor $\Gamma_{\mathbb{R}}$ attached to ζ here, the zeta function of $\mathbb{Q}$, is the simplest case of this product-over-places structure, and the functional equation of theorem 6.3.3 is its prototype.

With Λ continued to all of $\mathbb{C}$, the continuation of ζ itself follows by dividing out the gamma factor, and the symmetric functional equation (6.14) converts into the asymmetric form relating $\zeta(s)$ directly to $\zeta(1 - s)$.

Corollary 6.3.4: Continuation of ζ to the Whole Plane

The Riemann zeta function continues to a meromorphic function on all of $\mathbb{C}$ via

$$\zeta(s) = \frac{\pi^{s/2} \Lambda(s)}{\Gamma(s/2)} \quad (6.21)$$

whose only pole is the simple pole at $s = 1$, with residue 1. In asymmetric form the functional equation reads

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s) \quad (6.22)$$

for all $s \in \mathbb{C}$.

Proof:

The continuation comes from solving definition 6.3.1 for ζ ; the asymmetric form comes from substituting the symmetric functional equation and converting the gamma factors by reflection and duplication.

Step 1: The continuation and its single pole. Solving $\Lambda(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$ for ζ gives (6.21). The factor $\pi^{s/2}$ is entire and nowhere zero, and Λ is meromorphic on $\mathbb{C}$ with simple poles only at $s = 0, 1$ by theorem 6.3.3, so the right-hand side of (6.21) is meromorphic on $\mathbb{C}$. Its poles can only occur where Λ has a pole, and its zeros only where $1/\Gamma(s/2)$ has a zero, that is, where $\Gamma(s/2)$ has a pole. At $s = 1$ the factor $\Gamma(s/2) = \Gamma(1/2) = \sqrt{\pi}$ is finite and nonzero, so the simple pole of Λ at $s = 1$ passes to ζ ; the residue is $\text{Res}_{s=1} \zeta = \pi^{1/2} \text{Res}_{s=1} \Lambda / \Gamma(1/2) = \pi^{1/2} \cdot 1 / \sqrt{\pi} = 1$, recovering corollary 4.2.2. At $s = 0$ the factor $1/\Gamma(s/2)$ has a zero, because $\Gamma(s/2)$ has a pole there (theorem 6.1.2); this zero cancels the simple pole of Λ at $s = 0$, so ζ is holomorphic at the origin. No other poles of Λ exist, and $\Gamma(s/2) \neq 0$ everywhere by theorem 6.1.2, so $1/\Gamma(s/2)$ contributes no poles; the only pole of ζ is the simple pole at $s = 1$ with residue 1.

Step 2: Conversion to the asymmetric form. The symmetric equation (6.14) solved for $\zeta(s)$ reads

$$\zeta(s) = \pi^{s/2} \pi^{-(1-s)/2} \frac{\Gamma(\frac{1-s}{2})}{\Gamma(\frac{s}{2})} \zeta(1-s) = \pi^{s-1/2} \frac{\Gamma(\frac{1-s}{2})}{\Gamma(\frac{s}{2})} \zeta(1-s) \quad (6.23)$$

using $\frac{s}{2} - \frac{1-s}{2} = s - \frac{1}{2}$. The ratio of gamma values is reduced to elementary functions in two moves. First, the duplication formula (6.6) of theorem 6.1.4 at $z = s/2$, namely $\Gamma(z)\Gamma(z + \frac{1}{2}) = 2^{1-2z} \sqrt{\pi} \Gamma(2z)$, gives

$$\Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s+1}{2}\right) = 2^{1-s} \sqrt{\pi} \Gamma(s) \quad (6.24)$$

Second, the reflection formula (6.3) of theorem 6.1.3 at $z = (1+s)/2$, using $\sin(\pi \frac{1+s}{2}) = \cos(\frac{\pi s}{2})$, gives

$$\Gamma\left(\frac{1-s}{2}\right) \Gamma\left(\frac{1+s}{2}\right) = \frac{\pi}{\sin(\pi \frac{1+s}{2})} = \frac{\pi}{\cos(\frac{\pi s}{2})} \quad (6.25)$$

since $\Gamma(\frac{1-s}{2}) = \Gamma(1 - \frac{1+s}{2})$. Dividing (6.25) by (6.24) cancels the common factor $\Gamma(\frac{1+s}{2}) = \Gamma(\frac{s+1}{2})$ and isolates the required ratio:

$$\frac{\Gamma(\frac{1-s}{2})}{\Gamma(\frac{s}{2})} = \frac{\pi}{\cos(\frac{\pi s}{2})} \cdot \frac{1}{2^{1-s} \sqrt{\pi} \Gamma(s)} = \frac{\sqrt{\pi} 2^{s-1}}{\cos(\frac{\pi s}{2}) \Gamma(s)}$$

Substituting this into (6.23) and collecting the powers of π , with $\pi^{s-1/2} \cdot \pi^{1/2} = \pi^s$, gives

$$\zeta(s) = \frac{\pi^s 2^{s-1}}{\cos(\frac{\pi s}{2}) \Gamma(s)} \zeta(1-s) \quad (6.26)$$

It remains to clear $\Gamma(s)$ from the denominator. The reflection formula (6.3) gives $\Gamma(s)\Gamma(1-s) = \pi/\sin(\pi s)$, hence $1/\Gamma(s) = \sin(\pi s)\Gamma(1-s)/\pi$, and the double-angle identity $\sin(\pi s) = 2\sin(\frac{\pi s}{2})\cos(\frac{\pi s}{2})$ turns (6.26) into

$$\zeta(s) = \pi^s 2^{s-1} \cdot \frac{1}{\cos(\frac{\pi s}{2})} \cdot \frac{2\sin(\frac{\pi s}{2})\cos(\frac{\pi s}{2})\Gamma(1-s)}{\pi} \zeta(1-s)$$

The factor $\cos(\frac{\pi s}{2})$ cancels, and $2^{s-1} \cdot 2 = 2^s$ while $\pi^s/\pi = \pi^{s-1}$, leaving

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

which is (6.22). The identity was derived from the meromorphic equation (6.14) through identities valid wherever both sides are defined, and both sides of (6.22) are meromorphic on $\mathbb{C}$, so it holds for all s by the identity theorem ([16]), as we sought to show.

The asymmetric form (6.22) is the version of the functional equation most directly applied to values of ζ , since it expresses $\zeta(s)$ for s in the left half-plane in terms of $\zeta(1-s)$ for $1-s$ in the right half-plane, where the Dirichlet series converges. The three elementary factors record what the symmetric form hides inside the gamma factors. The factor $\zeta(1-s)$ supplies the values to be reflected; the factor $\Gamma(1-s)$ supplies the analytic continuation, contributing poles at the positive integers $s = 1, 2, 3, \dots$ that must be cancelled or accounted for; and the factor $\sin(\pi s/2)$ is the source of the trivial zeros, vanishing at the even integers $s = 0, \pm 2, \pm 4, \dots$. The interplay of these factors at the negative integers is exactly the computation of section 6.4: at $s = -2k$ the factor $\sin(\pi s/2)$ vanishes while the others stay finite, forcing $\zeta(-2k) = 0$, whereas at $s = 1 - 2k$ all three elementary factors are finite and nonzero, since $\sin(\pi(1-2k)/2) = \cos(\pi k) = (-1)^k$ and $\Gamma(1 - (1-2k)) = \Gamma(2k) = (2k-1)!$, so the equation expresses $\zeta(1-2k)$ as the nonzero rational value that, reflected, yields $\zeta(2k)$.

Example 6.3.5: The Functional Equation at $s = 2$

The asymmetric equation (6.22) at $s = 2$ reproduces the value $\zeta(-1) = -1/12$ from the known value $\zeta(2) = \pi^2/6$, the same computation as in example 6.3.2 carried out through the elementary factors. Setting $s = -1$ in (6.22), so that $1-s = 2$, gives

$$\zeta(-1) = 2^{-1} \pi^{-2} \sin\left(-\frac{\pi}{2}\right) \Gamma(2) \zeta(2)$$

The factors evaluate as $2^{-1} = \frac{1}{2}$, π^{-2} , $\sin(-\pi/2) = -1$, $\Gamma(2) = 1$ (example 6.1.5), and $\zeta(2) = \pi^2/6$ (example 4.1.3). Multiplying,

$$\zeta(-1) = \frac{1}{2} \cdot \frac{1}{\pi^2} \cdot (-1) \cdot 1 \cdot \frac{\pi^2}{6} = -\frac{1}{12}$$

the powers of π cancelling exactly, as they must for the rational value. The same equation read at an even negative integer shows the trivial zeros directly: at $s = -2$, the factor $\sin(\pi s/2) = \sin(-\pi) = 0$, while $2^{-2}\pi^{-3}$, $\Gamma(3) = 2$, and $\zeta(3)$ are all finite, so $\zeta(-2) = 0$. The single factor $\sin(\pi s/2)$ carries the vanishing.

6.4 Trivial Zeros and Special Values

The functional equation of section 6.3 continued $\zeta(s)$ to the whole plane and tied $\zeta(s)$ to $\zeta(1-s)$; this section reads off what that symmetry forces at the integers. The completed function $\Lambda(s) = \pi^{-s/2}\Gamma(s/2)\zeta(s)$ of definition 6.3.1 is holomorphic and finite away from its two poles at $s = 0$ and $s = 1$, while the gamma factor $\Gamma(s/2)$ carries poles of its own at the negative even integers. Where the gamma factor blows up but Λ stays finite, the only remaining factor, ζ , must vanish: these are the trivial zeros of ζ at $s = -2, -4, -6, \dots$. The same accounting, applied at the negative odd integers and the origin, pins ζ to the Bernoulli numbers of definition 1.5.1, giving the rational values $\zeta(0) = -1/2$ and $\zeta(1-2k) = -B_{2k}/(2k)$.

Reflecting these rational values across the line $\operatorname{Re}(s) = 1/2$ through the functional equation produces the values at the positive even integers, expressing each $\zeta(2k)$ as a rational multiple of π^{2k} . In particular $\zeta(2) = \pi^2/6$ and $\zeta(4) = \pi^4/90$, discharging the evaluations foreshadowed in example 4.1.3: what was cited there to external analysis is now a consequence of the functional equation, derived from the rational value $\zeta(1-2k)$ by reflection. The contrast with the odd values, which the same reflection sends to the vanishing $\zeta(-2k)$ and so leaves without any closed form, is the structural reason for the difficulty noted in chapter 4.

The trivial zeros come first, read directly off the pole structure of the completed function.

Corollary 6.4.1: The Trivial Zeros

For every integer $k \geq 1$,

$$\zeta(-2k) = 0$$

These are the *trivial zeros* of the Riemann zeta function. Together with the points of the critical strip $0 \leq \operatorname{Re}(s) \leq 1$, they account for every zero of ζ : outside the closed strip, the only zeros of ζ are the negative even integers $s = -2, -4, -6, \dots$.

Proof:

The argument reads the vanishing of $\zeta(-2k)$ off definition 6.3.1 and then locates all zeros outside the strip.

Step 1: Vanishing at the negative even integers. Fix $k \geq 1$ and set $s = -2k$. By theorem 6.3.3, the completed function $\Lambda(s) = \pi^{-s/2}\Gamma(s/2)\zeta(s)$ is meromorphic on $\mathbb{C}$ with its only poles the simple poles at $s = 0$ and $s = 1$; since $-2k \notin \{0, 1\}$, the value $\Lambda(-2k)$ is finite. Solving definition 6.3.1 for ζ as in (6.21) gives

$$\zeta(-2k) = \frac{\pi^{-k} \Lambda(-2k)}{\Gamma(-k)}$$

where $\Gamma(s/2) = \Gamma(-k)$ at $s = -2k$ and $\pi^{s/2} = \pi^{-k}$. The factor π^{-k} is finite, and $\Lambda(-2k)$ is finite by the preceding sentence. The denominator $\Gamma(-k)$, on the other hand, is infinite: Γ has a simple pole at the non-positive integer $-k$ (theorem 6.1.2). Equivalently, $1/\Gamma(s/2)$ has a zero at $s = -2k$, since $1/\Gamma$ is entire and vanishes exactly at the poles of Γ . A finite numerator times the zero $1/\Gamma(-k) = 0$ gives

$$\zeta(-2k) = \pi^{-k} \Lambda(-2k) \cdot \frac{1}{\Gamma(-k)} = \pi^{-k} \Lambda(-2k) \cdot 0 = 0$$

as claimed. The mechanism is the transfer recorded after theorem 6.1.2: the pole of the archimedean factor $\Gamma(s/2)$ forces the arithmetic factor ζ to vanish, so that the product Λ stays finite.

Step 2: These are the only zeros outside the strip. Suppose s_0 is a zero of ζ with $\operatorname{Re}(s_0) \notin [0, 1]$. The Euler product $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ of corollary 4.1.2 converges on $\operatorname{Re}(s) > 1$ and is a product of nonzero factors, none of which vanishes, so ζ has no zero with $\operatorname{Re}(s) > 1$; hence $\operatorname{Re}(s_0) < 0$. On the half-plane $\operatorname{Re}(s) < 0$ the asymmetric functional equation (6.22) of corollary 6.3.4 applies, reading

$$\zeta(s_0) = 2^{s_0} \pi^{s_0-1} \sin\left(\frac{\pi s_0}{2}\right) \Gamma(1-s_0) \zeta(1-s_0)$$

The reflected point has $\operatorname{Re}(1-s_0) > 1$, so $\zeta(1-s_0) \neq 0$ by the Euler-product argument just given. The factor $\Gamma(1-s_0)$ is finite and nonzero: $1-s_0$ has positive real part, away from the poles of Γ , and Γ has no zeros (theorem 6.1.2). The factors 2^{s_0} and π^{s_0-1} are nonzero. The product can therefore vanish only through the factor $\sin(\pi s_0/2)$, which is zero precisely when $s_0/2 \in \mathbb{Z}$, that is, when s_0 is an even integer. Combined with $\operatorname{Re}(s_0) < 0$, this forces $s_0 \in \{-2, -4, -6, \dots\}$. Every zero outside the closed strip is thus a negative even integer, and Step 1 shows each negative even integer is indeed a zero, completing the proof.

The trivial zeros are exactly the zeros the functional equation manufactures from the pole structure of the gamma factor; they carry no arithmetic content and are accounted for completely. What the corollary leaves open is the location of the remaining zeros, those inside the critical strip $0 \leq \operatorname{Re}(s) \leq 1$. The Euler product rules out zeros with $\operatorname{Re}(s) > 1$, the functional equation reflects this to rule out all zeros with $\operatorname{Re}(s) < 0$ except the trivial ones, and the non-vanishing on the line $\operatorname{Re}(s) = 1$ of theorem 4.4.2 (with its reflection $\operatorname{Re}(s) = 0$) pushes the boundary inward; the zeros that survive, the *nontrivial* zeros, all lie in the open strip $0 < \operatorname{Re}(s) < 1$. The Riemann Hypothesis (?? 5.5.6) asserts that they lie on the central line $\operatorname{Re}(s) = 1/2$, and it is the trivial zeros having been disposed of here that lets the conjecture be stated about the strip alone.

The same division of Λ by the gamma factor that produced the trivial zeros produces, at the negative *odd* integers and at the origin, finite values rather than zeros, because there the gamma factor $\Gamma(s/2)$ is finite. Those values are rational, and they are exactly the Bernoulli numbers.

Theorem 6.4.2: Values at the Negative Integers

The Riemann zeta function takes the value

$$\zeta(0) = -\frac{1}{2}$$

at the origin, and for every integer $k \geq 1$,

$$\zeta(1-2k) = -\frac{B_{2k}}{2k} \tag{6.27}$$

where B_{2k} is the Bernoulli number of definition 1.5.1. The remaining negative integer values are the trivial zeros $\zeta(-2k) = 0$ of corollary 6.4.1.

Proof:

The value at the origin is computed from the Laurent data of Λ and Γ at $s = 0$; the values at the negative odd integers come from the integral representation of theorem 2.4.5, deformed to a loop contour on which the Bernoulli numbers appear as residues.

Step 1: The value $\zeta(0) = -1/2$. Both $\Lambda(s)$ and $\Gamma(s/2)$ have simple poles at $s = 0$, and their quotient through (6.21) is finite there, so $\zeta(0)$ is computed as a ratio of residues. From (6.21),

$$\zeta(s) = \pi^{s/2} \frac{\Lambda(s)}{\Gamma(s/2)}$$

and ζ is holomorphic at $s = 0$ by corollary 6.3.4, so $\zeta(0) = \lim_{s \rightarrow 0} \zeta(s)$. The factor $\pi^{s/2} \rightarrow 1$ as $s \rightarrow 0$. For the quotient $\Lambda(s)/\Gamma(s/2)$, multiply numerator and denominator by s and take limits separately. The numerator has the simple pole $\text{Res}_{s=0} \Lambda = -1$ recorded in the proof of theorem 6.3.3, so $\lim_{s \rightarrow 0} s \Lambda(s) = -1$. For the denominator, the recursion (6.1) gives $\Gamma(s/2) = \Gamma(s/2+1)/(s/2) = (2/s) \Gamma(s/2+1)$, and $\Gamma(s/2+1) \rightarrow \Gamma(1) = 1$ as $s \rightarrow 0$ (example 6.1.5), so $\lim_{s \rightarrow 0} s \Gamma(s/2) = 2$. Therefore

$$\zeta(0) = \lim_{s \rightarrow 0} \pi^{s/2} \frac{s \Lambda(s)}{s \Gamma(s/2)} = 1 \cdot \frac{-1}{2} = -\frac{1}{2}$$

the cancellation of the factor s being legitimate because both one-sided limits exist and the denominator limit is nonzero.

Step 2: A loop representation of ζ . The continuation of corollary 6.3.4 is recast as a contour integral, starting from the integral representation $\Gamma(s)\zeta(s) = \int_0^\infty t^{s-1}/(e^t - 1) dt$ of theorem 2.4.5, valid for $\text{Re}(s) > 1$. Let H_ε be the *Hankel contour*: the path coming in from $+\infty$ along the upper side of the positive real axis, traversing the circle $|z| = \varepsilon$ once counterclockwise around the origin, and returning to $+\infty$ along the lower side. On H_ε the function $(-z)^{s-1} = e^{(s-1)\log(-z)}$ is defined with the branch of $\log(-z)$ that is real for $z < 0$, equivalently $\arg(-z) \in (-\pi, \pi)$; on the upper edge $\log(-z) = \log|z| - i\pi$ and on the lower edge $\log(-z) = \log|z| + i\pi$. The claim is the *Riemann–Hankel representation*

$$\zeta(s) = -\frac{\Gamma(1-s)}{2\pi i} \int_{H_\varepsilon} \frac{(-z)^{s-1}}{e^z - 1} dz \quad (6.28)$$

valid for all $s \in \mathbb{C}$ (the integral is independent of $\varepsilon \in (0, 2\pi)$, since the integrand is holomorphic in the punctured disc $0 < |z| < 2\pi$ between two such contours, and defines an entire function of s). To establish (6.28), fix $\text{Re}(s) > 1$ and shrink the radius $\varepsilon \rightarrow 0$. On the circle $|z| = \varepsilon$ the integrand is $O(\varepsilon^{\text{Re}(s)-1} \cdot \varepsilon^{-1}) = O(\varepsilon^{\text{Re}(s)-2})$ in modulus, since $1/(e^z - 1) = O(1/|z|)$ near the origin, and the circle has length $2\pi\varepsilon$; the contribution is $O(\varepsilon^{\text{Re}(s)-1}) \rightarrow 0$ as $\varepsilon \rightarrow 0$ because $\text{Re}(s) > 1$. The two straight edges survive. On the upper edge, $z = t$ with t running from ∞ down to 0 and $(-z)^{s-1} = e^{(s-1)(\log t - i\pi)} = t^{s-1} e^{-i\pi(s-1)}$; on the lower edge, $z = t$ with t running from 0 up to ∞ and $(-z)^{s-1} = t^{s-1} e^{i\pi(s-1)}$. The upper edge is traversed from ∞ to 0, contributing the negative of $e^{-i\pi(s-1)} \int_0^\infty t^{s-1}/(e^t - 1) dt$, while the lower edge contributes $+e^{i\pi(s-1)} \int_0^\infty t^{s-1}/(e^t - 1) dt$; their sum is

$$\int_{H_\varepsilon} \frac{(-z)^{s-1}}{e^z - 1} dz \xrightarrow{\varepsilon \rightarrow 0} (e^{i\pi(s-1)} - e^{-i\pi(s-1)}) \int_0^\infty \frac{t^{s-1}}{e^t - 1} dt = 2i \sin(\pi(s-1)) \Gamma(s) \zeta(s)$$

using theorem 2.4.5 for the surviving real integral. Now $\sin(\pi(s-1)) = -\sin(\pi s)$, and the reflection formula (6.3) gives $\sin(\pi s) \Gamma(s) = \pi / \Gamma(1-s)$, so the limiting value is

$$\int_{H_\varepsilon} \frac{(-z)^{s-1}}{e^z - 1} dz = 2i \sin(\pi(s-1)) \Gamma(s) \zeta(s) = -2i \sin(\pi s) \Gamma(s) \zeta(s) = -\frac{2\pi i \zeta(s)}{\Gamma(1-s)}$$

for $\operatorname{Re}(s) > 1$, which rearranges to (6.28). The integral over H_ε converges for every $s \in \mathbb{C}$, because away from the origin the contour stays a bounded distance from the poles of $1/(e^z - 1)$ at $z = 2\pi i n$ and the factor $1/(e^z - 1)$ decays as $\operatorname{Re}(z) \rightarrow +\infty$, so it defines an entire function of s ; multiplying by the meromorphic factor $-\Gamma(1-s)/(2\pi i)$ gives a meromorphic function agreeing with ζ on $\operatorname{Re}(s) > 1$, hence equal to the continued ζ on all of $\mathbb{C}$ by the identity theorem ([16]). This is (6.28).

Step 3: Collapse to a residue at a negative odd integer. Set $s = 1 - 2k$ in (6.28). The exponent $s - 1 = -2k$ is an even negative integer, so $(-z)^{s-1} = (-z)^{-2k} = z^{-2k}$ is single-valued, the branch ambiguity disappears, and the two edges of H_ε cancel: their integrands are now identical functions of t traversed in opposite directions. Only the circle $|z| = \varepsilon$ remains, traversed once counterclockwise, so the contour integral equals $2\pi i$ times the residue of the integrand at $z = 0$:

$$\int_{H_\varepsilon} \frac{z^{-2k}}{e^z - 1} dz = 2\pi i \operatorname{Res}_{z=0} \frac{z^{-2k}}{e^z - 1}$$

The residue is read from the generating function of definition 1.5.1. Writing $\frac{1}{e^z - 1} = \frac{1}{z} \cdot \frac{z}{e^z - 1} = \frac{1}{z} \sum_{m=0}^{\infty} \frac{B_m}{m!} z^m$, the integrand is

$$\frac{z^{-2k}}{e^z - 1} = z^{-2k-1} \sum_{m=0}^{\infty} \frac{B_m}{m!} z^m = \sum_{m=0}^{\infty} \frac{B_m}{m!} z^{m-2k-1}$$

whose residue, the coefficient of z^{-1} , is the term with $m = 2k$, namely $B_{2k}/(2k)!$. Substituting $s = 1 - 2k$ into the representation (6.28), with $\Gamma(1-s) = \Gamma(2k) = (2k-1)!$ (example 6.1.5) and the minus sign carried by (6.28),

$$\zeta(1-2k) = -\frac{(2k-1)!}{2\pi i} \cdot 2\pi i \frac{B_{2k}}{(2k)!} = -(2k-1)! \frac{B_{2k}}{(2k)!} = -\frac{B_{2k}}{2k}$$

the cancellation of $2\pi i$ against $2\pi i$ and of $(2k-1)!$ against $(2k)! = 2k \cdot (2k-1)!$ leaving the rational value, which is (6.27).

Step 4: Consistency check via the functional equation. The value (6.27) is confirmed independently from the asymmetric functional equation (6.22), which at $s = 1 - 2k$ reads, with $1 - s = 2k$,

$$\zeta(1-2k) = 2^{1-2k} \pi^{-2k} \sin\left(\frac{\pi(1-2k)}{2}\right) \Gamma(2k) \zeta(2k)$$

Here $\sin(\pi(1-2k)/2) = \sin(\pi/2 - \pi k) = \cos(\pi k) = (-1)^k$, and $\Gamma(2k) = (2k-1)!$. For $k = 1$ this gives $\zeta(-1) = 2^{-1} \pi^{-2} \cdot (-1) \cdot 1 \cdot \zeta(2) = -\frac{1}{2} \pi^{-2} \cdot (\pi^2/6) = -1/12$, matching $-B_2/(2 \cdot 1) = -(1/6)/2 = -1/12$ from (6.27). The general identity (6.27) is consistent with (6.22) for every k , as theorem 6.4.3 makes precise by inverting this relation, completing the proof.

The values $\zeta(1 - 2k) = -B_{2k}/(2k)$ are the arithmetic core of the negative-integer behavior. They are rational, and they are the same rational numbers that govern sums of powers and the Euler–Maclaurin expansion of chapter 1, which is the first hint that the zeta function at negative integers carries the combinatorics of the Bernoulli numbers rather than the transcendence of π . The vanishing of the odd-indexed Bernoulli numbers $B_3 = B_5 = \cdots = 0$ (proposition 1.5.2) is consistent with the picture: (6.27) involves only the even-indexed B_{2k} , never the odd ones, so the formula never references a vanishing Bernoulli number, and the trivial zeros at the negative even integers are a separate phenomenon coming from the gamma factor, not from any vanishing of B . The value $\zeta(0) = -1/2$ fits the pattern as the case “ $k = 0$ ” in the sense that $-B_0/0$ is replaced by the residue computation of Step 1, with $B_0 = 1$ entering through $\Gamma(s/2 + 1) \rightarrow 1$.

The functional equation now runs in the opposite direction. Having pinned ζ at the negative odd integers $1 - 2k$ to the rational values $-B_{2k}/(2k)$, reflecting across $\text{Re}(s) = 1/2$ pins ζ at the positive even integers $2k$, and the reflection introduces exactly the power of π that turns the rational value into the transcendental one.

Theorem 6.4.3: Values at the Positive Even Integers

For every integer $k \geq 1$,

$$\zeta(2k) = (-1)^{k+1} \frac{B_{2k} (2\pi)^{2k}}{2 (2k)!} \quad (6.29)$$

where B_{2k} is the Bernoulli number of definition 1.5.1. In particular $\zeta(2) = \pi^2/6$ and $\zeta(4) = \pi^4/90$, the values foreshadowed in example 4.1.3.

Proof:

The asymmetric functional equation (6.22) of corollary 6.3.4, evaluated at $s = 1 - 2k$, relates $\zeta(1 - 2k)$ to $\zeta(2k)$; substituting the value of $\zeta(1 - 2k)$ from theorem 6.4.2 and solving for $\zeta(2k)$ yields (6.29). The elementary factors in (6.22) are exactly those produced from the gamma factors of the symmetric equation (6.14) by the reflection formula (6.3) of theorem 6.1.3 and the duplication formula (6.6) of theorem 6.1.4, carried out in the proof of corollary 6.3.4; the computation below uses the asymmetric form directly, so those two identities enter through it. The equation (6.22) at $s = 1 - 2k$, with $1 - s = 2k$, reads

$$\zeta(1 - 2k) = 2^{1-2k} \pi^{-2k} \sin\left(\frac{\pi(1 - 2k)}{2}\right) \Gamma(2k) \zeta(2k) \quad (6.30)$$

The trigonometric factor simplifies by the addition formula:

$$\sin\left(\frac{\pi(1 - 2k)}{2}\right) = \sin\left(\frac{\pi}{2} - \pi k\right) = \cos(\pi k) = (-1)^k$$

and the gamma factor is $\Gamma(2k) = (2k - 1)!$ by example 6.1.5, since $2k$ is a positive integer. Substituting these and the value $\zeta(1 - 2k) = -B_{2k}/(2k)$ of (6.27) into (6.30),

$$-\frac{B_{2k}}{2k} = 2^{1-2k} \pi^{-2k} (-1)^k (2k - 1)! \zeta(2k)$$

Solving for $\zeta(2k)$ requires dividing by the factor $2^{1-2k} \pi^{-2k} (-1)^k (2k - 1)!$, which is nonzero, giving

$$\zeta(2k) = -\frac{B_{2k}}{2k} \cdot \frac{1}{2^{1-2k} \pi^{-2k} (-1)^k (2k - 1)!} = -\frac{B_{2k}}{2k} \cdot 2^{2k-1} \pi^{2k} (-1)^{-k} \frac{1}{(2k - 1)!}$$

using $1/2^{1-2k} = 2^{2k-1}$, $1/\pi^{-2k} = \pi^{2k}$, and $1/(-1)^k = (-1)^{-k} = (-1)^k$. Collecting the factorials with $2k \cdot (2k-1)! = (2k)!$ and the signs with $-(-1)^k = (-1)^{k+1}$,

$$\zeta(2k) = (-1)^{k+1} B_{2k} \frac{2^{2k-1} \pi^{2k}}{(2k)!}$$

Finally $2^{2k-1} \pi^{2k} = \frac{1}{2} 2^{2k} \pi^{2k} = \frac{1}{2} (2\pi)^{2k}$, which gives (6.29).

For the stated special cases, take $k = 1$ and $k = 2$. With $B_2 = 1/6$ (definition 1.5.1),

$$\zeta(2) = (-1)^2 \frac{B_2 (2\pi)^2}{2 \cdot 2!} = \frac{(1/6) \cdot 4\pi^2}{4} = \frac{\pi^2}{6}$$

and with $B_4 = -1/30$,

$$\zeta(4) = (-1)^3 \frac{B_4 (2\pi)^4}{2 \cdot 4!} = -\frac{(-1/30) \cdot 16\pi^4}{48} = \frac{16\pi^4}{1440} = \frac{\pi^4}{90}$$

recovering the values of example 4.1.3, completing the proof.

The even values are rational multiples of powers of π for a structural reason now visible in the proof: the functional equation ties $\zeta(2k)$ to the rational number $\zeta(1-2k) = -B_{2k}/(2k)$, and the elementary factors 2^{2k} , π^{2k} , and $1/(2k)!$ that mediate the reflection supply exactly one factor π^{2k} , turning the rational value into a rational multiple of π^{2k} . The transcendence of $\zeta(2k)$ is therefore inherited from π alone; no transcendence enters through the arithmetic side, which contributes only the rational B_{2k} . The odd values $\zeta(3), \zeta(5), \dots$ have no analogous expression precisely because the reflection sends them to the *other* family of negative-integer values: the functional equation pairs $\zeta(2k+1)$ with $\zeta(-2k)$, which is a trivial zero by corollary 6.4.1, so the reflection yields the identity $0 = 0$ rather than an evaluation. This is the exact mechanism behind the difficulty foreshadowed in chapter 4: the even values fall out because their mirror images are nonzero rationals, while the odd values resist because their mirror images vanish, leaving the functional equation silent. Whether $\zeta(2k+1)$ is even irrational is known only for $\zeta(3)$, Apéry's constant of example 4.1.3; the closed-form question is open for every odd argument.

The values at small integers are collected and verified below, showing both families, the rational negative-integer values and the π -power positive even values, side by side.

Example 6.4.4: The Special Values, Computed

The formulas (6.27) and (6.29), together with $\zeta(0) = -1/2$ from theorem 6.4.2, evaluate ζ at the first several integers from the Bernoulli numbers $B_2 = 1/6$, $B_4 = -1/30$, $B_6 = 1/42$ of definition 1.5.1.

1. *Negative integers.* From $\zeta(1-2k) = -B_{2k}/(2k)$:

$$\zeta(-1) = -\frac{B_2}{2} = -\frac{1/6}{2} = -\frac{1}{12}, \quad \zeta(-3) = -\frac{B_4}{4} = -\frac{-1/30}{4} = \frac{1}{120}$$

and $\zeta(-5) = -B_6/6 = -(1/42)/6 = -1/252$. The trivial zeros fill in the even arguments: $\zeta(-2) = \zeta(-4) = \dots = 0$ by corollary 6.4.1.

2. *Positive even integers.* From $\zeta(2k) = (-1)^{k+1} B_{2k} (2\pi)^{2k} / (2(2k)!)$, the case $k = 3$ with

$B_6 = 1/42$ gives

$$\zeta(6) = (-1)^4 \frac{B_6 (2\pi)^6}{2 \cdot 6!} = \frac{(1/42) \cdot 64\pi^6}{2 \cdot 720} = \frac{64\pi^6}{42 \cdot 1440} = \frac{64\pi^6}{60480} = \frac{\pi^6}{945}$$

alongside $\zeta(2) = \pi^2/6$ and $\zeta(4) = \pi^4/90$ from theorem 6.4.3.

Assembling the two families,

s	0	-1	-3	2	4	6
$\zeta(s)$	$-\frac{1}{2}$	$-\frac{1}{12}$	$\frac{1}{120}$	$\frac{\pi^2}{6}$	$\frac{\pi^4}{90}$	$\frac{\pi^6}{945}$

One even value is checked numerically against the defining series. For $\zeta(6)$, the closed form gives $\pi^6/945 = 961.389\dots/945 = 1.017343\dots$, while the partial sum $\sum_{n=1}^4 n^{-6} = 1 + 1/64 + 1/729 + 1/4096 = 1.017241\dots$ already agrees to three decimal places, the difference 1.02×10^{-4} matching the size of the tail $\sum_{n \geq 5} n^{-6}$, which is bounded by $\int_4^\infty t^{-6} dt = 1/(5 \cdot 4^5) \approx 1.95 \times 10^{-4}$. The notorious value $\zeta(-1) = -1/12$, which the divergent series $1 + 2 + 3 + \dots$ is sometimes said to “equal” in physics, is here simply the value of the analytically continued ζ at $s = -1$, the rational number $-B_2/2$ produced by the functional equation, with no summation of the divergent series involved.

6.5 The Critical Strip and the Riemann Hypothesis

The functional equation $\Lambda(s) = \Lambda(1-s)$ of theorem 6.3.3 does more than continue ζ to the whole plane; it imposes a reflection symmetry whose axis is the line $\text{Re}(s) = 1/2$. The trivial zeros of section 6.4, lying at the even negative integers, account for every zero in the left half-plane, and the half-plane $\text{Re}(s) \geq 1$ carries none by the non-vanishing on the line (theorem 4.4.2). What remains is a vertical strip of width one, the critical strip, into which the symmetry $s \leftrightarrow 1-s$ folds all the other zeros. These are the nontrivial zeros, the zeros that carry the arithmetic content of ζ through the explicit formula of box 5.5.5.

The functional equation arranges the nontrivial zeros into a configuration symmetric across the critical line, and a second symmetry, complex conjugation, pairs each zero with its reflection across the real axis. The two symmetries together confine the zeros to quadruples unless they fall on the critical line itself. The Riemann Hypothesis (?? 5.5.6) is the assertion that the quadruples always degenerate, that every nontrivial zero lies on the axis of symmetry. This section fixes the terminology, establishes the symmetries the functional equation forces, states the asymptotic count of the zeros in the strip, and introduces the Hardy Z -function, the real-valued companion of ζ on the critical line whose zeros are exactly the critical-line zeros of ζ .

Definition 6.5.1: The Critical Strip and the Critical Line

The *critical strip* is the closed vertical strip

$$\{s \in \mathbb{C} \mid 0 \leq \operatorname{Re}(s) \leq 1\}$$

The *critical line* is the vertical line $\operatorname{Re}(s) = 1/2$, the axis of symmetry of the functional equation. A *nontrivial zero* of ζ is a zero lying in the critical strip; equivalently (as proposition 6.5.3 shows the strip endpoints carry no zeros) a zero with $0 < \operatorname{Re}(s) < 1$. The nontrivial zeros of ζ are exactly the zeros of the entire function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ of box 5.5.5.

The qualifier “nontrivial” separates these zeros from the trivial zeros at $s = -2, -4, -6, \dots$ of corollary 6.4.1, which lie outside the strip and are completely understood. The identification of the nontrivial zeros with the zeros of ξ is what makes the strip the right object: the elementary prefactor $\frac{1}{2}s(s-1)$ removes the pole of ζ at $s = 1$ and the pole of the gamma factor at $s = 0$, and the gamma factor $\pi^{-s/2}\Gamma(s/2)$ has no zeros and supplies poles at $s = 0, -2, -4, \dots$ that cancel the trivial zeros of ζ one for one. What survives in ξ is an entire function whose only zeros are the nontrivial zeros of ζ , none of them at $s = 0$ or $s = 1$. Because ξ is entire and (by Stirling, theorem 6.1.4) of order one, it has infinitely many zeros, all inside the strip; the Hadamard product over them is what converts the explicit formula’s contour integral into a sum over ρ .

Example 6.5.2: The First Nontrivial Zero

The lowest nontrivial zero in the upper half-plane is

$$\rho_1 = \frac{1}{2} + i\gamma_1, \quad \gamma_1 = 14.134725\dots$$

a value with $\operatorname{Re}(\rho_1) = 1/2$ exactly, consistent with the Riemann Hypothesis. It lies in the interior of the critical strip, $0 < \frac{1}{2} < 1$, and not on either boundary line: it is not at $\operatorname{Re}(s) = 1$, in accordance with theorem 4.4.2, and not at $\operatorname{Re}(s) = 0$, where ξ is nonzero. By the conjugate symmetry of proposition 6.5.3 the point $\bar{\rho}_1 = \frac{1}{2} - i\gamma_1$ is also a zero, so the zeros nearest the real axis form the pair $\frac{1}{2} \pm 14.134725\dots i$. Since this zero sits on the critical line, the reflection $1 - \rho_1 = \frac{1}{2} - i\gamma_1$ coincides with $\bar{\rho}_1$, and the would-be quadruple of proposition 6.5.3 collapses to a conjugate pair; the numerical value 14.134725... is computed in [1, Ch. 15].

The two boundary lines of the strip and the region outside it must first be cleared of nontrivial zeros, and the symmetries that organize the zeros inside the strip must be established, before the count of theorem 6.5.4 can be stated. Both follow from the functional equation together with the non-vanishing on $\operatorname{Re}(s) = 1$.

Proposition 6.5.3: Symmetry of the Nontrivial Zeros

The nontrivial zeros of ζ lie in the open strip $0 < \operatorname{Re}(s) < 1$. They are symmetric under the two involutions

$$s \mapsto 1 - s \quad \text{and} \quad s \mapsto \bar{s}$$

that is, if ρ is a nontrivial zero then so are $1 - \rho$, $\bar{\rho}$, and $1 - \bar{\rho}$. Consequently the nontrivial zeros occur in quadruples $\{\rho, 1 - \rho, \bar{\rho}, 1 - \bar{\rho}\}$, which degenerate to a conjugate pair $\{\rho, \bar{\rho}\}$ precisely when ρ lies on the critical line $\operatorname{Re}(s) = 1/2$.

Proof:

The argument locates the zeros in the open strip, then derives each of the two symmetries, then reads off the quadruple structure.

Step 1: No nontrivial zeros with $\operatorname{Re}(s) \geq 1$. For $\operatorname{Re}(s) > 1$ the Euler product $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ of corollary 4.1.2 converges to a nonzero value, since each factor is nonzero and the product converges absolutely; hence $\zeta(s) \neq 0$ there. On the line $\operatorname{Re}(s) = 1$ itself, $\zeta(1 + it) \neq 0$ for every real $t \neq 0$ by theorem 4.4.2, and at $t = 0$ the point $s = 1$ is the pole of ζ (corollary 6.3.4), not a zero. Therefore ζ has no zeros in the closed half-plane $\operatorname{Re}(s) \geq 1$.

Step 2: No zeros with $\operatorname{Re}(s) \leq 0$ except the trivial ones. The asymmetric functional equation (6.22) of corollary 6.3.4 reads

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

Fix s with $\operatorname{Re}(s) < 0$, so that $\operatorname{Re}(1-s) > 1$. By Step 1 the factor $\zeta(1-s)$ is nonzero. The factors 2^s and π^{s-1} are nowhere zero, and $\Gamma(1-s)$ is nonzero everywhere, having no zeros (theorem 6.1.2); since $\operatorname{Re}(1-s) > 1 > 0$, the point $1-s$ is not a pole of Γ either, so $\Gamma(1-s)$ is finite and nonzero. The only factor that can vanish is $\sin(\pi s/2)$, which is zero exactly at the even integers; in the range $\operatorname{Re}(s) < 0$ these are $s = -2, -4, -6, \dots$, the trivial zeros of corollary 6.4.1. Hence in $\operatorname{Re}(s) < 0$ the function ζ vanishes only at the trivial zeros. On the line $\operatorname{Re}(s) = 0$ the same identity applies with $\operatorname{Re}(1-s) = 1$: the factor $\zeta(1-s) = \zeta(1-it)$ is nonzero for $t \neq 0$ by theorem 4.4.2, the factor $\sin(\pi s/2) = \sin(\pi it/2)$ is nonzero for $t \neq 0$, and $\Gamma(1-s)$ is finite and nonzero on this line; at $s = 0$ the factor $\sin(\pi s/2)$ vanishes, but $\zeta(1-s) = \zeta(1)$ is the pole, and the product is the finite value $\zeta(0) = -1/2$ (theorem 6.4.2), not a zero. Thus $\operatorname{Re}(s) = 0$ carries no zeros at all. Combining with Step 1, every zero other than the trivial zeros lies in the open strip $0 < \operatorname{Re}(s) < 1$, which is the asserted location of the nontrivial zeros.

Step 3: Symmetry under $s \mapsto 1-s$. Let ρ be a nontrivial zero, so $0 < \operatorname{Re}(\rho) < 1$ and $\zeta(\rho) = 0$. The gamma factor $\Gamma_{\mathbb{R}}(\rho) = \pi^{-\rho/2} \Gamma(\rho/2)$ is finite and nonzero at ρ , since $\Gamma(\rho/2)$ has its poles only at $\rho/2 = 0, -1, -2, \dots$, that is at $\rho = 0, -2, -4, \dots$, none of which lie in the open strip. Hence the completed function $\Lambda(\rho) = \Gamma_{\mathbb{R}}(\rho) \zeta(\rho) = 0$. By the functional equation (6.14), $\Lambda(1-\rho) = \Lambda(\rho) = 0$. The reflected point $1-\rho$ also lies in the open strip, since $\operatorname{Re}(1-\rho) = 1 - \operatorname{Re}(\rho) \in (0, 1)$, so its gamma factor $\Gamma_{\mathbb{R}}(1-\rho)$ is again finite and nonzero by the same reasoning. From $\Lambda(1-\rho) = \Gamma_{\mathbb{R}}(1-\rho) \zeta(1-\rho) = 0$ and $\Gamma_{\mathbb{R}}(1-\rho) \neq 0$ it follows that $\zeta(1-\rho) = 0$. Thus $1-\rho$ is a nontrivial zero.

Step 4: Symmetry under $s \mapsto \bar{s}$. On the half-plane $\operatorname{Re}(s) > 1$ the Dirichlet series gives $\zeta(\bar{s}) = \sum_{n \geq 1} n^{-\bar{s}} = \sum_{n \geq 1} \overline{n^{-s}} = \overline{\zeta(s)}$, the middle equality holding because $n^{-\bar{s}} = \overline{n^{-s}}$ for the positive integer base n . Both $\zeta(\bar{s})$ and $\overline{\zeta(s)}$ are holomorphic on $\mathbb{C} \setminus \{1\}$ (the continuation of corollary 6.3.4, composed with conjugation, is holomorphic), and they agree on the open set $\operatorname{Re}(s) > 1$, so by the identity theorem ([16]) the relation

$$\zeta(\bar{s}) = \overline{\zeta(s)} \tag{6.31}$$

holds for all $s \neq 1$. If ρ is a nontrivial zero then $\zeta(\bar{\rho}) = \overline{\zeta(\rho)} = \bar{0} = 0$, and $\bar{\rho}$ lies in the open strip because $\operatorname{Re}(\bar{\rho}) = \operatorname{Re}(\rho) \in (0, 1)$, so $\bar{\rho}$ is a nontrivial zero.

Step 5: The quadruple structure. The two involutions $s \mapsto 1 - s$ and $s \mapsto \bar{s}$ commute, their composite being $s \mapsto 1 - \bar{s}$. Applying them to a nontrivial zero ρ produces the four points ρ , $1 - \rho$, $\bar{\rho}$, and $1 - \bar{\rho}$, each a nontrivial zero by Steps 3 and 4. Writing $\rho = \beta + i\gamma$ with $0 < \beta < 1$, the four points are $\beta + i\gamma$, $(1 - \beta) - i\gamma$, $\beta - i\gamma$, and $(1 - \beta) + i\gamma$, the vertices of a rectangle centered at $\frac{1}{2}$ on the real axis, with sides parallel to the axes. This rectangle is nondegenerate, and the four zeros distinct, exactly when $\beta \neq 1/2$ and $\gamma \neq 0$. Conjugation pairs ρ with $\bar{\rho}$ symmetrically about the real axis, so a zero off the real axis ($\gamma \neq 0$) is never alone; the count along the line $\operatorname{Re}(s) = 1/2$ is the case where the reflection $s \mapsto 1 - s$ adds nothing new, since there $\beta = 1 - \beta$ forces $1 - \rho = \bar{\rho}$ and the quadruple collapses to the conjugate pair $\frac{1}{2} \pm i\gamma$. Thus a nontrivial zero ρ with $\gamma \neq 0$ generates a full quadruple of four distinct zeros unless it lies on the critical line, where it generates only the pair $\{\rho, \bar{\rho}\}$, as we sought to show.

The two symmetries have different sources, and the distinction is worth keeping in view. The conjugate symmetry $\zeta(\bar{s}) = \overline{\zeta(s)}$ of (6.31) is elementary, an instance of the Schwarz reflection principle that holds for any Dirichlet series with real coefficients; it places the zeros symmetrically about the real axis and is the reason the explicit formula of box 5.5.5 groups the zeros into conjugate pairs $\rho, \bar{\rho}$, so that the oscillatory terms $x^\rho/\rho + x^{\bar{\rho}}/\bar{\rho}$ combine into a real quantity. The reflection symmetry $s \mapsto 1 - s$ is the substantive one: it is the functional equation, and it places the zeros symmetrically about the critical line. A nontrivial zero off the line at $\beta + i\gamma$ with $\beta > 1/2$ forces a partner at $(1 - \beta) + i\gamma$ with $1 - \beta < 1/2$, on the opposite side of the line and equally far from it. The Riemann Hypothesis is the statement that this never happens, that the reflection $s \mapsto 1 - s$ fixes every nontrivial zero, which is possible only on the fixed-point line $\operatorname{Re}(s) = 1/2$.

The symmetry confines the zeros but does not count them. The distribution of the nontrivial zeros up to a given height is governed by an asymptotic formula, the foundation of every quantitative statement about the zeros and the starting point for the explicit formula's error analysis.

Theorem 6.5.4: The Zero-Counting Formula

Let $N(T)$ denote the number of nontrivial zeros $\rho = \beta + i\gamma$ of ζ with $0 < \gamma \leq T$, counted with multiplicity. Then

$$N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + O(\log T)$$

as $T \rightarrow \infty$.

The full proof is a careful application of the argument principle to ξ on a rectangle, together with Stirling's estimate for the gamma factor; the computation tracks the change in $\arg \xi$ along three sides of the rectangle and bounds the contribution of the fourth, and its several pages of estimation carry no conceptual turning point beyond the two inputs named. The argument is given in full in [1, Ch. 15] and [11, Ch. 14]; the key ideas and the origin of each term are recorded here.

Proof Key ideas:

Fix a large T avoiding the ordinates of zeros, and let R be the positively oriented boundary of the rectangle with corners $-1, 2, 2 + iT, -1 + iT$, chosen so that all nontrivial zeros with $0 < \gamma \leq T$ lie strictly inside (the strip $0 < \operatorname{Re}(s) < 1$ sits inside $-1 < \operatorname{Re}(s) < 2$) and none lies on R .

Step 1: The argument principle for ξ . The function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ is entire (definition 6.5.1), and the rectangle $R = [-1, 2] \times [0, T]$ lies in the upper half-plane, so its zeros inside R are exactly the nontrivial zeros of ζ with $0 < \gamma \leq T$. The argument principle ([16])

counts these as

$$N(T) = \frac{1}{2\pi} \Delta_R \arg \xi$$

the normalized total change in $\arg \xi$ around R . Two symmetries reduce this contour. The conjugate symmetry $\xi(\bar{s}) = \overline{\xi(s)}$, inherited from (6.31) and the real prefactor, together with the reflection $\xi(s) = \xi(1-s)$ gives $\xi(1-\bar{s}) = \overline{\xi(s)}$, so reflecting R across the line $\operatorname{Re}(s) = 1/2$ (the map $s \mapsto 1-\bar{s}$) conjugates ξ and leaves $|\Delta \arg \xi|$ unchanged; hence the change in $\arg \xi$ over the left half of R equals the change over the right half, and $\Delta_R \arg \xi = 2 \Delta_{R'} \arg \xi$, where R' is the right portion of the boundary running $\frac{1}{2} \rightarrow 2 \rightarrow 2+iT \rightarrow \frac{1}{2}+iT$. On the bottom segment $\frac{1}{2} \rightarrow 2$ the function ξ is real and positive, so $\arg \xi$ is constant there and that segment contributes nothing; the change over R' equals the change along the half-contour L running $2 \rightarrow 2+iT \rightarrow \frac{1}{2}+iT$. Assembling these reductions,

$$\pi N(T) = \Delta_L \arg \xi$$

the change in $\arg \xi$ along L .

Step 2: The gamma factor gives the main term. Split $\arg \xi = \arg(\frac{1}{2}s(s-1)) + \arg(\pi^{-s/2}\Gamma(s/2)) + \arg \zeta(s)$ and track each summand along the half-contour L . The change in $\arg(\pi^{-s/2}\Gamma(s/2))$ along L is evaluated by Stirling's estimate (6.7) of theorem 6.1.4: at the endpoint $1/2+iT$ the formula $\log \Gamma(s/2)$ contributes an imaginary part $\operatorname{Im} \log \Gamma(\frac{1}{4} + \frac{iT}{2})$ of size $\frac{T}{2} \log \frac{T}{2} - \frac{T}{2} + O(\log T)$, and the factor $\pi^{-s/2}$ contributes $-\frac{T}{2} \log \pi$. Combining and using $\log \frac{T}{2} - \log \pi = \log \frac{T}{2\pi}$ gives the change $\frac{T}{2} \log \frac{T}{2\pi} - \frac{T}{2} + O(\log T)$ along L ; dividing by the factor π from the relation $\pi N(T) = \Delta_L \arg \xi$ of Step 1 produces the main term

$$\frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi}$$

The elementary factor $\frac{1}{2}s(s-1)$ contributes a bounded change, of order $O(1)$, absorbed into the error.

Step 3: The zeta factor gives the error term. The remaining summand is $\frac{1}{\pi}$ times the change in $\arg \zeta$ along the half-contour L . On the segment $2 \rightarrow 2+iT$ this change is bounded, so up to $O(1)$ it equals the customary

$$S(T) = \frac{1}{\pi} \arg \zeta(\frac{1}{2} + iT)$$

the argument taken by continuous variation from $s=2$, where $\arg \zeta = 0$. The bound $S(T) = O(\log T)$ comes from controlling the number of sign changes of $\operatorname{Re} \zeta$ along the horizontal segment, which is in turn bounded by Jensen's inequality applied to ζ on a disc of fixed radius about $2+iT$, where ζ has size polynomial in T . This $O(\log T)$ is the entire error term. The full execution of all three steps, including the contour avoidance and the Jensen bound on $S(T)$, is carried out in [1, Ch. 15] and [11, Ch. 14], completing the proof.

The formula says the nontrivial zeros have density $\frac{1}{2\pi} \log \frac{T}{2\pi}$ near height T : the average gap between consecutive ordinates γ near T is $2\pi/\log T$, shrinking slowly as one climbs the critical strip. The main term is produced entirely by the archimedean data, the gamma factor and the power of π that complete ζ , and is therefore known unconditionally and exactly; the arithmetic of ζ enters only through the secondary term $S(T)$, whose true size is the subject of much of the finer theory. The Riemann Hypothesis is not needed for the count: $N(T)$ is the number of zeros in the strip regardless

of where in the strip they sit, and the formula holds whether or not they lie on the critical line. What the Riemann Hypothesis would add is the assertion that all $N(T)$ of these zeros have $\beta = 1/2$, refining a statement about how many zeros there are into a statement about where they are.

The count motivates a change of viewpoint. To study the zeros on the critical line directly, one restricts ζ to that line and removes the phase that the gamma factor introduces, producing a real-valued function whose sign changes locate the zeros.

Definition 6.5.5: The Hardy Z -Function

The *Riemann–Siegel theta function* (distinct from the Jacobi theta function of section 6.2, despite the shared symbol; the two never appear together) is

$$\theta(t) = \arg\left(\pi^{-it/2} \Gamma\left(\frac{1}{4} + \frac{it}{2}\right)\right)$$

the argument taken by the principal branch, continuous in t with $\theta(0) = 0$. The *Hardy Z -function* is

$$Z(t) = e^{i\theta(t)} \zeta\left(\frac{1}{2} + it\right)$$

defined for real t . It is real-valued, satisfies $|Z(t)| = |\zeta(\frac{1}{2} + it)|$, and its real zeros are exactly the zeros of ζ on the critical line.

Proof:

The three asserted properties are established in turn; the first is the substantive one and rests on the functional equation restricted to the critical line.

Step 1: $Z(t)$ is real-valued. Recall the real gamma factor $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma(s/2)$ of definition 6.3.1 and the completed function $\Lambda(s) = \Gamma_{\mathbb{R}}(s) \zeta(s)$. On the critical line $s = \frac{1}{2} + it$,

$$\Gamma_{\mathbb{R}}\left(\frac{1}{2} + it\right) = \pi^{-(1/2+it)/2} \Gamma\left(\frac{1}{4} + \frac{it}{2}\right) = \pi^{-1/4} \cdot \pi^{-it/2} \Gamma\left(\frac{1}{4} + \frac{it}{2}\right)$$

The factor $\pi^{-1/4}$ is a positive real number, so $\Gamma_{\mathbb{R}}(\frac{1}{2} + it)$ has the same argument as $\pi^{-it/2} \Gamma(\frac{1}{4} + \frac{it}{2})$, namely $\theta(t)$. Writing $\Gamma_{\mathbb{R}}(\frac{1}{2} + it) = |\Gamma_{\mathbb{R}}(\frac{1}{2} + it)| e^{i\theta(t)}$ therefore holds, and consequently

$$Z(t) = e^{i\theta(t)} \zeta\left(\frac{1}{2} + it\right) = \frac{\Gamma_{\mathbb{R}}(\frac{1}{2} + it)}{|\Gamma_{\mathbb{R}}(\frac{1}{2} + it)|} \zeta\left(\frac{1}{2} + it\right) = \frac{\Lambda(\frac{1}{2} + it)}{|\Gamma_{\mathbb{R}}(\frac{1}{2} + it)|} \quad (6.32)$$

using $\Gamma_{\mathbb{R}}(\frac{1}{2} + it) \zeta(\frac{1}{2} + it) = \Lambda(\frac{1}{2} + it)$. The denominator $|\Gamma_{\mathbb{R}}(\frac{1}{2} + it)|$ is a positive real number (the gamma factor has no zeros and no poles on the line, as $\frac{1}{4} + \frac{it}{2}$ is never a pole of Γ). It remains to show the numerator $\Lambda(\frac{1}{2} + it)$ is real. The functional equation (6.14) at $s = \frac{1}{2} + it$ gives $\Lambda(\frac{1}{2} + it) = \Lambda(1 - (\frac{1}{2} + it)) = \Lambda(\frac{1}{2} - it)$. The completed function inherits the Schwarz reflection $\Lambda(\bar{s}) = \overline{\Lambda(s)}$ from (6.31): indeed $\Gamma_{\mathbb{R}}(\bar{s}) = \pi^{-\bar{s}/2} \Gamma(\bar{s}/2) = \overline{\pi^{-s/2} \Gamma(s/2)} = \overline{\Gamma_{\mathbb{R}}(s)}$, since Γ has real Taylor coefficients and so commutes with conjugation, and multiplying by $\zeta(\bar{s}) = \overline{\zeta(s)}$ yields $\Lambda(\bar{s}) = \overline{\Lambda(s)}$. Applied at $s = \frac{1}{2} + it$, whose conjugate is $\frac{1}{2} - it$, this reads $\Lambda(\frac{1}{2} - it) = \overline{\Lambda(\frac{1}{2} + it)}$. Chaining the two relations,

$$\Lambda\left(\frac{1}{2} + it\right) = \Lambda\left(\frac{1}{2} - it\right) = \overline{\Lambda\left(\frac{1}{2} + it\right)}$$

so $\Lambda(\frac{1}{2} + it)$ equals its own conjugate and is real. By (6.32), $Z(t)$ is a real number divided by a positive real number, hence real.

Step 2: $|Z(t)| = |\zeta(\frac{1}{2} + it)|$. The phase factor has modulus one, $|e^{i\theta(t)}| = 1$, since $\theta(t)$ is real. Therefore $|Z(t)| = |e^{i\theta(t)}| |\zeta(\frac{1}{2} + it)| = |\zeta(\frac{1}{2} + it)|$ directly from the definition.

Step 3: The real zeros of Z are the critical-line zeros of ζ . For real t , the factor $e^{i\theta(t)}$ is nonzero, so $Z(t) = 0$ if and only if $\zeta(\frac{1}{2} + it) = 0$. Thus t is a real zero of Z exactly when $\frac{1}{2} + it$ is a zero of ζ on the critical line, with matching multiplicities since $e^{i\theta(t)}$ is a nonvanishing analytic factor. This is the asserted correspondence, completing the proof.

The Z -function trades a complex-valued function on a line for an equivalent real-valued one, and the gain is that real zeros can be detected by sign changes. Whenever $Z(t_1)$ and $Z(t_2)$ have opposite signs, the intermediate value theorem forces a zero of Z , hence a zero of ζ , between them on the critical line; this is the mechanism behind every numerical and theoretical verification that zeros lie on the line, including Hardy's theorem that infinitely many do. The factorization (6.32) shows Z is the completed function Λ on the line, stripped of the positive modulus of its gamma factor, so the zeros of Z , the zeros of Λ on the line, and the critical-line zeros of ζ all coincide. The Riemann–Siegel theta function $\theta(t)$ that supplies the phase grows like $\frac{t}{2} \log \frac{t}{2\pi} - \frac{t}{2}$ by the same Stirling estimate that produced the main term of theorem 6.5.4, and the count $N(T)$ is, up to the error $S(T)$, the number of times $\theta(t)$ sweeps through a multiple of π ; the zero-counting formula and the Z -function are thus two readings of the same gamma-factor asymptotics.

6.5.6 Note: The Riemann Hypothesis Revisited The Riemann Hypothesis (?? 5.5.6) was stated in chapter 5 as the assertion that every nontrivial zero has $\operatorname{Re}(s) = 1/2$. The functional equation now exhibits why this line is singled out: by proposition 6.5.3 the nontrivial zeros are symmetric across the critical line under $s \mapsto 1 - s$, and $\operatorname{Re}(s) = 1/2$ is the axis of that symmetry, the fixed-point set of the involution. The Riemann Hypothesis is therefore the statement that every nontrivial zero sits *on* the axis of symmetry, so that the quadruple $\{\rho, 1 - \rho, \bar{\rho}, 1 - \bar{\rho}\}$ always degenerates to the conjugate pair $\{\rho, \bar{\rho}\}$. The consequence for the distribution of primes is recorded in proposition 5.5.7: the hypothesis is equivalent to the error bound $\psi(x) = x + O(\sqrt{x}(\log x)^2)$, the sharpest possible, since the size of the zero-sum $\sum_{\rho} x^{\rho}/\rho$ in the explicit formula is governed by $\sup_{\rho} \operatorname{Re}(\rho)$. With the functional equation in hand, every ingredient of that explicit formula (box 5.5.5) is now established: the entire completed function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$, whose zeros are exactly the nontrivial zeros, and the product factorization over those zeros that converts the contour integral of $-\zeta'/\zeta$ into the sum over ρ .

Dirichlet Characters and L-Functions

The six preceding chapters built the analytic theory of the primes around a single function, the Riemann zeta function $\zeta(s) = \sum_{n \geq 1} n^{-s}$. Its Euler product (corollary 2.3.2) encoded unique factorization analytically; its pole at $s = 1$ (theorem 4.2.1) measured the abundance of the primes; its non-vanishing on the line $\operatorname{Re}(s) = 1$ (theorem 4.4.2) converted that abundance into the Prime Number Theorem (theorem 5.4.3); and its functional equation (theorem 6.3.3), assembled from the gamma and theta functions, fixed the symmetry of the whole picture. That theory counts all primes at once. It says nothing about how the primes distribute themselves among the residue classes modulo a fixed integer m , the question of whether the arithmetic progression $a, a + m, a + 2m, \dots$ with $\gcd(a, m) = 1$ contains infinitely many primes. To reach the residue classes one must split the prime-counting machinery according to congruence, and the device that performs the splitting is the Dirichlet character: a homomorphism of the unit group $(\mathbb{Z}/m\mathbb{Z})^\times$, extended to all of $\mathbb{Z}$ by zero on the integers sharing a factor with m . Twisting the zeta construction by such a character χ produces a Dirichlet L -function $L(s, \chi) = \sum_{n \geq 1} \chi(n) n^{-s}$, and the family of these functions, one for each character modulo m , resolves the single count $\sum_n n^{-s}$ into $\varphi(m)$ separate counts, one attached to each residue class.

This chapter is the character-twisted parallel of chapters 4 to 6. Each structural feature of ζ has its analogue for $L(s, \chi)$: an Euler product over the primes, a meromorphic continuation (entire when χ is nonprincipal, since the pole of ζ survives only in the principal twist), and a functional equation relating $L(s, \chi)$ to $L(1 - s, \bar{\chi})$, the reflection now mediated not only by the gamma factor but by a finite exponential sum, the Gauss sum, whose absolute value $\sqrt{m}$ supplies the constant in the symmetry. Above all there is one fact with no analogue in the zeta theory because it is automatic there, where the pole forces $\zeta(s) \rightarrow \infty$ as $s \rightarrow 1$: the non-vanishing $L(1, \chi) \neq 0$ for every nonprincipal character χ . This single inequality is what prevents the contributions of the nonprincipal characters from cancelling the main term, and it yields Dirichlet's theorem, that every arithmetic progression

$a \bmod m$ with $\gcd(a, m) = 1$ contains infinitely many primes, distributed with density $1/\varphi(m)$ among the primes (chapter 7 closes on this result). The development begins with the characters themselves, the finite group-theoretic objects on which the entire analytic edifice is raised.

7.1 Dirichlet Characters

The structure of the unit group $(\mathbb{Z}/m\mathbb{Z})^\times$ was settled in section 0.4: it is a finite abelian group of order $\varphi(m)$, cyclic precisely when $m \in \{1, 2, 4, p^k, 2p^k\}$ and a product of cyclic factors in general. The arithmetic functions of the previous chapters were defined directly on $\mathbb{Z}$, and the multiplicative ones, such as $n \mapsto n^{-s}$, were exactly the functions respecting the multiplication of $\mathbb{Z}$. To detect congruence classes one needs functions that also respect the multiplication of $(\mathbb{Z}/m\mathbb{Z})^\times$, and the homomorphisms from this finite abelian group to the multiplicative group $\mathbb{C}^\times$ are the natural such objects. Extended from the unit group to all integers by setting them to zero where they would otherwise be undefined, these homomorphisms become arithmetic functions in their own right, the Dirichlet characters. This section constructs them, counts them, records the two facts about them on which everything later depends (that they form a group dual to $(\mathbb{Z}/m\mathbb{Z})^\times$, and that they satisfy orthogonality relations), and isolates the refinements (the conductor and the parity) that the functional equation will require.

Definition 7.1.1: Dirichlet Character

Let $m \in \mathbb{Z}_{>0}$. A *Dirichlet character modulo m* is a function $\chi: \mathbb{Z} \rightarrow \mathbb{C}$ satisfying the following three conditions:

1. *Complete multiplicativity*: $\chi(ab) = \chi(a)\chi(b)$ for all $a, b \in \mathbb{Z}$;
2. *Periodicity*: $\chi(a + m) = \chi(a)$ for all $a \in \mathbb{Z}$;
3. *Support on the units*: $\chi(a) \neq 0$ if and only if $\gcd(a, m) = 1$.

Equivalently, χ arises from a group homomorphism $\tilde{\chi}: (\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ by setting $\chi(a) = \tilde{\chi}(a \bmod m)$ when $\gcd(a, m) = 1$ and $\chi(a) = 0$ otherwise. The integer m is the *modulus* of χ .

The three conditions and the homomorphism description are two presentations of the same object, and the equivalence is worth tracing once. A group homomorphism $\tilde{\chi}: (\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ extended by zero is completely multiplicative: if both a and b are coprime to m , then $\chi(ab) = \tilde{\chi}(ab) = \tilde{\chi}(a)\tilde{\chi}(b) = \chi(a)\chi(b)$, and if either shares a factor with m , then so does ab , so both sides vanish; it is periodic because its value depends only on the residue $a \bmod m$; and it is supported on the units by construction. Conversely, any χ satisfying the three conditions restricts to a function $(\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ (well defined by periodicity, nonzero by condition (iii)) that is multiplicative by condition (i), hence a group homomorphism. The values of a character are constrained: applying condition (i) to $a^{\varphi(m)}$ for $\gcd(a, m) = 1$ and using Euler's theorem (theorem 0.3.12), $a^{\varphi(m)} \equiv 1 \pmod{m}$, gives $\chi(a)^{\varphi(m)} = \chi(1) = 1$, so every nonzero value of χ is a $\varphi(m)$ -th root of unity and in particular has absolute value 1.

The passage from the group homomorphism $\tilde{\chi}$ to the arithmetic function χ is the move that makes character theory available to analytic number theory: a homomorphism of $(\mathbb{Z}/m\mathbb{Z})^\times$ knows nothing about the integers, but its extension by zero is a completely multiplicative function on $\mathbb{Z}$, the same class of function as $n \mapsto n^{-s}$, and so can be fed into a Dirichlet series. The remainder of this chapter is the study of the series $\sum_n \chi(n)n^{-s}$ so produced. The notational convention, followed throughout,

is to write χ for both the character on $\mathbb{Z}$ and the underlying homomorphism on $(\mathbb{Z}/m\mathbb{Z})^\times$, the meaning being fixed by the argument.

The count and the group law for these objects come directly from the duality theory of finite abelian groups, applied to $G = (\mathbb{Z}/m\mathbb{Z})^\times$.

Theorem 7.1.2: Characters as the Dual Group

Fix $m \in \mathbb{Z}_{>0}$ and write $G = (\mathbb{Z}/m\mathbb{Z})^\times$. Restriction to the units, $\chi \mapsto \tilde{\chi}$, is a bijection between the Dirichlet characters modulo m and the group homomorphisms $G \rightarrow \mathbb{C}^\times$, that is, the elements of the dual group $\hat{G} = \text{Hom}(G, \mathbb{C}^\times)$. Consequently:

1. there are exactly $\varphi(m)$ Dirichlet characters modulo m ;
2. under pointwise multiplication $(\chi_1\chi_2)(n) = \chi_1(n)\chi_2(n)$ the Dirichlet characters modulo m form a finite abelian group, with identity the principal character and inverse $\chi^{-1} = \bar{\chi}$ the complex conjugate;
3. this group is isomorphic to G itself; and
4. every character takes values in the $\varphi(m)$ -th roots of unity.

Proof:

The bijection in the opening sentence is the equivalence established after definition 7.1.1: extension by zero sends a homomorphism $G \rightarrow \mathbb{C}^\times$ to a Dirichlet character, restriction to the units sends a Dirichlet character to such a homomorphism, and the two operations are mutually inverse. Under this bijection pointwise multiplication of characters on $\mathbb{Z}$ corresponds to pointwise multiplication of homomorphisms on G , since the bijection is given by restriction and multiplication is computed pointwise on both sides; the bijection is therefore a group isomorphism between the Dirichlet characters modulo m and the dual group $\hat{G}$. It remains to import the structure of $\hat{G}$.

The group G is finite abelian (section 0.4). For a finite abelian group G the dual $\hat{G} = \text{Hom}(G, \mathbb{C}^\times)$ is a group under pointwise multiplication, with identity the trivial homomorphism $g \mapsto 1$ and inverse $\tilde{\chi}^{-1}(g) = \tilde{\chi}(g)^{-1} = \overline{\tilde{\chi}(g)}$, the last equality because $\tilde{\chi}(g)$ is a root of unity (every element of G has finite order, so its image does too), hence has modulus 1; and there is a (non-canonical) isomorphism $\hat{G} \cong G$, so in particular $|\hat{G}| = |G|$. These are the standard facts on characters of finite abelian groups; their proof is given in [13], the cyclic case by sending a generator to a chosen root of unity and the general case by the invariant factor decomposition $G \cong \prod_i \mathbb{Z}/d_i\mathbb{Z}$, under which $\hat{G} \cong \prod_i \widehat{\mathbb{Z}/d_i\mathbb{Z}} \cong \prod_i \mathbb{Z}/d_i\mathbb{Z} \cong G$.

Transporting these facts across the isomorphism establishes the numbered claims. The count $|\hat{G}| = |G| = \varphi(m)$ gives (1). The group law and the identification of the identity and inverse give (2), the principal character being the extension by zero of the trivial homomorphism (definition 7.1.3 below) and $\bar{\chi}$ the extension of $\tilde{\chi}$, which agrees with χ^{-1} on the units and is zero off them. The isomorphism $\hat{G} \cong G$ gives (3). Finally (4) restates the value constraint already noted after definition 7.1.1: for $\gcd(n, m) = 1$ the element $\tilde{\chi}(\bar{n})$ has order dividing $|G| = \varphi(m)$, so $\chi(n)^{\varphi(m)} = 1$, completing the proof.

The theorem fixes the population of characters and, more usefully, gives that population the structure

of a group isomorphic to $(\mathbb{Z}/m\mathbb{Z})^\times$. The isomorphism $\widehat{G} \cong G$ is not canonical, depending on a choice of generators, but its mere existence is what is used: it guarantees exactly $\varphi(m)$ characters, no more and no fewer, and this exact count is what makes the orthogonality relations below into clean equalities rather than inequalities. When $(\mathbb{Z}/m\mathbb{Z})^\times$ is cyclic, say generated by a primitive root g (section 0.4), a character is determined by the single value $\chi(g)$, which may be any $\varphi(m)$ -th root of unity, and the $\varphi(m)$ choices of $\chi(g)$ enumerate the $\varphi(m)$ characters; this is the situation for prime moduli and is the source of the concrete tables in example 7.1.7. For non-cyclic moduli a character is specified by its values on a set of generators of the invariant factors, one root of unity of the appropriate order for each factor.

Among the $\varphi(m)$ characters one is distinguished as the identity of the group, and it warrants a name because it plays the role for L -functions that ζ itself plays.

Definition 7.1.3: The Principal Character

The *principal character* modulo m , denoted χ_0 , is the Dirichlet character

$$\chi_0(n) = \begin{cases} 1 & \text{if } \gcd(n, m) = 1, \\ 0 & \text{if } \gcd(n, m) > 1 \end{cases}$$

the extension by zero of the trivial homomorphism $(\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$.

The principal character is the multiplicative indicator function of the integers coprime to m : it is 1 exactly where a character is allowed to be nonzero, and so records support without recording any finer information about the residue class. Its L -function will turn out to be the Riemann zeta function with the Euler factors at the primes dividing m removed (proposition 7.2.3), which is why χ_0 , alone among the characters modulo m , produces a function with a pole. Every other character distinguishes residue classes nontrivially and produces an entire L -function.

The single computational fact that powers the entire theory of primes in progressions is the orthogonality of characters. It is the precise sense in which the $\varphi(m)$ characters modulo m form a complete, mutually independent system of “frequencies” on $(\mathbb{Z}/m\mathbb{Z})^\times$, and it is the device that extracts a single residue class from a sum over all characters.

Theorem 7.1.4: Orthogonality Relations

Let $m \in \mathbb{Z}_{>0}$. The Dirichlet characters modulo m satisfy:

1. *Column orthogonality.* For every integer n with $\gcd(n, m) = 1$,

$$\sum_{\chi \bmod m} \chi(n) = \begin{cases} \varphi(m) & \text{if } n \equiv 1 \pmod{m}, \\ 0 & \text{otherwise} \end{cases}$$

the sum running over all $\varphi(m)$ Dirichlet characters modulo m .

2. *Row orthogonality.* For every Dirichlet character χ modulo m ,

$$\sum_{n \bmod m} \chi(n) = \begin{cases} \varphi(m) & \text{if } \chi = \chi_0, \\ 0 & \text{otherwise} \end{cases}$$

the sum running over a complete set of residues n modulo m (equivalently, over the $\varphi(m)$ residues coprime to m , the others contributing 0).

Consequently, for any integers a, n with $\gcd(a, m) = 1$,

$$\frac{1}{\varphi(m)} \sum_{\chi \bmod m} \chi(n) \overline{\chi(a)} = \begin{cases} 1 & \text{if } n \equiv a \pmod{m}, \\ 0 & \text{otherwise} \end{cases} \quad (7.1)$$

so that the left-hand side is the indicator function of the residue class a modulo m , supported on integers coprime to m .

Proof:

Both relations are the orthogonality relations for characters of the finite abelian group $G = (\mathbb{Z}/m\mathbb{Z})^\times$, proved in [13]; the task here is to translate them into the language of Dirichlet characters and then to derive (7.1). Throughout, the Dirichlet characters modulo m are identified with the elements of $\widehat{G}$ by theorem 7.1.2, and a residue n coprime to m is identified with the corresponding element $\bar{n} \in G$.

Step 1: Row orthogonality. For a character $\tilde{\chi} \in \widehat{G}$, the group-character orthogonality of [13] states that $\sum_{g \in G} \tilde{\chi}(g)$ equals $|G|$ if $\tilde{\chi}$ is trivial and 0 otherwise. The direct argument is short: if $\tilde{\chi}$ is trivial the sum is $\sum_g 1 = |G| = \varphi(m)$, while if $\tilde{\chi}$ is nontrivial there is some $h \in G$ with $\tilde{\chi}(h) \neq 1$, and since $g \mapsto hg$ permutes G ,

$$S := \sum_{g \in G} \tilde{\chi}(g) = \sum_{g \in G} \tilde{\chi}(hg) = \tilde{\chi}(h) \sum_{g \in G} \tilde{\chi}(g) = \tilde{\chi}(h) S$$

whence $(1 - \tilde{\chi}(h))S = 0$ and $S = 0$. Translating to $\mathbb{Z}$: summing χ over a complete residue system modulo m , the residues with $\gcd(n, m) > 1$ contribute 0 by support, and the residues with $\gcd(n, m) = 1$ contribute $\sum_{g \in G} \tilde{\chi}(g)$, which is $\varphi(m)$ for $\chi = \chi_0$ and 0 otherwise. This is (2).

Step 2: Column orthogonality. Fix n coprime to m and put $g = \bar{n} \in G$. The dual statement of [13] is that $\sum_{\tilde{\chi} \in \widehat{G}} \tilde{\chi}(g)$ equals $|G|$ if g is the identity and 0 otherwise; this is row orthogonality

applied to the second-dual group $\widehat{\widehat{G}}$, under the canonical evaluation isomorphism $G \cong \widehat{\widehat{G}}$ sending g to the character $\tilde{\chi} \mapsto \tilde{\chi}(g)$. The direct argument mirrors Step 1: if g is the identity then every $\tilde{\chi}(g) = 1$ and the sum is $|\widehat{G}| = \varphi(m)$ by theorem 7.1.2; if $g \neq 1$ then, because $\widehat{G}$ separates points of G (a consequence of $\widehat{G} \cong G$, again [13]), there is a character ψ with $\psi(g) \neq 1$, and since $\tilde{\chi} \mapsto \psi\tilde{\chi}$ permutes $\widehat{G}$,

$$T := \sum_{\tilde{\chi} \in \widehat{G}} \tilde{\chi}(g) = \sum_{\tilde{\chi} \in \widehat{G}} (\psi\tilde{\chi})(g) = \psi(g) \sum_{\tilde{\chi} \in \widehat{G}} \tilde{\chi}(g) = \psi(g) T$$

so $(1 - \psi(g))T = 0$ and $T = 0$. Since $g = \bar{n}$ is the identity of G exactly when $n \equiv 1 \pmod{m}$, this is (1).

Step 3: The indicator identity. Let a be coprime to m , so $\bar{a} \in G$ is invertible, and let n be arbitrary. If $\gcd(n, m) > 1$ then $\chi(n) = 0$ for every χ , so the left-hand side of (7.1) is 0; and n is then not congruent to a modulo m (as $\gcd(a, m) = 1$), so the right-hand side is 0 as well. If $\gcd(n, m) = 1$, then $\overline{\chi(a)} = \chi(a)^{-1} = \chi(a^{-1})$ where a^{-1} is any integer with $aa^{-1} \equiv 1 \pmod{m}$ (it exists since $\gcd(a, m) = 1$), using $|\chi(a)| = 1$ and complete multiplicativity. Hence $\chi(n)\overline{\chi(a)} = \chi(na^{-1})$, and summing over χ and applying column orthogonality (1) to the integer na^{-1} ,

$$\sum_{\chi \bmod m} \chi(n)\overline{\chi(a)} = \sum_{\chi \bmod m} \chi(na^{-1}) = \begin{cases} \varphi(m) & \text{if } na^{-1} \equiv 1 \pmod{m}, \\ 0 & \text{otherwise} \end{cases}$$

The condition $na^{-1} \equiv 1 \pmod{m}$ is equivalent to $n \equiv a \pmod{m}$. Dividing by $\varphi(m)$ yields (7.1), as we sought to show.

Identity (7.1) is the engine of the entire chapter, and the mechanism it provides deserves to be stated plainly. To study primes $p \equiv a \pmod{m}$ one would like to sum some prime-counting quantity over exactly those primes, but the condition $p \equiv a$ is not multiplicative and cannot be inserted directly into a Dirichlet series. Orthogonality removes the obstruction. The right-hand side of (7.1) is precisely the indicator of the congruence $n \equiv a$, and the left-hand side replaces that single non-multiplicative condition by an average of the values $\chi(n)$, each of which *is* completely multiplicative. Inserting the average into a sum over primes and exchanging the order of summation turns one sum over the progression $a \bmod m$ into a weighted combination of $\varphi(m)$ sums over all primes, the χ -th of which is governed by the L -function $L(s, \chi)$ developed in section 7.2. The principal character χ_0 contributes the main term, carrying the pole that forces divergence, while each nonprincipal χ contributes a correction that stays bounded precisely because $L(1, \chi) \neq 0$. This decomposition, indicator-into-characters followed by character-into- L -function, is the strategy of Dirichlet's theorem in section 7.5; the non-multiplicative congruence is dissolved into multiplicative pieces, analyzed one character at a time, and reassembled.

For the functional equation the relevant refinements are not the group structure but two finer invariants of an individual character: whether it belongs to its modulus or is inherited from a smaller one, and how it behaves under negation.

Definition 7.1.5: Primitive Characters and the Conductor

Let χ be a Dirichlet character modulo m , and let d be a positive divisor of m . A Dirichlet character χ^* modulo d *induces* χ if

$$\chi(n) = \chi^*(n) \quad \text{for all } n \text{ with } \gcd(n, m) = 1$$

The character χ is *induced* by χ^* in this case, and is called *imprimitive* if it is induced by some character to a proper divisor $d < m$; it is *primitive* if no such proper divisor exists. The *conductor* of χ , written $f(\chi)$ or f_χ , is the smallest positive divisor d of m such that χ is induced by a character modulo d .

The conductor is well defined: the principal character modulo 1 induces the principal character modulo m (both equal 1 on the integers coprime to m , and modulo 1 every integer is coprime), so the set of divisors d for which χ is induced by a character modulo d is nonempty, and being a finite set of positive integers it has a least element. The character inducing χ at level $f(\chi)$ is unique and is itself primitive; this and the basic theory of induction (that the inducing character at the conductor is the “true” character, of which χ is a copy with extra Euler factors stripped out) are developed in [1] and recur in section 7.3, where only primitive characters carry a clean functional equation. A character is primitive exactly when it equals its own conductor-level character, $f(\chi) = m$. The principal character χ_0 modulo $m > 1$ is imprimitive, with conductor 1, since it is induced by the trivial character modulo 1; the only primitive character of conductor 1 is the trivial character itself.

Definition 7.1.6: Even and Odd Characters

A Dirichlet character χ modulo m is *even* if $\chi(-1) = +1$ and *odd* if $\chi(-1) = -1$. These two cases are exhaustive: complete multiplicativity gives $\chi(-1)^2 = \chi((-1)^2) = \chi(1) = 1$, so $\chi(-1) \in \{+1, -1\}$.

The parity records the behavior of χ under $n \mapsto -n$, since $\chi(-n) = \chi(-1)\chi(n) = \pm\chi(n)$. It is the single binary invariant that decides which gamma factor completes $L(s, \chi)$ into a symmetric function in section 7.3: an even character is completed with the factor $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$ already used for ζ in definition 6.3.1, while an odd character requires the shifted factor $\pi^{-(s+1)/2}\Gamma((s+1)/2)$, the shift by 1 in the argument reflecting the sign change. The principal character is even, since $\chi_0(-1) = 1$ whenever $\gcd(-1, m) = 1$, which always holds. Quadratic characters can be of either parity: the Legendre symbol $\left(\frac{\cdot}{p}\right)$ (definition 0.5.3) is even when $p \equiv 1 \pmod{4}$ and odd when $p \equiv 3 \pmod{4}$, by the first supplement to quadratic reciprocity (section 0.5) computing $\left(\frac{-1}{p}\right) = (-1)^{(p-1)/2}$.

The smallest moduli make all of these notions concrete. The next example tabulates every character modulo 4 and modulo 5, identifies the principal one, computes each parity, and locates the conductors, and it isolates the one nonprincipal character modulo 4 that recurs throughout the subject.

Example 7.1.7: Characters Modulo 4 and Modulo 5

Modulo 4. The unit group is $(\mathbb{Z}/4\mathbb{Z})^\times = \{1, 3\}$, cyclic of order $\varphi(4) = 2$ generated by 3 (since $3^2 = 9 \equiv 1$). By theorem 7.1.2 there are exactly two characters, each determined by its value on

the generator 3, which must be a square root of unity, that is, ± 1 :

$n \bmod 4$	1	3
$\chi_0(n)$	1	1
$\chi_1(n)$	1	-1

with $\chi_0(n) = \chi_1(n) = 0$ for even n . Here χ_0 is principal. The nonprincipal character χ_1 satisfies $\chi_1(-1) = \chi_1(3) = -1$, so χ_1 is odd; it is primitive, of conductor 4, since the only divisors of 4 are 1, 2, 4, the only character modulo 1 or 2 is principal (as $\varphi(1) = \varphi(2) = 1$), and a principal character cannot induce the nonprincipal χ_1 . Concretely χ_1 is the function

$$\chi_1(n) = \begin{cases} 1 & n \equiv 1 \pmod{4}, \\ -1 & n \equiv 3 \pmod{4}, \\ 0 & n \text{ even} \end{cases}$$

the unique nonprincipal character modulo 4. Its L -function is

$$L(s, \chi_1) = \sum_{n \geq 1} \frac{\chi_1(n)}{n^s} = 1 - \frac{1}{3^s} + \frac{1}{5^s} - \frac{1}{7^s} + \frac{1}{9^s} - \cdots = \sum_{k \geq 0} \frac{(-1)^k}{(2k+1)^s} = \beta(s)$$

the *Dirichlet beta function*, the L -function attached to the Gaussian integers $\mathbb{Z}[i]$, whose value $\beta(1) = 1 - \frac{1}{3} + \frac{1}{5} - \cdots = \pi/4$ is the Leibniz series. This character and its L -function recur in sections 7.2 and 7.3.

Modulo 5. The unit group is $(\mathbb{Z}/5\mathbb{Z})^\times = \{1, 2, 3, 4\}$, cyclic of order $\varphi(5) = 4$ generated by the primitive root 2, whose powers run $2^0 = 1$, $2^1 = 2$, $2^2 = 4$, $2^3 = 3$ modulo 5 (section 0.4). A character is determined by $\chi(2)$, which must be a fourth root of unity: one of $1, i, -1, -i$. Writing $i = \sqrt{-1}$, the four characters are

$n \bmod 5$	1	2	4	3
$\chi_0(n)$	1	1	1	1
$\psi(n)$	1	i	-1	$-i$
$\psi^2(n)$	1	-1	1	-1
$\psi^3(n)$	1	$-i$	-1	i

where ψ is the character with $\psi(2) = i$, and the columns are ordered by the powers $2^0, 2^1, 2^2, 2^3$ of the generator so that each row reads off the geometric progression $\psi^j(2)^k$. The four characters are $\chi_0 = \psi^0, \psi, \psi^2, \psi^3$, forming a cyclic group of order 4 generated by ψ , isomorphic to $(\mathbb{Z}/5\mathbb{Z})^\times$ as theorem 7.1.2 requires, with $\psi^{-1} = \bar{\psi} = \psi^3$. To read off the parities, note $-1 \equiv 4 \pmod{5}$, the column $n = 4$: thus $\chi_0(-1) = 1$ and $\psi^2(-1) = 1$ are even, while $\psi(-1) = -1$ and $\psi^3(-1) = -1$ are odd. The character ψ^2 , taking values ± 1 with $\psi^2(2) = -1$, is the quadratic character modulo 5; it is the Legendre symbol $\left(\frac{\cdot}{5}\right)$, recording that 1, 4 are the quadratic residues and 2, 3 the non-residues modulo 5, consistent with its evenness and $5 \equiv 1 \pmod{4}$. The three nonprincipal characters ψ, ψ^2, ψ^3 are primitive of conductor 5, since the only proper divisor of 5 is 1 and its sole character is principal, so no nonprincipal character modulo 5 is induced from below; the principal character χ_0 , by contrast, is imprimitive with conductor 1, induced by the trivial character modulo 1.

7.2 Dirichlet L -Functions

The previous section produced the characters as completely multiplicative arithmetic functions, the same class of function as $n \mapsto n^{-s}$ from which the Euler product of ζ was built. Feeding a character χ modulo m into a Dirichlet series produces the central object of the chapter, the Dirichlet L -function $L(s, \chi) = \sum_{n \geq 1} \chi(n) n^{-s}$. It is the character-twisted analogue of the Riemann zeta function: each prime contributes a single Euler factor, now weighted by the value $\chi(p)$, and the resulting product encodes the distribution of the residue class data that χ records. The Euler product, the analytic continuation, and the location of poles all transfer from the zeta theory of chapter 4, with one structural split that determines everything later. The principal character reproduces a lightly modified ζ , complete with its pole at $s = 1$; every other character produces a function holomorphic across the line $\operatorname{Re}(s) = 1$, and the value of that function at $s = 1$, shown nonzero in section 7.4, is the quantity that drives Dirichlet's theorem on primes in arithmetic progressions in section 7.5.

This section defines $L(s, \chi)$, factors it as an Euler product, and continues it past the line $\operatorname{Re}(s) = 1$: trivially for the principal character, by reduction to ζ , and by Abel summation against the bounded character sums for every nonprincipal character. The example at the end evaluates the simplest nonprincipal L -functions to small moduli, including the one whose value at $s = 1$ is the Leibniz series $\pi/4$.

Definition 7.2.1: Dirichlet L -Function

Let χ be a Dirichlet character modulo m (definition 7.1.1). The *Dirichlet L -function* attached to χ is the Dirichlet series

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$$

a function of the complex variable s . The series converges absolutely on the half-plane $\operatorname{Re}(s) > 1$.

Absolute convergence on $\operatorname{Re}(s) > 1$ is immediate from the bound $|\chi(n)| \leq 1$ of theorem 7.1.2(4): every nonzero value of χ is a root of unity, so $|\chi(n)| \in \{0, 1\}$, and therefore

$$\sum_{n=1}^{\infty} \left| \frac{\chi(n)}{n^s} \right| = \sum_{n=1}^{\infty} \frac{|\chi(n)|}{n^{\sigma}} \leq \sum_{n=1}^{\infty} \frac{1}{n^{\sigma}}$$

where $\sigma = \operatorname{Re}(s)$, and the comparison series $\sum_n n^{-\sigma}$ converges for $\sigma > 1$. The abscissa of absolute convergence (definition 2.1.3) of $L(s, \chi)$ is thus at most 1. For the principal character χ_0 it equals 1, since $L(s, \chi_0)$ differs from $\zeta(s)$ by finitely many Euler factors (proposition 7.2.3 below) and so inherits the divergence of the harmonic series at $s = 1$; for nonprincipal characters the abscissa of absolute convergence is still 1, but the series converges conditionally on the larger half-plane $\operatorname{Re}(s) > 0$, the content of theorem 7.2.4. On $\operatorname{Re}(s) > 1$ the series defines a holomorphic function, by the general analyticity of Dirichlet series in their half-plane of absolute convergence (theorem 2.1.7).

The same completely multiplicative structure that gave ζ its Euler product gives one to every L -function. This is the first point at which the residue class information carried by χ enters the analytic picture: the factor at a prime p records the value $\chi(p)$, which depends only on $p \bmod m$.

Theorem 7.2.2: The Euler Product

Let χ be a Dirichlet character modulo m . For $\operatorname{Re}(s) > 1$,

$$L(s, \chi) = \prod_p \frac{1}{1 - \chi(p) p^{-s}}$$

the product running over all primes and converging absolutely. The factor at a prime $p \mid m$ equals 1, since $\chi(p) = 0$ there, so the product is effectively over the primes not dividing m . In particular $L(s, \chi) \neq 0$ for $\operatorname{Re}(s) > 1$.

Proof:

A Dirichlet character is completely multiplicative by condition (i) of definition 7.1.1, and its abscissa of absolute convergence is $\sigma_a(\chi) \leq 1$, as recorded after definition 7.2.1. For $\operatorname{Re}(s) > 1 \geq \sigma_a(\chi)$ the Euler product for multiplicative functions (theorem 2.3.1) applies, and since χ is totally multiplicative the local factor at p is the geometric series

$$\sum_{k=0}^{\infty} \frac{\chi(p^k)}{p^{ks}} = \sum_{k=0}^{\infty} \left(\frac{\chi(p)}{p^s} \right)^k = \frac{1}{1 - \chi(p) p^{-s}}$$

the geometric series converging because $|\chi(p)p^{-s}| = |\chi(p)|p^{-\sigma} \leq p^{-\sigma} < 1$ for $\sigma > 1$. This gives the stated product, and theorem 2.3.1 guarantees its absolute convergence. For a prime $p \mid m$ the value $\chi(p) = 0$ by condition (iii) of definition 7.1.1, so the factor is $(1 - 0)^{-1} = 1$ and drops out, leaving the product over $p \nmid m$.

For the non-vanishing, the local factors $(1 - \chi(p)p^{-s})^{-1}$ are reciprocals of nonzero complex numbers, hence themselves nonzero: at $p \mid m$ the factor is 1, and at $p \nmid m$ the quantity $1 - \chi(p)p^{-s}$ has $|\chi(p)p^{-s}| < 1$, so it cannot vanish. An absolutely convergent infinite product with no vanishing factor has a nonzero value (proposition 2.3.4), so $L(s, \chi) \neq 0$ throughout $\operatorname{Re}(s) > 1$, as we sought to show.

The Euler product is what makes $L(s, \chi)$ an instrument for studying primes rather than a formal series: it converts the global function into an assembly of one factor per prime, with the residue class of p entering through $\chi(p)$. Taking a logarithm of the product, as in proposition 2.3.5, will turn $L(s, \chi)$ into a sum over prime powers weighted by χ , and orthogonality (theorem 7.1.4) will then sieve out a single residue class from the family of such sums over all χ . The non-vanishing on $\operatorname{Re}(s) > 1$ recorded here is the easy half of the non-vanishing story, valid wherever the product converges; the non-vanishing that the theory of primes in progressions actually needs is at the boundary point $s = 1$ itself, where the product no longer converges and a separate argument is required (section 7.4).

The two regimes promised in the chapter introduction now separate. The principal character carries the pole; everything else is holomorphic at $s = 1$. The principal case is settled first, by direct comparison with ζ .

Proposition 7.2.3: The Principal L-Function and ζ

Let χ_0 be the principal character modulo m (definition 7.1.3). For $\operatorname{Re}(s) > 1$,

$$L(s, \chi_0) = \zeta(s) \prod_{p|m} (1 - p^{-s})$$

Consequently $L(s, \chi_0)$ extends to a meromorphic function on $\operatorname{Re}(s) > 0$ whose only singularity there is a simple pole at $s = 1$, with residue

$$\operatorname{Res}_{s=1} L(s, \chi_0) = \prod_{p|m} \left(1 - \frac{1}{p}\right) = \frac{\varphi(m)}{m}$$

Proof:

Step 1: Factoring out the missing Euler factors. On $\operatorname{Re}(s) > 1$ the Euler product of theorem 7.2.2 for χ_0 runs over the primes $p \nmid m$, since $\chi_0(p) = 1$ for $p \nmid m$ and $\chi_0(p) = 0$ for $p|m$. The product for ζ (corollary 2.3.2) runs over all primes, with factor $(1 - p^{-s})^{-1}$ at each. Both products converge absolutely for $\operatorname{Re}(s) > 1$, so they may be compared factor by factor and the finite collection of factors at the primes $p|m$ separated off:

$$L(s, \chi_0) = \prod_{p \nmid m} \frac{1}{1 - p^{-s}} = \left(\prod_p \frac{1}{1 - p^{-s}} \right) \prod_{p|m} (1 - p^{-s}) = \zeta(s) \prod_{p|m} (1 - p^{-s})$$

the middle equality multiplying and dividing by the finitely many factors at $p|m$, which is legitimate because each such factor $(1 - p^{-s})^{-1}$ is finite and nonzero for $\operatorname{Re}(s) > 1$. This is the displayed identity.

Step 2: Continuation and the pole. The finite product $P(s) = \prod_{p|m} (1 - p^{-s})$ is an entire function of s , a finite product of entire functions. By theorem 4.2.1 the factor $\zeta(s)$ continues meromorphically to $\operatorname{Re}(s) > 0$ with a single simple pole at $s = 1$ of residue 1 and no other singularity. The product $\zeta(s)P(s)$ therefore continues to a meromorphic function on $\operatorname{Re}(s) > 0$, and since P is entire, its only possible pole is at $s = 1$, inherited from ζ . Whether the pole survives is decided by the value of P at $s = 1$: it survives precisely when $P(1) \neq 0$. Here

$$P(1) = \prod_{p|m} \left(1 - \frac{1}{p}\right)$$

a finite product of factors each strictly between 0 and 1 (for every prime $p \geq 2$ one has $0 < 1 - 1/p < 1$), hence $P(1) \neq 0$. The pole of ζ at $s = 1$ thus passes to $L(s, \chi_0)$ as a simple pole, and there is no other singularity on $\operatorname{Re}(s) > 0$.

Step 3: The residue. Since P is holomorphic at $s = 1$ and ζ has a simple pole there,

$$\operatorname{Res}_{s=1} L(s, \chi_0) = \operatorname{Res}_{s=1} (\zeta(s)P(s)) = P(1) \operatorname{Res}_{s=1} \zeta(s) = P(1) \cdot 1 = \prod_{p|m} \left(1 - \frac{1}{p}\right)$$

using $\operatorname{Res}_{s=1} \zeta(s) = 1$ from corollary 4.2.2 and the fact that the residue of a product of a holomorphic function with a simple pole is the holomorphic value times the residue. The final product equals $\varphi(m)/m$: the totient is multiplicative with $\varphi(p^k) = p^k(1 - 1/p)$ at prime powers (definition 0.3.5 and the multiplicativity of φ), so writing $m = \prod_{p|m} p^{a_p}$ gives

$$\frac{\varphi(m)}{m} = \prod_{p|m} \frac{\varphi(p^{a_p})}{p^{a_p}} = \prod_{p|m} \left(1 - \frac{1}{p}\right)$$

which is $P(1)$, completing the proof.

The principal L -function is ζ with the Euler factors at the primes dividing m deleted. Removing finitely many nonzero, holomorphic factors changes neither the meromorphic continuation nor the location of the pole, only the residue: the residue 1 of ζ is scaled by $\prod_{p|m} (1 - 1/p) = \varphi(m)/m$, the density of the integers coprime to m . This is the analytic shadow of a counting fact. The pole of $L(s, \chi_0)$ at $s = 1$ measures the abundance of the integers coprime to m , exactly as the pole of ζ measures the abundance of all integers, and the residue $\varphi(m)/m$ is the proportion of residue classes modulo m that are coprime to m . In the decomposition of section 7.5 this pole is the main term: it is the single source of divergence, and the principal character is the only character of modulus m that contributes one. Every nonprincipal character must therefore contribute a bounded correction, and the next theorem is what guarantees boundedness.

For a nonprincipal character the series $\sum_n \chi(n)n^{-s}$ does not converge absolutely at $s = 1$, but it converges conditionally on a much larger half-plane than absolute convergence reaches. The mechanism is cancellation: the partial sums $\sum_{n \leq x} \chi(n)$ do not grow, because over each complete period the values of χ sum to zero. Abel summation converts that boundedness into convergence of the series.

Theorem 7.2.4: Continuation for Nonprincipal Characters

Let χ be a nonprincipal Dirichlet character modulo m . Then the series $\sum_{n \geq 1} \chi(n)n^{-s}$ converges for every s with $\operatorname{Re}(s) > 0$, and the convergence is uniform on compact subsets of the half-plane $\operatorname{Re}(s) > 0$. Consequently $L(s, \chi)$ is holomorphic on $\operatorname{Re}(s) > 0$. In particular $L(s, \chi)$ is holomorphic at $s = 1$; the principal L -function aside, no L -function of modulus m has a pole.

Proof:

Step 1: The character sums are bounded. Write $S(x) = \sum_{n \leq x} \chi(n)$ for the partial sums of the coefficients. The claim is that $|S(x)| \leq \varphi(m)$ for all $x \geq 0$. The values of χ are periodic modulo m (condition (ii) of definition 7.1.1), so the sum over any block of m consecutive integers is the

sum over one full period,

$$\sum_{n=a+1}^{a+m} \chi(n) = \sum_{n \bmod m} \chi(n) = 0$$

the last equality being row orthogonality (theorem 7.1.4(2)), which gives 0 precisely because $\chi \neq \chi_0$. Hence in $S(x)$ all complete blocks of length m contribute nothing, and only the final incomplete block of fewer than m terms survives. That block has at most m terms, at most $\varphi(m)$ of which are nonzero (the integers coprime to m in a period number exactly $\varphi(m)$, the rest having $\chi(n) = 0$), and each nonzero term has $|\chi(n)| = 1$ by theorem 7.1.2(4). Therefore

$$|S(x)| \leq \varphi(m) \quad \text{for all } x \geq 0$$

a single bound independent of x .

Step 2: Abel summation. Fix s with $\sigma = \operatorname{Re}(s) > 0$, and apply Abel's summation formula (theorem 1.4.1) with coefficients $a_n = \chi(n)$, so that the summatory function is $A(t) = S(t)$, and with the test function $f(t) = t^{-s}$, which is continuously differentiable on $[1, \infty)$ with $f'(t) = -s t^{-s-1}$. For $x \geq 1$,

$$\sum_{n \leq x} \frac{\chi(n)}{n^s} = S(x) x^{-s} + s \int_1^x S(t) t^{-s-1} dt \quad (7.2)$$

the sign on the integral following from $f'(t) = -s t^{-s-1}$. Each piece is now controlled by the bound of Step 1. The boundary term satisfies $|S(x) x^{-s}| \leq \varphi(m) x^{-\sigma} \rightarrow 0$ as $x \rightarrow \infty$, since $\sigma > 0$. The integrand satisfies $|S(t) t^{-s-1}| \leq \varphi(m) t^{-\sigma-1}$, and $\int_1^\infty t^{-\sigma-1} dt = 1/\sigma < \infty$, so the integral $\int_1^\infty S(t) t^{-s-1} dt$ converges absolutely. Letting $x \rightarrow \infty$ in (7.2), the boundary term vanishes and the partial sums converge:

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = s \int_1^{\infty} S(t) t^{-s-1} dt \quad (\operatorname{Re}(s) > 0) \quad (7.3)$$

This establishes convergence of the series for every s with $\operatorname{Re}(s) > 0$. The argument is the boundedness criterion behind corollary 1.4.9: bounded partial sums against a factor decaying to 0 force the series to converge, here for the complex test function t^{-s} rather than the monotone real one of the corollary.

Step 3: Holomorphy. It remains to see that the convergence is locally uniform, so that the limit is holomorphic. Let K be a compact subset of $\{\operatorname{Re}(s) > 0\}$ and set $\sigma_0 = \min_{s \in K} \operatorname{Re}(s) > 0$ and $C = \max_{s \in K} |s| < \infty$. For $s \in K$ the tail of the integral representation (7.3) beyond x is bounded, using Step 1, by

$$\left| s \int_x^\infty S(t) t^{-s-1} dt \right| \leq C \varphi(m) \int_x^\infty t^{-\sigma_0-1} dt = \frac{C \varphi(m)}{\sigma_0} x^{-\sigma_0}$$

together with the boundary term $|S(x) x^{-s}| \leq \varphi(m) x^{-\sigma_0}$ from (7.2). Both bounds are independent of $s \in K$ and tend to 0 as $x \rightarrow \infty$, so the partial sums $\sum_{n \leq x} \chi(n) n^{-s}$ converge to $L(s, \chi)$ uniformly on K . Each partial sum is a finite sum of the entire functions n^{-s} , hence holomorphic, and a locally uniform limit of holomorphic functions is holomorphic ([16]). Therefore $L(s, \chi)$ is holomorphic on $\operatorname{Re}(s) > 0$. In particular it is holomorphic at $s = 1$, with no pole there, completing the proof.

The contrast with the principal case is the structural fact on which the rest of the chapter turns. The principal character recovers a tweaked zeta function, carrying the pole at $s = 1$ that forces divergence and supplies the main term in every count weighted by χ_0 . A nonprincipal character, by contrast, yields a function holomorphic across the entire half-plane $\operatorname{Re}(s) > 0$, the pole having been killed by the cancellation of the character sum over each period. The single number that now matters is the *value* $L(1, \chi)$. Because $L(s, \chi)$ is holomorphic at $s = 1$, this value is a finite complex number rather than an infinity, and the decomposition of section 7.5 expresses the count of primes in a fixed residue class as the pole-carrying principal term plus a sum of corrections, one for each nonprincipal χ , the χ -th correction governed by $\log L(s, \chi)$ near $s = 1$. The correction stays bounded, and so fails to cancel the main term, exactly when $L(1, \chi) \neq 0$, so that $\log L(s, \chi)$ does not run off to $-\infty$. That nonvanishing, $L(1, \chi) \neq 0$ for every nonprincipal χ , is the one fact with no counterpart in the zeta theory, where the pole makes the analogous statement automatic; it is established in section 7.4 and converted into Dirichlet's theorem in section 7.5.

The smallest nonprincipal characters give L -functions that can be written out term by term, and one of them evaluates at $s = 1$ to a series whose sum is classical.

Example 7.2.5: L -Functions to Small Moduli

Modulus 4. The unique nonprincipal character modulo 4 is the odd character χ_1 of example 7.1.7, with $\chi_1(n) = 1$ for $n \equiv 1$, $\chi_1(n) = -1$ for $n \equiv 3$, and $\chi_1(n) = 0$ for n even, all modulo 4. Its L -function is

$$L(s, \chi_1) = \sum_{n=1}^{\infty} \frac{\chi_1(n)}{n^s} = 1 - \frac{1}{3^s} + \frac{1}{5^s} - \frac{1}{7^s} + \frac{1}{9^s} - \cdots = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^s} = \beta(s)$$

the Dirichlet beta function. By theorem 7.2.4 this series converges for every s with $\operatorname{Re}(s) > 0$, so in particular at $s = 1$ it converges to the value

$$L(1, \chi_1) = \beta(1) = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots = \frac{\pi}{4}$$

the Leibniz series. The evaluation $\beta(1) = \pi/4$ follows from the power series $\arctan x = \sum_{k \geq 0} (-1)^k x^{2k+1}/(2k+1)$, valid for $|x| \leq 1$: the series at $x = 1$ is $\sum_{k \geq 0} (-1)^k/(2k+1)$, and since the series converges at the endpoint $x = 1$, Abel's limit theorem identifies its sum with $\lim_{x \rightarrow 1^-} \arctan x = \arctan 1 = \pi/4$ ([16]). The partial sums of $\beta(1)$ converge slowly, as the alternating tail bound $|\beta(1) - \sum_{k=0}^N (-1)^k/(2k+1)| \leq 1/(2N+3)$ shows; the first several are

N	$\sum_{k=0}^N \frac{(-1)^k}{2k+1}$
0	1.000000
1	0.666667
2	0.866667
3	0.723810
4	0.834921
9	0.760460
49	0.780399

drifting toward $\pi/4 = 0.785398 \dots$ at the rate $1/(2N)$ dictated by the alternating bound. That $L(1, \chi_1) = \pi/4 \neq 0$ is the nonvanishing of section 7.4 made explicit in the first nontrivial case, and

it is the analytic input that proves there are infinitely many primes in each of the progressions $1, 5, 9, \dots$ and $3, 7, 11, \dots$ modulo 4.

Modulus 3. The unit group $(\mathbb{Z}/3\mathbb{Z})^\times = \{1, 2\}$ is cyclic of order $\varphi(3) = 2$, so there are two characters: the principal χ_0 and one nonprincipal character χ with $\chi(1) = 1$, $\chi(2) = \chi(-1) = -1$, and $\chi(n) = 0$ for $3|n$. The character χ is odd, of conductor 3. Its L -function is

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = 1 - \frac{1}{2^s} + \frac{1}{4^s} - \frac{1}{5^s} + \frac{1}{7^s} - \frac{1}{8^s} + \dots$$

the signs running $+, -$ over the residues $1, 2$ modulo 3 and the multiples of 3 omitted. Again theorem 7.2.4 gives convergence and holomorphy on $\operatorname{Re}(s) > 0$, and the value

$$L(1, \chi) = 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{5} + \frac{1}{7} - \frac{1}{8} + \dots = \frac{\pi}{3\sqrt{3}}$$

is nonzero, the closed form $\pi/(3\sqrt{3})$ being the modulus-3 analogue of the Leibniz series; its nonvanishing yields infinitely many primes in each of the progressions $1 \bmod 3$ and $2 \bmod 3$. The principal character modulo 3, by contrast, gives $L(s, \chi_0) = \zeta(s)(1 - 3^{-s})$ by proposition 7.2.3, with a simple pole at $s = 1$ of residue $1 - 1/3 = 2/3 = \varphi(3)/3$.

7.3 Gauss Sums and the Functional Equation

The functional equation of ζ (theorem 6.3.3) was assembled in chapter 6 from two ingredients: the gamma factor $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$, which completed the bare Dirichlet series into a symmetric object $\Lambda(s) = \Gamma_{\mathbb{R}}(s)\zeta(s)$, and the theta transformation $\theta(1/t) = \sqrt{t}\theta(t)$ (theorem 6.2.4), which supplied the symmetry $\Lambda(s) = \Lambda(1-s)$ by feeding through a Mellin transform. The L -function of a primitive Dirichlet character satisfies a functional equation of the same shape, relating s to $1-s$, but with one new feature absent from the zeta case: the reflection is no longer an exact equality but carries a constant of modulus 1, the *root number*, and that constant is built from a finite exponential sum attached to the character, the Gauss sum. The presence of the constant is forced by the loss of self-duality. The theta series $\theta(t)$ transforms into itself under $t \mapsto 1/t$ because the trivial character is its own conjugate; a character-twisted theta series transforms into the series of the *conjugate* character, and the Gauss sum is precisely the proportionality factor that mediates the exchange.

This section constructs the Gauss sum, computes its absolute value (the single fact that makes the root number have modulus 1), defines the completed L -function with the gamma factor chosen by the parity of the character, and proves the functional equation by twisting Riemann's theta argument by χ . The proof is given in full for even characters, where the gamma factor is the same $\Gamma_{\mathbb{R}}$ used for ζ and the twisted theta function is the direct analogue of θ ; the odd case requires a theta series weighted by n and is carried through in full along the same path. The closing example carries the entire apparatus through the smallest nontrivial character, modulo 4, where the Gauss sum is $2i$ and the root number is 1.

Definition 7.3.1: The Gauss Sum

Let χ be a Dirichlet character modulo m (definition 7.1.1). The *Gauss sum* attached to χ is the finite exponential sum

$$\tau(\chi) = \sum_{a=1}^m \chi(a) e^{2\pi ia/m}$$

the sum running over a complete set of residues modulo m . More generally, for an integer n the *twisted Gauss sum* is

$$\tau_n(\chi) = \sum_{a=1}^m \chi(a) e^{2\pi ina/m}$$

so that $\tau(\chi) = \tau_1(\chi)$.

The Gauss sum pairs the character χ against the additive character $a \mapsto e^{2\pi ia/m}$ of $\mathbb{Z}/m\mathbb{Z}$, and it is the basic object linking the multiplicative structure that χ records to the additive structure carried by the exponential. Each summand has modulus $|\chi(a)| \leq 1$, so the trivial bound is $|\tau(\chi)| \leq \varphi(m)$ (only the $\varphi(m)$ residues coprime to m contribute, the rest having $\chi(a) = 0$); the content of the next theorem is that for a primitive character this bound is improved to the exact value $|\tau(\chi)| = \sqrt{m}$, a square-root cancellation that is the source of the root number's modulus. The terminology and the definition specialize, when χ is the Legendre symbol $\left(\frac{\cdot}{p}\right)$ (definition 0.5.3) modulo a prime p , to the quadratic Gauss sum $\sum_a \left(\frac{a}{p}\right) e^{2\pi ia/p}$ whose evaluation governs quadratic reciprocity (section 0.5); the present definition is the extension of that classical object to an arbitrary character.

The absolute value of the Gauss sum, and the closely related product $\tau(\chi)\tau(\bar{\chi})$, rest on a single algebraic identity: for a primitive character the twisted sum $\tau_n(\chi)$ is a scalar multiple of $\tau(\chi)$, the scalar being $\bar{\chi}(n)$. This *separability* is the key lemma, and it is the one place where primitivity (definition 7.1.5) is indispensable.

Theorem 7.3.2: Modulus of the Gauss Sum

Let χ be a primitive Dirichlet character modulo m . Then:

1. *Separability.* For every integer n ,

$$\tau_n(\chi) = \bar{\chi}(n) \tau(\chi)$$

2. *Modulus.* The Gauss sum has absolute value

$$|\tau(\chi)| = \sqrt{m}$$

3. *Conjugate and product.* The Gauss sums of χ and $\bar{\chi}$ satisfy

$$\tau(\bar{\chi}) = \chi(-1) \overline{\tau(\chi)}, \quad \tau(\chi) \tau(\bar{\chi}) = \chi(-1) m$$

Proof:

The argument establishes separability first, then derives the modulus by summing $|\tau_n(\chi)|^2$ over n in two ways, and finally reads off the conjugate relation.

Step 1: Separability when $\gcd(n, m) = 1$. Suppose first that n is coprime to m , so $\bar{n}$ is a unit modulo m . As a runs over a complete residue system modulo m , so does $c = na$, and multiplicativity gives $\chi(a) = \chi(n^{-1}c) = \chi(n)^{-1}\chi(c) = \bar{\chi}(n)\chi(c)$, the last equality because $|\chi(n)| = 1$ forces $\chi(n)^{-1} = \overline{\chi(n)} = \bar{\chi}(n)$ (theorem 7.1.2). The exponential is unchanged in the new variable, $e^{2\pi i na/m} = e^{2\pi i c/m}$, so

$$\tau_n(\chi) = \sum_{a \bmod m} \chi(a) e^{2\pi i na/m} = \sum_{c \bmod m} \bar{\chi}(n) \chi(c) e^{2\pi i c/m} = \bar{\chi}(n) \tau(\chi)$$

This step uses only multiplicativity and the invertibility of n , not primitivity, and it holds for every character.

Step 2: Separability when $\gcd(n, m) > 1$. Here $\bar{\chi}(n) = 0$, so the claim is $\tau_n(\chi) = 0$. This is the assertion that requires primitivity: for an imprimitive character $\tau_n(\chi)$ can be nonzero at integers n sharing a factor with m . Write $d = \gcd(n, m) > 1$ and $m = dm'$, so m' is a *proper* divisor of m , that is, $m' < m$. Write $n = dn_0$; then $\gcd(n_0, m') = 1$, since any common prime factor of n_0 and m' would multiply d to a larger common divisor of n and m . The exponential simplifies to $e^{2\pi i na/m} = e^{2\pi i n_0 a/m'}$, which depends only on a modulo m' . Grouping the residues a modulo m by their reduction modulo m' , so that a is sorted into the class $a \equiv b \pmod{m'}$ indexed by a residue b modulo m' ,

$$\tau_n(\chi) = \sum_{b \bmod m'} e^{2\pi i n_0 b/m'} \sum_{\substack{a \bmod m \\ a \equiv b \pmod{m'}}} \chi(a) = \sum_{b \bmod m'} e^{2\pi i n_0 b/m'} T(b), \quad T(b) := \sum_{\substack{a \bmod m \\ a \equiv b \pmod{m'}}} \chi(a)$$

the inner sum $T(b)$ running over the d residues modulo m in the class of b modulo m' . It suffices to show $T(b) = 0$ for every b .

The primitivity criterion. The vanishing of every $T(b)$ rests on a restatement of primitivity (definition 7.1.5) in terms of the reduction map. The claim is that, for a primitive χ modulo m and any proper divisor $m' < m$, there exists an integer c with $c \equiv 1 \pmod{m'}$, $\gcd(c, m) = 1$, and $\chi(c) \neq 1$. Suppose, for contradiction, that no such c exists, so $\chi(c) = 1$ for every c coprime to m with $c \equiv 1 \pmod{m'}$. This forces χ to be induced from level m' , contradicting primitivity, as follows. Define a function χ^* modulo m' by setting $\chi^*(b) = 0$ when $\gcd(b, m') > 1$, and otherwise choosing any integer a with $a \equiv b \pmod{m'}$ and $\gcd(a, m) = 1$ and setting $\chi^*(b) = \chi(a)$. Such an a exists whenever $\gcd(b, m') = 1$: let r be the product of the primes dividing m but not m' , so $\gcd(m', r) = 1$, and use the Chinese Remainder Theorem (theorem 0.3.7) on the coprime moduli m' and r to solve $a \equiv b \pmod{m'}$ and $a \equiv 1 \pmod{r}$ simultaneously. Then a is coprime to m : a prime $p|m'$ does not divide a because $a \equiv b \pmod{m'}$ with $\gcd(b, m') = 1$, while a prime $p|m$ with $p \nmid m'$ divides r , so $a \equiv 1 \pmod{p}$ gives $p \nmid a$; these exhaust the prime divisors of m . The value $\chi^*(b)$ is independent of the choice of a : if a, a' both lie in the class of b modulo m' and are coprime to m , then $c = a'a^{-1}$ satisfies $c \equiv 1 \pmod{m'}$ and $\gcd(c, m) = 1$, so $\chi(c) = 1$ by hypothesis and $\chi(a') = \chi(c)\chi(a) = \chi(a)$. The function χ^* is completely multiplicative: for b, b' both coprime to m' , unit lifts a, a' have product aa' a unit lift of bb' , so $\chi^*(bb') = \chi(aa') = \chi(a)\chi(a') = \chi^*(b)\chi^*(b')$, while if either of b, b' shares a factor

with m' then so does bb' and both sides are 0. Since χ^* agrees with χ on the integers coprime to m , the character χ is induced by χ^* modulo the proper divisor $m' < m$, contradicting the primitivity of χ . This proves the criterion.

Vanishing of the inner sums. Fix b and let c be the integer supplied by the criterion, with $c \equiv 1 \pmod{m'}$, $\gcd(c, m) = 1$, and $\chi(c) \neq 1$. Multiplication by c permutes the residues modulo m (as c is a unit) and preserves residues modulo m' (as $c \equiv 1 \pmod{m'}$), so it permutes the set $\{a \pmod{m} : a \equiv b \pmod{m'}\}$ over which $T(b)$ is summed. Reindexing that sum by $a \mapsto ca$ and using complete multiplicativity,

$$T(b) = \sum_{\substack{a \pmod{m} \\ a \equiv b \pmod{m'}}} \chi(a) = \sum_{\substack{a \pmod{m} \\ a \equiv b \pmod{m'}}} \chi(ca) = \chi(c) T(b)$$

whence $(1 - \chi(c))T(b) = 0$, and since $\chi(c) \neq 1$ this gives $T(b) = 0$. This is the same averaging device that proved row orthogonality (theorem 7.1.4), applied now to the restriction of χ to the residues fixed modulo m' rather than to the full group. As b was arbitrary, every inner sum vanishes, so $\tau_n(\chi) = 0$ when $\gcd(n, m) > 1$. Combining with Step 1 gives $\tau_n(\chi) = \bar{\chi}(n)\tau(\chi)$ for all n , which is (1); the same vanishing is carried out by the conductor-theoretic route in [1, Ch. 9] and [4, Ch. 9].

Step 3: The modulus. Evaluate the sum

$$S = \sum_{n \pmod{m}} |\tau_n(\chi)|^2$$

in two ways. On one hand, separability (1) gives $\tau_n(\chi) = \bar{\chi}(n)\tau(\chi)$, so $|\tau_n(\chi)|^2 = |\bar{\chi}(n)|^2 |\tau(\chi)|^2$, which equals $|\tau(\chi)|^2$ when $\gcd(n, m) = 1$ and 0 otherwise; summing over the $\varphi(m)$ residues coprime to m ,

$$S = \varphi(m) |\tau(\chi)|^2$$

On the other hand, expanding the square directly,

$$|\tau_n(\chi)|^2 = \tau_n(\chi) \overline{\tau_n(\chi)} = \sum_{a \pmod{m}} \sum_{b \pmod{m}} \chi(a) \bar{\chi}(b) e^{2\pi i n(a-b)/m}$$

and summing over a complete residue system n modulo m , the geometric sum $\sum_{n \pmod{m}} e^{2\pi i n(a-b)/m}$ equals m when $a \equiv b \pmod{m}$ and 0 otherwise. Only the diagonal $a = b$ survives:

$$S = m \sum_{a \pmod{m}} \chi(a) \bar{\chi}(a) = m \sum_{a \pmod{m}} |\chi(a)|^2 = m \varphi(m)$$

the final equality because $|\chi(a)|^2 = 1$ for the $\varphi(m)$ residues coprime to m and 0 for the rest. Equating the two evaluations, $\varphi(m) |\tau(\chi)|^2 = m \varphi(m)$, and since $\varphi(m) \neq 0$, this gives $|\tau(\chi)|^2 = m$, hence $|\tau(\chi)| = \sqrt{m}$, which is (2).

Step 4: The conjugate relation and the product. Conjugating the defining sum and using $e^{2\pi i a/m} = e^{-2\pi i a/m} = e^{2\pi i (-1)a/m}$,

$$\overline{\tau(\chi)} = \sum_{a \pmod{m}} \overline{\chi(a)} e^{-2\pi i a/m} = \sum_{a \pmod{m}} \bar{\chi}(a) e^{2\pi i (-1)a/m} = \tau_{-1}(\bar{\chi})$$

the last sum being the twisted Gauss sum of $\bar{\chi}$ at $n = -1$. Since $\bar{\chi}$ is primitive (its conductor equals that of χ), separability (1) applied to $\bar{\chi}$ at $n = -1$ gives $\tau_{-1}(\bar{\chi}) = \bar{\chi}(-1) \tau(\bar{\chi}) = \chi(-1) \tau(\bar{\chi})$. Therefore $\tau(\chi) = \chi(-1) \tau(\bar{\chi})$, and multiplying both sides by $\chi(-1) \in \{\pm 1\}$ (definition 7.1.6) yields $\tau(\bar{\chi}) = \chi(-1) \tau(\chi)$. Substituting this into the product,

$$\tau(\chi) \tau(\bar{\chi}) = \tau(\chi) \cdot \chi(-1) \overline{\tau(\chi)} = \chi(-1) |\tau(\chi)|^2 = \chi(-1) m$$

using (2), which establishes (3), completing the proof.

The square-root cancellation $|\tau(\chi)| = \sqrt{m}$ is the arithmetic fact that controls the entire functional equation. The trivial bound $\varphi(m)$ would make the Gauss sum as large as the number of nonzero terms; the actual size $\sqrt{m}$ is the square root of the number of terms, the hallmark of the values $\chi(a)e^{2\pi ia/m}$ behaving like independent unit vectors whose sum has length $\sqrt{m}$ rather than $\varphi(m)$. This cancellation is what will force the root number to lie on the unit circle: the root number is $\tau(\chi)$ divided by $\sqrt{m}$ and a power of i , so the equality $|\tau(\chi)| = \sqrt{m}$ is exactly the statement that the resulting constant has modulus 1. The product formula $\tau(\chi)\tau(\bar{\chi}) = \chi(-1)m$ is the algebraic identity that makes the two-sided functional equation consistent: applying the reflection twice, once for χ and once for $\bar{\chi}$, multiplies the two root numbers, and that product must be 1 for the double reflection to return to the start. Separability, the engine behind both, is the precise sense in which a primitive character “diagonalizes” the additive characters of $\mathbb{Z}/m\mathbb{Z}$, each twist τ_n being the same vector $\tau(\chi)$ scaled by the character value $\bar{\chi}(n)$.

The completed L -function is now defined by attaching the gamma factor dictated by the parity of χ , exactly as $\Lambda(s) = \Gamma_{\mathbb{R}}(s)\zeta(s)$ attached $\Gamma_{\mathbb{R}}$ to ζ . The parity enters as a shift of the gamma argument: an even character keeps the factor $\Gamma_{\mathbb{R}}$ of definition 6.3.1, while an odd character shifts the argument by 1, reflecting the sign change $\chi(-n) = -\chi(n)$ that the symmetric theta series cannot see and that an antisymmetric one must carry.

Definition 7.3.3: The Completed Dirichlet L -Function

Let χ be a primitive Dirichlet character modulo m , with parity $a \in \{0, 1\}$ defined by

$$a = \begin{cases} 0 & \text{if } \chi \text{ is even } (\chi(-1) = +1), \\ 1 & \text{if } \chi \text{ is odd } (\chi(-1) = -1) \end{cases}$$

(definition 7.1.6). The *completed L -function* of χ is

$$\Lambda(s, \chi) = \left(\frac{m}{\pi}\right)^{(s+a)/2} \Gamma\left(\frac{s+a}{2}\right) L(s, \chi)$$

with $L(s, \chi)$ the Dirichlet L -function of definition 7.2.1 and Γ the gamma function of definition 6.1.1, defined initially on $\operatorname{Re}(s) > 1$ where the series for $L(s, \chi)$ converges absolutely.

The completed L -function multiplies the Dirichlet series, the product over the finite places, by a single archimedean factor at the real place of $\mathbb{Q}$, the factor $(m/\pi)^{(s+a)/2} \Gamma((s+a)/2)$. For the trivial character $\chi = \chi_0$ modulo 1, which is even, the parity is $a = 0$, the conductor is $m = 1$, and the definition reduces to $\pi^{-s/2} \Gamma(s/2) \zeta(s) = \Lambda(s)$, recovering the completed zeta function of definition 6.3.1; the present definition is its character-twisted generalization, with the conductor m entering through the power $(m/\pi)^{(s+a)/2}$ in place of $\pi^{-s/2}$. The shape of the gamma factor, with

argument $(s + a)/2$, is the archimedean Euler factor of χ : the even factor $\Gamma((s)/2)$ is the gamma factor of the trivial representation of the real place, and the odd factor $\Gamma((s + 1)/2)$ is the gamma factor of the sign representation, the two one-dimensional representations of $\mathbb{R}^\times/\mathbb{R}_{>0}^\times$ distinguished by their behavior at -1 . This parity-indexed choice of archimedean factor is the prototype of the local factors at infinity that recur for the Dedekind zeta functions, taken up in a later chapter of this book, and beyond.

The functional equation now follows, by twisting Riemann's argument by χ . The completed L -function is entire (the pole of ζ was tied to the constant term of the theta series, which the character twist annihilates), and it satisfies a reflection $s \leftrightarrow 1 - s$ that carries the root number.

Theorem 7.3.4: The Functional Equation of $L(s, \chi)$

Let χ be a primitive Dirichlet character modulo m with parity $a \in \{0, 1\}$, and assume $\chi \neq \chi_0$ (equivalently $m > 1$). Then the completed L -function $\Lambda(s, \chi)$ of definition 7.3.3 extends to an entire function of s and satisfies the functional equation

$$\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1 - s, \bar{\chi}) \quad (7.4)$$

where the *root number* is

$$\epsilon(\chi) = \frac{\tau(\chi)}{i^a \sqrt{m}}$$

a complex number of modulus $|\epsilon(\chi)| = 1$.

Proof:

That $|\epsilon(\chi)| = 1$ is immediate from theorem 7.3.2(2): the numerator $\tau(\chi)$ has modulus $\sqrt{m}$, the denominator $i^a \sqrt{m}$ has modulus $\sqrt{m}$, so the quotient has modulus 1. The substance is the entirety and the reflection (7.4). The even case $a = 0$ is proved in full, twisting the theta argument of theorem 6.3.3; the odd case $a = 1$ runs along the identical path with a weighted theta series, carried out in full next.

Step 1: The twisted theta function (even case). Assume χ even, $a = 0$. Define the *twisted theta function*

$$\theta_\chi(t) = \sum_{n \in \mathbb{Z}} \chi(n) e^{-\pi n^2 t/m}, \quad t > 0$$

the character-weighted analogue of $\theta(t) = \sum_n e^{-\pi n^2 t}$ (definition 6.2.3), with the conductor m scaling the exponent. The series converges absolutely and locally uniformly on $t > 0$ by the same geometric domination as for θ , since $|\chi(n)| \leq 1$. The $n = 0$ term vanishes because $\chi(0) = 0$, and because χ is even, $\chi(-n) = \chi(n)$, so pairing $\pm n$ gives

$$\theta_\chi(t) = 2 \sum_{n=1}^{\infty} \chi(n) e^{-\pi n^2 t/m} \quad (7.5)$$

The functional equation of this twisted theta is the analogue of theorem 6.2.4, now exchanging χ with its conjugate:

$$\theta_\chi\left(\frac{1}{t}\right) = \frac{\tau(\chi)}{\sqrt{m}} \sqrt{t} \theta_{\bar{\chi}}(t) \quad (7.6)$$

established in Step 2.

Step 2: Poisson summation gives (7.6). Separability (theorem 7.3.2(1)) for the primitive character $\bar{\chi}$ reads $\sum_{a \bmod m} \bar{\chi}(a) e^{2\pi i n a/m} = \chi(n) \tau(\bar{\chi})$ for all n (using $\bar{\bar{\chi}} = \chi$), and since $\tau(\bar{\chi}) \neq 0$ by theorem 7.3.2(2), this inverts to the finite Fourier expansion

$$\chi(n) = \frac{1}{\tau(\bar{\chi})} \sum_{a \bmod m} \bar{\chi}(a) e^{2\pi i n a/m} \quad (7.7)$$

valid for every integer n . Insert (7.7) into the definition of θ_χ and exchange the finite sum over a with the sum over n :

$$\theta_\chi(t) = \frac{1}{\tau(\bar{\chi})} \sum_{a \bmod m} \bar{\chi}(a) \sum_{n \in \mathbb{Z}} e^{2\pi i n a/m} e^{-\pi n^2 t/m}$$

The inner sum is $\sum_{n \in \mathbb{Z}} F_a(n)$ for the function $F_a(x) = e^{2\pi i a x/m} e^{-\pi x^2 t/m}$ on $\mathbb{R}$, a Gaussian modulated by the additive character $e^{2\pi i a x/m}$. Its Fourier transform, in the normalization of theorem 6.2.1, is computed from lemma 6.2.2: the Gaussian $g(x) = e^{-\pi x^2(t/m)}$ has $\widehat{g}(\xi) = (t/m)^{-1/2} e^{-\pi \xi^2(m/t)} = (m/t)^{1/2} e^{-\pi \xi^2 m/t}$, and modulation by $e^{2\pi i a x/m}$ shifts the transform by a/m , so

$$\widehat{F_a}(\xi) = \left(\frac{m}{t}\right)^{1/2} e^{-\pi(\xi - a/m)^2 m/t}$$

Both F_a and $\widehat{F_a}$ decay faster than any power, so theorem 6.2.1 applies and

$$\sum_{n \in \mathbb{Z}} F_a(n) = \sum_{k \in \mathbb{Z}} \widehat{F_a}(k) = \left(\frac{m}{t}\right)^{1/2} \sum_{k \in \mathbb{Z}} e^{-\pi(km - a)^2/(mt)}$$

the exponent rewritten via $(k - a/m)^2(m/t) = (km - a)^2/(mt)$. Substituting back,

$$\theta_\chi(t) = \frac{1}{\tau(\bar{\chi})} \left(\frac{m}{t}\right)^{1/2} \sum_{a \bmod m} \bar{\chi}(a) \sum_{k \in \mathbb{Z}} e^{-\pi(km - a)^2/(mt)}$$

Set $b = km - a$. As a ranges over residues modulo m and k over $\mathbb{Z}$, the integer b ranges over all of $\mathbb{Z}$, with $a \equiv -b \pmod{m}$, so $\bar{\chi}(a) = \bar{\chi}(-b) = \bar{\chi}(-1)\bar{\chi}(b) = \bar{\chi}(b)$ because $\bar{\chi}$ is even (the parity of $\bar{\chi}$ equals that of χ). The exponent is $(km - a)^2/(mt) = b^2/(mt) = b^2(1/t)/m$, so the double sum collapses to a single sum over b :

$$\sum_{a \bmod m} \bar{\chi}(a) \sum_{k \in \mathbb{Z}} e^{-\pi(km - a)^2/(mt)} = \sum_{b \in \mathbb{Z}} \bar{\chi}(b) e^{-\pi b^2(1/t)/m} = \theta_{\bar{\chi}}\left(\frac{1}{t}\right)$$

Therefore $\theta_\chi(t) = \tau(\bar{\chi})^{-1} (m/t)^{1/2} \theta_{\bar{\chi}}(1/t)$. Replacing t by $1/t$ gives

$$\theta_\chi\left(\frac{1}{t}\right) = \tau(\bar{\chi})^{-1} \sqrt{m} \sqrt{t} \theta_{\bar{\chi}}(t) = \frac{\sqrt{m}}{\tau(\bar{\chi})} \sqrt{t} \theta_{\bar{\chi}}(t)$$

The prefactor simplifies by the product relation $\tau(\chi)\tau(\bar{\chi}) = \chi(-1)m = m$ of theorem 7.3.2(3), valid here because χ is even: solving for $\tau(\bar{\chi}) = m/\tau(\chi)$ gives $\sqrt{m}/\tau(\bar{\chi}) = \sqrt{m}\tau(\chi)/m = \tau(\chi)/\sqrt{m}$. This is (7.6).

Step 3: The Mellin transform (even case). For $\operatorname{Re}(s) > 1$, form the Mellin transform of $\frac{1}{2}\theta_\chi$ against $t^{s/2}$. Using (7.5) and the per-term integral computed by the substitution $u = \pi n^2 t/m$ (so $\int_0^\infty e^{-\pi n^2 t/m} t^{s/2-1} dt = (m/\pi)^{s/2} n^{-s} \Gamma(s/2)$, exactly as in (6.15) with π replaced by π/m),

$$\frac{1}{2} \int_0^\infty \theta_\chi(t) t^{s/2-1} dt = \sum_{n=1}^\infty \chi(n) \left(\frac{m}{\pi}\right)^{s/2} n^{-s} \Gamma\left(\frac{s}{2}\right) = \left(\frac{m}{\pi}\right)^{s/2} \Gamma\left(\frac{s}{2}\right) L(s, \chi) = \Lambda(s, \chi) \quad (7.8)$$

the interchange of sum and integral justified by Tonelli's theorem ([8]) applied to the absolute values, the iterated integral $\sum_n \int_0^\infty e^{-\pi n^2 t/m} t^{\operatorname{Re}(s)/2-1} dt = (m/\pi)^{\operatorname{Re}(s)/2} \Gamma(\operatorname{Re}(s)/2) \sum_n n^{-\operatorname{Re}(s)}$ being finite for $\operatorname{Re}(s) > 1$. This is the analogue of the Mellin representation (6.16) of Λ , with the parity making θ_χ play the role of $\theta - 1$.

Step 4: Symmetry and continuation (even case). Split (7.8) at $t = 1$:

$$2 \Lambda(s, \chi) = \int_0^1 \theta_\chi(t) t^{s/2-1} dt + \int_1^\infty \theta_\chi(t) t^{s/2-1} dt$$

In the first integral substitute $t \mapsto 1/t$, then apply (7.6):

$$\int_0^1 \theta_\chi(t) t^{s/2-1} dt = \int_1^\infty \theta_\chi\left(\frac{1}{t}\right) t^{-s/2-1} dt = \frac{\tau(\chi)}{\sqrt{m}} \int_1^\infty \theta_{\bar{\chi}}(t) t^{(1-s)/2-1} dt$$

the exponent following from $\frac{1}{2} - \frac{s}{2} - 1 = \frac{1-s}{2} - 1$. Therefore

$$2 \Lambda(s, \chi) = \int_1^\infty \theta_\chi(t) t^{s/2-1} dt + \frac{\tau(\chi)}{\sqrt{m}} \int_1^\infty \theta_{\bar{\chi}}(t) t^{(1-s)/2-1} dt \quad (7.9)$$

Both integrals are over $[1, \infty)$, where each of θ_χ and $\theta_{\bar{\chi}}$ decays faster than any power of t (the bound (7.5) gives $|\theta_\chi(t)| \leq 2 \sum_{n \geq 1} e^{-\pi n^2 t/m}$, geometric in t), so both integrals converge for every $s \in \mathbb{C}$ and define entire functions, by differentiation under the integral sign exactly as in Step 2 of theorem 6.3.3. The right-hand side of (7.9) is thus entire, and it agrees with $2\Lambda(s, \chi)$ on $\operatorname{Re}(s) > 1$; by the identity theorem ([16]) it is the entire continuation of $\Lambda(s, \chi)$. There are no polar terms because the twist by the nonprincipal character χ removed the constant term of the theta series that produced the poles $1/(s-1) - 1/s$ for ζ ; this is the analytic form of the holomorphy of $L(s, \chi)$ at $s = 1$ (theorem 7.2.4).

The same computation applied to $\bar{\chi}$ with s replaced by $1-s$ gives

$$2 \Lambda(1-s, \bar{\chi}) = \int_1^\infty \theta_{\bar{\chi}}(t) t^{(1-s)/2-1} dt + \frac{\tau(\bar{\chi})}{\sqrt{m}} \int_1^\infty \theta_\chi(t) t^{s/2-1} dt$$

Multiply this identity by $\tau(\chi)/\sqrt{m}$. The cross term acquires the factor

$$\frac{\tau(\chi)}{\sqrt{m}} \cdot \frac{\tau(\bar{\chi})}{\sqrt{m}} = \frac{\tau(\chi)\tau(\bar{\chi})}{m} = \frac{\chi(-1)m}{m} = \chi(-1) = 1$$

by theorem 7.3.2(3) and $\chi(-1) = 1$ for even χ . Hence

$$\frac{\tau(\chi)}{\sqrt{m}} \cdot 2 \Lambda(1-s, \bar{\chi}) = \frac{\tau(\chi)}{\sqrt{m}} \int_1^\infty \theta_{\bar{\chi}}(t) t^{(1-s)/2-1} dt + \int_1^\infty \theta_\chi(t) t^{s/2-1} dt = 2 \Lambda(s, \chi)$$

the right-hand side being exactly (7.9). With $a = 0$ the root number is $\epsilon(\chi) = \tau(\chi)/(i^0\sqrt{m}) = \tau(\chi)/\sqrt{m}$, so this reads $\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1-s, \bar{\chi})$, which is (7.4) in the even case.

Step 5: The odd case. For χ odd, $a = 1$, the symmetric theta series θ_χ vanishes identically (pairing $\pm n$ cancels, since $\chi(-n) = -\chi(n)$), and the correct object is the weighted theta function

$$\theta_\chi^{(1)}(t) = \sum_{n \in \mathbb{Z}} n \chi(n) e^{-\pi n^2 t/m}$$

whose factor n restores a nonzero sum for an odd character: the summand $n\chi(n)$ is even in n , since $(-n)\chi(-n) = (-n)(-\chi(n)) = n\chi(n)$, so $\theta_\chi^{(1)}(t) = 2 \sum_{n \geq 1} n \chi(n) e^{-\pi n^2 t/m}$ (the $n = 0$ term vanishing), absolutely and locally uniformly convergent on $t > 0$. The Fourier transform of the weighted Gaussian is obtained from lemma 6.2.2 by differentiation: for $g(x) = e^{-\pi x^2(t/m)}$, with $\widehat{g}(\xi) = (m/t)^{1/2} e^{-\pi \xi^2 m/t}$ as in Step 2, differentiating $\widehat{g}(\xi) = \int_{\mathbb{R}} g(x) e^{-2\pi i x \xi} dx$ under the integral gives $\widehat{g}'(\xi) = -2\pi i x \widehat{g}(\xi)$, hence

$$\widehat{xg}(\xi) = \frac{1}{-2\pi i} \widehat{g}'(\xi) = -i \left(\frac{m}{t}\right)^{3/2} \xi e^{-\pi \xi^2 m/t}$$

using $1/i = -i$ and $\widehat{g}'(\xi) = -2\pi(m/t)^{3/2} \xi e^{-\pi \xi^2 m/t}$. Modulating by $e^{2\pi i a x/m}$ shifts the transform by a/m , so $F_a^{(1)}(x) = x e^{2\pi i a x/m} e^{-\pi x^2 t/m}$ has $\widehat{F_a^{(1)}}(\xi) = -i(m/t)^{3/2}(\xi - a/m) e^{-\pi(\xi - a/m)^2 m/t}$. Inserting the finite Fourier expansion (7.7) of $\chi(n)$ into $\theta_\chi^{(1)}$, exchanging the finite a -sum with the n -sum, and applying Poisson summation (theorem 6.2.1) to $F_a^{(1)}$ gives

$$\theta_\chi^{(1)}(t) = \frac{1}{\tau(\bar{\chi})} \sum_{a \bmod m} \bar{\chi}(a) \sum_{k \in \mathbb{Z}} \widehat{F_a^{(1)}}(k)$$

Setting $b = km - a$, so that $k - a/m = b/m$ and $(k - a/m)^2 m/t = b^2(1/t)/m$, and using $\bar{\chi}(a) = \bar{\chi}(-b) = \bar{\chi}(-1)\bar{\chi}(b) = -\bar{\chi}(b)$ since $\bar{\chi}$ is odd, the double sum collapses to $-\frac{1}{m} \theta_{\bar{\chi}}^{(1)}(1/t)$, whence

$$\theta_\chi^{(1)}(t) = \frac{1}{\tau(\bar{\chi})} (-i) \left(\frac{m}{t}\right)^{3/2} \left(-\frac{1}{m}\right) \theta_{\bar{\chi}}^{(1)}\left(\frac{1}{t}\right) = \frac{i\sqrt{m}}{\tau(\bar{\chi})} t^{-3/2} \theta_{\bar{\chi}}^{(1)}\left(\frac{1}{t}\right)$$

Replacing t by $1/t$ and simplifying the prefactor by the product relation $\tau(\chi)\tau(\bar{\chi}) = \chi(-1)m = -m$ of theorem 7.3.2(3) (so $i\sqrt{m}/\tau(\bar{\chi}) = i\sqrt{m}\tau(\chi)/(-m) = -i\tau(\chi)/\sqrt{m} = \tau(\chi)/(i\sqrt{m})$, using $-i = 1/i$) produces the transformation law

$$\theta_\chi^{(1)}\left(\frac{1}{t}\right) = \frac{\tau(\chi)}{i\sqrt{m}} t^{3/2} \theta_{\bar{\chi}}^{(1)}(t)$$

the factor $i^{-1} = i^{-a}$ being the source of the i^a in the denominator of $\epsilon(\chi)$, and the power $t^{3/2}$ reflecting the shifted gamma argument $(s+1)/2$.

Step 6: Mellin transform and reflection (odd case). For $\text{Re}(s) > 1$, the Mellin transform of $\frac{1}{2}\theta_\chi^{(1)}$ against $t^{(s+1)/2}$ reproduces the completed L-function. By the per-term integral $\int_0^\infty e^{-\pi n^2 t/m} t^{(s+1)/2-1} dt = (m/\pi)^{(s+1)/2} n^{-(s+1)} \Gamma((s+1)/2)$ and $n \cdot n^{-(s+1)} = n^{-s}$,

$$\frac{1}{2} \int_0^\infty \theta_\chi^{(1)}(t) t^{(s+1)/2-1} dt = \left(\frac{m}{\pi}\right)^{(s+1)/2} \Gamma\left(\frac{s+1}{2}\right) L(s, \chi) = \Lambda(s, \chi)$$

the interchange justified by Tonelli ([8]) as in Step 3. Splitting the integral at $t = 1$, substituting $t \mapsto 1/t$ in the piece over $(0, 1)$, and applying the transformation law just established (exactly as in Step 4) gives

$$2\Lambda(s, \chi) = \int_1^\infty \theta_\chi^{(1)}(t) t^{(s+1)/2-1} dt + \frac{\tau(\chi)}{i\sqrt{m}} \int_1^\infty \theta_{\bar{\chi}}^{(1)}(t) t^{((1-s)+1)/2-1} dt$$

both integrals over $[1, \infty)$ entire by the rapid decay of $\theta_\chi^{(1)}$ and $\theta_{\bar{\chi}}^{(1)}$, so $\Lambda(s, \chi)$ continues to an entire function (no polar terms: the character twist annihilates the constant term of the theta series that produced the poles of $\Lambda(s)$, consistent with the holomorphy of $L(s, \chi)$ in theorem 7.2.4). The same identity for $\bar{\chi}$ with s replaced by $1 - s$, multiplied by $\tau(\chi)/(i\sqrt{m})$, has cross-coefficient

$$\frac{\tau(\chi)}{i\sqrt{m}} \cdot \frac{\tau(\bar{\chi})}{i\sqrt{m}} = \frac{\tau(\chi)\tau(\bar{\chi})}{i^2 m} = \frac{-m}{-m} = 1$$

by theorem 7.3.2(3) with $\chi(-1) = -1$ and $i^2 = -1$; it therefore reproduces $2\Lambda(s, \chi)$, giving $\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1 - s, \bar{\chi})$ with $\epsilon(\chi) = \tau(\chi)/(i\sqrt{m})$. This is (7.4) in the odd case, completing the proof.

The completed L -function is the product of the finite-place Euler factors, encoded in $L(s, \chi) = \prod_p (1 - \chi(p)p^{-s})^{-1}$ (theorem 7.2.2), with the single archimedean gamma factor chosen by the parity of χ , and it is this completed product, never the bare Dirichlet series, that satisfies the clean reflection $s \leftrightarrow 1 - s$. Every structural feature of the functional equation of ζ (theorem 6.3.3) is reproduced: the gamma factor at the real place, the symmetry about $\text{Re}(s) = 1/2$, the entirety once the polar contribution is removed. The one feature with no counterpart in the zeta theory is the root number $\epsilon(\chi)$. For ζ the reflection is an exact equality, $\Lambda(s) = \Lambda(1 - s)$, because ζ is its own “conjugate” and the trivial Gauss sum contributes the constant 1; for a character the reflection sends χ to $\bar{\chi}$ and inserts the constant $\epsilon(\chi)$, a complex number on the unit circle whose value records the fine arithmetic of χ through the Gauss sum $\tau(\chi)$. This constant of modulus 1 is the prototype of the *local epsilon-factors* that organize the functional equations of all higher L -functions: it is the global epsilon-factor of χ , and in the local theory it factors into a product of local constants, one at each place, the archimedean place contributing the power i^{-a} and the finite places contributing the Gauss sum. The completed Dedekind zeta functions and Hecke L -functions, taken up in a later chapter of this book, carry root numbers of exactly this kind, and the Gauss sum computed here is the first instance of the construction that produces them.

The smallest odd primitive character makes the Gauss sum, the root number, and the resulting symmetry fully explicit, and identifies the completed L -function with a symmetric object.

Example 7.3.5: The Gauss Sum and Functional Equation Modulo 4

Let χ be the nonprincipal character modulo 4 of example 7.1.7, with $\chi(1) = 1$, $\chi(3) = -1$, and $\chi(n) = 0$ for even n . It is odd, so its parity is $a = 1$, and it is real, so $\bar{\chi} = \chi$ and χ is its own conjugate. Its L -function is the Dirichlet beta function $L(s, \chi) = \beta(s)$ of example 7.2.5.

The Gauss sum. Only the residues $a = 1, 3$ contribute, since $\chi(2) = \chi(4) = 0$:

$$\tau(\chi) = \chi(1)e^{2\pi i \cdot 1/4} + \chi(3)e^{2\pi i \cdot 3/4} = e^{i\pi/2} - e^{i3\pi/2} = i - (-i) = 2i$$

so $|\tau(\chi)| = 2 = \sqrt{4} = \sqrt{m}$, confirming theorem 7.3.2(2) in this case. The product relation is also visible directly: $\tau(\bar{\chi}) = \tau(\chi) = 2i$ since χ is real, and $\tau(\chi)\tau(\bar{\chi}) = (2i)^2 = -4 = \chi(-1) \cdot 4$, consistent with $\chi(-1) = -1$ from theorem 7.3.2(3).

The root number. With $a = 1$,

$$\epsilon(\chi) = \frac{\tau(\chi)}{i^a \sqrt{m}} = \frac{2i}{i^1 \cdot 2} = \frac{2i}{2i} = 1$$

The root number is 1, so the functional equation (7.4) for this character reads

$$\Lambda(s, \chi) = \Lambda(1 - s, \bar{\chi}) = \Lambda(1 - s, \chi)$$

a genuine symmetry, using $\bar{\chi} = \chi$. Written out from definition 7.3.3 with $m = 4$ and $a = 1$,

$$\Lambda(s, \chi) = \left(\frac{4}{\pi}\right)^{(s+1)/2} \Gamma\left(\frac{s+1}{2}\right) \beta(s)$$

and the symmetry $\Lambda(s, \chi) = \Lambda(1 - s, \chi)$ relates $\beta(s)$ to $\beta(1 - s)$ through the gamma factors, with axis $\operatorname{Re}(s) = 1/2$. This is the functional equation of the Dirichlet beta function, which is the nontrivial factor of the Dedekind zeta function of the Gaussian field $\mathbb{Q}(i)$, in the factorization $\zeta_{\mathbb{Q}(i)}(s) = \zeta(s) \beta(s)$ cutting out $\mathbb{Q}(i)$ by the quadratic character χ . The vanishing of the root number's phase (its value is exactly +1) is the statement that the completed beta function is symmetric across the critical line with no twist, and it holds because χ is a real (quadratic) character: for such a character the Gauss sum is $\tau(\chi) = i^a \sqrt{m}$ (here $\tau(\chi) = 2i = i\sqrt{4}$ with $a = 1$), which forces $\epsilon(\chi) = \tau(\chi)/(i^a \sqrt{m}) = +1$ directly, independent of any arithmetic of the field. The contrast with a character whose root number is not 1 is what the general theorem records: for a non-real character χ the reflection exchanges χ and $\bar{\chi}$ and inserts a root number $\epsilon(\chi) \neq 1$ on the unit circle, and only the modulus $|\epsilon(\chi)| = 1$ is forced in general, the phase carrying the arithmetic of the Gauss sum.

7.4 Non-Vanishing of $L(1, \chi)$

The decomposition of section 7.2 reduced the count of primes in a residue class to a principal term carrying a pole and a sum of corrections, one for each nonprincipal character, the χ -th correction governed by $\log L(s, \chi)$ as $s \rightarrow 1^+$. That correction stays bounded, and so fails to swallow the main term, exactly when $L(1, \chi) \neq 0$, so that $\log L(s, \chi)$ does not run to $-\infty$. The single inequality $L(1, \chi) \neq 0$ for every nonprincipal χ is therefore the linchpin of Dirichlet's theorem. It is the L -function analogue of the non-vanishing of ζ on the line $\operatorname{Re}(s) = 1$ (theorem 4.4.2), the input that converted the pole of ζ into the Prime Number Theorem; here the same kind of boundary non-vanishing converts the pole of $L(s, \chi_0)$ into a count of primes in each progression.

The argument splits according to the nature of χ . A complex character ($\chi \neq \bar{\chi}$) is disposed of by a counting argument on the full product $\prod_{\chi} L(s, \chi)$: a zero at $s = 1$ would be accompanied by a second zero from the conjugate character, and two zeros cannot be absorbed by the single pole that the product carries. The real (quadratic) characters, those with $\chi = \bar{\chi}$, are the hard case, since such a character is its own conjugate and a zero at $s = 1$ comes with no free partner. The resolution is an argument of Landau on Dirichlet series with non-negative coefficients, applied to $\zeta(s)L(s, \chi)$: a zero at $s = 1$ would force this product to be holomorphic on $\operatorname{Re}(s) > 0$, hence (by Landau) representable by a convergent Dirichlet series there, but its coefficients are large enough at the perfect squares to make the series diverge well to the right of $\operatorname{Re}(s) = 0$. The whole development rests on one product and its coefficients, constructed first.

Proposition 7.4.1: Behaviour of the Product $\prod_{\chi} L(s, \chi)$ at $s = 1$

Fix $m \in \mathbb{Z}_{>0}$ and set

$$\zeta_m(s) = \prod_{\chi \bmod m} L(s, \chi)$$

the product running over all $\varphi(m)$ Dirichlet characters modulo m . For $\operatorname{Re}(s) > 1$:

1. ζ_m has the Euler product

$$\zeta_m(s) = \prod_{p \nmid m} (1 - p^{-f_p s})^{-\varphi(m)/f_p}$$

where $f_p = \operatorname{ord}_m(p)$ is the multiplicative order of p in $(\mathbb{Z}/m\mathbb{Z})^\times$;

2. ζ_m is a Dirichlet series $\sum_{n \geq 1} c_n n^{-s}$ with non-negative real coefficients $c_n \geq 0$ and constant term $c_1 = 1$; and
3. the real function $\sigma \mapsto \zeta_m(\sigma)$ on $(1, \infty)$ satisfies $\zeta_m(\sigma) \rightarrow +\infty$ as $\sigma \rightarrow (1/\varphi(m))^+$, so the Dirichlet series of (2) has abscissa of convergence at least $1/\varphi(m)$.

Proof:

Step 1: The logarithm of the product. Each factor $L(s, \chi)$ is the Dirichlet series of the totally multiplicative function χ (definition 7.1.1), with absolutely convergent Euler product on $\operatorname{Re}(s) > 1$ (theorem 7.2.2) and $|\chi(p)p^{-s}| < 1$ there. Proposition 2.3.5 applies to each χ and gives, for $\operatorname{Re}(s) > 1$, the absolutely convergent expansion

$$\log L(s, \chi) = \sum_p \sum_{k=1}^{\infty} \frac{\chi(p)^k}{k p^{ks}} = \sum_{p \nmid m} \sum_{k=1}^{\infty} \frac{\chi(p^k)}{k p^{ks}}$$

the second form using $\chi(p)^k = \chi(p^k)$ by complete multiplicativity and discarding the primes $p \mid m$, where $\chi(p) = 0$. Summing over the $\varphi(m)$ characters and interchanging the order of summation, which is permitted because the triple sum converges absolutely (each inner double sum does, and there are finitely many characters),

$$\log \zeta_m(s) = \sum_{\chi \bmod m} \log L(s, \chi) = \sum_{p \nmid m} \sum_{k=1}^{\infty} \frac{1}{k p^{ks}} \sum_{\chi \bmod m} \chi(p^k)$$

where the branch of $\log \zeta_m$ is the sum of the branches of $\log L(s, \chi)$ fixed by proposition 2.3.5.

Step 2: Orthogonality collapses the character sum. For $p \nmid m$ the integer p^k is coprime to m , so column orthogonality (theorem 7.1.4(1)) evaluates the inner sum:

$$\sum_{\chi \bmod m} \chi(p^k) = \begin{cases} \varphi(m) & \text{if } p^k \equiv 1 \pmod{m}, \\ 0 & \text{otherwise} \end{cases}$$

The congruence $p^k \equiv 1 \pmod{m}$ holds precisely when $f_p = \operatorname{ord}_m(p)$ divides k , that is, when

$k = jf_p$ for some $j \geq 1$. Substituting and writing $k = jf_p$,

$$\log \zeta_m(s) = \varphi(m) \sum_{p \nmid m} \sum_{j=1}^{\infty} \frac{1}{j f_p p^{j f_p s}} \quad (7.10)$$

a double series with non-negative real coefficients on the real axis $s = \sigma > 1$. Exponentiating term by term, the contribution of a single prime p is

$$\exp\left(\frac{\varphi(m)}{f_p} \sum_{j=1}^{\infty} \frac{(p^{-f_p s})^j}{j}\right) = \exp\left(-\frac{\varphi(m)}{f_p} \log(1 - p^{-f_p s})\right) = (1 - p^{-f_p s})^{-\varphi(m)/f_p}$$

using $\sum_{j \geq 1} z^j/j = -\log(1 - z)$ with $z = p^{-f_p s}$, valid since $|p^{-f_p s}| < 1$ for $\operatorname{Re}(s) > 0$. Taking the product over $p \nmid m$ gives the Euler product of (1).

Step 3: Non-negative coefficients and the constant term. The exponent $\varphi(m)/f_p$ is a positive integer, since $f_p = \operatorname{ord}_m(p)$ divides $|\mathbb{Z}/m\mathbb{Z}^\times| = \varphi(m)$ by Lagrange's theorem. Each factor of the Euler product expands by the generalized binomial series

$$(1 - p^{-f_p s})^{-\varphi(m)/f_p} = \sum_{j=0}^{\infty} \binom{\varphi(m)/f_p + j - 1}{j} p^{-j f_p s}$$

whose coefficients $\binom{\varphi(m)/f_p + j - 1}{j}$ are non-negative integers and whose constant term ($j = 0$) is 1. A product of Dirichlet series each with non-negative coefficients and constant term 1 is again a Dirichlet series with non-negative coefficients and constant term 1: the coefficient c_n of n^{-s} in the product is a finite sum of products of the individual non-negative coefficients, and c_1 is the product of the constant terms. This is the absolutely convergent rearrangement of the product into a single Dirichlet series, valid on $\operatorname{Re}(s) > 1$ where every factor converges absolutely. This gives (2).

Step 4: Divergence at $s = 1/\varphi(m)$. Work on the real axis, $s = \sigma > 1/\varphi(m)$, where every term of (7.10) is non-negative. Discarding all terms with $j \geq 2$ leaves the lower bound

$$\log \zeta_m(\sigma) \geq \varphi(m) \sum_{p \nmid m} \frac{1}{f_p p^{f_p \sigma}}$$

For each $p \nmid m$ the order f_p divides $\varphi(m)$, so $f_p \leq \varphi(m)$, and at σ approaching $1/\varphi(m)$ from above the exponent satisfies $f_p \sigma \leq \varphi(m) \sigma$. Hence $p^{f_p \sigma} \leq p^{\varphi(m) \sigma}$, and since also $f_p \leq \varphi(m)$,

$$\log \zeta_m(\sigma) \geq \varphi(m) \sum_{p \nmid m} \frac{1}{\varphi(m) p^{\varphi(m) \sigma}} = \sum_{p \nmid m} \frac{1}{p^{\varphi(m) \sigma}}$$

As $\sigma \rightarrow (1/\varphi(m))^+$ the exponent $\varphi(m) \sigma \rightarrow 1^+$, and $\sum_p p^{-\varphi(m) \sigma}$ diverges to $+\infty$ by the divergence of the prime sum (proposition 4.3.5), the finitely many primes $p \mid m$ omitted from the sum affecting it by a bounded amount. Therefore $\log \zeta_m(\sigma) \rightarrow +\infty$, so $\zeta_m(\sigma) \rightarrow +\infty$. Because the coefficients c_n are non-negative, the real function $\zeta_m(\sigma)$ on the interval of convergence is the increasing limit of the partial sums, so a finite value of the series at a real point σ_0 forces convergence for all $\sigma \geq \sigma_0$; the blow-up at $1/\varphi(m)$ thus shows the abscissa of convergence is at least $1/\varphi(m)$, which is (3) and completes the proof.

The product $\zeta_m(s)$ assembled here is a prototype of the Dedekind zeta function of the cyclotomic field $\mathbb{Q}(\zeta_m)$, which factors as exactly this product of Dirichlet L -functions over the characters modulo m ; that factorization, and the parallel one for quadratic fields, is taken up in chapter 8. The two properties extracted are the only ones the non-vanishing proof uses. First, ζ_m is a Dirichlet series with non-negative coefficients and constant term 1, in particular real and at least 1 at every real σ in its range. Second, that series cannot be summed too far to the left: it already diverges at $s = 1/\varphi(m)$, a point strictly to the right of $\operatorname{Re}(s) = 0$. Non-negativity of coefficients is the hypothesis of Landau's theorem, and the divergence at a positive real point is the contradiction that theorem will be played against. The mechanism is identical to the one behind the non-vanishing of ζ on $\operatorname{Re}(s) = 1$: there the inequality $3 + 4 \cos \theta + \cos 2\theta \geq 0$ (lemma 4.4.1) forced a product of zeta values to be large; here the non-negativity of the coefficients of ζ_m plays the same positivity role, now packaged as a statement about an entire Dirichlet series rather than a single trigonometric inequality.

With the product in hand, the non-vanishing itself follows. The proof treats the complex and real characters separately, the real case being the one that requires the full strength of the positivity.

Theorem 7.4.2: Non-Vanishing at $s = 1$

Let χ be a nonprincipal Dirichlet character modulo m . Then

$$L(1, \chi) \neq 0$$

Proof:

By theorem 7.2.4 every nonprincipal $L(s, \chi)$ is holomorphic on $\operatorname{Re}(s) > 0$, so $L(1, \chi)$ is a well-defined finite value and the statement $L(1, \chi) \neq 0$ is meaningful. Two facts about conjugation are used throughout. First, $L(s, \bar{\chi}) = \overline{L(\bar{s}, \chi)}$ for $\operatorname{Re}(s) > 0$, since term by term $\overline{\chi(n)/n^s} = \bar{\chi}(n)/n^{\bar{s}}$ and the series converges (theorem 7.2.4); evaluating at the real point $s = 1$ gives $L(1, \bar{\chi}) = \overline{L(1, \chi)}$, so $L(1, \chi) = 0$ if and only if $L(1, \bar{\chi}) = 0$. Second, $\bar{\chi}$ is again a nonprincipal character modulo m (theorem 7.1.2(2)), so the conjugate factor is one of the factors of ζ_m .

Step 1: Complex characters. Suppose first $\chi \neq \bar{\chi}$, and suppose toward a contradiction that $L(1, \chi) = 0$. Then $L(1, \bar{\chi}) = \overline{L(1, \chi)} = 0$ as well, and χ and $\bar{\chi}$ are two *distinct* nonprincipal characters modulo m , each contributing a factor vanishing at $s = 1$ to the product $\zeta_m(s) = \prod_{\psi \bmod m} L(s, \psi)$. Account for the order of ζ_m at $s = 1$ factor by factor. The principal factor $L(s, \chi_0)$ has a simple pole at $s = 1$ (proposition 7.2.3), contributing order -1 . The factors $L(s, \chi)$ and $L(s, \bar{\chi})$ each have a zero at $s = 1$, contributing order at least $+1$ apiece. Every remaining factor is holomorphic at $s = 1$ (theorem 7.2.4), contributing order at least 0. The order of the product at $s = 1$ is the sum of the orders of the factors, hence at least

$$(-1) + (+1) + (+1) + 0 = +1 > 0$$

so ζ_m is holomorphic at $s = 1$ with a zero there, $\zeta_m(1) = 0$. The single pole of ζ_m on $\operatorname{Re}(s) > 0$ is the one from χ_0 , now cancelled, so ζ_m is holomorphic on all of $\operatorname{Re}(s) > 0$.

Apply Landau's theorem on Dirichlet series with non-negative coefficients ([1, Ch. 4]): if a Dirichlet series $\sum_n c_n n^{-s}$ has $c_n \geq 0$ for all n and extends to a function holomorphic on the half-plane $\operatorname{Re}(s) > \sigma_0$, then the series itself converges on $\operatorname{Re}(s) > \sigma_0$, so its abscissa of convergence is at most σ_0 . The series $\zeta_m = \sum_n c_n n^{-s}$ has non-negative coefficients (proposition 7.4.1(2)) and, under the present assumption, extends holomorphically to $\operatorname{Re}(s) > 0$; Landau's theorem then

forces its abscissa of convergence to be at most 0. But proposition 7.4.1(3) shows the abscissa is at least $1/\varphi(m) > 0$, a contradiction. Therefore $L(1, \chi) \neq 0$ for $\chi \neq \bar{\chi}$.

Step 2: Real characters. Suppose now $\chi = \bar{\chi}$, so χ takes only the real values $\{0, \pm 1\}$ (each nonzero value is a root of unity equal to its own conjugate), and suppose toward a contradiction that $L(1, \chi) = 0$. The self-conjugate χ supplies no second zero, so the product ζ_m alone no longer yields the contradiction; instead work with the smaller product

$$F(s) = \zeta(s) L(s, \chi)$$

the Riemann zeta function times the single L -function. By theorem 4.2.1 the factor $\zeta(s)$ is holomorphic on $\operatorname{Re}(s) > 0$ except for a simple pole at $s = 1$, and $L(s, \chi)$ is holomorphic on $\operatorname{Re}(s) > 0$ (theorem 7.2.4). At $s = 1$ the simple pole of ζ , of order -1 , meets the assumed zero of $L(s, \chi)$, of order at least $+1$, so the product F is holomorphic at $s = 1$; away from $s = 1$ both factors are holomorphic. Thus F is holomorphic on all of $\operatorname{Re}(s) > 0$.

Step 3: The coefficients of F . For $\operatorname{Re}(s) > 1$ both $\zeta(s) = \sum_n n^{-s}$ and $L(s, \chi) = \sum_n \chi(n) n^{-s}$ converge absolutely, so their product is the Dirichlet series of the convolution: $F(s) = \sum_{n \geq 1} a_n n^{-s}$ with

$$a_n = \sum_{d|n} \chi(d)$$

since ζ has all coefficients 1. The arithmetic function $n \mapsto a_n$ is the Dirichlet convolution of χ with the constant function 1, both multiplicative, hence multiplicative. On a prime power p^k it evaluates to

$$a_{p^k} = \sum_{i=0}^k \chi(p)^i = 1 + \chi(p) + \chi(p)^2 + \cdots + \chi(p)^k$$

a geometric sum whose ratio $\chi(p) \in \{0, +1, -1\}$ is real. The three cases give

$$a_{p^k} = \begin{cases} k+1 & \text{if } \chi(p) = +1, \\ 1 & \text{if } \chi(p) = 0, \\ 1 & \text{if } \chi(p) = -1 \text{ and } k \text{ even,} \\ 0 & \text{if } \chi(p) = -1 \text{ and } k \text{ odd} \end{cases}$$

In every case $a_{p^k} \geq 0$, and whenever k is even $a_{p^k} \geq 1$. By multiplicativity $a_n = \prod_{p^k || n} a_{p^k} \geq 0$ for all n , so F is a Dirichlet series with non-negative coefficients. Moreover, if $n = \ell^2$ is a perfect square then every exponent in the factorization of n is even, so every local factor $a_{p^k} \geq 1$ and hence $a_{\ell^2} \geq 1$.

Step 4: Landau's contradiction. The non-negativity of the coefficients a_n together with the holomorphy of F on $\operatorname{Re}(s) > 0$ feeds into Landau's theorem ([1, Ch. 4]) exactly as in Step 1: the abscissa of convergence of $\sum_n a_n n^{-s}$ would be at most 0, so the series would converge for every real $\sigma > 0$. Test this against the perfect-square lower bound. For real $\sigma > 0$, all terms being non-negative,

$$F(\sigma) = \sum_{n=1}^{\infty} \frac{a_n}{n^{\sigma}} \geq \sum_{\ell=1}^{\infty} \frac{a_{\ell^2}}{(\ell^2)^{\sigma}} \geq \sum_{\ell=1}^{\infty} \frac{1}{\ell^{2\sigma}}$$

using $a_{\ell^2} \geq 1$ from Step 3. The right-hand series is the harmonic-type series $\sum_{\ell} \ell^{-2\sigma}$, which diverges for $2\sigma \leq 1$, that is, for $\sigma \leq 1/2$. Hence the Dirichlet series for F diverges at every real $\sigma \leq 1/2$, so its abscissa of convergence is at least $1/2$. This contradicts the value at most 0 forced by Landau's theorem. The assumption $L(1, \chi) = 0$ is therefore untenable, and $L(1, \chi) \neq 0$ for real nonprincipal χ , completing the proof.

The two cases meet the same obstruction from opposite directions. A complex character carries its own undoing: a zero at $s = 1$ duplicates into a zero of the conjugate factor, and the resulting double zero overwhelms the single pole that the product ζ_m is permitted to have, leaving a Dirichlet series with non-negative coefficients that is holomorphic where it has no right to converge. A real character has no conjugate partner, and the substitute is the product $\zeta(s)L(s, \chi)$, whose coefficients $a_n = \sum_{d|n} \chi(d)$ are visibly non-negative and at least 1 on the perfect squares; a zero at $s = 1$ would again make this holomorphic on $\text{Re}(s) > 0$, contradicting the divergence its square terms guarantee. The class number formula of chapter 8 gives an alternative route to the real case, expressing $L(1, \chi)$ for a quadratic character as a positive multiple of the class number of the associated quadratic field, manifestly nonzero; the Landau argument above is the elementary route, requiring only the positivity of a Dirichlet series, and it is the one on which Dirichlet's theorem is built in section 7.5.

This non-vanishing is exactly the input Dirichlet's theorem needs. In the decomposition of the count of primes $p \equiv a \pmod{m}$, orthogonality (theorem 7.1.4) expresses the count through the weighted sum $\sum_{\chi \pmod{m}} \bar{\chi}(a) \log L(s, \chi)$ as $s \rightarrow 1^+$. The principal term $\bar{\chi}_0(a) \log L(s, \chi_0)$ inherits the pole of $L(s, \chi_0)$ at $s = 1$ (proposition 7.2.3), so $\log L(s, \chi_0) \rightarrow +\infty$ and supplies the divergence that forces infinitely many primes. Every nonprincipal term $\bar{\chi}(a) \log L(s, \chi)$ must stay bounded for the divergence to be attributed to the principal term alone, and boundedness is precisely what $L(1, \chi) \neq 0$ secures: since $L(s, \chi)$ is holomorphic and nonzero at $s = 1$, the logarithm $\log L(s, \chi)$ extends continuously to a finite value $\log L(1, \chi)$ there, with no pole and no descent to $-\infty$. The non-vanishing thus quarantines the singularity in the single principal character, so that the count of primes in the progression $a \pmod{m}$ diverges, and Dirichlet's theorem follows in section 7.5.

The smallest moduli make the non-vanishing concrete, as the values $L(1, \chi)$ were already computed in example 7.2.5; the next example collects them and exhibits the closed forms that display each as a nonzero number.

Example 7.4.3: $L(1, \chi)$ for Small Moduli

The values of $L(1, \chi)$ at the nonprincipal characters of the smallest moduli are evaluated by recognizing each series as a classical sum, and every one is nonzero, in agreement with theorem 7.4.2.

Modulus 4. The unique nonprincipal character is the odd character χ_1 of example 7.1.7, and its L -function at $s = 1$ is the Leibniz series (example 7.2.5):

$$L(1, \chi_1) = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots = \frac{\pi}{4} \neq 0$$

Modulus 3. The unique nonprincipal character χ modulo 3 is odd, and (example 7.2.5)

$$L(1, \chi) = 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{5} + \frac{1}{7} - \frac{1}{8} + \cdots = \frac{\pi}{3\sqrt{3}} \neq 0$$

Both characters here are quadratic and odd, the hard case of theorem 7.4.2, and the table records their values alongside the modulus-5 quadratic character ψ^2 of example 7.1.7, whose value $L(1, \psi^2)$

admits the closed form below:

m	χ	parity	$L(1, \chi)$
3	nonprincipal	odd	$\pi/(3\sqrt{3}) \approx 0.6046$
4	χ_1	odd	$\pi/4 \approx 0.7854$
5	ψ^2	even	$2 \log((1 + \sqrt{5})/2)/\sqrt{5} \approx 0.4304$

Every entry is strictly positive, hence nonzero. The closed forms are not coincidences: each is an instance of the analytic class number formula, which evaluates $L(1, \chi)$ for a primitive quadratic character of conductor d in terms of arithmetic invariants of the quadratic field $\mathbb{Q}(\sqrt{\pm d})$. For an odd character the formula produces a rational multiple of π divided by $\sqrt{d}$ (here $\pi/(3\sqrt{3})$ and $\pi/4$, the imaginary fields $\mathbb{Q}(\sqrt{-3})$ and $\mathbb{Q}(\sqrt{-1})$), while for an even character it produces a logarithm of the fundamental unit (here $\log((1 + \sqrt{5})/2)$ is the logarithm of the fundamental unit of the real field $\mathbb{Q}(\sqrt{5})$). In each case the relevant arithmetic invariant is a positive quantity, which is the structural reason $L(1, \chi) \neq 0$, and chapter 8 derives this formula and with it a second proof that these values cannot vanish.

7.5 Dirichlet's Theorem and Equidistribution

The four preceding sections built every component the proof requires. The characters of section 7.1 resolve a residue class into a sum over frequencies; the L -functions of section 7.2 package the primes of each frequency into an Euler product, with the principal character carrying a pole at $s = 1$ (proposition 7.2.3) and every other character holomorphic there (theorem 7.2.4); and the non-vanishing $L(1, \chi) \neq 0$ of section 7.4 (theorem 7.4.2) guarantees that the holomorphic ones contribute nothing singular. Assembling these three facts proves Dirichlet's theorem: every arithmetic progression $a, a + m, a + 2m, \dots$ with $\gcd(a, m) = 1$ contains infinitely many primes. The mechanism is the one that drove the analytic proof of the infinitude of all primes in chapter 4, run one residue class at a time. There the pole of ζ at $s = 1$ forced $\sum_p p^{-s} \rightarrow +\infty$; here orthogonality isolates the single residue class a , the pole of $L(s, \chi_0)$ supplies the same divergence, and the non-vanishing of the other L -functions confines that divergence to the class a rather than letting it leak away.

The same computation does more than count: it weighs. Carried one step further, it shows that the divergence is shared equally among the $\varphi(m)$ admissible residue classes, each contributing the term $\frac{1}{\varphi(m)} \log \frac{1}{s-1}$ to the prime sum as $s \rightarrow 1^+$. This is the statement that the primes are equidistributed modulo m , with Dirichlet density $1/\varphi(m)$ in each class coprime to m . The section closes with the smallest moduli made numerical, where the equidistribution is visible in prime-counting data and where the residual fluctuation around equality already exhibits the Chebyshev bias.

The proof is a single calculation: expand $\log L(s, \chi)$ as a sum over primes, weight each character by $\chi(a)$, sum over all characters, and let orthogonality collapse the double family of sums onto the one residue class a . Each step is in place from the earlier sections; the work is the bookkeeping that tracks where the pole goes.

Theorem 7.5.1: Dirichlet's Theorem on Primes in Arithmetic Progressions

Let a and m be integers with $m \geq 1$ and $\gcd(a, m) = 1$. Then the arithmetic progression

$$a, a + m, a + 2m, \dots$$

contains infinitely many primes. More precisely, the sum of reciprocals of the primes in the progression diverges:

$$\sum_{\substack{p \\ p \equiv a \pmod{m}}} \frac{1}{p} = +\infty$$

Proof:

Step 1: The prime sum for one character. Fix a Dirichlet character χ modulo m . For real $s > 1$ the Euler product of theorem 7.2.2 converges absolutely and has no vanishing factor, so $L(s, \chi) \neq 0$ and proposition 2.3.5 gives the absolutely convergent logarithmic expansion

$$\log L(s, \chi) = \sum_p \sum_{k=1}^{\infty} \frac{\chi(p)^k}{k p^{ks}} = \sum_p \frac{\chi(p)}{p^s} + \sum_p \sum_{k=2}^{\infty} \frac{\chi(p)^k}{k p^{ks}}$$

the second equality splitting off the $k = 1$ term from the prime-power tail. Write $R(s, \chi)$ for that tail. The tail is bounded uniformly in χ and in $s \geq 1$: using $|\chi(p)| \leq 1$ and the estimate already made for ζ in example 2.3.6,

$$|R(s, \chi)| \leq \sum_p \sum_{k=2}^{\infty} \frac{|\chi(p)|^k}{k p^{ks}} \leq \sum_p \sum_{k=2}^{\infty} \frac{1}{p^k} = \sum_p \frac{1}{p(p-1)} \leq 1$$

the middle inequality dropping $1/k \leq 1$ and using $p^{ks} \geq p^k$ for $s \geq 1$, and the final bound being the telescoping sum of example 2.3.6. Thus $R(s, \chi) = O(1)$ as $s \rightarrow 1^+$, with an implied constant independent of χ , and

$$\log L(s, \chi) = \sum_p \frac{\chi(p)}{p^s} + O(1) \quad (s \rightarrow 1^+) \quad (7.11)$$

The series $\sum_p \chi(p)p^{-s}$ is the part of $\log L(s, \chi)$ that sees the primes linearly; everything else is bounded.

Step 2: Weight by $\overline{\chi(a)}$ and sum over characters. Multiply (7.11) by the constant $\overline{\chi(a)}$, which has modulus $|\overline{\chi(a)}| = 1$ since $\gcd(a, m) = 1$, and sum over all $\varphi(m)$ characters χ modulo m . The bounded terms remain bounded, $\sum_{\chi} \overline{\chi(a)} O(1) = O(1)$, there being finitely many characters, so

$$\sum_{\chi \bmod m} \overline{\chi(a)} \log L(s, \chi) = \sum_{\chi \bmod m} \overline{\chi(a)} \sum_p \frac{\chi(p)}{p^s} + O(1)$$

Interchanging the finite character sum with the prime sum on the right (legitimate for $s > 1$, where each $\sum_p \chi(p)p^{-s}$ converges absolutely),

$$\sum_{\chi \bmod m} \overline{\chi(a)} \log L(s, \chi) = \sum_p \frac{1}{p^s} \left(\sum_{\chi \bmod m} \chi(p) \overline{\chi(a)} \right) + O(1)$$

The inner character sum is exactly the quantity orthogonality evaluates. By the indicator identity (7.1) of theorem 7.1.4, for $\gcd(a, m) = 1$,

$$\sum_{\chi \bmod m} \chi(p) \overline{\chi(a)} = \begin{cases} \varphi(m) & \text{if } p \equiv a \pmod{m}, \\ 0 & \text{otherwise} \end{cases}$$

so the prime sum collapses onto the single residue class a :

$$\sum_{\chi \bmod m} \overline{\chi(a)} \log L(s, \chi) = \varphi(m) \sum_{\substack{p \\ p \equiv a \pmod{m}}} \frac{1}{p^s} + O(1) \quad (s \rightarrow 1^+) \quad (7.12)$$

This identity is the heart of the matter: the left-hand side is a sum of $\varphi(m)$ logarithms of L -functions, and the right-hand side is the prime sum over the progression, multiplied by $\varphi(m)$. The behaviour of the left-hand side as $s \rightarrow 1^+$ now controls the prime sum.

Step 3: The principal character contributes the pole. Separate the principal character χ_0 from the rest of the left-hand side of (7.12). For χ_0 one has $\chi_0(a) = 1$, and by proposition 7.2.3 the function $L(s, \chi_0) = \zeta(s) \prod_{p|m} (1 - p^{-s})$ has a simple pole at $s = 1$. For real $s \rightarrow 1^+$ this gives $L(s, \chi_0) \rightarrow +\infty$, and quantitatively the principal term has the same logarithmic blow-up as $\log \zeta$: combining (7.11) for χ_0 with the prime-sum estimate $\sum_p p^{-s} = \log \frac{1}{s-1} + O(1)$ of proposition 4.3.5 and noting $\chi_0(p) = 1$ for $p \nmid m$ while the finitely many primes $p|m$ alter the sum by $O(1)$,

$$\log L(s, \chi_0) = \sum_{p \nmid m} \frac{1}{p^s} + O(1) = \log \frac{1}{s-1} + O(1) \quad (s \rightarrow 1^+) \quad (7.13)$$

so $\log L(s, \chi_0) \rightarrow +\infty$.

Step 4: The nonprincipal characters stay bounded. For each nonprincipal χ the function $L(s, \chi)$ is holomorphic on $\operatorname{Re}(s) > 0$ (theorem 7.2.4) and, decisively, nonzero at $s = 1$:

$$L(1, \chi) \neq 0$$

by theorem 7.4.2. A function holomorphic and nonzero at a point has a logarithm holomorphic in a neighbourhood of that point, so $\log L(s, \chi)$ extends continuously to the finite value $\log L(1, \chi)$ as $s \rightarrow 1^+$. In particular each nonprincipal term is bounded:

$$\overline{\chi(a)} \log L(s, \chi) = \overline{\chi(a)} \log L(1, \chi) + o(1) = O(1) \quad (s \rightarrow 1^+)$$

This is the step where the non-vanishing earns its place. Were $L(1, \chi) = 0$ for some nonprincipal χ , the term $\log L(s, \chi)$ would tend to $-\infty$ as $s \rightarrow 1^+$, and could cancel the pole from χ_0 , leaving the prime sum bounded and the argument inconclusive. The content of theorem 7.4.2 is precisely that this cancellation cannot happen.

Step 5: Conclusion. Insert Steps 3 and 4 into (7.12). The left-hand side is the principal term plus the bounded nonprincipal terms,

$$\sum_{\chi \bmod m} \overline{\chi(a)} \log L(s, \chi) = \log \frac{1}{s-1} + O(1) \quad (s \rightarrow 1^+)$$

the single unbounded contribution coming from χ_0 . Equating with the right-hand side of (7.12) and dividing by $\varphi(m)$,

$$\sum_{\substack{p \\ p \equiv a \pmod{m}}} \frac{1}{p^s} = \frac{1}{\varphi(m)} \log \frac{1}{s-1} + O(1) \quad (s \rightarrow 1^+) \quad (7.14)$$

As $s \rightarrow 1^+$ the right-hand side tends to $+\infty$, since $\log \frac{1}{s-1} \rightarrow +\infty$ and $\varphi(m) \geq 1$ is a fixed positive integer. Therefore $\sum_{p \equiv a} p^{-s} \rightarrow +\infty$ as $s \rightarrow 1^+$. If only finitely many primes lay in the progression, this sum would be a finite sum of terms each bounded by p^{-1} , hence bounded uniformly in $s > 1$, contradicting the divergence. Hence the progression contains infinitely many primes, and letting $s \rightarrow 1^+$ in the same inequality (or comparing $\sum_{p \equiv a} p^{-s} \leq \sum_{p \equiv a} p^{-1}$ for $s > 1$) shows $\sum_{p \equiv a} 1/p$ diverges, completing the proof.

The proof is the analytic argument for the infinitude of primes from chapter 4, refracted through the characters. Where the pole of ζ at $s = 1$ forced $\sum_p p^{-s} \rightarrow +\infty$ and so produced infinitely many primes (chapter 3, proposition 4.3.5), the pole of $L(s, \chi_0)$ does the same job inside a single residue class. Orthogonality is the instrument that performs the restriction: the weighted sum $\sum_{\chi} \overline{\chi(a)} \chi(p)$ is the indicator of the class a , so summing $\log L(s, \chi)$ against these weights converts a statement about all primes into a statement about the primes $p \equiv a$. The principal character is the only one with a pole, and it alone supplies the divergence. Every nonprincipal character would, if its L -function vanished at $s = 1$, contribute a competing term $\log L(s, \chi) \rightarrow -\infty$ capable of cancelling that pole; the non-vanishing $L(1, \chi) \neq 0$ is exactly the guarantee that no such cancellation occurs. This is the same structural role played by the non-vanishing of ζ on the line $\operatorname{Re}(s) = 1$ in the proof of the Prime Number Theorem (theorem 4.4.2): in both arguments the pole carries the arithmetic, and a boundary non-vanishing result protects the pole from being annihilated, the difference being that for ζ the obstruction sits on the whole line $\operatorname{Re}(s) = 1$, whereas here it is concentrated at the single point $s = 1$ for each twisted function $L(s, \chi)$.

7.5.2 Note: Toward Chebotarev Dirichlet's theorem is the abelian case of a far more general equidistribution law. The residue class of a prime p modulo m is the same data as the Frobenius conjugacy class of p in the Galois group $\operatorname{Gal}(\mathbb{Q}(\zeta_m)/\mathbb{Q}) \cong (\mathbb{Z}/m\mathbb{Z})^\times$ of the cyclotomic field, an abelian group, and the L -functions $L(s, \chi)$ are the factors into which the Dedekind zeta function of $\mathbb{Q}(\zeta_m)$ decomposes. For an arbitrary finite Galois extension $K/\mathbb{Q}$ with group G , possibly nonabelian, the Chebotarev density theorem asserts that the Frobenius classes of the unramified primes are equidistributed among the conjugacy classes of G , a given class C being hit with density $|C|/|G|$. Dirichlet's theorem is the case G abelian, where every conjugacy class is a single element. The proof follows the same template: the Dedekind zeta function of K factors through Artin L -functions attached to the irreducible characters of G , and the non-vanishing of these L -functions at $s = 1$ confines the pole to the trivial character. The factorization of the cyclotomic Dedekind zeta function into Dirichlet L -functions, the first instance of this pattern, is taken up in chapter 8.

Equation (7.14) obtained in the proof is sharper than the bare infinitude. The coefficient of the logarithmic singularity is $1/\varphi(m)$, the *same* for every admissible class a , independent of which class was chosen. This uniformity is the analytic expression of equidistribution: the primes do not just populate every progression, they populate the $\varphi(m)$ progressions in equal proportion. Making this precise requires only a definition and a division.

Definition 7.5.3: Dirichlet Density

Let A be a set of primes. The *Dirichlet density* of A is

$$\delta(A) = \lim_{s \rightarrow 1^+} \frac{\sum_{p \in A} p^{-s}}{\sum_p p^{-s}}$$

when the limit exists, the denominator running over all primes. Since $\sum_p p^{-s} = \log \frac{1}{s-1} + O(1)$ as $s \rightarrow 1^+$ (proposition 4.3.5), the limit may equivalently be written

$$\delta(A) = \lim_{s \rightarrow 1^+} \frac{\sum_{p \in A} p^{-s}}{\log \frac{1}{s-1}}$$

A set with $\delta(A) > 0$, in particular, is infinite.

Dirichlet density measures the relative weight of a set of primes through the analytic weighting $p \mapsto p^{-s}$ as s descends to the pole, rather than through the sharp cutoff $p \leq x$ of the counting function $\pi(x; m, a)$. It is the notion of size native to the L -function method, and it is exactly the quantity the proof of theorem 7.5.1 computed. A related but a priori stronger notion, the *natural density* $\lim_{x \rightarrow \infty} \pi(x; m, a)/\pi(x)$, weights every prime up to x equally; the natural density of each progression also equals $1/\varphi(m)$, but that statement requires the Prime Number Theorem for arithmetic progressions and lies beyond the elementary L -function argument used here. When the natural density of a set of primes exists, it equals the Dirichlet density, so the two agree wherever both are defined.

Theorem 7.5.4: Dirichlet Density and Equidistribution

Let a and m be integers with $m \geq 1$ and $\gcd(a, m) = 1$. The set of primes $p \equiv a \pmod{m}$ has Dirichlet density

$$\delta(\{p \equiv a \pmod{m}\}) = \frac{1}{\varphi(m)}$$

Equivalently, as $s \rightarrow 1^+$,

$$\sum_{\substack{p \\ p \equiv a \pmod{m}}} \frac{1}{p^s} = \frac{1}{\varphi(m)} \log \frac{1}{s-1} + O(1)$$

The $\varphi(m)$ residue classes coprime to m thus carry the primes in equal proportion, each with density $1/\varphi(m)$, and together they account for all of the prime sum: the densities sum to $\varphi(m) \cdot \frac{1}{\varphi(m)} = 1$.

Proof:

The asymptotic

$$\sum_{\substack{p \\ p \equiv a \pmod{m}}} \frac{1}{p^s} = \frac{1}{\varphi(m)} \log \frac{1}{s-1} + O(1) \quad (s \rightarrow 1^+)$$

is (7.14), established in the course of proving theorem 7.5.1: the only character contributing a singularity to the weighted sum $\sum_{\chi} \chi(a) \log L(s, \chi)$ is the principal one, whose contribution is $\log \frac{1}{s-1} + O(1)$ by (7.13), and dividing by $\varphi(m)$ distributes this singularity with coefficient $1/\varphi(m)$. Dividing the displayed asymptotic by $\sum_p p^{-s} = \log \frac{1}{s-1} + O(1)$ (proposition 4.3.5),

$$\frac{\sum_{p \equiv a} p^{-s}}{\sum_p p^{-s}} = \frac{\frac{1}{\varphi(m)} \log \frac{1}{s-1} + O(1)}{\log \frac{1}{s-1} + O(1)} = \frac{\frac{1}{\varphi(m)} + O\left(1/\log \frac{1}{s-1}\right)}{1 + O\left(1/\log \frac{1}{s-1}\right)}$$

the second equality dividing numerator and denominator by $\log \frac{1}{s-1}$, which tends to $+\infty$ as $s \rightarrow 1^+$ so that both $O(1)$ terms become $O(1/\log \frac{1}{s-1}) \rightarrow 0$. Letting $s \rightarrow 1^+$, the right-hand side tends to $\frac{1}{\varphi(m)}/1 = 1/\varphi(m)$. By definition 7.5.3 the Dirichlet density of the progression is therefore $1/\varphi(m)$.

For the final clause, the $\varphi(m)$ residue classes coprime to m partition the primes not dividing m , and the finitely many primes $p|m$ contribute $O(1)$ to every sum and so do not affect a density. Summing the densities over the $\varphi(m)$ admissible classes gives $\varphi(m) \cdot \frac{1}{\varphi(m)} = 1$, the density of the set of all primes, as we sought to show.

The coefficient $1/\varphi(m)$ is the analytic measure of the principle that, to the resolution the L -function method can detect, the primes prefer no residue class over another. The pole of $L(s, \chi_0)$ at $s = 1$, with residue $\varphi(m)/m$ (proposition 7.2.3), is the single source of the prime sum's divergence, and orthogonality distributes that one pole evenly across the $\varphi(m)$ admissible classes, giving each the coefficient $1/\varphi(m)$. The residue $\varphi(m)/m$ records that only $\varphi(m)$ of the m residue classes modulo m can contain more than one prime at all, the rest sharing a common factor with m ; among those $\varphi(m)$ classes the division is exactly equal. The equidistribution is a statement to leading order: it fixes the coefficient of $\log \frac{1}{s-1}$ but says nothing about the bounded $O(1)$ remainder, and it is in that remainder that the finer arithmetic of the primes modulo m resides, including the systematic bias the next example records.

The smallest moduli put the density $1/\varphi(m)$ within reach of direct computation, and the prime-counting data show both the equidistribution and the size of the fluctuation around it.

Example 7.5.5: Primes in Progressions mod 4 and mod 10

Modulus 4. Since $\varphi(4) = 2$, the admissible classes are 1 and 3, each of Dirichlet density $1/2$ by theorem 7.5.4. Every odd prime falls into one of them (the even prime $2 \equiv 2$ is the sole prime in a non-admissible class), and the two classes are the progressions

$$1, 5, 9, 13, 17, \dots \quad \text{and} \quad 3, 7, 11, 19, 23, \dots$$

Writing $\pi(x; 4, a) = \#\{p \leq x : p \equiv a \pmod{4}\}$ for the counting function of each class, the values

up to several powers of ten are:

x	$\pi(x; 4, 1)$	$\pi(x; 4, 3)$	$\pi(x; 4, 3) - \pi(x; 4, 1)$
10^2	11	13	2
10^3	80	87	7
10^4	609	619	10
10^5	4783	4808	25
10^6	39175	39322	147

The two counts track each other closely: at $x = 10^6$ the ratio $\pi(x; 4, 1)/\pi(x; 4, 3) = 39175/39322 = 0.9963\dots$ is near 1, and each count is close to half of $\pi(10^6) - 1 = 78497$ (subtracting the prime 2), namely $39175 + 39322 = 78497$, split as 0.4990 and 0.5010 of the total. This is the density $1/2$ visible in finite data.

Modulus 10. Since $\varphi(10) = 4$, the admissible classes are 1, 3, 7, 9, each of Dirichlet density $1/4$. These are the primes ending in the decimal digits 1, 3, 7, 9 respectively (the digits 0, 2, 4, 5, 6, 8 forcing a factor of 2 or 5, so 2 and 5 are the only primes outside these four classes). The counts are:

x	$\pi(x; 10, 1)$	$\pi(x; 10, 3)$	$\pi(x; 10, 7)$	$\pi(x; 10, 9)$
10^2	5	7	6	5
10^3	40	42	46	38
10^4	306	310	308	303
10^5	2387	2402	2411	2390
10^6	19617	19665	19621	19593

At $x = 10^6$ the four counts lie within 0.4% of their common mean 19624, and each is close to one quarter of $\pi(10^6) - 2 = 78496$ (subtracting 2 and 5), namely 19624, illustrating the density $1/4$ in each of the four classes.

7.5.6 Remark: The Chebyshev Bias The modulus-4 table shows the count $\pi(x; 4, 3)$ exceeding $\pi(x; 4, 1)$ at every listed x , with a difference that grows but remains small beside either count. This is the *Chebyshev bias*: primes congruent to 3 modulo 4 tend to lead those congruent to 1, despite both classes having density $1/2$. The bias is a phenomenon of the $O(1)$ remainder term in theorem 7.5.4, not of the leading $\frac{1}{\varphi(m)} \log \frac{1}{s-1}$ term, which is exactly equal for the two classes; the equidistribution theorem is silent about it. The bias does not contradict equidistribution and does not persist as a fixed inequality: the sign of $\pi(x; 4, 3) - \pi(x; 4, 1)$ does change for suitable x , the first sign change occurring near $x = 26,861$, and under the generalized Riemann hypothesis the set of x for which $\pi(x; 4, 3) > \pi(x; 4, 1)$ has logarithmic density approximately 0.9959, close to but not equal to 1. The explanation traces to the squares: the class 1 modulo 4 contains the squares of odd primes ($p^2 \equiv 1$ for p odd), giving that class a slight head start in the prime-power count $\psi(x; 4, a)$ that the prime count then echoes. A precise account belongs to the theory of the explicit formula and the distribution of zeros of $L(s, \chi)$; see [4, Ch. 4] for the analytic treatment.

The Dedekind Zeta Function and the Class Number Formula

chapters 4 to 7 built the analytic theory of the primes over a single base field, the field $\mathbb{Q}$ of rational numbers. The Riemann zeta function $\zeta(s) = \sum_{n \geq 1} n^{-s}$ packaged the rational primes through its Euler product, its pole at $s = 1$ measured their abundance, and twisting the construction by a Dirichlet character χ produced the family of L -functions $L(s, \chi)$ that resolved the primes among the residue classes modulo a fixed integer. Every object in that theory, the characters included, was attached to $\mathbb{Q}$: the Dirichlet characters are functions on $\mathbb{Z}$, and the splitting they detect is the splitting of integers into congruence classes. This chapter changes the base field. The field $\mathbb{Q}$ is replaced by an arbitrary number field K , a finite extension of $\mathbb{Q}$ of degree $d = [K : \mathbb{Q}] = r_1 + 2r_2$, where r_1 counts the real embeddings $K \hookrightarrow \mathbb{R}$ and r_2 counts the conjugate pairs of complex embeddings $K \hookrightarrow \mathbb{C}$; in place of $\mathbb{Z}$ stands the ring of integers $\mathcal{O}_K$, the integral closure of $\mathbb{Z}$ in K . The role formerly played by the rational primes is now played by the nonzero prime ideals of $\mathcal{O}_K$, and the central question becomes how a rational prime p decomposes among them.

The function that organizes this decomposition is the *Dedekind zeta function*

$$\zeta_K(s) = \sum_{\mathfrak{a}} (N\mathfrak{a})^{-s},$$

the sum running over the nonzero integral ideals $\mathfrak{a} \subseteq \mathcal{O}_K$, where $N\mathfrak{a} = [\mathcal{O}_K : \mathfrak{a}]$ is the absolute ideal norm. It is the correct generalization of $\zeta(s) = \zeta_{\mathbb{Q}}(s)$: its Euler product runs over the prime ideals $\mathfrak{p}$ of $\mathcal{O}_K$, so that the local factor at a rational prime p records, through the primes $\mathfrak{p}$ lying above p and their residue degrees, exactly how p splits in K . Like ζ , the function ζ_K extends meromorphically past its half-plane of convergence and carries a single simple pole at $s = 1$; but where the residue of ζ at $s = 1$ was simply 1, the residue of ζ_K is a single constant

$$\kappa_K = \text{Res}_{s=1} \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{|d_K|}}$$

assembled at once from all the basic arithmetic invariants of K : the class number $h_K = |\text{Cl}(K)|$, the regulator R_K , the discriminant d_K , and the number w_K of roots of unity in K . This identity, the *analytic class number formula*, is the central result of the chapter (section 8.3). The five sections develop it in order: section 8.1 defines ζ_K , establishes its convergence and Euler product, and works out the Gaussian field as a first example; section 8.2 carries out the geometry-of-numbers count of ideals of bounded norm that produces the pole; section 8.3 assembles the count into the residue κ_K and so proves the class number formula; section 8.4 factors ζ_K for quadratic and cyclotomic K into Dirichlet L -functions, recovering the nonvanishing $L(1, \chi) \neq 0$ as a corollary; and section 8.5 establishes the functional equation of ζ_K and the conductor-discriminant relation. Throughout the chapter, K denotes a number field of degree $d = r_1 + 2r_2$.

8.0.1 Remark: Algebraic Prerequisites From this chapter onward the book treats algebraic number theory as a co-requisite, drawing on it freely and citing [17] for the results it uses. The facts invoked are the structural backbone of the theory of $\mathcal{O}_K$: that $\mathcal{O}_K$ is a Dedekind domain, in which every nonzero ideal factors uniquely into prime ideals; the multiplicativity of the absolute ideal norm, $N(\mathfrak{a}\mathfrak{b}) = N\mathfrak{a} \cdot N\mathfrak{b}$ and $N((\alpha)) = |N_{K/\mathbb{Q}}(\alpha)|$; the finiteness of the ideal class group $\text{Cl}(K)$; Minkowski's geometry of numbers, including the canonical embedding $\iota: K \hookrightarrow \mathbb{R}^{r_1} \times \mathbb{C}^{r_2} \cong \mathbb{R}^d$ and its covolume computation together with Minkowski's convex-body bound; Dirichlet's unit theorem $\mathcal{O}_K^\times \cong \mu_K \times \mathbb{Z}^{r_1+r_2-1}$ and the regulator R_K ; and the Dedekind–Kummer theorem governing the splitting of a rational prime p in $\mathcal{O}_K$. None of these results is reproved here. They are stated where used and referenced to [17], which is the companion volume on the algebraic side.

8.1 The Dedekind Zeta Function

The Riemann zeta function admits two equivalent presentations: as the Dirichlet series $\sum_{n \geq 1} n^{-s}$ summed over positive integers, and as the Euler product $\prod_p (1 - p^{-s})^{-1}$ assembled from one factor per rational prime. The first encodes counting, the second encodes the multiplicative structure of $\mathbb{Z}$, and the equality of the two is the analytic form of unique factorization. The Dedekind zeta function of a number field K is built on the same two presentations, with the integers replaced by the nonzero integral ideals of $\mathcal{O}_K$ and the rational primes replaced by the nonzero prime ideals. The equivalence of the two presentations rests, exactly as over $\mathbb{Q}$, on unique factorization, which for $\mathcal{O}_K$ is the unique factorization of ideals into prime ideals supplied by the Dedekind property ([17, unique factorization of ideals]).

Definition 8.1.1: The Dedekind Zeta Function

Let K be a number field with ring of integers $\mathcal{O}_K$. The *Dedekind zeta function* of K is the series

$$\zeta_K(s) = \sum_{\mathfrak{a} \neq 0} (N\mathfrak{a})^{-s},$$

the sum running over the nonzero integral ideals $\mathfrak{a} \subseteq \mathcal{O}_K$, where $N\mathfrak{a} = [\mathcal{O}_K : \mathfrak{a}]$ is the absolute norm of $\mathfrak{a}$. Equivalently, grouping the ideals by their norm,

$$\zeta_K(s) = \sum_{n=1}^{\infty} \frac{a_K(n)}{n^s}, \quad a_K(n) = \# \{ \mathfrak{a} \subseteq \mathcal{O}_K \mid N\mathfrak{a} = n \},$$

so that ζ_K is the Dirichlet series with coefficients the ideal-counting function a_K .

The two forms agree term by term: each nonzero integral ideal $\mathfrak{a}$ contributes $(N\mathfrak{a})^{-s}$ to the first sum, and collecting the ideals of a fixed norm n contributes $a_K(n) n^{-s}$ to the second. The coefficients $a_K(n)$ are finite for every n , since an ideal of norm n contains the integer n (as $\mathfrak{a} \supseteq n\mathcal{O}_K$ whenever $N\mathfrak{a} = n$, because $[\mathcal{O}_K : \mathfrak{a}] = n$ kills the quotient) and there are only finitely many ideals containing the fixed nonzero ideal $n\mathcal{O}_K$; the second form is therefore a genuine Dirichlet series with finite coefficients.

The simplest case recovers the function with which the analytic theory began.

8.1.2 Remark: The Rational Case Recovers ζ For $K = \mathbb{Q}$ one has $\mathcal{O}_K = \mathbb{Z}$, and the nonzero integral ideals of $\mathbb{Z}$ are exactly the principal ideals $(n) = n\mathbb{Z}$ for $n \geq 1$, each with norm $N((n)) = [\mathbb{Z} : n\mathbb{Z}] = n$. Distinct positive integers give distinct ideals, so $a_{\mathbb{Q}}(n) = \# \{ (m) \mid m \geq 1, N((m)) = n \} = 1$ for every n , and the definition collapses to

$$\zeta_{\mathbb{Q}}(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \zeta(s),$$

the Riemann zeta function of chapter 4. The Dedekind zeta function is thus a strict generalization of ζ , reducing to it over the base field $\mathbb{Q}$.

The first analytic task is to locate the half-plane on which the defining series makes sense. The comparison is with a power of the Riemann zeta function, and the exponent is the degree d of the field.

Proposition 8.1.3: Absolute Convergence and Holomorphy on $\operatorname{Re}(s) > 1$

The series defining $\zeta_K(s)$ converges absolutely on the half-plane $\operatorname{Re}(s) > 1$ and defines a holomorphic function there. More precisely, for real $\sigma > 1$ the comparison

$$\sum_{\mathfrak{a} \neq 0} (N\mathfrak{a})^{-\sigma} \leq \zeta(\sigma)^d$$

holds, where $d = [K : \mathbb{Q}]$.

Proof:

Step 1: The coefficient function a_K is multiplicative. Let $m, n \geq 1$ be coprime. An ideal $\mathfrak{a}$ with $N\mathfrak{a} = mn$ factors uniquely into prime powers, $\mathfrak{a} = \prod_{\mathfrak{p}} \mathfrak{p}^{v_{\mathfrak{p}}}$, by the Dedekind property ([17, unique factorization of ideals]), and by multiplicativity of the absolute norm $N\mathfrak{a} = \prod_{\mathfrak{p}} (N\mathfrak{p})^{v_{\mathfrak{p}}}$ ([17, multiplicativity of the ideal norm]). Since $N\mathfrak{p} = p^{f_{\mathfrak{p}}}$ is a power of the rational prime p below $\mathfrak{p}$, grouping the prime ideals according to the rational prime they lie over partitions $\mathfrak{a}$ as a product $\mathfrak{a} = \mathfrak{b}\mathfrak{c}$ with $N\mathfrak{b} = m$ and $N\mathfrak{c} = n$, where $\mathfrak{b}$ collects the prime powers above the primes dividing m and $\mathfrak{c}$ those above the primes dividing n ; because $\gcd(m, n) = 1$, no rational prime divides both, so this splitting is well defined and unique. The map $\mathfrak{a} \mapsto (\mathfrak{b}, \mathfrak{c})$ is therefore a bijection between the ideals of norm mn and the pairs of ideals of norms m and n , which gives $a_K(mn) = a_K(m)a_K(n)$. Thus a_K is multiplicative.

Step 2: The comparison with $\zeta(\sigma)^d$. Fix a real number $\sigma > 1$. Every term $(N\mathfrak{a})^{-\sigma}$ is positive, so the sum may be rearranged freely; grouping the ideals by their factorization into prime powers and using $N\mathfrak{a} = \prod_{\mathfrak{p}} (N\mathfrak{p})^{v_{\mathfrak{p}}}$ gives, by unique factorization of ideals,

$$\sum_{\mathfrak{a} \neq 0} (N\mathfrak{a})^{-\sigma} = \prod_{\mathfrak{p}} \sum_{k=0}^{\infty} (N\mathfrak{p})^{-k\sigma} = \prod_{\mathfrak{p}} (1 - (N\mathfrak{p})^{-\sigma})^{-1},$$

each geometric series converging because $N\mathfrak{p} \geq 2$ forces $(N\mathfrak{p})^{-\sigma} < 1$. Group the prime ideals by the rational prime p below them:

$$\prod_{\mathfrak{p}} (1 - (N\mathfrak{p})^{-\sigma})^{-1} = \prod_p \prod_{\mathfrak{p}|p} (1 - (N\mathfrak{p})^{-\sigma})^{-1}$$

For a fixed rational prime p , the primes $\mathfrak{p}|p$ satisfy $\sum_{\mathfrak{p}|p} e_{\mathfrak{p}} f_{\mathfrak{p}} = d$ ([17, the degree identity $\sum_{\mathfrak{p}|p} e_{\mathfrak{p}} f_{\mathfrak{p}} = d$]), where $e_{\mathfrak{p}}$ and $f_{\mathfrak{p}}$ are the ramification index and residue degree; since each $e_{\mathfrak{p}}, f_{\mathfrak{p}} \geq 1$, there are at most d primes above p . Moreover $N\mathfrak{p} = p^{f_{\mathfrak{p}}} \geq p$, so $(N\mathfrak{p})^{-\sigma} \leq p^{-\sigma}$ and hence

$$(1 - (N\mathfrak{p})^{-\sigma})^{-1} \leq (1 - p^{-\sigma})^{-1}$$

The inner product over the at most d primes $\mathfrak{p}|p$ is therefore bounded by $(1 - p^{-\sigma})^{-d}$, and consequently

$$\sum_{\mathfrak{a} \neq 0} (N\mathfrak{a})^{-\sigma} = \prod_p \prod_{\mathfrak{p}|p} (1 - (N\mathfrak{p})^{-\sigma})^{-1} \leq \prod_p (1 - p^{-\sigma})^{-d} = \left(\prod_p (1 - p^{-\sigma})^{-1} \right)^d = \zeta(\sigma)^d,$$

the last equality being the Euler product for ζ (corollary 2.3.2). Since $\zeta(\sigma) < \infty$ for $\sigma > 1$, the displayed bound is finite.

Step 3: Absolute convergence and holomorphy on $\operatorname{Re}(s) > 1$. For any s with $\sigma = \operatorname{Re}(s) > 1$ and any ideal $\mathfrak{a}$ one has $|(N\mathfrak{a})^{-s}| = (N\mathfrak{a})^{-\sigma}$, so the bound of Step 2 gives $\sum_{\mathfrak{a}} |(N\mathfrak{a})^{-s}| = \sum_{\mathfrak{a}} (N\mathfrak{a})^{-\sigma} \leq \zeta(\sigma)^d < \infty$, establishing absolute convergence on $\operatorname{Re}(s) > 1$. In the Dirichlet-series form $\zeta_K(s) = \sum_{n \geq 1} a_K(n) n^{-s}$ this is absolute convergence of a Dirichlet series at every point of $\operatorname{Re}(s) > 1$. On any compact subset of the half-plane $\operatorname{Re}(s) > 1$ there is a real $\sigma_0 > 1$ with $\operatorname{Re}(s) \geq \sigma_0$ throughout, and then $|a_K(n) n^{-s}| \leq a_K(n) n^{-\sigma_0}$ with $\sum_n a_K(n) n^{-\sigma_0} \leq \zeta(\sigma_0)^d < \infty$

independent of s ; the Weierstrass M -test gives uniform convergence on the compact set. Each partial sum is holomorphic, so the locally uniform limit $\zeta_K(s)$ is holomorphic on $\operatorname{Re}(s) > 1$ ([16]; this is the convergence theory of Dirichlet series developed in chapter 2).

The product over prime ideals that appeared inside the convergence proof is the Euler product of ζ_K , and it deserves to be stated in its own right: it is the analytic form of unique factorization of ideals, the exact analogue for K of the Euler product of ζ (corollary 2.3.2).

Theorem 8.1.4: Euler Product

For $\operatorname{Re}(s) > 1$ the Dedekind zeta function factors as an absolutely convergent product over the nonzero prime ideals of $\mathcal{O}_K$,

$$\zeta_K(s) = \prod_{\mathfrak{p}} (1 - (N\mathfrak{p})^{-s})^{-1},$$

the product taken over all nonzero prime ideals $\mathfrak{p} \subseteq \mathcal{O}_K$.

Proof:

Step 1: Each local factor is a norm-weighted geometric series. Fix s with $\sigma = \operatorname{Re}(s) > 1$. For a nonzero prime ideal $\mathfrak{p}$ one has $|(N\mathfrak{p})^{-s}| = (N\mathfrak{p})^{-\sigma} \leq 2^{-\sigma} < 1$, so the geometric series sums to

$$(1 - (N\mathfrak{p})^{-s})^{-1} = \sum_{k=0}^{\infty} (N\mathfrak{p})^{-ks} = \sum_{k=0}^{\infty} (N\mathfrak{p}^k)^{-s},$$

the second equality using multiplicativity of the norm, $N\mathfrak{p}^k = (N\mathfrak{p})^k$ ([17, multiplicativity of the ideal norm]). The local factor is thus the sum of $(N\mathfrak{q})^{-s}$ over the powers $\mathfrak{q} = \mathfrak{p}^k$, $k \geq 0$, of the single prime $\mathfrak{p}$.

Step 2: Absolute convergence of the product. An infinite product $\prod_{\mathfrak{p}} (1 - (N\mathfrak{p})^{-s})^{-1}$ converges absolutely precisely when $\sum_{\mathfrak{p}} |(N\mathfrak{p})^{-s}| < \infty$. Grouping primes by the rational prime below them and using $N\mathfrak{p} \geq p$ together with the fact that at most d primes lie over each p (as in Step 2 of proposition 8.1.3),

$$\sum_{\mathfrak{p}} (N\mathfrak{p})^{-\sigma} = \sum_p \sum_{\mathfrak{p}|p} (N\mathfrak{p})^{-\sigma} \leq \sum_p d p^{-\sigma} = d \sum_p p^{-\sigma} < \infty$$

for $\sigma > 1$, since $\sum_p p^{-\sigma} \leq \zeta(\sigma) < \infty$. The product therefore converges absolutely.

Step 3: The product equals the sum. Because the product converges absolutely, its factors may be multiplied out and the resulting terms summed in any order. Multiplying the geometric series of Step 1 over all $\mathfrak{p}$, a generic term in the expansion is a finite product $\prod_{\mathfrak{p}} (N\mathfrak{p}^{k_{\mathfrak{p}}})^{-s}$ with $k_{\mathfrak{p}} \geq 0$ and all but finitely many $k_{\mathfrak{p}}$ equal to zero; by multiplicativity of the norm this equals $(N\mathfrak{a})^{-s}$ for the ideal $\mathfrak{a} = \prod_{\mathfrak{p}} \mathfrak{p}^{k_{\mathfrak{p}}}$. Unique factorization of ideals ([17, unique factorization in Dedekind domains]) asserts that the assignment $(k_{\mathfrak{p}})_{\mathfrak{p}} \mapsto \mathfrak{a} = \prod_{\mathfrak{p}} \mathfrak{p}^{k_{\mathfrak{p}}}$ is a bijection from the finitely-supported exponent sequences onto the nonzero integral ideals. Hence every nonzero

ideal $\mathfrak{a}$ arises exactly once, and the expansion of the product is precisely

$$\prod_{\mathfrak{p}} (1 - (N\mathfrak{p})^{-s})^{-1} = \sum_{\mathfrak{a} \neq 0} (N\mathfrak{a})^{-s} = \zeta_K(s),$$

the absolute convergence established in proposition 8.1.3 justifying the rearrangement into the single sum over ideals. (Equivalently, a_K is multiplicative by Step 1 of proposition 8.1.3, and the identity is the Euler product for a multiplicative Dirichlet coefficient, theorem 2.3.1, applied to the series $\sum_n a_K(n)n^{-s}$ with local factor $\sum_k a_K(p^k)p^{-ks}$ regrouped over the primes above p .)

As with the Riemann zeta function, the Euler product immediately yields nonvanishing throughout the region of absolute convergence: a convergent product of nonzero factors cannot vanish.

Corollary 8.1.5: Nonvanishing on $\operatorname{Re}(s) > 1$

$\zeta_K(s) \neq 0$ for every s with $\operatorname{Re}(s) > 1$.

Proof:

By theorem 8.1.4 the value $\zeta_K(s)$ is an absolutely convergent product over the prime ideals $\mathfrak{p}$, each factor $(1 - (N\mathfrak{p})^{-s})^{-1}$ being the reciprocal of the nonzero number $1 - (N\mathfrak{p})^{-s}$ (nonzero because $|(N\mathfrak{p})^{-s}| = (N\mathfrak{p})^{-\sigma} < 1$ for $\sigma > 1$). An absolutely convergent infinite product all of whose factors are nonzero has a nonzero value (proposition 2.3.4). Hence $\zeta_K(s) \neq 0$.

The convergence and nonvanishing established here mirror exactly the situation for ζ on $\operatorname{Re}(s) > 1$, and they are the easy part of the theory: both rest only on the Euler product, which lives inside the region of absolute convergence. The substantial analytic content of the chapter, the continuation of ζ_K across the line $\operatorname{Re}(s) = 1$ and the determination of its residue there, requires the geometric counting of ideals taken up in section 8.2. Before turning to it, the smallest nontrivial field illustrates how the Euler product encodes splitting, and it previews the factorization that section 8.4 establishes in general.

Example 8.1.6: The Gaussian Field $\mathbb{Q}(i)$

Take $K = \mathbb{Q}(i)$, the field of Gaussian rationals, with ring of integers $\mathcal{O}_K = \mathbb{Z}[i]$, the ring of Gaussian integers. The splitting of a rational prime p in $\mathbb{Z}[i]$ is governed by its residue modulo 4 ([17, splitting of primes in $\mathbb{Q}(i)$]):

1. if $p \equiv 1 \pmod{4}$, then p *splits*, $p\mathbb{Z}[i] = \mathfrak{p}\bar{\mathfrak{p}}$ with $\mathfrak{p} \neq \bar{\mathfrak{p}}$ two distinct primes each of norm $N\mathfrak{p} = N\bar{\mathfrak{p}} = p$;
2. if $p \equiv 3 \pmod{4}$, then p is *inert*, $p\mathbb{Z}[i]$ remains prime, a single prime of norm $N(p\mathbb{Z}[i]) = p^2$;
3. if $p = 2$, then p *ramifies*, $2\mathbb{Z}[i] = (1+i)^2$ (up to a unit), a single prime $(1+i)$ of norm $N((1+i)) = 2$.

The Euler factor of $\zeta_{\mathbb{Q}(i)}$ at p is the product of $(1 - (N\mathfrak{p})^{-s})^{-1}$ over the primes $\mathfrak{p} \mid p$, and the three cases give

$$p \equiv 1: (1 - p^{-s})^{-2}, \quad p \equiv 3: (1 - p^{-2s})^{-1}, \quad p = 2: (1 - 2^{-s})^{-1}$$

Let χ_{-4} denote the nontrivial Dirichlet character modulo 4 (the character written χ_1 in chapter 7), the character with $\chi_{-4}(p) = 1$ for $p \equiv 1 \pmod{4}$, $\chi_{-4}(p) = -1$ for $p \equiv 3 \pmod{4}$, and $\chi_{-4}(2) = 0$. Each Euler factor above factors as $(1 - p^{-s})^{-1}(1 - \chi_{-4}(p)p^{-s})^{-1}$:

- for $p \equiv 1 \pmod{4}$, $\chi_{-4}(p) = 1$ and $(1 - p^{-s})^{-1}(1 - p^{-s})^{-1} = (1 - p^{-s})^{-2}$;
- for $p \equiv 3 \pmod{4}$, $\chi_{-4}(p) = -1$ and $(1 - p^{-s})^{-1}(1 + p^{-s})^{-1} = (1 - p^{-2s})^{-1}$;
- for $p = 2$, $\chi_{-4}(2) = 0$ and $(1 - 2^{-s})^{-1}(1 - 0)^{-1} = (1 - 2^{-s})^{-1}$.

In every case the Euler factor of $\zeta_{\mathbb{Q}(i)}$ at p equals the product of the p -th Euler factors of $\zeta(s)$ and of $L(s, \chi_{-4})$. Taking the product over all primes and invoking the Euler products of ζ (corollary 2.3.2) and of $L(s, \chi_{-4})$ (theorem 7.2.2), valid on $\operatorname{Re}(s) > 1$, this factor-by-factor identity gives

$$\zeta_{\mathbb{Q}(i)}(s) = \zeta(s) L(s, \chi_{-4}), \quad \operatorname{Re}(s) > 1$$

The decomposition exhibits the residue class data of $\mathbb{Z}[i]$ (the splitting type of p) as the value $\chi_{-4}(p)$ of a Dirichlet character, the prototype of the factorization of ζ_K for quadratic fields established in section 8.4. It also connects the analytic invariants of the two threads: the pole of $\zeta_{\mathbb{Q}(i)}$ at $s = 1$ is the pole of ζ , and its residue is $\operatorname{Res}_{s=1} \zeta(s) \cdot L(1, \chi_{-4}) = L(1, \chi_{-4})$, the finite nonzero value $L(1, \chi_{-4}) = \pi/4$ computed in chapter 7. The class number formula of section 8.3 will recover this same residue from the invariants of $\mathbb{Q}(i)$ directly.

8.2 Ideal Counting and Lattice Point Geometry

The Euler product of theorem 8.1.4 presents $\zeta_K(s)$ as a Dirichlet series $\sum_{\mathfrak{a}} (N\mathfrak{a})^{-s} = \sum_{n \geq 1} a_K(n) n^{-s}$ whose coefficients $a_K(n) = \#\{\mathfrak{a} : N\mathfrak{a} = n\}$ count integral ideals of given norm. Whether this series has a pole at $s = 1$, and what its residue there is, is decided entirely by the average size of $a_K(n)$, that is by the growth of the summatory function

$$A_K(x) = \sum_{n \leq x} a_K(n) = \#\{\mathfrak{a} \subseteq \mathcal{O}_K \text{ integral} : N\mathfrak{a} \leq x\}$$

The principal object of this section is the asymptotic law governing $A_K(x)$, whose leading coefficient is exactly the constant κ_K of the notation contract, an expression assembling the class number, the regulator, the discriminant, the number of roots of unity, and the signature of K into a single number. The pole of ζ_K at $s = 1$ and its residue (extracted analytically in section 8.3) are then immediate consequences of this counting estimate, and with them the whole of the analytic class number formula.

The estimate is geometric in nature. An integral ideal of bounded norm is, after passing to a fixed representative of its class, a principal ideal generated by an algebraic integer of bounded field norm, and such generators are exactly the lattice points of the Minkowski embedding lying inside a dilate of a fixed region. Counting them is a problem in the geometry of numbers: the number of lattice points of a full lattice Λ inside a large homothetic copy tS of a fixed body S is asymptotically $\operatorname{vol}(S)/\operatorname{vol}(\mathbb{R}^d/\Lambda)$ times t^d , with an error controlled by the surface area of S . The same lattice-point principle underlies the elementary hyperbola counting of chapter 1, now carried out in dimension $d = [K : \mathbb{Q}]$ against the norm-form region rather than under a hyperbola in the plane.

The argument is organized into five steps. *Step 1* reduces the count to a single ideal class. *Step 2* converts counting in a class to counting principal ideals in a fixed ideal, modulo the action of the unit group. *Step 3* realizes that count geometrically through the Minkowski embedding. *Step 4* is the lattice-point counting estimate itself. *Step 5* evaluates the volume of the relevant region as $2^{r_1} \pi^{r_2} R_K$, the appearance of the regulator being the heart of the matter. The algebraic prerequisites (that $\mathcal{O}_K$ is a Dedekind domain with finite class group, the multiplicativity of the norm, the covolume of an ideal under the Minkowski embedding, and Dirichlet's unit theorem) are imported from [17].

Theorem 8.2.1: Asymptotics of the Ideal-Counting Function

Let K be a number field of degree $d = [K : \mathbb{Q}] = r_1 + 2r_2$, with r_1 real and r_2 conjugate pairs of complex embeddings. Then, as $x \rightarrow \infty$,

$$A_K(x) = \#\{\mathfrak{a} \subseteq \mathcal{O}_K \text{ integral} : N\mathfrak{a} \leq x\} = \kappa_K x + O(x^{1-1/d})$$

where

$$\kappa_K = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{|d_K|}}$$

with h_K the class number, R_K the regulator, d_K the discriminant, and w_K the number of roots of unity in K . The implied constant depends only on K .

The proof divides the count over the ideal class group and reduces, by the proposition stated immediately after, to a single class; that proposition carries the entire geometric content. The deduction of the theorem from the proposition is the bookkeeping of *Step 1*.

Proof:

Step 1 (reduction to one ideal class). The set of nonzero integral ideals of $\mathcal{O}_K$ is partitioned by the ideal class group $\text{Cl}(K)$, which is finite of order $h_K = |\text{Cl}(K)|$ ([17]). Since the norm $N\mathfrak{a}$ depends only on $\mathfrak{a}$,

$$A_K(x) = \sum_{C \in \text{Cl}(K)} \#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} \tag{8.1}$$

a sum of h_K terms. By proposition 8.2.2 below, each term equals

$$\#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} = \frac{2^{r_1} (2\pi)^{r_2} R_K}{w_K \sqrt{|d_K|}} x + O(x^{1-1/d})$$

with the same main term for every class C . Summing over the h_K classes multiplies the main term by h_K , and the sum of h_K error terms (the number of summands being the constant h_K) is again $O(x^{1-1/d})$. Hence

$$A_K(x) = h_K \cdot \frac{2^{r_1} (2\pi)^{r_2} R_K}{w_K \sqrt{|d_K|}} x + O(x^{1-1/d}) = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{|d_K|}} x + O(x^{1-1/d})$$

which is the asserted asymptotic with leading coefficient κ_K .

The whole of the work is therefore the per-class count, which is the next proposition. Its proof, occupying Steps 2 through 5, is the geometric core of the chapter.

Proposition 8.2.2: Counting in a Fixed Ideal Class

For every ideal class $C \in \text{Cl}(K)$, the number of integral ideals in C of norm at most x satisfies, as $x \rightarrow \infty$,

$$\#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} = \frac{2^{r_1}(2\pi)^{r_2} R_K}{w_K \sqrt{|d_K|}} x + O(x^{1-1/d})$$

with implied constant depending only on K (and not on C).

The proof relies on one external input from the geometry of numbers, the lattice-point estimate stated and proved next, after which Steps 2 through 5 assemble the proposition.

Lemma 8.2.3: Lipschitz Lattice-Point Principle

Let $\Lambda \subset \mathbb{R}^d$ be a full lattice and let $S \subset \mathbb{R}^d$ be a bounded set whose boundary ∂S is contained in the union of the images of finitely many Lipschitz maps $[0, 1]^{d-1} \rightarrow \mathbb{R}^d$. Then, as $t \rightarrow \infty$,

$$\#(\Lambda \cap tS) = \frac{\text{vol}(S)}{\text{vol}(\mathbb{R}^d/\Lambda)} t^d + O(t^{d-1})$$

the implied constant depending only on Λ and on the Lipschitz constants and number of the parametrizing maps.

Proof:

This is the Lipschitz lattice-point estimate of the geometry of numbers; a complete proof is given in [3, Ch. VI, §2]. The mechanism is the following. Tile $\mathbb{R}^d$ by the translates of a fundamental parallelepiped P of Λ , of volume $\text{vol}(P) = \text{vol}(\mathbb{R}^d/\Lambda)$, and to each lattice point $\lambda \in \Lambda$ attach the tile $\lambda + P$. Counting tiles whose corner λ lies in tS approximates the volume: the union of these tiles differs from tS only by tiles meeting the boundary $\partial(tS) = t\partial S$, so

$$|\#(\Lambda \cap tS) \cdot \text{vol}(P) - \text{vol}(tS)| \leq \text{vol}(P) \cdot \#\{\lambda \in \Lambda : (\lambda + P) \cap t\partial S \neq \emptyset\}$$

A tile meeting $t\partial S$ lies within bounded distance (the diameter of P) of that set. The hypothesis on ∂S bounds the number of such tiles: the image of $[0, 1]^{d-1}$ under a Lipschitz map with constant L , scaled by t , can be covered by $O((Lt)^{d-1})$ balls of unit radius, each meeting $O(1)$ tiles, so each of the finitely many boundary pieces accounts for $O(t^{d-1})$ tiles. Summing over the finitely many pieces, the boundary contributes $O(t^{d-1})$ tiles. Since $\text{vol}(tS) = \text{vol}(S)t^d$ by homogeneity of Lebesgue measure, dividing through by $\text{vol}(P)$ gives $\#(\Lambda \cap tS) = \text{vol}(S)t^d/\text{vol}(\mathbb{R}^d/\Lambda) + O(t^{d-1})$, as asserted.

Proof of proposition 8.2.2:

Fix the class C throughout.

Step 2 (from a class to principal ideals in a fixed ideal, modulo units). Choose, once and for all, an integral ideal $\mathfrak{b}$ lying in the inverse class C^{-1} ; such an ideal exists because every class contains integral ideals ([17]). The strategy is to clear the obstruction to principality by multiplying through by $\mathfrak{b}$. The map

$$\mathfrak{a} \mapsto \mathfrak{a}\mathfrak{b} \tag{8.2}$$

sends an integral ideal $\mathfrak{a} \in C$ to the integral ideal $\mathfrak{a}\mathfrak{b}$, whose class is $C \cdot C^{-1} = 1$, hence $\mathfrak{a}\mathfrak{b}$ is *principal*, say $\mathfrak{a}\mathfrak{b} = (\alpha)$. The generator α is nonzero, and since $\mathfrak{a}\mathfrak{b} \subseteq \mathfrak{b}$ one has $\alpha \in \mathfrak{b}$. Conversely, if $(\alpha) \subseteq \mathfrak{b}$ is a nonzero principal ideal with $\alpha \in \mathfrak{b}$, then $\mathfrak{b} | (\alpha)$ in the sense of Dedekind divisibility, so $(\alpha) = \mathfrak{a}\mathfrak{b}$ for a unique integral ideal $\mathfrak{a} = (\alpha)\mathfrak{b}^{-1}$, whose class is C . Thus (8.2) is a bijection

$$\{\mathfrak{a} \in C : \mathfrak{a} \text{ integral}\} \xrightarrow{\sim} \{(\alpha) : 0 \neq \alpha \in \mathfrak{b}\} / \sim \quad (8.3)$$

between integral ideals in C and nonzero principal ideals generated by elements of $\mathfrak{b}$, where on the right one counts principal *ideals*, that is, generators up to units: $(\alpha) = (\alpha')$ if and only if $\alpha' = u\alpha$ for some unit $u \in \mathcal{O}_K^\times$.

The bijection respects norms. By multiplicativity of the absolute norm and the identity $N((\alpha)) = |N_{K/\mathbb{Q}}(\alpha)|$ ([17]),

$$|N_{K/\mathbb{Q}}(\alpha)| = N((\alpha)) = N(\mathfrak{a}\mathfrak{b}) = N\mathfrak{a} \cdot N\mathfrak{b} \quad (8.4)$$

so the constraint $N\mathfrak{a} \leq x$ transcribes to $|N_{K/\mathbb{Q}}(\alpha)| \leq x N\mathfrak{b}$. Therefore

$$\#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} = \#\{\alpha \in \mathfrak{b} \setminus \{0\} : |N_{K/\mathbb{Q}}(\alpha)| \leq x N\mathfrak{b}\} / \sim \quad (8.5)$$

the quotient again being by the multiplicative action of $\mathcal{O}_K^\times$. The problem is now to count, up to units, the nonzero elements of $\mathfrak{b}$ whose field norm is bounded by $x N\mathfrak{b}$.

Step 3 (geometry of numbers). Counting elements of $\mathfrak{b}$ is a lattice problem under the Minkowski embedding. Let $\sigma_1, \dots, \sigma_{r_1}$ be the real embeddings of K and $\sigma_{r_1+1}, \dots, \sigma_{r_1+r_2}$ one from each conjugate pair of complex embeddings, and write

$$\iota : K \hookrightarrow V := \mathbb{R}^{r_1} \times \mathbb{C}^{r_2} \cong \mathbb{R}^d, \quad \iota(\alpha) = (\sigma_1(\alpha), \dots, \sigma_{r_1}(\alpha); \sigma_{r_1+1}(\alpha), \dots, \sigma_{r_1+r_2}(\alpha))$$

for the canonical (Minkowski) embedding. Under ι the ideal $\mathfrak{b}$ maps to a full lattice $\iota(\mathfrak{b}) \subset V \cong \mathbb{R}^d$ of covolume

$$\text{vol}(\mathbb{R}^d / \iota(\mathfrak{b})) = 2^{-r_2} \sqrt{|d_K|} N\mathfrak{b} \quad (8.6)$$

where $\mathbb{R}^d$ carries Lebesgue measure transported through the identification $\mathbb{C} \cong \mathbb{R}^2$ by $z \mapsto (\text{Re } z, \text{Im } z)$ ([17]; the factor 2^{-r_2} records that identification). On V the norm form

$$|N(v)| = \prod_{j=1}^{r_1} |x_j| \cdot \prod_{k=1}^{r_2} |z_k|^2, \quad v = (x_1, \dots, x_{r_1}; z_1, \dots, z_{r_2})$$

is the continuous extension of $\alpha \mapsto |N_{K/\mathbb{Q}}(\alpha)| = \prod_j |\sigma_j(\alpha)| \cdot \prod_k |\sigma_{r_1+k}(\alpha)|^2$ to all of V , and it is *homogeneous of degree d* : for $t > 0$,

$$|N(tv)| = t^{r_1} \cdot (t^2)^{r_2} |N(v)| = t^{r_1+2r_2} |N(v)| = t^d |N(v)| \quad (8.7)$$

since each real coordinate scales linearly and each complex coordinate contributes its squared modulus.

It remains to handle the quotient by units. By Dirichlet's unit theorem the unit group factors as $\mathcal{O}_K^\times \cong \mu_K \times \mathbb{Z}^{r_1+r_2-1}$, where μ_K is the finite cyclic group of roots of unity, of order w_K , and the free part has rank $r := r_1 + r_2 - 1$ ([17]). The torsion μ_K acts on $\iota(\mathfrak{b}) \setminus \{0\}$ freely, since a root of unity fixing a nonzero α would equal 1; it partitions the nonzero elements into orbits

of size exactly w_K . The free part acts through the logarithmic embedding, and one selects a fundamental domain for it on the norm-one hypersurface. Write

$$V^\times := \{v \in V : |N(v)| \neq 0\}$$

for the locus where no coordinate vanishes. There is a measurable *cone* $D \subseteq V^\times$ (closed under positive real scaling, $tD = D$ for $t > 0$) which is a fundamental domain for the action of the free units $\mathcal{O}_K^\times/\mu_K \cong \mathbb{Z}^r$ on $V^\times$, so that each orbit of the free units meets D in exactly one point; the construction of D and its cone property are recorded in [3, Ch. VI, §3] and [5, Ch. VII, §5]. Setting

$$S := \{v \in D : 0 < |N(v)| \leq 1\} \quad (8.8)$$

the cone property together with the homogeneity (8.7) gives, for every $Y > 0$, the homothety relation

$$\{v \in D : 0 < |N(v)| \leq Y\} = Y^{1/d} S \quad (8.9)$$

indeed $v \in D$ with $|N(v)| \leq Y$ if and only if $w := Y^{-1/d}v \in D$ (as D is a cone) with $|N(w)| = Y^{-1}|N(v)| \leq 1$, that is $w \in S$. The element-count of Step 2, taken up to the full unit group, is now a lattice-point count in such a region divided by w_K for the torsion: selecting the unique representative of each free-unit orbit inside D and one of the w_K torsion representatives,

$$\#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} = \frac{1}{w_K} \#(\iota(\mathfrak{b}) \cap (x N\mathfrak{b})^{1/d} S) \quad (8.10)$$

where (8.9) with $Y = x N\mathfrak{b}$ rewrites $\{v \in D : 0 < |N(v)| \leq x N\mathfrak{b}\}$ as $(x N\mathfrak{b})^{1/d} S$ and (8.5) supplies the norm bound. The region S is bounded, and its boundary is the union of finitely many images of Lipschitz maps from $[0, 1]^{d-1}$, by the explicit description of D through the logarithmic coordinates of [3, Ch. VI, §3]; this regularity is exactly the hypothesis of lemma 8.2.3. (In the rank-zero cases $r = 0$ the cone D is all of $V^\times$ and S is visibly a disk or an interval, as computed in Step 5; no unit theorem is needed there.)

Step 4 (lattice-point counting). Apply lemma 8.2.3 with the lattice $\Lambda = \iota(\mathfrak{b})$, the region S of (8.8), and the scaling parameter $t = (x N\mathfrak{b})^{1/d}$, so that $t^d = x N\mathfrak{b}$ and $t^{d-1} = (x N\mathfrak{b})^{(d-1)/d} = O(x^{1-1/d})$ as $x \rightarrow \infty$ (with $N\mathfrak{b}$ a constant depending only on C). Using the covolume (8.6),

$$\#(\iota(\mathfrak{b}) \cap (x N\mathfrak{b})^{1/d} S) = \frac{\text{vol}(S)}{2^{-r_2} \sqrt{|d_K|} N\mathfrak{b}} x N\mathfrak{b} + O(x^{1-1/d}) = \frac{\text{vol}(S)}{2^{-r_2} \sqrt{|d_K|}} x + O(x^{1-1/d}) \quad (8.11)$$

the factor $N\mathfrak{b}$ cancelling between numerator and covolume. This cancellation is the structural reason the count is independent of the choice of $\mathfrak{b}$, and (since Step 5 shows $\text{vol}(S)$ to depend only on K) independent of the class C . Dividing by w_K as in (8.10),

$$\#\{\mathfrak{a} \in C : N\mathfrak{a} \leq x\} = \frac{\text{vol}(S)}{w_K 2^{-r_2} \sqrt{|d_K|}} x + O(x^{1-1/d}) \quad (8.12)$$

Everything is now explicit except $\text{vol}(S)$, the volume of the norm-bounded fundamental region.

Step 5 (the volume constant). The claim that finishes the proof is

$$\text{vol}(S) = 2^{r_1} \pi^{r_2} R_K \quad (8.13)$$

The two rank-zero cases, where $r = r_1 + r_2 - 1 = 0$ and the regulator is by convention $R_K = 1$ (the empty determinant), are elementary and are done in full; they are also the cases of immediate use in section 8.3.

Case $\mathbb{Q}$ ($r_1 = 1, r_2 = 0, d = 1$). Here $V = \mathbb{R}$, the unit group is $\{\pm 1\}$ with $w_K = 2$ and free rank 0, so $D = \mathbb{R}^\times$ and the norm form is $|N(x)| = |x|$. The region is

$$S = \{x \in \mathbb{R} : 0 < |x| \leq 1\} = [-1, 0) \cup (0, 1]$$

of length $\text{vol}(S) = 2 = 2^1 = 2^{r_1}$, in agreement with (8.13) since $\pi^0 R_{\mathbb{Q}} = 1$. As a check, (8.12) then gives $\#\{\mathfrak{a} \subseteq \mathbb{Z} : N\mathfrak{a} \leq x\} = \frac{2}{2 \cdot 1 \cdot 1} x + O(1) = x + O(1)$, the count of positive integers at most x , as it must be.

Case imaginary quadratic ($r_1 = 0, r_2 = 1, d = 2$). Here $V = \mathbb{C}$, the free unit rank is 0, and $w_K \in \{2, 4, 6\}$ is the number of roots of unity. The cone D is all of $\mathbb{C}^\times$ and the norm form is $|N(z)| = |z|^2$, so

$$S = \{z \in \mathbb{C} : 0 < |z|^2 \leq 1\} = \{z \in \mathbb{C} : 0 < |z| \leq 1\}$$

the closed unit disk with the origin removed. Its area is that of the unit disk,

$$\text{vol}(S) = \pi \cdot 1^2 = \pi = \pi^1 = \pi^{r_2}$$

in agreement with (8.13) since $2^0 R_K = 1$. This is the geometric heart of the imaginary quadratic class number formula: the per-class count (8.12) becomes $\frac{\pi}{w_K 2^{-1} \sqrt{|d_K|}} x = \frac{2\pi}{w_K \sqrt{|d_K|}} x$, recovering the coefficient $2^{r_1} (2\pi)^{r_2} R_K / (w_K \sqrt{|d_K|})$ with $r_1 = 0, r_2 = 1, R_K = 1$.

General case. When $r = r_1 + r_2 - 1 \geq 1$ the volume (8.13) is computed by a change of variables to “norm-and-logarithm” coordinates adapted to the unit action, in which the integral over S separates into an angular part contributing the powers of 2 and π , a radial part of length 1 (from the norm constraint $0 < |N(v)| \leq 1$), and a part over the free-unit directions whose volume is by definition the covolume of the logarithmic unit lattice, namely the regulator R_K . In outline: pass each real coordinate to $(\text{sign } x_j, |x_j|)$ and each complex coordinate to polar form (θ_k, ρ_k) with $\rho_k = |z_k|$; the sign choices and angles θ_k contribute the factor $2^{r_1} \cdot (2\pi)^{r_2}$ on integration, after which the radial variables $(|x_j|, \rho_k)$ are transformed to $(\log |N(v)|; \ell_1, \dots, \ell_r)$, the ℓ_i being the coordinates of the logarithmic embedding along the unit directions. The fundamental domain D becomes a unit box in the ℓ -coordinates of volume equal to the covolume of the log-unit lattice, that is R_K ; the norm constraint $0 < |N(v)| \leq 1$ cuts out a single unit of length in the $\log |N(v)|$ direction; and the Jacobian of the substitution is constant, arranged so that a factor 2^{r_2} is absorbed against the $\rho_k d\rho_k d\theta_k$ area element of each complex place, leaving $2^{r_1} \pi^{r_2}$ in place of $2^{r_1} (2\pi)^{r_2}$. The complete Jacobian computation, including the precise normalization of the logarithmic coordinates and the verification that the resulting box has volume R_K , is carried out in [3, Ch. VI, §3] and, in the form adapted to the present residue, in [5, Ch. VII, §5]. The outcome is (8.13), consistent with both rank-zero cases above (where $R_K = 1$ and the box degenerates to a point).

Conclusion. Insert (8.13) into (8.12) and simplify the constant. The factor 2^{-r_2} in the denominator inverts to 2^{r_2} in the numerator, so

$$\frac{\text{vol}(S)}{w_K 2^{-r_2} \sqrt{|d_K|}} = \frac{2^{r_1} \pi^{r_2} R_K}{w_K 2^{-r_2} \sqrt{|d_K|}} = \frac{2^{r_1} 2^{r_2} \pi^{r_2} R_K}{w_K \sqrt{|d_K|}} = \frac{2^{r_1} (2\pi)^{r_2} R_K}{w_K \sqrt{|d_K|}} \quad (8.14)$$

the last equality from $2^{r_2}\pi^{r_2} = (2\pi)^{r_2}$. This is precisely the coefficient asserted in proposition 8.2.2. The leading coefficient depends only on the invariants of K , not on the class C ; the implied constant in the error term, which a priori depends on the chosen ideal $\mathfrak{b}$ (and hence on C), is made uniform across the classes by fixing one representative $\mathfrak{b}$ per class and taking the largest constant among the h_K choices.

The estimate just proved is the entire arithmetic input to the analytic class number formula. Two features deserve emphasis. First, the leading coefficient κ_K is genuinely a product of *all* the basic invariants of K at once: the signature (r_1, r_2) through the powers of 2 and 2π , the class number h_K from the number of summands in (8.1), the regulator R_K from the volume (8.13), the discriminant through the covolume (8.6), and the roots of unity w_K from the torsion of the unit group. No single step produces the formula; it assembles from the geometry of every step together. Second, the error exponent $1 - 1/d$ is dictated by the codimension-one boundary of the norm region: the surface of tS scales as t^{d-1} , and with $t = (xN\mathfrak{b})^{1/d}$ this is $x^{(d-1)/d} = x^{1-1/d}$. For $K = \mathbb{Q}$ the exponent degenerates to 0, recovering the trivial $A_{\mathbb{Q}}(x) = x + O(1)$; for higher degree the boundary error is a genuine $o(x)$ but no better at this level of generality.

A comparison with the rational case of chapter 1 situates the argument. There the counting was of lattice points (a, b) under the hyperbola $ab \leq x$, a two-dimensional region with main term $x \log x$ for the divisor sum; here the region is the d -dimensional norm body tS , its volume linear in x (after the d -th-root scaling) because the norm form is homogeneous of degree d , matching the ambient dimension exactly. The boundary-error principle is identical in both: a region with rectifiable boundary contains a number of lattice points equal to its volume divided by the covolume, up to a surface-order error. What is new over $\mathbb{Q}$ is the quotient by units, which forces the passage to a fundamental domain and is the precise point at which the regulator enters. In the next section this counting asymptotic is converted, by a direct Dirichlet-series argument, into the pole of ζ_K at $s = 1$ with residue κ_K , completing the analytic class number formula.

8.3 The Analytic Class Number Formula

The two preceding sections prepared the two halves of a single statement. section 8.1 introduced the Dedekind zeta function $\zeta_K(s) = \sum_{\mathfrak{a}} N\mathfrak{a}^{-s} = \sum_{n \geq 1} a_K(n) n^{-s}$ as the Dirichlet series whose coefficients $a_K(n) = \#\{\mathfrak{a} : N\mathfrak{a} = n\}$ count integral ideals by norm, and established its Euler product over the prime ideals of $\mathcal{O}_K$ (theorem 8.1.4). section 8.2 fixed the average size of those coefficients: the summatory function $A_K(x) = \sum_{n \leq x} a_K(n)$, the number of integral ideals of norm at most x , satisfies the asymptotic $A_K(x) = \kappa_K x + O(x^{1-1/d})$, where κ_K is the geometric constant assembled from the basic invariants of K (theorem 8.2.1). What remains is to read the analytic structure of ζ_K off this counting estimate, and to identify the constant κ_K in arithmetic terms.

The passage from the counting estimate to the analytic structure is purely formal, and it reproduces, for an arbitrary number field, the mechanism by which the meromorphic continuation of the Riemann zeta function was extracted from the elementary asymptotics of $\sum_{n \leq x} n^{-s}$ in theorem 4.2.1. A Dirichlet series with non-negative coefficients whose partial sums grow linearly, $A_K(x) = \kappa_K x + O(x^{1-\delta})$, is governed near $s = 1$ entirely by the leading coefficient κ_K : the series converges on $\operatorname{Re}(s) > 1$, continues across the line $\operatorname{Re}(s) = 1$, and acquires there a single simple pole whose residue is exactly κ_K . The error term, of size $O(x^{1-\delta})$, contributes only a holomorphic correction on the wider half-plane $\operatorname{Re}(s) > 1 - \delta$. Isolating this principle as a lemma, applicable to any series of this shape, and then feeding it the specific estimate of theorem 8.2.1, yields the analytic class number formula in two

strokes.

Lemma 8.3.1: From Counting to a Pole

Let $(a(n))_{n \geq 1}$ be a sequence of non-negative real numbers, and suppose its summatory function $A(x) = \sum_{n \leq x} a(n)$ satisfies

$$A(x) = \kappa x + O(x^{1-\delta})$$

for some constant κ and some $\delta \in (0, 1]$. Then the Dirichlet series $f(s) = \sum_{n \geq 1} a(n) n^{-s}$ converges absolutely on the half-plane $\operatorname{Re}(s) > 1$ and extends to a meromorphic function on the half-plane $\operatorname{Re}(s) > 1 - \delta$ whose only singularity there is a simple pole at $s = 1$ with residue κ . Explicitly, writing $A(x) = \kappa x + E(x)$ with $E(x) = O(x^{1-\delta})$, for all s with $\operatorname{Re}(s) > 1 - \delta$ and $s \neq 1$,

$$f(s) = \kappa + \frac{\kappa}{s-1} + s \int_1^\infty E(x) x^{-s-1} dx \quad (8.15)$$

Proof:

Since $A(x) = \kappa x + O(x^{1-\delta})$ with $\delta \leq 1$, the partial sums $A(x)$ are $O(x)$, so $a(n) = A(n) - A(n-1) = O(n)$; hence $|a(n)|n^{-\sigma} = O(n^{1-\sigma})$, and $\sum_n a(n)n^{-s}$ converges absolutely for $\operatorname{Re}(s) = \sigma > 2$. This crude bound is superseded below, where the integral representation will give convergence on the whole half-plane $\operatorname{Re}(s) > 1$; the present remark only secures a region on which f is defined as a series, so that the identity theorem may later be invoked. The argument has four stages: an integral representation of f valid for $\operatorname{Re}(s) > 1$; the splitting of that integral into a polar term and an error integral; the holomorphy of the error integral on $\operatorname{Re}(s) > 1 - \delta$; and the identification of the pole and its residue.

Step 1: The integral representation $f(s) = s \int_1^\infty A(x) x^{-s-1} dx$ for $\operatorname{Re}(s) > 1$. Fix s with $\sigma = \operatorname{Re}(s) > 1$. For each $n \geq 1$ the elementary identity

$$n^{-s} = s \int_n^\infty x^{-s-1} dx$$

holds, since $\sigma > 0$ makes the integral convergent and the fundamental theorem of calculus evaluates it to $[-x^{-s}]_n^\infty = n^{-s}$. Multiplying by $a(n) \geq 0$ and summing over n ,

$$f(s) = \sum_{n \geq 1} a(n) n^{-s} = \sum_{n \geq 1} a(n) s \int_n^\infty x^{-s-1} dx = s \int_1^\infty \left(\sum_{n \leq x} a(n) \right) x^{-s-1} dx = s \int_1^\infty A(x) x^{-s-1} dx$$

The interchange of summation and integration is justified by Tonelli's theorem ([8]): the summand-integrand $a(n) |x^{-s-1}| = a(n) x^{-\sigma-1}$ is non-negative, and the resulting iterated integral s replaced by σ gives $\sigma \int_1^\infty A(x) x^{-\sigma-1} dx$, which is finite because $A(x) = O(x)$ forces the integrand to be $O(x^{-\sigma})$ with $\sigma > 1$. The collapse of the double sum to $\sum_{n \leq x} a(n) = A(x)$ uses that the condition $n \leq x$ is exactly the range on which the inner integral $\int_n^\infty$ contributes to the outer integral $\int_1^\infty (\cdot) dx$ at the point x . This is partial summation in integral form, the same device that produced the representation of ζ in theorem 4.2.1.

Step 2: Splitting off the polar term. Substitute $A(x) = \kappa x + E(x)$ into the representation of Step 1. The two pieces integrate separately for $\operatorname{Re}(s) > 1$, where both converge absolutely. The polar

piece is

$$s \int_1^\infty \kappa x \cdot x^{-s-1} dx = \kappa s \int_1^\infty x^{-s} dx = \kappa s \cdot \frac{1}{s-1} = \kappa + \frac{\kappa}{s-1}$$

the integral $\int_1^\infty x^{-s} dx = 1/(s-1)$ being valid for $\operatorname{Re}(s) > 1$, and the last equality using $s/(s-1) = 1 + 1/(s-1)$. The error piece is $s \int_1^\infty E(x) x^{-s-1} dx$. Combining,

$$f(s) = \kappa + \frac{\kappa}{s-1} + s \int_1^\infty E(x) x^{-s-1} dx \quad (\operatorname{Re}(s) > 1) \quad (8.16)$$

which is (8.15) on the region $\operatorname{Re}(s) > 1$ where it was derived.

Step 3: The error integral is holomorphic on $\operatorname{Re}(s) > 1 - \delta$. Let $J(s) = \int_1^\infty E(x) x^{-s-1} dx$ denote the error integral. By hypothesis there is a constant $C > 0$ with $|E(x)| \leq C x^{1-\delta}$ for all $x \geq 1$, so the integrand is dominated by

$$|E(x) x^{-s-1}| = |E(x)| x^{-\sigma-1} \leq C x^{1-\delta} x^{-\sigma-1} = C x^{-\sigma-\delta}$$

where $\sigma = \operatorname{Re}(s)$. Fix a compact set $\mathcal{K} \subset \{\operatorname{Re}(s) > 1 - \delta\}$ and put $\sigma_0 = \min_{s \in \mathcal{K}} \operatorname{Re}(s)$, so that $\sigma_0 > 1 - \delta$, equivalently $\sigma_0 + \delta > 1$. Then for every $s \in \mathcal{K}$ and every $x \geq 1$ the bound $|E(x) x^{-s-1}| \leq C x^{-\sigma_0-\delta}$ holds, and

$$\int_1^\infty C x^{-\sigma_0-\delta} dx = \frac{C}{\sigma_0 + \delta - 1} < \infty$$

because $\sigma_0 + \delta - 1 > 0$. The dominating function $C x^{-\sigma_0-\delta}$ is independent of s and integrable on $[1, \infty)$, so $J(s)$ converges absolutely and uniformly for $s \in \mathcal{K}$. For each fixed $x \geq 1$ the integrand $E(x) x^{-s-1}$ is an entire function of s , and a parameter integral with holomorphic integrand and a locally uniform integrable bound is holomorphic in the parameter, by the standard theorem on holomorphy of integrals depending on a parameter ([16]). Since the half-plane $\operatorname{Re}(s) > 1 - \delta$ is exhausted by such compact sets $\mathcal{K}$, the function J is holomorphic on all of $\operatorname{Re}(s) > 1 - \delta$. Consequently $sJ(s)$ is holomorphic there as well, being the product of the entire function s with J .

Step 4: Continuation, the pole, and the residue. The right-hand side of (8.15),

$$R(s) := \kappa + \frac{\kappa}{s-1} + sJ(s)$$

is, by Step 3, meromorphic on $\operatorname{Re}(s) > 1 - \delta$: the term κ is constant, $sJ(s)$ is holomorphic, and $\kappa/(s-1)$ is rational with a single simple pole at $s = 1$. By (8.16) the functions R and f agree on $\operatorname{Re}(s) > 1$, and there f is holomorphic (the representation of Step 1 shows f extends holomorphically to $\operatorname{Re}(s) > 1$, where the series, the integral, and hence f all converge, refining the crude region $\operatorname{Re}(s) > 2$ of absolute convergence noted at the outset). The set $\{\operatorname{Re}(s) > 1\}$ is a nonempty open subset of the connected open set $\{\operatorname{Re}(s) > 1 - \delta\}$ and has accumulation points there, so by the identity theorem ([16]) R is the unique meromorphic extension of f to $\operatorname{Re}(s) > 1 - \delta$. Its only singularity is the simple pole of $\kappa/(s-1)$ at $s = 1$, the holomorphic term $sJ(s)$ being unable to cancel it. For the residue, $sJ(s)$ is holomorphic, hence finite, at $s = 1$, so

$$\operatorname{Res}_{s=1} f = \lim_{s \rightarrow 1} (s-1) f(s) = \lim_{s \rightarrow 1} \left[(s-1) \kappa + \kappa + (s-1) sJ(s) \right] = \kappa$$

the first and third terms vanishing in the limit. Thus the pole is simple with residue κ , and (8.15) holds throughout $\operatorname{Re}(s) > 1 - \delta$ with $s \neq 1$.

The lemma is stated for an abstract sequence so that the only field-specific input is the counting estimate. In particular, the non-negativity of the coefficients was used in Step 1 only to license Tonelli's theorem in the interchange; for a general (signed) Dirichlet series the same conclusion holds, by a routine absolute-convergence argument in place of Tonelli, but the present formulation is exactly adapted to the ideal-counting coefficients $a_K(n) \geq 0$. With the lemma in hand, the analytic class number formula is immediate.

Theorem 8.3.2: The Analytic Class Number Formula

Let K be a number field of degree $d = [K : \mathbb{Q}] = r_1 + 2r_2$, with r_1 real embeddings and r_2 conjugate pairs of complex embeddings. The Dedekind zeta function $\zeta_K(s)$ extends to a meromorphic function on the half-plane $\operatorname{Re}(s) > 1 - 1/d$ whose only singularity there is a simple pole at $s = 1$. Its residue at this pole is

$$\operatorname{Res}_{s=1} \zeta_K(s) = \lim_{s \rightarrow 1} (s-1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{|d_K|}} = \kappa_K \quad (8.17)$$

where $h_K = |\operatorname{Cl}(K)|$ is the class number, $R_K = \operatorname{Reg}(K)$ the regulator, w_K the number of roots of unity in K , and d_K the discriminant.

Proof:

The coefficients $a_K(n) = \#\{\mathfrak{a} : N\mathfrak{a} = n\}$ are non-negative integers, and by theorem 8.2.1 their summatory function satisfies $A_K(x) = \kappa_K x + O(x^{1-1/d})$, with the constant κ_K given by the right-hand side of (8.17). This is precisely the hypothesis of lemma 8.3.1 with $a(n) = a_K(n)$, $\kappa = \kappa_K$, and $\delta = 1/d \in (0, 1]$. The lemma then asserts that $\zeta_K(s) = \sum_n a_K(n) n^{-s}$ extends meromorphically to $\operatorname{Re}(s) > 1 - 1/d$ with a single simple pole at $s = 1$ of residue κ_K , which is the claim.

The residue (8.17) is the central identity of the chapter, and it deserves to be read for what it asserts. The left-hand side is a purely analytic quantity, the strength of the pole of a meromorphic function built from the Euler product over prime ideals; by theorem 8.1.4 it encodes, factor by factor, the splitting behavior of every rational prime in K . The right-hand side is a single number assembled from the coarsest arithmetic invariants of K all at once: the class number h_K , which measures the failure of $\mathcal{O}_K$ to be a principal ideal domain; the regulator R_K , which measures the volume of the lattice of units; the order w_K of the group of roots of unity; and the discriminant d_K , which measures the ramification. That a single analytic limit should equal a product of these otherwise unrelated quantities is the substance of the formula, and it is the prototype of a pervasive phenomenon: the leading behavior of an L -function at a distinguished point is a regulator of arithmetic data. The conjectures of Birch and Swinnerton-Dyer, and more generally the Beilinson and Bloch–Kato conjectures, are the modern descendants of this single residue computation.

The factor κ_K specializes, at $K = \mathbb{Q}$, to the residue of the Riemann zeta function, and the formula is consistent with the special L -values computed analytically in chapter 7. The following examples verify each arithmetic instance.

Example 8.3.3: Four Residues from Their Invariants

Each computation evaluates the right-hand side of (8.17) from the invariants of K , taken from [17]. Throughout, the regulator of a field with unit rank $r_1 + r_2 - 1 = 0$ is the empty product $R_K = 1$,

and for a real quadratic field the regulator is $R_K = \log \varepsilon$, where $\varepsilon > 1$ is the fundamental unit.

The rational field $K = \mathbb{Q}$. Here $r_1 = 1$, $r_2 = 0$, $h_K = 1$, $R_K = 1$ (unit rank 0), $w_K = 2$ (the roots of unity are ± 1), and $d_K = 1$, so $|d_K| = 1$. The formula gives

$$\kappa_{\mathbb{Q}} = \frac{2^1(2\pi)^0 \cdot 1 \cdot 1}{2 \cdot \sqrt{1}} = \frac{2}{2} = 1$$

and $\zeta_{\mathbb{Q}} = \zeta$ is the Riemann zeta function, so this recovers the simple pole of residue 1 established in corollary 4.2.2. The class number formula contains the residue of ζ as its degenerate case.

The Gaussian field $K = \mathbb{Q}(i)$. The ring of integers is $\mathcal{O}_K = \mathbb{Z}[i]$, with $r_1 = 0$, $r_2 = 1$, and discriminant $d_K = -4$, so $|d_K| = 4$. The unit group is $\{\pm 1, \pm i\}$, the full group of fourth roots of unity, so $w_K = 4$ and the unit rank $r_1 + r_2 - 1 = 0$ gives $R_K = 1$. The class number is $h_K = 1$, since $\mathbb{Z}[i]$ is a principal ideal domain. Hence

$$\kappa_{\mathbb{Q}(i)} = \frac{2^0(2\pi)^1 \cdot 1 \cdot 1}{4 \cdot \sqrt{4}} = \frac{2\pi}{4 \cdot 2} = \frac{2\pi}{8} = \frac{\pi}{4}$$

This value admits an independent check. The Dedekind zeta function factors as $\zeta_{\mathbb{Q}(i)}(s) = \zeta(s) L(s, \chi_{-4})$, the product of the Riemann zeta function with the Dirichlet L -function of the nontrivial character χ_{-4} modulo 4 (the factorization is proved in section 8.4). Taking residues at $s = 1$, where $L(s, \chi_{-4})$ is holomorphic, gives

$$\text{Res}_{s=1} \zeta_{\mathbb{Q}(i)}(s) = \left(\text{Res}_{s=1} \zeta(s) \right) \cdot L(1, \chi_{-4}) = 1 \cdot L(1, \chi_{-4})$$

and $L(1, \chi_{-4}) = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots = \pi/4$ is the Leibniz series evaluated in chapter 7. The two routes agree: $\kappa_{\mathbb{Q}(i)} = \pi/4$.

The field $K = \mathbb{Q}(\sqrt{-5})$. The ring of integers is $\mathcal{O}_K = \mathbb{Z}[\sqrt{-5}]$, with $r_1 = 0$, $r_2 = 1$. Since $-5 \equiv 3 \pmod{4}$, the discriminant is $d_K = 4 \cdot (-5) = -20$, so $|d_K| = 20$ and $\sqrt{|d_K|} = 2\sqrt{5}$. The only units are ± 1 , so $w_K = 2$ and the unit rank 0 gives $R_K = 1$. The class number is $h_K = 2$: the ideal $(2, 1 + \sqrt{-5})$ is non-principal, and $\text{Cl}(K) \cong \mathbb{Z}/2\mathbb{Z}$. Hence

$$\kappa_{\mathbb{Q}(\sqrt{-5})} = \frac{2^0(2\pi)^1 \cdot 2 \cdot 1}{2 \cdot 2\sqrt{5}} = \frac{4\pi}{4\sqrt{5}} = \frac{\pi}{\sqrt{5}}$$

This is the first field in the list with $h_K > 1$, and the class number $h_K = 2$ enters the residue directly as a factor of 2. The failure of unique factorization in $\mathbb{Z}[\sqrt{-5}]$, the historical origin of ideal theory, is visible analytically as a doubling of the residue relative to the principal case.

The real quadratic field $K = \mathbb{Q}(\sqrt{2})$. Here $r_1 = 2$, $r_2 = 0$, and the unit rank is $r_1 + r_2 - 1 = 1$, so the regulator is nontrivial and records the fundamental unit. Since $2 \equiv 2 \pmod{4}$, the discriminant is $d_K = 4 \cdot 2 = 8$, so $|d_K| = 8$ and $\sqrt{|d_K|} = 2\sqrt{2}$. The roots of unity are ± 1 , so $w_K = 2$, and the class number is $h_K = 1$. The fundamental unit is $\varepsilon = 1 + \sqrt{2}$, of norm $N_{K/\mathbb{Q}}(1 + \sqrt{2}) = 1 - 2 = -1$, so $R_K = \log(1 + \sqrt{2})$. Hence

$$\kappa_{\mathbb{Q}(\sqrt{2})} = \frac{2^2(2\pi)^0 \cdot 1 \cdot \log(1 + \sqrt{2})}{2 \cdot 2\sqrt{2}} = \frac{4 \log(1 + \sqrt{2})}{4\sqrt{2}} = \frac{\log(1 + \sqrt{2})}{\sqrt{2}}$$

In contrast to the three preceding fields, where the regulator was the empty product 1, here the residue is transcendental in a new way: it is a logarithm of an algebraic number, the fundamental unit. The regulator factor R_K is the contribution that distinguishes the real case, where there are infinitely many units, from the imaginary case, where there are only finitely many.

The two imaginary-quadratic computations are not isolated curiosities. The factor $2^{r_1}(2\pi)^{r_2}$ in the residue is, for an imaginary quadratic field, equal to 2π , and after division by $w_K\sqrt{|d_K|}$ it produces exactly a rational multiple of $\pi/\sqrt{|d_K|}$, namely the special value $L(1, \chi_{d_K})$ of the quadratic Dirichlet L -function attached to the field. This is the structural reason behind the closed forms tabulated in chapter 7: the value $L(1, \chi)$ of a primitive quadratic character was computed there by analytic means (Gauss sums and the Fourier expansion of the sawtooth function), and the class number formula now reproduces the same numbers from pure arithmetic, the class number, the units, and the discriminant of the associated quadratic field. The two derivations are independent, and their agreement is a theorem rather than a coincidence.

8.3.4 Remark: The Numerical Coincidence Made a Theorem The agreement observed for $\mathbb{Q}(i)$, that the arithmetically computed residue $\pi/4$ coincides with the analytically computed Leibniz value $L(1, \chi_{-4})$, is an instance of a general identity. For any imaginary quadratic field $K = \mathbb{Q}(\sqrt{D})$ of fundamental discriminant $D = d_K < 0$, the residue formula (8.17) reads

$$\frac{2\pi h_K}{w_K\sqrt{|D|}} = \text{Res}_{s=1}\zeta_K(s) = L(1, \chi_D)$$

where the last equality uses the factorization $\zeta_K(s) = \zeta(s)L(s, \chi_D)$ together with $\text{Res}_{s=1}\zeta(s) = 1$ from corollary 4.2.2. Equating the outer terms gives Dirichlet's class number formula in its classical form, $L(1, \chi_D) = 2\pi h_K/(w_K\sqrt{|D|})$, which expresses an analytic L -value as a positive rational multiple of $\pi/\sqrt{|D|}$ with the class number as numerator. The positivity of h_K is, in this light, the structural reason $L(1, \chi_D) \neq 0$, an arithmetic counterpart to the analytic non-vanishing proved in theorem 7.4.2. The factorization $\zeta_K = \zeta \cdot L(\cdot, \chi_D)$ that underlies the whole bridge is established, for both signs of the discriminant and with the real-quadratic regulator in place of the $\pi/\sqrt{|D|}$ factor, in section 8.4; that section turns the numerical agreement recorded here into the general theorem.

8.4 Factoring ζ_K and the Algebraic Non-Vanishing of $L(1, \chi)$

The class number formula of section 8.3 extracts a single transcendental constant, the residue κ_K , from the pole of ζ_K at $s = 1$, and reads off from it all the basic invariants of K at once. For a general number field that residue is the end of the story: ζ_K is an irreducible analytic object, not built from anything simpler. For the abelian fields, however, ζ_K is not irreducible at all. When $K/\mathbb{Q}$ is Galois with abelian Galois group, the Euler product of ζ_K regroups, prime by prime, into a product of Dirichlet L -functions, one factor for each character of the Galois group. This section carries out the regrouping in the two cases where it can be made completely explicit by hand: the quadratic fields, where ζ_K splits as $\zeta(s)L(s, \chi_D)$ for a single quadratic character, and the cyclotomic fields, where ζ_K splits as the product of $L(s, \chi)$ over all the Dirichlet characters of conductor dividing the cyclotomic level.

The factorization is not a formal coincidence. Reading it from left to right, it expresses the Dedekind zeta function, with its known pole, as a product of L -functions; reading it from right to left, it expresses each L -function as a ratio of zeta functions whose poles and residues are already in hand. The latter direction is the more consequential. For a quadratic field the identity $L(s, \chi_D) = \zeta_K(s)/\zeta(s)$ exhibits $L(s, \chi_D)$ as a quotient of two functions each with a simple pole of known residue at $s = 1$, so the quotient is holomorphic and nonzero there, and its value is forced to be the positive number κ_K .

This is a second, purely algebraic, proof that $L(1, \chi) \neq 0$ for a real primitive character, the subtle case of theorem 7.4.2, where the conjugate-character argument of chapter 7 collapses and the proof had to fall back on Landau's theorem applied to the auxiliary product $\zeta(s)L(s, \chi)$. The geometry of numbers, packaged into the residue κ_K , replaces the positivity argument entirely, and along the way produces Dirichlet's original closed formulas for $L(1, \chi_D)$ in terms of the class number and the fundamental unit.

The quadratic case rests on the splitting law for primes in a quadratic field, which the algebraic prerequisites record in the form of the Kronecker symbol.

Theorem 8.4.1: Quadratic Factorization

Let $K = \mathbb{Q}(\sqrt{m})$ be a quadratic field with fundamental discriminant $D = d_K$, and let $\chi_D = \left(\frac{D}{\cdot}\right)$ be the associated Kronecker symbol, a real primitive Dirichlet character modulo $|D|$ ([17]). Then for $\text{Re}(s) > 1$,

$$\zeta_K(s) = \zeta(s) L(s, \chi_D) \quad (8.18)$$

Proof:

Both sides of (8.18) are given on $\text{Re}(s) > 1$ by absolutely convergent Euler products: the left by the product of ζ_K over the prime ideals of $\mathcal{O}_K$ (theorem 8.1.4), the right by the product of ζ over the rational primes (corollary 2.3.2) times the product of $L(s, \chi_D)$ over the rational primes (theorem 7.2.2). Every nonzero prime ideal $\mathfrak{p}$ of $\mathcal{O}_K$ contains a unique rational prime p (namely the positive generator of $\mathfrak{p} \cap \mathbb{Z}$, a prime because $\mathcal{O}_K/\mathfrak{p}$ is a field of characteristic p), so the prime ideals of $\mathcal{O}_K$ are partitioned into the finite blocks $\{\mathfrak{p} : \mathfrak{p} | p\mathcal{O}_K\}$ indexed by the rational primes p . Because the Euler product of ζ_K converges absolutely on $\text{Re}(s) > 1$, its factors may be reordered and regrouped freely, and grouping the prime ideals of $\mathcal{O}_K$ according to the rational prime p they lie above turns the product into a product over rational primes,

$$\zeta_K(s) = \prod_p \prod_{\mathfrak{p} | p} (1 - N\mathfrak{p}^{-s})^{-1}$$

the inner product running over the finitely many prime ideals $\mathfrak{p}$ of $\mathcal{O}_K$ dividing $p\mathcal{O}_K$. In a quadratic field there are at most two such primes, and the norm of each is $N\mathfrak{p} = p^f$ with f the residue degree, since $\mathcal{O}_K/\mathfrak{p}$ is the finite field of $N\mathfrak{p}$ elements containing $\mathbb{F}_p$ with degree f ([17]). The two products on the right of (8.18) likewise factor over p , with the p -th factor

$$(1 - p^{-s})^{-1} (1 - \chi_D(p)p^{-s})^{-1}$$

It therefore suffices to prove, for each rational prime p , the local identity

$$\prod_{\mathfrak{p} | p} (1 - N\mathfrak{p}^{-s})^{-1} = (1 - p^{-s})^{-1} (1 - \chi_D(p)p^{-s})^{-1} \quad (8.19)$$

since taking the product of (8.19) over all p then yields (8.18), the rearrangement of the doubly indexed product being legitimate by absolute convergence. The factorization of $p\mathcal{O}_K$ in a quadratic field falls into exactly three types, governed by the value $\chi_D(p) \in \{+1, -1, 0\}$ through the quadratic splitting law (the Dedekind–Kummer theorem specialized to a quadratic field,

equivalently the law of quadratic reciprocity, [17] and section 0.5). Each type is checked in turn.

Step 1: p splits, $\chi_D(p) = +1$. Here $p\mathcal{O}_K = \mathfrak{p}_1\mathfrak{p}_2$ with $\mathfrak{p}_1 \neq \mathfrak{p}_2$ two distinct primes, each of residue degree 1, so $N\mathfrak{p}_1 = N\mathfrak{p}_2 = p$. The left side of (8.19) is

$$(1 - p^{-s})^{-1}(1 - p^{-s})^{-1} = (1 - p^{-s})^{-2}$$

The right side is $(1 - p^{-s})^{-1}(1 - (+1)p^{-s})^{-1} = (1 - p^{-s})^{-2}$. The two agree.

Step 2: p is inert, $\chi_D(p) = -1$. Here $p\mathcal{O}_K = \mathfrak{p}$ is prime, of residue degree 2, so $N\mathfrak{p} = p^2$. The left side of (8.19) is the single factor

$$(1 - p^{-2s})^{-1} = (1 - p^{-s})^{-1}(1 + p^{-s})^{-1}$$

using the factorization $1 - X^2 = (1 - X)(1 + X)$ with $X = p^{-s}$. The right side is $(1 - p^{-s})^{-1}(1 - (-1)p^{-s})^{-1} = (1 - p^{-s})^{-1}(1 + p^{-s})^{-1}$. The two agree.

Step 3: p ramifies, $\chi_D(p) = 0$. Ramification occurs precisely at the primes dividing the discriminant, $p|D$, which is exactly where $\chi_D(p) = 0$ (the Kronecker symbol vanishes on the primes dividing its modulus $|D|$). Here $p\mathcal{O}_K = \mathfrak{p}^2$ with $\mathfrak{p}$ of residue degree 1, so $N\mathfrak{p} = p$, and the left side of (8.19) is the single factor

$$(1 - p^{-s})^{-1}$$

The right side is $(1 - p^{-s})^{-1}(1 - 0 \cdot p^{-s})^{-1} = (1 - p^{-s})^{-1} \cdot 1 = (1 - p^{-s})^{-1}$. The two agree.

In all three cases the local identity (8.19) holds, and these cases are exhaustive. Taking the product over all rational primes p gives (8.18) on $\text{Re}(s) > 1$.

The three cases of the proof line up the three factorization types of a rational prime in K with the three values of the quadratic character: a split prime ($\chi_D(p) = +1$) contributes two factors of norm p , an inert prime ($\chi_D(p) = -1$) one factor of norm p^2 , and a ramified prime ($\chi_D(p) = 0$) one factor of norm p , and in each case the resulting Euler factor of ζ_K is exactly the product of the Euler factor of ζ with the Euler factor of $L(s, \chi_D)$. The identity (8.18) is the analytic incarnation of the splitting law: the entire arithmetic of how rational primes decompose in K is encoded in the single equation $\zeta_K = \zeta \cdot L(\cdot, \chi_D)$, with the character χ_D reading off the decomposition type from the residue of p modulo $|D|$.

Comparing Dirichlet-series coefficients in (8.18) records the splitting law one more way, now in terms of the ideal-counting function of section 8.2. On $\text{Re}(s) > 1$ the left side is $\sum_{n \geq 1} a_K(n)n^{-s}$ with $a_K(n) = \#\{\mathfrak{a} : N\mathfrak{a} = n\}$ the number of integral ideals of norm n (definition 8.1.1), while the right side is the Dirichlet product of $\zeta(s) = \sum_n n^{-s}$ with $L(s, \chi_D) = \sum_n \chi_D(n)n^{-s}$, whose n -th coefficient is the convolution $\sum_{d|n} \chi_D(d)$. Equating coefficients gives, for every $n \geq 1$,

$$a_K(n) = \sum_{d|n} \chi_D(d) \tag{8.20}$$

a closed formula for the number of ideals of norm n in a quadratic field as a divisor sum of the Kronecker symbol. This is the exact quadratic instance of the counting that section 8.2 performed by the geometry of numbers: the multiplicative function $n \mapsto a_K(n)$, which there was summed over $n \leq x$ to extract the residue κ_K , is here identified term by term with $\sum_{d|n} \chi_D(d)$, and its average size is governed by $L(1, \chi_D)$. In particular $a_K(p) = 1 + \chi_D(p)$ equals 2, 0, or 1 according as p splits, is inert, or ramifies, recovering the prime-by-prime splitting law from the coefficient identity.

The factorization is consistent with the pole structure already established for both sides, and the consistency is worth recording, since it is the mechanism behind the non-vanishing theorem to come. On $\operatorname{Re}(s) > 1$ the right side of (8.18) is the product of ζ , which continues to a function with a simple pole at $s = 1$ (theorem 4.2.1), and $L(s, \chi_D)$, which continues to a function holomorphic at $s = 1$ (theorem 7.2.4, χ_D being nonprincipal). The product thus has at $s = 1$ a pole of order at most one, and exactly one provided $L(1, \chi_D) \neq 0$. The left side, ζ_K , was shown in section 8.3 to have a simple pole at $s = 1$, of residue $\kappa_K \neq 0$. Equating the two descriptions, the simple pole of ζ_K must come from the simple pole of the ζ factor, and the holomorphic factor $L(s, \chi_D)$ cannot vanish at $s = 1$, for a zero there would cancel the pole and leave ζ_K holomorphic at $s = 1$, contradicting the pole of residue κ_K . This is exactly the non-vanishing argument, made quantitative in the next theorem.

The factorization, combined with the analytic facts already established for ζ and ζ_K , yields the non-vanishing of $L(1, \chi_D)$ as an immediate consequence, with no positivity argument at all.

Theorem 8.4.2: Algebraic Non-Vanishing of $L(1, \chi_D)$

For every fundamental discriminant D , the value $L(1, \chi_D)$ is finite, nonzero, and positive; in fact

$$L(1, \chi_D) = \kappa_K = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{|d_K|}} \quad (8.21)$$

where $K = \mathbb{Q}(\sqrt{m})$ is the quadratic field of discriminant $D = d_K$ and $\kappa_K = \operatorname{Res}_{s=1} \zeta_K(s)$.

Proof:

By theorem 8.4.1, on the half-plane $\operatorname{Re}(s) > 1$ one has $\zeta_K(s) = \zeta(s) L(s, \chi_D)$, and since $\zeta(s) \neq 0$ there (corollary 4.1.2), the factorization may be solved for $L(s, \chi_D)$:

$$L(s, \chi_D) = \frac{\zeta_K(s)}{\zeta(s)} \quad (\operatorname{Re}(s) > 1) \quad (8.22)$$

Each side of (8.22) continues meromorphically to a neighbourhood of $s = 1$: the numerator ζ_K by theorem 8.3.2, where it is shown to have a simple pole at $s = 1$ with residue $\operatorname{Res}_{s=1} \zeta_K = \kappa_K$, and the denominator ζ by theorem 4.2.1, with a simple pole at $s = 1$ of residue $\operatorname{Res}_{s=1} \zeta = 1$ (corollary 4.2.2). Write the Laurent expansions at $s = 1$,

$$\zeta_K(s) = \frac{\kappa_K}{s-1} + O(1), \quad \zeta(s) = \frac{1}{s-1} + O(1)$$

Since χ_D is a nonprincipal Dirichlet character (it is primitive of conductor $|D| > 1$, hence nonprincipal), the L -function $L(s, \chi_D)$ is holomorphic on $\operatorname{Re}(s) > 0$ (theorem 7.2.4); in particular it is holomorphic at $s = 1$, so the quotient on the right of (8.22) must be holomorphic at $s = 1$ as well. Computing the limit of that quotient as $s \rightarrow 1$, the two simple poles cancel:

$$L(1, \chi_D) = \lim_{s \rightarrow 1} \frac{\zeta_K(s)}{\zeta(s)} = \lim_{s \rightarrow 1} \frac{\kappa_K/(s-1) + O(1)}{1/(s-1) + O(1)} = \lim_{s \rightarrow 1} \frac{\kappa_K + O(s-1)}{1 + O(s-1)} = \frac{\kappa_K}{1} = \kappa_K$$

where the second equality multiplies numerator and denominator by $s - 1$. This is (8.21). The residue κ_K is strictly positive, being a product and quotient of the strictly positive quantities 2^{r_1} , $(2\pi)^{r_2}$, $h_K \geq 1$, $R_K > 0$, $w_K \geq 1$, and $\sqrt{|d_K|} > 0$, so $L(1, \chi_D) = \kappa_K > 0$. In particular $L(1, \chi_D) \neq 0$.

This is a second proof, independent of the one in chapter 7, that $L(1, \chi) \neq 0$ for a real primitive Dirichlet character χ . Every real primitive Dirichlet character is the Kronecker symbol χ_D of a unique fundamental discriminant D ([17]), so the present theorem 8.4.2 covers exactly the real-character case of theorem 7.4.2, which was the subtle case there: the case in which the conjugate-character argument of chapter 7 collapses and the proof had to fall back on Landau's theorem applied to $\zeta(s)L(s, \chi)$. The two proofs locate the non-vanishing in different places. The earlier proof derives it from the positivity of the coefficients of a Dirichlet series, an analytic input; the present one derives it from the strict positivity of the geometric constant κ_K , an input from the geometry of numbers, by exhibiting $L(1, \chi_D)$ as the ratio of the residue of ζ_K to the residue of ζ at $s = 1$. The complex-character case of theorem 7.4.2 is not reached this way, and remains the province of the conjugate-pair argument; only the real characters arise as χ_D .

Specializing (8.21) to the two signatures of a quadratic field recovers Dirichlet's original class number formulas. The signature is dictated by the sign of D : a negative discriminant gives an imaginary quadratic field, with no real embeddings and one pair of complex embeddings, while a positive discriminant gives a real quadratic field, with two real embeddings and no complex ones.

Corollary 8.4.3: Imaginary Quadratic Class Number Formula

Let $D < 0$ be a fundamental discriminant and $K = \mathbb{Q}(\sqrt{D})$ the corresponding imaginary quadratic field. Then

$$L(1, \chi_D) = \frac{2\pi h_K}{w_K \sqrt{|D|}} \quad (8.23)$$

where h_K is the class number and w_K the number of roots of unity in K .

Proof:

An imaginary quadratic field has signature $r_1 = 0$, $r_2 = 1$, since the two embeddings $K \hookrightarrow \mathbb{C}$ are a single conjugate pair of complex embeddings. The unit rank is $r_1 + r_2 - 1 = 0$, so $\mathcal{O}_K^\times$ is finite (Dirichlet's unit theorem, [17]) and the regulator of a rank-zero unit lattice is $R_K = 1$ by the convention of section 8.3. Substituting $r_1 = 0$, $r_2 = 1$, $R_K = 1$, and $|d_K| = |D|$ into (8.21),

$$L(1, \chi_D) = \frac{2^0 (2\pi)^1 h_K \cdot 1}{w_K \sqrt{|D|}} = \frac{2\pi h_K}{w_K \sqrt{|D|}}$$

which is (8.23).

Corollary 8.4.4: Real Quadratic Class Number Formula

Let $D > 0$ be a fundamental discriminant and $K = \mathbb{Q}(\sqrt{D})$ the corresponding real quadratic field. Then

$$L(1, \chi_D) = \frac{2 h_K \log \varepsilon_K}{\sqrt{D}} \quad (8.24)$$

where h_K is the class number and $\varepsilon_K > 1$ is the fundamental unit of K .

Proof:

A real quadratic field has signature $r_1 = 2, r_2 = 0$, since both embeddings $K \hookrightarrow \mathbb{R}$ are real. The unit rank is $r_1 + r_2 - 1 = 1$, so $\mathcal{O}_K^\times \cong \mu_K \times \mathbb{Z}$ with $\mu_K = \{\pm 1\}$ the only roots of unity in a real field, whence $w_K = 2$; the unit lattice has rank 1 and is generated modulo torsion by the fundamental unit $\varepsilon_K > 1$, so the regulator is the single entry $R_K = \log \varepsilon_K$ ([17] and section 8.3). Substituting $r_1 = 2, r_2 = 0, R_K = \log \varepsilon_K, w_K = 2$, and $|d_K| = D$ into (8.21),

$$L(1, \chi_D) = \frac{2^2 (2\pi)^0 h_K \log \varepsilon_K}{2\sqrt{D}} = \frac{4 h_K \log \varepsilon_K}{2\sqrt{D}} = \frac{2 h_K \log \varepsilon_K}{\sqrt{D}}$$

which is (8.24).

The two formulas exhibit the dichotomy between imaginary and real quadratic fields in its sharpest form. For $D < 0$ the value $L(1, \chi_D)$ is a rational multiple of $\pi/\sqrt{|D|}$, the transcendence coming entirely from the archimedean factor 2π of the complex place, and the arithmetic content is the class number h_K (the root-of-unity count w_K equals 2 except for the two fields $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt{-3})$, where it is 4 and 6). For $D > 0$ the value is instead a rational multiple of $\log \varepsilon_K/\sqrt{D}$, the transcendence now residing in the regulator $\log \varepsilon_K$, and the arithmetic content is the product $h_K \log \varepsilon_K$ of the class number with the regulator. The same residue κ_K produces both, the signature deciding which invariants survive.

The imaginary formula (8.23) matches the closed forms computed by hand in chapter 7 for the smallest moduli. Taking $D = -4$, the field is $K = \mathbb{Q}(i)$, with class number $h_K = 1$ and $w_K = 4$ roots of unity (the four powers of i), so

$$L(1, \chi_{-4}) = \frac{2\pi \cdot 1}{4 \cdot \sqrt{4}} = \frac{2\pi}{4 \cdot 2} = \frac{\pi}{4}$$

recovering the value of the Leibniz series $1 - \frac{1}{3} + \frac{1}{5} - \dots$ found in chapter 7, the L -function of the unique nonprincipal character modulo 4. Taking $D = -20$, the field is $K = \mathbb{Q}(\sqrt{-5})$, the standard example of a field with non-trivial class group: here $h_K = 2$ and $w_K = 2$, and $|D| = 20$, so

$$L(1, \chi_{-20}) = \frac{2\pi \cdot 2}{2 \cdot \sqrt{20}} = \frac{2\pi}{\sqrt{20}} = \frac{2\pi}{2\sqrt{5}} = \frac{\pi}{\sqrt{5}}$$

The factor of 2 over the naive guess $\pi/\sqrt{20}$ is exactly the class number $h_K = 2$, the analytic manifestation of the failure of unique factorization in $\mathbb{Z}[\sqrt{-5}]$. Two further values exhibit the exceptional root-of-unity counts and the real case. For $D = -3$ the field is $K = \mathbb{Q}(\sqrt{-3})$, which contains the sixth roots of unity, so $w_K = 6$ and $h_K = 1$, giving

$$L(1, \chi_{-3}) = \frac{2\pi \cdot 1}{6\sqrt{3}} = \frac{\pi}{3\sqrt{3}}$$

the closed form found for the unique nonprincipal character modulo 3 in chapter 7. For $D = 5$ the field is the real quadratic $K = \mathbb{Q}(\sqrt{5})$, with $h_K = 1$ and fundamental unit $\varepsilon_K = (1 + \sqrt{5})/2$ the golden ratio, so by (8.24)

$$L(1, \chi_5) = \frac{2 \cdot 1 \cdot \log((1 + \sqrt{5})/2)}{\sqrt{5}}$$

again matching the closed form recorded for the quadratic character modulo 5 in chapter 7, where the value appeared as a logarithm of the fundamental unit with no explanation; the explanation is the real-quadratic class number formula. The class number formula thus turns a transcendental special value of an L -function into a count of ideal classes (and, for $D > 0$, a regulator), and conversely computes the special value once the class number and fundamental unit, both finite and algorithmically computable from D , are known.

The quadratic factorization is the degree-two case of a general principle: the Dedekind zeta function of an abelian extension of $\mathbb{Q}$ factors as a product of Dirichlet L -functions indexed by the characters of the Galois group. The cyclotomic fields realize the principle in its cleanest form, and they are exactly the prototype promised in chapter 7, where the product $\prod_{\chi} L(s, \chi)$ over the characters modulo m was studied as a stand-in for the Dedekind zeta function of $\mathbb{Q}(\zeta_m)$.

Proposition 8.4.5: Cyclotomic Factorization

Let $m \geq 1$ and let $K = \mathbb{Q}(\zeta_m)$ be the m -th cyclotomic field, with Galois group $\text{Gal}(\mathbb{Q}(\zeta_m)/\mathbb{Q}) \cong (\mathbb{Z}/m\mathbb{Z})^\times$. Then for $\text{Re}(s) > 1$,

$$\zeta_K(s) = \prod_{\chi} L(s, \chi^*) \tag{8.25}$$

the product running over the Dirichlet characters χ modulo m (equivalently over the characters of $(\mathbb{Z}/m\mathbb{Z})^\times$), where χ^* denotes the primitive character of conductor dividing m that induces χ .

Proof:

Both sides of (8.25) are absolutely convergent Euler products on $\text{Re}(s) > 1$, the left over the prime ideals of $\mathcal{O}_K$ (theorem 8.1.4) and the right over the rational primes for each of the $\varphi(m)$ characters (theorem 7.2.2). Grouping the prime ideals of $\mathcal{O}_K$ over the rational prime p below them, it suffices to prove for each rational prime p the local identity

$$\prod_{\mathfrak{p}|p} (1 - N\mathfrak{p}^{-s})^{-1} = \prod_{\chi} (1 - \chi^*(p)p^{-s})^{-1} \tag{8.26}$$

the product over χ ranging over the characters modulo m , and then take the product over all p . The proof of the unramified case is given in full; the ramified case is reduced to a precise citation.

Step 1: The splitting of an unramified prime. Let $p \nmid m$. Such a prime is unramified in $\mathbb{Q}(\zeta_m)$, the discriminant of $\mathbb{Q}(\zeta_m)$ being divisible only by the primes dividing m ([17]), so the ramification index is $e = 1$. The splitting is governed by the order of the Frobenius. Under the isomorphism $\text{Gal}(\mathbb{Q}(\zeta_m)/\mathbb{Q}) \cong (\mathbb{Z}/m\mathbb{Z})^\times$, which sends an automorphism σ to the residue a with $\sigma(\zeta_m) = \zeta_m^a$, the Frobenius automorphism at p (well defined up to conjugacy, but the group is abelian, so

it is a single element) corresponds to the residue of p itself, since Frob_p acts as the p -power map modulo any prime above p . The decomposition group at p is the cyclic group generated by Frob_p , of order equal to the order of p in $(\mathbb{Z}/m\mathbb{Z})^\times$, and this order is the common residue degree of the primes above p . Thus the residue degree of every prime $\mathfrak{p}$ above p is

$$f = \text{ord}_m(p)$$

the multiplicative order of p in $(\mathbb{Z}/m\mathbb{Z})^\times$, and the number of primes above p is

$$g = \frac{\varphi(m)}{f}$$

since $efg = \varphi(m) = [\mathbb{Q}(\zeta_m) : \mathbb{Q}]$ with ramification index $e = 1$ ([17]). Each of the g primes $\mathfrak{p}$ above p has $N\mathfrak{p} = p^f$, so the left side of (8.26) is

$$\prod_{\mathfrak{p}|p} (1 - N\mathfrak{p}^{-s})^{-1} = (1 - p^{-fs})^{-g} \quad (8.27)$$

a single factor raised to the power g .

Step 2: The character side, via the values $\chi(p)$. For $p \nmid m$ every character χ modulo m has $\chi^*(p) = \chi(p)$, since the inducing character χ^* and χ agree on the integers coprime to m and p is one of them; the conductor bookkeeping affects only the primes dividing m . The right side of (8.26) is therefore $\prod_{\chi} (1 - \chi(p)p^{-s})^{-1}$, and the task is to evaluate the polynomial

$$\prod_{\chi} (1 - \chi(p)X), \quad X = p^{-s}$$

Consider the evaluation homomorphism

$$\text{ev}_p: (\widehat{\mathbb{Z}/m\mathbb{Z}})^\times \longrightarrow \mathbb{C}^\times, \quad \chi \longmapsto \chi(p)$$

from the character group to the multiplicative group of $\mathbb{C}$. Its image is a finite subgroup of $\mathbb{C}^\times$, hence a group of roots of unity; the image is exactly the group μ_f of f -th roots of unity, because $\chi(p)$ is an f -th root of unity for every χ (as $\chi(p)^f = \chi(p^f) = \chi(1) = 1$, using $p^f \equiv 1 \pmod{m}$ from $f = \text{ord}_m(p)$), and conversely every f -th root of unity is attained (the cyclic group $\langle p \rangle \subseteq (\mathbb{Z}/m\mathbb{Z})^\times$ generated by p has order f , and the characters of $(\mathbb{Z}/m\mathbb{Z})^\times$ restrict onto all f characters of $\langle p \rangle$, which take p to each f -th root of unity). The kernel of ev_p has order

$$|\ker \text{ev}_p| = \frac{|(\widehat{\mathbb{Z}/m\mathbb{Z}})^\times|}{|\mu_f|} = \frac{\varphi(m)}{f} = g$$

using $|(\widehat{\mathbb{Z}/m\mathbb{Z}})^\times| = \varphi(m)$ (theorem 7.1.2). By the orbit count for the surjection ev_p , every fibre has the same size g , so as χ ranges over all $\varphi(m)$ characters the value $\chi(p)$ runs through each f -th root of unity exactly g times.

Step 3: The Euler-factor identity for $p \nmid m$. Grouping the $\varphi(m)$ linear factors $1 - \chi(p)X$ by the common value $\chi(p) = \omega$, each f -th root of unity ω occurring g times,

$$\prod_{\chi} (1 - \chi(p)X) = \prod_{\omega \in \mu_f} (1 - \omega X)^g = \left(\prod_{\omega \in \mu_f} (1 - \omega X) \right)^g = (1 - X^f)^g$$

the last equality being the factorization of $1 - X^f = \prod_{\omega \in \mu_f} (1 - \omega X)$ into linear factors over the f -th roots of unity (the roots of $1 - X^f$ are exactly the inverses ω^{-1} , which range over μ_f as ω does). Taking reciprocals and substituting $X = p^{-s}$,

$$\prod_{\chi} (1 - \chi(p)p^{-s})^{-1} = (1 - p^{-fs})^{-g} \quad (8.28)$$

Comparing (8.28) with (8.27), both sides of the local identity (8.26) equal $(1 - p^{-fs})^{-g}$, so (8.26) holds for every unramified prime $p \nmid m$.

Step 4: The ramified primes and the conductor bookkeeping. For the finitely many primes $p \mid m$, the prime p ramifies in $\mathbb{Q}(\zeta_m)$, the equality $\chi^*(p) = \chi(p)$ can fail (an imprimitive χ has $\chi(p) = 0$ while its primitive inducing character χ^* , of conductor prime to p , may have $\chi^*(p) \neq 0$), and the matching of the local Euler factor of ζ_K at p with the product $\prod_{\chi} (1 - \chi^*(p)p^{-s})^{-1}$ requires tracking the conductors of the inducing characters together with the ramification of p . This local computation, and the verification that the primitive characters χ^* supplying the factors of (8.25) are exactly the primitive Dirichlet characters of conductor dividing m , are carried out in full in [14, Ch. 3]; the same factorization is established from the conductor-discriminant formula in [5, Ch. VII, (10.16)]. Granting the ramified local identity from these sources and combining it with the unramified identity (8.26) proved in Steps 1–3, the product over all rational primes p yields (8.25) on $\text{Re}(s) > 1$.

A small case makes the unramified mechanism concrete. Take $m = 5$, so $K = \mathbb{Q}(\zeta_5)$ has degree $\varphi(5) = 4$ and Galois group $(\mathbb{Z}/5\mathbb{Z})^\times$, cyclic of order 4 generated by 2; the four Dirichlet characters modulo 5 are $\chi_0, \psi, \psi^2, \psi^3$ in the notation of chapter 7, all primitive except the principal χ_0 (which has conductor 1), and (8.25) reads $\zeta_K(s) = \zeta(s)L(s, \psi)L(s, \psi^2)L(s, \psi^3)$. Consider three unramified primes. For $p = 11 \equiv 1 \pmod{5}$ the order is $f = \text{ord}_5(11) = 1$, so $g = 4$ and 11 splits into four primes of norm 11; on the character side every $\chi(11) = \chi(1) = 1$, so $\prod_{\chi} (1 - \chi(11)X) = (1 - X)^4$, matching $(1 - 11^{-s})^{-4}$. For $p = 2$, a generator, the order is $f = \text{ord}_5(2) = 4$, so $g = 1$ and 2 is inert with a single prime of norm $2^4 = 16$; the values $\chi(2)$ run once through each fourth root of unity $\{1, i, -1, -i\}$, so $\prod_{\chi} (1 - \chi(2)X) = \prod_{\omega^4=1} (1 - \omega X) = 1 - X^4$, matching $(1 - 2^{-4s})^{-1}$. For $p = 19 \equiv 4 \pmod{5}$ the order is $f = \text{ord}_5(19) = \text{ord}_5(4) = 2$ (since $4^2 = 16 \equiv 1$), so $g = 2$ and 19 splits into two primes of norm 19^2 ; the values $\chi(19)$ run twice through the square roots of unity $\{1, -1\}$, so $\prod_{\chi} (1 - \chi(19)X) = ((1 - X)(1 + X))^2 = (1 - X^2)^2$, matching $(1 - 19^{-2s})^{-2}$. In each case $efg = 4 = [\mathbb{Q}(\zeta_5) : \mathbb{Q}]$ and the local factors agree, as Steps 1–3 require.

The decomposition is again consistent with the pole at $s = 1$, and now the consistency runs the other way, from the non-vanishing to the residue. Among the $\varphi(m)$ factors of (8.25) exactly one comes from the trivial character: the primitive character inducing χ_0 modulo m is the trivial character modulo 1, whose L -function is ζ itself, contributing the lone simple pole at $s = 1$. Every other factor $L(s, \chi^*)$ is attached to a nontrivial primitive character, hence is holomorphic on $\text{Re}(s) > 0$ (theorem 7.2.4) and, by the non-vanishing $L(1, \chi^*) \neq 0$ (theorem 7.4.2), is also nonzero at $s = 1$. The product (8.25) therefore has a simple pole at $s = 1$, and taking residues,

$$\kappa_K = \text{Res}_{s=1} \zeta_K(s) = \left(\text{Res}_{s=1} \zeta(s) \right) \prod_{\chi^* \neq 1} L(1, \chi^*) = \prod_{\chi^* \neq 1} L(1, \chi^*)$$

the residue of ζ being 1 (corollary 4.2.2). For the cyclotomic field the class number formula thus equates the geometric residue $\kappa_K = 2^{r_1} (2\pi)^{r_2} h_K R_K / (w_K \sqrt{|d_K|})$ with a product of special values

$\prod_{\chi^* \neq 1} L(1, \chi^*)$ of Dirichlet L -functions; this is the analytic identity from which Dirichlet's class number formulas for cyclotomic fields, and the decomposition of the regulator into a determinant of logarithms of cyclotomic units, are derived in [14, Ch. 3 and Ch. 4].

The cyclotomic factorization (8.25) fulfills the forward reference of chapter 7, where the product $\prod_{\chi \bmod m} L(s, \chi)$ was introduced as the cyclotomic prototype and used, through the non-negativity of its coefficients, to drive the non-vanishing of $L(1, \chi)$. That product is, up to the conductor bookkeeping for the imprimitive characters at the ramified primes, the Dedekind zeta function $\zeta_{\mathbb{Q}(\zeta_m)}$, and the unramified Euler-factor identity proved above, $\prod_{\chi} (1 - \chi(p)X) = (1 - X^f)^g$ with $f = \text{ord}_m(p)$ and $g = \varphi(m)/f$, is precisely the identity that gave the prototype its Euler product $\prod_{p \nmid m} (1 - p^{-fs})^{-g}$ in chapter 7. The factorization places that prototype in its proper context: it is the Galois-theoretic decomposition of an abelian zeta function, the first instance of a pattern that, for a general abelian extension $K/\mathbb{Q}$, writes ζ_K as the product of the Dirichlet L -functions attached to the characters of $\text{Gal}(K/\mathbb{Q})$ (viewed as Dirichlet characters via the Artin reciprocity map of [9]), and that, for a nonabelian extension, becomes the factorization of ζ_K into Artin L -functions. In every such factorization the pole of ζ_K at $s = 1$, with its residue κ_K , is carried by the single trivial character, whose L -function is ζ , while every nontrivial factor is holomorphic and nonzero at $s = 1$; the non-vanishing of those factors is what confines the pole to one place and, read as the residue identity above, expresses the arithmetic constant κ_K as a product of the boundary values $L(1, \chi)$. This is the structural reason the non-vanishing of $L(1, \chi)$ and the analytic class number formula are two faces of the same fact, and it is the entry point, pursued for nonabelian G through the Chebotarev density theorem and Artin L -functions, to the general analytic theory of L -functions of number fields.

8.5 The Functional Equation for $\zeta_K(s)$

The analytic class number formula of section 8.3 extracted the residue of ζ_K at its single pole $s = 1$; it left untouched the behavior of ζ_K to the left of the line $\text{Re}(s) = 1$, where the defining series and Euler product of section 8.1 no longer converge. For $K = \mathbb{Q}$ the corresponding gap was closed in chapter 6 by the functional equation, which continued ζ to the whole plane and exhibited the reflection $s \leftrightarrow 1 - s$ as the governing symmetry. The same symmetry holds for every number field. Completing ζ_K by the archimedean factors attached to the real and complex places of K , and by a power of $|d_K|$ playing the role of the conductor, produces a function Λ_K invariant under $s \mapsto 1 - s$. This is *Hecke's functional equation*, the subject of the present section.

The general theorem is stated and its proof, a lattice-theoretic generalization of Riemann's theta argument, is deferred to the standard references by a precise citation. For abelian K the functional equation is recovered in full from the material already developed: the factorization $\zeta_K = \prod_{\chi} L(s, \chi)$ of section 8.4 reduces it to the functional equations of the Dirichlet L -functions established in chapter 7, and the reduction turns the bookkeeping of archimedean factors and conductors into two clean numerical matchings: the Legendre duplication formula pairs the gamma factors, and the conductor-discriminant formula multiplies the conductors to $|d_K|$.

Definition 8.5.1: The Completed Dedekind Zeta Function

Let K be a number field of degree $d = r_1 + 2r_2$, with r_1 real places and r_2 complex places, discriminant d_K , and Dedekind zeta function ζ_K of definition 8.1.1. Write the *archimedean factors*

$$\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right), \quad \Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$$

with Γ the gamma function of definition 6.1.1. The *completed Dedekind zeta function* of K is

$$\Lambda_K(s) = |d_K|^{s/2} \Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} \zeta_K(s) \quad (8.29)$$

defined initially on the half-plane $\operatorname{Re}(s) > 1$, where $\zeta_K(s)$ is given by its absolutely convergent Dirichlet series and each archimedean factor is holomorphic.

The factor $\Gamma_{\mathbb{R}}(s)$ is attached to each real place of K and is identical to the local factor at the real place of $\mathbb{Q}$ from definition 6.3.1; the factor $\Gamma_{\mathbb{C}}(s)$ is attached to each complex place, that is, to each conjugate pair of complex embeddings, and is the local factor at a complex place. The two are linked by the Legendre duplication formula: as verified in (8.35) below,

$$\Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s+1) = \Gamma_{\mathbb{C}}(s) \quad (8.30)$$

so that a complex place behaves, archimedean-factor-wise, as one real place at s together with one real place at $s+1$. The power $|d_K|^{s/2}$ is the conductor-type factor: it generalizes the trivial conductor 1 of $\mathbb{Q}$, whose discriminant is $d_{\mathbb{Q}} = 1$, and its presence is what makes the completed function symmetric rather than merely meromorphic.

8.5.2 Remark: Reduction to the Riemann case For $K = \mathbb{Q}$ one has $r_1 = 1$, $r_2 = 0$, $d_K = 1$, and $\zeta_K = \zeta$ (definition 8.1.1). The definition (8.29) then reads

$$\Lambda_{\mathbb{Q}}(s) = 1^{s/2} \Gamma_{\mathbb{R}}(s)^1 \Gamma_{\mathbb{C}}(s)^0 \zeta(s) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s) = \Lambda(s)$$

the completed zeta function of definition 6.3.1. Thus Λ_K is the number-field generalization of Λ , with the single real factor of $\mathbb{Q}$ replaced by r_1 real and r_2 complex factors and the trivial conductor replaced by $|d_K|^{1/2}$.

The functional equation for Λ_K is the following theorem of Hecke. Its proof generalizes the four-stage Riemann argument of theorem 6.3.3 from the lattice $\mathbb{Z} \subset \mathbb{R}$ to the Minkowski lattice $\iota(\mathcal{O}_K) \subset \mathbb{R}^d$, and is given in full in the references cited; the statement is recorded here, and the abelian case is proved independently in theorem 8.5.5.

Theorem 8.5.3: Hecke's Functional Equation

Let K be a number field and Λ_K its completed Dedekind zeta function (definition 8.5.1). Then Λ_K extends to a meromorphic function on all of $\mathbb{C}$, holomorphic everywhere except for simple poles at $s = 0$ and $s = 1$, and satisfies the functional equation

$$\Lambda_K(s) = \Lambda_K(1 - s) \quad (8.31)$$

for all $s \in \mathbb{C}$. Equivalently, ζ_K continues to a meromorphic function on $\mathbb{C}$, holomorphic except for a simple pole at $s = 1$, with $\zeta_K(s)$ recovered from $\Lambda_K(s)$ by dividing out the (nowhere-vanishing) factor $|d_K|^{s/2} \Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2}$.

Proof:

The full proof is Hecke's, generalizing the theta-and-Mellin argument of theorem 6.3.3. One forms the theta series $\Theta(t) = \sum_{x \in \iota(\mathcal{O}_K)} e^{-\pi t \langle x, x \rangle}$ of the Minkowski lattice $\iota(\mathcal{O}_K) \subset \mathbb{R}^d$ (more precisely, a sum over the lattice translates within each ideal class, weighted to build ζ_K), obtains its transformation law under $t \mapsto 1/t$ from the Poisson summation formula (theorem 6.2.1) applied over the lattice, the covolume $\text{vol}(\mathbb{R}^d / \iota(\mathcal{O}_K)) = 2^{-r_2} \sqrt{|d_K|}$ from [17] entering as the multiplicative constant in the dual-lattice term, and then takes a Mellin transform in t to recover $\Lambda_K(s)$. The self-duality of the theta transformation law produces the symmetry $s \leftrightarrow 1 - s$, and the constant term of the theta series produces the two simple poles, exactly as the constant term of ψ produced the poles of Λ in theorem 6.3.3. The complete argument, including the unit-group integration that handles the $r_1 + r_2 - 1$ regulator directions of [17], is carried out in [5, Ch. VII, §5–§6] and in [3, Ch. XIII–XIV]. The idelic reformulation, in which the functional equation becomes the Poisson summation formula on the adeles and the archimedean factors become local zeta integrals (Tate's thesis), is deferred to [9].

The completed Dedekind zeta function is symmetric with no phase: the constant relating $\Lambda_K(s)$ and $\Lambda_K(1 - s)$ is $+1$. In the language of theorem 7.3.4, the *root number* of ζ_K is $+1$, reflecting that ζ_K is self-dual (the trivial Hecke character is its own conjugate). This is the number-field counterpart of the bare reflection $\Lambda(s) = \Lambda(1 - s)$ for $\mathbb{Q}$, in contrast to the twisted reflection $\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1 - s, \bar{\chi})$ carrying a unit-modulus phase $\epsilon(\chi)$ for a nonprincipal Dirichlet character.

The conductor factor $|d_K|^{s/2}$ admits, for abelian K , an arithmetic decomposition matching the factorization of ζ_K into Dirichlet L -functions. This is the conductor-discriminant formula, which the next theorem records and verifies on the two families of section 8.4, deferring the general proof.

Theorem 8.5.4: Conductor-Discriminant Formula

Let $K/\mathbb{Q}$ be a finite abelian extension, and let X be the associated group of primitive Dirichlet characters, so that

$$\zeta_K(s) = \prod_{\chi \in X} L(s, \chi) \quad (8.32)$$

as in theorem 8.4.1 (quadratic case) and proposition 8.4.5 (cyclotomic case). Then the absolute discriminant of K is the product of the conductors of these characters:

$$|d_K| = \prod_{\chi \in X} f_\chi \quad (8.33)$$

where f_χ is the conductor of χ (definition 7.1.5).

Proof:

The identity (8.33) is verified below on the quadratic and cyclotomic families, which are the cases used in this chapter. The general statement, valid for every finite abelian $K/\mathbb{Q}$, is a consequence of the behavior of the discriminant under the conductor-ramification correspondence of class field theory; the complete proof is [5, Ch. VII, (11.9)], and the idelic account is in [9].

Verification, quadratic case. Let $K = \mathbb{Q}(\sqrt{D})$ for a fundamental discriminant D , so $d_K = D$ and $|d_K| = |D|$. By theorem 8.4.1 the character group is $X = \{\chi_0, \chi_D\}$, where χ_0 is the principal character (conductor $f_{\chi_0} = 1$, induced from level 1, by definition 7.1.5) and χ_D is the Kronecker symbol of D , a real primitive character modulo $|D|$ (so $f_{\chi_D} = |D|$, primitivity being recorded in section 8.4 from [17]). The product of conductors is

$$\prod_{\chi \in X} f_\chi = f_{\chi_0} f_{\chi_D} = 1 \cdot |D| = |D| = |d_K|$$

confirming (8.33).

Verification, cyclotomic case. Let $K = \mathbb{Q}(\zeta_p)$ for an odd prime p , of degree $\varphi(p) = p - 1$. By proposition 8.4.5 the character group X consists of the $p - 1$ Dirichlet characters modulo p . Of these, the principal character χ_0 has conductor 1. Each of the remaining $p - 2$ nonprincipal characters χ modulo p has conductor p : the only divisors of the prime p are 1 and p , the sole character modulo 1 is principal and cannot induce a nonprincipal χ , so no nonprincipal character modulo p is induced from a proper divisor, hence $f_\chi = p$ for each. The product of conductors is therefore

$$\prod_{\chi \in X} f_\chi = 1 \cdot \underbrace{p \cdots p}_{p-2 \text{ factors}} = p^{p-2}$$

On the other side, the discriminant of $\mathbb{Q}(\zeta_p)$ is $d_K = (-1)^{(p-1)/2} p^{p-2}$, so $|d_K| = p^{p-2}$ (the cyclotomic discriminant from [17]; see also [14, Ch. 2]). The two agree, confirming (8.33) in the cyclotomic case. For instance $\mathbb{Q}(\zeta_3) = \mathbb{Q}(\sqrt{-3})$ has $|d_K| = 3^1 = 3$, and $\mathbb{Q}(\zeta_5)$ has $|d_K| = 5^3 = 125$.

For abelian K the functional equation of theorem 8.5.3 need not invoke Hecke's general theorem. The factorization (8.32) expresses ζ_K through Dirichlet L -functions, each of which already carries

the functional equation of theorem 7.3.4, and assembling those reflections reconstitutes the reflection for Λ_K . The assembly rests on three matchings, which the next theorem makes precise and verifies: the gamma factors pair by the duplication formula (8.30), the character parities count the real and complex places, and the conductors multiply to $|d_K|$ by theorem 8.5.4.

Theorem 8.5.5: Abelian Consistency of the Functional Equation

Let $K/\mathbb{Q}$ be a finite abelian extension with associated group X of primitive Dirichlet characters, so that $\zeta_K(s) = \prod_{\chi \in X} L(s, \chi)$. Then the functional equation

$$\Lambda_K(s) = \Lambda_K(1-s)$$

holds, and follows from the functional equations of the Dirichlet L -functions $L(s, \chi)$, $\chi \in X$ (theorem 7.3.4), together with the conductor-discriminant formula (theorem 8.5.4), without recourse to Hecke's general theorem.

Proof:

The proof matches the completed object Λ_K against the product of the completed L -functions $\Lambda(s, \chi)$ of definition 7.3.3, then transports the reflection of each $\Lambda(s, \chi)$ through that product. The argument runs in four steps: the gamma-factor matching by duplication; the place-count matching of parities to r_1 and r_2 ; the conductor matching by theorem 8.5.4; and the assembly of the reflection with root number $+1$.

Step 1: The duplication identity. The completed L -function of a primitive character χ of conductor $m = f_\chi$ and parity $a \in \{0, 1\}$ is, by definition 7.3.3,

$$\Lambda(s, \chi) = \left(\frac{m}{\pi}\right)^{(s+a)/2} \Gamma\left(\frac{s+a}{2}\right) L(s, \chi) \quad (8.34)$$

For an even character ($a = 0$) the archimedean part of (8.34) is $(m/\pi)^{s/2} \Gamma(s/2) = m^{s/2} \Gamma_{\mathbb{R}}(s)$, and for an odd character ($a = 1$) it is $(m/\pi)^{(s+1)/2} \Gamma((s+1)/2) = m^{(s+1)/2} \Gamma_{\mathbb{R}}(s+1)$, in the notation of definition 8.5.1. The two real gamma factors combine into a complex gamma factor through the Legendre duplication formula (6.6). Indeed, applying (6.6) at argument $s/2$ gives $\Gamma(s/2) \Gamma((s+1)/2) = 2^{1-s} \sqrt{\pi} \Gamma(s)$, whence

$$\Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s+1) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \pi^{-(s+1)/2} \Gamma\left(\frac{s+1}{2}\right) = \pi^{-s-1/2} \cdot 2^{1-s} \sqrt{\pi} \Gamma(s) = 2^{1-s} \pi^{-s} \Gamma(s) = 2 (2\pi)^{-s} \Gamma(s) = \Gamma_{\mathbb{C}}(s) \quad (8.35)$$

which is the identity (8.30): one even plus one odd real gamma factor assemble into exactly one complex gamma factor.

Step 2: Parities count the archimedean places. The group X is closed under complex conjugation $\chi \mapsto \bar{\chi}$, since $K/\mathbb{Q}$ is Galois and X is the full group of primitive Dirichlet characters cut out by K , isomorphic to $\text{Gal}(K/\mathbb{Q})$ by the abelian reciprocity underlying theorem 8.4.1 and proposition 8.4.5; conjugation preserves the conductor, and $\bar{\chi}(-1) = \chi(-1) = \chi(-1)$ (the value lies in $\{\pm 1\}$), so it preserves parity as well. The count of even and odd characters is fixed by a structural dichotomy special to abelian fields. Complex conjugation is an element σ_∞ of the abelian group $\text{Gal}(K/\mathbb{Q})$, and it corresponds, under the identification of X with $\text{Gal}(K/\mathbb{Q})$, to the residue class of -1 , so that $\chi(\sigma_\infty) = \chi(-1)$ for every $\chi \in X$. Because σ_∞ is central

(the group is abelian), it acts in the same way on every archimedean embedding of K : either $\sigma_\infty = 1$, and K is totally real, with $r_2 = 0$ and $r_1 = [K : \mathbb{Q}]$; or σ_∞ has order 2, and K is totally imaginary, with $r_1 = 0$ and $[K : \mathbb{Q}] = 2r_2$. (No abelian K has both $r_1 > 0$ and $r_2 > 0$, since a central involution cannot fix one embedding while moving another.) The even characters are those trivial on σ_∞ , hence the characters of the quotient $\text{Gal}(K/\mathbb{Q})/\langle\sigma_\infty\rangle = \text{Gal}(K^+/\mathbb{Q})$, where K^+ is the maximal totally real subfield (the fixed field of σ_∞); thus the number of even characters is $[K^+ : \mathbb{Q}]$. In the totally real case $K^+ = K$, giving $[K^+ : \mathbb{Q}] = r_1 = r_1 + r_2$ even characters and no odd characters; in the totally imaginary case $[K^+ : \mathbb{Q}] = [K : \mathbb{Q}]/2 = r_2 = r_1 + r_2$ even characters, the other r_2 characters being odd. Either way,

$$\#\{\chi \in X : \chi(-1) = +1\} = r_1 + r_2, \quad \#\{\chi \in X : \chi(-1) = -1\} = r_2 \quad (8.36)$$

the even characters numbering $r_1 + r_2$ and the odd characters numbering r_2 ; this is the place decomposition recorded for abelian K in [5, Ch. VII, §5]. The two cases of the chapter make (8.36) explicit: for real quadratic $K = \mathbb{Q}(\sqrt{D})$, $D > 0$, both χ_0 and χ_D are even (as $\chi_D(-1) = +1$ for $D > 0$), giving $r_1 + r_2 = 2$ even and $r_2 = 0$ odd, consistent with $r_1 = 2$, $r_2 = 0$; for imaginary quadratic $K = \mathbb{Q}(\sqrt{D})$, $D < 0$, χ_0 is even and χ_D is odd (as $\chi_D(-1) = -1$ for $D < 0$), giving $r_1 + r_2 = 1$ even and $r_2 = 1$ odd, consistent with $r_1 = 0$, $r_2 = 1$.

Step 3: The product of completed L -functions. Form the product of (8.34) over all $\chi \in X$, separating its archimedean part from the L -functions. Using $(m/\pi)^{s/2}\Gamma(s/2) = m^{s/2}\Gamma_{\mathbb{R}}(s)$ for an even character and $(m/\pi)^{(s+1)/2}\Gamma((s+1)/2) = m^{(s+1)/2}\Gamma_{\mathbb{R}}(s+1)$ for an odd one (Step 1), and writing $m = \mathfrak{f}_\chi$, the archimedean parts multiply to

$$\prod_{\chi \in X} \left(\frac{\mathfrak{f}_\chi}{\pi} \right)^{(s+a_\chi)/2} \Gamma\left(\frac{s+a_\chi}{2}\right) = \left(\prod_{\chi \text{ even}} \mathfrak{f}_\chi^{s/2} \right) \left(\prod_{\chi \text{ odd}} \mathfrak{f}_\chi^{(s+1)/2} \right) \Gamma_{\mathbb{R}}(s)^{\#\{\text{even}\}} \Gamma_{\mathbb{R}}(s+1)^{\#\{\text{odd}\}} \quad (8.37)$$

By (8.36) there are $r_1 + r_2$ even characters and r_2 odd characters, so the gamma part of (8.37) is $\Gamma_{\mathbb{R}}(s)^{r_1+r_2} \Gamma_{\mathbb{R}}(s+1)^{r_2}$. Splitting off r_2 of the $\Gamma_{\mathbb{R}}(s)$ factors and pairing each with a $\Gamma_{\mathbb{R}}(s+1)$ through the duplication identity (8.35) converts r_2 of the products $\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1)$ into $\Gamma_{\mathbb{C}}(s)$, leaving r_1 copies of $\Gamma_{\mathbb{R}}(s)$:

$$\Gamma_{\mathbb{R}}(s)^{r_1+r_2} \Gamma_{\mathbb{R}}(s+1)^{r_2} = \Gamma_{\mathbb{R}}(s)^{r_1} (\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1))^{r_2} = \Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} \quad (8.38)$$

This is the gamma matching: each of the r_2 complex places of K arises as one even gamma factor paired with one odd gamma factor, exactly as $\Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{R}}(s+1) = \Gamma_{\mathbb{C}}(s)$ predicts, while the r_1 remaining even factors are the r_1 real places. The conductor part of (8.37) separates the exponent $s/2$ shared by all characters from the extra $1/2$ carried by each odd character:

$$\left(\prod_{\chi \text{ even}} \mathfrak{f}_\chi^{s/2} \right) \left(\prod_{\chi \text{ odd}} \mathfrak{f}_\chi^{(s+1)/2} \right) = \left(\prod_{\chi \in X} \mathfrak{f}_\chi \right)^{s/2} \cdot \prod_{\chi \text{ odd}} \mathfrak{f}_\chi^{1/2} = |d_K|^{s/2} \cdot c_K \quad (8.39)$$

where the conductor product $\prod_{\chi \in X} \mathfrak{f}_\chi = |d_K|$ is the conductor-discriminant formula (8.33), supplying the factor $|d_K|^{s/2}$, and

$$c_K = \prod_{\chi \text{ odd}} \mathfrak{f}_\chi^{1/2} > 0$$

is a positive constant independent of s . Combining the gamma matching (8.38), the conductor matching (8.39), and the factorization $\prod_{\chi \in X} L(s, \chi) = \zeta_K(s)$ of (8.32), the product of completed L -functions is

$$\prod_{\chi \in X} \Lambda(s, \chi) = c_K |d_K|^{s/2} \Gamma_{\mathbb{R}}(s)^{r_1} \Gamma_{\mathbb{C}}(s)^{r_2} \zeta_K(s) = c_K \Lambda_K(s) \quad (8.40)$$

by the definition (8.29) of Λ_K : the product of completed L -functions equals Λ_K up to the positive constant c_K . As a check, for imaginary quadratic $K = \mathbb{Q}(\sqrt{D})$ the sole odd character is χ_D of conductor $|D|$, so $c_K = \sqrt{|D|}$, while for real quadratic K there are no odd characters and $c_K = 1$.

Step 4: Reflection and root number. Each factor on the left of (8.40) satisfies a functional equation. For the principal character $\chi_0 \in X$, of conductor 1, the completed function is $\Lambda(s, \chi_0) = \Lambda(s)$ (definition 7.3.3) and the reflection is $\Lambda(s) = \Lambda(1-s)$ of theorem 6.3.3. For each nonprincipal $\chi \in X$, theorem 7.3.4 gives

$$\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1-s, \bar{\chi}), \quad \epsilon(\chi) = \frac{\tau(\chi)}{i^{a_\chi} \sqrt{f_\chi}}, \quad |\epsilon(\chi)| = 1 \quad (8.41)$$

Multiply (8.41) over all nonprincipal $\chi \in X$ and append the relation for χ_0 . Because X is closed under conjugation (Step 2), the assignment $\chi \mapsto \bar{\chi}$ permutes X , so on the right

$$\prod_{\chi \in X} \Lambda(1-s, \bar{\chi}) = \prod_{\chi \in X} \Lambda(1-s, \chi)$$

the product being merely reindexed. Therefore, writing $W = \prod_{\chi \in X} \epsilon(\chi)$ (with $\epsilon(\chi_0) = 1$),

$$\prod_{\chi \in X} \Lambda(s, \chi) = W \prod_{\chi \in X} \Lambda(1-s, \chi) \quad (8.42)$$

It remains to show $W = +1$. Partition X into characters fixed by conjugation and genuinely conjugate pairs. The principal character χ_0 is fixed by conjugation and has $\epsilon(\chi_0) = 1$, established above from the reflection $\Lambda(s) = \Lambda(1-s)$. A *nonprincipal* character fixed by conjugation, $\chi = \bar{\chi}$, is real, hence quadratic, and for such a character the root number is $\epsilon(\chi) = +1$. This is the sign theorem for the quadratic Gauss sum: for a real primitive character χ of parity a_χ the Gauss sum is exactly $\tau(\chi) = i^{a_\chi} \sqrt{f_\chi}$ (that is $\sqrt{f_\chi}$ for χ even and $i\sqrt{f_\chi}$ for χ odd), so $\epsilon(\chi) = \tau(\chi)/(i^{a_\chi} \sqrt{f_\chi}) = +1$. The modulus $|\tau(\chi)| = \sqrt{f_\chi}$ and the relation $\tau(\bar{\chi}) = \chi(-1)\overline{\tau(\chi)}$ of theorem 7.3.2 reduce this to determining a single sign (they give $\tau(\chi) \in \{\pm\sqrt{f_\chi}\}$ for χ even and $\tau(\chi) \in \{\pm i\sqrt{f_\chi}\}$ for χ odd); that the sign is $+$, the content of Gauss's theorem on the sign of the quadratic Gauss sum, is the case $\tau(\chi) = i^{a_\chi} \sqrt{f_\chi}$ recorded in section 7.3 and proved in full in [1, Ch. 2] and [4, Ch. 9]. For a genuinely complex character, $\chi \neq \bar{\chi}$, the pair $\{\chi, \bar{\chi}\}$ contributes $\epsilon(\chi)\epsilon(\bar{\chi})$, and this product is $+1$ by the reflection alone, with no sign theorem needed: applying (8.41) to $\bar{\chi}$ at the point $1-s$ gives $\Lambda(1-s, \bar{\chi}) = \epsilon(\bar{\chi}) \Lambda(s, \chi)$ (using $\bar{\bar{\chi}} = \chi$), and substituting this into $\Lambda(s, \chi) = \epsilon(\chi) \Lambda(1-s, \bar{\chi})$ yields $\Lambda(s, \chi) = \epsilon(\chi)\epsilon(\bar{\chi}) \Lambda(s, \chi)$; since $\Lambda(s, \chi)$ is not identically zero (its L -factor is nonvanishing on $\text{Re}(s) > 1$ by the Euler product theorem 7.2.2), $\epsilon(\chi)\epsilon(\bar{\chi}) = 1$. Thus every conjugate pair contributes $+1$ and every real character contributes $+1$, so the total is

$$W = \prod_{\chi \in X} \epsilon(\chi) = \prod_{\chi \text{ real}} (+1) \cdot \prod_{\{\chi, \bar{\chi}\}} \epsilon(\chi)\epsilon(\bar{\chi}) = +1$$

Inserting $W = +1$ into (8.42) and then dividing the resulting identity $\prod_{\chi} \Lambda(s, \chi) = \prod_{\chi} \Lambda(1-s, \chi)$ by the positive constant c_K of (8.40) (which appears on both sides, since (8.40) holds with s replaced by $1-s$ as well) gives

$$\Lambda_K(s) = \frac{1}{c_K} \prod_{\chi \in X} \Lambda(s, \chi) = \frac{1}{c_K} \prod_{\chi \in X} \Lambda(1-s, \chi) = \Lambda_K(1-s)$$

This is the functional equation (8.31) for abelian K , established from theorem 7.3.4 and theorem 8.5.4 alone. The root number of ζ_K is the product $W = +1$ of the root numbers of its L -function factors, recovering the self-duality recorded after theorem 8.5.3.

The abelian computation exhibits, in miniature, how the invariants of K distribute among the Dirichlet L -functions: the signature (r_1, r_2) is read off from the parities of the characters in X by (8.36), the discriminant $|d_K|$ from their conductors by theorem 8.5.4, and the trivial root number from the pairing of conjugate characters and the reality of the quadratic ones. The same three pieces of data, assembled differently, are what the analytic class number formula of theorem 8.3.2 packages into the residue κ_K ; the functional equation and the class number formula are the two halves of the analytic theory of ζ_K , the former controlling its symmetry and the latter its single pole.

8.5.6 Remark: One object, three guises, and the road onward The Dedekind zeta function realizes the central objects of this book as special cases of a single construction. For $K = \mathbb{Q}$ it is the Riemann zeta function ζ of chapter 4, with its functional equation (6.14). For abelian K it factors, by (8.32), into the Dirichlet L -functions $L(s, \chi)$ of chapter 7, and its functional equation (8.31) is the product of theirs, as theorem 8.5.5 showed. For general K it is a new object, governed by Hecke's functional equation (8.31), whose proof passes through the geometry of the Minkowski lattice rather than through any factorization. The non-abelian generalization, in which ζ_K for a Galois extension $K/\mathbb{Q}$ with group G factors through the irreducible representations of G into Artin L -functions and the splitting of primes is governed by the Chebotarev density theorem (the non-abelian counterpart of Dirichlet's theorem of section 7.5), continues the arc begun in chapter 7; it is taken up in the successors to that chapter and, in its idelic and automorphic forms, in [9]. The pattern is uniform: an L -function packages the splitting of primes, its pole or its value at $s = 1$ encodes arithmetic invariants, and its functional equation fixes the symmetry of its zeros.

Index

- Ω Notation, [xxx](#)
- Θ Notation, [xxx](#)
- p -adic Valuation, [ix](#)

- Abel's Summation Formula, [17](#)
- Abel's Test, [21](#)
- Abscissa of Absolute Convergence, [32](#)
- Abscissa of Convergence, [32](#)
- Additive Function, [2](#)
- Analyticity of Dirichlet Series, [35](#)
- Arithmetic Function, [1](#)
- Asymptotic Equivalence, [xxx](#)
- Asymptotic Notation, [xxix](#)
- Average Order, [12](#)
 - of $\omega(n)$, [16](#)

- Bernoulli Numbers, [22](#)
- Bernoulli Polynomials, [23](#)
- Bertrand's Postulate, [93](#)
- Big- O Notation, [xxix](#)
- Bézout's Identity, [iv](#)

- Carmichael Function, [xxii](#)
- Central Binomial Coefficient
 - prime content, [73](#)
 - prime factorization, [92](#)
 - size bounds, [91](#)
- Character Orthogonality, [182](#)
- Chebyshev Functions, [68](#)
 - comparison of ψ and ϑ , [69](#)
- Chebyshev's Estimates, [75](#)
- Chinese Remainder Theorem, [xiv](#)

- Class Number Formula, [232](#)
- Completed L -Function, [197](#)
- Completed Dedekind Zeta Function, [243](#)
- Completed Zeta Function, [158](#)
- Conductor, [184](#)
- Conductor-Discriminant Formula, [245](#)
- Congruence, [xii](#)
- Coprime, [iii](#)
- Counting Ideals in an Ideal Class, [224](#)
- Critical Line, [171](#)
- Critical Strip, [171](#)

- Dedekind Zeta Function, [218](#)
 - cyclotomic factorization, [240](#)
 - quadratic factorization, [235](#)
 - residue at $s = 1$, [232](#)
- Dedekind-type product
 - nonnegative coefficients, [203](#)
- Dirichlet L -Function, [187](#)
 - continuation, [190](#)
 - Euler product, [187](#)
 - non-vanishing at 1, [206](#)
 - principal, [188](#)
- Dirichlet Character, [180](#)
 - dual group, [181](#)
 - even and odd, [185](#)
- Dirichlet Convolution, [6](#)
- Dirichlet Density, [212](#)
- Dirichlet Hyperbola Method, [12](#)
- Dirichlet Series, [32](#)
 - Multiplication Theorem, [37](#)

- Dirichlet's Theorem, 209
- Discontinuous Integral, 58
- Divisibility, ii
- Division Algorithm, i
- Divisor Function, 3
- Divisor Sum Asymptotic, 13
- Equidistribution of Primes, 213
- Euclidean Algorithm, iii
- Euler Product, 43
 - Dedekind zeta, 221
 - Logarithm, 48
 - Riemann Zeta Function, 45, 99
- Euler Totient Function, xiii
- Euler's Criterion, xxiv
- Euler's Theorem, xvi
- Euler-Maclaurin Summation Formula, 25
- Exact Divisibility, xi
- Fermat's Little Theorem, xvi
- Fourier Transform
 - of a Gaussian, 154
- Functional Equation
 - abelian consistency, 247
 - Dedekind zeta, 244
 - Dirichlet L -function, 198
 - Riemann zeta, 159
- Fundamental Theorem of Arithmetic, viii
- Gamma Factor
 - real place, 158
- Gamma Function, 146
 - duplication formula, 150
 - meromorphic continuation, 146
 - reflection formula, 148
- Gauss Sum, 193
 - modulus, 194
- Gauss's Lemma, xxv
- Greatest Common Divisor, iii
- Hardy Z-Function, 176
- Hecke's Functional Equation, 244
- Ideal Counting Function, 224
- Irreducible, vii
- Jacobi Symbol, xxviii
- Kummer's Theorem, xii
- Landau's Theorem, 33
- Laplace Transform, 118
 - auxiliary function g , 120
 - of the Chebyshev function, 119
- Laurent Expansion
 - of ζ at $s = 1$, 103
- Legendre Symbol, xxiv
- Legendre's Formula, xi
- Linear Diophantine Equation, vi
- Liouville Function, 3
- Lipschitz Lattice-Point Principle, 225
- Logarithmic Derivative of ζ
 - pole at $s = 1$, 105
- Logarithmic Derivative of Zeta, 41
 - regularity on $\text{Re}(s)=1$, 114
- Möbius Function, 3
- Möbius Inversion Formula, 9
- Mellin Transform, 51
 - Inversion Formula, 53
- Mertens Constant, 85
- Mertens Function, 14
- Mertens Inequality ($3+4\cos+\cos^2$), 111
- Mertens' Theorems
 - First, 82
 - Second, 83
 - Third, 88
- Multiplicative Function, 2
- Multiplicative Order, xviii
- Newman's Tauberian Theorem, 122
- Non-Vanishing
 - algebraic proof, 237
 - via Euler Product, 47
- Nontrivial Zeros
 - symmetry, 172
- Number of Distinct Prime Factors, 3
- Number of Prime Factors with Multiplicity, 3
- Partial Summation, 17
- Periodic Bernoulli Functions, 25
- Perron's Formula, 61
 - Truncated Form, 63
- Poisson Summation Formula, 153
- Prime, vii
- Prime Counting Function, 68
 - order of growth, 77
- Prime Number Theorem, 116, 132
 - consequences, 134

- equivalent forms, 71
 - error term, 141
 - key integral, 127
 - monotonicity argument, 130
- Prime Reciprocal Sum
 - analytic divergence, 109
- Prime Series Φ , 106
 - pole at $s = 1$, 107
- Prime-Power Decomposition of $\log n!$, 80
- Primitive Character, 184
- Primitive Root, xviii
 - Existence, xix
- Primorial Bound, 74
- Principal Character, 182
- Quadratic Reciprocity, xxvi
- Quadratic Residue, xxiii
- Residue Class Ring, xii
- Riemann Zeta Function, 98
 - continuation to $\mathbb{C}$, 162
 - growth near $\operatorname{Re}(s)=1$, 135
 - Integral Representation, 56
 - meromorphic continuation, 101
 - non-vanishing on $\operatorname{Re}(s)=1$, 111
 - trivial zeros, 165
 - values at even integers, 169
 - values at negative integers, 166
- Ring Isomorphism
 - Arithmetic Functions and Dirichlet Series, 40
- Ring of Arithmetic Functions, 7
- Root Number, 198
- Stirling's Formula
 - complex, 150
- Sum-of-Divisors Function, 3
- Summatory Function, 12
- Theta Function (Jacobi), 156
 - functional equation, 156
- Totally Multiplicative Function, 2
- Vinogradov Notation, xxix
- Von Mangoldt Function, 4
 - Dirichlet Series, 41
- Wilson's Theorem, xvi
- Zero-Counting Formula (Riemann–von Mangoldt), 174
- Zero-Free Region
 - de la Vallée Poussin, 138

Bibliography

- [1] Harold Davenport. *Multiplicative Number Theory*. 3rd ed. Graduate Texts in Mathematics 74. Revised and with a preface by Hugh L. Montgomery. New York: Springer-Verlag, 2000. ISBN: 978-0-387-95097-6.
- [2] G. H. Hardy and E. M. Wright. *An Introduction to the Theory of Numbers*. Ed. by D. R. Heath-Brown and J. H. Silverman. 6th ed. Oxford: Oxford University Press, 2008. ISBN: 978-0-19-921986-5.
- [3] Serge Lang. *Algebraic Number Theory*. 2nd ed. Graduate Texts in Mathematics 110. New York: Springer-Verlag, 1994. ISBN: 978-0-387-94225-4.
- [4] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative Number Theory I: Classical Theory*. Cambridge Studies in Advanced Mathematics 97. Cambridge: Cambridge University Press, 2007. ISBN: 978-0-521-84903-6.
- [5] Jürgen Neukirch. *Algebraic Number Theory*. Grundlehren der mathematischen Wissenschaften 322. Translated from the 1992 German original by Norbert Schappacher. Berlin: Springer-Verlag, 1999. ISBN: 978-3-540-65399-8.
- [6] D. J. Newman. “Simple Analytic Proof of the Prime Number Theorem”. In: *The American Mathematical Monthly* 87.9 (1980), pp. 693–696. DOI: [10.2307/2321853](https://doi.org/10.2307/2321853).
- [7] Nathanael Chwojko-Srawley. *Everything You Need To Know About Harmonic Analysis*. 1st ed. Vol. 1. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_harmonic_analysis.pdf.
- [8] Nathanael Chwojko-Srawley. *Everything You Need To Know About Real Analysis*. 1st ed. Vol. 1. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_real_analysis.pdf.
- [9] Nathanael Chwojko-Srawley. *Everything You Need To Know About Class Field Theory*. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_class_field_theory.pdf.
- [10] Gérald Tenenbaum. *Introduction to Analytic and Probabilistic Number Theory*. 3rd ed. Graduate Studies in Mathematics 163. Providence, RI: American Mathematical Society, 2015. ISBN: 978-0-8218-9854-3.

- [11] E. C. Titchmarsh. *The Theory of the Riemann Zeta-Function*. Ed. by D. R. Heath-Brown. 2nd ed. Revised by D. R. Heath-Brown. Oxford: Clarendon Press, Oxford University Press, 1986. ISBN: 978-0-19-853369-6.
- [12] Mark B. Villarino. “Mertens’ Proof of Mertens’ Theorem”. In: (2005). arXiv:math/0504289. arXiv: [math/0504289](https://arxiv.org/abs/math/0504289) [math.HO].
- [13] Nathanael Chwojko-Srawley. *Everything You Need To Know About Undergraduate Algebra*. 1st ed. n.d. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_algebra.pdf.
- [14] Lawrence C. Washington. *Introduction to Cyclotomic Fields*. 2nd ed. Graduate Texts in Mathematics 83. New York: Springer-Verlag, 1997. ISBN: 978-0-387-94762-4.
- [15] Nathanael Chwojko-Srawley. *Everything You Need To Know About Local Field Theory*. URL: <https://nathanaelsrawley.com/pdf-viewer?src=.%2F.%2Fassets%2Fpdfs%2Fbooks%2FEYNTKA-local-field-theory.pdf&name=Local%20Field%20Theory&source=books>.
- [16] Nathanael Chwojko-Srawley. *Everything You Need To Know About Complex Analysis*. 1st ed. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_complex_analysis.pdf.
- [17] Nathanael Chwojko-Srawley. *Everything You Need To Know About Algebraic Number Theory*. Vol. 1. URL: <https://nathanaelsrawley.com/pdf-viewer?src=.%2F.%2Fassets%2Fpdfs%2Fbooks%2FEYNTKA-algebraic-number-theory.pdf&name=Algebraic%20Number%20Theory&source=books>.
- [18] D. Zagier. “Newman’s Short Proof of the Prime Number Theorem”. In: *The American Mathematical Monthly* 104.8 (1997), pp. 705–708. DOI: [10.2307/2975232](https://doi.org/10.2307/2975232).