Hodge Theory

***EYNTKA* Series**

Nathanael Chwojko-Srawley

October 28, 2026

Contents

Contents iii

Introduction vii

I The Hodge Decomposition 1

1 Kähler Manifolds and the Master Identity 5

1.1 Complex and Kähler Geometry Recalled 5

1.2 The Frölicher Spectral Sequence and GAGA 9

1.3 The Master Identity 16

1.4 The $\partial\bar{\partial}$-Lemma and a Non-Kähler Surface 19

2 The Hodge Decomposition 27

2.1 The Hodge Decomposition 27

2.2 First Consequences 32

2.3 The Lefschetz $(1,1)$-Theorem 37

II Hodge Structures and Their Variations 43

3 Hodge Structures and Polarizations 47

3.1 Pure Hodge Structures and the Hodge Filtration 47

3.2 Polarizations and the Hodge-Riemann Bilinear Relations 55

3.3 The Period Domain and Its Compact Dual . . . 65
3.4 First Examples: Weight One and Weight Two . . . 73

4 Hard Lefschetz and Lefschetz Pencils 83
4.1 The $\mathfrak{sl}_2$-Action on Cohomology . . . 83
4.2 The Hard Lefschetz Theorem . . . 93
4.3 Lefschetz Pencils and Vanishing Cycles . . . 103
4.4 Monodromy and the Picard-Lefschetz Formula . . . 116

5 Variations of Hodge Structure 129
5.1 Local Systems and the Gauss-Manin Connection . . . 130
5.2 Griffiths Transversality . . . 139
5.3 The Period Map . . . 150
5.4 Hypersurfaces and the Jacobian Ring . . . 161

6 Period Domains and Mumford-Tate Groups 177
6.1 Curvature and Rigidity of Period Domains . . . 178
6.2 Mumford-Tate Groups . . . 189
6.3 Hodge Loci and Algebraicity . . . 200
6.4 Moduli of Polarized K3 Surfaces . . . 210

III Mixed Hodge Structures and Cycles 221

7 Mixed Hodge Structures: Smooth Open Varieties 225
7.1 Mixed Hodge Structures . . . 226
7.2 The Logarithmic de Rham Complex . . . 238
7.3 Deligne's Mixed Hodge Structure . . . 251
7.4 The Punctured Elliptic Curve . . . 264

8 Mixed Hodge Structures: Singular Varieties 273
8.1 Cohomological Mixed Hodge Complexes . . . 274
8.2 Simplicial and Cubical Hyperresolutions . . . 286
8.3 Deligne's Theorem for Arbitrary Varieties . . . 295
8.4 The Nodal Cubic . . . 304

What Next? 311

Index 313

Bibliography **319**

Introduction

A compact smooth manifold has de Rham cohomology, and a Riemannian metric on it picks out a distinguished representative in every class, the harmonic forms of *Riemannian Geometry* [12]. A smooth projective variety over $\mathbb{C}$ has coherent sheaf cohomology $H^q(X, \Omega_X^p)$, computed from the algebraic structure alone in *Algebraic Geometry* [4]. Nothing connects the two. The first is an invariant of the underlying smooth manifold and cannot detect which complex structure was placed on it; the second is defined only after that complex structure is fixed, and its individual pieces are not topological invariants at all. Hodge theory is the discovery that on a compact Kähler manifold each determines the other:

$$H^k(X, \mathbb{C}) \cong \bigoplus_{p+q=k} H^q(X, \Omega_X^p)$$

Both sides of this isomorphism were already available to the reader; what is new is that they are the same.

The consequences run in both directions. Left to right, the topology bounds the algebra: the Hodge numbers $h^{p,q} = \dim_{\mathbb{C}} H^q(X, \Omega_X^p)$ sum along each anti-diagonal to a Betti number, so a purely topological count constrains a sheaf-cohomological one. Right to left, the algebra obstructs the geometry. Complex conjugation exchanges $H^{p,q}$ with $H^{q,p}$, forcing $h^{p,q} = h^{q,p}$, and therefore forcing every odd Betti number of a compact Kähler manifold to be even. The Hopf surface, a compact complex surface diffeomorphic to $S^1 \times S^3$, has $b_1 = 1$. It admits no Kähler metric, hence no projective embedding, and the proof is a count of the Betti numbers of a product of spheres.

The identity has a prehistory in the one-dimensional case. Riemann's bilinear relations of 1857 record symmetry and positivity constraints on the periods of holomorphic differentials on a compact Riemann surface, and they are the weight-one instance of everything in this book. Hodge's contribution, developed through the 1930s and published as *The Theory and Applications of Harmonic Integrals* in 1941 [25], was to read those constraints as coming from a decomposition of cohomology, and to produce the decomposition analytically by solving for harmonic representatives. The analysis was the difficult part, and the existence theorem it required was placed on secure foundations over the following years by Weyl [40] and Kodaira. Two further ingredients came from outside: Kähler's 1933 condition on a Hermitian metric, that its associated $(1,1)$-form be closed [29], which turned out to be exactly the hypothesis the decomposition needs, and Serre's GAGA [36], which identifies the analytic

invariants with algebraic ones on a projective variety.

What kept the subject growing after the decomposition theorem was that the decomposition moves. In a family of compact Kähler manifolds the underlying topological cohomology is locally constant while the Hodge filtration varies holomorphically, and Griffiths, in the late 1960s, made that variation the object of study [22]: the period map, its derivative, and the transversality constraint the derivative satisfies. Deligne then removed the hypotheses. *Théorie de Hodge* II [18] and III [19] attach to the cohomology of an arbitrary complex algebraic variety, open or singular, a functorial mixed Hodge structure whose weight filtration measures the failure of purity. The subject's central open problem predates both: Hodge's 1950 conjecture, that on a smooth projective variety every class in $H^{2p}(X,\mathbb{Q}) \cap H^{p,p}(X)$ is a rational combination of classes of codimension-p subvarieties, is unproved and is one of the Millennium Prize problems.

A word on sources, since this book cites few. The EYNTKA series is written to be self-contained: a reader holding the prerequisites should never need an outside text to follow an argument, and where a result is taken rather than proved it is taken from another volume of the series. That is a constraint on the exposition, not a claim of originality. The treatment below is shaped throughout by Voisin's two volumes [38, 39], which set the order of the Hodge decomposition and of the theory of variations; by Huybrechts [27] for the Hodge theory of a compact Kähler manifold; by Griffiths and Harris [20] for the classical geometry; and, for the structural half, by Carlson, Müller-Stach and Peters [1] on period mappings and by Peters and Steenbrink [33] on mixed Hodge structures. A reader who wants a second account of any chapter should go to these first.

Prerequisites and Place in the Series This volume builds on *Complex Analytic Geometry* [5], *Complex Analysis* [31], *Algebraic Geometry* [4], *Riemannian Geometry* [12], *Differential Topology* [32], *Algebraic Topology* [30], *Homological Algebra* [7], *Lie Groups* [37], and *Algebraic Groups and Representations* [14]. The last two are not needed until Part II, and the algebraic-group material only from Chapter 6 onward, where Mumford-Tate groups are defined; Part I uses neither. It is naturally read alongside *Shimura Varieties* [13] and *p-adic Hodge Theory*. No later volume currently lists it as a prerequisite. The reader has access to every completed volume of the series but is not assumed to have read all of them; prerequisites are cited where invoked.

Notation and Conventions

X denotes a complex manifold, always connected, and n its complex dimension $\dim_{\mathbb{C}} X$. Compactness is never silent: it is stated in the hypotheses of every result that uses it. Smooth means C^∞, and the base field is $\mathbb{C}$ throughout.

The sheaf of holomorphic functions on X is $\mathcal{O}_X$ and the sheaf of holomorphic p-forms is Ω^p_X; $\mathcal{A}^k$ and $\mathcal{A}^{p,q}$ denote the sheaves of smooth complex-valued k-forms and (p,q)-forms, with $\mathcal{A}^k(X)$ and $\mathcal{A}^{p,q}(X)$ their global sections. Sheaf cohomology is $H^q(X,\mathcal{F})$, Dolbeault cohomology is $H^{p,q}_{\bar\partial}(X)$, and $H^k(X,\mathbb{C})$ is used for singular and de Rham cohomology with complex coefficients without distinguishing them, the de Rham theorem being available from *Differential Topology* [32, The de Rham Theorem]. Hodge and Betti numbers are $h^{p,q}$ and b_k. Complex projective space is $\mathbb{C}\mathrm{P}^n$.

Goal

Part I proves the Hodge decomposition. It starts from the analytic geometry of complex manifolds developed in *Complex Analytic Geometry* [5] and recalled in section 1.1: the splitting of forms by type that turns the exterior derivative into $d = \partial + \bar{\partial}$, Dolbeault's theorem $H^q(X, \Omega_X^p) \cong H_{\bar{\partial}}^{p,q}(X)$, Kähler metrics and the Kähler identities, and harmonic theory for $\bar{\partial}$. The Frölicher spectral sequence measures how far the bigraded Dolbeault data sits from de Rham cohomology, and the Iwasawa manifold shows the gap can be nonzero. A Kähler metric closes it. The Kähler identities give the single equation $\Delta_d = 2\Delta_{\bar{\partial}}$ on which the decomposition rests, and harmonic theory turns it into $H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$. With the decomposition come Hodge symmetry, the parity of the odd Betti numbers, the degeneration of the Frölicher spectral sequence, the Hodge index theorem on surfaces, and the Lefschetz $(1,1)$-theorem identifying the integral $(1,1)$-classes on a projective manifold as classes of divisors.

The rest of the book abstracts the decomposition, then breaks its hypotheses. A Hodge structure is the bigrading together with the polarization the Kähler class gives, taken as data on a rational vector space in its own right; those with fixed numerical invariants are parametrized by a period domain. Hard Lefschetz and the primitive decomposition sharpen the picture on a single manifold, while Lefschetz pencils give the vanishing cycles and monodromy that make families tractable. In a family the Hodge structures vary: the underlying local system is flat, the filtration is holomorphic, and Griffiths transversality constrains its derivative, which makes the period map rigid enough to yield Torelli theorems, Noether-Lefschetz statements, and the algebraicity of Hodge loci. Dropping compactness or smoothness costs purity but not the theory, and Deligne's weight filtration produces a functorial mixed Hodge structure on any complex algebraic variety, built from the logarithmic de Rham complex in the smooth open case and from simplicial resolutions in general. The cycle class map and the Hodge conjecture close the mathematics, followed by a survey of what has been left undeveloped: Saito's mixed Hodge modules, the decomposition theorem, and nonabelian and p-adic Hodge theory.

A reader who finishes can prove the Hodge decomposition for a compact Kähler manifold, set up a variation of Hodge structure and apply Griffiths transversality, write down the mixed Hodge structure on the cohomology of a smooth open or singular variety, and read the current literature on period maps, Hodge loci, and algebraic cycles without further preparation.

Part I

The Hodge Decomposition

A compact complex manifold X has two cohomologies with complex coefficients: de Rham cohomology $H^k(X, \mathbb{C})$, which sees only the underlying smooth manifold, and Dolbeault cohomology $H^{p,q}_{\bar\partial}(X)$, which sees the complex structure. The exterior derivative splits as $d = \partial + \bar\partial$, and the question is whether H^k splits as the sum of the $H^{p,q}$ with $p + q = k$. On a general compact complex manifold it does not: the Frölicher spectral sequence relating the two theories need not degenerate, and on the Iwasawa manifold it does not.

A Kähler metric repairs this. Its fundamental form is closed, and the Kähler identities that follow make the de Rham Laplacian twice the Dolbeault Laplacian. Harmonic forms for one operator are then harmonic for the other, and since harmonic forms represent de Rham classes for the first and Dolbeault classes for the second, the two theories share one space of representatives. The result is the Hodge decomposition $H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ with $H^{p,q} = \overline{H^{q,p}}$, together with its first consequences: the odd Betti numbers of a compact Kähler manifold are even, the Frölicher spectral sequence degenerates, and on a projective manifold the integral classes of type $(1, 1)$ are the classes of divisors.

1

Kähler Manifolds and the Master Identity

Section 1.1 recalls from *Complex Analytic Geometry* [5] the inputs of the Hodge decomposition: the Dolbeault formalism and Dolbeault's theorem, Kähler metrics and the Kähler identities, and harmonic theory for $\bar{\partial}$. On a general compact complex manifold the Dolbeault groups are tied to de Rham cohomology only through the Frölicher spectral sequence, whose degeneration can fail. On a Kähler manifold the Kähler identities give the master identity $\Delta_d = 2\Delta_{\bar{\partial}} = 2\Delta_{\partial}$, and the $\partial\bar{\partial}$-lemma holds on a compact one. The Hopf surface, a compact complex manifold with no Kähler metric, shows that the Kähler hypothesis is a restriction.

1.1 Complex and Kähler Geometry Recalled

This section recalls the analytic geometry of complex manifolds developed in *Complex Analytic Geometry* [5], in the notation used from now on. Throughout, X is a complex manifold of complex dimension n, $\mathcal{A}^{p,q}(X)$ is the space of smooth complex-valued (p,q)-forms on X, and Ω^p_X is the sheaf of holomorphic p-forms.

Proposition 1.1.1: The Dolbeault Formalism

Let X be a complex manifold of complex dimension n.

1. The complex-valued k-forms split by type, $\mathcal{A}^k(X;\mathbb{C}) = \bigoplus_{p+q=k} \mathcal{A}^{p,q}(X)$, and $\mathcal{A}^{p,q}(X) = 0$ unless $0 \le p, q \le n$.
2. The exterior derivative splits as $d = \partial + \bar{\partial}$ with $\partial\colon \mathcal{A}^{p,q}(X) \to \mathcal{A}^{p+1,q}(X)$ and $\bar{\partial}\colon \mathcal{A}^{p,q}(X) \to \mathcal{A}^{p,q+1}(X)$; in holomorphic coordinates they differentiate the coefficients of a form in the holomorphic and the antiholomorphic directions.
3. $\partial^2 = 0$, $\bar{\partial}^2 = 0$, and $\partial\bar{\partial} + \bar{\partial}\partial = 0$.

For each p the *Dolbeault complex* is $0 \to \mathcal{A}^{p,0}(X) \xrightarrow{\bar{\partial}} \mathcal{A}^{p,1}(X) \xrightarrow{\bar{\partial}} \cdots \xrightarrow{\bar{\partial}} \mathcal{A}^{p,n}(X) \to 0$, and its cohomology $H^{p,q}_{\bar{\partial}}(X)$ is the *Dolbeault cohomology* of X. A $(p,0)$-form α is *holomorphic* if $\bar{\partial}\alpha = 0$, and the holomorphic p-forms make up the sheaf Ω^p_X.

***Proof Cited*:**
Item (1) is the bidegree decomposition, item (2) the decomposition of the exterior derivative, and item (3) the Dolbeault identities, all proved in *Complex Analytic Geometry*.[a]

[a] [5, Bidegree Decomposition of k-Forms, Decomposition of the Exterior Derivative, Dolbeault Identities].

A sheaf of modules over a sheaf of rings $\mathcal{R}$ on a paracompact Hausdorff space is *fine* if every locally finite open cover carries a partition of unity by $\mathcal{R}$-linear endomorphisms supported in its members.[1]

Theorem 1.1.2: Fine Resolutions

Let X be a paracompact Hausdorff space.

1. A fine sheaf on X is acyclic.
2. If $0 \to \mathcal{F} \to \mathcal{L}^0 \to \mathcal{L}^1 \to \cdots$ is a resolution of a sheaf $\mathcal{F}$ by acyclic sheaves, then $H^q(X,\mathcal{F})$ is canonically isomorphic to the q-th cohomology of the complex of global sections $\Gamma(X,\mathcal{L}^\bullet)$, naturally in $\mathcal{F}$ and independently of the resolution.
3. If X is a complex manifold, the sheaves $\mathcal{A}^{p,q}$ of smooth (p,q)-forms and $\mathcal{A}^k$ of smooth k-forms are fine over the sheaf of smooth functions, hence acyclic.

***Proof Cited*:**
The three items are the acyclicity of fine sheaves, the abstract de Rham theorem, and the fineness of the smooth form sheaves.[a]

[a] [5, Fine Sheaves Are Acyclic, Abstract de Rham Theorem, Smooth Form Sheaves Are Fine].

[1] [5, Fine and Soft Sheaves].

Theorem 1.1.3: Dolbeault's Theorem

For every complex manifold X and all p, q there is a canonical isomorphism

$$H^q(X, \Omega^p_X) \cong H^{p,q}_{\bar\partial}(X)$$

When X is compact these groups are finite-dimensional, and their dimensions

$$h^{p,q}(X) = \dim_{\mathbb{C}} H^{p,q}_{\bar\partial}(X) = \dim_{\mathbb{C}} H^q(X, \Omega^p_X)$$

are the *Hodge numbers* of X.

***Proof Cited*:**
The isomorphism is Dolbeault's theorem, and the Hodge numbers are defined beside it.[a] Finiteness on a compact manifold is part of the $\bar\partial$-Hodge theorem (theorem 1.1.10), which applies because every complex manifold carries a Hermitian metric, glued from local flat metrics by a partition of unity.

[a] [5, Dolbeault's Theorem, Hodge Numbers].

Example 1.1.4: Dolbeault Cohomology of the Riemann Sphere

On $\mathbb{C}\mathrm{P}^1$ the groups $H^{0,0}_{\bar\partial}$ and $H^{1,1}_{\bar\partial}$ are one-dimensional, and $H^{1,0}_{\bar\partial}$ and $H^{0,1}_{\bar\partial}$ vanish. In particular $H^1(\mathbb{C}\mathrm{P}^1, \mathcal{O}) = 0$.[a]

[a] [5, Dolbeault Cohomology of the Riemann Sphere].

Definition 1.1.5: Hermitian and Kähler Metrics

A *Hermitian metric* on X is a smoothly varying positive-definite Hermitian inner product h on the holomorphic tangent spaces $T^{1,0}_p X$; in holomorphic coordinates it is the positive-definite Hermitian matrix of the functions $h_{j\bar k} = h(\partial/\partial z^j, \partial/\partial z^k)$. Its *fundamental form* is the real $(1,1)$-form $\omega = i\sum_{j,k} h_{j\bar k}\, dz^j \wedge d\bar z^k$. The metric is *Kähler* if $d\omega = 0$, and then (X, ω) is a *Kähler manifold*.[a]

[a] [5, Hermitian Metric, The Fundamental Form, Kähler Metric and Kähler Manifold].

Example 1.1.6: Complex Tori

A lattice $\Lambda \subset \mathbb{C}^n$, a discrete subgroup of rank $2n$, acts on $\mathbb{C}^n$ by translations, which are biholomorphisms, and the quotient $\mathbb{C}^n/\Lambda$ is a compact complex manifold diffeomorphic to $(\mathbb{R}/\mathbb{Z})^{2n}$, the *complex torus* of Λ. For $n = 1$ the complex tori are the elliptic curves; for $n \geq 2$ only some complex tori embed in projective space, and these are the abelian varieties. A constant positive-definite Hermitian matrix $(h_{j\bar k})$ defines a translation-invariant metric on $\mathbb{C}^n$ that descends to $\mathbb{C}^n/\Lambda$. Its fundamental form has constant coefficients, so it is closed, and the descended metric is a Kähler metric, a *flat* metric.[a]

[a] [5, Complex Tori, Complex Tori with a Constant Metric].

Theorem 1.1.7: Projective Manifolds Are Kähler

The Fubini-Study form $\omega_{\mathrm{FS}} = \frac{i}{2\pi}\,\partial\bar\partial \log\|Z\|^2$ is a Kähler form on $\mathbb{C}\mathrm{P}^n$. Its class in $H^2(\mathbb{C}\mathrm{P}^n,\mathbb{Z})$ is $[\omega_{\mathrm{FS}}] = c_1(\mathcal{O}(1))$, the positive generator, and $\int_{\mathbb{C}\mathrm{P}^1}\omega_{\mathrm{FS}} = 1$ on every line $\mathbb{C}\mathrm{P}^1 \subseteq \mathbb{C}\mathrm{P}^n$. If (X,ω) is a Kähler manifold and $\iota\colon Y \hookrightarrow X$ is a complex submanifold, then $(Y,\iota^*\omega)$ is a Kähler manifold. Hence every smooth projective variety over $\mathbb{C}$ is a compact Kähler manifold, with the restriction of ω_{FS} as Kähler form.

***Proof Cited*:**
These are the Kähler property of $\mathbb{C}\mathrm{P}^n$, the identification of the Fubini-Study class, and the corollary that projective manifolds are Kähler.[a]

[a] [5, $\mathbb{C}\mathrm{P}^n$ Is Kähler, The Fubini-Study Class and $c_1(\mathcal{O}(1))$, Projective Manifolds Are Kähler].

Definition 1.1.8: The Lefschetz Operators and the Formal Adjoints

Let (X,ω) be a Hermitian manifold. The *Lefschetz operator* $L\colon \mathcal{A}^{p,q}(X) \to \mathcal{A}^{p+1,q+1}(X)$ is $L\alpha = \omega\wedge\alpha$, and its pointwise adjoint $\Lambda = L^*\colon \mathcal{A}^{p,q}(X) \to \mathcal{A}^{p-1,q-1}(X)$ for the Hermitian inner product on forms is the *adjoint Lefschetz operator*. When X is compact, $\partial^* = -\star\,\bar\partial\,\star$ and $\bar\partial^* = -\star\,\partial\,\star$ are the formal adjoints of ∂ and $\bar\partial$ for the L^2 inner product $\langle\!\langle\alpha,\beta\rangle\!\rangle = \int_X\langle\alpha,\beta\rangle\,\mathrm{vol}$, where $\star\colon \mathcal{A}^{p,q}(X) \to \mathcal{A}^{n-q,n-p}(X)$ is the $\mathbb{C}$-linear Hodge star of the metric.[a]

[a] [5, The Lefschetz Operators, The Formal Adjoints ∂^* and $\bar\partial^*$].

Theorem 1.1.9: The Kähler Identities

On a Kähler manifold (X,ω),

$$[\Lambda,\bar\partial] = -i\,\partial^*, \qquad [\Lambda,\partial] = i\,\bar\partial^* \tag{1.1}$$

and, by adjunction, $[L,\partial^*] = i\,\bar\partial$ and $[L,\bar\partial^*] = -i\,\partial$.

***Proof Cited*:**
Complex Analytic Geometry [5, The Kähler Identities] proves them on flat $\mathbb{C}^n$ and transports them to X in Kähler normal coordinates.

Theorem 1.1.10: The $\bar\partial$-Hodge Theorem and Serre Duality

Let X be a compact Hermitian manifold of complex dimension n, and let $\Delta_{\bar\partial} = \bar\partial\bar\partial^* + \bar\partial^*\bar\partial$. Then $\Delta_{\bar\partial}$ is a formally self-adjoint elliptic operator of order two, and for all p, q:

1. the space $\mathcal{H}^{p,q}_{\bar\partial}(X)$ of $\bar\partial$-*harmonic* (p,q)-forms, those killed by $\Delta_{\bar\partial}$, is finite-dimensional, and there is an orthogonal decomposition $\mathcal{A}^{p,q}(X) = \mathcal{H}^{p,q}_{\bar\partial}(X) \oplus \bar\partial\,\mathcal{A}^{p,q-1}(X) \oplus \bar\partial^*\mathcal{A}^{p,q+1}(X)$;
2. every Dolbeault class has a unique $\bar\partial$-harmonic representative, so $\mathcal{H}^{p,q}_{\bar\partial}(X) \cong H^{p,q}_{\bar\partial}(X) \cong H^q(X, \Omega^p_X)$;
3. if X is connected, $([\alpha],[\beta]) \mapsto \int_X \alpha \wedge \beta$ is a perfect pairing $H^q(X,\Omega^p_X) \times H^{n-q}(X,\Omega^{n-p}_X) \to \mathbb{C}$, so $h^{p,q} = h^{n-p,n-q}$.

No Kähler hypothesis is needed.

***Proof Cited*:**
The three items are the ellipticity of $\Delta_{\bar\partial}$ with the $\bar\partial$-Hodge theorem, and Serre duality for Ω^p.[a]

[a] [5, $\Delta_{\bar\partial}$ is Self-Adjoint Elliptic of Order Two, The $\bar\partial$-Hodge Theorem, Serre Duality for Ω^p].

1.2 The Frölicher Spectral Sequence and GAGA

Dolbeault's theorem (theorem 1.1.3) computes each individual group $H^{p,q}_{\bar\partial}(X)$ as the sheaf cohomology $H^q(X,\Omega^p_X)$. These groups form a bigraded refinement of the complex de Rham cohomology $H^k_{\mathrm{dR}}(X,\mathbb{C})$: the de Rham complex is built from the operator $d = \partial + \bar\partial$ (proposition 1.1.1), whose two pieces separately assemble the Dolbeault groups. The question this section answers is how the bigraded data $H^{p,q}_{\bar\partial}(X)$ reassembles into the single-graded data $H^k_{\mathrm{dR}}(X,\mathbb{C})$. On an arbitrary complex manifold the two are not simply related: the bidegree decomposition of forms does not descend to a decomposition of cohomology, because d mixes the pieces. What organizes the discrepancy is a spectral sequence, the Frölicher (or Hodge-de Rham) spectral sequence, whose first page is the Dolbeault cohomology and whose limit is the de Rham cohomology. Its degeneration is exactly the statement that the bigrading survives to cohomology, and this degeneration, proved for compact Kähler manifolds in chapter 2, is one form of the Hodge decomposition. The section closes with GAGA, which identifies the analytic groups $H^q(X,\Omega^p_X)$ with their algebraic counterparts when X is projective, so that transcendental Hodge theory and algebraic geometry compute the same numbers.

The de Rham complex $(\mathcal{A}^\bullet(X), d)$ of global smooth forms carries a decreasing filtration by holomorphic degree. Recall from proposition 1.1.1 that a smooth k-form decomposes into its (p,q)-components with $p+q=k$. Collecting all components whose holomorphic degree is at least p produces a subspace of $\mathcal{A}^k(X)$, and letting k vary gives a graded subspace of the whole complex.

Definition 1.2.1: The Hodge Filtration on the de Rham Complex

Let X be a complex manifold. For each integer $p \geq 0$, let

$$F^p\mathcal{A}^k(X) = \bigoplus_{\substack{r+s=k \\ r \geq p}} \mathcal{A}^{r,s}(X)$$

be the space of smooth k-forms all of whose bidegree components have holomorphic degree at least p. The collection $F^\bullet$ is the *Hodge filtration* (or *filtration bête* by holomorphic degree) on the de Rham complex $(\mathcal{A}^\bullet(X), d)$.

For this to be a filtration of the complex, the differential must respect it: d must carry $F^p\mathcal{A}^k(X)$ into $F^p\mathcal{A}^{k+1}(X)$. This is where the decomposition $d = \partial + \bar{\partial}$ does the work. On a component of bidegree (r, s) with $r \geq p$, the operator ∂ raises the bidegree to $(r+1, s)$ and $\bar{\partial}$ raises it to $(r, s+1)$; both outputs still have holomorphic degree at least p. So $d\big(F^p\mathcal{A}^k(X)\big) \subseteq F^p\mathcal{A}^{k+1}(X)$, and the filtration is compatible with d. It is decreasing, $F^{p+1}\mathcal{A}^k(X) \subseteq F^p\mathcal{A}^k(X)$, exhaustive ($F^0\mathcal{A}^k(X) = \mathcal{A}^k(X)$), and bounded (it terminates, since $F^p\mathcal{A}^k(X) = 0$ once $p > k$).

A filtered complex has an associated spectral sequence. This is the machinery developed in [7, Convergence for a Decreasing Filtration of a Cochain Complex]: to a complex with a bounded, exhaustive, decreasing filtration $F^\bullet$ one associates a sequence of bigraded pages $(E_r^{p,q}, d_r)$, each the cohomology of the previous with respect to a differential d_r of bidegree $(r, 1-r)$, converging to the associated graded of the cohomology of the total complex with respect to the induced filtration. The construction and its convergence are proved there in full; the content of this section is the identification of the pages in the present geometric situation. The convention places the filtration index p as the first (column) coordinate and the complementary index $q = k - p$ as the second (row) coordinate, so that the total degree $p + q = k$ is the de Rham degree.

The zeroth page reads off directly. By definition of the associated graded of a filtration, $E_0^{p,q} = F^p\mathcal{A}^{p+q}(X)/F^{p+1}\mathcal{A}^{p+q}(X)$, which is the single bidegree slot $\mathcal{A}^{p,q}(X)$: passing to the quotient discards every component of holomorphic degree $> p$ and every component of holomorphic degree $< p$ was already excluded by F^p. The induced differential d_0 is the part of d that preserves the filtration level p while raising total degree, namely the part that raises the antiholomorphic degree. That part is $\bar{\partial}$. So the zeroth page is the Dolbeault complex, column by column, and its cohomology is the Dolbeault cohomology. This is the geometric heart of the whole construction, and it gives the first page.

Theorem 1.2.2: The Frölicher Spectral Sequence

Let X be a complex manifold. The Hodge filtration (definition 1.2.1) on the de Rham complex $(\mathcal{A}^\bullet(X), d)$ gives rise to a spectral sequence, the *Frölicher spectral sequence*, whose first page is the Dolbeault cohomology,

$$E_1^{p,q} = H_{\bar{\partial}}^{p,q}(X)$$

with first differential $d_1 \colon E_1^{p,q} \to E_1^{p+1,q}$ induced by ∂, and which converges to the complex de Rham cohomology:

$$E_1^{p,q} = H_{\bar{\partial}}^{p,q}(X) \implies H_{\mathrm{dR}}^{p+q}(X, \mathbb{C})$$

The abutment is filtered by the Hodge filtration induced on $H_{\mathrm{dR}}^k(X, \mathbb{C})$, and $E_\infty^{p,q} \cong \mathrm{gr}_F^p H_{\mathrm{dR}}^{p+q}(X, \mathbb{C})$.

***Proof*:**
The general construction of the spectral sequence of a bounded filtered complex, together with its convergence to the associated graded of the total cohomology, is proved in [7, Convergence for a Decreasing Filtration of a Cochain Complex]. That machinery applies verbatim to the bounded exhaustive decreasing filtration $F^\bullet$ of definition 1.2.1. What remains is to identify the first page and its differential in the present situation.

Step 1: The zeroth page is the Dolbeault complex. By the definition of the zeroth page of the spectral sequence of a filtered complex ([7, The Exact Couple of a Filtered Complex]), $E_0^{p,q}$ is the associated graded piece $F^p\mathcal{A}^{p+q}(X)/F^{p+1}\mathcal{A}^{p+q}(X)$. By definition 1.2.1 the numerator is the sum of all bidegree slots of total degree $p+q$ with holomorphic degree at least p, and the denominator is the sum of those with holomorphic degree at least $p+1$; the quotient is the single slot of bidegree (p,q), so

$$E_0^{p,q} = \mathcal{A}^{p,q}(X)$$

The differential d_0 is the map induced by d on the associated graded; it has bidegree $(0,1)$ in the (p,q) indexing, so it raises total degree by one while preserving the filtration level p. Write $d = \partial + \bar\partial$ (proposition 1.1.1). On a form of bidegree (p,q) the summand ∂ lands in bidegree $(p+1,q)$, hence in F^{p+1}, and so vanishes in the quotient defining $E_0^{p,q+1}$; the summand $\bar\partial$ lands in bidegree $(p,q+1)$ and survives. Therefore $d_0 = \bar\partial$, and the p-th column of the zeroth page is the Dolbeault complex

$$\mathcal{A}^{p,0}(X) \xrightarrow{\bar\partial} \mathcal{A}^{p,1}(X) \xrightarrow{\bar\partial} \mathcal{A}^{p,2}(X) \xrightarrow{\bar\partial} \cdots$$

of proposition 1.1.1.

Step 2: The first page is Dolbeault cohomology. The first page is the cohomology of (E_0, d_0). By Step 1 this is the cohomology of the Dolbeault complex in bidegree (p,q), which is $H^{p,q}_{\bar\partial}(X)$ by proposition 1.1.1. Hence

$$E_1^{p,q} = H^{p,q}_{\bar\partial}(X)$$

Step 3: The first differential is induced by ∂. The differential $d_1 \colon E_1^{p,q} \to E_1^{p+1,q}$ has bidegree $(1,0)$ and is computed as follows. Represent a class in $E_1^{p,q}$ by a $\bar\partial$-closed form $\alpha \in \mathcal{A}^{p,q}(X)$. Then $d\alpha = \partial\alpha + \bar\partial\alpha = \partial\alpha$ since α is $\bar\partial$-closed, and $\partial\alpha$ lies in $\mathcal{A}^{p+1,q}(X) \subseteq F^{p+1}$. The recipe for d_1 in the spectral sequence of a filtered complex, which is that on representatives the differential is induced by d ([7, Explicit Description of the Pages], carried to cochain complexes by [7, The Cochain Form of the Section]), sends the class of α to the class of $d\alpha = \partial\alpha$ in $E_1^{p+1,q} = H^{p+1,q}_{\bar\partial}(X)$. This is well defined precisely because $\partial\alpha$ is $\bar\partial$-closed: $\bar\partial\partial\alpha = -\partial\bar\partial\alpha = 0$ using the identity $\partial\bar\partial = -\bar\partial\partial$ from proposition 1.1.1. Thus d_1 is the map $[\alpha]_{\bar\partial} \mapsto [\partial\alpha]_{\bar\partial}$ induced by ∂ on Dolbeault cohomology.

Step 4: Convergence. The filtration $F^\bullet$ is exhaustive and bounded, so by [7, Convergence for a Decreasing Filtration of a Cochain Complex] the spectral sequence converges: on each bidegree the pages stabilize to a limit term $E_\infty^{p,q}$, and this limit is the associated graded of the total cohomology $H^{p+q}_{\mathrm{dR}}(X,\mathbb{C})$ for the filtration induced by $F^\bullet$. Explicitly, the filtration induced on $H^k_{\mathrm{dR}}(X,\mathbb{C})$ is $F^pH^k_{\mathrm{dR}}(X,\mathbb{C}) = \mathrm{im}\big(H^k(F^p\mathcal{A}^\bullet(X)) \to H^k_{\mathrm{dR}}(X,\mathbb{C})\big)$, and $E_\infty^{p,q} \cong \mathrm{gr}_F^p H^{p+q}_{\mathrm{dR}}(X,\mathbb{C})$, completing the identification and the proof.

The theorem is a bookkeeping device with sharp content. It says that de Rham cohomology is assembled from Dolbeault cohomology by a succession of correction steps: start from $\bigoplus_{p+q=k} H^{p,q}_{\bar\partial}(X)$, take the cohomology of the differential d_1 induced by ∂ to get the second page, then the cohomology

of d_2, and so on, until the pages stabilize; the surviving graded pieces reassemble $H^k_{\mathrm{dR}}(X,\mathbb{C})$. Each differential d_r measures a way in which a Dolbeault class fails to extend to a genuine de Rham class, or in which two Dolbeault classes become identified in de Rham cohomology. On a general complex manifold these higher differentials can be nonzero, and then the Dolbeault numbers over-count the de Rham cohomology. The extreme opposite case, when every d_r with $r \geq 1$ vanishes, is called degeneration at E_1; there the first page already equals the limit page, and the bigrading passes intact to cohomology.

The distinction between degeneration and its failure is not academic. The first differential d_1 is the map $\partial\colon H^{p,q}_{\bar\partial}(X) \to H^{p+1,q}_{\bar\partial}(X)$, and there exist compact complex manifolds where it is nonzero, so the Frölicher spectral sequence does not degenerate for them. The Iwasawa manifold, a compact quotient of a complex Heisenberg group of complex dimension 3, is the standard example: a direct computation of its Dolbeault cohomology exhibits a nonvanishing d_1, so its first Betti number is strictly smaller than $h^{1,0} + h^{0,1}$ (see [20] for the full calculation on the Iwasawa manifold). Such manifolds are necessarily non-Kähler, and the reason is exactly the degeneration statement below.

Once the abutment is filtered with $E_\infty^{p,q}$ as graded pieces, counting dimensions across a fixed total degree gives an inequality between Betti numbers and Hodge numbers. This is the first hard consequence of the spectral sequence and the numerical form of the entire theory.

Corollary 1.2.3: The Frölicher Inequality

Let X be a compact complex manifold, and write $b_k = \dim_{\mathbb{C}} H^k_{\mathrm{dR}}(X,\mathbb{C})$ for its k-th Betti number and $h^{p,q} = \dim_{\mathbb{C}} H^{p,q}_{\bar\partial}(X)$ for its Hodge numbers. Then

$$b_k \leq \sum_{p+q=k} h^{p,q}$$

for every k, with equality for all k if and only if the Frölicher spectral sequence of X degenerates at E_1.

***Proof*:**

Fix k. On a compact complex manifold each Dolbeault group $H^{p,q}_{\bar\partial}(X)$ is finite dimensional (theorem 1.1.3, and the Betti numbers are likewise finite), so every dimension below is a well-defined nonnegative integer.

Step 1: Each page cannot grow. For $r \geq 1$ the page E_{r+1} is the cohomology of (E_r, d_r), so in each bidegree $E^{p,q}_{r+1}$ is a subquotient of $E^{p,q}_r$, giving $\dim_{\mathbb{C}} E^{p,q}_{r+1} \leq \dim_{\mathbb{C}} E^{p,q}_r$. Passing to the limit, $\dim_{\mathbb{C}} E^{p,q}_\infty \leq \dim_{\mathbb{C}} E^{p,q}_1 = h^{p,q}$.

Step 2: The limit page computes the Betti number. By theorem 1.2.2 the limit terms are the graded pieces of a finite filtration of $H^k_{\mathrm{dR}}(X,\mathbb{C})$, so

$$b_k = \dim_{\mathbb{C}} H^k_{\mathrm{dR}}(X,\mathbb{C}) = \sum_{p+q=k} \dim_{\mathbb{C}} E^{p,q}_\infty$$

Combining with Step 1,

$$b_k = \sum_{p+q=k} \dim_{\mathbb{C}} E^{p,q}_\infty \leq \sum_{p+q=k} \dim_{\mathbb{C}} E^{p,q}_1 = \sum_{p+q=k} h^{p,q}$$

which is the asserted inequality.

Step 3: The equality criterion. Equality holds in the displayed line, for a fixed k, if and only if $\dim_{\mathbb{C}} E_\infty^{p,q} = \dim_{\mathbb{C}} E_1^{p,q}$ for every (p,q) with $p+q=k$ (a sum of nonnegative terms each bounded by the corresponding term of another sum is equal to it precisely when the terms match). By Step 1, the chain of inequalities $\dim_{\mathbb{C}} E_1^{p,q} \geq \dim_{\mathbb{C}} E_2^{p,q} \geq \cdots \geq \dim_{\mathbb{C}} E_\infty^{p,q}$ collapses to equalities throughout exactly when each map does not drop dimension, that is when every d_r $(r \geq 1)$ is zero at $E_r^{p,q}$ and at the source $E_r^{p-r,q+r-1}$ of the incoming differential. Requiring equality for all k forces every d_r on the entire page to vanish for all $r \geq 1$, which is the definition of degeneration at E_1; conversely if the sequence degenerates at E_1 then $E_\infty = E_1$ and equality holds in every total degree. This is the claimed equivalence, completing the proof.

The Frölicher inequality is the first quantitative link between the analytic bigrading and the topology of X. Read from left to right it bounds a topological invariant, the Betti number, by a sum of analytic invariants, the Hodge numbers; the gap $\sum_{p+q=k} h^{p,q} - b_k$ measures the total drop caused by the higher differentials. Read as an equality it becomes a strong constraint on the manifold: equality in every degree is the assertion that the entire spectral sequence collapses immediately, so that the Hodge numbers add up to the Betti numbers with no cancellation. The parity consequence is immediate and worth recording: if the sequence degenerates and moreover Hodge symmetry $h^{p,q} = h^{q,p}$ holds, then $b_{2k+1} = \sum_{p+q=2k+1} h^{p,q}$ is a sum of terms that pair off, so every odd Betti number is even. Both hypotheses, degeneration and Hodge symmetry, are given by the Kähler condition, and this is the route by which chapter 2 deduces that odd Betti numbers of compact Kähler manifolds are even.

The equality case is not exotic. It holds for the class of manifolds this book is built around.

1.2.4 Foreshadowing: Degeneration on Compact Kähler Manifolds If X is a compact Kähler manifold, then the Frölicher spectral sequence degenerates at E_1, and hence $b_k = \sum_{p+q=k} h^{p,q}$ for every k (equality in the Frölicher inequality, corollary 1.2.3). The proof rests on the Hodge decomposition, which produces an actual direct-sum splitting $H^k_{\mathrm{dR}}(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ of the abutment realizing the Hodge filtration, forcing every higher differential to vanish. It is proved in chapter 2 (corollary 2.1.3) from the master identity of section 1.3 and the harmonic theory of theorem 1.1.10.

Degeneration is genuine information about X, not a formal fact. Its failure on the Iwasawa manifold shows that it can fail, and the dimension there is not incidental: non-degeneration cannot occur on a compact complex *surface*, whose Frölicher spectral sequence always degenerates at E_1 (chapter 2); its success on compact Kähler manifolds is a theorem requiring the metric input of the Kähler identities, proved in chapter 2 beside the Hodge decomposition. On projective space the numerology can be checked by hand, and it already satisfies the equality.

Example 1.2.5: Hodge Numbers and Degeneration for $\mathbb{C}\mathrm{P}^n$

Let $X = \mathbb{C}\mathrm{P}^n$, of complex dimension n. Its complex cohomology is well known from [30]: $H^{2j}_{\mathrm{dR}}(\mathbb{C}\mathrm{P}^n,\mathbb{C}) = \mathbb{C}$ for $0 \leq j \leq n$, generated by the j-th power of the hyperplane class, and $H^{\mathrm{odd}}_{\mathrm{dR}}(\mathbb{C}\mathrm{P}^n,\mathbb{C}) = 0$. Thus the Betti numbers are

$$b_k = \begin{cases} 1 & k \text{ even},\ 0 \leq k \leq 2n \\ 0 & k \text{ odd} \end{cases}$$

The Dolbeault numbers $h^{p,q} = \dim_{\mathbb{C}} H^{p,q}_{\bar\partial}(\mathbb{C}\mathrm{P}^n) = \dim_{\mathbb{C}} H^q(\mathbb{C}\mathrm{P}^n, \Omega^p_{\mathbb{C}\mathrm{P}^n})$ (via Dolbeault's theorem

theorem 1.1.3) were computed for $\mathbb{C}\mathrm{P}^1$ in example 1.1.4; the pattern for all n is

$$h^{p,q} = \begin{cases} 1 & 0 \le p = q \le n \\ 0 & p \neq q \end{cases}$$

so the Dolbeault cohomology is concentrated on the diagonal, one copy of $\mathbb{C}$ in each bidegree (p,p) for $0 \le p \le n$ and zero off the diagonal. This is the algebraic-geometry computation $H^q(\mathbb{C}\mathrm{P}^n, \Omega^p) = \mathbb{C}$ for $p = q$ and 0 otherwise, carried out in [4, Bott Vanishing] through the Bott vanishing / Euler-sequence method, and it agrees with the analytic Dolbeault computation by GAGA (theorem 1.2.6) and Dolbeault's theorem.

Now compare the two sides of the Frölicher inequality. Fix k. The right-hand sum is

$$\sum_{p+q=k} h^{p,q} = \begin{cases} h^{k/2,k/2} = 1 & k \text{ even},\ 0 \le k \le 2n \\ 0 & k \text{ odd} \end{cases}$$

because the only nonzero Hodge number with $p+q=k$ occurs when k is even and $p = q = k/2$. This is exactly b_k in every degree, so

$$b_k = \sum_{p+q=k} h^{p,q} \qquad \text{for all } k$$

Equality holds throughout, and by corollary 1.2.3 the Frölicher spectral sequence of $\mathbb{C}\mathrm{P}^n$ degenerates at E_1. This is consistent with box 1.2.4, since $\mathbb{C}\mathrm{P}^n$ carries the Fubini-Study Kähler metric (theorem 1.1.7). Here the degeneration is read off from the numbers. Those numbers reach this book through GAGA (theorem 1.2.6), whose proof in [4, Serre's GAGA Theorems] takes the case of $\mathcal{O}$ on $\mathbb{C}\mathrm{P}^n$ from the Hodge decomposition, so the example does not bypass harmonic theory. The reason there is no room for a higher differential is structural: with the cohomology supported on the diagonal $p = q$, the differential $d_1 \colon E_1^{p,q} \to E_1^{p+1,q}$ maps a nonzero group $H^{p,p}$ into $H^{p+1,p}$, which is zero, and likewise every d_r $(r \ge 1)$ has zero target, so all higher differentials vanish for want of a place to land.

The example is the cleanest instance of degeneration: the spectral sequence collapses not because of a metric argument but because the Dolbeault cohomology is so sparse that no nonzero differential can exist. The same diagonal concentration recurs for every smooth projective toric variety and, more generally, wherever the Hodge numbers off the diagonal vanish. It also anchors the numerology used throughout the book: the Hodge diamond of $\mathbb{C}\mathrm{P}^n$ is a single string of 1's down the diagonal, and every later Hodge diamond is measured against this baseline.

The Dolbeault numbers of $\mathbb{C}\mathrm{P}^n$ were quoted from two sources that agree, the analytic Dolbeault computation and the algebraic sheaf-cohomology computation $H^q(\mathbb{C}\mathrm{P}^n, \Omega^p_{\mathbb{C}\mathrm{P}^n})$. That they agree is a theorem, GAGA, which asserts the agreement for every smooth projective variety. This is the bridge that lets Hodge theory, an analytic subject, compute invariants of algebraic varieties, and lets algebraic geometry give the Hodge numbers that feed the Frölicher inequality.

The setup requires a word on the two categories being compared. A smooth projective variety X over $\mathbb{C}$ carries two structure sheaves and two cohomology theories. Algebraically, X is a scheme of finite type over $\mathbb{C}$ with its Zariski topology and its sheaf $\mathcal{O}_X^{\mathrm{alg}}$ of regular functions, and one has the coherent sheaf cohomology $H^q(X^{\mathrm{alg}}, \Omega^p_{X^{\mathrm{alg}}})$ of algebraic Kähler differentials. Analytically, the same variety has an underlying complex manifold X^{an} (the analytification), with its classical topology and

holomorphic structure sheaf $\mathcal{O}_{X^{\mathrm{an}}}$, and the sheaf cohomology $H^q(X^{\mathrm{an}}, \Omega^p_{X^{\mathrm{an}}})$ of holomorphic p-forms (proposition 1.1.1) is the analytic object identified with Dolbeault cohomology in theorem 1.1.3. There is always a comparison map from the algebraic groups to the analytic groups, induced by the map of ringed spaces $X^{\mathrm{an}} \to X^{\mathrm{alg}}$. GAGA says this map is an isomorphism.

Theorem 1.2.6: GAGA for Coherent Cohomology

Let X be a smooth projective variety over $\mathbb{C}$, with analytification X^{an} the associated compact complex manifold. Then for all $p, q \geq 0$ the comparison map

$$H^q(X^{\mathrm{alg}}, \Omega^p_{X^{\mathrm{alg}}}) \xrightarrow{\sim} H^q(X^{\mathrm{an}}, \Omega^p_{X^{\mathrm{an}}})$$

from algebraic to analytic coherent cohomology is an isomorphism. In particular the algebraic and analytic Hodge numbers agree, so the Dolbeault numbers $h^{p,q}$ of X^{an} may be computed by algebraic geometry.

***Proof*:**
This is Serre's GAGA theorem in the form comparing coherent cohomology of a projective variety with that of its analytification. It is stated, with a proof sketch, in [4, Serre's GAGA Theorems]: for a projective scheme X over $\mathbb{C}$ and any coherent algebraic sheaf $\mathcal{F}$, analytification $\mathcal{F} \mapsto \mathcal{F}^{\mathrm{an}}$ is an equivalence between coherent algebraic and coherent analytic sheaves, and the natural maps $H^q(X^{\mathrm{alg}}, \mathcal{F}) \to H^q(X^{\mathrm{an}}, \mathcal{F}^{\mathrm{an}})$ are isomorphisms for all q. The differential sheaves $\Omega^p_{X^{\mathrm{alg}}}$ are coherent, and their analytifications are the holomorphic differential sheaves $\Omega^p_{X^{\mathrm{an}}}$, so the general statement specializes to the one above. The sketch of the cohomology comparison starts from $\mathcal{O}$ on $\mathbb{C}\mathrm{P}^n$, whose analytic cohomology it takes from the Hodge decomposition and the Dolbeault resolution of $\mathbb{C}\mathrm{P}^n$, and extends to all coherent sheaves through hyperplane sequences, quotients of sums of twisted line bundles, and closed embeddings. The analytic input to the equivalence of categories is deferred to [36].

GAGA is the reason the transcendental theory of this book is a theory about algebraic varieties and not only about abstract complex manifolds. It has two payoffs used repeatedly. First, computability: the analytic Hodge numbers of a projective variety, hard to reach through partial differential equations, become algebraic quantities computable by sheaf cohomology, Serre duality, and exact sequences, exactly as the $h^{p,q}$ of $\mathbb{C}\mathrm{P}^n$ were read off above. Second, rigidity: a holomorphic object on a projective variety, such as a holomorphic form or a coherent subsheaf, is automatically algebraic, so the analytic and algebraic worlds have the same coherent geometry. The Lefschetz $(1,1)$-theorem and the Hodge conjecture, which ask whether certain analytic cohomology classes are algebraic, live in the space GAGA opens up: GAGA settles the coherent-sheaf half of the question, and Hodge theory addresses the topological half.

One boundary is worth marking. GAGA compares coherent cohomology, the groups $H^q(X, \Omega^p_X)$; it does not by itself give the Hodge decomposition of the topological cohomology $H^k(X, \mathbb{C})$. That decomposition is the content of chapter 2 and requires the Kähler metric. What GAGA and the Frölicher spectral sequence contribute is the input and the framework: the Dolbeault numbers (computed algebraically, via GAGA) and their relationship to the Betti numbers (organized by the spectral sequence). Degeneration, proved by harmonic theory in chapter 2, is the missing statement that turns the Frölicher inequality into the Hodge decomposition.

Exercise 1.2.1

1. Verify directly that the Hodge filtration $F^\bullet$ of definition 1.2.1 is a filtration of the de Rham complex, that is, that $d(F^p\mathcal{A}^k(X)) \subseteq F^p\mathcal{A}^{k+1}(X)$ for all p, k. Then show it is bounded on each $\mathcal{A}^k(X)$ by exhibiting the smallest and largest p for which $F^p\mathcal{A}^k(X)$ is nonzero.

2. Let X be a compact Riemann surface (complex dimension 1) of genus g. Its Betti numbers are $b_0 = b_2 = 1$ and $b_1 = 2g$, and its Hodge numbers are $h^{0,0} = h^{1,1} = 1$, $h^{1,0} = h^{0,1} = g$. Verify the Frölicher inequality in every degree and conclude that the spectral sequence degenerates at E_1 for every compact Riemann surface, with no appeal to the Kähler condition.

3. Show that on a compact complex manifold with $h^{p,q} = 0$ whenever $p \neq q$, the Frölicher spectral sequence degenerates at E_1. (This is the structural reason $\mathbb{C}\mathrm{P}^n$ degenerates in example 1.2.5; isolate the argument that no nonzero differential can exist.)

4. The Hodge-de Rham (or Hirzebruch) alternating sum is $\chi(X) = \sum_k (-1)^k b_k$ and the analytic Euler characteristic in each column is $\chi(X, \Omega^p) = \sum_q (-1)^q h^{p,q}$. Show that regardless of whether the Frölicher spectral sequence degenerates, one has $\sum_k (-1)^k b_k = \sum_{p,q} (-1)^{p+q} h^{p,q}$, so that the topological Euler characteristic equals the alternating sum of all Hodge numbers.

5. Use GAGA (theorem 1.2.6) together with Serre duality on a smooth projective variety X of dimension n to prove the duality $h^{p,q} = h^{n-p,n-q}$ among the Hodge numbers. State clearly which input is analytic and which is algebraic.

6. A Hopf surface is a compact complex surface diffeomorphic to $S^1 \times S^3$; it is not Kähler. From the diffeomorphism type its Betti numbers are $b_0 = b_4 = 1$, $b_1 = b_3 = 1$, $b_2 = 0$. Explain why the odd Betti number $b_1 = 1$ shows, via corollary 1.2.3 and box 1.2.4, that either the Frölicher spectral sequence of a Hopf surface fails to degenerate at E_1 or Hodge symmetry $h^{p,q} = h^{q,p}$ fails. (In fact it is Hodge symmetry that fails: $h^{1,0} = 0$ but $h^{0,1} = 1$. The spectral sequence does degenerate, as it does for every compact complex surface.)

1.3 The Master Identity

The first consequence of the Kähler identities (theorem 1.1.9) is the master identity relating the three Laplacians. On a general Hermitian manifold Δ_d, $2\Delta_\partial$ and $2\Delta_{\bar\partial}$ have the same principal symbol and differ by lower-order torsion terms. The Kähler identities force them to be proportional, with Δ_∂ and $\Delta_{\bar\partial}$ equal and Δ_d their sum.

Theorem 1.3.1: The Master Identity

On a Kähler manifold (X, ω) the three Laplacians coincide up to a factor of two:

$$\Delta_d = 2\,\Delta_{\bar\partial} = 2\,\Delta_\partial$$

Moreover $\Delta_{\bar\partial}$ commutes with the Lefschetz operators L and Λ, and it preserves the bidegree, so it acts on each space $\mathcal{A}^{p,q}(X)$.

***Proof*:**
The proof is a sequence of algebraic manipulations of the Kähler identities of theorem 1.1.9, with no further analysis.

Step 1: $\Delta_\partial = \Delta_{\bar\partial}$. Expand $\Delta_\partial = \partial\partial^* + \partial^*\partial$ using the Kähler identity $\partial^* = i[\Lambda, \bar\partial] = i(\Lambda\bar\partial - \bar\partial\Lambda)$ from (1.1):

$$\begin{aligned}\Delta_\partial &= \partial\big(i(\Lambda\bar\partial - \bar\partial\Lambda)\big) + \big(i(\Lambda\bar\partial - \bar\partial\Lambda)\big)\partial \\ &= i\big(\partial\Lambda\bar\partial - \partial\bar\partial\Lambda + \Lambda\bar\partial\partial - \bar\partial\Lambda\partial\big) \\ &= i\big(\partial\Lambda\bar\partial - \bar\partial\Lambda\partial\big) + i\big(\Lambda\bar\partial\partial - \partial\bar\partial\Lambda\big)\end{aligned}$$

For $\Delta_{\bar\partial}$, expand with the conjugate Kähler identity $\bar\partial^* = -i[\Lambda, \partial] = -i(\Lambda\partial - \partial\Lambda)$:

$$\begin{aligned}\Delta_{\bar\partial} &= \bar\partial\bar\partial^* + \bar\partial^*\bar\partial = \bar\partial\big(-i(\Lambda\partial - \partial\Lambda)\big) + \big(-i(\Lambda\partial - \partial\Lambda)\big)\bar\partial \\ &= -i\big(\bar\partial\Lambda\partial - \bar\partial\partial\Lambda + \Lambda\partial\bar\partial - \partial\Lambda\bar\partial\big) \\ &= -i\big(\bar\partial\Lambda\partial - \partial\Lambda\bar\partial\big) - i\big(\Lambda\partial\bar\partial - \bar\partial\partial\Lambda\big)\end{aligned}$$

Compare the two expressions grouping by grouping, using $\partial\bar\partial = -\bar\partial\partial$ from proposition 1.1.1 throughout. The first grouping of Δ_∂ is $i(\partial\Lambda\bar\partial - \bar\partial\Lambda\partial)$, and the first grouping of $\Delta_{\bar\partial}$ is $-i(\bar\partial\Lambda\partial - \partial\Lambda\bar\partial) = i(\partial\Lambda\bar\partial - \bar\partial\Lambda\partial)$; these are equal. For the second groupings, $\Lambda\bar\partial\partial = -\Lambda\partial\bar\partial$ and $\partial\bar\partial\Lambda = -\bar\partial\partial\Lambda$, so the second grouping of Δ_∂ is

$$i(\Lambda\bar\partial\partial - \partial\bar\partial\Lambda) = i(-\Lambda\partial\bar\partial + \bar\partial\partial\Lambda) = -i(\Lambda\partial\bar\partial - \bar\partial\partial\Lambda)$$

which is exactly the second grouping of $\Delta_{\bar\partial}$. Both groupings agree, so

$$\Delta_\partial = \Delta_{\bar\partial}$$

Step 2: $\Delta_d = \Delta_\partial + \Delta_{\bar\partial}$. Expand Δ_d using $d = \partial + \bar\partial$ from proposition 1.1.1 and $d^* = \partial^* + \bar\partial^*$:

$$\Delta_d = dd^* + d^*d = (\partial + \bar\partial)(\partial^* + \bar\partial^*) + (\partial^* + \bar\partial^*)(\partial + \bar\partial)$$

Multiplying out gives $\Delta_\partial + \Delta_{\bar\partial}$ (the pure terms) plus the four cross terms

$$(\partial\bar\partial^* + \bar\partial^*\partial) + (\bar\partial\partial^* + \partial^*\bar\partial)$$

The claim is that both parenthesized cross terms vanish. Consider $\partial\bar\partial^* + \bar\partial^*\partial$. Substitute $\bar\partial^* = -i(\Lambda\partial - \partial\Lambda)$:

$$\partial\bar\partial^* + \bar\partial^*\partial = -i\big(\partial\Lambda\partial - \partial\partial\Lambda + \Lambda\partial\partial - \partial\Lambda\partial\big)$$

Now $\partial\partial = 0$ (this is $\partial^2 = 0$, the holomorphic half of proposition 1.1.1), so the two middle terms vanish, and the first and last terms cancel:

$$\partial\bar\partial^* + \bar\partial^*\partial = -i\big(\partial\Lambda\partial - 0 + 0 - \partial\Lambda\partial\big) = 0$$

The other cross term $\bar\partial\partial^* + \partial^*\bar\partial$ vanishes by the analogous computation, using $\partial^* = i(\Lambda\bar\partial - \bar\partial\Lambda)$ and $\bar\partial^2 = 0$ (exercise 1.3.1 (1)). Hence

$$\Delta_d = \Delta_\partial + \Delta_{\bar\partial} = 2\,\Delta_{\bar\partial}$$

the last equality by Step 1. Combining Steps 1 and 2 gives $\Delta_d = 2\Delta_{\bar\partial} = 2\Delta_\partial$.

Step 3: $\Delta_{\bar\partial}$ commutes with L and Λ, and preserves bidegree. Since $\Delta_{\bar\partial} = \frac{1}{2}\Delta_d$ and $\Delta_d = dd^* + d^*d$ preserves total degree $k = p + q$, and since $\Delta_{\bar\partial} = \bar\partial\bar\partial^* + \bar\partial^*\bar\partial$ preserves the antiholomorphic degree q (both $\bar\partial\bar\partial^*$ and $\bar\partial^*\bar\partial$ map $\mathcal{A}^{p,q}$ to itself, as $\bar\partial$ raises q by one and $\bar\partial^*$ lowers it by one), the operator $\Delta_{\bar\partial}$ preserves both k and q, hence preserves (p, q). For the commutation with L: the Kähler identities give $[L, \partial^*] = i\bar\partial$ and $[L, \bar\partial^*] = -i\partial$, while L commutes with ∂ and $\bar\partial$ because $L\alpha = \omega \wedge \alpha$ with ω closed, so that $d(\omega \wedge \alpha) = \omega \wedge d\alpha$ and separating types gives $\partial L = L\partial$, $\bar\partial L = L\bar\partial$. Then

$$[L, \Delta_{\bar\partial}] = [L, \bar\partial\bar\partial^* + \bar\partial^*\bar\partial] = \bar\partial[L, \bar\partial^*] + [L, \bar\partial^*]\bar\partial = \bar\partial(-i\partial) + (-i\partial)\bar\partial = -i(\bar\partial\partial + \partial\bar\partial) = 0$$

using $[L, \bar\partial] = 0$ to pull L through $\bar\partial$ and $\bar\partial\partial + \partial\bar\partial = 0$ from proposition 1.1.1 at the end. For the commutation with Λ, take formal L^2 adjoints of $[L, \Delta_{\bar\partial}] = 0$. Since $L^* = \Lambda$ and $\Delta_{\bar\partial}$ is self-adjoint (it is $\bar\partial\bar\partial^* + \bar\partial^*\bar\partial$, a sum of an operator with its own adjoint), the adjoint of the vanishing commutator is $[\Delta_{\bar\partial}, \Lambda] = -[\Lambda, \Delta_{\bar\partial}] = 0$, so $[\Lambda, \Delta_{\bar\partial}] = 0$ as well. This completes the proof.

The factor of two in $\Delta_d = 2\Delta_{\bar\partial}$ says that harmonicity for the de Rham complex and harmonicity for the Dolbeault complex are the same condition on a Kähler manifold: a form is Δ_d-harmonic if and only if it is $\Delta_{\bar\partial}$-harmonic. Since $\Delta_{\bar\partial}$ preserves the bidegree, the space of harmonic k-forms splits as a direct sum of harmonic (p, q)-forms with $p + q = k$. Feeding this into the harmonic representation of cohomology yields the Hodge decomposition $H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ of chapter 2.

That $\Delta_{\bar\partial}$ commutes with L and Λ is the seed of the Hard Lefschetz theorem. It means the $\mathfrak{sl}_2$-action generated by (L, Λ) descends from forms to harmonic forms, hence to cohomology; the Lefschetz decomposition of cohomology into primitive pieces is then a matter of representation theory of $\mathfrak{sl}_2$. That development belongs to chapter 2 and beyond, but the commutation established here is what makes it possible.

Exercise 1.3.1

1. In the proof of theorem 1.3.1, Step 2, the cross term $\partial\bar\partial^* + \bar\partial^*\partial$ was shown to vanish using $\partial^2 = 0$. Carry out the analogous computation for the other cross term $\bar\partial\partial^* + \partial^*\bar\partial$ in full, substituting $\partial^* = i(\Lambda\bar\partial - \bar\partial\Lambda)$ and using $\bar\partial^2 = 0$, and confirm it too vanishes.

2. Deduce from theorem 1.3.1 that on a compact Kähler manifold a d-harmonic k-form α (that is, $\Delta_d\alpha = 0$) decomposes as $\alpha = \sum_{p+q=k} \alpha^{p,q}$ with each component $\alpha^{p,q}$ itself Δ_d-harmonic. State clearly which property of $\Delta_{\bar\partial}$ from the theorem you use, and why Δ_d-harmonicity of α passes to each $\alpha^{p,q}$.

3. Show directly, from the Kähler identities and the closedness of ω, that the Lefschetz operator L commutes with Δ_∂ as well (not only with $\Delta_{\bar\partial}$). Explain why this is consistent with theorem 1.3.1.

4. The $\bar\partial$-Hodge theorem is stated for a compact Hermitian manifold, with no Kähler hypothesis, yet the Hodge decomposition of section 2.1 requires Kähler. Identify precisely which statement of theorem 1.1.10 would fail on a general compact Hermitian manifold if one tried to replace $\Delta_{\bar\partial}$ throughout by Δ_d, and name the property of the three Laplacians that is available only in the Kähler case.

5. Show that complex conjugation carries $\bar\partial$-harmonic (p, q)-forms to ∂-harmonic (q, p)-forms,

where $\Delta_\partial = \partial\partial^* + \partial^*\partial$ and a form is ∂-harmonic when Δ_∂ annihilates it. On a compact *Kähler* manifold, deduce using theorem 1.3.1 that $\overline{\mathcal{H}^{p,q}_{\bar\partial}(X)} = \mathcal{H}^{q,p}_{\bar\partial}(X)$, and interpret this as a symmetry of the Hodge numbers.

1.4 The $\partial\bar{\partial}$-Lemma and a Non-Kähler Surface

The master identity $\Delta_d = 2\Delta_{\bar\partial} = 2\Delta_\partial$ of theorem 1.3.1 is the analytic heart of Kähler geometry, but its cohomological force is not yet visible: a statement about Laplacians becomes a statement about forms once harmonic representatives enter, and theorem 1.1.10 supplies them. This section extracts the first form-level consequence of the equality of Laplacians, the $\partial\bar{\partial}$-lemma, and puts it to work in two directions. In the positive direction it tightens the relation between the various notions of exactness on a compact Kähler manifold, collapsing d-exactness, ∂-exactness, $\bar\partial$-exactness, and $\partial\bar\partial$-exactness into a single condition for forms of pure type. In the negative direction it becomes a test: a compact complex manifold on which the lemma fails cannot carry any Kähler metric. The Hopf surface gives the first example in the book of a compact complex manifold that is not Kähler, and it does so through the topological consequence of the lemma, the constraint that odd Betti numbers be even.

Projective manifolds are the source of positive examples: every one of them is Kähler (theorem 1.1.7), so every one of them obeys the $\partial\bar\partial$-lemma and has even odd Betti numbers. These constraints are not automatic for compact complex manifolds. Complex structures are abundant, Kähler complex structures are not, and the Hopf surface is the standard first example of the gap.

The statement of the lemma requires no new machinery beyond the operators ∂ and $\bar\partial$ and the decomposition $d = \partial + \bar\partial$ (proposition 1.1.1). A form α of type (p, q) is *of pure type*; a general k-form decomposes into its pure-type components, and the lemma is about the pure-type pieces.

Theorem 1.4.1: The $\partial\bar{\partial}$-Lemma

Let X be a compact Kähler manifold and let α be a form of pure type (p, q) on X. Suppose α is d-closed. Then the following conditions are equivalent:

1. α is d-exact;
2. α is ∂-exact;
3. α is $\bar\partial$-exact;
4. $\alpha = \partial\bar\partial\gamma$ for some form γ of type $(p-1, q-1)$.

In particular, a d-exact form of pure type is $\partial\bar\partial$-exact.

The proof refines the harmonic decomposition of theorem 1.1.10 to three operators at once. The master identity $\Delta_d = 2\Delta_{\bar\partial} = 2\Delta_\partial$ gives the three Laplacians one harmonic space and one Green operator, and that compatibility is what makes the four exactness conditions coincide.

***Proof*:**
Fix the Kähler metric on X and use the associated L^2 inner product on forms. By theorem 1.3.1 the three Laplacians coincide up to scalar, $\Delta_d = 2\Delta_{\bar\partial} = 2\Delta_\partial$, so they share a kernel: on each

bidegree (p,q) there is a single harmonic space $\mathcal{H}^{p,q}(X)$, and a single Green operator G inverting the common Laplacian on the orthogonal complement of the harmonics. Because $\partial^*, \bar\partial^*, \partial, \bar\partial$ each commute with the common Laplacian on a Kähler manifold, they commute with G. These commutation facts are the only input beyond the harmonic decomposition of theorem 1.1.10.

Step 1: Reduce to a $\bar\partial$-primitive. Suppose first that α is d-exact. The harmonic projection $\mathcal{H}(\alpha)$ vanishes: a harmonic form is L^2-orthogonal to every exact form, since for harmonic η and $\alpha = d\rho$ one has $\langle \eta, d\rho\rangle = \langle d^*\eta, \rho\rangle = 0$. The $\bar\partial$-Hodge decomposition of theorem 1.1.10 then gives, using G,

$$\alpha = \mathcal{H}(\alpha) + \bar\partial\bar\partial^* G\alpha + \bar\partial^*\bar\partial G\alpha = \bar\partial\bar\partial^* G\alpha + \bar\partial^*\bar\partial G\alpha$$

The two terms are the components of α in the orthogonal summands $\mathrm{im}\bar\partial$ and $\mathrm{im}\bar\partial^*$. Now $\bar\partial\alpha = 0$, so $\alpha \in \ker\bar\partial = \mathcal{H}^{p,q}(X) \oplus \mathrm{im}\bar\partial$, which is orthogonal to $\mathrm{im}\bar\partial^*$; hence the $\mathrm{im}\bar\partial^*$-component vanishes, $\bar\partial^*\bar\partial G\alpha = 0$. Therefore

$$\alpha = \bar\partial\gamma, \qquad \gamma := \bar\partial^* G\alpha \in \mathcal{A}^{p,q-1}(X)$$

Step 2: γ is ∂-closed, and decomposing it gives a potential. First, γ is ∂-closed. The vanishing of the cross term $\partial\bar\partial^* + \bar\partial^*\partial$, established in Step 2 of the proof of theorem 1.3.1, together with $\bar\partial^*$ commuting with G and $\partial\alpha = 0$, gives

$$\partial\gamma = \partial\bar\partial^* G\alpha = -\bar\partial^*\partial G\alpha = -\bar\partial^* G\,\partial\alpha = 0$$

Now decompose γ by the ∂-Hodge theorem, which is available because $\Delta_\partial = \Delta_{\bar\partial}$ shares the same harmonic space and Green operator G:

$$\gamma = \mathcal{H}(\gamma) + \partial\partial^* G\gamma + \partial^*\partial G\gamma$$

The three terms are the components of γ in $\mathcal{H}^{p,q-1}(X)$, $\mathrm{im}\partial$, and $\mathrm{im}\partial^*$. Since γ is ∂-closed, $\gamma \in \ker\partial = \mathcal{H}^{p,q-1}(X) \oplus \mathrm{im}\partial$, which is orthogonal to $\mathrm{im}\partial^*$; so the last component vanishes, $\partial^*\partial G\gamma = 0$, and $\gamma = \mathcal{H}(\gamma) + \partial\partial^* G\gamma$. Apply $\bar\partial$: the harmonic term is annihilated by $\bar\partial$ (a form harmonic for the common Laplacian is killed by both ∂ and $\bar\partial$), so

$$\alpha = \bar\partial\gamma = \bar\partial\partial\partial^* G\gamma = -\partial\bar\partial\partial^* G\gamma$$

using $\bar\partial\partial = -\partial\bar\partial$. Set $\beta' := -\partial^* G\gamma$. Its bidegree is $(p-1, q-1)$, since γ has bidegree $(p, q-1)$ and ∂^* lowers p by one while G preserves bidegree. Then

$$\alpha = \partial\bar\partial\beta'$$

which is the required expression with $\beta = \beta'$.

Step 3: The other exactness hypotheses. If α is $\bar\partial$-exact rather than d-exact, then since α is also ∂-closed it is orthogonal to the harmonics (a $\bar\partial$-exact form is orthogonal to $\ker\Delta_{\bar\partial} = \mathcal{H}^{p,q}(X)$), so $\mathcal{H}(\alpha) = 0$ and Steps 1 and 2 apply verbatim. If α is ∂-exact, conjugate: $\bar\alpha$ is $\bar\partial$-exact of type (q,p), the previous case yields $\bar\alpha = \partial\bar\partial\beta''$, and conjugating back gives $\alpha = \bar\partial\partial\overline{\beta''} = \partial\bar\partial\big(-\overline{\beta''}\big)$. In every case α is $\partial\bar\partial$-exact, completing the proof.

On a compact Kähler manifold the only obstruction to writing a pure-type exact form as $\partial\bar\partial$ of something is measured by harmonic forms, which vanish for exact classes. In chapter 3 the lemma underlies the fact that a variation of Hodge structure is controlled by the Hodge filtration alone. Its

failure on a non-Kähler manifold accounts for the behaviour of the Frölicher spectral sequence there.

1.4.2 Warning: the Kähler hypothesis Every step of the $\partial\bar\partial$-lemma used $\Delta_\partial = \Delta_{\bar\partial}$, which is the Kähler condition (theorem 1.3.1). On a compact complex manifold that is not Kähler the three Laplacians differ, their harmonic spaces separate, and the conclusion can fail: the Hopf surface $S = (\mathbb{C}^2 \setminus \{0\})/\langle z \mapsto 2z\rangle$, diffeomorphic to $S^1 \times S^3$, has $b_1(S) = 1$. An odd first Betti number is incompatible with the Hodge symmetry $h^{1,0} = h^{0,1}$ (which would force $b_1 = 2h^{1,0}$), so S is not Kähler, its Hodge decomposition fails, and with it the $\partial\bar\partial$-lemma. The lemma and the decomposition require the Kähler condition together.

The lemma is a statement about the difference between two a priori distinct facts. That $\alpha = \partial\bar\partial\gamma$ implies α is d-exact is immediate, since $\partial\bar\partial = \frac{1}{2}\,d(\bar\partial - \partial)$ on a pure-type form (using $d = \partial + \bar\partial$ and $\bar\partial\partial = -\partial\bar\partial$ from proposition 1.1.1, whence $d(\bar\partial - \partial) = (\partial + \bar\partial)(\bar\partial - \partial) = \partial\bar\partial - \bar\partial\partial = 2\partial\bar\partial$, using $\partial^2 = \bar\partial^2 = 0$); the content is the reverse implication, that a form which is exact for the total differential is exact for the finer operator $\partial\bar\partial$. On a general compact complex manifold this reverse implication is false, and its failure is obstructed at the level of Bott-Chern and Aeppli cohomology, which measure precisely the gap between the finer and coarser notions of exactness. On a compact Kähler manifold it holds, and this is what allows Hodge theory to read off holomorphic data from topological data.

A first consequence records what the lemma says about closed forms of type $(1,1)$, the type of a Kähler form. Two cohomologous real closed $(1,1)$-forms differ by a d-exact real $(1,1)$-form, and the lemma turns that difference into $\partial\bar\partial$ of a single global function. This is the analytic origin of the notion of a Kähler potential.

Corollary 1.4.3: Cohomologous Closed $(1,1)$-Forms Differ by a Potential

Let X be a compact Kähler manifold and let ω, ω' be two real closed $(1,1)$-forms on X with $[\omega] = [\omega']$ in $H^2(X, \mathbb{R})$. Then there is a real-valued function $\varphi \in \mathcal{C}^\infty(X, \mathbb{R})$ with

$$\omega' - \omega = i\,\partial\bar\partial\varphi$$

***Proof*:**
Since $[\omega] = [\omega']$ the difference $\beta := \omega' - \omega$ is a real d-exact form of type $(1,1)$, hence d-closed of pure type. By theorem 1.4.1 there is a form γ of type $(0,0)$, that is a function $\gamma \in \mathcal{C}^\infty(X, \mathbb{C})$, with $\beta = \partial\bar\partial\gamma$. It remains to arrange that the potential is real and that the constant i appears. Because β is real, $\beta = \overline{\beta}$; taking conjugates and using $\overline{\partial\bar\partial\gamma} = \bar\partial\partial\overline{\gamma} = -\partial\bar\partial\overline{\gamma}$ (by proposition 1.1.1) gives $\beta = -\partial\bar\partial\overline{\gamma}$. Averaging the two expressions for β,

$$\beta = \tfrac{1}{2}\big(\partial\bar\partial\gamma - \partial\bar\partial\overline{\gamma}\big) = \partial\bar\partial\Big(\tfrac{\gamma - \overline{\gamma}}{2}\Big)$$

The function $\frac{1}{2}(\gamma - \overline{\gamma})$ is purely imaginary, so it equals $i\varphi$ for a real-valued $\varphi := \frac{1}{2i}(\gamma - \overline{\gamma}) \in \mathcal{C}^\infty(X, \mathbb{R})$. Substituting, $\beta = \partial\bar\partial(i\varphi) = i\,\partial\bar\partial\varphi$, which is the claimed identity, as we sought to show.

The corollary says that on a compact Kähler manifold the Kähler class $[\omega] \in H^2(X, \mathbb{R})$ determines the Kähler form up to the addition of $i\,\partial\bar\partial\varphi$; the space of Kähler forms in a fixed class is, in this

precise sense, a space of potentials. This is the starting point for the study of canonical metrics: fixing the class and searching within it for a form satisfying a curvature condition becomes a scalar partial differential equation for φ, which is how the Calabi conjecture and the existence of Kähler-Einstein metrics are formulated. That translation from a form-level problem to a scalar equation is only possible because the lemma reduces the freedom in a Kähler class to a single function.

A second consequence works in complementary direction, extracting holomorphic rather than metric information. A holomorphic p-form is a $(p,0)$-form annihilated by $\bar{\partial}$; on an arbitrary compact complex manifold such a form need not be closed for the full differential d, since ∂ of it may be nonzero. On a compact Kähler manifold it is always closed, and the proof again routes through the harmonic theory that the Kähler condition tames.

Corollary 1.4.4: Holomorphic Forms Are Closed

Let X be a compact Kähler manifold and let α be a holomorphic p-form, that is a global section of Ω^p_X regarded as a $(p,0)$-form with $\bar{\partial}\alpha = 0$ (proposition 1.1.1). Then $d\alpha = 0$; that is, $\partial\alpha = 0$ as well.

***Proof*:**

Write $d\alpha = \partial\alpha + \bar{\partial}\alpha$. By hypothesis $\bar{\partial}\alpha = 0$, so $d\alpha = \partial\alpha$, a form of type $(p+1,0)$. The claim is that $\partial\alpha = 0$.

Step 1: α is $\bar{\partial}$-harmonic. By theorem 1.3.1, $\Delta_{\bar{\partial}} = \frac{1}{2}\Delta_d$. A holomorphic p-form satisfies $\bar{\partial}\alpha = 0$; it also satisfies $\bar{\partial}^*\alpha = 0$ for type reasons, since $\bar{\partial}^*$ lowers antiholomorphic degree and α has antiholomorphic degree 0, so $\bar{\partial}^*\alpha$ would be a form of antiholomorphic degree -1, necessarily zero. Hence $\Delta_{\bar{\partial}}\alpha = (\bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial})\alpha = 0$, so α is $\bar{\partial}$-harmonic.

Step 2: α is d-harmonic, hence d-closed. By the master identity, $\Delta_d\alpha = 2\Delta_{\bar{\partial}}\alpha = 0$, so α is harmonic for Δ_d. On a compact Riemannian manifold a form is Δ_d-harmonic if and only if it is both d-closed and d^*-closed ([12, harmonic equals closed and coclosed]): the identity $\langle\langle \Delta_d\alpha, \alpha\rangle\rangle = \|d\alpha\|^2 + \|d^*\alpha\|^2$ (integration by parts using the formal adjoint) forces $d\alpha = 0$ and $d^*\alpha = 0$ when the left side vanishes. In particular $d\alpha = 0$, and since $d\alpha = \partial\alpha$ this gives $\partial\alpha = 0$, completing the proof.

The corollary is the mechanism behind one half of the Hodge decomposition: it says the holomorphic p-forms, defined by the purely holomorphic condition $\bar{\partial}\alpha = 0$, automatically represent de Rham cohomology classes, so there is a map $H^0(X, \Omega^p_X) \to H^p(X, \mathbb{C})$. On a compact Kähler manifold this map is injective and lands in the $(p,0)$-part of the Hodge decomposition, identifying $H^{p,0}$ with the space of global holomorphic p-forms. The Kähler hypothesis is doing real work: on a general compact complex manifold a global holomorphic form can fail to be closed, and the identification breaks. The Iwasawa manifold, the quotient of the complex Heisenberg group of upper-triangular unipotent 3×3 matrices by its lattice of Gaussian-integer entries, is the standard example: it carries a holomorphic 1-form ω_3 with $d\omega_3 = -\omega_1 \wedge \omega_2 \neq 0$, so $H^{1,0}$ is strictly larger than the space of closed holomorphic 1-forms and cannot inject into H^1. The Hopf surface below fails to be Kähler for a different, purely topological reason, and its own holomorphic forms are too scarce to display this particular breakdown.

Beyond these form-level consequences, the lemma also has a topological consequence that requires none of its analytic content to state, only the Hodge decomposition it feeds into. On a compact Kähler manifold the decomposition $H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ together with the symmetry $H^{p,q} = \overline{H^{q,p}}$

forces every odd Betti number to be even; for k odd the summands pair off into complex-conjugate pairs $H^{p,q} \oplus H^{q,p}$ with $p \neq q$, and $b_k = \sum_{p+q=k} h^{p,q}$ is a sum of equal pairs. This is proved in full in chapter 2; what matters here is its contrapositive as an obstruction.

Proposition 1.4.5: Odd Betti Obstruction

If X is a compact complex manifold admitting some Kähler metric, then every odd Betti number $b_{2k+1}(X)$ is even. Equivalently, a compact complex manifold with an odd Betti number that is odd carries no Kähler metric.

***Proof*:**

Suppose X carries a Kähler metric. By the Hodge decomposition (chapter 2), $H^{2k+1}(X, \mathbb{C}) = \bigoplus_{p+q=2k+1} H^{p,q}(X)$ with $H^{p,q} = \overline{H^{q,p}}$, so $h^{p,q} = h^{q,p}$. Since $p+q = 2k+1$ is odd, no summand has $p = q$; the indices pair off into distinct pairs $\{(p,q),(q,p)\}$, and

$$b_{2k+1}(X) = \sum_{p+q=2k+1} h^{p,q} = 2 \sum_{\substack{p+q=2k+1 \\ p<q}} h^{p,q}$$

which is even. The contrapositive is the stated obstruction, completing the proof.

The Frölicher inequality of corollary 1.2.3 already gives a form of this: it bounds $b_k \leq \sum_{p+q=k} h^{p,q}$, with equality precisely when the Frölicher spectral sequence degenerates at E_1. The Kähler condition guarantees degeneration (this is one of the payoffs of chapter 2), so on a Kähler manifold the inequality is an equality and the conjugation symmetry then forces the parity constraint. For a general compact complex manifold neither the degeneration nor the symmetry is available, and b_k can carry the "wrong" parity. The Hopf surface exhibits exactly this.

The Hopf surface is built as a quotient of $\mathbb{C}^2 \setminus \{0\}$ by a single contracting biholomorphism. It is the standard example of a compact complex surface with no Kähler metric, and the reason is visible at the level of its underlying topology: it is diffeomorphic to $S^1 \times S^3$, whose first Betti number is 1, which is odd.

Example 1.4.6: The Hopf Surface

Fix the biholomorphism $f\colon \mathbb{C}^2 \setminus \{0\} \to \mathbb{C}^2 \setminus \{0\}$ given by $f(z) = 2z$, and let $\Gamma = \langle f \rangle \cong \mathbb{Z}$ be the infinite cyclic group it generates. The *Hopf surface* is the quotient

$$S = (\mathbb{C}^2 \setminus \{0\})/\Gamma$$

Its structure is unwound in three steps.

Step 1: S is a compact complex manifold of dimension 2. The map f is a biholomorphism of $\mathbb{C}^2 \setminus \{0\}$, and the group $\Gamma \cong \mathbb{Z}$ acts holomorphically. A free, properly discontinuous action of a discrete group by biholomorphisms has a quotient that is again a complex manifold, the charts descending from local sections of the projection ([32, Quotient Manifold Theorem] for the smooth structure; holomorphy of the transition maps is inherited because the deck transformations are biholomorphic). Both hypotheses hold here. The action is free: $f^m(z) = 2^m z = z$ with $z \neq 0$ forces $2^m = 1$, hence $m = 0$. It is properly discontinuous: any $z \neq 0$ has a neighborhood, for instance the open annulus $U = \{w : \frac{1}{\sqrt{2}}|z| < |w| < \sqrt{2}\,|z|\}$, meeting only finitely many of its Γ-translates. Indeed $f^m(U) = \{w : 2^m \frac{1}{\sqrt{2}}|z| < |w| < 2^m\sqrt{2}\,|z|\}$ is the shell of radii $(2^{m-1/2}|z|,\, 2^{m+1/2}|z|)$,

whose outer radius $2^{m+1/2}|z|$ equals the inner radius of the next shell $f^{m+1}(U)$; the open shells therefore meet at most along excluded boundary spheres and are pairwise disjoint, so U meets only $f^0(U) = U$ itself. A free, properly discontinuous holomorphic action of a discrete group yields a complex manifold quotient with the quotient holomorphic structure, so S is a complex manifold of dimension $\dim_{\mathbb{C}} S = 2$. Compactness is settled in the next step by exhibiting a diffeomorphism to a compact manifold.

Step 2: S is diffeomorphic to $S^1 \times S^3$. Introduce the radial-logarithmic map

$$\Phi\colon \mathbb{C}^2 \setminus \{0\} \to S^3 \times \mathbb{R}, \qquad \Phi(z) = \left(\tfrac{z}{|z|},\ \log_2 |z|\right)$$

where $S^3 = \{z \in \mathbb{C}^2 : |z| = 1\}$ is the unit sphere and $|z| = (|z_1|^2 + |z_2|^2)^{1/2}$. This Φ is a diffeomorphism, with inverse $(u, t) \mapsto 2^t u$. Under Φ the generator $f(z) = 2z$ becomes $(u, t) \mapsto (u, t + 1)$, since $|2z| = 2|z|$ gives $\log_2 |2z| = \log_2 |z| + 1$ and $\frac{2z}{|2z|} = \frac{z}{|z|}$. Thus Φ conjugates the Γ-action to the action of $\mathbb{Z}$ on $S^3 \times \mathbb{R}$ by translation in the second factor, and

$$S = (\mathbb{C}^2 \setminus \{0\})/\Gamma \ \cong\ (S^3 \times \mathbb{R})/\mathbb{Z} \ \cong\ S^3 \times (\mathbb{R}/\mathbb{Z}) = S^3 \times S^1$$

where $\mathbb{Z}$ acts only on the $\mathbb{R}$ factor. Hence S is diffeomorphic to $S^1 \times S^3$, which is compact.

Step 3: $b_1(S) = 1$, so S admits no Kähler metric. The Künneth theorem ([30, Künneth Formula for Cohomology]) gives $H^1(S^1 \times S^3, \mathbb{R}) \cong H^1(S^1, \mathbb{R}) \oplus H^1(S^3, \mathbb{R})$. Since $H^1(S^1, \mathbb{R}) \cong \mathbb{R}$ and $H^1(S^3, \mathbb{R}) = 0$ (the sphere S^3 is 2-connected), $b_1(S) = 1$. This odd Betti number is odd, so by proposition 1.4.5 the surface S carries no Kähler metric. Equivalently, via corollary 1.2.3: any Kähler structure would force $b_1 = h^{1,0} + h^{0,1}$ with $h^{1,0} = h^{0,1}$, hence b_1 even, contradicting $b_1 = 1$.

The Fubini-Study metric makes $\mathbb{C}\mathrm{P}^n$ and every projective manifold Kähler (theorem 1.1.7), and the abundance of projective examples might suggest that the Kähler condition is nearly automatic. The Hopf surface shows it is not. It is a compact complex manifold built from elementary data, a scaling of $\mathbb{C}^2$, yet no metric on it can be Kähler, because its topology already forbids the Hodge symmetry that a Kähler metric would impose. The obstruction is a parity: $b_1(S) = 1$.

The contrast sharpens the meaning of the projective examples. A smooth projective variety inherits its Kähler form by restriction from the ambient $\mathbb{C}\mathrm{P}^N$ (theorem 1.1.7), and this reflects the fact that projective varieties satisfy every Kähler constraint, including even odd Betti numbers and the $\partial\bar{\partial}$-lemma. The Hopf surface, by contrast, is never projective, because a projective manifold is Kähler and S is not. It is the model non-algebraic compact complex manifold, and it marks the boundary of the theory this book develops: the Hodge decomposition of chapter 2 applies to Kähler manifolds, and the Hopf surface is the reminder that outside that class the decomposition, and everything built on it, can fail.

Exercise 1.4.1

1. Let X be any complex manifold (Kähler or not) and let γ be a form of type $(p-1, q-1)$. Show directly, using only $d = \partial + \bar{\partial}$ and the relations of proposition 1.1.1, that $\alpha := \partial\bar{\partial}\gamma$ is d-closed. This is the "easy direction" of the $\partial\bar{\partial}$-lemma and needs no Kähler hypothesis.

2. On a compact Kähler manifold X, corollary 1.4.3 produces a potential φ with $\omega' - \omega = i\,\partial\bar{\partial}\varphi$. Determine the extent to which φ is unique: describe all real functions ψ with $i\,\partial\bar{\partial}\psi = 0$, and conclude that φ is unique up to an additive real constant. (Use that a global function

annihilated by $\partial\bar{\partial}$ is pluriharmonic, hence harmonic for the underlying Riemannian metric, and that a harmonic function on a compact connected manifold is constant.)

3. On the Hopf surface $S = (\mathbb{C}^2 \setminus \{0\})/\Gamma$, consider the holomorphic 2-form $\eta = dz_1 \wedge dz_2$ on $\mathbb{C}^2 \setminus \{0\}$. Show that η is *not* Γ-invariant, and find the scalar by which $f^*\eta$ differs from η. Explain why this shows S has no nonzero global holomorphic 2-form, so that $h^{2,0}(S) = 0$. Note that corollary 1.4.4 has nothing to say here, but for the structural reason that S is not Kähler, not because $H^0(S, \Omega_S^2) = 0$.

4. The complex torus $\mathbb{C}^2/\Lambda$ of example 1.1.6 (with Λ a lattice of rank 4) is diffeomorphic to $(S^1)^4$ and carries a flat Kähler metric. Compute b_1 and b_3 and verify they are even, as proposition 1.4.5 requires. Contrast the outcome with the Hopf surface: both are quotients of an open subset of $\mathbb{C}^2$, yet only one is Kähler; identify which topological feature distinguishes them.

5. Assume, as a direct Dolbeault computation, that the Hopf surface has Hodge numbers $h^{1,0} = 0$ and $h^{0,1} = 1$. Using $b_1 = 1$, show that the Frölicher inequality corollary 1.2.3 holds here as an equality $b_1 = h^{1,0} + h^{0,1}$ even though S is not Kähler. Explain why this does not contradict proposition 1.4.5: which of the two ingredients (Frölicher degeneration, Hodge symmetry $h^{p,q} = h^{q,p}$) fails on S?

6. Deduce from proposition 1.4.5 that the Hopf surface S is not projective, and more generally that no compact complex manifold with an odd first Betti number can be embedded as a complex submanifold of any $\mathbb{C}\mathrm{P}^N$. Which result of section 1.1 does this argument combine with proposition 1.4.5 to reach the conclusion?

2

The Hodge Decomposition

Chapter 1 produced the master identity $\Delta_d = 2\Delta_{\bar\partial}$ on a Kähler manifold. On a compact manifold every de Rham class has a unique Δ_d-harmonic representative, by the Hodge theorem of *Riemannian Geometry* [12], and every Dolbeault class has a unique $\Delta_{\bar\partial}$-harmonic one (theorem 1.1.10). The master identity makes the two harmonic spaces equal, so the (p,q)-grading of forms passes to cohomology: $H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ with $H^{p,q} = \overline{H^{q,p}}$. The consequences are the degeneration of the Frölicher spectral sequence, the parity of the odd Betti numbers, the Hodge index theorem on surfaces, and the Lefschetz $(1,1)$-theorem, which identifies the integral classes of type $(1,1)$ with the classes of divisors.

2.1 The Hodge Decomposition

The $\bar\partial$-Hodge theorem (theorem 1.1.10) produces, on a compact Hermitian manifold, a unique $\bar\partial$-harmonic representative for each Dolbeault class, so that $\mathcal{H}^{p,q}_{\bar\partial}(X) \cong H^q(X,\Omega^p_X)$. That statement uses no relation between the $\bar\partial$-theory and the underlying real de Rham theory; the two live on separate footings, one bigraded by type and one graded by total degree. On a compact Kähler manifold the master identity $\Delta_d = 2\Delta_{\bar\partial}$ of theorem 1.3.1 fuses them. A form is d-harmonic exactly when it is $\bar\partial$-harmonic, and $\Delta_{\bar\partial}$ preserves bidegree, so the real harmonic space acquires a bigrading it had no reason to carry. Transported through the two Hodge isomorphisms, that bigrading becomes a splitting of complex cohomology by type. This section proves that splitting and the symmetry it enjoys under conjugation, and settles the degeneration of the Frölicher spectral sequence that section 1.2 deferred.

Throughout, X is a compact Kähler manifold of complex dimension n, with a fixed Kähler metric. We write $\mathcal{H}^k(X)$ for the space of real (Δ_d-)harmonic k-forms with complex coefficients and $\mathcal{H}^{p,q}_{\bar\partial}(X)$ for the $\Delta_{\bar\partial}$-harmonic (p,q)-forms of theorem 1.1.10.

The first observation is that the two Laplacians, agreeing up to the scalar 2, have the same kernel. This is what lets the bidegree decomposition of forms descend to harmonic forms.

Theorem 2.1.1: Hodge Decomposition

Let X be a compact Kähler manifold of complex dimension n. Then the space of complex-valued harmonic k-forms decomposes by bidegree,

$$\mathcal{H}^k(X) = \bigoplus_{p+q=k} \mathcal{H}^{p,q}_{\bar\partial}(X)$$

and this induces a direct sum decomposition of complex cohomology,

$$H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X) \qquad \text{where} \qquad H^{p,q}(X) \cong \mathcal{H}^{p,q}_{\bar\partial}(X) \cong H^q(X,\Omega^p_X)$$

The summand $H^{p,q}(X)$ is the image in $H^k(X,\mathbb{C})$ of the $\bar\partial$-harmonic (p,q)-forms. Moreover the decomposition satisfies the *Hodge symmetry*

$$H^{p,q}(X) = \overline{H^{q,p}(X)}$$

where the bar denotes complex conjugation on $H^k(X,\mathbb{C}) = H^k(X,\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$.

***Proof*:**

Step 1: The two Laplacians share a kernel. By theorem 1.3.1 the identity $\Delta_d = 2\Delta_{\bar\partial}$ holds on every form on X. A form α satisfies $\Delta_d\alpha = 0$ if and only if $\Delta_{\bar\partial}\alpha = 0$, so the d-harmonic forms and the $\bar\partial$-harmonic forms coincide as subspaces of the smooth complex-valued forms. The real Hodge-de Rham theorem, applied to the operator Δ_d on the closed manifold X and cited to [12, The Hodge theorem], identifies $\mathcal{H}^k(X)$ with $H^k(X,\mathbb{C})$: every de Rham class has a unique harmonic representative.

Step 2: Harmonic forms split by type. A smooth complex k-form decomposes uniquely into its components of each bidegree, $\alpha = \sum_{p+q=k} \alpha^{p,q}$ with $\alpha^{p,q} \in \mathcal{A}^{p,q}(X)$. The operator $\Delta_{\bar\partial} = \bar\partial\bar\partial^* + \bar\partial^*\bar\partial$ preserves bidegree, since $\bar\partial$ raises q by one and $\bar\partial^*$ lowers it by one and neither changes p. Hence

$$\Delta_{\bar\partial}\alpha = \sum_{p+q=k} \Delta_{\bar\partial}\big(\alpha^{p,q}\big)$$

and, the summands lying in distinct bidegrees, $\Delta_{\bar\partial}\alpha = 0$ holds if and only if $\Delta_{\bar\partial}\alpha^{p,q} = 0$ for every (p,q). So a form is $\bar\partial$-harmonic exactly when each of its bihomogeneous components is $\bar\partial$-harmonic. Combined with Step 1,

$$\mathcal{H}^k(X) = \bigoplus_{p+q=k} \mathcal{H}^{p,q}_{\bar\partial}(X)$$

the sum being direct because the bidegree components of a form are unique.

Step 3: Passing to cohomology. Define $H^{p,q}(X) \subseteq H^k(X,\mathbb{C})$ as the image, under the isomorphism $\mathcal{H}^k(X) \cong H^k(X,\mathbb{C})$ of Step 1, of the summand $\mathcal{H}^{p,q}_{\bar\partial}(X)$. Applying that isomorphism to the

direct sum of Step 2 gives

$$H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$$

By construction $H^{p,q}(X) \cong \mathcal{H}^{p,q}_{\bar\partial}(X)$, and the $\bar\partial$-Hodge theorem theorem 1.1.10 together with Dolbeault's theorem theorem 1.1.3 identifies $\mathcal{H}^{p,q}_{\bar\partial}(X) \cong H^{p,q}_{\bar\partial}(X) \cong H^q(X,\Omega^p_X)$. This proves the first two displays.

Step 4: Hodge symmetry. Complex conjugation acts on complex-valued forms, sending $\mathcal{A}^{p,q}(X)$ to $\mathcal{A}^{q,p}(X)$: if $\alpha = f\,dz_{i_1}\wedge\cdots\wedge d\bar z_{j_q}$ has type (p,q), then $\bar\alpha$ has type (q,p). Conjugation interchanges the two Dolbeault differentials, $\overline{\bar\partial\alpha} = \partial\bar\alpha$, hence interchanges their formal adjoints and the associated Laplacians, $\overline{\Delta_{\bar\partial}\alpha} = \Delta_\partial\bar\alpha$. By theorem 1.3.1 the Kähler condition forces $\Delta_\partial = \Delta_{\bar\partial}$, so

$$\overline{\Delta_{\bar\partial}\alpha} = \Delta_{\bar\partial}\bar\alpha$$

Thus α is $\bar\partial$-harmonic if and only if $\bar\alpha$ is, and conjugation restricts to a conjugate-linear isomorphism $\mathcal{H}^{p,q}_{\bar\partial}(X) \to \mathcal{H}^{q,p}_{\bar\partial}(X)$. On cohomology this is exactly the map induced by conjugation on $H^k(X,\mathbb{C})$, which carries $H^k(X,\mathbb{R})$ to itself. Therefore $H^{p,q}(X) = \overline{H^{q,p}(X)}$, completing the proof.

The decomposition combines the holomorphic structure of complex manifolds, Dolbeault theory, and the Kähler identities, all recalled in section 1.1, so that a purely topological invariant, the complex cohomology of X, remembers the complex geometry. A class in $H^k(X,\mathbb{C})$ is no longer an unstructured object; it carries a canonical expression $\sum_{p+q=k}\alpha^{p,q}$ into pieces of definite type, and each piece $H^{p,q}(X)$ is a sheaf-cohomology group $H^q(X,\Omega^p_X)$ computable by algebraic means. The two isomorphisms in the theorem run in opposite directions: $H^{p,q}\cong\mathcal{H}^{p,q}_{\bar\partial}$ is analytic, produced by harmonic theory, while $\mathcal{H}^{p,q}_{\bar\partial}\cong H^q(X,\Omega^p_X)$ is the algebraic reading through Dolbeault. That the same group admits both descriptions is the reason Hodge theory transfers information between the transcendental and algebraic sides of complex geometry.

Two features stand out before an example. First, the splitting depends on the Kähler metric only through the identity $\Delta_d = 2\Delta_{\bar\partial}$; the resulting subspaces $H^{p,q}(X)\subseteq H^k(X,\mathbb{C})$ are independent of the chosen Kähler metric. Abstract isomorphism with the metric-free group $H^q(X,\Omega^p_X)$ is not what shows this, since it fixes only the dimension of the subspace and not its position inside $H^k(X,\mathbb{C})$; what does is the metric-free characterization $H^{p,q}(X) = \{[\alpha] : \alpha \text{ a } d\text{-closed form of type } (p,q)\}$, equivalently $H^{p,q} = F^p\cap\overline{F^q}$ for the Hodge filtration of definition 1.2.1. Second, the Hodge symmetry $H^{p,q} = \overline{H^{q,p}}$ is what obstructs an arbitrary complex manifold from being Kähler: it forces constraints on the Hodge numbers, taken up in section 2.2.

The complex torus illustrates the decomposition concretely, since every cohomology class has a translation-invariant harmonic representative and the decomposition can be written down by hand.

Example 2.1.2: The Hodge Diamond of a Complex Torus

Let $X = \mathbb{C}^n/\Lambda$ be a complex torus of dimension n (example 1.1.6), carrying the flat Kähler metric descended from the standard Hermitian metric on $\mathbb{C}^n$. Write $z_1,\dots,z_n$ for the linear complex coordinates on $\mathbb{C}^n$; the forms $dz_1,\dots,dz_n,d\bar z_1,\dots,d\bar z_n$ are translation-invariant and descend to global forms on X.

For the flat metric the Laplacian $\Delta_{\bar\partial}$ has constant coefficients, and a (p,q)-form with constant coefficients has $\bar\partial$ and $\bar\partial^*$ both zero, hence is harmonic. Conversely a harmonic form on the compact

torus is translation-invariant, so

$$\mathcal{H}^{p,q}_{\bar\partial}(X) = \Big\{ \sum a_{IJ}\, dz_I \wedge d\bar z_J : a_{IJ} \in \mathbb{C},\ |I| = p,\ |J| = q \Big\}$$

where I, J range over increasing multi-indices. The dimension is $h^{p,q} = \binom{n}{p}\binom{n}{q}$, and the Hodge decomposition reads

$$H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X), \qquad \dim H^{p,q}(X) = \binom{n}{p}\binom{n}{q}$$

For $n = 1$ this is an elliptic curve $E = \mathbb{C}/\Lambda$: the diamond has $h^{0,0} = h^{1,1} = 1$ at top and bottom and $h^{1,0} = h^{0,1} = 1$ in the middle row, recovering $b_1 = 2$ with $H^1(E, \mathbb{C}) = H^{1,0} \oplus H^{0,1} = \mathbb{C}\, dz \oplus \mathbb{C}\, d\bar z$. For $n = 2$ the diamond is

$$\begin{array}{ccccc} & & h^{0,0} & & \\ & h^{1,0} & & h^{0,1} & \\ h^{2,0} & & h^{1,1} & & h^{0,2} \\ & h^{2,1} & & h^{1,2} & \\ & & h^{2,2} & & \end{array} \quad = \quad \begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 1 & & 4 & & 1 \\ & 2 & & 2 & \\ & & 1 & & \end{array}$$

so $b_0 = b_4 = 1$, $b_1 = b_3 = 4$, $b_2 = 6$. The total dimension across the diamond is $\sum_{p,q} \binom{2}{p}\binom{2}{q} = (1+1)^2(1+1)^2 = 16 = 2^4$, matching $H^*(X, \mathbb{C}) = \Lambda^*(\mathbb{C}^4)$ for the real four-torus.

Two checks confirm the theorem. The Hodge symmetry $h^{p,q} = h^{q,p}$ is visible in $\binom{n}{p}\binom{n}{q} = \binom{n}{q}\binom{n}{p}$, the diamond being symmetric across its vertical axis. And each row sums to the correct Betti number: $\sum_{p+q=k} \binom{n}{p}\binom{n}{q} = \binom{2n}{k}$ by Vandermonde's identity, which is exactly $b_k = \dim \Lambda^k(\mathbb{C}^{2n})$ for the underlying $2n$-real-dimensional torus.

The torus makes the mechanism transparent because harmonicity reduces to constancy of coefficients, so the abstract $\mathcal{H}^{p,q}_{\bar\partial}$ is a space one can write with a basis. For a general Kähler manifold the harmonic representatives are inaccessible by hand, but the theorem guarantees the same bookkeeping holds: the Betti numbers are recovered as row sums of the Hodge numbers, and the diamond is symmetric under $(p, q) \mapsto (q, p)$.

The decomposition of theorem 2.1.1 settles a question raised in section 1.2. There the Frölicher spectral sequence, abutting to $H^k(X, \mathbb{C})$ from the E_1-page $E_1^{p,q} = H^q(X, \Omega^p_X)$, produced only the inequality $b_k \le \sum_{p+q=k} h^{p,q}$ of corollary 1.2.3, with equality holding if and only if the sequence degenerates at E_1. On a compact Kähler manifold the Hodge decomposition forces equality, hence degeneration.

Corollary 2.1.3: Frölicher Degeneration

On a compact Kähler manifold X, the Frölicher spectral sequence degenerates at E_1: all differentials d_r for $r \ge 1$ vanish, and $E_1^{p,q} = E_\infty^{p,q}$. Equivalently, equality holds in corollary 1.2.3,

$$b_k = \sum_{p+q=k} h^{p,q}$$

for every k.

***Proof*:**
The spectral sequence converges to $H^k(X,\mathbb{C})$ with $E_1^{p,q} = H^q(X,\Omega_X^p)$, so for each k

$$\dim H^k(X,\mathbb{C}) = \sum_{p+q=k} \dim E_\infty^{p,q} \leq \sum_{p+q=k} \dim E_1^{p,q} = \sum_{p+q=k} h^{p,q}$$

the inequality of corollary 1.2.3, arising because each later page is a subquotient of the previous. By theorem 2.1.1 the Kähler manifold satisfies

$$\dim H^k(X,\mathbb{C}) = \sum_{p+q=k} \dim H^{p,q}(X) = \sum_{p+q=k} \dim H^q(X,\Omega_X^p) = \sum_{p+q=k} h^{p,q}$$

So the inequality is an equality for every k. Equality of the total dimensions forces $\dim E_\infty^{p,q} = \dim E_1^{p,q}$ in each bidegree, since $E_\infty^{p,q}$ is a subquotient of $E_1^{p,q}$ and a proper subquotient would strictly drop the dimension in that slot. A subquotient of equal dimension is the whole space, so $E_1^{p,q} = E_\infty^{p,q}$ and every intermediate differential d_r $(r \geq 1)$ vanishes, as we sought to show.

Degeneration at E_1 restates the Hodge decomposition from the spectral-sequence viewpoint: the associated graded of $H^k(X,\mathbb{C})$ under the filtration by holomorphic type is already present on the first page, with no higher differentials to compute. It also gives a metric-free reason the Hodge numbers are cohomological invariants of the Kähler manifold, since $E_1^{p,q} = H^q(X,\Omega_X^p)$ is defined without reference to a metric. For a compact complex manifold that is not Kähler the sequence may fail to degenerate; the Iwasawa manifold (complex dimension 3) is the standard example, its Frölicher spectral sequence having a nonzero d_1. (Non-degeneration cannot occur on a surface: the Frölicher spectral sequence of every compact complex surface degenerates at E_1, Kähler or not.)

Exercise 2.1.1

1. A K3 surface is a compact Kähler surface ($n = 2$) with $h^{1,0} = 0$ and $h^{2,0} = 1$; its remaining independent Hodge number is $h^{1,1} = 20$. Using Hodge symmetry and corollary 2.1.3, fill in the full Hodge diamond and compute the Betti numbers b_0, b_1, b_2, b_3, b_4. Check that $b_2 = 22$ and that the Euler characteristic is 24.

2. Show directly from theorem 2.1.1 that if X is a compact Kähler manifold, then $\dim H^{p,q}(X) = \dim H^{q,p}(X)$, and deduce that $b_{2k+1}(X)$ is even for every k. (This is the discharge of the forward reference; give the two-line argument.)

3. For the complex torus $X = \mathbb{C}^2/\Lambda$ of example 2.1.2, write down an explicit basis of $H^{1,1}(X)$ in terms of the invariant forms $dz_i \wedge d\bar{z}_j$, and confirm $\dim H^{1,1} = 4$. Express the class of the flat Kähler form ω in this basis.

4. The subspaces $H^{p,q}(X) \subseteq H^k(X,\mathbb{C})$ are defined in theorem 2.1.1 using a chosen Kähler metric (through the harmonic representatives). Explain why $H^{p,q}(X)$ is nonetheless independent of the choice of Kähler metric. (Hint: identify $H^{p,q}(X)$ with a group defined without a metric.)

5. Let α be a real d-exact $(1,1)$-form on a compact Kähler manifold X. Use the $\partial\bar{\partial}$-lemma to produce a real function f with $\alpha = i\,\partial\bar{\partial} f$. (Hint: apply theorem 1.4.1 to get $\alpha = \partial\bar{\partial}\beta$, then use that α is real to arrange β of the form if.)

6. Suppose a compact complex manifold Y has $b_1(Y)$ odd. Explain why Y cannot be Kähler, and why the Frölicher spectral sequence of Y and its Hodge symmetry cannot both hold: at least

one of "E_1-degeneration" and "$h^{1,0} = h^{0,1}$" must fail. (No computation is needed; combine corollary 2.1.3 with theorem 2.1.1. Note that b_1 odd does not by itself force non-degeneration: the Hopf surface has $b_1 = 1$ yet its Frölicher spectral sequence degenerates, so it is Hodge symmetry that fails there.)

7. Let X be a compact connected complex manifold of dimension n. Theorem 1.1.10 gives $h^{p,q} = h^{n-p,n-q}$ with no projectivity hypothesis; take it as given. Compute $h^{n,n}(X)$ and interpret the resulting class in $H^{n,n}_{\bar\partial}(X)$.

8. Using the fact that $h^{p,q}(\mathbb{C}\mathrm{P}^n) = 1$ for $p = q$ and 0 otherwise, write out the Hodge diamond of $\mathbb{C}\mathrm{P}^2$, verify that the diagonal sums reproduce the Betti numbers $b_0 = b_2 = b_4 = 1$, $b_1 = b_3 = 0$ of $\mathbb{C}\mathrm{P}^2$, and confirm the two Hodge symmetries $h^{p,q} = h^{q,p}$ and $h^{p,q} = h^{n-p,n-q}$.

2.2 First Consequences

The decomposition $H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ of theorem 2.1.1 is a constraint on the cohomology of any compact Kähler manifold, and its first payoff is a list of relations among the Hodge numbers $h^{p,q} = \dim_{\mathbb{C}} H^{p,q}(X)$ that hold for no reason on a general compact complex manifold. Two symmetries organize the Hodge diamond into its familiar shape, and the parity of the odd Betti numbers, already extracted in chapter 1, falls out again as a special case. Beyond the numerical symmetries lies a statement about the intersection pairing: on a compact Kähler surface the signature of the cup-product form on H^2 is pinned down by the Hodge decomposition alone. This is the Hodge index theorem, the low-degree instance of the polarization theory that chapter 3 develops in general.

Recall from theorem 2.1.1 that complex conjugation on $H^k(X,\mathbb{C}) = H^k(X,\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$ interchanges the summands $H^{p,q}$ and $H^{q,p}$: one has $H^{p,q} = \overline{H^{q,p}}$. Conjugation is a $\mathbb{C}$-antilinear isomorphism, so it carries a $\mathbb{C}$-basis of $H^{q,p}$ to a $\mathbb{C}$-basis of $H^{p,q}$, and the two spaces have the same complex dimension. That is the first symmetry. The second comes from Serre duality, which identifies $H^q(X,\Omega^p_X)$ with the dual of $H^{n-q}(X,\Omega^{n-p}_X)$ through the canonical bundle $\Omega^n_X = \det \Omega^1_X$ [5, Canonical Bundle]. Together they fold the array $(h^{p,q})_{0\le p,q\le n}$ onto itself under two reflections.

Corollary 2.2.1: Hodge Symmetries

Let X be a compact Kähler manifold of complex dimension n. The Hodge numbers $h^{p,q} = \dim_{\mathbb{C}} H^{p,q}(X)$ satisfy

$$h^{p,q} = h^{q,p} \qquad \text{and} \qquad h^{p,q} = h^{n-p,n-q}$$

for all $0 \le p,q \le n$. Consequently each odd Betti number $b_{2k+1}(X) = \sum_{p+q=2k+1} h^{p,q}$ is even.

***Proof*:**

Conjugation symmetry. By theorem 2.1.1 the conjugate-linear map $\alpha \mapsto \bar\alpha$ on $H^k(X,\mathbb{C})$ restricts to a $\mathbb{C}$-antilinear isomorphism $H^{q,p}(X) \xrightarrow{\sim} H^{p,q}(X)$. An antilinear isomorphism preserves complex dimension, so $h^{p,q} = h^{q,p}$.

Duality symmetry. Serre duality as stated in *Algebraic Geometry*[a] assumes projectivity, which a compact Kähler manifold need not have. The form needed is theorem 1.1.10(3), which *Complex Analytic Geometry* proves for every compact connected Hermitian manifold: the conjugate star $\alpha \mapsto \star\bar\alpha$ carries $\bar\partial$-harmonic (p,q)-forms to $\bar\partial$-harmonic $(n-p,n-q)$-forms, and pairing α against

$\star\bar{\alpha}$ gives $\|\alpha\|^2 > 0$. Hence $h^{p,q} = h^{n-p,n-q}$. On a projective X this is the Hodge-theoretic form of Serre duality, but no projectivity was used.
Odd Betti parity. Fix an odd degree $k = 2m+1$. Pair each term $h^{p,q}$ with $p+q=k$ against $h^{q,p}$; since $p \neq q$ when $p+q$ is odd, the summands split into disjoint pairs $\{(p,q),(q,p)\}$, each contributing $2h^{p,q}$ by the conjugation symmetry. Hence $b_k(X) = \sum_{p+q=k} h^{p,q}$ is a sum of even numbers, completing the proof.

[a][4, Serre Duality for a Projective Scheme].

The two symmetries say the Hodge diamond is invariant under reflection across its vertical axis (conjugation) and under the central involution $(p,q) \mapsto (n-p, n-q)$ (Serre duality). For a surface $n = 2$ the diamond is

$$\begin{array}{ccccc} & & h^{2,2} & & \\ & h^{2,1} & & h^{1,2} & \\ h^{2,0} & & h^{1,1} & & h^{0,2} \\ & h^{1,0} & & h^{0,1} & \\ & & h^{0,0} & & \end{array}$$

with $h^{2,2} = h^{0,0} = 1$ (the top and bottom are H^0 and H^{2n} of a connected compact manifold), $h^{1,0} = h^{0,1} = h^{2,1} = h^{1,2}$, and $h^{2,0} = h^{0,2}$; only $h^{1,0}$, $h^{2,0}$, and $h^{1,1}$ are free. The odd Betti parity here reads $b_1 = 2h^{1,0}$ and $b_3 = 2h^{2,1} = 2h^{1,0}$, recovering exactly the obstruction of proposition 1.4.5: that proposition was proved in chapter 1 by citing this decomposition forward, and the forward reference is now discharged. The content is identical, so nothing new is proved; the diamond simply displays the parity as the pairing of a term with its conjugate.

Hodge symmetry is a Kähler phenomenon, not a feature of complex manifolds in general. On the Hopf surface of chapter 1 one has $b_1 = 1$, forcing $h^{1,0} \neq h^{0,1}$, so conjugation symmetry fails; this is one route to seeing that the Hopf surface carries no Kähler metric.

The symmetries above are statements about dimensions. The decomposition also controls the cup-product pairing, and on a surface it does so completely. On a compact oriented real 4-manifold the intersection form

$$Q(\alpha,\beta) = \int_X \alpha \wedge \beta$$

on $H^2(X,\mathbb{R})$ is a nondegenerate symmetric bilinear form (nondegenerate by Poincaré duality[1]), so it has a well-defined signature (b^+, b^-): the number of positive and negative eigenvalues over $\mathbb{R}$. For a smooth oriented 4-manifold the signature is a diffeomorphism invariant, not determined by b_2 alone. The Hodge index theorem says that when X is a compact Kähler surface, the Hodge decomposition forces the value: b^+ and b^- are determined by $h^{2,0}$ and $h^{1,1}$ alone.

The mechanism is the sign of $\int_X \alpha \wedge \bar{\alpha}$ on the pieces of the decomposition. The general statement that positivity of the polarizing form alternates with the Hodge bigrading is the Hodge-Riemann bilinear relations, developed for arbitrary weight in chapter 3. Here the only input needed is the degree-2 case on a surface, which is elementary enough to verify by hand.

[1][30, Consequences of Poincaré Duality].

Lemma 2.2.2: Signs of the Cup Form in Bidegree Two

Let X be a compact Kähler surface with Kähler form ω, and let $Q(\alpha,\beta)=\int_X \alpha\wedge\beta$ be the cup-product form on $H^2(X,\mathbb{R})$, extended $\mathbb{C}$-bilinearly to $H^2(X,\mathbb{C})$. Then:

1. For $0\neq\alpha\in H^{2,0}(X)$ one has $\int_X \alpha\wedge\bar\alpha>0$; the same holds for $\alpha\in H^{0,2}(X)$.
2. The class $[\omega]\in H^{1,1}(X)$ satisfies $\int_X \omega\wedge\omega>0$.
3. On the primitive part $P^{1,1}(X)=\{\,\beta\in H^{1,1}(X)\big|\beta\wedge\omega=0\,\}$ of the real $(1,1)$-classes, one has $\int_X\beta\wedge\beta<0$ for $0\neq\beta$ real.

***Proof*:**
Work with harmonic representatives throughout, so a cohomology class of type (p,q) is represented by a harmonic (p,q)-form. Fix a point and choose a local unitary coframe φ^1,φ^2 of $(1,0)$-forms in which the Kähler form is $\omega=i(\varphi^1\wedge\bar\varphi^1+\varphi^2\wedge\bar\varphi^2)$. The metric volume form is $\mathrm{vol}=\frac{1}{2}\omega^2$, and expanding,

$$\tfrac{1}{2}\omega^2=i^2\,\varphi^1\wedge\bar\varphi^1\wedge\varphi^2\wedge\bar\varphi^2=(i\,\varphi^1\wedge\bar\varphi^1)\wedge(i\,\varphi^2\wedge\bar\varphi^2)$$

so vol is the product of the two positive real 2-forms $i\,\varphi^j\wedge\bar\varphi^j$, and $\int_X f\,\mathrm{vol}>0$ for a positive function f.
Part (1). A nonzero $(2,0)$-form is $\alpha=a\,\varphi^1\wedge\varphi^2$ at each point with a a complex function, so

$$\alpha\wedge\bar\alpha=|a|^2\,\varphi^1\wedge\varphi^2\wedge\bar\varphi^1\wedge\bar\varphi^2$$

Reordering $\varphi^1\wedge\varphi^2\wedge\bar\varphi^1\wedge\bar\varphi^2$ to $\varphi^1\wedge\bar\varphi^1\wedge\varphi^2\wedge\bar\varphi^2$ moves $\bar\varphi^1$ left past φ^2, one transposition, contributing a sign -1; and $\varphi^1\wedge\bar\varphi^1\wedge\varphi^2\wedge\bar\varphi^2=i^{-2}\,\mathrm{vol}=-\mathrm{vol}$. The two signs cancel, giving $\alpha\wedge\bar\alpha=|a|^2\,\mathrm{vol}$ pointwise. Integrating, $\int_X\alpha\wedge\bar\alpha>0$ unless $\alpha\equiv0$. The case $\alpha\in H^{0,2}$ is the conjugate statement.
Part (2). From $\omega^2=2\,\mathrm{vol}$ (the definition of the induced volume form on a Kähler surface) one gets $\int_X\omega\wedge\omega=2\int_X\mathrm{vol}=2\,\mathrm{Vol}(X)>0$.
Part (3). Let β be a real primitive $(1,1)$-form, $\beta\wedge\omega=0$. The claim is that the Hodge star, defined in definition 1.1.8, acts on such forms by $\star\beta=-\beta$. Primitivity says $\Lambda\beta=0$ (definition 1.1.8), which in the unitary coframe is the trace condition on the Hermitian matrix of β; the real primitive $(1,1)$-forms are spanned by

$$\gamma_1=i(\varphi^1\wedge\bar\varphi^1-\varphi^2\wedge\bar\varphi^2),\quad \gamma_2=\varphi^1\wedge\bar\varphi^2-\varphi^2\wedge\bar\varphi^1,\quad \gamma_3=i(\varphi^1\wedge\bar\varphi^2+\varphi^2\wedge\bar\varphi^1)$$

(each is real and has vanishing $\varphi^1\wedge\bar\varphi^1,\varphi^2\wedge\bar\varphi^2$ trace). On an oriented Riemannian 4-manifold $\star$ is an involution on 2-forms, giving the splitting $\Lambda^2=\Lambda^+\oplus\Lambda^-$ into self-dual and anti-self-dual parts; a general 2-form is a sum of the two and is *not* an eigenform, so this alone does not place γ_j in one summand. What does is a norm count. Write $\gamma_j=\gamma_j^+ +\gamma_j^-$; then $\gamma_j\wedge\gamma_j=(|\gamma_j^+|^2-|\gamma_j^-|^2)\,\mathrm{vol}$ while $\|\gamma_j\|^2=|\gamma_j^+|^2+|\gamma_j^-|^2$. Direct computation in the unitary coframe gives $\gamma_j\wedge\gamma_j=-2\,\mathrm{vol}$ and $\|\gamma_j\|^2=2$, and the two force $|\gamma_j^+|^2=0$. So each γ_j is anti-self-dual, $\star\gamma_j=-\gamma_j$. The sign is read off from $\gamma_j\wedge\star\gamma_j=\|\gamma_j\|^2\,\mathrm{vol}$, a positive multiple of vol: since $\gamma_1\wedge\gamma_1=-2\,\mathrm{vol}<0$ (and likewise $\gamma_2\wedge\gamma_2,\gamma_3\wedge\gamma_3<0$), each eigenvalue is -1. Hence $\star\beta=-\beta$ on the whole primitive space. Since β is real, $\bar\beta=\beta$, and

$$\int_X\beta\wedge\beta=-\int_X\beta\wedge\star\beta=-\int_X|\beta|^2\,\mathrm{vol}<0$$

for $\beta\not\equiv0$, completing the proof.

The three signs are the whole content: the cup form is positive on $H^{2,0} \oplus H^{0,2}$ (once one passes to a real basis), positive along the Kähler class inside $H^{1,1}$, and negative on the orthogonal complement of the Kähler class inside the real $(1,1)$-classes. Counting dimensions of the positive and negative eigenspaces gives the signature.

Theorem 2.2.3: Hodge Index Theorem

Let X be a compact Kähler surface. The cup-product form $Q(\alpha,\beta) = \int_X \alpha \wedge \beta$ on $H^2(X,\mathbb{R})$ has signature

$$(b^+, b^-) = \left(2h^{2,0} + 1,\ h^{1,1} - 1\right)$$

In particular its restriction to the real $(1,1)$-classes $H^{1,1}(X) \cap H^2(X,\mathbb{R})$ has signature $(1, h^{1,1} - 1)$: one positive direction, spanned by the Kähler class, and $h^{1,1} - 1$ negative directions.

***Proof*:**

Step 1: Orthogonality of the Hodge pieces. Extend Q $\mathbb{C}$-bilinearly to $H^2(X,\mathbb{C}) = H^{2,0} \oplus H^{1,1} \oplus H^{0,2}$. A product $\alpha \wedge \beta$ of a (p,q)-form and a (p',q')-form is a $(p+p', q+q')$-form, and only a $(2,2)$-form has nonzero integral over the surface X. So $Q(\alpha,\beta) = 0$ unless the bidegrees are complementary: Q pairs $H^{2,0}$ with $H^{0,2}$, pairs $H^{1,1}$ with itself, and kills every other combination. The decomposition is Q-orthogonal into the block $H^{2,0} \oplus H^{0,2}$ and the block $H^{1,1}$.

Step 2: The block $H^{2,0} \oplus H^{0,2}$. This complex subspace has complex dimension $2h^{2,0}$ and is defined over $\mathbb{R}$: complex conjugation swaps its two summands, so its real points form a real subspace $V \subseteq H^2(X,\mathbb{R})$ of real dimension $2h^{2,0}$, and every $w \in V$ is $w = \alpha + \bar{\alpha}$ for a unique $\alpha \in H^{2,0}$. On a surface $\alpha \wedge \alpha$ is a $(4,0)$-form, hence zero, and likewise $\bar{\alpha} \wedge \bar{\alpha} = 0$, so

$$Q(w,w) = \int_X (\alpha + \bar{\alpha}) \wedge (\alpha + \bar{\alpha}) = 2 \int_X \alpha \wedge \bar{\alpha}$$

which is > 0 for $\alpha \neq 0$ by part (1) of lemma 2.2.2. Thus $Q|_V$ is positive definite, and V contributes $(2h^{2,0}, 0)$ to the signature.

Step 3: The block $H^{1,1} \cap H^2(X,\mathbb{R})$. On the real $(1,1)$-classes Q is real-valued and nondegenerate. The Kähler class $[\omega]$ satisfies $Q([\omega],[\omega]) > 0$ by part (2) of lemma 2.2.2, so $\mathbb{R}[\omega]$ is a positive line. The Lefschetz operator $L = \omega \wedge (-)$ of definition 1.1.8 gives, in degree two on a surface, an orthogonal decomposition of the real $(1,1)$-classes into $\mathbb{R}[\omega]$ and the primitive part $P^{1,1} = \ker(\wedge\omega)$: any real $(1,1)$-class γ writes as $\gamma = \frac{Q(\gamma,[\omega])}{Q([\omega],[\omega])}[\omega] + \gamma_0$ with γ_0 primitive, and the two summands are Q-orthogonal because $Q([\omega],\gamma_0) = \int_X \omega \wedge \gamma_0 = 0$ by primitivity. Part (3) of lemma 2.2.2 makes Q negative definite on $P^{1,1}$, of dimension $h^{1,1} - 1$. So the real $(1,1)$-classes contribute $(1, h^{1,1} - 1)$.

Step 4: Assemble. By Step 1 the two blocks are Q-orthogonal, so their signatures add:

$$(b^+, b^-) = (2h^{2,0}, 0) + (1, h^{1,1} - 1) = (2h^{2,0} + 1,\ h^{1,1} - 1)$$

The restriction to the real $(1,1)$-classes is $(1, h^{1,1} - 1)$ from Step 3, as we sought to show.

The theorem earns its name from the negative-definite conclusion on the primitive $(1,1)$-classes: apart from the single positive direction of the Kähler class, the intersection form on divisor-type classes is negative definite. Since the Néron-Severi group $\mathrm{NS}(X)$ sits inside $H^{1,1}(X) \cap H^2(X,\mathbb{Z})$ (this is the Lefschetz $(1,1)$-theorem of the next section), the algebraic classes inherit signature $(1, \rho - 1)$ when X

is projective, where $\rho = \operatorname{rk} \mathrm{NS}(X)$ is the Picard number. Projectivity is needed and not automatic: the restriction of a form of signature $(1, h^{1,1}-1)$ to a subspace has a positive direction only if that subspace contains a class of positive square, which an ample class gives. On a non-projective Kähler surface $\mathrm{NS}(X)$ can be negative definite, or zero. For an algebraic surface with an ample class H, this recovers the classical Hodge index theorem: if a divisor class D satisfies $D \cdot H = 0$, then $D^2 \leq 0$, with equality only for D numerically trivial. The transcendental part $H^{2,0} \oplus H^{0,2}$ gives the extra positive directions $2h^{2,0}$, which is why a K3 surface, with $h^{2,0} = 1$, has $b^+ = 3$ rather than 1.

The value $b^- = h^{1,1} - 1$ also expresses the topological signature $\sigma(X) = b^+ - b^- = 2 + 2h^{2,0} - h^{1,1}$ purely through Hodge numbers. With $\chi_{\mathrm{top}}(X) = 2 - 4h^{1,0} + 2h^{2,0} + h^{1,1}$ (from $b_1 = 2h^{1,0}$, $b_2 = 2h^{2,0} + h^{1,1}$), the two combine to

$$\tfrac{1}{4}\big(\sigma(X) + \chi_{\mathrm{top}}(X)\big) = 1 - h^{1,0} + h^{2,0} = \chi(\mathcal{O}_X)$$

the holomorphic Euler characteristic $\chi(\mathcal{O}_X) = \sum_q (-1)^q h^{0,q}$. This is the topological form of Noether's formula: the alternating sum $\sum_q (-1)^q h^{0,q}$ is pinned down by the signature and the Euler characteristic together.

Example 2.2.4: The Intersection Form on a K3 and an Abelian Surface
Two surfaces show the two regimes of the index theorem.

1. *K3 surface.* A K3 surface X has $h^{1,0} = 0$ (so $b_1 = 0$) and $h^{2,0} = 1$, hence $b_2 = 2 \cdot 1 + h^{1,1} = 22$ forces $h^{1,1} = 20$. The Hodge diamond is

$$\begin{array}{ccccc} & & 1 & & \\ & 0 & & 0 & \\ 1 & & 20 & & 1 \\ & 0 & & 0 & \\ & & 1 & & \end{array}$$

   By theorem 2.2.3 the intersection form on $H^2(X, \mathbb{Z}) \cong \mathbb{Z}^{22}$ has signature $(2 \cdot 1 + 1,\ 20 - 1) = (3, 19)$. The intersection form is unimodular ([30, integral unimodularity of the intersection pairing]) and even. Those two facts do not by themselves pin down the lattice; the classification of indefinite even unimodular lattices, which this book does not prove, identifies it with $H^2(X, \mathbb{Z}) \cong U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$, the K3 lattice, where U is the hyperbolic plane and $E_8(-1)$ the negative E_8 lattice; the $(3, 19)$ signature is exactly $3 \cdot (1, 1) + 2 \cdot (0, 8)$. On the real $(1, 1)$-classes the signature is $(1, 19)$, and for a *projective* K3 the algebraic classes $\mathrm{NS}(X)$, of rank $\rho \leq 20$, carry signature $(1, \rho - 1)$.

2. *Abelian surface.* An abelian surface $X = \mathbb{C}^2/\Lambda$ has, by example 2.1.2 with $n = 2$, Hodge numbers $h^{1,0} = 2$, $h^{2,0} = 1$, $h^{1,1} = 4$, so $b_2 = 2 \cdot 1 + 4 = 6$. The intersection form on $H^2(X, \mathbb{Z}) \cong \mathbb{Z}^6$ has signature $(2 \cdot 1 + 1,\ 4 - 1) = (3, 3)$. Indeed $H^2(X, \mathbb{Z}) \cong \bigwedge^2 H^1(X, \mathbb{Z})$ with $H^1 \cong \mathbb{Z}^4$, and the cup product $\bigwedge^2 \mathbb{Z}^4 \to \bigwedge^4 \mathbb{Z}^4 \cong \mathbb{Z}$ is the standard split form of signature $(3, 3)$. On the real $(1, 1)$-classes, of dimension 4, the signature is $(1, 3)$.

Exercise 2.2.1

1. Let X be a compact Kähler surface. Using $b_1 = 2h^{1,0}$, $b_2 = 2h^{2,0} + h^{1,1}$, and $b_3 = b_1$, express the topological Euler characteristic $\chi_{\mathrm{top}}(X) = \sum_k (-1)^k b_k$ in terms of $h^{1,0}, h^{2,0}, h^{1,1}$, and show $\chi_{\mathrm{top}}(X) \equiv b_2 \equiv h^{1,1} \pmod 2$. (Do not expect χ_{top} itself to be even: $\mathbb{P}^2$ has $\chi_{\mathrm{top}} = 3$.)

2. Show that on any compact Kähler surface the topological signature $\sigma(X) = b^+ - b^-$ equals $2 + 2h^{2,0} - h^{1,1}$. Deduce that $\sigma(X) \equiv b_2 \pmod 2$.

3. The Hopf surface (a quotient of $\mathbb{C}^2 \setminus \{0\}$) is a compact complex surface with $b_1 = 1$. Show that it admits no Kähler metric, using only corollary 2.2.1.

4. Let X be a compact Kähler surface and H an ample class with $H^2 = d > 0$. For a real $(1,1)$-class D set $D_0 = D - \frac{D \cdot H}{d} H$. Show D_0 is primitive and deduce the classical inequality: if $D \cdot H = 0$ and $D \neq 0$ then $D^2 < 0$.

5. On a compact Kähler surface, use corollary 2.2.1 to show $h^{2,0} = h^{0,2} = p_g$ (the geometric genus $\dim H^0(X, \Omega_X^2)$) and $h^{1,0} = h^{0,1} = q$ (the irregularity $\dim H^0(X, \Omega_X^1)$), identifying $H^{2,0}(X)$ with $H^0(X, \Omega_X^2)$ and $H^{1,0}(X)$ with $H^0(X, \Omega_X^1)$.

6. For the abelian surface of example 2.2.4, exhibit an explicit basis of $H^2(X, \mathbb{R})$ in which the cup-product form is diagonal with entries $(+1, +1, +1, -1, -1, -1)$, starting from a real basis e_1, e_2, e_3, e_4 of $H^1(X, \mathbb{R})$.

2.3 The Lefschetz $(1,1)$-Theorem

The Hodge decomposition sorts $H^2(X, \mathbb{C})$ into the summands $H^{2,0}$, $H^{1,1}$, $H^{0,2}$; the topology alone cannot see this splitting, but the complex structure does. A holomorphic line bundle produces a class in $H^2(X, \mathbb{Z})$, its first Chern class, and this class always lands in the $(1,1)$-part. Lefschetz's theorem is the converse: on a compact Kähler manifold every integral class of type $(1,1)$ comes from a holomorphic line bundle, hence from a divisor. This section proves that converse through the exponential sequence, which is the exact sequence linking the additive sheaf $\mathcal{O}_X$ to the multiplicative sheaf $\mathcal{O}_X^*$ of nowhere-vanishing holomorphic functions.

The mechanism is a single connecting homomorphism. The holomorphic line bundles on X are classified by $H^1(X, \mathcal{O}_X^*)$, and the exponential sequence produces a map from this group to $H^2(X, \mathbb{Z})$ which is exactly c_1. The theorem describes the kernel and image of that map.

Proposition 2.3.1: The Exponential Sequence

Let X be a complex manifold. The map $\exp\colon \mathcal{O}_X \to \mathcal{O}_X^*$, $f \mapsto \exp(2\pi i f)$, fits into a short exact sequence of sheaves of abelian groups

$$0 \to \underline{\mathbb{Z}} \to \mathcal{O}_X \xrightarrow{\exp(2\pi i \cdot)} \mathcal{O}_X^* \to 0$$

where $\underline{\mathbb{Z}}$ is the constant sheaf and the multiplicative group structure is used on $\mathcal{O}_X^*$. The induced long exact sequence in cohomology contains

$$\cdots \to H^1(X, \mathcal{O}_X) \to H^1(X, \mathcal{O}_X^*) \xrightarrow{c_1} H^2(X, \mathbb{Z}) \to H^2(X, \mathcal{O}_X) \to \cdots$$

and the connecting homomorphism $c_1\colon H^1(X, \mathcal{O}_X^*) \to H^2(X, \mathbb{Z})$ sends the class of a holomorphic line bundle L to its first Chern class.

***Proof*:**
Step 1: Exactness of the sheaf sequence. The constant sheaf $\underline{\mathbb{Z}}$ includes into $\mathcal{O}_X$ as the locally constant integer-valued functions, and $\exp(2\pi i n) = 1$ for $n \in \mathbb{Z}$, so the composite $\underline{\mathbb{Z}} \to \mathcal{O}_X^*$ is trivial and the sequence is a complex. Injectivity of $\underline{\mathbb{Z}} \to \mathcal{O}_X$ is clear. For surjectivity of exp on stalks, let g be a nowhere-vanishing holomorphic function on a connected open U. Shrinking U to a polydisc neighborhood of any point, g admits a holomorphic logarithm: g has no zeros, so the holomorphic 1-form dg/g is closed, and on a simply connected domain it has a holomorphic primitive h with $d(ge^{-h}) = 0$, whence $g = c\,e^h$ for a constant $c \neq 0$; absorbing $\log c$ into h and dividing by $2\pi i$ gives a holomorphic f with $\exp(2\pi i f) = g$. Thus exp is surjective as a map of sheaves. Exactness in the middle is the observation that $\exp(2\pi i f) = 1$ forces f to be a locally constant integer, so the kernel of exp is exactly $\underline{\mathbb{Z}}$.

Step 2: The connecting map is c_1. A holomorphic line bundle L is given by a cocycle $(g_{\alpha\beta})$ of transition functions $g_{\alpha\beta} \in \mathcal{O}_X^*(U_\alpha \cap U_\beta)$ satisfying the cocycle condition $g_{\alpha\beta}g_{\beta\gamma}g_{\gamma\alpha} = 1$; this is precisely a Čech 1-cocycle representing a class in $H^1(X, \mathcal{O}_X^*)$.[a] The connecting homomorphism of the long exact sequence computes as follows. Choose logarithms $f_{\alpha\beta} = \frac{1}{2\pi i}\log g_{\alpha\beta} \in \mathcal{O}_X(U_\alpha \cap U_\beta)$ on the overlaps, so $\exp(2\pi i f_{\alpha\beta}) = g_{\alpha\beta}$. The failure of $(f_{\alpha\beta})$ to be a cocycle,

$$n_{\alpha\beta\gamma} = f_{\beta\gamma} - f_{\alpha\gamma} + f_{\alpha\beta}$$

is a 2-cocycle valued in $\underline{\mathbb{Z}}$ (it exponentiates to $g_{\beta\gamma}g_{\alpha\gamma}^{-1}g_{\alpha\beta} = 1$, so it is integer-valued), and its class in $H^2(X,\mathbb{Z})$ is by definition the image of $[L]$ under the connecting map. That integer 2-cocycle is the standard Čech representative of the first Chern class of L, completing the proof.

[a] [5, The Picard Group as Sheaf Cohomology].

The two boxed groups at the ends of the six-term stretch are the obstruction groups. Reading the sequence

$$H^1(X, \mathcal{O}_X^*) \xrightarrow{c_1} H^2(X,\mathbb{Z}) \xrightarrow{\iota} H^2(X, \mathcal{O}_X)$$

as an exact sequence, a class $\gamma \in H^2(X,\mathbb{Z})$ lies in the image of c_1 if and only if it dies in $H^2(X,\mathcal{O}_X)$. The map ι is the composite of the change-of-coefficients map $H^2(X,\mathbb{Z}) \to H^2(X,\mathbb{C})$ with the projection of $H^2(X,\mathbb{C})$ onto its $H^{0,2}$-summand, since Dolbeault's theorem (theorem 1.1.3) identifies $H^2(X,\mathcal{O}_X) = H^{0,2}(X)$. The Hodge decomposition turns the vanishing of $\iota(\gamma)$ into a statement about Hodge type, and that is the theorem.

Before stating it, one point of bookkeeping. The image of an integral class γ in $H^2(X,\mathbb{C})$ decomposes as $\gamma^{2,0} + \gamma^{1,1} + \gamma^{0,2}$. Because γ is real, complex conjugation fixes it, and conjugation exchanges $H^{2,0}$ with $H^{0,2}$ (theorem 2.1.1), so $\gamma^{2,0} = \overline{\gamma^{0,2}}$. Hence $\gamma^{0,2} = 0$ if and only if $\gamma^{2,0} = 0$, and both vanish exactly when γ is of pure type $(1,1)$.

Theorem 2.3.2: The Lefschetz $(1,1)$-Theorem

Let X be a compact Kähler manifold and $\gamma \in H^2(X,\mathbb{Z})$. The following are equivalent.

1. γ is the first Chern class $c_1(L)$ of a holomorphic line bundle L on X.
2. The image of γ in $H^2(X,\mathcal{O}_X)$ vanishes.
3. The image of γ in $H^2(X,\mathbb{C})$ lies in the summand $H^{1,1}(X)$; that is, γ is an integral class of type $(1,1)$.

***Proof*:**

(1) ⇔ (2). By proposition 2.3.1 the segment $H^1(X,\mathcal{O}_X^*) \xrightarrow{c_1} H^2(X,\mathbb{Z}) \xrightarrow{\iota} H^2(X,\mathcal{O}_X)$ is exact, so γ lies in the image of c_1 precisely when $\iota(\gamma)=0$. Since every element of $H^1(X,\mathcal{O}_X^*)$ is the class of a holomorphic line bundle,[a] the image of c_1 is exactly the set of Chern classes of line bundles, which is (1).

(2) ⇔ (3). Dolbeault's theorem (theorem 1.1.3) identifies $H^2(X,\mathcal{O}_X)=H^{0,2}(X)$, and under this identification ι is the composite

$$H^2(X,\mathbb{Z}) \to H^2(X,\mathbb{C}) = H^{2,0}(X)\oplus H^{1,1}(X)\oplus H^{0,2}(X) \to H^{0,2}(X)$$

of the coefficient map with projection onto the $(0,2)$-summand; the first equality is the Hodge decomposition (theorem 2.1.1), available because X is compact Kähler. Thus $\iota(\gamma)=0$ says $\gamma^{0,2}=0$. As noted before the statement, the image of γ is real, so $\gamma^{2,0}=\overline{\gamma^{0,2}}$, and $\gamma^{0,2}=0$ forces $\gamma^{2,0}=0$ as well. Hence $\gamma^{0,2}=0$ if and only if $\gamma=\gamma^{1,1}$ lies in $H^{1,1}(X)$, which is (3), as we sought to show.

[a] [5, The Picard Group as Sheaf Cohomology].

The equivalence of (1) and (3) is the usable form: it converts a transcendental condition (Hodge type, read off the complex structure) into an algebro-geometric one (representability by a line bundle). The Chern classes of line bundles are exactly the integral $(1,1)$-classes. Naming the group of such classes records this.

Definition 2.3.3: The Néron-Severi Group

Let X be a compact Kähler manifold. The *Néron-Severi group* of X is

$$\mathrm{NS}(X) = H^{1,1}(X)\cap H^2(X,\mathbb{Z})$$

the group of integral cohomology classes of type $(1,1)$, where the intersection is taken inside $H^2(X,\mathbb{C})$. Its rank is the *Picard number* $\rho(X)$.

Proposition 2.3.4: Bound on the Picard Number

For a compact Kähler manifold X, the Picard number satisfies $\rho(X)\le h^{1,1}(X)$.

***Proof*:**

On mapping into $H^2(X,\mathbb{C})$, the group $\mathrm{NS}(X)=H^{1,1}(X)\cap H^2(X,\mathbb{Z})$ lands inside $H^{1,1}(X)\cap H^2(X,\mathbb{R})$: its elements are real classes, so they lie in the conjugation-stable part of $H^{1,1}(X)$, and since $H^{1,1}=\overline{H^{1,1}}$, that part is a real form of $H^{1,1}(X)$, of real dimension $h^{1,1}$. As a subgroup of the lattice $H^2(X,\mathbb{Z})$, $\mathrm{NS}(X)$ is discrete, and a discrete subgroup of a real vector space of dimension $h^{1,1}$ has rank at most $h^{1,1}$. Hence $\rho(X)=\mathrm{rank}\mathrm{NS}(X)\le h^{1,1}(X)$.

By theorem 2.3.2 the Néron-Severi group is the image of $c_1\colon H^1(X,\mathcal{O}_X^*)\to H^2(X,\mathbb{Z})$, so it is the group of first Chern classes of holomorphic line bundles. On a projective manifold every holomorphic line bundle arises from a divisor. Combining the theorem with the divisor-to-Picard homomorphism[2]

[2] [5, Divisor Class to Picard Homomorphism].

gives the geometric reading in that case.

An element $D \in \mathrm{Div}(X)$ is a formal $\mathbb{Z}$-combination of irreducible hypersurfaces; the class map $D \mapsto \mathcal{O}_X(D)$ sends it to a holomorphic line bundle, and this descends to a homomorphism $\mathrm{Cl}(X) \to H^1(X, \mathcal{O}_X^*) = \mathrm{Pic}(X)$ from divisor classes to line bundles.[3] Composing with c_1, an integral $(1,1)$-class is the class of a divisor: on a projective manifold every holomorphic line bundle is $\mathcal{O}_X(D)$ for some divisor D, so $\mathrm{NS}(X)$ is exactly the group of divisor classes modulo those algebraically equivalent to zero. This is why the theorem is stated as "integral $(1,1)$-classes are divisor classes".

2.3.5 Warning: Kähler versus projective On a general compact Kähler manifold, $c_1(L)$ is an integral $(1,1)$-class for every line bundle L, and theorem 2.3.2 produces a line bundle from each such class. The further step "every line bundle comes from a divisor" needs a nonzero *meromorphic* section of L, equivalently that $L \otimes \mathcal{O}(mH)$ be globally generated for $m \gg 0$ with H ample, which writes $L = \mathcal{O}(D_1 - mH)$. That is available only when X is projective (equivalently, when X carries a Kähler class in $H^2(X,\mathbb{Z})$). A compact Kähler manifold with $\mathrm{NS}(X) = 0$, such as a generic complex torus (example 1.1.6), carries no divisors at all, consistent with an empty Néron-Severi group.

The running example shows these groups. A complex torus $X = \mathbb{C}^g/\Lambda$ (example 1.1.6) has $H^1(X, \mathcal{O}_X)$ and $H^2(X,\mathbb{Z})$ both computable from the lattice, so the Néron-Severi group can be described by hand. The case $g = 2$, an abelian surface, is the smallest one where the Picard number varies with the complex structure.

Example 2.3.6: The Néron-Severi Group of an Abelian Surface

Let $X = \mathbb{C}^2/\Lambda$ be a complex torus of dimension 2, with $\Lambda \cong \mathbb{Z}^4$ a lattice. Its integral cohomology is the exterior algebra on $H^1(X,\mathbb{Z}) = \mathrm{Hom}(\Lambda,\mathbb{Z}) \cong \mathbb{Z}^4$, so

$$H^2(X,\mathbb{Z}) = \bigwedge^2 \mathrm{Hom}(\Lambda,\mathbb{Z}) \cong \mathbb{Z}^6$$

The complex structure on $\mathbb{C}^2$ makes $\Lambda \otimes \mathbb{R} = \mathbb{C}^2$ a complex vector space, and the Hodge decomposition of $H^2(X,\mathbb{C})$ reads, in terms of the splitting of alternating $\mathbb{R}$-bilinear forms on $\mathbb{C}^2$ into their types,

$$H^{2,0} \oplus H^{1,1} \oplus H^{0,2}, \qquad h^{2,0} = 1, \quad h^{1,1} = 4, \quad h^{0,2} = 1$$

An integral class $\gamma \in H^2(X,\mathbb{Z})$ is an alternating form $E\colon \Lambda \times \Lambda \to \mathbb{Z}$. By theorem 2.3.2, $\gamma \in \mathrm{NS}(X)$ if and only if γ is of type $(1,1)$, and for an alternating form on the complex vector space $\mathbb{C}^2$ the $(1,1)$-condition is exactly the Riemann bilinear relation

$$E(iu, iv) = E(u,v) \qquad \text{for all } u, v \in \mathbb{C}^2$$

(the vanishing of the $(2,0)$ and $(0,2)$ parts of E is the invariance of E under the complex structure $J = i$). Thus

$$\mathrm{NS}(X) = \{\, E \in \bigwedge^2 \mathrm{Hom}(\Lambda,\mathbb{Z}) : E(iu,iv) = E(u,v) \,\}$$

The rank of this group depends on Λ. For a very general lattice the only integral J-invariant alternating form is 0, so $\rho(X) = 0$ and X has no divisors: it is a non-algebraic complex torus. Imposing the J-invariance on one form (a single nonzero solution together with a positivity

[3] [5, Line Bundle of a Divisor, Divisor Class to Picard Homomorphism].

condition) already forces X to be an abelian variety; the maximal Picard number for a 2-torus is $\rho = 4$, attained when X is isogenous to a product of elliptic curves with complex multiplication. The interpolation between these, and the Appell-Humbert description of line bundles by their Chern class E together with a semicharacter, is developed in [3]; the point here is that the Lefschetz theorem reduces the whole computation to counting integral solutions of the linear condition $E(iu, iv) = E(u, v)$.

The example exhibits the two extremes the theorem allows. When $\mathrm{NS}(X) = 0$ the manifold is Kähler but carries no line bundle with nonzero Chern class, so no divisor; when ρ is maximal the torus is an abelian variety with many divisors. Both are governed by the single linear condition that an integral class be of type $(1,1)$, and that condition is what theorem 2.3.2 extracts from the exponential sequence.

Exercise 2.3.1

1. Show that the exponential sequence $0 \to \underline{\mathbb{Z}} \to \mathcal{O}_X \to \mathcal{O}_X^* \to 0$ does not split as a sequence of sheaves of abelian groups when $X = \mathbb{C}^*$ (the punctured plane, a noncompact Riemann surface). (Hint: consider whether the identity function $z \in \mathcal{O}_X^*(X)$ admits a global holomorphic logarithm.)

2. Let X be a compact Kähler manifold. Using proposition 2.3.1, show that the kernel of $c_1 \colon H^1(X, \mathcal{O}_X^*) \to H^2(X, \mathbb{Z})$ is the quotient $H^1(X, \mathcal{O}_X)/\mathrm{im}\big(H^1(X, \mathbb{Z})\big)$, the group $\mathrm{Pic}^0(X)$ of line bundles with vanishing first Chern class. Identify this quotient as a complex torus using the Hodge decomposition of H^1.

3. Proposition 2.3.4 bounds the Picard number of a compact Kähler manifold by $h^{1,1}(X)$. Give an example (a very general complex torus of dimension ≥ 2 will do) where the inequality is strict.

4. Let $E = \mathbb{C}/\Lambda$ be an elliptic curve. Compute $H^2(E, \mathbb{Z})$ and show that $\mathrm{NS}(E) = H^2(E, \mathbb{Z}) \cong \mathbb{Z}$, so that every integral class of the top-degree cohomology of E is the class of a divisor. Identify the generator with the class of a single point.

5. Let $X = E_1 \times E_2$ be a product of two elliptic curves. Show that $\mathrm{NS}(X) = \mathbb{Z}F_1 \oplus \mathbb{Z}F_2 \oplus \mathrm{Hom}(E_1, E_2)$, where F_1, F_2 are the two fiber classes, and deduce that $\rho(X) = 2 + \mathrm{rank}\mathrm{Hom}(E_1, E_2)$. Conclude that $\rho(X) = 2$ for E_1, E_2 generic (non-isogenous), $\rho(X) = 3$ when E_1, E_2 are isogenous without complex multiplication, and $\rho(X) = 4$ when they are isogenous with complex multiplication.

6. State the Hodge conjecture for H^2 and observe that on a compact Kähler manifold it is a theorem, namely theorem 2.3.2. Explain in one or two sentences why the analogous statement in degree H^4 (for $(2,2)$-classes) is not accessible by the exponential-sequence argument.

Part II

Hodge Structures and Their Variations

Part I ended with a theorem about a single manifold: the complex cohomology of a compact Kähler X splits by type, and the splitting is an invariant of X. Read again, the conclusion barely mentions X. What it produces is a piece of linear algebra on a rational vector space, a conjugate-symmetric decomposition of its complexification together with a bilinear form given by the Kähler class, and every use of the geometry has been spent by the time that data appears.

This Part takes the linear algebra as the object. The gain is reach rather than economy. Structures of the same shape arise where no Kähler manifold is in sight, and the ones that do come from geometry satisfy constraints that a single manifold cannot exhibit: fix the numerical invariants and the polarized structures of that type are the points of a complex manifold, so a family of varieties traces a path in it and a degeneration is a limit. Questions that cannot even be posed about one X become questions about points and paths. Whether two varieties with the same cohomology are isomorphic is a question of injectivity; which classes are algebraic is a question about the locus where the structure acquires extra pieces.

The chapters that follow build the structure and the space that parametrizes it, prove the Lefschetz theorems that constrain it on a single manifold, and then let it move: a family of varieties gives a holomorphically varying structure whose derivative is constrained, and that constraint is the source of most of what is known about how varieties fit together in moduli.

3

Hodge Structures and Polarizations

The Hodge decomposition attaches to a compact Kähler manifold a bigrading of its complex cohomology. This chapter isolates that datum from the manifold that produced it. A *Hodge structure* is a conjugate-symmetric bigrading of the complexification of a rational vector space, and a *polarization* is the extra bilinear form that cup product against a Kähler class gives. The polarization is what makes the notion rigid: unpolarized structures of a given type form a set with no useful geometry, while polarized ones form a complex manifold.

Two features surface once the manifold is gone, and both are invisible while it is present. The bigrading turns out to be equivalent to a decreasing filtration, and it is the filtration, not the bigrading, that behaves well in families. And the polarized structures of a fixed numerical type are themselves the points of a complex manifold, the period domain, whose geometry the last two sections begin.

3.1 Pure Hodge Structures and the Hodge Filtration

The definition is read off from theorem 2.1.1. That theorem produces, on a compact Kähler manifold X, a decomposition $H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ satisfying $H^{p,q} = \overline{H^{q,p}}$, where the conjugation is the one fixing the real cohomology $H^k(X,\mathbb{R})$. Both halves matter. The decomposition alone is a bigrading of a complex vector space and carries no information beyond a list of dimensions; the conjugation symmetry is what ties it to a real, and in fact rational, structure underneath.

Definition 3.1.1: Pure Hodge Structure of Weight k

Let $V_{\mathbb{Z}}$ be a finitely generated free abelian group, $V_{\mathbb{Q}} = V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$ and $V_{\mathbb{C}} = V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$, and let $k \in \mathbb{Z}$. A *pure Hodge structure of weight k* on $V_{\mathbb{Z}}$ is a direct sum decomposition

$$V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$$

into complex subspaces, subject to the symmetry

$$\overline{V^{p,q}} = V^{q,p}$$

where the bar is the conjugation of $V_{\mathbb{C}} = V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ fixing $V_{\mathbb{R}}$. The integers $h^{p,q} = \dim_{\mathbb{C}} V^{p,q}$ are the *Hodge numbers* of the structure. The same definition with $V_{\mathbb{Q}}$ in place of $V_{\mathbb{Z}}$ defines a *rational* Hodge structure; $V_{\mathbb{Z}}$ is called the *lattice* and is retained because the geometric examples carry one.

The weight is part of the data, not a consequence of it: the same bigraded vector space underlies a weight-k structure and, after relabelling, structures of other weights. Only the sum $p+q$ is pinned, and pinning it is what makes the Hodge numbers of two structures comparable.

The symmetry deserves a second look, because it is the only place the rational structure enters. Conjugation on $V_{\mathbb{C}}$ is defined by $V_{\mathbb{R}}$, hence by $V_{\mathbb{Q}}$, hence by $V_{\mathbb{Z}}$; a bigrading of an abstract complex vector space has no conjugation to be symmetric under. So $\overline{V^{p,q}} = V^{q,p}$ is a compatibility between the bigrading and the lattice, and it immediately forces $h^{p,q} = h^{q,p}$, since conjugation is a conjugate-linear isomorphism and those preserve complex dimension.

Example 3.1.2: The Cohomology of a Compact Kähler Manifold

Let X be a compact Kähler manifold and let $V_{\mathbb{Z}} = H^k(X,\mathbb{Z})/(\text{torsion})$, so that $V_{\mathbb{C}} = H^k(X,\mathbb{C})$. Setting $V^{p,q} = H^{p,q}(X)$, theorem 2.1.1 gives both the decomposition and the symmetry $H^{p,q} = \overline{H^{q,p}}$, the conjugation there being the one fixing $H^k(X,\mathbb{R})$. So $H^k(X,\mathbb{Z})$ carries a pure Hodge structure of weight k, whose Hodge numbers are the $h^{p,q}(X)$ of theorem 1.1.3.

This is the example the definition was extracted from, and for the rest of the book it is the source of every geometric instance. The weight equals the cohomological degree, which is the reason the abstract theory indexes by weight at all.

Example 3.1.3: The Lowest Weights

The first two weights are already instructive. Both descriptions below assume the structure is *effective*, meaning $V^{p,q} = 0$ unless $p \geq 0$ and $q \geq 0$. Effectivity holds for the cohomology of a compact Kähler manifold, where p and q are form degrees, but it is not part of definition 3.1.1 and it fails after a Tate twist (definition 3.1.11); without it, weight alone constrains only the sum $p+q$.

1. *Weight* 0*, effective.* The only pair with $p+q=0$ and $p,q \geq 0$ is $(0,0)$, so $V_{\mathbb{C}} = V^{0,0}$ and the structure carries no information beyond $V_{\mathbb{Z}}$ itself. Effective weight-0 Hodge structures are just lattices.

2. *Weight* 1*, effective.* The available pairs are $(1,0)$ and $(0,1)$, giving $V_{\mathbb{C}} = V^{1,0} \oplus V^{0,1}$ with $\overline{V^{1,0}} = V^{0,1}$. The structure is therefore determined by the single subspace $V^{1,0}$, which must satisfy $V^{1,0} \oplus \overline{V^{1,0}} = V_{\mathbb{C}}$. Such a subspace is the same thing as a complex structure on the

real vector space $V_{\mathbb{R}}$: the projection $V_{\mathbb{R}} \hookrightarrow V_{\mathbb{C}} \to V^{1,0}$ along $V^{0,1}$ is an $\mathbb{R}$-linear isomorphism, and pulling back multiplication by i gives an endomorphism J of $V_{\mathbb{R}}$ with $J^2 = -\mathrm{id}$ (proposition 3.1.4). In particular $V_{\mathbb{R}}$ has even dimension and $h^{1,0} = h^{0,1} = \frac{1}{2}\dim_{\mathbb{R}} V_{\mathbb{R}}$.

The weight-1 case is the one that will reappear as the Hodge-theoretic description of complex tori in section 3.4, and the appearance of J is expected: it is the linear-algebraic form of an almost complex structure [5, Almost Complex Structure].

The weight-1 case of the example deserves to be a statement of record, since much of this book runs through it.

Proposition 3.1.4: Weight One Equals Complex Structure

Let $V_{\mathbb{R}}$ be a real vector space. Giving a subspace $V^{1,0} \subseteq V_{\mathbb{C}}$ with $V^{1,0} \oplus \overline{V^{1,0}} = V_{\mathbb{C}}$ is the same as giving an endomorphism J of $V_{\mathbb{R}}$ with $J^2 = -\mathrm{id}$; the correspondence sends J to its $(+i)$-eigenspace in $V_{\mathbb{C}}$. In particular $V_{\mathbb{R}}$ has even dimension, and an effective weight-1 Hodge structure on $V_{\mathbb{Z}}$ is the same as a complex structure on $V_{\mathbb{R}}$.

***Proof*:**

Given J with $J^2 = -\mathrm{id}$, extend it $\mathbb{C}$-linearly to $V_{\mathbb{C}}$. As for the tangent bundle of an almost complex manifold,[a] J satisfies $T^2 + 1 = (T - i)(T + i)$, which has distinct roots, so $V_{\mathbb{C}}$ splits as the direct sum of the $(\pm i)$-eigenspaces. Set $V^{1,0}$ to be the $(+i)$-eigenspace. Conjugation is a conjugate-linear automorphism of $V_{\mathbb{C}}$ commuting with J (because J is real), so it carries the $(+i)$-eigenspace to the $(-i)$-eigenspace. Hence $\overline{V^{1,0}}$ is the $(-i)$-eigenspace and $V^{1,0} \oplus \overline{V^{1,0}} = V_{\mathbb{C}}$.

Conversely, given such a $V^{1,0}$, the composite $V_{\mathbb{R}} \hookrightarrow V_{\mathbb{C}} \to V^{1,0}$, the second map being projection along $\overline{V^{1,0}}$, is $\mathbb{R}$-linear. It is injective because a real vector in its kernel lies in $\overline{V^{1,0}}$ and, being real, equals its own conjugate, hence lies in $V^{1,0} \cap \overline{V^{1,0}} = 0$. It is surjective by a dimension count: $\dim_{\mathbb{R}} V_{\mathbb{R}} = \dim_{\mathbb{C}} V_{\mathbb{C}} = 2\dim_{\mathbb{C}} V^{1,0} = \dim_{\mathbb{R}} V^{1,0}$. Transporting multiplication by i on $V^{1,0}$ back through this isomorphism defines J on $V_{\mathbb{R}}$ with $J^2 = -\mathrm{id}$, and the two constructions are inverse to each other by construction.

Since J makes $V_{\mathbb{R}}$ a complex vector space, $\dim_{\mathbb{R}} V_{\mathbb{R}}$ is even. Finally, an effective weight-1 Hodge structure on $V_{\mathbb{Z}}$ is exactly the datum of $V^{1,0}$ with $V^{1,0} \oplus \overline{V^{1,0}} = V_{\mathbb{C}}$, since $V^{0,1}$ is forced to be $\overline{V^{1,0}}$ by the symmetry. So it is the same as a complex structure on $V_{\mathbb{R}}$.

[a] [5, The $T^{1,0} \oplus T^{0,1}$ Decomposition].

For a single vector space the bigrading is the natural presentation. In a family it is not, and the reason is worth stating now even though the payoff is several chapters away: as a manifold varies, its subspaces $H^{p,q}$ move smoothly and no better, while certain sums of them move holomorphically. Those sums are the terms of a filtration.

Definition 3.1.5: The Hodge Filtration

Let V carry a pure Hodge structure of weight k. Its *Hodge filtration* is the decreasing chain of complex subspaces

$$F^pV_{\mathbb{C}} := \bigoplus_{r\geq p} V^{r,\,k-r}$$

so that $F^pV_{\mathbb{C}} \supseteq F^{p+1}V_{\mathbb{C}}$, with $F^pV_{\mathbb{C}} = V_{\mathbb{C}}$ for $p \leq \min\{r : V^{r,k-r} \neq 0\}$ and $F^pV_{\mathbb{C}} = 0$ for p large.

Nothing has been lost in passing from the bigrading to the filtration, because the bigrading can be read back off. The recovery formula is the content of the next theorem, and it is the identity that makes the two presentations interchangeable.

Theorem 3.1.6: Bigradings and Filtrations

Let $V_{\mathbb{Z}}$ be a finitely generated free abelian group and $k \in \mathbb{Z}$. The following data are equivalent.

1. A pure Hodge structure of weight k on $V_{\mathbb{Z}}$, that is, a decomposition $V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$ with $\overline{V^{p,q}} = V^{q,p}$.
2. A decreasing filtration $\cdots \supseteq F^pV_{\mathbb{C}} \supseteq F^{p+1}V_{\mathbb{C}} \supseteq \cdots$ of $V_{\mathbb{C}}$ by complex subspaces, which is exhaustive and separated ($F^pV_{\mathbb{C}} = V_{\mathbb{C}}$ for $p \ll 0$ and $F^pV_{\mathbb{C}} = 0$ for $p \gg 0$), satisfying

$$F^pV_{\mathbb{C}} \oplus \overline{F^{k-p+1}V_{\mathbb{C}}} = V_{\mathbb{C}} \qquad \text{for every } p \tag{3.1}$$

The passage from (1) to (2) is $F^pV_{\mathbb{C}} = \bigoplus_{r\geq p} V^{r,k-r}$, and the passage from (2) to (1) is

$$V^{p,q} = F^pV_{\mathbb{C}} \cap \overline{F^qV_{\mathbb{C}}}, \qquad p+q=k \tag{3.2}$$

The two constructions are mutually inverse.

***Proof*:**
Write F^p for $F^pV_{\mathbb{C}}$ throughout.

Step 1: a Hodge structure gives a filtration satisfying (3.1). Let $V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$ be a Hodge structure and set $F^p = \bigoplus_{r\geq p} V^{r,k-r}$. Decreasingness is clear, and the sum being finite gives exhaustiveness and separatedness. For (3.1), apply conjugation to the definition of F^{k-p+1} and use $\overline{V^{r,k-r}} = V^{k-r,r}$:

$$\overline{F^{k-p+1}} = \overline{\bigoplus_{r\geq k-p+1} V^{r,k-r}} = \bigoplus_{r\geq k-p+1} V^{k-r,\,r} = \bigoplus_{s\leq p-1} V^{s,\,k-s}$$

the last equality by the substitution $s = k-r$, which turns $r \geq k-p+1$ into $s \leq p-1$. So F^p collects the summands with first index at least p and $\overline{F^{k-p+1}}$ collects those with first index at most $p-1$. Every summand lies in exactly one of the two, so their sum is $V_{\mathbb{C}}$ and their intersection is 0.

Step 2: a filtration satisfying (3.1) *gives a Hodge structure.* Let $F^\bullet$ be as in (2) and define

$V^{p,q} := F^p \cap \overline{F^q}$ for $p+q=k$. The symmetry is immediate from the definition: conjugating,

$$\overline{V^{p,q}} = \overline{F^p \cap \overline{F^q}} = \overline{F^p} \cap F^q = V^{q,p}$$

since conjugation is an involution exchanging a subspace with its conjugate. It remains to prove that the $V^{p,q}$ sum to $V_{\mathbb{C}}$ and do so directly.

Fix p. The claim is

$$F^p = V^{p,k-p} \oplus F^{p+1} \tag{3.3}$$

Granting (3.3), descending induction on p from the top of the filtration gives $F^p = \bigoplus_{r \geq p} V^{r,k-r}$ for every p, and taking $p \ll 0$, where $F^p = V_{\mathbb{C}}$, gives the required decomposition of $V_{\mathbb{C}}$.

To prove (3.3), first note $V^{p,k-p} \cap F^{p+1} = 0$. An element of the intersection lies in F^{p+1}, and lies in $\overline{F^{k-p}}$ because $V^{p,k-p} = F^p \cap \overline{F^{k-p}}$. Applying (3.1) at $p+1$ gives $F^{p+1} \oplus \overline{F^{k-p}} = V_{\mathbb{C}}$, so the intersection is zero. For the sum, let $v \in F^p$. By (3.1) at $p+1$ write $v = a + b$ with $a \in F^{p+1}$ and $b \in \overline{F^{k-p}}$. Then $b = v - a$ lies in F^p, since $v \in F^p$ and $a \in F^{p+1} \subseteq F^p$; so $b \in F^p \cap \overline{F^{k-p}} = V^{p,k-p}$, and $v = b + a$ exhibits v in $V^{p,k-p} + F^{p+1}$. This proves (3.3).

Step 3: the constructions are mutually inverse. Step 1 computed $\overline{F^{k-p+1}} = \bigoplus_{s \leq p-1} V^{s,k-s}$ for the filtration attached to a Hodge structure, and the same substitution gives $\overline{F^q} = \bigoplus_{s \leq p} V^{s,k-s}$ when $q = k-p$. Intersecting with $F^p = \bigoplus_{r \geq p} V^{r,k-r}$ leaves exactly the single summand $V^{p,k-p}$, which is (3.2): starting from a Hodge structure, building its filtration and applying the recovery formula returns the original bigrading. Conversely, Step 2 built a Hodge structure from $F^\bullet$ and established $F^p = \bigoplus_{r \geq p} V^{r,k-r}$ along the way, so the filtration attached to that Hodge structure is $F^\bullet$ again. The two constructions are therefore mutually inverse, completing the proof.

Condition (3.1) is the whole content, and it is worth isolating what it says. A decreasing filtration on $V_{\mathbb{C}}$ is cheap: any flag of subspaces will do. What (3.1) requires is that each F^p be transverse to the conjugate of its complementary step, and transversality is an open condition. This is the first sign that the structures of a given numerical type form an open subset of a space of flags rather than a closed subvariety, a theme section 3.3 takes up.

Maps between Hodge structures are constrained by the bigrading, and the constraint has a consequence strong enough to be worth proving at once.

Definition 3.1.7: Morphism of Hodge Structures

Let V and W carry pure Hodge structures of weights k and l. A *morphism of Hodge structures* is a homomorphism $\varphi\colon V_{\mathbb{Z}} \to W_{\mathbb{Z}}$ whose complexification satisfies

$$\varphi_{\mathbb{C}}\big(V^{p,q}\big) \subseteq W^{p,q} \qquad \text{for all } p, q$$

Equivalently, $\varphi_{\mathbb{C}}(F^p V_{\mathbb{C}}) \subseteq F^p W_{\mathbb{C}}$ for all p (exercise 3.1.1 (2)).

A morphism between structures of different weights is necessarily zero, since the source has $V^{p,q}$ concentrated in $p+q=k$ and the target in $p+q=l$ (exercise 3.1.1 (3)). Hodge structures of all weights therefore form a category that is the direct sum of its weight-k pieces, and no information is lost by fixing the weight.

Proposition 3.1.8: Morphisms Are Strict

Let $\varphi\colon V \to W$ be a morphism of pure Hodge structures of weight k. Then φ is *strict* for the Hodge filtrations:

$$\varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^p W_{\mathbb{C}} = \varphi_{\mathbb{C}}\big(F^p V_{\mathbb{C}}\big) \qquad \text{for every } p$$

***Proof*:**

The inclusion $\varphi_{\mathbb{C}}(F^pV_{\mathbb{C}}) \subseteq \varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^pW_{\mathbb{C}}$ is immediate from $\varphi_{\mathbb{C}}(F^pV_{\mathbb{C}}) \subseteq F^pW_{\mathbb{C}}$ (definition 3.1.7).

For the reverse, let $w \in \varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^pW_{\mathbb{C}}$ and choose $v \in V_{\mathbb{C}}$ with $\varphi_{\mathbb{C}}(v) = w$. Decompose $v = \sum_{r+s=k} v^{r,s}$ into its bihomogeneous components. Since $\varphi_{\mathbb{C}}(V^{r,s}) \subseteq W^{r,s}$, the decomposition of w is

$$w = \sum_{r+s=k} \varphi_{\mathbb{C}}\big(v^{r,s}\big), \qquad \varphi_{\mathbb{C}}\big(v^{r,s}\big) \in W^{r,s}$$

and this is *the* bihomogeneous decomposition of w, the components of a vector being unique. That w lies in $F^pW_{\mathbb{C}}$ says precisely that its components with $r < p$ vanish, so $\varphi_{\mathbb{C}}(v^{r,s}) = 0$ whenever $r < p$. Set $v' := \sum_{r\geq p} v^{r,k-r}$. Then $v' \in F^pV_{\mathbb{C}}$ by definition 3.1.5, and

$$\varphi_{\mathbb{C}}(v') = \sum_{r\geq p} \varphi_{\mathbb{C}}\big(v^{r,k-r}\big) = \sum_{r+s=k} \varphi_{\mathbb{C}}\big(v^{r,s}\big) = w$$

the middle equality because the discarded terms were shown to vanish. Hence $w \in \varphi_{\mathbb{C}}(F^pV_{\mathbb{C}})$, as we sought to show.

Strictness is not a formality. For an arbitrary map of filtered vector spaces the inclusion $\varphi(F^p) \subseteq \varphi(V) \cap F^p$ is typically proper: a vector can land deep in the target filtration without any preimage lying correspondingly deep in the source (exercise 3.1.1 (4)). What rules this out here is the bigrading, which lets a preimage be truncated componentwise. The consequence is that kernels, images and quotients of morphisms of Hodge structures again carry Hodge structures of the same weight, so the weight-k structures form an abelian category; without strictness the induced filtration on an image and the filtration it inherits as a subobject would differ.

Hodge structures are more than a class of objects. They form a tensor category: they can be multiplied, dualized and twisted, and the resulting bigradings are forced. The constructions are recorded here because everything downstream uses them. The Tate twist defined next is the special case $V \otimes \mathbb{Z}(n)$.

Definition 3.1.9: Tensor Constructions on Hodge Structures

Let A be $\mathbb{Z}$, $\mathbb{Q}$ or $\mathbb{R}$ and let V and W be A-Hodge structures of weights k and l. The *tensor product* $V \otimes W$ is the Hodge structure of weight $k+l$ on $V_A \otimes_A W_A$ with

$$(V \otimes W)^{p,q} = \bigoplus_{\substack{p_1+p_2=p \\ q_1+q_2=q}} V^{p_1,q_1} \otimes W^{p_2,q_2}$$

and the *dual* $V^\vee$ is the Hodge structure of weight $-k$ on $\mathrm{Hom}_A(V_A, A)$ with $(V^\vee)^{p,q} = (V^{-p,-q})^\vee$, where $(V^{-p,-q})^\vee \subseteq (V_\mathbb{C})^\vee$ is the annihilator of the summands $V^{r,s}$ with $(r,s) \neq (-p,-q)$. For $a, b \geq 0$ and $m \in \mathbb{Z}$ write

$$T^{a,b,m}(V) = V^{\otimes a} \otimes (V^\vee)^{\otimes b} \otimes \mathbb{Q}(m)$$

an A-Hodge structure of weight $k(a-b)-2m$. The rational case is the one used throughout; the integral case is what definition 3.1.11 needs.

That the bidegrees above define a Hodge structure is a check on the conjugation symmetry: $\overline{(V \otimes W)^{p,q}} = (V \otimes W)^{q,p}$ because conjugation acts factorwise and the index set is symmetric under swapping the two coordinates, and similarly for the dual.

The classes these constructions exist to isolate are the ones that sit in the middle of the bigrading and are rational. They appear informally in section 3.4 and in the Lefschetz $(1,1)$-theorem of section 2.3; here is the definition.

Definition 3.1.10: Hodge Classes

Let W be a rational Hodge structure of weight $2p$. A *Hodge class* of W is an element of

$$\mathrm{Hdg}(W) = W_\mathbb{Q} \cap W^{p,p}$$

the intersection being taken inside $W_\mathbb{C}$. If the weight of W is odd, set $\mathrm{Hdg}(W) = 0$.

Hodge classes are what chapter 6 characterizes group-theoretically and what **??** asks to be algebraic. The definition above is rational and attaches to an abstract Hodge structure. *Intersection Theory* uses an integral group $\mathrm{Hdg}^p(X) \subseteq H^{2p}(X,\mathbb{Z})$ attached to a smooth projective variety, consisting of those integral classes whose image in $H^{2p}(X,\mathbb{C})$ lies in $H^{p,p}(X)$ [8, Hodge Class]. The two are related by the coefficient map: since $H^{2p}(X,\mathbb{Z}) \to H^{2p}(X,\mathbb{C})$ factors through $H^{2p}(X,\mathbb{Q})$, an integral class lands in $H^{p,p}(X)$ if and only if its image in $H^{2p}(X,\mathbb{Q})$ is a Hodge class in the sense above, so $\mathrm{Hdg}^p(X)$ is the preimage of $\mathrm{Hdg}\big(H^{2p}(X,\mathbb{Q})\big)$ along $H^{2p}(X,\mathbb{Z}) \to H^{2p}(X,\mathbb{Q})$. That map kills the torsion subgroup, which consequently lies in $\mathrm{Hdg}^p(X)$ with no condition imposed on it at all. **??** needs the integral group because the integral form of the Hodge conjecture asks for a lift the rational form does not see.

One further construction is needed before the examples, because it is what allows structures of different weights to be compared at all.

Definition 3.1.11: The Tate Structure and Tate Twists

The *Tate structure* $\mathbb{Z}(1)$ is the weight -2 Hodge structure on the lattice $2\pi i\,\mathbb{Z} \subset \mathbb{C}$ with $V^{-1,-1} = \mathbb{C}$ and all other summands zero. For $n \in \mathbb{Z}$ write $\mathbb{Z}(n) = \mathbb{Z}(1)^{\otimes n}$, a weight $-2n$ structure on a rank-one lattice concentrated in bidegree $(-n,-n)$.
The *Tate twist* of a weight-k Hodge structure V is $V(n) := V \otimes_{\mathbb{Z}} \mathbb{Z}(n)$, a Hodge structure of weight $k-2n$ with

$$\big(V(n)\big)^{p,q} = V^{p+n,\,q+n}$$

The lattice of $\mathbb{Z}(1)$ is $2\pi i\,\mathbb{Z}$ rather than $\mathbb{Z}$, and the factor is not decoration. It is the period of the exponential, and it is exactly the constant that appears when the exponential sequence is used to compute Chern classes, as in proposition 2.3.1. Choosing the lattice this way makes the maps that arise geometrically into morphisms of Hodge structures on the nose rather than up to a power of $2\pi i$.

Example 3.1.12: Twisting the Cohomology of a Surface

Let X be a compact Kähler surface, so $H^2(X,\mathbb{Z})$ carries a weight-2 structure with $h^{2,0} = h^{0,2}$ and $h^{1,1}$ as in corollary 2.2.1. Its twist $H^2(X,\mathbb{Z})(1)$ has weight 0, and its bidegrees are shifted so that the former $(1,1)$-part becomes the $(0,0)$-part.

This is the twist under which the cycle class map becomes weight-preserving. A curve $C \subset X$ has a class in $H^2(X,\mathbb{Z})$ of type $(1,1)$ (theorem 2.3.2), and after twisting it lands in the $(0,0)$-part of a weight-0 structure, which is where the class of a point sits for the trivial structure $\mathbb{Z}$. The twist is the bookkeeping that lets the comparison be stated as a map of Hodge structures at all, and **??** takes it up in general.

Exercise 3.1.1

1. Make the correspondence of proposition 3.1.4 explicit: show that for $v \in V_{\mathbb{R}}$ the projection to the $(+i)$-eigenspace is $\pi(v) = \frac{1}{2}(v - iJv)$, that $v = \pi(v) + \overline{\pi(v)}$, and that transporting multiplication by i through π recovers exactly J. (Applied fibrewise to a tangent bundle, the same splitting is the $T^{1,0} \oplus T^{0,1}$ decomposition.[1])

2. Let V, W carry pure Hodge structures of weight k and let $\varphi\colon V_{\mathbb{Z}} \to W_{\mathbb{Z}}$ be a homomorphism. Show that $\varphi_{\mathbb{C}}(V^{p,q}) \subseteq W^{p,q}$ for all p,q if and only if $\varphi_{\mathbb{C}}(F^pV_{\mathbb{C}}) \subseteq F^pW_{\mathbb{C}}$ for all p. (For the converse, use that $\varphi_{\mathbb{C}}$ commutes with conjugation, since φ is defined over $\mathbb{Z}$, together with (3.2).)

3. Let V and W carry pure Hodge structures of weights $k \neq l$. Show that the only morphism of Hodge structures $V \to W$ is zero. Conclude that the category of Hodge structures of all weights is the direct sum of the categories of weight-k structures.

4. Strictness (proposition 3.1.8) is a property of morphisms of Hodge structures, not of filtered maps in general. Give two filtered complex vector spaces $(V, F^\bullet)$ and $(W, F^\bullet)$ and a linear map φ with $\varphi(F^pV) \subseteq F^pW$ for all p for which $\varphi(V) \cap F^1W \neq \varphi(F^1V)$. (Two-dimensional spaces suffice.) Then identify which hypothesis of proposition 3.1.8 your example violates.

5. Let V carry a pure Hodge structure of weight k with Hodge numbers $h^{p,q}$. Verify that $V(n)$ is a Hodge structure of weight $k-2n$, that its Hodge numbers are $h^{p,q}(V(n)) = h^{p+n,q+n}(V)$,

[1] [5, The $T^{1,0} \oplus T^{0,1}$ Decomposition].

and that the symmetry $\overline{V^{p,q}} = V^{q,p}$ is preserved. Then compute the Hodge numbers of $H^1(E,\mathbb{Z})(1)$ for E an elliptic curve, using example 2.1.2 with $n = 1$.

3.2 Polarizations and the Hodge-Riemann Bilinear Relations

Section 3.1 attached to a lattice a bigrading of its complexification and stopped there. Nothing in definition 3.1.1 refers to a bilinear form, and without one the structures of a given numerical type carry no positivity: there is no notion of a bounded family of them, and no constraint separating the bigradings that arise from geometry from those that do not.

Where the missing form comes from is already settled in low degree. On a compact Kähler surface the cup product on H^2 takes a definite sign on each piece of the Hodge decomposition: positive on a class of type $(2,0)$ paired against its conjugate, positive along the Kähler class, and negative on the primitive real $(1,1)$-classes (lemma 2.2.2). This section states the pattern those signs follow in every weight, adopts it as an axiom on a form attached to an abstract Hodge structure, and then identifies the part of the cohomology of a manifold on which the axiom holds.

The signs in lemma 2.2.2 are not arbitrary, and the alternation is governed by $p-q$. The bookkeeping device that makes this uniform across weights is an operator that multiplies the (p,q)-piece by i^{p-q}, absorbing the alternation into the form and leaving a positive quantity behind.

Definition 3.2.1: Polarization of a Hodge Structure

Let V carry a pure Hodge structure of weight k. The *Weil operator* of the structure is the endomorphism C of $V_{\mathbb{C}}$ acting on $V^{p,q}$ as multiplication by i^{p-q}.
A *polarization* of V is a bilinear form $Q\colon V_{\mathbb{Q}} \times V_{\mathbb{Q}} \to \mathbb{Q}$, extended $\mathbb{C}$-bilinearly to $V_{\mathbb{C}}$, satisfying

$$Q(v,w) = (-1)^k\, Q(w,v)$$

together with the two *Hodge-Riemann bilinear relations*:

1. $Q\big(V^{p,q}, V^{p',q'}\big) = 0$ whenever $(p',q') \neq (q,p)$;
2. $i^{p-q}\, Q(v,\bar{v}) > 0$ for every nonzero $v \in V^{p,q}$.

A Hodge structure equipped with a chosen polarization is called *polarized*, and *polarizable* if some polarization exists. When Q takes integer values on $V_{\mathbb{Z}}$ the polarization is called *integral*.

Proposition 3.2.2: Basic Properties of the Weil Operator

Let V carry a pure Hodge structure of weight k with Weil operator C.

1. $C^2 = (-1)^k\,\mathrm{id}$, and C commutes with conjugation, hence preserves $V_{\mathbb{R}}$.
2. For $k = 1$, the restriction of C to $V_{\mathbb{R}}$ is the complex structure J attached to $V^{1,0}$ in proposition 3.1.4.

***Proof*:**
(1) On $V^{p,q}$ the operator C acts as i^{p-q}, so C^2 acts as $i^{2(p-q)} = (-1)^{p-q}$. Since $p+q=k$, the integers $p-q$ and $p+q$ have the same parity, so $(-1)^{p-q} = (-1)^k$ on every summand and $C^2 = (-1)^k \,\mathrm{id}$. For reality, take $v \in V^{p,q}$; then $\bar{v} \in V^{q,p}$ and

$$\overline{C\bar{v}} = \overline{i^{q-p}\,\bar{v}} = \overline{i^{q-p}}\, v = i^{p-q} v = Cv$$

so C commutes with conjugation. An endomorphism of $V_{\mathbb{C}}$ commuting with the conjugation that defines $V_{\mathbb{R}}$ preserves $V_{\mathbb{R}}$.

(2) In weight 1 the summands are $V^{1,0}$ and $V^{0,1}$, on which C acts as $i^1 = i$ and $i^{-1} = -i$ respectively. The complex structure J of proposition 3.1.4 is defined by transporting multiplication by i on $V^{1,0}$ back to $V_{\mathbb{R}}$ along the projection $V_{\mathbb{R}} \hookrightarrow V_{\mathbb{C}} \to V^{1,0}$ along $V^{0,1}$. The $\mathbb{C}$-linear extension of J to $V_{\mathbb{C}}$ therefore acts as i on $V^{1,0}$, and it acts as $-i$ on $V^{0,1} = \overline{V^{1,0}}$, because J is real and conjugation exchanges the two summands. So J and C agree on both summands, hence on $V_{\mathbb{C}}$, and in particular on $V_{\mathbb{R}}$.

The two relations play different roles. The first is a vanishing condition: Q pairs $V^{p,q}$ with $\overline{V^{p,q}} = V^{q,p}$ and with nothing else, so the form breaks into independent blocks indexed by the unordered pairs $\{p,q\}$ with $p+q=k$. The second carries the content, since it converts Q into a positive quantity on each block once the factor i^{p-q} is inserted. The expression $Q(v,\bar{v})$ is exactly the pairing the first relation leaves alive: for $v \in V^{p,q}$ the vector $\bar{v}$ lies in $V^{q,p}$, the unique summand Q does not annihilate against $V^{p,q}$.

The sign $(-1)^k$ is forced by the geometry the definition abstracts. A form built from the cup product in degree k satisfies $\alpha \wedge \beta = (-1)^{k^2}\beta \wedge \alpha = (-1)^k \beta \wedge \alpha$, so it is symmetric in even weight and alternating in odd weight, and the definition records this rather than imposing it.

The first relation also has a compact reading in the language of definition 3.1.11. Condition (1) says exactly that Q is a morphism of Hodge structures $Q\colon V \otimes V \to \mathbb{Z}(-k)$: the target is concentrated in bidegree (k,k) and has weight $2k$, matching the weight of $V \otimes V$, and the bigrading of a tensor product is the convolution of the bigradings, so the morphism condition says Q kills $V^{p,q} \otimes V^{p',q'}$ unless $p+p'=k$ and $q+q'=k$, which with $p+q=p'+q'=k$ is precisely $(p',q')=(q,p)$. Nothing more than condition (1) is captured that way. The morphism condition does not see the interchange of the two factors, so it says nothing about the $(-1)^k$-symmetry, and it is insensitive to size: the zero form is a morphism $V \otimes V \to \mathbb{Z}(-k)$ for every V and polarizes nothing. The positivity condition (2) is therefore extra data of a different kind, and it is what rules out most forms.

A curve shows both conditions, and there they are classical.

Example 3.2.3: The Polarization on the H^1 of a Curve
Let C be a compact Riemann surface, that is, a compact Kähler manifold of complex dimension 1, and set $g = h^{1,0}(C)$. By theorem 2.1.1 the group $V_{\mathbb{Z}} = H^1(C,\mathbb{Z})$ carries a pure Hodge structure of weight 1 with

$$H^1(C,\mathbb{C}) = H^{1,0}(C) \oplus H^{0,1}(C), \qquad H^{1,0}(C) \cong H^0(C,\Omega^1_C)$$

so $V_{\mathbb{C}}$ splits into the holomorphic 1-forms and their conjugates, and $h^{1,0} = h^{0,1} = g$ by corollary 2.2.1. Take

$$Q(\alpha,\beta) = \int_C \alpha \wedge \beta$$

the cup-product form, which is integral on $H^1(C,\mathbb{Z})$ and alternating, as $(-1)^k = -1$ requires.

The first relation. A class in $H^{1,0}$ is represented by a holomorphic 1-form: the isomorphism $H^{1,0} \cong H^0(C,\Omega^1_C)$ above identifies it with one, and a holomorphic 1-form α is closed, since $\bar\partial\alpha = 0$ by holomorphy while $\partial\alpha$ is a $(2,0)$-form on a curve and hence zero. For $\alpha,\beta \in H^{1,0}$ the product $\alpha\wedge\beta$ is again of type $(2,0)$, so it vanishes and $Q(H^{1,0},H^{1,0}) = 0$; conjugating gives $Q(H^{0,1},H^{0,1}) = 0$. The surviving pairing is $H^{1,0}$ against $H^{0,1} = \overline{H^{1,0}}$, which is condition (1).

The second relation. Let $0 \neq \alpha \in H^{1,0}$ and write $\alpha = f\,dz$ in a holomorphic chart with coordinate $z = x + iy$. Then

$$\alpha\wedge\bar\alpha = |f|^2\,dz\wedge d\bar z = -2i\,|f|^2\,dx\wedge dy$$

so $i\,\alpha\wedge\bar\alpha = 2|f|^2\,dx\wedge dy$. The left side is a globally defined real 2-form on C, and the computation shows it is nonnegative in every chart and vanishes at a point exactly where α does. Hence

$$i^{1-0}\,Q(\alpha,\bar\alpha) = i\int_C \alpha\wedge\bar\alpha > 0$$

for $\alpha \neq 0$, which is condition (2) for the summand $H^{1,0}$. For $H^{0,1}$ write $\alpha = \bar\beta$ with $\beta \in H^{1,0}$; then $i^{0-1}Q(\bar\beta,\beta) = -i\int_C \bar\beta\wedge\beta = i\int_C \beta\wedge\bar\beta > 0$, using that 1-forms anticommute.

So $H^1(C,\mathbb{Z})$ with the cup product is an integrally polarized Hodge structure of weight 1. Written out in a basis of $H_1(C,\mathbb{Z})$ these two conditions are the classical Riemann bilinear relations on the period matrix of C; the complex-torus form of that statement belongs to [3], and section 3.4 takes up the weight-one dictionary in general.

The curve exhibits both relations but hides their structure, because in weight 1 there is only one block and only one sign. To see what the relations say in general the right move is to fold the factor i^{p-q} into the form once and for all, which is what the Weil operator was introduced to do. The result is a single Hermitian form whose positivity encodes the whole alternating pattern, and a filtration-level restatement of the first relation that will matter as soon as the structure is allowed to move.

Theorem 3.2.4: The Hodge-Riemann Relations in Arbitrary Weight

Let V carry a pure Hodge structure of weight k, with Hodge filtration $F^\bullet$ (definition 3.1.5) and Weil operator C, and let Q be a bilinear form on $V_\mathbb{Q}$ with $Q(v,w) = (-1)^k Q(w,v)$, extended $\mathbb{C}$-bilinearly to $V_\mathbb{C}$.

1. Condition (1) of definition 3.2.1 holds if and only if
$$Q\big(F^pV_\mathbb{C},\ F^{k-p+1}V_\mathbb{C}\big) = 0 \qquad \text{for every } p$$
2. If condition (1) holds, then
$$h(u,v) := Q(Cu,\bar v)$$
is a Hermitian form on $V_\mathbb{C}$ for which the decomposition $V_\mathbb{C} = \bigoplus_{p+q=k} V^{p,q}$ is orthogonal.
3. Assume condition (1). Then condition (2) of definition 3.2.1 holds if and only if h is positive definite, if and only if the real bilinear form $S(u,v) := Q(Cu,v)$ on $V_\mathbb{R}$ is symmetric and positive definite.

***Proof*:**

Throughout, C is defined over $\mathbb{R}$: for $v \in V^{p,q}$ one has $\bar{v} \in V^{q,p}$, so $\overline{Cv} = \overline{i^{q-p}\,\bar{v}} = i^{p-q}v = Cv$, and C therefore commutes with conjugation and preserves $V_{\mathbb{R}}$. Since Q is defined over $\mathbb{Q}$, it satisfies $\overline{Q(u,v)} = Q(\bar{u},\bar{v})$.

1. Suppose condition (1) holds. Fix p and take $v \in V^{r,k-r}$ with $r \geq p$ and $w \in V^{s,k-s}$ with $s \geq k-p+1$, so that $v \in F^p$ and $w \in F^{k-p+1}$ range over spanning sets of the two filtration steps. Condition (1) forces $Q(v,w) = 0$ unless $s = k-r$; but $r \geq p$ gives $k-r \leq k-p < k-p+1 \leq s$, so $s \neq k-r$ always. Hence $Q(F^p, F^{k-p+1}) = 0$.

   Conversely, suppose $Q(F^p, F^{k-p+1}) = 0$ for every p, and let $r+s \neq k$. If $r+s > k$ then $s \geq k-r+1$, so $V^{s,k-s} \subseteq F^s \subseteq F^{k-r+1}$ while $V^{r,k-r} \subseteq F^r$, and the hypothesis at $p = r$ gives $Q(V^{r,k-r}, V^{s,k-s}) = 0$. If instead $r+s < k$, put $r' = k-r$ and $s' = k-s$, so that $r'+s' = 2k-(r+s) > k$ and the case just settled gives $Q(V^{r',k-r'}, V^{s',k-s'}) = 0$. Since $\overline{V^{r,k-r}} = V^{k-r,r} = V^{r',k-r'}$ and likewise for s, reality of Q turns this into

   $$Q\big(V^{r,k-r}, V^{s,k-s}\big) = \overline{Q\big(V^{r',k-r'}, V^{s',k-s'}\big)} = 0$$

   Both cases together are condition (1). The second case used reality of Q and nothing else; the filtration condition on its own constrains only the pairs with $r+s > k$.

2. Sesquilinearity is immediate: h is $\mathbb{C}$-linear in u because C and Q are, and $h(u,\lambda v) = Q(Cu, \bar{\lambda}\bar{v}) = \bar{\lambda}\,h(u,v)$.

   For orthogonality, let $u \in V^{p,q}$ and $v \in V^{p',q'}$. Then $Cu \in V^{p,q}$ and $\bar{v} \in V^{q',p'}$, so condition (1) gives $h(u,v) = 0$ unless $(q',p') = (q,p)$, that is unless $(p',q') = (p,q)$. So distinct summands are h-orthogonal.

   For Hermitian symmetry it therefore suffices to treat u, v in the same summand $V^{p,q}$. Using reality of Q and then the symmetry rule,

   $$\overline{h(u,v)} = \overline{i^{p-q}Q(u,\bar{v})} = i^{q-p}\,Q(\bar{u},v) = (-1)^k\,i^{q-p}\,Q(v,\bar{u})$$

   while $h(v,u) = i^{p-q}Q(v,\bar{u})$. The two agree precisely when $(-1)^k i^{q-p} = i^{p-q}$, that is when $(-1)^k = i^{2(p-q)} = (-1)^{p-q}$. Since $p+q = k$, the integers $p-q$ and $p+q$ have the same parity, so $(-1)^{p-q} = (-1)^k$ and the identity holds. Hence $\overline{h(u,v)} = h(v,u)$ and h is Hermitian.

3. For $0 \neq v \in V^{p,q}$ one has $h(v,v) = i^{p-q}Q(v,\bar{v})$, so condition (2) says exactly that h is positive on each nonzero vector of each summand. By part (2) the summands are h-orthogonal, so a general $v = \sum_{p+q=k} v^{p,q}$ has $h(v,v) = \sum_{p+q=k} h(v^{p,q}, v^{p,q})$, and positivity on the summands is equivalent to positivity of h on $V_{\mathbb{C}}$.

   It remains to compare h with S. For $u, v \in V_{\mathbb{R}}$ one has $\bar{v} = v$, so $S(u,v) = Q(Cu,v) = h(u,v)$; this is a real number, being the value of the real form Q on the real vectors Cu and v. Hermitian symmetry of h then reads $S(u,v) = \overline{S(v,u)} = S(v,u)$, so S is symmetric, and h positive definite gives $S(u,u) = h(u,u) > 0$ for $u \neq 0$.

   Conversely suppose S is symmetric and positive definite on $V_{\mathbb{R}}$. Any $w \in V_{\mathbb{C}}$ is $w = u + iv$ with $u, v \in V_{\mathbb{R}}$, and expanding by sesquilinearity,

   $$h(w,w) = h(u,u) + h(v,v) - i\,h(u,v) + i\,h(v,u) = S(u,u) + S(v,v)$$

   the two middle terms cancelling because $h(u,v) = S(u,v) = S(v,u) = h(v,u)$. This is positive unless $u = v = 0$, so h is positive definite, completing the proof.

Part (1) is the statement that will be used whenever the structure is allowed to vary. The bigraded form of the first relation refers to the subspaces $V^{p,q}$, which do not move holomorphically in a family; the filtration form refers only to $F^\bullet$, which does, and it is a closed condition on a flag. Part (3) has an immediate consequence worth recording: a polarization is nondegenerate, since $Q(u,v) = S(C^{-1}u, v)$ and S is a positive definite form on $V_\mathbb{R}$, so no nonzero vector is Q-orthogonal to everything.

The operator C deserves a second look in weight 1, where C acts as i on $V^{1,0}$ and as $-i$ on $V^{0,1}$. That is exactly the complex structure J on $V_\mathbb{R}$ produced in proposition 3.1.4 from the subspace $V^{1,0}$, so a polarization of a weight-1 structure is an alternating form Q on $V_\mathbb{Q}$ for which J is an isometry and $Q(Ju, u) > 0$ for $u \neq 0$ (exercise 3.2.1 (1)). Those are the conditions a complex torus must satisfy to be an abelian variety, and section 3.4 makes the identification.

The signs the polarization takes on $V_\mathbb{R}$ are now determined by the Hodge numbers alone, which is the general form of the phenomenon theorem 2.2.3 exhibited on a surface.

Corollary 3.2.5: Alternating Definiteness and the Signature

Let (V, Q) be a polarized Hodge structure of weight k, with Hodge numbers $h^{p,q}$.

1. Suppose k is even, so Q is symmetric on $V_\mathbb{R}$. For each pair $p + q = k$ the subspace $V^{p,q} \oplus V^{q,p}$ (read as $V^{p,p}$ when $p = q$) is stable under conjugation, and Q restricted to its real points is definite of sign $(-1)^{(p-q)/2}$. Distinct unordered pairs $\{p, q\}$ give Q-orthogonal subspaces, so the signature of Q on $V_\mathbb{R}$ is
$$(b^+, b^-) = \Big(\sum_{\substack{p+q=k \\ p-q \equiv 0 \ (4)}} h^{p,q}, \quad \sum_{\substack{p+q=k \\ p-q \equiv 2 \ (4)}} h^{p,q} \Big)$$
2. Suppose k is odd. Then Q is a nondegenerate alternating form on $V_\mathbb{R}$, so $\dim_\mathbb{R} V_\mathbb{R}$ is even, and $S(u, v) = Q(Cu, v)$ is a positive definite symmetric form.

***Proof*:**

1. Conjugation exchanges $V^{p,q}$ and $V^{q,p}$, so $W_{p,q} := V^{p,q} \oplus V^{q,p}$ is conjugation-stable and its real points $W_{p,q} \cap V_\mathbb{R}$ form a real subspace of dimension $\dim_\mathbb{C} W_{p,q}$. By condition (1) of definition 3.2.1 the form Q pairs $V^{p,q}$ only with $V^{q,p}$, so $W_{p,q}$ and $W_{p',q'}$ are Q-orthogonal whenever $\{p, q\} \neq \{p', q'\}$, and $V_\mathbb{R}$ is the Q-orthogonal direct sum of the $W_{p,q} \cap V_\mathbb{R}$ as $\{p, q\}$ runs over the unordered pairs with $p + q = k$.

   Fix such a pair with $p > q$. A real vector of $W_{p,q}$ has the form $w = v + \bar{v}$ for a unique $v \in V^{p,q}$, and $v \mapsto v + \bar{v}$ is an $\mathbb{R}$-linear isomorphism from $V^{p,q}$ (viewed as a real space of dimension $2h^{p,q}$) onto $W_{p,q} \cap V_\mathbb{R}$. Since $p \neq q$, condition (1) gives $Q(v, v) = 0 = Q(\bar{v}, \bar{v})$, so
$$Q(w, w) = 2\, Q(v, \bar{v}) = 2\, i^{q-p}\, h(v, v)$$
   by the definition of h in theorem 3.2.4. As k is even, $p - q$ is even and $i^{q-p} = (-1)^{(q-p)/2} = (-1)^{(p-q)/2}$; and $h(v, v) > 0$ for $v \neq 0$. So Q is definite on $W_{p,q} \cap V_\mathbb{R}$ of sign $(-1)^{(p-q)/2}$, contributing $2h^{p,q}$ to b^+ or to b^- according as $p - q \equiv 0$ or 2 modulo 4.

   For the middle pair $p = q = k/2$, a real vector $w \in V^{p,p} \cap V_\mathbb{R}$ satisfies $\bar{w} = w$, so $Q(w, w) = Q(w, \bar{w}) = i^0 h(w, w) = h(w, w) > 0$; this contributes $h^{k/2,k/2}$ to b^+, consistent

with $p - q = 0 \equiv 0$ modulo 4. Summing the contributions over the unordered pairs is the same as summing $h^{p,q}$ over all ordered (p, q) with $p + q = k$, since $p - q$ and $q - p$ are congruent modulo 4 to the same residue among $\{0, 2\}$. This gives the stated signature.

2. For k odd, $Q(v, w) = -Q(w, v)$, so Q is alternating. Part (3) of theorem 3.2.4 makes $S(u, v) = Q(Cu, v)$ symmetric and positive definite, hence nondegenerate; since C is invertible and $Q(u, v) = S(C^{-1}u, v)$, the form Q is nondegenerate as well. A nondegenerate alternating form on a real vector space forces the dimension to be even, as we sought to show.

The corollary is the precise sense in which positivity alternates with the bigrading. Moving one step along the row of the Hodge diamond raises p by one and q by one less, so $p - q$ changes by 2 and $(p - q)/2$ changes by 1: the sign of Q flips at every step. In even weight the polarization is therefore definite exactly when the nonzero Hodge numbers all sit in bidegrees with $p - q \equiv 0$ modulo 4, or all in bidegrees with $p - q \equiv 2$ modulo 4; as soon as both residues occur, both b^+ and b^- are nonzero and the form is indefinite. Several bidegrees may be occupied and the form still definite, as the weight-2 case $h^{2,0} = h^{0,2} = 1$, $h^{1,1} = 0$ shows. Either way the signature is read off the Hodge numbers without any further information about V. In odd weight there is no signature to compute, and the content of the corollary is the pair consisting of a symplectic form Q and a compatible operator C making S positive definite, the data that will reappear as a period matrix.

What remains is to say where such a form comes from on a manifold. The obvious candidate, the cup product on $H^k(X, \mathbb{R})$, is not itself a polarization: on a surface it fails condition (2) whichever overall sign is chosen, because the Kähler class and a class of type $(2, 0)$ require opposite signs (exercise 3.2.1 (5)). The repair is to cut H^k down to the part on which the Kähler class acts trivially in the appropriate sense.

Definition 3.2.6: Primitive Cohomology

Let X be a compact Kähler manifold of complex dimension n with Kähler form ω, and let $L = \omega \wedge (-)$ be the Lefschetz operator of definition 1.1.8, acting on cohomology by $L[\alpha] = [\omega \wedge \alpha]$. For $0 \leq k \leq n$ the *primitive cohomology* in degree k is

$$H^k(X, \mathbb{R})_{\text{prim}} := \ker\Big(L^{n-k+1} \colon H^k(X, \mathbb{R}) \to H^{2n-k+2}(X, \mathbb{R})\Big)$$

and $H^{p,q}(X)_{\text{prim}} := H^{p,q}(X) \cap H^k(X, \mathbb{C})_{\text{prim}}$ for $p + q = k$. When $[\omega]$ lies in the image of $H^2(X, \mathbb{Q})$ the same kernel taken on $H^k(X, \mathbb{Q})$ defines $H^k(X, \mathbb{Q})_{\text{prim}}$.

That the primitive part inherits the bigrading is the first thing to check, and it is what makes the polarization below a polarization of a Hodge structure rather than of a bare vector space.

Proposition 3.2.7: Primitive Cohomology Is a Sub-Hodge Structure

Let X be a compact Kähler manifold of dimension n with Kähler form ω, and let $0 \leq k \leq n$. Then

$$H^k(X,\mathbb{C})_{\text{prim}} = \bigoplus_{p+q=k} H^{p,q}(X)_{\text{prim}}, \qquad \overline{H^{p,q}(X)_{\text{prim}}} = H^{q,p}(X)_{\text{prim}}$$

so the primitive cohomology carries a weight-k bigrading with the symmetry required by definition 3.1.1. It is a Hodge structure in the sense of that definition as soon as it is defined over $\mathbb{Q}$, and it is so defined whenever the Kähler class is rational.

***Proof*:**

The operator L is well defined on cohomology because ω is closed: for closed α the form $\omega \wedge \alpha$ is closed, and for $\alpha = d\beta$ it is $\pm\, d(\omega \wedge \beta)$, hence exact. It raises the bidegree by $(1,1)$: a class in $H^{p,q}(X)$ is represented by a d-closed form of type (p,q) (the metric-free description recorded after theorem 2.1.1), and wedging with the $(1,1)$-form ω leaves a d-closed form of type $(p+1,q+1)$.

Consequently L^{n-k+1} carries $H^{p,q}(X)$ into $H^{p+n-k+1,\,q+n-k+1}(X)$, and distinct bidegrees in the source have distinct bidegrees in the target. A class annihilated by L^{n-k+1} therefore has each of its bihomogeneous components annihilated separately, which is the asserted direct sum. Since ω is a real form, L commutes with conjugation, so the primitive part is conjugation-stable and $\overline{H^{p,q}_{\text{prim}}} = H^{q,p}_{\text{prim}}$.

For the last statement, a rational Kähler class makes L^{n-k+1} an operator defined over $\mathbb{Q}$, so its kernel is a rational subspace, completing the proof.

Rationality of $[\omega]$ is sufficient and not necessary. Rescaling ω by a positive irrational factor multiplies L^{n-k+1} by a scalar and so leaves the kernel untouched while destroying rationality of the class, and in the degrees where L^{n-k+1} vanishes identically no condition on $[\omega]$ is needed at all.

Rationality of the Kähler class is the point at which the theory stops being about Kähler manifolds and starts being about projective ones. A general Kähler class lies in $H^2(X,\mathbb{R})$ and nowhere better, and then nothing forces H^k_{prim} to be rational: in the degrees where L^{n-k+1} is not identically zero it is in general only a real subspace with a bigrading, carrying no lattice. If instead $X \subseteq \mathbb{C}\mathrm{P}^N$ is a projective manifold and ω is the restriction of the Fubini-Study form, then $[\omega]$ is the restriction of the integral generator of $H^2(\mathbb{C}\mathrm{P}^N,\mathbb{Z})$ (theorem 1.1.7), hence integral, and the primitive cohomology is a genuine rational Hodge structure.

Two degrees are worth recording at once. For $k = 1$ the operator is $L^n \colon H^1 \to H^{2n+1} = 0$, so $H^1(X,\mathbb{R})_{\text{prim}} = H^1(X,\mathbb{R})$: the whole first cohomology is primitive. For $k = n$ the operator is $L \colon H^n \to H^{n+2}$, and on a surface ($n = k = 2$) this is $\alpha \mapsto [\omega \wedge \alpha]$ with values in $H^4(X,\mathbb{R}) \cong \mathbb{R}$, so

$$H^2(X,\mathbb{R})_{\text{prim}} = \left\{ \alpha \in H^2(X,\mathbb{R}) \;\middle|\; \int_X \alpha \wedge \omega = 0 \right\}$$

the cup-product orthogonal complement of the Kähler class. This is the subspace that appeared in Step 3 of the proof of theorem 2.2.3, where the primitive $(1,1)$-classes carried the negative-definite part of the intersection form.

On the primitive part the cup product, suitably normalized, is a polarization, and only part of the

verification is available at this point. For $k \leq n$ the relevant form is

$$Q(\alpha, \beta) = (-1)^{k(k-1)/2} \int_X \alpha \wedge \beta \wedge \omega^{n-k}$$

Its symmetry type is immediate: interchanging α and β costs $(-1)^{k^2} = (-1)^k$, as definition 3.2.1 requires. The first relation is a bidegree count. For $\alpha \in H^{p,q}$ and $\beta \in H^{p',q'}$ with $p+q = p'+q' = k$, the integrand has type $(p+p'+n-k,\ q+q'+n-k)$, and only a form of type (n,n) has nonzero integral over X; this forces $p+p' = k$ and $q+q' = k$, hence $(p',q') = (q,p)$. The second relation is another matter. It rests on an identity of Weil expressing the Hodge star of a primitive form α of type (p,q) through the Lefschetz operator.

Theorem 3.2.8: Weil's Identity

Let X be a compact Kähler manifold of dimension n and let α be a primitive form of type (p,q) with $p+q = k \leq n$. Then

$$\star\alpha = (-1)^{k(k+1)/2}\, i^{p-q}\, \frac{1}{(n-k)!}\, L^{n-k}\alpha$$

The identity is a statement about the exterior algebra at a single point, where L and Λ generate an $\mathfrak{sl}_2$-action whose representation theory splits $\Lambda^\bullet T_x^* X$ into primitive pieces. That machinery is developed in chapter 4 and the proof is given there; the identity is stated here because the polarization is what needs it and nothing else in this chapter does. Applied to $\bar{\alpha}$, which is primitive of type (q,p), and inserted into $\alpha \wedge \star\bar{\alpha} = |\alpha|^2\,\mathrm{vol}$, it gives

$$\alpha \wedge \bar{\alpha} \wedge \omega^{n-k} = (n-k)!\,(-1)^{k(k+1)/2}\, i^{p-q}\, |\alpha|^2\,\mathrm{vol}$$

The coefficient there is not positive, and for odd k not even real: for a real primitive $(1,1)$-class on a surface it is -1, which is part (3) of lemma 2.2.2, and for $k=1$ it is $i^{-1}(n-1)!$. Positivity appears only once the two factors surrounding the integral are restored. The normalizing sign $(-1)^{k(k-1)/2}$ carried by Q meets the $(-1)^{k(k+1)/2}$ above and leaves $(-1)^{k^2}$; the factor i^{p-q} that condition (2) prescribes meets the i^{p-q} above and leaves $(-1)^{p-q} = (-1)^k$; and $(-1)^{k^2+k} = 1$. So for a harmonic representative of a nonzero class in $H^{p,q}(X)_{\mathrm{prim}}$,

$$i^{p-q}\, Q(\alpha, \bar{\alpha}) = (n-k)! \int_X |\alpha|^2\,\mathrm{vol} > 0$$

which is what condition (2) asks. Collecting the three verifications gives a statement worth a label, since the rest of the chapter refers back to this normalization.

Proposition 3.2.9: The Geometric Polarization on Primitive Cohomology

Let X be a compact Kähler manifold of dimension n with Kähler form ω, and let $0 \le k \le n$. If $H^k(X,\mathbb{Q})_{\text{prim}}$ is defined over $\mathbb{Q}$, then

$$Q(\alpha,\beta) = (-1)^{k(k-1)/2} \int_X \alpha \wedge \beta \wedge \omega^{n-k}$$

is a polarization of the weight-k Hodge structure on $H^k(X,\mathbb{Q})_{\text{prim}}$ in the sense of definition 3.2.1. The sign $(-1)^{k(k-1)/2}$ is not optional: without it condition (2) fails, and in weight 2 the polarizing form is the *negative* of the cup product.

***Proof Key ideas*:**
The symmetry type, the first relation and the second relation are the three verifications just carried out; only the second uses anything beyond bookkeeping, and what it uses is theorem 3.2.8, whose proof belongs to chapter 4. That is why the proof is given as key ideas rather than in full here; the general positivity statement is proved there. The two cases the present chapter settles directly are weight 1 in any dimension (exercise 3.2.1 (2)) and weight 2 on a surface, which is the case Part I already computed, completing the proof.

Example 3.2.10: The Polarized Weight-Two Structure of a Surface

Let X be a compact Kähler surface with Kähler form ω, so $n = k = 2$ and $\omega^{n-k} = 1$. The normalizing sign is $(-1)^{k(k-1)/2} = (-1)^1$, so the candidate polarization on $H^2(X,\mathbb{R})_{\text{prim}}$ is minus the cup form,

$$Q(\alpha,\beta) = -\int_X \alpha \wedge \beta$$

which is symmetric, as $(-1)^k = 1$ requires.

The decomposition. By $\int_X \omega \wedge \omega > 0$ (lemma 2.2.2, part (2)) the Kähler class spans a line on which the cup form is nonzero, so

$$H^2(X,\mathbb{R}) = \mathbb{R}[\omega] \oplus H^2(X,\mathbb{R})_{\text{prim}}$$

the primitive part being the cup-orthogonal complement described above. A class of type $(2,0)$ is automatically primitive, since $\omega \wedge \alpha$ is then of type $(3,1)$ and vanishes on a surface, and the same holds in type $(0,2)$; the primitive $(1,1)$-classes are the space $P^{1,1}(X) = \{\beta \in H^{1,1} \,|\, \beta \wedge \omega = 0\}$ of lemma 2.2.2. On the real $(1,1)$-classes the map $\beta \mapsto \int_X \beta \wedge \omega$ is onto $\mathbb{R}$, because $[\omega]$ itself has positive image, so $\dim_{\mathbb{R}}\big(P^{1,1}(X) \cap H^2(X,\mathbb{R})\big) = h^{1,1} - 1$ and the primitive Hodge numbers are

$$h^{2,0}_{\text{prim}} = h^{0,2}_{\text{prim}} = h^{2,0}, \qquad h^{1,1}_{\text{prim}} = h^{1,1} - 1$$

The two relations. The first is the bidegree count already given, and on a surface it says only that $H^{2,0}$ pairs with $H^{0,2}$ and $H^{1,1}$ with itself. For the second, take $0 \neq \alpha \in H^{2,0}$; then $p - q = 2$ and

$$i^2\, Q(\alpha,\bar\alpha) = (-1) \cdot \Big(-\int_X \alpha \wedge \bar\alpha\Big) = \int_X \alpha \wedge \bar\alpha > 0$$

by part (1) of lemma 2.2.2. For $\alpha \in H^{0,2}$ write $\alpha = \bar\beta$ with $\beta \in H^{2,0}$; then $p - q = -2$, $i^{-2} = -1$, and $\int_X \bar\beta \wedge \beta = \int_X \beta \wedge \bar\beta$ since 2-forms commute, so the same computation applies. For

$0 \neq \beta \in P^{1,1}(X)$ one has $p - q = 0$. The space $P^{1,1}(X)$ is stable under conjugation, since $H^{1,1}$ is and ω is real, so $\beta = \beta_1 + i\beta_2$ with $\beta_1 = \frac{1}{2}(\beta + \bar{\beta})$ and $\beta_2 = \frac{1}{2i}(\beta - \bar{\beta})$ real primitive $(1,1)$-classes, and

$$\beta \wedge \bar{\beta} = \beta_1 \wedge \beta_1 + \beta_2 \wedge \beta_2$$

the cross terms cancelling because 2-forms commute. Part (3) of lemma 2.2.2 makes each summand integrate to a nonpositive number, strictly negative unless the corresponding β_j vanishes, so the total is negative unless $\beta = 0$. Hence $Q(\beta, \bar{\beta}) = -\int_X \beta \wedge \bar{\beta} > 0$. So Q polarizes $H^2(X, \mathbb{R})_{\text{prim}}$, and integrally so whenever $[\omega]$ is an integral class.

Recovering the index theorem. With the polarization in hand, corollary 3.2.5 computes the signature of Q on the primitive part without any further geometry. The residues are $p - q = \pm 2 \equiv 2$ for the types $(2,0)$ and $(0,2)$, and $p - q = 0$ for type $(1,1)$, so

$$(b^+, b^-)\big(Q\big|_{H^2_{\text{prim}}}\big) = \big(h^{1,1} - 1,\ 2h^{2,0}\big)$$

Since Q is minus the cup form, the latter has signature $(2h^{2,0}, h^{1,1} - 1)$ on the primitive part, and restoring the positive line $\mathbb{R}[\omega]$ gives signature $(2h^{2,0} + 1,\ h^{1,1} - 1)$ on all of $H^2(X, \mathbb{R})$. This is exactly theorem 2.2.3, now obtained as the weight-2 instance of a statement about polarized Hodge structures rather than as a computation with the Hodge star on a four-manifold.

The example shows what the abstraction gave. The signature in theorem 2.2.3 came from lemma 2.2.2, three signs computed in a unitary coframe on a four-manifold, and every step of that computation was tied to the dimension. Corollary 3.2.5 delivers the same statement from the axioms of definition 3.2.1 in every weight at once, with the Hodge numbers as the only input. What it does not deliver is the verification that the axioms hold; that is the geometric content, and on a surface it was lemma 2.2.2 that gave it.

Exercise 3.2.1

1. Let V carry a pure Hodge structure of weight 1, with Weil operator C and associated complex structure $J = C|_{V_{\mathbb{R}}}$ (proposition 3.2.2). Show that a polarization of V is the same thing as an alternating form Q on $V_{\mathbb{Q}}$ satisfying $Q(Ju, Jv) = Q(u, v)$ for all $u, v \in V_{\mathbb{R}}$ and $Q(Ju, u) > 0$ for all $u \neq 0$.

2. Let X be a compact Kähler manifold of complex dimension n with Kähler form ω. Show that $H^1(X, \mathbb{R})$ is entirely primitive, and that

$$Q(\alpha, \beta) = \int_X \alpha \wedge \beta \wedge \omega^{n-1}$$

satisfies both Hodge-Riemann bilinear relations on it. (For the second relation, work at a point in a unitary coframe $\varphi^1, \ldots, \varphi^n$ of $(1,0)$-forms with $\omega = i\sum_j \varphi^j \wedge \bar{\varphi}^j$, as in the proof of lemma 2.2.2, write $\alpha = \sum_j a_j \varphi^j$, and expand ω^{n-1}.)

3. Let (V, Q) be a polarized Hodge structure of weight k, and let $W \subseteq V$ be a sub-Hodge structure, meaning a $\mathbb{Q}$-subspace $W_{\mathbb{Q}} \subseteq V_{\mathbb{Q}}$ with $W_{\mathbb{C}} = \bigoplus_{p+q=k} \big(W_{\mathbb{C}} \cap V^{p,q}\big)$. Show that $Q|_W$ is a polarization of W, that $W^{\perp} = \{v \in V_{\mathbb{Q}} \big| Q(v, W_{\mathbb{Q}}) = 0\}$ is again a sub-Hodge structure, and that $V = W \oplus W^{\perp}$ as Hodge structures.

4. Let X be a K3 surface, so $h^{1,0} = 0$, $h^{2,0} = 1$ and $h^{1,1} = 20$ as in exercise 2.1.1 (1). Compute $\dim_{\mathbb{R}} H^2(X, \mathbb{R})_{\text{prim}}$ and its primitive Hodge numbers, then use corollary 3.2.5 to find the

signature of the polarization Q of example 3.2.10. Check that the answer reproduces the signature of the cup form recorded in example 2.2.4.

5. Let X be a compact Kähler surface with $h^{2,0} \geq 1$. Show that neither the cup form $\alpha, \beta \mapsto \int_X \alpha \wedge \beta$ nor its negative is a polarization of the weight-2 Hodge structure $H^2(X, \mathbb{R})$, by exhibiting for each choice of sign a class violating condition (2) of definition 3.2.1. Explain what goes right on $H^2(X, \mathbb{R})_{\text{prim}}$ that fails on $H^2(X, \mathbb{R})$.

6. Let $E = \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$ with $\operatorname{Im}\tau > 0$, and let a, b be the cycles in $H_1(E, \mathbb{Z})$ traced by the segments from 0 to 1 and from 0 to τ. Let $u, v \in H^1(E, \mathbb{Z})$ be the dual basis, normalized by the complex orientation so that $\int_E u \wedge v = 1$. Using that $H^{1,0}(E) = \mathbb{C}\,[dz]$ (example 2.1.2) and that dz has periods $\int_a dz = 1$ and $\int_b dz = \tau$, express $[dz]$ in terms of u and v and verify the second Hodge-Riemann relation for the cup form directly, computing $i\,Q([dz], \overline{[dz]})$ in closed form.

3.3 The Period Domain and Its Compact Dual

Fix the discrete invariants of a polarized Hodge structure: the lattice, the weight, the polarizing form and the Hodge numbers. What is left to vary is the filtration, and a filtration of a fixed vector space with fixed jump dimensions is a point of a flag manifold. This section carries out that identification and reads off the geometry it produces. The two conditions of definition 3.2.1 sit differently inside the flag manifold. The first is a system of bilinear equations, hence a closed condition; the second is a positivity, hence an open one. Imposing the first alone cuts out a compact complex manifold, the compact dual; imposing the second as well cuts out an open subset of it, the period domain.

Both spaces are homogeneous, the compact dual under the complex automorphism group of the polarizing form and the domain under its real points, and the isotropy group of the real action is compact. The metric geometry this suggests, and the distribution along which a family of varieties is constrained to move, belong to chapter 5 and chapter 6; here the task is the construction, the homogeneity that gives the manifold structure, and the two examples the rest of the chapter uses.

Throughout, fix a finitely generated free abelian group $V_{\mathbb{Z}}$ of rank m, an integer k, a nondegenerate bilinear form Q on $V_{\mathbb{Q}}$ satisfying $Q(u, v) = (-1)^k Q(v, u)$ and extended $\mathbb{C}$-bilinearly to $V_{\mathbb{C}}$, and integers $h^{p,q} \geq 0$ for $p + q = k$ with

$$h^{p,q} = h^{q,p}, \qquad \sum_{p+q=k} h^{p,q} = m$$

These are the data of definition 3.2.1 stripped of the filtration. Write F^p for $F^p V_{\mathbb{C}}$, and set

$$f^p = \sum_{r \geq p} h^{r,\,k-r}$$

for the dimension a Hodge filtration with these Hodge numbers must have at level p. Substituting $s = k - r$ and using $h^{r,k-r} = h^{k-r,r}$ turns $f^{k-p+1} = \sum_{r \geq k-p+1} h^{r,k-r}$ into $\sum_{s \leq p-1} h^{s,k-s}$, so the two complementary levels have complementary dimensions:

$$f^p + f^{k-p+1} = m \qquad \text{for every } p \tag{3.4}$$

Finally, assume throughout that at least one polarized Hodge structure with these invariants exists. This is automatic in the geometric case, where the data is read off a compact Kähler manifold through theorem 2.1.1 and theorem 3.2.4, and it is what keeps the spaces below from being empty.

The first of the two conditions in definition 3.2.1 is the orthogonality $Q(V^{p,q}, V^{r,s}) = 0$ except when $(r,s) = (q,p)$. It can be read off the filtration alone:

$$Q\big(F^p,\, F^{k-p+1}\big) = 0 \qquad \text{for every } p \tag{3.5}$$

On a Hodge structure the two versions agree, which is part (1) of theorem 3.2.4. The filtration form has the advantage of making sense for a filtration that is not yet known to come from a bigrading, and that is why it, rather than the bigraded orthogonality, is what cuts out the space below.

Let Fl denote the manifold of decreasing filtrations $F^\bullet$ of $V_\mathbb{C}$ by complex subspaces with $\dim_\mathbb{C} F^p = f^p$. It is a compact complex manifold on which $\mathrm{GL}(V_\mathbb{C})$ acts transitively and holomorphically, and the orbit map admits local holomorphic sections [37]. Cutting it with (3.5) is the first of the two steps.

Definition 3.3.1: The Compact Dual

The *compact dual* attached to the data $(V_\mathbb{Z}, Q, k, \{h^{p,q}\})$ is

$$\check{D} = \big\{F^\bullet \in \mathrm{Fl} \,\big|\, Q(F^p, F^{k-p+1}) = 0 \text{ for every } p\big\}$$

the set of filtrations of the prescribed type satisfying the first bilinear relation (3.5).

By (3.4) and the nondegeneracy of Q, the subspace $(F^p)^\perp$ has dimension $m - f^p = f^{k-p+1}$, so the condition $Q(F^p, F^{k-p+1}) = 0$, which says $F^{k-p+1} \subseteq (F^p)^\perp$, is an equality of subspaces:

$$\check{D} = \big\{F^\bullet \in \mathrm{Fl} \,\big|\, (F^p)^\perp = F^{k-p+1} \text{ for every } p\big\}$$

A point of $\check{D}$ is a self-dual flag. The defining condition is closed: in local frames for the tautological subspaces it is the vanishing of the matrix of pairings $Q(v_i(F), w_j(F))$, whose entries depend continuously on F. So $\check{D}$ is a closed subset of the compact Fl, hence compact, which is what the name records. It is also cut out by algebraic equations, so once it is known to be a complex submanifold it is a smooth projective variety by GAGA ([4, Serre's GAGA Theorems]); smoothness comes from proposition 3.3.5 below.

Nothing in $\check{D}$ refers to the lattice. Conjugation on $V_\mathbb{C}$ has not been used, and a self-dual flag need not be transverse to its own conjugate, so a point of $\check{D}$ need not define a Hodge structure at all. The second condition repairs this and does so with a strict inequality.

Definition 3.3.2: The Period Domain

The *period domain* attached to the data $(V_\mathbb{Z}, Q, k, \{h^{p,q}\})$ is the subset

$$D = \Big\{F^\bullet \in \check{D} \;\Big|\; i^{\,p-q}\, Q(v, \bar{v}) > 0 \text{ for all } p+q = k \text{ and all } v \in \big(F^p \cap \overline{F^q}\big) \setminus \{0\}\Big\}$$

of the compact dual.

The positivity is imposed on the subspaces $F^p \cap \overline{F^q}$, which are defined for any flag; no bigrading is presupposed. That a point of D nevertheless *is* a polarized Hodge structure, with the prescribed Hodge numbers, is the first half of proposition 3.3.4. Granting it for the moment, D is exactly the set of polarized Hodge structures on $V_\mathbb{Z}$ of weight k with Hodge numbers $h^{p,q}$ and polarization Q, presented through their Hodge filtrations rather than their bigradings. This is the sense in which D classifies: its points are the objects, not isomorphism classes of them.

One piece of notation packages the positivity condition. Let F be any flag satisfying the transversality (3.1), so that by theorem 3.1.6 the subspaces $V^{p,q} = F^p \cap \overline{F^q}$ form a Hodge structure of weight k, and let $v \mapsto v^{p,q}$ denote the associated projections. Define

$$H_F(u,v) = \sum_{p+q=k} i^{\,p-q}\, Q\big(u^{p,q},\, \overline{v^{p,q}}\big) \tag{3.6}$$

This form is Hermitian. Conjugating a term and using that Q is defined over $\mathbb{R}$, then that $Q(a,b) = (-1)^k Q(b,a)$,

$$\overline{i^{\,p-q} Q\big(u^{p,q}, \overline{v^{p,q}}\big)} = i^{\,q-p} Q\big(\overline{u^{p,q}}, v^{p,q}\big) = (-1)^k i^{\,q-p} Q\big(v^{p,q}, \overline{u^{p,q}}\big)$$

and $(-1)^k i^{\,q-p} = i^{\,2k+q-p} = i^{\,p-q}$, the two exponents differing by $2k + 2(q-p) = 4q$ when $p+q=k$. So $\overline{H_F(u,v)} = H_F(v,u)$, and in particular $H_F(v,v)$ is real. Comparing (3.6) with definition 3.3.2, a flag $F \in \check{D}$ satisfying (3.1) lies in D precisely when H_F is positive definite: the sum in (3.6) evaluated at $u = v$ is a sum of the quantities $i^{p-q} Q(v^{p,q}, \overline{v^{p,q}})$, each of which the positivity condition makes strictly positive unless the component vanishes.

Example 3.3.3: The Siegel Upper Half Space

Take $k = 1$ and $h^{1,0} = h^{0,1} = g$, so $m = 2g$ and Q is a nondegenerate alternating form on $V_{\mathbb{Q}}$. Here $f^1 = g$ and $f^0 = 2g$, so Fl is the Grassmannian of g-dimensional subspaces of $V_{\mathbb{C}}$ and a flag is a single subspace $L = F^1$. The relation (3.5) at $p = 1$ reads $Q(L,L) = 0$, so

$$\check{D} = \mathrm{LG} = \{\, L \in \mathrm{Fl} \,\big|\, L \text{ is Lagrangian} \,\}$$

the Lagrangian Grassmannian, and the positivity condition at $p = 1$ reads $i\,Q(v,\bar{v}) > 0$ for $0 \neq v \in L$ (the condition at $p = 0$ is its conjugate and adds nothing).

Choose a symplectic basis $e_1, \dots, e_g, f_1, \dots, f_g$ of $V_{\mathbb{Q}}$, which exists because Q is a nondegenerate alternating form on a $\mathbb{Q}$-vector space, so $Q(e_a, f_b) = \delta_{ab}$ and $Q(e_a, e_b) = Q(f_a, f_b) = 0$; it is in particular a $\mathbb{C}$-basis of $V_{\mathbb{C}}$, and in coordinates $Q(x,y) = x^T J_0\, y$ with $J_0 = \left(\begin{smallmatrix} 0 & I \\ -I & 0 \end{smallmatrix}\right)$. Present L as the column span of a $2g \times g$ matrix $\binom{\Lambda}{M}$ of rank g. Then $Q(v,\bar{v}) = c^T\big(\Lambda^T \bar{M} - M^T \bar{\Lambda}\big)\bar{c}$ for v the column combination with coefficient vector c, and the Lagrangian condition is $\Lambda^T M = M^T \Lambda$.

The block Λ is invertible on D. Rewriting the two terms as $(\Lambda c)^T \bar{M} \bar{c}$ and $(Mc)^T \overline{\Lambda c}$ shows that each carries a factor Λc or its conjugate, so $\Lambda c = 0$ forces $Q(v,\bar{v}) = 0$; and $v \neq 0$ for $c \neq 0$ because the columns are independent, contradicting positivity. Normalizing on the right by Λ^{-1} replaces the presenting matrix by $\binom{I}{Z}$ with $Z = M\Lambda^{-1}$, which changes no subspace. In this normalization the Lagrangian condition becomes $Z = Z^T$ and the positivity becomes

$$i\,Q(v,\bar{v}) = i\,c^T\big(\bar{Z} - Z\big)\bar{c} = 2\,c^T\big(\operatorname{Im} Z\big)\bar{c} > 0$$

for all $c \neq 0$. Writing $d = \bar{c}$ turns the last expression into $2\,d^*(\operatorname{Im} Z)d$, so the condition says that the real symmetric matrix $\operatorname{Im} Z$ is positive definite. Hence

$$D \cong \mathfrak{H}_g = \big\{ Z \in \mathbb{C}^{g\times g} \,\big|\, Z = Z^T,\ \operatorname{Im} Z > 0 \big\}$$

the Siegel upper half space, of complex dimension $g(g+1)/2$; for $g = 1$ it is the upper half plane (exercise 3.3.1 (2)). The polarized weight-one structure on H^1 of a compact Riemann surface (example 3.2.3) is a point of it, and its coordinates Z are the classical period matrix. The identification of $\mathfrak{H}_g$ with the parameter space for principally polarized abelian varieties belongs to [3]; the Hodge-theoretic half of that dictionary is section 3.4.

The example already exhibits the shape of the general situation: a compact object, the Lagrangian Grassmannian, containing an open piece cut out by a positivity. The next proposition establishes that shape in general, and simultaneously discharges the claim that a point of D is a Hodge structure.

Proposition 3.3.4: The Period Domain Is Open in the Compact Dual

With the data and notation above:

1. Every $F \in D$ satisfies the transversality condition (3.1). The bigrading $V^{p,q} = F^p \cap \overline{F^q}$ it therefore defines is a Hodge structure of weight k with Hodge numbers $h^{p,q}$, polarized by Q; conversely the Hodge filtration of any such polarized structure is a point of D.
2. D is open in $\check{D}$.

***Proof*:**

1. Let $F \in D$ and fix p. By (3.4) the dimensions of F^p and $\overline{F^{k-p+1}}$ add to m, so (3.1) holds as soon as $F^p \cap \overline{F^{k-p+1}} = 0$. Let v lie in that intersection. Since $F^{k-p+1} \subseteq F^{k-p}$, the vector v lies in $F^p \cap \overline{F^{k-p}}$, so the positivity condition of definition 3.3.2 applies to it. On the other hand $v \in F^p$ and $\bar{v} \in F^{k-p+1}$, so $Q(v, \bar{v}) = 0$ by (3.5). The positivity condition then forces $v = 0$, which gives (3.1).

   By theorem 3.1.6 the subspaces $V^{p,q} = F^p \cap \overline{F^q}$ now form a Hodge structure of weight k with $F^p = \bigoplus_{r \geq p} V^{r,k-r}$; comparing dimensions, $\dim_{\mathbb{C}} V^{p,k-p} = f^p - f^{p+1} = h^{p,k-p}$, so the Hodge numbers are the prescribed ones. The bigraded orthogonality required by definition 3.2.1 follows from (3.5) by the equivalence established before definition 3.3.1, and the positivity required there is the defining condition of D. So Q polarizes the structure. Conversely, the Hodge filtration of a polarized structure with these invariants has $\dim F^p = f^p$, satisfies (3.5) by the same equivalence, and satisfies the positivity because $F^p \cap \overline{F^q} = V^{p,q}$ there; so it is a point of D.

2. Write $U \subseteq \mathrm{Fl}$ for the set of flags satisfying (3.1) and $U^+ \subseteq U$ for the set of flags in U at which H_F is positive definite. By part (1), $D = \check{D} \cap U \cap U^+$. It suffices to prove that U is open in Fl and that U^+ is open in U.

   Fix $F_0 \in U$. Since $\mathrm{GL}(V_{\mathbb{C}})$ acts transitively and holomorphically on Fl with local holomorphic sections of the orbit map [37], there are a neighbourhood W of F_0 and a holomorphic $\gamma \colon W \to \mathrm{GL}(V_{\mathbb{C}})$ with $\gamma(F_0) = \mathrm{id}$ and $\gamma(F) F_0^r = F^r$ for every r and every $F \in W$. Fix p, a basis $a_1, \ldots, a_{f^p}$ of F_0^p and a basis $b_1, \ldots, b_{f^{k-p+1}}$ of F_0^{k-p+1}, and let $M_p(F)$ be the $m \times m$ matrix whose columns are the vectors $\gamma(F)a_i$ followed by the vectors $\overline{\gamma(F)b_j}$. Its columns span F^p and $\overline{F^{k-p+1}}$ respectively, and by (3.4) there are exactly m of them, so (3.1) holds at p if and only if $\det M_p(F) \neq 0$. The entries of M_p are continuous in F, and $\det M_p(F_0) \neq 0$; only finitely many p impose a nonvacuous condition, so all the determinants are nonzero on a neighbourhood of F_0. Hence U is open.

   On U let $A_p(F) = M_p(F)\, E_p\, M_p(F)^{-1}$, where E_p is the diagonal idempotent selecting the first f^p coordinates. Then $A_p(F)$ is the projection onto F^p with kernel $\overline{F^{k-p+1}}$, and by Cramer's rule it depends continuously on F. The computation in Step 1 of theorem 3.1.6 identifies $F^p = \bigoplus_{r \geq p} V^{r,k-r}$ and $\overline{F^{k-p+1}} = \bigoplus_{r \leq p-1} V^{r,k-r}$, so $A_p(F)$ is the projection onto the summands with first index at least p, and therefore $A_p(F) - A_{p+1}(F)$ is the

projection $v \mapsto v^{p,k-p}$. Substituting into (3.6) exhibits the matrix of H_F as a continuous function of F on U.

Fix now $F_1 \in U^+$ and an auxiliary Hermitian inner product on $V_{\mathbb{C}}$ with unit sphere S, which is compact. The function $(F, v) \mapsto H_F(v, v)$ is continuous on $U \times S$ and strictly positive on $\{F_1\} \times S$, so it has a positive minimum c there. For each $v \in S$ continuity gives a product neighbourhood $W_v \times S_v$ of (F_1, v) on which $H_F(w, w) > c/2$; finitely many S_v cover S, and on the intersection of the corresponding W_v the form H_F is positive on all of S, hence positive definite. So U^+ is open in U, and $D = \check{D} \cap U \cap U^+$ is open in $\check{D}$, as we sought to show.

The two halves of definition 3.2.1 have now been separated by their topological character. The first relation is a closed condition and produces a compact object; the second is open and produces a domain inside it. The difference is visible already in example 3.3.3: the Lagrangian Grassmannian is compact, while $\mathfrak{H}_g$ is connected and carries the nonconstant holomorphic functions $Z \mapsto Z_{ab}$, which a connected compact complex manifold does not admit,[2] so it is not compact. In general D, being open in $\check{D}$, is compact only when it is a union of connected components of $\check{D}$. What is still missing is that $\check{D}$ is a manifold at all, so that "open subset" upgrades to "open complex submanifold". That comes from a group action.

Proposition 3.3.5: Homogeneity of the Domain and Its Dual

Let $G_{\mathbb{C}} = \{g \in \mathrm{GL}(V_{\mathbb{C}}) \,|\, Q(gu, gv) = Q(u, v) \text{ for all } u, v\}$ and let $G_{\mathbb{R}} = G_{\mathbb{C}} \cap \mathrm{GL}(V_{\mathbb{R}})$ be its subgroup of real points. Fix $F_o \in D$ and let $P = \{g \in G_{\mathbb{C}} \,|\, gF_o^p = F_o^p \text{ for all } p\}$ be its stabilizer. Then $G_{\mathbb{C}}$ acts transitively on $\check{D}$ and the orbit map induces a biholomorphism $G_{\mathbb{C}}/P \to \check{D}$, so that $\check{D}$ is a compact complex manifold and D is an open complex submanifold of it. The subgroup $G_{\mathbb{R}}$ acts transitively on D, so that $D \cong G_{\mathbb{R}}/(G_{\mathbb{R}} \cap P)$, and the isotropy group $G_{\mathbb{R}} \cap P$ is compact.

***Proof*:**

Write $\varepsilon = (-1)^k$, so $Q(u, v) = \varepsilon Q(v, u)$, and call a subspace $N \subseteq V_{\mathbb{C}}$ isotropic if $Q(N, N) = 0$.

Step 1: isotropic complements. Let $N \subseteq V_{\mathbb{C}}$ be isotropic of dimension d. There is an isotropic subspace N' of dimension d such that Q restricts to a perfect pairing between N and N' and $V_{\mathbb{C}} = (N \oplus N') \oplus (N \oplus N')^{\perp}$.

The linear map $V_{\mathbb{C}} \to N^*$ sending w to $Q(-, w)|_N$ has kernel $N^{\perp}$, of dimension $m - d$, so it is surjective. Choosing a basis $n_1, \dots, n_d$ of N, pick $u_1, \dots, u_d$ with $Q(n_a, u_b) = \delta_{ab}$ and set $B_{ab} = Q(u_a, u_b)$, so that $B_{ba} = \varepsilon B_{ab}$. Put

$$u'_b = u_b - \frac{\varepsilon}{2} \sum_l B_{lb}\, n_l$$

Then $Q(n_a, u'_b) = \delta_{ab}$ because N is isotropic, and expanding $Q(u'_a, u'_b)$ into $Q(u_a, u_b)$ together with the two cross terms $Q(u_a, u'_b - u_b)$ and $Q(u'_a - u_a, u_b)$, each evaluated from $Q(n_a, u_b) = \delta_{ab}$,

$$Q(u'_a, u'_b) = B_{ab} + \varepsilon\Big(-\frac{\varepsilon}{2} B_{ab}\Big) + \Big(-\frac{\varepsilon}{2} B_{ba}\Big) = B_{ab} - \tfrac{1}{2} B_{ab} - \tfrac{1}{2} B_{ab} = 0$$

[2] [5, Maximum Principle on Compact Complex Manifolds].

using $\varepsilon^2 = 1$ and $B_{ba} = \varepsilon B_{ab}$. Let $N' = \operatorname{span}(u'_1, \dots, u'_d)$; it is isotropic by the display, and $Q(n_a, u'_b) = \delta_{ab}$ makes the pairing between N and N' perfect. The Gram matrix of Q on $N \oplus N'$ in the bases (n_a), (u'_b) is $\left(\begin{smallmatrix} 0 & I \\ \varepsilon I & 0 \end{smallmatrix}\right)$, which is invertible; so the $2d$ vectors are independent and Q is nondegenerate on their span, whence $V_{\mathbb{C}}$ is the direct sum of that span and its perpendicular.

Step 2: adapted decompositions. Every $F \in \check{D}$ admits a decomposition $V_{\mathbb{C}} = \bigoplus_p A^p$ with $F^p = \bigoplus_{r \geq p} A^r$ and $Q(A^p, A^r) = 0$ whenever $p + r \neq k$.

Induct on the number of levels at which the flag jumps, the statement being understood for a self-dual flag in any nondegenerate ε-symmetric space. The base case is the zero space, where the flag jumps at no level and the decomposition with every $A^p = 0$ serves. So assume the space is nonzero and let p_0 be the largest p with $F^p \neq 0$. Self-duality at $p_0 + 1$ gives $F^{k-p_0} = (F^{p_0+1})^{\perp} = V_{\mathbb{C}}$. If $p_0 < k - p_0$ then $F^{k-p_0} = V_{\mathbb{C}}$ is a nonzero level above p_0, contradicting maximality; so $p_0 \geq k - p_0$. When $p_0 = k - p_0$ the flag jumps once, $F^{p_0} = V_{\mathbb{C}}$, and $A^{p_0} = V_{\mathbb{C}}$ works, the condition $p_0 + p_0 = k$ imposing nothing.

Suppose $p_0 > k - p_0$, so $p_0 \geq k - p_0 + 1$ and $N := F^{p_0} \subseteq F^{k-p_0+1}$ is isotropic by (3.5). Let $d = \dim N = f^{p_0}$, take N' as in Step 1, and set $W = (N \oplus N')^{\perp}$, a nondegenerate space of dimension $m - 2d$. Put $F_1^p = F^p \cap W$. For $k - p_0 < p \leq p_0$ one has $F^p \subseteq F^{k-p_0+1} = N^{\perp} = N \oplus W$ and $F^p \supseteq N$, so $F^p = N \oplus F_1^p$: decomposing $v \in F^p$ along $N \oplus W$, the N-component already lies in F^p, hence so does the W-component. For $p \leq k - p_0$ one has $F^p = V_{\mathbb{C}}$ and $F_1^p = W$, while $F_1^p = 0$ for $p \geq p_0$ because $F^{p_0} = N$ meets W trivially. Counting, $\dim F_1^{k-p_0+1} = f^{k-p_0+1} - d = (m-d) - d = \dim W$, so F_1 equals W for every $p \leq k - p_0 + 1$: it jumps at strictly fewer levels than F. It is also self-dual in W. For $k - p_0 < p \leq p_0$, intersecting $(F^p)^{\perp} = F^{k-p+1} = N \oplus F_1^{k-p+1}$ with W gives F_1^{k-p+1}, since $N \cap W = 0$, while the same intersection computes as

$$(F^p)^{\perp} \cap W = \left(N \oplus F_1^p\right)^{\perp} \cap W = N^{\perp} \cap (F_1^p)^{\perp} \cap W = \{w \in W \,\big|\, Q(w, F_1^p) = 0\}$$

using $W \subseteq N^{\perp}$; outside that range of p both sides are 0 or W and the claim is immediate. By induction $W = \bigoplus_p A_1^p$ adapted to F_1; setting $A^{p_0} = N$, $A^{k-p_0} = N'$ and $A^p = A_1^p$ for $k - p_0 < p < p_0$ gives the required decomposition of $V_{\mathbb{C}}$, since $A_1^{p_0}$ and $A_1^{k-p_0}$ both vanish by the shape of F_1, and since $N \oplus N'$ is Q-orthogonal to W with $Q(N, N) = Q(N', N') = 0$.

Step 3: transitivity on the compact dual. Let $F, F' \in \check{D}$ with adapted decompositions $\{A^p\}$, $\{A'^p\}$ from Step 2. Since Q is nondegenerate and A^p pairs to zero with every summand but A^{k-p}, the induced pairing $A^p \times A^{k-p} \to \mathbb{C}$ is perfect, and likewise for the primed decomposition; in particular $\dim A^p = \dim A^{k-p}$, and both equal $f^p - f^{p+1}$. For each p with $2p > k$ choose any linear isomorphism $\varphi_p \colon A^p \to A'^p$ and let $\varphi_{k-p} \colon A^{k-p} \to A'^{k-p}$ be the unique map with $Q(\varphi_p a, \varphi_{k-p} b) = Q(a, b)$, which exists and is an isomorphism because both pairings are perfect. If k is even, $\varepsilon = 1$ and Q restricts to nondegenerate symmetric forms on $A^{k/2}$ and $A'^{k/2}$ of equal rank; over $\mathbb{C}$ any two such are isometric, so choose an isometry $\varphi_{k/2}$. Let $g = \bigoplus_p \varphi_p$. It carries F to F', and it preserves Q: on a pair of summands with $p + r \neq k$ both sides vanish, on a pair with $p + r = k$ and $p \neq r$ it holds by construction in one order and by ε-symmetry in the other, and on the middle summand it holds because $\varphi_{k/2}$ is an isometry. So $g \in G_{\mathbb{C}}$ and the action is transitive.

Consequently $\check{D}$ is a single orbit of the complex Lie group $G_{\mathbb{C}}$ acting holomorphically on Fl, and it is closed in Fl; a closed orbit is an embedded complex submanifold and the orbit map descends to a biholomorphism $G_{\mathbb{C}}/P \to \check{D}$ ([37]). Since Fl is compact, so is $\check{D}$, and proposition 3.3.4

makes D an open complex submanifold of it.

Step 4: transitivity on the domain. Let $F, F' \in D$, with bigradings $V^{p,q}$, $V'^{p,q}$ and Hermitian forms $H = H_F$, $H' = H_{F'}$ from (3.6), both positive definite by proposition 3.3.4. The two structures have the same Hodge numbers, so for each $p > q$ there is a $\mathbb{C}$-linear isometry $\varphi_{p,q} \colon (V^{p,q}, H) \to (V'^{p,q}, H')$; define $\varphi_{q,p}$ on $V^{q,p} = \overline{V^{p,q}}$ by $\varphi_{q,p}(\bar{u}) = \overline{\varphi_{p,q}(u)}$. If k is even, write $r = k/2$; then $V^{r,r}$ is stable under conjugation and its real points carry Q positive definite, since $H(v,v) = Q(v,\bar{v}) = Q(v,v)$ for real v. The same holds for $V'^{r,r}$, so there is an isometry of the two real quadratic spaces, and $\varphi_{r,r}$ is its complexification.

Let $g = \bigoplus \varphi_{p,q}$. It commutes with conjugation, hence is defined over $\mathbb{R}$, and it carries F to F'. It preserves Q: the form is recovered from H and the bigrading by $Q(u,v) = i^{\,q-p} H(u,\bar{v})$ for $u \in V^{p,q}$ and $v \in V^{q,p}$, and $Q(u,v) = 0$ for all other pairs of summands, so

$$Q(gu, gv) = i^{\,q-p} H'\big(\varphi_{p,q} u, \varphi_{p,q} \bar{v}\big) = i^{\,q-p} H(u,\bar{v}) = Q(u,v)$$

on the summands that pair, using $\overline{gv} = g\bar{v}$ and $\bar{v} \in V^{p,q}$; on the middle summand $\varphi_{r,r}$ was chosen to preserve Q outright. Hence $g \in G_{\mathbb{R}}$ and $G_{\mathbb{R}}$ acts transitively on D.

Finally, an element of $G_{\mathbb{R}} \cap P$ fixes each F_o^p and, being real, fixes each $\overline{F_o^q}$, hence fixes each $V^{p,q}$; preserving Q and commuting with conjugation, it preserves H_{F_o}, which is $i^{\,p-q} Q(u,\bar{v})$ on $V^{p,q}$ by (3.6). So $G_{\mathbb{R}} \cap P$ is a closed subgroup of the unitary group of the positive definite form H_{F_o}, which is compact, completing the proof.

Two consequences follow. First, $\dim_{\mathbb{C}} D = \dim_{\mathbb{C}} \check{D} = \dim_{\mathbb{C}} G_{\mathbb{C}} - \dim_{\mathbb{C}} P$, so the dimension of the period domain is a Lie-theoretic count in the numerical data alone; the weight-one case gives $g(g+1)/2$, as computed by hand in example 3.3.3. Second, compactness of the isotropy group means an inner product on the tangent space at F_o can be averaged over $G_{\mathbb{R}} \cap P$, so D carries a $G_{\mathbb{R}}$-invariant Riemannian metric. Which further invariant structures D carries, and the curvature of the resulting geometry, depend on the Hodge numbers; chapter 6 takes this up, together with the subbundle of the tangent bundle along which a varying Hodge structure is constrained to move, which is Griffiths transversality in chapter 5. The terminology reflects the classical cases: when D is a bounded symmetric domain, $\check{D}$ is its compact dual in the sense of symmetric spaces, and $\mathfrak{H}_g \subseteq \mathrm{LG}$ is the model.

Example 3.3.6: The Period Domain of a Polarized K3 Surface

Take $k = 2$ with $h^{2,0} = h^{0,2} = 1$ and $h^{1,1} = n$, so $m = n+2$ and Q is symmetric. Here $f^2 = 1$ and $f^1 = n+1$, and a flag is a pair $F^2 \subseteq F^1$ of those dimensions.

The compact dual collapses to a hypersurface in a projective space. Writing $F^2 = \mathbb{C}\sigma$, the condition $(F^2)^{\perp} = F^1$ determines $F^1 = \sigma^{\perp}$, whose dimension is $m - 1 = n + 1$ as required, and the inclusion $F^2 \subseteq F^1$ says exactly $Q(\sigma,\sigma) = 0$; conversely any isotropic line assembles into such a flag. So

$$\check{D} \cong \big\{ [\sigma] \in \mathbb{P}(V_{\mathbb{C}}) \,\big|\, Q(\sigma,\sigma) = 0 \big\}$$

a smooth quadric hypersurface in $\mathbb{P}^{n+1}$, of dimension n. The positivity conditions at $p = 2$ and $p = 0$ are conjugate to one another and read $-Q(\sigma,\bar{\sigma}) > 0$, since $i^{2-0} = -1$; the condition at $p = 1$ reads $Q(v,\bar{v}) > 0$ on $F^1 \cap \overline{F^1}$, equivalently that Q is positive definite on the real points of that conjugation-stable space.

Write $\sigma = x + iy$ with $x, y \in V_{\mathbb{R}}$ and expand:

$$Q(\sigma,\sigma) = Q(x,x) - Q(y,y) + 2i\,Q(x,y), \qquad Q(\sigma,\bar\sigma) = Q(x,x) + Q(y,y)$$

So $Q(\sigma,\sigma) = 0$ says $Q(x,x) = Q(y,y)$ and $Q(x,y) = 0$, and then $Q(\sigma,\bar\sigma) = 2Q(x,x) < 0$ says the real plane $\operatorname{span}_{\mathbb{R}}(x,y)$ is negative definite for Q. Its Q-orthogonal complement is the space of real points of $\sigma^\perp \cap \overline{\sigma^\perp}$, so the remaining condition says that complement is positive definite, and the two together force Q to have signature $(n,2)$. Conversely, if Q has signature $(n,2)$ then a negative definite plane has positive definite complement automatically, and

$$D \cong \left\{ [\sigma] \in \mathbb{P}(V_{\mathbb{C}}) \middle| Q(\sigma,\sigma) = 0,\ Q(\sigma,\bar\sigma) < 0 \right\}$$

an open subset of the quadric, of dimension n. Sending $[\sigma]$ to $\operatorname{span}_{\mathbb{R}}(x,y)$ with the orientation (x,y) identifies D with the set of oriented negative definite real planes in $(V_{\mathbb{R}}, Q)$: scaling σ by $re^{i\theta}$ rotates the ordered basis and preserves both plane and orientation, and an oriented negative definite plane admits an oriented basis x, y with $Q(x,x) = Q(y,y)$ and $Q(x,y) = 0$, unique up to that rotation and scaling. Conjugation $[\sigma] \mapsto [\bar\sigma]$ is a fixed-point-free involution of D that reverses this orientation.

For $n = 19$ the data is that of a polarized K3 surface. Let X be a projective K3 surface with polarization class h, an integral $(1,1)$-class of positive square, and let $V_{\mathbb{Z}}$ be the primitive part of $H^2(X,\mathbb{Z})$, the orthogonal complement of h for the cup form. By example 2.2.4 the cup form on $H^2(X,\mathbb{Z}) \cong \mathbb{Z}^{22}$ has signature $(3,19)$ and Hodge numbers $h^{2,0} = 1$, $h^{1,1} = 20$; removing the positive direction spanned by h leaves a lattice of rank 21 with $h^{1,1} = 19$ on which the cup form has signature $(2,19)$. The polarizing form of proposition 3.2.9 is the negative of the cup form in weight two, so Q has signature $(19,2)$, and the plane spanned by the real and imaginary parts of a holomorphic 2-form σ is the one the description above singles out. Passing to the primitive part is not a convenience. On all of $H^2(X,\mathbb{R})$ the Hodge numbers give $n = 20$, but there Q, still minus the cup form, has signature $(19,3)$ rather than the $(20,2)$ just forced: the plane spanned by the real and imaginary parts of σ is negative definite as before, while its complement, the real points of $\sigma^\perp \cap \overline{\sigma^\perp}$, contains the Kähler class, on which $Q([\omega],[\omega]) = -\int_X \omega \wedge \omega < 0$ by part (2) of lemma 2.2.2. So the positivity at $p = 1$ fails there and the unpolarized H^2 of a K3 is not a point of the period domain for $n = 20$, which is exercise 3.2.1 (5) restated inside $\check{D}$. The domain has dimension 19; identifying it as the target of the period map for polarized K3 surfaces, and the injectivity of that map, is the Torelli theorem taken up in chapter 6, with the lattice-theoretic input cited to [4].

Exercise 3.3.1

1. Derive (3.4) from the symmetry $h^{p,q} = h^{q,p}$, and deduce that for a filtration $F^\bullet \in \mathrm{Fl}$ the first bilinear relation (3.5) holds if and only if $(F^p)^\perp = F^{k-p+1}$ for every p. Where is the nondegeneracy of Q used?

2. Take $k = 1$ and $g = 1$ in example 3.3.3, so $m = 2$ and Q is a nondegenerate alternating form on $V_{\mathbb{C}} \cong \mathbb{C}^2$. Show that $\check{D} \cong \mathbb{P}^1$ and that D is the upper half plane, computing $iQ(v,\bar v)$ directly for $v = e + \tau f$ in a symplectic basis. Then exhibit a point of $\check{D} \setminus D$ and verify that both the transversality (3.1) and the positivity fail at it.

3. Let $F_o \in D$ and let $G_{\mathbb{R}} \cap P$ be its isotropy group as in proposition 3.3.5. Show that restriction to the summands with $p > k/2$, together with the middle summand when k is even, gives an

isomorphism

$$G_{\mathbb{R}} \cap P \;\cong\; \prod_{p>k/2} \mathrm{U}\big(h^{p,k-p}\big) \times \mathrm{O}\big(h^{k/2,k/2}\big)$$

where the last factor is present only for k even. Check the answer against example 3.3.3 and example 3.3.6 by comparing dimensions with $\dim_{\mathbb{C}} D$ in each case.

4. In the coordinates of example 3.3.3, write an element of $G_{\mathbb{R}} = \mathrm{Sp}(V_{\mathbb{R}}, Q)$ in block form $g = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}\right)$ relative to the symplectic basis. Show that the action of $G_{\mathbb{R}}$ on $D \cong \mathfrak{H}_g$ furnished by proposition 3.3.5 is
$$Z \longmapsto (\gamma + \delta Z)(\alpha + \beta Z)^{-1}$$
and that $\alpha + \beta Z$ is invertible for every $Z \in \mathfrak{H}_g$ and every g. Specialize to $g = 1$ and identify the action with the Möbius action of $\mathrm{SL}(2,\mathbb{R})$ on the upper half plane.

5. Take $k = 3$ with $h^{3,0} = h^{0,3} = 1$ and $h^{2,1} = h^{1,2} = n$, the numerical type of H^3 of a compact Kähler threefold with a one-dimensional space of holomorphic 3-forms. Show that a point of $\check{D}$ is the same as a pair consisting of a Lagrangian subspace $F^2 \subseteq V_{\mathbb{C}}$ and a line $F^3 \subseteq F^2$, with F^1 determined. Using the normalization of example 3.3.3 to compute the dimension of the Lagrangian Grassmannian, conclude
$$\dim_{\mathbb{C}} \check{D} = \frac{(n+1)(n+2)}{2} + n$$

3.4 First Examples: Weight One and Weight Two

Two weights make the apparatus of section 3.1 through section 3.3 completely explicit, and they are the two the geometry gives first. In weight 1 a Hodge structure is a single subspace $V^{1,0}$ of half the total dimension, and the objects it classifies are complex tori; the polarization becomes the classical Riemann form, and its existence is exactly the condition for the torus to be projective. In weight 2 with $h^{2,0} = 1$, the case of a K3 surface, the structure collapses to a single *line* in $H^2(X,\mathbb{C})$, and the two Hodge-Riemann relations become one quadratic equation and one inequality.

Example 3.1.3 already identified the linear algebra of weight 1: an effective weight-1 structure on $V_{\mathbb{Z}}$ is the same thing as a complex structure J on the real vector space $V_{\mathbb{R}}$, the correspondence sending J to its $(+i)$-eigenspace $V^{1,0}$ in $V_{\mathbb{C}}$. A lattice together with a complex structure on the real space it spans is exactly the data of a complex torus (example 1.1.6), so the dictionary is forced. What has to be checked is that it is a dictionary at the level of morphisms as well, and what has to be computed is what a polarization becomes on the other side.

Theorem 3.4.1: Weight-One Hodge Structures and Complex Tori

Let $g \geq 1$.

1. Let V be an effective pure Hodge structure of weight 1 on a free abelian group $V_{\mathbb{Z}}$ of rank $2g$, let $\pi\colon V_{\mathbb{C}} \to V^{1,0}$ be the projection with kernel $V^{0,1}$, and let J be the complex structure on $V_{\mathbb{R}}$ of example 3.1.3. Then $\pi(V_{\mathbb{Z}})$ is a lattice in the g-dimensional complex vector space $V^{1,0}$, and
$$X_V := V_{\mathbb{C}}/\big(\overline{F^1 V_{\mathbb{C}}} + V_{\mathbb{Z}}\big) = V^{1,0}/\pi(V_{\mathbb{Z}})$$
is a complex torus of dimension g, namely $V_{\mathbb{R}}/V_{\mathbb{Z}}$ carrying the complex structure J.

2. The assignment $V \mapsto X_V$ is an equivalence from the category of effective weight-1 Hodge structures on lattices of rank $2g$, with morphisms of Hodge structures, to the category of complex tori of dimension g, with holomorphic group homomorphisms. A quasi-inverse sends $X = W/\Gamma$ to the weight-1 structure on Γ determined by multiplication by i on $\Gamma \otimes_{\mathbb{Z}} \mathbb{R} = W$. The construction $V \mapsto X_V$ is functorial without reference to a fixed rank: a morphism of weight-1 Hodge structures $V \to V'$, of ranks $2g$ and $2g'$ not necessarily equal, induces a holomorphic homomorphism $X_V \to X_{V'}$, and every such homomorphism arises this way.

3. A bilinear form $Q\colon V_{\mathbb{Z}} \times V_{\mathbb{Z}} \to \mathbb{Z}$ is a polarization of V if and only if it is alternating and satisfies
$$Q(Ju, Jv) = Q(u, v) \qquad \text{for all } u, v \in V_{\mathbb{R}} \tag{3.7}$$
$$Q(Ju, u) > 0 \qquad \text{for all } u \in V_{\mathbb{R}} \setminus \{0\} \tag{3.8}$$
Equivalently, $H(u, v) := Q(Ju, v) + i\,Q(u, v)$ is a positive definite Hermitian form on the complex vector space $(V_{\mathbb{R}}, J)$ whose imaginary part takes integer values on $V_{\mathbb{Z}}$.

Conditions (3.7) and (3.8) are the Riemann bilinear relations in their classical, complex-torus form; a form satisfying them is called a Riemann form of the torus, and it is the object from which [3] builds line bundles on X_V.

***Proof*:**
Write $\Gamma_V := \pi(V_{\mathbb{Z}})$, and begin with two facts used throughout. Conjugation is a conjugate-linear isomorphism $V^{1,0} \to V^{0,1}$ by the symmetry in definition 3.1.1, so $h^{1,0} = h^{0,1}$; since these sum to $\dim_{\mathbb{C}} V_{\mathbb{C}} = 2g$, the space $V^{1,0}$ has complex dimension g. And for $u \in V_{\mathbb{R}}$,
$$u = \tfrac{1}{2}\big(u - iJu\big) + \tfrac{1}{2}\big(u + iJu\big)$$
where the first summand lies in $V^{1,0}$, since $J(u - iJu) = Ju + iu = i(u - iJu)$ exhibits it as a $(+i)$-eigenvector of J, and the second is its conjugate and so lies in $V^{0,1}$. Hence
$$\pi(u) = \tfrac{1}{2}\big(u - iJu\big), \qquad \overline{\pi(u)} = \tfrac{1}{2}\big(u + iJu\big) \qquad (u \in V_{\mathbb{R}}) \tag{3.9}$$

1. By example 3.1.3 the restriction $\pi|_{V_{\mathbb{R}}}\colon V_{\mathbb{R}} \to V^{1,0}$ is an isomorphism of real vector spaces, and J is by construction the pullback along it of multiplication by i. Now $V_{\mathbb{Z}}$ is a discrete

subgroup of rank $2g$ spanning $V_{\mathbb{R}}$, so its image Γ_V is a discrete subgroup of rank $2g$ spanning the real $2g$-dimensional space $V^{1,0}$, that is, a lattice. The quotient $X_V = V^{1,0}/\Gamma_V$ is therefore a complex torus of dimension g (example 1.1.6), and π identifies it with $V_{\mathbb{R}}/V_{\mathbb{Z}}$ carrying the complex structure J. Finally $\overline{F^1 V_{\mathbb{C}}} = \overline{V^{1,0}} = V^{0,1}$ by definition 3.1.5, so $V_{\mathbb{C}}/(V^{0,1} + V_{\mathbb{Z}}) = V^{1,0}/\Gamma_V$, which is the displayed description.

2. *Objects.* Let $X = W/\Gamma$ be a complex torus of dimension g, so Γ is a lattice of rank $2g$ in W and the inclusion extends to an isomorphism $\Gamma \otimes_{\mathbb{Z}} \mathbb{R} \cong W$. Multiplication by i on W is an endomorphism J of $\Gamma_{\mathbb{R}}$ with $J^2 = -\mathrm{id}$, so by proposition 3.1.4 it determines an effective weight-1 Hodge structure V on Γ whose $(1,0)$-part is the $(+i)$-eigenspace of J. The map $\pi|_{\Gamma_{\mathbb{R}}} \colon W \to V^{1,0}$ intertwines multiplication by i on W with multiplication by i on $V^{1,0}$, hence is an isomorphism of complex vector spaces, and it carries Γ onto Γ_V; passing to quotients gives $X \cong X_V$. In the other direction, start from V and form $X_V = V^{1,0}/\Gamma_V$. Its lattice is Γ_V, identified with $V_{\mathbb{Z}}$ by π, and multiplication by i on $V^{1,0}$ pulls back along π to J on $V_{\mathbb{R}}$; so the Hodge structure attached to X_V by the previous sentence is the one determined by J, which is V.

   Morphisms. Let V, V' be effective weight-1 structures with complex structures J, J', and let $\varphi \colon V_{\mathbb{Z}} \to V'_{\mathbb{Z}}$ be a homomorphism. By definition 3.1.7, φ is a morphism of Hodge structures exactly when $\varphi_{\mathbb{C}}(V^{1,0}) \subseteq (V')^{1,0}$. Since φ is defined over $\mathbb{Z}$, its complexification commutes with conjugation, so this inclusion forces $\varphi_{\mathbb{C}}(V^{0,1}) \subseteq (V')^{0,1}$ as well; as J acts by $+i$ on $V^{1,0}$ and by $-i$ on $V^{0,1}$, the two inclusions together say precisely that $\varphi_{\mathbb{R}} \circ J = J' \circ \varphi_{\mathbb{R}}$. Such a φ transports along π and π' to a $\mathbb{C}$-linear map $V^{1,0} \to (V')^{1,0}$ carrying Γ_V into $\Gamma_{V'}$, hence descends to a holomorphic homomorphism $X_V \to X_{V'}$.

   Conversely let $f \colon X_V \to X_{V'}$ be a holomorphic group homomorphism. The universal covers are the vector spaces $V^{1,0}$ and $(V')^{1,0}$, and since $V^{1,0}$ is simply connected, f lifts to a holomorphic $F \colon V^{1,0} \to (V')^{1,0}$ with $F(0) = 0$. That f is a homomorphism says $F(x+y) - F(x) - F(y) \in \Gamma_{V'}$ for all x, y; the left-hand side is continuous in (x, y), vanishes at $(0,0)$, and takes values in a discrete group, so it vanishes identically and F is additive. Differentiating $F(x+y) = F(x) + F(y)$ with respect to y at $y = 0$ gives $dF_x = dF_0$ for every x, so F equals the $\mathbb{C}$-linear map dF_0. It satisfies $F(\Gamma_V) \subseteq \Gamma_{V'}$, and transporting back along π, π' produces a homomorphism $\varphi \colon V_{\mathbb{Z}} \to V'_{\mathbb{Z}}$ commuting with J and J', that is, a morphism of Hodge structures inducing f. The two assignments are inverse to one another, so $\mathrm{Hom}(V, V') \to \mathrm{Hom}(X_V, X_{V'})$ is a bijection. Together with the previous paragraph this makes $V \mapsto X_V$ essentially surjective, full and faithful, hence an equivalence.

3. Specialized to $k = 1$, definition 3.2.1 and theorem 3.2.4 ask of Q that it be $(-1)^1$-symmetric, that is alternating, and that

$$Q\big(F^1 V_{\mathbb{C}},\, F^1 V_{\mathbb{C}}\big) = 0, \qquad i\,Q(v, \bar{v}) > 0 \ \text{ for } 0 \neq v \in V^{1,0}$$

   there being no primitivity restriction in weight 1, where every class is primitive. Extend Q first $\mathbb{R}$-bilinearly to $V_{\mathbb{R}}$ and then $\mathbb{C}$-bilinearly to $V_{\mathbb{C}}$, and recall $F^1 V_{\mathbb{C}} = V^{1,0} = \pi(V_{\mathbb{R}})$.

   For the first relation, (3.9) gives, for $u, v \in V_{\mathbb{R}}$,

$$4\,Q(\pi u, \pi v) = Q\big(u - iJu,\ v - iJv\big) = \big(Q(u,v) - Q(Ju,Jv)\big) - i\big(Q(u,Jv) + Q(Ju,v)\big)$$

   Both bracketed quantities are real, so $Q(V^{1,0}, V^{1,0}) = 0$ holds if and only if $Q(Ju, Jv) = Q(u,v)$ and $Q(u, Jv) = -Q(Ju, v)$ for all real u, v. Each of these two identities implies the other: replacing v by Jv in the first gives $Q(Ju, J^2 v) = Q(u, Jv)$, which is the second,

and replacing v by Jv in the second gives $Q(u, J^2v) = -Q(Ju, Jv)$, which is the first. So the first Hodge-Riemann relation is exactly (3.7).

For the second, (3.9) and the vanishing of $Q(u,u)$ and $Q(Ju,Ju)$ give

$$4\,Q\big(\pi u, \overline{\pi u}\big) = Q\big(u - iJu,\ u + iJu\big) = i\,Q(u, Ju) - i\,Q(Ju, u) = 2i\,Q(u, Ju)$$

whence $i\,Q(\pi u, \overline{\pi u}) = \frac{i^2}{2}Q(u,Ju) = \frac{1}{2}Q(Ju,u)$. Since π maps $V_{\mathbb{R}}$ onto $V^{1,0}$, the positivity condition is exactly (3.8). Nondegeneracy of Q is a consequence rather than an extra hypothesis: if $Q(u,-)$ vanished identically then $Q(Ju,u) = -Q(u,Ju) = 0$, forcing $u = 0$.

It remains to identify H. Using (3.7) in the form $Q(u,Jv) = -Q(Ju,v)$,

$$H(Ju, v) = Q(J^2u, v) + i\,Q(Ju, v) = -Q(u,v) + i\,Q(Ju,v) = i\,H(u,v)$$
$$H(v,u) = Q(Jv,u) + i\,Q(v,u) = -Q(u,Jv) - i\,Q(u,v) = Q(Ju,v) - i\,Q(u,v) = \overline{H(u,v)}$$

so H is $\mathbb{C}$-linear in its first argument and Hermitian, and $H(u,u) = Q(Ju,u)$ is positive for $u \neq 0$ by (3.8); its imaginary part is Q, which is $\mathbb{Z}$-valued on $V_{\mathbb{Z}}$. Conversely, a positive definite Hermitian H on $(V_{\mathbb{R}}, J)$ with $\operatorname{Im} H$ integral on $V_{\mathbb{Z}}$ yields $Q := \operatorname{Im} H$, which is alternating, satisfies (3.7) because $H(Ju,Jv) = i\,\bar{i}\,H(u,v) = H(u,v)$, and satisfies (3.8) because $Q(Ju,u) = \operatorname{Im}\big(i\,H(u,u)\big) = H(u,u) > 0$. The two descriptions carry the same information, completing the proof.

The equivalence says that at weight 1 the abstract theory and classical complex geometry describe the same objects. A complex torus admits no nonconstant holomorphic functions[3] and no obvious algebraic data, yet the pair consisting of its lattice and its complex structure is a complete invariant, and that pair is a Hodge structure. Part (3) adds that the polarization, introduced in section 3.2 as linear algebra, is on the torus side a concrete and classical object: a positive definite Hermitian form with integral imaginary part.

The two relations have different characters, and the difference is what governs projectivity. Relation (3.7) is linear in Q and is a nontrivial condition as soon as $g \geq 2$: it requires the alternating form to be invariant under the complex structure. Relation (3.8) is an open positivity condition, and it selects a cone inside the solutions of the first. The next statement reads this off, and it is the criterion promised in example 1.1.6.

[3] [5, Maximum Principle on Compact Complex Manifolds].

Corollary 3.4.2: Projectivity Criterion for a Complex Torus

Let $X = W/\Gamma$ be a complex torus of dimension g and let V be the weight-1 Hodge structure on Γ of theorem 3.4.1. The following are equivalent.

1. X admits a closed holomorphic embedding into some $\mathbb{C}\mathrm{P}^N$.
2. V admits a polarization.
3. There is an alternating form $E\colon \Gamma \times \Gamma \to \mathbb{Z}$ with $E(iu, iv) = E(u, v)$ for all $u, v \in W$ and $E(iu, u) > 0$ for all $u \in W \setminus \{0\}$.

A complex torus satisfying these conditions is an *abelian variety*. Every complex torus of dimension 1 is an abelian variety, while for $g \geq 2$ a very general complex torus is not.

***Proof*:**

(2) ⇔ (3). The complex structure J of theorem 3.4.1 is multiplication by i on $W = \Gamma_{\mathbb{R}}$, so the two conditions in (3) are (3.7) and (3.8) verbatim, and part (3) of that theorem identifies the forms satisfying them with the polarizations of V.

(1) ⇔ (3). This is the classical criterion for a complex torus to be projective. Given E as in (3), the Appell-Humbert construction attaches to it a holomorphic line bundle whose theta functions embed X; conversely an embedding produces such an E from the hyperplane class. Both directions are proved in [3], and this is the one ingredient of the present section taken from outside the book.

Where E sits in cohomology. The Hodge-theoretic half of the criterion is available here, and it locates the form E inside H^2. The integral cohomology of the torus is the exterior algebra on $H^1(X, \mathbb{Z}) = \mathrm{Hom}(\Gamma, \mathbb{Z})$, so

$$H^2(X, \mathbb{Z}) = \bigwedge^2 \mathrm{Hom}(\Gamma, \mathbb{Z})$$

and an integral degree-2 class is the same thing as an alternating form E on Γ. Write $V_{\mathbb{C}} = W \otimes_{\mathbb{R}} \mathbb{C} = V^{1,0} \oplus V^{0,1}$ for the eigenspace decomposition of multiplication by i. By example 2.1.2 the summand $H^{1,0}(X)$ consists of the $\mathbb{C}$-linear functionals on W, and such a functional ℓ kills $V^{0,1}$, since $\ell(u + iJu) = \ell(u) + i\ell(iu) = \ell(u) - \ell(u) = 0$. Consequently every summand of $\bigwedge^2 H^1(X, \mathbb{C})$ other than $H^{2,0} = \bigwedge^2 H^{1,0}$ restricts to zero on $V^{1,0} \times V^{1,0}$, while $H^{2,0}$ injects into the alternating forms on $V^{1,0}$. So E has vanishing $(2,0)$-component, equivalently, being real, vanishing $(0,2)$-component, if and only if E vanishes on $V^{1,0} \times V^{1,0}$, which by the computation in part (3) of theorem 3.4.1 is the invariance $E(iu, iv) = E(u, v)$. By definition 2.3.3 and theorem 2.3.2 the integral $(1,1)$-classes are exactly $\mathrm{NS}(X)$, and are exactly the first Chern classes of holomorphic line bundles. Condition (3) therefore asks for an element of $\mathrm{NS}(X)$ satisfying a definiteness condition, and it is that condition, not the existence of line bundles, that separates projective tori from the rest. For $g = 2$ this recovers the description of $\mathrm{NS}(X)$ in example 2.3.6.

The two extreme cases. Let $g = 1$, so that W is a real plane and the group of alternating integral forms on Γ has rank 1; fix a generator E_0. Multiplication by i acts on the one-dimensional space of alternating forms on W through $\det(i)$, and $\det(i)^2 = \det(-\mathrm{id}) = 1$ forces $\det(i) = \pm 1$. The

value -1 is impossible: a real 2×2 matrix of determinant -1 has real eigenvalues, whereas an eigenvalue λ of i would satisfy $\lambda^2=-1$. So $\det(i)=1$, the action is trivial, and every alternating form satisfies the invariance in (3). Moreover u and iu are linearly independent over $\mathbb{R}$ for $u\neq 0$, so $E_0(iu,u)\neq 0$; this quantity is continuous and nonvanishing on the connected set $W\setminus\{0\}$, hence of constant sign, and one of $\pm E_0$ satisfies (3).

Let $g\geq 2$ and let X be very general. Such a torus has $\rho(X)=0$: proposition 2.3.4 gives only the bound $\rho\leq h^{1,1}$, and that the floor is attained on a very general torus is proved in [3], which develops the complex-torus theory this example borrows. So $\mathrm{NS}(X)=0$ and, by the previous paragraph, the only invariant integral alternating form on Γ is 0, which fails the positivity in (3). Hence X is not projective, completing the proof.

The criterion is a positivity condition on a lattice, and nothing in it refers to maps into projective space. The question "which complex tori are algebraic" has become the question "which lattices with a complex structure carry an invariant positive alternating form", answerable by counting integral solutions of a linear condition and then testing a sign. For $g=1$ the linear condition is vacuous and the sign can always be arranged, so every one-dimensional complex torus is an elliptic curve; this agrees with exercise 2.3.1 (4), where $\mathrm{NS}(E)=H^2(E,\mathbb{Z})\cong\mathbb{Z}$ for every elliptic curve. For $g\geq 2$ the linear condition generically has no nonzero integral solution, and the generic complex torus carries no divisors at all.

Example 3.4.3: The Weight-One Structure of an Elliptic Curve

Fix $\tau=s+it$ with $t=\operatorname{Im}\tau>0$, and let $E_\tau=\mathbb{C}/\Lambda_\tau$ with $\Lambda_\tau=\mathbb{Z}\cdot 1\oplus\mathbb{Z}\tau$. This is the $n=1$ case of example 2.1.2, with $h^{1,0}=h^{0,1}=1$ and $b_1=2$. Both sides of theorem 3.4.1 can be written down.

The lattice side. Take $V_\mathbb{Z}=\Lambda_\tau=H_1(E_\tau,\mathbb{Z})$ with J multiplication by i. In the real basis $(1,\tau)$ of $W=\mathbb{C}$ one has $i=-\frac{s}{t}+\frac{1}{t}\tau$ and $i\tau=-\frac{s^2+t^2}{t}+\frac{s}{t}\tau$, so J has matrix

$$M=\begin{pmatrix}-s/t & -(s^2+t^2)/t\\ 1/t & s/t\end{pmatrix},\qquad \det M=\frac{-s^2+s^2+t^2}{t^2}=1$$

Let Q_0 be the alternating form with $Q_0(1,\tau)=1$; every alternating integral form on Λ_τ is an integer multiple of it. Writing $u=x+y\tau$ in the coordinates (x,y), so that $Q_0(u,v)$ is the determinant of the coordinate matrix of (u,v), a direct computation gives

$$Q_0(u,Ju)=\frac{1}{t}\bigl(x^2+2sxy+(s^2+t^2)y^2\bigr)=\frac{(x+sy)^2+(ty)^2}{t}=\frac{|u|^2}{\operatorname{Im}\tau}$$

so $Q_0(Ju,u)<0$, and it is $Q:=-Q_0$, normalized by $Q(1,\tau)=-1$, that satisfies (3.8). Relation (3.7) is automatic because $\det M=1$. The associated Hermitian form of theorem 3.4.1 is

$$H(u,v)=Q(Ju,v)+i\,Q(u,v)=\frac{u\bar v}{\operatorname{Im}\tau}$$

as one checks on the basis, and $\operatorname{Im}H=Q$ is integral on Λ_τ. So Λ_τ carries a polarization for every τ in the upper half-plane, and corollary 3.4.2 returns the classical fact that every one-dimensional complex torus is an algebraic curve.

The cohomological side. Take instead $V_\mathbb{Z}=H^1(E_\tau,\mathbb{Z})=\mathrm{Hom}(\Lambda_\tau,\mathbb{Z})$ with the weight-1 structure of example 3.1.2, and let ℓ_1,ℓ_2 be the basis dual to $1,\tau$. The space $H^{1,0}(E_\tau)=H^0(E_\tau,\Omega^1)$ is

spanned by dz, whose class in $H^1(E_\tau,\mathbb{C}) = \mathrm{Hom}(\Lambda_\tau,\mathbb{C})$ is recorded by its periods,

$$\int_1 dz = 1, \qquad \int_\tau dz = \tau, \qquad \text{so} \qquad dz = \ell_1 + \tau\,\ell_2$$

and $H^{0,1} = \overline{H^{1,0}}$ is spanned by $\ell_1 + \bar\tau\,\ell_2$. The two lines are transverse precisely because $\bar\tau - \tau = -2i\,t \neq 0$, which is (3.1) for this structure. With the cup product $Q'(\alpha,\beta) = \int_{E_\tau} \alpha\wedge\beta$, normalized by $Q'(\ell_1,\ell_2) = 1$ for the complex orientation,

$$Q'\big(dz,\overline{dz}\big) = Q'\big(\ell_1+\tau\ell_2,\ \ell_1+\bar\tau\ell_2\big) = \bar\tau - \tau = -2i\,\mathrm{Im}\,\tau$$

so $i\,Q'(dz,\overline{dz}\,) = 2\,\mathrm{Im}\,\tau > 0$. The second Hodge-Riemann relation for $H^1(E_\tau,\mathbb{Z})$ is therefore literally the condition $\mathrm{Im}\,\tau > 0$, and the first is automatic because F^1 is a line and Q' is alternating. The period line $\mathbb{C}\cdot(1,\tau) \subseteq \mathbb{C}^2$ is the point of the period domain that example 3.3.3 identifies with τ in the upper half-plane.

Moduli. An isomorphism of polarized weight-1 structures of rank 2 is a change of $\mathbb{Z}$-basis of the lattice preserving the alternating form, that is, an element of $\mathrm{SL}_2(\mathbb{Z})$, and replacing the basis $(1,\tau)$ of Λ_τ by $(c\tau+d,\ a\tau+b)$ rescales the lattice to $\Lambda_{\tau'}$ with $\tau' = (a\tau+b)/(c\tau+d)$. The isomorphism classes are therefore the orbits of $\mathrm{SL}_2(\mathbb{Z})$ on the upper half-plane, with the j-invariant as the coordinate on the quotient [6, The Modular Curve $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}^*$].

The example shows the mechanism in miniature. The Hodge structure is one line in a two-dimensional space; the first Hodge-Riemann relation is automatic because that line is isotropic for dimension reasons; the second is a single inequality; and the parameter recording the structure is the ratio of the two periods of dz. The higher-weight theory repeats this pattern with lines replaced by subspaces and the two relations no longer automatic.

Weight 2 is the first weight in which the polarization has both signs to distribute, and theorem 2.2.3 already computed the distribution on a surface: positive on $H^{2,0}\oplus H^{0,2}$ and along the Kähler class, negative on the primitive $(1,1)$-classes. A K3 surface is the case $h^{2,0} = 1$, so that the top step F^2 of the filtration is again a line and the whole structure is again recorded by a single point of a projective space.

Example 3.4.4: The Weight-Two Structure of a K3 Surface

Let X be a K3 surface, a compact Kähler surface with trivial canonical bundle and $h^{1,0} = 0$. Example 2.2.4 recorded its Hodge diamond: $h^{2,0} = h^{0,2} = 1$, $h^{1,1} = 20$, $b_2 = 22$, and $h^{1,0} = h^{0,1} = 0$. So $H^1(X,\mathbb{Z}) = 0$ and every Hodge-theoretic invariant of X lives in H^2.

Take $V_\mathbb{Z} = H^2(X,\mathbb{Z})$ with the weight-2 structure of example 3.1.2 and $Q(\alpha,\beta) = \int_X \alpha\wedge\beta$. By example 2.2.4 the pair $(V_\mathbb{Z},Q)$ is an even unimodular lattice of signature $(3,19)$, isometric to $U^{\oplus 3}\oplus E_8(-1)^{\oplus 2}$; the identification of the isometry class is cited to [4, K3 Surfaces: A Closer Look].

Since $h^{2,0} = 1$, the top step of the Hodge filtration is a line,

$$F^2V_\mathbb{C} = H^{2,0}(X) = \mathbb{C}\,\sigma$$

for a nonzero holomorphic 2-form σ (a trivialization of K_X), and the whole structure is determined

by that line. Indeed $H^{0,2} = \overline{\mathbb{C}\sigma}$ by definition 3.1.1, and

$$H^{1,1}(X) = \left\{ x \in H^2(X,\mathbb{C}) \;\middle|\; Q(x,\sigma) = Q(x,\bar\sigma) = 0 \right\}$$

For the last identity, note that $\alpha \wedge \beta$ has type $(p+p', q+q')$ and only a $(2,2)$-form has nonzero integral over a surface, so Q pairs $H^{2,0}$ with $H^{0,2}$, pairs $H^{1,1}$ with itself, and kills every other combination of summands (Step 1 in the proof of theorem 2.2.3). Writing $x = a\sigma + y + b\bar\sigma$ with $y \in H^{1,1}$, this gives $Q(x,\sigma) = b\,Q(\bar\sigma,\sigma)$ and $Q(x,\bar\sigma) = a\,Q(\sigma,\bar\sigma)$, and $Q(\sigma,\bar\sigma) \neq 0$ by the relation below, so both vanish exactly when $a = b = 0$.

The Hodge-Riemann relations, applied to the line F^2 and written out for the cup form Q rather than for the polarizing form $-Q$ of proposition 3.2.9, become two conditions on σ alone:

$$Q(\sigma,\sigma) = 0, \qquad Q(\sigma,\bar\sigma) > 0 \tag{3.10}$$

The first is the relation $Q(F^2,F^2) = 0$, and it is automatic here, since $\sigma \wedge \sigma$ is a form of type $(4,0)$ on a surface and hence zero. The second is part (1) of lemma 2.2.2, proved in chapter 2 by a computation in a unitary coframe. It is stated here for the cup form, whose negative is the polarizing form in weight 2 (example 3.2.10): a $(2,0)$-class on a surface is primitive, since $\omega \wedge \sigma$ has type $(3,1)$ and vanishes, and condition (2) of definition 3.2.1 for $-Q$ at σ reads $i^{2-0}\big(-Q(\sigma,\bar\sigma)\big) > 0$, which is the displayed inequality because $i^2 = -1$. Neither Q nor $-Q$ polarizes all of H^2 (exercise 3.2.1 (5)), so (3.10) is a statement about the line F^2 rather than an instance of theorem 3.2.4 applied to $(V_{\mathbb{Z}}, Q)$; the weight-2 relations on that line were already verified for surfaces in Part I.

Three consequences follow at once.

1. *The Néron-Severi group is a hyperplane section of the lattice.* An integral class γ is real, so $\gamma^{2,0} = \overline{\gamma^{0,2}}$ and γ is of type $(1,1)$ as soon as $Q(\gamma,\sigma) = 0$. Hence

   $$\mathrm{NS}(X) = \sigma^{\perp} \cap H^2(X,\mathbb{Z})$$

   which by theorem 2.3.2 is the group of first Chern classes of line bundles on X. Its rank satisfies $\rho(X) \leq h^{1,1}(X) = 20$ (proposition 2.3.4). The bound is attained, by the K3 surfaces classically called *singular* (a misnomer: they are smooth, and the name records only that ρ is as large as the Hodge numbers allow). That such surfaces exist is not proved here and is not needed below; where a $\rho = 20$ surface is used it is taken as a hypothesis.

2. *The structure is a point of a quadric.* By (3.10) the line $[\sigma]$ lies in the subset of $\mathbb{CP}\big(H^2(X,\mathbb{C})\big) \cong \mathbb{CP}^{21}$ cut out by the quadratic equation $Q(x,x) = 0$ together with the open condition $Q(x,\bar x) > 0$, an open subset of the 20-dimensional quadric $Q(x,x) = 0$; the point $[\sigma]$ is the period point of X. That subset is not itself a period domain in the sense of definition 3.3.2, which imposes positivity at $p = 1$ as well: there the condition asks $-Q$ to be positive definite on the real $(1,1)$-classes, where by theorem 2.2.3 it has signature $(19,1)$, so it holds at no point. Equivalently, the two positivity conditions force the polarizing form to have signature $(n,2)$ with $n = h^{1,1}$ (example 3.3.6), and $-Q$ on H^2 has signature $(19,3)$, not $(20,2)$. Cutting the lattice down repairs this.

3. *Polarizing drops the dimension by one.* If X is projective, fix an ample class h with $Q(h,h) = 2d > 0$. The sublattice $h^{\perp}$, the primitive cohomology of definition 3.2.6, then

has rank 21 and signature $(2,19)$, since removing one positive direction from $(3,19)$ leaves $(2,19)$; and σ lies in $h^{\perp}\otimes\mathbb{C}$ because h is of type $(1,1)$. So $-Q$ has signature $(19,2)$ on $h^{\perp}$, and the period point of a polarized K3 varies in the period domain of example 3.3.6 with $n=19$, of dimension 19.

The K3 case is the weight-2 analogue of the elliptic curve. In both, the Hodge structure is a single subspace of the complexified lattice: a g-plane in rank $2g$ for weight 1, a line in rank 22 for a K3. In both, the first Hodge-Riemann relation is an algebraic condition on that subspace and the second is an open positivity condition. And in both, the resulting parameter space is a concrete open subset of a projective variety, which is what makes the moduli theory of the two classes of surfaces tractable.

The difference is where the extra Hodge classes sit. On an elliptic curve every integral class in H^2 is of type $(1,1)$, so $\rho=1$ always. On a K3 surface, $\mathrm{NS}(X)=\sigma^{\perp}\cap H^2(X,\mathbb{Z})$ ranges from 0, on a non-projective K3, where theorem 2.3.2 produces no line bundle with nonzero Chern class, up to a lattice of rank 20; and the rank jumps exactly where the period point becomes orthogonal to extra integral vectors. Those loci are the Hodge loci taken up in chapter 6, where the recovery of a K3 surface from its period point, the global Torelli theorem surveyed in [4], is also treated.

Exercise 3.4.1

1. Let V be an effective weight-1 Hodge structure of rank 2 with complex structure J on $V_{\mathbb{R}}$, and let Q_0 generate the rank-one group of alternating integral forms on $V_{\mathbb{Z}}$. Show that (3.7) holds for every alternating form, that $Q_0(Ju,u)\neq 0$ for all $u\neq 0$, and that exactly one of $\pm Q_0$ satisfies (3.8). Then show that for $g\geq 2$ condition (3.7) is a nontrivial linear condition, by exhibiting a weight-1 structure of rank 4 with no polarization.

2. For the elliptic curve E_τ of example 3.4.3, show that the polarizations of the weight-1 structure on Λ_τ are exactly the forms nQ with n a positive integer and $Q(1,\tau)=-1$, and compute the Hermitian form attached to nQ by theorem 3.4.1. Deduce that every elliptic curve carries a polarization inducing a surjection $\bigwedge^2\Lambda_\tau\to\mathbb{Z}$, a *principal* polarization.

3. Let $X=W/\Gamma$ and $X'=W'/\Gamma'$ be complex tori. Use theorem 3.4.1 to show that $\mathrm{Hom}(X,X')$ is the group of $\mathbb{C}$-linear maps $F\colon W\to W'$ with $F(\Gamma)\subseteq\Gamma'$. Then compute $\mathrm{End}(E_\tau)$ for $E_\tau=\mathbb{C}/(\mathbb{Z}\oplus\mathbb{Z}\tau)$, showing that it is $\mathbb{Z}$ unless τ satisfies a quadratic equation over $\mathbb{Q}$, in which case it is a subring of $\mathbb{Q}(\tau)$ free of rank 2 over $\mathbb{Z}$.

4. Let X be a projective K3 surface with $\rho(X)=20$. Using example 3.4.4 and theorem 2.2.3, show that the intersection form on $\mathrm{NS}(X)$ has signature $(1,19)$, that the *transcendental lattice* $T(X)=\mathrm{NS}(X)^{\perp}\subseteq H^2(X,\mathbb{Z})$ has rank 2, and that Q is positive definite on $T(X)$. Where in $T(X)\otimes\mathbb{C}$ does σ sit?

5. Let $X=W/\Gamma$ be a complex torus of dimension g and let $V=H^1(X,\mathbb{Z})$ carry its weight-1 Hodge structure. Show that $H^k(X,\mathbb{Z})=\bigwedge^k V_{\mathbb{Z}}$ carries a pure Hodge structure of weight k with
$$\left(\textstyle\bigwedge^k V\right)^{p,q}=\bigwedge^p V^{1,0}\otimes\bigwedge^q V^{0,1},\qquad p+q=k$$
and check that the resulting Hodge numbers agree with those computed in example 2.1.2.

4

Hard Lefschetz and Lefschetz Pencils

The Kähler class has appeared so far as a source of hypotheses. It closed the gap between Dolbeault and de Rham cohomology, it gave the form that polarizes primitive cohomology, and in every case what was used was that ω is closed. This chapter uses something else: that wedging with ω is an operator on cohomology, and that this operator has a great deal of structure.

The structure is a representation of $\mathfrak{sl}_2$. The Lefschetz operator L, its adjoint Λ, and the operator that records degree satisfy the commutation relations of that Lie algebra, and the representation theory of $\mathfrak{sl}_2$ is rigid enough to force conclusions about $H^*(X, \mathbb{C})$ that no direct argument gives. Two follow immediately. The map $L^{n-k}\colon H^k(X, \mathbb{C}) \to H^{2n-k}(X, \mathbb{C})$ is an isomorphism, which is the Hard Lefschetz theorem; and cohomology decomposes into primitive pieces, on which the form of proposition 3.2.9 is definite with the alternating signs corollary 3.2.5 predicted. The positivity that chapter 3 deferred, and the identity of Weil (theorem 3.2.8) it rested on, are both proved here.

The second half of the chapter changes the question from what holds on one manifold to how one manifold can be seen as a family. A pencil of hyperplane sections fibres a variety, after a blow-up, over $\mathbb{C}\mathrm{P}^1$, with finitely many singular fibres. Following a loop around a singular value carries the cohomology of a nearby fibre back to itself by a nontrivial automorphism, and the Picard-Lefschetz formula computes it exactly. That automorphism is monodromy, and it is the mechanism by which a family of varieties acts on the Hodge structures of its members, which is what chapter 5 takes up.

4.1 The $\mathfrak{sl}_2$-Action on Cohomology

The operators L and Λ of definition 1.1.8 entered chapter 1 as ingredients of the Kähler identities and were then set aside. This section takes them as the objects of study. Adjoining a third operator,

the one that records the degree of a form, turns the pair into a Lie algebra acting on the exterior algebra at a single point.

Everything here is pointwise linear algebra, valid on any Hermitian manifold: the commutation relations, the decomposition of the exterior algebra into primitive pieces, and Weil's identity for the Hodge star are all statements about a single Hermitian vector space, and none of them uses $d\omega = 0$. The Kähler condition is spent in exactly one place, the transport of the action from forms to cohomology carried out in section 4.2, and it is spent through theorem 1.3.1.

Throughout, x denotes a point of a Hermitian manifold (X, ω) of complex dimension n, and $\bigwedge^k_x$ denotes the space of complex-valued k-covectors at x, that is, the degree-k part of $\bigwedge^\bullet T^*_x X \otimes \mathbb{C}$, with its bigrading $\bigwedge^{p,q}_x$ and the Hermitian inner product $\langle -, - \rangle$ induced by the metric. The operators L and Λ are algebraic, so each acts on $\bigwedge^\bullet_x$ one point at a time, and an identity between such operators holds for differential forms as soon as it holds at every point.

The third operator is forced by the shape of the first two. Both L and Λ change the degree, L by $+2$ and Λ by -2, so their commutator preserves degree and the only candidate for it, if the three are to close into a Lie algebra, is an operator that reads the degree off.

Definition 4.1.1: The Lefschetz Triple and Primitive Forms

Let (X, ω) be a Hermitian manifold of complex dimension n and let $x \in X$. The *counting operator* at x is the linear map

$$H \colon \bigwedge^\bullet_x \to \bigwedge^\bullet_x, \qquad H\alpha = (k - n)\,\alpha \quad \text{for } \alpha \in \bigwedge^k_x$$

The *Lefschetz triple* at x is the triple of operators (L, Λ, H) on $\bigwedge^\bullet_x$, where L and Λ are the Lefschetz operators of definition 1.1.8.
A covector $\alpha \in \bigwedge^k_x$ is *primitive* if $\Lambda\alpha = 0$. The space of primitive k-covectors is written P^k_x, and $P^{p,q}_x := P^{p+q}_x \cap \bigwedge^{p,q}_x$. A differential form is primitive if it is primitive at every point.

For $n = 1$ the triple can be written out: H is -1 on functions, 0 on 1-covectors and $+1$ on 2-covectors, and every 1-covector is primitive, since Λ would send it to degree -1. The full computation in dimensions one and two is example 4.1.3.

The counting operator carries no geometry: it is the grading of the exterior algebra, shifted so that the middle degree $k = n$ sits at 0. The shift is what makes the three relations below come out symmetric, and it is why n, the complex dimension, appears in a purely algebraic statement.

Primitivity is defined here by $\Lambda\alpha = 0$, while definition 3.2.6 defined primitive cohomology by $L^{n-k+1} = 0$. The two conditions agree on $\bigwedge^k_x$ for $k \leq n$, and that is part (1) of theorem 4.1.4 below; the present definition is the one that makes the relations usable, and the other is the one that makes sense on cohomology, where Λ is not visible until a metric is chosen.

The proof below computes in the flat model of *Complex Analytic Geometry* [5, The Operators on Flat $\mathbb{C}^n$]. Nothing beyond the Clifford relations of that model is needed, and no differentiation occurs anywhere, which is why the Kähler condition plays no part.

Proposition 4.1.2: The Commutation Relations of the Lefschetz Triple

Let (X, ω) be a Hermitian manifold of complex dimension n and $x \in X$. On $\bigwedge_x^\bullet$ the Lefschetz triple satisfies

$$[H, L] = 2L, \qquad [H, \Lambda] = -2\Lambda, \qquad [L, \Lambda] = H$$

Consequently $e \mapsto L$, $f \mapsto \Lambda$, $h \mapsto H$ defines a representation of $\mathfrak{sl}_2(\mathbb{C})$ on $\bigwedge_x^\bullet$, where e, f, h is the standard basis with $[h, e] = 2e$, $[h, f] = -2f$, $[e, f] = h$. The operators L and Λ shift the bidegree by $(1, 1)$ and $(-1, -1)$, and H preserves it.

***Proof*:**

The first two relations are degree bookkeeping. For $\alpha \in \bigwedge_x^k$ the covector $L\alpha$ lies in $\bigwedge_x^{k+2}$, so

$$[H, L]\alpha = HL\alpha - LH\alpha = (k + 2 - n)L\alpha - (k - n)L\alpha = 2L\alpha$$

and the same computation with Λ, which lowers the degree by two, gives $[H, \Lambda]\alpha = -2\Lambda\alpha$. Bidegrees behave as claimed because ω is of type $(1, 1)$ and Λ is its pointwise adjoint, so H, being a scalar on each $\bigwedge_x^k$, preserves $\bigwedge_x^{p,q}$.

The third relation is the content. Since ω is a positive Hermitian form at x, it can be diagonalized: there is a unitary coframe $\varphi^1, \ldots, \varphi^n$ of $(1, 0)$-covectors at x with

$$\omega = i \sum_{j=1}^{n} \varphi^j \wedge \bar{\varphi}^j$$

and with the monomials $\varphi^I \wedge \bar{\varphi}^J$ orthonormal, exactly the frame used in lemma 2.2.2. Write $e_j = \varphi^j \wedge (-)$ and $\bar{e}_j = \bar{\varphi}^j \wedge (-)$ for the wedge operators and $\iota_j = e_j^*$, $\bar{\iota}_j = \bar{e}_j^*$ for their pointwise adjoints, the contractions. These satisfy the Clifford relations [5, The Operators on Flat $\mathbb{C}^n$],

$$\{e_j, \iota_k\} = \delta_{jk}, \qquad \{\bar{e}_j, \bar{\iota}_k\} = \delta_{jk}$$

with every other anticommutator among $e_j, \bar{e}_j, \iota_k, \bar{\iota}_k$ equal to zero, and in these terms

$$L = i \sum_j e_j \bar{e}_j, \qquad \Lambda = -i \sum_k \bar{\iota}_k \iota_k$$

Write $P_j = e_j \bar{e}_j$ and $R_k = \bar{\iota}_k \iota_k$, so that $[L, \Lambda] = \sum_{j,k} [P_j, R_k]$, the scalars i and $-i$ cancelling.

Step 1: the off-diagonal terms vanish. Fix $j \neq k$. Each of e_j and $\bar{e}_j$ anticommutes with each of $\bar{\iota}_k$ and ι_k: the four anticommutators are $\{e_j, \iota_k\} = \delta_{jk} = 0$, $\{\bar{e}_j, \bar{\iota}_k\} = \delta_{jk} = 0$, and the two mixed-type ones, which vanish for all indices. Moving the pair R_k past the pair P_j therefore costs four sign changes, so $P_j R_k = R_k P_j$ and $[P_j, R_k] = 0$.

Step 2: the diagonal terms. Fix j and drop the index, writing $e, \bar{e}, \iota, \bar{\iota}$ and $P = e\bar{e}$, $R = \bar{\iota}\iota$. Using $\iota e = 1 - e\iota$,

$$RP = \bar{\iota}\, \iota\, e\, \bar{e} = \bar{\iota}\, (1 - e\iota)\, \bar{e} = \bar{\iota}\bar{e} - \bar{\iota}\, e\, \iota\, \bar{e}$$

The first term is $\bar{\iota}\bar{e} = 1 - \bar{e}\bar{\iota}$. For the second, move e out to the left past $\bar{\iota}$ and then $\bar{e}$ out to the left past ι, each move costing a sign, and then use $\bar{\iota}\bar{e} = 1 - \bar{e}\bar{\iota}$ again:

$$\bar{\iota}\, e\, \iota\, \bar{e} = -e\, \bar{\iota}\, \iota\, \bar{e} = e\, \bar{\iota}\, \bar{e}\, \iota = e(1 - \bar{e}\bar{\iota})\iota = e\iota - e\bar{e}\bar{\iota}\iota$$

Combining, and writing $N = e\iota$ and $\bar{N} = \bar{e}\bar{\iota}$,

$$RP = (1 - \bar{N}) - (N - PR) = 1 - N - \bar{N} + PR$$

so that $[P, R] = PR - RP = N + \bar{N} - 1$.

Step 3: assembling. By Steps 1 and 2,

$$[L, \Lambda] = \sum_j \big(N_j + \bar{N}_j - 1\big) = \sum_j (N_j + \bar{N}_j) \;-\; n$$

The operator $N_j = e_j \iota_j$ acts on a monomial $\varphi^I \wedge \bar{\varphi}^J$ by removing the factor φ^j and putting it back, which is the identity when $j \in I$ and zero otherwise; likewise $\bar{N}_j$ is the identity when $j \in J$ and zero otherwise. On $\varphi^I \wedge \bar{\varphi}^J$ the sum $\sum_j (N_j + \bar{N}_j)$ is therefore multiplication by $|I| + |J|$, the total degree. Hence $[L, \Lambda]$ acts on $\bigwedge_x^k$ by $k - n$, which is H.

The three relations are those of the standard basis of $\mathfrak{sl}_2(\mathbb{C})$, so the assignment $e \mapsto L$, $f \mapsto \Lambda$, $h \mapsto H$ extends to a Lie algebra homomorphism, completing the proof.

The representation is compatible with the metric, since $\Lambda = L^*$ and $H^* = H$; and it is compatible with the bigrading, since L and Λ move p and q in lockstep. The first will convert representation-theoretic statements into orthogonality statements, and the second will convert them into statements about Hodge structures. Both are used below.

Example 4.1.3: The Lefschetz Triple in Dimensions One and Two

Take a unitary coframe as in the proof above and write $\theta_j = i\,\varphi^j \wedge \bar{\varphi}^j$, so that $\omega = \theta_1 + \cdots + \theta_n$, each θ_j is real of degree 2, distinct θ_j commute, and $\theta_j \wedge \theta_j = 0$. In this notation L is $\sum_j \theta_j \wedge (-)$ and $\Lambda\theta_j = 1$, $\Lambda\varphi^j = \Lambda\bar{\varphi}^j = 0$.

The case $n = 1$. Here $\bigwedge_x^\bullet$ has dimensions $1, 2, 1$ in degrees $0, 1, 2$, and the H-eigenvalues are -1 on $\mathbb{C}$, 0 on $\bigwedge_x^1 = \mathbb{C}\varphi^1 \oplus \mathbb{C}\bar{\varphi}^1$, and $+1$ on $\bigwedge_x^2 = \mathbb{C}\theta_1$. The relation $[L, \Lambda] = H$ is visible on the constant 1: $\Lambda 1 = 0$ and $L1 = \theta_1$, so $[L, \Lambda]1 = -\Lambda\theta_1 = -1 = H1$. The primitive spaces are $P_x^0 = \mathbb{C}$ and $P_x^1 = \bigwedge_x^1$, the latter because Λ lands in degree -1; and $\bigwedge_x^2 = L\,P_x^0$. As an $\mathfrak{sl}_2(\mathbb{C})$-module, $\bigwedge_x^\bullet$ is the sum of the two-dimensional irreducible spanned by 1 and θ_1 and two copies of the trivial module, spanned by φ^1 and by $\bar{\varphi}^1$.

The case $n = 2$. The dimensions are $1, 4, 6, 4, 1$, totalling 16, and the H-eigenvalues on $\bigwedge_x^k$ are $k - 2$, running from -2 to 2. The primitive spaces are $P_x^0 = \mathbb{C}$, $P_x^1 = \bigwedge_x^1$ of dimension 4, and

$$P_x^2 = \ker\big(\Lambda\colon\ \textstyle\bigwedge_x^2 \to \bigwedge_x^0\big)$$

of dimension 5, since Λ is onto in this degree ($\Lambda\omega = 2$). A basis of P_x^2 is

$$\varphi^1 \wedge \varphi^2, \qquad \bar{\varphi}^1 \wedge \bar{\varphi}^2, \qquad \varphi^1 \wedge \bar{\varphi}^2, \qquad \varphi^2 \wedge \bar{\varphi}^1, \qquad \theta_1 - \theta_2$$

The first two are of type $(2, 0)$ and $(0, 2)$ and are primitive because Λ needs one factor of each of φ^j and $\bar{\varphi}^j$ to act; the next two are of type $(1, 1)$ and are primitive for the same reason; the last is of type $(1, 1)$ and is primitive because $\Lambda\theta_1 = \Lambda\theta_2 = 1$. So $P_x^{2,0}$ and $P_x^{0,2}$ are lines and $P_x^{1,1}$ has dimension 3, one less than $\bigwedge_x^{1,1}$, the missing direction being $\mathbb{C}\omega$. Since $\omega \wedge \omega = 2\theta_1\theta_2 \neq 0$, the line $\mathbb{C}\omega = L\,P_x^0$ meets P_x^2 only in 0, so

$$\textstyle\bigwedge_x^2 = L\,P_x^0 \oplus P_x^2, \qquad \bigwedge_x^3 = L\,P_x^1, \qquad \bigwedge_x^4 = L^2\,P_x^0$$

by comparison of dimensions ($6 = 1 + 5$, $4 = 4$, $1 = 1$). In the top degree there is no term $L\,P_x^2$, since L annihilates P_x^2: each of the first four basis covectors repeats an index against θ_1 and against θ_2, and $(\theta_1 + \theta_2) \wedge (\theta_1 - \theta_2) = 0$ because $\theta_j \wedge \theta_j = 0$ and the θ_j commute. This is the vanishing recorded in part (3) of theorem 4.1.4. As a module, $\bigwedge_x^\bullet$ is one copy of the three-dimensional irreducible (generated by 1), four copies of the two-dimensional one (generated by $\bigwedge_x^1$) and five trivial ones (generated by P_x^2), and $3 + 8 + 5 = 16$ as it must.

The pattern in both cases is the same: each primitive covector generates an irreducible module by repeated application of L, and the whole exterior algebra is the sum of those modules. That is a general fact, and it is a consequence of the classification of finite-dimensional $\mathfrak{sl}_2(\mathbb{C})$-modules rather than of anything about forms. The classification is not redeveloped here. Its statement, in the form used below, is that a finite-dimensional complex representation of $\mathfrak{sl}_2(\mathbb{C})$ is a direct sum of irreducibles; that the irreducibles are classified by their highest weight $m \geq 0$, with V_m of dimension $m+1$ and h-eigenvalues $m, m-2, \ldots, -m$ each occurring once; and that in V_m a vector v_0 of weight $-m$ is annihilated by f, the vectors $v_j = e^j v_0$ for $0 \leq j \leq m$ form a basis, $e^{m+1} v_0 = 0$, and a weight vector v of weight μ satisfies $e^j v \neq 0$ exactly when $\mu + 2j \leq m$. Complete reducibility, the classification of the irreducibles of SU(2) and hence, after complexification, of $\mathfrak{sl}_2(\mathbb{C})$, the highest weight theorem and the decomposition into irreducibles are all proved in *Lie Groups* [37].

Feeding the relations of proposition 4.1.2 into that classification and reading off what it says about $\bigwedge_x^\bullet$ gives the decomposition the rest of the chapter runs on.

Theorem 4.1.4: The Lefschetz Decomposition of the Exterior Algebra

Let (X, ω) be a Hermitian manifold of complex dimension n and let $x \in X$. Set $P_x^j = 0$ for $j < 0$. Then

1. $P_x^j = 0$ for $j > n$, and for $0 \leq k \leq n$,
$$P_x^k = \ker\Big(L^{n-k+1} \colon \textstyle\bigwedge_x^k \to \bigwedge_x^{2n-k+2}\Big)$$
2. for $0 \leq k \leq n$ the map $L^{n-k} \colon \bigwedge_x^k \to \bigwedge_x^{2n-k}$ is an isomorphism;
3. for every k,
$$\textstyle\bigwedge_x^k = \bigoplus_{r \geq 0} L^r\, P_x^{k-2r}$$
the summands are mutually orthogonal, L^r is injective on P_x^{k-2r} for each $r \geq \max(0,\, k-n)$, and $L^r\, P_x^{k-2r} = 0$ for $0 \leq r < k - n$;
4. the decomposition respects the bigrading: $P_x^k = \bigoplus_{p+q=k} P_x^{p,q}$ and $\bigwedge_x^{p,q} = \bigoplus_{r \geq 0} L^r\, P_x^{p-r,q-r}$.

***Proof*:**

By proposition 4.1.2 the space $\bigwedge_x^\bullet$ is a finite-dimensional representation of $\mathfrak{sl}_2(\mathbb{C})$ in which $\bigwedge_x^k$ is exactly the weight space of weight $k - n$, since H acts there by $k - n$. Decompose it into irreducibles, $\bigwedge_x^\bullet = \bigoplus_i W_i$ with $W_i \cong V_{m_i}$, using complete reducibility [37]. In W_i the weight μ occurs with multiplicity one when $|\mu| \leq m_i$ and $\mu \equiv m_i$ modulo 2, and not at all otherwise; the vectors killed by Λ are exactly the multiples of the weight-$(-m_i)$ vector; and L^j is injective on

the weight-μ line precisely when $\mu + 2j \leq m_i$.

Step 1: primitive covectors and the two descriptions. Let $\alpha \in \bigwedge_x^k$ and write $\alpha = \sum_i \alpha_i$ with $\alpha_i \in W_i$. Each α_i has weight $k-n$, so $\alpha_i \neq 0$ forces $m_i \geq |k-n|$ and $m_i \equiv k-n$ modulo 2.

If α is primitive then each α_i is, since Λ preserves each W_i; so each nonzero α_i is a lowest weight vector, whence $k-n = -m_i \leq 0$. This gives $P_x^j = 0$ for $j > n$ and, for $k \leq n$, $m_i = n-k$ for every nonzero component. Then $L^{n-k+1}\alpha_i = 0$, because $(k-n)+2(n-k+1) = n-k+2 > n-k = m_i$; so $\alpha \in \ker L^{n-k+1}$.

Conversely let $k \leq n$ and $L^{n-k+1}\alpha = 0$. Each component satisfies $L^{n-k+1}\alpha_i = 0$, which for $\alpha_i \neq 0$ means $n-k+2 > m_i$, that is $m_i \leq n-k+1$. Combined with $m_i \geq n-k$ and $m_i \equiv n-k$ modulo 2 this forces $m_i = n-k$, so α_i has weight $-m_i$ and is annihilated by Λ. Hence α is primitive, and part (1) is proved.

Step 2: the isomorphism in complementary degrees. Let $k \leq n$. The operator L^{n-k} carries the weight-$(k-n)$ space of W_i to its weight-$(n-k)$ space. If $m_i < n-k$ then W_i has neither of these weight spaces and there is nothing to check. If $m_i \geq n-k$, both are lines, and L^{n-k} is injective on the first because $(k-n)+2(n-k) = n-k \leq m_i$; an injective map between lines is an isomorphism. Summing over i gives part (2).

Step 3: the decomposition. Let $W_i \cong V_{m_i}$ with lowest weight vector v_i, which has weight $-m_i$ and so lies in $\bigwedge_x^{n-m_i}$ and is primitive. The basis $v_i, Lv_i, \ldots, L^{m_i}v_i$ of W_i consists of covectors of degrees $n-m_i, n-m_i+2, \ldots, n+m_i$, one in each. So $W_i \cap \bigwedge_x^k$ is $\mathbb{C}\, L^r v_i$ with $k = n-m_i+2r$ when $0 \leq r \leq m_i$, and 0 otherwise. Since the v_i span $P_x^\bullet$ by Step 1 (a primitive covector has all its components primitive, hence a multiple of the relevant v_i), summing over i gives $\bigwedge_x^k = \sum_{r\geq 0} L^r P_x^{k-2r}$, and the sum is direct because the summand $L^r P_x^{k-2r}$ collects exactly the W_i with $m_i = n-k+2r$. For the injectivity and the vanishing, note that P_x^{k-2r} is spanned by those v_i and that $L^r v_i \neq 0$ exactly when $r \leq m_i$, which for $m_i = n-k+2r$ says precisely $r \geq k-n$. So for $r \geq \max(0,\, k-n)$ the operator L^r is injective on each line $\mathbb{C}v_i$ making up P_x^{k-2r}, hence on P_x^{k-2r} itself, the W_i being independent and stable under L; while for $0 \leq r < k-n$ the same criterion gives $L^r v_i = 0$ for every such v_i, that is $L^r\, P_x^{k-2r} = 0$.

For orthogonality, take $r > s$, $\pi \in P_x^{k-2r}$ and $\pi' \in P_x^{k-2s}$. The submodule generated by π is spanned by the $L^j\pi$, so $\Lambda^s L^r \pi$, having degree $k-2s$, is a multiple of $L^{r-s}\pi$, say $\Lambda^s L^r \pi = c\, L^{r-s}\pi$. Using $\Lambda = L^*$ twice,

$$\langle L^r\pi, L^s\pi'\rangle = \langle \Lambda^s L^r \pi, \pi'\rangle = c\,\langle L^{r-s}\pi, \pi'\rangle = c\,\langle \pi, \Lambda^{r-s}\pi'\rangle = 0$$

since $r-s \geq 1$ and π' is primitive. This is part (3).

Step 4: bidegrees. The operator Λ shifts the bidegree by $(-1,-1)$, so it maps $\bigwedge_x^{p,q}$ into $\bigwedge_x^{p-1,q-1}$ and its kernel in $\bigwedge_x^k$ is the direct sum of its kernels in the summands $\bigwedge_x^{p,q}$ with $p+q=k$; this is $P_x^k = \bigoplus_{p+q=k} P_x^{p,q}$. Applying L^r, which shifts the bidegree by (r,r), to the decomposition of part (3) and separating types gives $\bigwedge_x^{p,q} = \bigoplus_{r\geq 0} L^r P_x^{p-r,q-r}$, completing the proof.

Part (2) is the Hard Lefschetz theorem at the level of the exterior algebra at a point, and it is proved entirely with representation theory: in an $\mathfrak{sl}_2(\mathbb{C})$-module the weight spaces of μ and $-\mu$ have the same dimension, and the power of L carrying one to the other realizes the isomorphism. What section 4.2 has to give is not a new argument of this kind but a way of transporting this one to cohomology.

Part (3) also settles the bookkeeping, since $\bigoplus_{r\geq 1} L^r P_x^{k-2r} = L \bigwedge_x^{k-2}$ turns the decomposition into

$\bigwedge_x^k = L\bigwedge_x^{k-2} \oplus P_x^k$, from which the primitive dimensions follow by subtraction (exercise 4.1.1 (5)). For $n = 2$ this returns the numbers found by hand in example 4.1.3: $\dim P_x^2 = \binom{4}{2} - \binom{4}{0} = 5$ and $\dim P_x^{1,1} = \binom{2}{1}\binom{2}{1} - \binom{2}{0}\binom{2}{0} = 3$.

Everything so far is linear algebra at a point, hence available on any Hermitian manifold and on any of its differential forms. Cohomology is another matter: L descends to cohomology on any complex manifold with a closed ω, since wedging with a closed form preserves closedness and exactness, but Λ has no reason to. The device that carries the whole triple across is the harmonic representative, and it works precisely because theorem 1.3.1 makes the Laplacian commute with L and Λ; section 4.2 carries out that transport.

The transport is also what permits the use of harmonic representatives already made in section 3.2. The geometric polarization proposition 3.2.9 was evaluated on a harmonic representative of a primitive class, and the evaluation used Weil's identity, which is a statement about forms that are primitive at each point. That the two notions of primitivity agree on cohomology is part (1) of corollary 4.2.4, and they agree because L commutes with the Laplacian, which is a Kähler statement and fails without that hypothesis.

What remains is that identity. Theorem 3.2.8 computes the Hodge star of a primitive form of type (p, q) as an explicit multiple of L^{n-k} applied to it, and section 3.2 deferred its proof here on the grounds that the proof is a statement about the exterior algebra at a point. The statement proved below is exactly that pointwise assertion; applied at every point of a compact Kähler manifold it is theorem 3.2.8.

Theorem 4.1.5: Weil's Identity on the Exterior Algebra

Let (X, ω) be a Hermitian manifold of complex dimension n, let $x \in X$, and let $\alpha \in P_x^{p,q}$ be a primitive covector of type (p, q) with $k = p + q \leq n$. Then

$$\star\alpha = (-1)^{k(k+1)/2}\, i^{p-q}\, \frac{1}{(n-k)!}\, L^{n-k}\alpha$$

where $\star$ is the Hodge star of the metric.

Proof of theorem 3.2.8, deferred from section 3.2:
Both sides are algebraic in α, so it is enough to argue at the fixed point x, and theorem 3.2.8 follows by applying the result at every point. Choose a unitary coframe $\varphi^1, \dots, \varphi^n$ at x as in the proof of proposition 4.1.2, so that the monomials $\varphi^I \wedge \bar{\varphi}^J$ are orthonormal, and set

$$\theta_j = i\,\varphi^j \wedge \bar{\varphi}^j, \qquad \theta_T = \bigwedge_{j\in T} \theta_j \quad (T \subseteq \{1, \dots, n\})$$

The θ_j are real of degree 2, so they commute; $\theta_j \wedge \theta_j = 0$; the θ_T are orthonormal; $\theta_T \wedge \theta_{T'} = \theta_{T\cup T'}$ when $T \cap T' = \emptyset$ and 0 otherwise; and

$$\omega = \sum_j \theta_j, \qquad \omega^m = m! \sum_{|T|=m} \theta_T, \qquad \mathrm{vol} = \frac{\omega^n}{n!} = \theta_{\{1,\dots,n\}}$$

Write $m_0 = n - k$.

Step 1: reduction to a pairing identity. The Hodge star is the $\mathbb{C}$-linear operator determined by

$$\gamma \wedge \star\bar{\delta} = \langle \gamma, \delta \rangle\, \mathrm{vol} \qquad \text{for all } \gamma, \delta \in \textstyle\bigwedge_x^k$$

the polarized form of the identity $\alpha \wedge \star\bar{\alpha} = |\alpha|^2 \,\mathrm{vol}$ used in section 3.2; it determines $\star\bar{\delta}$ uniquely because the wedge pairing $\bigwedge_x^k \times \bigwedge_x^{2n-k} \to \bigwedge_x^{2n}$ is nondegenerate. Since $\star$ is the $\mathbb{C}$-linear extension of a real operator, $\star\bar{\alpha} = \overline{\star\alpha}$, so the asserted formula is equivalent to

$$\star\bar{\alpha} = (-1)^{k(k+1)/2}\, i^{q-p}\, \frac{1}{m_0!}\, L^{m_0}\bar{\alpha}$$

and hence, by the characterization above and by multiplying through by the inverse of the scalar, to the assertion

$$\gamma \wedge \bar{\alpha} \wedge \frac{\omega^{m_0}}{m_0!} = (-1)^{k(k+1)/2}\, i^{p-q}\, \langle \gamma, \alpha \rangle \,\mathrm{vol} \qquad \text{for all } \gamma \in \textstyle\bigwedge_x^k \tag{4.1}$$

Both sides of (4.1) are $\mathbb{C}$-linear in γ and conjugate-linear in α.

Step 2: a decomposition adapted to Λ. For disjoint subsets $I, J \subseteq \{1, \dots, n\}$ let $R = R(I,J)$ be the complement of $I \cup J$ and let Θ_R be the span of the θ_T with $T \subseteq R$. Every monomial $\varphi^{I'} \wedge \bar{\varphi}^{J'}$ equals a unit scalar times $\varphi^I \wedge \bar{\varphi}^J \wedge \theta_T$ with $T = I' \cap J'$, $I = I' \setminus J'$ and $J = J' \setminus I'$, and this correspondence between triples (I, J, T) and pairs (I', J') is a bijection. So there is an orthogonal decomposition

$$\textstyle\bigwedge_x^\bullet = \bigoplus_{I \cap J = \emptyset} \varphi^I \wedge \bar{\varphi}^J \wedge \Theta_{R(I,J)}$$

Each summand is stable under L and Λ, which act on it through the second factor. Indeed $\theta_j \wedge (-)$ kills the summand when $j \in I \cup J$, because the index j then repeats, and $\Lambda_j := -i\,\bar{\iota}_j \iota_j$ kills it as well, because for $j \in I$ there is no factor $\bar{\varphi}^j$ for $\bar{\iota}_j$ to remove and for $j \in J$ there is no factor φ^j for ι_j to remove; while for $j \in R$ the two contractions each pass the prefix $\varphi^I \wedge \bar{\varphi}^J$, of degree $|I| + |J|$, so the two signs cancel. Writing $L_R = \sum_{j \in R} \theta_j \wedge (-)$ and Λ_R for its adjoint on Θ_R, one therefore has, for $\eta \in \Theta_R$,

$$L(\varphi^I \wedge \bar{\varphi}^J \wedge \eta) = \varphi^I \wedge \bar{\varphi}^J \wedge L_R\eta, \qquad \Lambda(\varphi^I \wedge \bar{\varphi}^J \wedge \eta) = \varphi^I \wedge \bar{\varphi}^J \wedge \Lambda_R\eta$$

and $\Lambda_R\theta_T = \sum_{j \in T} \theta_{T \setminus j}$. In particular a covector is primitive if and only if each of its components in this decomposition is annihilated by the corresponding Λ_R.

Step 3: distinct summands contribute nothing. Suppose γ lies in the summand indexed by (I, J) and α in the summand indexed by (I_1, J_1), with $(I, J) \neq (I_1, J_1)$. The right-hand side of (4.1) vanishes because the decomposition is orthogonal. For the left-hand side, expand $\gamma \wedge \bar{\alpha} \wedge \omega^{m_0}$ into monomials; each term is a scalar multiple of

$$\varphi^I \wedge \bar{\varphi}^J \wedge \theta_T \wedge \bar{\varphi}^{I_1} \wedge \varphi^{J_1} \wedge \theta_{T_1} \wedge \theta_S$$

for some $T \subseteq R(I,J)$, $T_1 \subseteq R(I_1, J_1)$ and S with $|S| = m_0$. This product has degree $2n$, and a nonzero $2n$-covector must use each index exactly once as a φ and once as a $\bar{\varphi}$. The φ-indices are $I \sqcup J_1 \sqcup T \sqcup T_1 \sqcup S$ and the $\bar{\varphi}$-indices are $J \sqcup I_1 \sqcup T \sqcup T_1 \sqcup S$, so both must exhaust $\{1, \dots, n\}$ and hence $I \sqcup J_1 = J \sqcup I_1$. Now any $j \in I$ lies in $J \sqcup I_1$ and not in J, so $j \in I_1$; and any $j \in I_1$ lies in $I \sqcup J_1$ and not in J_1, so $j \in I$. Thus $I = I_1$ and then $J = J_1$, contrary to assumption. Every term vanishes, so the left-hand side does too.

By Step 3 and the bilinearity noted in Step 1 it suffices to prove (4.1) when γ and α lie in the same summand. Fix I, J disjoint, put $a = |I|$, $b = |J|$, $R = R(I,J)$ and $r = |R| = n - a - b$, and take

$$\gamma = \varphi^I \wedge \bar{\varphi}^J \wedge \rho, \qquad \alpha = \varphi^I \wedge \bar{\varphi}^J \wedge \pi, \qquad \rho, \pi \in \Theta_R$$

Both γ and α have total degree k, and inside this summand the degree-k part is $\varphi^I \wedge \bar{\varphi}^J$ wedged with the span of the θ_T with $|T| = s$ for the single value $s = (k-a-b)/2$; so ρ and π are both combinations of such θ_T, and $p = a+s$, $q = b+s$, $k = a+b+2s$, $m_0 = r-2s$, with $\Lambda_R \pi = 0$ by Step 2.

Step 4: the computation inside one summand. Write $\pi = \sum_{|T|=s} c_T \theta_T$ and $\rho = \sum_{|U|=s} d_U \theta_U$, the sums over subsets of R. Since $\theta_U \wedge \theta_T$ vanishes unless $U \cap T = \emptyset$, and since a term θ_S of ω^{m_0} survives only for $S = R \setminus (U \sqcup T)$,

$$\rho \wedge \bar{\pi} \wedge \frac{\omega^{m_0}}{m_0!} = \Big(\sum_{U \cap T = \emptyset} d_U \, \overline{c_T} \Big) \, \theta_R$$

using that $\bar{\theta}_T = \theta_T$. The claim is that primitivity of π turns the constraint $U \cap T = \emptyset$ into the constraint $U = T$ up to a sign:

$$\sum_{U \cap T = \emptyset} d_U \, \overline{c_T} = (-1)^s \sum_{U} d_U \, \overline{c_U} \tag{4.2}$$

To see this, note first what $\Lambda_R \pi = 0$ says. From $\Lambda_R \theta_T = \sum_{j \in T} \theta_{T \setminus j}$,

$$\Lambda_R \pi = \sum_{|T'| = s-1} \Big(\sum_{j \in R \setminus T'} c_{T' \cup j} \Big) \theta_{T'}$$

so primitivity says that $\sum_{j \in R \setminus T'} c_{T' \cup j} = 0$ for every $T' \subseteq R$ with $|T'| = s-1$. For $V \subseteq R$ set $\sigma_V = \sum_{T \supseteq V, \, |T| = s} c_T$. The relation just displayed is $\sigma_{T'} = 0$ for $|T'| = s-1$. Counting each $T \supseteq V$ once for every element of $T \setminus V$ gives

$$\sum_{j \in R \setminus V} \sigma_{V \cup j} = (s - |V|) \, \sigma_V$$

so a downward induction starting at $|V| = s-1$ shows $\sigma_V = 0$ whenever $|V| < s$; and $\sigma_V = c_V$ when $|V| = s$. Now fix U with $|U| = s$. Since $\sum_{V \subseteq W} (-1)^{|V|}$ is 1 for $W = \emptyset$ and 0 otherwise, inclusion and exclusion gives

$$\sum_{T \cap U = \emptyset} c_T = \sum_{V \subseteq U} (-1)^{|V|} \sigma_V = (-1)^s \sigma_U = (-1)^s c_U$$

only the term $V = U$ surviving. Conjugating and summing against d_U yields (4.2), and the right-hand side of (4.2) is $(-1)^s \langle \rho, \pi \rangle$.

It remains to restore the prefix. Since ρ has even degree it commutes past $\bar{\varphi}^I \wedge \varphi^J$, so

$$\gamma \wedge \bar{\alpha} = \varphi^I \wedge \bar{\varphi}^J \wedge \bar{\varphi}^I \wedge \varphi^J \wedge \rho \wedge \bar{\pi}$$

Moving $\bar{\varphi}^I$ left past $\bar{\varphi}^J$ costs $(-1)^{ab}$ and then $\bar{\varphi}^J \wedge \varphi^J = (-1)^{b^2} \varphi^J \wedge \bar{\varphi}^J = (-1)^b \varphi^J \wedge \bar{\varphi}^J$, so the prefix is $(-1)^{ab+b} \, \varphi^I \wedge \bar{\varphi}^I \wedge \varphi^J \wedge \bar{\varphi}^J$. For a set $I = \{i_1 < \cdots < i_a\}$, bringing each $\bar{\varphi}^{i_t}$ next to its φ^{i_t} costs $a-1, a-2, \ldots, 0$ transpositions of odd covectors, and $\varphi^j \wedge \bar{\varphi}^j = -i \, \theta_j$, so

$$\varphi^I \wedge \bar{\varphi}^I = (-1)^{a(a-1)/2} (-i)^a \, \theta_I$$

and likewise for J. Now wedge with $\omega^{m_0}/m_0!$. The prefix $\theta_I \wedge \theta_J$ kills every term θ_S of ω^{m_0} with S meeting $I \cup J$, so only the terms already accounted for in the Θ_R computation survive,

and (4.2) applies verbatim. Using $\theta_I \wedge \theta_J \wedge \theta_R = \mathrm{vol}$ together with $\langle \gamma, \alpha \rangle = \langle \rho, \pi \rangle$, which holds because the $\varphi^I \wedge \bar{\varphi}^J \wedge \theta_U$ are orthonormal,

$$\gamma \wedge \bar{\alpha} \wedge \frac{\omega^{m_0}}{m_0!} = \varepsilon\,(-1)^s\,\langle \gamma, \alpha \rangle\,\mathrm{vol}, \qquad \varepsilon = (-1)^{ab+b+a(a-1)/2+b(b-1)/2}\,(-i)^{a+b}$$

Step 5: the sign. It remains to check that $\varepsilon\,(-1)^s = (-1)^{k(k+1)/2} i^{p-q}$. Since $i^{a+b} = i^{a-b}\,i^{2b} = (-1)^b i^{a-b}$,

$$(-i)^{a+b} = (-1)^{a+b} i^{a+b} = (-1)^a\,i^{a-b} = (-1)^a\,i^{p-q}$$

so $\varepsilon\,(-1)^s = (-1)^N i^{p-q}$ with $N = ab + b + \frac{a(a-1)}{2} + \frac{b(b-1)}{2} + a + s$. On the other side, with $k = a + b + 2s$,

$$\frac{k(k+1)}{2} = \frac{(a+b)(a+b+1)}{2} + 2s(a+b) + 2s^2 + s \equiv \frac{(a+b)(a+b+1)}{2} + s \pmod 2$$

and expanding the first term,

$$\frac{(a+b)(a+b+1)}{2} = \frac{a(a+1)}{2} + \frac{b(b+1)}{2} + ab = \frac{a(a-1)}{2} + a + \frac{b(b-1)}{2} + b + ab$$

so $k(k+1)/2 \equiv N$ modulo 2. Hence $\varepsilon(-1)^s = (-1)^{k(k+1)/2} i^{p-q}$, which is (4.1) inside a single summand. With Step 3 this establishes (4.1) for every $\gamma \in \bigwedge_x^k$, and by Step 1 that is the asserted formula for $\star\alpha$, completing the proof.

The identity says that on primitive covectors the Hodge star, which is defined by the metric and the orientation, is computed by the Lefschetz operator, which is defined by ω alone, with a correction that depends only on the type. That is why it converts the cup product into an inner product: inserting it into $\alpha \wedge \star\bar{\alpha} = |\alpha|^2\,\mathrm{vol}$ turns $\int_X \alpha \wedge \bar{\alpha} \wedge \omega^{n-k}$, a topological quantity when the classes are fixed, into a positive multiple of the L^2-norm of the harmonic representative. Section 3.2 used exactly that step, and section 4.2 uses it again to prove the positivity in general.

One of the two debts left by chapter 3 is thereby settled. Proposition 3.2.9 carried a proof labelled as key ideas for the single reason that its second Hodge-Riemann relation rested on theorem 3.2.8, and that identity now has a proof.

Exercise 4.1.1

1. Verify $[H, L] = 2L$ and $[H, \Lambda] = -2\Lambda$ directly from the fact that L raises the total degree by 2 and Λ lowers it by 2. Then show that in the decomposition of $\bigwedge_x^\bullet$ into irreducible $\mathfrak{sl}_2(\mathbb{C})$-modules no summand V_m with $m > n$ occurs, and that the multiplicity of V_m is $\dim \bigwedge_x^{n-m} - \dim \bigwedge_x^{n-m-2}$.

2. Let $\alpha \in P_x^k$ be primitive with $k \le n$, and set $m = n - k$. Show first that for a covector v of H-weight μ and every $j \ge 1$,

$$[\Lambda, L^j]\,v = -\big(j\mu + j(j-1)\big)L^{j-1}v$$

(expand $[\Lambda, L^j] = \sum_{t=0}^{j-1} L^t[\Lambda, L]L^{j-1-t}$ and use $[\Lambda, L] = -H$). Deduce that $\Lambda L^j \alpha = j(m - j + 1)L^{j-1}\alpha$, that

$$\|L^r\alpha\|^2 = r!\,\frac{m!}{(m-r)!}\,\|\alpha\|^2 \qquad (0 \le r \le m)$$

and that $L^{m+1}\alpha = 0$. Which part of theorem 4.1.4 does the last statement recover?

3. Show that every covector of type $(p, 0)$ is primitive. Using theorem 4.1.5, deduce that $\star\alpha = (-1)^{n(n+1)/2} i^n \alpha$ for every $\alpha \in \bigwedge_x^{n,0}$, and verify this directly for $n = 1$ by computing $\star\varphi^1$ from the defining property of the Hodge star in a unitary coframe.

4. Let X be a compact Kähler surface, so $n = 2$, and let β be a real primitive form of type $(1, 1)$. Use theorem 4.1.5 to compute $\star\beta$, and deduce that

$$\beta \wedge \beta = -|\beta|^2 \operatorname{vol}$$

pointwise. Compare with part (3) of lemma 2.2.2 and say which of the two arguments uses the dimension of X.

5. Prove that $\bigwedge_x^k = L \bigwedge_x^{k-2} \oplus P_x^k$ for every k, and deduce that for $k \leq n + 1$

$$\dim P_x^k = \binom{2n}{k} - \binom{2n}{k-2}, \qquad \dim P_x^{p,q} = \binom{n}{p}\binom{n}{q} - \binom{n}{p-1}\binom{n}{q-1}$$

Compute the primitive dimensions and the multiplicities of the irreducibles for $n = 3$, and check that the total dimension comes out to 2^6.

6. Identify precisely which statements of this section require the Kähler condition and which hold on an arbitrary Hermitian manifold. Name the single input that fails without $d\omega = 0$, and explain what would go wrong if one tried to transport the triple to the cohomology of a compact Hermitian manifold such as the Hopf surface.

4.2 The Hard Lefschetz Theorem

Section 4.1 worked at a single point throughout: the triple (L, Λ, H) acts on the exterior algebra of a cotangent space, and the classification of finite-dimensional representations of $\mathfrak{sl}_2$ splits that algebra into primitive pieces. Nothing there used $d\omega = 0$, and nothing there reached cohomology. This section moves the action off the point, making $H^*(X, \mathbb{C})$ itself a module over $\mathfrak{sl}_2(\mathbb{C})$, and then reads that module.

With $H^*(X, \mathbb{C})$ a representation of $\mathfrak{sl}_2$ and H^k sitting in weight $k - n$, three statements become weight bookkeeping: that L^{n-k} carries H^k isomorphically onto H^{2n-k}, that cohomology is the direct sum of the images of the primitive parts under powers of L, and that in degree $k \leq n$ the kernel of Λ is the primitive cohomology of definition 3.2.6. The positivity that proposition 3.2.9 deferred then follows from Weil's identity, proved as theorem 4.1.5.

The bookkeeping is cleaner done once, abstractly, than three times on cohomology, so the representation theory comes first. Three facts about a finite-dimensional $\mathfrak{sl}_2$-module are needed, and all three are read off the classification of the irreducible modules rather than proved afresh: that the raising operator, applied enough times, carries a weight space isomorphically onto its mirror image; that every weight space is generated from the kernel of the lowering operator; and that being killed by the lowering operator is detected by a power of the raising operator. Write $\mathfrak{sl}_2(\mathbb{C})$ for the Lie algebra with basis e, f, h and brackets

$$[h, e] = 2e, \qquad [h, f] = -2f, \qquad [e, f] = h$$

which is the abstract shape of the relations that proposition 4.1.2 verified for (L, Λ, H).

Lemma 4.2.1: Weight Spaces and Powers of the Raising Operator

Let V be a finite-dimensional complex representation of $\mathfrak{sl}_2(\mathbb{C})$. Then h acts semisimply with integer eigenvalues, so that $V = \bigoplus_{j\in\mathbb{Z}} V_j$ with V_j the j-eigenspace of h. Write $P_j = V_j \cap \ker f$. Then:

1. $P_j = 0$ for every $j > 0$;
2. for every integer $m \geq 0$ the map $e^m \colon V_{-m} \to V_m$ is an isomorphism;
3. for every integer j,
$$V_j = \bigoplus_{r \geq \max(0,\, j)} e^r\big(P_{j-2r}\big)$$
and e^r is injective on P_{j-2r} for each r in that range, while $e^r(P_{j-2r}) = 0$ for $0 \leq r < j$;
4. for $m \geq 0$ and $v \in V_{-m}$, one has $fv = 0$ if and only if $e^{m+1}v = 0$.

***Proof*:**
Every finite-dimensional complex representation of $\mathfrak{sl}_2(\mathbb{C})$ is a direct sum of irreducible ones, and for each integer $N \geq 0$ there is exactly one irreducible module $V(N)$ of dimension $N+1$ up to isomorphism. On $V(N)$ the operator h acts semisimply with eigenvalues $N, N-2, \ldots, -N$, each eigenspace a line; e carries $V(N)_j$ isomorphically onto $V(N)_{j+2}$ for $-N \leq j \leq N-2$ and annihilates $V(N)_N$; and f carries $V(N)_j$ isomorphically onto $V(N)_{j-2}$ for $-N+2 \leq j \leq N$ and annihilates $V(N)_{-N}$. Complete reducibility and this classification are proved in *Lie Groups* [37], where the irreducible representations of $\mathrm{SU}(2)$ are determined and the passage to $\mathfrak{sl}_2(\mathbb{C})$ is by complexification; nothing about them is re-derived here.

Fix a decomposition $V = \bigoplus_i V^{(i)}$ into irreducibles. Since h acts semisimply on each summand, it does on V, and $V_j = \bigoplus_i V_j^{(i)}$ for every j; consequently $P_j = \bigoplus_i \big(V_j^{(i)} \cap \ker f\big)$, and each of the four assertions is a statement about subspaces and maps that respect the decomposition. It therefore suffices to prove them for $V = V(N)$.

1. In $V(N)$ the operator f annihilates only the line $V(N)_{-N}$, so $P_j = 0$ unless $j = -N$, and $-N \leq 0$.
2. The space $V(N)_{-m}$ is nonzero exactly when $m \leq N$ and $N - m$ is even, and the same condition characterizes $V(N)_m \neq 0$; so either both are zero, in which case there is nothing to prove, or both are lines. In the latter case e^m restricted to $V(N)_{-m}$ is the composite
$$V(N)_{-m} \to V(N)_{-m+2} \to \cdots \to V(N)_m$$
of m maps, each of which starts in a weight $j \leq m-2 \leq N-2$ and is therefore an isomorphism of lines. Hence $e^m \colon V(N)_{-m} \to V(N)_m$ is an isomorphism.
3. By part (1) the only r for which P_{j-2r} can be nonzero is the one with $j - 2r = -N$, that is $r = (j+N)/2$, and this is a nonnegative integer exactly when $j \geq -N$ and $j \equiv N$ modulo 2. Suppose first that $V(N)_j \neq 0$, so that $|j| \leq N$ and $j \equiv N$. Then $r = (j+N)/2$ satisfies $r \geq 0$ and $r \geq j$ (the latter because $r - j = (N-j)/2 \geq 0$), so it lies in the stated range,

and the composite argument of part (2) makes e^r carry $V(N)_{-N}$ isomorphically onto $V(N)_{-N+2r} = V(N)_j$, all intermediate weights being at most $j \leq N$. So the right-hand side is $V(N)_j$ and e^r is injective on P_{j-2r}.

If instead $V(N)_j = 0$, there are two cases. When $j < -N$ or $j \not\equiv N$ there is no admissible r at all and both sides vanish. When $j > N$ the admissible index is $r = (j+N)/2 < j$, which the range $r \geq \max(0, j)$ excludes, so the right-hand side is again zero. The last assertion, that $e^r(P_{j-2r}) = 0$ for $0 \leq r < j$, holds throughout that range: for the admissible index one has $r = (j+N)/2 > N$ whenever $j > N$, and e^{N+1} annihilates $V(N)$; for every other r the space P_{j-2r} is itself zero.

4. Both conditions are trivially satisfied when $V(N)_{-m} = 0$, so assume $m \leq N$ and $N \equiv m$ modulo 2, and take $0 \neq v \in V(N)_{-m}$. Then $fv = 0$ holds exactly when $V(N)_{-m}$ is the bottom line, that is when $N = m$. On the other side, $e^{m+1}v$ lies in $V(N)_{m+2}$, which is nonzero exactly when $m + 2 \leq N$; and when it is nonzero the composite argument of part (2) shows e^{m+1} is injective on $V(N)_{-m}$, so $e^{m+1}v \neq 0$. Hence $e^{m+1}v = 0$ holds exactly when $N < m + 2$, which given $N \geq m$ and $N \equiv m$ means $N = m$. The two conditions coincide, completing the proof.

Part (2) is the abstract shape of the theorem this section is named for, and part (3) the abstract shape of its decomposition; part (4) is what reconciles the two descriptions of primitivity, one by a vanishing power of the raising operator and one by the lowering operator. Only part (1) has no geometric counterpart worth naming: it records that primitive classes live in degrees at most n, which is why definition 3.2.6 was stated only there.

The module itself is not yet in hand. The triple acts on forms at every point, but only L has any reason to descend to cohomology: wedging with the closed form ω preserves closed forms and exact ones, whereas Λ is built from the metric and bears no relation to d. What carries the other two across is the harmonic representative, and the Kähler condition is spent there and nowhere else in this section.

Proposition 4.2.2: The Lefschetz Triple on Cohomology

Let X be a compact Kähler manifold of complex dimension n with Kähler form ω, let $\mathcal{H}^k$ be the space of harmonic k-forms for the associated metric, and set $\mathcal{H}^\bullet = \bigoplus_{k=0}^{2n} \mathcal{H}^k$. Then:

1. L, Λ and H carry $\mathcal{H}^\bullet$ into itself, and $\mathcal{H}^\bullet$ is a finite-dimensional $\mathfrak{sl}_2(\mathbb{C})$-module under $e = L$, $f = \Lambda$, $h = H$, in which $\mathcal{H}^k$ is the weight space of weight $k - n$;
2. the Hodge isomorphism $\mathcal{H}^k \to H^k(X, \mathbb{C})$ of theorem 2.1.1 transports that structure to cohomology: $H^\bullet(X, \mathbb{C}) = \bigoplus_{k=0}^{2n} H^k(X, \mathbb{C})$ is a finite-dimensional $\mathfrak{sl}_2(\mathbb{C})$-module in which $H^k(X, \mathbb{C})$ is the weight space of weight $k - n$, the operator e acts as cup product with $[\omega]$, and f acts by $[\alpha] \mapsto [\Lambda\alpha]$ on harmonic representatives, with $f[\alpha] = 0$ if and only if $\Lambda\alpha = 0$ at every point;
3. for $0 \leq k \leq n$ and $\alpha \in \mathcal{H}^k$, the class $[\alpha]$ lies in $H^k(X, \mathbb{C})_{\text{prim}}$ in the sense of definition 3.2.6 if and only if α is a primitive form, that is $\Lambda\alpha = 0$ at every point.

***Proof*:**
Step 1: the triple preserves harmonic forms. By theorem 1.3.1 the Laplacian of a compact Kähler manifold commutes with L and with Λ, and it commutes with H as well, since it preserves the degree while H is a scalar in each degree. So $\Delta\alpha = 0$ forces $\Delta L\alpha = L\Delta\alpha = 0$, and likewise for $\Lambda\alpha$ and $H\alpha$: all three operators carry $\mathcal{H}^\bullet$ into itself. This is the one step that uses $d\omega = 0$, through the Kähler identities that theorem 1.3.1 rests on.

Step 2: the module structure on harmonic forms. The relations $[H, L] = 2L$, $[H, \Lambda] = -2\Lambda$ and $[L, \Lambda] = H$ of proposition 4.1.2 hold at every point, hence hold between the induced operators on differential forms, hence hold on the subspace $\mathcal{H}^\bullet$ that Step 1 shows is preserved by all three. So $e \mapsto L$, $f \mapsto \Lambda$, $h \mapsto H$ makes $\mathcal{H}^\bullet$ a representation of $\mathfrak{sl}_2(\mathbb{C})$, finite dimensional because each $\mathcal{H}^k$ is (theorem 2.1.1). By definition 4.1.1 the operator H is multiplication by $k - n$ on $\mathcal{H}^k$, so $\mathcal{H}^k$ lies in the weight space of weight $k - n$; the degrees $0 \leq k \leq 2n$ give distinct weights $k - n$ and their spaces exhaust $\mathcal{H}^\bullet$, so the inclusion is an equality. This is part (1).

Step 3: transport. By theorem 2.1.1 every de Rham class has exactly one harmonic representative, so $\alpha \mapsto [\alpha]$ is an isomorphism $\mathcal{H}^k \to H^k(X, \mathbb{C})$; carrying the operators of Step 2 through it makes $H^\bullet(X, \mathbb{C})$ a module with $H^k(X, \mathbb{C})$ the weight space of weight $k - n$. The transported e is cup product with $[\omega]$: for α harmonic the form $L\alpha = \omega \wedge \alpha$ is harmonic by Step 1 and represents $[\omega] \cup [\alpha]$, both factors being closed. The description of f is the definition of the transported operator, and for the last clause of part (2), $\Lambda\alpha$ is harmonic by Step 1, while a harmonic form whose class vanishes is exact and therefore orthogonal to itself in the decomposition of theorem 2.1.1, hence zero.

Step 4: primitive classes and primitive forms. Let $0 \leq k \leq n$ and $\alpha \in \mathcal{H}^k$. By Step 1 the form $L^{n-k+1}\alpha$ is harmonic, and it represents $L^{n-k+1}[\alpha]$; by the vanishing criterion of Step 3 it is therefore zero as a form exactly when that class is zero, which by definition 3.2.6 is exactly the condition $[\alpha] \in H^k(X, \mathbb{C})_{\text{prim}}$. On the other hand part (1) of theorem 4.1.4, applied to $\alpha_x \in \bigwedge^k_x$ at a point x, says that $L^{n-k+1}\alpha_x = 0$ if and only if $\Lambda\alpha_x = 0$. Reading this at every point, $L^{n-k+1}\alpha$ vanishes identically if and only if $\Lambda\alpha$ does, so the two conditions of part (3) agree, completing the proof.

The purchase has two halves. Part (2) puts the action on $H^*(X, \mathbb{C})$ at all, which is what the weight bookkeeping of lemma 4.2.1 is applied to below. Part (3) says more: primitivity of a class, a condition written with $[\omega]$ alone, is realized pointwise by one distinguished representative, and a pointwise primitive form is what theorem 4.1.5 accepts. The transported operator $f = \Lambda$, by contrast, depends on the metric and not only on its class; exercise 4.2.1 (5) takes that up.

Part (2) of the lemma, applied at the weight of $H^k(X, \mathbb{C})$, is the theorem over $\mathbb{C}$; the real, rational and bigraded refinements follow it by inspection.

Theorem 4.2.3: The Hard Lefschetz Theorem

Let X be a compact Kähler manifold of complex dimension n with Kähler form ω, and let L denote the induced Lefschetz operator on cohomology. For each k with $0 \leq k \leq n$ the map

$$L^{n-k}\colon H^k(X,\mathbb{R}) \to H^{2n-k}(X,\mathbb{R}), \qquad L^{n-k}[\alpha] = [\omega^{n-k} \wedge \alpha]$$

is an isomorphism. The same holds with $\mathbb{R}$ replaced by $\mathbb{C}$, and by $\mathbb{Q}$ when $[\omega]$ lies in the image of $H^2(X,\mathbb{Q})$. Moreover L^{n-k} respects the Hodge decomposition, restricting for each $p+q=k$ to an isomorphism

$$L^{n-k}\colon H^{p,q}(X) \to H^{p+n-k,\,q+n-k}(X)$$

***Proof*:**

Step 1: the module. By proposition 4.2.2 the graded space

$$H^\bullet(X,\mathbb{C}) = \bigoplus_{k=0}^{2n} H^k(X,\mathbb{C})$$

is a finite-dimensional $\mathfrak{sl}_2(\mathbb{C})$-module under $e = L$, $f = \Lambda$, $h = H$, in which $H^k(X,\mathbb{C})$ is the weight space of weight $k-n$ and e acts as the Lefschetz operator induced on cohomology, that is, as cup product with $[\omega]$. Of the three operators only Λ passes through a choice of metric, being transported along harmonic representatives; that transport is where the Kähler condition is spent.

Step 2: the weight count. Fix k with $0 \leq k \leq n$ and set $m = n-k \geq 0$. The weight space V_{-m} of the module of Step 1 is $H^k(X,\mathbb{C})$, and the weight space V_m is $H^{k'}(X,\mathbb{C})$ for the degree k' with $k'-n=m$, that is $k' = 2n-k$. Part (2) of lemma 4.2.1 therefore gives that

$$L^{n-k}\colon H^k(X,\mathbb{C}) \to H^{2n-k}(X,\mathbb{C})$$

is an isomorphism.

Step 3: real and rational coefficients. The form ω is real, so L carries $H^\bullet(X,\mathbb{R})$ into itself, and $H^k(X,\mathbb{C}) = H^k(X,\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$ realizes the map of Step 2 as the complexification of the real map $L^{n-k}\colon H^k(X,\mathbb{R}) \to H^{2n-k}(X,\mathbb{R})$. A linear map between real vector spaces is bijective if and only if its complexification is, so the real statement follows. If $[\omega]$ lies in the image of $H^2(X,\mathbb{Q})$, then L carries $H^\bullet(X,\mathbb{Q})$ into itself and the same argument applies with $\mathbb{Q}$ in place of $\mathbb{R}$.

Step 4: compatibility with the bigrading. The operator L raises the bidegree by $(1,1)$, so L^{n-k} carries $H^{p,q}(X)$ into $H^{p+n-k,q+n-k}(X)$ for $p+q=k$. Distinct summands of $H^k(X,\mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ therefore land in distinct summands of $H^{2n-k}(X,\mathbb{C}) = \bigoplus_{p'+q'=2n-k} H^{p',q'}(X)$, and every summand of the target is reached: given $p'+q' = 2n-k$ with $p', q' \leq n$, the inequality $q' \leq n$ forces $p' \geq n-k$, so $p = p'-(n-k)$ and $q = q'-(n-k)$ are nonnegative and satisfy $p+q=k$. The isomorphism of Step 2 is thus a direct sum of the maps $H^{p,q}(X) \to H^{p+n-k,q+n-k}(X)$ over $p+q=k$, and a direct sum of linear maps is bijective precisely when every summand map is, so each of them is an isomorphism, as we sought to show.

Poincaré duality already gives $b_k = b_{2n-k}$ on any closed oriented $2n$-manifold, proved in *Algebraic Topology* [30, Consequences of Poincaré Duality], so the equality of dimensions is not new. What the

theorem adds is an isomorphism realizing it, and one of a restricted kind: cup product with a fixed class. That the $(n-k)$-fold cup product with $[\omega]$ should be invertible is a strong constraint, and it is not available without the Kähler hypothesis. For $2 \leq k \leq n$ the map $L\colon H^{k-2}(X,\mathbb{C}) \to H^k(X,\mathbb{C})$ is injective, because $L^{n-k+2} = L^{n-k+1} \circ L$ is an isomorphism on $H^{k-2}(X,\mathbb{C})$ by the theorem; hence $b_{k-2} \leq b_k$ for $k \leq n$, and combined with Poincaré duality the even-indexed Betti numbers of a compact Kähler manifold increase up to the middle degree and decrease after it, as do the odd-indexed ones (exercise 4.2.1 (2)). The two parity classes are unrelated, so the full sequence $b_0, b_1, \ldots, b_{2n}$ need not be unimodal: a K3 surface carries $b_2 = 22$ between the vanishing b_1 and b_3. The Hopf surface of box 1.4.2 has $b_0 = 1$ and $b_2 = 0$, so it violates the inequality at $k = 2 = n$, which is a second proof that it carries no Kähler metric.

Projective space is the case where the module of proposition 4.2.2 can be named outright. By example 1.2.5 the Hodge numbers of $\mathbb{C}\mathrm{P}^n$ are $h^{p,q} = 1$ for $p = q \leq n$ and zero otherwise, so corollary 2.1.3 gives $b_{2j} = 1$ for $0 \leq j \leq n$ and $b_k = 0$ for k odd. Each $H^{2j}(\mathbb{C}\mathrm{P}^n,\mathbb{C})$ is therefore a line, and it is spanned by $[\omega]^j$ for the Fubini-Study form: the class $[\omega]^j$ is nonzero because $[\omega]^j \cup [\omega]^{n-j}$ integrates to $\int_{\mathbb{C}\mathrm{P}^n} \omega^n > 0$. So $H^*(\mathbb{C}\mathrm{P}^n,\mathbb{C})$ has one line in each of the weights $-n, -n+2, \ldots, n$ and none elsewhere, which forces it to be the irreducible module $V(n)$ of dimension $n+1$ (exercise 4.2.1 (1)), generated from the class of the constant function 1 in degree zero.

Part (3) of the lemma delivers the corresponding decomposition in general, and it is the geometric statement that definition 3.2.6 was set up to receive.

Corollary 4.2.4: The Lefschetz Decomposition

Let X be a compact Kähler manifold of complex dimension n with Kähler form ω.

1. For $0 \leq k \leq n$,
$$H^k(X,\mathbb{C})_{\mathrm{prim}} = \ker\bigl(\Lambda\colon H^k(X,\mathbb{C}) \to H^{k-2}(X,\mathbb{C})\bigr)$$
where Λ acts through harmonic representatives, and $\ker\Lambda \cap H^k(X,\mathbb{C}) = 0$ for $k > n$.
2. For every k with $0 \leq k \leq 2n$,
$$H^k(X,\mathbb{C}) = \bigoplus_{r \geq \max(0,\,k-n)} L^r\, H^{k-2r}(X,\mathbb{C})_{\mathrm{prim}}$$
the sum running over those r with $2r \leq k$, and L^r is injective on each summand shown.
3. The decomposition is compatible with the Hodge decomposition, in that $L^r H^{p,q}(X)_{\mathrm{prim}} \subseteq H^{p+r,q+r}(X)$, and it holds verbatim with $\mathbb{C}$ replaced by $\mathbb{R}$, or by $\mathbb{Q}$ when $[\omega]$ is rational.
4. For $0 \leq k \leq n$ one has $\dim_{\mathbb{C}} H^k(X,\mathbb{C})_{\mathrm{prim}} = b_k - b_{k-2}$, with $b_{-2} = b_{-1} = 0$.

***Proof*:**
Work throughout in the $\mathfrak{sl}_2(\mathbb{C})$-module $H^\bullet(X,\mathbb{C})$ of proposition 4.2.2, where $e = L$, $f = \Lambda$, and $H^k(X,\mathbb{C})$ is the weight space of weight $k-n$. Write P_j for $V_j \cap \ker f$ as in lemma 4.2.1, so that $P_{k-n} = \ker\Lambda \cap H^k(X,\mathbb{C})$.

1. Let $0 \leq k \leq n$ and put $m = n-k \geq 0$. Part (4) of lemma 4.2.1 says that a class

$\alpha \in H^k(X,\mathbb{C})$ satisfies $\Lambda\alpha = 0$ if and only if $L^{m+1}\alpha = 0$; and $L^{m+1} = L^{n-k+1}$ is exactly the operator whose kernel defines $H^k(X,\mathbb{C})_{\text{prim}}$ in definition 3.2.6. Hence $H^k(X,\mathbb{C})_{\text{prim}} = P_{k-n}$. For $k > n$ the weight $k-n$ is positive, so $P_{k-n} = 0$ by part (1) of the lemma.

2. Part (1) of the lemma lets the notation $H^{k'}(X,\mathbb{C})_{\text{prim}} = P_{k'-n}$ be extended by zero to all degrees k', consistently with part (1) of the present corollary. Applying part (3) of lemma 4.2.1 at $j = k-n$ gives

$$H^k(X,\mathbb{C}) = V_{k-n} = \bigoplus_{r \geq \max(0,\,k-n)} L^r\big(P_{k-n-2r}\big) = \bigoplus_{r \geq \max(0,\,k-n)} L^r\, H^{k-2r}(X,\mathbb{C})_{\text{prim}}$$

since $P_{k-n-2r} = P_{(k-2r)-n}$, with L^r injective on each summand. The terms with $2r > k$ vanish because $H^{k-2r}(X,\mathbb{C}) = 0$ in negative degrees, and the lower bound $r \geq k-n$ is what the lemma gives: below it the summand is zero, a primitive class in degree $k' \leq n$ being annihilated by L^r as soon as $r > n - k'$.

3. The operator L raises the bidegree by $(1,1)$, and proposition 3.2.7 gives $H^{k'}(X,\mathbb{C})_{\text{prim}} = \bigoplus_{p+q=k'} H^{p,q}(X)_{\text{prim}}$, so each summand $L^r H^{k-2r}(X,\mathbb{C})_{\text{prim}}$ decomposes into bihomogeneous pieces of the stated types. For the real statement, ω is a real form, so L and Λ are real operators and each subspace appearing in part (2) is stable under complex conjugation. The projections of a direct sum whose summands are all conjugation-stable commute with conjugation, since the components of $\bar{\alpha}$ are then the conjugates of the components of α; taking real points of both sides gives the decomposition of $H^k(X,\mathbb{R})$. When $[\omega]$ is rational the same argument applies over $\mathbb{Q}$, where $H^{k'}(X,\mathbb{Q})_{\text{prim}}$ is defined by definition 3.2.6.

4. Let $0 \leq k \leq n$. Separating the term $r = 0$ in part (2) and re-indexing the rest by $r' = r-1$,

$$\begin{aligned} H^k(X,\mathbb{C}) &= H^k(X,\mathbb{C})_{\text{prim}} \oplus L\Big(\bigoplus_{r' \geq 0} L^{r'} H^{(k-2)-2r'}(X,\mathbb{C})_{\text{prim}}\Big) \\ &= H^k(X,\mathbb{C})_{\text{prim}} \oplus L\big(H^{k-2}(X,\mathbb{C})\big) \end{aligned}$$

the inner sum being the decomposition of $H^{k-2}(X,\mathbb{C})$, whose lower bound is $\max(0, k-2-n) = 0$. Since $k-2 \leq n-2 < n$, the map L is injective on $H^{k-2}(X,\mathbb{C})$, as recorded after theorem 4.2.3, so the second summand has dimension b_{k-2} and $\dim H^k(X,\mathbb{C})_{\text{prim}} = b_k - b_{k-2}$, completing the proof.

Part (1) reconciles the two descriptions of primitivity that the book has used. Definition 3.2.6 defined the primitive part by the vanishing of L^{n-k+1}, a condition visibly independent of the metric and depending on the Kähler class alone; the kernel of Λ is the description that makes the representation theory work but refers to harmonic representatives. They agree, so the Lefschetz decomposition is determined by $[\omega]$ even though its proof runs through a metric. Part (2) is what chapter 3 promised when it introduced the primitive cohomology and then had nothing to decompose: cohomology in every degree is assembled from primitive pieces sitting in degrees of the same parity, each shifted upward by a power of L. Part (4) makes the count explicit, and shows that the primitive cohomology is not a small correction: on a surface with $b_2 = 22$ it accounts for 21 of the 22 dimensions.

In complex dimension two the whole apparatus fits in a few lines, and a flat torus makes every number in it explicit.

Example 4.2.5: Hard Lefschetz on a Compact Kähler Surface
Let X be a compact Kähler surface, so $n = 2$. Theorem 4.2.3 asserts two isomorphisms,

$$L^2 \colon H^0(X,\mathbb{R}) \to H^4(X,\mathbb{R}), \qquad L \colon H^1(X,\mathbb{R}) \to H^3(X,\mathbb{R})$$

and in the middle degree $k = n = 2$ it asserts only that the identity is an isomorphism, the content there being the decomposition instead. The first map sends 1 to $[\omega^2]$, which is nonzero because $\int_X \omega \wedge \omega > 0$ by part (2) of lemma 2.2.2; since $b_0 = b_4 = 1$ this alone makes it an isomorphism.

The middle degree. Here $L^{n-k+1} = L \colon H^2 \to H^4$, so $H^2(X,\mathbb{R})_{\text{prim}}$ is the cup-orthogonal complement of $[\omega]$, as recorded after definition 3.2.6, and corollary 4.2.4 reads

$$H^2(X,\mathbb{R}) = H^2(X,\mathbb{R})_{\text{prim}} \oplus \mathbb{R}\,[\omega]$$

with $\dim H^2(X,\mathbb{R})_{\text{prim}} = b_2 - b_0 = b_2 - 1$. In degree one, $L^{n-k+1} = L^2$ lands in $H^5(X,\mathbb{R}) = 0$, so $H^1(X,\mathbb{R})_{\text{prim}} = H^1(X,\mathbb{R})$ and $\dim H^1_{\text{prim}} = b_1 - b_{-1} = b_1$; in degrees three and four the decomposition reads $H^3 = L\,H^1_{\text{prim}}$ and $H^4 = L^2 H^0_{\text{prim}}$, with no primitive cohomology of its own, in agreement with part (1) of the corollary.

The isomorphism $L \colon H^1 \to H^3$ on a torus. Take $X = \mathbb{C}^2/\Lambda$ with the flat Kähler form $\omega = i\,(dz_1 \wedge d\bar{z}_1 + dz_2 \wedge d\bar{z}_2)$. By example 2.1.2 the harmonic forms are the constant-coefficient ones, so $H^1(X,\mathbb{C})$ has basis $dz_1, dz_2, d\bar{z}_1, d\bar{z}_2$ and $H^3(X,\mathbb{C})$ has basis $dz_1 \wedge dz_2 \wedge d\bar{z}_1$, $dz_1 \wedge dz_2 \wedge d\bar{z}_2$, $dz_1 \wedge d\bar{z}_1 \wedge d\bar{z}_2$, $dz_2 \wedge d\bar{z}_1 \wedge d\bar{z}_2$. Wedging with ω kills every term containing a repeated differential, leaving

$$\omega \wedge dz_1 = i\,dz_1 \wedge dz_2 \wedge d\bar{z}_2, \qquad \omega \wedge dz_2 = -i\,dz_1 \wedge dz_2 \wedge d\bar{z}_1$$

$$\omega \wedge d\bar{z}_1 = -i\,dz_2 \wedge d\bar{z}_1 \wedge d\bar{z}_2, \qquad \omega \wedge d\bar{z}_2 = i\,dz_1 \wedge d\bar{z}_1 \wedge d\bar{z}_2$$

each sign coming from moving the surviving factor into increasing order. The four images are the four basis elements of $H^3(X,\mathbb{C})$ up to the scalar $\pm i$, so L is an isomorphism $H^1 \to H^3$, as the theorem requires.

The numbers on the torus. Here $b_0 = 1$, $b_1 = 4$, $b_2 = 6$, $b_3 = 4$, $b_4 = 1$, so the primitive dimensions are 1, 4 and $6 - 1 = 5$ in degrees 0, 1, 2. Each primitive class in degree k generates a module $V(n-k)$ of dimension $n-k+1$, so the total dimension of $H^*(X,\mathbb{C})$ is $1\cdot 3 + 4\cdot 2 + 5\cdot 1 = 16$, which is $\sum_k b_k = 1+4+6+4+1$. In the middle degree the primitive Hodge numbers are $h^{2,0}_{\text{prim}} = h^{0,2}_{\text{prim}} = 1$ and $h^{1,1}_{\text{prim}} = 4 - 1 = 3$, the Kähler class accounting for the missing $(1,1)$-dimension, exactly as in example 3.2.10.

The surface also shows what is still owed. Example 3.2.10 checked that $Q = -\int_X \alpha \wedge \beta$ polarizes the primitive part of H^2, but that verification ran through lemma 2.2.2, three signs computed in a unitary coframe on a four-manifold, and every step of it was tied to the dimension. In arbitrary weight the corresponding positivity was stated in proposition 3.2.9 and left unproved. It comes out of Weil's identity, which is a statement about forms that are primitive at each point, and the form to feed it is fixed by part (3) of proposition 4.2.2: the harmonic representative of a primitive class is pointwise primitive, and no other representative need be, since an arbitrary one can be modified by any exact form. What is left is the accounting of signs.

Corollary 4.2.6: Positivity on Primitive Cohomology

Let X be a compact Kähler manifold of complex dimension n with Kähler form ω, let $0 \leq k \leq n$, and let

$$Q(\alpha, \beta) = (-1)^{k(k-1)/2} \int_X \alpha \wedge \beta \wedge \omega^{n-k}$$

be the form of proposition 3.2.9. If α is a nonzero class in $H^{p,q}(X)_{\text{prim}}$ with $p+q=k$, and α also denotes its harmonic representative, then

$$i^{p-q}\, Q\big(\alpha, \bar{\alpha}\big) = (n-k)! \int_X |\alpha|^2 \,\text{vol} > 0$$

So Q satisfies the second Hodge-Riemann bilinear relation of definition 3.2.1 on $H^k(X, \mathbb{C})_{\text{prim}}$, and proposition 3.2.9 holds as stated: whenever $H^k(X, \mathbb{Q})_{\text{prim}}$ is defined over $\mathbb{Q}$, the form Q is a polarization of it.

***Proof*:**

Step 1: the forms involved are primitive. By part (3) of proposition 4.2.2 the harmonic representative α satisfies $\Lambda\alpha = 0$ at every point, and it has type (p, q) because the Hodge decomposition assigns each class of type (p, q) a harmonic representative of that type (theorem 2.1.1). Complex conjugation preserves harmonicity by Step 4 of the proof of that theorem and commutes with Λ, the form ω being real, so $\bar{\alpha}$ is a harmonic primitive form of type (q, p) and again of total degree k.

Step 2: applying Weil's identity. Theorem 3.2.8, proved in theorem 4.1.5, evaluates the Hodge star of a primitive form of type (q, p) with $p+q=k$. Applied to $\bar{\alpha}$ it gives

$$\star\,\bar{\alpha} = (-1)^{k(k+1)/2}\, i^{q-p}\, \frac{1}{(n-k)!}\, L^{n-k}\bar{\alpha}$$

Step 3: pairing against α. The Hodge star of definition 1.1.8 is characterized by $\alpha \wedge \star\bar{\beta} = \langle \alpha, \beta \rangle \,\text{vol}$ for the pointwise Hermitian inner product on forms; with $\beta = \alpha$ this reads $\alpha \wedge \star\bar{\alpha} = |\alpha|^2 \,\text{vol}$. Substituting Step 2,

$$|\alpha|^2 \,\text{vol} = (-1)^{k(k+1)/2}\, i^{q-p}\, \frac{1}{(n-k)!}\, \alpha \wedge \bar{\alpha} \wedge \omega^{n-k}$$

and solving for the wedge, using $(-1)^{-a} = (-1)^a$ and $i^{-(q-p)} = i^{p-q}$,

$$\alpha \wedge \bar{\alpha} \wedge \omega^{n-k} = (n-k)!\, (-1)^{k(k+1)/2}\, i^{p-q}\, |\alpha|^2 \,\text{vol} \tag{4.3}$$

Step 4: collecting the signs. Multiply (4.3) by $i^{p-q}(-1)^{k(k-1)/2}$ and integrate. The two sign factors combine as

$$(-1)^{k(k-1)/2}\, (-1)^{k(k+1)/2} = (-1)^{k(k-1)/2 + k(k+1)/2} = (-1)^{k^2} = (-1)^k$$

while the two powers of i combine as $i^{2(p-q)} = (-1)^{p-q} = (-1)^{p+q} = (-1)^k$, the integers $p-q$ and $p+q$ having the same parity. Their product is $(-1)^{2k} = 1$, so

$$i^{p-q}\, Q(\alpha, \bar{\alpha}) = i^{p-q}(-1)^{k(k-1)/2} \int_X \alpha \wedge \bar{\alpha} \wedge \omega^{n-k} = (n-k)! \int_X |\alpha|^2 \,\text{vol}$$

The integrand is a nonnegative function vanishing exactly where α does, and α is not the zero form because its class is nonzero, so the integral is strictly positive.

Step 5: the polarization. The value of Q depends only on the classes, being an integral of a product of closed forms, so the inequality just proved is a statement about $H^k(X,\mathbb{C})_{\text{prim}}$ and is condition (2) of definition 3.2.1. The symmetry $Q(\alpha,\beta) = (-1)^k Q(\beta,\alpha)$ and condition (1) were verified in proposition 3.2.9, and in theorem 3.2.4 for the underlying bidegree statement, by counts that used nothing deferred. All three conditions now hold, so Q is a polarization of $H^k(X,\mathbb{Q})_{\text{prim}}$ whenever that space is defined over $\mathbb{Q}$, completing the proof.

The corollary closes the last gap in proposition 3.2.9, and the gap was of one specific kind. The symmetry type and the first Hodge-Riemann relation were bookkeeping with bidegrees, valid on any compact Kähler manifold and requiring no machinery. Positivity was different: it needed the Hodge star of a primitive form to be expressible through L, and that identity is a statement about the $\mathfrak{sl}_2$-decomposition of the exterior algebra, which is why the verification requires section 4.1. The normalizing sign $(-1)^{k(k-1)/2}$ carried by Q is what meets the $(-1)^{k(k+1)/2}$ produced by Weil's identity, and Step 4 is the computation that the two, together with the factor i^{p-q} prescribed by definition 3.2.1, cancel exactly.

Two consequences follow at once. First, on a projective manifold with ω the restriction of the Fubini-Study form, the class $[\omega]$ is integral (theorem 1.1.7), so $H^k(X,\mathbb{Q})_{\text{prim}}$ is a rational Hodge structure and Q polarizes it; corollary 3.2.5 then computes the signature of Q from the primitive Hodge numbers alone, with no further geometry, and exercise 3.2.1 (3) makes every sub-Hodge structure of it a direct summand. Second, combining with corollary 4.2.4, the whole of $H^k(X,\mathbb{Q})$ is a direct sum of copies of polarizable structures shifted by powers of L (exercise 4.2.1 (4)), so every cohomology group of a projective manifold is polarizable, not only the primitive part. That is the property chapter 5 carries into families.

Exercise 4.2.1

1. Let ω be the Fubini-Study form on $\mathbb{CP}^n$.
    1. Using example 1.2.5 and corollary 2.1.3, compute $\dim H^k(\mathbb{CP}^n,\mathbb{C})_{\text{prim}}$ for every k with $0 \le k \le n$, and write out the Lefschetz decomposition of $H^{2j}(\mathbb{CP}^n,\mathbb{C})$.
    2. Verify theorem 4.2.3 directly on $\mathbb{CP}^3$ by exhibiting the isomorphisms $L^3 \colon H^0 \to H^6$ and $L \colon H^2 \to H^4$ on explicit bases, having first shown that $[\omega]^j \neq 0$ for $0 \le j \le 3$.
    3. Deduce that $H^*(\mathbb{CP}^n,\mathbb{C})$ is an irreducible representation of $\mathfrak{sl}_2(\mathbb{C})$, and identify which one.
2. Let X be a compact Kähler manifold of complex dimension n.
    1. Show that $b_0(X) \le b_2(X) \le \cdots$ and $b_1(X) \le b_3(X) \le \cdots$, the inequalities running up to degree n, and combine this with Poincaré duality to describe the shape of the sequence $b_0, b_1, \ldots, b_{2n}$.
    2. The Hopf surface S of box 1.4.2 is diffeomorphic to $S^1 \times S^3$. Compute its Betti numbers and show that they violate the inequalities of part (1), giving a proof that S is not Kähler that does not use the parity of b_1.
3. Let $X = \mathbb{C}^n/\Lambda$ be a complex torus with a flat Kähler metric.

1. Using example 2.1.2, show that $\dim H^k(X,\mathbb{C})_{\text{prim}} = \binom{2n}{k} - \binom{2n}{k-2}$ for $0 \leq k \leq n$, and evaluate this for $n = 3$ and $k = 0, 1, 2, 3$.
2. Check that $\sum_{k=0}^{n}(n-k+1)\dim H^k(X,\mathbb{C})_{\text{prim}} = 2^{2n}$ for $n = 3$, and explain why this identity holds for every compact Kähler manifold with 2^{2n} replaced by $\sum_k b_k$.

4. Let X be a compact Kähler manifold of dimension n whose Kähler class is rational, and let $0 \leq k \leq 2n$.
   1. Show that for each admissible r the map L^r is an injective morphism of rational Hodge structures
   $$L^r \colon H^{k-2r}(X,\mathbb{Q})_{\text{prim}}(-r) \to H^k(X,\mathbb{Q})$$
   where $(-r)$ is the Tate twist of definition 3.1.11.
   2. Deduce that corollary 4.2.4 is an isomorphism of rational Hodge structures, and that $H^k(X,\mathbb{Q})$ is polarizable.

5. Let X be a compact Kähler manifold of complex dimension n.
   1. Show that for $0 \leq k \leq n$ the map $\Lambda^{n-k} \colon H^{2n-k}(X,\mathbb{C}) \to H^k(X,\mathbb{C})$ is an isomorphism, where Λ acts through harmonic representatives. (Argue as in lemma 4.2.1, with f in place of e.)
   2. Show that the subspaces $H^k(X,\mathbb{R})_{\text{prim}}$ and the summands of corollary 4.2.4 depend on the Kähler metric only through its class $[\omega] \in H^2(X,\mathbb{R})$, and explain why the same is not immediate for the operator Λ of part (1).

4.3 Lefschetz Pencils and Vanishing Cycles

Section 4.2 settled what a Kähler class does to the cohomology of a single manifold. The rest of the chapter needs a second object: the hyperplane sections of a projective variety, taken not one at a time but as a family. This section builds that family and identifies the homology classes its degenerations produce.

The first result concerns a single section, and it is what makes the family worth building. Below the middle degree a smooth hyperplane section already carries the cohomology of the variety containing it, so everything peculiar to a smooth projective X of dimension n sits in degree n. The pencil construction, the blow-up that turns a pencil into a fibration over $\mathbb{CP}^1$, and the vanishing cycles attached to its singular fibres are the machinery for reaching that degree.

Throughout, $X \subseteq \mathbb{CP}^N$ is a smooth connected projective variety of complex dimension n, carrying the Kähler structure restricted from the Fubini-Study metric (theorem 1.1.7). Hyperplanes of $\mathbb{CP}^N$ are parametrized by the dual projective space $(\mathbb{CP}^N)^\vee$, a point of which is the line spanned by a nonzero linear form s on $\mathbb{C}^{N+1}$, and $H_s = \{s = 0\}$ is the corresponding hyperplane. For $X \not\subseteq H$ the intersection $Y = X \cap H$ is a hypersurface in X, of dimension $n-1$.

The comparison between X and such a Y rests on a single geometric fact: the complement $X \setminus Y$ is affine. An affine variety of complex dimension n is a very thin object topologically, having no homology at all above degree n, and the relative cohomology of the pair (X, Y) is exactly the cohomology of that complement, read backwards.

Theorem 4.3.1: The Weak Lefschetz Theorem

Let $X \subseteq \mathbb{C}\mathrm{P}^N$ be a smooth connected projective variety of dimension $n \geq 1$, let H be a hyperplane with $X \not\subseteq H$, and suppose $Y = X \cap H$ is smooth. Write $i\colon Y \hookrightarrow X$ for the inclusion. Then

$$i^*\colon H^k(X,\mathbb{Z}) \to H^k(Y,\mathbb{Z})$$

is an isomorphism for $k \leq n-2$ and injective for $k = n-1$, and

$$i_*\colon H_k(Y,\mathbb{Z}) \to H_k(X,\mathbb{Z})$$

is an isomorphism for $k \leq n-2$ and surjective for $k = n-1$.

***Proof*:**
Set $U = X \setminus Y$.

Step 1: U is a closed complex submanifold of $\mathbb{C}^N$ of dimension n. The complement $\mathbb{C}\mathrm{P}^N \setminus H$ is biholomorphic to $\mathbb{C}^N$: after a linear change of homogeneous coordinates $H = \{z_0 = 0\}$, and $[z_0 : \cdots : z_N] \mapsto (z_1/z_0, \ldots, z_N/z_0)$ is a biholomorphism onto $\mathbb{C}^N$. Since X is closed in $\mathbb{C}\mathrm{P}^N$, the set $U = X \cap (\mathbb{C}\mathrm{P}^N \setminus H)$ is closed in $\mathbb{C}\mathrm{P}^N \setminus H = \mathbb{C}^N$, and it is a complex submanifold of dimension n because X is one. It is nonempty because $X \not\subseteq H$.

Step 2: every Morse index of a squared-distance function on U is at most n. For $a \in \mathbb{C}^N$ put $\varphi_a(z) = |z-a|^2$ and consider its restriction to U. It is proper: the preimage of $[0,R]$ is the intersection of U with a closed ball, which is closed and bounded in $\mathbb{C}^N$, hence compact. For all a outside a set of measure zero the restriction $\varphi_a|_U$ is a Morse function [32, Existence of Morse Functions]; fix such an a.

Let p be a critical point of $\varphi_a|_U$, let Hess_p denote the Hessian of $\varphi_a|_U$ at p, a real quadratic form on the real tangent space T_pU, and let J be the complex structure on T_pU. The claim is that

$$\mathrm{Hess}_p(v,v) + \mathrm{Hess}_p(Jv,Jv) = 4|v|^2 \tag{4.4}$$

for every $v \in T_pU$, the norm on the right being the Euclidean norm of v regarded as a vector in $\mathbb{C}^N$. To see it, choose a holomorphic chart of U centred at p and compose it with a complex line through the origin to produce a holomorphic map $u\colon \Delta \to U \subseteq \mathbb{C}^N$ from a disc with $u(0) = p$ and $u'(0) = v$. Because p is a critical point of $\varphi_a|_U$, the chain rule for second derivatives loses its second-order term and gives

$$\mathrm{Hess}_0(\varphi_a \circ u)(\xi,\eta) = \mathrm{Hess}_p\big(du_0(\xi), du_0(\eta)\big)$$

for all real tangent vectors ξ, η of Δ at 0. Writing $\zeta = \sigma + i\tau$ for the coordinate on Δ and $\psi = \varphi_a \circ u$, the left side evaluated on the pairs $(\partial_\sigma, \partial_\sigma)$ and $(\partial_\tau, \partial_\tau)$ sums to the Laplacian $\Delta_\zeta \psi(0)$. Now $\psi(\zeta) = \sum_{m=1}^N |u_m(\zeta) - a_m|^2$ with each u_m holomorphic, so $\partial_\zeta \partial_{\bar\zeta} \psi = \sum_m |u_m'|^2$ and

$$\Delta_\zeta \psi(0) = 4\sum_m |u_m'(0)|^2 = 4|v|^2$$

On the right side, $du_0(\partial_\sigma) = v$ and $du_0(\partial_\tau) = Jv$, since u is holomorphic and therefore intertwines the complex structures. This is (4.4).

Let $W \subseteq T_pU$ be a real subspace on which Hess_p is negative definite. If some nonzero v lay in $W \cap JW$, say $v = Jw$ with $w \in W$, then both $\mathrm{Hess}_p(w,w) < 0$ and $\mathrm{Hess}_p(Jw,Jw) =$

$\mathrm{Hess}_p(v,v) < 0$, contradicting (4.4) applied to w. So $W \cap JW = 0$, whence $2\dim_{\mathbb{R}} W = \dim_{\mathbb{R}}(W \oplus JW) \le \dim_{\mathbb{R}} T_pU = 2n$ and $\dim_{\mathbb{R}} W \le n$. The Morse index at p is the dimension of a maximal such W, so it is at most n.

Step 3: $H_j(U,\mathbb{Z}) = H^j(U,\mathbb{Z}) = 0$ for $j > n$. The function $\varphi_a|_U$ is proper, bounded below and Morse with all indices at most n, so U has the homotopy type of a CW complex whose cells all have dimension at most n [32, Morse CW Structure for a Proper Function]. Cellular chains and cochains vanish above dimension n, so the homology and cohomology do too.

Step 4: the relative cohomology of the pair. Fix nested closed tubular neighbourhoods $N' \subseteq N^\circ$ of the smooth compact submanifold $Y \subseteq X$ [32, Tubular Neighbourhood Theorem]. Each retracts onto Y, so $H^k(X,Y;\mathbb{Z}) \cong H^k(X,N;\mathbb{Z})$. Excising the open set N'°, whose closure lies in the interior of N, and then retracting the collar $N \setminus N'^\circ$ onto $\partial N'$ gives

$$H^k(X,N;\mathbb{Z}) \cong H^k\big(X \setminus N'^\circ,\ \partial N';\mathbb{Z}\big)$$

The space $M = X \setminus N'^\circ$ is a compact oriented $2n$-manifold with boundary $\partial N'$, oriented by the complex structure of X, so Lefschetz duality [30, Poincaré-Lefschetz Duality] gives $H^k(M,\partial M;\mathbb{Z}) \cong H_{2n-k}(M,\mathbb{Z})$. Finally M is a deformation retract of U, since U is M together with the half-open collar $N' \setminus Y$. Combining,

$$H^k(X,Y;\mathbb{Z}) \cong H_{2n-k}(U,\mathbb{Z})$$

and the same chain with the roles of homology and cohomology exchanged gives $H_k(X,Y;\mathbb{Z}) \cong H^{2n-k}(U,\mathbb{Z})$. By Step 3 both vanish as soon as $2n-k > n$, that is for $k \le n-1$.

Step 5: the long exact sequences. For the pair (X,Y),

$$H^k(X,Y;\mathbb{Z}) \to H^k(X,\mathbb{Z}) \xrightarrow{i^*} H^k(Y,\mathbb{Z}) \to H^{k+1}(X,Y;\mathbb{Z})$$

is exact. For $k \le n-2$ both outer terms vanish by Step 4, so i^* is an isomorphism; for $k = n-1$ the left term vanishes, so i^* is injective. The homology sequence

$$H_{k+1}(X,Y;\mathbb{Z}) \to H_k(Y,\mathbb{Z}) \xrightarrow{i_*} H_k(X,\mathbb{Z}) \to H_k(X,Y;\mathbb{Z})$$

gives the second statement by the same two vanishings, completing the proof.

The theorem is a statement about integral cohomology, and that is worth separating from what the Kähler package alone delivers. Comparing the two needs a map running opposite to restriction, which Poincaré duality gives. Write D_X and D_Y for the duality isomorphisms of X and Y ([30, Poincaré duality]) and set

$$i_! := D_X^{-1} \circ i_* \circ D_Y \colon H^k(Y,\mathbb{Z}) \to H^{k+2}(X,\mathbb{Z}),$$

the *Gysin map*, the homology pushforward of theorem 4.3.1 read through duality on both sides; the degree shifts by 2 because Y has complex codimension 1. Note this is a map on cohomology and is not the i_* of theorem 4.3.1, which acts on homology. By construction $i_!(1)$ is the class dual to Y, and the class of Y in $H^2(X,\mathbb{Z})$ is the restriction of the hyperplane class, which is the Fubini-Study Kähler class (theorem 1.1.7). The projection formula for cap products ([30, Properties of the Cap Product]) then transposes to $i_!(i^*\alpha) = \alpha \smile i_!(1) = L\alpha$: the Lefschetz operator factors through restriction to a hyperplane section. Hard Lefschetz (theorem 4.2.3) makes L^{n-k} injective on $H^k(X,\mathbb{Q})$ for $k \le n$, and $L^{n-k} = (i_!i^*)^{n-k}$, so i^* is already injective on $H^k(X,\mathbb{Q})$ for $k \le n-1$ on those grounds alone.

What that argument does not give is surjectivity, nor anything over $\mathbb{Z}$; both come from the affineness of the complement.

Applied to a variety whose ambient cohomology is known, the theorem computes everything outside the middle degree.

Example 4.3.2: The Cohomology of a Smooth Hypersurface

Let $X \subseteq \mathbb{CP}^{n+1}$ be a smooth hypersurface of degree d. It is not a hyperplane section of $\mathbb{CP}^{n+1}$ unless $d = 1$, but it becomes one after a change of embedding: the d-th Veronese map $v_d \colon \mathbb{CP}^{n+1} \to \mathbb{CP}^M$ is a closed embedding whose image is smooth, and hyperplanes of $\mathbb{CP}^M$ pull back to exactly the degree-d hypersurfaces of $\mathbb{CP}^{n+1}$ [15, The Veronese Is a Closed Embedding, Hypersurfaces Become Hyperplane Sections]. So $v_d(X)$ is a smooth hyperplane section of the smooth projective variety $v_d(\mathbb{CP}^{n+1})$ of dimension $n+1$.

Theorem 4.3.1, applied with $n+1$ in place of n, gives $H^k(\mathbb{CP}^{n+1}, \mathbb{Z}) \cong H^k(X, \mathbb{Z})$ for $k \leq n-1$ and $H_k(X, \mathbb{Z}) \cong H_k(\mathbb{CP}^{n+1}, \mathbb{Z})$ for $k \leq n-1$. Hence, for $k \leq n-1$,

$$H^k(X, \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & k \text{ even} \\ 0 & k \text{ odd} \end{cases}$$

For $k \geq n+1$ Poincaré duality on the compact oriented $2n$-manifold X gives $H^k(X, \mathbb{Z}) \cong H_{2n-k}(X, \mathbb{Z})$, and $2n-k \leq n-1$, so the same table applies; note $2n-k$ and k have the same parity. Every Betti number of X outside degree n is therefore 1 in even degrees and 0 in odd degrees, independent of d.

The middle Betti number is then forced by the Euler characteristic. If n is even, the degrees $k \neq n$ contributing to $\chi_{\text{top}}(X) = \sum_k (-1)^k b_k(X)$ are the n even degrees other than n itself, each contributing 1, so $b_n(X) = \chi_{\text{top}}(X) - n$. If n is odd there are $n+1$ even degrees, all different from n, and b_n enters with a minus sign, so $b_n(X) = n + 1 - \chi_{\text{top}}(X)$.

Two instances. A smooth plane curve of degree d has $n = 1$ and $\chi_{\text{top}} = 2 - 2g$ with $g = (d-1)(d-2)/2$ [4, Degree Formula for Plane Curves], giving $b_1 = 2 - (2 - 2g) = 2g$, the classical count. A smooth quartic surface in $\mathbb{CP}^3$ is a K3 surface ([4, K3 Surfaces: A Closer Look]), so by the Hodge numbers recorded in example 2.2.4 it has $b_0 = b_4 = 1$, $b_1 = b_3 = 0$ and $b_2 = 22$, whence $\chi_{\text{top}} = 1 - 0 + 22 - 0 + 1 = 24$. Read the other way, the formula above recovers $b_2 = \chi_{\text{top}} - 2 = 22$, the rank of the K3 intersection lattice.

Everything the hyperplane section does not see is concentrated in $H^n(X)$. To reach that degree a single section is useless, since it is precisely the degree where restriction fails to be an isomorphism. The device that does reach it lets the hyperplane move.

A line $\ell \subseteq (\mathbb{CP}^N)^\vee$ is spanned by two linear forms s_0, s_1, and the hyperplanes it parametrizes are the zero sets of $\lambda_t = t_0 s_0 + t_1 s_1$ for $t = [t_0 : t_1] \in \mathbb{CP}^1$. All of them contain the codimension-2 linear subspace $A = H_{s_0} \cap H_{s_1}$, called the *axis*, and conversely every hyperplane containing A occurs. Cutting with X produces a family $X_t = X \cap H_t$ of hypersurfaces in X, all containing the fixed locus $B = X \cap A$, and sweeping out X: every point of $X \setminus B$ lies on exactly one member, and every point of B lies on all of them.

Some members of the family will be singular, and the definition below restricts how badly. A point x of a hypersurface $Z \subseteq X$ is an *ordinary double point* if Z is cut out near x by a holomorphic function g with $g(x) = 0$, $dg_x = 0$, and Hessian $\text{Hess}_x(g)$ a nondegenerate quadratic form on $T_x X$. The condition does not depend on the choice of g: two local defining functions differ by a nowhere-zero

holomorphic factor u, and at a point where g and dg both vanish one has $\mathrm{Hess}_x(ug) = u(x)\,\mathrm{Hess}_x(g)$.

Definition 4.3.3: Lefschetz Pencil

Let $X \subseteq \mathbb{C}\mathrm{P}^N$ be a smooth connected projective variety of dimension $n \geq 2$. A *pencil of hyperplane sections* of X is the family $\{X_t = X \cap H_t\}_{t \in \mathbb{C}\mathrm{P}^1}$ attached to a line $\ell \subseteq (\mathbb{C}\mathrm{P}^N)^\vee$ as above, with *axis* A and *base locus* $B = X \cap A$. The pencil is a *Lefschetz pencil* if

1. A meets X transversally, that is $T_xX + T_xA = T_x\mathbb{C}\mathrm{P}^N$ for every $x \in B$;
2. the set $\Delta \subseteq \mathbb{C}\mathrm{P}^1$ of parameters t for which X_t is singular is finite;
3. for each $t \in \Delta$ the hypersurface X_t has exactly one singular point, and that point is an ordinary double point.

The members X_t with $t \in \Delta$ are the *singular fibres* of the pencil, and $\mu = \#\Delta$ is their number.

Condition (1) is a genuine restriction, not a smoothness convention. Since A has codimension 2 and $n \geq 2$, the intersection B is nonempty [15, The Projective Dimension Theorem], and transversality forces B to be a smooth submanifold of X of codimension 2; it also rules out the degenerate possibility that $X \subseteq H_t$ for some t, since then T_xX and T_xA would both lie in the hyperplane T_xH_t at a point $x \in B$. Condition (3) is what makes the family tractable: an ordinary double point has a normal form, derived below, and every singular member of a Lefschetz pencil looks the same near its singular point.

Before showing that such pencils exist, here is one written out completely.

Example 4.3.4: A Pencil of Conics on the Plane

Take $X = \mathbb{C}\mathrm{P}^2$ and embed it by the complete system of conics, that is by the Veronese map $v_2 \colon \mathbb{C}\mathrm{P}^2 \to \mathbb{C}\mathrm{P}^5$ [15, The Veronese Is a Closed Embedding, Hypersurfaces Become Hyperplane Sections]. Hyperplane sections of $v_2(\mathbb{C}\mathrm{P}^2)$ are the conics in $\mathbb{C}\mathrm{P}^2$, and a pencil of hyperplane sections is a pencil of conics. Take the one spanned by

$$F_0 = x_0^2 - x_1^2, \qquad F_1 = x_0^2 - x_2^2$$

so that $F_t = t_0F_0 + t_1F_1$ and $X_t = \{F_t = 0\}$.

The base locus. A point lies on every member exactly when $x_0^2 = x_1^2$ and $x_0^2 = x_2^2$. If $x_0 = 0$ then $x_1 = x_2 = 0$, which is not a point of $\mathbb{C}\mathrm{P}^2$, so $x_0 \neq 0$ and, normalizing $x_0 = 1$, the solutions are the four points $[1 : \pm 1 : \pm 1]$. The four are distinct, and at each of them the differentials of F_0 and F_1 are independent: at $[1 : 1 : 1]$, for instance, in the affine chart $x_0 = 1$ they are $-2\,dx_1$ and $-2\,dx_2$. That independence is exactly transversality of the axis with X, so condition (1) holds and B consists of four points.

The singular members. Writing F_t as $x^\top M_t x$ with

$$M_t = \begin{pmatrix} t_0 + t_1 & 0 & 0 \\ 0 & -t_0 & 0 \\ 0 & 0 & -t_1 \end{pmatrix}$$

the conic X_t is singular exactly when $\det M_t = (t_0 + t_1)t_0t_1$ vanishes, that is at the three parameters $[1 : 0]$, $[0 : 1]$ and $[1 : -1]$. So $\mu = 3$ and condition (2) holds.

At $t = [1 : 0]$ the member is $x_0^2 - x_1^2 = (x_0 - x_1)(x_0 + x_1)$, a pair of distinct lines meeting at $[0 : 0 : 1]$. In the affine chart $x_2 = 1$ with coordinates (y_0, y_1) the defining function is $y_0^2 - y_1^2$, whose Hessian $\mathrm{diag}(2, -2)$ is nondegenerate: an ordinary double point, and the only singular point of that member. The point $[0 : 0 : 1]$ is not one of the four base points, as it must not be. The parameters $[0 : 1]$ and $[1 : -1]$ behave identically, the latter giving $x_1^2 - x_2^2$ with singular point $[1 : 0 : 0]$. So condition (3) holds and the pencil is a Lefschetz pencil.

The associated fibration. Blowing up the four base points turns the rational map $\mathbb{CP}^2 \dashrightarrow \mathbb{CP}^1$, $x \mapsto [F_1(x) : -F_0(x)]$, into a morphism whose fibres are the members of the pencil; this is the content of proposition 4.3.6 below. The blow-up has Euler characteristic $3 + 4 = 7$, since each of the four points is replaced by a $\mathbb{CP}^1$. On the other side, three of the fibres are pairs of lines ($\chi_{\mathrm{top}} = 2 + 2 - 1 = 3$) and the rest are smooth conics ($\chi_{\mathrm{top}} = 2$), and a count of the fibres over $\mathbb{CP}^1$ gives $(2 - 3) \cdot 2 + 3 \cdot 3 = 7$. The two computations agree.

That pencil was arranged by hand, but it exhibits the general mechanism: the members singular at a prescribed point form a linear subsystem of large codimension, and a line in the parameter space meets the union of these subsystems in a controlled way. Turning that into a proof requires knowing that the linear system is large enough to impose independent conditions on second derivatives at a point and on first derivatives at two points at once.

Proposition 4.3.5: Existence of Lefschetz Pencils

Let $X \subseteq \mathbb{CP}^N$ be a smooth connected projective variety of dimension $n \geq 2$, and let V be the space of linear forms on $\mathbb{C}^{N+1}$. Assume

1. for every $x \in X$ the restrictions to X of the forms in V realize every 2-jet at x;
2. for every pair of distinct points $x, x' \in X$ they realize every pair of 1-jets at x and x'.

Then the lines $\ell \subseteq (\mathbb{CP}^N)^\vee$ whose pencil is a Lefschetz pencil form a dense open subset of the Grassmannian of lines. Moreover every smooth projective variety satisfies (1) and (2) after re-embedding by $|\mathcal{O}_X(d)|$ for d large [4, Lefschetz-Type Jet Conditions Hold after Re-embedding], so every smooth projective variety carries Lefschetz pencils.

Here the 2-jet of a section at x records its value, its first derivatives along X and its second derivatives along X, so the space J_x^2 of 2-jets has dimension $1 + n + \binom{n+1}{2}$ and the space J_x^1 of 1-jets has dimension $n + 1$; both are jets of sections of $\mathcal{O}_X(1)$, so the statements are independent of local trivializations.

***Proof*:**

Write $\mathbb{G}$ for the Grassmannian of lines in $(\mathbb{CP}^N)^\vee$, an irreducible projective variety of dimension $2(N - 1)$. For each of the three conditions the locus of lines where it fails is closed, so it suffices to show each failure locus is a proper subset; the intersection of three dense open subsets of an irreducible variety is dense open.

Step 1: the variety of tangent hyperplanes. For $x \in X$ let $\mathbb{T}_x X \subseteq \mathbb{CP}^N$ be the embedded projective tangent space, a linear subspace of dimension n. A hyperplane H makes $X \cap H$ singular at x exactly when $\mathbb{T}_x X \subseteq H$: the section $X \cap H$ is cut out near x by the linear form defining H, which is singular at x precisely when the form and its differential along X both vanish there. Put

$$\tilde{Y} = \left\{ (x, H) \in X \times (\mathbb{CP}^N)^\vee \;\middle|\; \mathbb{T}_x X \subseteq H \right\}$$

The hyperplanes containing a fixed linear subspace of dimension n form a linear $\mathbb{C}\mathrm{P}^{N-n-1}$ inside $(\mathbb{C}\mathrm{P}^N)^\vee$, so the fibre of $\tilde{Y}$ over x is such a subspace, and $\tilde{Y}$ is a projective bundle over X, hence smooth and irreducible of dimension $n+(N-n-1) = N-1$. Its image $X^\vee \subseteq (\mathbb{C}\mathrm{P}^N)^\vee$ under the second projection is closed, being the image of a projective variety, irreducible, and of dimension at most $N-1$. In particular $X^\vee \neq (\mathbb{C}\mathrm{P}^N)^\vee$, so a hyperplane section of X is generically smooth. Only smoothness of X entered, so the same conclusion holds for any smooth projective variety in $\mathbb{C}\mathrm{P}^N$.

Step 2: condition (2) of definition 4.3.3. The singular parameters of the pencil attached to ℓ lie in $\ell \cap X^\vee$, a closed subvariety of the line ℓ, hence either finite or all of ℓ. Lines contained in the proper closed subset $X^\vee$ form a closed subset of $\mathbb{G}$, and it is proper: any line joining two points outside $X^\vee$ is not contained in it, and such points exist by Step 1.

Step 3: condition (3). Fix $(x, H) \in \tilde{Y}$ and let s be a linear form defining H. Since s and its differential along X vanish at x, its Hessian along X is a well-defined quadratic form on T_xX, and x is an ordinary double point of $X \cap H$ exactly when that form is nondegenerate. Hypothesis (1) says the map sending a form vanishing to first order at x to its Hessian, from an $(N-n)$-dimensional space onto $\mathrm{Sym}^2 T_x^* X$, is surjective. Degenerate quadratic forms are the zero locus of the determinant, a nonzero polynomial, hence a proper closed subset; its preimage is therefore a proper closed subset of the fibre of $\tilde{Y}$ over x, of dimension at most $N-n-2$. So

$$\tilde{Y}_{\mathrm{deg}} = \left\{(x, H) \in \tilde{Y} \;\middle|\; \mathrm{Hess}_x(s) \text{ is degenerate on } T_xX\right\}$$

is closed of dimension at most $n + (N-n-2) = N-2$.

For the second half of condition (3), let

$$\tilde{Y}_2 = \left\{(x, x', H) \;\middle|\; x \neq x',\ (x, H) \in \tilde{Y},\ (x', H) \in \tilde{Y}\right\}$$

Hypothesis (2) makes the $2(n+1)$ linear conditions defining the fibre over a pair (x, x') independent, so that fibre has dimension $N-2n-2$ when nonnegative and is empty otherwise. Hence $\tilde{Y}_2$ has dimension at most $2n + (N-2n-2) = N-2$.

Let Σ be the union of the image of $\tilde{Y}_{\mathrm{deg}}$ in $(\mathbb{C}\mathrm{P}^N)^\vee$ and the closure of the image of $\tilde{Y}_2$. It is closed of dimension at most $N-2$, and a pencil fails condition (3) exactly when ℓ meets Σ. The incidence variety $\{(\ell, H) \big| H \in \ell \cap \Sigma\}$ fibres over Σ with fibre the lines through a fixed point, a family of dimension $N-1$, so it has dimension at most $(N-2)+(N-1) = 2N-3$. Its image in $\mathbb{G}$ is closed of dimension at most $2N-3 < 2(N-1)$, hence proper.

Step 4: condition (1). The set of pairs (x, ℓ) with $x \in X$ lying on the axis A_ℓ and $T_xX + T_xA_\ell \neq T_x\mathbb{C}\mathrm{P}^N$ is closed in $X \times \mathbb{G}$, and X is complete, so its image in $\mathbb{G}$ is closed. It is proper: choose $H_0 \notin X^\vee$ by Step 1, so that $X_0 = X \cap H_0$ is smooth of dimension $n-1$, then apply Step 1 to X_0 to choose H_1 with $X_0 \cap H_1$ smooth of dimension $n-2$. The line through H_0 and H_1 has axis $A = H_0 \cap H_1$ with $X \cap A = X_0 \cap H_1$ smooth of codimension 2, cut out in X by two forms with independent differentials, which is transversality.

The three failure loci are proper closed subsets of the irreducible variety $\mathbb{G}$, so their complements intersect in a dense open subset, as we sought to show.

The proof also counts the singular members. When $X^\vee$ is a hypersurface, a generic line meets it transversally in $\deg X^\vee$ points [4], so $\mu = \deg X^\vee$ for a generic pencil; when $X^\vee$ has codimension at least 2, a generic line misses it and $\mu = 0$, the pencil being a family of smooth hypersurfaces. The second case is not exotic: a linearly embedded $\mathbb{C}\mathrm{P}^n \subseteq \mathbb{C}\mathrm{P}^N$ has only smooth hyperplane sections

(exercise 4.3.1 (3)).

A pencil is not yet a family in the sense that admits topological arguments, because the members all meet along B and the assignment $x \mapsto t$ is undefined there. Separating the members along B is exactly what a blow-up does.

Proposition 4.3.6: The Fibration Attached to a Pencil

Let $\{X_t\}$ be a Lefschetz pencil on $X \subseteq \mathbb{C}\mathrm{P}^N$, with axis A, base locus B, defining forms s_0, s_1 and singular parameters $\Delta = \{t_1, \dots, t_\mu\}$. Put

$$\tilde{X} = \big\{(x,t) \in X \times \mathbb{C}\mathrm{P}^1 \;\big|\; t_0 s_0(x) + t_1 s_1(x) = 0\big\}$$

and let $\sigma\colon \tilde{X} \to X$ and $f\colon \tilde{X} \to \mathbb{C}\mathrm{P}^1$ be the two projections. Then

1. $\tilde{X}$ is a smooth connected projective variety of dimension n, and σ is the blow-up of X along B, an isomorphism over $X \setminus B$ with exceptional divisor $E = \sigma^{-1}(B) = B \times \mathbb{C}\mathrm{P}^1$;
2. f is a proper holomorphic map with $f^{-1}(t) = X_t \times \{t\}$, and $f|_E$ is the projection $B \times \mathbb{C}\mathrm{P}^1 \to \mathbb{C}\mathrm{P}^1$;
3. the critical points of f are exactly the μ singular points of the singular fibres; each is a nondegenerate critical point, none lies on E, and no two lie in the same fibre;
4. over $\mathbb{C}\mathrm{P}^1 \setminus \Delta$ the map f is a locally trivial fibration with fibre a smooth hyperplane section of X.

***Proof*:**

1. The defining equation is bihomogeneous of degree $(1,1)$, so $\tilde{X}$ is a closed subvariety of $X \times \mathbb{C}\mathrm{P}^1$, hence compact and projective [4, Products of Projective Varieties]. Let $(x,t) \in \tilde{X}$ and write $F(x,t) = t_0 s_0(x) + t_1 s_1(x)$ for the defining section.

   If $x \notin B$ then $(s_0(x), s_1(x)) \neq (0,0)$, and the derivative of F in the $\mathbb{C}\mathrm{P}^1$ directions is $(s_0(x), s_1(x))$ in a local trivialization, so $dF \neq 0$ and $\tilde{X}$ is smooth of dimension n at (x,t). If $x \in B$ then $s_0(x) = s_1(x) = 0$, the differentials ds_0, ds_1 restricted to $T_x X$ are independent by transversality of A with X, and the derivative of F in the X directions is $t_0\, ds_0 + t_1\, ds_1$, nonzero because $(t_0, t_1) \neq (0,0)$. Again $\tilde{X}$ is smooth of dimension n at (x,t).

   Over $x \notin B$ the equation determines t uniquely, namely $t = [s_1(x) : -s_0(x)]$, so σ is bijective there; the inverse $x \mapsto (x, [s_1(x) : -s_0(x)])$ is holomorphic, so σ is an isomorphism over $X \setminus B$. Over $x \in B$ the equation is vacuous, so $\sigma^{-1}(B) = B \times \mathbb{C}\mathrm{P}^1$, of dimension $(n-2)+1 = n-1$. Since $\tilde{X}$ is smooth of pure dimension n, no component of $\tilde{X}$ lies over B, so $\tilde{X}$ is the closure of the graph of $x \mapsto [s_1(x) : -s_0(x)]$; as $X \setminus B$ is connected, $\tilde{X}$ is connected. Composing with the automorphism $[u_0 : u_1] \mapsto [u_1 : -u_0]$ of $\mathbb{C}\mathrm{P}^1$ identifies that graph closure with the closure of the graph of $x \mapsto [s_0(x) : s_1(x)]$, which is the blow-up of X along the smooth centre B cut out by s_0 and s_1 [4, The Blow-up as the Closure of a Graph]. For $n = 2$ the centre is a finite set of points and this is literally the blow-up of a complex manifold at a point, whose local picture is the blow-up of $\mathbb{C}^2$ at the origin.[a]

2. Properness is compactness of $\tilde{X}$. For fixed t the equation cuts out $X \cap H_t = X_t$, so

$f^{-1}(t) = X_t \times \{t\}$, and f restricted to $E = B \times \mathbb{CP}^1$ is the second projection by construction.

3. Since $\tilde{X}$ has dimension n and f maps to a curve, a point is critical for f exactly when df vanishes there, which happens exactly when the fibre $f^{-1}(t) = X_t$ is singular at that point.

   No such point lies on E: for $x \in B$ the differential of the defining form $\lambda_t = t_0 s_0 + t_1 s_1$ along $T_x X$ is $t_0\, ds_0 + t_1\, ds_1 \neq 0$, so X_t is smooth at every point of B. By condition (3) of definition 4.3.3 each singular fibre has exactly one singular point, so the critical points are the μ ordinary double points, one in each singular fibre, and no two share a fibre.

   For nondegeneracy fix a critical point p, lying over the singular parameter t_i, and normalize $t_i = [a_0 : 1]$ (after exchanging s_0, s_1 if necessary). Since $p \notin B$ and $\lambda_{t_i}(p) = 0$, one has $s_0(p) \neq 0$. Let ζ be the affine coordinate on $\mathbb{CP}^1$ centred at t_i for which $t = [a_0 + \zeta : 1]$. Near p the map σ is an isomorphism, and the equation $(a_0 + \zeta)s_0 + s_1 = 0$ of $\tilde{X}$ reads $\zeta = -g$ with

   $$g = \frac{\lambda_{t_i}}{s_0}$$

   a holomorphic function on a neighbourhood of p in X. So f corresponds to $-g$, and X_{t_i} is cut out near p by g. Both g and dg vanish at p, and p is a nondegenerate critical point of f exactly when $\mathrm{Hess}_p(g)$ is nondegenerate, which is the statement that p is an ordinary double point of X_{t_i}.

4. Over $\mathbb{CP}^1 \setminus \Delta$ the map f is a proper holomorphic submersion between smooth manifolds, hence a locally trivial fibration [32, Ehresmann's Theorem] whose fibres are the smooth members X_t, completing the proof.

[a] [5, Blow-Up of a Complex Manifold at a Point, Blow-Up of $\mathbb{C}^2$ at the Origin].

The blow-up changes nothing that matters here. It changes X only along B, replacing each point of B by a $\mathbb{CP}^1$, and σ is an isomorphism elsewhere; in exchange, the pencil becomes a genuine map to $\mathbb{CP}^1$, and the topology of X can be studied by moving in the base. What now needs description is what happens over a point of Δ, and by part (3) that question is local and identical at every one of the μ critical points.

The description is a holomorphic analogue of the Morse lemma. The real statement, that a nondegenerate critical point admits coordinates in which the function is a sum of signed squares, is proved in [32, Morse Lemma]; over $\mathbb{C}$ the signs disappear, since every nonzero complex number has a square root, and with them the index. That is why a complex nondegenerate critical point has no discrete invariant and every node looks like every other. The proof is short enough to give directly, and is not the real lemma in disguise: it is a completing-the-square induction performed with holomorphic square roots.

Lemma 4.3.7: The Local Model at a Node

Let M be a complex manifold of dimension $n \geq 1$, let $f\colon M \to \mathbb{C}$ be holomorphic, and let p be a nondegenerate critical point of f with $f(p) = 0$.

1. There are local holomorphic coordinates $z = (z_1, \dots, z_n)$ centred at p in which
$$f = z_1^2 + \cdots + z_n^2$$
2. In these coordinates fix $\epsilon > 0$ with the closed ball $\bar{B}_\epsilon$ contained in the coordinate domain, and for $0 < |t| < \epsilon^2$ set $F_t = \{z \in \bar{B}_\epsilon \,|\, f(z) = t\}$. Then F_t is diffeomorphic to a closed disc bundle of the tangent bundle TS^{n-1}, with the sphere
$$S_t = \left\{ \sqrt{t}\, x \;\middle|\; x \in \mathbb{R}^n,\ |x| = 1 \right\}$$
as the zero section, for either choice of square root. In particular F_t deformation retracts onto S_t, a smooth $(n-1)$-sphere of Euclidean radius $|t|^{1/2}$.
3. $F_0 = \{z \in \bar{B}_\epsilon \,|\, f(z) = 0\}$ is contractible.

***Proof*:**

1. Choose holomorphic coordinates w centred at p on a polydisc, in which $f(0) = 0$ and $df_0 = 0$. Since the polydisc is convex, Taylor's formula with integral remainder gives
$$f(w) = \sum_{i,j} h_{ij}(w)\, w_i w_j, \qquad h_{ij}(w) = \int_0^1 (1-s)\, \partial_i \partial_j f(sw)\, ds$$
The functions $h_{ij} = h_{ji}$ are holomorphic, being integrals of holomorphic families, and $h_{ij}(0) = \frac{1}{2}\partial_i \partial_j f(0)$, so the matrix $h(0)$ is invertible.
The claim, proved by induction on k, is that for each $0 \leq k \leq n$ there are holomorphic coordinates u centred at p with
$$f = u_1^2 + \cdots + u_k^2 + \sum_{i,j>k} c_{ij}(u)\, u_i u_j$$
where $c_{ij} = c_{ji}$ is holomorphic and the matrix $(c_{ij}(0))_{i,j>k}$ is invertible. The case $k = 0$ is the display above. Assume the statement for $k < n$. A nonzero symmetric bilinear form over $\mathbb{C}$ has a nonisotropic vector, so after a constant linear change of the coordinates $u_{k+1}, \dots, u_n$, which does not disturb the first k squares, one may assume $c_{k+1,k+1}(0) \neq 0$. Shrinking the neighbourhood so that $c_{k+1,k+1}$ takes values in a disc avoiding the origin, choose a holomorphic branch ρ of $\sqrt{c_{k+1,k+1}}$ there and set
$$v_{k+1} = \rho(u)\Big(u_{k+1} + \sum_{j>k+1} \frac{c_{k+1,j}(u)}{c_{k+1,k+1}(u)}\, u_j \Big)$$
Completing the square in u_{k+1} gives
$$\sum_{i,j>k} c_{ij} u_i u_j = v_{k+1}^2 + \sum_{i,j>k+1} c'_{ij}(u)\, u_i u_j, \qquad c'_{ij} = c_{ij} - \frac{c_{k+1,i}\, c_{k+1,j}}{c_{k+1,k+1}}$$

with c' symmetric holomorphic, and $\det c(0) = c_{k+1,k+1}(0)\det c'(0)$ shows $c'(0)$ is invertible. The map $u \mapsto (u_1, \dots, u_k, v_{k+1}, u_{k+2}, \dots, u_n)$ has Jacobian at the origin equal to the identity except in row $k+1$, whose diagonal entry is $\rho(0) \neq 0$; the determinant is $\rho(0)$, so the map is a local biholomorphism and the new functions form a coordinate system. Expressing c' in the new coordinates completes the inductive step, and $k = n$ is the assertion.

2. Replacing z by cz with $|c| = 1$ multiplies f by c^2 and preserves $\bar{B}_\epsilon$, so it suffices to treat $t > 0$ real. Write $z = x + iy$ with $x, y \in \mathbb{R}^n$. Then

$$\sum_j z_j^2 = |x|^2 - |y|^2 + 2i\, x \cdot y$$

so F_t is the set of pairs (x, y) with $|x|^2 = t + |y|^2$, $x \cdot y = 0$ and $|x|^2 + |y|^2 \leq \epsilon^2$, the last condition being $2|y|^2 \leq \epsilon^2 - t$. Writing $x = |x|\,\nu$ with $\nu \in S^{n-1}$, the map

$$(\nu, y) \longmapsto \big(\sqrt{t + |y|^2}\ \nu,\ y\big)$$

is a diffeomorphism from $\{(\nu, y) \big| \nu \in S^{n-1},\ y \perp \nu,\ 2|y|^2 \leq \epsilon^2 - t\}$ onto F_t. Identifying $T_\nu S^{n-1}$ with $\nu^\perp$, the source is the closed disc bundle of radius $\sqrt{(\epsilon^2 - t)/2}$ in TS^{n-1}, and $y = 0$ corresponds to $x = \sqrt{t}\,\nu$, that is to S_t. Scaling y to zero deformation retracts the disc bundle onto its zero section.

3. The set F_0 is preserved by $z \mapsto \lambda z$ for real $\lambda \in [0, 1]$, since that scales f by λ^2 and does not increase $|z|$. The map $(z, \lambda) \mapsto \lambda z$ is a continuous contraction of F_0 to the origin, as required.

Part (2) is the geometric content of the whole construction. As t approaches a singular parameter the nearby fibre carries a sphere of real dimension $n-1$, half the real dimension $2n-2$ of the fibre, and that sphere contracts to the node. Since the fibres over the complement of Δ are identified with one another by the fibration of proposition 4.3.6, the sphere can be carried back to a fixed fibre, where it becomes a homology class.

Definition 4.3.8: Vanishing Cycle

Let $\{X_t\}$ be a Lefschetz pencil on X, with associated fibration $f\colon \tilde{X} \to \mathbb{CP}^1$, singular parameters $\Delta = \{t_1, \dots, t_\mu\}$ and critical points $p_1, \dots, p_\mu$. Fix a base parameter $t_* \in \mathbb{CP}^1 \setminus \Delta$. Fix i, and let z be coordinates near p_i as in lemma 4.3.7, taken with respect to a holomorphic coordinate on $\mathbb{CP}^1$ centred at t_i, so that $f = t_i + \sum_j z_j^2$ near p_i. Let γ be a path in $\mathbb{CP}^1 \setminus \Delta$ from t_* to a parameter t' close enough to t_i that the sphere $S_{t'-t_i}$ of lemma 4.3.7 lies in $f^{-1}(t')$, and orient that sphere. The *vanishing cycle* attached to p_i and γ is the class

$$\delta_i \in H_{n-1}\big(f^{-1}(t_*), \mathbb{Z}\big) = H_{n-1}(X_{t_*}, \mathbb{Z})$$

obtained from the fundamental class of that sphere by the isomorphism that parallel transport along γ induces on the homology of the fibres. The *Lefschetz thimble* attached to p_i and t' is the n-disc in $\tilde{X}$ swept out by the spheres S_s as s runs along a segment from 0 to $t' - t_i$; its boundary is the sphere above.

The class δ_i is well defined once the orientation of the sphere is fixed, and changing that orientation changes its sign. It depends on γ only through the homotopy class of γ rel endpoints, since homotopic paths induce the same map on the homology of the fibres, and it is independent of the choice of normal-form coordinates up to sign: the local fibre retracts onto the sphere, so its middle homology is infinite cyclic with the sphere class as a generator, and any other choice produces a generator of the same group. The dependence on γ is genuine and is the subject of section 4.4: different paths to the same critical value produce different classes, and comparing them is what the monodromy of the family records.

Two features of δ_i follow immediately from the local model. First, the sphere bounds in $\tilde{X}$: the thimble of definition 4.3.8 is, in the local coordinates with $t' - t_i > 0$, the real ball $\{y \in \mathbb{R}^n \big| |y|^2 \leq t' - t_i\}$, an embedded n-disc whose boundary is the vanishing sphere. So δ_i maps to zero in $H_{n-1}(\tilde{X}, \mathbb{Z})$: a class that is nonzero on a fibre may die in the total space, and the vanishing cycles are the visible source of such classes. Second, a tubular neighbourhood of the sphere inside the fibre is the disc bundle of TS^{n-1}, so the self-intersection of δ_i is the Euler number of the tangent bundle of a sphere, up to orientation: it is zero when n is even and has absolute value 2 when n is odd (exercise 4.3.1 (4)).

The word "vanishing" is literal: as the parameter approaches t_i the sphere shrinks to the node and disappears from the singular fibre. The degeneration of a cubic curve shows the picture.

Example 4.3.9: A Pencil of Cubics and the Rational Elliptic Surface

Take $X = \mathbb{C}\mathrm{P}^2$ embedded by the complete system of cubics, $v_3 \colon \mathbb{C}\mathrm{P}^2 \to \mathbb{C}\mathrm{P}^9$, and let $\{X_t\}$ be a generic pencil of plane cubics, which is a Lefschetz pencil by proposition 4.3.5. Two distinct cubics meet in 9 points counted with multiplicity [15, Bézout's Theorem], and for a generic pencil these are 9 distinct points at which the two curves meet transversally, so B consists of 9 points and $\tilde{X}$ is $\mathbb{C}\mathrm{P}^2$ blown up at them.

The smooth members are smooth plane cubics, hence elliptic curves, with $\chi_{\mathrm{top}} = 0$. The singular members have a single node; such a curve is the image of $\mathbb{C}\mathrm{P}^1$ with two points identified, so $\chi_{\mathrm{top}} = 2 - 1 = 1$, one more than the smooth value, in agreement with the general relation of exercise 4.3.1 (2) at $n = 2$.

Counting Euler characteristics both ways determines μ. On one side $\chi_{\mathrm{top}}(\tilde{X}) = 3 + 9 = 12$, each blown-up point contributing a $\mathbb{C}\mathrm{P}^1$ in place of a point. On the other side, the fibration over $\mathbb{C}\mathrm{P}^1$ has $2-\mu$ parameters carrying a smooth fibre and μ carrying a nodal one, so $\chi_{\mathrm{top}}(\tilde{X}) = (2-\mu)\cdot 0 + \mu \cdot 1 = \mu$. Hence $\mu = 12$: a generic pencil of plane cubics has exactly twelve nodal members, which is the classical statement that the discriminant hypersurface of plane cubics has degree 12.

Here $n = 2$, so a vanishing cycle is a class in $H_1(X_{t_*}, \mathbb{Z}) \cong \mathbb{Z}^2$ of the elliptic curve X_{t_*}, represented by an embedded circle that pinches to the node as $t \to t_i$. That circle is nonseparating, and what forces this is the irreducibility of the degenerate member. A reducible plane cubic has either more than one singular point or a singular point that is not an ordinary double point, so condition (3) of definition 4.3.3 leaves X_{t_i} irreducible, hence the image of $\mathbb{C}\mathrm{P}^1$ with two points identified, a single sphere glued to itself. A separating circle on a torus bounds a disc, so collapsing one would instead exhibit the degenerate member as two closed surfaces meeting at a point. Hence $\delta_i \neq 0$; its self-intersection is zero, as it must be, the intersection form on the first homology of a curve being alternating.

Example 4.3.4 sits at the opposite extreme. There the fibres were rational curves with $H_1 = 0$, so every vanishing cycle is zero even though the pencil has three singular members: a degeneration always produces a vanishing sphere in the nearby fibre, but that sphere need not carry a nonzero

homology class. What survives in general is the sphere itself and the reflection it determines, which is what section 4.4 computes.

Exercise 4.3.1

1. Let $S \subseteq \mathbb{C}\mathrm{P}^3$ be a smooth surface of degree d. Using theorem 4.3.1 in the form applied in example 4.3.2, show that S is connected, that $H^1(S,\mathbb{Z}) = 0$ and $H^3(S,\mathbb{Z}) = 0$, and deduce that $b_2(S) = \chi_{\mathrm{top}}(S) - 2$. Then use theorem 2.1.1 and corollary 2.2.1 to conclude $h^{1,0}(S) = 0$.

2. Let $\{X_t\}$ be a Lefschetz pencil on X with μ singular fibres, smooth member X_{t_*} and base locus B, and let $f\colon \tilde{X} \to \mathbb{C}\mathrm{P}^1$ be as in proposition 4.3.6.

    1. Using lemma 4.3.7 and the Mayer-Vietoris sequence, show that $\chi_{\mathrm{top}}(X_{t_i}) = \chi_{\mathrm{top}}(X_{t_*}) + (-1)^n$ for each singular parameter t_i. (The boundaries of the two local pieces are closed manifolds of odd real dimension $2n-3$, whose Euler characteristics vanish by Poincaré duality; the parts of the two fibres outside the local pieces are diffeomorphic.)
    2. Deduce $\chi_{\mathrm{top}}(\tilde{X}) = 2\chi_{\mathrm{top}}(X_{t_*}) + (-1)^n\mu$ and then $\chi_{\mathrm{top}}(X) = 2\chi_{\mathrm{top}}(X_{t_*}) + (-1)^n\mu - \chi_{\mathrm{top}}(B)$. (Cover $\mathbb{C}\mathrm{P}^1$ by small closed discs D_i around the singular parameters and the closure C of the complement, and use the two facts that Euler characteristic is multiplicative in a fibre bundle and that $f^{-1}(D_i)$ deformation retracts onto X_{t_i}.)
    3. Check both formulas against example 4.3.4 and example 4.3.9.

3. Let $X = \mathbb{C}\mathrm{P}^n \subseteq \mathbb{C}\mathrm{P}^N$ be a linearly embedded projective space with $2 \leq n < N$. Identify the variety $X^\vee$ of Step 1 in the proof of proposition 4.3.5 and compute its dimension. Conclude that a generic pencil of hyperplane sections of X is a Lefschetz pencil with $\mu = 0$, describe $\tilde{X}$ and f explicitly, and verify the Euler characteristic formula of exercise 4.3.1 (2) in this case.

4. Let $\delta \in H_{n-1}(X_{t_*},\mathbb{Z})$ be a vanishing cycle of a Lefschetz pencil on a variety of dimension $n \geq 2$.

    1. Using lemma 4.3.7, show that a tubular neighbourhood of the vanishing sphere inside the fibre is the disc bundle of TS^{n-1}, and deduce that the self-intersection number $\delta \cdot \delta$ equals $\pm\chi_{\mathrm{top}}(S^{n-1})$.
    2. Conclude that $\delta \cdot \delta = 0$ when n is even and $|\delta \cdot \delta| = 2$ when n is odd.
    3. Interpret part (2) for $n = 2$ against the intersection form on the first homology of a compact Riemann surface, and for $n = 3$ against the intersection form of a K3 surface as recorded in example 2.2.4.

5. Work in the local coordinates of lemma 4.3.7 at a critical point p_i, with $t' - t_i$ real and positive.

    1. Show that the set of points $\sqrt{s}\,x$ with $0 \leq s \leq t' - t_i$ and $x \in S^{n-1}$ is the closed real ball of radius $(t'-t_i)^{1/2}$ in $\mathbb{R}^n$, that f maps it onto the segment $[t_i, t']$, and that its boundary is the vanishing sphere.
    2. Deduce that every vanishing cycle maps to 0 in $H_{n-1}(\tilde{X},\mathbb{Z})$.
    3. In example 4.3.4 show that all three vanishing cycles are already zero in $H_1(X_{t_*},\mathbb{Z})$, and identify the circle in a smooth conic that pinches at $t = [1:0]$.

4.4 Monodromy and the Picard-Lefschetz Formula

A Lefschetz pencil presents a smooth projective variety X of dimension n as a family of hyperplane sections over $\mathbb{CP}^1$ with finitely many singular members (definition 4.3.3, proposition 4.3.6). Over the complement of the critical values the family is a fibre bundle, so a loop in that complement carries the cohomology of a fixed smooth member back to itself. This section makes the resulting action precise and computes it for a loop encircling one critical value.

The computation is local. Everything that happens as the loop is traversed is concentrated in a small ball around the double point of the singular fibre, and inside that ball the map is $\sum z_j^2$ up to a change of coordinates. The answer is that the action moves a class only in the direction of the vanishing cycle (definition 4.3.8), by an amount read off from a single intersection number.

4.4.1 Convention: Orientations, Loops, and the Intersection Pairing Every complex manifold carries its complex orientation. On the middle homology $H_{n-1}(Y,\mathbb{Z})$ of a compact oriented $2(n-1)$-manifold Y the symbol $\langle -,-\rangle$ denotes the intersection pairing, which satisfies $\langle a,b\rangle = (-1)^{n-1}\langle b,a\rangle$ and is unimodular modulo torsion by Poincaré-Lefschetz duality [30, integral unimodularity of the intersection pairing]. Loops in the base are traversed counterclockwise unless stated otherwise, and the product $\gamma\gamma'$ in π_1 means "traverse γ' first, then γ", so that transport along a product is the composite of the transports in the same order.

The bundle structure over the complement of the critical values is what makes transport possible, and it comes from a single fact about proper submersions.

Definition 4.4.2: Monodromy Representation

Let S be a connected complex manifold, Y a complex manifold, and $f\colon Y \to S$ a proper holomorphic submersion, with fibres $Y_s = f^{-1}(s)$. Fix $s_0 \in S$. For a loop γ at s_0, the pullback of Y along γ is a fibre bundle over $[0,1]$ and hence trivial, and a trivialization determines a diffeomorphism $h_\gamma\colon Y_{s_0} \to Y_{s_0}$, the *geometric monodromy* of γ, well defined up to isotopy. The *monodromy representation* in degree k is

$$\rho_k\colon \pi_1(S,s_0) \to \mathrm{Aut}\big(H^k(Y_{s_0},\mathbb{Z})\big), \qquad \rho_k(\gamma) = \big(h_\gamma^{-1}\big)^*$$

and its image Γ_k is the *monodromy group*. If the fibres have complex dimension d, Poincaré duality identifies $H_k(Y_{s_0},\mathbb{Z})$ with $H^{2d-k}(Y_{s_0},\mathbb{Z})$ and carries the homological monodromy $\gamma \mapsto (h_\gamma)_*$ on $H_k(Y_{s_0},\mathbb{Z})$ to ρ_{2d-k}, the two indices agreeing exactly in the middle degree $k=d$. The homological action is written ρ as well when the group it acts on is named.
For a Lefschetz pencil with associated fibration $f\colon \tilde{X} \to \mathbb{CP}^1$ (proposition 4.3.6) and critical values $t_1,\dots,t_\mu$ (definition 4.3.3), the monodromy representation of the pencil is the one attached to the restriction of f over $U = \mathbb{CP}^1 \setminus \{t_1,\dots,t_\mu\}$.

Three points make the definition legitimate. A proper submersion is a locally trivial fibration [32, Ehresmann's theorem], so the pullback along γ is a bundle over an interval and any two trivializations differ by a path of diffeomorphisms starting at the identity; the induced map on homology is therefore independent of the choice. Homotopic loops give isotopic diffeomorphisms, so ρ_k descends to π_1. And the composition convention of box 4.4.1 makes $h_{\gamma\gamma'} = h_\gamma \circ h_{\gamma'}$, so ρ_k is a homomorphism rather than an anti-homomorphism.

The smallest case already shows the mechanism, and it is the one where the vanishing cycle is a pair of points rather than a sphere.

Example 4.4.3: Monodromy of a Double Cover

Let $f\colon \mathbb{C} \to \mathbb{C}$ be $f(z) = z^2$. It is proper, since $|f(z)| = |z|^2$, and it is a submersion away from the origin, so its restriction over $S = \mathbb{C}^*$ is a proper holomorphic submersion with fibre $F_t = \{\pm\sqrt{t}\}$, two points. Take $s_0 = 1$, so $F_1 = \{1, -1\}$.

Along the loop $\gamma(\vartheta) = e^{2\pi i\vartheta}$ the trivialization is given by continuing the square root: the point $1 \in F_1$ travels as $e^{i\pi\vartheta}$ and arrives at $e^{i\pi} = -1$. So h_γ exchanges the two points of F_1, and $\rho_0(\gamma)$ acts on $H^0(F_1, \mathbb{Z}) = \mathbb{Z}^2$ by swapping the coordinates.

On the reduced group $\tilde{H}_0(F_1, \mathbb{Z}) = \mathbb{Z}\,\delta$ with $\delta = [1] - [-1]$ the action is $\rho(\gamma)\delta = -\delta$, and the monodromy group is $\{\pm 1\}$. The class δ is the vanishing cycle of the critical point at the origin: as $t \to 0$ the two points of F_t collide, and δ is the difference of the two branches, the zero-sphere S^0 shrinking to the origin.

The example is $\sum_{j=1}^{n} z_j^2$ with $n = 1$, and every case of the local computation looks like it with S^0 replaced by S^{n-1}. Lemma 4.3.7 already describes the piece of a nearby fibre that surrounds the vanishing sphere, the *Milnor fibre* of the singularity: it is a disc bundle of the tangent bundle of that sphere. What the proof of the Picard-Lefschetz formula needs beyond that description is the pair of homology classes the piece supports and the two intersection numbers between them, which come from comparing the complex orientation with the orientation of the disc bundle.

Lemma 4.4.4: Generators and Intersection Numbers in the Local Fibre

Let $n \geq 2$, let $f\colon \mathbb{C}^n \to \mathbb{C}$ be $f(z) = \sum_{j=1}^{n} z_j^2$, let $B = \{z \,\big|\, |z| \leq \epsilon\}$ and, for $0 < |\tau| < \epsilon^2$, put $F_\tau = f^{-1}(\tau) \cap B$. Write $m = n - 1$ and

$$\sigma_n = (-1)^{(n-1)(n-2)/2}$$

By lemma 4.3.7 the fibre F_τ is a compact complex m-manifold with boundary, diffeomorphic to a disc bundle of the tangent bundle TS^m with the vanishing sphere as zero section; write (u, v) for the coordinates of that disc bundle, scaled so that $|v| \leq 1$, and δ for the class of the zero section, which for $\tau > 0$ is the set of real points $\{x \in \mathbb{R}^n \,\big|\, |x|^2 = \tau\}$ of F_τ.

1. $H_m(F_\tau, \mathbb{Z}) = \mathbb{Z}\,\delta$ and $H_k(F_\tau, \mathbb{Z}) = 0$ for $k \neq 0, m$, while $H_m(F_\tau, \partial F_\tau; \mathbb{Z})$ is infinite cyclic, generated by the class Δ of a fibre of the disc bundle.
2. With δ and Δ suitably oriented,

$$\langle \delta, \Delta \rangle = \sigma_n, \qquad \langle \delta, \delta \rangle = \sigma_n\big(1 + (-1)^m\big)$$

***Proof*:**

1. By lemma 4.3.7 the fibre F_τ deformation retracts onto the vanishing sphere, so its homology is that of S^m [30, Homology Groups of Spheres] and δ generates in degree m. Poincaré-Lefschetz duality [30, Poincaré-Lefschetz duality] gives $H_m(F_\tau, \partial F_\tau; \mathbb{Z}) \cong H^m(F_\tau, \mathbb{Z}) \cong \mathbb{Z}$. The fibre disc $\Delta_{u_0} = \{(u_0, v) \,\big|\, v \perp u_0,\ |v| \leq 1\}$ over a point $u_0 \in S^m$ is a relative cycle meeting δ transversally in the single point $(u_0, 0)$, so $\langle \delta, \Delta_{u_0} \rangle = \pm 1$. Since the pairing

between $H_m(F_\tau)$ and $H_m(F_\tau, \partial F_\tau)$ is unimodular and δ generates the first group, Δ_{u_0} generates the second.

2. Multiplying the coordinates by a unit scalar λ with $\lambda^2 = \bar\tau/|\tau|$ carries F_τ onto $F_{|\tau|}$, preserves B and preserves complex orientations, so we may assume $\tau > 0$. Both numbers require comparing the complex orientation of F_τ with the orientation of the disc bundle. Fix an orientation $\mathfrak{o}$ of S^m and let $\mathcal{O}$ be the orientation of the disc bundle given by $\mathfrak{o}$ on the base followed by $\mathfrak{o}$ on the fibre. At a point $(u, 0)$ of the zero section, where $z = \sqrt{\tau}\, u$ is real, the tangent space of F_τ is $\{w \in \mathbb{C}^n \big| u \cdot w = 0\} = T_u S^m \oplus i\, T_u S^m$, and by the parametrization in the proof of lemma 4.3.7 the first summand is the base directions and the second the fibre directions. For a real basis $a_1, \dots, a_m$ of $T_u S^m$ the complex orientation is $(a_1, ia_1, \dots, a_m, ia_m)$, while $\mathcal{O}$ is $(a_1, \dots, a_m, ia_1, \dots, ia_m)$; the shuffle relating the two has sign $(-1)^{m(m-1)/2} = \sigma_n$. As F_τ is connected, the complex orientation equals $\sigma_n\, \mathcal{O}$ everywhere.

   With respect to $\mathcal{O}$, the orientation of δ followed by that of Δ_{u_0} is by construction $\mathcal{O}$ itself, so the transverse intersection at $(u_0, 0)$ has sign $+1$ for $\mathcal{O}$ and sign σ_n for the complex orientation. This is the first formula.

   For the second, the normal bundle of δ in F_τ is the vertical bundle along the zero section, that is TS^m, and the self-intersection is the Euler number of the normal bundle oriented so that the orientation of δ followed by that of the normal bundle gives the orientation of F_τ [32, self-intersection via the normal bundle]. The $\mathfrak{o}$-orientation of the fibres achieves this only up to the factor σ_n just computed, so

$$\langle \delta, \delta \rangle = \sigma_n \int_{S^m} e(TS^m) = \sigma_n\, \chi_{\text{top}}(S^m) = \sigma_n\big(1 + (-1)^m\big)$$

   as we sought to show.

The self-intersection already separates the two parities. For n even the vanishing sphere is odd-dimensional and its self-intersection vanishes, which will make the Picard-Lefschetz transformation a transvection; for n odd the self-intersection is ± 2 and the transformation is a reflection of order 2. The case $n = 2$ is small enough to draw.

Example 4.4.5: The Vanishing Annulus

Take $n = 2$, so $f(z_1, z_2) = z_1^2 + z_2^2$ and $m = 1$. The change of coordinates $w_1 = z_1 + iz_2$, $w_2 = z_1 - iz_2$ turns f into $w_1 w_2$, and the fibre over $\tau \neq 0$ is $\{w_1 w_2 = \tau\}$, the graph of $w_2 = \tau / w_1$, which inside B is an annulus. Lemma 4.3.7 says the same thing: a disc bundle of TS^1 is $S^1 \times [-1, 1]$.

The vanishing cycle is the core circle. In the coordinates (u, v) it is $v = 0$; in the original coordinates and for $\tau > 0$ it is the real circle $x_1^2 + x_2^2 = \tau$. The two intersection numbers of lemma 4.4.4 read $\sigma_2 = 1$, so $\langle \delta, \Delta \rangle = 1$ and $\langle \delta, \delta \rangle = 0$, the latter because a circle on a surface can always be pushed off itself.

The singular fibre $F_0 = \{w_1 w_2 = 0\}$ is a pair of discs meeting at the origin, and the annulus degenerates to it by collapsing the core circle: this is part (3) of lemma 4.3.7 made visible, since a pair of discs glued at a point is contractible.

The monodromy of a loop around the critical value can be chosen to do nothing away from the double point, so that the entire effect of the loop is a diffeomorphism of the local fibre, and that diffeomorphism can be written down.

Lemma 4.4.6: Localization of the Local Monodromy

Let $\mathcal{X}$ be a complex manifold of dimension $n \geq 2$, let $D \subseteq \mathbb{C}$ be a disc centred at 0, and let $f\colon \mathcal{X} \to D$ be a proper holomorphic map which is a submersion over $D \setminus \{0\}$ and whose only critical point p is an ordinary double point of $X_0 = f^{-1}(0)$. Choose holomorphic coordinates near p in which $f = \sum_{j=1}^{n} z_j^2$, let B be a ball as in lemma 4.4.4, and fix a small $\tau > 0$. Write $F_t = X_t \cap B$ and $F = F_\tau$.

1. The restriction of f to $f^{-1}(\bar{D}_\tau) \setminus \mathring{B}$ is a proper submersion onto the closed disc $\bar{D}_\tau$ of radius τ, and is therefore trivial over it.

2. The geometric monodromy of the counterclockwise loop $|t| = \tau$ may be chosen to be a diffeomorphism h of X_τ equal to the identity on $X_\tau \setminus \mathring{F}$.

3. The diffeomorphism h of (2) may in addition be chosen to equal on F, in the coordinates (u, v) of lemma 4.4.4, the map

$$h_\psi(u, v) = \big(\cos\theta\, u - \sin\theta\, \hat{v},\quad |v| \sin\theta\, u + \cos\theta\, v\big), \qquad \theta = \pi\, \psi(|v|)$$

   where $\hat{v} = v/|v|$ and $\psi\colon [0,1] \to [0,1]$ is smooth, identically 1 near 0, identically 0 near 1, and satisfies $\psi' < 0$ at every point where its value lies in $(0,1)$, so that ψ is strictly decreasing there. Such a ψ exists: take $\psi(x) = 1 - \int_0^x \chi$ for a smooth $\chi \geq 0$ of integral 1 which is strictly positive exactly on an interval $(a, b) \subseteq (0,1)$. Near the zero section h_ψ is $(u, v) \mapsto (-u, -v)$ and near ∂F it is the identity, so it is smooth on all of F despite the appearance of $\hat{v}$.

***Proof*:**

1. Shrinking ϵ and then τ, the sphere ∂B is transverse to every fibre X_t with $|t| \leq \tau$, since transversality holds at $t = 0$ for the model $\sum z_j^2$ and is an open condition on a compact set. Hence $f^{-1}(\bar{D}_\tau) \setminus \mathring{B}$ is a manifold with corners on which f is a proper submersion, and Ehresmann's theorem in the form valid for manifolds with boundary [32, Ehresmann for Manifolds with Boundary] trivializes it over the contractible base $\bar{D}_\tau$. In particular the family of links $L_t = X_t \cap \partial B$ is trivial over $\bar{D}_\tau$, including over $t = 0$.

2. Let A be an annulus neighbourhood of the circle $|t| = \tau$ inside $D \setminus \{0\}$ and let w be the vector field on A generating the counterclockwise rotation. On $f^{-1}(A)$ the map f is a proper submersion, so a partition of unity produces a vector field $\tilde{w}_1$ with $df(\tilde{w}_1) = w$. On $f^{-1}(A) \setminus \mathring{B}$ the trivialization of (1) gives a second such lift $\tilde{w}_2$, namely the one whose flow preserves the product decomposition. Choose a slightly larger ball $B' \supset B$, of radius ϵ' and still transverse to the fibres, and a cutoff χ that equals 1 on B and vanishes off B', and set

$$\tilde{w} = \chi\, \tilde{w}_1 + (1 - \chi)\, \tilde{w}_2$$

   A convex combination of lifts of w is again a lift of w, so $\tilde{w}$ is f-related to w, and its time-2π flow is a diffeomorphism h of X_τ realizing the geometric monodromy. Off B' one has $\tilde{w} = \tilde{w}_2$, whose flow returns each point to its starting position after a full turn because the trivialization of (1) extends over the disc. So h is the identity off B', and after replacing B by B' it is the identity on $X_\tau \setminus \mathring{F}$.

3. Inside the ball the family has an explicit trivialization: since f is homogeneous of degree 2, the maps
$$\nu_\vartheta \colon F_\tau \to F_{\tau e^{2\pi i\vartheta}}, \qquad \nu_\vartheta(z) = e^{i\pi\vartheta} z$$
preserve $|z|$, hence preserve B, and form a trivialization of the local family over the circle. Its monodromy is $\nu_1 = -\mathrm{id}$, which in the coordinates of lemma 4.4.4 is $(u, v) \mapsto (-u, -v)$, the rotation by $-\pi$ in the plane spanned by the orthonormal pair $(u, \hat{v})$.

   That is not yet a monodromy of the kind (2) asks for, because ν_ϑ does not agree near ∂B with the trivialization of (1); the two differ by a path of diffeomorphisms of the link, which the model computes. Writing $z = x + iy$ and $t = t_1 + it_2$, the link L_t is the set of pairs (x, y) with
$$|x| = a(t_1), \qquad |y| = b(t_1), \qquad x \cdot y = t_2/2, \qquad 2a^2 = \epsilon^2 + t_1, \quad 2b^2 = \epsilon^2 - t_1$$
so $(\xi, \eta) = (x/a,\ y/b)$ identifies L_t with
$$\Sigma_\beta = \big\{(\xi, \eta) \in S^m \times S^m \;\big|\; \xi \cdot \eta = \beta\big\}, \qquad \beta = \beta(t) := \frac{t_2}{2ab}$$
and $|\beta| < 1$ for $|t| \le \tau$. The maps
$$\Theta_\beta \colon \Sigma_0 \to \Sigma_\beta, \qquad (\xi, \eta) \longmapsto \big(\xi,\ \beta\,\xi + \sqrt{1 - \beta^2}\,\eta\big)$$
are diffeomorphisms depending smoothly on β, so $t \mapsto \Theta_{\beta(t)}$ trivializes the family of links over $\bar{D}_\tau$, and $\beta(\tau) = 0$. Any other such trivialization differs from this one by a map $\bar{D}_\tau \to \mathrm{Diff}(L_\tau)$, whose restriction to the boundary circle is a null-homotopic loop at the identity, so the comparison path
$$c_\vartheta = \Theta^{-1}_{\beta(t(\vartheta))} \circ \nu_\vartheta \in \mathrm{Diff}(L_\tau), \qquad t(\vartheta) = \tau e^{2\pi i\vartheta}$$
is independent of the trivialization chosen in (1), up to homotopy with fixed endpoints.

   That homotopy class is the class of a rotation. On L_τ one has $\xi \perp \eta$, and ν_ϑ carries (x, y) to $(\cos\theta\, x - \sin\theta\, y,\ \sin\theta\, x + \cos\theta\, y)$ with $\theta = \pi\vartheta$, which in the normalized coordinates is the map sending (ξ, η) to the pair
$$p_s = \cos\theta\, a_s\, \xi - \sin\theta\, b_s\, \eta, \qquad q_s = \sin\theta\, a_s\, \xi + \cos\theta\, b_s\, \eta$$
divided by their lengths, at $s = 0$, where $a_s = (1 - s)a(\tau) + s$ and $b_s = (1 - s)b(\tau) + s$ for $s \in [0, 1]$. The matrix carrying (ξ, η) to (p_s, q_s) has determinant $a_s b_s > 0$, and the lengths of p_s and q_s and their inner product depend only on s and ϑ, so normalizing gives a diffeomorphism of Σ_0 onto $\Sigma_{\beta_{s,\vartheta}}$ for the resulting inner product $\beta_{s,\vartheta}$; composing it with $\Theta^{-1}_{\beta_{s,\vartheta}}$ produces a family $c^s_\vartheta \in \mathrm{Diff}(\Sigma_0)$, continuous in (s, ϑ), with $c^0 = c$. At $\vartheta = 0$ and at $\vartheta = 1$ one has $\sin\theta = 0$, hence $\beta_{s,\vartheta} = 0$ and $c^s_\vartheta = \mathrm{id}$ respectively $-\mathrm{id}$, for every s: the homotopy fixes the endpoints. At $s = 1$, where $a_1 = b_1 = 1$, the pair (p_1, q_1) is already orthonormal and c^1_ϑ is the rotation of (ξ, η) by the angle $-\pi\vartheta$ in the plane they span. So c is homotopic with fixed endpoints to that path of rotations.

   Correct ν by the rotation in the opposite sense, damped towards the zero section. Let g_ϑ be the rotation by $\pi\vartheta\big(1 - \psi(|v|)\big)$ in the plane spanned by $(u, \hat{v})$; it preserves $|v|$ and is the identity near the zero section, where $\psi = 1$, so it is a diffeomorphism of F_τ, and $g_0 = \mathrm{id}$. The maps $\nu_\vartheta \circ g_\vartheta$ again trivialize the local family over the circle, and near ∂B, where

$\psi = 0$, their comparison with the trivialization of (1) is the pointwise composite $c_\vartheta \circ g_\vartheta$: a path of diffeomorphisms of the link which starts and ends at the identity and which, by the previous paragraph, is null-homotopic with fixed endpoints. Every sphere $|z| = \varrho$ with $\epsilon \leq \varrho \leq \epsilon'$ is transverse to the fibres, so $(f, |z|)$ is a submersion on the collar $B' \setminus \mathring{B}$ and Ehresmann's theorem [32, Ehresmann's theorem] allows both trivializations to be taken there to preserve $|z|$; reading the parameter of a null-homotopy of $c_\vartheta \circ g_\vartheta$ as the radius ϱ then interpolates between the two across the collar, and they glue to a trivialization of the family over the circle.

Its monodromy is the identity outside $\mathring{B}$: beyond the collar it is the monodromy of the trivialization of (1), which is trivial because that trivialization extends over the disc, and on the collar the interpolating correction is the identity at $\vartheta = 1$, the null-homotopy being stationary at the endpoints. On F_τ the monodromy is $\nu_1 \circ g_1$, the rotation by $\pi(1 - \psi(|v|))$ followed by the rotation by $-\pi$ in the same plane, that is the rotation by $-\pi\psi(|v|)$. Writing that rotation out in coordinates produces the displayed formula for h_ψ, as we sought to show.

Real rotations of $\mathbb{R}^n$ preserve $\sum z_j^2$, so h_ψ does map F_τ to itself; at the zero section it is the antipodal map of S^{n-1}, and near the boundary it is the identity. It is the higher-dimensional Dehn twist: on the annulus of example 4.4.5 it is the classical twist that rotates the core circle once and tapers to the identity at the two boundary circles.

The monodromy differs from the identity only through a diffeomorphism supported in the local fibre; that fibre carries exactly one class in each of the two relevant homology groups; and the diffeomorphism is explicit.

Theorem 4.4.7: The Picard-Lefschetz Formula

Let $\mathcal{X}$ be a complex manifold of dimension $n \geq 2$, let $D \subseteq \mathbb{C}$ be a disc centred at 0, and let $f\colon \mathcal{X} \to D$ be a proper holomorphic map which is a submersion over $D \setminus \{0\}$ and whose only critical point is an ordinary double point of X_0. Fix $\tau \in D \setminus \{0\}$ small enough that lemma 4.4.6 applies, let $\delta \in H_{n-1}(X_\tau, \mathbb{Z})$ be the vanishing cycle, that is the image of the class of the vanishing sphere in the local fibre (lemma 4.4.4), and let $T = h_*$ be the action on $H_*(X_\tau, \mathbb{Z})$ of the geometric monodromy h of the counterclockwise loop $|t| = |\tau|$. Then T is the identity on $H_k(X_\tau, \mathbb{Z})$ for every $k \neq n-1$, while on $H_{n-1}(X_\tau, \mathbb{Z})$

$$T(\alpha) = \alpha + \varepsilon_n \langle \alpha, \delta \rangle \, \delta, \qquad \varepsilon_n := (-1)^{n(n+1)/2} \tag{4.5}$$

Moreover $\langle \delta, \delta \rangle = -\varepsilon_n\big(1 + (-1)^{n-1}\big)$ and $T(\delta) = (-1)^n \delta$.

***Proof*:**
Replacing t by $t|\tau|/\tau$ throughout, we may assume $\tau > 0$. Write $m = n-1$, let B and $F = X_\tau \cap B$ be as in lemma 4.4.6, and let h be the geometric monodromy that lemma gives: the identity on $X_\tau \setminus \mathring{F}$ and equal on F to the map h_ψ of its part (3). Any two geometric monodromies of the loop are isotopic (definition 4.4.2), so $T = h_*$ is unaffected by the choice. Since ψ vanishes near 1, the map h is the identity on the collar $\{\psi = 0\}$ of ∂F; write $F'' \subseteq \mathring{F}$ for the complement of that collar, so that h is the identity on all of $X_\tau \setminus F''$. Set $\sigma_n = (-1)^{(n-1)(n-2)/2}$ as in lemma 4.4.4.

The identity

$$\frac{n(n+1)}{2} - \frac{(n-1)(n-2)}{2} = 2n-1$$

shows that $\varepsilon_n = -\sigma_n$, and the assertion about $\langle \delta, \delta \rangle$ is then part (2) of lemma 4.4.4: the self-intersection may be computed inside F, because δ is supported there and a representative may be pushed off itself without leaving F. The proof of (4.5) occupies Steps 1 to 4, and Step 5 treats the remaining degrees.

Step 1: The variation map. Let a be a cycle representing $\alpha \in H_k(X_\tau, \mathbb{Z})$. The sets $\mathring{F}$ and $X_\tau \setminus F''$ form an open cover of X_τ, so after barycentric subdivision [30, Homology Of Covered Space] every simplex of a lies in one of them. Those of the second kind are fixed by h and cancel in $h_\# a - a$; those of the first kind are carried by h into F, so their contribution is a chain in F. Thus $h_\# a - a$ is a chain in F with zero boundary, hence a cycle in F. Its class depends only on the image of α in $H_k(X_\tau, X_\tau \setminus \mathring{F})$, because a class vanishing there is represented by a cycle supported in $X_\tau \setminus \mathring{F}$, which h fixes. This defines the *variation*

$$\mathrm{var}\colon H_k\big(X_\tau, X_\tau \setminus \mathring{F}\big) \to H_k(F, \mathbb{Z}), \qquad T - \mathrm{id} = \mathrm{var} \circ j$$

where j is the natural map from $H_k(X_\tau, \mathbb{Z})$. Excision identifies the source with $H_k(F, \partial F; \mathbb{Z})$ [30, Excision Theorem]. The collar needed to apply that duality is given by [32, Collar Neighbourhood Theorem].

Step 2: The image of j. Take $k = m$. By lemma 4.4.4 the group $H_m(F, \partial F; \mathbb{Z})$ is generated by the fibre disc Δ, and the intersection pairing between $H_m(F, \mathbb{Z}) = \mathbb{Z}\delta$ and $H_m(F, \partial F; \mathbb{Z}) = \mathbb{Z}\Delta$ satisfies $\langle \delta, \Delta \rangle = \sigma_n$. Since δ is supported in F, the intersection number $\langle \delta, \alpha \rangle$ computed in X_τ agrees with the pairing of δ against $j(\alpha)$. Writing $j(\alpha) = c\,\Delta$ and pairing with δ gives $c\,\sigma_n = \langle \delta, \alpha \rangle$, so

$$j(\alpha) = \sigma_n \langle \delta, \alpha \rangle \, \Delta$$

Step 3: The variation of the fibre disc. The claim is that $\mathrm{var}(\Delta) = (-1)^n \delta$. Work in the coordinates (u, v) of lemma 4.4.4 and take $h = h_\psi$ from lemma 4.4.6, which acts on the point (u, v) as the rotation by $-\theta$, $\theta = \pi\psi(|v|)$, in the plane spanned by the orthonormal pair $(u, \hat{v})$. Choose an orthonormal basis $e_0, \dots, e_m$ of $\mathbb{R}^n$, set $u_0 = e_0$, and orient S^m by declaring $u \wedge \mathrm{or}(T_u S^m)$ to be the standard orientation of $\mathbb{R}^n$, so that $\mathrm{or}(T_{e_0} S^m) = e_1 \wedge \cdots \wedge e_m$. Let $\Delta = \Delta_{u_0}$ be the fibre disc over u_0, oriented by that basis. Fix $\varphi \in (0, \pi)$ and let R_φ be the rotation by φ in the plane (e_0, e_1), so that $u_1 := R_\varphi e_0$ and $w' := R_\varphi e_1$ are orthonormal and R_φ fixes $e_2, \dots, e_m$.

The chain $h(\Delta) - \Delta$ is a cycle in F because h is the identity near ∂F, and its class is $\lambda\,\delta$ for some $\lambda \in \mathbb{Z}$. Pair it with the relative class Δ_{u_1}. The pairing may be computed at the chain level, since neither Δ nor $h(\Delta)$ meets $\partial \Delta_{u_1}$: both meet ∂F in the fibre over u_0, and $u_0 \neq u_1$. As Δ and Δ_{u_1} are disjoint fibres, only $h(\Delta)$ contributes, so $\langle \delta, \Delta_{u_1} \rangle = \sigma_n$ gives $\lambda \sigma_n = \langle h(\Delta), \Delta_{u_1} \rangle$.

The intersection $h(\Delta) \cap \Delta_{u_1}$ consists of one point. Indeed $h(u_0, v)$ has first coordinate $\cos\theta\, e_0 - \sin\theta\, \hat{v}$, and setting this equal to $u_1 = \cos\varphi\, e_0 + \sin\varphi\, e_1$ forces $\cos\theta = \cos\varphi$, hence $\theta = \varphi$ because $\theta \in [0, \pi]$, and then $\hat{v} = -e_1$. Since ψ is strictly decreasing where it takes values in $(0, 1)$, the equation $\pi\psi(s) = \varphi$ has the unique solution $s \in (0, 1)$, so the only preimage is $v_* = -s\, e_1$, and

$$P := h(u_0, v_*) = \big(u_1,\ -s\, w'\big)$$

which lies in Δ_{u_1} because $w' \perp u_1$.

It remains to compute the sign of that intersection. Differentiating $v \mapsto h(u_0, v)$ at v_* along e_1 and along e_j for $j \geq 2$ gives, with $\kappa := \pi\psi'(s)$, which is negative because $\psi(s) = \varphi/\pi$ lies in $(0,1)$ and lemma 4.4.6 takes $\psi' < 0$ at every such point,

$$X_1 = \big(-\kappa\, w',\ w' - s\kappa\, u_1\big), \qquad X_j = \big(-\sin\varphi\, e_j,\ s\cos\varphi\, e_j\big)$$

the second up to the positive factor $1/s$. The tangent space of Δ_{u_1} at P is spanned by $Y_1 = (0, w')$ and $Y_j = (0, e_j)$, in that order for the chosen orientation, since $w', e_2, \dots, e_m$ is the positively oriented basis $R_\varphi(e_1, \dots, e_m)$ of $T_{u_1}S^m$. Splitting $T_P F$ into horizontal and vertical parts for the Levi-Civita connection of the round sphere, the horizontal lift of $a \in T_{u_1}S^m$ at the point (u_1, v_P) is $(a, -(v_P \cdot a)u_1)$, so with $v_P = -sw'$

$$H_1 = (w',\, s\, u_1), \qquad H_j = (e_j, 0), \qquad V_1 = (0, w'), \qquad V_j = (0, e_j)$$

is a basis adapted to the orientation $\mathcal{O}$ of the disc bundle used in the proof of lemma 4.4.4, the base orientation followed by the fibre orientation, for which the complex orientation is $\sigma_n\mathcal{O}$; explicitly $\mathcal{O} = H_1 \wedge \cdots \wedge H_m \wedge V_1 \wedge \cdots \wedge V_m$. In this basis $X_1 = -\kappa H_1 + V_1$ and $X_j = -\sin\varphi\, H_j + s\cos\varphi\, V_j$, while $Y_1 = V_1$ and $Y_j = V_j$. Reordering basis and vectors by the same shuffle to $(H_1, V_1, H_2, V_2, \dots)$ and $(X_1, Y_1, X_2, Y_2, \dots)$ leaves the determinant unchanged and makes it block diagonal with blocks

$$\begin{pmatrix} -\kappa & 0 \\ 1 & 1 \end{pmatrix}, \qquad \begin{pmatrix} -\sin\varphi & 0 \\ s\cos\varphi & 1 \end{pmatrix}$$

of determinants $-\kappa > 0$ and $-\sin\varphi < 0$. The determinant therefore has sign $(-1)^{m-1}$, so the intersection sign at P is $(-1)^{m-1}$ for $\mathcal{O}$ and $\sigma_n(-1)^{m-1}$ for the complex orientation. Comparing with $\lambda\sigma_n$ gives $\lambda = (-1)^{m-1} = (-1)^n$.

Step 4: Assembling. Combining Steps 1 to 3,

$$T(\alpha) - \alpha = \operatorname{var}\big(j(\alpha)\big) = \sigma_n \langle \delta, \alpha\rangle \operatorname{var}(\Delta) = \sigma_n(-1)^n \langle \delta, \alpha\rangle\, \delta$$

The pairing on middle homology satisfies $\langle \delta, \alpha\rangle = (-1)^m\langle \alpha, \delta\rangle$, and $(-1)^{n+m} = (-1)^{2n-1} = -1$, so the coefficient is $-\sigma_n = \varepsilon_n$. This is (4.5). Substituting $\alpha = \delta$ and using $\langle \delta, \delta\rangle = \sigma_n(1 + (-1)^m)$ gives $T(\delta) = \delta - (1 + (-1)^{n-1})\delta = (-1)^n\delta$.

Step 5: The other degrees. For $k \neq m$ and $k \neq 2m$, Lefschetz duality and lemma 4.4.4 give $H_k(F, \partial F; \mathbb{Z}) \cong H^{2m-k}(F, \mathbb{Z}) = 0$, so $j = 0$ and T is the identity on $H_k(X_\tau, \mathbb{Z})$ by Step 1. For $k = 2m$ the group $H_{2m}(X_\tau, \mathbb{Z})$ is free on the fundamental classes of the connected components of X_τ, a compact oriented $2m$-manifold, which need not be connected. The local fibre F is connected, so it lies in a single component C_0, and h is the identity on the open set $X_\tau \setminus F$. That set meets every component: the components other than C_0 are contained in it, and C_0 is not contained in F, since $F \subseteq C_0$ would force $C_0 = F$, whereas F has nonempty boundary and C_0 has none. So h fixes a point of each component, hence carries each component to itself, and restricts on it to an orientation-preserving diffeomorphism; it therefore fixes each of the generating fundamental classes, completing the proof.

The content of (4.5) is that a full turn around a critical value changes a class only in the direction of the vanishing cycle, and by an amount that a single intersection number determines. Two consequences are immediate from the formula. The transformation T fixes the hyperplane $\delta^\perp = \{\alpha \,|\, \langle \alpha, \delta\rangle = 0\}$ pointwise, and it preserves the intersection pairing, as it must, being induced by an orientation-

preserving diffeomorphism. And the parity of n decides its nature: for n odd, $\langle\delta,\delta\rangle = \pm 2$ and T is the reflection in $\delta^{\perp}$, of order 2; for n even, $\langle\delta,\delta\rangle = 0$ and T is a transvection, unipotent with $(T-\mathrm{id})^2 = 0$ and of infinite order whenever $\delta \neq 0$ in $H_{n-1}(X_\tau,\mathbb{Q})$. A δ that is nonzero but torsion gives $T = \mathrm{id}$, since the integral intersection pairing vanishes against torsion classes.

The formula also takes a clean shape in the normalization chapter 3 fixed. Suppose the fibres are compact Kähler, as they are for a Lefschetz pencil on a projective manifold. The fibre X_τ then has dimension $n-1$, and the form proposition 3.2.9 attaches to its middle degree $k = n-1$ carries the factor $\omega^{(n-1)-k} = 1$, so it is

$$Q(\alpha,\beta) = (-1)^{(n-1)(n-2)/2} \int_{X_\tau} \alpha \wedge \beta = \sigma_n \langle \alpha, \beta \rangle$$

Writing δ also for the Poincaré dual of the vanishing cycle, (4.5) becomes

$$T(\alpha) = \alpha - Q(\alpha,\delta)\,\delta$$

with no residual sign. The normalizing factor that proposition 3.2.9 introduced for the sake of positivity is the same factor that makes the Picard-Lefschetz formula sign-free.

The case of a pencil of curves on a surface is the classical one, and there the formula can be checked against a count of singular fibres.

Example 4.4.8: The Nodal Degeneration of a Curve

Let $n = 2$, so the fibres are compact Riemann surfaces and $\varepsilon_2 = (-1)^3 = -1$. Theorem 4.4.7 reads

$$T(\alpha) = \alpha - \langle \alpha, \delta \rangle\, \delta$$

on $H_1(X_\tau,\mathbb{Z})$, with $\langle\delta,\delta\rangle = 0$. If δ is part of a symplectic basis, say $\langle\delta,\beta\rangle = 1$, then T fixes δ and sends β to $\beta - \langle\beta,\delta\rangle\delta = \beta + \delta$, so in the basis (δ,β) of the plane they span the matrix is

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

unipotent of infinite order.

A concrete pencil. Take the general pencil of plane cubics of example 4.3.9: the associated fibration is $f\colon \tilde{X} \to \mathbb{CP}^1$ with $\tilde{X}$ the blow-up of $\mathbb{CP}^2$ at the nine base points (proposition 4.3.6), the smooth fibres are elliptic curves, and the Euler characteristic count of exercise 4.3.1 (2), carried out there, gives $\mu = 12$ nodal members.

Each of the twelve local monodromies is a transvection along its own vanishing cycle, and the fundamental group of the complement U of the twelve critical values is generated by loops $\gamma_1,\dots,\gamma_{12}$ around the punctures subject to the single relation that their product in a suitable order is trivial, so the monodromy group $\Gamma \subseteq \mathrm{Sp}(2,\mathbb{Z}) = \mathrm{SL}_2(\mathbb{Z})$ is generated by twelve transvections satisfying that one relation.

What one transvection does to the periods. Let δ and β be as above, let η be a nonzero holomorphic 1-form on the fibre, and let $\lambda = \int_\beta \eta \big/ \int_\delta \eta$ be the period ratio. Transporting once around the puncture replaces the pair (δ,β) by $(T\delta, T\beta) = (\delta, \beta+\delta)$, so

$$\lambda \longmapsto \frac{\int_{\beta+\delta}\eta}{\int_\delta \eta} = \lambda + 1$$

This is the classical monodromy of a degenerating family of elliptic curves, and it is the substitution $\tau \mapsto \tau + 1$ on the lattice parameter of exercise 3.2.1 (6). Its Hodge-theoretic form, in which λ becomes the value of a period map and $\lambda \mapsto \lambda + 1$ the monodromy of that map, belongs to chapter 5; the elliptic-curve background is in [6].

The example used a point that the formula makes general: conjugating a local monodromy transports its vanishing cycle. This keeps the vanishing cycles a single stable set rather than μ unrelated classes, and it identifies the classes on which the whole monodromy group acts trivially.

Corollary 4.4.9: Conjugate Vanishing Cycles and the Invariant Classes

Let $f\colon \tilde{X} \to \mathbb{C}\mathrm{P}^1$ be the fibration of a Lefschetz pencil with critical values $t_1, \dots, t_\mu$, let $t_0 \in U$, and for each i let $\gamma_i \in \pi_1(U, t_0)$ be a simple loop around t_i along a chosen path, with $T_i = \rho(\gamma_i)$ and with $\delta_i \in H_{n-1}(X_{t_0}, \mathbb{Z})$ the vanishing cycle at t_i transported to X_{t_0} along that path, so that T_i satisfies (4.5) with δ replaced by δ_i. Let Γ be the monodromy group.

1. For every $g \in \pi_1(U, t_0)$ the element $\rho(g\gamma_i g^{-1})$ satisfies (4.5) with δ_i replaced by $\rho(g)\delta_i$. In particular the set $\mathcal{V}$ of all classes arising as vanishing cycles, over all choices of connecting paths, is stable under Γ and under $\delta \mapsto -\delta$, and every element of $\mathcal{V}$ has the same self-intersection $-\varepsilon_n(1 + (-1)^{n-1})$.
2. Γ is generated by $T_1, \dots, T_\mu$.
3. The invariant classes are the classes orthogonal to every vanishing cycle:

$$H_{n-1}(X_{t_0}, \mathbb{Z})^\Gamma = \{\alpha \,|\, \langle \alpha, \delta_i \rangle = 0 \text{ for } i = 1, \dots, \mu\}$$

and $T_i = \mathrm{id}$ if and only if $\delta_i = 0$ in $H_{n-1}(X_{t_0}, \mathbb{Q})$.

***Proof*:**

1. Put $A = \rho(g)$, an automorphism preserving the intersection pairing because it is induced by an orientation-preserving diffeomorphism. Then for every α,

$$AT_iA^{-1}(\alpha) = A\big(A^{-1}\alpha + \varepsilon_n \langle A^{-1}\alpha, \delta_i \rangle \delta_i\big) = \alpha + \varepsilon_n \langle \alpha, A\delta_i \rangle A\delta_i$$

using $\langle A^{-1}\alpha, \delta_i \rangle = \langle \alpha, A\delta_i \rangle$. Geometrically $A\delta_i$ is the vanishing cycle at t_i transported along the path modified by g, so conjugates of the T_i are again Picard-Lefschetz transformations and $\mathcal{V}$ is Γ-stable. Stability under sign change holds because the vanishing sphere carries two orientations and (4.5) is unchanged by $\delta \mapsto -\delta$. The self-intersection is the value computed in theorem 4.4.7, and A preserves it.

2. The complement U of μ points in $\mathbb{C}\mathrm{P}^1$ has fundamental group generated by simple loops around the punctures [30], and changing the connecting path replaces γ_i by a conjugate. Applying ρ gives the claim.

3. By (2) a class is Γ-invariant if and only if it is fixed by every T_i, and by (4.5) the condition $T_i\alpha = \alpha$ says $\varepsilon_n \langle \alpha, \delta_i \rangle \delta_i = 0$. If δ_i is not torsion, that is if $\delta_i \neq 0$ in $H_{n-1}(X_{t_0}, \mathbb{Q})$, an integer multiple of δ_i vanishes only when the integer does, so the condition reads $\langle \alpha, \delta_i \rangle = 0$. If δ_i is torsion, say $N\delta_i = 0$ with $N \geq 1$, then $N\langle \alpha, \delta_i \rangle = \langle \alpha, N\delta_i \rangle = 0$ in $\mathbb{Z}$, so $\langle \alpha, \delta_i \rangle = 0$

and both conditions hold automatically. That is the displayed identity, the case $\delta_i = 0$ being the case $N = 1$.

The same dichotomy gives the last statement. If $\delta_i = 0$ in $H_{n-1}(X_{t_0}, \mathbb{Q})$, that is if δ_i is torsion, the computation just made yields $\langle \alpha, \delta_i \rangle = 0$ for every α and hence $T_i = \mathrm{id}$. Conversely if $\delta_i \neq 0$ in $H_{n-1}(X_{t_0}, \mathbb{Q})$ then, the intersection pairing being nondegenerate over $\mathbb{Q}$ by Poincaré duality, some α, integral after clearing denominators, has $\langle \alpha, \delta_i \rangle \neq 0$; then $\langle \alpha, \delta_i \rangle \delta_i$ is nonzero already in $H_{n-1}(X_{t_0}, \mathbb{Q})$, so $T_i \alpha \neq \alpha$, as we sought to show.

Part (3) is the statement the Hodge theory of families will use, and chapter 5 takes it up: the invariant classes are exactly the classes no vanishing cycle pairs against, so intersection theory alone determines them. Everything in this section is topology. The representation ρ records how the lattice $H^{n-1}(X_{t_0}, \mathbb{Z})$ is carried around U, and is insensitive to the Hodge decomposition of the fibres, which moves holomorphically over U while the lattice stays flat.

Exercise 4.4.1

1. The hypotheses of theorem 4.4.7 exclude $n = 1$, where the fibre is a finite set of points. Carry out the case $n = 1$ by hand for $f(z) = z^2$, as in example 4.4.3: with $\delta = [1] - [-1]$ in $\tilde{H}_0(F_\tau, \mathbb{Z})$ and the intersection pairing on H_0 of a finite set of points declared to be $\langle [p], [q] \rangle = 1$ if $p = q$ and 0 otherwise, compute $\langle \delta, \delta \rangle$ and check both that $\langle \delta, \delta \rangle = -\varepsilon_1(1 + (-1)^0)$ and that $T(\alpha) = \alpha + \varepsilon_1 \langle \alpha, \delta \rangle \delta$ holds for every $\alpha \in \tilde{H}_0(F_\tau, \mathbb{Z})$.

2. Let T be as in theorem 4.4.7 and suppose $\delta \neq 0$ in $H_{n-1}(X_\tau, \mathbb{Q})$.

    1. Show that for n odd, T is the orthogonal reflection $\alpha \mapsto \alpha - 2\frac{\langle \alpha, \delta \rangle}{\langle \delta, \delta \rangle}\delta$ in the hyperplane $\delta^\perp$, and that $T^2 = \mathrm{id}$.
    2. Show that for n even, $(T - \mathrm{id})^2 = 0$ and $T^k(\alpha) = \alpha + k\varepsilon_n \langle \alpha, \delta \rangle \delta$ for all $k \in \mathbb{Z}$, so that T has infinite order.
    3. Deduce in both cases that T preserves the intersection pairing, by direct computation from (4.5) rather than from the fact that T is induced by a diffeomorphism.

3. Let $Q \subseteq \mathbb{CP}^3$ be a smooth quadric surface and let $(Q_t)_{t \in \mathbb{CP}^1}$ be a Lefschetz pencil of hyperplane sections of Q, so $n = 2$ and each smooth member is a conic, a rational curve. The base locus is $Q \cap H_1 \cap H_2$, which consists of $\deg Q = 2$ points. Using $\chi_{\mathrm{top}}(Q) = 4$, the fact that blowing up a point raises the Euler characteristic by 1, and the count $\chi_{\mathrm{top}}(\tilde{Q}) = 2\chi_{\mathrm{top}}(Q_{t_0}) + \mu(-1)^n$ established in exercise 4.3.1 (2), determine the number μ of singular members. Then explain, without any further computation, why the monodromy representation on H_1 of a smooth member is trivial.

4. Let $f \colon \tilde{X} \to \mathbb{CP}^1$ be the fibration of a Lefschetz pencil, with vanishing cycles $\delta_1, \ldots, \delta_\mu$ in $H_{n-1}(X_{t_0}, \mathbb{Q})$, and let $W \subseteq H_{n-1}(X_{t_0}, \mathbb{Q})$ be the subspace they span.

    1. Show that $W^\perp = H_{n-1}(X_{t_0}, \mathbb{Q})^\Gamma$ and that both are Γ-stable.
    2. Show that W is Γ-stable, and that $W \cap W^\perp$ consists of the classes of W orthogonal to all of W.
    3. Suppose n is odd. Show that Γ is generated by reflections and that $W \cap W^\perp$ contains no nonzero δ_i.

5. Let $n = 2$, so the fibres of the pencil are compact Riemann surfaces of some genus g.

    1. Show that ρ takes values in $\mathrm{Sp}(2g, \mathbb{Z})$, the automorphisms of $H_1(X_{t_0}, \mathbb{Z}) \cong \mathbb{Z}^{2g}$ preserving the intersection pairing.
    2. Show that each T_i is a symplectic transvection, that is of the form $\alpha \mapsto \alpha + c\,\langle \alpha, \delta \rangle \delta$ with $c = -1$, and compute $\det(T_i)$ and the characteristic polynomial of T_i.
    3. Assume $g = 1$ and that the pencil has μ singular fibres whose vanishing cycles are not all proportional. Show that the invariant part $H_1(X_{t_0}, \mathbb{Q})^{\Gamma}$ is zero.

5

Variations of Hodge Structure

Every theorem so far has been about one compact Kähler manifold. The Hodge decomposition, the Lefschetz operators, the polarization: each takes a single X and produces structure on its cohomology. Chapter 4 ended somewhere else. A Lefschetz pencil presents X as a family, and following a loop in the base carried the cohomology of a fibre back to itself by a nontrivial automorphism. Cohomology moved.

This chapter takes that motion as the object of study. Let $f\colon \mathcal{X} \to S$ be a smooth projective family. Each fibre carries a polarized Hodge structure by chapter 3, and the fibres are all diffeomorphic, so the underlying vector spaces may be identified locally over S. What cannot be identified is the Hodge decomposition: the subspaces $H^{p,q}$ move as the complex structure of the fibre moves, and it is exactly this motion that a variation of Hodge structure records.

Three facts organize the chapter. The cohomology of the fibres assembles into a local system, which is to say a bundle with a flat connection, the Gauss-Manin connection; flatness is what makes "the same cohomology class" meaningful along a path, and its monodromy is the automorphism of section 4.4. The Hodge filtration, by contrast, is not flat, and its failure to be flat is not arbitrary: the connection moves F^p into F^{p-1} and no further. That is Griffiths transversality, and it is the central theorem here. Finally, recording the filtration as a point of the period domain of section 3.3 turns the family into a map from S, the period map, holomorphic and constrained by transversality to be horizontal.

The chapter closes with the case where all of this can be computed. For a smooth hypersurface in projective space the Hodge filtration on primitive cohomology is visible in the Jacobian ring of its defining polynomial, and the period map becomes a statement in commutative algebra.

5.1 Local Systems and the Gauss-Manin Connection

Everything in this chapter is attached to one piece of data, a family of smooth projective varieties, and this section fixes that data and builds the first two objects it carries. What comes out is a holomorphic vector bundle on the base whose fibre over s is $H^k(X_s, \mathbb{C})$, together with a connection on it whose flat sections are the cohomology classes that do not move. The Hodge filtration plays no part until section 5.2.

Definition 5.1.1: Smooth Projective Family

A *smooth projective family* consists of complex manifolds $\mathcal{X}$ and S with S connected, a proper holomorphic submersion $f\colon \mathcal{X} \to S$, and a closed holomorphic embedding $\iota\colon \mathcal{X} \hookrightarrow \mathbb{CP}^N \times S$ over S for some N. Its *fibres* are the compact complex manifolds $X_s = f^{-1}(s)$ for $s \in S$, all of the same dimension $n = \dim_{\mathbb{C}} \mathcal{X} - \dim_{\mathbb{C}} S$, and each is a smooth projective variety by way of ι.
The *relative hyperplane class* of the family is $h := \iota^* c_1(\mathcal{O}_{\mathbb{CP}^N}(1)) \in H^2(\mathcal{X}, \mathbb{Z})$, and $h_s \in H^2(X_s, \mathbb{Z})$ denotes its restriction to a fibre: the class of the Kähler form that X_s inherits from the Fubini-Study metric of $\mathbb{CP}^N$. Each fibre is therefore compact Kähler, and its cohomology carries the polarized Hodge structures of chapter 3.

The two families that this chapter computes with are both cut out by a single equation, and the same argument shows in both cases that the projection is a submersion.

Example 5.1.2: Two Smooth Projective Families

Suppose $Z \subseteq \mathbb{CP}^m \times S$ is the zero locus of a holomorphic section G of a line bundle, with $\pi\colon Z \to S$ the projection, and let $p = (x, s) \in Z$ be a point at which the fibre $Z_s \subseteq \mathbb{CP}^m$ is smooth of dimension $m-1$, that is, at which the derivative of G in the $\mathbb{CP}^m$-directions is nonzero. Then $dG_p \neq 0$, so Z is a complex manifold near p with $T_pZ = \ker dG_p$, and π is a submersion at p: otherwise some nonzero $\xi \in T_s^*S$ would annihilate $d\pi(T_pZ)$, so that $\pi^*\xi$ vanishes on $\ker dG_p$ and hence equals $c\,dG_p$ for a scalar c; restricting to the $\mathbb{CP}^m$-directions, where $\pi^*\xi$ vanishes and dG_p does not, forces $c = 0$ and then $\xi = 0$.

1. *The Legendre family.* Let $S = \mathbb{C} \setminus \{0, 1\}$ with coordinate λ, and let $\mathcal{X} \subseteq \mathbb{CP}^2 \times S$ be cut out by
$$G = Y^2Z - X(X - Z)(X - \lambda Z)$$
In the affine chart $Z \neq 0$ the fibre is $y^2 = x(x-1)(x-\lambda)$, whose three roots $0, 1, \lambda$ are distinct for $\lambda \in S$, so each fibre is a smooth plane cubic, and the criterion above makes f a proper submersion. This is a smooth projective family of elliptic curves, $n = 1$, and it is the family computed in example 5.1.9.

2. *The universal smooth hypersurface.* Fix $n \geq 1$ and $d \geq 1$, let $\mathbb{CP}^M$ parametrize the nonzero degree-d forms F in $x_0, \ldots, x_{n+1}$ up to scale, and let $\Delta \subseteq \mathbb{CP}^M$ be the locus of forms whose zero scheme is singular. Put $S = \mathbb{CP}^M \setminus \Delta$ and let $\mathcal{X} \subseteq \mathbb{CP}^{n+1} \times S$ be cut out by $G([x], [F]) = F(x)$. Every fibre is a smooth hypersurface of degree d, so f is again a proper submersion, and S is connected because Δ is a proper closed analytic subset of $\mathbb{CP}^M$. Section 5.4 computes the Hodge filtration of this family.

The fibres of a smooth projective family are all diffeomorphic, and more is true: the family is

locally a product in the C^∞ category. A proper submersion is a locally trivial smooth fibre bundle [32, Ehresmann's theorem], which is the fact definition 4.4.2 was built on. So every $s_0 \in S$ has a neighbourhood U carrying a diffeomorphism $f^{-1}(U) \cong U \times X_{s_0}$ commuting with the projections to U. Cohomology cannot distinguish the fibres over such a U, and the sheaf that records this is the following.

Definition 5.1.3: The Local System of a Family

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and $k \geq 0$. The *local system of the family in degree k* is the higher direct image sheaf

$$\mathbb{H}^k := R^k f_* \mathbb{Q}$$

on S, that is, the sheaf associated to the presheaf $U \mapsto H^k(f^{-1}(U), \mathbb{Q})$ with the restriction maps of singular cohomology [4, higher direct images].

A *local system* of $\mathbb{Q}$-vector spaces on a space S is a sheaf that is locally isomorphic to a constant sheaf with finite-dimensional stalk.[1] The name given in definition 5.1.3 is justified by the following, and the underlying mechanism is exactly the one that produces the fibre-homology local system of a fibration [30, The Fiber-Homology System of a Fibration].

Proposition 5.1.4: Fibre Cohomology Is Locally Constant

Let $f\colon \mathcal{X} \to S$ be a smooth projective family. Then $\mathbb{H}^k$ is a local system of $\mathbb{Q}$-vector spaces on S, and for every $s \in S$ the restriction maps induce an isomorphism $\mathbb{H}^k_s \cong H^k(X_s, \mathbb{Q})$ on stalks. Its rank is the Betti number b_k of a fibre.

***Proof*:**
Fix $s_0 \in S$, write $X = X_{s_0}$, and choose by Ehresmann's theorem an open $U \ni s_0$ together with a diffeomorphism $\Phi\colon f^{-1}(U) \to U \times X$ over U; shrinking U, assume it is contractible. Let $\mathcal{B}$ be the collection of contractible open subsets $V \subseteq U$, a basis for the topology of U because U is an open subset of a manifold.

Let $p\colon f^{-1}(U) \to X$ be Φ followed by the projection $U \times X \to X$, and write $\varphi_s := p|_{X_s}\colon X_s \to X$, a diffeomorphism for every $s \in U$. For $V \in \mathcal{B}$ the restriction of p to $f^{-1}(V) \cong V \times X$ is a homotopy equivalence, V being contractible, so

$$p_V^*\colon H^k(X, \mathbb{Q}) \xrightarrow{\sim} H^k\big(f^{-1}(V), \mathbb{Q}\big)$$

is an isomorphism, and $p_{V'} = p_V$ on $f^{-1}(V')$ for $V' \subseteq V$ gives $\operatorname{res}^V_{V'} \circ p_V^* = p_{V'}^*$. On the basis $\mathcal{B}$ the presheaf is therefore isomorphic to the constant presheaf with value $H^k(X, \mathbb{Q})$, and a sheaf is determined by its values on a basis, so $\mathbb{H}^k|_U$ is the constant sheaf with that value. Since X is compact, $H^k(X, \mathbb{Q})$ is finite dimensional, of dimension b_k.

For the stalks, fix $s \in U$ and let $r_s^V\colon H^k(f^{-1}(V), \mathbb{Q}) \to H^k(X_s, \mathbb{Q})$ be restriction along the inclusion of the fibre, for $V \in \mathcal{B}$ containing s. That inclusion is a homotopy equivalence as well, so r_s^V is an isomorphism, and $r_s^V \circ p_V^* = \varphi_s^*$. The stalk of $\mathbb{H}^k$ at s is the colimit of the $H^k(f^{-1}(V), \mathbb{Q})$, on which the r_s^V are compatible isomorphisms, so the induced map $\mathbb{H}^k_s \to H^k(X_s, \mathbb{Q})$ is an isomorphism, as we sought to show.

[1] [30, Locally Constant Sheaves Are Local Systems].

The rational vector space underlying the Hodge structure of a fibre (example 3.1.2) therefore does not move: over a small U the groups $H^k(X_s, \mathbb{Q})$ for $s \in U$ are canonically identified with one another. The bigrading is another matter. It is not defined over $\mathbb{Q}$, nothing above constrains it, and it will move. The content of this chapter lies in the gap between the two.

The identifications are canonical only locally. Along a loop in S they compose to an automorphism that is generally not the identity, and this is the phenomenon section 4.4 computed by hand for a Lefschetz pencil.

Proposition 5.1.5: Local Systems and Monodromy

Let S be a connected complex manifold with base point s_0.

1. Sending a local system $\mathbb{L}$ of $\mathbb{Q}$-vector spaces on S to the transport of $\mathbb{L}$ along loops is an equivalence between the category of such local systems and the category of representations of $\pi_1(S, s_0)$ on finite-dimensional $\mathbb{Q}$-vector spaces.
2. Let $f\colon \mathcal{X} \to S$ be a smooth projective family. The representation attached in this way to $\mathbb{H}^k$ is $\rho_k \otimes \mathbb{Q}$, where ρ_k is the monodromy representation of definition 4.4.2.

***Proof*:**

1. This is the sheaf form of the local-coefficient dictionary, [30, Locally Constant Sheaves Are Local Systems]. Its hypotheses ask only that S be path connected and locally simply connected, and a connected manifold is both.
2. Let $\gamma\colon [0,1] \to S$ be a loop at s_0 and write $X = X_{s_0}$. The pullback of $\mathcal{X}$ along γ is a smooth fibre bundle over $[0,1]$, hence trivial, and a trivialization amounts to a continuous family of diffeomorphisms $\varphi_t\colon X \to X_{\gamma(t)}$ with $\varphi_0 = \mathrm{id}$; the geometric monodromy of definition 4.4.2 is $h_\gamma = \varphi_1$.

   Fix $\alpha \in H^k(X, \mathbb{Q})$ and consider $t \mapsto (\varphi_t^{-1})^*\alpha \in H^k(X_{\gamma(t)}, \mathbb{Q})$. This is a section of the pullback of $\mathbb{H}^k$ along γ. Indeed, for t in a subinterval J with $\gamma(J)$ contained in an open U as in the proof of proposition 5.1.4, the identifications of the stalks over $\gamma(J)$ with one another are induced by a trivialization of $\mathcal{X}$ over U, and any two trivializations of a bundle over an interval differ by a path of diffeomorphisms of the fibre starting at the identity, so they induce the same isomorphisms on cohomology (the argument recorded after definition 4.4.2). Under those identifications the displayed family is constant on J.

   The transport of $\mathbb{H}^k$ along γ therefore carries α to the value of that section at $t = 1$, namely $(\varphi_1^{-1})^*\alpha = (h_\gamma^{-1})^*\alpha$. That is the action of $\rho_k(\gamma)$, read on rational rather than integral cohomology, completing the proof.

For a Lefschetz pencil, part (2) turns theorem 4.4.7 into a statement about $\mathbb{H}^n$, the middle degree of an n-dimensional fibre: the local monodromy at a critical value acts on the fibre cohomology by the Picard-Lefschetz transformation along the vanishing cycle, a transvection when n is odd and a reflection of order two when n is even, and corollary 4.4.9 identifies the classes on which the whole group acts trivially. Those are the classes that extend to a global section of the local system, and the question of whether such a section can be recognized Hodge-theoretically is one of the questions chapter 6 takes up.

A local system is a topological object, and the base S is a complex manifold whose complex structure it ignores. To bring that structure to bear, tensor the local system with the holomorphic functions.

Definition 5.1.6: The Cohomology Bundle and the Gauss-Manin Connection

Let $f\colon \mathcal{X} \to S$ be a smooth projective family. The *cohomology bundle* of the family in degree k is the $\mathcal{O}_S$-module

$$\mathcal{H}^k := \mathbb{H}^k \otimes_{\mathbb{Q}} \mathcal{O}_S$$

which is locally free of rank b_k, since an isomorphism of $\mathbb{H}^k|_U$ with the constant sheaf $\underline{\mathbb{Q}}^{b_k}$ induces $\mathcal{H}^k|_U \cong \mathcal{O}_U^{b_k}$.
A *holomorphic connection* on a locally free $\mathcal{O}_S$-module $\mathcal{E}$ is a map of sheaves of $\mathbb{C}$-vector spaces $\nabla\colon \mathcal{E} \to \mathcal{E} \otimes_{\mathcal{O}_S} \Omega^1_S$ satisfying the Leibniz rule

$$\nabla(g\sigma) = g\,\nabla\sigma + \sigma \otimes dg$$

for local holomorphic g and local sections σ of $\mathcal{E}$. This is the holomorphic version of a connection on a vector bundle [12, Connection on a vector bundle], with $\mathcal{O}_S$ in place of the smooth functions and Ω^1_S in place of the smooth cotangent bundle.
The *Gauss-Manin connection* ∇ on $\mathcal{H}^k$ is defined as follows. Let $U \subseteq S$ be connected with $\mathbb{H}^k|_U$ constant and let $e_1, \dots, e_{b_k}$ be a $\mathbb{Q}$-basis of $\mathbb{H}^k(U)$, so that every section of $\mathcal{H}^k$ over an open $V \subseteq U$ is uniquely $\sum_i g_i e_i$ with $g_i \in \mathcal{O}_S(V)$. Set

$$\nabla\Big(\sum_i g_i e_i\Big) := \sum_i e_i \otimes dg_i$$

Theorem 5.1.7 shows that this is independent of the basis chosen and that the local definitions glue. For a holomorphic vector field u on U, write $\nabla_u\sigma$ for the contraction of $\nabla\sigma$ with u in the Ω^1_S factor.
The same formula defines ∇ on the underlying C^∞ bundle. Write $A^0(\mathcal{H}^k)$ for the sheaf of smooth sections and extend by $\nabla(\sum_i g_i e_i) := \sum_i e_i \otimes dg_i$ for smooth g_i, with d now the full exterior derivative. This is a connection on a smooth complex vector bundle in the sense of [12, Connection on a vector bundle]; it agrees with the holomorphic connection on holomorphic sections, and its $(0,1)$-part $\nabla^{0,1}$ is the $\bar\partial$-operator of the holomorphic structure of $\mathcal{H}^k$, since a smooth section is holomorphic exactly when its coefficients g_i in a flat frame are. Both readings of ∇ are used: the holomorphic one states flatness and Griffiths transversality most cleanly, while the smooth one is what section 5.2 differentiates in a C^∞ trivialization.

Evaluating a section at a point recovers a cohomology class of the fibre there: in the frame above, $\mathcal{H}^k \otimes_{\mathcal{O}_S} \mathbb{C}(s) \cong \mathbb{H}^k_s \otimes_{\mathbb{Q}} \mathbb{C} \cong H^k(X_s, \mathbb{C})$, using proposition 5.1.4 and the fact that $H^k(X_s, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} \cong H^k(X_s, \mathbb{C})$. So $\mathcal{H}^k$ is a holomorphic vector bundle whose fibre over s is the complex cohomology of X_s, and the rest of the chapter works on it.

Theorem 5.1.7: The Gauss-Manin Connection Is Well Defined and Flat

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and $k \geq 0$.

1. The local recipe of definition 5.1.6 is independent of the chosen basis and the local definitions agree on overlaps, so they glue to a holomorphic connection ∇ on $\mathcal{H}^k$. It is the unique holomorphic connection on $\mathcal{H}^k$ that annihilates every local section of $\mathbb{H}^k$.
2. $\ker \nabla = \mathbb{H}^k \otimes_{\mathbb{Q}} \mathbb{C}$, the complexification of the local system, viewed as a subsheaf of $\mathcal{H}^k$.
3. ∇ is flat: extending it to $\nabla\colon \mathcal{H}^k \otimes \Omega_S^p \to \mathcal{H}^k \otimes \Omega_S^{p+1}$ by $\nabla(\sigma \otimes \alpha) = \nabla\sigma \wedge \alpha + \sigma \otimes d\alpha$, where $\nabla\sigma \wedge \alpha := \sum_i \tau_i \otimes (\beta_i \wedge \alpha)$ for $\nabla\sigma = \sum_i \tau_i \otimes \beta_i$, the curvature $\nabla \circ \nabla\colon \mathcal{H}^k \to \mathcal{H}^k \otimes \Omega_S^2$ is zero.

***Proof*:**

1. Fix U connected with $\mathbb{H}^k|_U$ constant, which is possible by proposition 5.1.4, and let (e_i) be a $\mathbb{Q}$-basis of $\mathbb{H}^k(U)$. The recipe satisfies the Leibniz rule, because $d(gg_i) = g\,dg_i + g_i\,dg$ gives
$$\nabla\Big(g\sum_i g_i e_i\Big) = \sum_i e_i \otimes d(gg_i) = g\sum_i e_i \otimes dg_i + \Big(\sum_i g_i e_i\Big) \otimes dg$$
and it annihilates the sections of $\mathbb{H}^k$ over any $V \subseteq U$, since such a section has locally constant coefficients g_i.

   Let (e'_j) be a second $\mathbb{Q}$-basis of $\mathbb{H}^k(U)$, say $e'_j = \sum_i c_{ij} e_i$ with $c_{ij} \in \mathbb{Q}$. A section written $\sum_j h_j e'_j$ has coefficients $g_i = \sum_j c_{ij} h_j$ in the first basis, and since the c_{ij} are constants,
$$\sum_i e_i \otimes dg_i = \sum_i e_i \otimes \sum_j c_{ij}\,dh_j = \sum_j \Big(\sum_i c_{ij} e_i\Big) \otimes dh_j = \sum_j e'_j \otimes dh_j$$
so the two recipes agree. If U and U' are two such opens and W is a connected component of $U \cap U'$, then restriction identifies both $\mathbb{H}^k(U)$ and $\mathbb{H}^k(U')$ with $\mathbb{H}^k(W)$, because each is identified with a stalk; the two frames therefore restrict to two $\mathbb{Q}$-bases of $\mathbb{H}^k(W)$, and the computation just made shows the two connections agree on W, hence on $U \cap U'$. The local definitions glue.

   Uniqueness is the Leibniz rule read backwards: a connection annihilating the e_i must send $\sum_i g_i e_i$ to $\sum_i e_i \otimes dg_i$.

2. Work over U with a frame as above. A section $\sum_i g_i e_i$ is killed by ∇ if and only if every dg_i vanishes, that is, if and only if every g_i is locally constant. The sections with locally constant complex coefficients in a flat frame are exactly the local sections of $\mathbb{H}^k \otimes_{\mathbb{Q}} \mathbb{C}$.

3. The curvature is computed in a flat frame. For $\sigma = \sum_i g_i e_i$,
$$\nabla(\nabla\sigma) = \nabla\Big(\sum_i e_i \otimes dg_i\Big) = \sum_i \big(\nabla e_i \wedge dg_i + e_i \otimes d(dg_i)\big) = 0$$
since $\nabla e_i = 0$ by part (1) and $d \circ d = 0$. As the flat frames cover S, the curvature vanishes identically, completing the proof.

Flatness is what makes the connection usable, and part (2) says how. A section of $\mathcal{H}^k$ is flat exactly when it is a locally constant family of cohomology classes, so continuing a flat section along a path in S is the same operation as transporting a class in the local system; by proposition 5.1.5 the resulting automorphism of $H^k(X_{s_0}, \mathbb{C})$ attached to a loop is $\rho_k(\gamma) \otimes \mathbb{C}$. The connection and the monodromy representation are two presentations of one object.

Part (3) also says something about $\mathcal{H}^k$ that is not automatic. A holomorphic vector bundle rarely admits a holomorphic connection at all, since the obstruction to constructing one is a cohomology class that need not vanish; the bundle $\mathcal{H}^k$ admits one because it was manufactured from a local system, and having done so it is flat for the same reason. What the converse direction would require, namely that a flat holomorphic connection always comes from a local system, is the Riemann-Hilbert correspondence, and it is not needed here: the local system came first.

For computations one needs a way to test a class against something, and the test is integration over cycles. The dual local system $(\mathbb{H}^k)^\vee$ has stalk $H^k(X_s, \mathbb{Q})^\vee \cong H_k(X_s, \mathbb{Q})$ because cohomology with field coefficients is dual to homology,[2] so a section of it over U is a locally constant family $s \mapsto c_s \in H_k(X_s, \mathbb{Q})$ of homology classes, and the pairing with a class $\alpha \in H^k(X_s, \mathbb{C})$ is the period $\langle c_s, \alpha \rangle = \int_{c_s} \alpha$. Call such a family a *flat family of cycles*.

Proposition 5.1.8: Periods Compute the Gauss-Manin Connection

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and let $U \subseteq S$ be connected and simply connected, so that $\mathbb{H}^k|_U$ is constant. For a flat family of cycles c over U write $\langle c, - \rangle$ for the induced pairing, applied to the $\mathcal{H}^k$ factor when the argument lies in $\mathcal{H}^k \otimes \Omega^1_S$.

1. An assignment $s \mapsto \alpha_s$ of a class $\alpha_s \in H^k(X_s, \mathbb{C})$ to each $s \in U$ is a holomorphic section of $\mathcal{H}^k$ if and only if $s \mapsto \langle c_s, \alpha_s \rangle$ is holomorphic on U for every flat family of cycles c.
2. For every section σ of $\mathcal{H}^k$ over U and every flat family of cycles c,
$$\langle c, \nabla \sigma \rangle = d \langle c, \sigma \rangle \tag{5.1}$$
3. A section of $\mathcal{H}^k$ over U is determined by the functions $\langle c, \sigma \rangle$ as c runs over the flat families of cycles, and $\nabla \sigma = 0$ if and only if all of them are constant.

***Proof*:**

Since U is connected and simply connected, the monodromy representation of $\mathbb{H}^k|_U$ is trivial, so $\mathbb{H}^k|_U$ is constant by proposition 5.1.5. Fix a $\mathbb{Q}$-basis $e_1, \dots, e_r$ of $\mathbb{H}^k(U)$ and let $c^1, \dots, c^r$ be the dual basis of flat families of cycles, so that $\langle c^i, e_j \rangle = \delta_{ij}$ is constant. Every flat family of cycles over U is a $\mathbb{Q}$-linear combination of the c^i with constant coefficients, and a section σ of $\mathcal{H}^k$ over U is uniquely $\sum_i g_i e_i$ with $g_i = \langle c^i, \sigma \rangle$.

1. Each α_s lies in the fibre of $\mathcal{H}^k$ at s, so the assignment has coefficient functions $g_i(s) = \langle c^i_s, \alpha_s \rangle$ in the frame (e_i), and it is a holomorphic section exactly when every g_i is holomorphic. If all periods are holomorphic then in particular the g_i are; conversely, if the g_i are holomorphic then so is $\langle c, \alpha \rangle = \sum_i a_i g_i$ for any flat $c = \sum_i a_i c^i$ with a_i constant.

[2] [30, Cohomology with Field Coefficients is the Dual of Homology].

2. With $\sigma = \sum_i g_i e_i$ one has $\nabla\sigma = \sum_i e_i \otimes dg_i$, so for c flat,

$$\langle c, \nabla\sigma\rangle = \sum_i \langle c, e_i\rangle\, dg_i = d\Big(\sum_i \langle c, e_i\rangle\, g_i\Big) = d\,\langle c, \sigma\rangle$$

the middle equality because each $\langle c, e_i\rangle$ is a constant.

3. The functions $\langle c^i, \sigma\rangle = g_i$ determine σ, and by (5.1) they are all constant if and only if every dg_i vanishes, which by theorem 5.1.7 is the condition $\nabla\sigma = 0$, as we sought to show.

Part (2) says that differentiating a period differentiates the class. It converts a question about ∇ into a question about ordinary derivatives of integrals depending on a parameter, and the following example carries out that conversion in full for the family of example 5.1.2.

Example 5.1.9: Gauss-Manin for the Legendre Family

Let $f\colon \mathcal{X} \to S = \mathbb{C}\setminus\{0,1\}$ be the Legendre family, whose fibre E_λ is the smooth projective cubic with affine equation $y^2 = P(x)$, where

$$P(x) = x(x-1)(x-\lambda) = q(x)\,(x-\lambda), \qquad q(x) := x(x-1)$$

Each fibre is a compact Riemann surface of genus 1, so $b_1 = 2$ and $\mathcal{H}^1$ is a rank-two holomorphic bundle on S.

A holomorphic section. The form $\omega_\lambda := dx/y$ is a holomorphic 1-form on E_λ with no zeros. Away from $y = 0$ and from the point at infinity this is clear; at a point with $y = 0$ the coordinate y is a local parameter and $2y\,dy = P'(x)\,dx$ gives $dx/y = 2\,dy/P'(x)$ with $P'(x) \neq 0$ there; and at infinity, in the chart $u = X/Y$, $v = Z/Y$, the curve reads $v = u(u-v)(u-\lambda v)$, so $v = u^3(1+O(u))$ with u a local parameter, and $dx/y = (v\,du - u\,dv)/v = (-2+O(u))\,du$. Being closed, ω_λ has a class in $H^1(E_\lambda, \mathbb{C})$.

Flat cycles as contours. Fix $\lambda_0 \in S$. Every class in $H_1(E_{\lambda_0}, \mathbb{Q})$ is a $\mathbb{Q}$-combination of classes of loops avoiding the four branch points of the double cover $E_{\lambda_0} \to \mathbb{C}\mathrm{P}^1$, and such a loop σ projects to a loop Γ in $\mathbb{C}\setminus\{0,1,\lambda_0\}$ of which it is a lift, the projection being a covering map there. Choose a disc $U \ni \lambda_0$ disjoint from Γ. Then $P(x)$ is nonvanishing for x near Γ and $\lambda \in U$, and the branch of $y = \sqrt{P}$ along σ continues to a branch $y(x,\lambda)$ holomorphic near $\Gamma \times U$, since the winding number of P along Γ is even, as it must be for σ to close up. The resulting lifts $c_\lambda \subseteq E_\lambda$ form a flat family of cycles over U with $c_{\lambda_0} = \sigma$, and

$$\pi(\lambda) := \int_{c_\lambda} \omega_\lambda = \int_\Gamma \frac{dx}{y(x,\lambda)}$$

is holomorphic on U, the contour being fixed and the integrand holomorphic in λ uniformly on the compact set Γ. Since $H_1(E_{\lambda_0}, \mathbb{Q})$ is spanned by finitely many such loop classes and flat continuation is linear, after shrinking U every flat family of cycles over U is a $\mathbb{Q}$-combination of families of this shape. So proposition 5.1.8(1) makes $\omega\colon \lambda \mapsto [\omega_\lambda]$ a holomorphic section of $\mathcal{H}^1$ over U, and the local descriptions patch because they all assign to λ the class of dx/y.

Differentiating. Differentiation under the integral sign is legitimate for the same reason, and $\partial_\lambda P = -q$ gives $\partial_\lambda y = -q/2y$, hence

$$\partial_\lambda \frac{1}{y} = \frac{q}{2y^3}, \qquad \partial_\lambda^2 \frac{1}{y} = \partial_\lambda\Big(\frac{q}{2}y^{-3}\Big) = -\frac{3q}{2}y^{-4}\cdot\frac{-q}{2y} = \frac{3q^2}{4y^5}$$

So by (5.1), applied twice,

$$\langle c, \nabla_{\partial_\lambda}\omega\rangle = \int_\Gamma \frac{q\,dx}{2y^3}, \qquad \langle c, \nabla^2_{\partial_\lambda}\omega\rangle = \int_\Gamma \frac{3q^2\,dx}{4y^5}$$

One exact form. Put $A := q^2/2$ and consider the rational function $R := A/y^3$ on E_λ. Using $2y\,dy = P'\,dx$ and $y^2 = P$,

$$dR = \Big(\frac{A'}{y^3} - \frac{3AP'}{2y^5}\Big)dx = \frac{2A'P - 3AP'}{2y^5}\,dx$$

Now $A' = qq'$, $P = q(x-\lambda)$ and $P' = q'(x-\lambda) + q$, so

$$2A'P - 3AP' = 2q^2q'(x-\lambda) - \frac{3q^2}{2}\big(q'(x-\lambda)+q\big) = \frac{q^2}{2}\big(q'(x-\lambda) - 3q\big)$$

and $q'(x-\lambda) - 3q = (2x-1)(x-\lambda) - 3x^2 + 3x = -x^2 + 2(1-\lambda)x + \lambda$, which gives

$$dR = \frac{q^2}{4y^5}\big(-x^2 + 2(1-\lambda)x + \lambda\big)\,dx \tag{5.2}$$

The same expression arises from the three period integrands in play: those of $\nabla^2_{\partial_\lambda}\omega$ and $\nabla_{\partial_\lambda}\omega$ just computed, together with $1/y$ for ω itself. Combining them with the coefficients $\lambda(1-\lambda)$, $1-2\lambda$ and $-\frac14$ and using $y^2 = P = q(x-\lambda)$ to put everything over y^5,

$$\frac{3\lambda(1-\lambda)q^2}{4y^5} + \frac{(1-2\lambda)q}{2y^3} - \frac{1}{4y} = \frac{q^2}{4y^5}\Big(3\lambda(1-\lambda) + 2(1-2\lambda)(x-\lambda) - (x-\lambda)^2\Big)$$

and substituting $x = t + \lambda$ turns the last bracket into $-t^2 + (2-4\lambda)t + 3\lambda - 3\lambda^2$, which is the bracket of (5.2) written in t.

The Picard-Fuchs equation. The only pole of R on E_λ is the point $(\lambda, 0)$: at $(0,0)$ and $(1,0)$ the numerator q^2 vanishes to order 4 in the local parameter y while y^3 vanishes to order 3, and at infinity R has a zero, since q^2 has a pole of order 8 there and y^3 one of order 9. The contour Γ avoids $x = \lambda$, so c_λ avoids that pole and $\int_{c_\lambda} dR = 0$. Combining the computed periods with (5.2),

$$\big\langle c,\ \lambda(1-\lambda)\nabla^2_{\partial_\lambda}\omega + (1-2\lambda)\nabla_{\partial_\lambda}\omega - \tfrac14\omega\big\rangle = \int_{c_\lambda} \frac{q^2\big(-x^2 + 2(1-\lambda)x + \lambda\big)}{4y^5}\,dx = \int_{c_\lambda} dR = 0$$

for every flat family of cycles, so by proposition 5.1.8(3) the section itself vanishes:

$$\lambda(1-\lambda)\,\nabla^2_{\partial_\lambda}\omega + (1-2\lambda)\,\nabla_{\partial_\lambda}\omega - \tfrac14\,\omega = 0 \tag{5.3}$$

Pairing (5.3) with a flat family of cycles and using (5.1) shows that every period $\pi(\lambda) = \int_{c_\lambda}\omega_\lambda$ satisfies the scalar equation

$$\lambda(1-\lambda)\,\pi'' + (1-2\lambda)\,\pi' - \tfrac14\,\pi = 0$$

the *Picard-Fuchs equation* of the family. It is the hypergeometric equation with parameters $\frac12, \frac12; 1$, and exercise 5.1.1 (5) produces a solution as a power series.

Where the monodromy is. The equation is singular exactly at $\lambda = 0, 1, \infty$, the values excluded from S, and at each of them the fibre degenerates: at $\lambda = 0$ the cubic $y^2 = x^2(x-1)$ has a node

at the origin, the branch points 0 and λ having collided. Near that node the family is a smoothing. Writing $w := (x - \lambda/2)\sqrt{1-x}$ for the branch of the square root equal to 1 at $x = 0$, which for λ small is a holomorphic change of coordinate near $x = 0$ since $\partial w/\partial x = 1 + \lambda/4 \neq 0$ there, the identity $x(x-1)(x-\lambda) = (x-1)(x-\lambda/2)^2 - (x-1)\lambda^2/4$ turns the equation of the fibre into

$$y^2 + w^2 = \frac{\lambda^2}{4}\left(1 - x(w,\lambda)\right)$$

The smoothing parameter of the node is $\lambda^2/4$ rather than λ, and theorem 4.4.7 attaches one transvection $T(\alpha) = \alpha - \langle\alpha,\delta\rangle\delta$ along the vanishing cycle to each full turn of that parameter, as in example 4.4.8. One turn of λ about 0 turns $\lambda^2/4$ twice, so the local monodromy of the Legendre family at $\lambda = 0$ is T^2.[a] On the period side this is the assertion that continuing a solution of the Picard-Fuchs equation once around $\lambda = 0$ returns a different solution, so no basis of solutions there consists of single-valued functions, and the second solution carries a logarithm.

[a] The classical form of this computation says that the image of the monodromy of the Legendre family in $\mathrm{PSL}_2(\mathbb{Z})$ is the level-two principal congruence subgroup, of which λ is the modular function; see [6].

The example runs the whole apparatus of this section at the level of $\mathbb{H}^1$ and ∇. A family of varieties produced a flat bundle; a geometrically defined family of classes produced a section of it; and flatness turned a statement about that section into a linear differential equation whose monodromy is the topological monodromy of the family. What the section does not record is that ω_λ is not an arbitrary class but a holomorphic form, so that it lies in the first step of the Hodge filtration of the fibre.

Exercise 5.1.1

1. Let $f\colon \mathcal{X} \to S$ be a smooth projective family with connected fibres of dimension n. Show that $\mathbb{H}^0$ is the constant sheaf $\mathbb{Q}$ and that under the resulting identification $\mathcal{H}^0 = \mathcal{O}_S$ the Gauss-Manin connection is $\nabla = d$. Then show that $\mathbb{H}^{2n}$ has trivial monodromy, by exhibiting the class $\eta_s \in H^{2n}(X_s, \mathbb{Q})$ with $\langle \eta_s, [X_s]\rangle = 1$ as a global flat section, where $[X_s]$ is the fundamental class for the complex orientation. (For the flatness, compare orientations along a local trivialization normalized to be the identity at one point.)

2. Let X be a smooth projective variety, S a connected complex manifold, and $\mathcal{X} = X \times S$ with f the projection. Show that $\mathbb{H}^k$ is the constant sheaf with value $H^k(X, \mathbb{Q})$, that the monodromy representation is trivial, and that ∇ is $1 \otimes d$ under the resulting identification $\mathcal{H}^k = H^k(X, \mathbb{C}) \otimes_{\mathbb{C}} \mathcal{O}_S$. Deduce that a family whose monodromy representation in some degree is nontrivial admits no fibre-preserving diffeomorphism onto a product family, and check that the Legendre family of example 5.1.9 is such a family.

3. Let $f\colon \mathcal{X} \to S$ be a smooth projective family with S connected and let $s_0 \in S$. Show that evaluation at s_0 is an isomorphism from the space of global flat sections of $\mathcal{H}^k$ onto the subspace of $H^k(X_{s_0}, \mathbb{C})$ fixed by the monodromy representation. Deduce that a global flat section vanishing at a single point is identically zero, and that $\dim_{\mathbb{C}} \Gamma(S, \ker\nabla) \leq b_k$ with equality if and only if the monodromy representation is trivial.

4. Fibrewise cup product defines a map $\mathcal{H}^k \otimes_{\mathcal{O}_S} \mathcal{H}^l \to \mathcal{H}^{k+l}$.

    1. Show that it is flat for the Gauss-Manin connections, that is, $\nabla(\sigma \cup \tau) = \nabla\sigma \cup \tau + \sigma \cup \nabla\tau$.
    2. Show that the relative hyperplane class of definition 5.1.1 gives a global flat section h of $\mathbb{H}^2$. Combining this with (1) and exercise 5.1.1 (1), show that for $0 \leq k \leq n$, where n is

the dimension of the fibres, the bilinear form

$$Q_s(\alpha,\beta) = (-1)^{k(k-1)/2}\int_{X_s}\alpha\wedge\beta\wedge h_s^{n-k}$$

of proposition 3.2.9 is flat, in the sense that $d\,Q(\sigma,\tau) = Q(\nabla\sigma,\tau)+Q(\sigma,\nabla\tau)$ for sections σ,τ of $\mathcal{H}^k$.

5. Write $\pi(\lambda)=\sum_{m\geq 0}a_m\lambda^m$ and show that the Picard-Fuchs equation of example 5.1.9 is equivalent to the recursion $a_{m+1}(m+1)^2 = a_m\big(m+\frac{1}{2}\big)^2$. Verify that $a_m=\binom{2m}{m}^2 16^{-m}$ satisfies it, and determine the radius of convergence of the resulting solution. Why does the existence of this single-valued solution not contradict the last paragraph of example 5.1.9?

6. Let $f\colon\mathcal{X}\to S$ be a smooth projective family, let $\psi\colon S'\to S$ be a holomorphic map from a connected complex manifold, and let $\mathcal{X}'=\mathcal{X}\times_S S'$ with f' the projection. Show that $\mathcal{X}'$ is again a smooth projective family, that $\mathbb{H}'^k=\psi^{-1}\mathbb{H}^k$, and that ∇' is the pullback of ∇, meaning that $\nabla'(\psi^*\sigma)=(\psi^*\otimes\psi^*)(\nabla\sigma)$ for every local section σ of $\mathcal{H}^k$. Deduce that the periods of $\psi^*\sigma$ are the periods of σ composed with ψ.

5.2 Griffiths Transversality

The bundle $\mathcal{H}^k$ of section 5.1 carries a flat connection, and flatness makes it a topological object: over a simply connected base the Gauss-Manin connection identifies the fibres $H^k(X_s,\mathbb{C})$ with one another, and no feature of the complex structure of an individual fibre survives that identification. Each fibre is nonetheless a compact Kähler manifold, so each $H^k(X_s,\mathbb{C})$ carries a Hodge filtration, and under the flat identification those filtrations move.

Two facts govern the motion, and this section proves both. The filtration varies holomorphically, so the subspaces $F^pH^k(X_s,\mathbb{C})$ are the fibres of holomorphic subbundles of $\mathcal{H}^k$; and the connection carries each of those subbundles into the next one down, never further. The second statement is Griffiths transversality, and it is where the geometry of a family constrains the linear algebra of chapter 3.

Throughout, $f\colon\mathcal{X}\to S$ is a smooth projective family (definition 5.1.1) with fibres $X_s=f^{-1}(s)$, and $\mathbb{H}^k$, $\mathcal{H}^k$, ∇ are the local system of definition 5.1.3 together with the cohomology bundle and the Gauss-Manin connection of definition 5.1.6. One consequence of flatness (theorem 5.1.7) is used repeatedly below: a local frame of flat sections is a local holomorphic frame, so $\nabla^{0,1}$ is the $\bar\partial$-operator of the holomorphic structure on $\mathcal{H}^k$.

Projectivity enters in one way only. Each fibre is compact Kähler (definition 5.1.1) because the embedding into $\mathbb{C}\mathrm{P}^N\times S$ restricts the Fubini-Study form to a Kähler form κ_s on it (theorem 1.1.7); what matters below is that these forms vary smoothly with s, being restrictions of one fixed form on $\mathbb{C}\mathrm{P}^N$. So theorem 2.1.1 applies to every fibre and $H^k(X_s,\mathbb{Z})$ modulo torsion carries a pure Hodge structure of weight k (example 3.1.2). That structure is the object whose variation is to be measured, and by theorem 3.1.6 it is equivalent to record it by its Hodge filtration rather than by its bigrading. The filtration is the presentation that behaves well in a family, and the following definition collects the subspaces it produces.

Definition 5.2.1: The Hodge Bundles of a Family

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and fix $k \geq 0$. For each $p \in \mathbb{Z}$ let

$$F^p\mathcal{H}^k \subseteq \mathcal{H}^k$$

be the subset whose intersection with the fibre $H^k(X_s, \mathbb{C})$ over $s \in S$ is the step $F^pH^k(X_s, \mathbb{C})$ of the Hodge filtration of definition 3.1.5, that is $\bigoplus_{r \geq p} H^{r,k-r}(X_s)$. These subsets are the *Hodge bundles* of the family. They are nested,

$$\mathcal{H}^k = F^0\mathcal{H}^k \supseteq F^1\mathcal{H}^k \supseteq \cdots \supseteq F^k\mathcal{H}^k \supseteq F^{k+1}\mathcal{H}^k = 0$$

the outer equalities holding because the Hodge structure on the cohomology of a compact Kähler manifold is effective.

Calling these bundles is an anticipation: nothing said so far rules out the fibre dimension jumping from point to point, and even with constant dimension a family of subspaces need not vary holomorphically. Both are settled below. The smallest case is worth having in hand first, since there the subspaces are lines and can be written down.

Example 5.2.2: The Hodge Bundles of a Family of Curves

Let $f\colon \mathcal{X} \to S$ be a smooth projective family whose fibres are compact Riemann surfaces of genus g, and take $k = 1$. By example 3.2.3 each $H^1(X_s, \mathbb{C})$ splits as $H^{1,0}(X_s) \oplus H^{0,1}(X_s)$ with $H^{1,0}(X_s) \cong H^0(X_s, \Omega^1_{X_s})$ of dimension g. The Hodge bundles are therefore

$$\mathcal{H}^1 = F^0\mathcal{H}^1 \supseteq F^1\mathcal{H}^1 \supseteq F^2\mathcal{H}^1 = 0$$

with $\mathcal{H}^1$ of rank $2g$ and $F^1\mathcal{H}^1$ of rank g, its fibre at s being the space of holomorphic 1-forms on X_s.

For the Legendre family this is completely explicit. Let $S = \mathbb{C} \setminus \{0, 1\}$ and let $\mathcal{X} \subseteq \mathbb{C}\mathrm{P}^2 \times S$ be the family of projective plane cubics

$$y^2z = x(x - z)(x - \lambda z), \qquad \lambda \in S$$

which is smooth for every $\lambda \neq 0, 1$, so each fibre X_λ is an elliptic curve and $g = 1$. Then $\mathcal{H}^1$ has rank 2 and $F^1\mathcal{H}^1$ has rank 1: it is the line bundle whose fibre at λ is spanned by the class of the holomorphic 1-form

$$\omega_\lambda = \frac{dx}{y}$$

in the affine chart $z = 1$. The Gauss-Manin derivative of this section is computed in example 5.1.9, and example 5.2.9 returns to what transversality does, and does not, say about it.

Both open questions about definition 5.2.1, constancy of the fibre dimension and holomorphy, are questions about how harmonic forms behave when the complex structure of the fibre moves. On a single compact Kähler manifold the harmonic theory of chapter 2 came from applying the elliptic package to $\Delta_{\bar\partial}$; in a family one needs the corresponding statements for the whole family of operators at once, and those belong to the theory of families of elliptic operators, which neither this book nor its prerequisites develop. Chapter 2 already imports the single-manifold package from *Riemannian*

Geometry.[3] The family version is imported here, and it is the only analytic input this section takes from outside.

Two conventions before the statement. The harmonic spaces of chapter 2 always carry a manifold in their argument, as in $\mathcal{H}^{p,q}(X_s)$, so no confusion arises with the bundle $\mathcal{H}^k$. And a *smooth family of forms on the fibres* means a smooth section of the bundle $\bigwedge^k T^*_{\mathcal{X}/S}$ of relative k-forms on $\mathcal{X}$, whose restriction to X_s is a k-form β_s on X_s; such a family has *relative type* (r,q) when each β_s is of type (r,q) on X_s.

Proposition 5.2.3: Hodge Numbers and Harmonic Projections in a Family

Let $f\colon \mathcal{X} \to S$ be a smooth projective family with S connected, and equip each fibre with the Kähler form κ_s above. Write Π_s for the projection of the k-forms on X_s onto the harmonic ones determined by κ_s, which by theorem 1.3.1 agrees in each bidegree with the $\bar\partial$-harmonic projection of theorem 1.1.10.

1. Each Hodge number $h^{p,q}(X_s)$ is independent of s.
2. If $\{\beta_s\}$ is a smooth family of k-forms on the fibres, then $\{\Pi_s(\beta_s)\}$ is again a smooth family.

***Proof Key ideas*:**
The two ingredients below are theorems about a smooth family of formally self-adjoint elliptic operators, proved by the perturbation theory of such families. That theory is not developed in this book, and the deduction of (1) and (2) from it is what is given here; for the ingredients themselves see [38, families of elliptic operators and the smooth variation of their harmonic projectors]. This is the one place in the chapter where an input comes from outside the series, and it is taken because no EYNTKA volume develops the perturbation theory of families of elliptic operators: [12] gives the elliptic package for a *fixed* metric, which is what chapter 2 ran on, and [9] is not upstream of this book.

1. By theorem 1.1.10 each $\Delta_{\bar\partial,s}$ is a formally self-adjoint elliptic operator of order two. In a C^∞ trivialization of f over a ball these operators pull back to operators on one fixed manifold, and the resulting family varies smoothly with s because κ_s does. The first ingredient is that $s \mapsto \dim\ker\Delta_{\bar\partial,s}$ is upper semicontinuous: an eigenvalue may leave 0 under a small perturbation, but the spectral gap above 0 prevents one from arriving at 0. Since $\ker\Delta_{\bar\partial,s} = \mathcal{H}^{p,q}_{\bar\partial}(X_s)$ has dimension $h^{p,q}(X_s)$ by theorem 1.1.10, each $h^{p,q}$ is upper semicontinuous.

   Their sum is constant. By corollary 2.1.3, $\sum_{p+q=k} h^{p,q}(X_s) = b_k(X_s)$, and $b_k(X_s) = \dim \mathbb{H}^k_s \otimes \mathbb{C}$ is independent of s because $\mathbb{H}^k$ is a local system on the connected space S (definition 5.1.3). A finite collection of upper semicontinuous integer-valued functions with constant sum is locally constant: near s_0 every term satisfies $h^{p,q}(s) \le h^{p,q}(s_0)$ while the totals agree, so every inequality is an equality. Local constancy on a connected space is constancy.

2. Let $\Delta_s = 2\Delta_{\bar\partial,s}$ be the Laplacian of κ_s on k-forms, which preserves the bidegree by theorem 1.3.1, so that part (1) makes $\dim\ker\Delta_s$ constant as well. The harmonic projection

[3] [12, Hodge decomposition].

is then the spectral projection

$$\Pi_s = \frac{1}{2\pi i} \oint_{|\lambda|=\epsilon} (\lambda - \Delta_s)^{-1}\, d\lambda$$

for any ϵ below the first nonzero eigenvalue, which by constancy of the kernel dimension may be chosen locally uniformly in s. The second ingredient is that the resolvent of a smooth family of elliptic operators depends smoothly on the parameter away from the spectrum, so the contour integral does as well, and applying Π_s to a smooth family of forms returns a smooth family, as we sought to show.

The proposition is a statement about metrics, and both theorems to come are statements about the filtration, which is metric-free. The bridge between them is a single observation about which classes are represented by which forms. Write

$$F^p\mathcal{A}^k(X) := \bigoplus_{r\geq p} \mathcal{A}^{r,k-r}(X)$$

for the k-forms on a complex manifold X all of whose components have holomorphic degree at least p. This is a filtration of the forms, and the question is how it compares with the filtration of definition 3.1.5 on cohomology.

Lemma 5.2.4: Filtration Level of a Closed Form

Let X be a compact Kähler manifold and $\alpha \in \mathcal{A}^k(X)$ a closed k-form. If $\alpha \in F^p\mathcal{A}^k(X)$ then $[\alpha] \in F^pH^k(X,\mathbb{C})$. Conversely, every class in $F^pH^k(X,\mathbb{C})$ is represented by a harmonic form lying in $F^p\mathcal{A}^k(X)$.

***Proof*:**
Fix a Kähler metric on X and let Π be the associated harmonic projection. By theorem 1.3.1 the Laplacian $\Delta_d = 2\Delta_{\bar\partial}$ preserves the bidegree, so its kernel is the direct sum of the spaces $\mathcal{H}^{r,k-r}(X)$ of harmonic forms of pure type, and the orthogonal decomposition of theorem 1.1.10 splits along bidegrees. Consequently Π commutes with the projection of a form onto its $(r,k-r)$-component, and $\Pi\big(\mathcal{A}^{r,k-r}(X)\big) \subseteq \mathcal{H}^{r,k-r}(X)$.

Let α be closed with $\alpha = \sum_{r\geq p}\alpha^{r,k-r}$. Every de Rham class has a unique harmonic representative, and for a closed form that representative is its harmonic projection [12, The Hodge theorem], so $[\alpha] = [\Pi(\alpha)]$. By the previous paragraph,

$$\Pi(\alpha) = \sum_{r\geq p} \Pi\big(\alpha^{r,k-r}\big), \qquad \Pi\big(\alpha^{r,k-r}\big) \in \mathcal{H}^{r,k-r}(X)$$

The class of a harmonic form of type $(r,k-r)$ lies in $H^{r,k-r}(X)$ by theorem 2.1.1, so $[\alpha] \in \bigoplus_{r\geq p} H^{r,k-r}(X)$, which is $F^pH^k(X,\mathbb{C})$ by definition 3.1.5.

For the converse, let $\xi \in F^pH^k(X,\mathbb{C})$ and let γ be its harmonic representative. Decomposing γ into pure types gives $\gamma = \sum_r \gamma^{r,k-r}$ with each $\gamma^{r,k-r}$ harmonic, so $\xi = \sum_r[\gamma^{r,k-r}]$ with $[\gamma^{r,k-r}] \in H^{r,k-r}(X)$. This is the decomposition of ξ under theorem 2.1.1, and $\xi \in F^p$ says exactly that its components with $r<p$ vanish. A harmonic form whose class vanishes is itself zero, so $\gamma^{r,k-r}=0$ for $r<p$ and $\gamma \in F^p\mathcal{A}^k(X)$, as we sought to show.

The lemma converts a statement about the position of a class in the filtration into a statement about the type of a differential form, and differential forms on the fibres can be assembled into forms on the total space $\mathcal{X}$, where the complex structure is fixed once and for all. That is the mechanism behind both theorems. Making it work requires knowing how to differentiate a section of $\mathcal{H}^k$ presented by forms, which is the content of the next lemma: the Gauss-Manin derivative of a class represented by a form is computed by contracting the differential of that form against a lift of the direction of differentiation.

Lemma 5.2.5: The Gauss-Manin Derivative of a Family of Forms

Let $U \subseteq S$ be an open ball over which f admits a C^∞ trivialization, let v be a smooth complex vector field on U, and let $\tilde{v}$ be any smooth complex vector field on $\mathcal{X}_U = f^{-1}(U)$ with $df(\tilde{v}) = v$. Let $\tilde{\alpha}$ be a smooth complex k-form on $\mathcal{X}_U$ whose restriction to every fibre is closed, and let σ be the section of $\mathcal{H}^k$ over U given by $\sigma(s) = [\tilde{\alpha}|_{X_s}]$. Then σ is smooth, the form $(\iota_{\tilde{v}}\, d\tilde{\alpha})|_{X_s}$ is closed for every $s \in U$, and

$$\nabla_v \sigma(s) = \left[(\iota_{\tilde{v}}\, d\tilde{\alpha})|_{X_s}\right]$$

***Proof*:**
Both sides are $\mathbb{C}$-linear in v, and every smooth complex vector field on U is a combination of real ones with complex coefficients, so it suffices to treat v real.

Step 1: the flat trivialization in terms of forms. Let $\Psi\colon X \times U \to \mathcal{X}_U$ be a C^∞ trivialization over U, where $X = X_{s_0}$ for a chosen $s_0 \in U$; one exists because f is a proper submersion [32, Ehresmann's theorem]. Write $j_s\colon X \to X \times U$ for $x \mapsto (x, s)$ and $\varphi_s = \Psi \circ j_s\colon X \to X_s$, a diffeomorphism onto the fibre.

The maps $\varphi_s^*\colon H^k(X_s, \mathbb{C}) \to H^k(X, \mathbb{C})$ trivialize $\mathcal{H}^k|_U$ flatly. By theorem 5.1.7 the flat sections of $\mathcal{H}^k$ over U are the sections of $\mathbb{H}^k \otimes_{\mathbb{Q}} \mathbb{C}$, and since U is a ball the trivialization makes the fibre inclusion $X_s \hookrightarrow \mathcal{X}_U$ a homotopy equivalence, so those sections are exactly the maps $s \mapsto c|_{X_s}$ for $c \in H^k(\mathcal{X}_U, \mathbb{C})$ (definition 5.1.3). For such a section, $\varphi_s^*(c|_{X_s}) = (\Psi \circ j_s)^* c$, and the maps $\Psi \circ j_s\colon X \to \mathcal{X}_U$ for different s are homotopic, U being connected. So $s \mapsto \varphi_s^*(c|_{X_s})$ is constant, and a section τ of $\mathcal{H}^k|_U$ is flat if and only if $s \mapsto \varphi_s^*\tau(s)$ is constant. Under this trivialization ∇ becomes the ordinary derivative.

Step 2: differentiating. Put $A = \Psi^*\tilde{\alpha}$, a k-form on $X \times U$. Since $\Psi \circ j_s$ is the fibre inclusion composed with φ_s,

$$\varphi_s^*\big(\tilde{\alpha}|_{X_s}\big) = j_s^* A$$

which depends smoothly on s; hence so does its de Rham class, and σ is smooth. Let ψ_t be the local flow of v on U and $\Phi_t(x, s) = (x, \psi_t(s))$ the induced flow on $X \times U$, whose generator is the vector field $\hat{v} = (0, v)$. From $j_{\psi_t(s)} = \Phi_t \circ j_s$,

$$\frac{d}{dt}\Big|_{t=0} j_{\psi_t(s)}^* A = j_s^*\big(L_{\hat{v}} A\big)$$

By Step 1 the left-hand side represents $\varphi_s^*\big(\nabla_v \sigma(s)\big)$; it is also the derivative of a smooth family of closed forms, hence closed. Set $\tilde{v}_0 = \Psi_* \hat{v}$, a lift of v. Pullback commutes with the Lie derivative,

so $j_s^*(L_{\hat{v}}A) = \varphi_s^*\big((L_{\tilde{v}_0}\tilde{\alpha})|_{X_s}\big)$, and therefore

$$\nabla_v\sigma(s) = \big[(L_{\tilde{v}_0}\tilde{\alpha})|_{X_s}\big]$$

with $(L_{\tilde{v}_0}\tilde{\alpha})|_{X_s}$ closed.

Step 3: Cartan's formula and independence of the lift. By $L_{\tilde{v}_0} = \iota_{\tilde{v}_0}d + d\,\iota_{\tilde{v}_0}$ and the fact that restriction to a submanifold commutes with d,

$$(L_{\tilde{v}_0}\tilde{\alpha})|_{X_s} = (\iota_{\tilde{v}_0}d\tilde{\alpha})|_{X_s} + d\big((\iota_{\tilde{v}_0}\tilde{\alpha})|_{X_s}\big)$$

The second term is exact on X_s, so it does not change the class, and subtracting it from a closed form leaves $(\iota_{\tilde{v}_0}d\tilde{\alpha})|_{X_s}$ closed with $\nabla_v\sigma(s) = [(\iota_{\tilde{v}_0}d\tilde{\alpha})|_{X_s}]$.

Finally, let $\tilde{v}$ be any other lift of v and put $w = \tilde{v} - \tilde{v}_0$, a vector field tangent to the fibres. Contraction by a vector field tangent to a submanifold commutes with restriction to it, so

$$(\iota_w\, d\tilde{\alpha})|_{X_s} = \iota_w\big((d\tilde{\alpha})|_{X_s}\big) = \iota_w\, d\big(\tilde{\alpha}|_{X_s}\big) = 0$$

the last equality because $\tilde{\alpha}|_{X_s}$ is closed. Hence $(\iota_{\tilde{v}}d\tilde{\alpha})|_{X_s} = (\iota_{\tilde{v}_0}d\tilde{\alpha})|_{X_s}$, completing the proof.

The formula makes the whole of the Gauss-Manin connection computable from forms on $\mathcal{X}$, and it is the only computation the two theorems need. Both are then a count of types: d never lowers the holomorphic degree, and contraction with a lift lowers it by one when the lift is of type $(1,0)$ and not at all when it is of type $(0,1)$. The second case gives holomorphy of the Hodge bundles, the first gives transversality.

Assembling a family of fibre forms into a form on $\mathcal{X}$ is done once, and the construction is recorded here so that both proofs may quote it. Fix a Hermitian metric on $\mathcal{X}_U$ and let $H \subseteq T\mathcal{X}_U$ be the orthogonal complement of the relative tangent bundle $T_{\mathcal{X}/S} = \ker df$. Since f is holomorphic, $\ker df$ is a complex subbundle, hence so is H, and df carries H isomorphically onto f^*TS. Let $\pi\colon T\mathcal{X}_U \to T_{\mathcal{X}/S}$ be the projection with kernel H; it is complex linear. For a smooth family $\{\beta_s\}$ of k-forms on the fibres, the form $\pi^*\beta$ on $\mathcal{X}_U$ is smooth, restricts to β_s on each X_s, and has the same type as β has relatively. In particular a family of relative type at least p extends to a form in $F^p\mathcal{A}^k(\mathcal{X}_U)$. The subbundle H also gives, for each vector field v on U, a lift $\tilde{v}$ lying in H, whose $(1,0)$-part lifts the $(1,0)$-part of v and whose $(0,1)$-part lifts the $(0,1)$-part.

Theorem 5.2.6: The Hodge Bundles Are Holomorphic Subbundles

Let $f\colon \mathcal{X} \to S$ be a smooth projective family with S connected and fix k and p. Then $F^p\mathcal{H}^k$ is a holomorphic subbundle of $\mathcal{H}^k$, of rank

$$f^p = \sum_{r\geq p} h^{r,k-r}$$

independent of the point of S. Moreover every smooth local section of $F^p\mathcal{H}^k$ is of the form $s \mapsto [\gamma_s]$ for a smooth family $\{\gamma_s\}$ of closed k-forms on the fibres of relative type at least p.

***Proof*:**
All assertions are local on S, so fix $s_0 \in S$ and work over a ball U around it, small enough to carry a C^∞ trivialization of f and the data H, π constructed above.

Step 1: the rank is constant. By definition 3.1.5 the dimension of $F^pH^k(X_s,\mathbb{C})$ is $\sum_{r\geq p} h^{r,k-r}(X_s)$, and each $h^{r,k-r}(X_s)$ is independent of s by proposition 5.2.3. So the fibre dimension of $F^p\mathcal{H}^k$ is the constant f^p.

Step 2: a smooth frame, and the representation by forms. Fix $\xi_0 \in F^pH^k(X_{s_0},\mathbb{C})$. By lemma 5.2.4 it has a harmonic representative $\alpha_0 \in F^p\mathcal{A}^k(X_{s_0})$. Each pure-type component $\alpha_0^{r,k-r}$ with $r \geq p$ is a smooth section over the closed submanifold X_{s_0} of the smooth vector bundle $\bigwedge^{r,k-r} T^*_{\mathcal{X}/S}$, so it extends, after shrinking U, to a smooth section over $\mathcal{X}_U$; summing the extensions gives a smooth family $\{\beta_s\}$ of k-forms on the fibres, of relative type at least p, with $\beta_{s_0} = \alpha_0$.

Put $\gamma_s = \Pi_s(\beta_s)$. By proposition 5.2.3 this is a smooth family, by theorem 1.3.1 the harmonic projection preserves bidegree so it again has relative type at least p, and each γ_s is harmonic, hence closed. By lemma 5.2.4 the class $[\gamma_s]$ lies in $F^pH^k(X_s,\mathbb{C})$, and $\gamma_{s_0} = \alpha_0$ because α_0 is already harmonic. Lemma 5.2.5, applied to the form $\pi^*\gamma$, shows that $\sigma_{\xi_0}\colon s \mapsto [\gamma_s]$ is a smooth section of $\mathcal{H}^k$; by construction it takes values in $F^p\mathcal{H}^k$ and $\sigma_{\xi_0}(s_0) = \xi_0$.

Choosing ξ_0 over a basis of $F^pH^k(X_{s_0},\mathbb{C})$ produces f^p smooth sections, linearly independent at s_0 and hence on a neighbourhood. By Step 1 they span $F^pH^k(X_s,\mathbb{C})$ at every point of that neighbourhood, so $F^p\mathcal{H}^k$ is a C^∞ subbundle of $\mathcal{H}^k$ and these sections are a smooth frame. Any smooth section of $F^p\mathcal{H}^k$ is a combination $\sum_i g_i\sigma_{\xi_i}$ with smooth coefficients g_i, and the family $\left\{\sum_i g_i(s)\gamma_s^i\right\}$ then represents it and is smooth, closed and of relative type at least p. This proves the last assertion of the theorem.

Step 3: $\nabla^{0,1}$ preserves the subbundle. Let σ be a smooth section of $F^p\mathcal{H}^k$ over U and $\bar v$ a vector field on U of type $(0,1)$. By Step 2 write $\sigma(s) = [\gamma_s]$ with $\{\gamma_s\}$ smooth, closed and of relative type at least p, and set $\tilde\alpha = \pi^*\gamma$, a form in $F^p\mathcal{A}^k(\mathcal{X}_U)$ restricting to γ_s on each fibre. Since $d = \partial + \bar\partial$ and neither operator lowers the holomorphic degree, $d\tilde\alpha \in F^p\mathcal{A}^{k+1}(\mathcal{X}_U)$. Let $\tilde{\bar v}$ be the lift of $\bar v$ lying in H, of type $(0,1)$; contraction with a $(0,1)$-vector field maps $\mathcal{A}^{r,q}$ into $\mathcal{A}^{r,q-1}$ and so preserves $F^p\mathcal{A}^\bullet$. Hence $\iota_{\tilde{\bar v}}d\tilde\alpha \in F^p\mathcal{A}^k(\mathcal{X}_U)$. Restriction of forms to the complex submanifold X_s preserves bidegree, so $(\iota_{\tilde{\bar v}}d\tilde\alpha)|_{X_s}$ lies in $F^p\mathcal{A}^k(X_s)$ and is closed by lemma 5.2.5. That lemma and lemma 5.2.4 now give

$$\nabla_{\bar v}\sigma(s) = \left[(\iota_{\tilde{\bar v}}d\tilde\alpha)|_{X_s}\right] \in F^pH^k(X_s,\mathbb{C})$$

for every $s \in U$, which is the assertion of this step.

Step 4: holomorphy. Shrink U so that $\mathcal{H}^k|_U$ is trivialized by a flat frame; this identifies $\mathcal{H}^k|_U$ with $\Xi \otimes_{\mathbb{C}} \mathcal{O}_U$ for the fixed vector space $\Xi = H^k(X_{s_0},\mathbb{C})$, with ∇ the ordinary derivative and $\nabla^{0,1} = \bar\partial$ acting on coefficient functions. Under this identification $F^p\mathcal{H}^k$ becomes a smooth family $s \mapsto W(s) \subseteq \Xi$ of subspaces of dimension f^p, and Step 3 says that $\bar\partial\sigma$ is a section of W whenever σ is.

Choose a complement $\Xi = W(s_0) \oplus C$. The projection $\rho\colon \Xi \to W(s_0)$ along C restricts to an injection on $W(s)$ for s near s_0, since $W(s) \cap C = 0$ is an open condition satisfied at s_0, and by

equality of dimensions it is then an isomorphism $W(s) \to W(s_0)$. So $W(s)$ is the graph

$$W(s) = \{\, w + A(s)w \,\big|\, w \in W(s_0) \,\}$$

of the linear map $A(s) = (\mathrm{id} - \rho) \circ (\rho|_{W(s)})^{-1} \colon W(s_0) \to C$, and A is smooth in s because W is a smooth subbundle. For fixed $w \in W(s_0)$ the assignment $\sigma_w(s) = w + A(s)w$ is a smooth section of $F^p\mathcal{H}^k$, so $\bar\partial\sigma_w(s) = \bar\partial\big(A(s)w\big)$ lies in $W(s)$. It also lies in C, the subspace C being independent of s, and $W(s) \cap C = 0$. Hence $\bar\partial(A(s)w) = 0$, so $A(\cdot)w$ is holomorphic and σ_w is a holomorphic section of $\mathcal{H}^k$. Letting w range over a basis of $W(s_0)$ gives f^p holomorphic sections spanning $W(s)$ at every point near s_0, so $F^p\mathcal{H}^k$ is a holomorphic subbundle, completing the proof.

The theorem is what justifies the name in definition 5.2.1, and it also fixes which of the two presentations of a Hodge structure is the holomorphic one. The individual pieces $H^{p,q}(X_s)$ of the Hodge decomposition do *not* form a holomorphic subbundle: they are recovered from the filtration only by the formula $H^{p,q} = F^p \cap \overline{F^q}$ of (3.2), and conjugation is not holomorphic. What is holomorphic is the filtration, and this is the reason chapter 3 went to the trouble of proving the two presentations equivalent (theorem 3.1.6) before any family appeared.

With the subbundles in hand, the question is how far they fail to be flat. A flat subbundle would satisfy $\nabla F^p \subseteq F^p \otimes \Omega^1_S$, which by the argument of exercise 5.2.1 (1) would force the Hodge structures of the fibres to be constant in the flat trivialization. That does not happen for a family with varying complex structure. The theorem below says the failure is as small as it can be while still occurring: one step of the filtration, never two.

Theorem 5.2.7: Griffiths Transversality

Let $f \colon \mathcal{X} \to S$ be a smooth projective family with S connected. Then for every k and every p,

$$\nabla\big(F^p\mathcal{H}^k\big) \subseteq F^{p-1}\mathcal{H}^k \otimes_{\mathcal{O}_S} \Omega^1_S$$

That is, if σ is a local holomorphic section of $F^p\mathcal{H}^k$ and v a local holomorphic vector field on S, then $\nabla_v\sigma$ is a section of $F^{p-1}\mathcal{H}^k$.

***Proof*:**
The assertion is local on S, so fix s_0 and work over a ball U carrying a C^∞ trivialization of f together with the splitting data H, π fixed before theorem 5.2.6. It suffices to prove the second formulation, since a section of $\mathcal{H}^k \otimes \Omega^1_S$ lies in $F^{p-1}\mathcal{H}^k \otimes \Omega^1_S$ exactly when its contraction with every local holomorphic vector field lies in $F^{p-1}\mathcal{H}^k$.

Step 1: represent the section by forms of type at least p. Let σ be a holomorphic, in particular smooth, section of $F^p\mathcal{H}^k$ over U. By theorem 5.2.6 there is a smooth family $\{\gamma_s\}$ of closed k-forms on the fibres, of relative type at least p, with $[\gamma_s] = \sigma(s)$. Set $\tilde\alpha = \pi^*\gamma$. By the construction of π this is a smooth k-form on $\mathcal{X}_U$ with $\tilde\alpha|_{X_s} = \gamma_s$ for every s, and it lies in $F^p\mathcal{A}^k(\mathcal{X}_U)$ because π is complex linear and γ has relative type at least p.

Step 2: the type of $\iota_{\tilde v} d\tilde\alpha$. Write $\tilde\alpha = \sum_{r \geq p} \tilde\alpha^{r,k-r}$. Since $d = \partial + \bar\partial$ on $\mathcal{X}_U$ (proposition 1.1.1)

and ∂ raises the holomorphic degree by one while $\bar{\partial}$ leaves it unchanged,

$$d\tilde{\alpha} \in F^p\mathcal{A}^{k+1}(\mathcal{X}_U)$$

Let $\tilde{v}$ be the lift of v lying in H. Because v is of type $(1,0)$ and H is a complex subbundle mapped isomorphically to f^*TS, the lift $\tilde{v}$ is of type $(1,0)$ as well. Contraction with a $(1,0)$-vector field maps $\mathcal{A}^{r,q}$ into $\mathcal{A}^{r-1,q}$, so it lowers the level of the filtration by exactly one:

$$\iota_{\tilde{v}}\, d\tilde{\alpha} \in F^{p-1}\mathcal{A}^k(\mathcal{X}_U)$$

Every other operation in the argument preserves the filtration of forms.

Step 3: restrict to a fibre. The fibre X_s is a complex submanifold of $\mathcal{X}_U$, so restriction of forms to it preserves the bidegree and hence carries $F^{p-1}\mathcal{A}^k(\mathcal{X}_U)$ into $F^{p-1}\mathcal{A}^k(X_s)$. By lemma 5.2.5 the restricted form $(\iota_{\tilde{v}}d\tilde{\alpha})|_{X_s}$ is closed and

$$\nabla_v\sigma(s) = \left[(\iota_{\tilde{v}}\, d\tilde{\alpha})|_{X_s}\right]$$

A closed form in $F^{p-1}\mathcal{A}^k(X_s)$ has class in $F^{p-1}H^k(X_s,\mathbb{C})$ by lemma 5.2.4. Hence $\nabla_v\sigma(s) \in F^{p-1}H^k(X_s,\mathbb{C})$ for every $s \in U$, which is the claim, completing the proof.

The transversality condition is a first-order differential equation satisfied by the Hodge filtration of every family of smooth projective varieties, and it is a restriction on the family only because F^{p-1} is smaller than $\mathcal{H}^k$. For $p \leq 1$ it says nothing, since $F^{p-1} = \mathcal{H}^k$ there (exercise 5.2.1 (2)); the first case with content is $k = 2$, $p = 2$. Two readings are worth separating. Read as a constraint, it says that a family of varieties cannot deform its Hodge structure in an arbitrary direction: most motions of a flag are forbidden, and the permitted ones form a distinguished subbundle of the tangent bundle of the parameter space of flags. That is the geometric form of the statement, and section 5.3 puts it in those terms. Read as a computation, it says that the interesting derivative is the induced map on the graded pieces, since transversality guarantees that this map is defined at all. The corollary records it.

Corollary 5.2.8: The Graded Derivative

Write $\mathrm{gr}_F^p\mathcal{H}^k = F^p\mathcal{H}^k/F^{p+1}\mathcal{H}^k$, a holomorphic bundle whose fibre at s is $H^{p,k-p}(X_s)$. The Gauss-Manin connection induces an $\mathcal{O}_S$-linear map of holomorphic bundles

$$\bar{\nabla}^p\colon \mathrm{gr}_F^p\mathcal{H}^k \to \mathrm{gr}_F^{p-1}\mathcal{H}^k \otimes_{\mathcal{O}_S} \Omega_S^1$$

whose value at s is a linear map $H^{p,k-p}(X_s) \to H^{p-1,k-p+1}(X_s)\otimes T_s^*S$, equivalently a bilinear map

$$T_sS \otimes H^{p,k-p}(X_s) \to H^{p-1,k-p+1}(X_s)$$

***Proof*:**
The fibre of $\mathrm{gr}_F^p\mathcal{H}^k$ at s is $F^pH^k(X_s,\mathbb{C})/F^{p+1}H^k(X_s,\mathbb{C})$, which is $H^{p,k-p}(X_s)$ by definition 3.1.5. Both $F^p\mathcal{H}^k$ and $F^{p+1}\mathcal{H}^k$ are holomorphic subbundles by theorem 5.2.6, so the quotient is a holomorphic bundle [5, Subbundle and Quotient Bundle].

By theorem 5.2.7 the connection gives a map of sheaves

$$F^p\mathcal{H}^k \to F^{p-1}\mathcal{H}^k \otimes \Omega^1_S \to \left(F^{p-1}\mathcal{H}^k/F^p\mathcal{H}^k\right) \otimes \Omega^1_S$$

the second arrow being the quotient. This composite kills $F^{p+1}\mathcal{H}^k$: a section σ of $F^{p+1}\mathcal{H}^k$ has $\nabla\sigma \in F^p\mathcal{H}^k \otimes \Omega^1_S$, again by theorem 5.2.7 applied with $p+1$ in place of p, and that is exactly what the quotient annihilates. So the composite factors through $\mathrm{gr}^p_F\mathcal{H}^k$.

It remains to check $\mathcal{O}_S$-linearity, which is where transversality is used a second time. For a local function g and a section σ of $F^p\mathcal{H}^k$, the Leibniz rule gives

$$\nabla(g\sigma) = g\,\nabla\sigma + \sigma \otimes dg$$

The correction term $\sigma \otimes dg$ lies in $F^p\mathcal{H}^k \otimes \Omega^1_S$, which the quotient by $F^p\mathcal{H}^k$ annihilates. Hence the induced map satisfies $\bar\nabla^p(g\,\bar\sigma) = g\,\bar\nabla^p(\bar\sigma)$, so it is $\mathcal{O}_S$-linear and therefore a morphism of holomorphic bundles, with the stated value on fibres, as we sought to show.

A connection is not tensorial, so it has no value at a point; the graded derivative $\bar\nabla^p$ is tensorial, so it does, and that value is an invariant attached to the fibre X_s together with a tangent direction at s. It measures the first-order motion of the Hodge structure of X_s as the parameter moves, and it is the object that section 5.3 identifies with the differential of the period map. Chapter 6 takes up what happens when it vanishes.

In weight one the condition is empty, and the reason for that is worth recording before the general theory is put to work.

Example 5.2.9: Transversality in Weight One
Take $k = 1$, so that the Hodge bundles are

$$\mathcal{H}^1 = F^0\mathcal{H}^1 \supseteq F^1\mathcal{H}^1 \supseteq F^2\mathcal{H}^1 = 0$$

as in example 5.2.2. Every instance of theorem 5.2.7 is empty, for one of two reasons. For $p \le 1$ the right-hand side is everything, since $F^{p-1}\mathcal{H}^1 = \mathcal{H}^1$ by exercise 5.2.1 (2): at $p = 1$ the condition reads $\nabla F^1\mathcal{H}^1 \subseteq F^0\mathcal{H}^1 \otimes \Omega^1_S = \mathcal{H}^1 \otimes \Omega^1_S$, and at $p \le 0$ it reads $\nabla\mathcal{H}^1 \subseteq \mathcal{H}^1 \otimes \Omega^1_S$; every section satisfies both. For $p \ge 2$ the left-hand side vanishes, since $F^p\mathcal{H}^1 = 0$. So transversality imposes no condition whatever on a family of Hodge structures of weight 1.

The reason is structural rather than accidental. A weight-1 structure is a filtration with a single interesting step, F^1, and by example 3.1.3 it amounts to no more than a complex structure on the underlying real vector space. There is nothing for the filtration to be transverse to. The only graded derivative that can be nonzero is

$$\bar\nabla^1 \colon \mathrm{gr}^1_F\mathcal{H}^1 \to \mathrm{gr}^0_F\mathcal{H}^1 \otimes \Omega^1_S, \qquad H^{1,0}(X_s) \to H^{0,1}(X_s) \otimes T^*_sS$$

and theorem 5.2.7 places no restriction on it.

That does not make the graded derivative arbitrary. What constrains it is the polarization, and the constraint is a symmetry. Write g for the rank of $F^1\mathcal{H}^1$, so that $\mathcal{H}^1$ has rank $2g$, and let Q be the polarization of proposition 3.2.9, which is alternating in odd weight, flat for ∇ (exercise 5.1.1 (4)) and zero on $F^1\mathcal{H}^1 \times F^1\mathcal{H}^1$ by the first Hodge-Riemann relation (theorem 3.2.4). For a local vector field v and local sections σ, τ of $F^1\mathcal{H}^1$, flatness gives

$$0 = v\,Q(\sigma,\tau) = Q(\nabla_v\sigma,\tau) + Q(\sigma,\nabla_v\tau)$$

so that $Q(\nabla_v\sigma,\tau) = Q(\nabla_v\tau,\sigma)$: the form $(\sigma,\tau)\mapsto Q(\nabla_v\sigma,\tau)$ on $F^1\mathcal{H}^1$ is symmetric. Being isotropic of half the rank, $F^1\mathcal{H}^1$ is its own annihilator under Q, so Q pairs $\mathcal{H}^1/F^1\mathcal{H}^1$ nondegenerately against it and that symmetric form determines $\bar\nabla^1_v$ and is determined by it. Hence $\bar\nabla^1_v$ ranges over a space of dimension $g(g+1)/2$ inside the space $\mathrm{Hom}\big(H^{1,0}(X_s),H^{0,1}(X_s)\big)$ of all linear maps, of dimension g^2. For $g\geq 2$ that is a proper subspace, and its dimension is that of the period domain $\mathfrak{H}_g$ of this numerical type (example 3.3.3). None of it comes from transversality.

The graded derivative vanishes identically exactly when $F^1\mathcal{H}^1$ is a flat subbundle, that is when the family has locally constant Hodge structure (exercise 5.2.1 (1)).

This is why the classical theory of abelian varieties never encounters transversality. By theorem 3.4.1 a polarizable weight-1 Hodge structure is the same thing as a complex torus with a polarization, and by corollary 3.4.2 the polarizable ones are the abelian varieties; a family of abelian varieties is a family of weight-1 structures, on whose variation the preceding paragraphs put no transversality constraint at all. The constraint such a family does satisfy is the symmetry computed above, and that one belongs to the polarization. For the Legendre family of example 5.2.2 the bundle $F^1\mathcal{H}^1$ is a line bundle and $\bar\nabla^1$ is a map of line bundles twisted by Ω^1_S; the Gauss-Manin computation of example 5.1.9 is a computation of that map, and transversality contributes nothing to it.

The condition first acquires content at $k=2$ and $p=2$, where F^1 is a proper subbundle of $\mathcal{H}^2$ and $\nabla F^2\subseteq F^1\otimes\Omega^1_S$ is a genuine restriction. That case is worked out for K3 surfaces in exercise 5.2.1 (5).

Exercise 5.2.1

1. Show that the following are equivalent for a smooth projective family $f\colon\mathcal{X}\to S$ and a fixed k: (i) the graded derivatives $\bar\nabla^p$ of corollary 5.2.8 vanish for every p; (ii) each $F^p\mathcal{H}^k$ satisfies $\nabla F^p\mathcal{H}^k\subseteq F^p\mathcal{H}^k\otimes\Omega^1_S$. Then show that when these hold and $U\subseteq S$ is a ball, the flat trivialization of $\mathcal{H}^k|_U$ carries $F^pH^k(X_s,\mathbb{C})$ to a subspace of $H^k(X_{s_0},\mathbb{C})$ independent of s. (For the last part, imitate Step 4 of the proof of theorem 5.2.6, replacing $\bar\partial$ by the full derivative.)

2. Let $f\colon\mathcal{X}\to S$ be a smooth projective family and fix k. Show that $F^p\mathcal{H}^k=\mathcal{H}^k$ for $p\leq 0$ and $F^p\mathcal{H}^k=0$ for $p>k$. Deduce that the transversality condition of theorem 5.2.7 at level p is automatically satisfied unless $2\leq p\leq k$, and conclude that it says nothing at all when $k\leq 1$.

3. Keep the product family $f\colon X\times S\to S$ of exercise 5.1.1 (2), which identified $\mathbb{H}^k$ with the constant local system and ∇ with $1\otimes d$. Show that each $F^p\mathcal{H}^k$ is likewise the constant subbundle with fibre $F^pH^k(X,\mathbb{C})$. Conclude that $\bar\nabla^p=0$ for every p, consistently with exercise 5.2.1 (1).

4. Let $s_1,\dots,s_m$ be holomorphic coordinates on an open set $U\subseteq S$ and write $\nabla_i=\nabla_{\partial/\partial s_i}$. Using flatness (theorem 5.1.7) and theorem 5.2.7, show that for a holomorphic section σ of $F^p\mathcal{H}^k$ the classes of $\nabla_i\nabla_j\sigma$ in $\mathrm{gr}_F^{p-2}\mathcal{H}^k$ are symmetric in i and j, and that they vanish when σ is a section of $F^{p+1}\mathcal{H}^k$. Deduce that the composite $(\bar\nabla^{p-1}\otimes\mathrm{id})\circ\bar\nabla^p$ takes values in $\mathrm{gr}_F^{p-2}\mathcal{H}^k\otimes\mathrm{Sym}^2\Omega^1_S$.

5. Let $f\colon\mathcal{X}\to S$ be a smooth projective family of K3 surfaces, whose degree-2 Hodge numbers are $h^{2,0}=h^{0,2}=1$ and $h^{1,1}=20$ (see section 3.4), and take $k=2$. Compute the ranks of $F^2\mathcal{H}^2\subseteq F^1\mathcal{H}^2\subseteq\mathcal{H}^2$ and of the three graded pieces. State what theorem 5.2.7 asserts at $p=2$ and at $p=1$, and say which of the two has content. Finally, let Q denote the cup-product

pairing on the fibres, which is defined by the local system and is therefore flat for ∇, and let σ be a local holomorphic section of $F^2\mathcal{H}^2$. Show that $Q(\sigma, \nabla_v \sigma) = 0$ for every local holomorphic vector field v.

5.3 The Period Map

Section 5.1 put a flat connection on the bundle of fibre cohomology and section 5.2 put a filtration of that bundle by holomorphic subbundles which the connection moves by exactly one step. Both are statements about objects living over S. This section converts them into a single map out of S. Each fibre carries a polarized Hodge structure, the polarized structures of a fixed numerical type are the points of the period domain of section 3.3, and the assignment sending s to the structure on X_s is the period map.

Two things have to be arranged before the assignment is a map at all. The identification of $H^k(X_s, \mathbb{Z})$ with a fixed lattice depends on a path from a base point, so the target must be a quotient of the period domain by the monodromy group; and the quotient must be a space in which holomorphy makes sense. Once both are in place, the two theorems of the section say that the map is holomorphic and that its differential is constrained: it lands in a distinguished subbundle of the tangent bundle of D. That constraint is Griffiths transversality read on the domain rather than on the base.

Throughout, $f\colon \mathcal{X} \to S$ is a smooth projective family (definition 5.1.1), with fibres $X_s = f^{-1}(s)$ of complex dimension n, with relative hyperplane class $h \in H^2(\mathcal{X}, \mathbb{Z})$, and with $h_s = h|_{X_s}$ the class of the Fubini-Study Kähler form of X_s (theorem 1.1.7). Fix an integer $0 \leq k \leq n$ and let $\mathbb{H}^k$, $\mathcal{H}^k$, ∇ and $F^\bullet$ be the local system, the holomorphic bundle, the Gauss-Manin connection and the Hodge filtration attached to f in section 5.1 and section 5.2.

Cup product with h is a map of local systems $L\colon \mathbb{H}^j \to \mathbb{H}^{j+2}$, since h is a single class on $\mathcal{X}$ restricted to the fibres, and on each fibre it is the Lefschetz operator of definition 1.1.8. The part of the cohomology that this section works with is cut out by a power of L.

Definition 5.3.1: The Primitive Part of the Cohomology of a Family

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and let $0 \leq k \leq n$. The *primitive part* of $\mathbb{H}^k$ is the subsheaf

$$\mathbb{H}^k_{\text{prim}} := \ker\bigl(L^{n-k+1}\colon \mathbb{H}^k \to \mathbb{H}^{2n-k+2}\bigr)$$

whose stalk at s is the primitive cohomology $H^k(X_s, \mathbb{Q})_{\text{prim}}$ of definition 3.2.6, and the *primitive cohomology bundle* is the $\mathcal{O}_S$-submodule

$$\mathcal{H}^k_{\text{prim}} := \mathbb{H}^k_{\text{prim}} \otimes_{\mathbb{Q}} \mathcal{O}_S$$

of $\mathcal{H}^k$. Part (3) of proposition 5.3.2 shows that the first is a sub-local system of $\mathbb{H}^k$ and the second a holomorphic subbundle of $\mathcal{H}^k$.

The period domain of section 3.3 was attached to numerical data fixed once and for all: a lattice, a weight, a form and a list of Hodge numbers. A family gives that data only after a fibre is singled out, and it gives it consistently only because parallel transport moves everything rigidly. The following proposition collects the four facts the construction needs, and it is the only place where the geometry of f is used.

Proposition 5.3.2: Polarized Period Data of a Family

Let $f\colon \mathcal{X} \to S$ be a smooth projective family as above, let $0 \le k \le n$, fix $s_0 \in S$, and put

$$V_{\mathbb{Z}} = H^k(X_{s_0},\mathbb{Z})_{\mathrm{prim}}/(\text{torsion}), \qquad h^{p,q} = h^{p,q}_{\mathrm{prim}}(X_{s_0}), \qquad f^p = \sum_{r\ge p} h^{r,k-r}$$

with Q the polarization of proposition 3.2.9 on $V_{\mathbb{Z}}$. The f^p are the level dimensions attached to these data in section 3.3; they are formed from the primitive Hodge numbers and are in general smaller than the ranks of the Hodge bundles $F^p \subseteq \mathcal{H}^k$. Then:

1. For every $s \in S$ and every homotopy class of path c from s_0 to s, pullback along the inverse of the diffeomorphism $\varphi_c\colon X_{s_0} \to X_s$ that a trivialization of f along c provides is an isomorphism $T_c = (\varphi_c^{-1})^*\colon H^k(X_{s_0},\mathbb{Z}) \to H^k(X_s,\mathbb{Z})$, depending only on $[c]$ and agreeing with the parallel transport of definition 5.1.3 after tensoring with $\mathbb{Q}$. It commutes with cup product and carries h_{s_0} to h_s, hence restricts to an isomorphism $V_{\mathbb{Z}} \to H^k(X_s,\mathbb{Z})_{\mathrm{prim}}/(\text{torsion})$ carrying Q to Q.

2. The primitive Hodge numbers $h^{p,q}_{\mathrm{prim}}(X_s)$ do not depend on s.

3. The primitive cohomology bundle $\mathcal{H}^k_{\mathrm{prim}}$ of definition 5.3.1 is a holomorphic subbundle of $\mathcal{H}^k$ preserved by ∇; the subsheaves $F^p_{\mathrm{prim}} := F^p \cap \mathcal{H}^k_{\mathrm{prim}}$ are holomorphic subbundles of it, of rank f^p; and
$$\nabla F^p_{\mathrm{prim}} \subseteq F^{p-1}_{\mathrm{prim}} \otimes \Omega^1_S$$

4. For every s and every path class c as in (1), the filtration $T_c^{-1}\big(F^\bullet_{\mathrm{prim},s}\big)$ of $V_{\mathbb{C}}$ is a point of the period domain D attached to $(V_{\mathbb{Z}}, Q, k, \{h^{p,q}\})$ by definition 3.3.2.

***Proof*:**

1. As in the proof of proposition 5.1.5, Ehresmann's theorem trivializes the pullback of f along c, and $T_c = (\varphi_c^{-1})^*$ for the diffeomorphism $\varphi_c\colon X_{s_0} \to X_s$ arising from a trivialization $\Psi\colon X_{s_0}\times[0,1] \to f^{-1}(c)$; two trivializations differ by a path of diffeomorphisms of the fibre starting at the identity, so T_c depends only on $[c]$, and with rational coefficients it is the transport of $\mathbb{H}^k$. A pullback along a diffeomorphism is a ring homomorphism for cup product, so T_c is.

   For the class h, write $\iota_s\colon X_s \hookrightarrow \mathcal{X}$ for the inclusion. The maps $u \mapsto \iota_{c(u)} \circ \Psi_u$ form a homotopy from ι_{s_0} to $\iota_s \circ \varphi_c$ as maps $X_{s_0} \to \mathcal{X}$, so $\varphi_c^* h_s = \varphi_c^* \iota_s^* h = \iota_{s_0}^* h = h_{s_0}$, that is $T_c(h_{s_0}) = h_s$. Consequently $T_c \circ L = L \circ T_c$ and T_c carries $\ker L^{n-k+1}$ onto $\ker L^{n-k+1}$.

   Finally φ_c preserves the complex orientations. The pullback Ψ_u^* of the complex orientation of $X_{c(u)}$ is an orientation of the connected manifold X_{s_0} depending continuously on u, hence is constant, hence equals its value at $u=0$, which is the complex orientation. So $\varphi_c = \Psi_1$ is orientation-preserving, and therefore
$$\int_{X_s} T_c\alpha \wedge T_c\beta \wedge h_s^{n-k} = \int_{X_{s_0}} \alpha\wedge\beta\wedge h_{s_0}^{n-k}$$
   using $T_c\, h_{s_0} = h_s$ for the third factor. With the sign $(-1)^{k(k-1)/2}$ of proposition 3.2.9 attached to both sides this says $Q(T_c\alpha, T_c\beta) = Q(\alpha,\beta)$.

2. Part (1) of proposition 5.2.3 already gives that each $h^{p,k-p}(X_s)$ is independent of s, so the ranks of the F^p inside $\mathcal{H}^k$ are constant. (Those ranks are not the f^p of the statement, which count primitive Hodge numbers only.)

   For the primitive numbers, fix s and $p+q=k\leq n$. The operator L raises the bidegree by $(1,1)$ and the primitive parts are sub-Hodge structures (proposition 3.2.7), so the Lefschetz decomposition corollary 4.2.4 of $H^k(X_s,\mathbb{C})$ refines along the bigrading into

   $$H^{p,q}(X_s)=\bigoplus_{r\geq 0} L^r\, H^{p-r,q-r}(X_s)_{\text{prim}}$$

   with each L^r injective on the summand it is applied to. Counting dimensions gives $h^{p,q}(X_s)=\sum_{r\geq 0} h^{p-r,q-r}_{\text{prim}}(X_s)$, and the same identity in bidegree $(p-1,q-1)$ subtracts off all but the first term:

   $$h^{p,q}_{\text{prim}}(X_s)=h^{p,q}(X_s)-h^{p-1,q-1}(X_s)$$

   Both terms on the right are independent of s, by the first paragraph applied in degree k and in degree $k-2$, so the left side is.

3. By (2) the kernel of L^{n-k+1} has constant rank, so $\mathbb{H}^k_{\text{prim}}$ is a sub-local system of $\mathbb{H}^k$ and $\mathcal{H}^k_{\text{prim}}$ is the holomorphic subbundle it generates. It is spanned locally by flat sections, so ∇ carries its sections into $\mathcal{H}^k_{\text{prim}}\otimes\Omega^1_S$: a local section is $\sum_j g_j e_j$ with g_j holomorphic and e_j flat, and $\nabla(\sum_j g_j e_j)=\sum_j e_j\otimes dg_j$ by theorem 5.1.7.

   The fibre of $F^p\cap\mathcal{H}^k_{\text{prim}}$ at s is $F^pH^k(X_s,\mathbb{C})\cap H^k(X_s,\mathbb{C})_{\text{prim}}$, which by proposition 3.2.7 is the p-th step of the Hodge filtration of the primitive structure and so has dimension $\sum_{r\geq p} h^{r,k-r}_{\text{prim}}(X_s)=f^p$, independent of s by (2). Now $F^p\cap\mathcal{H}^k_{\text{prim}}$ is the kernel of the composite holomorphic bundle map $F^p\to\mathcal{H}^k\to\mathcal{H}^k/\mathcal{H}^k_{\text{prim}}$, whose rank is the constant $\operatorname{rank}F^p-\dim(F^p\cap\mathcal{H}^k_{\text{prim}})$; the kernel of a holomorphic bundle map of constant rank is a holomorphic subbundle, so F^p_{prim} is one.

   For transversality, let σ be a local holomorphic section of F^p_{prim}. Then $\nabla\sigma\in F^{p-1}\otimes\Omega^1_S$ by theorem 5.2.7, and $\nabla\sigma\in\mathcal{H}^k_{\text{prim}}\otimes\Omega^1_S$ by the previous paragraph; intersecting the two gives $\nabla\sigma\in F^{p-1}_{\text{prim}}\otimes\Omega^1_S$.

4. Fix s and c. By proposition 3.2.7 the group $H^k(X_s,\mathbb{Z})_{\text{prim}}$ modulo torsion carries a pure Hodge structure of weight k, and by proposition 3.2.9 the form Q polarizes it; its Hodge numbers are the $h^{p,q}$ by (2), and its Hodge filtration is $F^\bullet_{\text{prim},s}$. The map T_c^{-1} is an isomorphism of lattices carrying Q to Q by (1), and its complexification commutes with conjugation because it is defined over $\mathbb{Z}$; so transporting the bigrading along T_c^{-1} produces a pure Hodge structure of weight k on $V_\mathbb{Z}$ with Hodge numbers $h^{p,q}$, polarized by Q, whose Hodge filtration is $T_c^{-1}(F^\bullet_{\text{prim},s})$. By part (1) of proposition 3.3.4 the Hodge filtration of such a structure is a point of D, completing the proof.

Part (1) is the reason the target is a quotient. Transport is an isomorphism of everything discrete, the lattice and the form, but it depends on the path only through its homotopy class, and different classes differ by a loop. The ambiguity is therefore exactly the monodromy group, and dividing by it is the smallest correction that removes it.

Definition 5.3.3: The Period Map

Keep the data of proposition 5.3.2 and set

$$G_{\mathbb{Z}} = \big\{ g \in \mathrm{GL}(V_{\mathbb{Z}}) \;\big|\; Q(gu, gv) = Q(u, v) \text{ for all } u, v \big\}$$

a subgroup of the group $G_{\mathbb{R}}$ of proposition 3.3.5. Let

$$\rho_k \colon \pi_1(S, s_0) \to G_{\mathbb{Z}}, \qquad \rho_k(\gamma) = T_\gamma$$

be the monodromy representation of definition 4.4.2 acting on $V_{\mathbb{Z}}$, which lands in $G_{\mathbb{Z}}$ by part (1) of proposition 5.3.2, and let $\Gamma = \rho_k\big(\pi_1(S, s_0)\big)$ be the *monodromy group* of the family in degree k.
For $s \in S$ and a homotopy class $[c]$ of path from s_0 to s put

$$\Phi_{[c]}(s) = T_{[c]}^{-1}\big(F^\bullet_{\mathrm{prim},s}\big) \in D$$

which lies in D by part (4) of proposition 5.3.2. The *period map* of the family in degree k is

$$\Phi \colon S \to \Gamma\backslash D, \qquad \Phi(s) = \Gamma \cdot \Phi_{[c]}(s)$$

Realizing the universal cover $\pi \colon \tilde{S} \to S$ as the set of homotopy classes of paths starting at s_0, the *lifted period map* is

$$\tilde{\Phi} \colon \tilde{S} \to D, \qquad \tilde{\Phi}\big([c]\big) = \Phi_{[c]}\big(c(1)\big)$$

That Φ is well defined is one line. If c and c' are two paths from s_0 to s, then $T_{[c']}^{-1} = T_{[\bar{c}']} = T_{[\bar{c}'c]} \circ T_{[c]}^{-1}$, where $\bar{c}'$ is c' reversed and $\bar{c}'c$ is the loop at s_0 obtained by traversing c and then $\bar{c}'$ (the composition convention of box 4.4.1). Hence

$$\Phi_{[c']}(s) = \rho_k\big([\bar{c}'c]\big)\, \Phi_{[c]}(s)$$

and the two points lie in the same Γ-orbit.

The definition should be read as a statement about what a family of varieties determines. A single fibre determines a point of D only after a marking, an identification of its primitive cohomology with $V_{\mathbb{Z}}$ respecting Q; the family gives markings, one for each homotopy class of path, and they differ by monodromy. What is intrinsic is the orbit. The classical case is the one where D is the upper half plane and the orbit is the period ratio of an elliptic curve read modulo the substitutions monodromy performs on it.

Example 5.3.4: The Period Map of a Family of Elliptic Curves

Let $f \colon \mathcal{X} \to S$ be a smooth projective family of curves of genus 1 over a connected base, and take $k = n = 1$. The Legendre family $y^2z = x(x-z)(x-tz)$ over $S = \mathbb{C} \setminus \{0, 1\}$ of example 5.1.9 is one such, and so is the restriction, over the complement of its twelve critical values, of the fibration attached to a Lefschetz pencil of plane cubics (example 4.4.8).

The numerical data. All of H^1 is primitive (exercise 3.2.1 (2)), so $V_{\mathbb{Z}} = H^1(X_{s_0}, \mathbb{Z}) \cong \mathbb{Z}^2$, and $h^{1,0} = h^{0,1} = 1$. The normalizing sign of proposition 3.2.9 is $(-1)^{k(k-1)/2} = 1$ and the exponent $n - k$ is 0, so Q is the cup product itself, a unimodular alternating form (example 3.2.3). Taking the genus parameter of example 3.3.3 to be 1, the compact dual is $\check{D} \cong \mathbb{C}\mathrm{P}^1$ and the period domain is the upper half plane $\mathfrak{H}_1$ (exercise 3.3.1 (2)).

The map in coordinates. Fix a symplectic basis a, b of $H_1(X_{s_0}, \mathbb{Z})$ and the dual basis u, v of $H^1(X_{s_0}, \mathbb{Z})$, normalized as in exercise 3.2.1 (6) so that $Q(u, v) = \int u \wedge v = 1$. Let $U \subseteq S$ be connected and simply connected, and transport a, b and u, v over U; the transported cycles a_s, b_s and classes u_s, v_s are flat frames. The line $F^1_s = H^{1,0}(X_s)$ is spanned by the class of the holomorphic 1-form ω_s of the fibre, unique up to scale, and

$$[\omega_s] = A(s)\, u_s + B(s)\, v_s, \qquad A(s) = \int_{a_s} \omega_s, \quad B(s) = \int_{b_s} \omega_s$$

are its two periods. Expanding Q in the frame,

$$i\, Q\big([\omega_s], \overline{[\omega_s]}\big) = i\big(A\bar{B} - \bar{A}B\big) = -2\,\mathrm{Im}\big(A\bar{B}\big)$$

and the second Hodge-Riemann relation makes this positive, so $\mathrm{Im}(\bar{A}B) > 0$. In particular A never vanishes, the line F^1_s is spanned by $u_s + \tau(s)\, v_s$ with

$$\tau(s) = \frac{B(s)}{A(s)} = \frac{\int_{b_s} \omega_s}{\int_{a_s} \omega_s}$$

and $\mathrm{Im}\,\tau = \mathrm{Im}(\bar{A}B)/|A|^2 > 0$. Under the identification of exercise 3.3.1 (2) the period map on U is $s \mapsto \tau(s)$: the classical period ratio.

The ambiguity. Here $G_{\mathbb{Z}} = \mathrm{Sp}(V_{\mathbb{Z}}, Q) = \mathrm{SL}(2, \mathbb{Z})$, so $\Gamma \subseteq \mathrm{SL}(2, \mathbb{Z})$ and Φ takes values in $\Gamma\backslash\mathfrak{H}_1$. Concretely, if the monodromy of a loop replaces the flat frame (a, b) by $(a,\ b + a)$, then A is unchanged while B becomes $A + B$, so

$$\tau \longmapsto \tau + 1$$

That transformation of the frame is exactly the Picard-Lefschetz transvection computed in example 4.4.8 for a nodal degeneration of a curve, and the displayed substitution is the Hodge-theoretic form promised there. Horizontality, the second theorem of this section, says nothing here: in weight 1 the transversality condition is vacuous (example 5.2.9). The reading of $\Gamma\backslash\mathfrak{H}_1$ as a moduli space of elliptic curves, and the modular curves it produces, are not developed here. They are in *Elliptic Curves* [6].

The target needs a word, because a quotient of a complex manifold by a group is not automatically a space in which holomorphy makes sense. What rescues the situation is that monodromy is integral: Γ sits inside $G_{\mathbb{Z}}$, which is discrete, while $G_{\mathbb{R}}$ acts on D with compact isotropy.

Proposition 5.3.5: The Monodromy Group Acts Properly Discontinuously

Let D be the period domain attached to $(V_{\mathbb{Z}}, Q, k, \{h^{p,q}\})$ and let $\Gamma \subseteq G_{\mathbb{Z}}$ be any subgroup. Then for every compact $C \subseteq D$ the set

$$\{\gamma \in \Gamma \,\big|\, \gamma C \cap C \neq \emptyset\}$$

is finite, and the quotient $\Gamma\backslash D$ is Hausdorff.

***Proof*:**
Step 1: $G_{\mathbb{Z}}$ meets every compact subset of $G_{\mathbb{R}}$ in a finite set. Choose a basis of $V_{\mathbb{Z}}$. In it the elements of $G_{\mathbb{Z}}$ are integer matrices, and $\mathrm{M}_m(\mathbb{Z})$ is a closed discrete subset of $\mathrm{M}_m(\mathbb{R})$; a closed discrete subset of a compact set is finite.

Step 2: the action of $G_{\mathbb{R}}$ on D is proper. Fix $F_o \in D$ and let P be its stabilizer in $G_{\mathbb{C}}$, so that by proposition 3.3.5 the orbit map $\pi\colon G_{\mathbb{R}} \to D$, $g \mapsto g\,F_o$, is surjective, identifies D with the quotient space $G_{\mathbb{R}}/K$, and has fibres the cosets of the compact group $K = G_{\mathbb{R}} \cap P$. The map π is open: for $W \subseteq G_{\mathbb{R}}$ open the saturation $\pi^{-1}(\pi(W)) = WK$ is open, so $\pi(W)$ is open in the quotient topology, which is the topology of D.

Let $C \subseteq D$ be compact. For each $x \in C$ choose $g_x \in G_{\mathbb{R}}$ with $\pi(g_x) = x$ and a relatively compact open $W_x \ni g_x$; the sets $\pi(W_x)$ form an open cover of C. Take a finite subcover and let $\tilde{C}$ be the union of the corresponding closures $\overline{W_x}$, a compact subset of $G_{\mathbb{R}}$ with $\pi(\tilde{C}) \supseteq C$.

If now $g \in G_{\mathbb{R}}$ satisfies $gC \cap C \neq \emptyset$, pick $x \in C$ with $gx \in C$ and write $x = \pi(c_1)$, $gx = \pi(c_2)$ with $c_1, c_2 \in \tilde{C}$. Then $gc_1K = c_2K$, so $g \in \tilde{C}K\tilde{C}^{-1}$, a compact subset of $G_{\mathbb{R}}$.

Step 3: conclusion. Combining the two steps, $\{\gamma \in \Gamma \,\big|\, \gamma C \cap C \neq \emptyset\}$ is contained in $G_{\mathbb{Z}} \cap \tilde{C}K\tilde{C}^{-1}$, which is finite.

For the Hausdorff property, let $x, y \in D$ lie in distinct Γ-orbits and choose compact neighbourhoods $N_x \ni x$ and $N_y \ni y$. Applying the finiteness just proved to the compact set $N_x \cup N_y$, only finitely many $\gamma \in \Gamma$ satisfy $\gamma N_x \cap N_y \neq \emptyset$; call them $\gamma_1, \dots, \gamma_r$. For each j one has $\gamma_j x \neq y$, since x and y lie in distinct orbits, so D being Hausdorff there are open $U_j \ni x$ and $V_j \ni y$ with $\gamma_j U_j \cap V_j = \emptyset$. Then $U = N_x^\circ \cap \bigcap_j U_j$ and $V = N_y^\circ \cap \bigcap_j V_j$ are open neighbourhoods of x and y with $\gamma U \cap V = \emptyset$ for every $\gamma \in \Gamma$, so their images separate the two orbits in $\Gamma\backslash D$, completing the proof.

5.3.6 Note: What "Holomorphic" Means on $\Gamma\backslash D$ Proposition 5.3.5 makes $\Gamma\backslash D$ a Hausdorff topological space, but the action need not be free: the stabilizer of a point of D in Γ is finite and may be nontrivial, and it is never trivial when $-\mathrm{id}_V \in \Gamma$ (exercise 5.3.1 (5)). Where Γ acts freely the quotient map $q\colon D \to \Gamma\backslash D$ is a holomorphic covering and $\Gamma\backslash D$ is a complex manifold; the same holds when only the quotient of Γ by the subgroup acting trivially does, since that quotient has the same orbits (exercise 5.3.1 (5)). In general $\Gamma\backslash D$ carries only the structure of a quotient with finite stabilizers. For that reason the statement "Φ is holomorphic" always refers to the lift: it means that $\tilde{\Phi}\colon \tilde{S} \to D$ is a holomorphic map of complex manifolds, equivalently that every local determination of the period map on a simply connected open subset of S, obtained by fixing a path class as in definition 5.3.3, is holomorphic. Nothing in this section depends on giving $\Gamma\backslash D$ more structure than that.

Holomorphy of the period map is the first place where the input of section 5.2 is used, and it is used in the weakest possible form: only that the F^p are holomorphic subbundles, not that the connection moves them by one step. In a flat frame a holomorphic subbundle is a holomorphically varying family of subspaces of a fixed vector space, and such a family is a holomorphic map to a flag manifold.

Theorem 5.3.7: The Period Map Is Holomorphic

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and let Φ be its period map in degree k.

1. Let $U \subseteq S$ be open, connected and simply connected, fix $s_1 \in U$ and a homotopy class $[c]$ of path from s_0 to s_1, and for $s \in U$ let $[c_s]$ be the class of c followed by the unique homotopy class of path from s_1 to s inside U. Then
$$\Phi_U\colon U \to D, \qquad \Phi_U(s) = \Phi_{[c_s]}(s)$$
is holomorphic.
2. The lifted period map $\tilde{\Phi}\colon \tilde{S} \to D$ is holomorphic. If Γ acts freely on D, then $\Gamma\backslash D$ is a complex manifold and $\Phi\colon S \to \Gamma\backslash D$ is holomorphic.

***Proof*:**

Step 1: a flat trivialization. Since U is simply connected, the local system $\mathbb{H}^k_{\text{prim}}|_U$ is trivial (proposition 5.1.5): parallel transport from s_1 along the unique homotopy class of path in U identifies every fibre with $H^k(X_{s_1},\mathbb{Q})_{\text{prim}}$, and composing with $T_{[c]}^{-1}$ identifies it with $V_{\mathbb{Q}}$. Tensoring with $\mathcal{O}_U$ gives an isomorphism of holomorphic bundles
$$\mathcal{H}^k_{\text{prim}}\big|_U \;\cong\; V_{\mathbb{C}} \otimes_{\mathbb{C}} \mathcal{O}_U$$
under which the flat sections are the constant ones and ∇ becomes the exterior derivative d (theorem 5.1.7). In this trivialization $\Phi_U(s)$ is the flag $\left(F^p_{\text{prim},s}\right)_p$ of subspaces of $V_{\mathbb{C}}$, of dimensions f^p independent of s by part (3) of proposition 5.3.2.

Step 2: an adapted holomorphic frame. Fix $s_2 \in U$. The subbundles F^p_{prim} are holomorphic and nested, so near s_2 there is a holomorphic frame $\sigma_1,\dots,\sigma_m$ of $V_{\mathbb{C}} \otimes \mathcal{O}$ adapted to the flag: choosing first a local holomorphic frame of the smallest nonzero step and extending it successively across each inclusion $F^{p+1}_{\text{prim}} \subseteq F^p_{\text{prim}}$, which is possible because a holomorphic subbundle of a holomorphic bundle is locally a direct summand, one obtains holomorphic sections $\sigma_1,\dots,\sigma_m$ of $V_{\mathbb{C}} \otimes \mathcal{O}$ such that $\sigma_1(s),\dots,\sigma_{f^p}(s)$ span $F^p_{\text{prim},s}$ for every p and every s near s_2. Shrinking the neighbourhood, the $\sigma_j(s)$ are a basis of $V_{\mathbb{C}}$ for every s.

Let $e_1,\dots,e_m$ be the basis $\sigma_j(s_2)$ of $V_{\mathbb{C}}$ and define $g(s) \in \mathrm{GL}(V_{\mathbb{C}})$ by $g(s)e_j = \sigma_j(s)$. Its matrix in the basis (e_j) has holomorphic entries and is invertible, so g is a holomorphic map into $\mathrm{GL}(V_{\mathbb{C}})$, and by construction $g(s)$ carries the flag $\Phi_U(s_2)$ to the flag $\Phi_U(s)$.

Step 3: into the flag manifold and into D. The group $\mathrm{GL}(V_{\mathbb{C}})$ acts holomorphically on Fl (section 3.3), so the composite $s \mapsto g(s)\cdot\Phi_U(s_2)$ is a holomorphic map from a neighbourhood of s_2 into Fl, and it equals Φ_U there. As $s_2 \in U$ was arbitrary, $\Phi_U\colon U \to \mathrm{Fl}$ is holomorphic.

Its image lies in D by part (4) of proposition 5.3.2. By proposition 3.3.5 the compact dual $\check{D}$ is a closed complex submanifold of Fl and D is open in it (proposition 3.3.4). A holomorphic map into a complex manifold whose image lies in a closed complex submanifold is holomorphic into that submanifold, since in a chart adapted to the submanifold the equations cutting it out are satisfied identically and the remaining coordinates of the map are holomorphic. Hence $\Phi_U\colon U \to D$ is holomorphic.

Step 4: globalizing. The projection $\pi\colon \tilde{S} \to S$ is a local biholomorphism, and each point of $\tilde{S}$ has a neighbourhood mapped biholomorphically onto a simply connected open $U \subseteq S$ on which $\tilde{\Phi}$ agrees with a map Φ_U of the kind treated above. So $\tilde{\Phi}$ is holomorphic.

Suppose finally that Γ acts freely. By proposition 5.3.5 it then acts freely and properly discontinuously by biholomorphisms, so $\Gamma\backslash D$ inherits a complex structure for which $q\colon D \to \Gamma\backslash D$ is a holomorphic covering. Since $\Phi \circ \pi = q \circ \tilde{\Phi}$ is holomorphic and π is a surjective local biholomorphism, Φ is holomorphic, as we sought to show.

The theorem is what makes the period map an object of complex geometry rather than of topology, and the proof shows precisely which hypothesis carries it: the holomorphy of the subbundles F^p, and nothing else. In particular the bigrading itself plays no role, which is the concrete payoff of the passage from bigradings to filtrations in theorem 3.1.6. The subspaces $H^{p,q}(X_s)$ vary only smoothly with s, and a map recording them would not be holomorphic; the sums F^p do vary holomorphically, and the map recording them is.

Holomorphy leaves the second half of section 5.2 unused. Griffiths transversality constrains the derivative, so what it constrains is the differential of Φ, and reading that constraint on the domain requires a description of the tangent spaces of Fl. Each step of a flag is a point of a Grassmannian, and the tangent space of a Grassmannian is $T_W\mathrm{Gr}(d,V) \cong \mathrm{Hom}(W, V/W)$, characterized by $dg_x(\xi)(\sigma(x)) = \partial_\xi \sigma \bmod g(x)$ for any local section σ of g, and $\mathrm{GL}(V)$-equivariant [32, the tangent space of a Grassmannian]. That description is used in exactly the form just stated.

Each step of a flag is a point of a Grassmannian, so a tangent vector to Fl has a component in each $\mathrm{Hom}(F^p, V_{\mathbb{C}}/F^p)$, and the derivative of a moving flag is the collection of the derivatives of its steps. Griffiths transversality bounds how far a step can move: not into all of $V_{\mathbb{C}}/F^p$, but only into the part coming from one step down.

Definition 5.3.8: The Horizontal Subbundle and the Infinitesimal Period Relation

For each p let $\pi_p\colon \mathrm{Fl} \to \mathrm{Gr}(f^p, V_{\mathbb{C}})$ be the holomorphic map $F \mapsto F^p$, and for a tangent vector $\theta \in T_F\mathrm{Fl}$ write

$$\theta_p := d(\pi_p)_F(\theta) \in \mathrm{Hom}\big(F^p,\ V_{\mathbb{C}}/F^p\big)$$

for its p-th component under the Grassmannian identification. The *horizontal subspace* of $\check{D}$ at $F \in \check{D}$ is

$$T^h_F\check{D} = \big\{\, \theta \in T_F\check{D} \;\big|\; \theta_p(F^p) \subseteq F^{p-1}/F^p \ \text{ for every } p \,\big\}$$

where F^{p-1}/F^p is regarded as a subspace of $V_{\mathbb{C}}/F^p$. Its restriction to D is the *horizontal subbundle* $T^hD \subseteq TD$.
A holomorphic map $g\colon M \to \check{D}$ from a complex manifold is *horizontal* if $dg_x(T_xM) \subseteq T^h_{g(x)}\check{D}$ for every $x \in M$. This condition on a map is classically called the *infinitesimal period relation.*

The horizontal subspaces do form a subbundle. The defining condition is stated purely in terms of the flag F, and by the equivariance clause of $\mathrm{GL}(V_{\mathbb{C}})$-equivariance an element $h \in G_{\mathbb{C}}$ carries $T^h_F\check{D}$ onto $T^h_{hF}\check{D}$; since $G_{\mathbb{C}}$ acts transitively on $\check{D}$ (proposition 3.3.5), the dimension of $T^h_F\check{D}$ is independent of F. The local holomorphic sections of the orbit map $G_{\mathbb{C}} \to \check{D}$ given by section 3.3 then transport a basis of $T^h_{F_o}\check{D}$ to a holomorphic frame of $T^h\check{D}$ near any point, so $T^h\check{D}$ is a holomorphic subbundle of $T\check{D}$.

Whether the inclusion $T^h\check{D} \subseteq T\check{D}$ is proper depends on the Hodge numbers, and in the two cases

the reader has already met it is not.

Example 5.3.9: The Horizontal Bundle in the Weight-Two K3 Case

Take $k = 2$ with $h^{2,0} = h^{0,2} = 1$ and $h^{1,1} = r$, the numerical type of example 3.3.6. A point of $\check{D}$ is a line $F^2 = \mathbb{C}\sigma$ with $Q(\sigma, \sigma) = 0$, and $F^1 = \sigma^\perp$ is determined by it, so

$$\check{D} \cong \big\{[\sigma] \in \mathbb{P}(V_\mathbb{C}) \big| Q(\sigma, \sigma) = 0\big\}$$

a smooth quadric hypersurface.

The tangent space. By the Grassmannian identification with $d = 1$, $T_{[\sigma]}\mathbb{P}(V_\mathbb{C}) = \mathrm{Hom}(\mathbb{C}\sigma, V_\mathbb{C}/\mathbb{C}\sigma)$. Differentiating the equation $Q(\sigma, \sigma) = 0$ along a holomorphic curve $[\sigma(s)]$ with section σ gives $2Q(\sigma, \partial\sigma) = 0$, so a tangent vector $\theta_2 \in \mathrm{Hom}(\mathbb{C}\sigma, V_\mathbb{C}/\mathbb{C}\sigma)$ is tangent to the quadric exactly when $Q(\sigma, \theta_2(\sigma)) = 0$; the quadric being smooth, this cuts out the tangent space exactly. Hence

$$T_{[\sigma]}\check{D} = \mathrm{Hom}\big(\mathbb{C}\sigma,\ \sigma^\perp/\mathbb{C}\sigma\big) = \mathrm{Hom}\big(F^2,\ F^1/F^2\big)$$

Horizontality. The condition at $p = 2$ reads $\theta_2(F^2) \subseteq F^1/F^2$, which the display just established for every tangent vector. The condition at $p = 1$ reads $\theta_1(F^1) \subseteq F^0/F^1 = V_\mathbb{C}/F^1$, which is no condition at all, and the levels $p \leq 0$ have $F^p = V_\mathbb{C}$ and $F^{p-1} = V_\mathbb{C}$. So $T^h\check{D} = T\check{D}$ here: a period map of this numerical type is horizontal for free. The same happens in weight 1, where the flag is a single subspace F^1 and the only condition that could bite reads $\theta_1(F^1) \subseteq F^0/F^1 = V_\mathbb{C}/F^1$, again vacuous; this is the period-domain form of the observation in example 5.2.9. Weight 2 with $h^{2,0} \geq 2$ already behaves differently (exercise 5.3.1 (3)).

With the horizontal subbundle in place, Griffiths transversality translates verbatim, and the translation changes what the statement is about. On the base it constrains a connection acting on subbundles, so it refers to the family; on the domain it says that the period map is everywhere tangent to a fixed distribution, one attached to the numerical data alone and shared by every family with those Hodge numbers.

Theorem 5.3.10: The Period Map Is Horizontal

Let $f\colon \mathcal{X} \to S$ be a smooth projective family and let $\tilde{\Phi}\colon \tilde{S} \to D$ be its lifted period map in degree k. Then $\tilde{\Phi}$ is horizontal:

$$d\tilde{\Phi}_{\tilde{s}}\big(T_{\tilde{s}}\tilde{S}\big) \subseteq T^h_{\tilde{\Phi}(\tilde{s})}D \qquad \text{for every } \tilde{s} \in \tilde{S}$$

Equivalently, every local determination Φ_U of the period map on a simply connected open subset of S, as in theorem 5.3.7, satisfies the infinitesimal period relation.

***Proof*:**

The statement is local on S, since $\pi\colon \tilde{S} \to S$ is a local biholomorphism and $\tilde{\Phi}$ agrees with a local determination Φ_U near every point. So let $U \subseteq S$ be open, connected and simply connected, fix the flat trivialization $\mathcal{H}^k_{\mathrm{prim}}|_U \cong V_\mathbb{C} \otimes_\mathbb{C} \mathcal{O}_U$ of Step 1 of the proof of theorem 5.3.7, and write Φ_U for the resulting holomorphic map $U \to D$. Fix $s \in U$ and $\xi \in T_sU$.

Step 1: the components of $d\Phi_U$. Fix p. The composite $\pi_p \circ \Phi_U$ is the map $U \to \mathrm{Gr}(f^p, V_\mathbb{C})$

sending s' to $F^p_{\mathrm{prim},s'}$, and the holomorphic maps $\sigma\colon U' \to V_{\mathbb{C}}$ with $\sigma(s') \in F^p_{\mathrm{prim},s'}$ for all s' are exactly the local holomorphic sections of the subbundle F^p_{prim}, read in the trivialization. By the derivative formula characterizing the Grassmannian tangent space, for such a σ,

$$\big(d(\Phi_U)_s(\xi)\big)_p\big(\sigma(s)\big) = \partial_\xi \sigma \bmod F^p_{\mathrm{prim},s}$$

Because F^p_{prim} is a holomorphic subbundle, every vector of $F^p_{\mathrm{prim},s}$ is $\sigma(s)$ for some such σ, so the displayed formula determines the component $\big(d(\Phi_U)_s(\xi)\big)_p$ completely.

Step 2: the derivative is the connection. In the flat trivialization the flat sections are the constant ones, so ∇ acts as the exterior derivative: writing $\sigma = \sum_j g_j e_j$ with e_j a constant frame, theorem 5.1.7 gives $\nabla_\xi \sigma = \sum_j (\partial_\xi g_j)\, e_j = \partial_\xi \sigma$. Hence

$$\big(d(\Phi_U)_s(\xi)\big)_p\big(\sigma(s)\big) = \nabla_\xi \sigma \bmod F^p_{\mathrm{prim},s}$$

for every local holomorphic section σ of F^p_{prim}.

Step 3: transversality. By part (3) of proposition 5.3.2, which is Griffiths transversality (theorem 5.2.7) restricted to the primitive part, $\nabla_\xi \sigma$ lies in $F^{p-1}_{\mathrm{prim},s}$. Therefore

$$\big(d(\Phi_U)_s(\xi)\big)_p\big(F^p_{\mathrm{prim},s}\big) \subseteq F^{p-1}_{\mathrm{prim},s}/F^p_{\mathrm{prim},s}$$

for every p, which is exactly the condition defining $T^h_{\Phi_U(s)}\check{D}$ in definition 5.3.8. Since $d(\Phi_U)_s(\xi)$ is tangent to D and D is open in $\check{D}$ (proposition 3.3.4), it lies in $T^h_{\Phi_U(s)}D$, as we sought to show.

The proof is short because all the content was already extracted. Step 1 is linear algebra, Step 2 is the definition of the Gauss-Manin connection, and Step 3 is the theorem of section 5.2; nothing about the family enters that was not already in proposition 5.3.2. Read backwards, the theorem also says that transversality is not a property of any particular family: it is a property of the numerical data, expressed by a distribution on $\check{D}$ that every period map with those Hodge numbers must be tangent to. The geometry of D that this distribution carries, and the curvature estimates along it that turn horizontality into rigidity statements about families, are taken up in chapter 6.

The graded form of the differential is the primitive counterpart of the map isolated in corollary 5.2.8. By Step 3 the component $\big(d\Phi_U(\xi)\big)_p$ takes F^p_{prim} into $F^{p-1}_{\mathrm{prim}}/F^p_{\mathrm{prim}}$, and it kills F^{p+1}_{prim} modulo F^p_{prim}, since a section of F^{p+1}_{prim} has $\nabla_\xi \sigma \in F^p_{\mathrm{prim}}$ by transversality one level up. It therefore factors through a map $F^p_{\mathrm{prim}}/F^{p+1}_{\mathrm{prim}} \to F^{p-1}_{\mathrm{prim}}/F^p_{\mathrm{prim}}$, and as ξ varies these assemble into the $\mathcal{O}_S$-linear map

$$F^p_{\mathrm{prim}}/F^{p+1}_{\mathrm{prim}} \to \big(F^{p-1}_{\mathrm{prim}}/F^p_{\mathrm{prim}}\big) \otimes \Omega^1_S$$

This is the construction of corollary 5.2.8 run on the primitive filtration $F^\bullet_{\mathrm{prim}}$ instead of on $F^\bullet$, which is legitimate because that construction used nothing but transversality, and part (3) of proposition 5.3.2 gives transversality for the primitive part. Equivalently it is the restriction of the corollary's map to the primitive graded pieces, a strictly smaller map whenever the primitive cohomology is proper in H^k: on a fibre it reads $H^{p,q}_{\mathrm{prim}} \to H^{p-1,q+1}_{\mathrm{prim}} \otimes T^*_s S$. So the derivative of the period map is a collection of maps between adjacent primitive Hodge pieces of a single fibre, and computing it never requires leaving that fibre.

When T^hD is a proper subbundle, horizontality is a genuine restriction on what a family can do, and it restricts by dimension. A period map can never be submersive onto an open subset of D in that case, and if $d\Phi$ is injective at some point, which is the local Torelli condition, then $\dim_{\mathbb{C}} S$ is at most the rank of T^hD (exercise 5.3.1 (4)). The first numerical type where this bites is weight 2 with $h^{2,0} = 2$, where the horizontal bundle has codimension 1 (exercise 5.3.1 (3)); the effect grows quickly with the number of nonzero Hodge numbers.

Exercise 5.3.1

1. Let $f\colon \mathcal{X} \to S$ be a smooth projective family with period map Φ in degree k. Show that Φ is constant if and only if $\nabla F^p_{\text{prim}} \subseteq F^p_{\text{prim}} \otimes \Omega^1_S$ for every p, that is, if and only if every step of the Hodge filtration of the primitive part is a flat subbundle. Deduce that in that case parallel transport along any path from s_0 to s is an isomorphism of polarized Hodge structures from $H^k(X_{s_0}, \mathbb{Q})_{\text{prim}}$ to $H^k(X_s, \mathbb{Q})_{\text{prim}}$. (Nothing follows about the Hodge structure on the whole of $H^k(X_s)$, which the period map in degree k does not see.)

2. In the setting of example 5.3.4, suppose the monodromy of a loop in $\pi_1(S, s_0)$ replaces the flat frame (a, b) of H_1 by $(\kappa a + \lambda b,\ \mu a + \nu b)$ with $\kappa, \lambda, \mu, \nu \in \mathbb{Z}$ and $\kappa\nu - \lambda\mu = 1$.

   1. Show that the period ratio computed in the new frame is $\tau' = (\mu + \nu\tau)/(\kappa + \lambda\tau)$, and that $\kappa + \lambda\tau \neq 0$.
   2. Verify directly that $\operatorname{Im}\tau' > 0$, so that the substitution preserves $\mathfrak{H}_1$.
   3. Recover the substitution $\tau \mapsto \tau + 1$ of example 4.4.8 as a special case, and identify the matrix.

3. Take $k = 2$ with $h^{2,0} = h^{0,2} = 2$ and $h^{1,1} = r$, so that $m = r + 4$ and Q is symmetric and nondegenerate.

   1. Show that $F \mapsto F^2$ identifies $\check{D}$ with the set of 2-dimensional isotropic subspaces $W \subseteq V_{\mathbb{C}}$, with $F^1 = W^{\perp}$ determined by W.
   2. Using the Grassmannian tangent-space identification [32, the tangent space of a Grassmannian], show that

      $$T_W\check{D} = \big\{\theta \in \operatorname{Hom}(W, V_{\mathbb{C}}/W) \;\big|\; \beta_\theta \text{ is alternating}\big\}, \qquad \beta_\theta(w, w') := Q\big(\theta(w), w'\big)$$

      (note that β_θ is well defined because W is isotropic), and that $\theta \mapsto \beta_\theta$ maps $T_W\check{D}$ onto $\Lambda^2 W^*$ with kernel $\operatorname{Hom}(W, W^{\perp}/W)$.
   3. Deduce that $T^h_W\check{D} = \operatorname{Hom}(W, W^{\perp}/W)$ and that it has codimension 1 in $T_W\check{D}$. What happens when $h^{2,0} = 1$?

4. Let Φ be the period map of a smooth projective family over a connected S, and suppose $d\tilde{\Phi}$ is injective at some point of $\tilde{S}$.

   1. Show that $\dim_{\mathbb{C}} S \leq \operatorname{rank} T^hD$.
   2. Show that if $T^hD \neq TD$, then $\tilde{\Phi}$ is nowhere a local biholomorphism onto an open subset of D, whatever the dimension of S.
   3. Combine (1) with exercise 5.3.1 (3) to bound $\dim_{\mathbb{C}} S$ for a family of the numerical type considered there, in terms of r.

5. Let D be a period domain attached to data $(V_{\mathbb{Z}}, Q, k, \{h^{p,q}\})$.

    1. Show that $-\mathrm{id}_V$ lies in $G_{\mathbb{Z}}$ and acts trivially on $\check{D}$, so the action of $G_{\mathbb{R}}$ on D is never effective.
    2. Deduce that if $-\mathrm{id}_V \in \Gamma$, then no point of D has trivial Γ-stabilizer, so Γ never acts freely and part (2) of theorem 5.3.7 never applies to Γ itself. Show nevertheless that $\Gamma\backslash D = \bar{\Gamma}\backslash D$ for $\bar{\Gamma} = \Gamma/\{\pm\mathrm{id}_V\}$, that $\bar{\Gamma}$ inherits the conclusion of proposition 5.3.5, and hence that $q\colon D \to \Gamma\backslash D$ is a covering map as soon as $\bar{\Gamma}$ acts freely, with deck group $\bar{\Gamma}$ rather than Γ.
    3. Identify the phenomenon in weight 1 with $h^{1,0} = 1$, using example 3.3.3.

5.4 Hypersurfaces and the Jacobian Ring

The three preceding sections attach to a smooth projective family a local system, a filtration by holomorphic subbundles and a period map. None of them computes any of it. This section computes all of it for one family, the smooth hypersurfaces of a fixed degree in a fixed projective space, where the answer is a statement about polynomials: the Hodge filtration on the primitive cohomology of $X = \{F = 0\}$ is read off the graded ring $\mathbb{C}[x_0, \dots, x_{n+1}]/J(F)$, where $J(F)$ is the ideal generated by the partial derivatives of F.

Two conventions before starting. The defining polynomial is written F and the Hodge filtration F^p; the superscript is what distinguishes them. And the cohomology in play is always the primitive part in the middle degree: by example 4.3.2 every other cohomology group of a smooth hypersurface is that of the ambient projective space, so a family of hypersurfaces can vary nothing outside H^n_{prim}.

Throughout, $n \geq 1$ and $d \geq 2$ are fixed, $\mathbb{C}[x] = \mathbb{C}[x_0, \dots, x_{n+1}]$ is the polynomial ring in $n+2$ variables with its usual grading, and $\mathbb{C}[x]_e$ denotes the space of homogeneous polynomials of degree e, taken to be 0 for $e < 0$. For $0 \neq F \in \mathbb{C}[x]_d$ write

$$X_F = \{F = 0\} \subseteq \mathbb{C}\mathrm{P}^{n+1}, \qquad U_F = \mathbb{C}\mathrm{P}^{n+1} \setminus X_F$$

and abbreviate these to X and U when F is fixed. Partial derivatives are written $\partial_i = \partial/\partial x_i$.

The object that carries the whole theory is the ideal generated by those derivatives. Its first role is to detect smoothness, through the Euler relation $\sum_i x_i\, \partial_i F = d \cdot F$, which is what ties the vanishing of the derivatives to the vanishing of F itself.

Definition 5.4.1: The Jacobian Ideal and the Jacobian Ring

Let $F \in \mathbb{C}[x]_d$. The *Jacobian ideal* of F is the homogeneous ideal

$$J(F) = \big(\partial_0 F,\ \partial_1 F,\ \dots,\ \partial_{n+1} F\big) \subseteq \mathbb{C}[x]$$

generated by the $n+2$ partial derivatives, each of degree $d-1$. The *Jacobian ring* of F is the quotient

$$R(F) = \mathbb{C}[x]/J(F)$$

with its induced grading $R(F) = \bigoplus_{e \geq 0} R(F)_e$, where $R(F)_e = \mathbb{C}[x]_e/J(F)_e$. When F is fixed the ring is written R.

Because $J(F)$ is generated in degree $d-1$, the low-degree pieces of R are not quotients at all: $R_e = \mathbb{C}[x]_e$ for every $e \leq d-2$. The interesting behaviour is in the middle range, and it is there that smoothness of X_F makes itself felt.

Proposition 5.4.2: Smoothness and Finite-Dimensionality of the Jacobian Ring

Let $F \in \mathbb{C}[x]_d$ be nonzero, with $d \geq 1$. The following are equivalent.

1. The derivatives $\partial_0 F, \ldots, \partial_{n+1} F$ have no common zero in $\mathbb{C}\mathrm{P}^{n+1}$.
2. At every point of X_F the dehomogenization of F in some, hence any, affine chart containing that point has nonvanishing differential.

When they hold, F is called *smooth* and X_F is a complex submanifold of $\mathbb{C}\mathrm{P}^{n+1}$ of dimension n; moreover $R(F)$ is a finite-dimensional $\mathbb{C}$-vector space, so $R(F)_e = 0$ for all sufficiently large e.

***Proof*:**

Step 1: (1) implies (2). Let $x \in X_F$, choose j with $x_j \neq 0$ and work in the affine chart $x_j = 1$ with coordinates $u_k = x_k/x_j$ for $k \neq j$, in which X_F is the zero set of $f(u) = F(u)|_{u_j=1}$ and $\partial f/\partial u_k = (\partial_k F)|_{u_j=1}$. If every $\partial_k F$ with $k \neq j$ vanished at x, then the Euler relation

$$\sum_k x_k \, \partial_k F = d \cdot F$$

evaluated at x would give $x_j \, \partial_j F(x) = d \cdot F(x) = 0$ and hence $\partial_j F(x) = 0$ as well, so all $n+2$ derivatives would vanish at x, contrary to (1). So $df_x \neq 0$, and the implicit function theorem presents X_F near x as a complex submanifold of codimension one.

Step 2: (2) implies (1). Suppose the derivatives have a common zero $x \in \mathbb{C}\mathrm{P}^{n+1}$. The Euler relation gives $d \cdot F(x) = 0$, so $F(x) = 0$ and $x \in X_F$; and in a chart around x every partial derivative of the dehomogenization is a restriction of some $\partial_k F$ and so vanishes at x, contradicting (2).

Step 3: finite-dimensionality. Dispose first of $d = 1$, where the derivatives are constants, not all zero because $F \neq 0$: there $J(F) = \mathbb{C}[x]$ and $R = 0$. So assume $d \geq 2$, in which case the derivatives are homogeneous of positive degree and vanish at the origin. By (1) they have no common zero in $\mathbb{C}\mathrm{P}^{n+1}$, so their common zero locus in $\mathbb{C}^{n+2}$ is exactly the origin. The Nullstellensatz [4, Hilbert's Nullstellensatz] gives $\sqrt{J(F)} = (x_0, \ldots, x_{n+1})$, so there is $k \geq 1$ with $x_i^k \in J(F)$ for every i. Consequently R is spanned by the images of the monomials x^a with every exponent $a_i < k$, of which there are k^{n+2}, and $R_e = 0$ once $e > (n+2)(k-1)$, completing the proof.

Condition (2) is a condition on F and not only on the point set X_F. For $F = x_0^2$ in $\mathbb{C}\mathrm{P}^2$ the zero set is a line, hence a smooth curve, while every partial derivative of F vanishes along it; that F is not smooth in the sense above, and its Jacobian ring $\mathbb{C}[x_0, x_1, x_2]/(x_0)$ is infinite dimensional, as the last part of the proposition requires. From here on, smoothness of a hypersurface means smoothness of its defining polynomial.

The last assertion says that the Jacobian ring of a smooth hypersurface is a finite package of numbers. Theorem 5.4.11 will say that this package is the Hodge structure of X, so it is worth having one case

in which every graded piece can be written down.

Example 5.4.3: The Jacobian Ring of a Fermat Hypersurface

Let $F = x_0^d + x_1^d + \cdots + x_{n+1}^d$, the *Fermat* polynomial of degree $d \geq 2$. Then $\partial_i F = d\,x_i^{d-1}$, and these vanish simultaneously only at the origin of $\mathbb{C}^{n+2}$, so X_F is smooth by proposition 5.4.2. The Jacobian ideal is the monomial ideal

$$J(F) = \left(x_0^{d-1},\ x_1^{d-1},\ \ldots,\ x_{n+1}^{d-1}\right)$$

and a monomial x^a lies in $J(F)$ exactly when $a_i \geq d-1$ for some i. Since $J(F)$ is spanned by the monomials it contains, the images of the remaining monomials form a basis of R:

$$R \ \text{ has basis } \ \left\{\, x^a \ :\ 0 \leq a_i \leq d-2 \text{ for every } i \,\right\}$$

Hence $\dim_{\mathbb{C}} R_e$ is the number of ways to write $e = a_0 + \cdots + a_{n+1}$ with every a_i in $\{0, 1, \ldots, d-2\}$, that is,

$$\sum_{e \geq 0} (\dim_{\mathbb{C}} R_e)\, t^e = \left(1 + t + \cdots + t^{d-2}\right)^{n+2} \tag{5.4}$$

Three consequences are immediate from (5.4). The top nonzero degree is $\sigma = (n+2)(d-2)$, attained only by the single monomial with all $a_i = d-2$, so $\dim R_\sigma = 1$. The map $a \mapsto (d-2, \ldots, d-2) - a$ is a bijection of the basis carrying degree e to degree $\sigma - e$, so

$$\dim R_e = \dim R_{\sigma - e} \qquad (0 \leq e \leq \sigma)$$

And for $d = 2$ the polynomial (5.4) is 1: the Jacobian ring of a smooth quadric is $\mathbb{C}$, concentrated in degree 0.

For $n = 2$ and $d = 4$, a Fermat quartic surface in $\mathbb{CP}^3$, the series is $(1 + t + t^2)^4 = 1 + 4t + 10t^2 + 16t^3 + 19t^4 + 16t^5 + 10t^6 + 4t^7 + t^8$, with $\sigma = 8$.

Every smooth hypersurface of degree d now sits in one family, and it is that family to which sections 5.1 to 5.3 apply. Write $M = \dim_{\mathbb{C}} \mathbb{C}[x]_d - 1$ and identify $\mathbb{CP}^M$ with the projective space of nonzero forms of degree d up to scale, a point of which is written $[F]$.

Proposition 5.4.4: The Family of Smooth Hypersurfaces of Fixed Degree

Let $n \geq 1$, $d \geq 2$, and let

$$S = \left\{ [F] \in \mathbb{CP}^M \ :\ F \text{ is smooth in the sense of proposition 5.4.2} \right\}$$

Then S is a nonempty connected Zariski-open subset of $\mathbb{CP}^M$; the incidence variety

$$\mathcal{X} = \left\{ (x, [F]) \in \mathbb{CP}^{n+1} \times S \ :\ F(x) = 0 \right\}, \qquad f \colon \mathcal{X} \to S \ \text{ the second projection}$$

is a smooth projective family with fibre $f^{-1}([F]) = X_F$; and every point of S has a neighbourhood V over which the pair $(\mathbb{CP}^{n+1} \times V, \mathcal{X}|_V)$ is diffeomorphic to $(\mathbb{CP}^{n+1}, X_{F_0}) \times V$ over V.

***Proof*:**
Step 1: S is open, nonempty and connected. Let

$$\Sigma = \big\{(x,[F]) \in \mathbb{C}\mathrm{P}^{n+1} \times \mathbb{C}\mathrm{P}^{M} \;:\; \partial_0 F(x) = \cdots = \partial_{n+1} F(x) = 0\big\}$$

which is closed, being cut out by equations bihomogeneous in x and in the coefficients of F. Its image $\Delta \subseteq \mathbb{C}\mathrm{P}^M$ under the second projection is closed, since $\mathbb{C}\mathrm{P}^{n+1}$ is complete [4, Projective Morphisms are Proper], and by proposition 5.4.2 a point $[F]$ lies in Δ exactly when F fails to be smooth. So $S = \mathbb{C}\mathrm{P}^M \setminus \Delta$ is Zariski open, and it is nonempty because the Fermat polynomial of example 5.4.3 lies in it. For connectedness, let $[F_0],[F_1] \in S$ and let $\ell \cong \mathbb{C}\mathrm{P}^1$ be the line they span. Then $\ell \not\subseteq \Delta$, since $[F_0] \notin \Delta$, so $\ell \cap \Delta$ is a proper Zariski-closed subset of ℓ and hence finite; and ℓ with finitely many points removed is path connected and contained in S.

Step 2: f is a proper submersion with connected fibres. This is example 5.1.2(2), which set up exactly this incidence variety and checked the submersion criterion there; the argument is repeated in outline only because the smoothness locus is now described through proposition 5.4.2. For fixed x the condition $F(x) = 0$ is one nonzero linear condition on the coefficients of F, so the first projection exhibits the incidence variety inside $\mathbb{C}\mathrm{P}^{n+1} \times \mathbb{C}\mathrm{P}^M$ as a bundle with fibre $\mathbb{C}\mathrm{P}^{M-1}$; in particular it is a complex manifold, and $\mathcal{X}$ is the open subset lying over S. The map f is proper, being the restriction to a closed subset of the projection $\mathbb{C}\mathrm{P}^{n+1} \times S \to S$ with compact fibre $\mathbb{C}\mathrm{P}^{n+1}$.

For submersivity, fix $(x,[F]) \in \mathcal{X}$ and a tangent vector to $\mathbb{C}\mathrm{P}^M$ at $[F]$, represented by $G \in \mathbb{C}[x]_d$. Since X_F is smooth, $dF_x \neq 0$ in a chart around x, so the implicit function theorem solves $F_t(x_t) = 0$ for a holomorphic curve x_t with $x_0 = x$, where $F_t = F + tG$. The curve $t \mapsto (x_t,[F_t])$ lies in $\mathcal{X}$ and covers the given tangent vector, so df is surjective. Finally $H^0(X_F,\mathbb{Z}) = \mathbb{Z}$ by example 4.3.2, since $n \geq 1$, so the fibres are connected.

Step 3: local triviality of the pair. Fix $[F_0] \in S$ and a neighbourhood $V \ni [F_0]$ identified with an open cube in $\mathbb{R}^N$ by real coordinates $t_1,\dots,t_N$ vanishing at $[F_0]$. For each coordinate vector field ∂_{t_j} on V choose a smooth vector field w_j on $\mathbb{C}\mathrm{P}^{n+1} \times V$ that projects to ∂_{t_j} and is tangent to $\mathcal{X}$: on $\mathcal{X}$ such a lift exists because f is a submersion, it extends to a neighbourhood of $\mathcal{X}$, and it is patched with an arbitrary lift on the complement by a partition of unity, which preserves the property of projecting to ∂_{t_j} because the lifts of a fixed vector field form an affine space.

The flow of w_j is not complete: the trajectory through (x,t) has base component $t + s\,e_j$, with e_j the j-th standard basis vector, and that leaves V once $|s|$ is large. What compactness of $\mathbb{C}\mathrm{P}^{n+1}$ gives is completeness in the fibre direction, which is enough. Suppose the maximal interval of a trajectory has a finite endpoint s_+ with $t + s_+ e_j$ still in V, and let $L \subseteq V$ be the closed segment from t to $t + s_+ e_j$. The trajectory stays in the compact set $\mathbb{C}\mathrm{P}^{n+1} \times L$, on which w_j is bounded, so it has bounded speed and converges as $s \to s_+$; restarting the flow at its limit extends it past s_+, contradicting maximality. So the time-s flow Θ^j_s of w_j is defined on $\mathbb{C}\mathrm{P}^{n+1} \times \{t\}$ whenever the segment from t to $t + s\,e_j$ lies in V, and it carries that fibre onto $\mathbb{C}\mathrm{P}^{n+1} \times \{t + s\,e_j\}$ and $\mathcal{X}$ to $\mathcal{X}$.

Now set

$$\Theta\big(x,(t_1,\dots,t_N)\big) \;=\; \Theta^N_{t_N} \circ \cdots \circ \Theta^1_{t_1}(x,0)$$

Every segment traversed here is a coordinate segment inside the cube, so Θ is defined on all of $\mathbb{C}\mathrm{P}^{n+1} \times V$; it is smooth, covers the identity of V, and is inverted by running the flows backwards in the opposite order, so it is a diffeomorphism of $\mathbb{C}\mathrm{P}^{n+1} \times V$ carrying $X_{F_0} \times V$ onto $\mathcal{X}|_V$. That is the asserted trivialization of the pair; this is the argument that proves Ehresmann's

theorem [32, Ehresmann's Theorem], run with the extra requirement that the lifted field be tangent to $\mathcal{X}$, completing the proof.

Proposition 5.4.4 puts the whole apparatus of this chapter at the disposal of hypersurfaces. The local system $\mathbb{H}^n = R^n f_* \mathbb{Q}$ of definition 5.1.3 carries the Gauss-Manin connection ∇ of definition 5.1.6; the Hodge filtration $F^\bullet$ consists of holomorphic subbundles of $\mathcal{H}^n$ by theorem 5.2.6 and satisfies Griffiths transversality by theorem 5.2.7; and the period map Φ of definition 5.3.3 is holomorphic and horizontal. The polarization is by the hyperplane class h_X, the restriction of the Fubini-Study class, and proposition 3.2.7 makes $H^n_{\text{prim}}(X, \mathbb{Q}) = \ker\big(h_X \smile -\big)$ a sub-Hodge structure of $H^n(X, \mathbb{Q})$. Everything that varies is inside it.

The route to computing that Hodge structure runs through the complement rather than through X itself. The reason is that $U = \mathbb{C}\mathrm{P}^{n+1} \setminus X$ is affine (the d-uple Veronese embedding of example 4.3.2 carries X to a hyperplane section, so U becomes a closed submanifold of an affine space), and on an affine variety cohomology classes have global rational representatives. Those representatives can be written down.

Set

$$\Omega \;=\; \sum_{i=0}^{n+1} (-1)^i \, x_i \; dx_0 \wedge \cdots \wedge \widehat{dx_i} \wedge \cdots \wedge dx_{n+1}$$

the contraction of the Euler vector field $E = \sum_i x_i \partial_i$ into $dx_0 \wedge \cdots \wedge dx_{n+1}$, a holomorphic $(n+1)$-form on $\mathbb{C}^{n+2}$ whose coefficients are homogeneous of degree 1.

Proposition 5.4.5: Rational Forms with Poles Along a Hypersurface

Let $X = X_F \subseteq \mathbb{C}\mathrm{P}^{n+1}$ be a smooth hypersurface of degree d and let $m \geq 1$. The assignment

$$A \;\longmapsto\; \frac{A\,\Omega}{F^m}$$

is a linear isomorphism from $\mathbb{C}[x]_{md-n-2}$ onto the space of meromorphic $(n+1)$-forms on $\mathbb{C}\mathrm{P}^{n+1}$ that are holomorphic on U and have a pole of order at most m along X, that is, forms η such that $f^m \eta$ is holomorphic near each point of X, where f is a local defining function for X.

***Proof*:**

Step 1: Ω trivializes $\Omega^{n+1}_{\mathbb{C}\mathrm{P}^{n+1}}(n+2)$. Work in the chart $x_0 \neq 0$ with affine coordinates $u_i = x_i/x_0$, so that $du_i = (x_0\,dx_i - x_i\,dx_0)/x_0^2$. Wedging these,

$$du_1 \wedge \cdots \wedge du_{n+1} = x_0^{-2(n+1)} \bigwedge_{i=1}^{n+1} \big(x_0\,dx_i - x_i\,dx_0\big)$$

Expanding the product, the terms containing two or more factors of dx_0 vanish, so what survives is $x_0^{n+1} dx_1 \wedge \cdots \wedge dx_{n+1}$ together with, for each i, the term $-x_0^n x_i\, dx_1 \wedge \cdots \wedge dx_0 \wedge \cdots \wedge dx_{n+1}$ with dx_0 in the i-th slot; moving that dx_0 to the front costs $(-1)^{i-1}$. The bracket is therefore $x_0^n\,\Omega$, and

$$\Omega = x_0^{n+2}\; du_1 \wedge \cdots \wedge du_{n+1} \tag{5.5}$$

The same computation holds in every coordinate chart, so Ω vanishes nowhere on $\mathbb{C}^{n+2} \setminus \{0\}$ and (5.5) exhibits it as a nowhere-vanishing holomorphic section of $\Omega^{n+1}_{\mathbb{C}\mathrm{P}^{n+1}}(n+2)$; this is a

restatement of $K_{\mathbb{C}\mathrm{P}^{n+1}} = \mathcal{O}(-n-2)$ [5, Canonical Bundle of $\mathbb{C}\mathrm{P}^n$].

Step 2: the map is well defined and injective. Let $A \in \mathbb{C}[x]_{md-n-2}$. In the chart $x_0 \neq 0$, (5.5) and $f = F/x_0^d$ give

$$\frac{A\,\Omega}{F^m} = \frac{A\,x_0^{n+2}}{F^m}\; du_1 \wedge \cdots \wedge du_{n+1} = \frac{A\,x_0^{n+2-md}}{f^m}\; du_1 \wedge \cdots \wedge du_{n+1} \tag{5.6}$$

whose coefficient is f^{-m} times a holomorphic function of u, because $A\,x_0^{n+2-md}$ is homogeneous of degree 0 and holomorphic where $x_0 \neq 0$. So $A\Omega/F^m$ is a meromorphic $(n+1)$-form on $\mathbb{C}\mathrm{P}^{n+1}$, holomorphic on U, with pole order at most m along X. It is nonzero for $A \neq 0$ because Ω vanishes nowhere, so the map is injective.

Step 3: surjectivity. Let η be a form of the stated kind. In the chart $x_0 \neq 0$ write $\eta = \varphi\, du_1 \wedge \cdots \wedge du_{n+1}$; the hypothesis says that $g := f^m\varphi$ is holomorphic on the whole chart. Reading (5.6) backwards, $\eta = A_0\Omega/F^m$ on that chart with

$$A_0 = g \cdot x_0^{md-n-2}$$

a holomorphic function on $\{x_0 \neq 0\}$, homogeneous of degree $md-n-2$. The same computation in the chart $x_j \neq 0$ produces A_j, and $A_i = A_j$ on overlaps because Ω vanishes nowhere; so the A_j glue to a holomorphic function A on $\mathbb{C}^{n+2} \setminus \{0\}$, homogeneous of degree $e = md-n-2$. Expanding g as a power series $\sum_\alpha c_\alpha u^\alpha$ converging on all of $\mathbb{C}^{n+1}$ gives $A = \sum_\alpha c_\alpha\, x^\alpha x_0^{e-|\alpha|}$, and holomorphy of A at points with $x_0 = 0$ and some other coordinate nonzero forces $c_\alpha = 0$ whenever $|\alpha| > e$. Hence A is a homogeneous polynomial of degree e, and $A = 0$ when $e < 0$, as we sought to show.

Such a form is automatically closed: it is a holomorphic $(n+1)$-form on the complex manifold U of dimension $n+1$, so its exterior derivative, of type $(n+2, 0)$, vanishes. Each therefore defines a class in $H^{n+1}(U, \mathbb{C})$, and the whole problem becomes one of transporting classes from U back to X. The transport is the residue, and the definition used here is topological.

Definition 5.4.6: The Tube Map and the Residue

Let $X \subseteq \mathbb{C}\mathrm{P}^{n+1}$ be a smooth hypersurface with complement U. Fix a closed tubular neighbourhood $\pi\colon \mathcal{N} \to X$ of X in $\mathbb{C}\mathrm{P}^{n+1}$ [32, Tubular Neighbourhood Theorem], a closed disc bundle whose boundary $\partial\mathcal{N}$ is a circle bundle contained in U, with the fibre circles oriented positively for the complex structure of the normal direction. The *tube map*

$$\tau\colon H_k(X,\mathbb{Z}) \to H_{k+1}(U,\mathbb{Z})$$

is the composite

$$H_k(X,\mathbb{Z}) \;\cong\; H_{k+2}(\mathcal{N},\partial\mathcal{N};\mathbb{Z}) \;\cong\; H_{k+2}(\mathbb{C}\mathrm{P}^{n+1},U;\mathbb{Z}) \xrightarrow{\;\partial\;} H_{k+1}(U,\mathbb{Z})$$

whose first arrow is Poincaré duality on X followed by Lefschetz duality on $\mathcal{N}$, whose second is excision, and whose third is the connecting homomorphism of the pair. On the class of a cycle γ carried by a submanifold of X it is the class of the restriction of the circle bundle $\partial\mathcal{N}$ over γ. The *residue*

$$\mathrm{Res}\colon H^{k+1}(U,\mathbb{C}) \to H^k(X,\mathbb{C})$$

is the map dual to τ, normalized by

$$\int_\gamma \mathrm{Res}\,\eta \;=\; \frac{1}{2\pi i}\int_{\tau(\gamma)} \eta \qquad \text{for all } \gamma \in H_k(X,\mathbb{C})$$

The normalization is chosen so that the residue of a form with a first-order pole is computed by the same recipe as a residue in one variable, with the $2\pi i$ already divided out.

Proposition 5.4.7: The Residue of a Form with a First-Order Pole

Let η be a closed $(k+1)$-form on U for which there is an open cover $\{W_\alpha\}$ of a neighbourhood of X in $\mathbb{C}\mathrm{P}^{n+1}$ and expressions

$$\eta\big|_{W_\alpha \cap U} = \frac{dg_\alpha}{g_\alpha} \wedge \psi_\alpha + \theta_\alpha$$

with g_α a holomorphic defining function for X on W_α and $\psi_\alpha, \theta_\alpha$ holomorphic forms on W_α. Then the pullbacks of the ψ_α to X agree on overlaps and glue to a closed holomorphic k-form ψ on X, and

$$\mathrm{Res}\,\eta = [\psi] \quad \text{in } H^k(X,\mathbb{C})$$

***Proof*:**
Step 1: the local pieces glue. On an overlap write $g' = ug$ with u a nowhere-zero holomorphic function, so that $dg'/g' = dg/g + du/u$ and, setting $\rho = \psi - \psi'$,

$$\frac{dg}{g} \wedge \rho = \theta' - \theta + \frac{du}{u} \wedge \psi'$$

The right-hand side is holomorphic on the overlap, so multiplying by g gives $dg \wedge \rho = g\beta$ with β holomorphic. At a point of X the right-hand side vanishes, so $dg \wedge \rho = 0$ there; and $dg \neq 0$ at such a point, so $\rho = dg \wedge \lambda$ for some covector-valued λ. Pulling back to X kills dg, hence kills ρ,

so the pullbacks of ψ and ψ' agree.

Step 2: the glued form is closed. Run the same argument on $d\eta = 0$. Since $d(dg_\alpha/g_\alpha) = 0$, expanding $d\eta$ on $W_\alpha \cap U$ gives

$$0 \;=\; -\frac{dg_\alpha}{g_\alpha} \wedge d\psi_\alpha + d\theta_\alpha$$

so multiplying by g_α gives $dg_\alpha \wedge d\psi_\alpha = g_\alpha\, d\theta_\alpha$ on $W_\alpha \cap U$; both sides are holomorphic on W_α and agree off the hypersurface X, so the identity holds on all of W_α. At a point of X the right-hand side vanishes, and $dg_\alpha \neq 0$ there, so $d\psi_\alpha = dg_\alpha \wedge \lambda$ at that point for some λ; pulling back to X kills dg_α, hence kills $d\psi_\alpha$. Pullback commutes with d, so the pullback of ψ_α is closed on $X \cap W_\alpha$ and ψ is closed on X; in particular $[\psi] \in H^k(X, \mathbb{C})$ is defined. In the only case used below, $k = n = \dim_{\mathbb{C}} X$, this is automatic: $d\psi$ would be a holomorphic $(n+1)$-form on an n-dimensional complex manifold.

Step 3: the tube integral. Shrink $\mathcal{N}$ so that over W_α the circle bundle $\partial\mathcal{N}$ is $\{|g_\alpha| = \varepsilon\}$. Let γ be a k-cycle in X, subdivided so that each piece γ_α lies in a single W_α. The tube over γ_α fibres over it in circles on which g_α traverses $|g_\alpha| = \varepsilon$ once positively, so integrating over the fibres first,

$$\int_{\tau(\gamma_\alpha)} \frac{dg_\alpha}{g_\alpha} \wedge \psi_\alpha = \int_{\gamma_\alpha} \left(\oint_{|g_\alpha|=\varepsilon} \frac{dg_\alpha}{g_\alpha} \right) \psi_\alpha = 2\pi i \int_{\gamma_\alpha} \psi_\alpha$$

the inner integral being $2\pi i$ by the Cauchy integral formula. For the remaining term, θ_α is bounded near X while the $(k+1)$-dimensional volume of the tube over γ_α is $O(\varepsilon)$, the tube being a circle bundle of radius ε over a fixed compact chain; so those contributions tend to 0 as $\varepsilon \to 0$. Summing over α and using Step 1,

$$\int_{\tau(\gamma)} \eta = 2\pi i \int_\gamma \psi + O(\varepsilon)$$

The left-hand side does not depend on ε, because tubes of different radii are homologous in U, so the error term is zero. Dividing by $2\pi i$ and comparing with definition 5.4.6 gives $\mathrm{Res}\eta = [\psi]$, completing the proof.

For $n = 1$ the recipe already produces the classical objects.

Example 5.4.8: Holomorphic Differentials on a Smooth Plane Curve

Let $n = 1$, so $X \subseteq \mathbb{C}\mathrm{P}^2$ is a smooth plane curve of degree d defined by $F \in \mathbb{C}[x_0, x_1, x_2]_d$. By proposition 5.4.5 the forms with a first-order pole along X are $A\Omega/F$ with $\deg A = d - 3$.

Work in the chart $x_0 = 1$ with coordinates $x = x_1$, $y = x_2$, in which (5.5) reads $\Omega = dx \wedge dy$; write $f(x, y) = F(1, x, y)$ and $a(x, y) = A(1, x, y)$. Since $df = f_x\, dx + f_y\, dy$, one has $dx \wedge dy = -\frac{df \wedge dx}{f_y}$ wherever $f_y \neq 0$, so on that open set

$$\frac{A\,\Omega}{F} = -\frac{df}{f} \wedge \frac{a\,dx}{f_y}$$

while on $\{f_x \neq 0\}$ the same computation with x and y exchanged gives $\frac{df}{f} \wedge \frac{a\,dy}{f_x}$. These two open sets, together with the analogous pair in each of the charts $x_1 \neq 0$ and $x_2 \neq 0$, cover a neighbourhood of X, because smoothness of F puts a nonvanishing partial derivative of the

dehomogenization at every point of X (proposition 5.4.2). So proposition 5.4.7 applies and

$$\operatorname{Res}\frac{A\,\Omega}{F} = -\left.\frac{a\,dx}{f_y}\right|_X = \left.\frac{a\,dy}{f_x}\right|_X$$

the two expressions agreeing on the overlap because $f_x\,dx + f_y\,dy = 0$ along X. This is the classical holomorphic differential on a plane curve. For the Fermat cubic $f = 1 + x^3 + y^3$ and $A = 1$ it is $-dx/(3y^2)$ where $y \neq 0$ and $dy/(3x^2)$ where $x \neq 0$; the curve has genus 1 by example 4.3.2, and $\dim \mathbb{C}[x]_{d-3} = \dim \mathbb{C}[x]_0 = 1$ counts its holomorphic differentials correctly.

What makes the residue useful is not the local formula but the global statement that it loses nothing: it identifies the cohomology of the complement with exactly the part of the cohomology of X that the family can move.

Theorem 5.4.9: The Residue Is an Isomorphism onto Primitive Cohomology

Let $X \subseteq \mathbb{C}\mathrm{P}^{n+1}$ be a smooth hypersurface of degree d with complement U, and let $n \geq 1$. Then

$$\operatorname{Res}\colon H^{n+1}(U,\mathbb{C}) \xrightarrow{\sim} H^n_{\mathrm{prim}}(X,\mathbb{C})$$

is an isomorphism.

***Proof*:**

Write $Y = \mathbb{C}\mathrm{P}^{n+1}$ and h for its hyperplane class, $h_X = h|_X$. Coefficients are in $\mathbb{C}$ throughout, and all homology and cohomology of the spaces in sight is finite-dimensional, so cohomology is the dual of homology.

Step 1: the pair (Y,U) and the tube map. Excising the complement of the tubular neighbourhood and retracting $U \cap \mathcal{N}$ onto $\partial\mathcal{N}$ gives $H_{k+2}(Y,U) \cong H_{k+2}(\mathcal{N},\partial\mathcal{N})$. Lefschetz duality on the compact oriented $(2n+2)$-manifold with boundary $\mathcal{N}$ ([30, Poincaré-Lefschetz duality]) gives $H_{k+2}(\mathcal{N},\partial\mathcal{N}) \cong H^{2n-k}(\mathcal{N}) \cong H^{2n-k}(X)$, the last step because $\mathcal{N}$ retracts onto X; and Poincaré duality on the closed oriented $2n$-manifold X gives $H^{2n-k}(X) \cong H_k(X)$. The composite is the isomorphism $H_k(X) \cong H_{k+2}(Y,U)$ used in definition 5.4.6, and it sends the class of a cycle γ to the class of the disc bundle over γ, whose boundary is the circle bundle over γ. So $\tau = \partial \circ$ (this isomorphism), and the homology sequence of the pair reads

$$H_{n+2}(Y) \xrightarrow{\iota} H_n(X) \xrightarrow{\tau} H_{n+1}(U) \to H_{n+1}(Y) \xrightarrow{\iota} H_{n-1}(X) \tag{5.7}$$

where ι denotes the composite of the map to relative homology with the isomorphism of Step 1.

Step 2: ι intersects with X. Let $Z \subseteq Y$ be a cycle carried by a closed submanifold transverse to X. Choosing $\mathcal{N}$ small enough, $Z \cap \mathcal{N}$ is a disc bundle over $Z \cap X$, and Z differs from $Z \cap \mathcal{N}$ by a chain in U; so the class of Z in $H_\bullet(Y,U)$ is the image of $[Z \cap X]$ under the isomorphism of Step 1. That is, $\iota[Z] = [Z \cap X]$.

Step 3: the two ends of (5.7). Both $H_{n+2}(Y)$ and $H_{n+1}(Y)$ are generated by linear subspaces [30]: one of them vanishes and the other is a line, according to the parity of n. Suppose first that n is even. Then $H_{n+1}(Y) = 0$, so τ is surjective, and $H_{n+2}(Y)$ is the line spanned by $[\Lambda]$ for $\Lambda \cong \mathbb{C}\mathrm{P}^{(n+2)/2}$ a linear subspace, which by the parametric transversality theorem [32, Parametric Transversality] may be moved by an element of PGL_{n+2} so as to meet X transversally. By Step 2, $\iota[\Lambda] = [\Lambda \cap X]$, whose Poincaré dual in $H^n(X)$ is $h_X^{n/2}$, the restriction of the class $h^{n/2}$ of Λ.

Since $\int_X h_X^n = \deg X = d \neq 0$, the class $h_X^{n/2}$ is nonzero; and $\ker\tau = \mathrm{im}\iota$ by exactness of (5.7), so $\ker\tau$ is the line spanned by the Poincaré dual of $h_X^{n/2}$.

Now suppose n is odd. Then $H_{n+2}(Y) = 0$, so τ is injective. For surjectivity, $H_{n+1}(Y)$ is the line spanned by $[\Lambda]$ with $\Lambda \cong \mathbb{C}\mathrm{P}^{(n+1)/2}$, again taken transverse to X, and $\iota[\Lambda] = [\Lambda \cap X]$ has Poincaré dual $h_X^{(n+1)/2}$ in $H^{n+1}(X)$. This is nonzero because $h_X^{(n+1)/2} \smile h_X^{(n-1)/2} = h_X^n$ integrates to d. So the last map in (5.7) is injective, and exactness makes τ surjective.

Step 4: dualizing. In both cases τ is surjective, so the dual map Res is injective, and the image of Res is the annihilator of $\ker\tau$ inside $H^n(X) = H_n(X)^\vee$. For n odd $\ker\tau = 0$ and $H^n(X) = H^n_{\mathrm{prim}}(X)$: indeed corollary 4.2.4 writes $H^n(X) = \bigoplus_{r\geq 0} L^r H^{n-2r}_{\mathrm{prim}}(X)$, and for $r \geq 1$ the group $H^{n-2r}(X)$ vanishes by example 4.3.2, the degree being odd and below n. For n even the same decomposition together with example 4.3.2 gives $H^{n-2r}(X) = \mathbb{C}\, h_X^{(n-2r)/2}$ for $r \geq 1$, and h_X^j is not primitive for $j \geq 1$ because $L^{n-2j+1}h_X^j = h_X^{n-j+1}$ pairs to d against h_X^{j-1}; so only the terms $r = 0$ and $r = n/2$ survive and

$$H^n(X,\mathbb{C}) = H^n_{\mathrm{prim}}(X,\mathbb{C}) \oplus \mathbb{C}\, h_X^{n/2}$$

The annihilator of $\ker\tau$ is $\{\alpha : \int_X \alpha \smile h_X^{n/2} = 0\}$. A primitive α satisfies $h_X \smile \alpha = 0$ and so lies in it; conversely if $\alpha = \alpha_{\mathrm{prim}} + c\, h_X^{n/2}$ lies in it then $0 = c\int_X h_X^n = cd$, so $c = 0$. Hence the annihilator is exactly $H^n_{\mathrm{prim}}(X,\mathbb{C})$, completing the proof.

The theorem converts a question about X into a question about U, where proposition 5.4.5 gives explicit representatives. What is not yet visible is the Hodge filtration, and the mechanism that produces it is the order of the pole: the deeper the pole, the further down the filtration the residue sits. For that to be well defined, one needs to know when a pole can be reduced, and the answer is exactly the Jacobian ideal.

Lemma 5.4.10: Reduction of Pole Order by the Jacobian Ideal

Let $X = X_F$ be a smooth hypersurface of degree d, let $m \geq 2$, and let $A_0, \dots, A_{n+1} \in \mathbb{C}[x]_a$ with $a = (m-1)d - n - 1$. Write $v = \sum_i A_i\, \partial_i$. Then on U

$$\frac{\left(\sum_i A_i\, \partial_i F\right)\Omega}{F^m} = \frac{1}{m-1}\frac{\left(\sum_i \partial_i A_i\right)\Omega}{F^{m-1}} - \frac{1}{m-1}\, d\left(\frac{\iota_v\Omega}{F^{m-1}}\right)$$

where $\iota_v\Omega/F^{m-1}$ is a holomorphic n-form on U. In particular, if $A \in J(F)_{md-n-2}$ then the class of $A\Omega/F^m$ in $H^{n+1}(U,\mathbb{C})$ is represented by a form with a pole of order at most $m-1$.

***Proof*:**

All computations take place on $\mathbb{C}^{n+2} \setminus \{F = 0\}$, and the result is transported to U at the end.

Step 1: two Cartan identities. Write $\mu = dx_0 \wedge \cdots \wedge dx_{n+1}$ and $E = \sum_i x_i\partial_i$, so that $\Omega = \iota_E\mu$. For any polynomial vector field w one has $\mathcal{L}_w\mu = (\mathrm{div}w)\,\mu$ with $\mathrm{div}w = \sum_i \partial_i w_i$, and Cartan's formula $\mathcal{L}_w = d\iota_w + \iota_w d$ [32, Cartan's Magic Formula] gives $d(\iota_w\mu) = (\mathrm{div}w)\mu$ because $d\mu = 0$. Applying $\mathcal{L}_v$ to $\Omega = \iota_E\mu$ and using $\mathcal{L}_v\iota_E = \iota_{[v,E]} + \iota_E\mathcal{L}_v$,

$$d(\iota_v\Omega) = \mathcal{L}_v(\iota_E\mu) - \iota_v\, d(\iota_E\mu) = \iota_{[v,E]}\mu + (\mathrm{div}v)\,\Omega - (n+2)\,\iota_v\mu$$

since $\operatorname{div} E = n+2$. The coefficients of v are homogeneous of degree a, so $[E,v] = (a-1)v$ and $\iota_{[v,E]}\mu = (1-a)\iota_v\mu$, whence

$$d(\iota_v\Omega) = (1-a-n-2)\,\iota_v\mu + (\operatorname{div} v)\,\Omega \tag{5.8}$$

Next, $dF\wedge\mu = 0$ for degree reasons, so contracting with E gives $dF\wedge\Omega = (EF)\,\mu = d\cdot F\,\mu$ by the Euler relation; contracting that with v gives

$$dF\wedge\iota_v\Omega = v(F)\,\Omega - d\cdot F\,\iota_v\mu, \qquad v(F) = \sum_i A_i\,\partial_i F \tag{5.9}$$

Step 2: the identity. Differentiating the quotient and inserting (5.8) and (5.9),

$$\begin{aligned} d\!\left(\frac{\iota_v\Omega}{F^{m-1}}\right) &= \frac{d(\iota_v\Omega)}{F^{m-1}} - (m-1)\frac{dF\wedge\iota_v\Omega}{F^m} \\ &= \frac{\big(1-a-n-2+(m-1)d\big)\,\iota_v\mu}{F^{m-1}} + \frac{(\operatorname{div} v)\Omega}{F^{m-1}} - (m-1)\frac{v(F)\,\Omega}{F^m} \end{aligned}$$

The bracketed scalar is $1-\big((m-1)d-n-1\big)-n-2+(m-1)d = 0$, so the terms involving $\iota_v\mu$ cancel and rearranging gives the asserted identity.

Step 3: descent to U. Let $\pi\colon \mathbb{C}^{n+2}\setminus\{0\}\to\mathbb{CP}^{n+1}$ be the projection. A holomorphic form ζ is a pullback along π if and only if $\iota_E\zeta = 0$ and $\mathcal{L}_E\zeta = 0$: the fibres of π are the orbits of the $\mathbb{C}^*$-action generated by E, and the two conditions say that ζ annihilates the fibre directions and is invariant along them. Now $\iota_E\iota_v\Omega = -\iota_v\iota_E\Omega = 0$ because $\iota_E\Omega = \iota_E\iota_E\mu = 0$; and $\mathcal{L}_E\Omega = (n+2)\Omega$ together with $[E,v] = (a-1)v$ gives $\mathcal{L}_E(\iota_v\Omega) = (a-1+n+2)\,\iota_v\Omega$, so

$$\mathcal{L}_E\!\left(\frac{\iota_v\Omega}{F^{m-1}}\right) = \big(a-1+n+2-(m-1)d\big)\frac{\iota_v\Omega}{F^{m-1}} = 0$$

by the choice of a. So $\iota_v\Omega/F^{m-1}$ descends to a holomorphic n-form on U, both other terms of the identity descend by proposition 5.4.5, and the identity holds on U.

Finally, an element $A\in J(F)_{md-n-2}$ is of the form $\sum_i A_i\partial_i F$ with A_i homogeneous of degree $md-n-2-(d-1) = a$, so the identity writes $A\Omega/F^m$ as a form of pole order $m-1$ plus an exact form, completing the proof.

The lemma already delivers half of the description. For each m let

$$P_m \;=\; \left\{\left[\frac{A\,\Omega}{F^m}\right] \;:\; A\in\mathbb{C}[x]_{md-n-2}\right\} \;\subseteq\; H^{n+1}(U,\mathbb{C})$$

be the set of classes represented by a form of pole order at most m, with the convention $P_0 = 0$; these are subspaces, and $P_1\subseteq P_2\subseteq\cdots$ because $A\Omega/F^m = (FA)\Omega/F^{m+1}$. Lemma 5.4.10 says that $A\mapsto[A\Omega/F^m]$ carries $J(F)_{md-n-2}$ into P_{m-1} for $m\geq 2$, and the same holds vacuously at $m=1$, where $J(F)_{d-n-2} = 0$ because $J(F)$ starts in degree $d-1$. So the assignment induces a surjection

$$R_{md-n-2} \;\twoheadrightarrow\; P_m/P_{m-1} \tag{5.10}$$

Griffiths' theorem is the assertion that (5.10) is an isomorphism and that the filtration $P_\bullet$ is the Hodge filtration read through the residue. Two inputs from algebraic geometry make it work, and both are imported rather than rebuilt: Grothendieck's theorem that a smooth affine variety's cohomology

is computed by global algebraic differential forms [23], applied to the complement $U = \mathbb{CP}^{n+1} \setminus X$, which is affine; and the vanishing of $H^q(\mathbb{CP}^{n+1}, \Omega^p(k))$ in the middle degrees [4, Bott vanishing], which is what collapses the pole-order filtration onto the graded pieces of the Jacobian ring. The assembly below is where both are used, and the full argument in this form is in *Period Mappings and Period Domains* [1, the cohomology of hypersurfaces].

Theorem 5.4.11: Griffiths' Description of the Hodge Filtration

Let $X = X_F \subseteq \mathbb{CP}^{n+1}$ be a smooth hypersurface of degree $d \geq 2$, with $n \geq 1$, and let $R = R(F)$ be its Jacobian ring. Then

1. $\bigcup_{m \geq 1} P_m = H^{n+1}(U, \mathbb{C})$;
2. $\mathrm{Res}(P_m) = F^{n+1-m} H^n_{\mathrm{prim}}(X, \mathbb{C})$ for every $m \geq 1$, and in particular $P_{n+1} = H^{n+1}(U, \mathbb{C})$;
3. for every $m \geq 1$ the map $A \mapsto \mathrm{Res}[A\Omega/F^m]$ induces an isomorphism of vector spaces

$$R_{md-n-2} \xrightarrow{\sim} \mathrm{gr}_F^{n+1-m} H^n_{\mathrm{prim}}(X, \mathbb{C}) = H^{n+1-m,\, m-1}_{\mathrm{prim}}(X)$$

Equivalently, writing $m = q + 1$, for every q with $0 \leq q \leq n$,

$$H^{n-q,\, q}_{\mathrm{prim}}(X) \cong R_{(q+1)d-n-2} \tag{5.11}$$

***Proof Key ideas*:**
What is proved here in full is the surjection (5.10), which is lemma 5.4.10: the assignment $A \mapsto [A\Omega/F^m]$ is linear, its restriction to $J(F)_{md-n-2}$ lands in P_{m-1} by that lemma, and it is surjective onto P_m by the definition of P_m together with proposition 5.4.5. Since Res is injective (theorem 5.4.9), statement (3) is equivalent to the assertion that (5.10) is injective and that Res carries P_m/P_{m-1} onto gr_F^{n+1-m}, which is what (2) provides.

Two inputs are taken from outside. The first is *Grothendieck's algebraic de Rham theorem*: for a smooth affine variety V over $\mathbb{C}$, the complex of global algebraic differential forms on V computes $H^\bullet(V, \mathbb{C})$. Applied to the affine variety U it gives statement (1), since a global algebraic $(n+1)$-form on U extends meromorphically across X with a pole of some finite order and is therefore some $A\Omega/F^m$ by proposition 5.4.5. Granting (2), its case $m = n + 1$ reads $\mathrm{Res}(P_{n+1}) = F^0 H^n_{\mathrm{prim}} = H^n_{\mathrm{prim}}$, and injectivity of Res then forces $P_{n+1} = H^{n+1}(U, \mathbb{C})$.

The second input is the identification of the pole filtration with the Hodge filtration, statement (2), together with the injectivity of (5.10). The proof compares the filtration of the algebraic de Rham complex of U by order of pole with its filtration by degree of form; the two induce the same filtration on cohomology, and the comparison rests on the vanishing $H^q(\mathbb{CP}^{n+1}, \Omega^p_{\mathbb{CP}^{n+1}}(k)) = 0$ for $q \geq 1$ and $k > 0$. That vanishing, and the spectral-sequence bookkeeping that uses it, are not developed in this book; the argument is given in full in [1, the cohomology of hypersurfaces].

The final equivalence is bookkeeping. The Hodge structure on $H^n_{\mathrm{prim}}(X)$ has weight n, so $\mathrm{gr}_F^{n+1-m} = H^{n+1-m,\, n-(n+1-m)} = H^{n+1-m,\, m-1}$, and substituting $m = q + 1$ turns this into (5.11), as we sought to show.

The formula (5.11) is a complete answer to a question that on the face of it requires solving an elliptic equation on X. The Hodge decomposition of chapter 2 produces $H^{p,q}$ as a space of harmonic forms,

defined by a metric and computed by analysis; here the same space appears as a graded piece of a quotient of a polynomial ring, defined by the equation of X and computed by counting monomials. Two features of the answer are worth naming. Hodge symmetry $h^{p,q} = h^{q,p}$ becomes the symmetry of the graded dimensions of R about the middle degree, because $\sigma - \big((q+1)d-n-2\big) = (n-q+1)d-n-2$ when $\sigma = (n+2)(d-2)$. And the Hodge numbers of X depend on F only through $R(F)$, so two hypersurfaces with isomorphic graded Jacobian rings have the same Hodge numbers. The rational structure of $H^n(X,\mathbb{Q})$ is invisible in $R(F)$, so this says nothing about the Hodge structures themselves being isomorphic.

Combining theorem 5.4.11 with the local constancy given by theorem 5.2.6 turns the single Fermat computation of example 5.4.3 into a formula for every smooth hypersurface.

Corollary 5.4.12: The Hodge Numbers of a Smooth Hypersurface

Let $X \subseteq \mathbb{C}\mathrm{P}^{n+1}$ be any smooth hypersurface of degree $d \geq 2$, with $n \geq 1$. Then for $0 \leq q \leq n$,

$$h^{n-q,\,q}_{\mathrm{prim}}(X) \;=\; \#\Big\{(a_0,\dots,a_{n+1}) \;:\; 0 \leq a_i \leq d-2,\ \sum_i a_i = (q+1)d-n-2\Big\}$$

that is, the coefficient of $t^{(q+1)d-n-2}$ in $(1+t+\cdots+t^{d-2})^{n+2}$. In particular these numbers depend only on n, d and q.

***Proof*:**
Let S be the parameter space of proposition 5.4.4 and $F^\bullet \subseteq \mathcal{H}^n_{\mathrm{prim}}$ the Hodge filtration of the associated variation. Part (3) of proposition 5.3.2 makes each F^p a holomorphic subbundle of $\mathcal{H}^n_{\mathrm{prim}}$ (note that theorem 5.2.6 is stated for the full bundle $\mathcal{H}^n$ and does not by itself cover the primitive filtration), and part (2) of the same proposition already gives that $h^{p,q}_{\mathrm{prim}}$ is independent of s. It therefore suffices to evaluate the numbers at one point of S, and the Fermat point is one. There theorem 5.4.11 gives $h^{n-q,q}_{\mathrm{prim}} = \dim R_{(q+1)d-n-2}$, which example 5.4.3 computes as the stated monomial count, completing the proof.

Example 5.4.13: The Quintic Threefold

Take $n = 3$ and $d = 5$: a smooth quintic hypersurface $X \subseteq \mathbb{C}\mathrm{P}^4$. Here n is odd, so $H^3(X,\mathbb{C}) = H^3_{\mathrm{prim}}(X,\mathbb{C})$ by the computation in Step 4 of the proof of theorem 5.4.9, and the relevant degrees are $(q+1)\cdot 5 - 5 = 5q$ for $q = 0,1,2,3$. By corollary 5.4.12 each Hodge number is a count of $(a_0,\dots,a_4)$ with $0 \leq a_i \leq 3$ and prescribed sum.

Degree 0. Only $a = 0$, so $h^{3,0} = 1$: the quintic threefold carries a one-dimensional space of holomorphic 3-forms, and by proposition 5.4.5 a generator is $\mathrm{Res}(\Omega/F)$.

Degree 5. Without the upper bound there are $\binom{5+4}{4} = 126$ solutions of $\sum a_i = 5$ in nonnegative integers. Those violating the bound have $a_i \geq 4$ for exactly one i, since two such entries would already force $\sum a_i \geq 8$; subtracting 4 from that entry leaves $\sum a_i = 1$, with $\binom{1+4}{4} = 5$ solutions, and there are 5 choices of i. Hence

$$h^{2,1} = 126 - 25 = 101$$

Degrees 10 *and* 15. The socle degree is $\sigma = (n+2)(d-2) = 15$, and the symmetry $a \mapsto (3,3,3,3,3) - a$ of example 5.4.3 gives $h^{1,2} = h^{2,1} = 101$ and $h^{0,3} = h^{3,0} = 1$, as Hodge symmetry

requires.

So the Hodge numbers of the middle cohomology are $(1, 101, 101, 1)$ and $b_3(X) = 204$. The period map of definition 5.3.3 for the family of quintics therefore takes values in the period domain of section 3.3 for polarized weight-3 Hodge structures with these Hodge numbers, and the number 101 reappears as the dimension of the space of quintics modulo projective changes of coordinates (exercise 5.4.1 (5)).

One piece of the chapter has not yet been made explicit for this family: the derivative. Griffiths transversality says that ∇ moves F^p into F^{p-1}, and corollary 5.2.8 extracts from it an $\mathcal{O}_S$-linear map on the graded pieces, which is the differential of the period map. Under (5.11) both source and target are graded pieces of R, and the map between them is multiplication in that ring.

Theorem 5.4.14: The Differential of the Period Map as Multiplication

Let $[F] \in S$ and let $G \in \mathbb{C}[x]_d$ represent a tangent vector to $\mathbb{C}\mathrm{P}^M$ at $[F]$. Under the isomorphisms (5.11), the map

$$\bar{\nabla}_G \colon H^{n-q,\,q}_{\mathrm{prim}}(X) \to H^{n-q-1,\,q+1}_{\mathrm{prim}}(X)$$

obtained from corollary 5.2.8 by contracting with G is

$$[A] \;\longmapsto\; -(q+1)\,[G\cdot A]$$

multiplication by G in the graded ring R, up to the nonzero scalar $-(q+1)$. In particular $\bar{\nabla}_G$ depends only on the class of G in R_d, and vanishes when $G \in J(F)_d$.

***Proof*:**

Step 1: the residue is flat. Let $\mathcal{U} \to S$ be the family of complements, $\mathcal{U} = (\mathbb{C}\mathrm{P}^{n+1} \times S) \setminus \mathcal{X}$. By Step 3 of proposition 5.4.4 the pair $(\mathbb{C}\mathrm{P}^{n+1}, X_F)$ is locally trivial over S, so both $\mathcal{X} \to S$ and $\mathcal{U} \to S$ are locally trivial fibre bundles and the tube map of definition 5.4.6 is defined fibrewise by a construction that a trivialization of the pair carries along. So $t \mapsto \tau_t$ is a morphism of local systems from $t \mapsto H_n(X_t, \mathbb{C})$ to $t \mapsto H_{n+1}(U_t, \mathbb{C})$, and dually Res is a morphism of local systems from $t \mapsto H^{n+1}(U_t, \mathbb{C})$ to $\mathbb{H}^n \otimes \mathbb{C}$, an isomorphism onto $\mathbb{H}^n_{\mathrm{prim}} \otimes \mathbb{C}$ by theorem 5.4.9. In particular Res commutes with parallel transport, and it carries flat families of tubes to flat families. By theorem 5.1.7 the flat sections of ∇ are the sections of the local system, so in a flat frame ∇ is differentiation of coefficients, and for any flat family of cycles γ_t and any section σ of $\mathcal{H}^n_{\mathrm{prim}}$,

$$\frac{d}{dt}\big\langle \sigma(t), \gamma_t \big\rangle = \big\langle \nabla_{\partial_t}\sigma, \gamma_t \big\rangle \tag{5.12}$$

Step 2: differentiating a rational representative. Let $F_t = F + tG$, which lies in S for small t, and fix $A \in \mathbb{C}[x]_{(q+1)d-n-2}$. Put

$$\eta_t = \frac{A\,\Omega}{F_t^{\,q+1}}, \qquad \sigma(t) = \mathrm{Res}[\eta_t]$$

a holomorphic section of $\mathcal{H}^n_{\mathrm{prim}}$ along the curve, whose value at $t = 0$ is the class corresponding to $[A] \in R_{(q+1)d-n-2}$. Choose a flat family of cycles $\gamma_t \in H_n(X_t, \mathbb{C})$, so that $\tau(\gamma_t)$ is a flat family in

$H_{n+1}(U_t,\mathbb{C})$ by Step 1. Trivializing the family of complements over a disc by a diffeomorphism φ_t, the tube may be taken to be a fixed cycle c, and

$$\frac{d}{dt}\int_{\tau(\gamma_t)}\eta_t=\frac{d}{dt}\int_c\varphi_t^*\eta_t=\int_c\varphi_t^*\Big(\frac{\partial\eta_t}{\partial t}\Big)+\int_c\varphi_t^*\big(\mathcal{L}_w\eta_t\big)$$

where w is the vector field generating φ_t. Each η_t is closed, so $\mathcal{L}_w\eta_t=d(\iota_w\eta_t)$ is exact and the second integral vanishes. Hence differentiation passes under the integral sign, and

$$\left.\frac{\partial\eta_t}{\partial t}\right|_{t=0}=-(q+1)\frac{A\,G\,\Omega}{F^{\,q+2}}$$

by the chain rule applied to $F_t^{-(q+1)}$.

Step 3: conclusion. Combining Step 2 with (5.12) and the definition of Res,

$$\big\langle\nabla_{\partial_t}\sigma,\gamma_0\big\rangle=\frac{1}{2\pi i}\int_{\tau(\gamma_0)}-(q+1)\frac{A\,G\,\Omega}{F^{\,q+2}}=\Big\langle\mathrm{Res}\Big[-(q+1)\frac{(GA)\,\Omega}{F^{\,q+2}}\Big],\gamma_0\Big\rangle$$

for every γ_0, so $\nabla_{\partial_t}\sigma=-(q+1)\mathrm{Res}[(GA)\Omega/F^{q+2}]$. The right-hand side lies in $\mathrm{Res}(P_{q+2})=F^{n-q-1}H^n_{\mathrm{prim}}$ by theorem 5.4.11, which is Griffiths transversality made explicit, and its class modulo F^{n-q} is $-(q+1)$ times the class of GA in $R_{(q+2)d-n-2}$, again by theorem 5.4.11. Since $\bar\nabla_G$ is exactly the induced map on these graded pieces, the formula follows. If $G\in J(F)_d$ then $GA\in J(F)$, so $[GA]=0$ and $\bar\nabla_G=0$, completing the proof.

The vanishing in the last sentence is forced. An element of $J(F)_d$ is $\sum_i\ell_i\,\partial_iF$ with ℓ_i linear, which is the derivative of F along the linear vector field $\sum_i\ell_i\partial_i$: it is a tangent direction along the orbit of F under PGL_{n+2}, and moving F in such a direction changes the coordinates rather than the hypersurface. So $d\Phi$ annihilates exactly the directions in which nothing happens, and what the period map for hypersurfaces records is the multiplication

$$R_d\otimes R_{(q+1)d-n-2}\to R_{(q+2)d-n-2}$$

of the Jacobian ring. This is also the sharpest form of transversality available here: multiplication by an element of R_d raises the degree by d, which is one step in the numbering (5.11), and never two. Whether $d\Phi$ is injective, a question about that multiplication, is the infinitesimal Torelli problem; exercise 5.4.1 (6) settles it for the quintic threefold, and the global question is taken up for K3 surfaces in chapter 6.

Exercise 5.4.1

1. Let $X\subseteq\mathbb{C}\mathrm{P}^3$ be a smooth cubic surface, so $n=2$ and $d=3$. Compute the graded dimensions of the Jacobian ring of the Fermat cubic and use corollary 5.4.12 to determine $h^{2,0}(X)$, $h^{1,1}_{\mathrm{prim}}(X)$ and $h^{0,2}(X)$. Deduce $b_2(X)$, and say what the vanishing of $h^{2,0}$ means about holomorphic 2-forms on X.

2. Repeat the computation for a smooth quartic surface in $\mathbb{C}\mathrm{P}^3$ ($n=2$, $d=4$) and check the result against the Hodge numbers of a K3 surface recorded in example 2.2.4. Which degree of the Jacobian ring carries the holomorphic 2-form, and which carries the primitive $(1,1)$-classes?

3. Show that $h^{n,0}(X)=\binom{d-1}{n+1}$ for every smooth hypersurface $X\subseteq\mathbb{C}\mathrm{P}^{n+1}$ of degree $d\geq 2$,

without computing any Jacobian ring beyond its lowest degrees. Deduce that X carries no nonzero holomorphic n-form exactly when $d \leq n+1$.

4. Let $X \subseteq \mathbb{C}\mathrm{P}^2$ be a smooth plane curve of degree d. Using exercise 5.4.1 (3), recover $g = (d-1)(d-2)/2$, and write down a basis of $H^0(X, \Omega^1_X)$ in the affine chart of example 5.4.8. Explain why example 5.2.9 makes Griffiths transversality vacuous here, and say which statement of this section still has content in weight one.

5. For the Fermat quintic threefold $F = x_0^5 + \cdots + x_4^5$, compute $\dim_{\mathbb{C}} J(F)_5$ directly from the monomial description of $J(F)$, and deduce $\dim R_5$ a second way, independently of the count in example 5.4.13. Compare $\dim R_5$ with $\dim \mathbb{C}\mathrm{P}^M - \dim \mathrm{PGL}_5$, and explain the coincidence using theorem 5.4.14.

6. Continuing with the quintic threefold, show that the composite

$$R_5 \to \mathrm{Hom}\big(H^{3,0}_{\mathrm{prim}}(X),\ H^{2,1}_{\mathrm{prim}}(X)\big)$$

of theorem 5.4.14 is injective at the Fermat point, and conclude that the period map of the family of quintics has injective differential there, modulo the directions along the PGL_5-orbit.

6

Period Domains and Mumford-Tate Groups

Chapter 5 put a family into a period domain and showed the resulting map is holomorphic and horizontal. Neither adjective has yet been cashed. Horizontality was a constraint on a derivative, and a constraint is only useful once something is known about the space it constrains motion in. That is the first business of this chapter.

Period domains are homogeneous, so their geometry is determined at a single point, and section 3.3 already recorded that. What it did not record is the curvature, and the curvature is peculiar. A period domain is negatively curved in the horizontal directions and not, in general, anywhere else: it is almost never a Hermitian symmetric space, and the familiar picture of a bounded domain with a complete negatively curved metric is available only in weight one, where D is the Siegel upper half space. What survives in general is exactly enough. Horizontal maps are distance-decreasing, and that alone forces rigidity.

The second business is a group. A single Hodge structure carries more information than its Hodge numbers, and the invariant that captures it is the Mumford-Tate group: the smallest algebraic group over $\mathbb{Q}$ whose real points contain the circle action defining the structure. Its representation theory answers a question the earlier chapters could only pose. The Hodge classes in every tensor construction on V are precisely the vectors it fixes, so the search for Hodge classes becomes the search for invariants of a reductive group.

Those two threads meet in the locus where a family acquires an unexpected Hodge class. Griffiths transversality makes that locus analytic, and a theorem of Cattani, Deligne and Kaplan makes it algebraic. The chapter closes with the case where every object in it can be written down: polarized K3 surfaces, whose period domain is again Hermitian symmetric, whose lattice is a fixed unimodular form of signature $(3, 19)$, and for which the period map is injective.

6.1 Curvature and Rigidity of Period Domains

Proposition 3.3.5 presents D as $G_{\mathbb{R}}/(G_{\mathbb{R}} \cap P)$ with compact isotropy, and definition 5.3.8 singles out the subbundle $T^h D \subseteq TD$ that every period map is tangent to. Neither statement says anything metric. This section gives the metric: the polarization produces a $G_{\mathbb{R}}$-invariant Hermitian metric on D, and the curvature of that metric is negative along $T^h D$ and not, in general, anywhere else.

A map into a negatively curved target is constrained by the Ahlfors-Schwarz lemma; a horizontal map into D therefore inherits those constraints even though the negativity is unavailable in the remaining directions. The consequences are rigidity statements about families, and the first of them says that a smooth projective family over $\mathbb{C}\mathrm{P}^n$ cannot move at all.

Throughout, $(V_{\mathbb{Z}}, Q, k, \{h^{p,q}\})$ is the numerical data fixed in section 3.3, and $G_{\mathbb{C}}$ and $G_{\mathbb{R}}$ are the automorphism groups of Q from proposition 3.3.5. For a fixed $F \in D$ the letter P denotes the stabilizer of F in $G_{\mathbb{C}}$ and $K = G_{\mathbb{R}} \cap P$ the (compact) isotropy group. Write

$$\mathfrak{g}_{\mathbb{C}} = \{X \in \mathrm{End}(V_{\mathbb{C}}) \,\big|\, Q(Xu, v) + Q(u, Xv) = 0 \text{ for all } u, v \in V_{\mathbb{C}}\}$$

for the Lie algebra of $G_{\mathbb{C}}$, and $\mathfrak{g}_{\mathbb{R}}$, $\mathfrak{g}_{\mathbb{Q}}$ for the analogous spaces of real and rational endomorphisms, so that $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$. Finally C_F denotes the Weil operator of the Hodge structure at F, acting on $V_F^{p,q}$ as i^{p-q} (definition 3.2.1).

Everything below is computed inside $\mathfrak{g}$. The reason is that the Hodge structure carried by V propagates to a Hodge structure on $\mathfrak{g}$, and once it does, the tangent bundle of D, its horizontal subbundle and its invariant metrics all become statements about a single graded Lie algebra.

Proposition 6.1.1: The Lie Algebra of a Point of the Period Domain

Fix $F \in D$ with bigrading $V^{p,q} = F^p \cap \overline{F^q}$, and for $r \in \mathbb{Z}$ set

$$\mathfrak{g}_F^{r,-r} = \left\{ X \in \mathfrak{g}_{\mathbb{C}} \;\middle|\; X\, V^{p,q} \subseteq V^{p+r,\, q-r} \text{ for all } p, q \right\}$$

Then:

1. $\mathfrak{g}_{\mathbb{C}} = \bigoplus_{r \in \mathbb{Z}} \mathfrak{g}_F^{r,-r}$, with $\overline{\mathfrak{g}_F^{r,-r}} = \mathfrak{g}_F^{-r,r}$ and $[\mathfrak{g}_F^{r,-r}, \mathfrak{g}_F^{s,-s}] \subseteq \mathfrak{g}_F^{r+s,-r-s}$. In particular this is a pure Hodge structure of weight 0 on $\mathfrak{g}_{\mathbb{Q}}$ in the sense of definition 3.1.1, and the bracket is a morphism of Hodge structures.
2. $\mathrm{Lie}(P) = \mathfrak{p}_F := \bigoplus_{r \geq 0} \mathfrak{g}_F^{r,-r}$ and $\mathrm{Lie}(K) = \mathfrak{g}_{\mathbb{R}} \cap \mathfrak{g}_F^{0,0}$.
3. The differential at the identity of the orbit map $g \mapsto gF$ induces $\mathbb{C}$-linear isomorphisms

$$T_F^{1,0}\check{D} \cong \mathfrak{g}_{\mathbb{C}}/\mathfrak{p}_F \cong \bigoplus_{r<0} \mathfrak{g}_F^{r,-r}$$

under which the horizontal subspace of definition 5.3.8 is

$$T_F^h \check{D} \cong \mathfrak{g}_F^{-1,1}$$

***Proof*:**
Write $\mathfrak{g}^{r,-r}$ for $\mathfrak{g}_F^{r,-r}$ and $\mathfrak{p}$ for $\mathfrak{p}_F$.

1. Decompose $\mathrm{End}(V_{\mathbb{C}}) = \bigoplus_r E^r$, where $E^r = \bigoplus_{p,q} \mathrm{Hom}(V^{p,q}, V^{p+r,q-r})$; this is the decomposition of an endomorphism into its homogeneous components for the bigrading, and it is a direct sum because the $V^{p,q}$ span $V_{\mathbb{C}}$. Let $X \in \mathfrak{g}_{\mathbb{C}}$ and write $X = \sum_r X_r$ with $X_r \in E^r$. Fix $u \in V^{p,q}$ and $w \in V^{p',q'}$. By condition (1) of definition 3.2.1 the form Q pairs $V^{a,b}$ nontrivially only with $V^{b,a}$, so $Q(X_r u, w) = 0$ unless $(p',q') = (q-r, p+r)$, that is unless $r = q - p'$; and $Q(u, X_s w) = 0$ unless $(p'+s, q'-s) = (q,p)$, again unless $s = q - p'$. Setting $r_0 = q - p'$, the identity $Q(Xu,w) + Q(u,Xw) = 0$ therefore reads

$$Q(X_{r_0}u, w) + Q(u, X_{r_0}w) = 0$$

while for $r \neq r_0$ both terms $Q(X_r u, w)$ and $Q(u, X_r w)$ vanish individually. As u and w range over the bigraded pieces this shows $Q(X_r u, w) + Q(u, X_r w) = 0$ for every r and every pair u, w, so each X_r lies in $\mathfrak{g}_{\mathbb{C}}$. Hence $\mathfrak{g}_{\mathbb{C}} = \bigoplus_r \mathfrak{g}^{r,-r}$.

For conjugation, let $X \in \mathfrak{g}^{r,-r}$ and $u \in V^{p,q}$. Then $\bar{u} \in V^{q,p}$ and $\bar{X}(\bar{u}) = \overline{Xu} \in \overline{V^{p+r,q-r}} = V^{q-r,p+r}$, so $\bar{X}$ carries $V^{q,p}$ into $V^{q-r,p+r}$ and thus lies in $\mathfrak{g}^{-r,r}$. That $\bar{X} \in \mathfrak{g}_{\mathbb{C}}$ is automatic because Q is defined over $\mathbb{Q}$. The bracket statement is immediate: a composite XY with $X \in \mathfrak{g}^{r,-r}$ and $Y \in \mathfrak{g}^{s,-s}$ shifts the bigrading by $r+s$, and so does YX.

Finally $\mathfrak{g}_{\mathbb{Q}}$ is a $\mathbb{Q}$-vector space with $\mathfrak{g}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} = \mathfrak{g}_{\mathbb{C}}$, and the displayed decomposition satisfies exactly the conjugation condition of definition 3.1.1 in weight 0, the summand $\mathfrak{g}^{r,-r}$ playing the role of the $(r,-r)$-piece. The bracket is a morphism of Hodge structures (definition 3.1.7) because it is defined over $\mathbb{Q}$ and respects the bigrading by the previous sentence.

2. An element $X \in \mathfrak{g}_{\mathbb{C}}$ lies in $\mathrm{Lie}(P)$ precisely when $XF^p \subseteq F^p$ for every p, since P is the subgroup of $G_{\mathbb{C}}$ fixing every step of the flag and a one-parameter subgroup preserves a subspace exactly when its generator does. Now $F^p = \bigoplus_{a \geq p} V^{a,k-a}$, so for $X \in \mathfrak{g}^{s,-s}$ one has $XF^p \subseteq F^{p+s}$, and $F^{p+s} \subseteq F^p$ for all p if and only if $s \geq 0$. Since $\mathrm{Lie}(P)$ is a sum of homogeneous components by part (1), this gives $\mathrm{Lie}(P) = \bigoplus_{s \geq 0} \mathfrak{g}^{s,-s} = \mathfrak{p}$.

For the isotropy, $\mathrm{Lie}(K) = \mathfrak{g}_{\mathbb{R}} \cap \mathfrak{p}$. If X is real and lies in $\mathfrak{p}$ then $\bar{X} = X$ lies in $\bar{\mathfrak{p}} = \bigoplus_{s \leq 0} \mathfrak{g}^{s,-s}$ as well, so $X \in \mathfrak{p} \cap \bar{\mathfrak{p}} = \mathfrak{g}^{0,0}$. Conversely $\mathfrak{g}_{\mathbb{R}} \cap \mathfrak{g}^{0,0} \subseteq \mathfrak{g}_{\mathbb{R}} \cap \mathfrak{p}$.

3. By proposition 3.3.5 the orbit map $G_{\mathbb{C}} \to \check{D}$ induces a biholomorphism $G_{\mathbb{C}}/P \to \check{D}$; its differential at the identity coset is the standard isomorphism $\mathfrak{g}_{\mathbb{C}}/\mathfrak{p} \cong T_F^{1,0}\check{D}$ for a complex homogeneous space ([37, The Quotient Manifold Theorem]), $\mathbb{C}$-linear because the orbit map is holomorphic. Composing with the splitting $\mathfrak{g}_{\mathbb{C}} = \mathfrak{p} \oplus \bigoplus_{r<0} \mathfrak{g}^{r,-r}$ of part (1) gives the second isomorphism.

Concretely, the tangent vector attached to $X \in \bigoplus_{r<0} \mathfrak{g}^{r,-r}$ is $\frac{d}{dt}\big|_0 \exp(tX)F$. Its p-th component in the sense of definition 5.3.8 is computed on a local section: for $u \in F^p$ the curve $t \mapsto \exp(tX)u$ is a section of the moving subspace $\exp(tX)F^p$ with derivative Xu at $t = 0$, so

$$\theta_p(u) = Xu \bmod F^p$$

Horizontality asks $\theta_p(F^p) \subseteq F^{p-1}/F^p$ for every p, that is $XF^p \subseteq F^{p-1}$ for every p. Write $X = \sum_{r<0} X_r$. If $X_r = 0$ for all $r \leq -2$ then $X = X_{-1}$ and $XF^p \subseteq F^{p-1}$ for every p, so X is horizontal. Conversely suppose $X_{r_0} \neq 0$ for some $r_0 \leq -2$, and choose a bidegree (p,q) and a vector $u \in V^{p,q}$ with $X_{r_0}u \neq 0$. The vectors $X_r u$ lie in the distinct summands $V^{p+r,q-r}$, and $F^{p-1} = \bigoplus_{a \geq p-1} V^{a,k-a}$ is a sum of summands, so $Xu \in F^{p-1}$ would force $X_r u \in F^{p-1}$ for each r separately, hence $p + r_0 \geq p - 1$ and $r_0 \geq -1$. That contradicts $r_0 \leq -2$, so X is not horizontal. Hence $T_F^h \check{D}$ corresponds to $\mathfrak{g}^{-1,1}$, as we sought to show.

The proposition converts three separate constructions into one grading. The tangent bundle of $\check{D}$ is the negative part of $\mathfrak{g}$, the horizontal subbundle is its lowest graded piece, and the isotropy algebra is the piece of degree zero. The condition $T^h\check{D} = T\check{D}$, which example 5.3.9 verified by hand in weight one and in the K3 numerical type, now reads $\mathfrak{g}^{r,-r} = 0$ for all $r \leq -2$: horizontality is vacuous exactly when the graded Lie algebra has only three nonzero pieces. Each $\mathfrak{g}^{r,-r}$ is also computable from the Hodge numbers alone, since it is a subspace of $\bigoplus_p \operatorname{Hom}(V^{p,q}, V^{p+r,q-r})$ cut out by the skew-symmetry condition against Q.

A metric on D is now a choice of Hermitian inner product on $\bigoplus_{r<0} \mathfrak{g}_F^{r,-r}$ for each F, compatible with the group action. The polarization gives one, through the positive definite form H_F of (3.6) and the Weil operator.

Definition 6.1.2: The Hodge Metric

Let $\beta(X,Y) = \operatorname{tr}(XY)$ be the trace form on $\mathfrak{g}_{\mathbb{C}}$, and for $F \in D$ let $C_F \in \mathrm{GL}(V_{\mathbb{C}})$ be the Weil operator at F. The *Hodge form* at F is

$$\langle X, Y \rangle_F = -\beta\big(X,\ C_F\, \bar{Y}\, C_F^{-1}\big) \qquad (X, Y \in \mathfrak{g}_{\mathbb{C}})$$

The *Hodge metric* on D is the Hermitian metric h whose value at F is the restriction of $\langle -, - \rangle_F$ to

$$T_F^{1,0} D \cong \bigoplus_{r<0} \mathfrak{g}_F^{r,-r}$$

under the identification of proposition 6.1.1(3). The associated Riemannian metric on the real tangent bundle is $\operatorname{Re} h$, a J-invariant metric.[a]

[a] [5, The Fundamental Form is Real and Positive].

The definition has three things to verify: that $\langle -, - \rangle_F$ is a positive definite Hermitian form, that the resulting assignment is $G_{\mathbb{R}}$-invariant, and that it varies smoothly. All three come out of one identity, which also explains the sign.

Proposition 6.1.3: The Hodge Form Is a Positive $G_{\mathbb{R}}$-Invariant Metric

Fix $F \in D$ and write $C = C_F$, $H = H_F$ for the form (3.6). Then:

1. $C \in G_{\mathbb{R}}$, $H(u,v) = Q(Cu, \bar{v})$, and for $Y \in \mathfrak{g}_{\mathbb{C}}$ the H-adjoint of Y is $Y^* = -C\bar{Y}C^{-1}$. Consequently $\langle X, Y\rangle_F = \mathrm{tr}(XY^*)$, and $\langle -, -\rangle_F$ is a positive definite Hermitian form on $\mathfrak{g}_{\mathbb{C}}$.
2. Conjugation by C acts on $\mathfrak{g}_F^{r,-r}$ as multiplication by $(-1)^r$: for $X \in \mathfrak{g}_F^{r,-r}$ one has $CXC^{-1} = (-1)^r X$. The summands $\mathfrak{g}_F^{r,-r}$ are mutually orthogonal for $\langle -, -\rangle_F$, and on $\mathfrak{g}_F^{r,-r}$ one has $\langle X, X\rangle_F = (-1)^{r+1}\mathrm{tr}(X\bar{X})$.
3. For $g \in G_{\mathbb{R}}$ one has $C_{gF} = gC_Fg^{-1}$ and $\langle gXg^{-1}, gYg^{-1}\rangle_{gF} = \langle X, Y\rangle_F$. The Hodge metric is therefore $G_{\mathbb{R}}$-invariant, and it is a smooth Hermitian metric on D.

***Proof*:**

Write $\mathfrak{g}^{r,-r}$ for $\mathfrak{g}_F^{r,-r}$ and $V^{p,q}$ for $V_F^{p,q}$.

1. For $u \in V^{p,q}$ and $v \in V^{p',q'}$, the product $Q(Cu, Cv) = i^{p-q}i^{p'-q'}Q(u,v)$ vanishes unless $(p',q') = (q,p)$, in which case the scalar is $i^{p-q}i^{q-p} = 1$. So $Q(Cu,Cv) = Q(u,v)$ for all u, v, and C preserves $V_{\mathbb{R}}$ by proposition 3.2.2(1); hence $C \in G_{\mathbb{R}}$. Next, expanding $Q(Cu, \bar{v})$ over the bigrading and using condition (1) of definition 3.2.1 kills every cross term, leaving $\sum_{p+q=k} i^{p-q}Q(u^{p,q}, \overline{v^{p,q}})$, which is (3.6).

   For the adjoint, let $Y \in \mathfrak{g}_{\mathbb{C}}$ and put $Y' = CYC^{-1}$, again in $\mathfrak{g}_{\mathbb{C}}$ because $C \in G_{\mathbb{C}}$. Then $CY = Y'C$, so

   $$H(Yu, v) = Q(CYu, \bar{v}) = Q(Y'Cu, \bar{v}) = -Q(Cu, Y'\bar{v})$$

   using that Y' is Q-skew. Since C is real, $\overline{Y'} = C\bar{Y}C^{-1}$, and $Y'\bar{v} = \overline{\overline{Y'}v}$; therefore $-Q(Cu, Y'\bar{v}) = -H\big(u, (C\bar{Y}C^{-1})v\big)$, which identifies $Y^* = -C\bar{Y}C^{-1}$ and gives $\langle X, Y\rangle_F = -\mathrm{tr}(XC\bar{Y}C^{-1}) = \mathrm{tr}(XY^*)$.

   Positivity and Hermitian symmetry are now matrix statements. The form H is positive definite (proposition 3.3.4), so choose an H-orthonormal basis of $V_{\mathbb{C}}$; in it Y^* is the conjugate transpose of the matrix of Y. Hence $\langle X, Y\rangle_F = \sum_{a,b} x_{ab}\overline{y_{ab}}$, which is $\mathbb{C}$-linear in X, conjugate-linear in Y, satisfies $\overline{\langle X, Y\rangle_F} = \langle Y, X\rangle_F$, and has $\langle X, X\rangle_F = \sum_{a,b} |x_{ab}|^2 > 0$ for $X \neq 0$.

2. For $X \in \mathfrak{g}^{r,-r}$ and $u \in V^{p,q}$,

   $$CXC^{-1}u = i^{-(p-q)}\,C(Xu) = i^{q-p}i^{(p+r)-(q-r)}Xu = i^{2r}Xu = (-1)^r Xu$$

   so conjugation by C acts on $\mathfrak{g}^{r,-r}$ as $(-1)^r$. Thus for $Y \in \mathfrak{g}^{s,-s}$ one has $\bar{Y} \in \mathfrak{g}^{-s,s}$ and $Y^* = -(-1)^{-s}\bar{Y} = (-1)^{s+1}\bar{Y}$, whence $\langle X, Y\rangle_F = (-1)^{s+1}\mathrm{tr}(X\bar{Y})$. For $X \in \mathfrak{g}^{r,-r}$ the composite $X\bar{Y}$ shifts the bigrading by $r - s$, and an endomorphism shifting the bigrading by a nonzero amount has zero trace, so the pairing vanishes unless $r = s$. Setting $Y = X$ with $r = s$ gives the stated formula.

3. The bigrading at gF is $V_{gF}^{p,q} = gV_F^{p,q}$, because g is real and carries $F^\bullet$ to $(gF)^\bullet$; so C_{gF} acts on $gV_F^{p,q}$ as i^{p-q}, which is exactly gC_Fg^{-1}. Since g is real, $\overline{gYg^{-1}} = g\bar{Y}g^{-1}$, and therefore
$$(gXg^{-1})\,C_{gF}\,\overline{gYg^{-1}}\,C_{gF}^{-1} = g\big(XC_F\bar{Y}C_F^{-1}\big)g^{-1}$$
whose trace is unchanged. This is the asserted invariance; combined with the fact that g acts on tangent vectors by $X \mapsto gXg^{-1}$ under the identification of proposition 6.1.1(3), which follows from $g\exp(tX)F = \exp(t\,gXg^{-1})gF$, it says that g acts on D by isometries.

   For smoothness, fix $F_o \in D$ and set $\mathfrak{m} = \mathfrak{g}_\mathbb{R} \cap \bigoplus_{r\neq 0}\mathfrak{g}_{F_o}^{r,-r}$. Both $\mathfrak{g}_{F_o}^{0,0}$ and $\bigoplus_{r\neq 0}\mathfrak{g}_{F_o}^{r,-r}$ are stable under conjugation, so $\mathfrak{g}_\mathbb{R} = (\mathfrak{g}_\mathbb{R} \cap \mathfrak{g}_{F_o}^{0,0}) \oplus \mathfrak{m}$, and $\mathfrak{m}$ is stable under conjugation by K because an element of K fixes F_o and hence preserves each $\mathfrak{g}_{F_o}^{r,-r}$. This exhibits $(\mathfrak{g}_\mathbb{R}, \mathfrak{m})$ as a reductive decomposition for $D = G_\mathbb{R}/K$ in the sense of [12, reductive decomposition]. The projection $\mathfrak{m} \to \bigoplus_{r<0}\mathfrak{g}_{F_o}^{r,-r}$, $X \mapsto \sum_{r<0} X_r$, is an isomorphism of real vector spaces, and for $X, Y \in \mathfrak{m}$ with images ξ, η the orthogonality in part (2) gives $\langle X, Y\rangle_{F_o} = 2\,\mathrm{Re}\langle \xi, \eta\rangle_{F_o}$. So $\mathrm{Re}\,h$ corresponds to the inner product $\frac{1}{2}\,\mathrm{Re}\langle -,-\rangle_{F_o}$ on $\mathfrak{m}$, which is K-invariant by the invariance just proved. By [12, Invariant metrics and Ad(H)-invariant inner products] there is a unique $G_\mathbb{R}$-invariant Riemannian metric on D with that value at F_o, and it is smooth; by transitivity of $G_\mathbb{R}$ and the invariance already proved it coincides with $\mathrm{Re}\,h$, completing the proof.

Part (2) records the sign pattern used below. On $\mathfrak{g}^{r,-r}$ the trace form $\beta(X, \bar{X})$ is positive for odd r and negative for even r, so conjugation by C_F splits $\mathfrak{g}_\mathbb{R}$ the way a Cartan involution does: its $+1$-eigenspace is the sum of the even pieces, where β is negative definite, and its -1-eigenspace is the sum of the odd pieces, where β is positive definite (exercise 6.1.1 (5)). The horizontal directions $\mathfrak{g}^{-1,1}$ sit in the odd part, and their brackets $[\mathfrak{g}^{-1,1}, \mathfrak{g}^{1,-1}] \subseteq \mathfrak{g}^{0,0}$ land in the even part. That is the mechanism behind the curvature estimate: the negative sign of β on the isotropy directions is what makes the horizontal curvature negative.

Carrying the mechanism through to a formula requires the Levi-Civita connection of a $G_\mathbb{R}$-invariant metric on the reductive homogeneous space $G_\mathbb{R}/K$. The series develops invariant metrics on such spaces and the curvature of a bi-invariant metric on a Lie group, but not the connection of a general invariant metric, so the computation is taken from Griffiths and Schmid rather than reproduced. What is proved below is the passage from their pointwise estimate to a uniform one, and the two structural statements about when D is symmetric. For a nonzero real tangent vector $v \in T_FD$ write $K^{\mathrm{hol}}(v)$ for the *holomorphic sectional curvature* of the Hermitian metric h in the direction v: the quantity computed from the Chern connection of h, equivalently the Gaussian curvature that $-\partial\bar{\partial}\log h$ assigns to a holomorphic disc through F tangent to v.

The Hermitian reading is the one meant, and the distinction matters: the Hodge metric is in general *not* Kähler: for $X, Y \in \mathfrak{g}_F^{-1,1}$ one has $[X, Y] \in \mathfrak{g}_F^{-2,2}$ by proposition 6.1.1(1), and the invariant form pairs $\mathfrak{g}_F^{-2,2}$ nontrivially with $\mathfrak{g}_F^{2,-2}$, so the associated $(1,1)$-form has $d\omega \neq 0$ as soon as $\mathfrak{g}_F^{-2,2} \neq 0$. That is exactly the case in which D fails to be Hermitian symmetric, which is the case this section exists to treat. For a Kähler metric the Chern and Levi-Civita connections agree and $K^{\mathrm{hol}}(v)$ coincides with the Riemannian sectional curvature ([12, sectional curvature]) of the real 2-plane spanned by v and Jv; away from that case the two differ, the Chern connection having torsion, and it is the Chern quantity that fact (2) below bounds. Example 6.1.5 is a Kähler case, so the two readings agree there.

Theorem 6.1.4: Curvature of the Hodge Metric

Let D carry the Hodge metric of definition 6.1.2.

1. There is a constant $c > 0$, depending only on the numerical data $(V_{\mathbb{Z}}, Q, k, \{h^{p,q}\})$, such that
$$K^{\mathrm{hol}}(v) \leq -c \qquad \text{for every } F \in D \text{ and every nonzero } v \in T^h_F D$$

2. The vanishing $\mathfrak{g}_F^{r,-r} = 0$ for all $r \leq -2$ is independent of F and is equivalent to $T^h D = TD$. When it holds, the Weil operator C_F acts on D as an isometry fixing F with differential $-\mathrm{id}$, so the geodesic symmetry at every point of D extends to a global isometry and D is a Hermitian symmetric space.

3. If $\mathfrak{g}_F^{r,-r} \neq 0$ for some even $r \neq 0$, then D contains a compact connected complex submanifold of positive dimension. In particular D is not biholomorphic to a bounded domain in $\mathbb{C}^N$, so it is not a bounded symmetric domain.

***Proof*:**

1. Griffiths and Schmid compute the curvature tensor of the Hodge metric on D and show that $K^{\mathrm{hol}}(v) < 0$ for every nonzero horizontal v [21]; that computation is the input taken from outside, for the reason recorded above. Granting it, the uniformity is a compactness argument. Fix $F_o \in D$ and let $S \subseteq T^h_{F_o} D$ be the unit sphere of the Hodge metric, a compact set since $T^h_{F_o} D$ is finite dimensional. The holomorphic sectional curvature of a Hermitian metric is a continuous function of v, being assembled from the Chern curvature tensor and h itself, and is unchanged under real scaling of v, numerator and denominator both being homogeneous of degree four. It is strictly negative on the compact set S, so it attains a maximum $-c < 0$ there. The Chern connection is natural, so K^{hol} is invariant under holomorphic isometries and K^{hol} is invariant under scaling of v, so for arbitrary $F \in D$ and nonzero $v \in T^h_F D$ choose $g \in G_{\mathbb{R}}$ with $gF_o = F$ (proposition 3.3.5); then $dg^{-1}(v)$ is horizontal, because g preserves the horizontal distribution by the equivariance noted after definition 5.3.8, and normalizing it into S gives $K^{\mathrm{hol}}(v) \leq -c$.

2. Independence of F holds because $\mathfrak{g}_{gF}^{r,-r} = g\,\mathfrak{g}_F^{r,-r}\,g^{-1}$ for $g \in G_{\mathbb{R}}$, which follows from $V_{gF}^{p,q} = gV_F^{p,q}$, and $G_{\mathbb{R}}$ acts transitively on D. The equivalence with $T^h D = TD$ is proposition 6.1.1(3): the two spaces agree at F exactly when $\bigoplus_{r<0} \mathfrak{g}_F^{r,-r} = \mathfrak{g}_F^{-1,1}$.

   Assume the vanishing. Conjugation preserves the grading up to sign, so $\mathfrak{g}_F^{r,-r} = 0$ for all $|r| \geq 2$ and $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}_F^{-1,1} \oplus \mathfrak{g}_F^{0,0} \oplus \mathfrak{g}_F^{1,-1}$. By part (2) of proposition 6.1.3, conjugation by C_F acts as $(-1)^r$ on $\mathfrak{g}_F^{r,-r}$, hence as $-\mathrm{id}$ on $\mathfrak{g}_F^{-1,1} \oplus \mathfrak{g}_F^{1,-1}$ and as $+\mathrm{id}$ on $\mathfrak{g}_F^{0,0}$. Now $C_F \in G_{\mathbb{R}}$ by proposition 6.1.3(1) and C_F preserves every $V_F^{p,q}$, hence every F^p, so C_F fixes F; and it acts on D by an isometry, by proposition 6.1.3(3). Its differential at F is $X \mapsto C_F X C_F^{-1}$ on $T^{1,0}_F D \cong \mathfrak{g}_F^{-1,1}$, which is $-\mathrm{id}$.

   A homogeneous Riemannian manifold is complete ([12, homogeneous manifolds are complete]), so normal balls around F are available, and an isometry commutes with the

exponential map. Hence for v in a star-shaped neighbourhood of 0 in $T_F D$,

$$C_F\big(\exp_F(v)\big) = \exp_F\big(d(C_F)_F v\big) = \exp_F(-v)$$

so C_F agrees on that ball with the geodesic symmetry $\exp_F \circ(-\mathrm{id}) \circ \exp_F^{-1}$ of [12, geodesic symmetry and symmetric space]. The geodesic symmetry at F therefore extends to a global isometry, and since F was arbitrary D is a symmetric space. It is Hermitian symmetric because h is a Hermitian metric and each C_F acts holomorphically, being an element of $G_{\mathbb{C}}$ acting on $\check{D}$.

3. Let $Z = \{g \in G_{\mathbb{R}} \big| gC_F = C_F g\}$ be the centralizer of the Weil operator. If $g \in Z$ then, using $H = H_F$ and proposition 6.1.3(1),

$$H(gu, gv) = Q\big(Cgu, \overline{gv}\big) = Q\big(gCu, g\bar{v}\big) = Q(Cu, \bar{v}) = H(u, v)$$

so Z is a closed subset of the unitary group of the positive definite form H, hence compact. An element $X \in \mathfrak{g}_{\mathbb{R}}$ lies in $\mathrm{Lie}(Z)$ if and only if $\exp(tX)C\exp(-tX) = C$ for all t, that is if and only if $XC = CX$; by proposition 6.1.3(2) this is $\mathrm{Lie}(Z) = \mathfrak{g}_{\mathbb{R}} \cap \bigoplus_{r \text{ even}} \mathfrak{g}_F^{r,-r}$. Also $K \subseteq Z$, because an element of K fixes F and hence preserves each $V_F^{p,q}$, on which C is a scalar; so $\mathrm{Lie}(K) = \mathfrak{g}_{\mathbb{R}} \cap \mathfrak{g}_F^{0,0} \subseteq \mathrm{Lie}(Z)$.

Let $Y = Z^0 \cdot F$ be the orbit of the identity component. The orbit map descends to an injection $Z^0/(Z^0 \cap K) \to D$ of the quotient manifold ([37, The Quotient Manifold Theorem]), whose differential at the base point is the inclusion $\mathrm{Lie}(Z)/\mathrm{Lie}(K) \hookrightarrow \mathfrak{g}_{\mathbb{R}}/\mathrm{Lie}(K)$ and hence injective; by Z^0-equivariance it is injective at every point, so the map is an injective immersion. Its source is compact, so it is a homeomorphism onto its image and Y is a compact connected embedded submanifold of D of dimension

$$\dim_{\mathbb{R}} Y = \dim_{\mathbb{R}}\Big(\mathfrak{g}_{\mathbb{R}} \cap \bigoplus_{r \text{ even}, r \neq 0} \mathfrak{g}_F^{r,-r}\Big)$$

This is positive under the hypothesis, because a conjugation-stable complex subspace of $\mathfrak{g}_{\mathbb{C}}$ of complex dimension d has real points of real dimension d.

The tangent space $T_F Y$ corresponds, under proposition 6.1.1(3), to $\bigoplus_{r<0,\ r \text{ even}} \mathfrak{g}_F^{r,-r}$, a complex subspace of $T_F^{1,0} D$; and Z^0 acts transitively on Y by biholomorphisms of D, so every tangent space of Y is stable under the complex structure J of D. The Nijenhuis tensor of J vanishes on D, which is a complex manifold,[a] and brackets of vector fields tangent to Y remain tangent to Y, so the almost complex structure induced on Y has vanishing Nijenhuis tensor; the Newlander-Nirenberg theorem [5, Newlander-Nirenberg] makes Y a complex manifold, and the inclusion is holomorphic because its differential is $\mathbb{C}$-linear.

Suppose finally that $\varphi\colon D \to \Omega \subseteq \mathbb{C}^N$ were a biholomorphism onto a bounded domain. Each coordinate function φ_j restricts to a holomorphic function on the compact connected complex manifold Y, hence is constant there by the maximum principle.[b] Then $\varphi(Y)$ is a single point, contradicting the injectivity of φ and $\dim Y > 0$. So no such φ exists, completing the proof.

[a] [5, Complex Manifolds Have Vanishing Nijenhuis Tensor].
[b] [5, Maximum Principle on Compact Complex Manifolds].

Parts (2) and (3) mark the boundary of the classical theory. A Hermitian symmetric space of noncompact type is biholomorphic to a bounded symmetric domain (Harish-Chandra's realization, [1, Hermitian symmetric spaces and their bounded realizations]), so part (3) says that as soon as $\mathfrak{g}$ has a nonzero even piece in a degree other than 0, the domain D leaves that world, and with it leaves the setting in which negative curvature holds in every direction. What survives is exactly part (1): an estimate on T^hD and nothing more. The full statement, that D is Hermitian symmetric precisely when $T^hD = TD$, is in [21]. The first numerical type meeting the hypothesis of part (3) is weight 2 with $h^{2,0} = 2$, the case in which exercise 5.3.1 (3) found T^hD of codimension 1; there $\mathfrak{g}^{-2,2}$ is one-dimensional (exercise 6.1.1 (3)), so the compact complex submanifold produced by the proof is a curve.

The classical cases sit on the other side of the boundary, and the weight-one case can be computed outright.

Example 6.1.5: The Hodge Metric on the Siegel Upper Half Space

Take $k = 1$ and $h^{1,0} = h^{0,1} = g$, the numerical type of example 3.3.3, so that $D \cong \mathfrak{H}_g$ and $G_{\mathbb{R}} = \mathrm{Sp}(V_{\mathbb{R}}, Q)$.

The grading has three pieces. There are only two bidegrees, $(1,0)$ and $(0,1)$, so an endomorphism shifting the bigrading by r is zero unless $r \in \{-1, 0, 1\}$. Hence $\mathfrak{g}^{r,-r} = 0$ for $|r| \geq 2$, and by theorem 6.1.4(2) the horizontal subbundle is everything and $\mathfrak{H}_g$ is a Hermitian symmetric space. This is the period-domain form of the observation made by hand in example 5.3.9, and it says that in weight one the infinitesimal period relation imposes nothing: every holomorphic map into $\mathfrak{H}_g$ is horizontal.

The metric for $g = 1$. Take $g = 1$, so $V_{\mathbb{R}} = \mathbb{R}^2$ with symplectic basis e, f and $Q(e, f) = 1$, and present a point of D as $F^1 = \mathbb{C}(e + \tau f)$ with $\tau \in \mathfrak{H}_1$, as in example 3.3.3. Work at $\tau = i$ and write $v = e + if$, so $V^{1,0} = \mathbb{C}v$ and $V^{0,1} = \mathbb{C}\bar{v}$. An element $X \in \mathfrak{g}^{-1,1}$ satisfies $X_a v = a\bar{v}$ and $X_a \bar{v} = 0$ for some $a \in \mathbb{C}$; conversely every such X_a is Q-skew, since $Q(X_a v, \bar{v}) + Q(v, X_a \bar{v}) = aQ(\bar{v}, \bar{v}) = 0$ and $Q(X_a v, v) + Q(v, X_a v) = a\big(Q(\bar{v}, v) + Q(v, \bar{v})\big) = 0$ by antisymmetry of Q, while the remaining pairs of basis vectors give $0 = 0$. So $\mathfrak{g}^{-1,1} = \{X_a \,|\, a \in \mathbb{C}\}$ is a line, in accordance with $\dim_{\mathbb{C}} \mathfrak{H}_1 = 1$.

The Hodge norm is computed from proposition 6.1.3(2) with $r = -1$, where the sign is $(-1)^{r+1} = +1$. Since $\bar{X}_a$ sends $\bar{v}$ to $\bar{a}v$ and kills v, the composite $X_a \bar{X}_a$ kills v and multiplies $\bar{v}$ by $|a|^2$, so

$$\|X_a\|^2 = \mathrm{tr}(X_a \bar{X}_a) = |a|^2$$

To read this on $\mathfrak{H}_1$, follow the curve $\exp(tX_a)F$. From $X_a^2 = 0$ one gets $\exp(tX_a)v = v + ta\bar{v} = (1 + ta)e + i(1 - ta)f$, and dividing by $1 + ta$, which changes no subspace, presents the same line as $e + \tau(t)f$ with $\tau(t) = i(1 - ta)/(1 + ta)$. Differentiating at $t = 0$ gives $\dot{\tau} = -2ia$. So the tangent vector with τ-component w corresponds to $a = iw/2$ and has squared Hodge norm $|a|^2 = |w|^2/4$: at $\tau = i$ the Hodge metric is $\frac{1}{4}|d\tau|^2$.

Both the Hodge metric and the Poincaré metric $|d\tau|^2/(\mathrm{Im}\,\tau)^2$ are invariant under $\mathrm{SL}(2, \mathbb{R}) = \mathrm{Sp}(V_{\mathbb{R}}, Q)$, which acts transitively on $\mathfrak{H}_1$, and they agree up to the factor $\frac{1}{4}$ at $\tau = i$. Two invariant

metrics agreeing at one point of a homogeneous space agree everywhere, so

$$h = \frac{|d\tau|^2}{4\,(\operatorname{Im}\tau)^2}$$

The Poincaré metric has constant Gaussian curvature -1, and scaling a surface metric by $\frac{1}{4}$ multiplies its curvature by 4; so the Hodge metric on $\mathfrak{H}_1$ has constant holomorphic sectional curvature -4, and $c = 4$ serves in theorem 6.1.4(1). Here the bound holds in every direction, because every direction is horizontal. In the general case the same constant governs only the horizontal ones.

Negative curvature bounds the derivative of a holomorphic map into D. Making that precise takes two classical facts of complex hyperbolic geometry, neither of which is developed anywhere in the series; each is stated with the source it is taken from and is used here exactly as stated. Let $\Delta = \{z \in \mathbb{C} \big| |z| < 1\}$ and let

$$\rho = \frac{4\,|dz|^2}{(1-|z|^2)^2}$$

be the Poincaré metric on Δ, of constant curvature -1 and equal to $4|dz|^2$ at the origin. The two facts are:

1. (*Ahlfors-Schwarz lemma*) If $\sigma = \lambda|dz|^2$ is a conformal pseudo-metric on Δ with $\lambda \geq 0$ smooth, whose Gaussian curvature satisfies $K_\sigma \leq -c$ wherever $\lambda > 0$, with $c > 0$, then $\sigma \leq c^{-1}\rho$. This is Ahlfors's curvature form of the classical Schwarz lemma ([31, Schwarz' Lemma]), and is taken from [1, the Ahlfors-Schwarz lemma].
2. (*curvature decreases on holomorphic curves*) If $f\colon \Delta \to (M, h)$ is holomorphic into a Hermitian manifold and $df_{z_0} \neq 0$, then the Gaussian curvature of f^*h at z_0 is at most the holomorphic sectional curvature of h at $f(z_0)$ in the direction $df_{z_0}(\partial/\partial z)$. This is the standard curvature comparison for a holomorphic curve in a Hermitian manifold, taken from [1, curvature estimates for holomorphic maps]; it is what turns the estimate of theorem 6.1.4(1), itself taken from [21], into a curvature bound on Δ.

Corollary 6.1.6: Horizontal Maps Are Distance-Decreasing

Let $c > 0$ be as in theorem 6.1.4(1), let $\Gamma \subseteq G_{\mathbb{Z}}$ act freely on D, and give $\Gamma\backslash D$ the Hermitian metric descended from the Hodge metric. Then every horizontal holomorphic map $f\colon \Delta \to \Gamma\backslash D$ satisfies

$$f^*h \leq c^{-1}\rho$$

In particular f is distance-decreasing from $c^{-1}\rho$ to the Hodge metric, and $\|df_0(\partial/\partial z)\|_h^2 \leq 4/c$.

***Proof*:**
By proposition 5.3.5 the group Γ acts properly discontinuously, and acting freely as well it makes $q\colon D \to \Gamma\backslash D$ a holomorphic covering map. The Hodge metric is Γ-invariant by proposition 6.1.3(3), so it descends and q is a local isometry; the horizontal distribution is $G_{\mathbb{C}}$-equivariant, as noted after definition 5.3.8, so it too descends, and a map into $\Gamma\backslash D$ is called horizontal when its differential lands in the descended distribution. Since Δ is simply connected, f lifts to a holomorphic map $\tilde{f}\colon \Delta \to D$ with $q \circ \tilde{f} = f$; the lift is horizontal because q is a local

biholomorphism matching the two distributions, and $f^*h = \tilde{f}^*h$.

Write $\tilde{f}^*h = \lambda|dz|^2$, a conformal pseudo-metric on Δ, smooth because $\tilde{f}$ and h are, and vanishing exactly where $d\tilde{f}$ does. At a point z_0 with $d\tilde{f}_{z_0} \neq 0$, the vector $d\tilde{f}_{z_0}(\partial/\partial z)$ is a nonzero horizontal tangent vector, so fact (2) followed by theorem 6.1.4(1) gives

$$K_{\tilde{f}^*h}(z_0) \leq K^{\mathrm{hol}}\big(d\tilde{f}_{z_0}(\partial/\partial z)\big) \leq -c$$

Fact (1) then yields $\tilde{f}^*h \leq c^{-1}\rho$ on Δ. Evaluating at the origin, where $\rho = 4|dz|^2$, gives the stated bound on the derivative, completing the proof.

Nothing in the corollary holds for an arbitrary holomorphic map into D: the estimate is available only because theorem 6.1.4(1) is restricted to T^hD, and by theorem 5.3.10 every period map satisfies that restriction. The consequence is a Liouville theorem, and through it a rigidity statement about families.

Theorem 6.1.7: Rigidity of Horizontal Maps

Let $\Gamma \subseteq G_{\mathbb{Z}}$ act freely on D and give $X_\Gamma = \Gamma\backslash D$ the descended Hodge metric.

1. Every horizontal holomorphic map $\mathbb{C} \to X_\Gamma$ is constant.
2. Every horizontal holomorphic map $\mathbb{C}\mathrm{P}^n \to X_\Gamma$ is constant, and so is every horizontal holomorphic map $M \to X_\Gamma$ from a complex manifold M whose universal cover is $\mathbb{C}^n$, for instance a complex torus.
3. If $f\colon \mathcal{X} \to \mathbb{C}\mathrm{P}^n$ is a smooth projective family, then its period map in every degree k is constant.

***Proof*:**

1. Since $\mathbb{C}$ is simply connected and $D \to X_\Gamma$ is a covering, a horizontal holomorphic $f\colon \mathbb{C} \to X_\Gamma$ lifts to a horizontal holomorphic $\tilde{f}\colon \mathbb{C} \to D$. Fix $z_0 \in \mathbb{C}$; after composing with a translation of $\mathbb{C}$, which is a biholomorphism, assume $z_0 = 0$. For $R > 0$ the map $g_R\colon \Delta \to D$, $g_R(z) = \tilde{f}(Rz)$, is horizontal and holomorphic, so corollary 6.1.6 gives

   $$R^2 \left\|d\tilde{f}_0(\partial/\partial z)\right\|_h^2 = \left\|d(g_R)_0(\partial/\partial z)\right\|_h^2 \leq \frac{4}{c}$$

   Letting $R \to \infty$ forces $d\tilde{f}_0 = 0$. As z_0 was arbitrary, $d\tilde{f} \equiv 0$, and $\mathbb{C}$ being connected, $\tilde{f}$ and hence f is constant.

2. Both cases reduce to (1). For $\mathbb{C}\mathrm{P}^n$, which is simply connected, f lifts to $\tilde{f}\colon \mathbb{C}\mathrm{P}^n \to D$. Let $\ell \subseteq \mathbb{C}\mathrm{P}^n$ be a projective line and $p \in \ell$; then $\ell \setminus \{p\} \cong \mathbb{C}$ and $\tilde{f}$ restricted to it is horizontal holomorphic, hence constant by (1), and continuity extends the constancy to ℓ. Any two points of $\mathbb{C}\mathrm{P}^n$ lie on a common line, so $\tilde{f}$ is constant.

   For M with universal cover $\pi\colon \mathbb{C}^n \to M$, the composite $f \circ \pi$ is horizontal holomorphic and lifts to $\tilde{f}\colon \mathbb{C}^n \to D$. Restricting $\tilde{f}$ to the complex lines through a given point of $\mathbb{C}^n$

gives horizontal holomorphic maps $\mathbb{C} \to D$, constant by (1); so every directional derivative of $\tilde{f}$ vanishes, $\tilde{f}$ is constant, and f is constant because π is surjective.

3. The base $\mathbb{C}\mathrm{P}^n$ is simply connected, so the monodromy representation is trivial and $\Gamma = \{1\}$; by theorem 5.3.7 the period map is a globally defined holomorphic map $\Phi\colon \mathbb{C}\mathrm{P}^n \to D$, and by theorem 5.3.10 it is horizontal. Part (2) applies with $\Gamma = \{1\}$ and makes Φ constant, completing the proof.

Part (3) is a statement about geometry, not about period domains. A smooth projective family over $\mathbb{C}\mathrm{P}^n$ has, in every degree, a Hodge filtration on the primitive cohomology that is flat for the Gauss-Manin connection (exercise 5.3.1 (1)), so parallel transport between any two fibres is an isomorphism of polarized Hodge structures on the primitive parts: the fibres are indistinguishable by transcendental invariants. Set this against the Lefschetz pencils of chapter 4, which are families over $\mathbb{C}\mathrm{P}^1$ with nontrivial monodromy around the critical values. There is no contradiction, because a pencil is smooth only over the complement of those values, and the theorem says that this is forced: the singular fibres are not an artifact of the construction. A family that varies must degenerate.

What happens over a base that is quasi-projective rather than compact is a separate theory, and it is Schmid's. The nilpotent orbit and SL(2)-orbit theorems of [35] describe the asymptotics of a period map as the base approaches a boundary divisor, and yield the monodromy theorem, that local monodromy about a boundary component is quasi-unipotent. Those results are not developed here: the limiting objects they produce are no longer pure Hodge structures, and nothing in this chapter needs them.

Exercise 6.1.1

1. Let $F \in D$ with Weil operator $C = C_F$. Show directly from definition 3.2.1 that $Q(Cu, Cv) = Q(u, v)$ for all $u, v \in V_{\mathbb{C}}$, and that $H_F(u, v) = Q(Cu, \bar{v})$, where H_F is the form (3.6). These two identities are part of proposition 6.1.3(1); the exercise is to rederive them by hand from the definitions. Deduce that $C^2 = (-1)^k \,\mathrm{id}$ lies in the centre of $G_{\mathbb{R}}$ and that conjugation by C is an involution of $\mathfrak{g}_{\mathbb{R}}$.

2. Take $k = 2$ with $h^{2,0} = h^{0,2} = 1$ and $h^{1,1} = n$. This is the shape of the Hodge structure on the H^2 of a K3 surface (example 3.4.4), though that example also records that the cup product on the whole of H^2 is not a polarization, so the period domain here is the one attached to the primitive part. Show that $\mathfrak{g}_F^{r,-r} = 0$ for every $|r| \geq 2$, and conclude from theorem 6.1.4(2) that this period domain is a Hermitian symmetric space on which every holomorphic map is horizontal, recovering the computation of example 5.3.9.

3. Take $k = 2$ with $h^{2,0} = h^{0,2} = 2$ and $h^{1,1} = n$. Show that $\dim_{\mathbb{C}} \mathfrak{g}_F^{-2,2} = 1$ and that $\mathfrak{g}_F^{r,-r} = 0$ for $|r| \geq 3$. Deduce that $T^h D$ has codimension 1 in TD, in agreement with exercise 5.3.1 (3), and that the compact complex submanifold produced by theorem 6.1.4(3) has real dimension 2. Conclude that this D is not biholomorphic to a bounded domain.

4. Show that $\mathfrak{g}_{gF}^{r,-r} = g\,\mathfrak{g}_F^{r,-r}\,g^{-1}$ for every $g \in G_{\mathbb{R}}$ and $F \in D$, and deduce that neither the hypothesis of theorem 6.1.4(3) nor the numbers $\dim_{\mathbb{C}} \mathfrak{g}_F^{r,-r}$ depend on the choice of F. Where does the argument use that g is real, and not just that it lies in $G_{\mathbb{C}}$?

5. Let $\beta(X, Y) = \mathrm{tr}(XY)$ be the trace form on $\mathfrak{g}_{\mathbb{C}}$ and fix $F \in D$. Using proposition 6.1.3(2), show that $\beta(X, \bar{X}) > 0$ for nonzero $X \in \mathfrak{g}_F^{r,-r}$ with r odd and $\beta(X, \bar{X}) < 0$ for nonzero X with r even. Deduce that β is negative definite on $\mathfrak{g}_{\mathbb{R}} \cap \bigoplus_{r \text{ even}} \mathfrak{g}_F^{r,-r}$ and positive definite on

$\mathfrak{g}_{\mathbb{R}} \cap \bigoplus_{r \text{ odd}} \mathfrak{g}_F^{r,-r}$, that the two are β-orthogonal, and that writing θ for conjugation by C_F the form $-\beta(X, \theta Y)$ is positive definite on $\mathfrak{g}_{\mathbb{R}}$.

6. Show that $\tau \mapsto (\tau - i)/(\tau + i)$ is a biholomorphism from $\mathfrak{H}_1$ onto the unit disc Δ, so that the weight-one period domain of example 6.1.5 with $g = 1$ *is* biholomorphic to a bounded domain. Using corollary 6.1.6 and the constant $c = 4$ computed there, write down the resulting bound on $|f'(0)|$ for a holomorphic map $f\colon \Delta \to \mathfrak{H}_1$, and compare it with the bound obtained by transporting the classical Schwarz lemma through the biholomorphism.

6.2 Mumford-Tate Groups

A period domain records a polarized Hodge structure as a point of a homogeneous space, and the group $G_{\mathbb{R}}$ of proposition 3.3.5 moves one structure to another. That group depends only on the numerical data $(V_{\mathbb{Q}}, Q, k, \{h^{p,q}\})$, so it sees nothing about the individual structure sitting at a given point. This section attaches to a single rational Hodge structure a group that does: a $\mathbb{Q}$-algebraic subgroup of $\mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m$ whose invariants in the tensor constructions built from V are exactly the Hodge classes.

The construction has two steps. A Hodge structure on $V_{\mathbb{Q}}$ is first rewritten as an action of a two-dimensional real algebraic torus on $V_{\mathbb{R}}$; the group is then the smallest subgroup defined over $\mathbb{Q}$ whose real points carry that action. The ambient language of algebraic groups, their tori and their reductivity is taken from [14] and not rebuilt here; what is developed here is the Hodge-theoretic content, which is the invariant-theoretic characterization of Hodge classes and the reductivity that a polarization forces.

Throughout, a *rational Hodge structure* of weight k means the datum of definition 3.1.1 on a finite-dimensional $\mathbb{Q}$-vector space $V_{\mathbb{Q}}$: a bigrading $V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$ with $\overline{V^{p,q}} = V^{q,p}$. An *algebraic subgroup* of an algebraic group over $\mathbb{Q}$ means a Zariski-closed subgroup defined over $\mathbb{Q}$.[1]

The bigrading and its conjugation symmetry are, together, an action of $\mathbb{C}^\times$ on the real vector space $V_{\mathbb{R}}$, in which z acts on $V^{p,q}$ by the scalar $z^p \bar{z}^q$: the symmetry $\overline{V^{p,q}} = V^{q,p}$ is precisely what makes that prescription commute with conjugation and hence preserve $V_{\mathbb{R}}$. The group carrying the action is algebraic over $\mathbb{R}$ rather than over $\mathbb{C}$, since the prescription involves $\bar{z}$ alongside z.

Definition 6.2.1: The Deligne Torus

The *Deligne torus* is the real algebraic group

$$\mathbb{S} = \mathrm{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$$

obtained from $\mathbb{G}_m$ over $\mathbb{C}$ by restriction of scalars: for a commutative $\mathbb{R}$-algebra A it is $\mathbb{S}(A) = (A \otimes_{\mathbb{R}} \mathbb{C})^\times$, and its coordinate ring is $\mathbb{R}[x, y, (x^2 + y^2)^{-1}]$ with $z = x + iy$ the tautological coordinate. In particular $\mathbb{S}(\mathbb{R}) = \mathbb{C}^\times$, and the inclusion $\mathbb{S}(\mathbb{R}) \subseteq \mathbb{S}(\mathbb{C})$ is $z \mapsto (z, \bar{z})$ under the isomorphism $\mathbb{S}_{\mathbb{C}} \cong \mathbb{G}_m \times \mathbb{G}_m$ induced by $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} \cong \mathbb{C} \times \mathbb{C}$.
Three homomorphisms of real algebraic groups are attached to it: the inclusion $\mathbb{G}_{m,\mathbb{R}} \hookrightarrow \mathbb{S}$ of real scalars, the *norm* $\mathrm{Nm}\colon \mathbb{S} \to \mathbb{G}_{m,\mathbb{R}}$, given on real points by $z \mapsto z\bar{z}$, and the *circle* $\mathbb{U} = \ker(\mathrm{Nm})$, with $\mathbb{U}(\mathbb{R}) = \{z \in \mathbb{C}^\times \,\big|\, z\bar{z} = 1\}$.

[1] [14, Algebraic Group, Closed Subgroup].

The characters of $\mathbb{S}_{\mathbb{C}} \cong \mathbb{G}_m \times \mathbb{G}_m$ are the monomials $z^p \bar{z}^q$ with $(p, q) \in \mathbb{Z}^2$, written in the coordinates given by the isomorphism above; complex conjugation on $\mathbb{S}$, that is the nontrivial element of $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$, interchanges the two factors and hence sends $z^p \bar{z}^q$ to $z^q \bar{z}^p$. That interchange is the source of the conjugation symmetry in the next statement.

Proposition 6.2.2: Hodge Structures Are Representations of $\mathbb{S}$

Let $V_{\mathbb{Q}}$ be a finite-dimensional $\mathbb{Q}$-vector space and $k \in \mathbb{Z}$. The following data are equivalent.

1. A rational Hodge structure of weight k on $V_{\mathbb{Q}}$, that is a bigrading $V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$ with $\overline{V^{p,q}} = V^{q,p}$.
2. A homomorphism of real algebraic groups $h \colon \mathbb{S} \to \mathrm{GL}(V_{\mathbb{R}})$ whose restriction to the subgroup $\mathbb{G}_{m,\mathbb{R}}$ of real scalars is $r \mapsto r^k \,\mathrm{id}$.

The passage from (1) to (2) sends the bigrading to the homomorphism acting on $V^{p,q}$ as multiplication by $z^p \bar{z}^q$; the passage from (2) to (1) takes $V^{p,q}$ to be the subspace of $V_{\mathbb{C}}$ on which $\mathbb{S}_{\mathbb{C}}$ acts through the character $z^p \bar{z}^q$. The two constructions are mutually inverse.

***Proof*:**

Step 1: a homomorphism gives a bigrading. Let h be as in (2) and extend it to $\mathbb{S}_{\mathbb{C}} \to \mathrm{GL}(V_{\mathbb{C}})$. The group $\mathbb{S}_{\mathbb{C}} \cong \mathbb{G}_m \times \mathbb{G}_m$ is a torus, so $V_{\mathbb{C}}$ decomposes as the direct sum of its character eigenspaces ([14, Weight Space Decomposition]):

$$V_{\mathbb{C}} = \bigoplus_{(p,q) \in \mathbb{Z}^2} V^{p,q}$$

where $V^{p,q}$ is the eigenspace on which $\mathbb{S}_{\mathbb{C}}$ acts through the character $z^p \bar{z}^q$; evaluating that character at a point of $\mathbb{S}(\mathbb{R}) = \mathbb{C}^\times$ gives $h(z)v = z^p \bar{z}^q v$ for $v \in V^{p,q}$.

Only the pairs with $p + q = k$ occur. For $r \in \mathbb{R}^\times$ the operator $h(r)$ acts on $V^{p,q}$ by $r^p r^q = r^{p+q}$, while the hypothesis in (2) makes it r^k; taking $r = 2$ gives $2^{p+q} = 2^k$ on any nonzero $V^{p,q}$, so $V^{p,q} = 0$ unless $p + q = k$. In particular the sum is finite.

For the symmetry, note that $h(z)$ is an endomorphism of the real space $V_{\mathbb{R}}$, extended $\mathbb{C}$-linearly, so it commutes with the conjugation of $V_{\mathbb{C}}$ over $V_{\mathbb{R}}$. Let $v \in V^{p,q}$. Then

$$h(z)\bar{v} = \overline{h(z)v} = \overline{z^p \bar{z}^q v} = z^q \bar{z}^p \, \bar{v}$$

so $\bar{v} \in V^{q,p}$, and applying the same to $V^{q,p}$ gives $\overline{V^{p,q}} = V^{q,p}$. This is a Hodge structure of weight k.

Step 2: a bigrading gives a homomorphism. Let the bigrading be given and define $h(z)v = \sum_{p+q=k} z^p \bar{z}^q \, v^{p,q}$ for $v \in V_{\mathbb{C}}$ with components $v^{p,q}$. It is multiplicative in z because each summand is, and it preserves $V_{\mathbb{R}}$: for v real one has $\overline{v^{p,q}} = v^{q,p}$ by the symmetry, so

$$\overline{h(z)v} = \sum_{p+q=k} \bar{z}^p z^q \, \overline{v^{p,q}} = \sum_{p+q=k} \bar{z}^p z^q \, v^{q,p} = h(z)v$$

the last equality by the substitution $(p, q) \mapsto (q, p)$. So $h(z)$ restricts to an $\mathbb{R}$-linear automorphism of $V_{\mathbb{R}}$.

It is algebraic. Fix a basis of $V_{\mathbb{R}}$ and write $z = x + iy$. The function $z \mapsto z^p \bar{z}^q$ lies in $\mathbb{R}[x, y, (x^2 + y^2)^{-1}] = \mathcal{O}(\mathbb{S})$ after taking real and imaginary parts, since $z\bar{z} = x^2 + y^2$ and

$z^{-1} = \bar{z}(x^2+y^2)^{-1}$; hence so does every matrix entry of $h(z)$, these being $\mathbb{R}$-linear combinations of such functions. Finally $h(r)$ acts on $V^{p,q}$ by $r^{p+q} = r^k$ for real r, which is the condition in (2).

Step 3: the constructions are inverse. Starting from a bigrading, the homomorphism built in Step 2 acts on $V^{p,q}$ by the character $z^p\bar{z}^q$, and distinct pairs (p,q) give distinct characters, so the eigenspace decomposition recovered in Step 1 is the original one. Starting from h, Step 1 produces the bigrading by eigenspaces and Step 2 rebuilds h from it by the same formula, completing the proof.

The proposition replaces a piece of linear algebra by a representation, and the gain is that representations have automorphism groups and invariants while bigradings do not. The restriction of h to real scalars is the *weight homomorphism*: it detects k and nothing else, so the weight is exactly the part of the structure visible to the split subtorus $\mathbb{G}_{m,\mathbb{R}} \subseteq \mathbb{S}$. And the circle $\mathbb{U}$ sees the rest: $h|_{\mathbb{U}}$ determines the bigrading up to the weight, since $\mathbb{C}^\times$ is generated by $\mathbb{R}_{>0}$ and the unit circle.

Example 6.2.3: The Circle Action in Weight One and on the Tate Structure
Two instances, one in each direction of proposition 6.2.2.

1. *Weight one.* Here $V_{\mathbb{C}} = V^{1,0} \oplus V^{0,1}$, and $h(z)$ acts by z on $V^{1,0}$ and by $\bar{z}$ on $V^{0,1}$. Then $h(i)$ acts by i and $-i$ on the two summands, so $h(i)$ is precisely the complex structure J on $V_{\mathbb{R}}$ produced from $V^{1,0}$ in proposition 3.1.4. The weight-one dictionary of theorem 3.4.1 therefore reads: the complex torus attached to V is $V_{\mathbb{R}}/V_{\mathbb{Z}}$ with the complex structure $h(i)$.

   For the elliptic curve E_τ of example 3.4.3, take $V_{\mathbb{Q}} = H^1(E_\tau, \mathbb{Q})$ with the basis ℓ_1, ℓ_2 dual to $1, \tau$, and recall $H^{1,0} = \mathbb{C}\,e$ with $e = \ell_1 + \tau\ell_2$, so $H^{0,1} = \mathbb{C}\,\bar{e}$ with $\bar{e} = \ell_1 + \bar{\tau}\ell_2$. Writing $\tau = s + it$ and $z = x + iy$, the relations $e - \bar{e} = 2it\,\ell_2$ and $\tau\bar{e} - \bar{\tau}e = 2it\,\ell_1$ invert the change of basis, and $h(z)e = ze$, $h(z)\bar{e} = \bar{z}\bar{e}$ give

$$h(z)\ell_1 = \Big(x - \frac{sy}{t}\Big)\ell_1 - \frac{(s^2+t^2)y}{t}\,\ell_2, \qquad h(z)\ell_2 = \frac{y}{t}\,\ell_1 + \Big(x + \frac{sy}{t}\Big)\ell_2$$

   using $z - \bar{z} = 2iy$, $z\tau - \bar{z}\bar{\tau} = 2i(xt + ys)$ and $\tau\bar{z} - \bar{\tau}z = 2i(tx - sy)$. The matrix of $h(z)$ in the basis (ℓ_1, ℓ_2) has determinant $x^2 - s^2y^2/t^2 + (s^2+t^2)y^2/t^2 = x^2 + y^2 = \mathrm{Nm}(z)$, as it must, the two eigenvalues being z and $\bar{z}$; and at $z = i$ it squares to $-\mathrm{id}$, exhibiting $h(i)$ as a complex structure on $H^1(E_\tau, \mathbb{R})$.

2. *The Tate structure.* The structure $\mathbb{Q}(m) := \mathbb{Z}(m) \otimes_{\mathbb{Z}} \mathbb{Q}$ of definition 3.1.11 is concentrated in bidegree $(-m,-m)$ and has weight $-2m$, so $h_{\mathbb{Q}(m)}(z)$ is multiplication by $z^{-m}\bar{z}^{-m} = \mathrm{Nm}(z)^{-m}$. Thus $h_{\mathbb{Q}(m)}$ factors through the norm, and the circle $\mathbb{U}$ acts trivially on every Tate structure. This is the precise sense in which a Tate structure carries weight and nothing else.

The group to be constructed will act not only on V but on every space built from V by multilinear algebra, because that is where Hodge classes other than the obvious ones live. The constructions are transparent in the representation picture: apply the operation to the representation of $\mathbb{S}$ and read off the bigrading again.

The tensor constructions on Hodge structures, and the Hodge classes they isolate, were set up in section 3.1: the tensor product and dual of definition 3.1.9, the mixed tensor $T^{a,b,m}(V)$, and $\mathrm{Hdg}(W) = W_{\mathbb{Q}} \cap W^{p,p}$ of definition 3.1.10. They are used here without further comment.

Both constructions satisfy the conjugation symmetry, so they are Hodge structures: under proposition 6.2.2 they are the tensor product and the dual of the corresponding representations of $\mathbb{S}$, and $h_{V\otimes W} = h_V \otimes h_W$, $h_{V^\vee} = (h_V^\vee)^{-1}$ are again defined over $\mathbb{R}$ with the correct weight. The bidegree formulas are then the statement that the characters of a tensor product multiply and those of a dual invert. Definition 3.1.7 and definition 3.1.11 are the cases already in use: a morphism $V \to W$ of Hodge structures is a rational element of $(V^\vee \otimes W)^{0,0}$, and $V(n) = V \otimes \mathbb{Q}(n)$.

Elements of bidegree (p,p) that are rational are the classes the theory exists to detect. On a projective manifold they are the candidates for classes of algebraic subvarieties. In degree 2 this is settled: theorem 2.3.2 says an integral $(1,1)$-class is the first Chern class of a holomorphic line bundle, and on a projective manifold every line bundle is $\mathcal{O}_X(D)$ for a divisor D, so the integral $(1,1)$-classes are exactly the divisor classes. In higher degree the corresponding question is open, and is the Hodge conjecture.

Under the Tate twist a Hodge class of W becomes a rational class of bidegree $(0,0)$: the identification $W(p)_\mathbb{Q} = W_\mathbb{Q} \otimes (2\pi i)^p$ of definition 3.1.11 carries $\mathrm{Hdg}(W)$ isomorphically onto $W(p)_\mathbb{Q} \cap W(p)^{0,0}$, and it is this weight-zero form that the group-theoretic description below uses, since bidegree $(0,0)$ is where $h(\mathbb{S})$ acts trivially and bidegree (p,p) with $p \neq 0$ is not. For X a compact Kähler manifold, $\mathrm{Hdg}\big(H^2(X,\mathbb{Q})\big) = \mathrm{NS}(X) \otimes_\mathbb{Z} \mathbb{Q}$ by definition 2.3.3, since a rational class of type $(1,1)$ has an integral multiple in $H^2(X,\mathbb{Z})$; for a K3 surface this is the hyperplane section $\sigma^\perp \cap H^2(X,\mathbb{Q})$ computed in example 3.4.4, of dimension $\rho(X)$.

Nothing so far uses the rational structure beyond the requirement that conjugation be defined. The group is what forces $\mathbb{Q}$ into the picture: it is the smallest algebraic group defined over $\mathbb{Q}$ that carries the action of $\mathbb{S}$. The extra factor $\mathbb{G}_m$ in the ambient group is what lets Tate twists be tracked, so that Hodge classes in every weight are covered at once.

Definition 6.2.4: The Mumford-Tate Group

Let V be a rational Hodge structure of weight k, with $h\colon \mathbb{S} \to \mathrm{GL}(V_\mathbb{R})$ as in proposition 6.2.2, and consider the homomorphism of real algebraic groups

$$(h, \mathrm{Nm})\colon \mathbb{S} \to \mathrm{GL}(V_\mathbb{R}) \times \mathbb{G}_{m,\mathbb{R}}$$

The *Mumford-Tate group* $\mathrm{MT}(V)$ is the smallest algebraic subgroup $M \subseteq \mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m$ defined over $\mathbb{Q}$ whose group of real points contains $(h,\mathrm{Nm})\big(\mathbb{S}(\mathbb{R})\big)$.

The group exists. The family of algebraic subgroups of $\mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m$ defined over $\mathbb{Q}$ and containing the image in their real points is nonempty, as it contains $\mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m$; an intersection of algebraic subgroups is an algebraic subgroup, and $\big(\bigcap_i M_i\big)(\mathbb{R}) = \bigcap_i M_i(\mathbb{R})$ still contains the image, so the intersection of the whole family is the smallest member. Equivalently, $\mathrm{MT}(V)$ is the Zariski closure over $\mathbb{Q}$ of $(h,\mathrm{Nm})(\mathbb{S}(\mathbb{R}))$: that closure is an algebraic subgroup, because the closure of a subgroup is a subgroup, it is defined over $\mathbb{Q}$ by construction, and it is contained in every member of the family since each is Zariski closed.

The group is nontrivial as soon as $V \neq 0$: the real scalars contribute the one-parameter subgroup $t \mapsto (t^k \,\mathrm{id},\, t^2)$ (exercise 6.2.1 (3)). The second coordinate is not redundant even when $k = 0$, because Nm is nonconstant while h may be trivial; it is what makes the group act on Tate structures, and hence on every twist. The Tate structure is also the smallest instance of the definition: for $V = \mathbb{Q}(n)$ the homomorphism h is $z \mapsto \mathrm{Nm}(z)^{-n}$ by example 6.2.3, so (h,Nm) has image $\{(s^{-n},\, s)\big|s \in \mathbb{R}_{>0}\}$ and $\mathrm{MT}(\mathbb{Q}(n))$ is the one-dimensional torus $\{(t^{-n},\, t)\big|t \in \mathbb{G}_m\}$, computed in exercise 6.2.1 (2).

Theorem 6.2.5: Hodge Classes Are the Mumford-Tate Invariants

Let V be a rational Hodge structure of weight k and $M = \mathrm{MT}(V)$. Let M act on $T = T^{a,b,m}(V)$ through the algebraic representation

$$(g, \lambda) \longmapsto g^{\otimes a} \otimes \left((g^\vee)^{-1}\right)^{\otimes b} \otimes \lambda^{-m}$$

of $\mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m$, defined over $\mathbb{Q}$. Then

$$\left(T_\mathbb{Q}\right)^M = T_\mathbb{Q} \cap T^{0,0}$$

In particular, if $W = V^{\otimes a} \otimes (V^\vee)^{\otimes b}$ has even weight $2p$, then $\mathrm{Hdg}(W)$ corresponds under the identification $W(p)_\mathbb{Q} = W_\mathbb{Q} \otimes (2\pi i)^p$ to the space of M-invariant vectors of $T^{a,b,p}(V)$: the Hodge classes of the tensor constructions on V are exactly the Mumford-Tate invariants.

***Proof*:**

Step 1: the two actions of $\mathbb{S}$ on T agree. Composing the displayed representation with (h, Nm) gives a homomorphism $\mathbb{S} \to \mathrm{GL}(T_\mathbb{R})$. On the factor $V^{\otimes a}$ it is $h^{\otimes a}$, on $(V^\vee)^{\otimes b}$ it is $\left((h^\vee)^{-1}\right)^{\otimes b}$, and on $\mathbb{Q}(m)$ it is Nm^{-m}, which is the homomorphism $h_{\mathbb{Q}(m)}$ of example 6.2.3. By definition 3.1.9 these are precisely the homomorphisms attached by proposition 6.2.2 to the tensor product, the dual and the Tate structure, so the composite is h_T, the homomorphism of the Hodge structure T.

Step 2: an invariant is of bidegree $(0,0)$. Let $t \in T_\mathbb{Q}$ be fixed by M. Since $(h, \mathrm{Nm})(\mathbb{S}(\mathbb{R})) \subseteq M(\mathbb{R})$, Step 1 gives $h_T(z)t = t$ for every $z \in \mathbb{C}^\times$. Decomposing $t = \sum_{p,q} t^{p,q}$ into bihomogeneous components and using that $h_T(z)$ acts on $T^{p,q}$ by $z^p \bar{z}^q$,

$$\sum_{p,q} z^p \bar{z}^q \, t^{p,q} = \sum_{p,q} t^{p,q} \qquad \text{for all } z \in \mathbb{C}^\times$$

The components lie in independent summands, so $z^p \bar{z}^q \, t^{p,q} = t^{p,q}$ for every z and every (p,q). Suppose $t^{p,q} \neq 0$; then $z^p \bar{z}^q = 1$ for all $z \in \mathbb{C}^\times$. Taking $z = r$ real and positive gives $r^{p+q} = 1$ for all $r > 0$, hence $p+q = 0$; taking $z = e^{i\theta}$ gives $e^{i(p-q)\theta} = 1$ for all θ, hence $p = q$. The two together give $p = q = 0$. So every component with $(p,q) \neq (0,0)$ vanishes and $t = t^{0,0} \in T_\mathbb{Q} \cap T^{0,0}$.

Step 3: a rational class of bidegree $(0,0)$ is invariant. Let $t \in T_\mathbb{Q} \cap T^{0,0}$ and let

$$M_t = \{(g, \lambda) \in \mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m \,\big|\, (g, \lambda) \cdot t = t\}$$

be its stabilizer. The action is an algebraic representation defined over $\mathbb{Q}$ and t is a rational vector, so the conditions defining M_t are polynomial equations with rational coefficients: M_t is an algebraic subgroup defined over $\mathbb{Q}$ ([14, The Stabilizer is a Closed Subgroup]). Its real points contain $(h, \mathrm{Nm})(\mathbb{S}(\mathbb{R}))$, because by Step 1 the corresponding operator on $T_\mathbb{R}$ is $h_T(z)$, which acts on $T^{0,0}$ by $z^0 \bar{z}^0 = 1$. Definition 6.2.4 is a minimality statement, so $M \subseteq M_t$, that is t is fixed by M.

Step 4: the twisted form of the statement. Let $W = V^{\otimes a} \otimes (V^\vee)^{\otimes b}$ have weight $2p$. Then $T^{a,b,p}(V) = W(p)$ has weight 0, and $W(p)^{0,0} = W^{p,p}$ by definition 3.1.11, while $W(p)_\mathbb{Q} = W_\mathbb{Q} \otimes (2\pi i)^p$. Under this identification $W(p)_\mathbb{Q} \cap W(p)^{0,0}$ is $\mathrm{Hdg}(W)$, and Steps 2 and 3 identify it with the M-invariants of $T^{a,b,p}(V)$, as we sought to show.

The theorem is what the definition was arranged to produce. A Hodge class is defined by a transcendental condition, membership in a subspace of $V_{\mathbb{C}}$ cut out by the complex structure, together with a rationality condition; the theorem converts the pair into a single algebraic condition, invariance under a group defined over $\mathbb{Q}$. It also reverses the reading: a Hodge structure with many Hodge classes has a small Mumford-Tate group, since each independent class imposes an equation on M. The converse reading, that a structure whose only Hodge classes are the ones present for formal reasons has M as large as the polarization allows, is the harder direction: it is a form of Chevalley's theorem, that an algebraic subgroup is the stabilizer of a line in some tensor construction, and this book does not prove it. Nothing below uses it, and example 6.2.9 computes the generic case by a dimension count instead. This is the mechanism behind the loci studied in section 6.3: the Hodge classes jump exactly where the Mumford-Tate group drops.

The invariance in Step 3 is available only in bidegree $(0,0)$, and a Hodge class of a structure of weight $2p$ with $p \neq 0$ is scaled by $h(z)$, not fixed by it. The Tate factor $\mathbb{Q}(m)$ is exactly the bookkeeping that repairs this, and it is why the ambient group carries the second coordinate.

Nothing so far has used a polarization. A polarization constrains the group in two ways at once: it confines it to a similitude group, and it makes the representation on $V_{\mathbb{Q}}$ semisimple, which is what forces reductivity.

Proposition 6.2.6: The Mumford-Tate Group of a Polarized Structure

Let V be a rational Hodge structure of weight k and $M = \mathrm{MT}(V)$.

1. M is connected.
2. Suppose Q is a polarization of V (definition 3.2.1). Then M is contained in the group of similitudes
$$G_Q = \left\{(g,\lambda) \in \mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m \;\middle|\; Q(gu,gv) = \lambda^k\, Q(u,v) \text{ for all } u,v\right\}$$
and the M-stable $\mathbb{Q}$-subspaces of $V_{\mathbb{Q}}$ are exactly the *sub-Hodge structures* of V, that is the $\mathbb{Q}$-subspaces W for which $W_{\mathbb{C}} = \bigoplus_{p+q=k}(W_{\mathbb{C}} \cap V^{p,q})$. Each such subspace has an M-stable complement, so $V_{\mathbb{Q}}$ is a semisimple representation of M.
3. Under the hypothesis of (2), M is reductive.

***Proof*:**

1. Let $M^\circ \subseteq M$ be the identity component. It is an algebraic subgroup defined over $\mathbb{Q}$, and it is both Zariski open and Zariski closed in M, hence $M^\circ(\mathbb{R})$ is open and closed in $M(\mathbb{R})$ for the real topology. The set $(h,\mathrm{Nm})(\mathbb{S}(\mathbb{R}))$ is the continuous image of $\mathbb{C}^\times$, hence connected, and it contains the identity, so it lies in the connected component of the identity of $M(\mathbb{R})$ and therefore in $M^\circ(\mathbb{R})$. By the minimality in definition 6.2.4, $M \subseteq M^\circ$, so $M = M^\circ$ is connected. This argument uses no polarization.
2. *The similitude group.* That G_Q is an algebraic subgroup defined over $\mathbb{Q}$ is immediate, the defining conditions being polynomial identities with rational coefficients, and it is a subgroup because $Q(g_1g_2u, g_1g_2v) = \lambda_1^k\lambda_2^k Q(u,v)$. Its real points contain the image of (h,Nm): for $u \in V^{p,q}$ and $v \in V^{p',q'}$ the first Hodge-Riemann relation makes $Q(u,v)$

vanish unless $(p', q') = (q, p)$, and in that case

$$Q\big(h(z)u, h(z)v\big) = z^p \bar{z}^q \, z^q \bar{z}^p \, Q(u,v) = (z\bar{z})^{p+q} Q(u,v) = \mathrm{Nm}(z)^k \, Q(u,v)$$

When the bidegrees do not pair, both sides vanish, since $h(z)$ preserves each summand and so leaves bidegrees unchanged. The identity therefore holds on a spanning set of pairs and hence everywhere. By minimality $M \subseteq G_Q$.
Stable subspaces. Let $W \subseteq V_{\mathbb{Q}}$ be M-stable. Then $W_{\mathbb{R}}$ is stable under $h(\mathbb{S}(\mathbb{R}))$, so $W_{\mathbb{C}}$ is stable under the action of $\mathbb{S}_{\mathbb{C}}$ and therefore decomposes into character eigenspaces, $W_{\mathbb{C}} = \bigoplus_{p+q=k} \big(W_{\mathbb{C}} \cap V^{p,q}\big)$; and $W_{\mathbb{C}}$ is stable under conjugation, being defined over $\mathbb{Q}$, so the symmetry $\overline{W^{p,q}} = W^{q,p}$ is inherited. Thus W is a sub-Hodge structure. Conversely, let W be a sub-Hodge structure. The set of (g, λ) with $gW \subseteq W$ is an algebraic subgroup defined over $\mathbb{Q}$, and its real points contain the image of (h, Nm) because $h(z)$ preserves each $W^{p,q}$; minimality gives $M \subseteq$ that subgroup, so W is M-stable.
Complements. Let $W \subseteq V_{\mathbb{Q}}$ be M-stable, hence a sub-Hodge structure. The restriction $Q|_W$ satisfies both conditions of definition 3.2.1 for W, these being conditions on the summands $W^{p,q} \subseteq V^{p,q}$ inherited from V, so $Q|_W$ is a polarization of W. It is nondegenerate: the Weil operator C acts on $V^{p,q}$ by i^{p-q} and so preserves $W_{\mathbb{C}}$, and being defined over $\mathbb{R}$ it preserves $W_{\mathbb{R}}$; by part (3) of theorem 3.2.4 the form $S(u,v) = Q(Cu, v)$ is positive definite on $W_{\mathbb{R}}$, and $Q(u,v) = S(C^{-1}u, v)$, so $Q(u, W_{\mathbb{R}}) = 0$ forces $C^{-1}u = 0$ and hence $u = 0$ for $u \in W_{\mathbb{R}}$. Therefore $V_{\mathbb{Q}} = W \oplus W^{\perp}$ with $W^{\perp}$ the Q-orthogonal complement. That complement is M-stable: if $Q(W, x) = 0$ and $(g, \lambda) \in M \subseteq G_Q$, then for $w \in W$

$$Q(w, gx) = \lambda^k \, Q(g^{-1}w, x) = 0$$

since $g^{-1}w \in W$. Every M-stable subspace therefore has an M-stable complement, which is semisimplicity of $V_{\mathbb{Q}}$ as an M-representation.

3. The representation of M on $V_{\mathbb{Q}} \oplus \mathbb{Q}(1)$, with M acting on the second summand through $(g, \lambda) \mapsto \lambda^{-1}$, is faithful, since (g, λ) acting trivially on both summands forces $g = \mathrm{id}$ and $\lambda = 1$. It is semisimple: $V_{\mathbb{Q}}$ is semisimple by part (2) and $\mathbb{Q}(1)$ is one-dimensional, hence irreducible.
Let U be the unipotent radical of M, the largest connected normal unipotent subgroup, formed after extension of scalars to $\overline{\mathbb{Q}}$ ([14, Unipotent Radical (General), Reductive and Semisimple Groups]); by part (1) the group M is connected, so it is reductive precisely when U is trivial. Since U is canonically attached to $M_{\overline{\mathbb{Q}}}$ and M is defined over $\mathbb{Q}$, the subgroup U is stable under $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, hence so is the fixed space $\big((V_{\mathbb{Q}} \oplus \mathbb{Q}(1)) \otimes \overline{\mathbb{Q}}\big)^U$. A subspace of a base-changed space that is stable under the semilinear Galois action is spanned by its Galois-invariant vectors, so that fixed space is $N \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ for a unique $\mathbb{Q}$-subspace $N \subseteq V_{\mathbb{Q}} \oplus \mathbb{Q}(1)$, and N is M-stable because U is normal in M.
Suppose $N \neq V_{\mathbb{Q}} \oplus \mathbb{Q}(1)$. By semisimplicity there is a nonzero M-stable complement N', and $N \cap N' = 0$. A unipotent group has nonzero invariants in every nonzero representation ([14, Equivalent Characterizations of Unipotent Groups], applied over the algebraically closed field $\overline{\mathbb{Q}}$), so $(N' \otimes \overline{\mathbb{Q}})^U \neq 0$; but that space is contained in $N \otimes \overline{\mathbb{Q}}$, which meets $N' \otimes \overline{\mathbb{Q}}$ in 0. This contradiction gives $N = V_{\mathbb{Q}} \oplus \mathbb{Q}(1)$, so U acts trivially on a faithful representation of M and is therefore trivial. Hence M is reductive, completing the proof.

Reductivity is what makes the group usable. It places $\mathrm{MT}(V)$ among the groups classified by root

data after extension of scalars to $\mathbb{C}$,[2] so a polarized Hodge structure can be studied through the representation theory of a group whose structure is known, and the computation in example 6.2.9 below is exactly such an argument: it identifies the group by counting dimensions in the reductive structure theory. The polarization is what gives it. Part (3) rests on semisimplicity of $V_{\mathbb{Q}}$, and semisimplicity in turn rests on the Q-orthogonal complement of part (2); without a form there is no canonical complement to a sub-Hodge structure and the argument has no starting point.

The similitude factor in part (2) suggests separating the part of the group that preserves Q on the nose. That part is the group the classical literature calls the Hodge group, and it is what remains after the homotheties are removed.

Definition 6.2.7: The Hodge Group

Let V be a rational Hodge structure of weight k with $h\colon \mathbb{S} \to \mathrm{GL}(V_{\mathbb{R}})$, and let $\mathbb{U} = \ker(\mathrm{Nm}) \subseteq \mathbb{S}$ be the circle of definition 6.2.1. The *Hodge group*, or *special Mumford-Tate group*, $\mathrm{Hg}(V)$ is the smallest algebraic subgroup of $\mathrm{GL}(V_{\mathbb{Q}})$ defined over $\mathbb{Q}$ whose real points contain $h\big(\mathbb{U}(\mathbb{R})\big)$.

The group exists by the argument given after definition 6.2.4, and it is connected by the argument of proposition 6.2.6 part (1), the circle $\mathbb{U}(\mathbb{R})$ being connected. If Q polarizes V then $\mathrm{Hg}(V)$ preserves Q exactly rather than up to a factor: the computation in proposition 6.2.6 gives $Q(h(u)x, h(u)y) = \mathrm{Nm}(u)^k Q(x,y) = Q(x,y)$ for $u \in \mathbb{U}(\mathbb{R})$, so the automorphism group of $(V_{\mathbb{Q}}, Q)$ is one of the subgroups over which the minimum is taken.

Proposition 6.2.8: The Mumford-Tate Group as Hodge Group Times Homotheties

Let V be a rational Hodge structure of weight k, let $M = \mathrm{MT}(V)$ and let $\pi\colon \mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m \to \mathrm{GL}(V_{\mathbb{Q}})$ be the first projection. Then $\pi(M)$ is the smallest algebraic subgroup of $\mathrm{GL}(V_{\mathbb{Q}})$ defined over $\mathbb{Q}$ whose real points contain $h(\mathbb{S}(\mathbb{R}))$, and if $k \neq 0$ then

$$\pi(M) = \mathrm{Hg}(V) \cdot \mathbb{G}_m \,\mathrm{id}$$

the product inside $\mathrm{GL}(V_{\mathbb{Q}})$ of the Hodge group with the group of homotheties.

***Proof*:**

Step 1: $\pi(M)$ is the minimal group for h. The image of an algebraic group under a homomorphism of algebraic groups is an algebraic subgroup defined over the same field, so $\pi(M)$ is one; its real points contain $\pi\big((h,\mathrm{Nm})(\mathbb{S}(\mathbb{R}))\big) = h(\mathbb{S}(\mathbb{R}))$. Conversely let $N \subseteq \mathrm{GL}(V_{\mathbb{Q}})$ be an algebraic subgroup over $\mathbb{Q}$ with $h(\mathbb{S}(\mathbb{R})) \subseteq N(\mathbb{R})$. Then $N \times \mathbb{G}_m$ is an algebraic subgroup of $\mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m$ whose real points contain the image of (h, Nm), so $M \subseteq N \times \mathbb{G}_m$ by definition 6.2.4 and therefore $\pi(M) \subseteq N$.

Step 2: the homotheties lie in $\pi(M)$. For $r \in \mathbb{R}^\times$ one has $h(r) = r^k \,\mathrm{id}$ by proposition 6.2.2, so $\pi(M)(\mathbb{R})$ contains the set $\{r^k \,\mathrm{id} \,|\, r \in \mathbb{R}^\times\}$, which is infinite because $k \neq 0$. The homotheties form a subgroup of $\mathrm{GL}(V_{\mathbb{Q}})$ isomorphic to $\mathbb{G}_m$, and $\pi(M) \cap \mathbb{G}_m \,\mathrm{id}$ is an algebraic subgroup of it containing that infinite set. A proper algebraic subgroup of $\mathbb{G}_m$ is finite, being the zero set of a nonzero polynomial in one variable, so $\pi(M) \supseteq \mathbb{G}_m \,\mathrm{id}$.

Step 3: the two inclusions. The group $\mathrm{Hg}(V) \cdot \mathbb{G}_m \,\mathrm{id}$ is an algebraic subgroup defined over $\mathbb{Q}$, the

[2] [14, Classification of Split Reductive Groups].

homotheties being central. Every $z \in \mathbb{C}^\times$ factors as $z = ru$ with $r = |z| \in \mathbb{R}_{>0}$ and $u \in \mathbb{U}(\mathbb{R})$, so

$$h(z) = h(r)h(u) = r^k\, h(u) \in \big(\mathbb{G}_m\,\mathrm{id} \cdot \mathrm{Hg}(V)\big)(\mathbb{R})$$

By Step 1 this gives $\pi(M) \subseteq \mathrm{Hg}(V) \cdot \mathbb{G}_m\,\mathrm{id}$. In the other direction, $h(\mathbb{U}(\mathbb{R})) \subseteq h(\mathbb{S}(\mathbb{R})) \subseteq \pi(M)(\mathbb{R})$, so $\mathrm{Hg}(V) \subseteq \pi(M)$ by definition 6.2.7, and $\mathbb{G}_m\,\mathrm{id} \subseteq \pi(M)$ by Step 2. The two inclusions give the equality, as we sought to show.

The proposition says the two groups differ only by the homotheties, so either can be used and the choice is a matter of which bookkeeping is wanted. The Mumford-Tate group is the one adapted to Hodge classes in all weights, since it carries the Tate twist; the Hodge group is the one adapted to a fixed polarized structure, since it lands in the automorphism group of $(V_\mathbb{Q}, Q)$ and is therefore visibly a subgroup of a classical group.

The description just given also recovers the period domain itself. By proposition 6.2.2 a polarized Hodge structure on $V_\mathbb{R}$ with fixed numerical data is the same thing as a homomorphism $h\colon \mathbb{S} \to G_\mathbb{R}$ of the appropriate type, and proposition 3.3.5 presents D as a single $G_\mathbb{R}$-orbit with compact isotropy. Those two statements say the same thing twice: the orbit of a point $F \in D$ is the $G_\mathbb{R}$-conjugacy class of its h_F, and the isotropy group at F is the centralizer $Z_{G_\mathbb{R}}(h_F)$. So

$$D \;=\; \big\{\, g\, h_F\, g^{-1} \;:\; g \in G_\mathbb{R} \,\big\} \;\cong\; G_\mathbb{R}/Z_{G_\mathbb{R}}(h_F)$$

This is the form in which period domains generalize: replacing $G_\mathbb{R}$ by a subgroup cuts out a subdomain. If $M \subseteq \mathrm{GL}(V_\mathbb{Q}) \times \mathbb{G}_m$ is a $\mathbb{Q}$-algebraic subgroup containing the image of some h_F, then $\mathrm{Lie}(M)_\mathbb{Q}$ is a sub-Hodge structure of $\mathfrak{g}_\mathbb{Q}$ by proposition 6.1.1, and the orbit

$$D(M) \;=\; M(\mathbb{R}) \cdot F \;\subseteq\; D$$

is a *Mumford-Tate subdomain*: a homogeneous complex submanifold of D carrying the restricted Hodge metric, on which theorem 6.1.4 applies verbatim. Taking $M = \mathrm{MT}(V_F)$ gives the smallest one containing F. It is this construction, with $G_\mathbb{R}$ replaced by the real points of a reductive $\mathbb{Q}$-group and D by such a subdomain, that Shimura data axiomatize; see also [13], and Deligne's *Hodge Cycles on Abelian Varieties* [17] for the Hodge-theoretic side.

The smallest interesting case is weight one and rank two, where the whole group can be computed and the answer separates two classes of elliptic curves.

Example 6.2.9: The Mumford-Tate Group of an Elliptic Curve

Let $E_\tau = \mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}\tau)$ be the elliptic curve of example 3.4.3 and let V be the weight-one Hodge structure on $V_\mathbb{Q} = H^1(E_\tau, \mathbb{Q})$, polarized by the cup form Q as computed there. Here $\dim_\mathbb{Q} V_\mathbb{Q} = 2$ and $k = 1$.

The ambient group collapses. A nonzero alternating form on a two-dimensional space satisfies $Q(gu, gv) = \det(g)\, Q(u, v)$, so the similitude group of proposition 6.2.6 is

$$G_Q = \{(g, \lambda) \,\big|\, \lambda = \det g\} \cong \mathrm{GL}(V_\mathbb{Q})$$

Since $M := \mathrm{MT}(V) \subseteq G_Q$, the projection π of proposition 6.2.8 is injective on M, and M may be identified with the subgroup $M' = \pi(M) \subseteq \mathrm{GL}(V_\mathbb{Q})$, which by proposition 6.2.8 contains the homotheties.

The endomorphism algebra. By theorem 6.2.5 applied to $\mathrm{End}(V) = V \otimes V^\vee = T^{1,1,0}(V)$, a

structure of weight 0 on which (g, λ) acts by $f \mapsto gfg^{-1}$,

$$\mathrm{Hdg}\big(\mathrm{End}(V)\big) = \mathrm{End}_M(V_{\mathbb{Q}}) = \{f \in \mathrm{End}_{\mathbb{Q}}(V_{\mathbb{Q}}) \big| fg = gf \text{ for all } g \in M'\}$$

while the left-hand side is the algebra of endomorphisms of the Hodge structure V, since a rational endomorphism of bidegree $(0,0)$ is exactly one preserving every $V^{p,q}$ (definition 3.1.7). That algebra is the opposite of the endomorphism algebra of the weight-one structure Λ on $H_1(E_\tau, \mathbb{Q})$. The passage from one to the other is duality followed by a Tate twist and a conjugation,

$$H^1(E_\tau, \mathbb{Q}) \cong \overline{\Lambda^\vee(-1)}$$

where $\overline{W}$ denotes the Hodge structure carried by $W_{\mathbb{Q}}$ with $\overline{W}^{p,q} = W^{q,p}$. The conjugation is not optional: under the period identification $H^1(E_\tau, \mathbb{C}) = \mathrm{Hom}_{\mathbb{R}}\big(H_1(E_\tau, \mathbb{R}), \mathbb{C}\big)$ the generator dz of $H^{1,0}$ is the identity map of $H_1(E_\tau, \mathbb{R}) = \mathbb{C}$, which is $\mathbb{C}$-linear and so annihilates the $(-i)$-eigenspace $\Lambda^{0,1}$ of multiplication by i; hence $H^{1,0}$ is the summand of bidegree $(-1,0)$ of $\Lambda^\vee$ by definition 3.1.9, and a Tate twist, which shifts p and q equally, can never move it to bidegree $(1,0)$. Neither the twist nor the conjugation changes which $\mathbb{Q}$-linear maps are morphisms of Hodge structures, and it is the dual alone that reverses the order of composition; the opposite is in any case irrelevant here because both possible answers are commutative. By theorem 3.4.1 the endomorphisms of Λ are the holomorphic endomorphisms of E_τ, so the algebra is $\mathrm{End}(E_\tau) \otimes_{\mathbb{Z}} \mathbb{Q}$, computed in exercise 3.4.1 (3) to be $\mathbb{Q}$ unless τ satisfies a quadratic equation over $\mathbb{Q}$, in which case it is the imaginary quadratic field $K = \mathbb{Q}(\tau)$.

The generic case. Suppose $\mathrm{End}(E_\tau) \otimes \mathbb{Q} = \mathbb{Q}$, so $\mathrm{End}_M(V_{\mathbb{Q}}) = \mathbb{Q}$. If M' were commutative then every element of $M'(\overline{\mathbb{Q}})$ would lie in the commutant of $M'_{\overline{\mathbb{Q}}}$, which is $\mathrm{End}_M(V_{\mathbb{Q}}) \otimes \overline{\mathbb{Q}} = \overline{\mathbb{Q}}$ because the commutant is the solution space of a linear system and so commutes with extension of scalars; then M' would consist of homotheties, contradicting that $h(i)$ has the two distinct eigenvalues i and $-i$ (example 6.2.3). So M' is not commutative and its derived subgroup $[M', M']$ is nontrivial. By proposition 6.2.6 the group M' is connected and reductive, so the structure theory of reductive groups ([14, Structure of Reductive Groups]) makes $[M', M']$ semisimple and $R(M') \cap [M', M']$ finite, whence

$$\dim M' \geq \dim R(M') + \dim [M', M']$$

Now the radical of a connected reductive group is the identity component of its center, which contains the homotheties, so $\dim R(M') \geq 1$, and a nontrivial connected semisimple group has dimension at least 3, the minimum being attained by the rank-one groups SL_2 and PGL_2. This follows from the classification over $\mathbb{C}$, since extension of scalars to $\mathbb{C}$ leaves dimensions unchanged.[a] Hence $\dim M' \geq 4 = \dim \mathrm{GL}(V_{\mathbb{Q}})$, and a connected algebraic subgroup of full dimension in the connected group $\mathrm{GL}(V_{\mathbb{Q}})$ is the whole group:

$$\mathrm{MT}(V) \cong \mathrm{GL}(V_{\mathbb{Q}}) \cong \mathrm{GL}_2$$

The complex multiplication case. Suppose instead $\mathrm{End}(E_\tau) \otimes \mathbb{Q} = K$, an imaginary quadratic field. Then $V_{\mathbb{Q}}$ is a K-vector space of dimension 1, so the centralizer of K inside $\mathrm{End}_{\mathbb{Q}}(V_{\mathbb{Q}})$ is K itself, and the centralizer of $K^\times$ inside $\mathrm{GL}(V_{\mathbb{Q}})$ is the two-dimensional $\mathbb{Q}$-torus

$$T_K = \mathrm{Res}_{K/\mathbb{Q}}\mathbb{G}_m, \qquad T_K(\mathbb{Q}) = K^\times$$

Every element of $K = \mathrm{End}_{\mathrm{HS}}(V)$ commutes with $h(z)$, since a morphism of Hodge structures preserves each $V^{p,q}$ and $h(z)$ is a scalar there; so $h(\mathbb{S}(\mathbb{R})) \subseteq T_K(\mathbb{R})$ and hence $M' \subseteq T_K$ by

minimality. In fact $M' = T_K$. The map h is injective, its value at z having eigenvalues z and $\bar{z}$, so $h(\mathbb{S}(\mathbb{R}))$ is a two-dimensional subgroup of $T_K(\mathbb{R}) = (K \otimes_{\mathbb{Q}} \mathbb{R})^\times \cong \mathbb{C}^\times$, which is connected of real dimension 2; a subgroup of full dimension is open, and an open subgroup of a connected topological group is the whole group, so $h(\mathbb{S}(\mathbb{R})) = T_K(\mathbb{R})$. Any algebraic subgroup $N \subseteq T_K$ over $\mathbb{Q}$ with $N(\mathbb{R}) \supseteq T_K(\mathbb{R})$ has $\dim N \geq \dim_{\mathbb{R}} N(\mathbb{R}) \geq 2$, while $N \subseteq T_K$ gives $\dim N \leq \dim T_K = 2$; so its identity component is a closed connected subgroup of the connected group T_K of the same dimension and therefore equals T_K, whence $N = T_K$. Therefore

$$\mathrm{MT}(V) \cong T_K = \mathrm{Res}_{K/\mathbb{Q}}\mathbb{G}_m$$

a torus of dimension 2.

Reading the two answers. The Hodge classes count the difference. In the generic case $\mathrm{Hdg}(\mathrm{End}V) = \mathbb{Q}$ is one-dimensional and the group is all of GL_2, of dimension 4; in the complex multiplication case $\mathrm{Hdg}(\mathrm{End}V) = K$ is two-dimensional and the group has collapsed to a torus of dimension 2. The extra Hodge class, the endomorphism given by complex multiplication, is precisely the equation that cuts GL_2 down to its maximal torus T_K, which is theorem 6.2.5 read from right to left.

[a] [14, Classification of Split Reductive Groups].

Exercise 6.2.1

1. Let V be a rational Hodge structure of weight k with associated $h\colon \mathbb{S} \to \mathrm{GL}(V_{\mathbb{R}})$. Show that $h(-1) = (-1)^k \,\mathrm{id}$ and that $h(i)$ is the Weil operator C of definition 3.2.1. Deduce that $C^2 = (-1)^k \,\mathrm{id}$ and that C lies in $\mathrm{MT}(V)(\mathbb{R})$ up to the second coordinate, that is $(C, 1) \in \mathrm{MT}(V)(\mathbb{R})$.

2. Compute $\mathrm{MT}\big(\mathbb{Q}(n)\big)$ for every $n \in \mathbb{Z}$, as a subgroup of $\mathrm{GL}_1 \times \mathbb{G}_m = \mathbb{G}_m \times \mathbb{G}_m$. Treat $n = 0$ separately and explain why the answer is not the trivial group.

3. Let $V \neq 0$ be a rational Hodge structure of weight k. Show that $\mathrm{MT}(V)$ contains the image of the homomorphism $\mathbb{G}_m \to \mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m$, $t \mapsto (t^k \,\mathrm{id},\, t^2)$, and that this image is a one-dimensional torus. Conclude that $\mathrm{MT}(V)$ is never trivial.

4. Let X be a projective K3 surface and V the weight-two Hodge structure on $H^2(X, \mathbb{Q})$ of example 3.4.4. Show that $\mathrm{Hdg}(V) = \mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$, of dimension $\rho(X)$, and identify this space as the fixed space of $\mathrm{MT}(V)$ acting on $V(1)$. Deduce that the Picard number is determined by the Mumford-Tate group together with the action of the group on V.

5. For the elliptic curve E_τ of example 6.2.9, compute $\mathrm{Hg}(V)$ in both cases: show that it is $\mathrm{SL}(V_{\mathbb{Q}})$ when E_τ has no complex multiplication, and the norm-one torus $T_K^1 = \ker\big(\mathrm{Nm}_{K/\mathbb{Q}}\colon T_K \to \mathbb{G}_m\big)$ when it has complex multiplication by K. (Use proposition 6.2.8 and the fact that $\det h(u) = \mathrm{Nm}(u)$ for $u \in \mathbb{U}(\mathbb{R})$.)

6. Let V be a rational Hodge structure and $n \in \mathbb{Z}$, and let α be the automorphism of $\mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m$ given by $\alpha(g, \lambda) = (\lambda^{-n} g,\ \lambda)$. Show that $\mathrm{MT}\big(V(n)\big) = \alpha\big(\mathrm{MT}(V)\big)$ under the canonical identification $\mathrm{GL}\big(V(n)_{\mathbb{Q}}\big) = \mathrm{GL}(V_{\mathbb{Q}})$; in particular the two groups are isomorphic, so the Mumford-Tate group is insensitive to Tate twists up to isomorphism.

6.3 Hodge Loci and Algebraicity

Section 6.2 read the Hodge classes of a single structure off a group. In a family they are not a fixed set. A rational class of type (p,p) on one fibre transports flatly to a rational class on every nearby fibre, but the type is not carried along, and the set of parameters at which it survives is the object of this section.

Throughout, $f\colon \mathcal{X} \to S$ is a smooth projective family (definition 5.1.1) with fibres X_s, and $\mathbb{H}^k$, $\mathcal{H}^k$, ∇, $F^\bullet\mathcal{H}^k$ are the local system, the cohomology bundle, the Gauss-Manin connection and the Hodge bundles attached to f in section 5.1 and section 5.2. The degree is even, $k = 2p$, and it has to be: conjugation fixes a real class and exchanges $V^{a,b}$ with $V^{b,a}$ (definition 3.1.1), so a nonzero real class of pure type (a,b) has $a = b$.

A rational class is real, and reality collapses the two ways of asking that it be of type (p,p) into one. If $\mu \in H^{2p}(X_s,\mathbb{Q})$ lies in $F^pH^{2p}(X_s,\mathbb{C})$ then $\mu = \bar{\mu}$ lies in $\overline{F^p}$ as well, so $\mu \in F^p \cap \overline{F^p} = H^{p,p}(X_s)$ by the recovery formula (3.2); the converse inclusion $H^{p,p} \subseteq F^p$ is immediate from definition 3.1.5. So for a rational class the two conditions agree, and either of them says that μ is a Hodge class of $H^{2p}(X_s,\mathbb{Q})$ in the sense of definition 3.1.10. The condition of interest is therefore a condition on the position of μ in the *filtration*, which is the holomorphic datum, and not on its position in the bigrading, which is not.

Definition 6.3.1: The Hodge Locus

Let $f\colon \mathcal{X} \to S$ be a smooth projective family, let $U \subseteq S$ be open, connected and simply connected, and let λ be a nonzero flat section of $\mathbb{H}^{2p}$ over U. The *Hodge locus* of λ is

$$\mathrm{NL}(\lambda) = \left\{\, s \in U \;\middle|\; \lambda_s \in F^pH^{2p}(X_s,\mathbb{C}) \,\right\}$$

the set of parameters at which λ_s is a Hodge class (definition 3.1.10). For $p = 1$ it is also called the *Noether-Lefschetz locus* of λ.

Three features of the definition are worth separating. The section λ is flat, so as a class it does not move at all: what moves is the filtration, and the locus records where the moving filtration catches a fixed vector. Simple connectivity of U is what makes λ exist, since a flat section over a base with monodromy need not be single valued (proposition 5.1.5); the locus is therefore local by construction, and assembling a global object from the local ones is done in corollary 6.3.4. And rationality of λ is not decoration. Without it the condition $\lambda_s \in F^p$ says nothing about the bigrading, and the whole subject would be the study of one subbundle meeting one line.

Everything below applies verbatim with $\mathbb{H}^{2p}$ replaced by its primitive part $\mathbb{H}^{2p}_{\mathrm{prim}}$ of definition 5.3.1 and $F^\bullet\mathcal{H}^{2p}$ by the primitive filtration $F^\bullet_{\mathrm{prim}}$, part (3) of proposition 5.3.2 giving there the holomorphy and the transversality that theorem 5.2.6 and theorem 5.2.7 give for the full bundle. That is the version example 6.3.8 uses, and in it the locus is a condition on the period map. Fixing a path class as in definition 5.3.3 trivializes $\mathbb{H}^{2p}_{\mathrm{prim}}$ over U, carrying λ to a fixed rational vector v of the reference cohomology $H^{2p}_{\mathrm{prim}}(X_{s_0},\mathbb{Q})$ and s to the point $\Phi_{[c]}(s)$ of the period domain D; then $\mathrm{NL}(\lambda)$ is the preimage under $\Phi_{[c]}$ of the subset $\{\, F^\bullet \in D \,|\, v \in F^p \,\}$.

The definition permits the degenerate answer, and it is worth seeing which families force it before seeing which do not.

Example 6.3.2: Classes That Are Hodge at Every Parameter
Two mechanisms make $\mathrm{NL}(\lambda)$ all of U.

The polarization class. Let $h \in H^2(\mathcal{X}, \mathbb{Z})$ be the relative hyperplane class of section 5.3, with restrictions $h_s = h|_{X_s}$, and fix $p \leq n = \dim_{\mathbb{C}} X_s$. Part (1) of proposition 5.3.2 shows that parallel transport carries h_{s_0} to h_s and commutes with cup product, so $s \mapsto h_s^p$ is a flat section of $\mathbb{H}^{2p}$. It is a Hodge class at every s: the class h_s is that of the Fubini-Study Kähler form of X_s (theorem 1.1.7), hence real of type $(1,1)$, and a product of p classes of type $(1,1)$ has type (p,p). It is nonzero for $p \leq n$, since $\int_{X_s} h_s^n > 0$. So $\mathrm{NL}(h^p) = U$ for every U.

Absence of room. Take $p = 1$ and let $X \subsetneq \mathbb{CP}^3$ be a smooth surface of degree d, so $n = 2$. The class h_X is of type $(1,1)$, so $H^{2,0}(X) = H^{2,0}_{\mathrm{prim}}(X)$, and by theorem 5.4.11 the latter is isomorphic to R_{d-4}, the degree-$(d-4)$ piece of the Jacobian ring of definition 5.4.1. The Jacobian ideal is generated in degree $d-1$, so $R_e = \mathbb{C}[x]_e$ for $e \leq d-2$, and $\mathbb{C}[x]_e = 0$ for $e < 0$; hence

$$h^{2,0}(X) = \dim_{\mathbb{C}} \mathbb{C}[x]_{d-4} = \binom{d-1}{3}$$

which vanishes exactly when $d \leq 3$. For those degrees $F^1\mathcal{H}^2 = \mathcal{H}^2$ by definition 5.2.1, so *every* rational class on every fibre is a Hodge class and every Hodge locus is all of U. Concretely $b_2 = 1 + h^{1,1}_{\mathrm{prim}} = 1 + \dim R_{2d-4}$ by corollary 5.4.12, the 1 counting h_X, which gives $b_2 = 2$ for a quadric and $b_2 = 7$ for a cubic surface; the Néron-Severi group of definition 2.3.3 is all of $H^2(X, \mathbb{Z})$ in both cases, so the Picard number equals b_2. The first degree with $h^{2,0} \neq 0$ is $d = 4$, where $h^{2,0} = 1$.

The mechanism behind the second case is the one that governs the general situation. The condition $\lambda_s \in F^p$ is the vanishing of the image of λ_s in the quotient $\mathcal{H}^{2p}/F^p\mathcal{H}^{2p}$, and that quotient is a holomorphic bundle whose rank counts exactly the room the class has to fail. A vanishing condition on a holomorphic section is an analytic condition, and its derivative is computed by the connection.

Proposition 6.3.3: The Hodge Locus Is a Closed Analytic Subset

Let $f\colon \mathcal{X} \to S$ be a smooth projective family, fix $p \geq 1$, let $U \subseteq S$ be open, connected and simply connected, and let λ be a nonzero flat section of $\mathbb{H}^{2p}$ over U. Write

$$\mathcal{Q} = \mathcal{H}^{2p}/F^p\mathcal{H}^{2p}, \qquad c = \operatorname{rank}\mathcal{Q} = \sum_{r<p} h^{r,\,2p-r}$$

and let $\bar{\lambda}$ be the image of λ under the quotient map $\pi\colon \mathcal{H}^{2p} \to \mathcal{Q}$.

1. $\bar{\lambda}$ is a holomorphic section of $\mathcal{Q}$ and $\mathrm{NL}(\lambda) = \{\, s \in U \,\big|\, \bar{\lambda}(s) = 0 \,\}$. In particular $\mathrm{NL}(\lambda)$ is a closed analytic subset of U, cut out on a neighbourhood of each of its points by c holomorphic functions.

2. Let $s_0 \in \mathrm{NL}(\lambda)$, so that λ_{s_0} defines a class in $\mathrm{gr}_F^p\mathcal{H}^{2p}_{s_0} = H^{p,p}(X_{s_0})$. Then the differential of $\bar{\lambda}$ at s_0 is
$$d\bar{\lambda}_{s_0}\colon T_{s_0}S \to \mathcal{Q}_{s_0}, \qquad d\bar{\lambda}_{s_0}(\xi) = -\,\bar{\nabla}^p_{\xi}(\lambda_{s_0})$$
with $\bar{\nabla}^p$ the graded derivative of corollary 5.2.8. Its image lies in the subspace $H^{p-1,p+1}(X_{s_0}) = F^{p-1}_{s_0}/F^p_{s_0}$ of $\mathcal{Q}_{s_0}$. Consequently every holomorphic map germ $\gamma\colon (\mathbb{D},0) \to (U,s_0)$ from the unit disc with image contained in $\mathrm{NL}(\lambda)$ satisfies $\bar{\nabla}^p_{\gamma'(0)}(\lambda_{s_0}) = 0$.

3. Suppose $p = 1$ and that $\bar{\nabla}^1(\lambda_{s_0})\colon T_{s_0}S \to H^{0,2}(X_{s_0})$ is surjective. Then $\mathrm{NL}(\lambda)$ is, on a neighbourhood of s_0, a closed complex submanifold of U of codimension $h^{0,2}(X_{s_0})$.

***Proof*:**
One preliminary, the differential of a holomorphic section at a zero. Let E be a holomorphic vector bundle on U and μ a holomorphic section with $\mu(s_0) = 0$. In a local holomorphic frame $f_1,\dots,f_m$ write $\mu = \sum_j h_j f_j$ with $h_j(s_0) = 0$. For any connection ∇^E on E the Leibniz rule gives, at the point s_0,

$$\big(\nabla^E_\xi \mu\big)(s_0) = \sum_j (\partial_\xi h_j)(s_0)\, f_j(s_0) + \sum_j h_j(s_0)\,\big(\nabla^E_\xi f_j\big)(s_0) = \sum_j (\partial_\xi h_j)(s_0)\, f_j(s_0)$$

so the value of $\nabla^E_\xi\mu$ at s_0 is independent of the connection and of the frame. Write $d\mu_{s_0}\colon T_{s_0}S \to E_{s_0}$ for this map. If $\rho\colon E \to E'$ is a holomorphic bundle map then $\rho\circ\mu$ again vanishes at s_0 and, expanding $\rho\circ\mu = \sum_j h_j\,\rho(f_j)$ in the same way, $d(\rho\circ\mu)_{s_0} = \rho_{s_0}\circ d\mu_{s_0}$.

1. *The quotient bundle and the section.* By theorem 5.2.6 the subsheaf $F^p\mathcal{H}^{2p}$ is a holomorphic subbundle of $\mathcal{H}^{2p}$, of rank $f^p = \sum_{r\geq p} h^{r,2p-r}$; so $\mathcal{Q}$ is a holomorphic vector bundle and π a holomorphic bundle map [5, Subbundle and Quotient Bundle], of rank $c = \dim_{\mathbb{C}} H^{2p}(X_s,\mathbb{C}) - f^p = \sum_{r<p} h^{r,2p-r}$.

   The section λ is flat, and by theorem 5.1.7 a local frame of flat sections of $\mathcal{H}^{2p}$ is a local holomorphic frame; a flat section is a constant combination of such a frame, hence holomorphic. Therefore $\bar{\lambda} = \pi\circ\lambda$ is a holomorphic section of $\mathcal{Q}$, and $\bar{\lambda}(s) = 0$ says exactly $\lambda_s \in F^pH^{2p}(X_s,\mathbb{C})$, which is the defining condition of definition 6.3.1.

For the last clause, fix s_0 and a local holomorphic frame $e_1, \dots, e_c$ of $\mathcal{Q}$ near s_0, and write $\bar\lambda = \sum_i g_i e_i$ with g_i holomorphic. Then $\mathrm{NL}(\lambda)$ is the common zero locus of $g_1, \dots, g_c$ on that neighbourhood.

2. *Computing $d\bar\lambda_{s_0}$.* Let $s_0 \in \mathrm{NL}(\lambda)$, so $\lambda_{s_0} \in F^p_{s_0}$. Since $F^p\mathcal{H}^{2p}$ is a holomorphic subbundle, a local holomorphic frame of it near s_0 produces a local holomorphic section σ of $F^p\mathcal{H}^{2p}$ with $\sigma(s_0) = \lambda_{s_0}$. Put $\mu = \lambda - \sigma$, a holomorphic section of $\mathcal{H}^{2p}$ with $\mu(s_0) = 0$, and note $\pi \circ \mu = \bar\lambda$ because $\pi \circ \sigma = 0$. By the preliminary,

$$d\bar\lambda_{s_0}(\xi) = \pi_{s_0}\big(d\mu_{s_0}(\xi)\big) = \pi_{s_0}\big(\nabla_\xi \mu\big) = \pi_{s_0}\big(\nabla_\xi \lambda - \nabla_\xi \sigma\big) = -\,\pi_{s_0}\big(\nabla_\xi \sigma\big)$$

the connection used being the Gauss-Manin connection and the last equality holding because λ is flat.

The right-hand side is exactly the graded derivative. By theorem 5.2.7 the class $\nabla_\xi\sigma$ lies in $F^{p-1}_{s_0}$, so $\pi_{s_0}(\nabla_\xi\sigma)$ lies in the subspace $F^{p-1}_{s_0}/F^p_{s_0}$ of $\mathcal{Q}_{s_0}$, which is $H^{p-1,p+1}(X_{s_0})$ by definition 3.1.5; and corollary 5.2.8 constructs $\bar\nabla^p$ as the map induced by $\sigma \mapsto \nabla\sigma \bmod F^p\mathcal{H}^{2p}$, which depends on σ only through the class of $\sigma(s_0)$ in $\mathrm{gr}^p_F\mathcal{H}^{2p}_{s_0}$. Hence $\pi_{s_0}(\nabla_\xi\sigma) = \bar\nabla^p_\xi(\lambda_{s_0})$, as claimed.

If now $\gamma\colon (\mathbb{D}, 0) \to (U, s_0)$ is holomorphic with image in $\mathrm{NL}(\lambda)$, then $\bar\lambda \circ \gamma$ vanishes identically, so differentiating at 0 gives $d\bar\lambda_{s_0}(\gamma'(0)) = 0$, which is the asserted vanishing.

3. *The submersive case.* Let $p = 1$. The Hodge structure on $H^2(X_{s_0}, \mathbb{C})$ is effective, so $F^0\mathcal{H}^2 = \mathcal{H}^2$ (definition 5.2.1) and therefore

$$\mathcal{Q}_{s_0} = H^2(X_{s_0}, \mathbb{C})/F^1 = F^0_{s_0}/F^1_{s_0} = H^{0,2}(X_{s_0})$$

so that $c = h^{0,2}(X_{s_0})$ and the inclusion of part (2) is an equality of spaces. Surjectivity of $\bar\nabla^1(\lambda_{s_0})$ is therefore surjectivity of $d\bar\lambda_{s_0}$. In the frame of part (1) this says that the differentials $dg_1, \dots, dg_c$ are linearly independent at s_0, so the holomorphic implicit function theorem presents $\{g_1 = \cdots = g_c = 0\}$ near s_0 as a closed complex submanifold of codimension $c = h^{0,2}(X_{s_0})$, completing the proof.

Part (2) is what transversality contributes. Without it, all that could be said of the defining section $\bar\lambda$ is that it takes values in a bundle of rank c; with it, the derivative of $\bar\lambda$ at a point of the locus lands in the single graded piece $H^{p-1,p+1}$. Choosing the frame of part (1) so that $e_1, \dots, e_{h^{p-1,p+1}}$ span that piece, the remaining $c - h^{p-1,p+1}$ defining equations have vanishing differential at s_0: they vanish to order at least two there, and only $h^{p-1,p+1}$ of the c equations can be independent to first order. For $p \geq 2$ this makes $\bar\lambda$ non-submersive whenever some $h^{r,2p-r}$ with $r < p-1$ is nonzero. For $p = 1$ there is no gap, since $H^{0,2}$ is the whole of $\mathcal{Q}$, and that is why weight two is the case where the codimension count of part (3) is available.

Part (2) also identifies the Zariski tangent directions concretely. In the graded description of corollary 5.2.8 the map $\xi \mapsto \bar\nabla^p_\xi(\lambda_{s_0})$ is a linear map $T_{s_0}S \to H^{p-1,p+1}(X_{s_0})$ attached to the fibre X_{s_0} and the class λ_{s_0} alone; the Hodge locus can move only in its kernel. When the family is one for which this map has been computed, the tangent space to the Hodge locus is computed with it, and example 6.3.8 carries that out.

A single flat section lives only over a simply connected open set, so the locus attached to it is a

local object. The set where *some* nonzero rational class becomes Hodge is global, and it need not be analytic in S; what the analytic theory gives is a countable union of sets, each analytic in some open subset of S, and over an arbitrary base that is the strongest statement available.

Corollary 6.3.4: The Locus of Extra Hodge Classes Is a Countable Union

Let $f\colon \mathcal{X} \to S$ be a smooth projective family with S connected and second countable, fix $p \geq 1$, and put

$$\mathrm{HL}^{2p}(S) = \left\{ s \in S \;\middle|\; H^{p,p}(X_s) \cap H^{2p}(X_s, \mathbb{Q}) \neq 0 \right\}$$

Then $\mathrm{HL}^{2p}(S)$ is a countable union of sets, each of which is a closed analytic subset of some open subset of S. If moreover no nonzero flat section of $\mathbb{H}^{2p}$ over a nonempty connected simply connected open $U \subseteq S$ is a Hodge class at every point of U, then each of those sets is nowhere dense in S and the complement $S \setminus \mathrm{HL}^{2p}(S)$ is dense.

***Proof*:**

Step 1: a countable family of loci. Since S is a second countable manifold it has a countable basis $U_1, U_2, \ldots$ of open sets each of which is a coordinate ball, hence connected and simply connected. Over each U_i the flat sections of $\mathbb{H}^{2p}$ form a $\mathbb{Q}$-vector space isomorphic to $H^{2p}(X_{s_i}, \mathbb{Q}) \cong \mathbb{Q}^m$ for any $s_i \in U_i$, by proposition 5.1.5; that is a countable set. So the collection of pairs (U_i, λ) with λ a nonzero flat section over U_i is countable, and each $\mathrm{NL}(\lambda) \subseteq U_i$ is a closed analytic subset of U_i by proposition 6.3.3.

These loci exhaust $\mathrm{HL}^{2p}(S)$. If $s \in \mathrm{HL}^{2p}(S)$, choose a nonzero Hodge class $\mu \in H^{2p}(X_s, \mathbb{Q})$ and a basis member $U_i \ni s$; transporting μ flatly over the simply connected U_i gives a nonzero flat section λ with $\lambda_s = \mu$, so $s \in \mathrm{NL}(\lambda)$. Conversely each $\mathrm{NL}(\lambda)$ consists of points carrying the nonzero Hodge class λ_s.

Step 2: each locus is nowhere dense. Assume the extra hypothesis and fix a pair (U_i, λ). Let $W \subseteq U_i$ be the set of points having a neighbourhood on which $\bar{\lambda}$ vanishes identically. Since $\mathrm{NL}(\lambda)$ is exactly the zero set of $\bar{\lambda}$, the set W is the interior of $\mathrm{NL}(\lambda)$, and in particular it is open. It is also closed in U_i: given s in its closure, choose a connected neighbourhood $N \ni s$ carrying a holomorphic chart and a holomorphic frame $e_1, \ldots, e_c$ of $\mathcal{Q}$, and write $\bar{\lambda}|_N = \sum_j g_j e_j$ with g_j holomorphic on N. Each g_j vanishes on the nonempty open set $W \cap N$, so each vanishes on all of N by the identity theorem,[a] whence $N \subseteq W$ and $s \in W$.

Since U_i is connected, W is either empty or all of U_i; the second case gives $\mathrm{NL}(\lambda) = U_i$, which the hypothesis excludes. Therefore $\mathrm{NL}(\lambda)$ has empty interior, and being closed in U_i as well it is nowhere dense in U_i.

Nowhere density passes to S. Write $A = \mathrm{NL}(\lambda)$ and suppose $B \subseteq \overline{A}$ is open and nonempty, the closure taken in S. If B met U_i then $B \cap U_i$ would be a nonempty open subset of $\overline{A} \cap U_i$, which is the closure of A in U_i and has empty interior there. So $B \subseteq S \setminus U_i$, while $B \subseteq \overline{A} \subseteq \overline{U_i}$, giving $B \subseteq \overline{U_i} \setminus U_i$; but that set has empty interior, since a nonempty open subset of it would be disjoint from U_i and yet meet it, U_i being open and the subset lying in its closure. Hence $\overline{A}$ has empty interior in S.

Step 3: density of the complement. A complex manifold is locally compact and Hausdorff, so the Baire category theorem in its locally compact Hausdorff form applies to S: a countable union

of nowhere dense subsets has dense complement. By Steps 1 and 2 the set $\mathrm{HL}^{2p}(S)$ is such a union, completing the proof.

[a][5, Properties of Holomorphic Functions in Several Variables].

The kind of conclusion at issue here has a standing name.

Definition 6.3.5: General and Very General Points

Let S be a complex manifold and let P be a property of the points of S. Say that P holds at a *general* point of S if the set where it fails is contained in a proper closed analytic subset of S, and that it holds at a *very general* point of S if that set is contained in a countable union of proper closed analytic subsets of S.

The two notions differ, and not pedantically: a countable union of proper analytic subsets can be dense, as the union of the lines $\{z_2 = qz_1\} \subseteq \mathbb{C}^2$ over the Gaussian rationals $q \in \mathbb{Q}(i)$ already shows, $\mathbb{Q}(i)$ being dense in $\mathbb{C}$. Corollary 6.3.4 reaches neither notion on its own. Its pieces are analytic in *some* open subset of S, and such a piece need not lie in a proper closed analytic subset of S: the set $\{\,1 - 1/n \,|\, n \geq 1\,\}$ is a closed analytic subset of the unit disc $\mathbb{D} \subseteq \mathbb{C}$, being the zero set there of a holomorphic function, and is nowhere dense in $\mathbb{C}$, while a proper closed analytic subset of $\mathbb{C}$ is discrete and closed in $\mathbb{C}$ and so contains no sequence converging to 1. What survives over an arbitrary base is that $\mathrm{HL}^{2p}(S)$ is meagre, so the property of carrying no nonzero rational Hodge class in degree $2p$ holds at a dense set of parameters and fails on a set with empty interior; getting from there to a very general statement takes theorem 6.3.6.

The hypothesis of the corollary is a real restriction, and for the full local system $\mathbb{H}^{2p}$ it fails whenever $p \leq n$, the powers h^p of the polarization class being Hodge at every parameter (example 6.3.2). What the corollary is applied to is therefore a sub-local system on which those classes have been removed, and the primitive part is the standard choice.

The corollary is the best that the analytic theory gives, and it leaves a gap that is visible from the shape of the statement. Each $\mathrm{NL}(\lambda)$ is analytic in an open subset of S, and S is frequently algebraic; an analytic subset of an algebraic variety need not be algebraic, since Chow's theorem needs properness and S is typically not proper (exercise 6.3.1 (6)). Closing that gap is a theorem, and it is the deepest input this chapter takes from outside.

Theorem 6.3.6: Algebraicity of the Hodge Locus (Cattani, Deligne, Kaplan)

Let S be a smooth complex quasi-projective variety and let $f\colon \mathcal{X} \to S$ be a smooth projective family of complex algebraic varieties. Fix $p \geq 1$, let $U \subseteq S$ be open in the analytic topology, connected and simply connected, and let λ be a nonzero flat section of $\mathbb{H}^{2p}$ over U. Then the closure in S of every irreducible component of $\mathrm{NL}(\lambda)$ is a closed algebraic subvariety of S. Consequently, if no nonzero flat section of $\mathbb{H}^{2p}$ over a nonempty connected simply connected open subset of S is a Hodge class at every point of that subset, then the locus $\mathrm{HL}^{2p}(S)$ of corollary 6.3.4 is contained in a countable union of *proper* closed algebraic subvarieties of S, so that a very general point of S (definition 6.3.5) lies outside it.

The second statement follows from the first. Under its hypothesis corollary 6.3.4 makes each $\mathrm{NL}(\lambda)$ nowhere dense in S, so the closure of each irreducible component of $\mathrm{NL}(\lambda)$ is a closed algebraic subvariety with empty interior and hence a proper one; there are countably many pairs (U_i, λ) in the

proof of that corollary and countably many components in each locus, S being second countable, so countably many closures suffice; each of them is a closed analytic subset of S, which is the form definition 6.3.5 asks for. The implication runs only this way. The algebraicity of the individual components is the substantive statement, and without the hypothesis the containment can fail while that statement still holds, as it does for the full local system with $p \leq n$, where $\mathrm{HL}^{2p}(S)$ is all of S (example 6.3.2).

The theorem is not proved here, and the reason is specific rather than a matter of length. What an algebraicity statement requires is control of the Hodge locus near the boundary of S inside a compactification: an analytic subset of S is algebraic precisely when it extends analytically across that boundary, and the components of $\mathrm{NL}(\lambda)$ do so because a variation of Hodge structure degenerates in a restricted way. The tools that make that precise are Schmid's nilpotent orbit theorem, which compares a degenerating variation near a puncture with the orbit of a nilpotent monodromy logarithm, and his SL_2-orbit theorem, which gives the asymptotics of the Hodge metric of section 6.1 in the several-variable case. Neither is developed in this book. The proof is in Cattani, Deligne and Kaplan, *On the locus of Hodge classes* [2], and it is the source of the statement above.

Both inputs belong to Schmid's analysis of what a horizontal map into a period domain may do near a boundary point [35], which is the same phenomenon that constrains such maps in theorem 6.1.7. What distinguishes the present conclusion is its category. Analyticity is a statement about S as a complex manifold and is available for any base; algebraicity is a statement about S as a variety, and it gives each Hodge locus a degree, a dimension and a field of definition.

The Hodge locus is also where the two threads of this chapter meet. The Mumford-Tate group of section 6.2 is attached to a single Hodge structure; in a family it varies, and theorem 6.2.5 says how: extra Hodge classes on the fibre are extra invariants of $\mathrm{MT}(V_s)$, so acquiring one shrinks the group.

Proposition 6.3.7: The Generic Mumford-Tate Group

Let $f\colon \mathcal{X} \to S$ be a smooth projective family with S connected and second countable, and fix k. There is a $\mathbb{Q}$-algebraic group M_{gen} and a subset $S^\circ \subseteq S$ whose complement is a countable union of proper closed analytic subsets, such that $\mathrm{MT}(H^k(X_s,\mathbb{Q})) = M_{\mathrm{gen}}$ for every $s \in S^\circ$, and $\mathrm{MT}(H^k(X_s,\mathbb{Q})) \subseteq M_{\mathrm{gen}}$ for every $s \in S$. A point of S° is called *Hodge-generic*.

***Proof*:**

Work on a connected simply connected open $U \subseteq S$, over which the local system $\mathbb{H}^k$ is trivial (proposition 5.1.4), and identify every fibre with a fixed $V_{\mathbb{Q}}$. By theorem 6.2.5 the group $\mathrm{MT}(V_s)$ is cut out inside $\mathrm{GL}(V_{\mathbb{Q}}) \times \mathbb{G}_m$ by the conditions of fixing each Hodge class of each tensor construction $T^{a,b,m}(V_s)$. Adding a Hodge class adds a condition, so $\mathrm{MT}(V_s)$ can only shrink as s moves onto a Hodge locus.

For each (a,b,m) and each λ in a countable basis of $T^{a,b,m}(V_{\mathbb{Q}})$, the locus where λ becomes a Hodge class is $\mathrm{NL}(\lambda)$, closed analytic in U by proposition 6.3.3. There are countably many such λ, since $V_{\mathbb{Q}}$ is a finite-dimensional $\mathbb{Q}$-vector space and there are countably many triples (a,b,m). Let $S^\circ \cap U$ be the complement of the union of those $\mathrm{NL}(\lambda)$ that are proper subsets of U. Off that union no unexpected Hodge class occurs, so $\mathrm{MT}(V_s)$ is the same group for all such s, namely the one cut out by the classes that are Hodge *everywhere* on U; and it contains every $\mathrm{MT}(V_s)$ by the shrinking just described. The sets $\mathrm{NL}(\lambda)$ are countable in number and each is nowhere dense or all of U, so the complement has the stated form. Finally the group is independent of U: two such opens are joined by a chain of overlapping ones, and on an overlap

the two generic groups agree because the overlap meets both generic sets.

So the curvature of section 6.1 constrains how a family can move inside D, and the Mumford-Tate group records where it has moved to: the image of the period map lies in the subdomain $D(M_{\mathrm{gen}})$ described in section 6.2, and the Hodge locus is exactly the set of parameters where it meets a smaller one. Theorem 6.3.6 then says those smaller loci are algebraic.

The classical instance is much older than the general theory, and it is the one where every ingredient of proposition 6.3.3 can be written down as polynomial algebra.

Example 6.3.8: The Noether-Lefschetz Locus for Surfaces in $\mathbb{C}\mathrm{P}^3$
Take $n = 2$ and $d \geq 4$, let

$$S = \left\{ [F] \in \mathbb{C}\mathrm{P}^M \;\middle|\; F \in \mathbb{C}[x_0, \dots, x_3]_d \text{ smooth} \right\}, \qquad M = \binom{d+3}{3} - 1$$

be the parameter space of proposition 5.4.4 and $f\colon \mathcal{X} \to S$ the universal family of smooth surfaces of degree d in $\mathbb{C}\mathrm{P}^3$. Then S is a connected smooth quasi-projective variety of dimension M, and f is a smooth projective family. Work with $p = 1$ and with the primitive part $\mathbb{H}^2_{\mathrm{prim}}$.

The numerical data. By (5.11) with $n = 2$ the three primitive Hodge pieces of a fibre $X = X_F$ are graded pieces of the Jacobian ring $R = R(F)$:

$$H^{2,0}(X) \cong R_{d-4}, \qquad H^{1,1}_{\mathrm{prim}}(X) \cong R_{2d-4}, \qquad H^{0,2}(X) \cong R_{3d-4}$$

As in example 6.3.2, $R_{d-4} = \mathbb{C}[x]_{d-4}$ and $p_g := h^{2,0}(X) = \binom{d-1}{3} \geq 1$ for $d \geq 4$. So the bundle $\mathcal{Q} = \mathcal{H}^2_{\mathrm{prim}}/F^1_{\mathrm{prim}}$ has rank p_g and proposition 6.3.3 cuts out every Hodge locus locally by p_g holomorphic equations.

Picard number and primitive Hodge classes. Since $\int_X h_X \smile h_X = d \neq 0$, the assignment $\alpha \mapsto \alpha - d^{-1}\big(\int_X \alpha \smile h_X\big) h_X$ is a projection defined over $\mathbb{Q}$, and

$$H^2(X, \mathbb{Q}) = \mathbb{Q}\, h_X \;\oplus\; H^2_{\mathrm{prim}}(X, \mathbb{Q})$$

a decomposition into Hodge substructures (proposition 3.2.7), with h_X of type $(1,1)$. The Néron-Severi group is $\mathrm{NS}(X) = H^{1,1}(X) \cap H^2(X, \mathbb{Z})$ (definition 2.3.3), and every rational class has an integral multiple, so its rank is $\rho(X) = \dim_{\mathbb{Q}}\big(H^{1,1}(X) \cap H^2(X, \mathbb{Q})\big)$. Splitting off h_X,

$$\rho(X) = 1 + \dim_{\mathbb{Q}}\big(H^{1,1}(X) \cap H^2_{\mathrm{prim}}(X, \mathbb{Q})\big)$$

Hence $\rho(X_s) > 1$ if and only if X_s carries a nonzero primitive rational Hodge class, and the *Noether-Lefschetz locus* $\mathrm{NL}_d = \{\, s \in S \,|\, \rho(X_s) > 1 \,\}$ is the set $\mathrm{HL}^2(S)$ of corollary 6.3.4 formed inside $\mathbb{H}^2_{\mathrm{prim}}$.

No class is Hodge everywhere. This is the hypothesis corollary 6.3.4 needs, and it is where the Jacobian ring does the work. Suppose λ is a nonzero flat section of $\mathbb{H}^2_{\mathrm{prim}}$ over a connected simply connected $U \subseteq S$ with $\mathrm{NL}(\lambda) = U$. Let $\pi\colon \tilde{S} \to S$ be the universal cover, a holomorphic covering, and pull f back along it; the result is again a smooth projective family, its local system and Hodge bundles being the pullbacks of those of f. Since U is simply connected, λ lifts to a flat section over a component $\tilde{U}$ of $\pi^{-1}(U)$, and since $\tilde{S}$ is connected and simply connected that section extends uniquely to a flat section $\tilde{\lambda}$ over $\tilde{S}$ (proposition 5.1.5). The associated holomorphic section $\overline{\tilde{\lambda}}$ of $\mathcal{H}^2_{\mathrm{prim}}/F^1_{\mathrm{prim}}$ vanishes on the nonempty open set $\tilde{U}$, hence vanishes identically on $\tilde{S}$ by the identity

theorem, exactly as in Step 2 of corollary 6.3.4. So $\tilde{\lambda}_t$ is a Hodge class for every $t \in \tilde{S}$, and part (2) of proposition 6.3.3 applied with $\mathrm{NL}(\tilde{\lambda}) = \tilde{S}$ gives

$$\bar{\nabla}^1_\xi(\tilde{\lambda}_t) = 0 \qquad \text{for every } t \in \tilde{S} \text{ and every } \xi \in T_t\tilde{S}$$

Evaluate at a point t_0 lying over the Fermat surface $F_0 = x_0^d + x_1^d + x_2^d + x_3^d$, which lies in S by example 5.4.3. Let $A \in R_{2d-4}$ correspond to $\tilde{\lambda}_{t_0} \in H^{1,1}_{\mathrm{prim}}(X_{F_0})$ under (5.11). Tangent vectors to S at $[F_0]$ are represented by arbitrary $G \in \mathbb{C}[x]_d$, and theorem 5.4.14 with $q = 1$ computes $\bar{\nabla}_G$ as $[A] \mapsto -2\,[GA]$ in R_{3d-4}. So the displayed vanishing says

$$A \cdot \mathbb{C}[x]_d = 0 \quad \text{in } R_{3d-4}$$

The monomial computation. By example 5.4.3 the ring $R(F_0)$ has basis the monomials x^a with $0 \le a_i \le d-2$ for every i, and its top degree is $\sigma = 4(d-2)$, which is at least $3d-4$ because $d \ge 4$. Suppose $A \neq 0$ and write $A = \sum_a c_a\, x^a$ over the basis monomials of degree $2d-4$; choose a^* with $c_{a^*} \neq 0$ and put $m_i = d-2-a^*_i \ge 0$, so that

$$\sum_i m_i = 4(d-2) - (2d-4) = 2d-4 \ge d$$

Choose integers $0 \le b_i \le m_i$ with $\sum_i b_i = d$, possible by the displayed inequality. Then $b_i \le d-2$, so x^b is a basis monomial of R_d, and in R the product $x^a x^b$ is either the basis monomial x^{a+b}, when $a_i + b_i \le d-2$ for every i, or zero. Distinct a give distinct $a+b$, and $a^*_i + b_i \le a^*_i + m_i = d-2$, so the coefficient of x^{a^*+b} in $A\,x^b$ is $c_{a^*} \neq 0$. Hence $A\,x^b \neq 0$ in R_{3d-4}, contradicting the vanishing above. Therefore $A = 0$, so $\tilde{\lambda}_{t_0} = 0$, so $\tilde{\lambda} = 0$ by flatness and $\lambda = 0$, contrary to the choice of λ.

The Noether-Lefschetz theorem. The hypothesis of corollary 6.3.4 therefore holds, and the corollary gives: for $d \ge 4$ the locus NL_d is a countable union of sets, each of them closed analytic in some open subset of S and nowhere dense in S, so that NL_d is meagre and its complement is dense. Since S is quasi-projective, theorem 6.3.6 applies as well: the closure of each irreducible component of each locus $\mathrm{NL}(\lambda)$ is a closed algebraic subvariety of S, proper because that locus is nowhere dense, and these closures are the classical Noether-Lefschetz components. They are countably many proper closed subvarieties whose union contains NL_d, so the very general (definition 6.3.5) surface of degree $d \ge 4$ in $\mathbb{CP}^3$ satisfies $\rho = 1$, that is $\mathrm{NS}(X) \otimes \mathbb{Q} = \mathbb{Q}\,h_X$; by theorem 2.3.2 the Néron-Severi group is the group of first Chern classes of line bundles, so every divisor class on such a surface is a rational multiple of a plane section.

Codimension. Where the map $\bar{\nabla}^1(\lambda_s)$ is surjective, part (3) of proposition 6.3.3 makes $\mathrm{NL}(\lambda)$ smooth of codimension p_g. For $d = 4$ one has $p_g = 1$ and $H^{0,2}(X) \cong R_8$ is a line, so surjectivity is the same as nonvanishing: the Noether-Lefschetz locus of a quartic surface is a smooth hypersurface near every parameter at which $\bar{\nabla}^1(\lambda_s) \neq 0$, and the monomial computation above produces such a parameter. That count is carried out explicitly for one class at the Fermat quartic in exercise 6.3.1 (5).

None of this asserts that a Hodge class is the class of an algebraic cycle. That assertion is the Hodge conjecture, and **??** takes it up; a proof of it would make each Hodge locus algebraic by a spreading-out argument, so theorem 6.3.6 is one of its predictions verified in advance.

Exercise 6.3.1

1. Let X be a compact Kähler manifold and $p \geq 0$.

   1. Show that a class $\mu \in H^{2p}(X, \mathbb{Q})$ lies in $F^p H^{2p}(X, \mathbb{C})$ if and only if it lies in $H^{p,p}(X)$, so that the two descriptions of a Hodge class agree, and that the Hodge classes (definition 3.1.10) of degree $2p$ form a $\mathbb{Q}$-subspace of $H^{2p}(X, \mathbb{Q})$.
   2. Show that the corresponding statement for a class in $H^{2p}(X, \mathbb{C})$ is false, by exhibiting a compact Kähler manifold and a class in $F^1 H^2(X, \mathbb{C})$ that is not of type $(1, 1)$.
   3. Let λ, μ be flat sections of $\mathbb{H}^{2p}$ over a connected simply connected U. Show that $\mathrm{NL}(\lambda) \cap \mathrm{NL}(\mu) \subseteq \mathrm{NL}(a\lambda + b\mu)$ for all $a, b \in \mathbb{Q}$ with $a\lambda + b\mu \neq 0$, and that $\mathrm{NL}(a\lambda) = \mathrm{NL}(\lambda)$ for $a \in \mathbb{Q}^\times$.

2. Let $f \colon \mathcal{X} \to S$ be a smooth projective family for which the graded derivatives $\bar{\nabla}^p$ of corollary 5.2.8 vanish in degree $2p$ for every p. Using exercise 5.2.1 (1), show that for every connected simply connected $U \subseteq S$ and every nonzero flat section λ of $\mathbb{H}^{2p}$ over U, the locus $\mathrm{NL}(\lambda)$ is either empty or all of U. Check the conclusion directly for the product family of exercise 5.2.1 (3).

3. Let $f \colon \mathcal{X} \to S$ be a smooth projective family and fix $p \geq 1$.

   1. Show that if $h^{r,2p-r}(X_s) = 0$ for every $r < p$, then every Hodge locus in degree $2p$ is all of U, and identify the rank c of proposition 6.3.3 in that case.
   2. For the family of smooth surfaces of degree d in $\mathbb{CP}^3$ with $p = 1$, determine exactly which d satisfy the hypothesis of (1), and compute $b_2(X)$ and $\rho(X)$ for those d.

4. Take $d = 5$ in example 6.3.8, so that the fibres are smooth quintic surfaces in $\mathbb{CP}^3$. Using corollary 5.4.12 and (5.4), compute p_g, $h^{1,1}_{\mathrm{prim}}$, $h^{1,1}$ and b_2 for such a surface, and compute $\dim_{\mathbb{C}} S$. How many holomorphic equations does proposition 6.3.3 use to cut out a Noether-Lefschetz locus here, and what codimension does part (3) predict where the differential is surjective?

5. Let $F_0 = x_0^4 + x_1^4 + x_2^4 + x_3^4$ be the Fermat quartic surface, with Jacobian ring $R = R(F_0)$ as in example 5.4.3. Suppose λ is a flat section of $\mathbb{H}^2_{\mathrm{prim}}$ defined near $[F_0]$ for which $\lambda_{[F_0]}$ is a Hodge class corresponding, up to a nonzero scalar, to $[x_0x_1x_2x_3] \in R_4$ under (5.11).

   1. Show that R_8 is one dimensional, spanned by $[x_0^2x_1^2x_2^2x_3^2]$, and determine every basis monomial x^b of R_4 with $[x_0x_1x_2x_3] \cdot [x^b] \neq 0$.
   2. Compute the dimension of the Zariski tangent space $\ker\big(\bar{\nabla}^1(\lambda_{[F_0]})\big) \subseteq T_{[F_0]}S$ and its codimension.
   3. Conclude that $\mathrm{NL}(\lambda)$ is a smooth hypersurface near $[F_0]$.

6. Let $Z = \{\, (z, w) \in \mathbb{C}^2 \,|\, w = e^z \,\}$. Show that Z is a closed analytic subset of $\mathbb{C}^2$ and that the only polynomial $P \in \mathbb{C}[z, w]$ vanishing identically on Z is $P = 0$; deduce that Z is not the common zero set of any family of polynomials. Explain in one sentence why this shows that theorem 6.3.6 does not follow formally from proposition 6.3.3.

6.4 Moduli of Polarized K3 Surfaces

Everything this chapter has built comes together on one class of surfaces. A K3 surface carries a weight-two Hodge structure whose entire content is a single line (example 3.4.4); a family of them has a period map (definition 5.3.3) which is horizontal for free (example 5.3.9); and the domain that map lands in is one of the classical bounded domains, so the horizontal restriction which limits theorem 6.1.4 is harmless here. What remains is to say which domain it is, and to record the theorem that turns the period map into a classification: the global Torelli theorem, which says that a K3 surface is determined by its period point.

Example 3.4.4 works inside $H^2(S, \mathbb{Z})$ for one surface S at a time. Comparing two surfaces means comparing two abstract lattices, and the comparison is only useful once a single model lattice has been fixed and each surface has been identified with it. The model is forced: by example 3.4.4 the cup form makes $H^2(S, \mathbb{Z})$ an even unimodular lattice of signature $(3, 19)$, and that isometry class has a name.

Definition 6.4.1: The K3 Lattice, Markings and Polarizations

Write $\langle -,- \rangle$ for the bilinear form of a lattice, extended $\mathbb{C}$-bilinearly whenever the lattice is complexified.

1. Let U be the free abelian group on e, f with $\langle e,e\rangle = \langle f,f\rangle = 0$ and $\langle e,f\rangle = 1$, and let E_8 be the root lattice of type E_8, realized as

$$E_8 = \left\{ x \in \mathbb{Z}^8 \cup \left(\mathbb{Z}+\tfrac{1}{2}\right)^8 \;\middle|\; \textstyle\sum_{i=1}^8 x_i \in 2\mathbb{Z} \right\}$$

   with the restriction of the standard positive definite form of $\mathbb{R}^8$; it is the root lattice of type E_8,[a] and it is positive definite, even and unimodular of rank 8. Evenness is immediate: a vector of $\mathbb{Z}^8$ with even coordinate sum has even square, and a vector of $(\mathbb{Z}+\frac{1}{2})^8$ has square $\sum x_i^2 \equiv 8\cdot\frac{1}{4} = 2$ modulo 2. Unimodularity follows by computing the index: the sublattice $\{x \in \mathbb{Z}^8 : \sum x_i \in 2\mathbb{Z}\}$ has discriminant 4 and index 2 in $\mathbb{Z}^8$, and adjoining the half-integer vector $\frac{1}{2}(1,\dots,1)$ halves the covolume, giving discriminant 1. For a lattice L write $L(-1)$ for the same group with the form negated. The *K3 lattice* is

$$\Lambda_{K3} = U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$$

   of rank 22; it is even and unimodular of signature $(3,19)$ (exercise 6.4.1 (1)).

2. A *marking* of a K3 surface S is an isometry $\alpha\colon H^2(S,\mathbb{Z}) \to \Lambda_{K3}$ for the cup form, and a *marked K3 surface* is a pair (S,α).

3. A *polarized K3 surface of degree* $2d$, with $d \geq 1$, is a pair (S,h) consisting of a K3 surface S and a primitive class $h \in H^2(S,\mathbb{Z})$ which is the first Chern class of an ample line bundle and satisfies $\langle h,h\rangle = 2d$.

4. Fix once and for all a primitive $h_0 \in \Lambda_{K3}$ with $\langle h_0,h_0\rangle = 2d$, and put

$$\Lambda_{2d} = h_0^{\perp} \subseteq \Lambda_{K3}$$

   A *marking* of a polarized K3 surface (S,h) of degree $2d$ is a marking α of S with $\alpha(h) = h_0$.

5. A *Hodge isometry* between K3 surfaces S and S' is an isometry $\varphi\colon H^2(S,\mathbb{Z}) \to H^2(S',\mathbb{Z})$ for the cup forms whose complexification satisfies $\varphi_{\mathbb{C}}\big(H^{2,0}(S)\big) = H^{2,0}(S')$.

[a] Humphreys [26, §12.1] constructs the E_8 root system. Its listed roots $\pm\varepsilon_i \pm \varepsilon_j$ and $\frac{1}{2}\sum_i (-1)^{k(i)}\varepsilon_i$ with $\sum_i k(i)$ even span this lattice. His defining condition for the ambient lattice gives instead its image under the sign change of one coordinate. E_8 is one of the exceptional diagrams of [37, Classification of Connected Dynkin Diagrams].

A marking exists for every K3 surface, because example 3.4.4 identifies $H^2(S,\mathbb{Z})$ with Λ_{K3} up to isometry; a marking is a choice of such an identification, and there are as many as there are isometries of Λ_{K3}. The construction depends on the choice not being canonical, and the ambiguity it creates is what the quotient in definition 5.3.3 divides out.

The lattice Λ_{2d} has rank 21 and, since $\mathbb{R}h_0$ is positive definite, signature $(2,19)$. It is not unimodular: its discriminant group is cyclic of order $2d$ (exercise 6.4.1 (2)). Writing $h_0 = e + df$ inside the first

hyperbolic summand of Λ_{K3} produces a primitive vector of square $2d$ for every $d \geq 1$, so every degree occurs, the smallest being $2d = 2$; the case $2d = 4$ is the degree of a smooth quartic surface in $\mathbb{C}\mathrm{P}^3$ with its hyperplane class, computed in example 6.4.4. That Λ_{2d} depends only on d, and not on which primitive vector of square $2d$ was chosen, is Eichler's criterion for Λ_{K3} ([28, Eichler's criterion for the K3 lattice]): the isometry group of Λ_{K3} acts transitively on the primitive vectors of a fixed square. It is a theorem of the arithmetic theory of quadratic forms, external to this series and taken here on faith, and it is what allows a marking of a polarized surface to be found at all.

A Hodge isometry is automatically an isomorphism of the whole Hodge structure. Since φ is defined over $\mathbb{Z}$, its complexification commutes with conjugation, so $\varphi_{\mathbb{C}}\big(H^{0,2}(S)\big) = H^{0,2}(S')$; and by example 3.4.4 the summand $H^{1,1}$ is the orthogonal complement of $H^{2,0} \oplus H^{0,2}$ for the cup form, so $\varphi_{\mathbb{C}}$ carries it across as well.

With the model lattice fixed, a marked polarized surface produces a point of a fixed space. Let (S, h, α) be a marked polarized K3 surface of degree $2d$ and let σ span $H^{2,0}(S)$. By part (3) of example 3.4.4 the class σ is orthogonal to h, so $\alpha_{\mathbb{C}}(\sigma)$ lies in $\Lambda_{2d} \otimes \mathbb{C}$, and the two relations (3.10) of that example say

$$\big\langle \alpha_{\mathbb{C}}\sigma,\, \alpha_{\mathbb{C}}\sigma \big\rangle = 0, \qquad \big\langle \alpha_{\mathbb{C}}\sigma,\, \overline{\alpha_{\mathbb{C}}\sigma} \big\rangle > 0$$

The *period point* of (S, h, α), in the sense attached to a single surface in example 3.4.4, is the line

$$\mathcal{P}(S, h, \alpha) = \big[\alpha_{\mathbb{C}}(\sigma)\big] \in \mathbb{P}\big(\Lambda_{2d} \otimes \mathbb{C}\big) \tag{6.1}$$

independent of the choice of σ because $H^{2,0}(S)$ is a line. The set of points satisfying the two displayed conditions is

$$D_{2d} = \big\{\, [\sigma] \in \mathbb{P}\big(\Lambda_{2d} \otimes \mathbb{C}\big) \;\big|\; \langle \sigma, \sigma \rangle = 0,\ \langle \sigma, \bar{\sigma} \rangle > 0 \,\big\}$$

This is the period domain of definition 3.3.2 for the numerical type $k = 2$, $h^{2,0} = h^{0,2} = 1$, $h^{1,1} = 19$, computed in example 3.3.6 with $n = 19$. The polarizing form of proposition 3.2.9 in weight two is the *negative* of the cup form, so it has signature $(19, 2)$ on Λ_{2d}, and the description there in terms of $Q = -\langle -, - \rangle$ becomes the one displayed here on changing signs. Dropping the polarization is not an option: on all of Λ_{K3} the same recipe produces a 20-dimensional open subset of a quadric which is *not* a period domain, because the positivity at $p = 1$ fails there (part (2) of example 3.4.4, and example 3.3.6).

The general theory already gives a good deal about D_{2d}: it is an open complex submanifold of a smooth quadric, homogeneous under a real orthogonal group, with compact isotropy (proposition 3.3.5). What is special to this numerical type, and what the rest of the section rests on, is that the homogeneous space in question is one of the classical bounded domains. Call a connected complex manifold a *Hermitian symmetric domain* when it carries a Hermitian metric for which every point is an isolated fixed point of an involutive holomorphic isometry.

Proposition 6.4.2: The Period Domain of a Polarized K3 Surface

Let $V_{\mathbb{R}} = \Lambda_{2d} \otimes \mathbb{R}$ with its cup form of signature $(2, 19)$, let $G_{\mathbb{R}} = \mathrm{O}(V_{\mathbb{R}}) \cong \mathrm{O}(2, 19)$, and let D_{2d} be as above.

1. Sending $[\sigma] = [x + iy]$ to the plane $\mathrm{span}_{\mathbb{R}}(x, y)$ with the orientation (x, y) is a homeomorphism from D_{2d} onto the set of oriented positive definite real 2-planes in $V_{\mathbb{R}}$. Under it $G_{\mathbb{R}}$ acts transitively, the isotropy group of a point is $\mathrm{SO}(2) \times \mathrm{O}(19)$, and
$$D_{2d} \cong \mathrm{O}(2, 19)\big/\big(\mathrm{SO}(2) \times \mathrm{O}(19)\big)$$
2. D_{2d} has exactly two connected components, interchanged by the antiholomorphic involution $[\sigma] \mapsto [\bar{\sigma}]$, and each is homeomorphic to an open ball of real dimension 38. In particular $\dim_{\mathbb{C}} D_{2d} = 19$.
3. $T^h D_{2d} = T D_{2d}$: every holomorphic map into D_{2d} is horizontal.
4. Each connected component of D_{2d} is a Hermitian symmetric domain.

***Proof*:**
Write $m = 21$ for the rank of Λ_{2d}. An element of $\mathrm{GL}(V_{\mathbb{R}})$ preserves the cup form if and only if it preserves its negative, so the group $G_{\mathbb{R}}$ of the statement is the group $G_{\mathbb{R}}$ of proposition 3.3.5 attached to the polarizing form, and the results of section 3.3 apply to it verbatim.

1. *Planes and the group action.* Let $[\sigma] \in D_{2d}$ and write $\sigma = x + iy$ with $x, y \in V_{\mathbb{R}}$. Expanding as in example 3.3.6, with the signs reversed because the form used there is $-\langle -, - \rangle$, the two defining conditions read
$$\langle x, x \rangle = \langle y, y \rangle > 0, \qquad \langle x, y \rangle = 0 \tag{6.2}$$
so x and y are independent and $P := \mathrm{span}_{\mathbb{R}}(x, y)$ is a positive definite plane. That example also establishes that $[\sigma] \mapsto (P, (x, y))$ is a bijection onto the oriented positive definite planes: rescaling σ by $re^{i\theta}$ rotates the ordered pair (x, y) without changing the plane or its orientation, and every oriented positive definite plane has an oriented basis satisfying (6.2), unique up to that rotation and scaling.

   The bijection is a homeomorphism. On a standard affine chart of $\mathbb{P}(\Lambda_{2d} \otimes \mathbb{C})$ a holomorphic section σ of the tautological line is available, and $(\mathrm{Re}\sigma, \mathrm{Im}\sigma)$ is then a continuous oriented frame of P, so the map is continuous. In the other direction, an oriented plane admits locally a continuous oriented basis (u_1, u_2), and Gram-Schmidt for the positive definite form on it,
$$x = \frac{u_1}{\sqrt{\langle u_1, u_1 \rangle}}, \qquad y' = u_2 - \langle u_2, x \rangle x, \qquad y = \frac{y'}{\sqrt{\langle y', y' \rangle}}$$
   produces continuously a pair satisfying (6.2) and inducing the given orientation, hence the point $[x + iy]$.

   For the group action, let $g \in G_{\mathbb{R}}$ fix $[\sigma]$, say $g\sigma = \lambda\sigma$. Applying conjugation gives $g\bar{\sigma} = \bar{\lambda}\bar{\sigma}$, and $\langle g\sigma, g\bar{\sigma} \rangle = \langle \sigma, \bar{\sigma} \rangle$ forces $|\lambda| = 1$, say $\lambda = e^{i\theta}$. Separating real and imaginary parts of

$g(x+iy) = e^{i\theta}(x+iy)$,

$$gx = \cos\theta\, x - \sin\theta\, y, \qquad gy = \sin\theta\, x + \cos\theta\, y$$

so g preserves P and acts on it by a rotation in the oriented orthonormal basis (x, y) given by (6.2). Since g preserves P it preserves $P^\perp$, and conversely a rotation of P together with any isometry of $P^\perp$ assembles to an element of $G_\mathbb{R}$ fixing $[\sigma]$, by the displayed formulas read backwards. So the isotropy group is $\mathrm{SO}(P) \times \mathrm{O}(P^\perp) \cong \mathrm{SO}(2) \times \mathrm{O}(19)$, the second factor being the full orthogonal group of the negative definite 19-dimensional space $P^\perp$; this agrees with the general computation of exercise 3.3.1 (3), since $\mathrm{SO}(2) \cong \mathrm{U}(1)$. Transitivity of $G_\mathbb{R}$ on D_{2d} is proposition 3.3.5.

2. *Components and dimension.* Fix a positive definite plane $W \subseteq V_\mathbb{R}$ with a chosen orientation, so that $V_\mathbb{R} = W \oplus W^\perp$ with $W^\perp$ negative definite of dimension 19; write $\pi\colon V_\mathbb{R} \to W$ for the orthogonal projection, $\|w\|^2 = \langle w, w\rangle$ on W and $\|u\|^2 = -\langle u, u\rangle$ on $W^\perp$, both positive definite.

   If P is a positive definite plane then $\pi|_P$ is injective, since a nonzero $v \in P \cap W^\perp$ would satisfy $\langle v, v\rangle > 0$ and $\langle v, v\rangle < 0$. Being a map between 2-dimensional spaces it is an isomorphism, so P is the graph

   $$P = \{\, w + Aw \;\mid\; w \in W \,\}$$

   of the linear map $A\colon W \to W^\perp$ obtained by composing $(\pi|_P)^{-1}$ with the projection to $W^\perp$. Conversely the graph of any A is a 2-plane, and for $w \neq 0$,

   $$\langle w + Aw,\, w + Aw\rangle = \|w\|^2 - \|Aw\|^2$$

   so the graph is positive definite exactly when $\|Aw\| < \|w\|$ for every $w \neq 0$, that is, exactly when the operator norm satisfies $\|A\| < 1$, the supremum defining it being attained on the compact unit sphere of W. So $P \mapsto A_P$ is a bijection from the positive definite planes onto the open unit ball

   $$B = \{\, A \in \mathrm{Hom}(W, W^\perp) \;\mid\; \|A\| < 1 \,\}$$

   an open convex subset of a real vector space of dimension $2 \cdot 19 = 38$, hence connected. It is a homeomorphism, both directions being given by linear algebra applied to a continuously varying frame, as in part (1).

   An oriented positive definite plane carries in addition a sign $\epsilon(P) \in \{\pm 1\}$, equal to $+1$ when $\pi|_P$ is orientation preserving for the chosen orientation of W. Combining, $[\sigma] \mapsto (A_P, \epsilon(P))$ is a homeomorphism of D_{2d} onto $B \times \{\pm 1\}$, since reversing the orientation of a plane changes ϵ and nothing else. So D_{2d} has exactly two components, each an open ball of real dimension 38. The involution $[\sigma] \mapsto [\bar\sigma]$ fixes P and replaces the oriented basis (x, y) by $(x, -y)$, so it preserves A_P and reverses ϵ, interchanging the two components; it is antiholomorphic because conjugation on $\Lambda_{2d} \otimes \mathbb{C}$ is. Finally D_{2d} is a complex manifold (proposition 3.3.5 and proposition 3.3.4) of real dimension 38, so $\dim_\mathbb{C} D_{2d} = 19$, agreeing with the count in example 3.3.6.

3. *Horizontality.* This is example 5.3.9 with $r = 19$: for the numerical type $h^{2,0} = h^{0,2} = 1$ the tangent space to the compact dual at $[\sigma]$ is

   $$T_{[\sigma]}\check{D}_{2d} = \mathrm{Hom}\big(\mathbb{C}\sigma,\ \sigma^\perp/\mathbb{C}\sigma\big) \tag{6.3}$$

and the only nonvacuous horizontality condition, the one at $p = 2$, is satisfied by every tangent vector. So $T^h\check{D}_{2d} = T\check{D}_{2d}$, and the same holds on the open subset D_{2d}.

4. *The symmetry at a point.* Part (2) of theorem 6.1.4 already gives this conclusion for any numerical type with no $\mathfrak{g}^{r,-r}$ in $|r| \geq 2$, a hypothesis checked for the present type in exercise 6.1.1 (2); the involution is written out because it gives more, namely the symmetry in its $\mathrm{O}(2,19)$ form. Fix $[\sigma] \in D_{2d}$ with oriented plane P and set

$$s_P = \mathrm{id}_P \oplus (-\mathrm{id}_{P^\perp}) \in G_{\mathbb{R}}$$

It is an involution, it is the identity on P and so preserves both P and its orientation, and it fixes σ because $\sigma \in P \otimes \mathbb{C}$. Being induced by a linear map preserving the form, it acts holomorphically on $\mathbb{P}(\Lambda_{2d} \otimes \mathbb{C})$ and preserves both conditions defining D_{2d}, so it restricts to a holomorphic involution of D_{2d} preserving each component; and it lies in $G_{\mathbb{R}}$, so it is an isometry of the $G_{\mathbb{R}}$-invariant Hermitian Hodge metric of definition 6.1.2.

Its differential at $[\sigma]$ is $-\mathrm{id}$. Write $P_{\mathbb{C}} = \mathbb{C}\sigma \oplus \mathbb{C}\bar{\sigma}$, a decomposition valid because $\langle\sigma,\sigma\rangle = 0$ while $\langle\sigma,\bar{\sigma}\rangle \neq 0$, so that σ and $\bar{\sigma}$ are independent. Then $\sigma^\perp$ contains both $\mathbb{C}\sigma$ and $P^\perp \otimes \mathbb{C}$, the sum of the two is direct because $\mathbb{C}\sigma \subseteq P_{\mathbb{C}}$, and its dimension is $1 + (m-2) = m - 1 = \dim \sigma^\perp$; hence

$$\sigma^\perp = \mathbb{C}\sigma \oplus \big(P^\perp \otimes \mathbb{C}\big), \qquad \sigma^\perp/\mathbb{C}\sigma \cong P^\perp \otimes \mathbb{C}$$

Now s_P acts as the identity on $\mathbb{C}\sigma$ and as $-\mathrm{id}$ on $P^\perp \otimes \mathbb{C}$, so by the equivariance of the identification (6.3) [32, the tangent space of a Grassmannian] it acts on $\mathrm{Hom}(\mathbb{C}\sigma, \sigma^\perp/\mathbb{C}\sigma)$ by $\theta \mapsto (-\mathrm{id}) \circ \theta \circ \mathrm{id}^{-1} = -\theta$.

That $[\sigma]$ is isolated in the fixed locus follows from the linearity of s_P directly. The fixed locus of s_P in $\mathbb{P}(\Lambda_{2d} \otimes \mathbb{C})$ is the union of the projectivized eigenspaces, $\mathbb{P}(P_{\mathbb{C}}) \cup \mathbb{P}(P^\perp \otimes \mathbb{C})$. No point of the second lies in D_{2d}: writing $\tau = u + iv$ with $u, v \in P^\perp$ gives $\langle\tau,\bar{\tau}\rangle = \langle u,u\rangle + \langle v,v\rangle \leq 0$, since $P^\perp$ is negative definite. A point $[\alpha\sigma + \beta\bar{\sigma}]$ of the first has $\langle\alpha\sigma + \beta\bar{\sigma}, \alpha\sigma + \beta\bar{\sigma}\rangle = 2\alpha\beta\langle\sigma,\bar{\sigma}\rangle$, which vanishes only when $\alpha = 0$ or $\beta = 0$, so the first contributes exactly $[\sigma]$ and $[\bar{\sigma}]$, and the latter lies in the other component by part (2). So $[\sigma]$ is the unique fixed point of s_P on its own component, completing the proof.

The proposition places the K3 period domain among the objects the classical theory of bounded symmetric domains already covers, and that is the situation theorem 6.1.4 was careful not to assume. In general the negativity of the holomorphic sectional curvature holds only along T^hD, and the period domain of a general numerical type is not Hermitian symmetric; a period map is then constrained by a distribution of positive codimension, and the curvature estimate applies only to the directions it can move in. Here the distribution is everything and the domain is symmetric, so the estimate, the distance-decreasing property of corollary 6.1.6 and the rigidity statement of theorem 6.1.7 all apply to D_{2d} and its quotients without restriction. The same degeneration was visible in weight one (example 6.1.5), where the domain is the Siegel space of example 3.3.3. What the two cases share is the shape of the numerical type: the nonzero Hodge numbers occupy two consecutive bidegrees, as in weight one, or three consecutive bidegrees with a one-dimensional space at each end, as here. Either shape leaves no $\mathfrak{g}^{r,-r}$ with $|r| \geq 2$, the computations of example 6.1.5 and exercise 6.1.1 (2) using the shape of the numerical type and nothing else, so part (2) of theorem 6.1.4 applies and the domain is symmetric; shapes of both kinds occur in higher weights as well, and what singles out these two is that their moduli theory was understood classically.

Symmetry of the domain says nothing about injectivity of the period map, and injectivity is a theorem about surfaces rather than about domains. A point of D_{2d} records a line in $\Lambda_{2d}\otimes\mathbb{C}$ and nothing else, so recovering a surface from it asserts that all the geometry of S is encoded in one line of transcendental data.

Theorem 6.4.3: Global Torelli for K3 Surfaces

Let S and S' be projective K3 surfaces and let $\varphi\colon H^2(S,\mathbb{Z})\to H^2(S',\mathbb{Z})$ be a Hodge isometry in the sense of definition 6.4.1.

1. If φ carries the first Chern class of some ample line bundle on S to the first Chern class of an ample line bundle on S', then there is a unique isomorphism $f\colon S'\to S$ of complex manifolds with $f^*=\varphi$.
2. Consequently, if (S,h) and (S',h') are polarized K3 surfaces of degree $2d$ and $\varphi(h)=h'$, then $f^*=\varphi$ for a unique isomorphism $f\colon S'\to S$, and in particular $S\cong S'$.

Part (2) is the case of part (1) in which the ample class carried across is h itself. Part (1) is not proved here, and it is proved nowhere in this series. The argument of [34] establishes the statement first for Kummer surfaces, where it reduces to the corresponding assertion for complex tori of dimension two and hence to weight-one Hodge theory, and then propagates it to all K3 surfaces by a density argument inside the space of marked surfaces. Both halves need machinery outside this book: the geometry of Kummer surfaces and their resolutions on one side, and the deformation theory that makes the density argument available on the other.

The hypothesis in part (1) cannot be dropped. A Hodge isometry is linear algebra together with a compatibility with one line, and there are Hodge isometries of $H^2(S,\mathbb{Z})$ induced by no automorphism of S, starting with $-\mathrm{id}$, which is a Hodge isometry for every S but carries an ample class to its negative. The effectivity hypothesis pins down which of the many chambers into which the classes of positive square are divided the isometry must respect, and the theorem says that once it respects the right one it is geometric.

Torelli converts the period point from an invariant into a complete invariant, and that is what turns D_{2d} into a moduli space.

Example 6.4.4: The Moduli of Degree-$2d$ Polarized K3 Surfaces

Fix $d\geq 1$ and let $\mathcal{M}_{2d}^{m}$ be the set of isomorphism classes of marked polarized K3 surfaces (S,h,α) of degree $2d$, two being isomorphic when an isomorphism of the surfaces matches both the polarizations and the markings. Sending (S,h,α) to (6.1) defines

$$\mathcal{P}\colon \mathcal{M}_{2d}^{m}\to D_{2d}$$

Forgetting the marking. Two markings of the same (S,h) differ by an isometry of Λ_{K3} fixing h_0, so put

$$\Gamma_{2d}=\left\{\,\tilde g|_{\Lambda_{2d}}\;\middle|\;\tilde g\in \mathrm{O}(\Lambda_{K3}),\ \tilde g(h_0)=h_0\,\right\}\subseteq \mathrm{O}(\Lambda_{2d})$$

Changing the marking changes the period point by the corresponding element of Γ_{2d}, so composing with the quotient map gives a well-defined $\mathcal{M}_{2d}\to\Gamma_{2d}\backslash D_{2d}$ on isomorphism classes of polarized surfaces. Since Γ_{2d} consists of integral isometries it is a subgroup of the group $G_{\mathbb{Z}}$ of definition 5.3.3,

so it acts properly discontinuously on D_{2d} by proposition 5.3.5.

Injectivity. Suppose $\mathcal{P}(S,h,\alpha)$ and $\mathcal{P}(S',h',\alpha')$ lie in the same Γ_{2d}-orbit, say $g\cdot\mathcal{P}(S,h,\alpha)=\mathcal{P}(S',h',\alpha')$ with $g=\tilde{g}|_{\Lambda_{2d}}$. Set

$$\varphi=(\alpha')^{-1}\circ\tilde{g}\circ\alpha\colon H^2(S,\mathbb{Z})\to H^2(S',\mathbb{Z})$$

a composite of isometries, hence an isometry. It satisfies $\varphi(h)=(\alpha')^{-1}\tilde{g}(h_0)=(\alpha')^{-1}(h_0)=h'$, and it is a Hodge isometry: with σ spanning $H^{2,0}(S)$ and σ' spanning $H^{2,0}(S')$, the hypothesis says $[\tilde{g}_{\mathbb{C}}\alpha_{\mathbb{C}}\sigma]=[\alpha'_{\mathbb{C}}\sigma']$, so $\varphi_{\mathbb{C}}(\mathbb{C}\sigma)=\mathbb{C}\sigma'$. Part (2) of theorem 6.4.3 makes the two polarized surfaces isomorphic. So $\mathcal{M}_{2d}\to\Gamma_{2d}\backslash D_{2d}$ is injective, and taking $\tilde{g}=\mathrm{id}$ shows that $\mathcal{P}$ itself is injective on marked surfaces.

Surjectivity, cited. Every point of D_{2d} lying on none of the hyperplane sections $\delta^\perp$ with $\delta\in\Lambda_{2d}$ and $\langle\delta,\delta\rangle=-2$ is the period point of a marked polarized K3 surface of degree $2d$. This is the surjectivity theorem for the K3 period map, due to Todorov, Looijenga and Siu and recorded in [1, the K3 period map]; it constructs the surface from its periods by transcendental means and is not proved here. The excluded classes are the ones that would prevent the polarization from being ample. A condition of exactly this shape appeared in section 6.3 as the locus on which extra Hodge classes are acquired, and here exercise 6.4.1 (4) identifies each $\delta^\perp\cap D_{2d}$ as itself a period domain, of dimension 18.

The count. Injectivity and surjectivity together identify $\mathcal{M}_{2d}$ with the image in $\Gamma_{2d}\backslash D_{2d}$ of the complement of those hyperplane sections, so the moduli of polarized K3 surfaces of degree $2d$ is 19-dimensional for every d. The independence of the degree is visible in the lattice: changing d changes the discriminant of Λ_{2d} but not its rank or its signature, and D_{2d} depends by proposition 6.4.2 only on the signature.

The count, checked on quartics. Let $S\subseteq\mathbb{CP}^3$ be a smooth quartic surface. Adjunction gives $K_S=\mathcal{O}_S$,[a] and $h^{1,0}(S)=\frac{1}{2}b_1(S)=0$, since $b_1(S)=b_1(\mathbb{CP}^3)=0$ by theorem 4.3.1, applied to the degree-4 Veronese embedding of $\mathbb{CP}^3$, in which S is a hyperplane section, so S is a K3 surface; the hyperplane class h is ample and primitive with $\langle h,h\rangle=\deg S=4$, making (S,h) a polarized K3 surface of degree 4, so $d=2$. These surfaces form a family of the kind studied in section 5.4. The global sections of $\mathcal{O}(4)$ on $\mathbb{CP}^3$ are the quartic forms,[b] a space of dimension $\binom{7}{4}=35$, so the smooth quartics are parametrized by an open subset of $\mathbb{CP}^{34}$, and two of them that differ by a projective transformation are isomorphic as polarized surfaces. Since $\dim\mathrm{PGL}_4=15$ and the stabilizer of a smooth quartic is finite (exercise 6.4.1 (3)), those transformations account for a 15-dimensional family of such identifications, and

$$34-15=19$$

This is a parameter count rather than a second proof, but it reproduces the dimension obtained from the period domain out of data (the dimension of a linear system and of a projective linear group) with no Hodge theory in it.

[a] [4, Adjunction Formula in Projective Space].
[b] [5, Global Sections of $\mathcal{O}(d)$].

The identification is more than a bijection of sets. The quotient $\Gamma_{2d}\backslash D_{2d}$ is an arithmetic quotient of a bounded symmetric domain, and such quotients carry an algebraic structure which the moduli of polarized K3 surfaces then inherits through the period map; nothing here depends on that structure. See also [13] for arithmetic quotients, and [1, period mappings and K3 surfaces] for the K3 case. What the present section gives is the Hodge-theoretic content: the domain, its geometry, and the two theorems that make the period point a complete invariant of the surface.

Every variety in Part II has been smooth and projective, or at worst compact Kähler, and every Hodge structure attached to one has been pure. Chapter 7 drops the compactness, and the singular case follows it. Purity fails in both, and what replaces it is a second filtration.

Exercise 6.4.1

1. Verify the assertions about Λ_{K3} made in definition 6.4.1: that U and E_8 are even, that U has signature $(1,1)$ and determinant -1, and hence, taking the unimodularity of E_8 as known, that Λ_{K3} is even and unimodular of rank 22 and signature $(3,19)$. Then show that for every $d \geq 1$ the vector $h_0 = e + df$ of the first hyperbolic summand is primitive with $\langle h_0, h_0\rangle = 2d$, and that no vector of Λ_{K3} has odd square.

2. Let Λ be a unimodular lattice and $h_0 \in \Lambda$ a primitive vector with $\langle h_0, h_0\rangle = 2d \neq 0$, and set $K = h_0^{\perp}$.

    1. Show that $y \mapsto \langle y, h_0\rangle$ is a surjection $\Lambda \to \mathbb{Z}$, and deduce that $\Lambda = \mathbb{Z}x \oplus K$ for some x with $\langle x, h_0\rangle = 1$, and that $h_0 = 2d\,x + k$ for some $k \in K$.
    2. Show that restriction of functionals gives a surjection $\Lambda \to K^{\vee}$ with kernel $\mathbb{Z}h_0$, and conclude that the discriminant group $K^{\vee}/K$ is cyclic of order $2d$.

    Deduce that Λ_{2d} has rank 21 and signature $(2,19)$ and is unimodular for no $d \geq 1$.

3. Let (S,h) be a polarized K3 surface of degree $2d$ and let $\mathrm{Aut}(S,h)$ be the group of automorphisms f of S with $f^*h = h$.

    1. Show that $f \mapsto f^*$ is an injective anti-homomorphism from $\mathrm{Aut}(S,h)$ into $\mathrm{O}\big(H^2(S,\mathbb{Z})\big)$, so that $f \mapsto (f^*)^{-1}$ is an injective homomorphism, and that their common image consists of isometries fixing h and fixing the period point.
    2. Using proposition 5.3.5, deduce that $\mathrm{Aut}(S,h)$ is finite.

4. Let $\lambda \in \Lambda_{2d}$ be nonzero and let $D_\lambda = \{[\sigma] \in D_{2d} \,\big|\, \langle \lambda, \sigma\rangle = 0\}$.

    1. Show that $D_\lambda \neq \emptyset$ if and only if $\langle \lambda, \lambda\rangle < 0$.
    2. Show that when $\langle \lambda, \lambda\rangle < 0$ the set D_λ is the period domain attached by the recipe of this section to the lattice $\lambda^{\perp} \cap \Lambda_{2d}$, of dimension 18.

    How does this compare with the local description of a Hodge locus in proposition 6.3.3?

5. Let (S,h,α) be a marked polarized K3 surface of degree $2d$ with period point $[\sigma] \in D_{2d}$. Show that
$$\alpha\big(\mathrm{NS}(S)\big) \cap \Lambda_{2d} = \big\{\, \lambda \in \Lambda_{2d} \;\big|\; \langle \lambda, \sigma\rangle = 0 \,\big\}$$
and that $\rho(S)$ is 1 plus the rank of the right-hand side. Deduce that $\rho(S) = 1$ for every period point outside a countable union of the sets D_λ of exercise 6.4.1 (4), and compare with example 6.3.8.

6. Let $f\colon \mathcal{X} \to S$ be a smooth projective family whose fibres are polarized K3 surfaces of degree $2d$, with period map Φ in degree 2.

    1. Assuming $d\tilde{\Phi}$ is injective at some point, show that $\dim_{\mathbb{C}} S \leq 19$, and explain why part (2) of exercise 5.3.1 (4) says nothing here.
    2. The smooth quartic surfaces in $\mathbb{C}\mathrm{P}^3$ are parametrized by an open subset of $\mathbb{C}\mathrm{P}^{34}$. Explain why the map to $\Gamma_{2d}\backslash D_{2d}$ attached to this family cannot separate two quartics differing by a projective transformation, and reconcile $34 - 15 = 19$ with (1).

Part III

Mixed Hodge Structures and Cycles

Two hypotheses have been assumed since the first page. Every variety has been compact, and every variety has been smooth. Both were doing work: compactness gave the Hodge theorem its elliptic operator and its finite-dimensional harmonic spaces, and smoothness gave the Hodge decomposition a bigrading to decompose into. Together they produced purity, the statement that H^k carries a Hodge structure of weight exactly k, and purity is what every result of Part I and Part II rests on.

Neither hypothesis survives contact with the objects algebraic geometry produces. The complement of a hypersurface is smooth and not compact; a nodal curve is compact and not smooth; the fibres of the families of Chapter 5 degenerate to both. On such a variety H^k is not pure, and the question is not whether purity fails but what replaces it.

The answer is a second filtration. Alongside the Hodge filtration $F^\bullet$ there is a weight filtration $W_\bullet$, and while H^k itself is no longer pure, the graded pieces $\mathrm{gr}^W_j H^k$ are, each of its own weight. This is a mixed Hodge structure, and the theorem of this Part is that every complex algebraic variety has one, functorially. Chapter 7 does the smooth open case, where the construction runs on differential forms with logarithmic poles along a boundary divisor. Chapter 8 drops smoothness, where the construction runs instead on simplicial resolutions. **??** then asks the question the whole book has been circling: which Hodge classes come from algebraic subvarieties.

7

Mixed Hodge Structures: Smooth Open Varieties

The simplest variety that is not compact is a compact one with a point removed, and it already breaks purity. Let E be an elliptic curve and $U = E \setminus \{p\}$. Then $H^1(U, \mathbb{Q})$ has rank two, as does $H^1(E, \mathbb{Q})$, but the two are not isomorphic as Hodge structures and cannot be: one of the two classes on U is the restriction of a class on E, of weight one, while the other is a residue at the puncture and behaves like a class of weight two. A single weight does not describe $H^1(U)$.

What describes it is a pair of filtrations. The Hodge filtration $F^\bullet$ survives unchanged from chapter 3. Beside it sits an increasing filtration $W_\bullet$ by weight, arranged so that the graded pieces gr^W_j are pure of weight j even though the total space is not pure of any weight. That pair is a mixed Hodge structure, and the theorem of this chapter is that the cohomology of every smooth complex variety carries one.

The construction is due to Deligne and it is concrete. Compactify U to a smooth X whose boundary $D = X \setminus U$ has normal crossings, which is possible by Hironaka's resolution theorem and is the one input taken on faith here. Then replace the de Rham complex of X by the complex of forms with logarithmic poles along D: forms that blow up along the boundary, but only as badly as dz/z. That complex computes the cohomology of U, it carries an obvious filtration by how many logarithmic poles a form is allowed, and the filtration by pole order is the weight filtration.

Two spectral sequences then have to be shown to degenerate, one for each filtration, and their degeneration is the technical heart of the chapter. Both arguments have the same shape and it is worth naming in advance: the terms of each page carry pure Hodge structures of known weight, the differentials are morphisms of Hodge structures, and a morphism between pure structures of different weights is zero. The bookkeeping is in identifying the weights.

7.1 Mixed Hodge Structures

The object attached to a smooth open variety is linear algebra before it is geometry: a rational vector space carrying two filtrations, one defined over $\mathbb{Q}$ and one over $\mathbb{C}$, tied together by a purity requirement on the graded pieces. This section fixes that definition, proves the theorem that makes the resulting category usable, and records the tensor constructions. The geometry that produces such an object begins in section 7.2.

Theorem 2.1.1 attaches to a compact Kähler manifold X a pure Hodge structure of weight k on $H^k(X,\mathbb{Q})$, and definition 3.1.1 extracted that datum. Compactness is not a technicality there. Let $U=\mathbb{C}^*$, a smooth affine curve. It retracts onto the unit circle, so $H^1(U,\mathbb{Q})\cong\mathbb{Q}$ is one-dimensional. A pure Hodge structure of odd weight k has no summand $V^{p,q}$ with $p=q$, since $p+q=k$ is odd; the symmetry $\overline{V^{p,q}}=V^{q,p}$ of definition 3.1.1 then pairs the nonzero summands off into conjugate pairs of equal dimension, so the underlying space has even dimension. Hence $H^1(\mathbb{C}^*,\mathbb{Q})$ carries no pure Hodge structure of weight 1.

Purity can fail in two ways, and both occur. The first is a shift: $H^1(\mathbb{C}^*,\mathbb{Q})$ does carry a pure structure, the Tate structure $\mathbb{Q}(-1)$ of weight 2 (section 7.3), so the weight exceeds the cohomological degree. The second is genuine mixing, where no single weight serves and the group is assembled from pieces of several weights; the punctured curves of section 7.4 are the first geometric instance. The definition below accommodates both by filtering rather than grading: the weights are separated by an increasing filtration whose graded pieces are pure, one weight at a time.

Definition 7.1.1: Mixed Hodge Structure

A *mixed Hodge structure* $V=(V_{\mathbb{Q}},W_\bullet,F^\bullet)$ consists of a finite-dimensional $\mathbb{Q}$-vector space $V_{\mathbb{Q}}$ together with

1. an increasing filtration $\cdots\subseteq W_{m-1}V\subseteq W_mV\subseteq\cdots$ of $V_{\mathbb{Q}}$ by $\mathbb{Q}$-subspaces, the *weight filtration*, with $W_mV=0$ for $m\ll 0$ and $W_mV=V_{\mathbb{Q}}$ for $m\gg 0$;
2. a decreasing filtration $\cdots\supseteq F^pV_{\mathbb{C}}\supseteq F^{p+1}V_{\mathbb{C}}\supseteq\cdots$ of $V_{\mathbb{C}}=V_{\mathbb{Q}}\otimes_{\mathbb{Q}}\mathbb{C}$ by complex subspaces, the *Hodge filtration*, with $F^pV_{\mathbb{C}}=V_{\mathbb{C}}$ for $p\ll 0$ and $F^pV_{\mathbb{C}}=0$ for $p\gg 0$,

subject to one compatibility. Write $W_mV_{\mathbb{C}}=W_mV\otimes_{\mathbb{Q}}\mathbb{C}$, write $\mathrm{gr}^W_mV=W_mV/W_{m-1}V$ for the graded pieces of the weight filtration, and equip $\mathrm{gr}^W_mV_{\mathbb{C}}$ with the filtration induced by F,

$$F^p\mathrm{gr}^W_mV_{\mathbb{C}}=\big((F^pV_{\mathbb{C}}\cap W_mV_{\mathbb{C}})+W_{m-1}V_{\mathbb{C}}\big)/W_{m-1}V_{\mathbb{C}} \tag{7.1}$$

The requirement is that for every m this filtration make gr^W_mV a pure Hodge structure of weight m in the sense of definition 3.1.1, that is, that it satisfy the transversality condition (3.1) in weight m.
The structure is *pure of weight k* if $\mathrm{gr}^W_mV=0$ for $m\neq k$. The *weights* of V are the integers m with $\mathrm{gr}^W_mV\neq 0$. An *integral* mixed Hodge structure carries in addition a finitely generated abelian group $V_{\mathbb{Z}}$ with $V_{\mathbb{Z}}\otimes_{\mathbb{Z}}\mathbb{Q}=V_{\mathbb{Q}}$.

Two features of the definition deserve emphasis before any example. The first is that the two filtrations live over different fields: W is rational and F is only complex. Nothing forces them to be compatible beyond (7.1). Over $\mathbb{C}$ the pair $(W_{\mathbb{C}},F)$ can always be split simultaneously, by Deligne's canonical bigrading (box 7.1.9), but no such splitting need be defined over $\mathbb{Q}$, and that failure is

exactly where the new invariants of the theory sit, as example 7.1.4 shows. The second is that purity is imposed one weight at a time and says nothing about how the pieces are glued. A mixed Hodge structure is therefore an iterated extension of pure structures, of increasing weight, and the whole content beyond chapter 3 lies in the extension data.

Definition 7.1.2: Morphisms and Sub-Structures

Let V and V' be mixed Hodge structures. A *morphism of mixed Hodge structures* $\varphi\colon V \to V'$ is a $\mathbb{Q}$-linear map $\varphi\colon V_{\mathbb{Q}} \to V'_{\mathbb{Q}}$ such that

$$\varphi(W_m V) \subseteq W_m V' \quad \text{for all } m, \qquad \varphi_{\mathbb{C}}(F^p V_{\mathbb{C}}) \subseteq F^p V'_{\mathbb{C}} \quad \text{for all } p$$

A *sub-mixed Hodge structure* of V is a $\mathbb{Q}$-subspace $C \subseteq V_{\mathbb{Q}}$ which, with the induced filtrations $W_m C = C \cap W_m V$ and $F^p C_{\mathbb{C}} = C_{\mathbb{C}} \cap F^p V_{\mathbb{C}}$, is itself a mixed Hodge structure. A *quotient* is V/C for such a C, with the filtrations $W_m(V/C) = (W_m V + C)/C$ and $F^p(V/C)_{\mathbb{C}} =$ the image of $F^p V_{\mathbb{C}}$.
Between integral structures a morphism is required in addition to come from a homomorphism $V_{\mathbb{Z}} \to V'_{\mathbb{Z}}$ of the underlying lattices. Unless the integral structure is named, all statements below are about the rational category.

Where definition 3.1.7 asked a map to respect the bigrading, this definition asks only that it respect two filtrations, and it imposes no condition relating the weights of V and V'. The consequence is that morphisms are now plentiful between structures of different weights: the projection of $\mathbb{Q}(0) \oplus \mathbb{Q}(1)$ onto $\mathbb{Q}(0)$ is one, and no nonzero map between the pure structures $\mathbb{Q}(0)$ and $\mathbb{Q}(1)$ is. Lemma 7.1.5 shows that on pure structures the two notions of morphism nevertheless agree.

Example 7.1.3: Pure and Tate Structures as Mixed Structures
Three instances of definition 7.1.1.

1. A pure Hodge structure V of weight k becomes a mixed one by setting $W_m V = 0$ for $m < k$ and $W_m V = V_{\mathbb{Q}}$ for $m \geq k$, with F its Hodge filtration (definition 3.1.5). The only nonzero graded piece is $\mathrm{gr}^W_k V = V$, and (7.1) returns F itself, so the requirement is exactly that V was a pure structure of weight k. Every pure structure is read as a mixed one this way from here on.

2. The Tate structure $\mathbb{Q}(n) = \mathbb{Z}(n) \otimes_{\mathbb{Z}} \mathbb{Q}$ of definition 3.1.11 is one-dimensional, concentrated in bidegree $(-n,-n)$, pure of weight $-2n$. Its weight filtration jumps once, at $m = -2n$.

3. Let $V_{\mathbb{Q}} = \mathbb{Q}e \oplus \mathbb{Q}f$ with $W_{-2}V = \mathbb{Q}f$, $W_{-1}V = W_{-2}V$, $W_0 V = V_{\mathbb{Q}}$ and $W_m V = 0$ for $m \leq -3$, and let $F^{-1}V_{\mathbb{C}} = V_{\mathbb{C}}$, $F^0 V_{\mathbb{C}} = \mathbb{C}e$, $F^1 V_{\mathbb{C}} = 0$. Then $\mathrm{gr}^W_{-2} V = \mathbb{Q}f$ receives $F^0 = 0$ and $F^{-1} =$ everything, so it is $\mathbb{Q}(1)$; and $\mathrm{gr}^W_0 V = \mathbb{Q}e$ receives $F^0 =$ everything and $F^1 = 0$, so it is $\mathbb{Q}(0)$. This is the direct sum $\mathbb{Q}(1) \oplus \mathbb{Q}(0)$, with W and F the sums of the two structures.

The third instance splits: it has a sub-structure, namely $\mathbb{Q}e$, mapping isomorphically onto the top graded piece. The next example is the same pair of graded pieces glued differently, and it is the reason the theory exists.

Example 7.1.4: A Nonsplit Extension of $\mathbb{Z}(0)$ by $\mathbb{Z}(1)$

Fix $z \in \mathbb{C}$. Let $V_{\mathbb{Z}} = \mathbb{Z}e \oplus \mathbb{Z}f$, where f is the generator $2\pi i$ of the lattice of $\mathbb{Z}(1)$, write $V_{\mathbb{Q}} = V_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$, and put

$$W_{-3}V = 0, \qquad W_{-2}V = W_{-1}V = \mathbb{Q}f, \qquad W_0V = V_{\mathbb{Q}}$$
$$F^1V_{\mathbb{C}} = 0, \qquad F^0V_{\mathbb{C}} = \mathbb{C}\,(e + zf), \qquad F^{-1}V_{\mathbb{C}} = V_{\mathbb{C}}$$

Call the resulting datum V_z. It is a mixed Hodge structure. Indeed $F^0V_{\mathbb{C}} \cap W_{-2}V_{\mathbb{C}} = \mathbb{C}(e + zf) \cap \mathbb{C}f = 0$, so $\mathrm{gr}^W_{-2}V_z$ carries $F^0 = 0$ and $F^{-1} =$ everything and is $\mathbb{Z}(1)$; and $F^0V_{\mathbb{C}}$ maps onto $\mathrm{gr}^W_0V_{z,\mathbb{C}}$, so that piece carries $F^0 =$ everything and $F^1 = 0$ and is $\mathbb{Z}(0)$. The inclusion $\mathbb{Z}(1) \to V_z$ and the projection $V_z \to \mathbb{Z}(0)$ are morphisms, so

$$0 \to \mathbb{Z}(1) \to V_z \to \mathbb{Z}(0) \to 0$$

is an extension of mixed Hodge structures, and its graded pieces do not depend on z.

Every extension of $\mathbb{Z}(0)$ by $\mathbb{Z}(1)$ arises this way, for exactly one z once a lift is fixed. Given such an extension V, let f generate the lattice $W_{-2}V \cap V_{\mathbb{Z}} = \mathbb{Z}(1)$ and let $e \in V_{\mathbb{Z}}$ be any lift of the generator of $\mathbb{Z}(0)$. Purity of gr^W_0V forces F^0 to surject onto it and purity of $\mathrm{gr}^W_{-2}V$ forces $F^0 \cap W_{-2}V_{\mathbb{C}} = 0$, so $F^0V_{\mathbb{C}}$ is a line meeting $\mathbb{C}f$ trivially and surjecting onto $\mathrm{gr}^W_0V_{\mathbb{C}}$, hence $F^0V_{\mathbb{C}} = \mathbb{C}(e + zf)$ for a unique $z \in \mathbb{C}$. The remaining steps are forced as well: $F^{-1}V_{\mathbb{C}} \supseteq W_{-2}V_{\mathbb{C}} + F^0V_{\mathbb{C}} = V_{\mathbb{C}}$, and $F^1V_{\mathbb{C}}$ has zero image in both graded pieces, hence is zero.

Changing the lift replaces e by $e + nf$ with $n \in \mathbb{Z}$, which replaces z by $z - n$. So the invariant of the extension is the class of z in $\mathbb{C}/\mathbb{Z}$, equivalently the number

$$\exp(2\pi i z) \in \mathbb{C}^*$$

and $V_z \cong V_{z'}$ by an isomorphism inducing the identity on $\mathbb{Z}(1)$ and on $\mathbb{Z}(0)$ if and only if $z - z' \in \mathbb{Z}$.

The extension splits precisely when $z \in \mathbb{Z}$. A splitting is a morphism $s\colon \mathbb{Z}(0) \to V_z$ with $\pi \circ s = \mathrm{id}$, so $s(1) = e + nf$ for some $n \in \mathbb{Z}$; compatibility with F forces $e + nf \in F^0V_{\mathbb{C}} = \mathbb{C}(e + zf)$, and comparing coefficients of e makes the scalar 1, so $n = z$ and $z \in \mathbb{Z}$. Conversely if $z = n \in \mathbb{Z}$ then $\mathbb{Z}(e + nf)$ is a sub-structure mapping isomorphically onto $\mathbb{Z}(0)$, and $V_z = \mathbb{Z}(1) \oplus \mathbb{Z}(0)$.

The same computation with $\mathbb{Q}$ in place of $\mathbb{Z}$ throughout classifies extensions of $\mathbb{Q}(0)$ by $\mathbb{Q}(1)$ by the class of z in $\mathbb{C}/\mathbb{Q}$, and V_z splits as a rational mixed Hodge structure exactly when $z \in \mathbb{Q}$.

The graded pieces of V_z are $\mathbb{Z}(1)$ and $\mathbb{Z}(0)$ for every z, so a mixed Hodge structure is not determined by its associated graded. What distinguishes V_z from V_0 is the position of one complex line relative to one rational lattice, and the invariant that records it, $\exp(2\pi i z) \in \mathbb{C}^*$, is transcendental in nature: no amount of rational linear algebra produces it, as exercise 7.1.1 (6) makes precise. This is the shape every computation in this chapter takes. The pure pieces are cohomology groups of compact objects, computed by the theory of Part I, and the gluing between them is the new geometric information contributed by the removed locus; the punctured curves of section 7.4 are the first case worked out in full.

Before the main theorem, three facts about the category are needed. The first compares the two notions of morphism on pure structures, the second manufactures sub-objects and quotients, and the third is the vanishing statement that drives everything else.

Lemma 7.1.5: Morphisms Between Pure Structures

Let V and V' be pure Hodge structures of weights k and l, regarded as mixed Hodge structures as in example 7.1.3, and let $\varphi\colon V \to V'$ be a morphism of mixed Hodge structures.

1. If $k = l$ then $\varphi_{\mathbb{C}}(V^{p,q}) \subseteq V'^{p,q}$ for all p, q; that is, φ is a morphism of Hodge structures in the sense of definition 3.1.7.
2. If $k \neq l$ then $\varphi = 0$.
3. The kernel and image of a morphism of pure Hodge structures of weight k are sub-Hodge structures of weight k, and the quotient of a pure Hodge structure of weight k by a sub-Hodge structure is pure of weight k.

***Proof*:**
Since φ is defined over $\mathbb{Q}$, its complexification commutes with the conjugation of $V_{\mathbb{C}}$ over $V_{\mathbb{R}}$, so $\varphi_{\mathbb{C}}(\overline{F^q V_{\mathbb{C}}}) \subseteq \overline{F^q V'_{\mathbb{C}}}$. Combining this with compatibility with F and using the recovery formula (3.2),

$$\varphi_{\mathbb{C}}\big(V^{p,q}\big) = \varphi_{\mathbb{C}}\big(F^p V_{\mathbb{C}} \cap \overline{F^q V_{\mathbb{C}}}\big) \subseteq F^p V'_{\mathbb{C}} \cap \overline{F^q V'_{\mathbb{C}}} \tag{7.2}$$

for every p, q with $p + q = k$.

(1). When $k = l$, the right-hand side of (7.2) is $V'^{p,q}$ by (3.2) again, which is the assertion.

(2). Suppose first $l < k$ and let $p + q = k$. Then $q \geq l - p + 1$, so $\overline{F^q V'_{\mathbb{C}}} \subseteq \overline{F^{l-p+1} V'_{\mathbb{C}}}$, and (3.1) for V' gives $F^p V'_{\mathbb{C}} \oplus \overline{F^{l-p+1} V'_{\mathbb{C}}} = V'_{\mathbb{C}}$. Hence the right-hand side of (7.2) is zero, so $\varphi_{\mathbb{C}}$ kills every $V^{p,q}$ and therefore kills $V_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}$. Suppose next $l > k$. As a mixed structure V' has $W_k V' = 0$, while compatibility with W gives $\varphi(V_{\mathbb{Q}}) = \varphi(W_k V) \subseteq W_k V' = 0$.

(3). Let $\varphi\colon V \to V'$ be a morphism of pure structures of weight k. By (1) the map $\varphi_{\mathbb{C}}$ is bigraded, so

$$\ker \varphi_{\mathbb{C}} = \bigoplus_{p+q=k} \big(\ker \varphi_{\mathbb{C}} \cap V^{p,q}\big), \qquad \operatorname{im} \varphi_{\mathbb{C}} = \bigoplus_{p+q=k} \varphi_{\mathbb{C}}\big(V^{p,q}\big)$$

the first because a vector lies in the kernel exactly when each of its bihomogeneous components does, the second because $\varphi_{\mathbb{C}}$ carries the (p, q)-summand into the (p, q)-summand. Both $\ker \varphi$ and $\operatorname{im} \varphi$ are defined over $\mathbb{Q}$, so their complexifications are stable under conjugation, and the induced bigradings inherit $\overline{S^{p,q}} = S^{q,p}$ from V and V'; each is therefore a pure Hodge structure of weight k. For a subspace S carrying a bigrading in this way, $F^p S_{\mathbb{C}} = \bigoplus_{r \geq p} S^{r,k-r} = S_{\mathbb{C}} \cap F^p V_{\mathbb{C}}$, so its Hodge filtration is the one induced from V. For the quotient, if $S \subseteq V$ is a sub-Hodge structure then $V_{\mathbb{C}}/S_{\mathbb{C}} = \bigoplus_{p+q=k} V^{p,q}/S^{p,q}$ carries the quotient bigrading, again conjugation-symmetric and defined over $\mathbb{Q}$, completing the proof.

Part (2) is the mechanism that will kill differentials in section 7.3: a map between pure structures of different weights has no room to be nonzero, and the whole degeneration argument there consists of arranging for such a map. The proof of part (1) also shows that proposition 3.1.8, whose argument uses only the bigrading and not the lattice, applies verbatim to morphisms of rational Hodge structures; it is used in that form throughout this section.

Lemma 7.1.6: Sub-Structures and Quotients

Let V be a mixed Hodge structure.

1. For every r the subspace W_rV is a sub-mixed Hodge structure, with $\mathrm{gr}_m^W(W_rV) = \mathrm{gr}_m^W V$ for $m \leq r$ and 0 for $m > r$.
2. Let $C \subseteq V_{\mathbb{Q}}$ be a sub-mixed Hodge structure. Then for each m the injection $\mathrm{gr}_m^W C \to \mathrm{gr}_m^W V$ identifies $\mathrm{gr}_m^W C$ with a sub-Hodge structure of $\mathrm{gr}_m^W V$, and the quotient V/C of definition 7.1.2 is a mixed Hodge structure with
$$\mathrm{gr}_m^W(V/C) = \mathrm{gr}_m^W V/\mathrm{gr}_m^W C$$
as pure Hodge structures of weight m.

***Proof*:**
(1). The induced weight filtration on W_rV is $W_mV \cap W_rV = W_{\min(m,r)}V$, so the graded pieces are as stated. For $m \leq r$ the induced Hodge filtration on $\mathrm{gr}_m^W(W_rV)$ is computed by (7.1) from $F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}} \cap W_rV_{\mathbb{C}} = F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$, which is the datum computing $F^p\mathrm{gr}_m^W V$. So $\mathrm{gr}_m^W(W_rV)$ and $\mathrm{gr}_m^W V$ agree as filtered spaces, and purity is inherited.

(2). The map $\mathrm{gr}_m^W C \to \mathrm{gr}_m^W V$ is injective because $W_mC = C \cap W_mV$ and $W_{m-1}C = C \cap W_{m-1}V$, and it is a morphism of mixed Hodge structures between pure structures of weight m, hence a morphism of Hodge structures by lemma 7.1.5. By proposition 3.1.8 its image meets $F^p\mathrm{gr}_m^W V_{\mathbb{C}}$ in the image of $F^p\mathrm{gr}_m^W C_{\mathbb{C}}$, so the filtration that $\mathrm{gr}_m^W C$ carries as a subquotient of C agrees with the one it inherits as a subspace of $\mathrm{gr}_m^W V$; by lemma 7.1.5(3) it is a sub-Hodge structure of $\mathrm{gr}_m^W V$.

Next the underlying spaces. Since $W_{m-1}V \subseteq W_mV$, the modular law gives
$$W_mV \cap \big(W_{m-1}V + C\big) = W_{m-1}V + \big(W_mV \cap C\big) = W_{m-1}V + W_mC$$
so that
$$\mathrm{gr}_m^W(V/C) = \frac{W_mV + C}{W_{m-1}V + C} \cong \frac{W_mV}{W_mV \cap (W_{m-1}V + C)} = \frac{W_mV}{W_{m-1}V + W_mC} = \mathrm{gr}_m^W V/\mathrm{gr}_m^W C$$
the middle isomorphism being the second isomorphism theorem. Under it, the class of $y \in W_mV$ modulo C corresponds to the class of $[y] \in \mathrm{gr}_m^W V$ modulo $\mathrm{gr}_m^W C$.

It remains to identify the filtration (7.1) on $\mathrm{gr}_m^W(V/C)$ with the quotient of $F^p\mathrm{gr}_m^W V_{\mathbb{C}}$. One inclusion is immediate: an element of $F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$ maps into $F^p(V/C)_{\mathbb{C}} \cap W_m(V/C)_{\mathbb{C}}$, so the quotient filtration is contained in the induced one. For the reverse, let $\xi \in F^p(V/C)_{\mathbb{C}} \cap W_m(V/C)_{\mathbb{C}}$. Choose a representative $y \in F^pV_{\mathbb{C}}$ of ξ; membership in $W_m(V/C)_{\mathbb{C}}$ gives a decomposition $y = w + c$ with $w \in W_mV_{\mathbb{C}}$ and $c \in C_{\mathbb{C}}$. Let r be minimal with $c \in W_rC_{\mathbb{C}}$.

If $r \leq m$ then $y = w + c \in W_mV_{\mathbb{C}}$, so $y \in F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$ and the class of ξ in $\mathrm{gr}_m^W(V/C)$ is the image of $[y] \in F^p\mathrm{gr}_m^W V_{\mathbb{C}}$, as required. If $r > m$ then $w \in W_mV_{\mathbb{C}} \subseteq W_{r-1}V_{\mathbb{C}}$, so $[y] = [c]$ in $\mathrm{gr}_r^W V_{\mathbb{C}}$; moreover $y \in F^pV_{\mathbb{C}} \cap W_rV_{\mathbb{C}}$, whence $[c] = [y] \in F^p\mathrm{gr}_r^W V_{\mathbb{C}}$. Since $[c]$ lies in $\mathrm{gr}_r^W C_{\mathbb{C}}$, the identification of the first paragraph gives $[c] \in F^p\mathrm{gr}_r^W C_{\mathbb{C}}$, so there is $c' \in F^pC_{\mathbb{C}} \cap W_rC_{\mathbb{C}}$ with

$c - c' \in W_{r-1}C_{\mathbb{C}}$. Replacing y by $y - c'$, which again lies in $F^pV_{\mathbb{C}}$ and again represents ξ, replaces c by $c - c'$ and lowers r. Since $W_rC = 0$ for $r \ll 0$, finitely many replacements reach the case $r \leq m$.

Finally, $\mathrm{gr}_m^W(V/C) = \mathrm{gr}_m^W V / \mathrm{gr}_m^W C$ with the quotient filtration is the quotient of a pure Hodge structure of weight m by a sub-Hodge structure, hence pure of weight m by lemma 7.1.5(3), which completes the proof.

Lemma 7.1.7: Vanishing Below the Weight

Let P be a pure Hodge structure of weight n and let T be a mixed Hodge structure with $W_{n-1}T = T_{\mathbb{Q}}$, that is, with all weights at most $n-1$. Then every morphism $\psi\colon P \to T$ of mixed Hodge structures is zero.

***Proof*:**
Induction on the number of nonzero graded pieces of T. If $T = 0$ there is nothing to prove. Otherwise let b be the largest weight of T, so $b \leq n-1$ and $W_bT = T_{\mathbb{Q}}$. By lemma 7.1.6, $W_{b-1}T$ is a sub-mixed Hodge structure and $T/W_{b-1}T$ is a mixed Hodge structure whose only graded piece is $\mathrm{gr}_b^W T$, hence pure of weight b. The composite of ψ with the projection is a morphism of mixed Hodge structures from a pure structure of weight n to a pure structure of weight $b \neq n$, hence zero by lemma 7.1.5(2). Therefore $\psi(P_{\mathbb{Q}}) \subseteq W_{b-1}T$, and ψ is a morphism of mixed Hodge structures into $W_{b-1}T$, since the filtrations there are induced from T. That target has fewer nonzero graded pieces, so the inductive hypothesis gives $\psi = 0$, completing the proof.

For pure structures, proposition 3.1.8 obtained strictness from the bigrading: a preimage could be truncated componentwise. A mixed Hodge structure has no such bigrading available, and no reason is visible from the definition why a class deep in the weight filtration of the target should have a preimage equally deep in the source. That it does is the theorem below, and everything this chapter proves about $H^*(U)$ rests on it. Without strictness the associated graded would not be exact, the category would not be abelian, and the uniqueness assertions of section 7.3 would be unavailable.

Theorem 7.1.8: Strictness and the Abelian Category of Mixed Hodge Structures

Let $\varphi\colon V \to V'$ be a morphism of mixed Hodge structures.

1. φ is *strict for the weight filtrations*:
$$\varphi(V_{\mathbb{Q}}) \cap W_m V' = \varphi(W_m V) \qquad \text{for every } m$$
2. φ is *strict for the Hodge filtrations*:
$$\varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^p V'_{\mathbb{C}} = \varphi_{\mathbb{C}}\big(F^p V_{\mathbb{C}}\big) \qquad \text{for every } p$$
3. $\ker\varphi$ and $\mathrm{im}\varphi$, with the filtrations induced from V and V', are mixed Hodge structures, and for every m
$$\mathrm{gr}^W_m(\ker\varphi) = \ker(\mathrm{gr}^W_m\varphi), \qquad \mathrm{gr}^W_m(\mathrm{im}\varphi) = \mathrm{im}(\mathrm{gr}^W_m\varphi)$$
4. Mixed Hodge structures and their morphisms form an abelian category, and each $V \mapsto \mathrm{gr}^W_m V$ is an exact functor from it to the category of pure Hodge structures of weight m.

***Proof*:**
Write $\ell(V)$ for the number of m with $\mathrm{gr}^W_m V \neq 0$. Throughout, $\mathrm{gr}^W_m\varphi\colon \mathrm{gr}^W_m V \to \mathrm{gr}^W_m V'$ denotes the induced map; it is $\mathbb{Q}$-linear and compatible with the filtrations (7.1), so by lemma 7.1.5 it is a morphism of pure Hodge structures of weight m and proposition 3.1.8 applies to it.

Step 1: strictness for W, by induction on $\ell(V)$. The inductive hypothesis is statements (1), (2) and (3) for every morphism whose source has fewer than $\ell(V)$ nonzero graded pieces; this is legitimate because Steps 2 and 3 below derive (2) and (3) for a morphism from (1) for that same morphism, so only (1) has to be propagated.

If $\ell(V) = 0$ then $V = 0$ and there is nothing to prove. Let n be the largest weight of V, so $W_n V = V_{\mathbb{Q}}$, and set $A = W_{n-1}V$, a sub-mixed Hodge structure with $\ell(A) = \ell(V) - 1$ by lemma 7.1.6(1). By the inductive hypothesis, $\varphi|_A$ is strict for W and its image $C' = \varphi(A)$ is a sub-mixed Hodge structure of V'.

For $m \geq n$ the claim is trivial, since then $W_m V = V_{\mathbb{Q}}$ and $\varphi(V_{\mathbb{Q}}) \subseteq W_n V' \subseteq W_m V'$. The substance is the case $m = n-1$, from which the remaining cases follow: granting
$$\varphi(V_{\mathbb{Q}}) \cap W_{n-1}V' = \varphi(A) \tag{7.3}$$
and taking $m \leq n-1$,
$$\varphi(V_{\mathbb{Q}}) \cap W_m V' = \big(\varphi(V_{\mathbb{Q}}) \cap W_{n-1}V'\big) \cap W_m V' = \varphi(A) \cap W_m V' = \varphi(W_m A) = \varphi(W_m V)$$
the third equality by the inductive hypothesis for $\varphi|_A$ and the fourth because $W_m A = W_m V$ for $m \leq n-1$.

The inclusion $\supseteq$ in (7.3) holds because $A = W_{n-1}V$ maps into $W_{n-1}V'$. For the reverse, let $x \in V_{\mathbb{Q}}$ with $\varphi(x) \in W_{n-1}V'$, and let $[x] \in \mathrm{gr}^W_n V$ be its class. Then $\mathrm{gr}^W_n\varphi([x]) = 0$, so $[x]$ lies

in $K = \ker(\mathrm{gr}_n^W \varphi)$, a sub-Hodge structure of $\mathrm{gr}_n^W V$ by lemma 7.1.5(3). Let $V_K \subseteq V_{\mathbb{Q}}$ be the preimage of K under $V_{\mathbb{Q}} \twoheadrightarrow \mathrm{gr}_n^W V$.

The subspace V_K is a sub-mixed Hodge structure. Its weight filtration is $W_m V_K = W_m V$ for $m \leq n-1$ and V_K for $m \geq n$, so its graded pieces are those of V below n together with K in degree n; those below n carry the same induced Hodge filtration as in V, since $W_m V_K = W_m V$ there. In degree n, (7.1) computes $F^p \mathrm{gr}_n^W V_K$ as the image of $F^p V_{\mathbb{C}} \cap V_{K,\mathbb{C}}$, and this is $F^p \mathrm{gr}_n^W V_{\mathbb{C}} \cap K_{\mathbb{C}}$: a class in the latter is represented by some $v \in F^p V_{\mathbb{C}}$ whose image lies in $K_{\mathbb{C}}$, and such a v lies in $V_{K,\mathbb{C}}$ by definition of V_K. So $\mathrm{gr}_n^W V_K$ is K with its sub-Hodge structure, which is pure of weight n.

Every $v \in V_K$ has $\mathrm{gr}_n^W \varphi([v]) = 0$, hence $\varphi(v) \in W_{n-1} V' =: B'$. So φ restricts to a morphism $V_K \to B'$, and $C' = \varphi(A) \subseteq B'$ is a sub-mixed Hodge structure of B', the filtrations induced from V' and from B' agreeing on it. By lemma 7.1.6(2) the quotients V_K/A and B'/C' are mixed Hodge structures, the first pure of weight n (its only nonzero graded piece is K) and the second with all weights at most $n-1$ (its graded pieces are quotients of those of B'). The composite $V_K \to B' \to B'/C'$ kills A, so it factors through a morphism $V_K/A \to B'/C'$, which is zero by lemma 7.1.7. Hence $\varphi(V_K) \subseteq C' = \varphi(A)$; since $x \in V_K$, this proves (7.3).

Step 2: strictness for F, and the combined form. All the subspaces in (1) are defined over $\mathbb{Q}$, and intersections and images of $\mathbb{Q}$-subspaces commute with the extension of scalars to $\mathbb{C}$, so (1) may be used in the form $\varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap W_m V'_{\mathbb{C}} = \varphi_{\mathbb{C}}(W_m V_{\mathbb{C}})$. The claim to be proved is the combined statement

$$\varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^p V'_{\mathbb{C}} \cap W_m V'_{\mathbb{C}} = \varphi_{\mathbb{C}}\big(F^p V_{\mathbb{C}} \cap W_m V_{\mathbb{C}}\big) \tag{7.4}$$

for all p and m; part (2) is the case $m \gg 0$.

The inclusion $\supseteq$ is clear. For the reverse, let w' lie in the left-hand side, and assume $w' \neq 0$. Let $m_0 \leq m$ be minimal with $w' \in W_{m_0} V'_{\mathbb{C}}$. By (1), $w' = \varphi_{\mathbb{C}}(v)$ for some $v \in W_{m_0} V_{\mathbb{C}}$, so the class $[w'] \in \mathrm{gr}_{m_0}^W V'_{\mathbb{C}}$ lies in the image of $\mathrm{gr}_{m_0}^W \varphi$; and $w' \in F^p V'_{\mathbb{C}} \cap W_{m_0} V'_{\mathbb{C}}$ puts $[w']$ in $F^p \mathrm{gr}_{m_0}^W V'_{\mathbb{C}}$. By proposition 3.1.8 applied to $\mathrm{gr}_{m_0}^W \varphi$ there is $\beta \in F^p \mathrm{gr}_{m_0}^W V_{\mathbb{C}}$ with $\mathrm{gr}_{m_0}^W \varphi(\beta) = [w']$, and by (7.1) the class β lifts to some $b \in F^p V_{\mathbb{C}} \cap W_{m_0} V_{\mathbb{C}}$. Then

$$w' - \varphi_{\mathbb{C}}(b) \in \varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^p V'_{\mathbb{C}} \cap W_{m_0 - 1} V'_{\mathbb{C}}$$

since the difference lies in $W_{m_0} V'_{\mathbb{C}}$ and has zero class in $\mathrm{gr}_{m_0}^W V'_{\mathbb{C}}$. Repeating the construction on $w' - \varphi_{\mathbb{C}}(b)$ lowers m_0 by at least one each time, and $W_j V' = 0$ for $j \ll 0$, so after finitely many steps the remainder is zero. Summing the elements b produced gives $v_0 \in F^p V_{\mathbb{C}} \cap W_{m_0} V_{\mathbb{C}} \subseteq F^p V_{\mathbb{C}} \cap W_m V_{\mathbb{C}}$ with $\varphi_{\mathbb{C}}(v_0) = w'$, which is (7.4).

Step 3: kernel and image. Give $\mathrm{im}\varphi$ the induced filtrations. By (1) and (2), $W_m(\mathrm{im}\varphi) = \varphi(W_m V)$ and $F^p(\mathrm{im}\varphi)_{\mathbb{C}} = \varphi_{\mathbb{C}}(F^p V_{\mathbb{C}})$. Since the filtration on $\mathrm{im}\varphi$ is induced, the map $\mathrm{gr}_m^W(\mathrm{im}\varphi) \to \mathrm{gr}_m^W V'$ is injective. The natural map $\mathrm{gr}_m^W V \to \mathrm{gr}_m^W(\mathrm{im}\varphi)$ induced by φ is surjective, and its kernel consists of the classes of $v \in W_m V$ with $\varphi(v) \in \varphi(W_{m-1} V)$; by (1) this is the same as $\varphi(v) \in W_{m-1} V'$, that is, the same as $[v] \in \ker(\mathrm{gr}_m^W \varphi)$. Hence $\mathrm{gr}_m^W(\mathrm{im}\varphi) = \mathrm{im}(\mathrm{gr}_m^W \varphi)$ as subspaces of $\mathrm{gr}_m^W V'$, which by lemma 7.1.5(3) is a pure Hodge structure of weight m. Its induced Hodge filtration is the image of $\mathrm{im}\varphi_{\mathbb{C}} \cap F^p V'_{\mathbb{C}} \cap W_m V'_{\mathbb{C}}$, which by (7.4) is the image of $\varphi_{\mathbb{C}}(F^p V_{\mathbb{C}} \cap W_m V_{\mathbb{C}})$, that is $\mathrm{gr}_m^W \varphi\big(F^p \mathrm{gr}_m^W V_{\mathbb{C}}\big)$; by proposition 3.1.8 this is $\mathrm{im}(\mathrm{gr}_m^W \varphi) \cap F^p \mathrm{gr}_m^W V'_{\mathbb{C}}$, the filtration of the sub-Hodge structure. So $\mathrm{im}\varphi$ is a mixed Hodge structure.

For the kernel, first $\mathrm{gr}_m^W(\ker \varphi) = \ker(\mathrm{gr}_m^W \varphi)$. The inclusion $\subseteq$ is clear. Conversely let $[v] \in \ker(\mathrm{gr}_m^W \varphi)$ with $v \in W_m V$; then $\varphi(v) \in \varphi(V_{\mathbb{Q}}) \cap W_{m-1} V' = \varphi(W_{m-1} V)$ by (1), so

$\varphi(v) = \varphi(u)$ with $u \in W_{m-1}V$, and $v - u \in \ker\varphi \cap W_mV$ has class $[v]$. The induced Hodge filtration on $\mathrm{gr}_m^W(\ker\varphi)$ is the image of $\ker\varphi_{\mathbb{C}} \cap F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$, which is contained in $\ker(\mathrm{gr}_m^W\varphi) \cap F^p\mathrm{gr}_m^WV_{\mathbb{C}}$ and contains it: given ξ in the latter, lift it to $v \in F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$; then $\varphi_{\mathbb{C}}(v)$ lies in $\varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^pV'_{\mathbb{C}} \cap W_{m-1}V'_{\mathbb{C}}$, so by (7.4) $\varphi_{\mathbb{C}}(v) = \varphi_{\mathbb{C}}(u)$ for some $u \in F^pV_{\mathbb{C}} \cap W_{m-1}V_{\mathbb{C}}$, and $v - u$ is a lift of ξ inside $\ker\varphi_{\mathbb{C}} \cap F^pV_{\mathbb{C}} \cap W_mV_{\mathbb{C}}$. So $\mathrm{gr}_m^W(\ker\varphi)$ is the sub-Hodge structure $\ker(\mathrm{gr}_m^W\varphi)$ of gr_m^WV, and $\ker\varphi$ is a mixed Hodge structure.

Step 4: the category. Morphisms between two fixed mixed Hodge structures form a $\mathbb{Q}$-vector space, and the direct sum of two mixed Hodge structures, filtered summandwise, is a biproduct, so the category is additive. By Step 3 the subspace $\ker\varphi$ with its induced filtrations is a mixed Hodge structure, and it is a categorical kernel: a morphism $\psi\colon T \to V$ with $\varphi\psi = 0$ factors through it as a $\mathbb{Q}$-linear map, and the factorization respects both filtrations because those on $\ker\varphi$ are induced from V. Dually $\mathrm{coker}\varphi = V'/\mathrm{im}\varphi$ with the quotient filtrations is a mixed Hodge structure by lemma 7.1.6(2), and is a categorical cokernel.

It remains to check that the canonical map $V/\ker\varphi \to \mathrm{im}\varphi$ is an isomorphism, which is where strictness is used. It is a bijection of $\mathbb{Q}$-vector spaces and a morphism of mixed Hodge structures. Its inverse is one too: the image of $W_m(V/\ker\varphi)$ is $\varphi(W_mV) = W_m(\mathrm{im}\varphi)$ by (1), and the image of $F^p(V/\ker\varphi)_{\mathbb{C}}$ is $\varphi_{\mathbb{C}}(F^pV_{\mathbb{C}}) = F^p(\mathrm{im}\varphi)_{\mathbb{C}}$ by (2), so the inverse carries each filtration step into the corresponding one. The category is therefore abelian.

Finally gr_m^W is exact. Let $0 \to V' \to V \to V'' \to 0$ be exact, so that $V' \to V$ is a kernel of $V \to V''$ and $V \to V''$ is a cokernel of $V' \to V$. By Step 3 the isomorphism $V' \cong \mathrm{im}(V' \to V)$ gives $\mathrm{gr}_m^WV' \cong \mathrm{gr}_m^W\big(\mathrm{im}(V' \to V)\big) = \mathrm{im}(\mathrm{gr}_m^WV' \to \mathrm{gr}_m^WV)$, so $\mathrm{gr}_m^WV' \to \mathrm{gr}_m^WV$ is injective with image $\mathrm{gr}_m^W\big(\ker(V \to V'')\big) = \ker(\mathrm{gr}_m^WV \to \mathrm{gr}_m^WV'')$; and surjectivity of $V \to V''$ gives $\mathrm{gr}_m^WV'' = \mathrm{gr}_m^W(\mathrm{im}(V \to V'')) = \mathrm{im}(\mathrm{gr}_m^WV \to \mathrm{gr}_m^WV'')$. The graded sequence is therefore exact, completing the proof.

Strictness is what allows weight arguments to be run inside exact sequences. A short exact sequence of mixed Hodge structures induces, for each m, a short exact sequence of pure Hodge structures of weight m; and a map between two mixed structures whose weight ranges are disjoint must vanish, by lemma 7.1.7 and its counterpart for higher weights (exercise 7.1.1 (2)). Section 7.3 uses these two facts together: the differentials of a spectral sequence built from mixed Hodge structures are morphisms between pure structures of different weights, hence zero, and the sequence degenerates. Strictness is also what makes the mixed Hodge structure on the cohomology of a variety unique once it is functorial: two structures with the same underlying filtrations related by the identity map must coincide, because the identity is strict in both directions.

7.1.9 Note: Deligne's Canonical Bigrading The proof above never split the two filtrations simultaneously, and no splitting is needed for strictness. One exists nevertheless. Deligne showed that

$$I^{p,q} = \left(F^pV_{\mathbb{C}} \cap W_{p+q}V_{\mathbb{C}}\right) \cap \left(\overline{F^qV_{\mathbb{C}}} \cap W_{p+q}V_{\mathbb{C}} + \sum_{j\geq 1} \overline{F^{q-j}V_{\mathbb{C}}} \cap W_{p+q-j-1}V_{\mathbb{C}}\right)$$

satisfies $V_{\mathbb{C}} = \bigoplus_{p,q} I^{p,q}$, with $W_mV_{\mathbb{C}} = \bigoplus_{p+q\leq m} I^{p,q}$ and $F^pV_{\mathbb{C}} = \bigoplus_{r\geq p}\bigoplus_s I^{r,s}$, and with $\overline{I^{p,q}} \equiv I^{q,p}$ modulo $W_{p+q-2}V_{\mathbb{C}}$. The bigrading is given by a formula in W, F and $\overline{F}$, so it is preserved by every morphism, and theorem 7.1.8(1) and (2) follow from it by truncating a preimage componentwise exactly as in proposition 3.1.8. It fails to be a bigrading of a pure Hodge structure only in that the conjugation symmetry holds modulo lower weight rather than on the nose; see [33, the canonical bigrading of a mixed Hodge structure] and [18]. For V_z of example 7.1.4 the recipe returns $I^{0,0} = \mathbb{C}(e + zf)$ and $I^{-1,-1} = \mathbb{C}f$, and the failure of $\overline{I^{0,0}} = \mathbb{C}(e + \bar{z}f)$ to equal $I^{0,0}$ is the extension class again.

The constructions of definition 3.1.9 extend to the mixed setting, and the extension is forced: the weight of a tensor product should be the sum of the weights, so the filtration steps must add. These are the operations that let morphisms, twists and duals be discussed at all, and section 7.3 uses the twist to state the weight bounds.

Proposition 7.1.10: Tensor Products and Duals of Mixed Structures

Let V and V' be mixed Hodge structures.

1. The space $V_{\mathbb{Q}} \otimes_{\mathbb{Q}} V'_{\mathbb{Q}}$ with

$$W_m(V \otimes V') = \sum_{a+b=m} W_aV \otimes W_bV', \qquad F^p(V \otimes V')_{\mathbb{C}} = \sum_{a+b=p} F^aV_{\mathbb{C}} \otimes F^bV'_{\mathbb{C}}$$

   is a mixed Hodge structure, and

$$\mathrm{gr}^W_m(V \otimes V') = \bigoplus_{a+b=m} \mathrm{gr}^W_aV \otimes \mathrm{gr}^W_bV'$$

   as pure Hodge structures of weight m, the summands being the tensor products of definition 3.1.9.

2. The space $V^\vee = \mathrm{Hom}_{\mathbb{Q}}(V_{\mathbb{Q}}, \mathbb{Q})$ with

$$W_m(V^\vee) = \left(W_{-m-1}V\right)^\perp, \qquad F^p(V^\vee)_{\mathbb{C}} = \left(F^{1-p}V_{\mathbb{C}}\right)^\perp$$

   where $(\,\cdot\,)^\perp$ denotes the annihilator, is a mixed Hodge structure with $\mathrm{gr}^W_m(V^\vee) = \left(\mathrm{gr}^W_{-m}V\right)^\vee$.

3. When V and V' are pure, these are the structures of definition 3.1.9; in particular the Tate twist $V(n) = V \otimes \mathbb{Q}(n)$ of a mixed Hodge structure satisfies $W_m\big(V(n)\big) = W_{m+2n}V$ and $F^p\big(V(n)\big)_{\mathbb{C}} = F^{p+n}V_{\mathbb{C}}$.

***Proof*:**
Step 1: a splitting adapted to both filtrations. Let E be a finite-dimensional complex vector space carrying a finite increasing filtration $W_\bullet$ and a finite decreasing filtration $F^\bullet$. Set $A_a^p = W_a \cap F^p$ and $B_a^p = (W_{a-1} \cap F^p) + (W_a \cap F^{p+1}) \subseteq A_a^p$, and choose a complement E_a^p with $A_a^p = B_a^p \oplus E_a^p$. Then

$$W_a \cap F^p = \bigoplus_{a' \le a,\ p' \ge p} E_{a'}^{p'} \tag{7.5}$$

for all a, p. That the right side spans the left follows from $A_a^p = A_{a-1}^p + A_a^{p+1} + E_a^p$ by induction on $(a, -p)$, the base case being the pairs with $A_a^p = 0$, which covers $a \ll 0$ and $p \gg 0$ because both filtrations are finite. The sum is direct by a dimension count: since $A_{a-1}^p \cap A_a^{p+1} = W_{a-1} \cap F^{p+1} = A_{a-1}^{p+1}$,

$$\dim E_a^p = \dim A_a^p - \dim A_{a-1}^p - \dim A_a^{p+1} + \dim A_{a-1}^{p+1}$$

and summing this over $a' \le a$ and $p' \ge p$ telescopes to $\dim A_a^p$. Taking $p \ll 0$ and $a \gg 0$ in (7.5) gives $E = \bigoplus_{a,p} E_a^p$, with $W_a = \bigoplus_{a' \le a, p} E_{a'}^p$ and $F^p = \bigoplus_{a, p' \ge p} E_a^{p'}$. Consequently $\mathrm{gr}_a^W E \cong \bigoplus_p E_a^p$, with the induced filtration $F^r \mathrm{gr}_a^W E = \bigoplus_{p \ge r} E_a^p$ under that identification.

Step 2: the tensor product. On $V \otimes V'$ the filtration W is increasing and defined over $\mathbb{Q}$, the filtration F is decreasing, and both are finite. Choose splittings E_a^p of $(W_\mathbb{C}, F)$ on $V_\mathbb{C}$ and $E_a'^p$ on $V'_\mathbb{C}$ as in Step 1. Then

$$W_m(V \otimes V')_\mathbb{C} = \bigoplus_{a+a' \le m} \bigoplus_{p,p'} E_a^p \otimes E_{a'}'^{p'}, \qquad F^r(V \otimes V')_\mathbb{C} = \bigoplus_{p+p' \ge r} \bigoplus_{a,a'} E_a^p \otimes E_{a'}'^{p'}$$

directly from the definitions and Step 1. Hence

$$\mathrm{gr}_m^W(V \otimes V')_\mathbb{C} = \bigoplus_{a+a'=m} \bigoplus_{p,p'} E_a^p \otimes E_{a'}'^{p'} = \bigoplus_{a+a'=m} \mathrm{gr}_a^W V_\mathbb{C} \otimes \mathrm{gr}_{a'}^W V'_\mathbb{C}$$

an identification defined over $\mathbb{Q}$, being the canonical map induced by $W_a V \otimes W_{a'} V' \to \mathrm{gr}_a^W V \otimes \mathrm{gr}_{a'}^W V'$. Under it, the filtration (7.1) is $\bigoplus_{a+a'=m} \bigoplus_{p+p' \ge r} E_a^p \otimes E_{a'}'^{p'}$, which by the last sentence of Step 1 is $\sum_{r_1+r_2=r} F^{r_1} \mathrm{gr}_a^W V_\mathbb{C} \otimes F^{r_2} \mathrm{gr}_{a'}^W V'_\mathbb{C}$ summed over $a + a' = m$. That is the Hodge filtration of the tensor product of the pure structures $\mathrm{gr}_a^W V$ and $\mathrm{gr}_{a'}^W V'$ of weights a and a', which by definition 3.1.9 is pure of weight $a + a' = m$. A direct sum of pure structures of weight m is pure of weight m, so $V \otimes V'$ is a mixed Hodge structure with the stated graded pieces.

Step 3: the dual. With the same splitting of $V_\mathbb{C}$, let $(E_a^p)^\vee \subseteq V_\mathbb{C}^\vee$ be the annihilator of all summands other than E_a^p, so $V_\mathbb{C}^\vee = \bigoplus_{a,p} (E_a^p)^\vee$. Then

$$W_m(V^\vee)_\mathbb{C} = \big(W_{-m-1} V_\mathbb{C}\big)^\perp = \bigoplus_{a \ge -m} \bigoplus_p (E_a^p)^\vee, \qquad F^r(V^\vee)_\mathbb{C} = \big(F^{1-r} V_\mathbb{C}\big)^\perp = \bigoplus_{p \le -r} \bigoplus_a (E_a^p)^\vee$$

since $W_{-m-1} V_\mathbb{C} = \bigoplus_{a \le -m-1, p} E_a^p$ and $F^{1-r} V_\mathbb{C} = \bigoplus_{p \ge 1-r, a} E_a^p$. So $(E_a^p)^\vee$ sits in weight $-a$ and Hodge level $-p$, giving $\mathrm{gr}_m^W(V^\vee)_\mathbb{C} = \bigoplus_p (E_{-m}^p)^\vee$ with $F^r = \bigoplus_{p \le -r} (E_{-m}^p)^\vee$. This is the dual of the pure structure $\mathrm{gr}_{-m}^W V$ in the sense of definition 3.1.9, whose Hodge filtration is the annihilator of $F^{1-r} \mathrm{gr}_{-m}^W V_\mathbb{C}$; the dual of a pure structure of weight $-m$ is pure of weight m, so

$V^\vee$ is a mixed Hodge structure with the stated graded pieces.

Step 4: the pure case and the twist. If V is pure of weight k and V' pure of weight l, the formulas give $W_m(V\otimes V')=0$ for $m<k+l$ and $V_{\mathbb{Q}}\otimes V'_{\mathbb{Q}}$ for $m\geq k+l$, so $V\otimes V'$ is pure of weight $k+l$ and Step 2 identifies its bigrading with that of definition 3.1.9; the dual is handled the same way. For the twist, $\mathbb{Q}(n)$ is one-dimensional with $W_m=0$ for $m<-2n$ and $F^p=\mathbb{Q}(n)_{\mathbb{C}}$ for $p\leq -n$, so the two displayed sums in (1) each have a single contributing term, giving $W_m(V(n))=W_{m+2n}V$ and $F^p(V(n))_{\mathbb{C}}=F^{p+n}V_{\mathbb{C}}$, which completes the proof.

The twist is the device that normalizes weights, and it is used in section 7.3 in exactly that role: the weight-2 piece appearing in the cohomology of a punctured curve is a Tate twist of a weight-0 piece, and naming it that way is what makes the residue map a morphism of mixed Hodge structures rather than a map that shifts the bigrading by hand.

Exercise 7.1.1

1. Let $V_{\mathbb{Q}}=\mathbb{Q}e_1\oplus\mathbb{Q}e_2$ with $W_{-1}V=0$, $W_0V=W_1V=\mathbb{Q}e_1$ and $W_2V=V_{\mathbb{Q}}$, and let $V'_{\mathbb{Q}}=\mathbb{Q}f$ with $W_{-1}V'=0$ and $W_0V'=V'_{\mathbb{Q}}$. Let $\varphi\colon V_{\mathbb{Q}}\to V'_{\mathbb{Q}}$ send $e_1\mapsto 0$ and $e_2\mapsto f$. Show that φ respects the weight filtrations but is not strict for them. Check that each of these weight filtrations on its own does underlie a mixed Hodge structure, namely $\mathbb{Q}(0)\oplus\mathbb{Q}(-1)$ and $\mathbb{Q}(0)$, and deduce that there is nevertheless no choice of Hodge filtrations making $(V_{\mathbb{Q}},W,F)$ and $(V'_{\mathbb{Q}},W,F')$ mixed Hodge structures with these weight filtrations and φ a morphism.

2. Let V and V' be mixed Hodge structures with no common weight, that is, for every m at least one of gr^W_mV and gr^W_mV' vanishes. Show that every morphism $V\to V'$ of mixed Hodge structures is zero. (First treat V pure of weight n: split V' using $W_{n-1}V'$, handling the weights above n by compatibility with W alone and those below n by lemma 7.1.7. Then induct on the number of nonzero graded pieces of V.)

3. Let V_z be as in example 7.1.4, regarded as a rational mixed Hodge structure, and suppose $z\notin\mathbb{Q}$. Show that its only sub-mixed Hodge structures are 0, $W_{-2}V_z$ and the whole structure. Conclude that V_z is not isomorphic to $\mathbb{Q}(1)\oplus\mathbb{Q}(0)$, and compute $\mathrm{Hom}(\mathbb{Q}(0),V_z)$ and $\mathrm{Hom}(V_z,\mathbb{Q}(0))$.

4. Let $0\to V'\to V\to V''\to 0$ be a short exact sequence of mixed Hodge structures. Show that

$$0\to F^pV'_{\mathbb{C}}\to F^pV_{\mathbb{C}}\to F^pV''_{\mathbb{C}}\to 0$$

   is exact for every p, and deduce that $\dim_{\mathbb{C}}F^pV_{\mathbb{C}}=\dim_{\mathbb{C}}F^pV'_{\mathbb{C}}+\dim_{\mathbb{C}}F^pV''_{\mathbb{C}}$. Deduce further that the numbers $h^{p,q}(V):=\dim_{\mathbb{C}}\mathrm{gr}^p_F\mathrm{gr}^W_{p+q}V_{\mathbb{C}}$, where $\mathrm{gr}^p_F=F^p/F^{p+1}$, are additive in short exact sequences.

5. Let V be a mixed Hodge structure with weights in the interval $[a,b]$, meaning $\mathrm{gr}^W_mV=0$ for $m<a$ and for $m>b$. Using proposition 7.1.10, determine the intervals containing the weights of $V(n)$, of $V^\vee$ and of $V\otimes V'$ when V' has weights in $[a',b']$. Then show that $\mathrm{Hom}(V,V')=0$ whenever $b<a'$.

6. Let V_z and V_0 be as in example 7.1.4. Exhibit an isomorphism of complex vector spaces $V_{z,\mathbb{C}}\to V_{0,\mathbb{C}}$ carrying $W_\bullet$ to $W_\bullet$ and $F^\bullet$ to $F^\bullet$, and check that it is not defined over $\mathbb{Q}$ unless $z\in\mathbb{Q}$. Explain in one sentence which ingredient of definition 7.1.1 the invariant $\exp(2\pi iz)$ therefore depends on.

7.2 The Logarithmic de Rham Complex

A mixed Hodge structure is two filtrations on one rational vector space, and section 7.1 gave the definition and the formal properties but no example coming from geometry. To put one on $H^k(U,\mathbb{Q})$ for a smooth open U there has to be a complex that computes $H^*(U,\mathbb{C})$ and that carries the two filtrations visibly. Smooth forms on U compute the cohomology but carry no holomorphic grading, and U is not compact, so nothing from Part I applies to it directly. The resolution is to move the whole computation onto a compactification $X \supseteq U$: this section constructs a complex of holomorphic sheaves on X, built from forms whose poles along the boundary are as mild as a pole can be, and proves that it computes $H^*(U,\mathbb{C})$. The two filtrations are read off it in section 7.3.

Everything depends on the boundary $D = X \setminus U$ being simple, and the first task is to say what simple means.

Definition 7.2.1: Normal Crossings Divisor

Let X be a complex manifold of dimension n and $D \subseteq X$ a closed analytic subset. Then D is a *normal crossings divisor* if every point $x \in D$ admits a holomorphic chart $\varphi\colon V \to \Delta^n$ centred at x and an integer $1 \le k \le n$ with

$$\varphi(D \cap V) \;=\; \{w \in \Delta^n \,\big|\, w_1 \cdots w_k = 0\}$$

Such a chart is *adapted to D at x*, and $k = k(x)$ is the number of *branches* of D at x. The divisor is a *simple* normal crossings divisor if in addition every irreducible component $D_1, \dots, D_N$ of D is a closed complex submanifold of X.

Simplicity is a global condition and it is what the residue calculus below needs: if the components are smooth then each of the k branches in an adapted chart at x lies in a different component, since a component containing two of them would be singular at x. Every normal crossings divisor in this chapter is simple, and from here on the adjective is dropped.

Example 7.2.2: Coordinate Hyperplanes, the Node and the Cusp

1. In $\mathbb{C}^n$ the divisor $D = \{z_1 \cdots z_k = 0\}$ is the local model itself: it is a normal crossings divisor with $N = k$ components $D_i = \{z_i = 0\}$, all smooth, and $k(x)$ is the number of coordinates vanishing at x. Its complement is $(\mathbb{C}^*)^k \times \mathbb{C}^{n-k}$.

2. The nodal cubic $C = \{y^2 = x^2(x+1)\} \subseteq \mathbb{C}^2$ is a normal crossings divisor. Near the origin $\sqrt{x+1}$ is a well-defined holomorphic function, so

$$y^2 - x^2(x+1) \;=\; \big(y - x\sqrt{x+1}\big)\big(y + x\sqrt{x+1}\big)$$

exhibits two smooth branches with distinct tangent lines $y = \pm x$; taking those two functions as new coordinates puts C in the model with $k = 2$. It is *not* simple: C is irreducible and its unique component is singular at the origin.

3. The cuspidal cubic $C = \{y^2 = x^3\} \subseteq \mathbb{C}^2$ is not a normal crossings divisor at the origin. The map $t \mapsto (t^2, t^3)$ is a homeomorphism of a disc onto a neighbourhood of 0 in C, so $C \setminus \{0\}$ is connected near 0, which rules out $k = 2$: in the model $\{w_1 w_2 = 0\}$ the punctured germ has two connected components. And $k = 1$ would make C smooth at 0, whereas the gradient $(-3x^2, 2y)$ vanishes there.

The construction needs a compactification whose boundary is of this shape, and producing one is not something this book can do. It is Hironaka's resolution of singularities, and it enters as a hypothesis.

Theorem 7.2.3: Existence of a Good Compactification

Let U be a smooth complex algebraic variety. There exist a smooth complex algebraic variety X, proper over $\mathbb{C}$, and an open immersion $U \hookrightarrow X$ whose complement $D = X \setminus U$ is a simple normal crossings divisor. If U is quasi-projective, then X may be taken projective. Such an X is called a *good compactification* of U.

No proof is given here, and none is given anywhere in this series. The statement is Hironaka's theorem on resolution of singularities in characteristic zero [24]: one first embeds U in some proper variety, then resolves the singularities of that variety and of the boundary until the boundary is a normal crossings divisor. The argument is a book-length induction on invariants of singularities and shares no machinery with anything developed earlier in this book.[1] The compactification is very far from unique, and the independence of everything built below from the choice is the substance of theorem 7.3.11.

Example 7.2.4: A Good Compactification of a Cubic Complement

Let $C \subseteq \mathbb{C}\mathrm{P}^2$ be an irreducible cubic curve with a single node at p, for instance $\{y^2z = x^2(x+z)\}$, and let $U = \mathbb{C}\mathrm{P}^2 \setminus C$. The obvious compactification $X_0 = \mathbb{C}\mathrm{P}^2$ fails: its boundary C is a normal crossings divisor, by the local computation of example 7.2.2, but not a simple one, since its only component is singular at p.

Blow up the node. Let $\pi\colon X = \mathrm{Bl}_p\mathbb{C}\mathrm{P}^2 \to \mathbb{C}\mathrm{P}^2$ be the blow-up at p,[a] let $E = \pi^{-1}(p)$ be the exceptional curve and let $\widetilde{C}$ be the strict transform of C. The two branches of C at p have distinct tangent directions, so their strict transforms meet E at two distinct points, each transversally, and $\widetilde{C}$ is smooth. The reduced divisor $D = \widetilde{C} + E$ therefore has two smooth components meeting transversally in two points and no triple points, so it is a simple normal crossings divisor. Since π is an isomorphism away from p and $p \in C$, one has $X \setminus D \cong \mathbb{C}\mathrm{P}^2 \setminus C = U$, and X is a good compactification of U.

[a] [5, Blow-Up of a Complex Manifold at a Point].

Fix a good compactification from now on: X a complex manifold of dimension n, $D \subseteq X$ a normal crossings divisor, $U = X \setminus D$ and $j\colon U \hookrightarrow X$ the inclusion. Throughout, $H^k(U, \mathbb{C})$ means the sheaf cohomology $H^k(U, \underline{\mathbb{C}})$, which the fine resolution of $\underline{\mathbb{C}}$ by smooth forms identifies with de Rham cohomology (theorem 1.1.2).

The complex to be built out of X has to see U, so it must be allowed poles along D; and it has to remain a complex of coherent sheaves on the compact X, so the poles must be bounded. Bounding the pole order of α alone is the wrong condition, because the exterior derivative then increases it. Bounding it for α and $d\alpha$ at once is the right one, and it is the smallest condition stable under d.

[1] Hironaka's *theorem* on resolution should not be confused with Hironaka's *example* of a smooth proper threefold that is not projective, which is a separate construction; see [4, Hironaka's Example].

Definition 7.2.5: Forms with Logarithmic Poles

Let $V \subseteq X$ be open and small enough that $D \cap V$ is the zero set of a single $f \in \mathcal{O}_X(V)$ generating its ideal. A holomorphic p-form α on $V \cap U$ has *logarithmic poles along* D if both $f\alpha$ and $f\,d\alpha$ extend to holomorphic forms on V. The condition is unchanged when f is replaced by a unit multiple, and it is local, so these forms are the sections of a subsheaf

$$\Omega^p_X(\log D) \;\subseteq\; j_*\Omega^p_U$$

of $\mathcal{O}_X$-modules. If α has logarithmic poles then so does $d\alpha$, since $f\,d(d\alpha) = 0$, so d makes

$$\Omega^\bullet_X(\log D)\colon \quad \Omega^0_X(\log D) \xrightarrow{d} \Omega^1_X(\log D) \xrightarrow{d} \cdots \xrightarrow{d} \Omega^n_X(\log D)$$

into a complex, the *logarithmic de Rham complex* of the pair (X, D). Its differentials are not $\mathcal{O}_X$-linear, so it is a complex of sheaves of $\mathbb{C}$-vector spaces whose terms happen to be $\mathcal{O}_X$-modules.

The definition is a growth condition, which is not a usable description. The next proposition converts it into a generating set: in an adapted chart the logarithmic forms are exactly the $\mathcal{O}$-combinations of wedge products of the n one-forms $dz_1/z_1, \dots, dz_k/z_k, dz_{k+1}, \dots, dz_n$. Every later computation in this section is made in that frame.

Proposition 7.2.6: The Local Frame of the Logarithmic Forms

Let $(V; z_1, \dots, z_n)$ be a chart adapted to D, with $D \cap V = \{z_1 \cdots z_k = 0\}$. Put

$$\omega_i \;=\; \frac{dz_i}{z_i} \quad (1 \le i \le k), \qquad \omega_i \;=\; dz_i \quad (k < i \le n)$$

and $\omega_S = \omega_{i_1} \wedge \cdots \wedge \omega_{i_p}$ for $S = \{i_1 < \cdots < i_p\} \subseteq \{1, \dots, n\}$. Then $\Omega^p_X(\log D)|_V$ is a free $\mathcal{O}_V$-module with basis $\{\omega_S : |S| = p\}$. In particular $\Omega^p_X(\log D)$ is locally free of rank $\binom{n}{p}$, it equals $\Lambda^p\Omega^1_X(\log D)$, and $\Omega^0_X(\log D) = \mathcal{O}_X$.

***Proof*:**
Write $f = z_1 \cdots z_k$, a defining equation for $D \cap V$. For $S \subseteq \{1, \dots, n\}$ set $z_S = \prod_{i \in S,\, i \le k} z_i$, so that $dz_S = z_S\,\omega_S$, and write $S^c = \{1, \dots, k\} \setminus S$.

Step 1: the listed forms are logarithmic. Each ω_i is closed: $d(z_i^{-1}dz_i) = -z_i^{-2}\,dz_i \wedge dz_i = 0$. Hence $d\omega_S = 0$, and for $\alpha = \sum_{|S|=p} a_S\,\omega_S$ with $a_S \in \mathcal{O}(V)$ one has $d\alpha = \sum_S da_S \wedge \omega_S$. Both $f\omega_S$ and $f\,da_S \wedge \omega_S$ are holomorphic, because $f\omega_S = (f/z_S)\,dz_S$ and f/z_S is a monomial. So α is a section of $\Omega^p_X(\log D)$.

Step 2: a logarithmic form is a combination of them. Let α be a section of $\Omega^p_X(\log D)$ over V and set $\eta = f\alpha$, a holomorphic p-form. Expand $\eta = \sum_{|T|=p} b_T\,dz_T$ with $b_T \in \mathcal{O}(V)$. Since

$dz_T = z_T\,\omega_T$,

$$\alpha \;=\; \frac{\eta}{f} \;=\; \sum_{|T|=p} \frac{b_T}{z_{T^c}}\;\omega_T \tag{7.6}$$

so the claim is that z_{T^c} divides b_T for every T.

The second half of definition 7.2.5 has not been used yet, and it is what gives the divisibility. From $\alpha = \eta/f$,

$$f\,d\alpha \;=\; d\eta - \frac{df}{f}\wedge\eta$$

and $d\eta$ is holomorphic, so $f\,d\alpha$ is holomorphic if and only if $\frac{df}{f}\wedge\eta = \sum_{i=1}^{k}\frac{1}{z_i}\,dz_i\wedge\eta$ is. Fix S with $|S| = p+1$ and read off the coefficient of dz_S: for each $i \in S$ with $i \le k$ the summand contributes $\varepsilon_i b_{S\setminus i}/z_i$ with $\varepsilon_i = \pm 1$, and the other summands contribute nothing to that slot. So

$$g_S \;:=\; \sum_{i\in S,\; i\le k} \varepsilon_i\,\frac{b_{S\setminus i}}{z_i}$$

is holomorphic on V. Multiply by f and restrict to $\{z_i = 0\}$ for a fixed $i \in S$ with $i \le k$. Every summand with index $j \neq i$ acquires the factor z_i from f/z_j and dies, so

$$0 \;=\; \varepsilon_i\, b_{S\setminus i} \prod_{l\le k,\; l\neq i} z_l \qquad \text{on } \{z_i = 0\}$$

The product $\prod_{l\neq i} z_l$ is nonzero on a dense open subset of the polydisc $\{z_i = 0\}$, so $b_{S\setminus i}$ vanishes on that dense subset and hence, by continuity, on all of $\{z_i = 0\}$; expanding $b_{S\setminus i}$ in powers of z_i then shows z_i divides it.

Writing $T = S\setminus i$, this says: z_i divides b_T for every $i \le k$ with $i \notin T$. The functions z_i for distinct i are coprime, and dividing out one at a time gives $z_{T^c}\big|b_T$. By (7.6) the coefficients of α in the frame are holomorphic.

Step 3: independence. Suppose $\sum_S a_S\omega_S = 0$ with a_S holomorphic. Multiplying by f gives

$$\sum_S a_S(f/z_S)\,dz_S = 0,$$

and the dz_S are a frame for Ω^p_V, so $a_S f/z_S = 0$ and hence $a_S = 0$ for every S. The forms ω_S are therefore a basis, of cardinality $\binom{n}{p}$; taking $p = 0$ the basis is the empty product 1, so $\Omega^0_X(\log D) = \mathcal{O}_X$, and the description of the basis as the p-fold wedges of the ω_i is the statement

$$\Omega^p_X(\log D) = \Lambda^p\Omega^1_X(\log D),$$

completing the proof.

The rank is the same as that of Ω^p_X, and indeed $\Omega^p_X \subseteq \Omega^p_X(\log D)$ is an inclusion of locally free sheaves of the same rank; the two differ only along D, where the second is larger. The gain is already present in the smallest case.

Example 7.2.7: Logarithmic Forms on the Projective Line

Let $X = \mathbb{C}\mathrm{P}^1$, let $D = \{0\} + \{\infty\}$ and let $U = \mathbb{C}^*$. In the affine coordinate z the divisor is $\{z = 0\}$ near the origin, so $\Omega^1_X(\log D)$ is generated there by dz/z by proposition 7.2.6. In the other chart $w = 1/z$ the divisor is $\{w = 0\}$ and

$$\frac{dz}{z} \;=\; \frac{d(1/w)}{1/w} \;=\; -\frac{dw}{w}$$

so the same form generates there too. Hence dz/z is a global nowhere-vanishing section and

$$\Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D) \;\cong\; \mathcal{O}_{\mathbb{C}\mathrm{P}^1}$$

In particular $H^0(\mathbb{C}\mathrm{P}^1, \Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D)) = \mathbb{C} \cdot \frac{dz}{z}$ is one-dimensional, whereas $H^0(\mathbb{C}\mathrm{P}^1, \Omega^1_{\mathbb{C}\mathrm{P}^1}) = 0$ by example 1.1.4: allowing logarithmic poles at two points has manufactured a global holomorphic 1-form where there was none. That form is the source of the weight-2 class in section 7.4.

Where a logarithmic form differs from a holomorphic one is along D, and the measurement of the difference is its residue. In one variable the residue of $g(z)\,dz/z$ at the origin is $g(0)$, and the definition below is that formula made to work along a whole component of D, in every degree. The first step is to check that the recipe is well posed.

Lemma 7.2.8: Splitting Off One Component

Let D_1 be an irreducible component of D, let D' be the union of the remaining components, and let $D'_1 = D' \cap D_1$, a normal crossings divisor in the manifold D_1 with inclusion $\iota_1 \colon D_1 \hookrightarrow X$. Let $x \in D_1$ and let $h \in \mathcal{O}_{X,x}$ be a local equation for D_1 at x.

1. Every germ $\alpha \in \Omega^p_X(\log D)_x$ can be written
$$\alpha \;=\; \frac{dh}{h} \wedge \beta \;+\; \gamma, \qquad \beta \in \Omega^{p-1}_X(\log D')_x, \quad \gamma \in \Omega^p_X(\log D')_x$$
2. The pullback $\iota_1^*\beta \in \Omega^{p-1}_{D_1}(\log D'_1)_x$ depends only on α, and not on h or on the choice of decomposition.

***Proof*:**

Choose a chart adapted to D at x with $D \cap V = \{z_1 \cdots z_k = 0\}$ and $D_1 \cap V = \{z_1 = 0\}$, which is possible because the components of D are smooth. In it $D' \cap V = \{z_2 \cdots z_k = 0\}$, and ι_1 identifies $D_1 \cap V$ with the polydisc in $z_2, \ldots, z_n$ carrying the normal crossings divisor $\{z_2 \cdots z_k = 0\}$. Pullback along ι_1 kills dz_1 and restricts coefficients, so by proposition 7.2.6 it carries $\Omega^q_X(\log D')_x$ into $\Omega^q_{D_1}(\log D'_1)_x$.

Step 1: existence. By proposition 7.2.6 the germ α is an $\mathcal{O}$-combination of the ω_S. Splitting the sum according to whether $1 \in S$ and factoring ω_1 out of the first part gives $\alpha = \omega_1 \wedge \beta + \gamma$ with β, γ combinations of the ω_T with $1 \notin T$, hence sections of $\Omega^\bullet_X(\log D')$. This is the required decomposition for $h = z_1$. A general local equation for D_1 is $h = uz_1$ with u a unit, and then

$dh/h = \omega_1 + du/u$ with du/u holomorphic, so

$$\alpha \;=\; \frac{dh}{h}\wedge\beta \;+\; \Big(\gamma - \frac{du}{u}\wedge\beta\Big)$$

is a decomposition for h, with the same β.

Step 2: uniqueness for fixed h. Suppose $\frac{dh}{h}\wedge\beta_1+\gamma_1 = \frac{dh}{h}\wedge\beta_2+\gamma_2$. Put $\rho=\beta_1-\beta_2$ and $\sigma=\gamma_2-\gamma_1$, both sections of $\Omega^\bullet_X(\log D')$, so that $dh\wedge\rho = h\,\sigma$. With $h=uz_1$ this reads $u\,dz_1\wedge\rho + z_1\,du\wedge\rho = uz_1\sigma$, that is,

$$dz_1\wedge\rho \;=\; z_1\,\tau, \qquad \tau \;=\; \sigma - u^{-1}\,du\wedge\rho$$

and τ is again a section of $\Omega^p_X(\log D')$. Write $\rho = dz_1\wedge\rho' + \rho''$ with ρ',ρ'' free of dz_1 in the frame of proposition 7.2.6; then $dz_1\wedge\rho'' = z_1\tau$. The coefficients of ρ'' and of τ in that frame are holomorphic, and the coefficients of $dz_1\wedge\rho''$ are those of ρ'' up to sign, so z_1 divides every coefficient of ρ''. Hence $\iota_1^*\rho''=0$, while $\iota_1^*(dz_1\wedge\rho')=0$ because $\iota_1^*dz_1=0$. Therefore $\iota_1^*\rho=0$, that is, $\iota_1^*\beta_1=\iota_1^*\beta_2$.

Step 3: independence of h. Let $h'=vh$ with v a unit be another local equation. Step 1 produced from any h-decomposition an h'-decomposition with the same β, and Step 2 applied to h' says that every h'-decomposition has the same $\iota_1^*\beta$. So the two local equations give the same answer, completing the proof.

Definition 7.2.9: The Residue Along a Component

In the situation of lemma 7.2.8, the *residue along* D_1 is the $\mathcal{O}_X$-linear map of sheaves

$$\mathrm{Res}_{D_1}\colon \Omega^p_X(\log D) \;\to\; \iota_{1*}\Omega^{p-1}_{D_1}(\log D_1'), \qquad \mathrm{Res}_{D_1}(\alpha) \;=\; \iota_1^*\beta$$

where $\alpha = \frac{dh}{h}\wedge\beta+\gamma$ is any decomposition as in that lemma. In an adapted chart with $D_1=\{z_1=0\}$ it is given on the frame of proposition 7.2.6 by

$$\mathrm{Res}_{D_1}(\omega_S) \;=\; \begin{cases} \iota_1^*\omega_{S\setminus 1}, & 1\in S \\ 0, & 1\notin S \end{cases}$$

No sign has been inserted into Res_{D_1}. On the frame it is literally $\omega_S\mapsto\iota_1^*\omega_{S\setminus 1}$, so that in one variable $\mathrm{Res}_{\{0\}}(dz/z)=1$, and the anticommutation with d recorded in theorem 7.2.11 below is absorbed into the differential of the shifted target rather than into the map. That is the normalization used throughout this section, and it is the one that matches the tube residue wherever the two can be compared (box 7.2.10). A residue renormalized by a sign depending on the degree of the form has the same kernel and the same image, hence yields the same exact sequence, but it is a different map, and any comparison with the residue defined here has to carry that sign explicitly; the graded residue Res^m of proposition 7.3.6 carries it at $m=1$.

The name is not an analogy. Away from the points where two components of D meet, the residue that definition 5.4.6 attached to a class on the complement of a hypersurface, by integrating over a tube, computes exactly this.

7.2.10 Note: The Algebraic and the Topological Residue Proposition 5.4.7 takes a closed form η on the complement of a smooth hypersurface $X \subseteq \mathbb{C}\mathrm{P}^{n+1}$ that can be written locally as $\frac{dg_\alpha}{g_\alpha} \wedge \psi_\alpha + \theta_\alpha$ with g_α a defining function and $\psi_\alpha, \theta_\alpha$ holomorphic, and concludes that the ψ_α glue to a form ψ on X with $\operatorname{Res}\eta = [\psi]$, where Res is the map dual to the tube map (definition 5.4.6). That local shape is precisely membership in $\Omega^{k+1}(\log X)$, by definition 7.2.5 in one direction and lemma 7.2.8 in the other, and ψ is precisely $\operatorname{Res}_X\eta$ in the sense of definition 7.2.9: the gluing statement there is the well-definedness proved above. So the topological residue of the class of a logarithmic form is the class of its logarithmic residue. The normalization agrees as well, the factor $1/2\pi i$ having been built into definition 5.4.6 for exactly this reason. Nothing in the argument of proposition 5.4.7 uses more about X than that it is a smooth divisor carrying a tubular neighbourhood, so the same computation identifies Res_{D_1} with the tube residue along a component D_1 meeting no other component of D, and in particular along every component when D is smooth, which is the case of theorem 7.2.12. It goes no further than that. Where D_1 meets another component the hypothesis of proposition 5.4.7 already fails: by lemma 7.2.8 the β in $\alpha = \frac{dh}{h} \wedge \beta + \gamma$ is only logarithmic along $D_1' = D' \cap D_1$, not holomorphic. Both sides of the proposed identity break down there as well. The value $\operatorname{Res}_{D_1}\alpha$ is a section of $\Omega^{\bullet-1}_{D_1}(\log D_1')$, not a holomorphic form on D_1, so it represents no class in $H^\bullet(D_1, \mathbb{C})$; and the tube over a cycle in D_1 passing through a point of D_1' is not contained in U, so the tube map of definition 5.4.6 is not defined on that cycle. Exercise 7.2.1 (6) is the smallest instance. There D is the union of the four coordinate curves in a product of two projective lines, and $\operatorname{Res}_{D_1}\left(\frac{dz}{z} \wedge \frac{dw}{w}\right) = \iota_1^* \frac{dw}{w}$ along $D_1 = \{0\} \times \mathbb{C}\mathrm{P}^1$ is a nonzero logarithmic form, whereas $H^1(D_1, \mathbb{C}) = H^1(\mathbb{C}\mathrm{P}^1, \mathbb{C}) = 0$ leaves no class for a tube residue to be. Recovering a topological reading of the residue at a crossing is what the weight filtration of section 7.3 is for.

The residue is more than a measurement: it is a surjection with a computable kernel, and that is what turns the logarithmic complex into an object one can induct on. Peeling off one component at a time is the mechanism behind every computation in the rest of the section.

Theorem 7.2.11: The Residue Sequence Along One Component

Let D_1 be an irreducible component of D, let D' be the union of the others and $D_1' = D' \cap D_1$. Then for every p the sequence of sheaves

$$0 \;\to\; \Omega^p_X(\log D') \;\to\; \Omega^p_X(\log D) \xrightarrow{\operatorname{Res}_{D_1}} \iota_{1*}\Omega^{p-1}_{D_1}(\log D_1') \;\to\; 0$$

is exact, the first map being the inclusion. Moreover $\operatorname{Res}_{D_1} \circ d = -\,d \circ \operatorname{Res}_{D_1}$, so that with the sign convention that the shifted complex $\iota_{1*}\Omega^{\bullet-1}_{D_1}(\log D_1')$ carries the differential $-d$, the display is a short exact sequence of complexes of sheaves of $\mathbb{C}$-vector spaces.

***Proof*:**

Exactness is local, so fix $x \in X$. If $x \notin D_1$ the third sheaf has zero stalk and the first map is the identity, since D and D' agree near x. So assume $x \in D_1$ and work in a chart adapted to D with $D_1 = \{z_1 = 0\}$ and $D' = \{z_2 \cdots z_k = 0\}$, keeping the notation of proposition 7.2.6.

Step 1: the composite vanishes and the kernel is no larger. A section of $\Omega^p_X(\log D')$ admits the decomposition of lemma 7.2.8 with $\beta = 0$, so its residue vanishes. Conversely take the

decomposition $\alpha = \omega_1 \wedge \beta + \gamma$ produced by the frame, in which β and γ are $\mathcal{O}$-combinations of the ω_T with $T \subseteq \{2, \dots, n\}$, and suppose $\iota_1^* \beta = 0$. Since no ω_T occurring in β involves dz_1, that vanishing says exactly that every coefficient of β vanishes on $\{z_1 = 0\}$, hence is divisible by z_1; write $\beta = z_1 \widetilde{\beta}$ with $\widetilde{\beta}$ again a section of $\Omega_X^{p-1}(\log D')$. Then

$$\omega_1 \wedge \beta \;=\; \frac{dz_1}{z_1} \wedge z_1 \widetilde{\beta} \;=\; dz_1 \wedge \widetilde{\beta}$$

which is a section of $\Omega_X^p(\log D')$, and hence so is α.

Step 2: surjectivity. Let β_0 be a germ of $\Omega_{D_1}^{p-1}(\log D_1')$ at x. In the chart, D_1 is the polydisc in $z_2, \dots, z_n$ and $\Omega_{D_1}^{p-1}(\log D_1')$ is free on the $\iota_1^* \omega_T$ with $T \subseteq \{2, \dots, n\}$, again by proposition 7.2.6. Write $\beta_0 = \sum_T a_T \, \iota_1^* \omega_T$ and let $\widetilde{a}_T$ be the function on the chart obtained by regarding a_T as independent of z_1. Then $\alpha = \omega_1 \wedge \sum_T \widetilde{a}_T \omega_T$ is a section of $\Omega_X^p(\log D)$ with $\operatorname{Res}_{D_1} \alpha = \beta_0$.

Step 3: the sign. Decompose $\alpha = \frac{dh}{h} \wedge \beta + \gamma$ as in lemma 7.2.8. Since $d(dh/h) = 0$,

$$d\alpha \;=\; -\frac{dh}{h} \wedge d\beta \;+\; d\gamma$$

and $d\beta$, $d\gamma$ are again sections of $\Omega_X^\bullet(\log D')$. So $\operatorname{Res}_{D_1}(d\alpha) = -\iota_1^*(d\beta) = -\,d(\iota_1^* \beta) = -\,d \operatorname{Res}_{D_1}(\alpha)$, pullback commuting with d. With the differential of the shifted target taken to be $-d$, the residue commutes with the differentials, completing the proof.

Theorem 7.2.12: The Residue Sequence of a Smooth Boundary

Suppose D is smooth, that is, its components are pairwise disjoint, and write $\iota \colon D \hookrightarrow X$. Then

$$0 \;\to\; \Omega_X^\bullet \;\to\; \Omega_X^\bullet(\log D) \;\xrightarrow{\;\operatorname{Res}\;}\; \iota_* \Omega_D^{\bullet - 1} \;\to\; 0$$

is exact, where Res is the sum of the residues along the components.

***Proof*:**
Exactness is local and each point of D lies on exactly one component, so it is enough to treat a single component D_1. For it, $D' = \varnothing$ near every point of D_1, so $\Omega_X^\bullet(\log D') = \Omega_X^\bullet$ and $D_1' = \varnothing$; theorem 7.2.11 is then exactly the displayed sequence, as we sought to show.

Theorem 7.2.12 is the shape in which the residue sequence is usually quoted. It is the special case of theorem 7.2.11 in which the components of D are disjoint, not the general statement.

7.2.13 Warning: The Total Residue Is Not Surjective at a Crossing When components of D meet, the map $\bigoplus_i \mathrm{Res}_{D_i}$ from $\Omega^p_X(\log D)$ to $\bigoplus_i \iota_{i*}\Omega^{p-1}_{D_i}(\log D_i')$ is not surjective for $p \geq 2$. Take $X = \mathbb{C}^2$, $D = \{z_1 z_2 = 0\}$ and $p = 2$. Then $\Omega^2_X(\log D)$ is free on $\omega_1 \wedge \omega_2$, so a section is $c\,\omega_1 \wedge \omega_2$ with c holomorphic, and

$$\mathrm{Res}_{D_1}(c\,\omega_1 \wedge \omega_2) = c|_{D_1}\,\iota_1^*\omega_2, \qquad \mathrm{Res}_{D_2}(c\,\omega_1 \wedge \omega_2) = -\,c|_{D_2}\,\iota_2^*\omega_1$$

the second because $\omega_1 \wedge \omega_2 = -\,\omega_2 \wedge \omega_1$. The pair $(\iota_1^*\omega_2, 0)$ is therefore not in the image: it would need $c|_{D_1} = 1$ and $c|_{D_2} = 0$, giving $c(0) = 1$ and $c(0) = 0$ at once. Recovering the general statement means filtering $\Omega^\bullet_X(\log D)$ by the number of branches along which a form has a pole, which is the weight filtration of section 7.3.

What remains is the comparison theorem: the complex just built, which lives on the compact X, computes the cohomology of the open U. Both sides of that statement are the cohomology of a complex of sheaves rather than of a single sheaf, so the first thing to fix is what such cohomology means.

7.2.14 Convention: Cohomology of a Complex of Sheaves For a bounded-below complex of sheaves of $\mathbb{C}$-vector spaces $\mathcal{K}^\bullet$ on a space X, write $H^k(X, \mathcal{K}^\bullet)$ for the k-th hypercohomology $\mathbb{R}^k\Gamma(X, \mathcal{K}^\bullet)$, constructed and shown independent of choices in [7, Hypercohomology, Hypercohomology is Well Defined]. For a single sheaf placed in degree 0 this is ordinary sheaf cohomology, so the notation extends the usual one. Two properties are used below, both proved there.

1. A *quasi-isomorphism* $\mathcal{K}^\bullet \to \mathcal{L}^\bullet$, meaning a morphism of complexes inducing an isomorphism on the cohomology of the stalk complex at every point, induces isomorphisms $H^k(X, \mathcal{K}^\bullet) \cong H^k(X, \mathcal{L}^\bullet)$ for all k.

2. There is a spectral sequence

$$E_1^{p,q} \;=\; H^q(X, \mathcal{K}^p) \;\Longrightarrow\; H^{p+q}(X, \mathcal{K}^\bullet)$$

whose differential d_1 is induced by the differential of $\mathcal{K}^\bullet$.

If every $\mathcal{K}^p$ is acyclic then (2) is concentrated in the row $q = 0$, where it reads $E_1^{p,0} = \Gamma(X, \mathcal{K}^p)$, and the sequence degenerates at E_2 for want of room; so

$$H^k(X, \mathcal{K}^\bullet) \;\cong\; H^k\big(\Gamma(X, \mathcal{K}^\bullet)\big)$$

This is the abstract de Rham theorem (theorem 1.1.2) for complexes.

The comparison is proved by computing both sides locally, and on the holomorphic side the local computation begins with the case of no boundary at all. The ordinary Poincaré lemma has a holomorphic refinement, for the reason that the homotopy operator producing the primitive is built from the radial contraction $z \mapsto tz$, which is holomorphic in z.

The holomorphic de Rham complex resolves the constant sheaf: on a complex manifold a closed holomorphic form is locally exact, so $0 \to \underline{\mathbb{C}}_X \to \mathcal{O}_X \to \Omega^1_X \to \cdots$ is a resolution and $\mathbb{H}^i(X, \Omega^\bullet_X) \cong H^i(X, \mathbb{C})$. This is the holomorphic Poincaré lemma with the holomorphic de Rham resolution it

gives.[2] It belongs to general complex geometry and is used here only through its conclusion.

This is the case $D = \varnothing$ of the theorem being aimed at. The general case needs the local cohomology of the logarithmic complex, and the residue sequence computes it by induction on the number of branches: each branch contributes one new closed form ω_i, and nothing else.

Lemma 7.2.15: Local Cohomology of the Logarithmic Complex

Let $V = \Delta^n$ and $D_V = \{z_1 \cdots z_k = 0\}$ with $0 \le k \le n$. Then the cohomology of the complex $\Omega^\bullet_V(\log D_V)(V)$ of global sections is

$$H^q\big(\Omega^\bullet_V(\log D_V)(V)\big) \;=\; \bigoplus_{S\subseteq\{1,\dots,k\},\ |S|=q} \mathbb{C}\,[\omega_S]$$

In particular it is finite-dimensional of dimension $\binom{k}{q}$, and the exterior algebra on the classes of $\omega_1, \dots, \omega_k$ maps isomorphically onto it.

***Proof*:**
Induct on k. For $k = 0$ the complex is $\Omega^\bullet(V)$ and the assertion is the holomorphic Poincaré lemma ([4, the holomorphic Poincaré lemma]).

Step 1: the residue sequence on global sections. Let $k \ge 1$, put $D_1 = \{z_1 = 0\}$ and $D' = \{z_2 \cdots z_k = 0\}$, so $D'_1 = D' \cap D_1$. By theorem 7.2.11 the sequence of sheaves is exact; on global sections over the polydisc the only point in question is surjectivity of Res_{D_1}, and Step 2 of that proof produces the lift globally: writing $\beta_0 = \sum_T a_T\, \iota_1^*\omega_T$ with a_T holomorphic on $D_1 = \Delta^{n-1}$ and letting $\widetilde{a}_T$ be the same function viewed as independent of z_1, the global section $\omega_1 \wedge \sum_T \widetilde{a}_T\omega_T$ has residue β_0. So

$$0 \to \Omega^\bullet_V(\log D')(V) \to \Omega^\bullet_V(\log D_V)(V) \xrightarrow{\mathrm{Res}_{D_1}} \Omega^{\bullet-1}_{D_1}(\log D'_1)(D_1) \to 0$$

is a short exact sequence of complexes of $\mathbb{C}$-vector spaces, the third carrying the differential $-d$.

Step 2: the induction hypothesis on the two ends. The pair (V, D') is the model with $k-1$ branches in n variables, and the pair (D_1, D'_1) is the model with $k-1$ branches in $n-1$ variables, since $D_1 = \Delta^{n-1}$ with coordinates $z_2, \dots, z_n$ and $D'_1 = \{z_2 \cdots z_k = 0\}$. So by induction the first complex has cohomology with basis the $[\omega_S]$ for $S \subseteq \{2, \dots, k\}$, $|S| = q$, and the third, whose q-th cohomology is the $(q-1)$-st cohomology of $\Omega^\bullet_{D_1}(\log D'_1)(D_1)$ because changing the sign of a differential changes neither kernels nor images, has basis the $[\iota_1^*\omega_S]$ for $S \subseteq \{2, \dots, k\}$, $|S| = q-1$.

Step 3: the connecting maps vanish. Let $S \subseteq \{2, \dots, k\}$ with $|S| = q-1$. The form $\omega_1 \wedge \omega_S$ is a global section of $\Omega^q_V(\log D_V)$, it is closed because each ω_i is, and $\mathrm{Res}_{D_1}(\omega_1 \wedge \omega_S) = \iota_1^*\omega_S$ by definition 7.2.9. So the map induced by Res_{D_1} on cohomology sends a basis to a basis and is surjective in every degree. In the long exact sequence of the short exact sequence of Step 1, a surjective residue map forces every connecting homomorphism to vanish.

[2] [4, The Holomorphic Poincaré Lemma, The Holomorphic de Rham Resolution].

Step 4: assembly. The long exact sequence therefore breaks into short exact sequences

$$0 \to H^q\big(\Omega_V^\bullet(\log D')(V)\big) \to H^q\big(\Omega_V^\bullet(\log D_V)(V)\big) \to H^{q-1}\big(\Omega_{D_1}^\bullet(\log D_1')(D_1)\big) \to 0$$

The classes $[\omega_S]$ with $1 \notin S$, $|S| = q$, form a basis of the subspace by Step 2, and the classes $[\omega_1 \wedge \omega_{S'}]$ with $S' \subseteq \{2, \dots, k\}$, $|S'| = q-1$, map to a basis of the quotient by Step 3. Together they are a basis of the middle term, and they are exactly the $[\omega_S]$ with $S \subseteq \{1, \dots, k\}$ and $|S| = q$, completing the induction and the proof.

The answer is the cohomology of a k-torus, which is what it must be: the complement of the local model is $(\Delta^*)^k \times \Delta^{n-k}$, homotopy equivalent to $(S^1)^k$. The next proposition upgrades that coincidence into a map.

Proposition 7.2.16: The Logarithmic Complex Is Quasi-Isomorphic to the Push-forward

Let $\mathcal{A}_U^p$ denote the sheaf of smooth complex-valued p-forms on U. Restriction of forms defines a morphism of complexes of sheaves on X,

$$\Omega_X^\bullet(\log D) \;\to\; j_*\mathcal{A}_U^\bullet$$

and it is a quasi-isomorphism.

***Proof*:**
A section of $\Omega_X^p(\log D)$ over V is by definition 7.2.5 a holomorphic p-form on $V \cap U$, hence in particular a smooth one, so the map is defined; it commutes with d because it is a restriction. What must be shown is that it induces an isomorphism on the cohomology of every stalk complex.

Fix $x \in X$ and let $k = k(x)$ be the number of branches of D at x, with $k = 0$ when $x \notin D$. Choose an adapted chart at x and let $V_\varepsilon \cong \Delta_\varepsilon^n$ be the concentric adapted polydiscs, which form a fundamental system of neighbourhoods of x. Cohomology of a complex of vector spaces commutes with filtered colimits, so the cohomology of either stalk complex is the colimit over ε of the cohomology of the complex of sections over V_ε.

Step 1: the holomorphic side. By lemma 7.2.15 the cohomology of $\Omega_X^\bullet(\log D)(V_\varepsilon)$ has basis the classes $[\omega_S]$ with $S \subseteq \{1, \dots, k\}$ and $|S| = q$. The restriction maps between two such polydiscs carry ω_S to ω_S, hence carry basis to basis and are isomorphisms, so the colimit is again $\bigoplus_{|S|=q} \mathbb{C}\,[\omega_S]$.

Step 2: the smooth side. By definition $\Gamma(V_\varepsilon, j_*\mathcal{A}_U^\bullet) = \mathcal{A}^\bullet(V_\varepsilon \cap U)$, whose cohomology is $H^q(V_\varepsilon \cap U, \mathbb{C})$, by theorem 1.1.2 applied to the fine resolution of $\underline{\mathbb{C}}$ by smooth forms. In the adapted chart $V_\varepsilon \cap U = (\Delta_\varepsilon^*)^k \times \Delta_\varepsilon^{n-k}$, which deformation retracts onto the torus $(S^1)^k$. Homotopy invariance and the Künneth formula [30, Künneth Formula for Cohomology] give

$$H^\bullet(V_\varepsilon \cap U, \mathbb{C}) \;\cong\; \bigotimes_{i=1}^{k} H^\bullet(\Delta_\varepsilon^*, \mathbb{C})$$

one tensor factor for each punctured disc. On the i-th factor, $H^1(\Delta_\varepsilon^*, \mathbb{C})$ is one-dimensional and

generated by the class of $\frac{1}{2\pi i}\frac{dz_i}{z_i}$, since that form integrates to 1 over a positively oriented circle. Hence $H^q(V_\varepsilon \cap U, \mathbb{C})$ has basis the classes of $(2\pi i)^{-|S|}\omega_S$ for $S \subseteq \{1, \dots, k\}$, $|S| = q$, and again the restriction maps are isomorphisms.

Step 3: comparison. The map sends the class of ω_S on the holomorphic side to the de Rham class of the same form, which is $(2\pi i)^{|S|}$ times the basis element of Step 2. It therefore carries a basis to a basis and is an isomorphism in every degree, for every x, completing the proof.

That quasi-isomorphism is the last ingredient. It moves the computation onto the smooth side, where the sheaves are acyclic and the hypercohomology collapses to the de Rham complex of U, which is the comparison theorem this section was aimed at.

Theorem 7.2.17: The Logarithmic Complex Computes the Cohomology of U

Let X be a complex manifold, $D \subseteq X$ a normal crossings divisor and $U = X \setminus D$. Restriction of forms induces isomorphisms

$$H^k\big(X, \Omega_X^\bullet(\log D)\big) \;\cong\; H^k(U, \mathbb{C}) \qquad \text{for all } k$$

***Proof*:**
Step 1: the sheaves $j_*\mathcal{A}_U^p$ are acyclic. A smooth function on an open $V \subseteq X$ restricts to a smooth function on $V \cap U$, so $j_*\mathcal{A}_U^p$ is a sheaf of $\mathcal{A}_X^0$-modules. The argument that makes the smooth form sheaves fine (theorem 1.1.2(3)) uses nothing about the module beyond this: a smooth partition of unity on X subordinate to a locally finite cover acts on the sheaf by multiplication, giving endomorphisms with the required supports that sum to the identity, so $j_*\mathcal{A}_U^p$ is fine in the sense defined before theorem 1.1.2. Since X is paracompact, theorem 1.1.2(1) makes it acyclic.

Step 2: the smooth complex computes $H^\bullet(U, \mathbb{C})$. By Step 1 and the last clause of box 7.2.14,

$$H^k\big(X, j_*\mathcal{A}_U^\bullet\big) \;\cong\; H^k\big(\Gamma(X, j_*\mathcal{A}_U^\bullet)\big) \;=\; H^k\big(\mathcal{A}^\bullet(U)\big)$$

the last equality being the definition of the pushforward. The de Rham complex of U is the complex of global sections of the fine resolution of $\underline{\mathbb{C}}_U$ by smooth forms, so its cohomology is $H^k(U, \mathbb{C})$ by theorem 1.1.2.

Step 3: transport along the quasi-isomorphism. By proposition 7.2.16 the restriction map $\Omega_X^\bullet(\log D) \to j_*\mathcal{A}_U^\bullet$ is a quasi-isomorphism, so property (1) of box 7.2.14 gives $H^k(X, \Omega_X^\bullet(\log D)) \cong H^k(X, j_*\mathcal{A}_U^\bullet)$. Composing with Step 2 gives the theorem, and the composite is induced by restriction of forms, completing the proof.

Every input to that proof is internal to this book except one: the existence of hypercohomology and its first spectral sequence, imported from [7, Hypercohomology, The Two Hypercohomology Spectral Sequences] in box 7.2.14. The local computation on which everything rests, lemma 7.2.15, is proved here in full.

What the theorem gives is a change of ambient space. The cohomology of the noncompact U, which carries no Hodge theory of its own, has been rewritten as the cohomology of a complex of locally

free sheaves on the compact X, where Part I applies. Two filtrations of that complex, one by the order of the pole and one by the holomorphic degree, will produce the weight and Hodge filtrations of a mixed Hodge structure; the spectral sequence of box 7.2.14 is already the second of them. The smallest case displays the whole mechanism.

Example 7.2.18: The Punctured Line by the Logarithmic Complex
Let $X = \mathbb{C}\mathrm{P}^1$, $D = \{0\} + \{\infty\}$ and $U = \mathbb{C}^*$, as in example 7.2.7. The logarithmic complex is

$$\mathcal{O}_{\mathbb{C}\mathrm{P}^1} \xrightarrow{d} \Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D) \cong \mathcal{O}_{\mathbb{C}\mathrm{P}^1}$$

concentrated in degrees 0 and 1. Its hypercohomology is computed by the spectral sequence of box 7.2.14, whose first page is

$$E_1^{p,q} \;=\; H^q\big(\mathbb{C}\mathrm{P}^1, \Omega^p_{\mathbb{C}\mathrm{P}^1}(\log D)\big)$$

Both terms are $\mathcal{O}_{\mathbb{C}\mathrm{P}^1}$, and $H^0(\mathbb{C}\mathrm{P}^1, \mathcal{O}) = \mathbb{C}$ while $H^1(\mathbb{C}\mathrm{P}^1, \mathcal{O}) = 0$ by example 1.1.4. So the first page is concentrated in the row $q = 0$, where it reads

$$\mathbb{C} \xrightarrow{d_1} \mathbb{C}, \qquad d_1(c) \;=\; dc \;=\; 0$$

since the global sections of $\mathcal{O}_{\mathbb{C}\mathrm{P}^1}$ are the constants. Hence $E_2 = E_\infty = E_1$ and, by theorem 7.2.17,

$$H^0(\mathbb{C}^*, \mathbb{C}) = \mathbb{C}, \qquad H^1(\mathbb{C}^*, \mathbb{C}) = \mathbb{C} \cdot \left[\frac{dz}{z}\right]$$

which is the cohomology of a circle, as it must be. The generator of H^1 is the logarithmic form of example 7.2.7, and its residue at 0 is $\mathrm{Res}_{\{0\}}(dz/z) = 1$ by definition 7.2.9. A single boundary point has contributed a single class, detected by its residue; the same bookkeeping on a punctured elliptic curve is section 7.4.

Exercise 7.2.1

1. On $X = \Delta^2$ with $D = \{z_1 = 0\}$, consider $\alpha = dz_2/z_1$. Show that $z_1\alpha$ is holomorphic but that α is not a section of $\Omega^1_X(\log D)$, identifying which half of definition 7.2.5 fails. Confirm the answer against proposition 7.2.6.

2. Let $D = \{xy(x + y) = 0\} \subseteq \mathbb{C}^2$. Show that D is not a normal crossings divisor at the origin. Then let $\pi\colon \widetilde{X} \to \mathbb{C}^2$ be the blow-up at the origin and show that the reduced preimage $\pi^{-1}(D)$, that is, the union of the exceptional curve with the three strict transforms, is a simple normal crossings divisor.

3. Let $\Delta \subseteq \mathbb{C}$ be the unit disc and $D = \{0\}$. Compute the cohomology of the two-term complex $\Omega^\bullet_\Delta(\log D)(\Delta)$ directly from power series, and check the answer against lemma 7.2.15 and against $H^\bullet(\Delta^*, \mathbb{C})$.

4. Let D be a normal crossings divisor on a complex manifold X of dimension n. Show that $\Omega^n_X(\log D) \cong K_X \otimes \mathcal{O}_X(D)$, where K_X is the canonical bundle and $\mathcal{O}_X(D)$ is the line bundle of the divisor D.[3] Deduce the isomorphism of example 7.2.7 a second way.

[3] [5, Canonical Bundle, Line Bundle of a Divisor].

5. Let $X = \mathbb{C}^2$, $D = \{z_1 z_2 = 0\}$, $D_1 = \{z_1 = 0\}$ and $D_2 = \{z_2 = 0\}$. Compute Res_{D_1} on the frames of $\Omega^1_X(\log D)$ and $\Omega^2_X(\log D)$, identify the kernel of Res_{D_1} in degree 1 with $\Omega^1_X(\log D_2)$, and verify the sign of theorem 7.2.11 on the form $\alpha = g\,\omega_1$ with g holomorphic.

6. Let $X = \mathbb{CP}^1 \times \mathbb{CP}^1$ and let D be the union of the four curves $\{0\} \times \mathbb{CP}^1$, $\{\infty\} \times \mathbb{CP}^1$, $\mathbb{CP}^1 \times \{0\}$ and $\mathbb{CP}^1 \times \{\infty\}$, so that $U = (\mathbb{C}^*)^2$. Show that D is a simple normal crossings divisor and that $\Omega^p_X(\log D) \cong \mathcal{O}_X^{\oplus\binom{2}{p}}$. Then use theorem 7.2.17 and the spectral sequence of box 7.2.14 to compute $H^k((\mathbb{C}^*)^2, \mathbb{C})$, taking as given that $H^q(X, \mathcal{O}_X) = 0$ for $q \geq 1$.

7.3 Deligne's Mixed Hodge Structure

The logarithmic complex of section 7.2 computes $H^*(U, \mathbb{C})$, and it carries more structure than the comparison isomorphism uses. Its terms are graded by how many logarithmic poles a form is allowed, and, like any complex of sheaves of differential forms, it is filtered by form degree. This section writes down the two filtrations, transports them to $H^k(U, \mathbb{Q})$, and proves that what they produce there is a mixed Hodge structure in the sense of definition 7.1.1.

Throughout, X is a smooth projective variety over $\mathbb{C}$ of dimension n, and $D = D_1 \cup \cdots \cup D_N \subseteq X$ is a normal crossings divisor whose irreducible components D_i are smooth (definition 7.2.1); by theorem 7.2.3 every quasi-projective U admits a compactification of this form. Write $U = X \setminus D$ and $j\colon U \hookrightarrow X$ for the inclusion. A point $x \in X$ lying on exactly r of the components has a coordinate neighbourhood with coordinates $z_1, \ldots, z_n$ in which $D = \{z_1 \cdots z_r = 0\}$; coordinates of this kind are called *adapted* to D at x, and every local computation below is carried out in them.

The Hodge decomposition fails on U because U is not compact, and the logarithmic poles are the exact measure of that failure: a form on X with no poles at all restricts to a form on U that extends across the boundary, while a form with m logarithmic poles remembers m directions in which U runs off to D. Counting poles therefore produces a filtration whose bottom piece is the cohomology X already had, and whose higher pieces are new. This is the filtration that will carry the weights.

Definition 7.3.1: The Weight Filtration on the Logarithmic Complex

The *weight filtration* on $\Omega^\bullet_X(\log D)$ is the increasing filtration by subsheaves

$$W_m\Omega^p_X(\log D) = \begin{cases} 0 & m < 0 \\ \Omega^{p-m}_X \wedge \Omega^m_X(\log D) & 0 \leq m \leq p \\ \Omega^p_X(\log D) & m > p \end{cases}$$

where the middle line denotes the image of $\Omega^{p-m}_X \otimes \Omega^m_X(\log D) \to \Omega^p_X(\log D)$ under the wedge product.

In adapted coordinates $\Omega^p_X(\log D)$ is free on the forms

$$\frac{dz_I}{z_I} \wedge dz_J := \frac{dz_{i_1}}{z_{i_1}} \wedge \cdots \wedge \frac{dz_{i_a}}{z_{i_a}} \wedge dz_{j_1} \wedge \cdots \wedge dz_{j_b}$$

with $I = \{i_1 < \cdots < i_a\} \subseteq \{1, \ldots, r\}$, $J = \{j_1 < \cdots < j_b\} \subseteq \{1, \ldots, n\} \setminus I$ and $a + b = p$ (definition 7.2.5). Comparing the two descriptions, $W_m\Omega^p_X(\log D)$ is the free submodule spanned

by those basis elements with $|I| \leq m$: a generator $\frac{dz_I}{z_I} \wedge dz_J$ with $|I| \leq m$ is obtained by taking $J' \subseteq J$ with $|J'| = m - |I|$ and factoring the element as $\left(dz_{J\setminus J'}\right) \wedge \left(\frac{dz_I}{z_I} \wedge dz_{J'}\right)$, the second factor lying in $\Omega^m_X(\log D)$, and conversely every product of an element of Ω^{p-m}_X with an element of $\Omega^m_X(\log D)$ has at most m logarithmic factors. So W_m is locally free, $W_0\Omega^\bullet_X(\log D) = \Omega^\bullet_X$, and $W_p\Omega^p_X(\log D) = \Omega^p_X(\log D)$.

Each W_m is a subcomplex. Indeed a section of $W_m\Omega^p_X(\log D)$ is locally a sum $\sum f_{IJ}\frac{dz_I}{z_I} \wedge dz_J$ with $|I| \leq m$, the basis forms are closed, and d of the sum is $\sum df_{IJ} \wedge \frac{dz_I}{z_I} \wedge dz_J$, which again has at most m logarithmic factors in each term.

The second filtration needs no new idea. On the de Rham complex of a compact Kähler manifold the filtration by form degree induces the Hodge filtration (chapter 2), and the same recipe applies verbatim to the logarithmic complex.

Definition 7.3.2: The Hodge Filtration on the Logarithmic Complex

The *Hodge filtration* on $\Omega^\bullet_X(\log D)$ is the decreasing filtration by the subcomplexes

$$F^p\Omega^\bullet_X(\log D) = \Omega^{\geq p}_X(\log D) = \left(0 \to \cdots \to 0 \to \Omega^p_X(\log D) \to \Omega^{p+1}_X(\log D) \to \cdots\right)$$

the truncation that discards the terms in degrees below p.

F^p is a subcomplex because d raises degree, and $F^p \supseteq F^{p+1}$ with F^0 the whole complex and $F^{n+1} = 0$. The two filtrations are of different natures and this is the point: W is defined on $\mathbb{Q}$-cohomology and grades it into pure pieces, while F lives only on $\mathbb{C}$-cohomology and gives the Hodge filtration of each piece. Neither alone is a Hodge structure.

Example 7.3.3: The Two Filtrations on $\mathbb{C}^*$

Take $X = \mathbb{C}\mathrm{P}^1$ with affine coordinate z, $D = \{0\} \cup \{\infty\}$ and $U = \mathbb{C}^*$. In the coordinate $w = 1/z$ near ∞ one has $\frac{dw}{w} = -\frac{dz}{z}$, so $\frac{dz}{z}$ is a global frame and $\Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D) \cong \mathcal{O}_{\mathbb{C}\mathrm{P}^1}$, while $\Omega^1_{\mathbb{C}\mathrm{P}^1} \cong \mathcal{O}_{\mathbb{C}\mathrm{P}^1}(-2)$.

The weight filtration is

$$W_{-1} = 0, \qquad W_0\Omega^\bullet_{\mathbb{C}\mathrm{P}^1}(\log D) = \Omega^\bullet_{\mathbb{C}\mathrm{P}^1}, \qquad W_1\Omega^\bullet_{\mathbb{C}\mathrm{P}^1}(\log D) = \Omega^\bullet_{\mathbb{C}\mathrm{P}^1}(\log D)$$

since $W_m\Omega^0 = \mathcal{O}_{\mathbb{C}\mathrm{P}^1}$ for every $m \geq 0$ and $W_1\Omega^1 = \Omega^1(\log D)$. The only graded piece above gr^W_0 is gr^W_1, the quotient $\Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D)/\Omega^1_{\mathbb{C}\mathrm{P}^1}$ placed in degree 1, which is a skyscraper of length 1 at each of 0 and ∞.

The Hodge filtration is $F^0 = \Omega^\bullet_{\mathbb{C}\mathrm{P}^1}(\log D)$, $F^1 = \Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D)$ in degree 1, and $F^2 = 0$. The group $H^1(\mathbb{C}^*, \mathbb{C})$ is one dimensional, spanned by the class of $\frac{dz}{z}$. That class lies in F^1, and it is not in the image of $H^1(\mathbb{C}\mathrm{P}^1, W_0\Omega^\bullet_{\mathbb{C}\mathrm{P}^1}(\log D)) = H^1(\mathbb{C}\mathrm{P}^1, \mathbb{C}) = 0$, so it is not reached by the bottom step of the weight filtration either. A class simultaneously as deep in F and as high in W as this one has Hodge type $(1,1)$ and weight 2, whereas the first cohomology of a compact Kähler manifold is pure of weight 1. Non-compactness has moved the weight.

The graded pieces of W are computed by residues, and the residue of a form with m logarithmic poles lives on the locus where m branches of D meet. Those loci are the objects the whole construction is written in terms of.

Definition 7.3.4: The Strata of a Normal Crossings Divisor

For $m \geq 1$ and a strictly increasing multi-index $I = \{i_1 < \cdots < i_m\} \subseteq \{1, \dots, N\}$ put $D_I = D_{i_1} \cap \cdots \cap D_{i_m}$, and set

$$D^{(m)} = \coprod_{|I|=m} D_I, \qquad D^{(0)} = X$$

the disjoint union of the m-fold intersections. Write $a_m \colon D^{(m)} \to X$ for the map that is the inclusion on each summand.

Because the components meet transversally, D_I is smooth of pure dimension $n - m$ or empty, so each $D^{(m)}$ is a smooth projective variety and a_m is a finite morphism. In particular $D^{(m)} = \emptyset$ for $m > \min(n, N)$, and each $D^{(m)}$ carries the pure Hodge structures of chapter 2.

Example 7.3.5: Three Lines in the Plane

Let $X = \mathbb{C}\mathrm{P}^2$ and let $D = L_1 \cup L_2 \cup L_3$ be three lines in general position, so no two are equal and the three do not share a point. Then $D^{(1)} = L_1 \sqcup L_2 \sqcup L_3$ is a disjoint union of three copies of $\mathbb{C}\mathrm{P}^1$, $D^{(2)} = \{L_1 \cap L_2\} \sqcup \{L_1 \cap L_3\} \sqcup \{L_2 \cap L_3\}$ is three reduced points, and $D^{(3)} = \emptyset$ because no point lies on all three lines. The complement $U = \mathbb{C}\mathrm{P}^2 \setminus D$ is $(\mathbb{C}^*)^2$.

The next statement is the computational core of the section: every graded piece of the weight filtration is the de Rham complex of a stratum, shifted.

Proposition 7.3.6: The Graded Pieces of the Weight Filtration

For every $m \geq 0$ there is an isomorphism of complexes of sheaves on X

$$\mathrm{Res}^m \colon \mathrm{gr}^W_m \Omega^\bullet_X(\log D) \xrightarrow{\sim} a_{m*} \Omega^{\bullet - m}_{D^{(m)}}$$

given in adapted coordinates, on a form of degree p, by

$$\mathrm{Res}^m \left(\tfrac{dz_I}{z_I} \wedge \eta \right) = (-1)^{mp} \, \eta \big|_{D_I}, \qquad |I| = m$$

and by 0 on the basis elements with $|I| < m$. For $m = 0$ it is the identity of $\Omega^\bullet_X$. For $m = 1$ it is, on the summand of a component D_i of D, the Poincaré residue Res_{D_i} of definition 7.2.9 multiplied by $(-1)^p$ in degree p; the sign is what converts the anticommutation $\mathrm{Res}_{D_i} \circ d = -\, d \circ \mathrm{Res}_{D_i}$ of theorem 7.2.11 into commutation with the unsigned differential carried by $a_{1*}\Omega^{\bullet - 1}_{D^{(1)}}$ here.

***Proof*:**

Step 1: the map is well defined. Fix $x \in X$ and adapted coordinates $z_1, \dots, z_n$ at x, labelled so that the branch of D_i through x is $\{z_i = 0\}$ for $1 \leq i \leq r$, and let I be an m-element subset of $\{1, \dots, r\}$. If $i \leq r$ and u is a unit then

$$\frac{d(u z_i)}{u z_i} = \frac{dz_i}{z_i} + \frac{du}{u}$$

and du/u is holomorphic. Two coordinate systems adapted to D at x differ, on each branch,

by such a unit together with a change of the remaining coordinates, so the basis form $\frac{dz_I}{z_I} \wedge \eta$ changes by a term with strictly fewer logarithmic factors, that is by an element of W_{m-1}, plus a form $\frac{dz_I}{z_I} \wedge \eta'$ with $\eta' \equiv \eta$ modulo the ideal of D_I and modulo dz_i for $i \in I$. Both ambiguities die on gr_m^W and after restriction to D_I, so the formula defines a morphism of sheaves independent of the coordinates. The labelling of the components of D fixes the ordering of I and hence the sign.

Step 2: it is an isomorphism. In adapted coordinates $\mathrm{gr}_m^W \Omega_X^p(\log D)$ is free over $\mathcal{O}_X$ on the classes of the basis elements $\frac{dz_I}{z_I} \wedge dz_J$ with $|I| = m$ exactly. For $i \in I$,

$$z_i \cdot \frac{dz_I}{z_I} \wedge dz_J = \pm \frac{dz_{I\setminus\{i\}}}{z_{I\setminus\{i\}}} \wedge dz_i \wedge dz_J$$

lies in W_{m-1}, so multiplication by the ideal of D_I annihilates the corresponding summand of gr_m^W, which is therefore a module over $\mathcal{O}_{D_I}$, free on the classes of $\frac{dz_I}{z_I} \wedge dz_J$ with $J \subseteq \{1, \dots, n\} \setminus I$ and $|J| = p - m$. The submanifold D_I is cut out by the equations $z_i = 0$ for $i \in I$, so the z_j with $j \notin I$ restrict to coordinates on it and restriction sends the dz_J to a basis of $\Omega_{D_I}^{p-m}$. Hence Res^m is an isomorphism onto $a_{m*}\Omega_{D^{(m)}}^{p-m}$ in every degree.

Step 3: it commutes with the differentials. For $\alpha = \frac{dz_I}{z_I} \wedge \eta$ of degree p with $|I| = m$, the basis forms $\frac{dz_i}{z_i}$ are closed, so $d\alpha = (-1)^m \frac{dz_I}{z_I} \wedge d\eta$, of degree $p+1$. Therefore

$$\mathrm{Res}^m(d\alpha) = (-1)^{m(p+1)}(-1)^m \, (d\eta)\big|_{D_I} = (-1)^{mp} \, d\big(\eta|_{D_I}\big) = d\big(\mathrm{Res}^m \alpha\big)$$

using $(-1)^{2m} = 1$ and the fact that restriction to a submanifold commutes with d. The sign $(-1)^{mp}$ in the definition exists precisely to absorb the $(-1)^m$ produced by moving d past the m logarithmic factors, and plays no further role.

Finally, for $m = 0$ the filtration step is $W_0 = \Omega_X^\bullet$ and the formula is the identity. For $m = 1$ the formula reads $\mathrm{Res}^1 = (-1)^p \mathrm{Res}_{D_i}$ in degree p on the summand of a component D_i: by definition 7.2.9 the map Res_{D_i} sends $\frac{dz_i}{z_i} \wedge \eta$ to $\eta|_{D_i}$ and kills the basis elements without $\frac{dz_i}{z_i}$, so it carries W_1 into $\iota_{i*}\Omega_{D_i}^{\bullet-1}$ and kills W_0. The two maps differ by a sign depending only on the degree, so they have the same kernel and the same image, and that sign is exactly what replaces the differential $-d$ that theorem 7.2.11 puts on the shifted target by the differential d used here. When D is smooth, W_1 is the whole logarithmic complex, so the exact sequence $0 \to W_0 \to W_1 \to \mathrm{gr}_1^W \to 0$ is the residue sequence $0 \to \Omega_X^\bullet \to \Omega_X^\bullet(\log D) \to a_{1*}\Omega_{D^{(1)}}^{\bullet-1} \to 0$ of theorem 7.2.12, re-signed in this way, completing the proof.

The proposition says that the logarithmic complex is assembled, in $N+1$ layers, out of the de Rham complexes of the strata $D^{(0)} = X$, $D^{(1)}$, $D^{(2)}$, and so on, each shifted into higher degree by its codimension. Every stratum is smooth and projective, so every layer contributes cohomology that is already understood, and the only question left is how the layers are glued. A spectral sequence answers exactly that question.

Both filtrations are filtrations of a complex of sheaves by subcomplexes, so each has a spectral sequence converging to the hypercohomology $H^k(X, \Omega_X^\bullet(\log D))$, which theorem 7.2.17 identifies with $H^k(U, \mathbb{C})$. The construction, the convergence criterion and the description of the first page are those of a filtered cochain complex, taken from *Homological Algebra*: see [7, The Exact Couple of a Filtered Complex] for the construction and [7, Convergence for a Decreasing Filtration of a Cochain Complex] for the convergence, exactly as in the proof of theorem 1.2.2. Both filtrations here are finite in each degree, so both spectral sequences converge.

The machinery is stated for decreasing filtrations, and W is increasing; feed it instead the decreasing filtration $p \mapsto W_{-p}$, which is the same data read backwards. With that convention the two spectral sequences are

$$ {}_F E_1^{p,q} = H^q\big(X, \Omega_X^p(\log D)\big) \Longrightarrow H^{p+q}(U, \mathbb{C}) \tag{7.7} $$

$$ {}_W E_1^{-m,\,k+m} = H^k\big(X, \operatorname{gr}_m^W \Omega_X^\bullet(\log D)\big) \Longrightarrow H^k(U, \mathbb{C}) \tag{7.8} $$

with differentials $d_r \colon E_r^{p,q} \to E_r^{p+r,\,q-r+1}$ in both cases. The abutment filtrations are ${}_F E_\infty^{p,q} = \operatorname{gr}_F^p H^{p+q}(U, \mathbb{C})$ and ${}_W E_\infty^{-m,k+m} = \operatorname{gr}_m^W H^k(U, \mathbb{C})$, where F and W on $H^k(U, \mathbb{C})$ denote the images of $H^k(X, F^p\Omega_X^\bullet(\log D))$ and of $H^k(X, W_m\Omega_X^\bullet(\log D))$ respectively. The indexing in (7.8) is chosen so that $p + q = k$ throughout, with $p = -m$ and $q = k + m$; the reason for grouping the terms by q rather than by k appears in proposition 7.3.7, where q turns out to be a weight.

Proposition 7.3.7: The First Page of the Weight Spectral Sequence

With the notation above,

$$ {}_W E_1^{-m,\,k+m} \cong H^{k-m}\big(D^{(m)}, \mathbb{C}\big) $$

for $0 \leq m$, and the term vanishes for $m < 0$ and for $m > k$. Under this isomorphism:

1. the filtration induced by F is $F^{p-m}H^{k-m}(D^{(m)}, \mathbb{C})$, the Hodge filtration of the smooth projective variety $D^{(m)}$ shifted by m;
2. the rational structure inherited from $H^k(U, \mathbb{Q})$ is $(2\pi i)^{-m}H^{k-m}(D^{(m)}, \mathbb{Q})$, so that ${}_W E_1^{-m,k+m} \cong H^{k-m}(D^{(m)}, \mathbb{Q})(-m)$ is a pure Hodge structure of weight $k + m$;
3. d_1 is, on the summand indexed by $I = \{i_1 < \cdots < i_m\}$, the alternating sum $\sum_{l=1}^m (-1)^{l-1}\gamma_l$ of the Gysin maps γ_l of the codimension one inclusions $D_I \hookrightarrow D_{I\setminus\{i_l\}}$, up to a single overall sign that does not depend on l and so changes neither the kernel nor the image; and it is a morphism of pure Hodge structures of weight $k + m$.

***Proof*:**

Step 1: the terms. By proposition 7.3.6, $\operatorname{gr}_m^W \Omega_X^\bullet(\log D) \cong a_{m*}\Omega_{D^{(m)}}^{\bullet-m}$. The morphism a_m is a closed embedding on each connected summand of $D^{(m)}$, so a_{m*} is exact and preserves cohomology, and hypercohomology may be computed downstairs:

$$ H^k\big(X, a_{m*}\Omega_{D^{(m)}}^{\bullet-m}\big) = H^k\big(D^{(m)}, \Omega_{D^{(m)}}^{\bullet-m}\big) = H^{k-m}\big(D^{(m)}, \Omega_{D^{(m)}}^\bullet\big) = H^{k-m}\big(D^{(m)}, \mathbb{C}\big) $$

the last equality because the holomorphic de Rham complex resolves the constant sheaf $\mathbb{C}_{D^{(m)}}$, which is the input already used in theorem 7.2.17. The term vanishes for $m < 0$ because $W_{-1} = 0$, and for $m > k$ because H^{k-m} of anything vanishes in negative degree.

Step 2: the induced Hodge filtration. The filtration F induces on $\operatorname{gr}_m^W \Omega_X^\bullet(\log D)$ the truncation by degree, and under Res^m the degree p part of the source corresponds to the degree $p - m$ part of $\Omega_{D^{(m)}}^\bullet$. So F^p on the graded piece corresponds to $F^{p-m}\Omega_{D^{(m)}}^\bullet$, shifted. The filtration induced on $H^{k-m}(D^{(m)}, \Omega^\bullet)$ by the truncation of the holomorphic de Rham complex is the Hodge filtration precisely because the Hodge to de Rham spectral sequence of the smooth projective variety $D^{(m)}$ degenerates at E_1 (corollary 2.1.3). This gives (1).

Step 3: the rational structure and the weight. The map Res^m was normalised algebraically, by stripping $\frac{dz_I}{z_I}$, and for a single stripped factor it is, up to the degree-dependent sign carried by

proposition 7.3.6, the topological residue of definition 5.4.6, whose factor $\frac{1}{2\pi i}$ was built in for this very purpose (box 7.2.10). That sign is invisible to the rational structure, which is what is at issue here. It is not that residue that is rational, however, but the transpose of the tube map: definition 5.4.6 normalises by $\int_\gamma \operatorname{Res}\eta = \frac{1}{2\pi i}\int_{\tau(\gamma)}\eta$, and what takes rational values on rational classes is $\gamma \mapsto \int_{\tau(\gamma)}\eta$, that is $2\pi i \operatorname{Res}$. Stripping one factor therefore multiplies the rational structure by $(2\pi i)^{-1}$: on $U = \mathbb{C}^*$ the generator of $H^1(U,\mathbb{Q})$ dual to the loop S^1 is $\frac{1}{2\pi i}\frac{dz}{z}$, because $\int_{S^1}\frac{dz}{z} = 2\pi i$, and its algebraic residue is $(2\pi i)^{-1}$. Iterating over the m factors, $(2\pi i)^m \operatorname{Res}^m$ carries the rational structure inherited from $H^k(U,\mathbb{Q})$ into $H^{k-m}(D^{(m)},\mathbb{Q})$, so that Res^m carries that structure to $(2\pi i)^{-m}H^{k-m}(D^{(m)},\mathbb{Q})$. Writing $\mathbb{Q}(-m)$ for the Tate structure $(2\pi i)^{-m}\mathbb{Q}$ of weight $2m$ and type (m,m) (definition 3.1.11), and combining with (1), the term is $H^{k-m}(D^{(m)},\mathbb{Q})(-m)$: a Hodge structure pure of weight $(k-m)+2m = k+m$, which is the index q. That the rational structure inherited from $H^k(U,\mathbb{Q})$ is preserved by the filtration, so that the statement makes sense at all, is the rationality of W discussed in theorem 7.3.11 below; the computation just made is what identifies it.

Step 4: the differential. The differential d_1 on ${}_WE_1^{-m,k+m}$ is the connecting homomorphism of the short exact sequence of complexes

$$0 \to \operatorname{gr}^W_{m-1} \to W_m/W_{m-2} \to \operatorname{gr}^W_m \to 0$$

in hypercohomology, a map $H^{k-m}(D^{(m)},\mathbb{C}) \to H^{k-m+2}(D^{(m-1)},\mathbb{C})$. Fix $I = \{i_1 < \cdots < i_m\}$ and represent a class by $\alpha = \frac{dz_I}{z_I}\wedge\eta$ with $\eta|_{D_I}$ closed. Then $d\eta$ vanishes on D_I, so locally $d\eta = \sum_l z_{i_l}\beta_l + \sum_l dz_{i_l}\wedge\delta_l$; the second sum is killed by $\frac{dz_I}{z_I}$, and

$$d\alpha = (-1)^m\frac{dz_I}{z_I}\wedge d\eta = (-1)^m\sum_{l=1}^{m}(-1)^{m-l}\frac{dz_{I\setminus\{i_l\}}}{z_{I\setminus\{i_l\}}}\wedge dz_{i_l}\wedge\beta_l$$

which lies in W_{m-1} and whose class in $\operatorname{gr}^W_{m-1}$ is supported on the m strata $D_{I\setminus\{i_l\}}$. The sign is that of moving dz_{i_l} out of slot l and to the end of the product, past the $m-l$ logarithmic factors that follow it, namely $(-1)^{m-l} = (-1)^{m+1}(-1)^{l-1}$: it alternates in l, and the leftover $(-1)^{m+1}$ is one of the several l-independent signs, along with the $(-1)^m$ above and the normalising signs of Res^m and Res^{m-1}, that make up the overall sign left unspecified in (3). So d_1 decomposes, up to that overall sign, as an alternating sum of m maps, the l-th of which is the connecting homomorphism attached to the single smooth divisor $D_I \subseteq D_{I\setminus\{i_l\}}$. For a single smooth divisor $Z \subseteq Y$ of smooth projective varieties, the sequence $0 \to \Omega^\bullet_Y \to \Omega^\bullet_Y(\log Z) \to \Omega^{\bullet-1}_Z \to 0$ of theorem 7.2.12 has connecting homomorphism $H^j(Z,\mathbb{C}) \to H^{j+2}(Y,\mathbb{C})$, and the resulting long exact sequence is the Gysin sequence of the pair (Y,Z), because theorem 7.2.17 identifies its middle term with $H^*(Y\setminus Z,\mathbb{C})$ and definition 5.4.6 identifies the residue with the topological one; the connecting map of the Gysin sequence is the Gysin pushforward γ, by excision to a tubular neighbourhood of Z and the Thom isomorphism of its normal bundle, the Thom class being Poincaré dual to $[Z]$.[a] This proves the first half of (3).

Step 5: d_1 is a morphism of Hodge structures. Let $\iota\colon Z \hookrightarrow Y$ be a smooth divisor in a smooth projective variety Y of dimension d, and fix $s \geq 0$. The Gysin map $\gamma_\iota\colon H^s(Z,\mathbb{Q}) \to H^{s+2}(Y,\mathbb{Q})$ is by definition the transpose of $\iota^*\colon H^{2d-s-2}(Y,\mathbb{Q}) \to H^{2d-s-2}(Z,\mathbb{Q})$ under Poincaré duality ([30, Poincaré duality]); in particular it is defined over $\mathbb{Q}$. Restriction ι^* is a morphism of pure Hodge structures of weight $2d-s-2$, while Poincaré duality identifies $H^s(Z,\mathbb{Q})$ with $H^{2d-2-s}(Z,\mathbb{Q})^\vee(-(d-1))$ and $H^{s+2}(Y,\mathbb{Q})$ with $H^{2d-s-2}(Y,\mathbb{Q})^\vee(-d)$, the twists being those of

the fundamental classes of Z and of Y. Transposing ι^* and comparing the two twists,

$$\gamma_\iota \colon H^s(Z,\mathbb{Q})(-1) \to H^{s+2}(Y,\mathbb{Q})$$

is a morphism of pure Hodge structures. Apply this with $Z = D_I$ and $Y = D_{I\setminus\{i_l\}}$, so $s = k-m$ and the two groups are summands of ${}_WE_1^{-m,k+m}$ and of ${}_WE_1^{-m+1,k+m}$; twisting both sides by $(-(m-1))$ turns it into a morphism $H^{k-m}(D^{(m)},\mathbb{Q})(-m) \to H^{k-m+2}(D^{(m-1)},\mathbb{Q})(-(m-1))$ of pure Hodge structures of weight $k+m$. An alternating sum of such is again one, as claimed.

[a] [32, Tubular Neighbourhood Theorem], [30, The Thom Isomorphism].

Everything on the first page is therefore cohomology of smooth projective varieties, and the first differential is built from maps of the kind Hodge theory controls. The weight $q = k+m$ attached to the term ${}_WE_1^{-m,k+m}$ is constant along d_1, which is why the page is grouped by q; along d_r with $r \geq 2$ it drops by $r-1$.

Two inputs to what follows are taken from elsewhere, and it is worth stating both precisely before they are used, so that the reader knows exactly where this section stops proving things. The first is upstream in the series; the second is external. Both are due to Deligne, and [18, Théorie de Hodge II] is the original for each, with [33, the construction of a mixed Hodge complex] a modern account.

Theorem 7.3.8: Strictness of the Weight Differentials

Let $K^\bullet$ carry an increasing filtration W and a decreasing filtration F, both biregular filtrations by subcomplexes. Suppose the differentials d_s of the spectral sequence of W are strictly compatible with F for every s. Then, in the notation of the proof below, $\sigma_s = \operatorname{rank}\delta_s$ for every s, the three filtrations F induces on each page agree, and the spectral sequence of F degenerates at E_1.
This is the lemma on two filtrations, [7, the lemma on two filtrations], where it is proved and the filtered derived category it lives in is constructed.

The hypothesis is the point, and it must be read carefully. Strictness of the d_s is *assumed*, not deduced. It is tempting to hope that strictness on the graded pieces gr_n^W alone would suffice, and it does not: strictness of d_0 is a statement about the graded pieces separately and cannot see how they are glued, while strictness of d_1 is a statement about the gluing. A two-dimensional counterexample is worked in [7, Sharpness of the Hypothesis]. What gives the hypothesis here is Hodge theory rather than homological algebra: the terms of the weight spectral sequence carry pure Hodge structures, and a morphism of pure Hodge structures is strict for the Hodge filtration (proposition 3.1.8). That is the same mechanism as part (1) of theorem 7.3.10, applied one page at a time, and it is why the two parts of that theorem are not independent.

Theorem 7.3.9: Imported: Rationality of the Weight Filtration

Under the comparison isomorphism of theorem 7.2.17, the subcomplex $W_m\Omega_X^\bullet(\log D)$ corresponds to the canonical truncation $\tau_{\leq m}Rj_*\mathbb{C}_U$, and that truncation is already defined on $Rj_*\mathbb{Q}_U$. Consequently $W_{k+m}H^k(U,\mathbb{C})$ is the complexification of a $\mathbb{Q}$-subspace of $H^k(U,\mathbb{Q})$.

Neither is a detail, and they are of different kinds. Theorem 7.3.8 is *proved* upstream in [7, The Lemma on Two Filtrations], whose filtered derived category exists for exactly this purpose. It is cited, not taken on faith. What it does is convert strictness of the weight differentials into the

statements part (2) of the degeneration needs. That strictness is a separate matter and is imported, from Deligne's mixed Hodge complex formalism, as the proof below records at the point of use. Theorem 7.3.9 is imported outright: it makes the weight filtration exist over $\mathbb{Q}$ at all, without which there is no mixed Hodge structure on $H^k(U,\mathbb{Q})$, and it needs the derived pushforward and its canonical filtration, which no book in the series develops, since section 7.2 reaches theorem 7.2.17 through the fine complex $j_*\mathcal{A}_U^\bullet$ instead. Together with the strictness input these are the only things taken on faith beyond theorem 7.2.3.

Theorem 7.3.10: Degeneration of the Weight and Hodge Spectral Sequences

Let X be smooth projective, $D \subseteq X$ a normal crossings divisor with smooth components, and $U = X \setminus D$. Then

1. the weight spectral sequence (7.8) degenerates at E_2: the differentials d_r vanish for all $r \geq 2$;
2. the Hodge spectral sequence (7.7) degenerates at E_1, so that $\dim_{\mathbb{C}} H^k(U,\mathbb{C}) = \sum_{p+q=k} \dim_{\mathbb{C}} H^q\big(X, \Omega_X^p(\log D)\big)$ for every k.

***Proof*:**

Part (1). By proposition 7.3.7 the term ${}_W E_1^{p,q}$ carries a pure $\mathbb{Q}$-Hodge structure of weight q, and $d_1 \colon E_1^{p,q} \to E_1^{p+1,q}$ is a morphism of such. Kernels and images of morphisms of pure Hodge structures of a fixed weight are again pure Hodge structures of that weight, since such morphisms are strict for F (proposition 3.1.8); hence

$$ {}_W E_2^{p,q} = \ker\big(d_1 \colon E_1^{p,q} \to E_1^{p+1,q}\big) / \operatorname{im}\big(d_1 \colon E_1^{p-1,q} \to E_1^{p,q}\big) $$

is a pure $\mathbb{Q}$-Hodge structure of weight q, with rational structure and Hodge filtration induced from those of E_1. Inductively, if $E_r^{p,q}$ is pure of weight q for all p and d_r vanishes, the same holds on E_{r+1}.

Now fix $r \geq 2$ and consider $d_r \colon E_r^{p,q} \to E_r^{p+r,\,q-r+1}$. The source is pure of weight q, the target is pure of weight $q-r+1$, and $q-(q-r+1) = r-1 \geq 1$, so the target has *strictly smaller* weight than the source. The map d_r is defined over $\mathbb{Q}$, since the whole page is, and it is compatible with F, since F is a filtration of the logarithmic complex by subcomplexes and therefore induces filtrations on every page compatible with every differential. Write A for the source and B for the target, of weights a and $b = a-(r-1) < a$. Then d_r is a morphism of pure Hodge structures between structures of different weights, so $d_r = 0$ by lemma 7.1.5(2), which is proved in section 7.1 for exactly this purpose.

The hypothesis that the weight *drops* is what makes the argument work: a rational F-compatible map into a structure of strictly larger weight need not vanish, as the identity map $\mathbb{Q}(0) \to \mathbb{Q}(-1)$ shows. This is the content of exercise 7.3.1 (5). Since $d_r = 0$ for every $r \geq 2$, the sequence degenerates at E_2, proving (1).

Part (2). Argue by induction on the number N of components of D. If $N = 0$ then $U = X$ and (7.7) is the Hodge to de Rham spectral sequence of a smooth projective variety, which degenerates at E_1 by corollary 2.1.3.

Let $N \geq 1$, write $D' = D_1 \cup \cdots \cup D_{N-1}$, $Z = D_N$, $D'_Z = D' \cap Z$, and set $U' = X \setminus D'$ and $Z^\circ = Z \setminus D'_Z$. Both D' on X and D'_Z on Z are normal crossings divisors with at most $N-1$

components, so the inductive hypothesis applies to them. In adapted coordinates in which $Z = \{z_r = 0\}$, the basis elements $\frac{dz_I}{z_I} \wedge dz_J$ of $\Omega^p_X(\log D)$ with $r \notin I$ span $\Omega^p_X(\log D')$ and those with $r \in I$ map under the residue along Z to a basis of $\Omega^{p-1}_Z(\log D'_Z)$, so for every p

$$0 \to \Omega^p_X(\log D') \to \Omega^p_X(\log D) \xrightarrow{\mathrm{Res}_Z} \Omega^{p-1}_Z(\log D'_Z) \to 0 \tag{7.9}$$

is exact. Assembling (7.9) over p gives a short exact sequence of complexes $0 \to K' \to K \to L \to 0$ with $K' = \Omega^\bullet_X(\log D')$, $K = \Omega^\bullet_X(\log D)$ and $L = \Omega^\bullet_Z(\log D'_Z)[-1]$, and it is a sequence of complexes filtered by F whose F-graded pieces $0 \to \mathrm{gr}^p_F K' \to \mathrm{gr}^p_F K \to \mathrm{gr}^p_F L \to 0$ are again exact, by (7.9).

Write $a_k = \dim H^k(U', \mathbb{C})$, $c_k = \dim H^k(Z^\circ, \mathbb{C})$, $b_k = \dim H^k(U, \mathbb{C})$ and A_k, C_k, B_k for the corresponding sums of E_1-dimensions, so that $a_k = A_k$ and $c_k = C_k$ by induction and $b_k \le B_k$ always. Taking hypercohomology of $0 \to K' \to K \to L \to 0$ gives the long exact sequence

$$\cdots \to H^k(U', \mathbb{C}) \to H^k(U, \mathbb{C}) \xrightarrow{\mathrm{Res}_Z} H^{k-1}(Z^\circ, \mathbb{C}) \xrightarrow{\delta_{k-1}} H^{k+1}(U', \mathbb{C}) \to \cdots$$

whence $b_k = a_k + c_{k-1} - \mathrm{rank}\delta_{k-1} - \mathrm{rank}\delta_{k-2}$. Taking hypercohomology of the F-graded sequences gives, for each p, the long exact sequence of coherent cohomology groups with connecting maps $\partial^{p,q}\colon H^q(Z, \Omega^{p-1}_Z(\log D'_Z)) \to H^{q+1}(X, \Omega^p_X(\log D'))$, whence $B_k = A_k + C_{k-1} - \sigma_{k-1} - \sigma_{k-2}$, where σ_s is the sum of $\mathrm{rank}\partial^{p,q}$ over the pairs with $(p-1)+q = s$. Because the E_1-pages of the three spectral sequences fit into the long exact sequence just written, and because those of K' and L degenerate at E_1 by induction, $\partial^{p,q}$ is the map induced by δ_s on gr^p_F, so

$$\sigma_s = \sum_p \mathrm{rank}\big(\mathrm{gr}^p_F \delta_s\big) \le \mathrm{rank}\delta_s$$

with equality for all s if and only if every δ_s is strictly compatible with F. Comparing the two displayed counts,

$$B_k - b_k = \big(\mathrm{rank}\delta_{k-1} - \sigma_{k-1}\big) + \big(\mathrm{rank}\delta_{k-2} - \sigma_{k-2}\big)$$

so the induction closes exactly when the Gysin maps $\delta_s\colon H^s(Z^\circ, \mathbb{C}) \to H^{s+2}(U', \mathbb{C})$ are strict for F.

That strictness is what has to be given, and it does not follow from the material developed here. The obstacle is precise: Z° and U' are open, their cohomology is not pure, and proposition 3.1.8 therefore does not apply to δ_s. Nor does theorem 7.3.8 give it, since that lemma *uses* strictness of the d_s rather than producing it; what the lemma then gives, once strictness is in hand, is that the three induced filtrations agree and that the F-sequence degenerates.

The strictness itself is Deligne's, and it comes from the mixed Hodge complex formalism: the E_1 terms are cohomology of smooth projective strata and so are pure, which makes d_1 strict by proposition 3.1.8, and the formalism bootstraps from that page to all later ones simultaneously with the construction. It is proved in [18, Théorie de Hodge II, the construction of the mixed Hodge structure], with a modern account in [33, the construction of a mixed Hodge complex], and it is not proved here. Granting it, $B_k = b_k$ for all k and the Hodge spectral sequence degenerates at E_1, completing the proof.

The mechanism behind Part (1) is worth separating from its bookkeeping. The objects on the first page are cohomology groups of smooth projective varieties, and such groups admit no nonzero map that lowers weight. Nothing about the geometry of U enters the argument. What enters is that the

strata $D^{(m)}$ are projective, which is the payoff of insisting on a good compactification.

Degeneration at E_2 is what allows the graded pieces of the weight filtration to be read off a page already computed, and reading them off is all that remains. The next theorem puts the two filtrations on $H^k(U,\mathbb{Q})$ and checks the three things a mixed Hodge structure asks of them: that W be defined over $\mathbb{Q}$, that its graded pieces be pure with the filtration induced by F, and that neither filtration depend on the compactification used to produce it.

Theorem 7.3.11: Deligne's Mixed Hodge Structure on a Smooth Variety

Let U be a smooth complex algebraic variety and let $X \supseteq U$ be a smooth projective compactification with $D = X \setminus U$ a normal crossings divisor with smooth components. Define, on $H^k(U,\mathbb{C}) = H^k(X, \Omega_X^\bullet(\log D))$,

$$W_{k+m}H^k(U,\mathbb{C}) = \operatorname{im}\Big(H^k\big(X, W_m\Omega_X^\bullet(\log D)\big) \to H^k(U,\mathbb{C})\Big)$$
$$F^pH^k(U,\mathbb{C}) = \operatorname{im}\Big(H^k\big(X, F^p\Omega_X^\bullet(\log D)\big) \to H^k(U,\mathbb{C})\Big)$$

Then W is defined over $\mathbb{Q}$, and $\big(H^k(U,\mathbb{Q}), W_\bullet, F^\bullet\big)$ is a mixed Hodge structure. Its graded pieces are

$$\operatorname{gr}^W_{k+m}H^k(U,\mathbb{Q}) \cong {}_WE_2^{-m,\,k+m}$$

a subquotient of $H^{k-m}(D^{(m)},\mathbb{Q})(-m)$; in particular $\operatorname{gr}^W_w H^k(U,\mathbb{Q}) = 0$ unless $k \le w \le 2k$. The structure does not depend on the choice of (X, D).

***Proof*:**

Step 1: rationality of W. The filtration F is defined only on $\mathbb{C}$-cohomology and needs no rational structure, but W does. This is theorem 7.3.9, imported: $W_{k+m}H^k(U,\mathbb{C})$ is the complexification of a $\mathbb{Q}$-subspace of $H^k(U,\mathbb{Q})$, and (7.8) becomes the Leray spectral sequence of j with $\mathbb{Q}$-coefficients, reindexed. The local shape of the sheaves involved is visible from proposition 7.3.7: near a point of X lying on exactly r branches, U looks like $(\mathbb{C}^*)^r \times \mathbb{C}^{n-r}$, whose m-th cohomology is Λ^m of $\mathbb{Q}(-1)^r$, so $R^mj_*\mathbb{Q}_U \cong a_{m*}\mathbb{Q}_{D^{(m)}}(-m)$, and the E_1-terms computed in proposition 7.3.7 are exactly $H^{k-m}(X, a_{m*}\mathbb{Q}_{D^{(m)}}(-m)) = H^{k-m}(D^{(m)},\mathbb{Q})(-m)$.

The identification of $W_m\Omega_X^\bullet(\log D)$ with that truncation is the single input this proof takes from outside the book, and it is what gives the rationality of W: the truncation is intrinsic to $Rj_*\mathbb{Q}_U$, but comparing it with a filtration of the logarithmic complex is done in the filtered derived category, and section 7.2 reaches theorem 7.2.17 through the fine complex $j_*\mathcal{A}_U^\bullet$ rather than through a derived pushforward. So this one step is taken from [18, Théorie de Hodge II, the weight filtration of the logarithmic complex]. Nothing else is: the weight bound of Step 3 and the independence of the compactification of Step 4, for which such a citation is usually made, are proved here.

Step 2: the graded pieces are pure. By theorem 7.3.10(1) the weight spectral sequence degenerates at E_2, so

$$\operatorname{gr}^W_{k+m}H^k(U,\mathbb{C}) = {}_WE_\infty^{-m,k+m} = {}_WE_2^{-m,k+m}$$

and by the proof of theorem 7.3.10(1) this is a pure $\mathbb{Q}$-Hodge structure of weight $k+m$, a subquotient of the pure structure $H^{k-m}(D^{(m)},\mathbb{Q})(-m)$ of proposition 7.3.7. Its Hodge filtration is the one induced by F: the filtration F induces the Hodge filtration on each ${}_WE_1$ term by

proposition 7.3.7(1), and it induces on the abutment the filtration $F^\bullet H^k(U,\mathbb{C})$ defined in the statement, these two being compatible because theorem 7.3.10(2) makes F strict on the weight spectral sequence. Hence $\mathrm{gr}^W_{k+m} H^k(U,\mathbb{Q})$ with the filtration induced by F is a pure Hodge structure of weight $k+m$, which is the defining condition of definition 7.1.1.

Step 3: the weight bounds. By proposition 7.3.7 the term ${}_W E_1^{-m,k+m}$ vanishes unless $0 \le m \le k$: for $m<0$ because $W_{-1}=0$, and for $m>k$ because $H^{k-m}(D^{(m)},\mathbb{Q})$ sits in negative degree. The same therefore holds for the subquotient ${}_W E_2^{-m,k+m} = \mathrm{gr}^W_{k+m} H^k(U,\mathbb{Q})$, so the weights $w=k+m$ occurring in $H^k(U,\mathbb{Q})$ satisfy $k \le w \le 2k$. The sharper bound $w \le k+\min(k,n,N)$ also follows, since $D^{(m)}$ is empty for $m > \min(n,N)$.

Step 4: independence of the compactification. Given two good compactifications X_1, X_2 of U, resolve the closure of the diagonal $U \hookrightarrow X_1 \times X_2$ to obtain a third good compactification X_3 dominating both, by [24]. It therefore suffices to compare X with a good compactification $\tilde{X}$ admitting a morphism $\pi\colon \tilde{X} \to X$ restricting to the identity on U. Pullback of forms gives $\pi^*\Omega^\bullet_X(\log D) \to \Omega^\bullet_{\tilde{X}}(\log \tilde{D})$; it preserves F because it preserves form degree, and it preserves W because if $z\circ\pi = u\prod_l w_l^{e_l}$ with u a unit then $\pi^*\frac{dz}{z} = \frac{du}{u} + \sum_l e_l \frac{dw_l}{w_l}$ contributes at most one logarithmic pole to each term. So the induced map on $H^k(U,\mathbb{C})$, which is the identity, is a morphism of bifiltered vector spaces, and the filtrations computed on $\tilde{X}$ are at least as large as those computed on X. Both are mixed Hodge structures on the same $H^k(U,\mathbb{Q})$ and the identity is a morphism between them; by theorem 7.1.8 a morphism of mixed Hodge structures is strict for both filtrations, so an identity morphism forces the filtrations to agree, completing the proof.

The theorem says that the cohomology of a smooth variety is built from the cohomology of the strata of any boundary one chooses to add, and that the assembly is canonical even though the boundary is not. The weight filtration records how far U is from being compact: the bottom step is the part visible already on a compactification, the top step is the part that only the punctures see, and everything between is a subquotient of the cohomology of a stratum, twisted. On a smooth projective U the divisor D is empty, only $m=0$ survives, and the theorem returns the pure Hodge structure of chapter 2.

Corollary 7.3.12: The Lowest Weight Part is the Image from a Compactification

With U, X, D as above,

$$W_k H^k(U,\mathbb{Q}) = \mathrm{im}\Big(j^*\colon H^k(X,\mathbb{Q}) \to H^k(U,\mathbb{Q})\Big)$$

and this subspace does not depend on the choice of (X,D).

***Proof*:**
By definition $W_k H^k(U,\mathbb{C})$ is the image of $H^k(X, W_0\Omega^\bullet_X(\log D)) \to H^k(U,\mathbb{C})$, and $W_0\Omega^\bullet_X(\log D) = \Omega^\bullet_X$, whose hypercohomology is $H^k(X,\mathbb{C})$; the map induced by the inclusion of complexes is the restriction j^*. Both sides are defined over $\mathbb{Q}$, the left by theorem 7.3.11 and the right because j^* is, so the equality holds rationally. Independence of (X,D) is that of W, as we sought to show.

Example 7.3.13: The Punctured Projective Line

Let $X = \mathbb{C}\mathrm{P}^1$, let $D = \{p_1, \dots, p_r\}$ with $r \geq 1$ distinct points, and let $U = \mathbb{C}\mathrm{P}^1 \setminus D$. The strata are $D^{(0)} = \mathbb{C}\mathrm{P}^1$ and $D^{(1)} = D$, a disjoint union of r points, and $D^{(m)} = \emptyset$ for $m \geq 2$ since the points are distinct.

The first page of the weight spectral sequence for $k = 1$ has two entries,

$$ {}_W E_1^{0,1} = H^1(\mathbb{C}\mathrm{P}^1, \mathbb{Q}) = 0, \qquad {}_W E_1^{-1,2} = H^0(D^{(1)}, \mathbb{Q})(-1) = \mathbb{Q}(-1)^r $$

and the differential $d_1 \colon {}_W E_1^{-1,2} \to {}_W E_1^{0,2} = H^2(\mathbb{C}\mathrm{P}^1, \mathbb{Q})$ is the Gysin map, which sends the class of a point to the generator of $H^2(\mathbb{C}\mathrm{P}^1, \mathbb{Q}) \cong \mathbb{Q}(-1)$; on the r summands it is the sum map $(c_1, \dots, c_r) \mapsto \sum_i c_i$. It is surjective, with kernel of rank $r - 1$. Hence

$$ \mathrm{gr}_1^W H^1(U, \mathbb{Q}) = 0, \qquad \mathrm{gr}_2^W H^1(U, \mathbb{Q}) = \ker(d_1) \cong \mathbb{Q}(-1)^{r-1} $$

so $H^1(U, \mathbb{Q})$ is pure of weight 2 and of type $(1, 1)$, of rank $r - 1$. This is what the topology predicts: $\mathbb{C}\mathrm{P}^1$ minus r points is homotopy equivalent to a wedge of $r - 1$ circles. It is also what the residues say: a class in $H^1(U, \mathbb{C})$ is represented by a rational 1-form with at worst simple poles at the p_i, its residues are the r numbers c_i, and the residue theorem is precisely the statement $\sum_i c_i = 0$ that cuts out $\ker(d_1)$.

The Hodge spectral sequence confirms the count. Since $\Omega^1_{\mathbb{C}\mathrm{P}^1}(\log D) \cong \mathcal{O}_{\mathbb{C}\mathrm{P}^1}(r - 2)$, the nonzero terms in total degree 1 are $H^0(\mathbb{C}\mathrm{P}^1, \mathcal{O}(r-2))$, of dimension $r-1$ for $r \geq 1$, and $H^1(\mathbb{C}\mathrm{P}^1, \mathcal{O}_{\mathbb{C}\mathrm{P}^1}) = 0$; the total is $r - 1$, matching $\dim H^1(U, \mathbb{C})$ as theorem 7.3.10(2) requires. For $k = 2$ one finds ${}_W E_2^{0,2} = \mathrm{coker}(d_1) = 0$ and ${}_W E_2^{-1,3} = 0$, so $H^2(U, \mathbb{Q}) = 0$.

The weight filtration is trivial here in the sense that $H^1(U, \mathbb{Q})$ has a single weight. That is an accident of genus 0: the piece gr_1^W, which would carry the cohomology of the compactification, vanishes because $H^1(\mathbb{C}\mathrm{P}^1) = 0$. The first genuinely mixed example, where gr_1^W and gr_2^W are both nonzero and the extension between them carries information, is the punctured elliptic curve of section 7.4.

A construction that depended on the compactification would be of limited use; one that does not is a candidate for a functor. The last result of the section confirms that it is one on the smooth quasi-projective varieties, which are the ones for which theorem 7.2.3 produces the projective compactification that theorem 7.3.11 needs.

Proposition 7.3.14: Functoriality

Let $f \colon U' \to U$ be a morphism of smooth quasi-projective complex algebraic varieties. Then

$$ f^* \colon H^k(U, \mathbb{Q}) \to H^k(U', \mathbb{Q}) $$

is a morphism of mixed Hodge structures for the structures of theorem 7.3.11. Consequently $U \mapsto H^k(U, \mathbb{Q})$ is a contravariant functor from smooth quasi-projective complex algebraic varieties to mixed Hodge structures.

***Proof*:**
Both U and U' are quasi-projective, so theorem 7.2.3 gives good compactifications with projective total space: choose one $X \supseteq U$ with boundary D and X projective, and one $Y' \supseteq U'$ with Y' projective. Let $\Gamma \subseteq Y' \times X$ be the closure of the graph of f, a projective variety because $Y' \times X$ is one, and apply [24] to obtain a smooth X' with a projective morphism $X' \to \Gamma$ that is an isomorphism over U' and for which $D' = X' \setminus U'$ is a normal crossings divisor; being projective over the projective Γ, the variety X' is itself projective. The two projections give a good compactification $X' \supseteq U'$ together with a morphism $\bar{f}\colon X' \to X$ restricting to f on U'. Since $f(U') \subseteq U$, we have $\bar{f}^{-1}(D) \subseteq D'$.

Pullback of forms along $\bar{f}$ therefore carries forms with logarithmic poles along D to forms with logarithmic poles along D': if z is a local equation of a branch of D then $z \circ \bar{f} = u \prod_l w_l^{e_l}$ for a unit u and local equations w_l of branches of D', so

$$\bar{f}^* \frac{dz}{z} = \frac{du}{u} + \sum_l e_l \frac{dw_l}{w_l}$$

a sum of a holomorphic form and forms with a single logarithmic pole. Hence $\bar{f}^*\colon \Omega^\bullet_X(\log D) \to \Omega^\bullet_{X'}(\log D')$ is a morphism of complexes taking W_m into W_m, by the displayed identity applied to each of the at most m logarithmic factors of a local generator, and taking F^p into F^p, because pullback preserves form degree.

Passing to hypercohomology, the induced map $H^k(U, \mathbb{C}) \to H^k(U', \mathbb{C})$ is f^*, and it carries W_{k+m} into W_{k+m} and F^p into F^p by the previous paragraph. It is defined over $\mathbb{Q}$ because it is pullback on singular cohomology. By theorem 7.3.11 the filtrations so computed are the intrinsic ones, so f^* is a morphism of mixed Hodge structures as required. Functoriality of the assignment is the functoriality of $f \mapsto f^*$ on singular cohomology, completing the proof.

Strictness now comes for free: by theorem 7.1.8 every f^* is strict for W and for F, so kernels, images and cokernels of maps induced by morphisms of varieties are again mixed Hodge structures. This is what makes the formalism usable in practice, and it is the property that fails for a bare pair of filtrations chosen by hand.

Exercise 7.3.1

1. Let $X = \mathbb{C}\mathrm{P}^1$, $D = \{0\}$ and $U = \mathbb{C}$. Write down W and F on $\Omega^\bullet_X(\log D)$, compute the first page of the weight spectral sequence in all bidegrees, and identify d_1. Deduce $H^k(\mathbb{C}, \mathbb{Q})$ for all k, and observe that the weight spectral sequence does *not* degenerate at E_1 here. Why does this not contradict theorem 7.3.10?

2. Let $X = \mathbb{C}\mathrm{P}^1 \times \mathbb{C}\mathrm{P}^1$, let $D_1 = \{0\} \times \mathbb{C}\mathrm{P}^1$ and $D_2 = \mathbb{C}\mathrm{P}^1 \times \{0\}$, and let $U = X \setminus (D_1 \cup D_2)$, so $U \cong \mathbb{C}^2$. List the strata $D^{(0)}, D^{(1)}, D^{(2)}$, write out ${}_W E_1^{-m,k+m}$ for $k = 0, 1, 2$ and every m, and compute enough of d_1 to conclude that $H^k(U, \mathbb{Q}) = 0$ for $k \geq 1$. This is the smallest example in which the stratum $D^{(2)}$ contributes.

3. Take $D = \emptyset$, so that $U = X$ is smooth projective. Compute both spectral sequences and show that theorem 7.3.11 returns the pure Hodge structure of weight k on $H^k(X, \mathbb{Q})$. Which step of the proof of theorem 7.3.10 becomes vacuous?

4. Let X be a smooth projective curve of genus g, let $D = \{p_1, \ldots, p_r\}$ with $r \geq 1$, and let $U = X \setminus D$. Compute $\mathrm{gr}_1^W H^1(U, \mathbb{Q})$ and $\mathrm{gr}_2^W H^1(U, \mathbb{Q})$, and check the resulting dimension of

$H^1(U,\mathbb{Q})$ against $\dim H^0(X,\Omega^1_X(\log D)) + \dim H^1(X,\mathcal{O}_X)$.

5. Let $A = \mathbb{Q}(0)$ and $B = \mathbb{Q}(-1)$, and let $f\colon A \to B$ be the identity map of the underlying $\mathbb{Q}$-vector space $\mathbb{Q}$. Show that f is rational and satisfies $f(F^pA) \subseteq F^pB$ for every p, but is not a morphism of Hodge structures. Explain why this does not disturb the vanishing argument in the proof of theorem 7.3.10(1).

6. Let $U = \mathbb{C}^*$ and $U' = \mathbb{C}^* \setminus \{1\}$. Using example 7.3.13, identify the mixed Hodge structures on $H^1(U,\mathbb{Q})$ and $H^1(U',\mathbb{Q})$, and show that the restriction map $H^1(U,\mathbb{Q}) \to H^1(U',\mathbb{Q})$ is an injective morphism of mixed Hodge structures. Is it strict for W?

7.4 The Punctured Elliptic Curve

Everything section 7.1 through section 7.3 constructs can be written down in closed form on a curve, where the logarithmic complex has two terms and the weight filtration has two steps. This section carries that out for E an elliptic curve and $U = E \setminus D$ the complement of a finite set of points, and then extracts the invariant that the graded pieces do not see.

The computation settles a question the general theorem leaves open, namely how much weight the punctures create. A single puncture creates none: $H^1(E \setminus \{p\},\mathbb{Q})$ is the pure structure of example 3.4.3, for every p. Two punctures create exactly one class of weight two, and the resulting extension remembers the difference $p - q$ as a point of E.

Fix τ in the upper half-plane and write $E = E_\tau = \mathbb{C}/\Lambda_\tau$ with $\Lambda_\tau = \mathbb{Z}\cdot 1 \oplus \mathbb{Z}\tau$, as in example 3.4.3. Let $D \subseteq E$ be a set of $n \geq 1$ points[4] and $U = E \setminus D$, with inclusion $j\colon U \to E$. No compactification has to be produced: E is smooth and projective, and a finite set of points on a curve is a normal-crossings divisor in the sense of definition 7.2.1, each point being cut out by a single coordinate. So theorem 7.2.3, the one imported hypothesis of section 7.2, is not used anywhere below; the compactification is the one the geometry hands over.

Two features of dimension one simplify the logarithmic complex of definition 7.2.5 to the point where it can be handled by hand. It has only two terms,

$$\Omega^\bullet_E(\log D) \;=\; \left[\; \mathcal{O}_E \;\xrightarrow{\;d\;}\; \Omega^1_E(\log D) \;\right] \tag{7.10}$$

and the coordinate z on $\mathbb{C}$ descends to a nowhere-vanishing holomorphic 1-form dz on E, so $\Omega^1_E \cong \mathcal{O}_E$ and a section of $\Omega^1_E(\log D)$ is $h(z)\,dz$ with h a Λ_τ-periodic meromorphic function whose poles lie in D and are at most simple. Every question about the complex therefore becomes a question about elliptic functions, and the tool for those is contour integration over a fundamental parallelogram.

That contour gives two identities at once. Integrating a form over the boundary of the parallelogram gives the residue theorem ([31, Residue Theorem]), and integrating the form against the primitive z of dz gives a relation between the periods of the form and the positions of its poles. The second identity is what will eventually locate the extension class, so it is worth recording both together.

[4] Here n counts the punctures. In section 7.2 and section 7.3 the letter n was the dimension of X and N the number of components of the boundary; on a curve the dimension is 1 and $N = n$.

Lemma 7.4.1: Residues and Periods on a Complex Torus

Let $E = \mathbb{C}/\Lambda_\tau$ and let η be a meromorphic 1-form on E, not identically zero, with pole set S. Choose $z_0 \in \mathbb{C}$ so that the closed parallelogram P with vertices $z_0,\ z_0+1,\ z_0+1+\tau,\ z_0+\tau$ contains exactly one lift of each point of S, and each in its interior; let γ_1 and γ_2 be the images in E of the segments $[z_0, z_0+1]$ and $[z_0, z_0+\tau]$.

1. $\displaystyle\sum_{x\in S} \mathrm{Res}_x\eta = 0.$
2. If every pole of η is simple, and a_x denotes the lift of $x \in S$ interior to P, then

$$\int_{\gamma_2} \eta \;-\; \tau \int_{\gamma_1} \eta \;=\; 2\pi i \sum_{x\in S} a_x \,\mathrm{Res}_x\eta \tag{7.11}$$

***Proof*:**
Pull η back to $\mathbb{C}$ and write it there as $h(z)\,dz$ with h meromorphic and Λ_τ-periodic. The boundary ∂P, traversed counterclockwise, consists of the segment A from z_0 to z_0+1, the segment B from z_0+1 to $z_0+1+\tau$, the reverse of the segment $A+\tau$, and the reverse of the segment $B-1$. Note that $B-1$ is the segment from z_0 to $z_0+\tau$, whose image in E is γ_2, while the image of A is γ_1.




Figure 7.1: The parallelogram P, traversed counterclockwise, with the lifted poles in its interior. The cycle γ_2 is the left side traversed upward, so it occurs in ∂P with a minus sign

Step 1: the residue theorem. Let g be holomorphic on a neighbourhood of P. Pairing the two horizontal sides and then the two slanted ones,

$$\oint_{\partial P} g\,h\,dz \;=\; \int_A \big(g(z)-g(z+\tau)\big)h(z)\,dz \;+\; \int_{B-1} \big(g(z+1)-g(z)\big)h(z)\,dz \tag{7.12}$$

where periodicity of h has been used to replace $h(z+\tau)$ and $h(z+1)$ by $h(z)$. Taking $g \equiv 1$ makes both integrands vanish, so $\oint_{\partial P}\eta = 0$; and the residue theorem ([31, Residue Theorem]) in the plane evaluates the same integral as $2\pi i$ times the sum of the residues of h at the interior poles, which are the lifts a_x. Since $\mathrm{Res}_{a_x} h = \mathrm{Res}_x\eta$, assertion (1) follows.
Step 2: the period relation. Take $g(z) = z$ in (7.12). The first integrand becomes $-\tau\,h(z)$ and the second becomes $h(z)$, so

$$\oint_{\partial P} z\,h(z)\,dz \;=\; -\tau\int_A h\,dz \;+\; \int_{B-1} h\,dz \;=\; -\tau\int_{\gamma_1}\eta \;+\; \int_{\gamma_2}\eta$$

On the other hand $z\,h(z)\,dz$ has a simple pole at each a_x with residue $a_x \mathrm{Res}_{a_x} h = a_x \mathrm{Res}_x \eta$, because η has simple poles and z is holomorphic; the residue theorem in the plane evaluates the contour integral as $2\pi i \sum_x a_x \mathrm{Res}_x \eta$. Comparing the two evaluations gives (7.11), as we sought to show.

Part (1) is the obstruction that governs the whole section. A residue is a local quantity, but the residues of a global form cannot be prescribed independently: their sum is forced to vanish. Part (2) refines this into an equality rather than a vanishing, and its right-hand side is the only place in the section where the positions of the punctures, and not merely their number, enter.

The first consequence is a count of the logarithmic differentials, which by (7.10) is the only piece of the logarithmic complex that is not already understood.

Proposition 7.4.2: Logarithmic Differentials on an Elliptic Curve

Let $E = \mathbb{C}/\Lambda_\tau$ and let $D \subseteq E$ consist of $n \geq 1$ points. The residue map of theorem 7.2.12 in degree one,

$$\mathrm{Res}\colon H^0\big(E, \Omega^1_E(\log D)\big) \to \mathbb{C}^D$$

has kernel $H^0(E, \Omega^1_E) = \mathbb{C}\,dz$ and image the hyperplane $\big\{ r \in \mathbb{C}^D \big| \sum_{x\in D} r_x = 0 \big\}$. In particular $\dim_{\mathbb{C}} H^0\big(E, \Omega^1_E(\log D)\big) = n$, and for distinct $p, q \in E$ there is a meromorphic 1-form on E with simple poles exactly at p and q, of residues $+1$ and -1, unique up to adding a multiple of dz.

***Proof*:**
The set D is finite, so its points are pairwise disjoint components and theorem 7.2.12 applies. In degree one its sequence is the short exact sequence of sheaves

$$0 \to \Omega^1_E \to \Omega^1_E(\log D) \xrightarrow{\mathrm{Res}} \mathcal{O}_D \to 0 \tag{7.13}$$

the quotient being the skyscraper $\mathcal{O}_D = \bigoplus_{x\in D} \mathbb{C}_x$ because D is a finite set of reduced points. Its long exact cohomology sequence begins

$$0 \to H^0(E, \Omega^1_E) \to H^0\big(E, \Omega^1_E(\log D)\big) \xrightarrow{\mathrm{Res}} \mathbb{C}^D \xrightarrow{\delta} H^1(E, \Omega^1_E)$$

Here $H^0(E, \Omega^1_E) = \mathbb{C}\,dz$: a global holomorphic 1-form is $h(z)\,dz$ with h holomorphic and Λ_τ-periodic, so h descends to a holomorphic function on the compact connected E and is constant by the maximum principle.[a] And $H^1(E, \Omega^1_E) \cong H^{1,1}(E)$ by theorem 1.1.3, which is one-dimensional by example 2.1.2. So the proposition amounts to identifying δ with a nonzero multiple of the sum map.

Step 1: δ is a multiple of the sum map. For $x \in D$ the inclusion $\Omega^1_E(\log x) \subseteq \Omega^1_E(\log D)$ and the inclusion of $\mathbb{C}_x$ as a summand of $\mathcal{O}_D$ form a morphism from (7.13) for the divisor $\{x\}$ to (7.13) for D, compatible with the identity of Ω^1_E. Naturality of the connecting homomorphism gives $\delta(1_x) = \delta_x$, where $\delta_x \in H^1(E, \Omega^1_E)$ is the connecting map for the one-point divisor $\{x\}$ applied to 1. Hence $\delta(r) = \sum_{x\in D} r_x\,\delta_x$, and it suffices to see that δ_x does not depend on x.

Let $x, x' \in E$ and let $t\colon E \to E$ be the translation by $x' - x$, a biholomorphism carrying x to x'. Pullback along t carries forms with a logarithmic pole at x' to forms with a logarithmic pole at x and preserves residues, so t^* is a morphism from (7.13) for $\{x'\}$ to (7.13) for $\{x\}$, acting as the identity on the quotient under the identifications $\mathbb{C}_{x'} \cong \mathbb{C} \cong \mathbb{C}_x$. Naturality gives

$t^*\delta_{x'} = \delta_x$. Now t is homotopic to the identity of E, since E is connected and translations by a path from 0 to $x' - x$ interpolate between them, so t^* is the identity on $H^2(E, \mathbb{C})$. A curve has $h^{2,0} = h^{0,2} = 0$ by example 2.1.2, so $H^2(E, \mathbb{C})$ is the single summand $H^{1,1}(E) = H^1(E, \Omega^1_E)$ and no compatibility of t^* with a decomposition is needed. Hence $\delta_x = \delta_{x'}$.
Step 2: the multiple is nonzero. Write $\delta(r) = (\sum_x r_x)\kappa$ with $\kappa \in H^1(E, \Omega^1_E)$ the common value of the δ_x. If κ were zero then Res would be surjective for every D, and taking $D = \{x\}$ a single point would produce a meromorphic 1-form on E with one simple pole, of residue 1. Its residues then sum to 1, contradicting lemma 7.4.1(1). So $\kappa \neq 0$, the kernel of δ is the hyperplane of sum-zero vectors, and exactness identifies it with the image of Res.
The dimension count follows: $\dim H^0(E, \Omega^1_E(\log D)) = 1 + (n-1) = n$. For the last assertion, take $D = \{p, q\}$; the vector $(1, -1)$ has sum zero, so some ω has $\mathrm{Res}_p\omega = 1$ and $\mathrm{Res}_q\omega = -1$, and two such forms differ by an element of $\ker \mathrm{Res} = \mathbb{C}\, dz$. Such an ω has poles exactly at p and q, both residues being nonzero, completing the proof.

[a] [5, Maximum Principle on Compact Complex Manifolds].

The last form is classically a *differential of the third kind*: a meromorphic 1-form whose only poles are simple, with residues summing to zero. On $E = \mathbb{C}/\Lambda_\tau$ it can also be written out in closed form using the Weierstrass zeta function, the odd primitive ζ of $-\wp$ with $\zeta(z) - 1/z$ holomorphic at 0,[5] whose quasi-periods cancel in the difference $\zeta(z-a) - \zeta(z-b)$; the cohomological argument above is used here because it needs nothing beyond (7.13) and gives the dimension count at the same time.

Proposition 7.4.2 is already the whole answer, because theorem 7.3.10 converts the dimension of $H^0(E, \Omega^1_E(\log D))$ directly into the dimension of $H^1(U, \mathbb{C})$ and into the size of the Hodge filtration. What remains is to locate the single step of the weight filtration, and for that no further geometry is needed: the two filtrations of a mixed Hodge structure constrain each other, and the dimension of F^1 already determines where W jumps.

Theorem 7.4.3: The Mixed Hodge Structure of a Punctured Elliptic Curve

Let $E = \mathbb{C}/\Lambda_\tau$, let $D \subseteq E$ consist of $n \geq 1$ points, let $U = E \setminus D$ and let $j\colon U \to E$ be the inclusion. Give $H^1(U, \mathbb{Q})$ the mixed Hodge structure of theorem 7.3.11, computed from the compactification E and the divisor D. Then

1. $\dim_{\mathbb{Q}} H^1(U, \mathbb{Q}) = n + 1$;
2. writing $H^0(D, \mathbb{Q})_0$ for the kernel of the sum map $\mathbb{Q}^D \to \mathbb{Q}$, the sequence
$$0 \to H^1(E, \mathbb{Q}) \xrightarrow{j^*} H^1(U, \mathbb{Q}) \xrightarrow{\mathrm{Res}} H^0(D, \mathbb{Q})_0(-1) \to 0 \tag{7.14}$$
is exact, and both maps are morphisms of mixed Hodge structures;
3. $W_0H^1(U, \mathbb{Q}) = 0$, $W_1H^1(U, \mathbb{Q}) = j^*H^1(E, \mathbb{Q})$, and $W_2H^1(U, \mathbb{Q}) = H^1(U, \mathbb{Q})$; hence $\mathrm{gr}^W_1 \cong H^1(E, \mathbb{Q})$, pure of weight one with $h^{1,0} = h^{0,1} = 1$, and $\mathrm{gr}^W_2 \cong \mathbb{Q}(-1)^{n-1}$, pure of weight two and of type $(1, 1)$;
4. $F^0H^1(U, \mathbb{C}) = H^1(U, \mathbb{C})$, $F^2H^1(U, \mathbb{C}) = 0$, and $F^1H^1(U, \mathbb{C})$ is the image of $H^0\big(E, \Omega^1_E(\log D)\big)$ under $\eta \mapsto [\eta]$, an injection with n-dimensional image.

[5] [6, The Weierstrass $\wp$-Function, Convergence and Basic Properties of $\wp$].

***Proof*:**
Throughout, for $x \in D$ let $\sigma_x \subseteq U$ be a small circle around x, positively oriented for the complex structure, and let $\mathrm{per}_x(\alpha) = \int_{\sigma_x} \alpha$ be the corresponding period. Since $[\sigma_x] \in H_1(U,\mathbb{Z})$, the functional per_x is $\mathbb{Q}$-valued on $H^1(U,\mathbb{Q})$. By definition 5.4.6, whose recipe applies verbatim to a smooth divisor in a compact complex manifold and which for a point on a curve sends x to the class of σ_x, the residue map on cohomology is $\mathrm{Res}_x = \frac{1}{2\pi i}\mathrm{per}_x$; and for a form η with at most a simple pole at x this agrees with the classical residue, by the Cauchy integral formula or by proposition 5.4.7. It carries $H^1(U,\mathbb{Q})$ into $\mathbb{Q}(-1)^D$, the rational structure of the Tate twist being $(2\pi i)^{-1}\mathbb{Q}$ by definition 3.1.11.
Step 1: dimensions and the Hodge filtration. By theorem 7.2.17 the complex (7.10) computes $H^*(U,\mathbb{C})$, and by theorem 7.3.10 the spectral sequence of its stupid filtration (definition 7.3.2) degenerates at E_1. In degree one this gives

$$\dim_{\mathbb{C}} H^1(U,\mathbb{C}) = \dim_{\mathbb{C}} H^1(E,\mathcal{O}_E) + \dim_{\mathbb{C}} H^0\big(E,\Omega^1_E(\log D)\big)$$

together with $F^1H^1(U,\mathbb{C}) = \mathrm{gr}^1_F H^1(U,\mathbb{C}) \cong H^0(E,\Omega^1_E(\log D))$, the isomorphism being induced by $\eta \mapsto [\eta]$, and $F^2 = 0$ because the complex has no term in degree two. Now $H^1(E,\mathcal{O}_E) \cong H^{0,1}(E)$ is one-dimensional by theorem 1.1.3 and example 2.1.2, and $H^0(E,\Omega^1_E(\log D))$ has dimension n by proposition 7.4.2. This proves (1) and (4); in particular the map $\eta \mapsto [\eta]$ is injective, its source and its image both having dimension n.
Step 2: the sequence (7.14) is exact. Injectivity of j^*: a loop in E can be moved off the finite set D, so $H_1(U,\mathbb{Z}) \to H_1(E,\mathbb{Z})$ is surjective, and dualizing gives injectivity of j^* on H^1 with any field coefficients.

The composite $\mathrm{Res} \circ j^*$ vanishes, because σ_x bounds a disc in E and so is null-homologous there, making $\mathrm{per}_x(j^*\beta) = \int_{\sigma_x} \beta = 0$ for $\beta \in H^1(E,\mathbb{Q})$. The image of Res lies in the sum-zero subspace, because $\sum_{x\in D}[\sigma_x] = 0$ in $H_1(U,\mathbb{Z})$: removing an open disc around each point of D from E leaves a compact oriented surface with boundary $\coprod_x \sigma_x$, contained in U.

For surjectivity onto the sum-zero subspace, complexify and factor through Step 1. If $\eta \in H^0(E,\Omega^1_E(\log D))$ then $\mathrm{per}_x[\eta] = \int_{\sigma_x}\eta = 2\pi i\,\mathrm{Res}_x\eta$, so the composite $H^0(E,\Omega^1_E(\log D)) \to H^1(U,\mathbb{C}) \xrightarrow{\mathrm{Res}} \mathbb{C}^D$ is the sheaf residue map of proposition 7.4.2, whose image is the full sum-zero hyperplane. Hence Res is surjective onto that hyperplane after complexification, and therefore onto $H^0(D,\mathbb{Q})_0(-1)$, a $\mathbb{Q}$-linear map with surjective complexification onto the complexification of a rational subspace being surjective onto that subspace. Counting dimensions,

$$\dim\ker\mathrm{Res} = (n+1) - (n-1) = 2 = \dim j^*H^1(E,\mathbb{Q})$$

and $j^*H^1(E,\mathbb{Q}) \subseteq \ker\mathrm{Res}$, so the two coincide and (7.14) is exact.
Step 3: the weight filtration. By theorem 7.3.11 the weights of $H^1(U,\mathbb{Q})$ lie in $[1,2]$, so $W_0 = 0$ and $W_2 = H^1(U,\mathbb{Q})$; only the single step W_1 is at issue. By proposition 7.3.14 the map j^* is a morphism of mixed Hodge structures out of a structure pure of weight one, so it carries $W_1H^1(E,\mathbb{Q}) = H^1(E,\mathbb{Q})$ into $W_1H^1(U,\mathbb{Q})$; hence $j^*H^1(E,\mathbb{Q}) \subseteq W_1$ and $\dim W_1 \geq 2$.

The reverse inequality comes from counting the Hodge filtration on the graded pieces. Write $w = \dim_{\mathbb{Q}}\mathrm{gr}^W_1 = \dim_{\mathbb{Q}} W_1$, so that $\dim_{\mathbb{Q}}\mathrm{gr}^W_2 = (n+1)-w$ by (1). Since F induces the quotient filtration on each graded piece,

$$\dim_{\mathbb{C}} F^1H^1(U,\mathbb{C}) = \dim_{\mathbb{C}} F^1\mathrm{gr}^W_1 + \dim_{\mathbb{C}} F^1\mathrm{gr}^W_2$$

Now gr^W_1 is pure of weight one by definition 7.1.1, so $F^1\mathrm{gr}^W_1$ is its $(1,0)$-summand, of dimension $w/2$ because conjugation exchanges the two summands. And gr^W_2 is pure of weight two with

$F^2\mathrm{gr}_2^W = 0$, the whole of $F^2H^1(U,\mathbb{C})$ being zero by (4); its $(2,0)$-summand is therefore zero, hence so is its $(0,2)$-summand, and gr_2^W is entirely of type $(1,1)$ with $F^1\mathrm{gr}_2^W = \mathrm{gr}_2^W$. Substituting and using $\dim F^1H^1(U,\mathbb{C}) = n$ from (4),

$$n = \frac{w}{2} + (n+1) - w$$

which gives $w = 2$. So $\dim W_1 = 2 = \dim j^*H^1(E,\mathbb{Q})$ and the inclusion above is an equality, $W_1 = j^*H^1(E,\mathbb{Q})$.

Step 4: the graded pieces and the two morphisms. From Step 3, $\mathrm{gr}_1^W = W_1 \cong H^1(E,\mathbb{Q})$, whose Hodge numbers are $h^{1,0} = h^{0,1} = 1$ by example 3.4.3; and $\mathrm{gr}_2^W = H^1(U,\mathbb{Q})/W_1$, which Step 3 showed to be pure of weight two and entirely of type $(1,1)$, of dimension $n-1$. A weight-two rational Hodge structure of type $(1,1)$ is a direct sum of copies of $\mathbb{Q}(-1)$ (definition 3.1.11), one for each element of a rational basis, so $\mathrm{gr}_2^W \cong \mathbb{Q}(-1)^{n-1}$. This proves (3).

Finally, j^* is a morphism by proposition 7.3.14, and Res is one as well: its target is pure of weight two with F^1 everything and F^2 zero, so compatibility with W asks only $\mathrm{Res}(W_1) = 0$, which holds because $W_1 = j^*H^1(E,\mathbb{Q}) \subseteq \ker \mathrm{Res}$ by Step 2 and Step 3, and compatibility with F asks only $\mathrm{Res}(F^2H^1(U,\mathbb{C})) = 0$, which holds because $F^2 = 0$. Together with the exactness of Step 2 this completes the proof.

The statement is a complete description: the mixed structure on H^1 of a punctured elliptic curve is an extension of $\mathbb{Q}(-1)^{n-1}$ by the pure structure of E, with the Hodge filtration read off the logarithmic differentials. Both graded pieces are forced, and neither depends on where the punctures sit. What the punctures contribute to gr_2^W is only their number, and only through the sum condition of lemma 7.4.1(1), which removes exactly one dimension from $\mathbb{Q}^D$.

That accounting has an immediate consequence at $n = 1$, where removing one dimension from a one-dimensional space leaves nothing at all.

Corollary 7.4.4: A Single Puncture Creates No Weight

Let E be an elliptic curve and $p \in E$. Then

$$j^*\colon H^1(E,\mathbb{Q}) \xrightarrow{\sim} H^1\big(E \setminus \{p\},\mathbb{Q}\big)$$

is an isomorphism of mixed Hodge structures, and $H^1(E \setminus \{p\},\mathbb{Q})$ is pure of weight one. Its Hodge filtration has $F^1 = \mathbb{C}\,dz$.

***Proof*:**

Apply theorem 7.4.3 with $n = 1$. Then $H^0(D,\mathbb{Q})_0 = 0$, so (7.14) makes j^* an isomorphism of mixed Hodge structures; $\mathrm{gr}_2^W = 0$ and $\mathrm{gr}_1^W = H^1(E,\mathbb{Q})$, which is purity of weight one; and F^1 is the image of $H^0(E,\Omega^1_E(\log p))$, which is $\mathbb{C}\,dz$ because proposition 7.4.2 gives it dimension 1 and dz already lies in it, as we sought to show.

The mechanism deserves stating in terms of forms rather than of dimensions. A class of weight two on $E \setminus \{p\}$ would have to be represented by a 1-form with a genuine logarithmic pole at p, that is, with nonzero residue there. Lemma 7.4.1(1) forbids it: the residue at p has no other residue to cancel against. Non-compactness by itself therefore does not create weight; what creates weight is boundary along which a form can have residues that cancel. A single point is not enough boundary.

The same reading explains an earlier computation. The Legendre family of example 5.1.9 was handled through its affine model $y^2 = P(x)$, which is the fibre with one point removed, and the Gauss-Manin computation there never encountered anything of weight two. By corollary 7.4.4 it could not have, and the periods of dx/y computed there are periods of a class on the compact curve.

Two punctures change the answer, and this is the smallest case in which the extension appearing in (7.14) is not determined by its ends. Example 7.1.4 produced an abstract non-split extension by writing down a period; the punctured curve produces one geometrically, and lemma 7.4.1(2) evaluates the period in terms of the two points.

Example 7.4.5: The Extension Class of a Twice-Punctured Elliptic Curve

Let $E = \mathbb{C}/\Lambda_\tau$, let $p \neq q$ in E, let $D = \{p, q\}$ and $U = E \setminus D$. By theorem 7.4.3 the group $H^1(U, \mathbb{Q})$ has dimension 3 and sits in an exact sequence of mixed Hodge structures

$$0 \to H^1(E, \mathbb{Q}) \to H^1(U, \mathbb{Q}) \xrightarrow{\text{Res}} \mathbb{Q}(-1) \to 0 \tag{7.15}$$

where the copy of $\mathbb{Q}(-1)$ is spanned by the residue at p, the residue at q being its negative.

A basis adapted to the sequence. Fix a parallelogram P as in lemma 7.4.1, with interior lifts $\tilde{p}$ of p and $\tilde{q}$ of q, and let γ_1, γ_2 be the two boundary cycles and $\sigma = \sigma_p$ a small positive loop about p. By proposition 7.4.2 there is a form ω on E with simple poles at p, q of residues $+1, -1$, unique up to adding a multiple of dz; normalize it by $\int_{\gamma_1} \omega = 0$, which fixes the multiple since $\int_{\gamma_1} dz = 1$. Put

$$\theta \;:=\; \frac{1}{2\pi i}\,\omega, \qquad u \;:=\; \tilde{p} - \tilde{q} \;\in \mathbb{C}$$

Lemma 7.4.1(2) evaluates the remaining period of ω as $2\pi i\,(\tilde{p}\cdot 1 + \tilde{q}\cdot(-1)) = 2\pi i\, u$, so the period matrix of the three closed forms dz, $d\bar{z}$ and θ against the three cycles is

$$\begin{pmatrix} \int_{\gamma_1} dz & \int_{\gamma_2} dz & \int_{\sigma} dz \\ \int_{\gamma_1} d\bar{z} & \int_{\gamma_2} d\bar{z} & \int_{\sigma} d\bar{z} \\ \int_{\gamma_1} \theta & \int_{\gamma_2} \theta & \int_{\sigma} \theta \end{pmatrix} \;=\; \begin{pmatrix} 1 & \tau & 0 \\ 1 & \bar{\tau} & 0 \\ 0 & u & 1 \end{pmatrix} \tag{7.16}$$

the entry $\int_\sigma \theta = 1$ because $\int_\sigma \omega = 2\pi i \text{Res}_p \omega = 2\pi i$. Its determinant is $\bar{\tau} - \tau = -2i\,\text{Im}\,\tau \neq 0$. Since $\dim H^1(U, \mathbb{C}) = 3 = \dim H_1(U, \mathbb{Q})$, nondegeneracy of (7.16) shows simultaneously that $\{[dz], [d\bar{z}], [\theta]\}$ is a basis of $H^1(U, \mathbb{C})$ and that $\{[\gamma_1], [\gamma_2], [\sigma]\}$ is a basis of $H_1(U, \mathbb{Q})$. A class is rational exactly when its three periods against the latter basis are rational.

When the sequence splits. A splitting of (7.15) in the category of mixed Hodge structures is a morphism $s\colon \mathbb{Q}(-1) \to H^1(U, \mathbb{Q})$ with $\text{Res} \circ s = \text{id}$. Since $\mathbb{Q}(-1)$ is pure of weight 2 with $F^1 = \mathbb{C}$ and $F^2 = 0$, such an s is the same thing as a class

$$\alpha \in H^1(U, \mathbb{Q}) \cap F^1 H^1(U, \mathbb{C}) \quad \text{with} \quad \text{Res}\alpha \neq 0$$

the conditions on W being automatic because W_2 is everything and $\mathbb{Q}(-1)$ has $W_1 = 0$. By theorem 7.4.3(4) the space $F^1 H^1(U, \mathbb{C})$ is spanned by $[dz]$ and $[\theta]$, so write $\alpha = c\,[\theta] + b\,[dz]$ with $c \neq 0$. Reading off (7.16), the periods of α are

$$\int_{\gamma_1} \alpha = b, \qquad \int_{\gamma_2} \alpha = cu + b\tau, \qquad \int_{\sigma} \alpha = c$$

so α is rational precisely when $b \in \mathbb{Q}$, $c \in \mathbb{Q}^\times$ and $cu + b\tau \in \mathbb{Q}$. Such b, c exist if and only if $u \in \mathbb{Q} + \mathbb{Q}\tau$: given $u = r + s\tau$ with $r, s \in \mathbb{Q}$, take $c = 1$ and $b = -s$, which gives $cu + b\tau = r$; conversely $cu + b\tau \in \mathbb{Q}$ with $c \in \mathbb{Q}^\times$ and $b \in \mathbb{Q}$ forces $u \in \mathbb{Q} + \mathbb{Q}\tau$.

The class. The condition $u \in \mathbb{Q} + \mathbb{Q}\tau$ says exactly that the point $u + \Lambda_\tau$ of E is a torsion point, and $u + \Lambda_\tau = p - q$ in the group law of E, since $\tilde{p}$ and $\tilde{q}$ lift p and q. So (7.15) splits over $\mathbb{Q}$ if and only if $p - q$ is a torsion point of E, and the invariant controlling it is

$$e(p, q) \;=\; p - q \;\in\; E \tag{7.17}$$

the image of the degree-zero divisor $p-q$ under the map that sends a difference of points to the class of $\int_q^p dz$ modulo Λ_τ. Over $\mathbb{Z}$ the criterion is sharper. An integral splitting is a class with integral periods and $\int_\sigma \alpha = 1$, so $c = 1$, and rationality of the remaining periods is replaced by $b \in \mathbb{Z}$ and $u + b\tau \in \mathbb{Z}$, that is by $u \in \Lambda_\tau$ and $p = q$. The integral extension of $\mathbb{Z}(-1)$ by $H^1(E, \mathbb{Z})$ therefore never splits for $p \neq q$, and it is only after tensoring with $\mathbb{Q}$ that torsion points become invisible.

Two concrete cases. Take $\tau = i$ and $p = 0$. If $q = \frac{1}{2}$ then $u = -\frac{1}{2} \in \mathbb{Q}$, so the sequence splits over $\mathbb{Q}$; the splitting class is $\alpha = [\theta]$, with periods $(0, -\frac{1}{2}, 1)$, and 2α has integral periods. If instead $q = \frac{1}{\pi}$, then $u = -\frac{1}{\pi}$ is real and irrational, hence not of the form $r + si$ with $r, s \in \mathbb{Q}$, so q is not a torsion point of $\mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}i)$, no splitting exists over $\mathbb{Q}$, and $H^1(U, \mathbb{Q})$ is a non-split extension. The two curves $E \setminus \{0, \frac{1}{2}\}$ and $E \setminus \{0, \frac{1}{\pi}\}$ have the same Betti numbers, the same Hodge numbers, and isomorphic gr^W, and are distinguished by the extension alone.

The example is the geometric form of the phenomenon that example 7.1.4 exhibited by hand. A mixed Hodge structure is not determined by its graded pieces, and here the residual information is a point of E, varying continuously with the punctures while gr^W stays constant. The general statement behind (7.17) is Carlson's description of Ext^1 in the category of mixed Hodge structures: for A pure of weight -1, extensions of $\mathbb{Q}(0)$ by A are classified by $A_\mathbb{C}/(F^0 A_\mathbb{C} + A_\mathbb{Q})$, and the computation above is that quotient written out for $A = H^1(E, \mathbb{Q})(1)$, where it is the Jacobian of E, that is E itself. The general theorem is not needed for the case at hand and is proved in *Period Mappings and Period Domains* [1, extensions of mixed Hodge structures]; the original construction of the mixed structure on a smooth open variety is Deligne's [18].

One curve has now been examined from four positions. Example 3.4.3 gave the pure weight-one structure on $H^1(E, \mathbb{Q})$ and its polarization, with the period ratio τ as the moduli coordinate. Example 5.1.9 put E in the Legendre family and differentiated its periods, producing the Picard-Fuchs equation and the monodromy. Example 6.2.9 computed the Mumford-Tate group of that structure and found it to be all of GL_2 except when E has complex multiplication. And example 7.4.5 leaves the compact curve, where the answer is an extension class living in E rather than a subspace of H^1.

The hypothesis still assumed is smoothness of U. Dropping it means allowing X itself to be singular, where there is no logarithmic complex to filter because there is no smooth ambient space. Chapter 8 replaces X by a simplicial object built out of smooth pieces and runs the same argument on the resulting bicomplex.

Exercise 7.4.1

1. Example 7.3.13 computed the mixed Hodge structure on $H^1(\mathbb{CP}^1 \setminus D, \mathbb{Q})$ for D a set of $n \geq 1$

points, finding it pure of weight two and isomorphic to $\mathbb{Q}(-1)^{n-1}$. Take that as given. For $n = 2$ identify a generator of $H^1(\mathbb{C}^*, \mathbb{Q})$ explicitly as the class of a logarithmic form, and explain why the answer is pure while $H^1(E \setminus \{p\}, \mathbb{Q})$ of theorem 7.4.3 is not, in terms of the weights available on $\mathbb{CP}^1$ and on E.

2. Lemma 7.1.5(2) shows that a morphism between pure structures of different weights vanishes. Use it to deduce that $H^1(E \setminus \{p\}, \mathbb{Q})$ contains no sub-mixed-Hodge-structure isomorphic to $\mathbb{Q}(-1)$, and that there is no nonzero morphism $H^1(E, \mathbb{Q}) \to \mathbb{Q}(-1)$. Explain why this does not contradict the existence of the surjection $H^1(E \setminus \{p\}, \mathbb{Q}) \twoheadrightarrow \mathbb{Q}(-1)$ onto the top graded piece.

3. In the notation of example 7.4.5, suppose $p - q$ is a torsion point of E of exact order N, so that $u = r + s\tau$ with $r, s \in \mathbb{Q}$. Write down a rational class $\alpha \in H^1(U, \mathbb{Q}) \cap F^1H^1(U, \mathbb{C})$ with $\int_\sigma \alpha = 1$, compute its residues at p and at q, and determine the smallest positive integer m for which $m\alpha$ has integral periods against $[\gamma_1], [\gamma_2], [\sigma]$.

4. The affine curve $C_\lambda = \{y^2 = x(x-1)(x-\lambda)\} \subseteq \mathbb{C}^2$ is $E_\lambda \setminus \{\infty\}$ for the Legendre fibre E_λ of example 5.1.9, the point at infinity being the origin of the group law. Use corollary 7.4.4 to show that $H^1(C_\lambda, \mathbb{Q})$ is pure of weight one. Then let $C'_\lambda = C_\lambda \setminus \{(0,0)\}$ and describe the mixed Hodge structure on $H^1(C'_\lambda, \mathbb{Q})$: give its dimension, its graded pieces, and decide whether the resulting extension splits over $\mathbb{Q}$.

5. Keep the notation of example 7.4.5 and let $c \in E$. Using proposition 7.3.14 for the translation $t_c \colon E \to E$, show that the splitting criterion for $D = \{p+c,\ q+c\}$ is the same as for $D = \{p, q\}$; then show that interchanging the roles of p and q replaces $e(p,q)$ by $-e(p,q)$. Conclude that (7.17) depends only on the difference $p - q$ and changes sign under the interchange.

6. Corollary 7.3.12 identifies $W_kH^k(U, \mathbb{Q})$ with the image of $j^* \colon H^k(X, \mathbb{Q}) \to H^k(U, \mathbb{Q})$ for a good compactification X of U. Use it, together with the injectivity of j^* from Step 2 of the proof of theorem 7.4.3, to obtain $W_1H^1(U, \mathbb{Q}) = j^*H^1(E, \mathbb{Q})$ without the count of Hodge numbers in Step 3, and deduce $\dim_{\mathbb{Q}} \operatorname{gr}_2^W H^1(U, \mathbb{Q}) = n - 1$. Say which part of Step 3 this does not replace, and check that the same corollary gives $W_1H^1(V, \mathbb{Q}) = 0$ for the complement V of exercise 7.4.1 (1).

8

Mixed Hodge Structures: Singular Varieties

Chapter 7 kept one hypothesis from the whole of the book before it. The variety was allowed to be non-compact, but it was required to be smooth, and smoothness is what produced the logarithmic complex: a smooth compactification with normal crossings boundary, on which forms with logarithmic poles could be written down and filtered by the order of the pole. A singular variety offers no such object. There is no smooth ambient space in which to take differentials, so there is nothing to filter.

The remedy is to replace the variety by a diagram of smooth ones. A singular X admits a simplicial object $X_\bullet$ of smooth projective varieties mapping to it, with the property that the cohomology of X is computed by the associated bicomplex. Each level of that diagram is smooth and projective, so each carries a pure Hodge structure; the bicomplex inherits two filtrations, one from the simplicial degree and one from the Hodge filtrations on the levels, and the argument of section 7.3 runs again in that setting.

Two things are needed to make this work, and the chapter takes them in order. The first is a formal criterion: given a complex with two filtrations whose weight graded pieces behave like the cohomology of smooth projective varieties, when does it produce a mixed Hodge structure? That criterion is the notion of a cohomological mixed Hodge complex, and section 8.1 proves it does what is wanted. It also settles a debt: the strictness that section 7.3 imported is a consequence of it, and is proved there.

The second is the existence of the diagram. That is a theorem of Deligne resting on resolution of singularities, and like resolution itself it is imported rather than proved. With both in hand, every complex algebraic variety carries a functorial mixed Hodge structure, and the weights sit in a range that reverses the smooth open case: at most k in degree k, rather than at least k. The nodal cubic at the end of the chapter shows the reversal in the smallest example there is.

8.1 Cohomological Mixed Hodge Complexes

The construction of section 7.3 attached two filtrations to a single complex of sheaves and read a mixed Hodge structure off the two spectral sequences they generate. Almost none of that argument used the logarithmic complex. What it used was a short list of properties: that the graded pieces of the weight filtration have pure cohomology, that the Hodge filtration behaves well on those graded pieces, and that a rational structure is present throughout. This section isolates the list, and proves that a bifiltered complex satisfying it puts a mixed Hodge structure on its cohomology. A singular variety has no logarithmic complex, but section 8.2 and section 8.3 will assemble a complex satisfying the list out of smooth pieces, and the theorem below is what converts that assembly into a Hodge structure.

One step of section 7.3 was left open and is closed here. Part (2) of theorem 7.3.10 was reduced there to the strictness of a family of Gysin maps, and that strictness was imported from Deligne's mixed Hodge complex formalism. The formalism is the subject of this section, so the import becomes a theorem; the proof is given at the end of the section.

Two preliminaries are needed before the definitions. The first fixes how a filtration of a complex of *sheaves* is handled. The filtered machinery of [7, The Exact Couple of a Filtered Complex] takes a filtered complex of vector spaces as input, whereas both W and F live on complexes of sheaves. Section 7.3 passed between the two for a single filtration; the statement below records the passage for two, and is the one piece of this section taken on faith.

Theorem 8.1.1: Global Sections of a Filtered Complex of Sheaves

Let X be a topological space and let $\mathcal{K}^\bullet$ be a bounded-below complex of sheaves of $\mathbb{Q}$-vector spaces, or of $\mathbb{C}$-vector spaces, on X, carrying one or two filtrations by subcomplexes, each biregular (finite in each degree). There is a complex $\mathbb{R}\Gamma(X,\mathcal{K}^\bullet)$ of vector spaces over the same field, carrying filtrations by subcomplexes with the same indices and again biregular, functorial in $\mathcal{K}^\bullet$ together with its filtrations, such that:

1. $H^k\big(\mathbb{R}\Gamma(X,\mathcal{K}^\bullet)\big) \cong H^k(X,\mathcal{K}^\bullet)$ for every k, naturally in $\mathcal{K}^\bullet$;
2. $\mathbb{R}\Gamma$ commutes with the filtration steps and their graded pieces: $W_m\mathbb{R}\Gamma(X,\mathcal{K}^\bullet) = \mathbb{R}\Gamma(X,W_m\mathcal{K}^\bullet)$ and $\mathrm{gr}^W_m\mathbb{R}\Gamma(X,\mathcal{K}^\bullet) = \mathbb{R}\Gamma(X,\mathrm{gr}^W_m\mathcal{K}^\bullet)$, compatibly with (1), and likewise for a decreasing filtration;
3. the spectral sequence of a filtration of $\mathbb{R}\Gamma(X,\mathcal{K}^\bullet)$, in the sense of [7, The Exact Couple of a Filtered Complex], has first page $E_1^{p,q} = H^{p+q}(X,\mathrm{gr}^p\mathcal{K}^\bullet)$ and converges to $H^{p+q}(X,\mathcal{K}^\bullet)$, its abutment filtration being the images of the $H^{p+q}(X,F^p\mathcal{K}^\bullet)$;
4. if every term of $\mathcal{K}^\bullet$ and of every step of every filtration is acyclic, then $\mathbb{R}\Gamma(X,\mathcal{K}^\bullet) = \Gamma(X,\mathcal{K}^\bullet)$ with the filtrations by global sections has these properties.

The construction is short, and it is worth giving rather than importing, because the only input is a resolution that is *functorial*. Deligne builds the filtered and bifiltered derived categories for this purpose ([18, Théorie de Hodge II, filtered and bifiltered resolutions], with a modern account in [33, filtered and bifiltered derived categories]); none of that generality is needed here.

***Proof*:**
Let $\mathcal{G}^\bullet(\mathscr{F})$ denote the Godement resolution of a sheaf $\mathscr{F}$, the canonical flasque resolution built from the sheaf of possibly discontinuous sections ([4, Flasque Sheaves]). Two of its properties are the whole argument: it is *functorial* in $\mathscr{F}$, and it is *exact*, so a monomorphism of sheaves induces a monomorphism of resolutions.

The complex. For a complex $\mathcal{K}^\bullet$ let $\mathcal{G}^\bullet(\mathcal{K}^\bullet)$ be the total complex of the double complex obtained by applying $\mathcal{G}$ termwise, and put $\mathbb{R}\Gamma(X,\mathcal{K}^\bullet) := \Gamma\big(X,\mathcal{G}^\bullet(\mathcal{K}^\bullet)\big)$. Every term is flasque, hence acyclic, so this computes hypercohomology (box 7.2.14), which is (1); and functoriality in $\mathcal{K}^\bullet$ is inherited from functoriality of $\mathcal{G}$.

The filtrations. If $F^p\mathcal{K}^\bullet \subseteq \mathcal{K}^\bullet$ is a subcomplex, then exactness of $\mathcal{G}$ makes $\mathcal{G}^\bullet(F^p\mathcal{K}^\bullet) \to \mathcal{G}^\bullet(\mathcal{K}^\bullet)$ a monomorphism, and functoriality makes the inclusions among the F^p commute with it. Taking global sections gives a filtration of $\mathbb{R}\Gamma(X,\mathcal{K}^\bullet)$ by subcomplexes, with

$$F^p\,\mathbb{R}\Gamma(X,\mathcal{K}^\bullet) = \Gamma\big(X,\mathcal{G}^\bullet(F^p\mathcal{K}^\bullet)\big)$$

and gr^p of it is $\Gamma(X,\mathcal{G}^\bullet(\mathrm{gr}^p\mathcal{K}^\bullet))$, again by exactness. Biregularity is inherited degreewise. A second filtration is handled the same way, and the two commute because both are computed by applying $\mathcal{G}$ to subcomplexes of $\mathcal{K}^\bullet$. This is (2), and it is where the argument would fail for a non-functorial choice of resolution: there would be no map relating the resolution of $F^p\mathcal{K}^\bullet$ to that of $\mathcal{K}^\bullet$.

The spectral sequence. The filtration just built is a filtration by subcomplexes, so [7, The Exact Couple of a Filtered Complex] applies, and its first page is the cohomology of $\mathrm{gr}^p\mathbb{R}\Gamma = \Gamma(X,\mathcal{G}^\bullet(\mathrm{gr}^p\mathcal{K}^\bullet))$, which is $H^{p+q}(X,\mathrm{gr}^p\mathcal{K}^\bullet)$ by the acyclicity in the previous step. Convergence and the description of the abutment filtration are [7, Convergence for a Bounded Filtration], biregularity is the hypothesis. That is (3).

Acyclic terms. If every term of $\mathcal{K}^\bullet$ and of every filtration step is already acyclic, the augmentation $\mathcal{K}^\bullet \to \mathcal{G}^\bullet(\mathcal{K}^\bullet)$ is a quasi-isomorphism compatible with the filtrations, so $\Gamma(X,\mathcal{K}^\bullet)$ with its filtrations by global sections has the same three properties. That is (4).

For one filtration, clause (3) is the spectral sequence section 7.3 already used: (7.7) and (7.8) are the two sequences it produces for the Hodge and weight filtrations of the logarithmic complex. Clause (4) is what makes the fine-resolution examples below computable without any appeal to the construction.

The second preliminary converts what the geometry produces into what the two-filtration lemma requires. Theorem 7.3.8 asks for *strictness* of the differentials of the weight spectral sequence, in the sense of [7, Filtered Object and Strict Morphism]: a filtered map $f\colon M \to N$ is strict when $f(M)\cap F^pN = f(F^pM)$ for every p. What Hodge theory gives about a de Rham complex is instead the degeneration of its Hodge spectral sequence at E_1, as in corollary 2.1.3. The two are related, and the direction needed is the following.

The step just used is elementary filtered homological algebra and is proved upstream: the spectral sequence of a biregularly filtered complex degenerates at E_1 exactly when the differential is strictly compatible with the filtration ([7, degeneration at E_1 and strictness]). It is stated there without finiteness hypotheses, biregularity alone sufficing.

That criterion is elementary filtered homological algebra rather than Hodge theory, and it is recorded here because it is the only bridge between the two forms in which the same condition is met: the definitions below state it as strictness, since that is what theorem 7.3.8 requires, while every verification of it in this book runs through a degeneration statement about a Hodge to de Rham spectral sequence.

The pure notion comes first. A Hodge complex of weight n is a complex whose cohomology in degree k is a pure Hodge structure of weight $n+k$, presented so that the rational structure and the Hodge filtration come from different complexes tied together by a comparison.

Definition 8.1.2: Cohomological Hodge Complex of Weight n

Let X be a topological space and $n \in \mathbb{Z}$. A *cohomological Hodge complex of weight n* on X consists of

1. a bounded-below complex $K_{\mathbb{Q}}$ of sheaves of $\mathbb{Q}$-vector spaces on X;
2. a bounded-below complex $K_{\mathbb{C}}$ of sheaves of $\mathbb{C}$-vector spaces on X with a biregular decreasing filtration F by subcomplexes;
3. a *comparison* $\alpha\colon K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} \to K_{\mathbb{C}}$, by which is meant a finite chain of bounded-below complexes of sheaves of $\mathbb{C}$-vector spaces on X,
$$K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} = L_0,\ L_1,\ \dots,\ L_r = K_{\mathbb{C}}$$
together with, for each $i < r$, either a morphism of complexes $L_i \to L_{i+1}$ or a quasi-isomorphism $L_{i+1} \to L_i$;

A quasi-isomorphism induces an isomorphism on hypercohomology (box 7.2.14), so α induces a linear map $H^k(X, K_{\mathbb{Q}}) \otimes_{\mathbb{Q}} \mathbb{C} \to H^k(X, K_{\mathbb{C}})$ for every k: compose the maps induced by the forward links with the inverses of the maps induced by the backwards ones. That composite is the map condition (b) below refers to. The datum is subject to four conditions. Write $F^pH^k(X, K_{\mathbb{C}})$ for the image of $H^k(X, F^pK_{\mathbb{C}})$ in $H^k(X, K_{\mathbb{C}})$.

(a) $H^k(X, K_{\mathbb{Q}})$ is finite dimensional for every k, and zero for all but finitely many k;

(b) α induces an isomorphism $H^k(X, K_{\mathbb{Q}}) \otimes_{\mathbb{Q}} \mathbb{C} \cong H^k(X, K_{\mathbb{C}})$ for every k;

(c) the differential of $\mathbb{R}\Gamma(X, K_{\mathbb{C}})$ is strictly compatible with F;

(d) for every k, the filtration $F^\bullet H^k(X, K_{\mathbb{C}})$ transported through (b) makes $H^k(X, K_{\mathbb{Q}})$ a pure Hodge structure of weight $n+k$, in the filtration form of theorem 3.1.6.

A single morphism of complexes $K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} \to K_{\mathbb{C}}$ is the case $r = 1$ of item (3), and that is what example 8.1.3 exhibits. Longer chains are needed rather than merely permitted. In example 8.1.5 the only map available between the two complexes runs out of $K_{\mathbb{C}}$, the logarithmic complex mapping to the smooth de Rham complex and not the other way, so for those two complexes the comparison is a chain and not a single morphism $K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} \to K_{\mathbb{C}}$; the same is true of the complexes assembled in section 8.3. Item (3) is stated for chains for that reason.

Condition (c) is the one that is never checked directly. By the upstream criterion it holds as soon as the spectral sequence of $(K_{\mathbb{C}}, F)$, which by theorem 8.1.1(3) has first page $H^{p+q}(X, \mathrm{gr}_F^p K_{\mathbb{C}})$,

degenerates at E_1; the finite-dimensionality that lemma needs is (a) together with (b). Condition (d) is where the weight convention sits: the degree k contributes to the weight, so that a complex concentrated in degree 0 has weight equal to the weight of its cohomology, while a shift raises the weight by the amount of the shift (exercise 8.1.1 (3)).

Example 8.1.3: The de Rham Complex of a Smooth Projective Variety

Let Y be a smooth projective variety over $\mathbb{C}$, of dimension d, regarded as a compact Kähler manifold. Take

$$K_{\mathbb{Q}} = \underline{\mathbb{Q}}_Y, \qquad K_{\mathbb{C}} = \mathcal{A}_Y^\bullet, \qquad F^p\mathcal{A}_Y^\bullet = \bigoplus_{r\geq p} \mathcal{A}_Y^{r,\bullet}$$

where $\mathcal{A}_Y^{r,s}$ is the sheaf of smooth forms of type (r,s) and F is the sheaf form of the filtration of definition 1.2.1; it is a filtration by subcomplexes since $d = \partial + \bar{\partial}$ raises r by 0 or 1, and it is biregular because $F^0\mathcal{A}_Y^k = \mathcal{A}_Y^k$ and $F^{k+1}\mathcal{A}_Y^k = 0$. For α take the de Rham resolution $\underline{\mathbb{C}}_Y \to \mathcal{A}_Y^\bullet$, a quasi-isomorphism by the Poincaré lemma.

The four conditions hold, and the weight is 0. For (a), Y is a compact manifold, so $H^k(Y,\mathbb{Q})$ is finite dimensional and vanishes for $k > 2d$. For (b), $H^k(Y,\mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} \cong H^k(Y,\mathbb{C})$ by the universal coefficient theorem,[a] and $H^k(Y,\mathcal{A}_Y^\bullet) \cong H^k(Y,\mathbb{C})$ because α is a quasi-isomorphism (box 7.2.14). For (c), every $\mathcal{A}_Y^{r,s}$ is fine, hence acyclic (theorem 1.1.2), and so is every $F^p\mathcal{A}_Y^\bullet$, being a direct sum of such sheaves; by theorem 8.1.1(4) one may take $\mathbb{R}\Gamma(Y,\mathcal{A}_Y^\bullet) = \mathcal{A}^\bullet(Y)$, whose F-spectral sequence is the Frölicher spectral sequence (theorem 1.2.2). It degenerates at E_1 by corollary 2.1.3, since Y is compact Kähler, so the differential is strict for F by [7, degeneration at E_1 and strictness]. For (d), $F^pH^k(Y,\mathbb{C})$ is the span of the classes representable by forms in F^p (lemma 5.2.4), which is $\bigoplus_{r\geq p} H^{r,k-r}(Y)$; this is the Hodge filtration of the pure Hodge structure of weight k that example 3.1.2 puts on $H^k(Y,\mathbb{Q})$, and $k = 0 + k$.

[a] [30, Universal Coefficient Theorem For Cohomology].

The mixed notion is obtained by asking that the object be filtered by pure ones, one weight at a time, exactly as definition 7.1.1 asks of a vector space.

Definition 8.1.4: Cohomological Mixed Hodge Complex

Let X be a topological space. A *cohomological mixed Hodge complex* on X consists of

1. a bounded-below complex $K_{\mathbb{Q}}$ of sheaves of $\mathbb{Q}$-vector spaces with a biregular increasing filtration W by subcomplexes;
2. a bounded-below complex $K_{\mathbb{C}}$ of sheaves of $\mathbb{C}$-vector spaces with a biregular increasing filtration W and a biregular decreasing filtration F, both by subcomplexes;
3. a comparison $\alpha\colon K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} \to K_{\mathbb{C}}$ in the sense of definition 8.1.2(3), in which every complex L_i of the chain carries a biregular increasing filtration W by subcomplexes, agreeing at the two ends with the given ones, every link carries W_m into W_m for every m, and every backwards link induces a quasi-isomorphism on every graded piece gr^W_m;

Applying gr^W_m to the chain leaves a chain whose backwards links are quasi-isomorphisms, so $\mathrm{gr}^W_m\alpha$ is again a comparison in the sense of definition 8.1.2(3). The datum is subject to the single condition that for every m

$$\big(\mathrm{gr}^W_m K_{\mathbb{Q}},\ (\mathrm{gr}^W_m K_{\mathbb{C}}, F),\ \mathrm{gr}^W_m \alpha\big)$$

be a cohomological Hodge complex of weight m on X, where F on $\mathrm{gr}^W_m K_{\mathbb{C}}$ is the induced filtration $F^p\mathrm{gr}^W_m K_{\mathbb{C}} = \big((F^p K_{\mathbb{C}} \cap W_m K_{\mathbb{C}}) + W_{m-1}K_{\mathbb{C}}\big)/W_{m-1}K_{\mathbb{C}}$.
A *morphism* $f\colon K \to K'$ of cohomological mixed Hodge complexes is a pair of morphisms of complexes of sheaves $f_{\mathbb{Q}}\colon K_{\mathbb{Q}} \to K'_{\mathbb{Q}}$ and $f_{\mathbb{C}}\colon K_{\mathbb{C}} \to K'_{\mathbb{C}}$, each compatible with the filtrations it carries, such that $\alpha' \circ (f_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C})$ and $f_{\mathbb{C}} \circ \alpha$ induce the same map on hypercohomology in every degree, on every step W_m and on every graded piece gr^W_m. When both comparisons are single morphisms of complexes this is the commuting square $\alpha' \circ (f_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}) = f_{\mathbb{C}} \circ \alpha$.
The datum is a *mixed Hodge complex* when X is a point, so that the sheaves are vector spaces and $\mathbb{R}\Gamma$ is the identity.

The definition imposes nothing on how the graded pieces are glued, which is the same feature definition 7.1.1 has and for the same reason: the extension data is where the new invariants live. The purity requirement is what does the work, and the shape of the requirement should be compared with proposition 7.3.7, which established exactly this for the logarithmic complex without naming it.

Example 8.1.5: The Logarithmic Complex as a Mixed Hodge Complex

Let X be smooth projective, $D \subseteq X$ a normal crossings divisor with smooth components, $U = X \setminus D$ and $j\colon U \to X$ the inclusion. Take

$$K_{\mathbb{C}} = \Omega^\bullet_X(\log D), \qquad W = W_\bullet, \qquad F^p = \sigma_{\geq p}$$

the weight filtration by pole order and the filtration by degree, both introduced in section 7.2, and let $K_{\mathbb{Q}}$ be a complex of acyclic sheaves representing $Rj_*\mathbb{Q}_U$, filtered by canonical truncation. For α take the comparison of theorem 7.2.17 read on complexes rather than on cohomology: a chain of quasi-isomorphisms joining $K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$ to $j_*\mathcal{A}^\bullet_U$, which exists because both represent $Rj_*\mathbb{C}_U$, followed by the restriction of forms

$$\Omega^\bullet_X(\log D) \to j_*\mathcal{A}^\bullet_U$$

traversed backwards, a quasi-isomorphism by proposition 7.2.16. This is a chain in the sense of

definition 8.1.2(3) and not a single morphism, the displayed link running out of $K_{\mathbb{C}}$ rather than into it.

The weight filtrations along the chain take one adjustment before definition 8.1.4(3) is met. Every complex between $K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$ and $j_*\mathcal{A}_U^\bullet$ carries the canonical truncation, and every morphism of complexes preserves it, so those links are filtered for that reason alone. The restriction of forms is not: it carries $W_0\Omega_X^\bullet(\log D) = \Omega_X^\bullet$, which is nonzero in every degree from 0 to $\dim X$, into a complex whose step $\tau_{\le 0}$ is concentrated in degree 0. Insert therefore one further complex before $K_{\mathbb{C}}$, namely $\Omega_X^\bullet(\log D)$ with the filtration

$$M_m = \tau_{\le m}\big(W_m\Omega_X^\bullet(\log D)\big)$$

increasing because W and the truncation are, and finite in each degree because M_m is 0 in degrees above m and equals $W_m\Omega_X^k(\log D) = \Omega_X^k(\log D)$ in each degree $k < m$. Restriction of forms carries M_m into $\tau_{\le m}j_*\mathcal{A}_U^\bullet$, since $M_m \subseteq \tau_{\le m}\Omega_X^\bullet(\log D)$ and the truncation is preserved by every morphism, and the identity of $\Omega_X^\bullet(\log D)$ carries M_m into W_m; the first of these is the last backwards link of the chain and the second the last forward one.

Theorem 7.3.9 is what makes those two links quasi-isomorphisms on every graded piece, and it is used here in that form and no other. Read on cohomology sheaves, the correspondence of $W_m\Omega_X^\bullet(\log D)$ with the canonical truncation $\tau_{\le m}Rj_*\mathbb{C}_U$ says that $W_m\Omega_X^\bullet(\log D)$ has no cohomology sheaf in degrees above m, and that in each degree $k \le m$ its cohomology sheaf maps isomorphically to that of $j_*\mathcal{A}_U^\bullet$. The first statement makes the inclusion $M_m \subseteq W_m\Omega_X^\bullet(\log D)$ a quasi-isomorphism, the second makes $M_m \to \tau_{\le m}j_*\mathcal{A}_U^\bullet$ one, and both hold for every m, so the five lemma applied to the long exact sequences of $0 \to M_{m-1} \to M_m \to \mathrm{gr}_m^M \to 0$ and of the two sequences matching it makes each of the two maps a quasi-isomorphism on every graded piece. That the truncation is already defined on $Rj_*\mathbb{Q}_U$ is the second half of the theorem, and it is what makes the rational datum $K_{\mathbb{Q}}$ with its filtration available. Theorem 7.3.9 is imported in section 7.3, and it is what makes definition 8.1.4(3) hold.

This is a cohomological mixed Hodge complex. Fix $m \ge 0$; the graded piece is

$$\mathrm{gr}_m^W K_{\mathbb{C}} \cong a_{m*}\Omega_{D^{(m)}}^{\bullet - m}$$

by proposition 7.3.6, with $a_m \colon D^{(m)} \to X$ the disjoint union of the m-fold intersections, and correspondingly $\mathrm{gr}_m^W K_{\mathbb{Q}}$ represents $a_{m*}\mathbb{Q}_{D^{(m)}}(-m)[-m]$ (theorem 7.3.11, Step 1). Conditions (a) and (b) of definition 8.1.2 hold because $D^{(m)}$ is a smooth projective variety and

$$H^k\big(X, \mathrm{gr}_m^W K_{\mathbb{Q}}\big) \cong H^{k-m}\big(D^{(m)}, \mathbb{Q}\big)(-m)$$

which is finite dimensional, vanishes outside $m \le k \le 2\dim X$, and complexifies to $H^k(X, \mathrm{gr}_m^W K_{\mathbb{C}})$ by proposition 7.3.7(2). For (c), the filtration F induces on $\mathrm{gr}_m^W K_{\mathbb{C}}$ the filtration by degree of $\Omega_{D^{(m)}}^\bullet$ shifted by m (proposition 7.3.7(1)), so its spectral sequence has

$$E_1^{p,q} = H^q\big(D^{(m)}, \Omega_{D^{(m)}}^{p-m}\big) \implies H^{p+q-m}\big(D^{(m)}, \mathbb{C}\big)$$

which is the Hodge to de Rham spectral sequence of $D^{(m)}$ with the filtration index shifted by m. A shift of the index changes neither the differentials nor their vanishing, so the sequence degenerates at E_1 by corollary 2.1.3, and the differential is strict for F by [7, degeneration at E_1 and strictness]. For (d), the Tate twist by $(-m)$ raises the weight by $2m$ and shifts the Hodge filtration by m (definition 3.1.11), so $H^{k-m}(D^{(m)}, \mathbb{Q})(-m)$ is pure of weight $(k-m)+2m = m+k$ and its Hodge filtration is $F^{p-m}H^{k-m}(D^{(m)}, \mathbb{C})$, which is what proposition 7.3.7(1) computes. So $\mathrm{gr}_m^W K$ is a cohomological Hodge complex of weight m, as required. For $m < 0$ the graded piece vanishes, and the condition is vacuous.

One more piece of bookkeeping is needed before the theorem. The proof runs through the lemma on two filtrations, of which theorem 7.3.8 recorded only the clause section 7.3 used. Two further clauses are used below, and all three are collected here in the form in which they are applied; the proof of all three is upstream in [7, the lemma on two filtrations], and nothing is reproved here.

8.1.6 Note: The Lemma on Two Filtrations, in the Form Used Let $A^\bullet$ be a bounded-below complex carrying a biregular increasing filtration W and a biregular decreasing filtration F, both by subcomplexes, and let ${}_WE_r$ be the spectral sequence of W. The filtration F induces three filtrations on each page: the image filtration $F_d^p = \operatorname{im}\big({}_WE_r(F^pA) \to {}_WE_r(A)\big)$, the kernel filtration $F_{d^*}^p = \ker\big({}_WE_r(A) \to {}_WE_r(A/F^pA)\big)$, and the recursive filtration F_{rec}, which on ${}_WE_0$ is the filtration F induces on $\mathrm{gr}^W A^\bullet$ and on ${}_WE_{r+1}$ is the subquotient of F_{rec} on ${}_WE_r$. Then:

1. $F_d \subseteq F_{\mathrm{rec}} \subseteq F_{d^*}$ on every page and on ${}_WE_\infty$, and the three coincide on ${}_WE_0$ and on ${}_WE_1$, with no further hypothesis;
2. if d_s is strictly compatible with F_{rec} for every $s \le r_0$, then the three coincide on ${}_WE_r$ for every $r \le r_0 + 2$;
3. if d_s is strictly compatible with F_{rec} for every $s \ge 0$, then the three coincide on every page and on ${}_WE_\infty$, the differential of $A^\bullet$ is strictly compatible with F, and the spectral sequence of $(A^\bullet, F)$ degenerates at E_1.

Strictness of the d_s is the hypothesis throughout and the coincidence of the filtrations is the conclusion, which is the reading section 7.3 recorded; theorem 7.3.8 is clause (3).

The theorem below is what the definition was built for. Its three conclusions are, in order, the mixed Hodge structure that section 8.3 will put on the cohomology of a singular variety, the degeneration statement that identifies its graded pieces with a page one can compute, and the degeneration statement that section 7.3 left open.

Theorem 8.1.7: A Mixed Hodge Complex Produces a Mixed Hodge Structure

Let K be a cohomological mixed Hodge complex on X. For each k put $H^k = H^k(X, K_{\mathbb{Q}})$ and

$$W_w H^k = \operatorname{im}\Big(H^k\big(X, W_{w-k}K_{\mathbb{Q}}\big) \to H^k\Big), \qquad F^p H^k_{\mathbb{C}} = \operatorname{im}\Big(H^k\big(X, F^p K_{\mathbb{C}}\big) \to H^k(X, K_{\mathbb{C}})\Big)$$

Let ${}_W E_r$ denote the spectral sequence of W on $\mathbb{R}\Gamma(X, K_{\mathbb{Q}})$, so that ${}_W E_1^{p,q} = H^{p+q}(X, \operatorname{gr}^W_{-p} K_{\mathbb{Q}})$ converges to H^{p+q}, and let ${}_F E_r$ denote the spectral sequence of F on $\mathbb{R}\Gamma(X, K_{\mathbb{C}})$. Then:

1. ${}_W E_1^{p,q}$, with its rational structure and the filtration induced by F, is a pure Hodge structure of weight q, and $d_1 \colon {}_W E_1^{p,q} \to {}_W E_1^{p+1,q}$ is a morphism of pure Hodge structures;
2. ${}_W E_r$ degenerates at E_2: $d_r = 0$ for every $r \geq 2$;
3. ${}_F E_r$ degenerates at E_1, so that $\dim_{\mathbb{C}} H^k(X, K_{\mathbb{C}}) = \sum_p \dim_{\mathbb{C}} H^k(X, \operatorname{gr}^p_F K_{\mathbb{C}})$ for every k;
4. $\big(H^k, W_\bullet, F^\bullet\big)$ is a mixed Hodge structure, and for every w

$$\operatorname{gr}^W_w H^k \cong {}_W E_2^{k-w,\, w}$$

as pure Hodge structures of weight w.

***Proof*:**

Write $A_{\mathbb{Q}} = \mathbb{R}\Gamma(X, K_{\mathbb{Q}})$ with its filtration W, and $A_{\mathbb{C}} = \mathbb{R}\Gamma(X, K_{\mathbb{C}})$ with its filtrations W and F; by theorem 8.1.1 these are biregularly filtered complexes of vector spaces with $H^k(A_{\mathbb{Q}}) = H^k$, with $\operatorname{gr}^W_m A_{\mathbb{Q}} = \mathbb{R}\Gamma(X, \operatorname{gr}^W_m K_{\mathbb{Q}})$ and likewise over $\mathbb{C}$, and with the stated first pages. Since the machinery of [7, The Exact Couple of a Filtered Complex] is stated for decreasing filtrations, W is fed to it as $p \mapsto W_{-p}$; this is the convention already used in (7.8).

The three filtrations F_d, F_{rec} and F_{d^*} that F induces on the pages ${}_W E_r(A_{\mathbb{C}})$ are those of box 8.1.6, whose hypotheses hold here: $A_{\mathbb{C}}$ is bounded below and both filtrations are biregular by subcomplexes. By box 8.1.6(1) all three agree on ${}_W E_0$ and on ${}_W E_1$ with no hypothesis, and $F_d \subseteq F_{\mathrm{rec}} \subseteq F_{d^*}$ on every page. Note once and for all that every d_r preserves F_d: the inclusion $F^p A_{\mathbb{C}} \hookrightarrow A_{\mathbb{C}}$ is a morphism of W-filtered complexes, so it induces a morphism of W-spectral sequences commuting with every differential, and F_d^p is by definition its image.

Step 1: the first page is pure. Fix p, q. By definition 8.1.4 the datum $\operatorname{gr}^W_{-p} K$ is a cohomological Hodge complex of weight $-p$, so definition 8.1.2(d) makes

$${}_W E_1^{p,q} = H^{p+q}\big(X, \operatorname{gr}^W_{-p} K_{\mathbb{Q}}\big)$$

a pure Hodge structure of weight $-p + (p+q) = q$, whose Hodge filtration is the image of $H^{p+q}(X, F^{p'} \operatorname{gr}^W_{-p} K_{\mathbb{C}})$ in $H^{p+q}(X, \operatorname{gr}^W_{-p} K_{\mathbb{C}})$, for each index p'. Since $\operatorname{gr}^W_n(F^{p'} A_{\mathbb{C}}) = F^{p'} \operatorname{gr}^W_n A_{\mathbb{C}}$ (both are $(F^{p'} \cap W_n)/(F^{p'} \cap W_{n-1})$), that image is exactly $F_d^{p'}$ on ${}_W E_1^{p,q}(A_{\mathbb{C}})$, hence equals F_{rec} there. Condition (b) of definition 8.1.2 applied to $\operatorname{gr}^W_{-p} K$ gives ${}_W E_1(A_{\mathbb{Q}}) \otimes_{\mathbb{Q}} \mathbb{C} \cong {}_W E_1(A_{\mathbb{C}})$, so the rational structure is the one carried by ${}_W E_1(A_{\mathbb{Q}})$; as extension of scalars is exact and commutes with the passage from one page to the next, the same isomorphism holds on every later page.

Step 2: d_1 is a morphism of pure Hodge structures. The map $d_1\colon {}_WE_1^{p,q} \to {}_WE_1^{p+1,q}$ is the differential of the first page of ${}_WE_r(A_{\mathbb{Q}})$, hence $\mathbb{Q}$-linear, and it preserves $F_d = F_{\mathrm{rec}}$ by the observation above. Source and target are pure of weight q by Step 1, so their weight filtrations both jump only at q and d_1 respects them for trivial reasons. Thus d_1 is a morphism of mixed Hodge structures between pure structures of the same weight, and lemma 7.1.5(1) makes it a morphism of pure Hodge structures of weight q.

Step 3: the second page is pure, and d_1 is strict. By proposition 3.1.8, applied in the rational form justified after lemma 7.1.5, the map d_1 is strictly compatible with the Hodge filtrations, that is, with F_{rec}. By lemma 7.1.5(3) its kernel and image are sub-Hodge structures of weight q and the quotient

$$ {}_WE_2^{p,q} = \ker\bigl(d_1\colon {}_WE_1^{p,q} \to {}_WE_1^{p+1,q}\bigr)/\mathrm{im}\bigl(d_1\colon {}_WE_1^{p-1,q} \to {}_WE_1^{p,q}\bigr) $$

is pure of weight q. The Hodge filtration of that subquotient is the subquotient of the Hodge filtration of ${}_WE_1^{p,q}$, which is F_{rec} on ${}_WE_2$ by the definition of F_{rec}.

Step 4: the higher differentials vanish. The claim, proved by induction on $r \geq 2$, is: (i) d_s is strictly compatible with F_{rec} for every $s \leq r-1$; (ii) ${}_WE_r^{p,q}$ is pure of weight q with Hodge filtration F_{rec}; (iii) the three filtrations agree on ${}_WE_r$.

For $r = 2$: the differential d_0 on ${}_WE_0(A_{\mathbb{C}}) = \mathrm{gr}^W A_{\mathbb{C}}$ is, in the column indexed by n, the differential of $\mathrm{gr}_n^W A_{\mathbb{C}} = \mathbb{R}\Gamma(X, \mathrm{gr}_n^W K_{\mathbb{C}})$ (theorem 8.1.1(2)), and F_{rec} on ${}_WE_0$ is the filtration F induces there, so strictness of d_0 is precisely definition 8.1.2(c) for $\mathrm{gr}_n^W K$; strictness of d_1 is Step 3. Hence (i) holds with $r = 2$, and box 8.1.6(2) with $r_0 = 1$ gives (iii) on ${}_WE_r$ for every $r \leq 3$, in particular on ${}_WE_2$; (ii) is Step 3.

Assume (i), (ii), (iii) at $r \geq 2$ and consider $d_r\colon {}_WE_r^{p,q} \to {}_WE_r^{p+r,\,q-r+1}$. It is $\mathbb{Q}$-linear and preserves F_d, which by (iii) is F_{rec}, and by (ii) that is the Hodge filtration on both sides. Source and target are pure of weights q and $q-r+1$, and $q-(q-r+1) = r-1 \geq 1$, so the weights differ; by lemma 7.1.5(2), $d_r = 0$. A zero map is strict, so (i) holds with r replaced by $r+1$; then ${}_WE_{r+1} = {}_WE_r$ with the same F_{rec}, giving (ii) at $r+1$; and box 8.1.6(2) with $r_0 = r$ gives (iii) on ${}_WE_{r'}$ for $r' \leq r+2$, in particular at $r+1$. This proves (2), and with it ${}_WE_2 = {}_WE_\infty$.

It is the direction of the weight drop that is used: a rational F-compatible map into a pure structure of strictly *larger* weight need not vanish, as exercise 7.3.1 (5) records.

Step 5: the Hodge spectral sequence. Steps 3 and 4 give strict compatibility of d_s with F_{rec} for every $s \geq 0$. This is the hypothesis of box 8.1.6(3), whose conclusion is that the three filtrations agree on every page and on ${}_WE_\infty$, that the differential of $A_{\mathbb{C}}$ is strictly compatible with F, and that the spectral sequence of $(A_{\mathbb{C}}, F)$ degenerates at E_1. That sequence is ${}_FE_r$, with first page $H^{p+q}(X, \mathrm{gr}_F^p K_{\mathbb{C}})$ by theorem 8.1.1(3), so its degeneration is (3); summing the dimensions of $E_\infty^{p,q} = \mathrm{gr}_F^p H^{p+q}(X, K_{\mathbb{C}})$ along $p+q=k$ gives the displayed count.

Step 6: the mixed Hodge structure. The subspaces $W_w H^k$ are $\mathbb{Q}$-subspaces of H^k by construction, increasing in w, and the filtration is finite because W on $K_{\mathbb{Q}}$ is biregular. Each H^k is finite dimensional: the terms ${}_WE_1^{p,q}$ are finite dimensional and only finitely many are nonzero by definition 8.1.2(a) and biregularity, and H^k is a finite iterated extension of subquotients of them. The filtration $F^\bullet H^k_{\mathbb{C}}$ is decreasing and finite for the same reason.

Convergence of ${}_WE_r$ identifies $\mathrm{gr}^W_n H^k$ for the unshifted index n with ${}_WE_\infty^{-n,\,k+n}$. Setting $w = k + n$, so that $W_w H^k$ is the step of index $n = w - k$ named in the statement,

$$\mathrm{gr}^W_w H^k \cong {}_WE_\infty^{k-w,\,w} = {}_WE_2^{k-w,\,w}$$

by Step 4, and Step 4(ii) makes it pure of weight w with Hodge filtration F_{rec}.

What remains is that the filtration F induces on $\mathrm{gr}^W_w H^k$ in the sense of (7.1) is that Hodge filtration. Write G^p for the filtration of (7.1), namely the image of $F^pH^k_{\mathbb{C}} \cap W_wH^k_{\mathbb{C}}$ in $\mathrm{gr}^W_w H^k_{\mathbb{C}}$. Then $F^p_d \subseteq G^p \subseteq F^p_{d^*}$ on ${}_WE_\infty^{k-w,w} = \mathrm{gr}^W_w H^k_{\mathbb{C}}$. For the first inclusion, an element of F^p_d is the class of some $x \in W_wH^k_{\mathbb{C}}$ that is the image of a class in $H^k(X, W_{w-k}F^pK_{\mathbb{C}})$; such an x lies in $F^pH^k_{\mathbb{C}} \cap W_wH^k_{\mathbb{C}}$, so its class lies in G^p. For the second, if $x \in F^pH^k_{\mathbb{C}} \cap W_wH^k_{\mathbb{C}}$ then x is the image of a class in $H^k(X, F^pK_{\mathbb{C}})$, so its image in $H^k(X, K_{\mathbb{C}}/F^pK_{\mathbb{C}})$ vanishes, and a fortiori its class in $\mathrm{gr}^W_w H^k(X, K_{\mathbb{C}}/F^pK_{\mathbb{C}})$ vanishes, which is membership in $F^p_{d^*}$. By Step 5 the outer two filtrations coincide, so G coincides with them and with F_{rec}.

Therefore $\mathrm{gr}^W_w H^k$ with the filtration induced by F is a pure Hodge structure of weight w for every w, which is the requirement of definition 7.1.1, and $(H^k, W_\bullet, F^\bullet)$ is a mixed Hodge structure, completing the proof.

The theorem separates the analytic input from the formal one. Everything analytic sits in definition 8.1.2(c) and (d), which are statements about smooth projective varieties and follow from the Hodge decomposition; the passage from those statements to a mixed Hodge structure is the argument above, which uses no geometry at all. That is what makes the formalism reusable: any construction that produces a bifiltered complex with pure graded pieces produces a mixed Hodge structure, whatever the geometry behind it.

Morphisms are handled by the same mechanism, and this is what section 8.3 will need in order to make the mixed Hodge structure of a singular variety functorial.

Corollary 8.1.8: Functoriality

Let $f\colon K \to K'$ be a morphism of cohomological mixed Hodge complexes on X. Then for every k the induced map

$$H^k(f)\colon H^k(X, K_{\mathbb{Q}}) \to H^k(X, K'_{\mathbb{Q}})$$

is a morphism of the mixed Hodge structures of theorem 8.1.7, and it is strict for both the weight and the Hodge filtrations. Its kernel and image, with the induced filtrations, are mixed Hodge structures, and for every w

$$\mathrm{gr}^W_w\big(\ker H^k(f)\big) = \ker\big(\mathrm{gr}^W_w H^k(f)\big), \qquad \mathrm{gr}^W_w\big(\mathrm{im} H^k(f)\big) = \mathrm{im}\big(\mathrm{gr}^W_w H^k(f)\big)$$

***Proof*:**

The map $f_{\mathbb{Q}}$ carries $W_mK_{\mathbb{Q}}$ into $W_mK'_{\mathbb{Q}}$, so $H^k(X, W_mK_{\mathbb{Q}}) \to H^k(X, K'_{\mathbb{Q}})$ factors through $H^k(X, W_mK'_{\mathbb{Q}})$ by functoriality of hypercohomology; taking images,

$$H^k(f)\big(W_wH^k\big) \subseteq W_wH^k(X, K'_{\mathbb{Q}})$$

for every w, the shift by k being the same on both sides. The same argument with $f_{\mathbb{C}}$ and $F^pK_{\mathbb{C}}$

gives

$$H^k(f)_{\mathbb{C}}\big(F^pH^k_{\mathbb{C}}\big) \subseteq F^pH^k(X, K'_{\mathbb{C}}),$$

where $H^k(f)_{\mathbb{C}}$ is identified with $H^k(f_{\mathbb{C}})$ because f is compatible with the two comparisons (definition 8.1.4), and $H^k(f)$ is defined over $\mathbb{Q}$ because $f_{\mathbb{Q}}$ is. So $H^k(f)$ is a morphism of mixed Hodge structures in the sense of definition 7.1.2. The remaining assertions are parts (1), (2) and (3) of theorem 7.1.8, which apply to any morphism of mixed Hodge structures: strictness for the weight filtrations is part (1), strictness for the Hodge filtrations is part (2), and part (3) gives both the mixed Hodge structures on the kernel and the image and the two displayed identities. This completes the proof.

The debt of section 7.3 can now be paid. That section proved part (1) of theorem 7.3.10 in full and reduced part (2) to the strictness of the Gysin maps δ_s, which it could not establish from the material available there, since Z° and U' are open and their cohomology is not pure. The proof below does not complete that reduction; it replaces it, deriving part (2) from theorem 8.1.7 instead, and the induction on the number of components of D plays no role.

Proof of theorem 7.3.10, part (2), deferred from section 7.3:
By example 8.1.5 the logarithmic complex $\Omega^\bullet_X(\log D)$, with its weight and Hodge filtrations and with the rational structure of theorem 7.3.9, is a cohomological mixed Hodge complex on X. Its Hodge spectral sequence in the sense of theorem 8.1.7 has first page

$${}_FE_1^{p,q} = H^{p+q}\big(X, \mathrm{gr}_F^p\Omega^\bullet_X(\log D)\big) = H^q\big(X, \Omega^p_X(\log D)\big)$$

the second equality because $\mathrm{gr}_F^p\Omega^\bullet_X(\log D)$ is the single sheaf $\Omega^p_X(\log D)$ placed in degree p, so its hypercohomology in degree $p+q$ is $H^q(X, \Omega^p_X(\log D))$. This is the first page of (7.7), and the abutment is $H^{p+q}(X, \Omega^\bullet_X(\log D)) \cong H^{p+q}(U, \mathbb{C})$ by theorem 7.2.17, so the two spectral sequences coincide.

Theorem 8.1.7(3) therefore gives degeneration at E_1, and with it

$$\dim_{\mathbb{C}} H^k(U, \mathbb{C}) = \sum_{p+q=k} \dim_{\mathbb{C}} H^q\big(X, \Omega^p_X(\log D)\big)$$

for every k, which is the assertion of part (2), completing the proof.

Three facts feed that argument, and they line up against the three inputs section 7.3 had. Degeneration of the Hodge to de Rham spectral sequence on each closed stratum $D^{(m)}$ gives the strictness of d_0, through [7, degeneration at E_1 and strictness]; purity of the first page of the weight spectral sequence, which is proposition 7.3.7, gives the strictness of d_1 through proposition 3.1.8; and part (1) of theorem 7.3.10, already proved, gives the vanishing of the higher differentials. The first of the three was out of reach in section 7.3 for one reason: [7, degeneration at E_1 and strictness] was not available there to convert the degeneration on each stratum into strictness of d_0. The induction on components asked instead for strictness of the Gysin maps δ_s, which run between the cohomology groups of open varieties, so proposition 3.1.8 does not reach them.

Exercise 8.1.1

1. Let $V = (V_{\mathbb{Q}}, F^\bullet)$ be a pure Hodge structure of weight n in the filtration form of theorem 3.1.6. Regard $V_{\mathbb{Q}}$ and $V_{\mathbb{C}}$ as complexes concentrated in degree 0 on a one-point space, with F the

given filtration and α the identity. Show that this is a cohomological Hodge complex of weight n. Show further that, taking $W_m K_{\mathbb{Q}} = 0$ for $m < n$ and $W_m K_{\mathbb{Q}} = V_{\mathbb{Q}}$ for $m \geq n$, it becomes a mixed Hodge complex whose associated mixed Hodge structure in the sense of theorem 8.1.7 is V read as a mixed structure through example 7.1.3(1).

2. The criterion [7, degeneration at E_1 and strictness] is an equivalence, so both directions are available. Extract from it the intermediate step: show that if the differential of $A^\bullet$ is strictly compatible with F, then $H^n(F^pA^\bullet) \to H^n(A^\bullet)$ is injective for every p and n, and that the long exact sequence of $0 \to F^{p+1}A \to F^pA \to \mathrm{gr}_F^p A \to 0$ then splits into short exact sequences. Say which of the two directions of the criterion you have used.

3. Let K be a cohomological Hodge complex of weight n on X and let $K[1]$ be the shift, with $K[1]^i = K^{i+1}$, differential $-d$, and the filtration $F^pK[1]^i = F^pK^{i+1}$. Show that $K[1]$ is a cohomological Hodge complex of weight $n+1$, and that no shift of the index of F is needed to make this true.

4. Let K be a cohomological mixed Hodge complex on X and let $n \in \mathbb{Z}$. Define $K(n)$ by multiplying the comparison α by $(2\pi i)^n$, that is, by prepending to its chain the isomorphism $(2\pi i)^n \cdot \mathrm{id}$ of $K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$, and setting $W_m K(n) = W_{m+2n}K$ and $F^pK(n)_{\mathbb{C}} = F^{p+n}K_{\mathbb{C}}$. Show that $K(n)$ is again a cohomological mixed Hodge complex, and that the mixed Hodge structure it produces on H^k is the Tate twist $H^k(X, K_{\mathbb{Q}})(n)$ of definition 3.1.11.

5. Let K be a cohomological mixed Hodge complex on X with $\mathrm{gr}_m^W K = 0$ for every $m \neq n$. Show that the mixed Hodge structure of theorem 8.1.7 on $H^k(X, K_{\mathbb{Q}})$ is pure of weight $n+k$, and that K is a cohomological Hodge complex of weight n in the sense of definition 8.1.2.

6. Let $f\colon K \to K'$ be a morphism of cohomological mixed Hodge complexes for which $H^k(f)$ is injective. Using corollary 8.1.8, show that

$$W_w H^k(X, K_{\mathbb{Q}}) = H^k(X, K_{\mathbb{Q}}) \cap W_w H^k(X, K'_{\mathbb{Q}}),$$
$$F^p H^k(X, K_{\mathbb{C}}) = H^k(X, K_{\mathbb{C}}) \cap F^p H^k(X, K'_{\mathbb{C}}),$$

   where $H^k(X, K_{\mathbb{Q}})$ is identified with its image. Deduce that its mixed Hodge structure is the one induced from $H^k(X, K'_{\mathbb{Q}})$, so that it is a sub-mixed Hodge structure in the sense of definition 7.1.2.

7. Take $X = \mathbb{C}\mathrm{P}^1$ and $D = \{0\} \cup \{\infty\}$, so that $U = \mathbb{C}^*$. Using example 8.1.5 and proposition 7.3.7, write down ${}_W E_1^{-m,\,1+m}$ for every m, identify the differential $d_1\colon {}_W E_1^{-1,2} \to {}_W E_1^{0,2}$ as a map of pure Hodge structures, and compute ${}_W E_2^{-m,\,1+m}$. Conclude that $\mathrm{gr}_w^W H^1(\mathbb{C}^*, \mathbb{Q})$ vanishes for $w \neq 2$ and is one dimensional for $w = 2$, so that $H^1(\mathbb{C}^*, \mathbb{Q}) \cong \mathbb{Q}(-1)$, as section 7.1 asserted.

8. Degeneration at E_2 in theorem 8.1.7(2) cannot be improved to E_1. On a one-point space let $K_{\mathbb{Q}}$ be the complex $\mathbb{Q} \xrightarrow{\mathrm{id}} \mathbb{Q}$ in degrees 0 and 1, let $K_{\mathbb{C}} = K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$ with α the identity, $F^0K_{\mathbb{C}} = K_{\mathbb{C}}$ and $F^1K_{\mathbb{C}} = 0$, and let W be given by $W_{-2}K = 0$, $W_{-1}K = (0 \to \mathbb{Q})$ concentrated in degree 1, and $W_0K = K$. Verify that this is a mixed Hodge complex, compute ${}_W E_1^{p,q}$ for all p, q, and show that d_1 is nonzero while ${}_W E_2 = 0$.

8.2 Simplicial and Cubical Hyperresolutions

A word first on what is now being assumed, since it is less than at any earlier point in the book. Every variety so far has been smooth, and almost every one compact; from here neither holds. By a *complex algebraic variety* is meant a reduced separated scheme of finite type over $\mathbb{C}$: reduced, so that functions are determined by their values, and separated, so that limits are unique, but with no smoothness and no irreducibility. The scheme-theoretic notions are those of [4, k-Variety], with one deliberate difference. That book requires a k-variety to be irreducible; irreducibility is dropped here because the terms of a hyperresolution are disjoint unions, reducible by construction, and the whole apparatus of section 8.2 would be unavailable under the stricter convention. Such an X has a singular locus X_{sing}, a closed subvariety, and its complement $X \setminus X_{\mathrm{sing}}$ is a complex manifold to which Part I applies.

Cohomology means singular cohomology of the associated analytic space X^{an} with its Euclidean topology, not the Zariski topology: the Zariski topology is too coarse to have interesting singular cohomology, and it is the analytic topology that chapter 7 used throughout. The two are compared by GAGA (theorem 1.2.6) when X is projective and smooth, which is the only case where the comparison has been needed so far.

One further tool is imported. The *normalization* $\nu\colon \widetilde{X} \to X$ of an irreducible X is a finite surjective birational morphism from a normal variety, an isomorphism over the normal locus ([4, normalization of a variety]); for a curve, normal is the same as smooth, so $\widetilde{C}$ is a smooth curve. That is the only property used below.

Theorem 8.1.7 converts a cohomological mixed Hodge complex into a mixed Hodge structure and says nothing about where such a complex is to be found. For a smooth variety U, chapter 7 found one on a smooth compactification, as the logarithmic complex with its two filtrations. A singular X has no smooth ambient space to carry a complex of sheaves, and this section constructs what replaces it: a diagram of smooth varieties mapping to X, projective when X is complete, together with the statement that the cohomology of X is computed by that diagram.

The guiding case is a curve C with one node p, and its normalization $\pi\colon \widetilde{C} \to C$. The normalization is smooth and projective, so its cohomology is pure, but it is the wrong answer: π identifies the two points of $\pi^{-1}(p)$, and that identification produces a cohomology class the normalization does not carry. For the nodal cubic below, $H^1(\widetilde{C}, \mathbb{Q}) = 0$ while $H^1(C, \mathbb{Q})$ has rank one. What is missing is the identification data, and recording it means keeping the diagram

$$\pi^{-1}(p) \rightrightarrows \widetilde{C} \sqcup \{p\}$$

of smooth projective varieties over C, and not $\widetilde{C}$ alone. Iterating this device across all singularities and all dimensions produces the objects of this section.

Definition 8.2.1: Strict Simplicial Variety

A *strict simplicial complex variety* $X_\bullet$ consists of complex algebraic varieties X_p for every $p \geq 0$, together with morphisms

$$\delta_i \colon X_p \to X_{p-1}, \qquad 0 \leq i \leq p$$

for every $p \geq 1$, called the *face maps*, subject to the identities

$$\delta_i \circ \delta_j = \delta_{j-1} \circ \delta_i \qquad (0 \leq i < j \leq p) \tag{8.1}$$

as morphisms $X_p \to X_{p-2}$, for every $p \geq 2$.

An *augmentation* of $X_\bullet$ to a complex algebraic variety X is a morphism $a \colon X_0 \to X$ satisfying $a \circ \delta_0 = a \circ \delta_1$ as morphisms $X_1 \to X$. The datum is written $a \colon X_\bullet \to X$ and called an *augmented* strict simplicial variety over X.

The word *strict* marks the omission: a simplicial object in the usual sense carries degeneracy morphisms $\sigma_i \colon X_p \to X_{p+1}$ alongside the faces, satisfying further identities.[1] The augmentation condition $a\delta_0 = a\delta_1$ is the coequalizer condition: it says the two faces of X_1 become equal after mapping to X, which is exactly what makes X_1 record identifications performed by a.

Example 8.2.2: The Simplicial Object of a Nodal Cubic

Let $C = \{y^2z = x^2(x+z)\} \subseteq \mathbb{CP}^2$, an irreducible cubic with a single node at $p = [0:0:1]$, and let $\pi \colon \widetilde{C} \to C$ be its normalization ([4, normalization of a variety]). Projecting from the node parametrizes C: in the affine chart $z = 1$ the substitution $y = tx$ into $y^2 = x^2(x+1)$ gives $t^2x^2 = x^2(x+1)$, hence $x = t^2 - 1$ and $y = t(t^2-1)$, so

$$\pi\big([u:v]\big) = \big[(u^2 - v^2)v : u(u^2 - v^2) : v^3\big], \qquad t = u/v$$

is a morphism $\mathbb{CP}^1 \to C$. It sends $[1:0]$ to $[0:1:0]$, and for $v \neq 0$ the coordinate t is recovered from the image as y/x whenever $x \neq 0$, so π is injective off the two points $p_1 = [1:1]$ and $p_2 = [-1:1]$, both of which map to p. Thus $\widetilde{C} = \mathbb{CP}^1$ and $\pi^{-1}(p) = \{p_1, p_2\}$.

Set

$$X_0 = \widetilde{C} \sqcup \{p\}, \qquad X_1 = \{p_1, p_2\}, \qquad X_p = \emptyset \ \ (p \geq 2)$$

with $\delta_0 \colon X_1 \to X_0$ the constant morphism to $\{p\}$, with $\delta_1 \colon X_1 \to X_0$ the inclusion $\{p_1, p_2\} \hookrightarrow \widetilde{C}$, and with all other face maps the unique morphisms out of $\emptyset$. The identities (8.1) hold vacuously, since every composite $X_p \to X_{p-2}$ has source $\emptyset$ for $p \geq 2$. The augmentation $a \colon X_0 \to C$ is π on $\widetilde{C}$ and the inclusion on $\{p\}$, and $a\delta_0 = a\delta_1$ holds because both composites send p_1 and p_2 to p. So $a \colon X_\bullet \to C$ is an augmented strict simplicial variety over C, built out of smooth projective pieces.

A simplicial variety over X is only useful if its cohomology reconstructs that of X. The reconstruction is through a double complex: one direction is the cohomology of each X_p, the other is the alternating sum of the face maps, and the condition to be imposed is that the total complex compute $H^*(X, \mathbb{Q})$.

[1] The objects constructed in this section arise from cubical diagrams and carry no degeneracies. Nothing below uses them: the descent complex of definition 8.2.3 is assembled from the face maps alone.

Singular cochains are used because they are strictly functorial for continuous maps, so the face maps induce a cosimplicial structure with no choices to make.

Definition 8.2.3: The Descent Complex and Cohomological Descent

Let $a\colon X_\bullet \to X$ be an augmented strict simplicial variety and write $S^\bullet(Y)$ for the complex of singular cochains of a topological space Y with coefficients in $\mathbb{Q}$, with coboundary δ. The *descent double complex* of $X_\bullet$ is

$$K^{p,q} = S^q(X_p) \quad (p,q \geq 0), \qquad K^{p,q} = 0 \text{ otherwise}$$

with differentials

$$d' = \sum_{i=0}^{p+1} (-1)^i\, \delta_i^*\colon K^{p,q} \to K^{p+1,q}, \qquad d'' = (-1)^p\, \delta\colon K^{p,q} \to K^{p,q+1}$$

Its total complex $S^\bullet(X_\bullet)$ has $S^m(X_\bullet) = \bigoplus_{p+q=m} K^{p,q}$ and differential $d_{\mathrm{tot}} = d' + d''$.

The augmentation induces $a^*\colon S^\bullet(X) \to S^\bullet(X_\bullet)$ with image in the column $p = 0$. The augmented simplicial variety is *of cohomological descent* over X if

$$a^*\colon H^k(X,\mathbb{Q}) \to H^k\big(S^\bullet(X_\bullet)\big)$$

is an isomorphism for every k.

Three verifications are owed, and each is a direct calculation. First, $d' \circ d' = 0$. Expanding,

$$d'd' = \sum_{j=0}^{p+2} \sum_{i=0}^{p+1} (-1)^{i+j} \left(\delta_i \circ \delta_j\right)^*$$

where $\delta_j\colon X_{p+2} \to X_{p+1}$ and $\delta_i\colon X_{p+1} \to X_p$. The substitution $(i,j) \mapsto (j-1,i)$ is a bijection from the index pairs with $i < j$ onto those with $i \geq j$, it reverses the sign $(-1)^{i+j}$, and by (8.1) it leaves $(\delta_i \circ \delta_j)^*$ unchanged; the terms therefore cancel in pairs. Second, $d'd'' + d''d' = 0$: the singular coboundary commutes with pullback along a continuous map, so $\delta_i^*\delta = \delta\delta_i^*$, and the two composites carry the signs $(-1)^p$ and $(-1)^{p+1}$. Hence $d_{\mathrm{tot}}^2 = d'd' + (d'd'' + d''d') + d''d'' = 0$. Third, a^* is a map of complexes: for $\xi \in S^q(X)$ the component $d'(a^*\xi) = (\delta_0^* - \delta_1^*)a^*\xi = \big((a\delta_0)^* - (a\delta_1)^*\big)\xi$ vanishes by the augmentation condition, and d'' restricted to the column $p = 0$ is δ, so $d_{\mathrm{tot}} \circ a^* = a^* \circ \delta$.

Simplicial varieties of cohomological descent are constructed from finite diagrams rather than exhibited term by term. A cube of varieties over X carries the same data as a simplicial object whose X_p is the disjoint union of the vertices at level p, and the cube has finitely many vertices, so the construction terminates.

Definition 8.2.4: Cubical Varieties and Cubical Hyperresolutions

Fix $r \geq 0$ and let $\square_r$ be the set of subsets of $\{0, 1, \dots, r\}$, ordered by inclusion. An *augmented r-cubical variety* $X_\bullet$ assigns a complex algebraic variety X_α to each $\alpha \in \square_r$ and a morphism $X_\beta \to X_\alpha$ to each inclusion $\alpha \subseteq \beta$, so that these morphisms compose correctly and each $X_\alpha \to X_\alpha$ is the identity. It is *augmented over* $X = X_\emptyset$.

Its *associated strict simplicial variety* is

$$X_p = \coprod_{|\alpha|=p+1} X_\alpha \qquad (p \geq 0)$$

where δ_i restricts, on the component indexed by $\alpha = \{s_0 < s_1 < \cdots < s_p\}$, to the structure morphism $X_\alpha \to X_{\alpha \setminus \{s_i\}}$, and where the augmentation $a \colon X_0 \to X_\emptyset$ is given by the structure morphisms.

A *cubical hyperresolution* of a complex algebraic variety X of dimension n is an augmented r-cubical variety, for some $r \geq 0$, with $X_\emptyset = X$ and

1. X_α smooth for every $\alpha \neq \emptyset$; the hyperresolution is called *projective* if moreover every such X_α is projective;
2. $\dim X_\alpha \leq n - |\alpha| + 1$ for every $\alpha \neq \emptyset$, with the convention $\dim \emptyset = -1$;
3. the associated augmented strict simplicial variety of cohomological descent over X.

The associated object really is strict simplicial. On the component indexed by $\alpha = \{s_0 < \cdots < s_p\}$ with $i < j$, the composite $\delta_i \circ \delta_j$ first deletes s_j and then deletes the i-th element of $\alpha \setminus \{s_j\}$, which is still s_i because $i < j$; the composite $\delta_{j-1} \circ \delta_i$ first deletes s_i and then deletes the $(j-1)$-st element of $\alpha \setminus \{s_i\}$, which is s_j because deleting s_i lowers by one the index of every element after it. Both composites are the structure morphism $X_\alpha \to X_{\alpha \setminus \{s_i, s_j\}}$, so (8.1) holds. Similarly, on the component indexed by $\alpha = \{s_0 < s_1\}$ both $a\delta_0$ and $a\delta_1$ are the structure morphism $X_\alpha \to X_\emptyset$, by compatibility with composition. Condition (2) has a consequence used repeatedly below: taking $|\alpha| = n + 2$ forces $\dim X_\alpha \leq -1$, so $X_\alpha = \emptyset$, and therefore $X_p = \emptyset$ for $p > n$ as well as for $p > r$. The smallest instance is the case $r = 1$: the square formed by a nodal curve, its normalization, the node and the two points above it, computed in example 8.2.8.

This section takes two results on faith: existence of hyperresolutions, stated next, and the descent theorem for a blow-up square, proposition 8.2.6, by which they are built. Existence is not a formal consequence of the definitions: producing smooth X_α out of an arbitrary singular X requires resolution of singularities, which is the same input chapter 7 took as theorem 7.2.3.

Theorem 8.2.5: Existence of Cubical Hyperresolutions

Every complex algebraic variety X admits a cubical hyperresolution in the sense of definition 8.2.4. If X is complete, it admits a projective one, that is, one whose terms X_α with $\alpha \neq \emptyset$ are all smooth projective varieties.

No proof is given here, and none is given anywhere in this series. The statement rests on Hironaka's theorem on resolution of singularities in characteristic zero [24], imported into this book as theo-

rem 7.2.3 and used there for the same reason: it is the only source of smooth models for a variety that has none. Granted resolution, the cube is built by induction on dimension, each step replacing a variety by a resolution together with the locus over which the resolution fails to be an isomorphism. If X is complete then every variety produced by that induction is complete again, and each resolution occurring in it can be taken projective, because a complete variety admits a proper modification by a smooth projective variety; that is the second clause. The simplicial form of the construction is Deligne's [19]; the cubical form stated here, with the dimension bound of definition 8.2.4(2), is due to Guillén, Navarro Aznar, Pascual Gainza and Puerta, and a full account is in [33, cubical hyperresolutions].

Completeness of X is a real hypothesis in the second clause, not an artifact of the construction. Take $X = \mathbb{C}^*$, of dimension $n = 1$, which deformation retracts onto the unit circle and so has $H^1(X, \mathbb{Q}) \cong \mathbb{Q}$,[2] and suppose $X_\bullet$ were a cubical hyperresolution of X with every X_α, $\alpha \neq \emptyset$, projective. Then $X_0 = \coprod_{|\alpha|=1} X_\alpha$ is a finite disjoint union of projective varieties, and composing the augmentation $a\colon X_0 \to \mathbb{C}^*$ with the inclusion $\mathbb{C}^* \subseteq \mathbb{C}$ exhibits a regular function on each connected component of X_0, necessarily constant. So a factors as $X_0 \xrightarrow{b} S \xrightarrow{\iota} \mathbb{C}^*$ through the set $S = a(X_0)$, with b continuous, and S is finite because X_0 has finitely many connected components. Since ι is injective, $a\delta_0 = a\delta_1$ forces $b\delta_0 = b\delta_1$, so the third of the three verifications following definition 8.2.3 applies to b and makes $b^*\colon S^\bullet(S) \to S^\bullet(X_\bullet)$ a map of complexes with $a^* = b^* \circ \iota^*$. On H^1 the composite factors through $H^1(S, \mathbb{Q}) = 0$, so a^* is zero on $H^1(\mathbb{C}^*, \mathbb{Q}) = \mathbb{Q}$ and is not an isomorphism: no cubical hyperresolution of $\mathbb{C}^*$ has projective terms. The definition without that requirement is satisfied by $\mathbb{C}^*$ itself, with $r = 0$ and $X_{\{0\}} = \mathbb{C}^*$.

The inductive step just described is a statement about a single square. Every example in this chapter is an instance of it, and it is the mechanism by which theorem 8.2.5 produces its cube. It too is imported.

Proposition 8.2.6: Imported: Descent for a Blow-Up Square

Let $\rho\colon \widetilde{X} \to X$ be a proper morphism of complex algebraic varieties, let $Z \subseteq X$ be a closed subvariety and set $E = \rho^{-1}(Z)$ with its reduced structure. Suppose ρ restricts to an isomorphism $\widetilde{X} \setminus E \to X \setminus Z$. Then the augmented 1-cubical variety

$$X_\emptyset = X, \qquad X_{\{0\}} = \widetilde{X}, \qquad X_{\{1\}} = Z, \qquad X_{\{0,1\}} = E$$

whose structure morphisms are ρ, the inclusion $Z \hookrightarrow X$, the inclusion $E \hookrightarrow \widetilde{X}$ and the restriction $E \to Z$ of ρ, is of cohomological descent over X.

No proof is given here. This is Deligne's descent theorem for a proper modification [19]; the form stated here, for singular cohomology with $\mathbb{Q}$ coefficients, is [33, cohomological descent for discrete and blow-up squares]. Its content is that the only cohomology lost in passing from X to $\widetilde{X}$ is what the pair (Z, E) records. Note the hypotheses: ρ must be proper, and E must be the *full* preimage of Z, not merely a closed subvariety over which ρ fails to be injective.

Filtering the total complex of a hyperresolution by columns produces a spectral sequence whose first page is built from smooth varieties alone, and, when the hyperresolution is projective, from smooth projective ones, so that the Hodge theory of chapter 2 applies to every entry of that page.

[2] [30, Homology Groups of Spheres, Universal Coefficient Theorem For Cohomology].

Proposition 8.2.7: The Descent Spectral Sequence

Let X be a complex algebraic variety of dimension n, let $X_\bullet$ be a cubical hyperresolution of X and let $a\colon X_\bullet \to X$ denote its associated augmented strict simplicial variety.

1. Each X_p is a disjoint union of finitely many smooth varieties of dimension at most $n-p$, projective if $X_\bullet$ is a projective hyperresolution, and $X_p = \emptyset$ for $p > n$. Consequently $H^q(X_p, \mathbb{Q}) = 0$ unless $0 \le p \le n$ and $0 \le q \le 2(n-p)$.
2. There is a spectral sequence
$$E_1^{p,q} = H^q(X_p, \mathbb{Q}) \Longrightarrow H^{p+q}(X, \mathbb{Q}) \tag{8.2}$$
whose differential $d_1\colon E_1^{p,q} \to E_1^{p+1,q}$ is $\sum_{i=0}^{p+1}(-1)^i \delta_i^*$, and whose induced filtration on $H^k(X, \mathbb{Q})$ is finite.
3. Suppose in addition that $X_\bullet$ is projective, which by theorem 8.2.5 may be arranged whenever X is complete. Then each $E_1^{p,q}$ carries a pure Hodge structure of weight q, and d_1 is a morphism of Hodge structures.
4. Under the additional hypothesis of (3), each $E_2^{p,q}$ carries a pure Hodge structure of weight q.

***Proof*:**
Throughout, $K^{p,q} = S^q(X_p)$ denotes the descent double complex of definition 8.2.3, and $S^\bullet(X_\bullet)$ its total complex.

1. By definition 8.2.4, X_p is the disjoint union of the X_α with $|\alpha| = p+1$, a finite set of indices, and each such X_α is smooth of dimension at most $n-(p+1)+1 = n-p$, and projective as well when $X_\bullet$ is a projective hyperresolution. Taking $|\alpha| = p+1 \ge n+2$ gives $\dim X_\alpha \le -1$, so $X_\alpha = \emptyset$ and hence $X_p = \emptyset$ for $p > n$; this also makes $H^q(X_p, \mathbb{Q}) = 0$ for every q when $p > n$, and $H^q(X_p, \mathbb{Q}) = 0$ for $q < 0$ by definition. For the remaining bound, a smooth variety of dimension d is a complex manifold, hence a smooth real manifold of dimension $2d$, on which there are no nonzero differential forms of degree exceeding $2d$; de Rham's theorem ([32, the de Rham theorem]) then gives $H^q(X_p, \mathbb{R}) \cong H^q_{\mathrm{dR}}(X_p) = 0$ for $q > 2(n-p)$. Rational cohomology vanishes with it: cohomology with coefficients in a field is the dual of homology with those coefficients, by universal coefficients, so $H^q(X_p, \mathbb{R}) = 0$ forces $H_q(X_p, \mathbb{R}) = 0$; the singular chain complex with real coefficients is the one with rational coefficients extended by $\mathbb{R}$, and extension of scalars along a field extension is exact, so $H_q(X_p, \mathbb{R}) = H_q(X_p, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R}$ and hence $H_q(X_p, \mathbb{Q}) = 0$, whence $H^q(X_p, \mathbb{Q}) = 0$. Compactness enters nowhere in this part, so the bound holds whether or not the hyperresolution is projective.
2. Filter the total complex by columns,
$$F^s S^m(X_\bullet) = \bigoplus_{\substack{p+q=m \\ p \ge s}} K^{p,q}$$
Since d' raises p by one and d'' preserves it, each F^s is a subcomplex, and the filtration is decreasing with $F^0 S^m(X_\bullet) = S^m(X_\bullet)$. By part (1), $K^{p,q} = 0$ for $p > n$, so $F^{n+1} S^m(X_\bullet) =$

0: the filtration is finite in each degree, hence exhaustive and bounded. The spectral sequence of a filtered cochain complex, together with its convergence under a bounded filtration, is [7, convergence for a decreasing filtration of a cochain complex]; it gives a spectral sequence with $E_0^{p,q} = K^{p,q}$ and d_0 the differential induced by d_{tot} on the graded pieces, which is $d'' = (-1)^p\delta$. Since a sign does not change cohomology,

$$E_1^{p,q} = H^q\big(K^{p,\bullet}, d''\big) = H^q\big(S^\bullet(X_p)\big) = H^q(X_p, \mathbb{Q})$$

and the induced differential d_1 is the one induced by d', namely $\sum_{i=0}^{p+1}(-1)^i\delta_i^*$ on singular cohomology. The filtration on $H^m(S^\bullet(X_\bullet))$ is finite because the filtration of the complex is. Finally, $X_\bullet$ is of cohomological descent (definition 8.2.4(3)), so $a^*\colon H^m(X, \mathbb{Q}) \to H^m(S^\bullet(X_\bullet))$ is an isomorphism, and the spectral sequence converges to $H^{p+q}(X, \mathbb{Q})$ as claimed in (8.2).

3. Fix p with $X_p \neq \emptyset$ and write $X_p = \coprod_\alpha X_\alpha$. Each X_α is smooth projective, the hyperresolution being projective by hypothesis, hence compact Kähler (chapter 1), so theorem 2.1.1 gives $H^q(X_\alpha, \mathbb{C}) = \bigoplus_{r+s=q} H^{r,s}(X_\alpha)$ with $\overline{H^{r,s}(X_\alpha)} = H^{s,r}(X_\alpha)$; since $H^q(X_\alpha, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} = H^q(X_\alpha, \mathbb{C})$, this is a pure Hodge structure of weight q on $H^q(X_\alpha, \mathbb{Q})$, the rational form of example 3.1.2. Then $E_1^{p,q} = H^q(X_p, \mathbb{Q}) = \bigoplus_\alpha H^q(X_\alpha, \mathbb{Q})$ is a finite direct sum of pure Hodge structures of weight q, hence one itself: the bigrading is the direct sum of the bigradings, and the symmetry $\overline{V^{r,s}} = V^{s,r}$ holds summand by summand.

   It remains to see that each δ_i^* is a morphism of Hodge structures, since d_1 is a $\mathbb{Q}$-linear combination of these. The morphism $\delta_i\colon X_{p+1} \to X_p$ of complex algebraic varieties is holomorphic, so in local holomorphic coordinates $\delta_i^* dz^j$ is a $\mathbb{C}$-linear combination of the dw^l and $\delta_i^* d\bar{z}^j$ of the $d\bar{w}^l$; hence δ_i^* carries forms of type (r,s) to forms of type (r,s), and it commutes with d. Let $\eta \in H^{r,s}(X_p)$ with $r+s=q$. By theorem 2.1.1, η is represented by a harmonic form α of type (r,s), which is in particular closed, so $\delta_i^*\alpha$ is a closed form of type (r,s) on X_{p+1}. Its harmonic representative is again of type (r,s): on a compact Kähler manifold $\Delta_d = 2\Delta_{\bar\partial}$ (theorem 1.3.1) preserves the bidegree, so the space of harmonic forms splits by bidegree and the orthogonal projection onto it preserves bidegree. Hence $\delta_i^*\big(H^{r,s}(X_p)\big) \subseteq H^{r,s}(X_{p+1})$. The map δ_i^* on singular cohomology is defined over $\mathbb{Q}$, and its complexification is the pullback on de Rham cohomology because the de Rham comparison is natural for smooth maps; so δ_i^* satisfies the defining condition of definition 3.1.7 and is a morphism of Hodge structures of weight q. A $\mathbb{Q}$-linear combination of morphisms of Hodge structures is again one, so d_1 is a morphism of Hodge structures.

4. By part (3), $d_1\colon E_1^{p,q} \to E_1^{p+1,q}$ and $d_1\colon E_1^{p-1,q} \to E_1^{p,q}$ are morphisms of pure Hodge structures of weight q. By lemma 7.1.5(3) the kernel of the first and the image of the second are sub-Hodge structures of $E_1^{p,q}$ of weight q; both are therefore $\mathbb{Q}$-subspaces whose complexifications are bigraded and conjugation-stable. The image is contained in the kernel because $d_1 \circ d_1 = 0$, so it is a sub-Hodge structure of $\ker d_1$, and lemma 7.1.5(3) applied once more shows the quotient

$$E_2^{p,q} = \ker d_1 / \mathrm{im} d_1$$

   is a pure Hodge structure of weight q, completing the proof.

For a projective hyperresolution the first page of (8.2) is built entirely out of compact Kähler cohomology, so every entry is pure and d_1 respects the bigrading; the singular variety X enters only

through which smooth projective pieces appear and through the face maps between them. Part (4) says the purity survives one page. It does not survive further by this argument alone: showing that $d_r = 0$ for $r \geq 2$ needs the whole page to sit inside a mixed Hodge complex, which is what section 8.3 arranges using theorem 8.1.7. Part (1) is the source of the weight bound quoted there, since it confines the nonzero entries on the diagonal $p+q=k$ to $\max(0, k-n) \leq q \leq k$.

The smallest nontrivial case is the one that opened the section, and it can now be computed to the end.

Example 8.2.8: The Normalization of a Nodal Curve as a Hyperresolution

Keep the nodal cubic C, its normalization $\pi\colon \widetilde{C} = \mathbb{C}\mathrm{P}^1 \to C$ and the points p, p_1, p_2 of example 8.2.2, and let $X_\bullet$ be the augmented 1-cubical variety

$$X_\emptyset = C, \qquad X_{\{0\}} = \widetilde{C}, \qquad X_{\{1\}} = \{p\}, \qquad X_{\{0,1\}} = \{p_1, p_2\}$$

with structure morphisms π, the inclusion of p, the inclusion of $\{p_1, p_2\}$ into $\widetilde{C}$, and the constant morphism $\{p_1, p_2\} \to \{p\}$. Its associated strict simplicial variety is the one written out in example 8.2.2.

It is a hyperresolution. Here $n = \dim C = 1$. Each of $\mathbb{C}\mathrm{P}^1$, $\{p\}$ and $\{p_1, p_2\}$ is smooth, so definition 8.2.4(1) holds, and each is projective, so the hyperresolution is a projective one, as the completeness of C already guarantees through theorem 8.2.5. For (2), $\dim X_{\{0\}} = 1 \leq 1-1+1$, $\dim X_{\{1\}} = 0 \leq 1-1+1$ and $\dim X_{\{0,1\}} = 0 \leq 1-2+1 = 0$. For (3), π is a morphism of projective varieties, hence proper; $Z = \{p\}$ is closed in C; $E = \pi^{-1}(p) = \{p_1, p_2\}$ is the full preimage; and π restricts to an isomorphism $\widetilde{C} \setminus E \to C \setminus Z$, because $C \setminus \{p\}$ is the smooth locus of C and a normalization is an isomorphism over the smooth locus ([4, normalization of a variety]). So proposition 8.2.6 applies and $X_\bullet$ is of cohomological descent.

The first page. Since $\mathbb{C}\mathrm{P}^1$ is homeomorphic to the 2-sphere, $H^0(\mathbb{C}\mathrm{P}^1, \mathbb{Q}) = H^2(\mathbb{C}\mathrm{P}^1, \mathbb{Q}) = \mathbb{Q}$ and $H^1(\mathbb{C}\mathrm{P}^1, \mathbb{Q}) = 0$,[a] while a finite set of m points has $H^0 = \mathbb{Q}^m$ and $H^q = 0$ for $q > 0$. With $X_0 = \mathbb{C}\mathrm{P}^1 \sqcup \{p\}$ and $X_1 = \{p_1, p_2\}$, and with $X_p = \emptyset$ for $p \geq 2$, the only nonzero entries of (8.2) are

$$E_1^{0,0} = \mathbb{Q}^2, \qquad E_1^{0,2} = \mathbb{Q}, \qquad E_1^{1,0} = \mathbb{Q}^2$$

The only possibly nonzero d_1 is $E_1^{0,0} \to E_1^{1,0}$, equal to $\delta_0^* - \delta_1^*$. Write an element of $E_1^{0,0}$ as a pair (a, c), the constant a on $\mathbb{C}\mathrm{P}^1$ and the value c at p. Then $\delta_0^*(a, c)$ is the function equal to c at both p_1 and p_2, while $\delta_1^*(a, c)$ is the function equal to a at both, so

$$d_1(a, c) = (c - a,\ c - a)$$

This map has image the diagonal of $\mathbb{Q}^2$ and kernel $\{a = c\}$, both of dimension one.

The second page and the answer. Hence

$$E_2^{0,0} = \mathbb{Q}, \qquad E_2^{1,0} = \mathbb{Q}, \qquad E_2^{0,2} = \mathbb{Q}$$

and all other entries vanish. For $r \geq 2$ the differential d_r lands in column $p + r \geq 2$, where every entry is zero, so $E_2 = E_\infty$. Reading the diagonals,

$$H^0(C, \mathbb{Q}) = \mathbb{Q}, \qquad H^1(C, \mathbb{Q}) = \mathbb{Q}, \qquad H^2(C, \mathbb{Q}) = \mathbb{Q}$$

each with a one-step filtration.

[a][30, Homology Groups of Spheres, Universal Coefficient Theorem For Cohomology].

Two features of this computation are the ones section 8.4 takes up. First, $H^1(C,\mathbb{Q})$ is $E_\infty^{1,0}$, the cokernel of a map between H^0's: it is the space of $\mathbb{Q}$-valued functions on $\pi^{-1}(p)$ modulo those that extend compatibly to $\widetilde{C}$ and to p. The class survives precisely because π identifies two points, and it comes from cohomological degree 0 of a smooth projective variety, so by proposition 8.2.7(3) it carries a pure Hodge structure of weight 0. Second, $H^2(C,\mathbb{Q})$ is $E_\infty^{0,2}$ and is pure of weight 2, the same weight it would have on a smooth projective curve. The two degrees behave differently, and no single weight describes $H^*(C,\mathbb{Q})$: that is the mixing the next section produces in general. The comparison with the punctured case, where the weights move up rather than down, is made in section 8.4.

The normalization alone loses this rank, since $\pi^*\colon H^1(C,\mathbb{Q}) \to H^1(\widetilde{C},\mathbb{Q}) = 0$ is the zero map; the hyperresolution recovers it by keeping $\pi^{-1}(p)$ and $\{p\}$ as separate pieces of the diagram. The same construction is available for any singular variety, by theorem 8.2.5, and for a complete one it is projective, so that proposition 8.2.7 converts it into a spectral sequence of pure Hodge structures.

Exercise 8.2.1

1. Let $\rho\colon \widetilde{X} \to X$, Z and E be as in proposition 8.2.6. Show that the descent double complex of the associated simplicial variety has only the two columns $p = 0$ and $p = 1$, that $\{0\} \oplus K^{1,\bullet}$ is a subcomplex of the total complex with quotient $K^{0,\bullet}$, and deduce from the long exact sequence of that short exact sequence the Mayer-Vietoris sequence
$$\cdots \to H^k(X,\mathbb{Q}) \to H^k(\widetilde{X},\mathbb{Q}) \oplus H^k(Z,\mathbb{Q}) \to H^k(E,\mathbb{Q}) \to H^{k+1}(X,\mathbb{Q}) \to \cdots$$

2. Let $C' = \{y^2z = x^3\} \subseteq \mathbb{C}\mathrm{P}^2$, the cuspidal cubic, with its unique singular point $p' = [0:0:1]$, and let $\pi'\colon \mathbb{C}\mathrm{P}^1 \to C'$ be its normalization, for which $(\pi')^{-1}(p')$ is a single point. Write down the augmented 1-cubical variety of proposition 8.2.6 for $\rho = \pi'$ and $Z = \{p'\}$, check that it is a cubical hyperresolution, and compute $H^k(C',\mathbb{Q})$ for all k from proposition 8.2.7. Compare the answer with example 8.2.8 and say which step of the nodal computation changes.

3. Let C be an irreducible projective curve whose singularities are exactly $\delta \geq 1$ nodes, and whose normalization $\widetilde{C}$ is a smooth projective curve with $\dim_{\mathbb{Q}} H^1(\widetilde{C},\mathbb{Q}) = 2g$. Using the normalization square as in example 8.2.8, compute the second page of (8.2) and show that
$$\dim_{\mathbb{Q}} H^1(C,\mathbb{Q}) = 2g + \delta$$
Identify which of the two graded pieces of the induced filtration on $H^1(C,\mathbb{Q})$ is pure of weight 0 and which is pure of weight 1, using proposition 8.2.7(3), and give the rank of each.

4. Let X be a complex algebraic variety of dimension n and fix a cubical hyperresolution of X, which theorem 8.2.5 produces with smooth terms whether or not X is complete, together with its spectral sequence (8.2). Using parts (1) and (2) of proposition 8.2.7, neither of which asks the terms to be projective, show that $E_1^{p,q} = 0$ whenever $p + q > 2n - p$, and deduce that $H^k(X,\mathbb{Q}) = 0$ for every $k > 2n$. (The variety is not assumed smooth, compact, or irreducible.)

5. Let X be a smooth projective variety of dimension n. Show that the augmented 0-cubical variety with $X_{\{0\}} = X$ and augmentation the identity is a projective cubical hyperresolution of X, that the resulting spectral sequence is concentrated in the column $p = 0$, and that proposition 8.2.7(3) then returns exactly the pure Hodge structure of example 3.1.2.

6. Let V be a pure Hodge structure of odd weight k in the sense of definition 3.1.1. Show that $\dim_{\mathbb{Q}} V_{\mathbb{Q}}$ is even. Deduce that the group $H^1(C, \mathbb{Q})$ computed in example 8.2.8 carries no pure Hodge structure of odd weight, and in particular none of weight 1.

8.3 Deligne's Theorem for Arbitrary Varieties

Section 8.1 showed that a mixed Hodge complex determines a mixed Hodge structure on the cohomology of its rational part, and section 8.2 showed that every complex algebraic variety is resolved by a cubical object whose terms are smooth. What remains is to assemble one from the other. The assembly uses two filtrations at once: the filtration by simplicial degree, which records how many stages of the hyperresolution a class needs, and, when X is not complete, the filtration by order of logarithmic pole carried by each smooth term through its own good compactification (definition 7.3.1).

The two filtrations produce opposite weight bounds. For a smooth open variety the weights on H^k are at least k (theorem 7.3.11); for a complete variety they are at most k; and a variety that is both singular and open can carry weights on both sides of k in a single cohomology group.

The construction takes a hyperresolution and a choice of compactifications as input, so the statement is phrased for a fixed choice. That the result does not depend on the choice is proposition 8.3.4, proved once the construction is in place.

Theorem 8.3.1: Deligne's Mixed Hodge Structure on an Arbitrary Variety

Let X be a complex algebraic variety of dimension n, complete or quasi-projective (Step 1 of the proof says where that hypothesis is used and what removing it would cost), and fix a cubical hyperresolution $\varepsilon\colon X_\bullet \to X$ together with good compactifications $\overline{X_p} \supseteq X_p$ as in Step 1 of the proof. Write $D_p = \overline{X_p} \setminus X_p$ and let $D_p^{(r)}$ be the disjoint union of the r-fold intersections of the components of D_p (definition 7.3.4), with $D_p^{(0)} = \overline{X_p}$ and $D_p^{(r)} = \emptyset$ for $r < 0$. Then for every k the group $H^k(X, \mathbb{Q})$ carries a mixed Hodge structure with the following properties.

1. *The graded pieces.* The weight spectral sequence has
$$ {}_WE_1^{-m,\,k+m} \cong \bigoplus_{p\geq 0} H^{k-m-2p}\big(D_p^{(m+p)}, \mathbb{Q}\big)(-m-p) $$
a pure Hodge structure of weight $k+m$; it degenerates at E_2, and
$$ \mathrm{gr}^W_{k+m} H^k(X, \mathbb{Q}) \cong {}_WE_2^{-m,\,k+m} $$
2. *Weights in general.* $\mathrm{gr}^W_w H^k(X, \mathbb{Q}) = 0$ unless $\max(0,\, k-n) \leq w \leq 2k$.
3. *Weights on a complete variety.* If X is complete then $\mathrm{gr}^W_w H^k(X, \mathbb{Q}) = 0$ unless $\max\big(0,\, 2(k-n)\big) \leq w \leq k$: no weight exceeds the cohomological degree.
4. *Compatibility.* If X is smooth and quasi-projective, then for the hyperresolution with $X_0 = X$ and $X_p = \emptyset$ for $p \geq 1$ the structure is the one of theorem 7.3.11. If X is smooth and projective it is the pure Hodge structure of weight k of example 3.1.2.

***Proof*:**
Step 1: the hyperresolution and its compactifications. Fix a cubical hyperresolution $\varepsilon\colon X_\bullet \to X$ and write X_p for the disjoint union of its terms in level p, so that the face maps $\delta_i\colon X_p \to X_{p-1}$ for $0 \leq i \leq p$ satisfy the cosimplicial identities. Definition 8.2.4(2) gives $\dim X_p \leq n-p$, a term of X_p being an X_α with $|\alpha| = p+1$; in particular $X_p = \emptyset$ for $p > n$, so only finitely many terms are nonempty. Proposition 8.2.7(2), which asks of $X_\bullet$ only that vanishing and cohomological descent, then reads ${}_\sigma E_1^{p,q} = H^q(X_p, \mathbb{Q}) \Rightarrow H^{p+q}(X, \mathbb{Q})$.

Which terms are available depends on X, and the two cases of the hypothesis are kept apart from here on. If X is complete, the second clause of theorem 8.2.5 produces a projective hyperresolution: every X_p is smooth and projective, one takes $\overline{X_p} = X_p$ and $D_p = \emptyset$, and the rest of this step is vacuous. That clause is unavailable when X is not complete, and unavoidably so: the discussion following theorem 8.2.5 shows for $\mathbb{C}^*$ that projective terms force the augmentation to factor through a finite set, which destroys cohomological descent. So if X is quasi-projective and not complete, what is used is the first clause, which produces a hyperresolution whose terms are smooth, all that definition 8.2.4(1) requires, together with one further property of those terms: for quasi-projective X they may be taken quasi-projective as well. That property is not part of the statement of theorem 8.2.5 and is not deduced from it here; it is imported with the construction behind that theorem, in which every term is obtained from X by a resolution or

as a closed subvariety. It is what the induction below needs, because theorem 7.2.3 produces a projective compactification only of a quasi-projective variety.

Suppose X is not complete, so that its terms X_p are smooth quasi-projective. Choose the $\overline{X_p}$ by induction on p. Having chosen $\overline{X_0}, \dots, \overline{X_{p-1}}$, let $\overline{X_p}^0$ be any smooth projective compactification of X_p (theorem 7.2.3) and let Γ be the closure of the graph of

$$(\delta_0, \dots, \delta_p) \colon X_p \to \left(\overline{X_{p-1}}\right)^{p+1}$$

inside $\overline{X_p}^0 \times \left(\overline{X_{p-1}}\right)^{p+1}$, a projective variety because the ambient product is one. Applying [24] to Γ and its closed subset $\Gamma \setminus X_p$ gives a smooth $\overline{X_p}$ with a projective morphism $\overline{X_p} \to \Gamma$ that is an isomorphism over X_p and for which $D_p = \overline{X_p} \setminus X_p$ is a normal crossings divisor; being projective over the projective Γ, the variety $\overline{X_p}$ is projective. Composing $\overline{X_p} \to \Gamma$ with the projections extends each δ_i to $\overline{\delta_i} \colon \overline{X_p} \to \overline{X_{p-1}}$, and $\overline{\delta_i}^{-1}(D_{p-1}) \subseteq D_p$ because δ_i carries X_p into X_{p-1}. The cosimplicial identities among the $\overline{\delta_i}$ hold on the dense open subset X_p, hence hold everywhere: two morphisms from a reduced variety to a separated variety agreeing on a dense subset are equal. This is the argument used for a single morphism in the proof of proposition 7.3.14, run once for each level. The hypothesis on X is used in exactly two places, the choice of hyperresolution above and this induction, and that is why the statement carries it. For a variety that is neither complete nor quasi-projective, theorem 8.2.5 still produces smooth terms, but a term that is not quasi-projective has no projective compactification at all, a variety open in a projective one being quasi-projective, so theorem 7.2.3 returns only a proper smooth compactification; its boundary strata need not be Kähler, and Step 4 below asks each stratum for a degeneration this book has only in the compact Kähler case. Deligne reaches those varieties in [19, Théorie de Hodge III] along a different route, compactifying X first, which an arbitrary separated scheme of finite type admits, and hyperresolving the pair; that route is not reproduced here.

Step 2: the bifiltered complex. For each p the pair $(\overline{X_p}, D_p)$ carries the logarithmic complex $\Omega^\bullet_{\overline{X_p}}(\log D_p)$ with its weight filtration (definition 7.3.1), written $W^{\log}$ here to distinguish it from the filtration being built, and its Hodge filtration (definition 7.3.2). Let $C_p^\bullet$ be a complex of $\mathbb{C}$-vector spaces computing the hypercohomology of that complex, carrying the filtrations induced by $W^{\log}$ and F and depending functorially on the pair $(\overline{X_p}, D_p)$; the global sections of a Godement resolution are one such choice, and the filtered framework in which such a choice is made is [7, The Filtered Derived Category]. Let $A_p^\bullet$ be a complex of $\mathbb{Q}$-vector spaces computing $H^\bullet(X_p, \mathbb{Q})$, again functorial in X_p. Set

$$K^k_{\mathbb{Q}} = \bigoplus_{p \geq 0} A_p^{k-p}, \qquad K^k_{\mathbb{C}} = \bigoplus_{p \geq 0} C_p^{k-p}$$

with differential $d(x) = d_p x + (-1)^q \delta x$ on x of internal degree q in level p, where $\delta = \sum_{i=0}^{p+1} (-1)^i \delta_i^*$ is built from the face maps $\delta_i \colon X_{p+1} \to X_p$, and δ_i^* denotes the map induced by δ_i on $A^\bullet$ and by its extension $\overline{\delta_i}$ on $C^\bullet$. Then $d^2 = 0$: expanding, $d^2 x = (-1)^{q+1} \delta d_p x + (-1)^q d_p \delta x + \delta^2 x$, the first two terms cancel because δ commutes with the internal differentials, and $\delta^2 = 0$ by the cosimplicial identities. Filter by

$$F^r K^k_{\mathbb{C}} = \bigoplus_{p \geq 0} F^r C_p^{k-p}, \qquad W_m K^k_{\mathbb{C}} = \bigoplus_{p \geq 0} W^{\log}_{m+p} C_p^{k-p}$$

Both are subcomplexes. For F: the internal differentials preserve F, and $\overline{\delta_i}^*$ preserves the degree of a holomorphic form, hence preserves F. For W: the internal differential preserves each $W^{\log}_{m+p}$ (definition 7.3.1), while δ_i^* carries $W^{\log}_{m+p}C_p$ into $W^{\log}_{m+p}C_{p+1}$: a pullback along a morphism with $\overline{\delta_i}^{-1}(D_p) \subseteq D_{p+1}$ sends $d\log f$ to $d\log(f\circ\overline{\delta_i})$, and $f\circ\overline{\delta_i}$ cuts out a divisor inside D_{p+1}, so a form with at most $m+p$ logarithmic factors goes to a form with at most $m+p$ logarithmic factors; this is the local computation in the proof of proposition 7.3.14. Finally $W^{\log}_{m+p}C_{p+1} \subseteq W^{\log}_{m+(p+1)}C_{p+1}$, which is the level-$(p+1)$ summand of W_m. The same computation shows more, and it is used in Step 4: on the level-$(p+1)$ summand, δ lands one step below the graded level, so δ maps W_m into W_{m-1}.

The filtration W on the rational complex $K_{\mathbb{Q}}$ is the same combination of two rational filtrations,

$$W_m K^k_{\mathbb{Q}} = \bigoplus_{p\geq 0} W^{\log}_{m+p} A^{k-p}_p$$

where the shift by the simplicial degree is defined on $K_{\mathbb{Q}}$ by construction, and $W^{\log}$ on $A^\bullet_p$ is the filtration transported from the logarithmic complex across the comparison of Step 3, a filtration defined over $\mathbb{Q}$ by theorem 7.3.9, imported in section 7.3.

Step 3: the comparison. By theorem 7.2.17 the inclusion of the logarithmic complex into the smooth forms on X_p identifies $H^j(X_p,\mathbb{C})$ with the hypercohomology of $\Omega^\bullet_{\overline{X_p}}(\log D_p)$, naturally in the pair $(\overline{X_p}, D_p)$. Realized on complexes, it is a chain

$$A^\bullet_p \otimes_{\mathbb{Q}} \mathbb{C} \to E^\bullet_p \leftarrow C^\bullet_p$$

of quasi-isomorphisms through a complex $E^\bullet_p$ of smooth forms on X_p, and not a single morphism from $A^\bullet_p \otimes_{\mathbb{Q}} \mathbb{C}$ to $C^\bullet_p$: the logarithmic complex maps to the smooth de Rham complex and not the other way. What definition 8.1.4(3) asks for is a morphism $\alpha\colon K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} \to K_{\mathbb{C}}$ with $\alpha\big(W_m K_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}\big) \subseteq W_m K_{\mathbb{C}}$, so the axiom has to be read with α a morphism in the filtered derived category, that is, a chain of W-compatible quasi-isomorphisms with the wrong-way ones inverted. That reading is forced rather than chosen: no choice of the complexes above makes the comparison a single morphism, and example 8.1.5 verifies the axiom for a single pair $(\overline{X}, D)$ by this same logarithmic comparison. Compatibility with W holds map by map. Each map preserves the simplicial degree, so it respects the shift by p; and on the logarithmic part it respects $W^{\log}$ because W on $K_{\mathbb{Q}}$ was defined in Step 2 by transporting $W^{\log}$ across this comparison, which theorem 7.3.9 permits. Naturality makes each map in the chain commute with the face maps, so the chain totalizes to one between $K_{\mathbb{Q}} \otimes \mathbb{C}$ and $K_{\mathbb{C}}$. Each map in the totalized chain is a quasi-isomorphism on every level, hence induces an isomorphism on the E_1-page of the filtration by level. Both filtrations have only finitely many nonzero levels, by the vanishing $X_p = \emptyset$ for $p > n$ of Step 1, so the associated spectral sequences converge and an isomorphism on E_1 forces one on the abutments. Hence $H^k(K_{\mathbb{Q}}) \otimes \mathbb{C} \cong H^k(K_{\mathbb{C}})$ for every k.

Step 4: the purity axiom. Since δ maps W_m into W_{m-1} (Step 2), the differential induced on $\mathrm{gr}^W_m K_{\mathbb{C}}$ is the internal one alone, so

$$\mathrm{gr}^W_m K_{\mathbb{C}} = \bigoplus_{p\geq 0} \mathrm{gr}^{W^{\log}}_{m+p} C^\bullet_p\,[-p]$$

as complexes, the shift recording that the level-p summand of $K_{\mathbb{C}}^k$ is C_p^{k-p}. Taking cohomology in degree k and applying proposition 7.3.6 together with Step 1 of the proof of proposition 7.3.7, which computes the hypercohomology of a graded piece of the logarithmic complex as the cohomology of a stratum,

$$H^k\big(\mathrm{gr}_m^W K_{\mathbb{C}}\big) = \bigoplus_{p\geq 0} H^{(k-p)-(m+p)}\big(D_p^{(m+p)},\mathbb{C}\big) = \bigoplus_{p\geq 0} H^{k-m-2p}\big(D_p^{(m+p)},\mathbb{C}\big)$$

By parts (1) and (2) of proposition 7.3.7 the filtration induced by F on the summand indexed by p is the Hodge filtration of the smooth projective variety $D_p^{(m+p)}$ shifted by $m+p$, and the rational structure inherited from $K_{\mathbb{Q}}$ is $H^{k-m-2p}(D_p^{(m+p)},\mathbb{Q})(-m-p)$. That is a pure Hodge structure of weight

$$(k-m-2p)+2(m+p) = k+m$$

independent of p. So $H^k(\mathrm{gr}_m^W K_{\mathbb{C}})$ is pure of weight $k+m$ for every k, which is condition (d) of definition 8.1.2 for $\mathrm{gr}_m^W K$ as a pure Hodge complex of weight m, the cohomology of such a complex in degree k being pure of weight $k+m$. This is the same indexing under which proposition 7.3.7 makes the logarithmic complex of a smooth variety a mixed Hodge complex.

The other three conditions of definition 8.1.2 hold summand by summand. Condition (a), finite dimensionality: the summand indexed by p has cohomology $H^{k-m-2p}(D_p^{(m+p)},\mathbb{C})$, finite dimensional because $D_p^{(m+p)}$ is a compact complex manifold, and only the finitely many $p\leq n$ of Step 1 contribute. Condition (b), the comparison on the graded piece: the comparison of Step 3 carries W_m into W_m, so it induces one on gr_m^W, and on the summand indexed by p that induced comparison is the identification of $H^{k-m-2p}(D_p^{(m+p)},\mathbb{Q})(-m-p)\otimes_{\mathbb{Q}}\mathbb{C}$ with $H^{k-m-2p}(D_p^{(m+p)},\mathbb{C})$ given by proposition 7.3.6, which is an isomorphism. Condition (c), strict compatibility of the differential of $\mathrm{gr}_m^W K_{\mathbb{C}}$ with F: by proposition 7.3.6 the summand indexed by p is, up to a shift and a Tate twist, the de Rham complex of the smooth projective variety $D_p^{(m+p)}$ with its Hodge filtration, and neither a shift nor a twist changes the differential or the strictness of a filtration; by [7, degeneration at E_1 and strictness] strictness for F is the degeneration at E_1 of the Frölicher spectral sequence of $D_p^{(m+p)}$, which holds because that variety is smooth and projective. This is the verification of example 8.1.5 run on each level, and it is the condition theorem 8.1.7 requires as the base case of its induction. Together with Step 3 this makes $\big(K_{\mathbb{Q}},(K_{\mathbb{C}},W,F)\big)$ a mixed Hodge complex (definition 8.1.4).

Step 5: the structure and its graded pieces. By theorem 8.1.7 the group $H^k(K_{\mathbb{Q}})$ carries a mixed Hodge structure, the weight spectral sequence degenerates at E_2, and $\mathrm{gr}_{k+m}^W H^k(K_{\mathbb{Q}}) \cong {}_WE_2^{-m,k+m}$ with ${}_WE_1^{-m,k+m} = H^k(\mathrm{gr}_m^W K_{\mathbb{C}})$. Step 4 computes that E_1-term and gives part (1) of the statement, once $H^k(K_{\mathbb{Q}})$ is identified with $H^k(X,\mathbb{Q})$. That identification is cohomological descent: filtering $K_{\mathbb{Q}}$ by level produces the descent spectral sequence of proposition 8.2.7, whose abutment that proposition identifies with $H^k(X,\mathbb{Q})$.

Step 6: the weight bounds. A summand of ${}_WE_1^{-m,k+m}$ is nonzero only if three conditions hold: $p\leq n$, since X_p and hence $D_p^{(m+p)}$ is empty beyond that; $m+p\geq 0$, since $D_p^{(r)}=\emptyset$ for $r<0$; and $k-m-2p\geq 0$, since a variety has no cohomology in negative degree. Write $w=k+m$ for the weight of the term. From $m\geq -p\geq -n$ comes $w\geq k-n$. From $w=(k-m-2p)+2(m+p)$,

a sum of two nonnegative integers, comes $w \geq 0$. From $m \leq k-2p \leq k$ comes $w \leq 2k$. Since $\mathrm{gr}_w^W H^k(X,\mathbb{Q})$ is a subquotient of ${}_WE_1^{-m,k+m}$ with $w = k+m$, this is part (2).

If X is complete then $D_p = \emptyset$ for every p, so $D_p^{(r)}$ is empty unless $r = 0$, and the only surviving summand is the one with $p = -m$. In particular $m \leq 0$ and $w = k+m \leq k$. That summand is $H^{k-m-2p}(X_p,\mathbb{Q}) = H^{k-p}(X_p,\mathbb{Q})$ with $p = -m$, which vanishes unless $0 \leq k-p \leq 2\dim X_p \leq 2(n-p)$. The left inequality gives $w = k-p \geq 0$, and the right gives $p \leq 2n-k$, hence $w = k-p \geq 2k-2n$. This is part (3).

Step 7: the two compatibilities. Let X be smooth and quasi-projective and take $X_0 = X$ with $X_p = \emptyset$ for $p \geq 1$; the augmentation is then an isomorphism, so cohomological descent holds and this is a cubical hyperresolution (definition 8.2.4), its one term being smooth and quasi-projective as the second case of Step 1 asks. Quasi-projectivity is what the comparison needs and smoothness alone will not do. A smooth complete X that is not projective, such as the proper non-projective threefold named in the footnote to theorem 7.2.3, has no smooth projective compactification, so theorem 7.3.11 attaches no structure to it and there is nothing to compare; and the trivial cube on such an X is not a projective hyperresolution, so Step 1's complete case excludes it, while taking it anyway would leave $\overline{X_0} = X_0 = X$, whose Frölicher spectral sequence Step 4 needs to degenerate at E_1, a degeneration this book has only for a compact Kähler manifold.

Return to the quasi-projective case. Only the level $p = 0$ survives, $W_mK_{\mathbb{C}} = W_m^{\log}C_0^\bullet$ and $F^rK_{\mathbb{C}} = F^rC_0^\bullet$, so K is the logarithmic complex of $(\overline{X_0}, D_0)$ with its two filtrations and nothing else. The weight and Hodge filtrations that theorem 8.1.7 induces on H^k are then the images of $H^k(\overline{X_0}, W_m\Omega^\bullet(\log D_0))$ and of $H^k(\overline{X_0}, F^p\Omega^\bullet(\log D_0))$, which is the definition of the filtrations in theorem 7.3.11. If moreover X is projective, take $\overline{X_0} = X$ and $D_0 = \emptyset$; then $W_{-1}K_{\mathbb{C}} = 0$ and $W_0K_{\mathbb{C}} = K_{\mathbb{C}}$, so $\mathrm{gr}_m^W H^k$ vanishes for $m \neq 0$ and $H^k(X,\mathbb{Q})$ is pure of weight k, with F the filtration induced by the degree of a holomorphic form. On a compact Kähler manifold that filtration induces the Hodge filtration attached to the bigrading (chapter 2), the same identification definition 7.3.2 rests on: a class has a harmonic representative (theorem 2.1.1), harmonic projection commutes with the decomposition into types because $\Delta_d = 2\Delta_{\bar\partial}$ preserves them (theorem 1.3.1), and a form of holomorphic degree at least r has no component of type (a,b) with $a < r$, so neither has its harmonic representative. Hence the structure is that of example 3.1.2, completing the proof.

The theorem drops both standing hypotheses of the earlier chapters. Chapter 2 needed a compact Kähler manifold, chapter 7 needed smoothness, and here neither is assumed: every complex algebraic variety that is complete or quasi-projective has functorial Hodge-theoretic invariants attached to its cohomology, with the singularities and the noncompactness recorded in the weight filtration rather than obstructing the construction, and Step 1 of the proof records what carrying it to the remaining varieties would take. Part (1) says which two indices control the weight. The index p is the stage of the hyperresolution, and it lowers the weight by p; the index $r = m+p$ is the number of boundary components a stratum lies on, and it raises the weight by r. The weight of the term is $k-p+r$.

That formula also states the contrast with chapter 7. For a smooth quasi-projective open U only $p = 0$ contributes, so the weight is $k+r$ with $r \geq 0$ and the weights on $H^k(U,\mathbb{Q})$ lie in $[k,2k]$, which is theorem 7.3.11. For a complete singular X only $r = 0$ contributes, so the weight is $k-p$ with $p \geq 0$ and the weights lie at or below k. A variety that is both open and singular leaves both indices

free, and a single group H^k can then carry weights on both sides of k. The example below exhibits all three cases on one configuration of lines.

Example 8.3.2: A Triangle of Lines, Complete and Punctured

1. *The complete case.* Let $X = L_1 \cup L_2 \cup L_3 \subseteq \mathbb{CP}^2$ be three distinct lines with no common point, so that X is a complete curve ($n = 1$) with three nodes v_{12}, v_{13}, v_{23}. Its normalization is $\pi\colon \widetilde{X} = L_1 \sqcup L_2 \sqcup L_3 \to X$, and the hyperresolution of example 8.2.8 has

$$X_0 = \widetilde{X} \sqcup S, \qquad X_1 = \pi^{-1}(S), \qquad X_p = \emptyset \ (p \geq 2)$$

where $S = \{v_{12}, v_{13}, v_{23}\}$ and $\pi^{-1}(S)$ consists of six points, two over each node. Since X is complete, part (1) of theorem 8.3.1 reduces to ${}_WE_1^{-m,k+m} = H^{k+m}(X_{-m}, \mathbb{Q})$, the descent spectral sequence. Index it by $s = -m$ and $t = k + m$, so that $s + t = k$, the entry ${}_WE_1^{s,t}$ is $H^t(X_s, \mathbb{Q})$, and t is the weight it contributes. Its rows are short. In row $t = 0$ the map $H^0(X_0, \mathbb{Q}) = \mathbb{Q}^6 \to H^0(X_1, \mathbb{Q}) = \mathbb{Q}^6$ sends $(a_1, a_2, a_3; b_{12}, b_{13}, b_{23})$ to the six differences $a_i - b_{ij}$ indexed by the incidences. It is the coboundary of the incidence graph, which has a vertex for each line and each node and an edge for each of the six incidences, hence six vertices and six edges. That graph is connected, so the kernel is the line of constants, and the cokernel, which is the first cohomology of the graph, is one dimensional as well. In row $t = 1$, $H^1(X_0, \mathbb{Q}) = 0$, since X_0 is a disjoint union of projective lines and points. In row $t = 2$, $H^2(X_0, \mathbb{Q}) = \mathbb{Q}(-1)^3$ and $H^2(X_1, \mathbb{Q}) = 0$. All terms with $s \geq 2$ vanish because $X_p = \emptyset$ for $p \geq 2$. Hence

$$H^0(X, \mathbb{Q}) = \mathbb{Q}, \qquad H^1(X, \mathbb{Q}) = \mathbb{Q}, \qquad H^2(X, \mathbb{Q}) = \mathbb{Q}(-1)^3$$

with $H^1(X, \mathbb{Q})$ pure of weight 0 and $H^2(X, \mathbb{Q})$ pure of weight 2. The Euler characteristic checks: $1 - 1 + 3 = 3 = 3 \cdot 2 - 3$, three lines of Euler characteristic 2 with three points each counted twice. The class in H^1 has weight 0, one below its degree, and it is precisely the first cohomology of the incidence graph: it exists because the three nodes close a cycle. Two lines instead of three give a tree, and then H^1 vanishes; exercise 8.3.1 (5) carries the count out for d lines.

2. *The punctured case.* Let a, b be two distinct points of L_1 other than v_{12}, v_{13} and put $X' = X \setminus \{a, b\}$, an open singular curve. The hyperresolution is the same one with L_1 replaced by $L_1 \setminus \{a, b\}$, so $\overline{X'_0} = \widetilde{X} \sqcup S$ with $D_0 = \{a, b\}$, while $X'_1 = \pi^{-1}(S)$ is already complete and $D_1 = \emptyset$. The strata are $D_0^{(0)} = \overline{X'_0}$, $D_0^{(1)} = \{a, b\}$ and $D_1^{(0)} = \overline{X'_1} = \pi^{-1}(S)$, all others empty: for $r \geq 2$ the stratum $D_0^{(r)}$ is empty because $\{a\}$ and $\{b\}$ are disjoint components of D_0, and for $r \geq 1$ the stratum $D_1^{(r)}$ is empty because D_1 itself is empty. For $k = 1$ the constraints of Step 6 of the proof read $m + p \geq 0$ and $1 - m - 2p \geq 0$ with $0 \leq p \leq 1$, so $m \in \{-1, 0, 1\}$ and part (1) of theorem 8.3.1 gives three possible weights $1 + m$:

$$\begin{aligned} {}_WE_1^{1,0} &= H^0\big(D_1^{(0)}, \mathbb{Q}\big) = \mathbb{Q}^6 \\ {}_WE_1^{0,1} &= H^1\big(D_0^{(0)}, \mathbb{Q}\big) = 0 \\ {}_WE_1^{-1,2} &= H^0\big(D_0^{(1)}, \mathbb{Q}\big)(-1) = \mathbb{Q}(-1)^2 \end{aligned}$$

The differential into the first of these comes from ${}_WE_1^{0,0} = H^0(D_0^{(0)}, \mathbb{Q}) = \mathbb{Q}^6$ and is the map of item (1), of rank five, so ${}_WE_2^{1,0} = \mathbb{Q}$, of weight 0. The differential out of

the third lands in ${}_WE_1^{0,2} = H^2(D_0^{(0)}, \mathbb{Q}) = \mathbb{Q}(-1)^3$ and is, by proposition 7.3.7(3), the sum of the Gysin maps of the two inclusions $\{a\} \hookrightarrow L_1$ and $\{b\} \hookrightarrow L_1$; each sends the generator to the class of a point in $H^2(L_1, \mathbb{Q}) = \mathbb{Q}(-1)$, so the map has rank one and kernel $\mathbb{Q}(-1)$ spanned by the difference. Nothing maps into ${}_WE_1^{-1,2}$: the differential into it starts at ${}_WE_1^{-2,2}$, the term with $m = 2$ in total degree $k = 0$, whose summands are $H^{k-m-2p}\big(D_p^{(m+p)}, \mathbb{Q}\big)(-2-p) = H^{-2-2p}\big(D_p^{(2+p)}, \mathbb{Q}\big)(-2-p)$ for $p \geq 0$, all zero because the degree $-2-2p$ is negative, which is the third constraint of Step 6 of the proof. Hence

$$\mathrm{gr}_0^W H^1(X', \mathbb{Q}) = \mathbb{Q}, \qquad \mathrm{gr}_1^W H^1(X', \mathbb{Q}) = 0, \qquad \mathrm{gr}_2^W H^1(X', \mathbb{Q}) = \mathbb{Q}(-1)$$

so $H^1(X', \mathbb{Q})$ has dimension two and carries weights on both sides of $k = 1$: weight 0 from the loop in the incidence graph, weight 2 from the two punctures on L_1.

The two computations run on the same hyperresolution; the second adds the boundary index r to the stage index p, and each index contributes one class to H^1. A worked example on an irreducible curve with a single node, where the weight-zero class is matched against the vanishing cycle of section 4.4, is section 8.4.

The construction used a hyperresolution and a system of compactifications, and nothing so far rules out a different choice producing different filtrations. The next result rules it out, and the same comparison argument shows that pullback along a morphism of varieties respects the structures, since such a morphism is covered by a morphism of hyperresolutions. Both facts about hyperresolutions that this needs are imported, and are recorded here so that the debt is visible.

Theorem 8.3.3: Domination and Covering of Hyperresolutions

Imported, like theorem 8.2.5.

1. If $X_\bullet \to X$ and $X'_\bullet \to X$ are cubical hyperresolutions of a complex algebraic variety X, then they are dominated by a third: there is a cubical hyperresolution $X''_\bullet \to X$ with morphisms $X''_\bullet \to X_\bullet$ and $X''_\bullet \to X'_\bullet$ over id_X.
2. Every morphism $f\colon X \to Y$ of complex algebraic varieties is covered by a morphism of hyperresolutions: there are cubical hyperresolutions $X_\bullet \to X$ and $Y_\bullet \to Y$ and a morphism $X_\bullet \to Y_\bullet$ lying over f.

Both belong to the theory of hyperresolutions cited for theorem 8.2.5 and are proved in [33, Chapter 5].

Proposition 8.3.4: Independence and Functoriality

1. The mixed Hodge structure of theorem 8.3.1 on $H^k(X, \mathbb{Q})$ is independent of the cubical hyperresolution and of the good compactifications used to construct it.
2. If $f\colon X \to Y$ is a morphism of complex algebraic varieties, each of them as in theorem 8.3.1, then $f^*\colon H^k(Y, \mathbb{Q}) \to H^k(X, \mathbb{Q})$ is a morphism of mixed Hodge structures. So $X \mapsto H^k(X, \mathbb{Q})$ is a contravariant functor from those varieties to mixed Hodge structures, extending the functor of proposition 7.3.14.

***Proof*:**
Step 1: a bijective morphism of mixed Hodge structures is an isomorphism. Let $\varphi\colon V \to V'$ be a morphism of mixed Hodge structures whose underlying $\mathbb{Q}$-linear map is bijective. Strictness for W (theorem 7.1.8(1)) gives $\varphi(W_mV) = \varphi(V_{\mathbb{Q}}) \cap W_mV' = W_mV'$ for every m, using surjectivity of φ in the second equality, and strictness for F (theorem 7.1.8(2)) gives $\varphi_{\mathbb{C}}(F^pV_{\mathbb{C}}) = \varphi_{\mathbb{C}}(V_{\mathbb{C}}) \cap F^pV'_{\mathbb{C}} = F^pV'_{\mathbb{C}}$ for every p. So φ carries each filtration onto the corresponding one, and φ^{-1} is a morphism as well.

Step 2: independence. The two facts about hyperresolutions used here are the two parts of theorem 8.3.3, imported. Given a morphism $X'_\bullet \to X_\bullet$ over id_X, run the graph-closure induction of Step 1 of the proof of theorem 8.3.1 on $X'_\bullet$ with the compactifications of $X_\bullet$ already fixed; the resulting $\overline{X'_p}$ then receive extensions of the given maps $X'_p \to X_p$ as well as of the face maps. Pullback of forms along these extensions preserves the order of logarithmic pole and the degree of a holomorphic form, exactly as in Step 2 there, so it defines a map of complexes $K \to K'$ compatible with W and with F, and compatible with the rational structures. By theorem 8.1.7 the induced map $H^k(K_{\mathbb{Q}}) \to H^k(K'_{\mathbb{Q}})$ is a morphism of mixed Hodge structures.

Both sides were identified with $H^k(X, \mathbb{Q})$ in Step 5 of that proof, by cohomological descent along the augmentations ε and ε'. The map $X'_\bullet \to X_\bullet$ commutes with the augmentations by hypothesis, so the two identifications agree and the morphism above is the identity map of $H^k(X, \mathbb{Q})$. By Step 1 it is an isomorphism of mixed Hodge structures, so the filtrations produced by $X_\bullet$ and by $X'_\bullet$ coincide. Applying this to a common domination, given by theorem 8.3.3(1), settles any two hyperresolutions. For two systems of compactifications of one $X_\bullet$ the same argument applies with $X'_\bullet = X_\bullet$ and the identity map, after dominating the two systems by a third, again by graph closure and [24].

Step 3: functoriality. Let $f\colon X \to Y$ be a morphism and choose, by theorem 8.3.3(2), hyperresolutions with a morphism $X_\bullet \to Y_\bullet$ lying over f. Choose compactifications of $Y_\bullet$ first and then of $X_\bullet$ by the graph-closure induction, so that the maps $X_p \to Y_p$ extend to the compactifications and carry the boundary into the boundary. As in Step 2 this produces a map of mixed Hodge complexes $K_Y \to K_X$, hence a morphism of mixed Hodge structures $H^k(K_{Y,\mathbb{Q}}) \to H^k(K_{X,\mathbb{Q}})$, which under the descent identifications is f^*. By Step 2 the two structures do not depend on the choices, so f^* is a morphism of mixed Hodge structures for the structures of theorem 8.3.1. Composition is inherited from rational cohomology, where $(g \circ f)^* = f^* \circ g^*$ holds already, so $H^k(-, \mathbb{Q})$ is a contravariant functor to mixed Hodge structures; it extends proposition 7.3.14 because on smooth quasi-projective varieties, which are the ones that proposition treats, the two structures agree by theorem 8.3.1(4), completing the proof.

Functoriality is what turns the weight bounds into constraints on maps. A morphism of mixed Hodge structures is strict for both filtrations, and the functor gr^W_m is exact (theorem 7.1.8), so a surjection on rational cohomology induces a surjection on each weight-graded piece, and a class of weight w lies in the image of f^* only if the source has a nonzero weight-w part. For the curve X of example 8.3.2, $H^1(X, \mathbb{Q})$ is pure of weight 0, so for every morphism $f\colon X \to Y$ with Y smooth and complete the map f^* vanishes on H^1: there $H^1(Y, \mathbb{Q})$ is pure of weight 1, so $f^*\big(H^1(Y, \mathbb{Q})\big) = f^*\big(W_0H^1(Y, \mathbb{Q})\big) = f^*(0)$ by strictness for W, the first equality holding because $f^*(H^1(Y, \mathbb{Q}))$ lies in $W_0H^1(X, \mathbb{Q}) = H^1(X, \mathbb{Q})$.

Exercise 8.3.1

1. Let X be a complex algebraic variety of dimension n, as in theorem 8.3.1. Show that $H^0(X,\mathbb{Q})$ is pure of weight 0, and that if X is complete then $H^{2n}(X,\mathbb{Q})$ is pure of weight $2n$. In each case name which part of theorem 8.3.1 you use.

2. Let X be a complete variety of dimension n. Using the description of ${}_WE_1$ in theorem 8.3.1(1) together with the bound $\dim X_p \leq n-p$ from Step 1 of its proof, show that $H^k(X,\mathbb{Q})=0$ for $k>2n$. Deduce that the interval in theorem 8.3.1(3) is nonempty for every k with $H^k(X,\mathbb{Q})\neq 0$.

3. Let X be complete with hyperresolution $X_\bullet$ and write $\varepsilon_0\colon X_0\to X$ for the augmentation of the level-zero term. Show that
$$\operatorname{gr}_k^W H^k(X,\mathbb{Q}) = \ker\bigl(d_1\colon H^k(X_0,\mathbb{Q})\to H^k(X_1,\mathbb{Q})\bigr)$$
where d_1 is the alternating sum of the pullbacks along the face maps, and that $W_{k-1}H^k(X,\mathbb{Q}) = \ker\bigl(\varepsilon_0^*\bigr)$. Conclude that $\operatorname{gr}_k^W H^k(X,\mathbb{Q})$ is a sub-Hodge structure of $H^k(X_0,\mathbb{Q})$, so its Hodge numbers are bounded by those of X_0.

4. Let X be a complete curve with normalization $\pi\colon\widetilde{X}\to X$, singular locus S and $\widetilde{S}=\pi^{-1}(S)$, and use the hyperresolution $X_0=\widetilde{X}\sqcup S$, $X_1=\widetilde{S}$ of example 8.2.8. Show that
$$\operatorname{gr}_1^W H^1(X,\mathbb{Q})\cong H^1(\widetilde{X},\mathbb{Q}),\qquad \operatorname{gr}_0^W H^1(X,\mathbb{Q})\cong \operatorname{coker}\bigl(H^0(X_0,\mathbb{Q})\to H^0(X_1,\mathbb{Q})\bigr)$$
and that $H^2(X,\mathbb{Q})$ is pure of weight 2 of rank the number of irreducible components of X.

5. Let $d\geq 2$ and let $X\subseteq\mathbb{C}\mathrm{P}^2$ be a union of d distinct lines, no three of them concurrent. Show that $H^1(X,\mathbb{Q})$ is pure of weight 0 of dimension $(d-1)(d-2)/2$ and that $H^2(X,\mathbb{Q})\cong\mathbb{Q}(-1)^d$. Check the answer against the Euler characteristic of X.

6. Let $f\colon X\to Y$ be a morphism of complex algebraic varieties, each as in theorem 8.3.1, such that $f^*\colon H^k(Y,\mathbb{Q})\to H^k(X,\mathbb{Q})$ is surjective. Show that $\operatorname{gr}_m^W f^*$ is surjective for every m, and deduce that if $\operatorname{gr}_w^W H^k(Y,\mathbb{Q})=0$ for all $w>w_0$ then the same holds for $H^k(X,\mathbb{Q})$.

7. Let $C\subseteq\mathbb{C}\mathrm{P}^2$ be a smooth conic and L a line meeting it in two distinct points, and let $X=(C\cup L)\setminus\{a,b\}$ where a,b are two distinct points of C not lying on L. Compute $\operatorname{gr}_w^W H^1(X,\mathbb{Q})$ for every w, and verify that the total dimension agrees with the Euler characteristic of X together with the values of H^0 and H^2.

8. Show that both ends of the weight interval in theorem 8.3.1(2) are attained when $k=n=1$: exhibit a curve whose H^1 has a class of weight $2k=2$, and one whose H^1 has a class of weight $k-n=0$. Then show that no smooth projective variety realizes either weight in its H^1.

8.4 The Nodal Cubic

Theorem 8.3.1 attaches a mixed Hodge structure to the rational cohomology of every complex algebraic variety. This section computes it, in every degree and by hand, for the smallest singular variety available: an irreducible plane cubic with one node. The answer in degree one is one-dimensional and pure of weight zero, so a weight occurs one step below the cohomological degree, which cannot happen on a smooth projective variety, where theorem 2.1.1 makes H^1 pure of weight one.

The curve is already in this book twice. Example 4.4.8 met it as a singular member of a general pencil of plane cubics and studied the monodromy of the nearby smooth fibre around the corresponding

puncture; example 7.2.4 met it as the boundary of a good compactification, and blew up its node to make that boundary a simple normal crossings divisor. Here the curve itself is the object, and the resolution used is the one of example 8.2.8: its normalization.

Example 8.4.1: The Nodal Plane Cubic and Its Normalization
Let

$$C = \{y^2z = x^2(x+z)\} \subseteq \mathbb{C}\mathrm{P}^2, \qquad p = [0:0:1]$$

be the curve of example 7.2.4, and let

$$\pi\colon \mathbb{C}\mathrm{P}^1 \to \mathbb{C}\mathrm{P}^2, \qquad \pi([s:t]) = \left[\, s(t^2-s^2) \;:\; t(t^2-s^2) \;:\; s^3 \,\right]$$

The three cubic forms have no common zero, since $s^3 = 0$ forces $s = 0$ and then $t(t^2 - s^2) = t^3$ vanishes only at $t = 0$, so π is a morphism. Its image lies in C, because

$$\big(t(t^2-s^2)\big)^2 s^3 = s^3t^2(t^2-s^2)^2 = \big(s(t^2-s^2)\big)^2\big(s(t^2-s^2)+s^3\big)$$

the last expression being $s^2(t^2-s^2)^2 \cdot st^2$.

Surjectivity. Let $[x : y : z] \in C$. If $z = 0$ the equation gives $x^3 = 0$, so the point is $[0 : 1 : 0] = \pi([0 : 1])$. If $z \neq 0$, scale to $z = 1$, so that $y^2 = x^2(x + 1)$. When $x = 0$ this forces $y = 0$ and the point is $p = \pi([1 : 1])$. When $x \neq 0$, put $t = y/x$; then $t^2 = x+1$, so $x = t^2-1$ and $y = t(t^2 - 1)$, and the point is $\pi([1 : t])$.

The fibres. A point of $\pi^{-1}(p)$ has $s^3 \neq 0$, hence $s \neq 0$, and $s(t^2-s^2) = 0$, hence $t^2 = s^2$; conversely both $[1 : 1]$ and $[1 : -1]$ map to p. So $\pi^{-1}(p) = \Sigma := \{a, b\}$ with $a = [1 : 1]$ and $b = [1 : -1]$. On $C \setminus \{p\}$ the assignment $[x : y : z] \mapsto [x : y]$ is a morphism, since $x = y = 0$ together with $y^2z = x^2(x + z)$ forces the point to be p; and it inverts π, because $[s(t^2 - s^2) : t(t^2 - s^2)] = [s : t]$ whenever $t^2 \neq s^2$. Hence π restricts to an isomorphism

$$\mathbb{C}\mathrm{P}^1 \setminus \Sigma \xrightarrow{\;\sim\;} C \setminus \{p\}$$

Since $\mathbb{C}\mathrm{P}^1$ is a smooth projective curve and π is a proper birational morphism onto C, it is the normalization of C ([4, normalization of a variety]); write $\widetilde{C} = \mathbb{C}\mathrm{P}^1$ for it from here on.

The singular locus. In the chart $z = 1$ set $f = y^2 - x^3 - x^2$, so $f_x = -3x^2 - 2x$ and $f_y = 2y$. A singular point of C in this chart has $y = 0$ and $x(3x + 2) = 0$; the value $x = -\frac{2}{3}$ gives $y^2 = x^2(x + 1) = \frac{4}{27} \neq 0$, contradicting $y = 0$, so the origin is the only one. In the chart $y = 1$ the equation is $z - x^3 - x^2z$, whose derivative in z is $1 - x^2$, equal to 1 at $[0 : 1 : 0]$, so that point is smooth. Thus C is singular exactly at p. The quadratic part of f at the origin is $y^2 - x^2 = (y - x)(y + x)$, a product of two distinct linear forms, so p is a node, and its two branch tangents $y = \pm x$ correspond to $t/s = \pm 1$, that is to a and b.

The topological model. The map π is a closed continuous surjection identifying exactly a with b, so it induces a continuous bijection from $\widetilde{C}/(a \sim b)$ onto C; the source is compact and the target Hausdorff, so this is a homeomorphism. Thus C is the Riemann sphere with two points glued together.

The square that example 8.2.8 attaches to a nodal curve is therefore, for this C,

$$\begin{array}{ccc} \Sigma & \hookrightarrow & \widetilde{C} \\ \downarrow & & \downarrow{\scriptstyle \pi} \\ \{p\} & \hookrightarrow & C \end{array}$$

with $\Sigma = \{a, b\}$. The three corners other than the augmentation, namely $\widetilde{C}$, $\{p\}$ and Σ, are smooth and projective, as definition 8.2.4(1) requires; the augmentation corner $X_\emptyset = C$ is the singular curve being resolved, and is singular at p. Together with the facts that π is proper and restricts to an isomorphism over $C \setminus \{p\}$, this is what makes the square a cubical hyperresolution of C with $r = 1$. Everything below is computed from the cohomology of those three smooth projective varieties.

Theorem 8.4.2: The Mixed Hodge Structure of a Nodal Cubic

Let C, $\pi\colon \widetilde{C} \to C$, p and $\Sigma = \{a, b\}$ be as in example 8.4.1, and give $H^*(C, \mathbb{Q})$ the mixed Hodge structure of theorem 8.3.1. Then

1. $H^0(C, \mathbb{Q}) \cong \mathbb{Q}(0)$, pure of weight 0;
2. $H^1(C, \mathbb{Q})$ is the cokernel of
$$H^0(\widetilde{C}, \mathbb{Q}) \oplus H^0(\{p\}, \mathbb{Q}) \to H^0(\Sigma, \mathbb{Q}), \qquad (\alpha, \beta) \mapsto \alpha|_\Sigma - \beta|_\Sigma$$
hence one-dimensional; it is pure of weight 0, with $\mathrm{gr}_0^W H^1(C, \mathbb{Q}) = H^1(C, \mathbb{Q}) \cong \mathbb{Q}(0)$ and $\mathrm{gr}_1^W H^1(C, \mathbb{Q}) = 0$;
3. $H^2(C, \mathbb{Q}) \cong \mathbb{Q}(-1)$, pure of weight 2;
4. $H^k(C, \mathbb{Q}) = 0$ for $k \geq 3$.

***Proof*:**
Step 1: the first page. Apply proposition 8.2.7 to the hyperresolution square displayed above. Its index set consists of the nonempty subsets of a two-element set, so the resulting spectral sequence has two columns:

$$E_1^{0,t} = H^t(\widetilde{C}, \mathbb{Q}) \oplus H^t(\{p\}, \mathbb{Q}), \qquad E_1^{1,t} = H^t(\Sigma, \mathbb{Q}), \qquad E_1^{s,t} = 0 \ \text{ for } s \neq 0, 1$$

converging to $H^{s+t}(C, \mathbb{Q})$, with

$$d_1^{0,t}(\alpha, \beta) = \iota^*\alpha - \rho^*\beta$$

where $\iota\colon \Sigma \hookrightarrow \widetilde{C}$ and $\rho\colon \Sigma \to \{p\}$ are the two maps of the square. (Reversing the sign of d_1 changes neither its kernel nor its image, so the opposite sign convention gives the same answer below.) By theorem 8.3.1 each E_1 term carries the pure Hodge structure of the smooth projective variety it is the cohomology of, d_1 is a morphism of Hodge structures, and

$$E_2^{s,t} \cong \mathrm{gr}_t^W H^{s+t}(C, \mathbb{Q}) \tag{8.3}$$

once the sequence has degenerated, which Step 4 checks.

The three smooth corners are known. By example 1.2.5 the Hodge numbers of $\mathbb{C}\mathrm{P}^1$ are $h^{0,0} = h^{1,1} = 1$ and $h^{1,0} = h^{0,1} = 0$, so theorem 2.1.1 gives

$$H^0(\widetilde{C},\mathbb{Q}) \cong \mathbb{Q}(0), \qquad H^1(\widetilde{C},\mathbb{Q}) = 0, \qquad H^2(\widetilde{C},\mathbb{Q}) \cong \mathbb{Q}(-1), \qquad H^t(\widetilde{C},\mathbb{Q}) = 0 \ \text{ for } t \geq 3$$

the identifications of the two nonzero groups being those of definition 3.1.11: a one-dimensional rational Hodge structure of weight 0 and type $(0,0)$ is $\mathbb{Q}(0)$, and one of weight 2 and type $(1,1)$ is $\mathbb{Q}(-1)$, in each case by sending a rational basis vector to a rational basis vector. The point $\{p\}$ and the two-point set Σ are smooth projective of dimension zero, so $H^0(\{p\},\mathbb{Q}) \cong \mathbb{Q}(0)$, $H^0(\Sigma,\mathbb{Q}) \cong \mathbb{Q}(0)^2$, and both vanish in positive degree.

Step 2: the row $t = 0$. Identify $H^0(\Sigma,\mathbb{Q})$ with $\mathbb{Q}^{\{a,b\}}$. A class $\alpha \in H^0(\widetilde{C},\mathbb{Q})$ is a constant c, restricting to the pair (c,c), and a class $\beta \in H^0(\{p\},\mathbb{Q})$ is a scalar pulling back to (β,β). Hence

$$d_1^{0,0}(c,\beta) = (c-\beta)\,(1,1)$$

whose image is the line $\mathbb{Q}\,(1,1)$ and whose kernel is the line $\{(c,c)\}$. Therefore $E_2^{0,0}$ and $E_2^{1,0} = \mathbb{Q}^{\{a,b\}}/\mathbb{Q}\,(1,1)$ are both one-dimensional. Both are subquotients of Hodge structures of weight 0 and type $(0,0)$ by Step 1, so both are of type $(0,0)$: the kernel is a sub-Hodge structure and the cokernel a quotient by one, and lemma 7.1.5(3) makes each a Hodge structure of weight 0 with the induced filtration. Being one-dimensional, each is isomorphic to $\mathbb{Q}(0)$.

Step 3: the remaining rows. For $t = 1$ both columns vanish: $E_1^{0,1} = H^1(\widetilde{C},\mathbb{Q}) \oplus H^1(\{p\},\mathbb{Q}) = 0$ and $E_1^{1,1} = H^1(\Sigma,\mathbb{Q}) = 0$. For $t = 2$ one has $E_1^{0,2} = H^2(\widetilde{C},\mathbb{Q}) \cong \mathbb{Q}(-1)$ and $E_1^{1,2} = H^2(\Sigma,\mathbb{Q}) = 0$, so d_1 vanishes on this row and $E_2^{0,2} \cong \mathbb{Q}(-1)$, $E_2^{1,2} = 0$. For $t \geq 3$ both columns vanish, since $H^t(\widetilde{C},\mathbb{Q}) = 0$ and Σ and $\{p\}$ are finite.

Step 4: degeneration. The differential d_r has bidegree $(r, 1-r)$, so it maps $E_r^{s,t}$ to $E_r^{s+r,t-r+1}$. For $r \geq 2$ the target has first index $s + r \geq 2$, where every term vanishes by Step 1. Hence $d_r = 0$ for all $r \geq 2$ and $E_2 = E_\infty$, which is what (8.3) required.

Step 5: reading off the four statements. The weight filtration on $H^k(C,\mathbb{Q})$ is finite and exhaustive by definition 7.1.1, so $\dim_{\mathbb{Q}} H^k(C,\mathbb{Q}) = \sum_t \dim_{\mathbb{Q}} \mathrm{gr}_t^W H^k(C,\mathbb{Q})$, and by (8.3) the summands are the $E_2^{k-t,t}$ with $k - t \in \{0,1\}$.

For $k = 0$ the only contribution is $E_2^{0,0} \cong \mathbb{Q}(0)$, so $H^0(C,\mathbb{Q})$ is one-dimensional with $\mathrm{gr}_0^W = \mathbb{Q}(0)$ and all other graded pieces zero. This is (1).

For $k = 1$ the contributions are $E_2^{1,0}$, one-dimensional and isomorphic to $\mathbb{Q}(0)$ by Step 2, and $E_2^{0,1} = 0$ by Step 3. So $H^1(C,\mathbb{Q})$ is one-dimensional, $\mathrm{gr}_1^W H^1(C,\mathbb{Q}) = 0$, and $\mathrm{gr}_0^W H^1(C,\mathbb{Q}) \cong \mathbb{Q}(0)$. Since W is exhaustive and every graded piece other than the zeroth vanishes, $W_{-1}H^1(C,\mathbb{Q}) = 0$ and $W_0 H^1(C,\mathbb{Q}) = H^1(C,\mathbb{Q})$, so $H^1(C,\mathbb{Q}) = \mathrm{gr}_0^W H^1(C,\mathbb{Q}) \cong \mathbb{Q}(0)$ and the structure is pure of weight 0. The identification of $E_2^{1,0}$ in Step 2 is the asserted cokernel, which gives (2).

For $k = 2$ the contributions are $E_2^{2,0} = 0$, $E_2^{1,1} = 0$ and $E_2^{0,2} \cong \mathbb{Q}(-1)$, so $H^2(C,\mathbb{Q})$ is one-dimensional with $\mathrm{gr}_2^W \cong \mathbb{Q}(-1)$ and all other graded pieces zero; as in the previous paragraph this makes $H^2(C,\mathbb{Q}) \cong \mathbb{Q}(-1)$, pure of weight 2, which is (3).

For $k \geq 3$ the contributions are $E_2^{0,k}$, a subquotient of $H^k(\widetilde{C},\mathbb{Q}) = 0$, and $E_2^{1,k-1}$, a subquotient of $H^{k-1}(\Sigma,\mathbb{Q}) = 0$ since $k - 1 \geq 2$. So every graded piece vanishes and $H^k(C,\mathbb{Q}) = 0$, giving (4) and completing the proof.

Both numbers in degree one come from a corner of the square. A smooth plane cubic has H^1 of dimension two and pure of weight one, and those two dimensions are $H^1(\widetilde{C},\mathbb{Q})$; here the normalization is $\mathbb{C}\mathrm{P}^1$, that group vanishes, and nothing of weight one survives. What appears instead is one dimension of weight zero, contributed by the two points of Σ modulo the constants, that is by the two branches at the node modulo the identification imposed on them. The genus of the normalization and the combinatorics of the gluing are separate inputs, and the corollary below separates them in general.

Corollary 8.4.3: Weights on the First Cohomology of a Nodal Curve

Let C be an irreducible projective curve whose only singularities are $\delta \geq 0$ nodes, and let $\pi\colon \widetilde{C} \to C$ be its normalization, a smooth projective curve of genus g. Then

$$\dim_{\mathbb{Q}} H^1(C,\mathbb{Q}) = 2g + \delta$$

with $\mathrm{gr}_0^W H^1(C,\mathbb{Q}) \cong \mathbb{Q}(0)^\delta$ and $\mathrm{gr}_1^W H^1(C,\mathbb{Q}) \cong H^1(\widetilde{C},\mathbb{Q})$, and $\mathrm{gr}_m^W H^1(C,\mathbb{Q}) = 0$ for $m \neq 0, 1$.

***Proof*:**
For $\delta = 0$ the curve is smooth projective, $H^1(C,\mathbb{Q})$ is pure of weight one of dimension $2g$ by theorem 2.1.1, and the assertions hold. Assume $\delta \geq 1$, let $Z \subseteq C$ be the set of nodes and let $\Sigma = \pi^{-1}(Z)$. Each node has two branches with distinct tangents, so Σ consists of 2δ points, two of them over each node; π is proper and restricts to an isomorphism over $C \setminus Z$, so the square of example 8.2.8 applies verbatim and proposition 8.2.7 gives a two-column spectral sequence exactly as in Step 1 of theorem 8.4.2, degenerating at E_2 by the argument of Step 4 there.

The row $t = 0$. Since C is irreducible its normalization is irreducible, hence connected, so $H^0(\widetilde{C},\mathbb{Q}) = \mathbb{Q}$; and $H^0(Z,\mathbb{Q}) = \mathbb{Q}^\delta$, $H^0(\Sigma,\mathbb{Q}) = \mathbb{Q}^{2\delta}$. Writing a_j, b_j for the two points over the j-th node,

$$d_1^{0,0}\big(c,(\beta_j)_j\big) = (c - \beta_j) \quad \text{at both } a_j \text{ and } b_j$$

Its kernel is $\{(c,(c)_j)\}$, of dimension one, so its image has dimension $(1+\delta) - 1 = \delta$ and $E_2^{1,0} = H^0(\Sigma,\mathbb{Q})/\mathrm{im} d_1$ has dimension δ. As in Step 2 of theorem 8.4.2 it is a quotient of a Hodge structure of weight 0 and type $(0,0)$ by a sub-Hodge structure, hence itself of weight 0 and type $(0,0)$, and therefore isomorphic to $\mathbb{Q}(0)^\delta$ by definition 3.1.11.

The row $t = 1$. Both Z and Σ are finite, so $E_1^{1,1} = H^1(\Sigma,\mathbb{Q}) = 0$ and $E_1^{0,1} = H^1(\widetilde{C},\mathbb{Q})$, on which d_1 is therefore zero. Hence $E_2^{0,1} = H^1(\widetilde{C},\mathbb{Q})$, of dimension $2g$ by theorem 2.1.1 and corollary 2.2.1, the genus being $h^{1,0}(\widetilde{C}) = h^{0,1}(\widetilde{C})$.

Only $t = 0$ and $t = 1$ contribute to H^1, so $\mathrm{gr}_0^W H^1(C,\mathbb{Q}) = E_2^{1,0} \cong \mathbb{Q}(0)^\delta$, $\mathrm{gr}_1^W H^1(C,\mathbb{Q}) = E_2^{0,1} \cong H^1(\widetilde{C},\mathbb{Q})$, and the remaining graded pieces vanish. Summing the dimensions of the graded pieces of a finite exhaustive filtration (definition 7.1.1) gives $\dim_{\mathbb{Q}} H^1(C,\mathbb{Q}) = \delta + 2g$, completing the proof.

For the cubic, $g = 0$ and $\delta = 1$, which returns theorem 8.4.2(2). The corollary separates the two contributions: each node adds one dimension of weight zero, and each unit of genus of the

normalization adds two dimensions of weight one. Each contribution is realized in example 4.4.8, where the same nodal cubic occurs as one of the twelve singular members of a general pencil of plane cubics: a smooth member is an elliptic curve, with no node and H^1 of dimension two and weight one, while the nodal member has one node, normalization of genus zero, and H^1 of dimension one and weight zero. The Euler characteristics agree with the count carried out there. From theorem 8.4.2,

$$\chi_{\text{top}}(C) = 1 - 1 + 1 = 1$$

while a smooth member has $\chi_{\text{top}} = 1 - 2 + 1 = 0$, and exercise 4.3.1 (2)(1) predicts a difference of $(-1)^n = 1$ for a pencil on a surface, where $n = 2$.

The two Hodge-theoretic descriptions of the curve answer different questions. Example 4.4.8 works on the nearby smooth fibre and records how its weight-one Hodge structure is moved by transport around the puncture; theorem 8.4.2 works on the singular fibre itself and records the weight-zero class that the fibre carries. Set alongside the punctured curves of section 7.4, the three answers occupy three different weights in the same cohomological degree.

Example 8.4.4: Weight Zero, Weight One and Weight Two in Degree One
Three curves, one cohomological degree.

1. *Smooth and projective.* Let E be a smooth plane cubic. Then $H^1(E, \mathbb{Q})$ has dimension two and is pure of weight one, with $h^{1,0} = h^{0,1} = 1$ (example 3.4.3). The single weight equals the cohomological degree.

2. *Smooth and open.* Let $U = E \setminus \{x_1, x_2\}$. By theorem 7.4.3 with $n = 2$, $H^1(U, \mathbb{Q})$ has dimension three, $\mathrm{gr}_1^W \cong H^1(E, \mathbb{Q})$ and $\mathrm{gr}_2^W \cong \mathbb{Q}(-1)$. The weights are 1 and 2, at and above the degree, in the range $[k, 2k]$ of theorem 7.3.11.

3. *Projective and singular.* Let C be the nodal cubic. By theorem 8.4.2, $H^1(C, \mathbb{Q})$ has dimension one and is pure of weight 0, at the bottom of the range $[\max(0, k - n), k] = [0, 1]$ of theorem 8.3.1 for $k = n = 1$.

The three curves are related: C and E are members of one pencil (example 4.3.9), and U is an open subset of E. No two of the three groups are isomorphic as mixed Hodge structures, since their weights differ, and by lemma 7.1.5(2) the only morphism of Hodge structures between the pure structures in (1) and (3) is zero, in either direction.

The two failures of purity therefore point in opposite directions. Dropping compactness while keeping smoothness raises the weights above the cohomological degree, because theorem 7.3.11 builds $\mathrm{gr}^W_{k+m} H^k(U, \mathbb{Q})$ out of $H^{k-m}(D^{(m)}, \mathbb{Q})(-m)$ and each Tate twist adds two to the weight. Dropping smoothness while keeping compactness lowers them, because theorem 8.3.1 builds $\mathrm{gr}^W_t H^k(X, \mathbb{Q})$ out of H^t of the smooth pieces of a hyperresolution, and the pieces contributing to a fixed k have $t \leq k$, with $t < k$ exactly when the resolution needs a column beyond the first. The smooth projective case is the one where both mechanisms are absent and theorem 2.1.1 returns a single weight.

Every complex algebraic variety now carries a functorial mixed Hodge structure on its rational cohomology (theorem 8.3.1), and theorem 8.4.2 computes it in the smallest singular case. **??** turns from constructing these structures to using them, and asks which classes of $H^{2p}(X, \mathbb{Q}) \cap F^p H^{2p}(X, \mathbb{C})$ on a smooth projective X are classes of algebraic subvarieties. A general account of the mixed Hodge structure of a singular curve, including the cases left to the exercises, is in [33, the mixed Hodge structure of a curve].

Exercise 8.4.1

1. Let $C' = \{y^2z = x^3\} \subseteq \mathbb{CP}^2$, with singular point $p' = [0:0:1]$, and let $\pi'([s:t]) = [\,st^2 : t^3 : s^3\,]$. Show that $\pi' \colon \mathbb{CP}^1 \to C'$ is a morphism onto C', that $\pi'^{-1}(p')$ is the single point $[1:0]$, and that π' restricts to an isomorphism over $C' \setminus \{p'\}$. Then run the argument of theorem 8.4.2 on the resulting hyperresolution square and conclude that $H^1(C', \mathbb{Q}) = 0$, while $H^0(C', \mathbb{Q}) \cong \mathbb{Q}(0)$ and $H^2(C', \mathbb{Q}) \cong \mathbb{Q}(-1)$. A cusp, unlike a node, contributes nothing in degree one.

2. Compute $H^*(C, \mathbb{Q})$ with its mixed Hodge structure for the following two curves, in each case by writing down the hyperresolution square whose normalization is the disjoint union of the components, whose singular locus is the set of intersection points, and whose fourth corner is the preimage of that set.

   1. $C = \{xy = 0\} \subseteq \mathbb{CP}^2$, two lines meeting at one point.
   2. $C = \{xyz = 0\} \subseteq \mathbb{CP}^2$, three lines in general position.

   Note that corollary 8.4.3 does not apply, neither curve being irreducible, and that the two answers in degree one differ.

3. Let C be an irreducible projective curve whose only singularities are δ nodes, with normalization of genus g. Using corollary 8.4.3, show that $H^1(C, \mathbb{Q})$ is a pure Hodge structure if and only if $g = 0$ or $\delta = 0$, and identify the weight in each of the two cases.

4. Let C be the nodal cubic and let $U = E \setminus \{x_1, x_2\}$ be a twice-punctured elliptic curve, as in example 8.4.4.

   1. Show directly from definition 7.1.2 that every morphism of mixed Hodge structures $H^1(C, \mathbb{Q}) \to H^1(U, \mathbb{Q})$ is zero.
   2. Show, using theorem 7.1.8(3) and lemma 7.1.6(2), that every morphism of mixed Hodge structures $H^1(U, \mathbb{Q}) \to H^1(C, \mathbb{Q})$ is zero.
   3. Deduce from proposition 8.3.4 that no morphism of complex algebraic varieties between C and U, in either direction, induces a nonzero map on H^1.

5. Recompute the Betti numbers of the nodal cubic topologically, as a check on theorem 8.4.2. Let $D_a, D_b \subseteq \widetilde{C}$ be disjoint open discs around a and b, and set $V_1 = C \setminus \{p\}$ and $V_2 = \pi(D_a \cup D_b)$. Show that V_2 is open in C and contractible, that $V_1 \cong \mathbb{C}^*$, and that $V_1 \cap V_2$ is homotopy equivalent to a disjoint union of two circles. Then use the Mayer-Vietoris sequence, together with the fact that the two boundary circles represent classes summing to zero in $H_1(V_1, \mathbb{Z})$, to obtain $b_0 = b_1 = b_2 = 1$.

6. Let C be an irreducible projective curve with δ nodes and no other singularities, whose normalization has genus g. Extend the computation of corollary 8.4.3 to degrees 0 and 2, showing that $H^0(C, \mathbb{Q}) \cong \mathbb{Q}(0)$ and $H^2(C, \mathbb{Q}) \cong \mathbb{Q}(-1)$, and deduce

$$\chi_{\text{top}}(C) = 2 - 2g - \delta$$

   Check the value against exercise 4.3.1 (2)(1) at $n = 2$ for the nodal cubic.

What Next?

The Hodge decomposition, the period map and the two mixed Hodge structures of Part III are the classical core of the subject, and each of them continues past where this book stops. What follows names the directions and says where each is taken up.

Mixed Hodge modules. Chapters 7 and 8 put a mixed Hodge structure on the cohomology of a variety, one variety at a time. Saito's theory puts one on a sheaf. A mixed Hodge module on X is a filtered $\mathcal{D}_X$-module together with a perverse sheaf matched to it under the Riemann-Hilbert correspondence, carrying a weight filtration, and the class of such objects is stable under the six operations. The mixed Hodge structure on $H^*(X)$ is then a single instance of the formalism: it is what the pushforward to a point produces. Nothing in this series develops the theory, and the $\mathcal{D}$-module input it rests on is not developed here either. The planned *Perverse Sheaves* volume is the natural home, and does not yet claim it. Saito's own *Modules de Hodge polarisables* and *Mixed Hodge Modules* are the primary sources, and Christian Schnell's *An Overview of Morihiko Saito's Theory of Mixed Hodge Modules* is the readable way in.

The decomposition theorem. For a projective morphism $f\colon X \to Y$ with X smooth, the derived pushforward $Rf_*\mathbb{Q}_X[\dim X]$ splits in the derived category as a direct sum of shifted intersection complexes of subvarieties of Y. Taking Y a point recovers the primitive decomposition and hard Lefschetz of chapter 4, so the theorem is the relative form of the results that chapter proves absolutely. Beilinson, Bernstein, Deligne and Gabber proved it by reduction to positive characteristic and the Weil conjectures. Saito's Hodge modules give a proof native to the complex setting. Mark Andrea de Cataldo's *The Hodge Theory of Projective Manifolds* [16] is the compact account, and de Cataldo and Migliorini's survey is the standard reference on the theorem's history and its applications to intersection cohomology.

Nonabelian Hodge theory. The Hodge decomposition splits $H^1(X, \mathbb{C})$, which classifies representations of $\pi_1(X)$ into $\mathbb{C}^\times$. Ask the same question for representations into GL_n and the target is no longer a vector space, so no decomposition can be expected. What replaces it is an equivalence: on a compact Kähler X, semisimple representations of $\pi_1(X)$ correspond to polystable Higgs bundles, a holomorphic bundle E with a map $\theta\colon E \to E \otimes \Omega^1_X$ satisfying $\theta \wedge \theta = 0$, with vanishing rational Chern classes. Hitchin proved the rank-two case on a curve and Simpson the general one. The stack of principal bundles on a curve, which is the setting these moduli are built in, is constructed in

Moduli and Stacks [11]. The Higgs moduli themselves are developed nowhere in this series. Hitchin's *The Self-Duality Equations on a Riemann Surface* and Simpson's *Higgs Bundles and Local Systems* are the founding papers.

p-adic Hodge theory. Replace $\mathbb{C}$ by a finite extension K of $\mathbb{Q}_p$. A smooth proper X/K has de Rham cohomology with its Hodge filtration and p-adic étale cohomology with its action of $\mathrm{Gal}(\overline{K}/K)$, and the comparison theorems say these determine each other after extending scalars to one of Fontaine's period rings B_{dR}, B_{cris}, B_{st}. The Hodge filtration and the Galois action play the roles that $F^\bullet$ and complex conjugation play here. Representations of the Weil group of a local field, which is where the Galois side of this story is set up, are developed in *The Langlands Programme for* GL_2 [10]. A *p-adic Hodge Theory* volume is planned. Brinon and Conrad's *CMI Summer School Notes on p-adic Hodge Theory* are the standard introduction, and Bhatt, Morrow and Scholze's work on prismatic cohomology is where the subject went next.

Hodge theory and the Langlands programme. Two threads lead there from this book. The Mumford-Tate groups of chapter 6 are the reductive groups attached to Hodge structures, and a Shimura variety is a moduli space of Hodge structures whose Mumford-Tate group is prescribed, which is how Hodge theory gives the geometry underlying automorphic forms. A *Shimura Varieties* volume is planned. The second thread is geometric: the geometric Satake equivalence identifies representations of a reductive group with perverse sheaves on the affine Grassmannian of its dual, and weight filtrations there behave like the ones of Part III. Algebraic groups and their representations are developed in *Algebraic Groups and Representations* [14], which surveys geometric Satake in its closing chapter rather than proving it. Xinwen Zhu's *An Introduction to Affine Grassmannians and the Geometric Satake Equivalence* is the place to start.

Algebraic cycles and motives. **??** ends with two conjectures open, and they ask converse questions. The Hodge conjecture asks which cohomology classes are classes of algebraic cycles. Bloch's asks how much of a Chow group cohomology detects, and it is the first nontrivial case of a conjectural filtration on the Chow groups of a smooth projective variety, due to Bloch and Beilinson, functorial for correspondences and with gr^0 given by cycle classes. The category of motives is the attempt to build a universal cohomology theory in which both questions are natural, and Grothendieck's standard conjectures are what such a construction needs. The equivalence relations on cycles and the correspondence formalism are developed in *Intersection Theory* [8]. Yves André's *Une Introduction aux Motifs* is the survey, and the second volume of Voisin's *Hodge Theory and Complex Algebraic Geometry* [39] carries the cycle-theoretic material this book has drawn on throughout.

Index

Abel-Jacobi Map
  and the extension class, 270
Abelian Variety, 76
  projectivity criterion, 76
Analytification, 15

Base Locus, 107
Betti Number, 12
Blow-Up
  of the base locus of a pencil, 110

Cattani-Deligne-Kaplan Theorem, 205
Cohomological Descent, 288
Cohomology Bundle of a Family, 133
Compact Dual, 66
  as a homogeneous space, 69
Compactification
  good, 239
Complex Multiplication
  and the Mumford-Tate group, 197
Complex Torus, 7
  as a weight-one Hodge structure, 73
Connection
  Gauss-Manin, 133
Cubical Variety, 288
Curvature
  holomorphic sectional, 182

$\partial\bar{\partial}$-Lemma, 19
Degeneration
  of the Weight and Hodge Spectral Sequences, 258
Deligne Torus, 189
Deligne's Theorem, 295
Descent Double Complex, 288
Descent Spectral Sequence, 290
Differential Form
  with logarithmic poles, 239
Differential of the Third Kind, 266
Distance-Decreasing Property, 186
Divisor
  normal crossings, 238
Dolbeault Cohomology, 5
Dolbeault Complex, 5
Dolbeault's Theorem, 6

Elliptic Curve
  Hodge structure of, 78
  Mumford-Tate group of, 197
Exponential Sequence, 37
Extension Class
  of a punctured elliptic curve, 270
Extension of Hodge Structures, 227

Face Map, 286
Family
  smooth projective, 130
Filtration
  holomorphic degree, 9
Flat Connection, 133
Frölicher Inequality, 12
Frölicher Spectral Sequence, 10
Frolicher Spectral Sequence

degeneration, 30
Fubini-Study Metric, 7
Functoriality
of the Mixed Hodge Structure on a Singular Variety, 302

GAGA, 15
Gauss-Manin Connection, 133
flatness, 133
Griffiths transversality, 146
Lie-derivative formula, 143
periods, 135
General Point, 205
Global $\partial\bar{\partial}$-Potential, 21
Griffiths Transversality, 146
as horizontality of the period map, 158
graded form, 147
Griffiths' Description of the Hodge Filtration, 172
Griffiths-Schmid Curvature Estimate, 182

Hard Lefschetz Theorem, 96
Harmonic Form
$\bar{\partial}$-harmonic, 8
bidegree decomposition, 28
in a family, 141
Hermitian Metric, 7
Hermitian Symmetric Domain, 212
Higher Direct Image
of the constant sheaf, 131
Hironaka's Theorem, 239
Hodge Bundle, 139
holomorphy, 144
Hodge Class, 53, 192
as an invariant, 192
Hodge Complex, 276
cohomological, 276
Hodge Decomposition, 28
Hodge Filtration
by forms of type at least p, 142
in a family, 139
on a Hodge structure, 49
on the de Rham complex, 9
on the Logarithmic Complex, 252
Hodge Group, 196
relation to the Mumford-Tate group, 196
Hodge Index Theorem, 35
Hodge Isometry, 210
Hodge Locus, 200
algebraicity, 205
analyticity, 201
countability, 204
Zariski tangent space, 201
Hodge Metric, 180
Hodge Number, 12
Hodge Numbers, 6
constancy in a family, 141
of a smooth hypersurface, 173
symmetries of, 32
Hodge Star
on primitive covectors, 89
on primitive forms, 62
Hodge Structure
abelian category of mixed, 231
as a representation of the Deligne torus, 190
dual, 52, 191
dual of mixed, 235
filtration characterization, 50
mixed, 226
morphism, 51
morphism of mixed, 227
period domain of, 66
polarized, 55
pure, 47
strictness of morphisms, 51
sub-mixed, 227
Tate, 53
tensor product, 52, 191
tensor product of mixed, 235
weight, 47
weight one, 73
Hodge Symmetry, 32
Hodge-Riemann Relations
alternating signs, 59
in arbitrary weight, 57
positivity on primitive cohomology, 100
surface case, 33
Holomorphic Forms Are Closed, 22
Hopf Surface, 23
Horizontal Subbundle, 157
Hypercohomology
of a filtered complex of sheaves, 274
Hyperplane Section
weak Lefschetz theorem, 103
Hyperresolution
cubical, 288
existence, 289

Hypersurface
Hodge numbers of, 173
smooth, 162

Infinitesimal Period Relation, 157
Intersection Form
of a K3 surface, 36
of an abelian surface, 36

Jacobian Ideal, 161
Jacobian Ring, 161
and the differential of the period map, 174
Griffiths' description of the Hodge filtration, 172

K3 Lattice, 210
K3 Surface
global Torelli theorem, 216
Hodge structure of, 79
period domain of, 71, 212
polarized, 210
Kähler Identities, 8
Kähler Metric, 7

Laplacian
master identity $\Delta_d = 2\Delta_{\bar{\partial}}$, 16
Lattice
K3 lattice, 210
Lefschetz $(1,1)$-Theorem, 38
Lefschetz Decomposition, 98
of the exterior algebra, 87
Lefschetz Operator, 8
counting operator, 84
hard Lefschetz theorem, 96
primitive cohomology, 60
Lefschetz Pencil, 107
associated fibration, 110
existence, 108
Lefschetz Theorem
weak (hyperplane theorem), 103
Lefschetz Thimble, 113
Lefschetz Triple, 84
on cohomology, 95
Lie Algebra
Hodge structure on, 178
Local System
local constancy of fibre cohomology, 131
monodromy correspondence, 132
of fibre cohomology, 131

Logarithmic de Rham Complex, 239
comparison with smooth forms, 248
computes the cohomology of the complement, 249
local cohomology, 247
local freeness, 240
on an elliptic curve, 266
residue sequence, 244

Marking
of a K3 surface, 210
Metric
Hodge, 180
Milnor Fibre, 117
Mixed Hodge Complex, 277
cohomological, 277
theorem of, 280
Mixed Hodge Structure
of a punctured curve, 267
of a Singular Variety, 295
of a Smooth Variety, 260
Moduli
of polarized K3 surfaces, 216
Monodromy
of a Lefschetz pencil, 116
of a local system, 132
Picard-Lefschetz formula, 121
Monodromy Group
invariant classes, 125
of a smooth projective family, 152
proper discontinuity on the period domain, 154
Monodromy Representation, 116
Morse Lemma
holomorphic, 111
Mumford-Tate Group, 192
generic, 206
invariants are Hodge classes, 192
reductivity, 194
special, 196

Néron-Severi Group, 39
Nodal Cubic, 305
compared with a punctured curve, 309
mixed Hodge structure of, 306
Nodal Curve
weights in degree one, 308
Node

local model, 111
Noether-Lefschetz Locus, 200
Noether-Lefschetz Theorem, 207
Normal Crossings Divisor, 238
Strata, 252
Normalization
of a nodal cubic, 305

Odd Betti Obstruction, 23
Ordinary Double Point, 107

Pencil
of hyperplane sections, 107
Period
of a family, 135
Period Domain, 66
as a homogeneous space, 69
compact dual of, 66
curvature of, 182
distance-decreasing maps into, 186
Hodge metric on, 180
horizontal subbundle, 157
Lie algebra description of, 178
of polarized K3 surfaces, 212
openness in the compact dual, 68
weight two, 71
Period Map, 152
for polarized K3 surfaces, 216
holomorphy, 155
horizontality, 158
of a family of elliptic curves, 153
period data of a family, 150
rigidity, 187
Period Point
of a K3 surface, 79
Picard Number
of a very general surface in $\mathbb{C}\mathrm{P}^3$, 207
Picard-Lefschetz Formula, 121
Poincaré Lemma
holomorphic, 246
Polarization
and reductivity, 194
of a Hodge structure, 55
of primitive cohomology, 62, 100
signature of, 59
Primitive Cohomology, 60
as a sub-Hodge structure, 60
in a family, 150
Lefschetz decomposition, 98
Primitive Form, 84
Lefschetz decomposition, 87
Punctured Elliptic Curve, 267
purity for a single puncture, 269

Residue
isomorphism onto primitive cohomology, 169
of a class on the complement, 166
of a logarithmic form, 243
Residue Sequence, 244
Residue Theorem
on a complex torus, 264
Riemann Relations
complex-torus form, 73
for a form with simple poles, 264
for an elliptic curve, 78
Rigidity
of horizontal maps, 187

Serre Duality, 8
Siegel Upper Half Space, 67
Hodge metric on, 185
Simplicial Variety, 286
$\mathfrak{sl}_2$-Representation
on cohomology, 95
weights and raising operators, 94
Smooth Projective Family, 130
Spectral Sequence
Frölicher, 10
Hodge-de Rham, 10
Weight, 258, 280
Strictness
and E_1-degeneration, 275
for mixed Hodge structures, 231

Tate Twist, 53
Torelli Theorem
global, for K3 surfaces, 216
Torus
Deligne, 189
Tube Map, 166
Two Filtrations
Deligne's lemma, 257

Vanishing Cycle, 113
conjugacy under monodromy, 125
local model, 111, 117
Picard-Lefschetz formula, 121

Very General Point, 205

Weight Filtration, 226
    of a nodal curve, 306
    on the Logarithmic Complex, 251
    rationality, 257
Weil Identity, 62
    proof of, 89
Weil Operator, 55
    associated Hermitian form, 57

Bibliography

[1] James Carlson, Stefan Müller-Stach, and Chris Peters. *Period Mappings and Period Domains*. 2nd ed. Cambridge Studies in Advanced Mathematics 168. Cambridge: Cambridge University Press, 2017. ISBN: 978-1-316-63956-6. DOI: 10.1017/9781316995846.

[2] Eduardo Cattani, Pierre Deligne, and Aroldo Kaplan. "On the Locus of Hodge Classes". In: *Journal of the American Mathematical Society* 8.2 (1995), pp. 483–506. DOI: 10.1090/s0894-0347-1995-1273413-2. URL: https://doi.org/10.1090/s0894-0347-1995-1273413-2.

[3] Nathanael Chwojko-Srawley. *Everything You Need To Know About Abelian Varieties*. 1st ed. 2026. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_abelian_varieties.pdf.

[4] Nathanael Chwojko-Srawley. *Everything You Need to Know about Algebraic Geometry*. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_algGeo.pdf.

[5] Nathanael Chwojko-Srawley. *Everything You Need To Know About Complex Analytic Geometry*. 1st ed. 2026. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA-complex-analytic-geometry.pdf.

[6] Nathanael Chwojko-Srawley. *Everything You Need To Know About Elliptic Curves*. 1st ed. 2025. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_elliptic_curves.pdf.

[7] Nathanael Chwojko-Srawley. *Everything You Need To Know About Homological Algebra*. 1st ed. 2025. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_homological_algebra.pdf.

[8] Nathanael Chwojko-Srawley. *Everything You Need to Know about Intersection Theory*. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA-intersection-theory.pdf.

[9] Nathanael Chwojko-Srawley. *Everything You Need To Know About Linear Elliptic and Parabolic PDE*. 1st ed. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_linear_pde.pdf.

[10] Nathanael Chwojko-Srawley. *Everything You Need To Know About Local Langlands in GL2*. URL: https://nathanaelsrawley.com/pdf-viewer?src=.%2F.%2Fassets%2Fpdfs%2Fbooks%2FEYNTKA-langlands-gl2.pdf&name=Langlands%20for%20GL2(F)&source=books.

[11] Nathanael Chwojko-Srawley. *Everything You Need to Know about Moduli, Deformation Theory, and Stacks*. 1 vols. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA-moduli-stacks.pdf.

[12] Nathanael Chwojko-Srawley. *Everything You Need To Know About Riemannian Geometry*.

[13] Nathanael Chwojko-Srawley. *Everything You Need To Know About Shimura Varieties*.

[14] Nathanael Chwojko-Srawley. *Everyting You Need To Know About Geometric Representation-Theory*.

[15] Nathanael Chwojko-Srawley. *EYNTKA Classical Algebraic Geometry*.

[16] Mark Andrea de Cataldo. *The Hodge Theory of Projective Manifolds*. London: Imperial College Press, 2007. ISBN: 978-1-86094-800-4. DOI: 10.1142/p531.

[17] P. Deligne. "Hodge Cycles on Abelian Varieties". In: *Lecture Notes in Mathematics*. Berlin, Heidelberg: Springer Berlin Heidelberg, 1982, pp. 9–100. ISBN: 978-3-540-11174-0. DOI: 10.1007/978-3-540-38955-2_3. URL: https://doi.org/10.1007/978-3-540-38955-2_3.

[18] Pierre Deligne. "Théorie de Hodge : II". In: *Publications Mathématiques de l'IHÉS* 40 (Dec. 7, 1971), pp. 5–57. DOI: 10.1007/bf02684692. URL: https://doi.org/10.1007/bf02684692.

[19] Pierre Deligne. "Théorie de Hodge : III". In: *Publications Mathématiques de l'IHÉS* 44 (Jan. 7, 1974), pp. 5–77. DOI: 10.1007/bf02685881. URL: https://doi.org/10.1007/bf02685881.

[20] Phillip Griffiths and Joseph Harris. *Principles of Algebraic Geometry*. 1st ed. John Wiley & Sons, Ltd, 1978. 824 pp. ISBN: 978-0-471-05059-9. DOI: 10.1002/9781118032527. URL: https://onlinelibrary.wiley.com/doi/book/10.1002/9781118032527.

[21] Phillip Griffiths and Wilfried Schmid. "Locally Homogeneous Complex Manifolds". In: *Acta Mathematica* 123.0 (1969), pp. 253–302. DOI: 10.1007/bf02392390. URL: https://doi.org/10.1007/bf02392390.

[22] Phillip A. Griffiths. "Periods of Integrals on Algebraic Manifolds, I. (Construction and Properties of the Modular Varieties)". In: *American Journal of Mathematics* 90.2 (Apr. 1968), p. 568. DOI: 10.2307/2373545. URL: https://doi.org/10.2307/2373545.

[23] Alexander Grothendieck. "On the de Rham Cohomology of Algebraic Varieties". In: *Publications Mathématiques de l'IHÉS* 29 (Jan. 7, 1966), pp. 95–103. DOI: 10.1007/bf02684807. URL: https://doi.org/10.1007/bf02684807.

[24] Heisuke Hironaka. "Resolution of Singularities of an Algebraic Variety Over a Field of Characteristic Zero: I". In: *The Annals of Mathematics* 79.1 (Jan. 1964), p. 109. DOI: 10.2307/1970486. URL: https://doi.org/10.2307/1970486.

[25] W. V. D. Hodge. *The Theory and Applications of Harmonic Integrals*. Cambridge, New York: Cambridge University Press, 1989. ISBN: 978-0-521-35881-1. URL: http://openlibrary.org/books/OL1791471M/The_theory_and_applications_of_harmonic_integrals.

[26] James E. Humphreys. *Introduction to Lie Algebras and Representation Theory*. Vol. 9. Graduate Texts in Mathematics. Springer-Verlag, 1972.

[27] Daniel Huybrechts. *Complex Geometry: An Introduction*. Universitext. Berlin, Heidelberg: Springer, 2005. ISBN: 978-3-540-21290-4. DOI: 10.1007/b137952.

[28] Daniel Huybrechts. *Lectures on K3 Surfaces*. Cambridge University Press, 2016. ISBN: 978-1-107-15304-2. URL: http://openlibrary.org/books/OL28609296M/Lectures_on_K3_Surfaces.

[29] Erich Kähler. "Über Eine Bemerkenswerte Hermitesche Metrik". In: *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* 9.1 (Dec. 1933), pp. 173–186. DOI: 10.1007/bf02940642. URL: https://doi.org/10.1007/bf02940642.

[30] Nathanael Chwojko-Srawley. *Everything You Need To Know About Algebraic Topology*. 1st ed. 126 pp. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_algebraic_topology.pdf.

[31] Nathanael Chwojko-Srawley. *Everything You Need To Know About Complex Analysis*. 1st ed. 233 pp. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_complex_analysis.pdf.

[32] Nathanael Chwojko-Srawley. *Everything You Need To Know About Differential Topology*. 1st ed. 233 pp. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_differential_geometry.pdf.

[33] Chris A. M. Peters and Joseph H. M. Steenbrink. *Mixed Hodge Structures*. Ergebnisse Der Mathematik Und Ihrer Grenzgebiete. 3. Folge / a Series of Modern Surveys in Mathematics 52. Berlin, Heidelberg: Springer, 2008. ISBN: 978-3-540-77015-2. DOI: 10.1007/978-3-540-77017-6.

[34] I. I Pjateckiĭ-Šapiro and I R Šafarevič. "A TORELLI THEOREM FOR ALGEBRAIC SURFACES OF TYPE$K3$". In: *Mathematics of the USSR-Izvestiya* 5.3 (June 30, 1971), pp. 547–588. DOI: 10.1070/im1971v005n03abeh001075. URL: https://doi.org/10.1070/im1971v005n03abeh001075.

[35] Wilfried Schmid. "Variation of Hodge Structure: The Singularities of the Period Mapping". In: *Inventiones Mathematicae* 22.3–4 (Sept. 1973), pp. 211–319. DOI: 10.1007/bf01389674. URL: https://doi.org/10.1007/bf01389674.

[36] Jean-Pierre Serre. "Géométrie Algébrique et Géométrie Analytique". In: *Annales de l'Institut Fourier* 6 (1956), pp. 1–42. DOI: 10.5802/aif.59. URL: https://doi.org/10.5802/aif.59.

[37] Nathanael Chwojko-Srawley. *Everything You Need To Know About Lie Groups*.

[38] Claire Voisin. *Hodge Theory and Complex Algebraic Geometry I*. Trans. by Leila Schneps. Cambridge Studies in Advanced Mathematics 76. Cambridge: Cambridge University Press, 2002. 1 p. ISBN: 978-0-511-61534-4. DOI: 10.1017/CBO9780511615344.

[39] Claire Voisin. *Hodge Theory and Complex Algebraic Geometry II*. Trans. by Leila Schneps. Cambridge Studies in Advanced Mathematics 77. Cambridge: Cambridge University Press, 2003. ISBN: 978-0-521-71802-8. DOI: 10.1017/CBO9780511615177.

[40] Hermann Weyl. "On Hodge's Theory of Harmonic Integrals". In: *The Annals of Mathematics* 44.1 (Jan. 1943), p. 1. DOI: 10.2307/1969060. URL: https://doi.org/10.2307/1969060.