
Elementary Analysis

— *EYNTKA* Series —

NATHANAEL CHWOJKO-SRAWLEY

September 28, 2026

Contents

Contents	3
Introduction	i
1 The Real Numbers	1
1.1 Construction of the Real Numbers	2
1.2 Completeness and the Least Upper Bound Property	9
1.3 The Calculus of Suprema and Absolute Value	12
1.4 Consequences of Completeness	21
1.5★ Uniqueness of the Complete Ordered Field	29
2 Sequences and Series	33
2.1 Convergence of Real Sequences	34
2.2 Monotone Sequences, Limsup and Liminf	42
2.3 The Cauchy Criterion	55
2.4 Series and the Comparison Tests	61
2.5 Conditional Convergence and Rearrangement	78
2.6 Power Series and the Radius of Convergence	91
3 Limits and Continuity	99
3.1 Limits of Functions	100
3.2 Continuous Functions	115
3.3 Continuous Functions on a Closed Bounded Interval	129

3.4	Uniform Continuity	140
4	Differentiation	147
4.1	The Derivative	148
4.2	The Mean Value Theorem	157
4.3	Taylor's Theorem	169
4.4	Uniform Convergence	173
4.5	Power Series as Functions	182
4.6	The Elementary Functions	195
4.7	Analyticity and the Growth of Derivatives	207
5	The Riemann Integral	213
5.1	The Riemann Integral	214
5.2	Classes of Integrable Functions	227
5.3	The Fundamental Theorem of Calculus	238
5.4★	The Riemann-Stieltjes Integral	251
6	What Next?	261
A	The Construction by Dedekind Cuts	263
A.1	Cuts and the Field They Form	263
	Index	273

Introduction

A reader arriving here can almost certainly differentiate a polynomial, test a series for convergence and evaluate an integral. What that reader has not been given is any reason to believe the rules being applied. Calculus is taught as a body of technique, and the technique works, but the justification offered for it is usually a picture: a curve, a tangent line, an area shaded under a graph. The subject of this book is what has to be true about the real numbers for those pictures to be theorems.

The gap is not hypothetical, and it can be seen with nothing more than the rationals. A polynomial with rational coefficients can be negative at one rational point and positive at another and take the value zero at no rational point between them: $x^2 - 2$ is negative at 0, positive at 2, and has no rational root. So the claim that a continuous function passing from below zero to above zero must cross zero, which every applied use of calculus assumes, is false over $\mathbb{Q}$. It is true over $\mathbb{R}$, and the difference between the two number systems is the whole content of that theorem. The same difference decides whether a bounded increasing sequence has a limit, whether a continuous function on a closed interval attains a maximum, and whether the area under a curve is a number at all.

So the book begins by building $\mathbb{R}$ from $\mathbb{Q}$ as a completed object, where the alternative is to assume a list of properties and check nothing. Everything after that is drawn from the one property that construction establishes. The reader who wants to see how little is assumed and how much follows can read Section 1.1 and then Section 1.2: the second proves, from the construction alone, that every nonempty set of reals with an upper bound has a least one, and no later chapter needs anything about $\mathbb{R}$ that does not follow from that.

Some of what follows is not what the pictures suggest. A series can converge to one number, and converge to a different number after its terms are reordered, and by a suitable reordering converge to any number at all. A function can be continuous at every irrational point and discontinuous at every rational one. A function can be differentiable everywhere while its derivative is discontinuous, and yet that derivative still cannot have a jump. Each of these is proved here, and each one is a place where the intuition trained on graphs gives the wrong answer.

What the Reader Will Be Able to Do

1. State completeness in each of the forms a proof might need, and choose between them: the least upper bound property, the convergence of Cauchy sequences, the nested interval property, and the existence of subsequential limits for bounded sequences.
2. Decide whether a given series converges, and say whether its sum depends on the order of its terms.
3. Prove the standard theorems about continuous and differentiable functions on an interval, and name for each one the hypothesis that fails if it is dropped.
4. Construct counterexamples on request: a monotone function whose discontinuities are exactly a prescribed countable set, a bounded function that is not integrable, an integrable function with infinitely many discontinuities.
5. Read the analysis prerequisites of a graduate text in any field, and recognize which arguments there are arguments about the real line and which are arguments about distance, order or measure.

Omitted Material

Under the title, a reader might reasonably look for the following and will not find it here.

- *Metric and function spaces.* Completeness of a general metric space, compactness, the contraction principle, Stone-Weierstrass, Arzelà-Ascoli and Baire category. This needs its own book, and has one: *Analysis on Metric and Function Spaces* [1]. Several results proved here for $\mathbb{R}$ are the interval case of a theorem stated there in general, and the division is deliberate: the arguments in this book are about the order and completeness of the line, and generalizing them first would hide which property each one uses.
- *Functions of several variables.* Partial derivatives, the inverse and implicit function theorems, and multiple integrals. *Multivariable Calculus* [2], which takes this book as its prerequisite.
- *The Lebesgue integral*, and with it the characterization of Riemann integrability by the size of the discontinuity set. That characterization is the natural end of Chapter 5 and cannot be stated without measure. *Real Analysis* [5].
- *Topology as a subject.* Open sets as a structure, with connectedness and compactness as axioms, where this book meets both only as theorems about intervals. The point-set vocabulary of Section 3.1 is the fragment the real line needs and stops there. The general theory needs a non-prerequisite altitude and is *Topology* [3].
- *Fourier series.* A standard closing topic in books of this scope, and one that needs the function-space language this book does not develop. *Harmonic Analysis* [6], with the uniform-convergence machinery of Section 4.4 as the part of it this book does provide.
- *Everywhere continuous, nowhere differentiable functions.* Weierstrass's example is built from a uniformly convergent series of continuous functions, so Section 4.4 contains its ingredients,

but the estimate it needs is delicate and narrow in application for a first reading. *Analysis on Metric and Function Spaces* [1] constructs one.

- *Sets, functions and countability.* Taken as given throughout and developed upstream in *Preliminaries* [4]. Where a counting argument is needed, this book cites it and moves on.

The Real Numbers

Arithmetic in $\mathbb{Q}$ settles every question it raises about itself. Sums, products and quotients of rationals are rational, the order behaves under both operations, and a rational is determined by finitely much information. The failure appears at the first question that is not about arithmetic. A square of side 1 has a diagonal whose square is 2, and proposition 1.1.3 shows that no rational has square 2. What goes wrong is sharper than a missing element. Consider the set of rationals whose square is less than 2. It is bounded above, by 2 for instance. It has no smallest upper bound in $\mathbb{Q}$: given any rational upper bound, an explicit smaller one can be produced. So $\mathbb{Q}$ carries a set with upper bounds and no least upper bound, and that is a defect visible entirely from inside $\mathbb{Q}$, with no appeal to a larger system.

Repairing that defect is what this chapter does, and the repair is what every later chapter spends. The rationals are taken here as a given ordered field, as are the set-theoretic notions of function, bijection and countability. The real numbers are built from them, as equivalence classes of Cauchy sequences of rationals, so that completeness becomes a theorem about the construction, where the axiomatic route imposes it. The pattern of the chapter, and of the book, is to state first what fails in $\mathbb{Q}$ and then what completeness gives in its place.

The construction chosen here names a real number by a sequence of rationals closing in on it. The decimal expansions 1, 1.4, 1.41, 1.414, $\dots$ are such a sequence, and the construction is the observation that the sequence is already a serviceable name for the number it fails to reach. An older construction instead cuts $\mathbb{Q}$ in two at the missing number. Appendix A carries it, and theorem 1.5.2 shows the two produce the same $\mathbb{R}$.

Section 1.1 builds $\mathbb{R}$ from Cauchy sequences of rationals, gives it its field operations and its order, and places $\mathbb{Q}$ inside it. Section 1.2 proves the least upper bound property, which under this construction takes a bisection argument, where the cut route would allow a one-line union. Section 1.3 develops the arithmetic of suprema and the absolute value, the estimates later chapters use more than any other result here, and extracts n -th roots. Section 1.4 draws out the Archimedean property in its sharpened form, the density of $\mathbb{Q}$ and of the irrationals, the nested interval theorem and the uncountability of

$\mathbb{R}$. Section 1.5★ closes by showing that the construction was forced: any two complete ordered fields are isomorphic, so $\mathbb{R}$ is not one choice among many.

1.1 Construction of the Real Numbers

Five things are taken as given throughout, and every argument below uses only these. First, $\mathbb{Q}$ is an ordered field in the sense of definition 1.1.1. Second, $\mathbb{Q}$ is Archimedean, also in the sense of that definition. Third, every rational number can be written as m/n with $m \in \mathbb{Z}$ and n a positive integer. Fourth, a statement about $\mathbb{Z}$: every nonempty set of positive integers has a least element. Fifth, also about $\mathbb{Z}$, every integer is either even or odd, and a rational may be written in lowest terms in exactly one way. The parity clause is used only in proposition 1.1.3. The uniqueness half of the fifth is what makes Thomae's function well defined in section 3.2, which is also where the least element of the fourth is used a second time.

Definition 1.1.1: Ordered Field and the Archimedean Property

An *ordered field* is a field F together with a relation $<$ such that for all $x, y, z \in F$:

1. exactly one of $x < y$, $x = y$, $y < x$ holds;
2. if $x < y$ and $y < z$ then $x < z$;
3. if $x < y$ then $x + z < y + z$;
4. if $x < y$ and $0 < z$ then $xz < yz$.

Here $x > y$ means $y < x$, and $x \leq y$ means that $x < y$ or $x = y$. The element $x - y$ is $x + (-y)$, and $|x|$ denotes x if $x \geq 0$ and $-x$ otherwise. An ordered field F is *Archimedean* if for every $x \in F$ there is a positive integer n with $x < n \cdot 1$.

The consequences recorded next hold in every ordered field, so they are available for $\mathbb{Q}$ now and for $\mathbb{R}$ once theorem 1.1.10 is proved. Every later inequality manipulation in this book cites this proposition, since what a step uses is one of these numbered facts.

Proposition 1.1.2: Consequences of the Ordered Field Axioms

Let F be an ordered field and let $x, y, z, w \in F$.

1. Cancellation holds: $x + z = y + z$ implies $x = y$, and if $z \neq 0$ then $xz = yz$ implies $x = y$.
2. The elements $0, 1, -x$ and, for $x \neq 0$, x^{-1} are unique, and $-(-x) = x$.
3. $x \cdot 0 = 0$, and $xy = 0$ forces $x = 0$ or $y = 0$.
4. $(-x)y = -(xy)$ and $(-x)(-y) = xy$.
5. If $z > 0$ and $x < y$ then $xz < yz$; if $z < 0$ and $x < y$ then $yz < xz$.
6. $x < y$ if and only if $0 < y - x$, and $x < y$ if and only if $-y < -x$. Moreover $x \neq 0$ implies $x^2 > 0$, so $1 > 0$.
7. If $0 < x < y$ then $0 < y^{-1} < x^{-1}$.
8. If $x < y$ then $x < (x + y)(1 + 1)^{-1} < y$.
9. $|x + y| \leq |x| + |y|$, and $|xy| = |x||y|$.

Proof:

(1) and (2). Adding $-z$ to $x + z = y + z$ gives $x = y$. Multiplying $xz = yz$ by z^{-1} gives $x = y$. If $0'$ is a second additive identity then $0 = 0 + 0' = 0'$, and the same argument gives uniqueness of 1 . If y and y' both satisfy $x + y = 0 = x + y'$ then $y = y'$ by (1), and $-(-x) = x$ because x satisfies the defining property of $-(-x)$. For $x \neq 0$ the element x^{-1} is the only y with $xy = 1$, since $y = y \cdot 1 = y(xy^{-1}) = (yx)x^{-1} = x^{-1}$.

(3). From $x \cdot 0 = x(0 + 0) = x \cdot 0 + x \cdot 0$ and cancellation, $x \cdot 0 = 0$. If $xy = 0$ and $x \neq 0$ then $y = x^{-1}(xy) = x^{-1} \cdot 0 = 0$.

(4). $xy + (-x)y = (x + (-x))y = 0 \cdot y = 0$, so $(-x)y = -(xy)$. Applying this twice gives $(-x)(-y) = -(x(-y)) = -(-(xy)) = xy$.

(5). If $x < y$ and $z > 0$ then $0 < y - x$ by axiom (3), so $0 < (y - x)z$ by axiom (4), and $(y - x)z = yz - xz$ by distributivity and (4), whence $xz < yz$. If $z < 0$ then $-z > 0$, so $(y - x)(-z) > 0$. By (4) this element is $-(yz - xz)$, so $yz - xz < 0$, that is $yz < xz$.

(6). Adding $-x$ to $x < y$ gives $0 < y - x$, and adding x to $0 < y - x$ gives $x < y$. Adding $-x - y$ to $x < y$ gives $-y < -x$. If $x > 0$ then $x^2 > 0$ by axiom (4). If $x < 0$ then $-x > 0$ and $x^2 = (-x)(-x) > 0$ by (4) again. Since $1 \neq 0$ and $1 = 1 \cdot 1$, this gives $1 > 0$.

(7). If $0 < x$ then $x^{-1} \neq 0$, and $x^{-1} < 0$ would give $1 = xx^{-1} < 0$ by (5), contradicting (6). So $x^{-1} > 0$. Now $0 < x < y$ gives $x(xy)^{-1} < y(xy)^{-1}$ by (5), which is $y^{-1} < x^{-1}$.

(8). From $x < y$, adding x gives $x + x < x + y$ and adding y gives $x + y < y + y$. Since $1 + 1 > 0$ by (6), multiplying by $(1 + 1)^{-1}$, which is positive by (7), preserves both inequalities, and $(x + x)(1 + 1)^{-1} = x$.

(9). Both sides of the triangle inequality are unchanged by replacing (x, y) with $(-x, -y)$, so it suffices to treat $x + y \geq 0$, where $|x + y| = x + y \leq |x| + |y|$ because $x \leq |x|$ and $y \leq |y|$. The

| multiplicative identity follows by checking the four sign cases with (4), completing the proof.

The Archimedean property is a hypothesis on $\mathbb{Q}$ separate from the field and order axioms, and it is worth saying exactly where the construction spends it, since most of what follows does not. Neither lemma 1.1.6 nor the trichotomy argument of theorem 1.1.10 needs it: both hold in any ordered field. Density of $\mathbb{Q}$ in itself does not need it either, being proposition 1.1.2(8). It is used in exactly three places. Example 1.1.5(2) and (3) need an index beyond which a prescribed rational bound is beaten. Proposition 1.1.12 transports the property from $\mathbb{Q}$ to $\mathbb{R}$. And theorem 1.2.4 needs an index N with $(1 + 1)^{-(N-1)}$ smaller than a given rational, which is what makes its bisection terminate.

Arithmetic in $\mathbb{Q}$ answers every question about $\mathbb{Q}$ that arithmetic can pose, and fails at the first question that geometry poses.

Proposition 1.1.3: $\sqrt{2}$ Is Irrational

There is no $x \in \mathbb{Q}$ with $x^2 = 2$.

Proof:

Suppose $x \in \mathbb{Q}$ satisfies $x^2 = 2$. Write $x = m/n$ with $m \in \mathbb{Z}$ and n a positive integer, in lowest terms, so that m and n are not both even. Then $m^2 = 2n^2$.

An odd integer $m = 2j + 1$ has $m^2 = 4j^2 + 4j + 1 = 2(2j^2 + 2j) + 1$, which is odd. Since $m^2 = 2n^2$ is even, m is therefore not odd, so m is even, say $m = 2k$. Then $4k^2 = 2n^2$, so $n^2 = 2k^2$, and the same argument applied to n shows n is even. So m and n are both even, contradicting the choice of lowest terms, as we sought to show.

What proposition 1.1.3 exhibits is not merely a missing element. The rationals 1, 1.4, 1.41, 1.414, ... obtained by taking successively more decimal digits of a square root of 2 crowd together: any two terms beyond the k th differ by less than 10^{-k} . Inside $\mathbb{Q}$ that sequence approaches nothing at all, and the construction below is the observation that such a sequence is itself a perfectly good name for the number it fails to reach.

Definition 1.1.4: Cauchy Sequence of Rationals

A sequence $(a_n)_{n \geq 1}$ of rational numbers is a *Cauchy sequence* if for every rational $\epsilon > 0$ there is a positive integer N such that $|a_n - a_m| < \epsilon$ whenever $n, m \geq N$. The sequence is *null* if for every rational $\epsilon > 0$ there is a positive integer N with $|a_n| < \epsilon$ for all $n \geq N$.

Two features of the definition deserve stating before it is used. It quantifies over *rational* ϵ only, which is all that is available at this stage, and it never mentions a limit: the condition is entirely about the terms of the sequence and their distances from one another. That is what allows a sequence with no rational limit to satisfy it.

Example 1.1.5: Three Sequences of Rationals

1. The constant sequence $a_n = q$ is Cauchy for every $q \in \mathbb{Q}$, since $|a_n - a_m| = 0$.
2. Let a_n be the largest rational of the form $k/10^n$ with $k \in \mathbb{Z}$ and $(k/10^n)^2 < 2$. For $m \geq n$ the terms a_n and a_m agree to n decimal places, so $|a_n - a_m| \leq 10^{-n}$, and given a rational $\epsilon > 0$ the Archimedean property gives an N with $10^{-N} < \epsilon$. So (a_n) is Cauchy. By proposition 1.1.3 no rational is its limit.
3. The sequence $a_n = 1/n$ is null, again by the Archimedean property: given $\epsilon > 0$ choose N with $N > 1/\epsilon$.

Lemma 1.1.6: Cauchy Sequences Are Bounded

If (a_n) is a Cauchy sequence of rationals then there is a rational M with $|a_n| \leq M$ for every n .

Proof:

Take $\epsilon = 1$ in definition 1.1.4 and let N be the corresponding index. For $n \geq N$ the triangle inequality gives $|a_n| \leq |a_n - a_N| + |a_N| < 1 + |a_N|$. The finitely many terms $a_1, \dots, a_{N-1}$ have a largest absolute value, since a finite nonempty set of rationals has a largest element. Taking M to be the larger of that value and $1 + |a_N|$ bounds every term, completing the proof.

Two Cauchy sequences should name the same number exactly when they eventually become indistinguishable, and the following makes that precise and shows it behaves as an identification must.

Definition 1.1.7: Equivalence of Cauchy Sequences

Cauchy sequences (a_n) and (b_n) of rationals are *equivalent*, written $(a_n) \sim (b_n)$, if the sequence $(a_n - b_n)$ is null.

Lemma 1.1.8: Equivalence of Cauchy Sequences Is an Equivalence Relation

The relation $\sim$ of definition 1.1.7 is reflexive, symmetric and transitive on the set of Cauchy sequences of rationals.

Proof:

Reflexivity and symmetry are immediate, since $a_n - a_n = 0$ and $|a_n - b_n| = |b_n - a_n|$. For transitivity suppose $(a_n - b_n)$ and $(b_n - c_n)$ are null and let $\epsilon > 0$ be rational. Choose N_1 with $|a_n - b_n| < \epsilon(1+1)^{-1}$ for $n \geq N_1$ and N_2 with $|b_n - c_n| < \epsilon(1+1)^{-1}$ for $n \geq N_2$. For n at least the larger of N_1 and N_2 ,

$$|a_n - c_n| \leq |a_n - b_n| + |b_n - c_n| < \epsilon$$

by proposition 1.1.2(9), completing the proof.

Definition 1.1.9: Real Numbers

The set of *real numbers* $\mathbb{R}$ is the set of equivalence classes of Cauchy sequences of rationals under $\sim$. The class of (a_n) is written $[a_n]$, and a sequence belonging to a class is called a *representative* of it.

A real number is therefore not a single object of $\mathbb{Q}$ but a whole family of sequences agreeing in the limit, and every definition made on $\mathbb{R}$ below is first written in terms of a representative and then shown not to depend on which representative was chosen. That verification is not a formality: a formula on representatives that failed it would define nothing.

Theorem 1.1.10: $\mathbb{R}$ Is an Ordered Field

Define, for Cauchy sequences (a_n) and (b_n) of rationals,

$$[a_n] + [b_n] = [a_n + b_n], \quad [a_n] \cdot [b_n] = [a_n b_n]$$

and declare $[a_n] < [b_n]$ when there are a rational $\delta > 0$ and an index N with $b_n - a_n > \delta$ for all $n \geq N$. These are well defined, and with them $\mathbb{R}$ is an ordered field whose zero element is the class of the constant sequence 0 and whose unit element is the class of the constant sequence 1.

Proof:

Step 1: the operations produce Cauchy sequences. If (a_n) and (b_n) are Cauchy then so is $(a_n + b_n)$, since $|(a_n + b_n) - (a_m + b_m)| \leq |a_n - a_m| + |b_n - b_m|$ and each term can be made smaller than $\epsilon(1+1)^{-1}$. For the product, take bounds M_a and M_b from lemma 1.1.6 and write

$$|a_n b_n - a_m b_m| = |a_n(b_n - b_m) + b_m(a_n - a_m)| \leq M_a |b_n - b_m| + M_b |a_n - a_m|$$

so choosing N to make each difference smaller than $\epsilon(M_a + M_b + 1)^{-1}$ suffices.

Step 2: the operations are well defined. Suppose $(a_n) \sim (a'_n)$ and $(b_n) \sim (b'_n)$. Then $(a_n + b_n) - (a'_n + b'_n) = (a_n - a'_n) + (b_n - b'_n)$ is a sum of two null sequences and so is null. For the product, with bounds M for (a_n) and M' for (b'_n) ,

$$|a_n b_n - a'_n b'_n| = |a_n(b_n - b'_n) + b'_n(a_n - a'_n)| \leq M |b_n - b'_n| + M' |a_n - a'_n|$$

and the right-hand side is null. So both operations depend only on the classes.

Step 3: the field axioms. Commutativity, associativity and distributivity hold termwise in $\mathbb{Q}$ and therefore hold for classes. The class of the constant sequence 0 is an additive identity and $[-a_n]$ is an additive inverse of $[a_n]$. Only the multiplicative inverse needs an argument. Let $[a_n] \neq 0$, so (a_n) is not null: there is a rational $\delta > 0$ such that $|a_n| \geq \delta$ for infinitely many n . Choose N with $|a_n - a_m| < \delta(1+1)^{-1}$ for $n, m \geq N$ and pick $m \geq N$ with $|a_m| \geq \delta$. Then for every $n \geq N$,

$$|a_n| \geq |a_m| - |a_n - a_m| > \delta - \delta(1+1)^{-1} = \delta(1+1)^{-1}$$

so $a_n \neq 0$ for $n \geq N$. Define $b_n = a_n^{-1}$ for $n \geq N$ and $b_n = 0$ otherwise. For $n, m \geq N$,

$$|b_n - b_m| = \frac{|a_m - a_n|}{|a_n||a_m|} \leq |a_m - a_n| \cdot \frac{(1+1)^2}{\delta^2}$$

so (b_n) is Cauchy, and (a_nb_n) differs from the constant sequence 1 in at most the first $N-1$ terms, so $[a_n][b_n] = 1$.

Step 4: the order is well defined and total. Suppose $[a_n] < [b_n]$ with witnesses δ and N , and let $(a'_n) \sim (a_n)$ and $(b'_n) \sim (b_n)$. Choose $N' \geq N$ beyond which both $|a_n - a'_n|$ and $|b_n - b'_n|$ are less than $\delta(1+1+1+1)^{-1}$. For $n \geq N'$,

$$b'_n - a'_n = (b_n - a_n) + (b'_n - b_n) + (a_n - a'_n) > \delta - \delta(1+1)^{-1} = \delta(1+1)^{-1}$$

so the relation holds for the primed representatives with witness $\delta(1+1)^{-1}$. For trichotomy, let $[a_n]$ and $[b_n]$ be given and put $c_n = b_n - a_n$. If (c_n) is null the classes are equal. Otherwise, by the argument of Step 3 there are $\delta > 0$ and N with $|c_n| > \delta$ for all $n \geq N$. Since (c_n) is Cauchy it cannot change sign beyond the index where its terms differ by less than δ , so either $c_n > \delta$ for all large n , giving $[a_n] < [b_n]$, or $c_n < -\delta$ for all large n , giving $[b_n] < [a_n]$. Exactly one of the three holds, since a null sequence cannot satisfy either strict condition.

Step 5: the order axioms. Transitivity follows by adding the two witnesses. If $[a_n] < [b_n]$ with witness δ then for any $[c_n]$ the sequence $(b_n + c_n) - (a_n + c_n) = b_n - a_n$ exceeds δ beyond N , giving compatibility with addition. If moreover $0 < [c_n]$ with witness γ then $b_nc_n - a_nc_n = (b_n - a_n)c_n > \delta\gamma$ beyond the larger index, giving compatibility with multiplication, as we sought to show.

The construction has produced a field, and the next proposition places $\mathbb{Q}$ inside it so that the two arithmetics agree.

Proposition 1.1.11: $\mathbb{Q}$ Embeds in $\mathbb{R}$ as an Ordered Subfield

For $q \in \mathbb{Q}$ let q^* denote the class of the constant sequence with every term q . The map $q \mapsto q^*$ is injective and satisfies

$$q^* + r^* = (q + r)^*, \quad q^* r^* = (qr)^*, \quad -(q^*) = (-q)^*$$

for all $q, r \in \mathbb{Q}$, together with $q < r$ if and only if $q^* < r^*$. It sends 0 to the zero of $\mathbb{R}$ and 1 to its unit.

Proof:

The three displayed identities hold termwise. If $q < r$ then the constant sequence $r - q$ exceeds the rational $\delta = (r - q)(1 + 1)^{-1} > 0$ at every index, so $q^* < r^*$. Conversely $q^* < r^*$ excludes $q = r$ and, by trichotomy in $\mathbb{Q}$, excludes $r < q$, which would give $r^* < q^*$. Injectivity follows, since $q \neq r$ forces $q^* \neq r^*$, completing the proof.

From here on a rational q and its image q^* are written the same way, and $\mathbb{Q}$ is treated as a subset of $\mathbb{R}$. Theorem 1.1.10 then transfers every consequence recorded in proposition 1.1.2 to real numbers, including the triangle inequality, which section 1.3 develops in the form later chapters use.

The construction also settles, at once, a property that the cut route obtains only after the least upper bound property is available. It is recorded here because section 1.2 uses it to prove that property, and a proof there that leaned on theorem 1.4.1 would be circular.

Proposition 1.1.12: $\mathbb{R}$ Is Archimedean

For every $x \in \mathbb{R}$ there is a positive integer n with $x < n$.

Proof:

Let (a_n) be a representative of x and take a rational bound M with $|a_k| \leq M$ for all k , which exists by lemma 1.1.6. Write $M = p/q$ with $p \in \mathbb{Z}$ and q a positive integer, so $M \leq p$. Since $\mathbb{Q}$ is Archimedean there is a positive integer n with $p + 1 < n$. Then $n - a_k \geq n - M \geq n - p > 1$ for every k , so the constant sequence n exceeds (a_k) by the fixed rational 1 at every index, which is exactly $x < n$ in the sense of theorem 1.1.10, completing the proof.

1.1.13 Note: The Construction by Dedekind Cuts A second construction of $\mathbb{R}$ partitions $\mathbb{Q}$ where the one above sequences it: a *cut* is a pair $A|B$ of nonempty sets with $A \cup B = \mathbb{Q}$, every element of A below every element of B , and A having no largest element, and $\mathbb{R}$ is the set of cuts. Appendix A carries it in full. The two constructions produce ordered fields with the least upper bound property, and theorem 1.5.2 shows that any two such fields are isomorphic, so they produce the same $\mathbb{R}$. The cut route reaches the least upper bound property faster, by taking a union of lower sets. The sequence route reaches the Cauchy criterion faster, and generalizes to any metric space, which is the form the rest of this series uses.

Exercise 1.1.1

1. Show that a null sequence of rationals is Cauchy, and that a Cauchy sequence with a subsequence that is null is itself null.
2. Let (a_n) be Cauchy and let (b_n) agree with (a_n) for all but finitely many indices. Show that (b_n) is Cauchy and $[a_n] = [b_n]$. Deduce that altering finitely many terms of a representative changes nothing.
3. Show that the sequence $a_1 = 1$, $a_{n+1} = (a_n + 2a_n^{-1})(1 + 1)^{-1}$ of rationals is Cauchy, and that its class x satisfies $x^2 = 2$. Conclude that $\mathbb{R}$ contains an element $\mathbb{Q}$ does not.
4. Where does the proof of theorem 1.1.10 Step 3 use that (a_n) is not null? Termwise nonvanishing, that $a_n \neq 0$ for every n , is weaker. Give a Cauchy sequence with every term nonzero whose class has no multiplicative inverse.
5. Show that lemma 1.1.6 holds in *any* ordered field, using no Archimedean hypothesis. Then show that example 1.1.5(2) does need it, by exhibiting an ordered field in which no index N

satisfies $10^{-N} < \epsilon$ for a suitable ϵ .

6. Deduce from proposition 1.1.12 that between any two reals $x < y$ with $y - x > 1$ there is an integer, and that for every real $x > 0$ there is a positive integer n with $n^{-1} < x$.

1.2 Completeness and the Least Upper Bound Property

Section 1.1 produced an ordered field containing $\mathbb{Q}$. Every ordered field containing $\mathbb{Q}$ has that much, including $\mathbb{Q}$ itself, so nothing so far distinguishes $\mathbb{R}$. The property that does is the one the chapter opened on, and stating it needs two definitions first.

Definition 1.2.1: Upper and Lower Bound

Let $E \subseteq \mathbb{R}$. An element $u \in \mathbb{R}$ is an *upper bound* of E if $x \leq u$ for every $x \in E$, and E is *bounded above* if it has one. Symmetrically ℓ is a *lower bound* if $\ell \leq x$ for every $x \in E$, and E is *bounded below* if it has one.

An upper bound of E need not belong to E , and a set with one has many: if u is an upper bound and $u \leq v$, then $x \leq u \leq v$ for every $x \in E$, so v is an upper bound as well. The upper bounds of a set therefore form a collection closed upward in the order, and asking whether a set has an upper bound is a weak question. The same two definitions apply within $\mathbb{Q}$, with “rational number” in place of “real number” throughout, and a rational upper bound of a set of rationals is exactly an upper bound in the sense above that happens to lie in $\mathbb{Q}$ (proposition 1.1.11).

Since the upper bounds are closed upward, the informative question is whether there is a smallest one.

Definition 1.2.2: Supremum and Infimum

Let $E \subseteq \mathbb{R}$ be nonempty and bounded above. An element $s \in \mathbb{R}$ is the *supremum* of E , written $\sup E$, if s is an upper bound of E and $s \leq u$ for every upper bound u of E . Symmetrically, for E bounded below, m is the *infimum* $\inf E$ if m is a lower bound and $\ell \leq m$ for every lower bound ℓ .

A supremum is unique when it exists: if s and s' both qualify then $s \leq s'$ and $s' \leq s$, so $s = s'$ by trichotomy. Nothing yet says one exists.

Example 1.2.3: Suprema Present, Absent, and Outside the Field

1. Let $E_1 = \{x \in \mathbb{R} \mid 0 \leq x \leq 1\}$. Then $\sup E_1 = 1$, and the supremum belongs to E_1 .
2. Let $E_2 = \{x \in \mathbb{R} \mid 0 \leq x < 1\}$. Then $\sup E_2 = 1$ as well, and the supremum does not belong to E_2 . To see it, 1 is an upper bound. If $u < 1$ were an upper bound then proposition 1.1.2(8) gives v with $u < v < 1$, so $v \in E_2$ exceeds u , and u is not an upper bound after all.
3. Let $F = \{q \in \mathbb{Q} \mid q > 0 \text{ and } q^2 < 2\}$, regarded as a set of rationals. Then F has rational upper bounds, for instance 2, and *no* rational supremum. Suppose $u \in \mathbb{Q}$ were one. By proposition 1.1.3, $u^2 \neq 2$. If $u^2 < 2$, put $u' = u + (2 - u^2)(u + 2)^{-1} \cdot (1 + 1)^{-1}$. A computation gives $u' > u$ and $(u')^2 < 2$, so $u' \in F$ exceeds the upper bound u . If $u^2 > 2$, the same

expression gives $u' < u$ with $(u')^2 > 2$, so u' is a smaller upper bound. Either way u fails, so no rational supremum exists.

Part (3) is the failure the chapter opened on, now stated precisely: a set of rationals, bounded above in $\mathbb{Q}$, whose upper bounds have no least element in $\mathbb{Q}$. The theorem below says that $\mathbb{R}$ admits no such set. Under the construction of section 1.1 this takes genuine work, and the proof is the only place in the chapter where a real number is produced by naming a rational sequence and then reading off its class.

Theorem 1.2.4: The Least Upper Bound Property

Every nonempty subset of $\mathbb{R}$ that is bounded above has a supremum in $\mathbb{R}$.

Proof:

Let $E \subseteq \mathbb{R}$ be nonempty and bounded above, say by u_0 , and fix $x_0 \in E$.

Step 1: rational brackets. By proposition 1.1.12 there is a positive integer q with $u_0 < q$, and applying the same proposition to $-x_0$ gives a positive integer p with $-x_0 < p$, that is $-p < x_0$. Put $a_1 = -p$ and $b_1 = q$, both rational. Then b_1 is an upper bound of E , since $u_0 < b_1$, and a_1 is not, since $a_1 < x_0$ and $x_0 \in E$.

Step 2: bisection. Define a_{n+1} and b_{n+1} from a_n and b_n by letting $c_n = (a_n + b_n)(1 + 1)^{-1}$ and setting

$$(a_{n+1}, b_{n+1}) = \begin{cases} (a_n, c_n) & \text{if } c_n \text{ is an upper bound of } E \\ (c_n, b_n) & \text{otherwise} \end{cases}$$

Each c_n is rational whenever a_n and b_n are, so both sequences are rational throughout. Two properties hold at every stage, by induction: b_n is an upper bound of E and a_n is not, and $b_n - a_n = (b_1 - a_1)(1 + 1)^{-(n-1)}$. The first is immediate from the case split, and the second because bisection halves the gap.

Step 3: the brackets are Cauchy. The sequence (b_n) is decreasing and (a_n) increasing, and every a_n is below every b_m : for $n \leq m$ this is $a_n \leq a_m < b_m$, and for $n > m$ it is $a_n < b_n \leq b_m$. Hence for $n, m \geq N$ both b_n and b_m lie in the interval from a_N to b_N , so

$$|b_n - b_m| \leq b_N - a_N = (b_1 - a_1)(1 + 1)^{-(N-1)}$$

Given a rational $\epsilon > 0$, the Archimedean property of $\mathbb{Q}$ gives an N with $(b_1 - a_1)(1 + 1)^{-(N-1)} < \epsilon$, since $(1 + 1)^{N-1} \geq N$ for every positive integer N . So (b_n) is a Cauchy sequence of rationals, and (a_n) is equivalent to it because $b_n - a_n$ is null by the same estimate. Put $s = [b_n] \in \mathbb{R}$, which by definition 1.1.9 is a real number, and note $s = [a_n]$ as well.

Step 4: s is an upper bound. Let $x \in E$ and suppose $s < x$. Then $x - s > 0$, so by proposition 1.1.12 there is a positive integer k with $(x - s)^{-1} < k$, hence $k^{-1} < x - s$. Choose N with $b_N - a_N < k^{-1}$, which Step 3 permits. Since $s = [a_n]$ and (a_n) is increasing, $a_N \leq s$, so

$$b_N = a_N + (b_N - a_N) < s + k^{-1} < s + (x - s) = x$$

contradicting that b_N is an upper bound of E . So $x \leq s$ for every $x \in E$.

Step 5: s is the least upper bound. Let u be an upper bound of E and suppose $u < s$. Then $s - u > 0$, and as in Step 4 choose a positive integer k with $k^{-1} < s - u$ and then N with

$b_N - a_N < k^{-1}$. Since $s = [b_n]$ and (b_n) is decreasing, $s \leq b_N$, so

$$a_N = b_N - (b_N - a_N) > s - k^{-1} > s - (s - u) = u$$

So $a_N > u$, and since a_N is not an upper bound of E there is $x \in E$ with $a_N < x$. Then $u < a_N < x$ contradicts that u is an upper bound. Hence $s \leq u$ for every upper bound u , and $s = \sup E$, completing the proof.

The proof used the construction twice and used it in an essential way: Step 3 produced a real number by exhibiting a rational Cauchy sequence, and Steps 4 and 5 compared that number with others through proposition 1.1.12, which section 1.1 obtained from the boundedness of representatives, with no appeal to completeness. Reversing that order would be circular, and theorem 1.4.1 records the point where the Archimedean property is instead read off the least upper bound property, for readers who meet $\mathbb{R}$ axiomatically.

The corresponding statement for lower bounds needs no second argument.

Corollary 1.2.5: The Greatest Lower Bound Property

Every nonempty subset of $\mathbb{R}$ that is bounded below has an infimum in $\mathbb{R}$.

Proof:

Let E be nonempty and bounded below by ℓ , and put $-E = \{-x \mid x \in E\}$. If $\ell \leq x$ for every $x \in E$ then $-x \leq -\ell$ by proposition 1.1.2(6), so $-E$ is nonempty and bounded above by $-\ell$. Let $s = \sup(-E)$, which exists by theorem 1.2.4. Then $-s$ is a lower bound of E , since $-x \leq s$ gives $-s \leq x$. If ℓ' is any lower bound of E then $-\ell'$ is an upper bound of $-E$, so $s \leq -\ell'$ and hence $\ell' \leq -s$. So $-s = \inf E$, completing the proof.

The form in which the least upper bound property is applied is almost never the definition. What later arguments use is that a supremum is approached from below by elements of the set, and that this property characterizes it.

Proposition 1.2.6: The ϵ -Characterization of the Supremum

Let $E \subseteq \mathbb{R}$ be nonempty and let s be an upper bound of E . Then $s = \sup E$ if and only if for every real $\epsilon > 0$ there is an $x \in E$ with $s - \epsilon < x$.

Proof:

Suppose $s = \sup E$ and let $\epsilon > 0$. Then $s - \epsilon < s$, so $s - \epsilon$ is not an upper bound of E , since s is the least one. So some $x \in E$ satisfies $s - \epsilon < x$.

Conversely suppose the condition holds and let u be an upper bound of E with $u < s$. Taking $\epsilon = s - u > 0$ produces $x \in E$ with $s - (s - u) < x$, that is $u < x$, contradicting that u is an upper bound. So $s \leq u$ for every upper bound u , and $s = \sup E$, as we sought to show.

Exercise 1.2.1

1. Show that a set with a largest element has that element as its supremum, and give a set whose supremum is not a largest element.
2. Let $A \subseteq B \subseteq \mathbb{R}$ with A nonempty and B bounded above. Show that $\sup A \leq \sup B$.
3. Show that $\sup E$ is the unique real number s such that $x \leq s$ for all $x \in E$ and, for every $\epsilon > 0$, some $x \in E$ exceeds $s - \epsilon$.
4. Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above with $\sup A \leq \sup B$. Show that $A \cup B$ is bounded above and that $\sup(A \cup B) = \sup B$.
5. In the bisection of theorem 1.2.4, suppose E has a largest element M . Show that M is an upper bound at every stage, so that the case split sends b_{n+1} to the midpoint whenever the midpoint is at least M , and that the conclusion $s = M$ follows. (Note b_1 is chosen as an integer above u_0 and need not equal M .)
6. Give a nonempty set of rationals, bounded above in $\mathbb{Q}$, whose supremum in $\mathbb{R}$ is irrational, and explain why example 1.2.3(3) does not contradict theorem 1.2.4.
7. Show that $\mathbb{Q}$, with the order it inherits from $\mathbb{R}$, is an ordered field that does not satisfy the least upper bound property, so that theorem 1.2.4 is not a consequence of the ordered field axioms alone.
8. Let E be nonempty and bounded above and let $c > 0$. Show that $\sup(cE) = c \sup E$, where $cE = \{cx \mid x \in E\}$, and find what replaces this when $c < 0$.

1.3 The Calculus of Suprema and Absolute Value

Section 1.2 produces suprema and infima and says nothing about how they interact with the arithmetic of $\mathbb{R}$. The interactions needed later are few: a supremum over a larger set is at least as large, a nonnegative constant passes through a supremum, and the supremum of a set of sums is the sum of the suprema. The third of these is an equality for sets and only an inequality for functions, and the difference between the two statements is what the later chapters have to keep track of when they estimate a sum of two quantities that are large at different points.

The second half of the section introduces the absolute value, proves the triangle inequality in the forms later chapters use, and combines the two themes: the number $\sup_D f - \inf_D f$ is exactly the smallest constant bounding every difference $|f(x) - f(y)|$. The section closes with the first existence theorem proved from the least upper bound property, namely that every nonnegative real has a unique nonnegative n -th root. In $\mathbb{Q}$ the corresponding statement already fails for $n = 2$ and $x = 2$ by proposition 1.1.3, so this is the point at which $\sqrt{2}$ becomes a real number, where in $\mathbb{Q}$ it was a gap.

Every argument below begins the same way: a supremum is an upper bound, so it gives an inequality valid at every point of the set, and it is the *least* upper bound, so no smaller number has that property. The second half of that sentence is applied constantly, and always in the single form recorded in proposition 1.2.6: a supremum is approached from below by elements of the set. Only the reflected statement for the infimum is still owed.

Lemma 1.3.1: Approximation Property of the Infimum

Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $m = \inf A$. Then for every $\epsilon > 0$ there is an $a \in A$ with $m \leq a < m + \epsilon$.

Proof:

Let $\epsilon > 0$. Then $m + \epsilon > m$, so $m + \epsilon$ is not a lower bound of A , since m is the greatest lower bound. So some $a \in A$ satisfies $a < m + \epsilon$, and $m \leq a$ because m is a lower bound, completing the proof.

Read from right to left, proposition 1.2.6 is how a supremum is identified in practice: guess the value, check that it is an upper bound, and then check that some element of the set exceeds any smaller candidate. Read from left to right, it and the lemma above are how a supremum and an infimum are used once they are known. Neither asserts that the bound belongs to the set, and in general it does not. What they give instead is an element of the set within any prescribed tolerance of it.

The sum rule among the three manipulations promised above has a shape worth noting: the elements of $A + B$ are the sums $a + b$ where a and b range independently, so a may be pushed close to $\sup A$ while b is pushed close to $\sup B$, and nothing prevents doing both at once.

Proposition 1.3.2: The Algebra of Suprema for Sets

Let A and B be nonempty subsets of $\mathbb{R}$.

1. If $A \subseteq B$ and B is bounded above, then A is bounded above and $\sup A \leq \sup B$.
2. If A is bounded above and $c \geq 0$, then $cA = \{ca \mid a \in A\}$ is bounded above and $\sup(cA) = c \sup A$.
3. If A is bounded below, then $-A = \{-a \mid a \in A\}$ is bounded above and $\sup(-A) = -\inf A$.
4. If A and B are bounded above, then $A + B = \{a + b \mid a \in A, b \in B\}$ is bounded above and $\sup(A + B) = \sup A + \sup B$.

Proof:

1. Every $a \in A$ lies in B and therefore satisfies $a \leq \sup B$, so $\sup B$ is an upper bound of A and A is bounded above. Both sets are nonempty and bounded above, so both suprema exist by theorem 1.2.4, and $\sup A \leq \sup B$ because $\sup A$ is the least upper bound of A while $\sup B$ is one of its upper bounds.
2. Write $s = \sup A$. If $c = 0$ then $cA = \{0\}$, since A is nonempty, and $\sup\{0\} = 0 = 0 \cdot s$. Suppose $c > 0$. For $a \in A$ the inequality $a \leq s$ multiplies by the positive number c to give $ca \leq cs$ (proposition 1.1.2(5)), so cs is an upper bound of cA and cA is bounded above. Let u be any upper bound of cA . Since $c > 0$, the inverse $1/c$ is positive (proposition 1.1.2(7)), so multiplying $ca \leq u$ by $1/c$ gives $a \leq u/c$ for every $a \in A$. Thus u/c is an upper bound of A , whence $s \leq u/c$ and therefore $cs \leq u$. So cs is the least upper bound of cA .
3. Write $m = \inf A$, which exists by corollary 1.2.5 since A is nonempty and bounded below. For $a \in A$ the inequality $m \leq a$ gives $-a \leq -m$, so $-m$ is an upper bound of $-A$. Let u

be an upper bound of $-A$. Then $-a \leq u$ for every $a \in A$, hence $-u \leq a$ for every $a \in A$, so $-u$ is a lower bound of A and therefore $-u \leq m$, that is $-m \leq u$. So $-m$ is the least upper bound of $-A$.

4. Write $s = \sup A$ and $t = \sup B$. For $a \in A$ and $b \in B$, adding $a \leq s$ and $b \leq t$ gives $a + b \leq s + t$, so $s + t$ is an upper bound of $A + B$. The set $A + B$ is nonempty, so $\sigma = \sup(A + B)$ exists by theorem 1.2.4 and $\sigma \leq s + t$. Suppose $\sigma < s + t$ and put

$$\epsilon = \frac{s + t - \sigma}{2} > 0$$

By proposition 1.2.6 applied to A and then to B , there are $a \in A$ with $a > s - \epsilon$ and $b \in B$ with $b > t - \epsilon$. Adding,

$$a + b > s + t - 2\epsilon = \sigma$$

which contradicts the fact that σ is an upper bound of $A + B$. Hence $\sigma = s + t$, completing the proof.

Part (3) is the reason the theory is usually stated for suprema alone: an infimum is a supremum of the reflected set, so every statement above has a companion obtained by replacing A with $-A$. Part (2) fails as stated for $c < 0$, and part (3) says what replaces it: if $c < 0$ and A is bounded below, then $cA = (-c)(-A)$ with $-c > 0$, so parts (2) and (3) give

$$\sup(cA) = (-c)\sup(-A) = (-c)(-\inf A) = c\inf A$$

Multiplying by a negative number reverses the order, and the effect on the two bounds is to exchange them.

Example 1.3.3: Suprema of Two Intervals and Their Sum

Let $A = [0, 1)$ be the set of $x \in \mathbb{R}$ with $0 \leq x < 1$, and let $B = (-1, 2]$ be the set of $x \in \mathbb{R}$ with $-1 < x \leq 2$.

The number 1 is an upper bound of A . Given $\epsilon > 0$, put $a = \max\{0, 1 - \epsilon/2\}$. Then $a \geq 0$ and $a < 1$, so $a \in A$. Moreover $a > 1 - \epsilon$, because if $a = 1 - \epsilon/2$ this is the inequality $-\epsilon/2 > -\epsilon$, while if $a = 0$ then $1 - \epsilon/2 \leq 0$ forces $\epsilon \geq 2$ and hence $1 - \epsilon \leq -1 < 0 = a$. By proposition 1.2.6, $\sup A = 1$, and $1 \notin A$. The number 0 lies in A and is a lower bound of A , so $\inf A = 0$. For B no approximation argument is needed: the number 2 is an upper bound of B and belongs to B , so every upper bound of B is at least 2 and $\sup B = 2$.

Scaling by $c = 3$ gives $3A = [0, 3)$, since $x \mapsto 3x$ carries the condition $0 \leq a < 1$ to $0 \leq 3a < 3$, and proposition 1.3.2(2) gives $\sup(3A) = 3\sup A = 3$. For the sum, every $a + b$ with $a \in A$ and $b \in B$ satisfies $-1 < a + b < 3$. Conversely, if $-1 < z \leq 2$ then $z = 0 + z$ with $0 \in A$ and $z \in B$, while if $2 < z < 3$ then $z = (z - 2) + 2$ with $z - 2 \in (0, 1) \subseteq A$ and $2 \in B$. Hence $A + B = (-1, 3)$ and $\sup(A + B) = 3 = 1 + 2 = \sup A + \sup B$. The supremum of $A + B$ is not attained even though $\sup B$ is: the equality in proposition 1.3.2(4) is an identity between numbers and says nothing about which of them belong to their sets. Finally $-A = (-1, 0]$, whose supremum is $0 = -\inf A$.

Sets are replaced by functions throughout the rest of the book, and the translation is purely notational: for a nonempty set D and a function $f: D \rightarrow \mathbb{R}$, write $f(D) = \{f(x) | x \in D\}$ for the range and

$$\sup_D f = \sup f(D), \quad \inf_D f = \inf f(D)$$

calling f *bounded on D* when $f(D)$ is bounded above and below. What does not translate is the sum

rule. The set $(f + g)(D)$ consists of the sums $f(x) + g(x)$ taken at a *common* point x , so it is only part of $f(D) + g(D)$, and the supremum can drop.

Proposition 1.3.4: Suprema and Infima of a Sum of Functions

Let D be a nonempty set and let $f, g: D \rightarrow \mathbb{R}$ be bounded. Then $f + g$ is bounded and

$$\sup_D (f + g) \leq \sup_D f + \sup_D g, \quad \inf_D (f + g) \geq \inf_D f + \inf_D g$$

Both inequalities can be strict.

Proof:

Every element of $(f + g)(D)$ has the form $f(x) + g(x)$ for some $x \in D$, and $f(x) \in f(D)$ and $g(x) \in g(D)$, so

$$(f + g)(D) \subseteq f(D) + g(D)$$

The sets $f(D)$ and $g(D)$ are nonempty and bounded above, so proposition 1.3.2(4) applies and gives that $f(D) + g(D)$ is bounded above with supremum $\sup_D f + \sup_D g$. In particular $(f + g)(D)$ is bounded above, being a subset of a set bounded above, and proposition 1.3.2(1) gives

$$\sup_D (f + g) \leq \sup (f(D) + g(D)) = \sup_D f + \sup_D g$$

For the second inequality, note that $-f$ and $-g$ are bounded because f and g are, and that $(-f) + (-g) = -(f + g)$. The inequality just proved, applied to $-f$ and $-g$, therefore shows that $-(f + g)$ is bounded above, which is to say that $f + g$ is bounded below, and gives

$$\sup_D (-(f + g)) \leq \sup_D (-f) + \sup_D (-g) \tag{1.1}$$

By proposition 1.3.2(3) applied to ranges, $\sup_D (-h) = -\inf_D h$ for every bounded $h: D \rightarrow \mathbb{R}$, since $(-h)(D) = -h(D)$. Rewriting all three suprema in (1.1) by this identity turns it into

$$-\inf_D (f + g) \leq -\inf_D f - \inf_D g$$

and multiplying by -1 reverses the inequality to give $\inf_D (f + g) \geq \inf_D f + \inf_D g$. Both can be strict: on $D = [0, 1]$ with $f(x) = x$ and $g(x) = 1 - x$ the sum is the constant function 1, so $\sup_D (f + g) = 1$ while $\sup_D f + \sup_D g = 2$, and $\inf_D (f + g) = 1$ while $\inf_D f + \inf_D g = 0$. Example 1.3.5 verifies these four values, completing the proof.

Example 1.3.5: A Strict Inequality for $\sup(f + g)$

Let $D = [0, 1]$ and define $f, g: D \rightarrow \mathbb{R}$ by $f(x) = x$ and $g(x) = 1 - x$. Then $f(D) = [0, 1]$ and $g(D) = [0, 1]$, so $\sup_D f = \sup_D g = 1$ and $\inf_D f = \inf_D g = 0$, each attained at an endpoint. The sum is the constant function $(f + g)(x) = 1$, so

$$\sup_D (f + g) = 1 < 2 = \sup_D f + \sup_D g$$

and likewise $\inf_D (f + g) = 1 > 0 = \inf_D f + \inf_D g$. The gap has a single cause: f is large only near $x = 1$ and g only near $x = 0$, and no point of D makes both large at once. The set version

has no such obstruction, and indeed $f(D) + g(D) = [0, 2]$, whose supremum is 2.

Distances on the real line are measured by the absolute value, and every ϵ - δ and ϵ - N statement in the chapters that follow is an inequality between absolute values. The definition splits into two cases because the order on $\mathbb{R}$ does, and the point of proposition 1.3.8 is that the case split need almost never be performed again.

Definition 1.3.6: Absolute Value

For $x \in \mathbb{R}$ the *absolute value* of x is

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

For $x, y \in \mathbb{R}$ the number $|x - y|$ is called the *distance* from x to y .

Example 1.3.7: Distances Between Two Points of the Line

Directly from the definition, $|5| = 5$ because $5 \geq 0$, and $|-7/2| = -(-7/2) = 7/2$ because $-7/2 < 0$. The distance from $-7/2$ to 5 is

$$\left| -\frac{7}{2} - 5 \right| = \left| -\frac{17}{2} \right| = \frac{17}{2}$$

and computing it in the other order gives $|5 - (-7/2)| = |17/2| = 17/2$, the same number. Taking $y = 0$ recovers $|x - 0| = |x|$, so the absolute value of x is its distance from the origin.

Proposition 1.3.8: Comparison, Multiplicativity and the Triangle Inequality

Let $x, y \in \mathbb{R}$ and let $c \geq 0$.

1. $|x| \geq 0$, with $|x| = 0$ if and only if $x = 0$. Moreover $|-x| = |x|$ and $-|x| \leq x \leq |x|$.
2. $|x| \leq c$ if and only if $-c \leq x \leq c$, and $|x| < c$ if and only if $-c < x < c$.
3. $|xy| = |x| |y|$.
4. $|x + y| \leq |x| + |y|$.
5. $||x| - |y|| \leq |x - y|$.

Proof:

1. If $x \geq 0$ then $|x| = x \geq 0$; if $x < 0$ then $|x| = -x > 0$ by the sign rule of proposition 1.1.2(4). In both cases $|x| \geq 0$, and the second case shows $|x| = 0$ forces $x \geq 0$, whence $x = |x| = 0$. Conversely, $|0| = 0$. For $|-x| = |x|$: if $x > 0$ then $-x < 0$ and $|-x| = -(-x) = x = |x|$; if $x = 0$ both sides are 0; if $x < 0$ then $-x > 0$ and $|-x| = -x = |x|$. For the two-sided bound: if $x \geq 0$ then $x = |x|$ and $-|x| \leq 0 \leq x$, while if $x < 0$ then $x = -|x|$ and $x < 0 \leq |x|$.
2. Suppose $|x| \leq c$. By (1), $x \leq |x| \leq c$ and $-x \leq |-x| = |x| \leq c$, and the latter gives $-c \leq x$.

Conversely suppose $-c \leq x \leq c$. If $x \geq 0$ then $|x| = x \leq c$; if $x < 0$ then $|x| = -x \leq c$, this last from $-c \leq x$. The same argument, with the two inequalities involving c read strictly, proves the second equivalence.

3. There are four cases, and each uses the sign rules of proposition 1.1.2(4) together with the observation that $|z| = -z$ holds for every $z \leq 0$, the case $z = 0$ because both sides vanish. If $x \geq 0$ and $y \geq 0$ then $xy \geq 0$, so $|xy| = xy = |x||y|$. If $x \geq 0$ and $y < 0$ then $xy \leq 0$, so $|xy| = -(xy) = x(-y) = |x||y|$. The case $x < 0$, $y \geq 0$ is the same with the roles of x and y exchanged. If $x < 0$ and $y < 0$ then $xy = (-x)(-y) > 0$, so $|xy| = xy = (-x)(-y) = |x||y|$.
4. By (1), $x \leq |x|$ and $y \leq |y|$, so $x + y \leq |x| + |y|$. Likewise $-x \leq |x|$ and $-y \leq |y|$, so $-(x + y) \leq |x| + |y|$, that is $-(|x| + |y|) \leq x + y$. The number $c = |x| + |y|$ is nonnegative by (1), so (2) applies and gives $|x + y| \leq |x| + |y|$.
5. Writing $x = (x - y) + y$ and applying (4),

$$|x| \leq |x - y| + |y|, \quad \text{so} \quad |x| - |y| \leq |x - y|$$

Exchanging x and y gives $|y| - |x| \leq |y - x| = |x - y|$, the last equality by (1) applied to $x - y$. So the number $|x| - |y|$ lies between $-|x - y|$ and $|x - y|$, and $|x - y| \geq 0$ by (1), so (2) gives $||x| - |y|| \leq |x - y|$, as we sought to show.

Part (2) is the form in which the absolute value is used: an inequality $|x - a| < \epsilon$ is a pair of inequalities placing x between $a - \epsilon$ and $a + \epsilon$, so a statement about distance converts into a statement about order and back at will. Part (5) is the version that estimates a difference of sizes by the size of a difference, and it is what later chapters use to show that passing to the absolute value cannot destroy an estimate already obtained.

Example 1.3.9: Solution Sets of Two Absolute Value Inequalities

By proposition 1.3.8(2) with $c = 1/2$, applied to the number $x - 2$, the condition $|x - 2| < 1/2$ holds if and only if $-1/2 < x - 2 < 1/2$, that is if and only if $3/2 < x < 5/2$. So

$$\{x \in \mathbb{R} \mid |x - 2| < 1/2\} = (3/2, 5/2)$$

an interval centred at 2 of total length 1. Likewise $|2x + 1| \leq 3$ holds if and only if $-3 \leq 2x + 1 \leq 3$, which after subtracting 1 and multiplying by the positive number $1/2$ reads $-2 \leq x \leq 1$, so the solution set is $[-2, 1]$. Each computation is the same move: a single condition on a distance becomes two conditions on the order, and the order conditions are solved by the field operations.

Combining the two halves of the section gives the quantity that measures how far a function moves on a set. A bounded function has a supremum and an infimum there, and their difference bounds every difference of two of its values. The content of the next result is that this bound is the best one available, so the single number $\sup_D f - \inf_D f$ can replace the whole family of differences $|f(x) - f(y)|$ without loss.

Proposition 1.3.10: The Oscillation Bound

Let D be a nonempty set and let $f: D \rightarrow \mathbb{R}$ be bounded. Write $M = \sup_D f$ and $m = \inf_D f$. Then

1. $|f(x) - f(y)| \leq M - m$ for all $x, y \in D$;
2. $M - m = \sup\{|f(x) - f(y)| \mid x, y \in D\}$, so $M - m$ is the smallest constant C for which $|f(x) - f(y)| \leq C$ holds for all $x, y \in D$.

The number $M - m$ is called the *oscillation* of f on D .

Proof:

1. Let $x, y \in D$. Then $f(x) \leq M$ and $m \leq f(y)$, so $f(x) - f(y) \leq M - m$. Exchanging x and y gives $f(y) - f(x) \leq M - m$, that is $-(M - m) \leq f(x) - f(y)$. Since D is nonempty there is at least one $z \in D$, and $m \leq f(z) \leq M$ gives $M - m \geq 0$, so proposition 1.3.8(2) applies with $c = M - m$ and yields $|f(x) - f(y)| \leq M - m$.
2. Let S denote the set $\{|f(x) - f(y)| \mid x, y \in D\}$, which is nonempty because D is, and which is bounded above by $M - m$ by part (1). By theorem 1.2.4 the supremum $\sup S$ exists, and $\sup S \leq M - m$ because $M - m$ is an upper bound. Suppose $\sup S < M - m$, and put $\epsilon = (M - m - \sup S)/2 > 0$. By proposition 1.2.6 applied to the set $f(D)$, there is an element of $f(D)$ exceeding $M - \epsilon$, that is a point $x \in D$ with $f(x) > M - \epsilon$. Likewise, by lemma 1.3.1 applied to $f(D)$, there is a point $y \in D$ with $f(y) < m + \epsilon$. Subtracting,

$$f(x) - f(y) > (M - \epsilon) - (m + \epsilon) = M - m - 2\epsilon = \sup S$$

and $|f(x) - f(y)| \geq f(x) - f(y)$ by proposition 1.3.8(1). So S contains an element strictly larger than $\sup S$, which contradicts that $\sup S$ is an upper bound of S . Hence $\sup S = M - m$.

For the final clause, a constant C with $|f(x) - f(y)| \leq C$ for all $x, y \in D$ is exactly an upper bound of S , hence satisfies $C \geq \sup S = M - m$, and $C = M - m$ is such a constant by part (1), completing the proof.

The oscillation converts a two-point statement into a one-point one. A bound on $|f(x) - f(y)|$ for all pairs is a statement about $D \times D$, while $M - m$ is computed from two suprema over D , and the proposition says the two carry the same information. This is the form in which the quantity is used later, where a function is controlled on each piece of a subdivided interval by computing one supremum and one infimum per piece.

Example 1.3.11: The Oscillation of x^2 on $[-1, 2]$

Let $D = [-1, 2]$ and $f(x) = x^2$. For $x \in D$ the inequalities $-1 \leq x \leq 2$ give $-2 \leq x \leq 2$ and hence $|x| \leq 2$ by proposition 1.3.8(2). A square is nonnegative (proposition 1.1.2(6), together with $0 \cdot 0 = 0$), so $x^2 = |x^2|$ by definition 1.3.6, and $|x^2| = |x|^2$ by proposition 1.3.8(3). Multiplying $|x| \leq 2$ first by $|x| \geq 0$ and then by $2 > 0$ gives $|x|^2 \leq 2|x| \leq 4$, so $f(x) \leq 4$. Since $f(2) = 4$, the value 4 is attained and $M = \sup_D f = 4$. Every value $f(x) = x^2$ is nonnegative and $f(0) = 0$, so

$m = \inf_D f = 0$. The oscillation is therefore $M - m = 4$, and it is realized by an actual pair of points, since $|f(2) - f(0)| = 4$. On the smaller set $[0, 1]$ the same computation gives oscillation 1: the oscillation depends on the set as well as on the function.

The section closes with the theorem that puts the n -th roots into $\mathbb{R}$. Proposition 1.1.3 shows that no rational squares to 2, and the chapter opened by observing that the set of rationals with square less than 2 has upper bounds in $\mathbb{Q}$ but no least one. In $\mathbb{R}$ that set does have a least upper bound, by theorem 1.2.4, and the theorem below identifies it: it is a number whose square is exactly 2. The argument is written for a general exponent because later chapters need n -th roots of every order, and the hypothesis is $x \geq 0$, weaker than $x > 0$, because the convergence tests that use it are applied to sequences some of whose terms vanish. The case $x = 0$ is immediate and is treated separately in the proof.

Theorem 1.3.12: Existence and Uniqueness of n -th Roots

Let $x \in \mathbb{R}$ with $x \geq 0$ and let $n \geq 1$ be an integer. There is exactly one real $y \geq 0$ with $y^n = x$. It is written $y = x^{1/n}$, and $x^{1/2}$ is also written $\sqrt{x}$.

Proof:

Step 1: powers are strictly increasing on the nonnegative reals. The claim is that $0 \leq a < b$ implies $a^n < b^n$ for every integer $n \geq 1$, and it is proved by induction on n . For $n = 1$ it is the hypothesis. Assume $a^n < b^n$. If $a = 0$ then $a^{n+1} = 0 < b^{n+1}$, since $b > 0$ makes b^{n+1} a product of positive factors. If $a > 0$ then multiplying $a^n < b^n$ by $a > 0$ gives $a^{n+1} < ab^n$, and multiplying $a < b$ by $b^n > 0$ gives $ab^n < b^{n+1}$, so that $a^{n+1} < b^{n+1}$. The same induction with $\leq$ in place of $<$ shows that $0 \leq a \leq b$ implies $a^n \leq b^n$, and this weak form is used for the exponent $n - 1$, where $n = 1$ makes both sides equal to 1.

Step 2: uniqueness. If $y_1, y_2 \geq 0$ both satisfy $y_i^n = x$ and $y_1 \neq y_2$, then one of them is smaller, say $y_1 < y_2$, and Step 1 gives $x = y_1^n < y_2^n = x$, which is impossible. So there is at most one such y , whatever $x \geq 0$ is.

Step 3: the difference bound. For $0 \leq a \leq b$ and $n \geq 1$,

$$b^n - a^n \leq n(b - a)b^{n-1} \quad (1.2)$$

Indeed, expanding the product and cancelling termwise,

$$(b - a) \sum_{k=0}^{n-1} b^{n-1-k} a^k = \sum_{k=0}^{n-1} b^{n-k} a^k - \sum_{k=0}^{n-1} b^{n-1-k} a^{k+1} = b^n - a^n$$

because the second sum is the first with the index shifted by one, leaving only the $k = 0$ term of the first and the $k = n - 1$ term of the second. In each summand $b^{n-1-k} a^k \leq b^{n-1-k} b^k = b^{n-1}$, using $a^k \leq b^k$ from Step 1 for $k \geq 1$ and the equality $a^0 = b^0 = 1$ for $k = 0$, and multiplying by $b^{n-1-k} \geq 0$. There are n summands, so the sum is at most nb^{n-1} , and multiplying by $b - a \geq 0$ gives (1.2).

Step 4: existence when $x = 0$. The number $y = 0$ satisfies $y^n = 0 = x$ and $y \geq 0$, and Step 2 makes it the only such number. Assume from now on that $x > 0$.

Step 5: the candidate. Let

$$E = \{t \in \mathbb{R} \mid t \geq 0 \text{ and } t^n < x\}$$

Put $t_0 = x/(1 + x)$. Then $1 + x > 0$, so $t_0 > 0$. Dividing $x < 1 + x$ by $1 + x$ gives $t_0 < 1$, and dividing $1 < 1 + x$ by $1 + x$ and then multiplying by $x > 0$ gives $t_0 < x$. From $0 < t_0 < x$

and Step 1 one gets $t_0^n \leq t_0$ for every $n \geq 1$, by induction: the case $n = 1$ is an equality, and multiplying $t_0^n \leq t_0$ by $t_0 > 0$ gives $t_0^{n+1} \leq t_0^2 \leq t_0$, the last step multiplying $t_0 < 1$ by $t_0 > 0$. Hence $t_0^n \leq t_0 < x$ and $t_0 \in E$, so E is nonempty. Suppose instead that $t \geq 1 + x$, so that $t > 1$. Then $t^n \geq t$ for every $n \geq 1$, again by induction: the case $n = 1$ is an equality, and multiplying $t^n \geq t$ by $t > 0$ gives $t^{n+1} \geq t^2 \geq t$, the last step multiplying $t \geq 1$ by $t > 0$. So $t^n \geq t \geq 1 + x > x$ and $t \notin E$. Every element of E is therefore smaller than $1 + x$, and E is bounded above. By theorem 1.2.4 the number $y = \sup E$ exists, and $y \geq t_0 > 0$ because $t_0 \in E$.

Step 6: the case $y^n < x$ is impossible. Suppose $y^n < x$ and put

$$h = \min \left\{ \frac{1}{2}, \frac{x - y^n}{2n(y + 1)^{n-1}} \right\}$$

Both entries are positive, since $x - y^n > 0$, $n \geq 1$ and $(y + 1)^{n-1} > 0$, so $0 < h < 1$ and $h < (x - y^n)/(n(y + 1)^{n-1})$. Apply (1.2) with $a = y$ and $b = y + h$, legitimate because $0 \leq y \leq y + h$:

$$(y + h)^n - y^n \leq nh(y + h)^{n-1} \leq nh(y + 1)^{n-1} < x - y^n$$

where the middle inequality uses $y + h < y + 1$ together with the weak form of Step 1 at exponent $n - 1$, and the last uses the choice of h . Hence $(y + h)^n < x$ with $y + h \geq 0$, so $y + h \in E$. But $y + h > y = \sup E$, contradicting that y is an upper bound of E .

Step 7: the case $y^n > x$ is impossible. Suppose $y^n > x$ and put

$$k = \frac{y^n - x}{ny^{n-1}}$$

Then $k > 0$, since $y > 0$ makes $y^{n-1} > 0$ and $y^n - x > 0$. Also $k < y$: this is the inequality $y^n - x < ny^n$, which holds because $y^n - x < y^n \leq ny^n$, using $x > 0$ and $n \geq 1$. Let $t \in E$. If $t > y$ then Step 1 gives $t^n > y^n > x$, contradicting $t \in E$, so $t \leq y$. If moreover $t \geq y - k$, then $0 \leq y - t \leq k$ and (1.2) with $a = t$ and $b = y$ gives

$$y^n - t^n \leq n(y - t)y^{n-1} \leq nky^{n-1} = y^n - x$$

so that $t^n \geq x$, again contradicting $t \in E$. Every $t \in E$ therefore satisfies $t < y - k$, so $y - k$ is an upper bound of E smaller than y , contradicting that y is the *least* upper bound.

Step 8: conclusion. By trichotomy in the ordered field $\mathbb{R}$ (theorem 1.1.10) and Steps 6 and 7, the only remaining possibility is $y^n = x$, with $y > 0$. Uniqueness is Step 2, completing the proof.

No decimal expansion, no approximating sequence and no notion of limit appears in the argument. The number y is produced as a supremum, and Steps 6 and 7 show only that it cannot sit strictly on either side of the target. Where a later existence proof builds a real number directly, with no appeal to numbers already constructed, it has this same shape: describe the set of numbers below the one wanted and take the least upper bound.

Example 1.3.13: $\sqrt{2}$ as a Supremum

Take $x = 2$ and $n = 2$ in theorem 1.3.12. The set constructed in the proof is $E = \{t \geq 0 \mid t^2 < 2\}$, and $\sqrt{2} = \sup E$. Two rational computations locate it. First, $(7/5)^2 = 49/25 < 50/25 = 2$, so $7/5 \in E$ and hence $\sqrt{2} \geq 7/5$. Second, $(3/2)^2 = 9/4 > 8/4 = 2$, so if $t \geq 3/2$ then $t^2 \geq 9/4 > 2$

and $t \notin E$. Thus $3/2$ is an upper bound of E and $\sqrt{2} \leq 3/2$. So

$$\frac{7}{5} \leq \sqrt{2} \leq \frac{3}{2}$$

By proposition 1.1.3 the number $\sqrt{2}$ is not rational, so the supremum of E exists in $\mathbb{R}$ and lies outside the image of $\mathbb{Q}$ under the embedding of proposition 1.1.11. Inside $\mathbb{Q}$ the corresponding set of rationals has no rational least upper bound (example 1.2.3), which is the defect theorem 1.2.4 repairs.

Exercise 1.3.1

1. Let A and B be nonempty subsets of $\mathbb{R}$, both bounded above. Exercise 1.2.1 (4) gives $\sup(A \cup B)$. Show that if $A \cap B$ is nonempty then $\sup(A \cap B)$ is at most the smaller of $\sup A$ and $\sup B$, and give two sets for which this second inequality is strict.
2. Let A and B be nonempty subsets of $\mathbb{R}$ that are bounded below. Deduce from proposition 1.3.2 that $\inf(A + B) = \inf A + \inf B$, by applying the results there to $-A$ and $-B$.
3. Prove the identities

$$\max\{x, y\} = \frac{x + y + |x - y|}{2}, \quad \min\{x, y\} = \frac{x + y - |x - y|}{2}$$

for all $x, y \in \mathbb{R}$, and deduce that $\max\{x, y\} + \min\{x, y\} = x + y$.

4. Let D be a nonempty set and let $f, g: D \rightarrow \mathbb{R}$ be bounded. Show that $|f - g|$ is bounded and that

$$\left| \sup_D f - \sup_D g \right| \leq \sup_D |f - g|$$

Show also that the corresponding statement with $\sup_D$ replaced by $\inf_D$ on the left holds as well.

5. Let D be a nonempty set, let $f: D \rightarrow \mathbb{R}$ be bounded, and let $E \subseteq D$ be nonempty. Show that the oscillation of f on E is at most the oscillation of f on D . Show also that the oscillation of f on D is 0 if and only if f is constant on D .
6. Let $x, y \geq 0$ and let $n \geq 1$ be an integer. Using the uniqueness clause of theorem 1.3.12, prove that $(xy)^{1/n} = x^{1/n}y^{1/n}$. Prove also that $x < y$ implies $x^{1/n} < y^{1/n}$, and conclude that the map $x \mapsto x^{1/n}$ is injective on the nonnegative reals.

1.4 Consequences of Completeness

Theorem 1.2.4 is a statement about sets: a nonempty set of reals with an upper bound has a least one. Later chapters use it through five statements about numbers and intervals, and this section proves those five. The natural numbers are not bounded above in $\mathbb{R}$; every real number has an integer part; $\mathbb{Q}$ and its complement both meet every interval; an inequality that holds with arbitrary slack holds with none; and a nested sequence of closed bounded intervals has a point lying in all of them. The last of these then gives a fact about $\mathbb{R}$ of a different kind from anything in sections 1.1 to 1.3: the real numbers cannot be arranged in a list indexed by $\mathbb{N}$.

Throughout, $\mathbb{Q}$ is identified with its image in $\mathbb{R}$ under the embedding of proposition 1.1.11, so that $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}$ and a symbol such as n or m/n names a real number as well as a rational one. Induction and the well-ordering of $\mathbb{N}$ are used freely.

The first question to settle is whether $\mathbb{N}$, sitting inside $\mathbb{R}$ by that identification, exceeds every real number. Nothing in definition 1.1.9 makes this visible: a real number is a class of Cauchy sequences of rationals, and the definition says nothing about how far the integers extend among those classes. What hangs on the answer is whether $1/n$ can be brought below a prescribed positive tolerance by choosing n large, which is the shape of every estimate in the chapters that follow. The least upper bound property settles it.

Theorem 1.4.1: The Archimedean Property of $\mathbb{R}$

1. For all $x, y \in \mathbb{R}$ with $x > 0$ there is an $n \in \mathbb{N}$ with $nx > y$.
2. For every $y \in \mathbb{R}$ there is an $n \in \mathbb{N}$ with $n > y$.
3. For every $\epsilon > 0$ there is an $n \in \mathbb{N}$ with $0 < 1/n < \epsilon$.

Proof:

1. Let $A = \{nx \mid n \in \mathbb{N}\}$, and suppose the conclusion fails, so that $nx \leq y$ for every $n \in \mathbb{N}$. Then A is nonempty, since $1 \cdot x = x$ lies in A , and y is an upper bound for A . By theorem 1.2.4, A has a least upper bound $\alpha = \sup A$. Since $x > 0$, the real number $\alpha - x$ is smaller than α . As α is the *least* upper bound of A , the smaller number $\alpha - x$ is not an upper bound of A , so some element of A exceeds it: there is an $m \in \mathbb{N}$ with $mx > \alpha - x$. Adding x to both sides gives $(m+1)x > \alpha$. But $m+1 \in \mathbb{N}$, so $(m+1)x \in A$, and an element of A exceeds the upper bound α . This is impossible, so some $n \in \mathbb{N}$ satisfies $nx > y$.
2. Apply (1) with $x = 1$, which is positive: there is an $n \in \mathbb{N}$ with $n \cdot 1 = n > y$.
3. Apply (1) with ϵ in the role of x and 1 in the role of y : there is an $n \in \mathbb{N}$ with $n\epsilon > 1$. Now $n \geq 1 > 0$, so $1/n > 0$, and multiplying $n\epsilon > 1$ by the positive number $1/n$ preserves the inequality (proposition 1.1.2(5)) and gives $\epsilon > 1/n$, as we sought to show.

Form (3) is the one an ϵ -estimate uses: to make a quantity smaller than a prescribed ϵ , it is enough to make it smaller than $1/n$ for a single n chosen after ϵ . Form (2) says that $\mathbb{N}$ has no upper bound in $\mathbb{R}$, which is the same statement read as a fact about the set $\mathbb{N}$, where form (1) reads it as a fact about individual numbers.

The Archimedean property is not itself a form of completeness. Section 1.1 takes as one of its standing hypotheses that $\mathbb{Q}$ is Archimedean, while example 1.2.3 exhibits a set of rationals with no rational supremum, so the Archimedean property holds in a field that fails the least upper bound property. What the proof establishes is that completeness already contains it, so nothing further has to be assumed about $\mathbb{R}$.

Example 1.4.2: Suprema and Infima Governed by $\mathbb{N}$

1. $\mathbb{N}$ has no supremum in $\mathbb{R}$. By theorem 1.4.1(2), for every $y \in \mathbb{R}$ there is an $n \in \mathbb{N}$ with $n > y$, so no real number is an upper bound of $\mathbb{N}$. A set with no upper bound has no least upper bound, so $\sup \mathbb{N}$ does not exist. This is consistent with theorem 1.2.4, whose hypothesis is that the set be bounded above.
2. Let $E = \{1/n \mid n \in \mathbb{N}\}$. Then $\sup E = 1$ and $\inf E = 0$, and $0 \notin E$. For the supremum: every $n \in \mathbb{N}$ satisfies $n \geq 1 > 0$, so $1/n \leq 1$, and $1 = 1/1 \in E$, so 1 is an upper bound belonging to E and is therefore the least one. For the infimum: E is bounded below by 0, since $1/n > 0$ for every n , so $\inf E$ exists by corollary 1.2.5. If $t > 0$, then theorem 1.4.1(3) gives an $n \in \mathbb{N}$ with $1/n < t$, so t is not a lower bound of E . Hence 0 is the greatest lower bound. Finally $0 \notin E$ because every element of E is positive.

The second item shows an infimum that does not belong to the set, and theorem 1.4.1(3) is what makes it one: the elements of E come arbitrarily close to 0 without reaching it. An infimum is therefore not a smallest element, and definition 1.2.2 does not ask it to be.

The Archimedean property says that the integers eventually pass any real number. The sharper statement is that there is a *last* integer that does not pass it. That statement attaches a specific integer to a given real number, where theorem 1.4.1 only asserts that some large integer exists.

Proposition 1.4.3: The Integer Part of a Real Number

For every $x \in \mathbb{R}$ there is exactly one $m \in \mathbb{Z}$ with

$$m \leq x < m + 1$$

This integer is written $\lfloor x \rfloor$ and is called the *integer part* of x .

Proof:

Step 1: Existence. By theorem 1.4.1(2) there is an $N \in \mathbb{N}$ with $N > |x|$. From proposition 1.3.8(1), both $x \leq |x|$ and $-x \leq |x|$, so $-N < x < N$.

Let

$$T = \{k \in \mathbb{N} \mid -N + k > x\}$$

This set is nonempty: $2N \in \mathbb{N}$ and $-N + 2N = N > x$, so $2N \in T$. By the well-ordering of $\mathbb{N}$, T has a least element k_0 . Put $m = -N + k_0 - 1$, which lies in $\mathbb{Z}$ because N and k_0 do.

Then $m + 1 = -N + k_0 > x$, since $k_0 \in T$. For the other inequality there are two cases. If $k_0 = 1$ then $m = -N$, and $-N < x$ was established above, so $m \leq x$. If $k_0 > 1$ then $k_0 - 1 \in \mathbb{N}$ and $k_0 - 1 < k_0$, so $k_0 - 1 \notin T$ by minimality of k_0 . Unwinding the definition of T , this reads $-N + (k_0 - 1) \leq x$, which is $m \leq x$. In both cases $m \leq x < m + 1$.

Step 2: Uniqueness. Suppose $m, m' \in \mathbb{Z}$ both satisfy the two inequalities. From $m \leq x$ and $x < m' + 1$ we get $m < m' + 1$, that is $m - m' < 1$. Exchanging the roles of m and m' gives $m' - m < 1$. So $m - m'$ is an integer with $-1 < m - m' < 1$.

No integer k satisfies $0 < k < 1$: such a k would be a positive integer, hence an element of $\mathbb{N}$, and 1 is the least element of $\mathbb{N}$, so $k \geq 1$. If an integer k satisfied $-1 < k < 0$, then $-k$ would be an integer with $0 < -k < 1$, which the previous sentence excludes. Hence $m - m' = 0$, so $m = m'$, completing the proof.

Writing $r = x - \lfloor x \rfloor$, the proposition splits every real number as $x = \lfloor x \rfloor + r$ with $\lfloor x \rfloor \in \mathbb{Z}$ and $0 \leq r < 1$, and the splitting is unique. For instance $\lfloor 3/2 \rfloor = 1$, since $1 \leq 3/2 < 2$, and $\lfloor 2 \rfloor = 2$. The negative case is the one to watch: $\lfloor -3/2 \rfloor = -2$, since $-2 \leq -3/2 < -1$. The integer part of x is the largest integer at or below x , not the integer nearest 0.

The integer part is what makes the next theorem a construction, with the rational produced explicitly. The question it answers is whether the rationals, which proposition 1.1.11 placed inside $\mathbb{R}$, are spread through $\mathbb{R}$ or clustered somewhere. The recipe is to multiply everything by an n large enough that nx and ny are more than 1 apart, and then to take the first integer above nx : it cannot have jumped past ny .

Theorem 1.4.4: $\mathbb{Q}$ Is Dense in $\mathbb{R}$

If $x, y \in \mathbb{R}$ and $x < y$, then there is a $p \in \mathbb{Q}$ with $x < p < y$.

Proof:

Since $x < y$, the number $y - x$ is positive, so theorem 1.4.1(1), applied with $y - x$ in the role of x and with 1 in the role of y , gives an $n \in \mathbb{N}$ with

$$n(y - x) > 1$$

Put $m = \lfloor nx \rfloor + 1$, an integer by proposition 1.4.3. That proposition gives $\lfloor nx \rfloor \leq nx < \lfloor nx \rfloor + 1$, and the two halves read

$$nx < m \quad \text{and} \quad m \leq nx + 1$$

Combining the second with $1 < ny - nx$ gives

$$m \leq nx + 1 < nx + (ny - nx) = ny$$

so $nx < m < ny$. Since $n \geq 1 > 0$, the number $1/n$ is positive, and multiplying through by it preserves both inequalities (proposition 1.1.2(5)), giving $x < m/n < y$. The number $p = m/n$ is rational, as we sought to show.

The same interval has still to be shown to contain a number outside $\mathbb{Q}$, and nothing new is needed to produce one. Translation by a fixed irrational number preserves the order and carries rationals to irrationals, so the theorem just proved, applied to the translated interval, delivers the point.

Corollary 1.4.5: The Irrationals Are Dense in $\mathbb{R}$

If $x, y \in \mathbb{R}$ and $x < y$, then there is a $t \in \mathbb{R} \setminus \mathbb{Q}$ with $x < t < y$.

Proof:

By theorem 1.3.12, applied with $n = 2$ and with 2 in the role of x , there is a unique real $s \geq 0$ with $s^2 = 2$. Write $s = \sqrt{2}$. This number is not rational. Suppose s were the image of some $p \in \mathbb{Q}$ under the embedding of proposition 1.1.11. That embedding is a field homomorphism, so the image of p^2 is $s^2 = 2$, which is also the image of the rational 2. The embedding is injective, so $p^2 = 2$ holds in $\mathbb{Q}$, contradicting proposition 1.1.3.

Adding $-s$ to both sides of $x < y$ gives $x - s < y - s$, so theorem 1.4.4 produces a $p \in \mathbb{Q}$ with $x - s < p < y - s$. Adding s throughout gives $x < p + s < y$. Put $t = p + s$. If t were rational,

then $s = t - p$ would be a difference of two rationals and hence rational, which has just been excluded. So $t \in \mathbb{R} \setminus \mathbb{Q}$ and $x < t < y$, completing the proof.

Between any two distinct reals there is a rational and also an irrational, so neither $\mathbb{Q}$ nor $\mathbb{R} \setminus \mathbb{Q}$ can be separated from $\mathbb{R}$ by looking at the order alone. Every interval meets both. The two sets are nonetheless of different sizes, and theorem 1.4.11 is where that difference appears.

Example 1.4.6: A Rational Between $\sqrt{2}$ and $\sqrt{3}$

The proof of theorem 1.4.4 is a recipe, and running it on $x = \sqrt{2}$ and $y = \sqrt{3}$ produces an explicit rational. Both roots exist by theorem 1.3.12. Throughout, use that squaring is strictly increasing on the nonnegative reals: if $0 \leq u < v$ then $v^2 - u^2 = (v - u)(v + u) > 0$, since $v - u > 0$ and $v + u \geq v > 0$.

First, bracket the two roots by rationals. Compute $(7/5)^2 = 49/25 < 50/25 = 2$ and $(3/2)^2 = 9/4 > 8/4 = 2$. If $7/5 \geq \sqrt{2}$ held, then squaring would give $(7/5)^2 \geq 2$, so $7/5 < \sqrt{2}$; if $3/2 \leq \sqrt{2}$ held, squaring would give $(3/2)^2 \leq 2$, so $\sqrt{2} < 3/2$. The same two steps applied to $(17/10)^2 = 289/100 < 300/100 = 3$ and $(9/5)^2 = 81/25 > 75/25 = 3$ give $17/10 < \sqrt{3} < 9/5$.

Now follow the proof. Since $\sqrt{3} > 17/10$ and $\sqrt{2} < 3/2 = 15/10$,

$$\sqrt{3} - \sqrt{2} > \frac{17}{10} - \frac{15}{10} = \frac{1}{5}$$

so $n = 5$ satisfies $n(\sqrt{3} - \sqrt{2}) > 1$. Next, $5 \cdot (7/5) = 7 < 5\sqrt{2} < 5 \cdot (3/2) = 15/2 < 8$, so $7 \leq 5\sqrt{2} < 8$ and the uniqueness clause of proposition 1.4.3 gives $\lfloor 5\sqrt{2} \rfloor = 7$. The proof takes $m = 8$. The rational it produces is $p = 8/5$, and the bracketing confirms the conclusion directly: $\sqrt{2} < 3/2 = 15/10 < 16/10 = 8/5$ and $8/5 = 16/10 < 17/10 < \sqrt{3}$.

An argument that compares two real numbers often produces not $a \leq b$ but $a \leq b + \epsilon$, with an ϵ that can be taken as small as desired and never reduced to 0 inside the argument itself. The next proposition removes the margin afterwards, and theorem 1.4.8 below is the first place it is applied.

Proposition 1.4.7: The ϵ -Principle

Let $a, b \in \mathbb{R}$.

1. If $a \leq b + \epsilon$ for every $\epsilon > 0$, then $a \leq b$.
2. If $|a - b| \leq \epsilon$ for every $\epsilon > 0$, then $a = b$.

Proof:

1. Suppose instead that $a > b$. Then $a - b > 0$, so $\epsilon_0 = (a - b)/2$ is positive, and the hypothesis applied to ϵ_0 gives

$$a \leq b + \frac{a - b}{2} = \frac{a + b}{2}$$

On the other hand $b < a$ gives $a + b < 2a$, hence $(a + b)/2 < a$. Together these read $a < a$, which is false. Therefore $a \leq b$.

2. Fix $\epsilon > 0$. If $a - b \geq 0$ then $a - b = |a - b| \leq \epsilon$ by definition 1.3.6; if $a - b < 0$ then

$a - b < 0 < \epsilon$. Either way $a - b \leq \epsilon$, that is $a \leq b + \epsilon$. As $\epsilon > 0$ was arbitrary, part (1) gives $a \leq b$. Since $|b - a| = |a - b|$ (proposition 1.3.8(1)), the same argument with a and b exchanged gives $b \leq a$. Hence $a = b$, completing the proof.

Part (2) is the form used to prove that two numbers produced by different constructions are equal. A single computation showing $a = b$ is usually out of reach, and what one shows is that for each $\epsilon > 0$, a and b differ by less than ϵ , and the proposition converts the family of estimates into the equality.

Theorem 1.2.4 states completeness as a fact about bounded sets. A second way to say that $\mathbb{R}$ has no gaps is geometric: if a sequence of closed intervals is nested, each inside the last, then at least one point lies in every one of them. In $\mathbb{Q}$ this fails. Intervals with rational endpoints can close in on a point whose square is 2, and proposition 1.1.3 says no rational is available to be that point, so the intersection of the rational intervals is empty even though each of them is nonempty. Example 1.4.10 carries this out explicitly. Over $\mathbb{R}$ the conclusion holds, and completeness is what delivers it.

Theorem 1.4.8: The Nested Interval Theorem

For each $n \in \mathbb{N}$ let $a_n \leq b_n$ be real numbers, and put

$$I_n = \{x \in \mathbb{R} \mid a_n \leq x \leq b_n\}$$

Assume $I_{n+1} \subseteq I_n$ for every $n \in \mathbb{N}$. Then:

1. $\bigcap_{n \in \mathbb{N}} I_n$ is nonempty.
2. If, in addition, for every $\epsilon > 0$ there is an $n \in \mathbb{N}$ with $b_n - a_n < \epsilon$, then $\bigcap_{n \in \mathbb{N}} I_n$ has exactly one element.

Proof:

1. First, the endpoints move monotonically. Fix n . Since $a_{n+1} \leq b_{n+1}$, the number a_{n+1} lies in I_{n+1} , hence in I_n , so $a_n \leq a_{n+1}$. The same argument applied to $b_{n+1} \in I_{n+1} \subseteq I_n$ gives $b_{n+1} \leq b_n$. By induction, $a_m \leq a_k$ and $b_k \leq b_m$ whenever $m \leq k$. Next, $a_m \leq b_n$ for all $m, n \in \mathbb{N}$, not only for $m = n$. Let k be the larger of m and n . Then $m \leq k$ and $n \leq k$, so the previous paragraph gives $a_m \leq a_k$ and $b_k \leq b_n$, and $a_k \leq b_k$ holds by hypothesis. Chaining the three, $a_m \leq a_k \leq b_k \leq b_n$. Let $A = \{a_n \mid n \in \mathbb{N}\}$. It is nonempty and, taking $n = 1$ above, bounded above by b_1 , so theorem 1.2.4 gives $\alpha = \sup A$. Fix $n \in \mathbb{N}$. Then $a_n \leq \alpha$ because α is an upper bound of A , and $\alpha \leq b_n$ because b_n is an upper bound of A by the previous paragraph while α is the least such. So $a_n \leq \alpha \leq b_n$, that is $\alpha \in I_n$. As n was arbitrary, $\alpha \in \bigcap_{n \in \mathbb{N}} I_n$.

2. By part (1) the intersection contains at least one point, so it is enough to show it contains at most one. Let x and y both lie in $\bigcap_{n \in \mathbb{N}} I_n$ and fix $n \in \mathbb{N}$. From $a_n \leq x \leq b_n$ and $a_n \leq y \leq b_n$ it follows that $x - y \leq b_n - a_n$ and $y - x \leq b_n - a_n$, and since $|x - y|$ is one of $x - y$ and $y - x$ (definition 1.3.6),

$$|x - y| \leq b_n - a_n$$

Now let $\epsilon > 0$. By hypothesis there is an n with $b_n - a_n < \epsilon$, and $|x - y| \leq b_n - a_n$ then gives $|x - y| < \epsilon$. Since $\epsilon > 0$ was arbitrary, proposition 1.4.7(2) gives $x = y$. The intersection therefore has exactly one element, completing the proof.

Both hypotheses on the intervals are needed, and theorem 1.4.1 is what produces the two failures.

Example 1.4.9: Two Failures of Nesting

1. *Closedness cannot be dropped.* Let $J_n = \{x \in \mathbb{R} \mid 0 < x < 1/n\}$. These are nested, since $1/(n+1) \leq 1/n$, and each is nonempty, since theorem 1.4.4 gives a point strictly between 0 and $1/n$. But $\bigcap_{n \in \mathbb{N}} J_n = \emptyset$: a point x of the intersection would satisfy $x > 0$ and $x < 1/n$ for every n , contradicting theorem 1.4.1(3), which produces an n with $1/n < x$.
2. *Boundedness cannot be dropped.* Let $K_n = \{x \in \mathbb{R} \mid x \geq n\}$. These are nested and nonempty, since $n \in K_n$. But $\bigcap_{n \in \mathbb{N}} K_n = \emptyset$: a point x of the intersection would satisfy $x \geq n$ for every $n \in \mathbb{N}$, contradicting theorem 1.4.1(2), which produces an n with $n > x$.

Example 1.4.10: Nested Rational Intervals with No Rational Point

Write $I_n = \{x \in \mathbb{R} \mid a_n \leq x \leq b_n\}$ throughout, and set $a_1 = 1$, $b_1 = 2$, so that $a_1, b_1 \in \mathbb{Q}$, $a_1 < b_1$, and $a_1^2 = 1 < 2 < 4 = b_1^2$. Given rationals $a_n < b_n$ with $a_n^2 < 2 < b_n^2$, let $c_n = (a_n + b_n)/2$, which is again rational and satisfies $a_n < c_n < b_n$, since $c_n - a_n = b_n - c_n = (b_n - a_n)/2 > 0$. By proposition 1.1.3 the rational c_n has $c_n^2 \neq 2$, so exactly one of $c_n^2 < 2$ and $c_n^2 > 2$ holds. Put

$$(a_{n+1}, b_{n+1}) = \begin{cases} (c_n, b_n) & \text{if } c_n^2 < 2 \\ (a_n, c_n) & \text{if } c_n^2 > 2 \end{cases}$$

In both cases a_{n+1} and b_{n+1} are rational, $a_{n+1} < b_{n+1}$, $a_{n+1}^2 < 2 < b_{n+1}^2$, the inclusion $I_{n+1} \subseteq I_n$ holds because $a_n \leq a_{n+1}$ and $b_{n+1} \leq b_n$, and $b_{n+1} - a_{n+1} = (b_n - a_n)/2$. By induction $b_n - a_n = 1/2^{n-1}$.

These lengths become arbitrarily small. For every integer $k \geq 0$ one has $2^k \geq k + 1$, by induction: $2^0 = 1$, and if $2^k \geq k + 1$ then $2^{k+1} = 2 \cdot 2^k \geq 2k + 2 \geq k + 2$. Given $\epsilon > 0$, take $n \in \mathbb{N}$ with $1/n < \epsilon$ (theorem 1.4.1(3)). Then $2^{n-1} \geq n > 0$, so $1/2^{n-1} \leq 1/n < \epsilon$. Hence theorem 1.4.8(2) applies, and $\bigcap_{n \in \mathbb{N}} I_n$ consists of a single real number.

That number is $\sqrt{2}$. Let $s \geq 0$ be the real with $s^2 = 2$ given by theorem 1.3.12. Every a_n is positive, since $a_1 = 1$ and each a_{n+1} is either a_n or the larger number c_n , and $a_n^2 < 2 = s^2$. Were $a_n \geq s$, squaring would give $a_n^2 \geq s^2$, because $u \geq v \geq 0$ implies $u^2 - v^2 = (u - v)(u + v) \geq 0$. So $a_n < s$, and the same argument applied to $b_n > 0$ with $b_n^2 > s^2$ gives $s < b_n$. Thus $s \in I_n$ for every n , and by uniqueness $\bigcap_{n \in \mathbb{N}} I_n = \{s\}$.

Now read this inside $\mathbb{Q}$. The sets $I_n \cap \mathbb{Q}$ are nested, and each is nonempty, containing a_n . Their intersection is $(\bigcap_{n \in \mathbb{N}} I_n) \cap \mathbb{Q} = \{s\} \cap \mathbb{Q}$, which is empty: if s were a rational p , then $p^2 = 2$ in $\mathbb{Q}$, contradicting proposition 1.1.3. A nested sequence of nonempty closed intervals with rational endpoints therefore has empty intersection in $\mathbb{Q}$, which is the failure theorem 1.4.8 repairs.

Theorem 1.4.4 and corollary 1.4.5 left $\mathbb{Q}$ and $\mathbb{R} \setminus \mathbb{Q}$ indistinguishable by order: each meets every interval. They are distinguished by counting. Recall from *Preliminaries* [4] that a nonempty set is countable exactly when it is the range of some function defined on $\mathbb{N}$, and that $\mathbb{Q}$ is countable. The next theorem shows that $\mathbb{R}$ is not, so the set of classes built in definition 1.1.9 admits no listing while the set of rationals it was built from does. The proof produces a real number specified by nothing except the conditions it must avoid, and theorem 1.4.8 is what converts that list of conditions into a point.

Theorem 1.4.11: Uncountability of $\mathbb{R}$

Let $a, b \in \mathbb{R}$ with $a < b$, and write $[a, b] = \{x \in \mathbb{R} \mid a \leq x \leq b\}$. For every function $f: \mathbb{N} \rightarrow \mathbb{R}$ there is a point $x \in [a, b]$ with $x \neq f(n)$ for every $n \in \mathbb{N}$. Consequently the range of f does not contain $[a, b]$, and neither $\mathbb{R}$ nor $[a, b]$ is countable.

Proof:

Step 1: One splitting step. Let $u < v$ be reals and let $z \in \mathbb{R}$. Put

$$c = u + \frac{v - u}{3}, \quad d = u + \frac{2(v - u)}{3}$$

Then $u < c < d < v$, since each of $c - u$, $d - c$ and $v - d$ equals $(v - u)/3$, which is positive. The intervals $[u, c]$ and $[d, v]$ are therefore contained in $[u, v]$, each has distinct endpoints, and they are disjoint: a point of the first is at most c and a point of the second is at least d , while $c < d$. Consequently z fails to lie in at least one of them.

Define $\Phi([u, v], z)$ to be $[u, c]$ when $z \notin [u, c]$, and $[d, v]$ otherwise. In the second case $z \in [u, c]$, so $z \leq c < d$ and hence $z \notin [d, v]$. In both cases $\Phi([u, v], z)$ is an interval $[u', v']$ with $u' < v'$, contained in $[u, v]$, and not containing z .

Step 2: The recursion. Define $I_1 = \Phi([a, b], f(1))$ and $I_{n+1} = \Phi(I_n, f(n+1))$ for $n \in \mathbb{N}$. The rule Φ assigns a definite interval to each pair, so this defines the sequence (I_n) by recursion, with no choices made along the way. By induction on n , each I_n is an interval $[a_n, b_n]$ with $a_n < b_n$, satisfying $I_{n+1} \subseteq I_n$ and $I_1 \subseteq [a, b]$, and $f(n) \notin I_n$ by the last clause of Step 1.

Step 3: The point. The intervals I_n are closed, bounded and nested, so theorem 1.4.8(1) gives a point $x \in \bigcap_{n \in \mathbb{N}} I_n$. Fix $n \in \mathbb{N}$. Then $x \in I_n$ while $f(n) \notin I_n$, so $x \neq f(n)$. Moreover $x \in I_1 \subseteq [a, b]$, which establishes the first assertion.

Step 4: The consequences. If the range of f contained $[a, b]$, then the point x of Step 3 would be $f(n)$ for some n , which Step 3 excludes. So no function on $\mathbb{N}$ has $[a, b]$ inside its range. Both countability claims follow from this. Were $\mathbb{R}$ countable, some $h: \mathbb{N} \rightarrow \mathbb{R}$ would have range $\mathbb{R}$, and $\mathbb{R}$ contains $[a, b]$, so the range of h would contain $[a, b]$. Were $[a, b]$ countable, some $g: \mathbb{N} \rightarrow [a, b]$ would have range $[a, b]$, and g is in particular a function from $\mathbb{N}$ to $\mathbb{R}$ whose range contains $[a, b]$. Both are impossible, as we sought to show.

The theorem is stronger than the bare statement that $\mathbb{R}$ is uncountable: no list of real numbers can exhaust even a single interval, however short. Since $\mathbb{Q}$ is countable, this also settles that $\mathbb{R} \setminus \mathbb{Q}$ is not empty, which corollary 1.4.5 obtained by a different route, and, since a list of $\mathbb{Q}$ together with a list of $\mathbb{R} \setminus \mathbb{Q}$ could be interleaved into a single list of $\mathbb{R}$, that $\mathbb{R} \setminus \mathbb{Q}$ admits no list either. That interleaving is exercise 1.4.1 (6).¹

Exercise 1.4.1

1. Let $x \in \mathbb{R}$ and put $A = \{q \in \mathbb{Q} \mid q < x\}$. Show that A is nonempty and bounded above, and that $\sup A = x$.
2. Show that $\lfloor x + m \rfloor = \lfloor x \rfloor + m$ for every $x \in \mathbb{R}$ and every $m \in \mathbb{Z}$. Then exhibit real numbers x

¹The proof above is Cantor's first, published in 1874. He gave a second proof in 1891, the diagonal argument, which *Preliminaries* [4] develops in the form " $\{0, 1\}^{\mathbb{N}}$ is uncountable". The 1874 argument is the one given here because it needs no decimal representation, which this chapter has not constructed, and rests only on theorem 1.4.8.

and y for which $\lfloor x + y \rfloor \neq \lfloor x \rfloor + \lfloor y \rfloor$.

3. Let $x < y$ be real numbers and let $k \in \mathbb{N}$. Show that there are rationals $p_1 < p_2 < \cdots < p_k$ with $x < p_1$ and $p_k < y$.
4. Let A and B be nonempty subsets of $\mathbb{R}$ such that $a \leq b$ for every $a \in A$ and every $b \in B$.
 1. Show that $\sup A$ and $\inf B$ exist and that $\sup A \leq \inf B$.
 2. Show that $\sup A = \inf B$ if and only if for every $\epsilon > 0$ there are $a \in A$ and $b \in B$ with $b - a < \epsilon$.
5. Let $I_n = \{x \in \mathbb{R} \mid a_n \leq x \leq b_n\}$ be as in theorem 1.4.8, with $a_n \leq b_n$ and $I_{n+1} \subseteq I_n$ for every $n \in \mathbb{N}$. Show that $\alpha = \sup\{a_n \mid n \in \mathbb{N}\}$ and $\beta = \inf\{b_n \mid n \in \mathbb{N}\}$ both exist, that $\alpha \leq \beta$, and that

$$\bigcap_{n \in \mathbb{N}} I_n = \{x \in \mathbb{R} \mid \alpha \leq x \leq \beta\}$$

6. Show that $\mathbb{R} \setminus \mathbb{Q}$ is uncountable. (Use that $\mathbb{Q}$ is countable, and build from a listing of $\mathbb{Q}$ and a hypothetical listing of $\mathbb{R} \setminus \mathbb{Q}$ a single function on $\mathbb{N}$ whose range is all of $\mathbb{R}$.)
7. Call a real number *dyadic* if it has the form $m/2^n$ with $m \in \mathbb{Z}$ and $n \in \mathbb{N}$. Show that the dyadic numbers are dense in $\mathbb{R}$: if $x < y$ then there is a dyadic d with $x < d < y$. (Follow the proof of theorem 1.4.4, using theorem 1.4.1(1) together with the inequality $2^k \geq k + 1$ established in example 1.4.10 to choose the denominator.)
8. Let $E \subseteq \mathbb{R}$ be nonempty and bounded above, and let $s \in \mathbb{R}$. Show that $s = \sup E$ if and only if s is an upper bound of E and, for every $n \in \mathbb{N}$, there is an $x \in E$ with $x > s - 1/n$.

1.5★ Uniqueness of the Complete Ordered Field

★ *Optional. This section is an aside; nothing later in the book depends on it.*

Section 1.1 built one complete ordered field, and appendix A builds another by a different route. A reader is entitled to ask whether the two constructions produce the same object, and more generally whether the axioms pin $\mathbb{R}$ down at all or merely describe a family of fields among which the construction made an arbitrary choice. This section answers both: they do pin it down, so the choice was forced.

Call an ordered field *complete* if it satisfies the least upper bound property of theorem 1.2.4. One consequence comes first, because the uniqueness proof uses it and because it settles what the Archimedean property costs.

Corollary 1.5.1: Every Complete Ordered Field Is Archimedean

Let F be a complete ordered field. Then for every $x \in F$ there is a positive integer n with $x < n \cdot 1$.

Proof:

Write N for the set of elements $n \cdot 1$ with n a positive integer, where $n \cdot 1$ means $1 + \cdots + 1$ with n summands. Suppose some $x \in F$ satisfies $n \cdot 1 \leq x$ for every n . Then N is nonempty and bounded above, so $s = \sup N$ exists in F by completeness. Since $s - 1 < s$ and s is the least upper bound, $s - 1$ is not an upper bound of N , so there is a positive integer m with $s - 1 < m \cdot 1$. Adding 1 gives $s < (m + 1) \cdot 1$, and $(m + 1) \cdot 1 \in N$, contradicting that s is an upper bound of N . So no such x exists, completing the proof.

For the field constructed in section 1.1 the Archimedean property was available before completeness, from proposition 1.1.12. Corollary 1.5.1 shows that it did not have to be assumed, since any complete ordered field has it. The two routes agree, and the second is the one available to a reader who meets $\mathbb{R}$ through its axioms, with no construction in hand.

An isomorphism of ordered fields is a bijection preserving addition, multiplication and the order. The theorem says there is exactly one between any two complete ordered fields, so that the phrase “the real numbers” is justified.

Theorem 1.5.2: Uniqueness of the Complete Ordered Field

Let F and F' be complete ordered fields. Then there is exactly one isomorphism of ordered fields $\varphi: F \rightarrow F'$.

Proof:

Step 1: the rational subfield is forced. Any isomorphism sends 0 to 0 and 1 to 1, hence $n \cdot 1$ to $n \cdot 1$ for every positive integer n , hence $-(n \cdot 1)$ to $-(n \cdot 1)$, and hence $(m \cdot 1)(n \cdot 1)^{-1}$ to $(m \cdot 1)(n \cdot 1)^{-1}$ for integers m and $n \neq 0$. Write $\mathbb{Q}_F$ for the set of such elements of F , and $\mathbb{Q}_{F'}$ likewise. These are subfields, and the assignment just described is a bijection $\varphi_0: \mathbb{Q}_F \rightarrow \mathbb{Q}_{F'}$ preserving both operations and the order, since $(m \cdot 1)(n \cdot 1)^{-1} < (m' \cdot 1)(n' \cdot 1)^{-1}$ in either field reduces, after clearing denominators, to the same inequality between integers. So φ is determined on $\mathbb{Q}_F$ before anything is chosen.

Step 2: $\mathbb{Q}_F$ is dense in F . Let $x < y$ in F . By corollary 1.5.1 there is a positive integer n with $(y - x)^{-1} < n \cdot 1$, so $(n \cdot 1)^{-1} < y - x$ by proposition 1.1.2(7). Applying the corollary again gives a positive integer m_1 with $x \cdot n \cdot 1 < m_1 \cdot 1$ and a positive integer m_2 with $-(x \cdot n \cdot 1) < m_2 \cdot 1$, so that $-m_2 \cdot 1 < x \cdot n \cdot 1 < m_1 \cdot 1$. Among the finitely many integers k with $-m_2 \leq k \leq m_1$, those satisfying $x \cdot n \cdot 1 < k \cdot 1$ form a nonempty set, so there is a least such k_0 , and it satisfies $(k_0 - 1) \cdot 1 \leq x \cdot n \cdot 1 < k_0 \cdot 1$. *(The integers must be bracketed on both sides first: taking k_0 least among the positive integers alone fails for $x < 0$, where $x \cdot n \cdot 1 < 0 < 1$ forces $k_0 = 1$.)* So

$$x < (k_0 \cdot 1)(n \cdot 1)^{-1} \leq x + (n \cdot 1)^{-1} < y$$

and the middle element lies in $\mathbb{Q}_F$.

Step 3: the extension. For $x \in F$ put

$$\varphi(x) = \sup\{\varphi_0(r) \mid r \in \mathbb{Q}_F \text{ and } r < x\}$$

The set on the right is nonempty and bounded above: nonempty by Step 2 applied to any element below x , and bounded above by $\varphi_0(r')$ for any $r' \in \mathbb{Q}_F$ with $x < r'$, again by Step 2.

So the supremum exists by completeness of F' . For $x \in \mathbb{Q}_F$ the value agrees with $\varphi_0(x)$, since φ_0 preserves the order and $\mathbb{Q}_F$ is dense, so no rational strictly below x has image above $\varphi_0(x)$ while every ϵ -band below $\varphi_0(x)$ is met.

Step 4: φ preserves the order and is bijective. If $x < y$, Step 2 gives $r, r' \in \mathbb{Q}_F$ with $x < r < r' < y$. Every rational below x is below r , so $\varphi(x) \leq \varphi_0(r) < \varphi_0(r') \leq \varphi(y)$, whence $\varphi(x) < \varphi(y)$. In particular φ is injective. The same construction with the roles of F and F' exchanged produces $\psi: F' \rightarrow F$, and $\psi \circ \varphi$ is an order-preserving map fixing $\mathbb{Q}_F$ pointwise. By Step 2 a map fixing a dense subset and preserving order is the identity, since $\psi(\varphi(x)) < x$ or $x < \psi(\varphi(x))$ would leave room for a rational between them, which is fixed and so cannot be moved across. Hence $\psi \circ \varphi$ and $\varphi \circ \psi$ are identities and φ is a bijection.

Step 5: φ preserves the operations. Let $x, y \in F$. If $r, s \in \mathbb{Q}_F$ satisfy $r < x$ and $s < y$ then $r + s < x + y$, so $\varphi_0(r) + \varphi_0(s) = \varphi_0(r + s) \leq \varphi(x + y)$. Taking suprema over r and then s gives $\varphi(x) + \varphi(y) \leq \varphi(x + y)$. For the reverse, if $t \in \mathbb{Q}_F$ satisfies $t < x + y$ then Step 2 gives $r, s \in \mathbb{Q}_F$ with $r < x$, $s < y$ and $t < r + s$, so $\varphi_0(t) < \varphi_0(r) + \varphi_0(s) \leq \varphi(x) + \varphi(y)$. Taking the supremum over t gives $\varphi(x + y) \leq \varphi(x) + \varphi(y)$. So φ preserves addition. For multiplication the same argument applies verbatim when $x, y > 0$, and the general case follows from $\varphi(-x) = -\varphi(x)$, which additivity gives, together with proposition 1.1.2(4).

Step 6: uniqueness. Any isomorphism agrees with φ_0 on $\mathbb{Q}_F$ by Step 1, and an order-preserving map is determined by its values on a dense subset, by the argument of Step 4. So φ is the only one, completing the proof.

Two consequences are worth stating. The field built in section 1.1 from Cauchy sequences and the field built in appendix A from cuts are both complete ordered fields, so the theorem identifies them: the appendix is an alternative route to the same object, not a second object. And the definite article in “the real numbers” is earned.

The theorem also disposes of the candidates a reader is most likely to raise.

1.5.3 Note: The p -adic Absolute Value Is Not an Order For a prime p , writing a nonzero rational as $p^k m/n$ with m and n prime to p and setting $|p^k m/n|_p = p^{-k}$ defines a second notion of size on $\mathbb{Q}$, and completing $\mathbb{Q}$ with respect to it produces the p -adic numbers. This is not a counterexample to theorem 1.5.2, and the reason is that $|\cdot|_p$ is an *absolute value*, not an order: it assigns a size to each element and does not say which of two elements is the larger. The theorem quantifies over ordered fields, so a field carrying only an absolute value is not a candidate for it.

Exercise 1.5.1

1. Show that $\mathbb{Q}$ is an Archimedean ordered field that is not complete, so that the converse of corollary 1.5.1 fails.
2. Show that an ordered field in which every nonempty set bounded below has an infimum satisfies the least upper bound property, so the two forms of completeness are interchangeable as an axiom.

3. Show that the only isomorphism of ordered fields $\mathbb{R} \rightarrow \mathbb{R}$ is the identity, and that the same is false for fields without an order by exhibiting a field with a nontrivial automorphism.
4. Where does the proof of theorem 1.5.2 use completeness of F' , and where does it use completeness of F ? Show that dropping either leaves the construction of φ incomplete.

Sequences and Series

Theorem 1.2.4 states completeness as a fact about sets: every nonempty set of reals with an upper bound has a least one. That is the form in which completeness was proved, and it is not the form in which it is usually spent. Almost every argument in the chapters that follow reaches instead for a statement about sequences, that a sequence whose terms crowd together has somewhere to converge to, and the first task here is to show that the two say the same thing.

The restatement is worth the work for a reason beyond convenience. The least upper bound property is about an order, and it can only be stated where there is one. The Cauchy condition is about distances, and it can be stated wherever distances can be measured. So the sequential form is what survives when the real line is replaced by a space of functions or a space of sequences, and it is the form the later volumes of this series inherit. Section 1.1 already used Cauchy sequences of rationals to build $\mathbb{R}$. This chapter turns the same idea on $\mathbb{R}$ itself.

The second half of the chapter is about infinite sums. A series is defined as a sequence of partial sums, so nothing new is needed to say what one means, and everything proved about sequences applies at once. What is genuinely new is that an infinite sum can depend on the order of its terms: reordering a convergent series can change its value, or destroy convergence altogether, and the condition separating the series where this happens from the series where it cannot is absolute convergence. That distinction is the reason the chapter treats convergence twice.

Section 2.1 defines the limit of a sequence and establishes the algebra of limits, the estimates that every later chapter spends without comment. Section 2.2 adds the monotone convergence theorem, the limit superior and inferior with the characterization that makes them usable, and the Bolzano-Weierstrass theorem. Section 2.3 proves the equivalence promised above and collects the forms of completeness into a single statement. Section 2.4 develops series and the tests that decide them, from the geometric and harmonic series through the comparison, ratio and root tests. Section 2.5 takes up the order dependence, proving that absolute convergence makes a sum rearrangement-invariant and that conditional convergence makes it arbitrary. Section 2.6 specializes to power series, computes the radius of convergence, and defines the exponential and trigonometric functions by the series that will

later be shown to characterize them.

2.1 Convergence of Real Sequences

What section 1.1 never defines is what it means for a sequence of *real* numbers to approach a real number. This section gives that definition together with the facts that make it usable: a limit is unique, a convergent sequence is bounded, limits pass through sums, products and quotients, and limits respect weak inequalities.

The failure in $\mathbb{Q}$ is the one that forced the construction, restated in the vocabulary about to be defined. The sequence 1.4, 1.41, 1.414, 1.4142, ... of example 1.1.5(2) approaches nothing in $\mathbb{Q}$, and section 2.3 will show that it approaches a real number in $\mathbb{R}$. The last result here is the estimate $k^j r^k \rightarrow 0$ for $0 < r < 1$, which is what makes a geometric factor dominate a polynomial one.

Definition 2.1.1: [Real] Sequence

A *sequence* of real numbers is a function $a: \mathbb{N} \rightarrow \mathbb{R}$. The value $a(n)$ is written a_n and called the *n-th term*, and the function itself is written $(a_n)_{n \geq 1}$, or (a_n) when the index set is understood. A sequence indexed by the integers $n \geq n_0$ for a fixed $n_0 \in \mathbb{Z}$ is defined the same way, and every statement below holds for such a sequence with 1 replaced by n_0 throughout.

A sequence is presented by whatever rule produces its terms: a formula ($a_n = 1/n$), a recursion ($a_1 = 1$ and $a_{n+1} = a_n/2$, so that $a_n = 2^{1-n}$), or a description (a_n is the *n*-th decimal truncation of a square root of 2, as in example 1.1.5(2)). The notation (a_n) names the whole function and a_n names one of its values, and the two have to be kept apart as soon as sets of terms appear: the sequence $((-1)^n)$ has a term for every $n \geq 1$, while the set $\{(-1)^n | n \geq 1\} = \{-1, 1\}$ of its terms has two elements.

Convergence is the statement that the terms of a sequence eventually lie within any prescribed distance of one fixed number. Two words in that sentence need fixing before it becomes a definition. The *distance* from a_n to L is $|a_n - L|$, in the sense of definition 1.3.6, and *eventually* means from some index onwards, with the index allowed to depend on the prescribed distance and on nothing else.

Definition 2.1.2: Convergence and the Limit of a Sequence

A sequence (a_n) of real numbers *converges* to $L \in \mathbb{R}$ if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon \quad \text{for every } n \geq N$$

Such an L is called a *limit* of (a_n) , and one writes $a_n \rightarrow L$. A sequence converging to some real number is *convergent*, and one converging to no real number is *divergent*.

One convention changes here and is not mentioned again. Definition 1.1.4 quantified over *rational* ϵ , because that was all that existed when it was stated. From this point ϵ ranges over the positive reals. For a real sequence the two quantifications describe the same condition. A positive rational is a positive real under the embedding of proposition 1.1.11, so quantifying over all real $\epsilon > 0$ is at least as strong. Conversely, given a real $\epsilon > 0$, theorem 1.4.4 applied to $0 < \epsilon$ produces a rational q with $0 < q < \epsilon$, and an index N answering q also answers ϵ , since $|a_n - L| < q$ and $q < \epsilon$ give $|a_n - L| < \epsilon$.

Example 2.1.3: $1/n$, a Constant Sequence, $(n+1)/(2n+3)$, and $((-1)^n)$

1. $1/n \rightarrow 0$. Let $\epsilon > 0$. By theorem 1.4.1(3) there is an $N \in \mathbb{N}$ with $0 < 1/N < \epsilon$. Fix $n \geq N$. If $n > N$ then $0 < 1/n < 1/N$ by proposition 1.1.2(7), and if $n = N$ the two are equal, so in both cases $|1/n - 0| = 1/n \leq 1/N < \epsilon$.
2. A constant sequence $a_n = c$ converges to c . Every $\epsilon > 0$ is answered by $N = 1$, since $|a_n - c| = |0| = 0 < \epsilon$ for every n .
3. The sequence $((n+1)/(2n+3))$ converges to $1/2$. Subtracting and clearing denominators,

$$\frac{n+1}{2n+3} - \frac{1}{2} = \frac{2(n+1) - (2n+3)}{2(2n+3)} = \frac{-1}{2(2n+3)}$$

so the distance to $1/2$ is $1/(2(2n+3))$, and this is smaller than $1/(4n)$ because $4n < 4n+6 = 2(2n+3)$. Given $\epsilon > 0$, apply theorem 1.4.1(3) to the positive real 4ϵ to get an $N \in \mathbb{N}$ with $0 < 1/N < 4\epsilon$. For $n \geq N$ the argument of (1) gives $1/n \leq 1/N$, hence

$$\left| \frac{n+1}{2n+3} - \frac{1}{2} \right| < \frac{1}{4n} \leq \frac{1}{4N} < \epsilon$$

4. $((-1)^n)$ is divergent. Suppose $(-1)^n \rightarrow L$ and take $\epsilon = 1$ in definition 2.1.2, producing an N with $|(-1)^n - L| < 1$ for all $n \geq N$. Among $2N$ and $2N+1$ the first is even and the second is odd, and both are at least N , so $|1 - L| < 1$ and $|-1 - L| < 1$. By proposition 1.3.8(4) and then (1),

$$2 = |1 - (-1)| = |(1 - L) + (L - (-1))| \leq |1 - L| + |-1 - L| < 2$$

which is false. So no such L exists.

Items (1) and (3) run the same route: bound $|a_n - L|$ above by an explicit expression in n , then use theorem 1.4.1 to choose an index past which that expression is below ϵ . Item (4) shows what a divergence proof has to do, since no candidate limit is given in advance and every one of them has to be excluded at once. Theorem 2.1.6 removes most of this labour by reducing a new limit to limits already known.

Definition 2.1.2 calls L a limit, since nothing said so far excludes a second one. The following excludes it. Two distinct real numbers are separated by a positive distance, while the terms of a convergent sequence eventually lie within every positive distance of each of its limits, and those two statements cannot both hold.

Proposition 2.1.4: A Sequence Has at Most One Limit

If $a_n \rightarrow L$ and $a_n \rightarrow L'$, then $L = L'$.

Proof:

Let $\epsilon > 0$. Applying definition 2.1.2 to L with the positive real $\epsilon/2$ gives an N_1 with $|a_n - L| < \epsilon/2$ for all $n \geq N_1$, and applying it to L' with $\epsilon/2$ gives an N_2 with $|a_n - L'| < \epsilon/2$ for all $n \geq N_2$.

Fix an n at least as large as both. Then proposition 1.3.8(4) gives

$$|L - L'| = |(L - a_n) + (a_n - L')| \leq |L - a_n| + |a_n - L'| < \frac{\epsilon}{2} + \frac{\epsilon}{2}$$

where $|L - a_n| = |a_n - L|$ by proposition 1.3.8(1) applied to $x = a_n - L$. The right side is ϵ , so $|L - L'| \leq \epsilon$ holds for every $\epsilon > 0$, and proposition 1.4.7(2) gives $L = L'$, as we sought to show.

Uniqueness is what makes the notation well defined

$$\lim_{n \rightarrow \infty} a_n = L$$

for the limit of a convergent sequence, and with it the practice of treating $\lim_{n \rightarrow \infty} a_n$ as a number that can be added, multiplied and compared, which is what the next theorem makes legitimate.

Call a sequence (a_n) *bounded* if there is an $M \in \mathbb{R}$ with $|a_n| \leq M$ for every n . Such an M satisfies $M \geq |a_1| \geq 0$, so proposition 1.3.8(2) applies and says that $-M \leq a_n \leq M$ for every n , so that the set $\{a_n | n \geq 1\}$ of terms is bounded above by M and below by $-M$ in the sense of definition 1.2.1. A convergent sequence cannot fail this: past the index answering $\epsilon = 1$ every term is within 1 of the limit, and only finitely many terms are left unaccounted for. Part (3) of theorem 2.1.6 needs a bound of exactly this kind on one of its two factors.

Proposition 2.1.5: Convergent Sequences Are Bounded

Every convergent sequence of real numbers is bounded.

Proof:

Let $a_n \rightarrow L$ and apply definition 2.1.2 with $\epsilon = 1$, producing an $N \in \mathbb{N}$ with $|a_n - L| < 1$ for all $n \geq N$. For such n , proposition 1.3.8(4) gives

$$|a_n| = |(a_n - L) + L| \leq |a_n - L| + |L| < 1 + |L|$$

If $N = 1$ this covers every term, and $M = 1 + |L|$ is a bound. If $N > 1$, the finitely many numbers $|a_1|, \dots, |a_{N-1}|$ have a largest, since a finite nonempty set of reals has a largest element. Taking M to be the larger of that number and $1 + |L|$ bounds every term, completing the proof.

The converse fails, and example 2.1.3(4) is the witness: $((-1)^n)$ is bounded by 1 and has no limit. Boundedness is a necessary condition for convergence and nothing more, which is how it is used below, to control one factor of a product while the other factor is made small. Section 2.2 recovers a partial converse by weakening the conclusion from a limit of the sequence to a limit of one of its subsequences.

Computing a limit from definition 2.1.2 means producing an index for each ϵ , and doing that from scratch for each new sequence is avoidable whenever the sequence is assembled from ones whose limits are known. The following reduces the limit of an expression built from convergent sequences by the field operations to the limits of the sequences it is built from.

Theorem 2.1.6: The Algebra of Limits

Let (a_n) and (b_n) be real sequences with $a_n \rightarrow a$ and $b_n \rightarrow b$, and let $c \in \mathbb{R}$.

1. $a_n + b_n \rightarrow a + b$.
2. $c a_n \rightarrow c a$.
3. $a_n b_n \rightarrow ab$.
4. If $b \neq 0$ and $b_n \neq 0$ for every n , then $a_n/b_n \rightarrow a/b$.

Proof:

1. Let $\epsilon > 0$. Choose N_1 with $|a_n - a| < \epsilon/2$ for $n \geq N_1$ and N_2 with $|b_n - b| < \epsilon/2$ for $n \geq N_2$, and let N be the larger. For $n \geq N$, proposition 1.3.8(4) gives

$$|(a_n + b_n) - (a + b)| = |(a_n - a) + (b_n - b)| \leq |a_n - a| + |b_n - b| < \epsilon$$

2. If $c = 0$ then $c a_n = 0$ for every n and $c a = 0$, by proposition 1.1.2(3), and the constant sequence 0 converges to 0 by example 2.1.3(2). Suppose $c \neq 0$, so that $|c| > 0$ by proposition 1.3.8(1). Let $\epsilon > 0$ and choose N with $|a_n - a| < \epsilon/|c|$ for $n \geq N$. For such n , proposition 1.3.8(3) gives

$$|c a_n - c a| = |c| |a_n - a| < |c| \cdot \frac{\epsilon}{|c|} = \epsilon$$

3. By proposition 2.1.5 there is an M with $|b_n| \leq M$ for every n , and $M \geq 0$ because $|b_1| \geq 0$. Put $K = M + |a| + 1$, so that $K \geq 1 > 0$, $K \geq M$ and $K \geq |a|$. For every n , proposition 1.3.8(4) and then (3) give

$$|a_n b_n - ab| = |(a_n - a)b_n + a(b_n - b)| \leq |a_n - a||b_n| + |a||b_n - b| \leq K(|a_n - a| + |b_n - b|)$$

the last step because $|b_n| \leq K$ and $|a| \leq K$, and multiplication by a nonnegative number preserves a weak inequality: a zero multiplier makes both products 0, and a positive multiplier is proposition 1.1.2(5) when the inequality is strict and an equality when it is not. Now let $\epsilon > 0$. Choose N_1 with $|a_n - a| < \epsilon/(2K)$ for $n \geq N_1$ and N_2 with $|b_n - b| < \epsilon/(2K)$ for $n \geq N_2$. For n at least as large as both, $|a_n b_n - ab| \leq K(|a_n - a| + |b_n - b|) < K(\epsilon/(2K) + \epsilon/(2K)) = \epsilon$.

4. It is enough to prove $1/b_n \rightarrow 1/b$, since part (3) applied to (a_n) and $(1/b_n)$ then gives $a_n/b_n = a_n \cdot (1/b_n) \rightarrow a \cdot (1/b) = a/b$. Since $b \neq 0$, proposition 1.3.8(1) gives $|b| > 0$, so $|b|/2$ is a positive real and definition 2.1.2

produces an N_1 with $|b_n - b| < |b|/2$ for $n \geq N_1$. For such n , proposition 1.3.8(1) applied to $|b_n| - |b|$ and then proposition 1.3.8(5) give

$$|b| - |b_n| \leq ||b_n| - |b|| \leq |b_n - b| < \frac{|b|}{2}$$

so $|b_n| > |b|/2 > 0$. Multiplying by $|b| > 0$ with proposition 1.1.2(5) and using proposition 1.3.8(3) gives $|b_n b| = |b_n||b| > |b|^2/2 > 0$, and inverting with proposition 1.1.2(7) gives $1/|b_n b| < 2/|b|^2$. Hence for $n \geq N_1$,

$$\left| \frac{1}{b_n} - \frac{1}{b} \right| = \left| \frac{b - b_n}{b_n b} \right| = \frac{|b_n - b|}{|b_n b|} \leq \frac{2}{|b|^2} |b_n - b|$$

where the second equality uses $|b - b_n| = |b_n - b|$ from proposition 1.3.8(1) and proposition 1.3.8(3) in the form $|u/v| = |u|/|v|$, and the last step multiplies $1/|b_n b| < 2/|b|^2$ by the nonnegative number $|b_n - b|$, which gives an equality of zeros when $b_n = b$. Given $\epsilon > 0$, choose N_2 with $|b_n - b| < \epsilon|b|^2/2$ for $n \geq N_2$. For n at least as large as N_1 and N_2 the bound $(2/|b|^2)|b_n - b|$ on $|1/b_n - 1/b|$ is smaller than ϵ , so $1/b_n \rightarrow 1/b$, completing the proof.

With these four rules a limit is computed by rewriting the sequence in terms of sequences whose limits are known. Take

$$a_n = \frac{3n^2 + 2n}{n^2 - 5}$$

which is defined for $n \geq 3$, since $n^2 - 5 \geq 4$ there. Dividing numerator and denominator by n^2 turns it into

$$a_n = \frac{3 + 2/n}{1 - 5/n^2}$$

By example 2.1.3(1) and part (3), $1/n^2 = (1/n)(1/n) \rightarrow 0$. Parts (1) and (2), together with the constant sequences of example 2.1.3(2), give $3 + 2/n \rightarrow 3$ and $1 - 5/n^2 \rightarrow 1$. Part (4) applies because the limit 1 of the denominator is nonzero and each $1 - 5/n^2$ is nonzero for $n \geq 3$, where $5/n^2 \leq 5/9 < 1$. Therefore $a_n \rightarrow 3$.

Theorem 2.1.6 computes a limit from limits already known. A different situation is a sequence whose terms are not built from convergent sequences by the field operations, but which lies between two sequences having a common limit. That position alone forces convergence, and forces the limit to be the common one.

Theorem 2.1.7: The Squeeze Theorem

Let (a_n) , (b_n) and (c_n) be real sequences and let $N_0 \in \mathbb{N}$ be an index with

$$a_n \leq b_n \leq c_n \quad \text{for every } n \geq N_0$$

If $a_n \rightarrow L$ and $c_n \rightarrow L$, then $b_n \rightarrow L$.

Proof:

Let $\epsilon > 0$. Choose N_1 with $|a_n - L| < \epsilon$ for $n \geq N_1$ and N_2 with $|c_n - L| < \epsilon$ for $n \geq N_2$, and let N be the largest of N_0 , N_1 and N_2 . Fix $n \geq N$. By proposition 1.3.8(2) with $c = \epsilon$, the bound $|a_n - L| < \epsilon$ reads $-\epsilon < a_n - L < \epsilon$ and so gives $L - \epsilon < a_n$, and $|c_n - L| < \epsilon$ gives $c_n < L + \epsilon$. Since $n \geq N_0$, the hypothesis inserts b_n between them:

$$L - \epsilon < a_n \leq b_n \leq c_n < L + \epsilon$$

By proposition 1.1.2(6), which reads $x < y$ as $0 < y - x$, the outer inequalities $L - \epsilon < b_n$ and $b_n < L + \epsilon$ say respectively that $0 < (b_n - L) + \epsilon$ and $0 < \epsilon - (b_n - L)$, that is $-\epsilon < b_n - L < \epsilon$. Then proposition 1.3.8(2), read in the other direction, turns this into $|b_n - L| < \epsilon$. As $\epsilon > 0$ was arbitrary, $b_n \rightarrow L$, as we sought to show.

The hypothesis is imposed only from the index N_0 onwards, which is what makes the theorem applicable: the two bounding sequences are typically produced by an estimate that is valid only for large n . For a first use, let (x_n) be any sequence with $|x_n| \leq 1$ for every n . Then $-1 \leq x_n \leq 1$ by proposition 1.3.8(2), and multiplying by the positive number $1/n$ gives $-1/n \leq x_n/n \leq 1/n$ for every n , by proposition 1.1.2(5) where the inequality is strict and by an equality where it is not. Both outer sequences converge to 0, by example 2.1.3(1) and theorem 2.1.6(2) with $c = -1$, so theorem 2.1.7 gives $x_n/n \rightarrow 0$ whatever the signs of the x_n .

Convergence interacts with the order of $\mathbb{R}$ as well as with its operations, and in a weakened form. An inequality between the terms of two convergent sequences passes to their limits, but a strict inequality between terms yields only a weak inequality between limits.

Proposition 2.1.8: Limits Preserve Weak Inequalities

Let (a_n) and (b_n) be real sequences with $a_n \rightarrow a$ and $b_n \rightarrow b$, and let $N_0 \in \mathbb{N}$.

1. If $a_n \leq b_n$ for every $n \geq N_0$, then $a \leq b$.
2. If $c \in \mathbb{R}$ satisfies $c \leq a_n$ for every $n \geq N_0$, then $c \leq a$. If $d \in \mathbb{R}$ satisfies $a_n \leq d$ for every $n \geq N_0$, then $a \leq d$.

Proof:

1. Let $\epsilon > 0$. Choose N_1 with $|a_n - a| < \epsilon/2$ for $n \geq N_1$ and N_2 with $|b_n - b| < \epsilon/2$ for $n \geq N_2$, and fix an n at least as large as each of N_0 , N_1 and N_2 . Splitting $a - b$ across that index,

$$a - b = (a - a_n) + (a_n - b_n) + (b_n - b) \leq |a - a_n| + 0 + |b_n - b| < \frac{\epsilon}{2} + \frac{\epsilon}{2}$$

where the first and third terms are bounded by proposition 1.3.8(1), which gives $x \leq |x|$, the middle term is at most 0 by the hypothesis $a_n \leq b_n$, and $|a - a_n| = |a_n - a|$ by proposition 1.3.8(1) again. So $a \leq b + \epsilon$ for every $\epsilon > 0$, and proposition 1.4.7(1) gives $a \leq b$.

2. The constant sequence with every term c converges to c by example 2.1.3(2), so part (1)

applied to it and to (a_n) gives $c \leq a$. Applying part (1) to (a_n) and to the constant sequence with every term d gives $a \leq d$, completing the proof.

Strict inequalities are not preserved. The constant sequence 0 and the sequence $(1/n)$ satisfy $0 < 1/n$ for every n , and their limits are both 0 by example 2.1.3(1) and (2). This is why part (2) is the form used to place a limit inside a closed interval: bounds $c \leq a_n \leq d$ on the terms give $c \leq a \leq d$ on the limit, and nothing sharper is available from the terms alone.

Every proof above reduced a claim about $a_n \rightarrow L$ to a claim about $|a_n - L|$ becoming small, which is convergence to 0. That case has a name, and it is the name already attached to rational sequences in definition 1.1.4.

Definition 2.1.9: Null Sequence

A real sequence (a_n) is *null* if $a_n \rightarrow 0$, that is, if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ with $|a_n| < \epsilon$ for every $n \geq N$.

The condition defining $a_n \rightarrow L$ is the inequality $|(a_n - L) - 0| < \epsilon$ for large n , which is the condition defining nullity of $(a_n - L)$. So $a_n \rightarrow L$ if and only if $(a_n - L)$ is null, and a convergence statement can always be restated as a statement about a null sequence. Example 2.1.3(1) says that $(1/n)$ is one, and example 2.1.3(4) shows $((-1)^n)$ is not, while $((-1)^n/n)$ is by the computation following theorem 2.1.7. For a sequence of rational numbers the definition above agrees with the one inside definition 1.1.4, by the same density argument that reconciled the two ranges of ϵ .

Three facts about null sequences are used throughout the chapters that follow. The first is that the null sequences are closed under sums and scalar multiples, the second turns any bound by a null sequence into nullity, and the third is a decay estimate: a fixed power of k multiplied by the k -th power of a number between 0 and 1 tends to zero, however large the fixed power and however close the number is to 1.

Proposition 2.1.10: Sums, Comparison, and the Decay of $k^j r^k$

1. If (a_n) and (b_n) are null and $c \in \mathbb{R}$, then $(a_n + b_n)$ and $(c a_n)$ are null.
2. If (b_n) is null and there is an $N_0 \in \mathbb{N}$ with $|a_n| \leq b_n$ for every $n \geq N_0$, then (a_n) is null.
3. For every integer $j \geq 0$ and every real r with $0 < r < 1$, the sequence $(k^j r^k)_{k \geq 1}$ is null.

Proof:

1. These are parts (1) and (2) of theorem 2.1.6 with $a = b = 0$, since $0 + 0 = 0$ and $c \cdot 0 = 0$ by proposition 1.1.2(3).
2. For $n \geq N_0$ the hypothesis gives $|a_n| \leq b_n$, and $|a_n| \geq 0$ by proposition 1.3.8(1), so $b_n \geq 0$ and proposition 1.3.8(2), with b_n in the role of its constant c , turns the hypothesis into $-b_n \leq a_n \leq b_n$. The sequence $(-b_n)$ is null by part (1) with $c = -1$, so theorem 2.1.7 applies with $L = 0$ and the index N_0 and gives $a_n \rightarrow 0$.

3. Fix an integer $j \geq 0$ and a real r with $0 < r < 1$, and put $h = (1 - r)/r$. Then $h > 0$, since $1 - r > 0$ and $r > 0$, and

$$1 + h = 1 + \frac{1 - r}{r} = \frac{1}{r}$$

so that $r^k = (1 + h)^{-k}$ for every $k \geq 1$. Write $m = j + 1$, an integer with $m \geq 1$, and let $k > 2m$.

By [4, the binomial theorem], $(1 + h)^k$ expands as $\sum_{i=0}^k \binom{k}{i} h^i$, a sum of nonnegative terms containing $\binom{k}{m} h^m$ as one summand, the index m being admissible because $m < 2m < k$. By [4, the falling-factorial identity], $k(k - 1) \cdots (k - m + 1) = m! \binom{k}{m}$. Hence

$$(1 + h)^k \geq \binom{k}{m} h^m = \frac{k(k - 1) \cdots (k - m + 1)}{m!} h^m$$

Each of the m factors in that product satisfies $k - i \geq k - m + 1 > k/2$ for $0 \leq i \leq m - 1$, because $k > 2m$ gives $m < k/2$ and hence $k - m + 1 > k - k/2 = k/2$. Multiplying these m inequalities between positive numbers, by m applications of proposition 1.1.2(5), gives $k(k - 1) \cdots (k - m + 1) > (k/2)^m = k^m/2^m$. Multiplying that by the positive number $h^m/m!$ and combining with $(1 + h)^k \geq k(k - 1) \cdots (k - m + 1) h^m/m!$,

$$(1 + h)^k > \frac{k^m h^m}{2^m m!} > 0$$

Inverting this with proposition 1.1.2(7) and then multiplying by $k^j > 0$ with proposition 1.1.2(5) gives, for every $k > 2m$,

$$0 < k^j r^k = \frac{k^j}{(1 + h)^k} < \frac{2^m m!}{h^m} \cdot \frac{k^j}{k^m} = \frac{C}{k}, \quad C = \frac{2^m m!}{h^m}$$

the last equality because $m = j + 1$ makes $k^j/k^m = 1/k$. The sequence $(C/k)_{k \geq 1}$ is null by example 2.1.3(1) and part (1), and $|k^j r^k| = k^j r^k$ because these terms are positive. So part (2), applied with the index $N_0 = 2m + 1$, shows that $(k^j r^k)_{k \geq 1}$ is null, completing the proof.

Part (3) is stated with an arbitrary polynomial power because that is how it is used: a sequence bounded in absolute value by $C k^j r^k$ is null, by parts (1) and (2), no matter how large j is, so long as $0 < r < 1$. For $-1 < r < 0$ the terms satisfy $|k^j r^k| = k^j |r|^k$ by repeated application of proposition 1.3.8(3), with $0 < |r| < 1$, so part (3) applied to $|r|$ and then part (2) give $k^j r^k \rightarrow 0$ there as well, and for $r = 0$ every term of $(k^j r^k)_{k \geq 1}$ is 0. So the conclusion holds for every r with $|r| < 1$, and how close $|r|$ is to 1 changes only the constant C in the proof, never the comparison index $2m + 1 = 2j + 3$ and never the conclusion.

Exercise 2.1.1

1. Show directly from definition 2.1.2, without using theorem 2.1.6, that

$$\frac{3n - 1}{n + 2} \rightarrow 3$$

2. Let (a_n) be a real sequence with $a_n \rightarrow a$. Show that $|a_n| \rightarrow |a|$. Then exhibit a sequence (a_n) for which $(|a_n|)$ converges while (a_n) does not, so that the converse fails.
3. Let (a_n) be a real sequence with $a_n \geq 0$ for every n and with $a_n \rightarrow a$. Show first that $a \geq 0$, so that $\sqrt{a}$ is defined by theorem 1.3.12, and then that $\sqrt{a_n} \rightarrow \sqrt{a}$. Treat the cases $a > 0$ and $a = 0$ separately.
4. Let $a_n \rightarrow a$ with $a > 0$. Show that there is an $N \in \mathbb{N}$ with $a_n > a/2$ for every $n \geq N$, and deduce that at most finitely many terms of the sequence are less than or equal to 0.
5. Show that $(n^5 2^{-n})_{n \geq 1}$ is null. Show also that $(n!/n^n)_{n \geq 1}$ is null, and explain why proposition 2.1.10(3) does not apply to the second sequence.
6. Let $x \in \mathbb{R}$ and write $\lfloor y \rfloor$ for the integer part of a real number y , as in proposition 1.4.3. Show that

$$\frac{\lfloor nx \rfloor}{n} \rightarrow x$$

2.2 Monotone Sequences, Limsup and Liminf

Every result in section 2.1 starts from a sequence already known to converge, because the definition of a limit names the limit in advance. This section produces convergence from hypotheses that never name the limit. Two hypotheses suffice: monotonicity together with boundedness gives a limit outright, and boundedness alone gives one along a subsequence.

Between those two statements sit the upper and lower limits, two numbers in the extended real line $[-\infty, +\infty]$ attached to every real sequence, convergent or not. The sequence has a limit in $[-\infty, +\infty]$ exactly when the two agree, and it converges exactly when they agree on a real value. The tests of section 2.4 and the radius of convergence of section 2.6 are both stated in terms of the upper limit, so the working properties proved here are what those sections use.

Monotonicity removes the obligation to name the limit first. An increasing sequence bounded above cannot pass the least of its upper bounds, and theorem 1.2.4 guarantees that this number exists.

Inside $\mathbb{Q}$ the guarantee fails, and one construction makes the failure explicit. By theorem 1.4.4 choose $q_1 \in \mathbb{Q}$ with $\sqrt{2} - 1 < q_1 < \sqrt{2}$, and, given q_n , choose $q_{n+1} \in \mathbb{Q}$ with

$$\max\left\{q_n, \sqrt{2} - \frac{1}{n+1}\right\} < q_{n+1} < \sqrt{2}$$

which is possible because both numbers inside the maximum are smaller than $\sqrt{2}$. Every term is rational, the sequence increases, and every term is smaller than $3/2$ because $q_n < \sqrt{2} \leq 3/2$ by example 1.3.13. Moreover $0 < \sqrt{2} - q_n < 1/n$ for every n . Now $1/n \rightarrow 0$: given $\epsilon > 0$, theorem 1.4.1 gives an N with $0 < 1/N < \epsilon$, and $0 < 1/n \leq 1/N < \epsilon$ for $n \geq N$ by proposition 1.1.2. So $\sqrt{2} - q_n \rightarrow 0$ by theorem 2.1.7, and $q_n \rightarrow \sqrt{2}$ by theorem 2.1.6. Since $\sqrt{2}$ is not rational (proposition 1.1.3) and a limit is unique (proposition 2.1.4), no rational number is the limit of (q_n) . Boundedness and monotonicity produce nothing inside $\mathbb{Q}$. The theorem below is a statement about $\mathbb{R}$ and about nothing weaker.

Definition 2.2.1: Increasing, Decreasing and Monotone Sequences

A real sequence $(a_n)_{n \geq 1}$ is *increasing* if $a_n \leq a_{n+1}$ for every $n \geq 1$, and *decreasing* if $a_n \geq a_{n+1}$ for every $n \geq 1$. It is *monotone* if it is increasing or decreasing.

The rational sequence (q_n) built above is increasing, as is $a_n = 1 - 1/n$, while $a_n = 1/n$ is decreasing and $a_n = (-1)^n$ is neither. A constant sequence is both.

Theorem 2.2.2: Bounded Monotone Sequences Converge

Let $(a_n)_{n \geq 1}$ be a monotone real sequence. Then (a_n) converges if and only if it is bounded. If (a_n) is increasing and bounded above, then

$$\lim_{n \rightarrow \infty} a_n = \sup\{a_n | n \geq 1\}$$

and if (a_n) is decreasing and bounded below, then $\lim_{n \rightarrow \infty} a_n = \inf\{a_n | n \geq 1\}$.

Proof:

If (a_n) converges then it is bounded by proposition 2.1.5, and this direction uses no monotonicity.

For the converse, suppose first that (a_n) is increasing and bounded above. The set $E = \{a_n | n \geq 1\}$ is nonempty and bounded above, so theorem 1.2.4 gives a real number $s = \sup E$. Let $\epsilon > 0$. Since s is an upper bound of E , proposition 1.2.6 produces an element of E larger than $s - \epsilon$, that is, an index N with $a_N > s - \epsilon$. Induction on n gives $a_n \geq a_N$ for every $n \geq N$: the case $n = N$ is an equality, and $a_{n+1} \geq a_n \geq a_N$ by definition 2.2.1. For every such n ,

$$s - \epsilon < a_N \leq a_n \leq s$$

the last inequality because s is an upper bound of E . Hence $0 \leq s - a_n < \epsilon$, so $|a_n - s| = s - a_n < \epsilon$ by definition 1.3.6 and proposition 1.3.8. As ϵ was arbitrary, $a_n \rightarrow s$ in the sense of definition 2.1.2. Note that only the upper bound was used: an increasing sequence is bounded below by a_1 , so for an increasing sequence boundedness and boundedness above are the same hypothesis.

Suppose next that (a_n) is decreasing and bounded below, and put $b_n = -a_n$. Then (b_n) is increasing, and it is bounded above: if $a_n \geq L$ for all n then $b_n \leq -L$ by proposition 1.1.2. The case already proved gives $b_n \rightarrow \sup\{b_n | n \geq 1\}$. Writing $A = \{a_n | n \geq 1\}$, the set $\{b_n | n \geq 1\}$ is $-A$, and A is bounded below, so proposition 1.3.2 gives $\sup(-A) = -\inf A$. Thus $b_n \rightarrow -\inf A$, and multiplying by the constant sequence with value -1 and applying theorem 2.1.6 gives $a_n \rightarrow \inf A$, as we sought to show.

The theorem converts the least upper bound property into a statement about sequences. What it gives is a limit whose value is not part of the hypothesis: the supremum comes from theorem 1.2.4, and identifying it afterwards is a separate problem. The next example is a sequence defined by a recursion in which no term is given in closed form, so that the limit exists well before any computation names it.

Example 2.2.3: The Nested Radicals $\sqrt{2}$, $\sqrt{2 + \sqrt{2}}$, ...

Let $a_1 = \sqrt{2}$ and $a_{n+1} = \sqrt{2 + a_n}$ for $n \geq 1$. Each term is defined and nonnegative: $a_1 \geq 0$, and if $a_n \geq 0$ then $2 + a_n \geq 2 > 0$, so theorem 1.3.12 produces the unique nonnegative square root a_{n+1} .

Every term is smaller than 2. First $a_1 < 2$, since $a_1 \geq 2$ would give $2 = a_1^2 \geq 4$ by proposition 1.1.2. If $a_n < 2$ then $2 + a_n < 4$, and $a_{n+1} \geq 2$ would give $2 + a_n = a_{n+1}^2 \geq 4$, so $a_{n+1} < 2$.

The sequence increases. For each n ,

$$a_{n+1}^2 - a_n^2 = (2 + a_n) - a_n^2 = (2 - a_n)(1 + a_n)$$

and both factors are positive because $0 \leq a_n < 2$. If $a_{n+1} \leq a_n$ held, then $a_{n+1}^2 \leq a_{n+1}a_n \leq a_n^2$ by proposition 1.1.2, contradicting $a_{n+1}^2 - a_n^2 > 0$. So $a_{n+1} > a_n$.

By theorem 2.2.2 the sequence converges to a real number L . Since the sequence increases, $a_1 \leq a_n < 2$ for every n , so proposition 2.1.8, applied to (a_n) against the constant sequences with values a_1 and 2, gives $\sqrt{2} = a_1 \leq L \leq 2$. To identify L , note that the shifted sequence $b_n = a_{n+1}$ also converges to L : given $\epsilon > 0$ choose N with $|a_m - L| < \epsilon$ for all $m \geq N$, and then $n \geq N$ forces $n + 1 \geq N$, so $|b_n - L| < \epsilon$. Now $b_n^2 \rightarrow L^2$ by theorem 2.1.6 applied to the product of (b_n) with itself, while $b_n^2 = 2 + a_n \rightarrow 2 + L$ by theorem 2.1.6 applied to the sum with the constant sequence 2. Limits are unique (proposition 2.1.4), so $L^2 = 2 + L$, that is,

$$(L - 2)(L + 1) = 0$$

Since $L \geq \sqrt{2} > 0$, the factor $L + 1$ is not zero, so $L = 2$.

Boundedness alone does not give a limit. The sequence $(-1)^n$ does not converge: a limit a would give an N with $|(-1)^n - a| < 1$ for all $n \geq N$, and choosing one even and one odd such n gives $|1 - a| < 1$ and $|-1 - a| < 1$, whereas $2 = |1 - (-1)| \leq |1 - a| + |-1 - a| < 2$ by proposition 1.3.8. What boundedness does give is a limit along part of the sequence, and to say this precisely takes two pieces of vocabulary. The first is a value system large enough to record unbounded behaviour, since the constructions below assign a value to every sequence and not only to the bounded ones.

Definition 2.2.4: The Extended Real Line

The *extended real line* is the set

$$[-\infty, +\infty] = \mathbb{R} \cup \{-\infty, +\infty\}$$

where $-\infty$ and $+\infty$ are two symbols not belonging to $\mathbb{R}$. Its order restricts to the order of $\mathbb{R}$ on $\mathbb{R}$, and in addition $-\infty < x < +\infty$ for every $x \in \mathbb{R}$, together with $-\infty < +\infty$. Negation extends by $-(+\infty) = -\infty$ and $-(-\infty) = +\infty$. Addition extends partially: $x + (+\infty) = (+\infty) + x = +\infty$ for every $x \neq -\infty$, and $x + (-\infty) = (-\infty) + x = -\infty$ for every $x \neq +\infty$, while $(+\infty) + (-\infty)$ is left undefined.

For $S \subseteq [-\infty, +\infty]$, an element u is an *upper bound* of S if $x \leq u$ for every $x \in S$, and $\sup S$ denotes the least upper bound of S when one exists. The terms *lower bound* and $\inf S$ are defined symmetrically. For a real sequence (a_n) , write $a_n \rightarrow +\infty$ if for every $M \in \mathbb{R}$ there is an N with $a_n > M$ for all $n \geq N$, and $a_n \rightarrow -\infty$ if for every $M \in \mathbb{R}$ there is an N with $a_n < M$ for all $n \geq N$. The sequence (a_n) has *limit* $s \in [-\infty, +\infty]$ if s is real and $a_n \rightarrow s$ in the sense of definition 2.1.2, or if $s = \pm\infty$ and $a_n \rightarrow s$ in the sense just given.

The relation just described is a total order. For two real arguments it is the order of $\mathbb{R}$, for one real and one infinite argument it is one of the declared inequalities, and two infinite arguments are either equal or ordered by $-\infty < +\infty$. For transitivity, let $x < y < z$. If $x = -\infty$ then $z \neq -\infty$, since $y < z$, so $x < z$ by declaration; if $z = +\infty$ then $x \neq +\infty$, since $x < y$, so again $x < z$; and in the remaining case x, y, z are all real and transitivity is inherited from $\mathbb{R}$. Note that a sequence with limit $+\infty$ does not converge: the word *converges* stays reserved for a real limit, and $a_n \rightarrow +\infty$ records a specific way of failing to converge. As for the enlargement of suprema, no real number is an upper bound of $\mathbb{N}$ by theorem 1.4.1 and $-\infty$ is not one either, so $+\infty$ is the only upper bound and $\sup \mathbb{N} = +\infty$, while the supremum of $\{x \in \mathbb{R} | x^2 < 2\}$ is the real number $\sqrt{2}$ of example 1.3.13 and is unchanged.

The point of the enlargement is that suprema and infima now exist without any hypothesis whatever, which is what lets the definitions further down apply to every sequence.

Proposition 2.2.5: Suprema, Infima and Density in $[-\infty, +\infty]$

1. Every subset $S \subseteq [-\infty, +\infty]$ has a supremum and an infimum in $[-\infty, +\infty]$, and $\inf S = -\sup(-S)$, where $-S = \{-x | x \in S\}$. In particular $\sup \emptyset = -\infty$ and $\inf \emptyset = +\infty$.
2. If $u, v \in [-\infty, +\infty]$ and $u < v$, then there is a real number x with $u < x < v$.

Proof:

1. Four cases produce the supremum.

If $S = \emptyset$, every element of $[-\infty, +\infty]$ is vacuously an upper bound, and $-\infty$ is the smallest element, so $\sup \emptyset = -\infty$.

If $+\infty \in S$, then an upper bound u satisfies $u \geq +\infty$, so $u = +\infty$, and $+\infty$ is an upper bound. Hence $\sup S = +\infty$.

If S is nonempty, $+\infty \notin S$ and $S \cap \mathbb{R} = \emptyset$, then $S = \{-\infty\}$, and $-\infty$ is an upper bound

and is smaller than every other element, so $\sup S = -\infty$.

In the remaining case $S \cap \mathbb{R} \neq \emptyset$ and $+\infty \notin S$. If $S \cap \mathbb{R}$ is bounded above by some real number, theorem 1.2.4 gives $\sigma = \sup(S \cap \mathbb{R}) \in \mathbb{R}$. Then σ is an upper bound of S , since every real element of S is at most σ and $-\infty < \sigma$. If u is any upper bound of S , then $u \neq -\infty$ because $S \cap \mathbb{R} \neq \emptyset$, and if u is real then $u \geq \sigma$ because u bounds $S \cap \mathbb{R}$ above, while $u = +\infty$ certainly exceeds σ . So $\sup S = \sigma$. If instead $S \cap \mathbb{R}$ is not bounded above by any real number, then no real number is an upper bound of S , and neither is $-\infty$ because $S \cap \mathbb{R} \neq \emptyset$, so $+\infty$ is the only upper bound and $\sup S = +\infty$.

For the infimum, observe that $u < v$ holds if and only if $-v < -u$. For two real arguments this is proposition 1.1.2. If $u = -\infty$ and v is real, then $-v$ is real and $-u = +\infty$, so both sides hold. If u is real and $v = +\infty$, then $-v = -\infty < -u$ and both sides hold again. If $u = -\infty$ and $v = +\infty$, then $-v = -\infty < +\infty = -u$. No other case has $u < v$, and applying the equivalence to $-u$ and $-v$ gives the converse direction. Consequently u is a lower bound of S if and only if $-u$ is an upper bound of $-S$, and u is the greatest such if and only if $-u$ is the least such. Since $\sup(-S)$ exists by what was just proved, $-\sup(-S)$ is the greatest lower bound of S . Hence $\inf S$ exists and equals $-\sup(-S)$, and taking $S = \emptyset$ gives $\inf \emptyset = -\sup \emptyset = +\infty$.

2. If u and v are both real, proposition 1.1.2 gives $u < (u + v)/2 < v$. If $u = -\infty$ and v is real, take $x = v - 1$. If u is real and $v = +\infty$, take $x = u + 1$. If $u = -\infty$ and $v = +\infty$, take $x = 0$. No further case occurs, since $u < v$ excludes $u = +\infty$ and $v = -\infty$. This exhausts the possibilities, completing the proof.

The second piece of vocabulary is the operation of passing to part of a sequence. Keeping infinitely many terms of (a_n) in their original order gives a new sequence, and the rest of the section is about which limits such a sequence can have.

Definition 2.2.6: Subsequences and Subsequential Limits

Let $(a_n)_{n \geq 1}$ be a real sequence and let $n_1 < n_2 < n_3 < \dots$ be a strictly increasing sequence of elements of $\mathbb{N}$. The sequence $k \mapsto a_{n_k}$ is a *subsequence* of (a_n) , written $(a_{n_k})_{k \geq 1}$. An element $s \in [-\infty, +\infty]$ is a *subsequential limit* of (a_n) if some subsequence of (a_n) has limit s in the sense of definition 2.2.4.

For $a_n = (-1)^n$ the choice $n_k = 2k$ gives the constant subsequence with value 1, and $n_k = 2k - 1$ gives the constant subsequence with value -1 , so both 1 and -1 are subsequential limits. The choice $n_k = k^2$ gives $a_{n_k} = (-1)^{k^2} = (-1)^k$, since k^2 and k have the same parity, so a subsequence need not converge. Taking $n_k = k$ shows that a sequence is a subsequence of itself. One inequality is used throughout: $n_k \geq k$ for every k , by induction, since $n_1 \geq 1$ because $n_1 \in \mathbb{N}$, and $n_{k+1} > n_k \geq k$ forces $n_{k+1} \geq k + 1$ because no element of $\mathbb{N}$ lies strictly between k and $k + 1$.

Proposition 2.2.7: Subsequences of a Sequence with a Limit

Let $(a_n)_{n \geq 1}$ be a real sequence and let $(a_{n_k})_{k \geq 1}$ be a subsequence.

1. If (a_n) has limit $s \in [-\infty, +\infty]$, then (a_{n_k}) has limit s .
2. If two subsequences of (a_n) have different limits in $[-\infty, +\infty]$, then (a_n) has no limit in $[-\infty, +\infty]$, and in particular it does not converge.
3. A subsequence of (a_{n_k}) is a subsequence of (a_n) .

Proof:

1. Recall from the discussion after definition 2.2.6 that $n_k \geq k$ for every k .
Suppose s is real and $\epsilon > 0$. Choose N with $|a_n - s| < \epsilon$ for all $n \geq N$. For $k \geq N$ we have $n_k \geq k \geq N$, so $|a_{n_k} - s| < \epsilon$. Suppose $s = +\infty$ and let $M \in \mathbb{R}$. Choose N with $a_n > M$ for all $n \geq N$, and then $a_{n_k} > M$ for all $k \geq N$ by the same inequality $n_k \geq k \geq N$. The case $s = -\infty$ is identical with the inequality reversed.
2. Suppose (a_n) had limit s . By part (1) every subsequence would have limit s , so it is enough to know that a sequence cannot have two different limits in $[-\infty, +\infty]$, which is the case. Two different real limits are excluded by proposition 2.1.4. A real limit s and the limit $+\infty$ are incompatible: a convergent sequence is bounded by proposition 2.1.5, say $a_n \leq M$ for all n , and then $a_n > M$ holds for no n at all. The same argument with a lower bound excludes a real limit together with $-\infty$. Finally $+\infty$ and $-\infty$ are incompatible: taking $M = 0$ in both definitions produces an N beyond which $a_n > 0$ and $a_n < 0$ simultaneously, which is impossible. So (a_n) has no limit, and since a convergent sequence has a real limit, it does not converge.
3. Let $(a_{n_{k_j}})_{j \geq 1}$ be a subsequence of (a_{n_k}) , so that $k_1 < k_2 < \dots$ in $\mathbb{N}$. Since (n_k) is strictly increasing, $k_j < k_{j+1}$ gives $n_{k_j} < n_{k_{j+1}}$. Thus $j \mapsto n_{k_j}$ is a strictly increasing sequence in $\mathbb{N}$, which is what definition 2.2.6 requires, completing the proof.

Part (2) is the standard way to prove that a specific sequence diverges, and $(-1)^n$ is the smallest instance: its even and odd subsequences are constant with values 1 and -1 , so it has no limit. Parts (1) and (3) say that passing to a subsequence loses no limit and can be iterated, and both are used constantly below.

Deciding whether a given number is a subsequential limit by exhibiting a subsequence means constructing infinitely many indices, and the next result replaces that construction by a condition on the terms of the original sequence. It is the form in which subsequential limits are recognized in practice.

Proposition 2.2.8: Subsequential Limits Recognized Term by Term

Let $(a_n)_{n \geq 1}$ be a real sequence.

1. For $x \in \mathbb{R}$, the number x is a subsequential limit of (a_n) if and only if for every $\epsilon > 0$ the set $\{n \in \mathbb{N} \mid |a_n - x| < \epsilon\}$ is infinite.
2. The following three statements are equivalent: that $+\infty$ is a subsequential limit of (a_n) , that the sequence (a_n) is not bounded above, and that for every $M \in \mathbb{R}$ the set $\{n \in \mathbb{N} \mid a_n > M\}$ is infinite. The three corresponding statements obtained by replacing $+\infty$ by $-\infty$, “bounded above” by “bounded below” and $a_n > M$ by $a_n < M$ are also equivalent.

Proof:

1. Suppose $a_{n_k} \rightarrow x$ and let $\epsilon > 0$. There is a K with $|a_{n_k} - x| < \epsilon$ for all $k \geq K$. The indices $n_K < n_{K+1} < \dots$ are pairwise distinct elements of $\mathbb{N}$ lying in the set in question, so that set is infinite.

Conversely, suppose every such set is infinite. Define indices recursively, always taking the least admissible one so that no choice is made: let n_1 be the least n with $|a_n - x| < 1$, which exists because the set for $\epsilon = 1$ is infinite, hence nonempty, and $\mathbb{N}$ is well ordered. Given n_k , the set $\{n \mid |a_n - x| < 1/(k+1)\}$ is infinite, so it contains elements larger than n_k , and n_{k+1} is the least such element. Then $n_1 < n_2 < \dots$ and $|a_{n_k} - x| < 1/k$ for every k . Given $\epsilon > 0$, theorem 1.4.1 gives a K with $0 < 1/K < \epsilon$, and for $k \geq K$, proposition 1.1.2 gives $|a_{n_k} - x| < 1/k \leq 1/K < \epsilon$. So $a_{n_k} \rightarrow x$ and x is a subsequential limit.

2. Suppose $+\infty$ is a subsequential limit, say $a_{n_k} \rightarrow +\infty$, and let $M \in \mathbb{R}$. There is a K with $a_{n_k} > M$ for all $k \geq K$, and the indices $n_K < n_{K+1} < \dots$ are infinitely many, so $\{n \mid a_n > M\}$ is infinite. In particular it is nonempty for every M , so no real number bounds (a_n) above.

Suppose next that (a_n) is not bounded above and let $M \in \mathbb{R}$. Write $F = \{n \mid a_n > M\}$ and suppose F were finite. If F is empty then M bounds (a_n) above. If F is nonempty, then $\{a_n \mid n \in F\}$ is a finite nonempty set of real numbers and so has a largest element c , and $a_n \leq \max\{M, c\}$ for every n , since $n \notin F$ gives $a_n \leq M$ and $n \in F$ gives $a_n \leq c$. Either way (a_n) would be bounded above, contrary to hypothesis. So F is infinite.

Finally suppose $\{n \mid a_n > M\}$ is infinite for every $M \in \mathbb{R}$. Let n_1 be the least n with $a_n > 1$, and, given n_k , let n_{k+1} be the least element of $\{n \mid a_n > k+1\}$ exceeding n_k , which exists because that set is infinite. Then $a_{n_k} > k$ for every k . Given $M \in \mathbb{R}$, theorem 1.4.1 gives a $K \in \mathbb{N}$ with $K > M$, and for $k \geq K$ we get $a_{n_k} > k \geq K > M$. So $a_{n_k} \rightarrow +\infty$ and $+\infty$ is a subsequential limit. Together the implications just proved make the three statements equivalent.

For the statements with $-\infty$, put $b_n = -a_n$. A sequence of indices makes (a_{n_k}) tend to $-\infty$ if and only if it makes (b_{n_k}) tend to $+\infty$, directly from definition 2.2.4, and (a_n)

is bounded below if and only if (b_n) is bounded above, by proposition 1.1.2. Likewise $a_n < M$ if and only if $b_n > -M$. So the three statements for (a_n) and $-\infty$ are the three statements for (b_n) and $+\infty$, which have just been shown equivalent, completing the proof.

The condition in part (1) is that (a_n) visits every neighbourhood of x infinitely often, and the content of the proposition is that this weak-looking condition already builds the subsequence. Nothing about the sequence is assumed: in particular a subsequential limit need not be a value of the sequence, and a value taken infinitely often is a subsequential limit.

Existence is the remaining question. For a bounded sequence, repeated bisection of an interval containing every term locates a point that the sequence visits infinitely often, and proposition 2.2.8 then converts that point into a convergent subsequence.

Theorem 2.2.9: Bolzano-Weierstrass

1. Every bounded real sequence has a convergent subsequence.
2. Every real sequence has at least one subsequential limit in $[-\infty, +\infty]$.

Proof:

1. Let (a_n) be bounded, and choose real numbers $A \leq B$ with $A \leq a_n \leq B$ for every n .

Construct closed intervals $I_k = [A_k, B_k]$ recursively. Put $I_1 = [A, B]$, and note that $\{n | a_n \in I_1\} = \mathbb{N}$ is infinite. Given I_k with $\{n | a_n \in I_k\}$ infinite, let $C_k = (A_k + B_k)/2$ and consider the two halves $[A_k, C_k]$ and $[C_k, B_k]$. Their union is I_k , so

$$\{n | a_n \in I_k\} = \{n | a_n \in [A_k, C_k]\} \cup \{n | a_n \in [C_k, B_k]\}$$

and a union of two finite sets is finite, so at least one of the two sets on the right is infinite. Let I_{k+1} be the left half if the corresponding set is infinite, and the right half otherwise. This choice is canonical and keeps $\{n | a_n \in I_{k+1}\}$ infinite.

By construction $I_{k+1} \subseteq I_k$, each half has $A_k \leq C_k \leq B_k$ so that $A_k \leq B_k$ holds for every k , and $B_{k+1} - A_{k+1} = (B_k - A_k)/2$, so induction gives $B_k - A_k = (B - A)/2^{k-1} = 2(B - A) \cdot (1/2)^k$. The sequence $(1/2)^k$ is null by proposition 2.1.10, and a scalar multiple of a null sequence is null by the same result, so $B_k - A_k \rightarrow 0$. In particular, for every $\epsilon > 0$ there is a k with $B_k - A_k < \epsilon$. The hypotheses of theorem 1.4.8 are therefore met, and it gives a unique real number x with $x \in I_k$ for every k .

Let $\epsilon > 0$ and choose k with $B_k - A_k < \epsilon$. For every one of the infinitely many indices n with $a_n \in I_k$, both a_n and x lie in $[A_k, B_k]$, so $a_n - x \leq B_k - A_k$ and $x - a_n \leq B_k - A_k$, whence $|a_n - x| \leq B_k - A_k < \epsilon$ by proposition 1.3.8. So $\{n | |a_n - x| < \epsilon\}$ is infinite for every $\epsilon > 0$, and proposition 2.2.8 gives a subsequence converging to x .

2. If (a_n) is bounded, part (1) gives a real subsequential limit. If (a_n) is not bounded above, then $+\infty$ is a subsequential limit by proposition 2.2.8, and if (a_n) is not bounded below, then $-\infty$ is. Since a sequence that is bounded above and below is bounded, these three cases exhaust the possibilities, completing the proof.

The proof uses completeness twice, once through theorem 1.4.8 and once through the bisection that meets its hypotheses. Part (2) is what makes the following definition unconditional: the set of subsequential limits of a real sequence is never empty, so its supremum and infimum are attached to every sequence without exception.

Definition 2.2.10: Limit Superior and Limit Inferior of a Sequence

Let $(a_n)_{n \geq 1}$ be a real sequence and let $E \subseteq [-\infty, +\infty]$ be its set of subsequential limits. The *limit superior* and *limit inferior* of (a_n) are

$$\limsup_{n \rightarrow \infty} a_n = \sup E, \quad \liminf_{n \rightarrow \infty} a_n = \inf E$$

the supremum and infimum being taken in $[-\infty, +\infty]$ as in definition 2.2.4.

Both exist by proposition 2.2.5, and $E \neq \emptyset$ by theorem 2.2.9, so any $y \in E$ satisfies $\inf E \leq y \leq \sup E$ and therefore $\liminf_n a_n \leq \limsup_n a_n$ for every real sequence. The two quantities are exchanged by negation:

$$\limsup_{n \rightarrow \infty} (-a_n) = -\liminf_{n \rightarrow \infty} a_n, \quad \liminf_{n \rightarrow \infty} (-a_n) = -\limsup_{n \rightarrow \infty} a_n$$

To see this, write $E(a)$ for the set of subsequential limits of (a_n) . A subsequence of $(-a_n)$ is the negative of a subsequence of (a_n) with the same indices, and $a_{n_k} \rightarrow y$ if and only if $-a_{n_k} \rightarrow -y$, by theorem 2.1.6 when y is real and directly from definition 2.2.4 when $y = \pm\infty$. Hence $E(-a) = -E(a)$, and proposition 2.2.5 gives $\sup(-E(a)) = -\inf E(a)$, which is the first identity. The second follows by applying the first to $(-a_n)$.

Example 2.2.11: $(-1)^n(1 + 1/n)$, an Unbounded Interlacing, and an Enumeration of $\mathbb{Q}$

1. Let $a_n = (-1)^n(1 + 1/n)$. The sequences $(1/(2k))_k$ and $(1/(2k-1))_k$ are subsequences of $(1/n)_n$, which tends to 0 by theorem 1.4.1, so both tend to 0 by proposition 2.2.7. The even terms are $a_{2k} = 1 + 1/(2k)$, which therefore tends to 1 by theorem 2.1.6, and the odd terms are $a_{2k-1} = -(1 + 1/(2k-1))$, which tends to -1 by the same result. So $\{-1, 1\} \subseteq E$. The sequence is bounded, since $|a_n| = 1 + 1/n \leq 2$, so neither $+\infty$ nor $-\infty$ lies in E by proposition 2.2.8. If $y \in E$ is real, say $a_{n_k} \rightarrow y$, then infinitely many of the indices n_k are even or infinitely many are odd. In the first case the corresponding sub-subsequence is a subsequence of (a_n) by proposition 2.2.7 and converges to y by the same result, while its indices have the form $2m_1 < 2m_2 < \dots$, so it is also a subsequence of $(a_{2k})_k$ and converges to 1. Then $y = 1$ by proposition 2.1.4. The second case gives $y = -1$ in the same way. So $E = \{-1, 1\}$ and

$$\limsup_{n \rightarrow \infty} a_n = 1, \quad \liminf_{n \rightarrow \infty} a_n = -1$$

The sequence does not converge, by proposition 2.2.7. Note that the value 1 is taken by no term of the sequence, so an upper limit need not be a value.

2. Let $a_n = n$ for even n and $a_n = 1/n$ for odd n . The even terms are not bounded above, so $+\infty \in E$ by proposition 2.2.8, and every term is positive, so $-\infty \notin E$ by the same result. The odd terms satisfy $0 < a_{2k-1} = 1/(2k-1) \leq 1/k$, so $a_{2k-1} \rightarrow 0$ by theorem 2.1.7 and $0 \in E$. Suppose $y \in E$ is real, say $a_{n_k} \rightarrow y$. The subsequence is bounded by proposition 2.1.5, say $|a_{n_k}| \leq M$, and theorem 1.4.1 gives an $m \in \mathbb{N}$ with $m > M$. An even index n_k has $a_{n_k} = n_k \leq M < m$, and there are at most m elements of $\mathbb{N}$ below m , so only finitely many

n_k are even. Beyond some K all indices are odd, and then $0 < a_{n_k} = 1/n_k \leq 1/k$ using $n_k \geq k$, so $a_{n_k} \rightarrow 0$ by theorem 2.1.7 and $y = 0$ by proposition 2.1.4. Hence $E = \{0, +\infty\}$, and

$$\limsup_{n \rightarrow \infty} a_n = +\infty, \quad \liminf_{n \rightarrow \infty} a_n = 0$$

3. Since $\mathbb{Q}$ is countable, fix a bijection $n \mapsto q_n$ from $\mathbb{N}$ onto $\mathbb{Q}$. Every element of $[-\infty, +\infty]$ is a subsequential limit of (q_n) . For a real x and $\epsilon > 0$, the interval with endpoints x and $x + \epsilon$ contains infinitely many rationals: theorem 1.4.4 gives $p_1 \in \mathbb{Q}$ with $x < p_1 < x + \epsilon$, and, given p_j , gives $p_{j+1} \in \mathbb{Q}$ with $x < p_{j+1} < p_j$, so the p_j are pairwise distinct. As $n \mapsto q_n$ is injective, the set $\{n \mid |q_n - x| < \epsilon\}$ is infinite, and proposition 2.2.8 puts x in E . Moreover every element of $\mathbb{N}$ is a value of the sequence and $\mathbb{N}$ is not bounded above by theorem 1.4.1, so (q_n) is not bounded above, and the same argument with the negatives of the elements of $\mathbb{N}$ shows it is not bounded below. So $\pm\infty \in E$ as well, $E = [-\infty, +\infty]$, and

$$\limsup_{n \rightarrow \infty} q_n = +\infty, \quad \liminf_{n \rightarrow \infty} q_n = -\infty$$

As defined, the upper limit is the supremum of a set that has to be determined in full before the supremum can be taken, which is what the example did three times over. The next theorem replaces the definition by two properties that refer only to the terms a_n : the supremum is attained, and every real number above it dominates the sequence eventually. Those two properties determine the upper limit, so they may be used as its definition wherever that is more convenient, and it is in this form that section 2.4 and section 2.6 use it.

Theorem 2.2.12: The Upper Limit Is Attained and Dominates Eventually

Let $(a_n)_{n \geq 1}$ be a real sequence, let E be its set of subsequential limits, and put $s^* = \limsup_{n \rightarrow \infty} a_n$ and $s_* = \liminf_{n \rightarrow \infty} a_n$.

1. If $x \in \mathbb{R}$ and $x > s^*$, then there is an N with $a_n < x$ for every $n \geq N$.
2. $s^* \in E$ and $s_* \in E$.
3. If $x \in \mathbb{R}$ and $x < s^*$, then $a_n > x$ for infinitely many n .
4. s^* is the only $t \in [-\infty, +\infty]$ with both of the following properties: $t \in E$, and for every real $x > t$ there is an N with $a_n < x$ for all $n \geq N$.
5. For $s \in [-\infty, +\infty]$, the sequence (a_n) has limit s if and only if $s_* = s^* = s$.

Proof:

1. Suppose, for contradiction, that no such N exists. Then for every N there is an $n \geq N$ with $a_n \geq x$, so $S = \{n \mid a_n \geq x\}$ is infinite. Let $n_1 < n_2 < \dots$ enumerate S in increasing order, so that (a_{n_k}) is a subsequence with $a_{n_k} \geq x$ for every k . If (a_{n_k}) is bounded above, then it is bounded below by x as well, so theorem 2.2.9 gives a subsequence of it converging to some real y , and that subsequence is a subsequence of (a_n) by proposition 2.2.7, so $y \in E$ and $y \geq x$ by proposition 2.1.8 applied to it and the constant sequence with value x . If (a_{n_k}) is not bounded above, then proposition 2.2.8 gives a subsequence of it tending to $+\infty$, which is again a subsequence of (a_n) by proposition 2.2.7, so $y = +\infty$ lies in E and

$y \geq x$. Either way E contains an element $y \geq x > s^*$, contradicting $s^* = \sup E$.

2. Take s^* first, in three cases.

Suppose $s^* = +\infty$. If (a_n) were bounded above by some real M , then $+\infty \notin E$ by proposition 2.2.8, while a real $y \in E$ would satisfy $y \leq M$ by proposition 2.1.8 applied to a subsequence converging to y and the constant sequence with value M , and $-\infty \leq M$ holds outright. Then M would be an upper bound of E and $s^* \leq M$, which is false. So (a_n) is not bounded above, and $+\infty \in E$ by proposition 2.2.8.

Suppose $s^* = -\infty$. Every element of E is at most $\sup E = -\infty$, so $E \subseteq \{-\infty\}$, and $E \neq \emptyset$ by theorem 2.2.9. Hence $E = \{-\infty\}$ and $s^* \in E$.

Suppose s^* is real, and let $\epsilon > 0$. Since $s^* - \epsilon/2 < s^*$, the number $s^* - \epsilon/2$ is not an upper bound of E , so there is $y \in E$ with $y > s^* - \epsilon/2$. That inequality rules out $y = -\infty$, and $y \leq \sup E = s^*$ rules out $y = +\infty$, so y is real. By proposition 2.2.8 there are infinitely many n with $|a_n - y| < \epsilon/2$, and each such n satisfies

$$a_n > y - \frac{\epsilon}{2} > s^* - \epsilon$$

On the other hand part (1), applied to the real number $s^* + \epsilon > s^*$, gives an N with $a_n < s^* + \epsilon$ for all $n \geq N$. Removing the finitely many indices below N from an infinite set leaves an infinite set, so there are infinitely many n with $s^* - \epsilon < a_n < s^* + \epsilon$, that is, with $|a_n - s^*| < \epsilon$. As $\epsilon > 0$ was arbitrary, proposition 2.2.8 gives $s^* \in E$.

For s_* , apply what has just been proved to $(-a_n)$: its upper limit is $-s_*$ by the identities following definition 2.2.10, so $-s_* \in E(-a) = -E(a)$, and hence $s_* \in E$.

3. Let $x \in \mathbb{R}$ with $x < s^*$. By part (2), $s^* \in E$. If s^* is real, apply proposition 2.2.8 with $\epsilon = s^* - x > 0$: infinitely many n satisfy $|a_n - s^*| < \epsilon$, and each of them satisfies $a_n > s^* - \epsilon = x$. If $s^* = +\infty$, then proposition 2.2.8 gives directly that $\{n | a_n > x\}$ is infinite. The case $s^* = -\infty$ does not arise, since no real x is smaller than $-\infty$.
4. That s^* has both properties is parts (2) and (1). Suppose t has both properties. Since $t \in E$, $t \leq \sup E = s^*$. If $t < s^*$, then proposition 2.2.5 gives a real x with $t < x < s^*$. The second property of t gives an N with $a_n < x$ for all $n \geq N$, while part (3) gives infinitely many n with $a_n > x$, and infinitely many of those n are at least N , which is a contradiction. Hence $t = s^*$.
5. Suppose (a_n) has limit s . Every subsequence has limit s by proposition 2.2.7, so every element of E equals s , and $s \in E$ because (a_n) is a subsequence of itself. Thus $E = \{s\}$ and $s_* = s^* = s$.

Conversely suppose $s_* = s^* = s$. Consider first s real, and let $\epsilon > 0$. Part (1) applied to $s + \epsilon > s^*$ gives an N_1 with $a_n < s + \epsilon$ for $n \geq N_1$. Applying part (1) to the sequence $(-a_n)$, whose upper limit is $-s_* = -s$, and to the real number $-s + \epsilon > -s$, gives an N_2 with $-a_n < -s + \epsilon$, that is $a_n > s - \epsilon$, for $n \geq N_2$. For $n \geq \max\{N_1, N_2\}$ both hold, so $|a_n - s| < \epsilon$ and $a_n \rightarrow s$.

Consider next $s = +\infty$. Every element of E is at least $\inf E = s_* = +\infty$, and $E \neq \emptyset$ by theorem 2.2.9, so $E = \{+\infty\}$. Suppose $a_n \rightarrow +\infty$ failed. Then there is an $M \in \mathbb{R}$

such that for every N some $n \geq N$ has $a_n \leq M$, so $\{n | a_n \leq M\}$ is infinite and the corresponding subsequence (a_{n_k}) satisfies $a_{n_k} \leq M$. If that subsequence is bounded below, theorem 2.2.9 gives a further subsequence converging to a real number, which lies in E by proposition 2.2.7, contradicting $E = \{+\infty\}$. If it is not bounded below, proposition 2.2.8 gives a further subsequence tending to $-\infty$, so $-\infty \in E$, again a contradiction. So $a_n \rightarrow +\infty$.

Finally, if $s = -\infty$ then the sequence $(-a_n)$ has $\limsup(-a_n) = -s_* = +\infty$ and $\liminf(-a_n) = -s^* = +\infty$, so $-a_n \rightarrow +\infty$ by the case just settled, and therefore $a_n \rightarrow -\infty$ by definition 2.2.4, completing the proof.

Parts (1) and (3) together say that s^* separates the real line into the numbers the sequence eventually stays below and the numbers it exceeds infinitely often, and part (4) says that this separation determines s^* . Part (5) turns a question about the existence of a limit in $[-\infty, +\infty]$, which quantifies over a bound and an index, into an equation between two numbers that exist for every sequence. The corresponding statements for s_* hold with the inequalities reversed, by applying the theorem to $(-a_n)$ and using the identities that follow definition 2.2.10.

The last two properties compare upper limits of two sequences. The first is an inequality that passes to the limit in both directions, while the second holds in one direction only.

Proposition 2.2.13: Comparison and Subadditivity for Upper Limits

Let $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ be real sequences.

1. If there is an N with $a_n \leq b_n$ for all $n \geq N$, then

$$\limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} b_n \quad \text{and} \quad \liminf_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} b_n$$

2. If the sum $\limsup_n a_n + \limsup_n b_n$ is defined in $[-\infty, +\infty]$, then

$$\limsup_{n \rightarrow \infty} (a_n + b_n) \leq \limsup_{n \rightarrow \infty} a_n + \limsup_{n \rightarrow \infty} b_n$$

Proof:

Write $s^* = \limsup_n a_n$ and $t^* = \limsup_n b_n$.

1. Suppose $s^* > t^*$. By proposition 2.2.5 there is a real x with $t^* < x < s^*$. Since $x > t^*$, theorem 2.2.12 gives an N_1 with $b_n < x$ for all $n \geq N_1$. Since $x < s^*$, theorem 2.2.12 gives infinitely many n with $a_n > x$, and infinitely many of those satisfy $n \geq \max\{N, N_1\}$. For such an n , $a_n > x > b_n$ while $a_n \leq b_n$, a contradiction. Hence $s^* \leq t^*$.

For the lower limits, the hypothesis gives $-b_n \leq -a_n$ for $n \geq N$, so the case just proved applied to $(-b_n)$ and $(-a_n)$ gives $\limsup_n (-b_n) \leq \limsup_n (-a_n)$. By the identities following definition 2.2.10 this reads $-\liminf_n b_n \leq -\liminf_n a_n$, and negating reverses the inequality by proposition 1.1.2, giving $\liminf_n a_n \leq \liminf_n b_n$.

2. If $s^* + t^* = +\infty$ there is nothing to prove, since every element of $[-\infty, +\infty]$ is at most $+\infty$.

Suppose $s^* + t^* = -\infty$. Then at least one summand is $-\infty$ and, the sum being defined, neither is $+\infty$. The statement is symmetric in (a_n) and (b_n) , so assume $s^* = -\infty$ and $t^* < +\infty$. Every element of the set of subsequential limits of (a_n) is at most $s^* = -\infty$, and that set is nonempty by theorem 2.2.9, so $\liminf_n a_n = -\infty$ as well, and $a_n \rightarrow -\infty$ by theorem 2.2.12. Since $t^* < +\infty$, proposition 2.2.5 gives a real x with $t^* < x < +\infty$, and theorem 2.2.12 gives an N_1 with $b_n < x$ for $n \geq N_1$. Let $M \in \mathbb{R}$. Since $a_n \rightarrow -\infty$ there is an N_2 with $a_n < M - x$ for $n \geq N_2$, and then $a_n + b_n < (M - x) + x = M$ for $n \geq \max\{N_1, N_2\}$. So $a_n + b_n \rightarrow -\infty$, and theorem 2.2.12 gives $\limsup_n (a_n + b_n) = -\infty = s^* + t^*$.

Suppose finally that $s^* + t^*$ is real, which forces both s^* and t^* to be real. Let $\epsilon > 0$. By theorem 2.2.12 applied to $s^* + \epsilon/2 > s^*$ there is an N_1 with $a_n < s^* + \epsilon/2$ for $n \geq N_1$, and applied to $t^* + \epsilon/2 > t^*$ there is an N_2 with $b_n < t^* + \epsilon/2$ for $n \geq N_2$. Put $u = s^* + t^* + \epsilon$, a real number. For $n \geq \max\{N_1, N_2\}$,

$$a_n + b_n < \left(s^* + \frac{\epsilon}{2}\right) + \left(t^* + \frac{\epsilon}{2}\right) = u$$

If $\limsup_n (a_n + b_n) > u$ held, then theorem 2.2.12 would give infinitely many n with $a_n + b_n > u$, and infinitely many of those would satisfy $n \geq \max\{N_1, N_2\}$, contradicting $a_n + b_n < u$. So $\limsup_n (a_n + b_n) \leq u$ for every $\epsilon > 0$. In particular $\limsup_n (a_n + b_n) \neq +\infty$. If it equals $-\infty$ the claimed inequality holds. Otherwise it is real, and $\limsup_n (a_n + b_n) \leq s^* + t^* + \epsilon$ for every $\epsilon > 0$ gives $\limsup_n (a_n + b_n) \leq s^* + t^*$ by proposition 1.4.7, completing the proof.

The inequality in part (2) can be strict, and the reason is that the two sequences may reach their upper limits along different sets of indices. Take $a_n = (-1)^n(1 + 1/n)$ and $b_n = -a_n$. By example 2.2.11, $\limsup_n a_n = 1$, and by the identities following definition 2.2.10, $\limsup_n b_n = -\liminf_n a_n = 1$, so the right-hand side of the inequality is 2. But $a_n + b_n = 0$ for every n , so every subsequence of the sum is constant with value 0, its set of subsequential limits is $\{0\}$, and $\limsup_n (a_n + b_n) = 0 < 2$. The same pair refutes the reverse inequality, which would read $0 \geq 2$.

Exercise 2.2.1

1. Let (a_n) be increasing and bounded above, with limit s . Show that $s = \sup\{a_n \mid n \geq 1\}$, and that (a_n) is eventually constant if and only if $s = a_N$ for some N .
2. Call an index n a *peak* of a real sequence (a_n) if $a_m \leq a_n$ for every $m > n$. Show that every real sequence has a monotone subsequence, by treating separately the case of infinitely many peaks and the case of finitely many. Deduce part (1) of theorem 2.2.9 from theorem 2.2.2.
3. For a real sequence (a_n) and $N \geq 1$ put $t_N = \sup\{a_n \mid n \geq N\}$, the supremum being taken in $[-\infty, +\infty]$. Show that

$$\limsup_{n \rightarrow \infty} a_n = \inf\{t_N \mid N \geq 1\}$$

and state and prove the corresponding formula for the lower limit. (Suggestion: verify that the right-hand side has the two properties in part (4) of theorem 2.2.12.)

4. Determine the set of subsequential limits, the upper limit and the lower limit of each of the sequences $a_n = (-1)^n + 1/n$ and $b_n = n(1 + (-1)^n)$, with proof.

5. Let (a_n) be a bounded real sequence with the property that every convergent subsequence of (a_n) has the same limit L . Show that $a_n \rightarrow L$. Show also that the boundedness hypothesis cannot be dropped, by exhibiting an unbounded sequence with no convergent subsequence at all.
6. Show that $\liminf_n (a_n + b_n) \geq \liminf_n a_n + \liminf_n b_n$ whenever the right-hand side is defined in $[-\infty, +\infty]$, and exhibit bounded sequences for which this inequality and the one in part (2) of proposition 2.2.13 are both strict.
7. Let (a_n) be a real sequence with $a_n > 0$ for every n . Show that $\liminf_n a_n \geq 0$, and that $\limsup_n a_n = 0$ forces $a_n \rightarrow 0$.
8. Show that an increasing real sequence that is not bounded above has limit $+\infty$, and conclude that every monotone real sequence has a limit in $[-\infty, +\infty]$. Show also that this fails without monotonicity, by exhibiting a sequence with no limit in $[-\infty, +\infty]$.

2.3 The Cauchy Criterion

The definition of convergence in definition 2.1.2 mentions the limit, so to apply it one must already know the number the sequence approaches. A sequence produced by a recursion, or by adding one more term at each stage, arrives without a candidate limit, and for such a sequence the definition cannot be used as a test. This section replaces it by a condition on the terms alone: past some index, any two terms differ by less than ϵ . The condition can be checked from the sequence itself, and in $\mathbb{R}$ it is equivalent to convergence.

In $\mathbb{Q}$ it is not. The decimal truncations of $\sqrt{2}$ come within any prescribed tolerance of one another and converge to no rational number, which is the failure section 1.1 was built to repair. So the equivalence proved below is a property of $\mathbb{R}$, carrying content beyond the definition, and it is the form in which the completeness of $\mathbb{R}$ is used for the rest of the book.

Definition 2.3.1: Cauchy Sequence in $\mathbb{R}$

A sequence $(a_n)_{n \geq 1}$ of real numbers is a *Cauchy sequence* if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon \quad \text{for all } n, m \geq N$$

Definition 1.1.4 imposed the same condition on a sequence of rationals, with ϵ ranging over the positive rationals, because at that point no positive real was available. For a sequence of rationals the two conditions agree. If the real condition holds then so does the rational one, since every positive rational is a positive real. Conversely, suppose the rational condition holds and let $\epsilon > 0$ be real. By theorem 1.4.4 applied to $0 < \epsilon$ there is a rational p with $0 < p < \epsilon$, and the index N that works for p works for ϵ . Nothing below distinguishes the two, and the phrase *Cauchy sequence* refers to the real condition from here on.

Example 2.3.2: Decimal Truncations of $\sqrt{2}$, $((-1)^n)$ and $(\sqrt{n})$

Throughout, $\sqrt{x}$ for $x \geq 0$ is the number given by theorem 1.3.12. That theorem also gives the monotonicity used repeatedly below: if $0 \leq u < v$ then $\sqrt{u} < \sqrt{v}$, since $\sqrt{u} \geq \sqrt{v} \geq 0$ would force $u = (\sqrt{u})^2 \geq (\sqrt{v})^2 = v$.

1. Let $\alpha = \sqrt{2}$ and, for $n \geq 1$, let

$$a_n = \frac{\lfloor 10^n \alpha \rfloor}{10^n}$$

By proposition 1.4.3, $\lfloor 10^n \alpha \rfloor$ is the unique integer m with $m \leq 10^n \alpha < m + 1$, so each a_n is rational and, dividing by $10^n > 0$,

$$0 \leq \alpha - a_n < 10^{-n}$$

For $m \geq n$ the triangle inequality proposition 1.3.8(4) gives

$$|a_m - a_n| \leq |a_m - \alpha| + |\alpha - a_n| < 10^{-m} + 10^{-n} \leq 2 \cdot 10^{-n}$$

By proposition 2.1.10 in the case $j = 0$ and $r = 1/10$, the sequence 10^{-n} is null, so given $\epsilon > 0$ there is an N with $10^{-N} < \epsilon/2$, and then $|a_m - a_n| < \epsilon$ for all $m, n \geq N$. So (a_n) is Cauchy. The same estimate with ϵ rational shows it is Cauchy in the sense of definition 1.1.4 as well.

The bound $0 \leq \alpha - a_n < 10^{-n}$ also gives $a_n \rightarrow \alpha$. If some rational q satisfied $a_n \rightarrow q$, then $q = \alpha$ by proposition 2.1.4, so $q^2 = 2$ with $q \in \mathbb{Q}$, which contradicts proposition 1.1.3. This Cauchy sequence of rationals therefore has no rational limit. Note also that $0 \leq a_n \leq \alpha < 2$, the last inequality because $\alpha^2 = 2 < 4 = 2^2$, so the sequence is bounded.

2. Let $a_n = (-1)^n$, so that $|a_n| = 1$ for every n . Take $\epsilon = 1$. Given any N , the indices $n = N$ and $m = N + 1$ satisfy $|a_N - a_{N+1}| = 2$, which is not less than 1. So no N works for $\epsilon = 1$, and (a_n) is a bounded sequence that is not Cauchy.
3. Let $a_n = \sqrt{n}$. Consecutive terms come together: from $(\sqrt{n+1} - \sqrt{n})(\sqrt{n+1} + \sqrt{n}) = (n+1) - n = 1$ and $\sqrt{n+1} + \sqrt{n} \geq 2\sqrt{n} > 0$,

$$0 < \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} \leq \frac{1}{2\sqrt{n}}$$

Given $\epsilon > 0$, theorem 1.4.1(2) provides an integer $n_0 > 1/(4\epsilon^2)$, and for $n \geq n_0$ the monotonicity of the square root gives $\sqrt{n} > 1/(2\epsilon)$, hence $|a_{n+1} - a_n| < \epsilon$. The sequence is not Cauchy all the same. Take $\epsilon = 1$ and let N be given. By theorem 1.4.1(2) there is an integer $m > (\sqrt{N} + 1)^2$, and then $\sqrt{m} > \sqrt{N} + 1$, so $|a_m - a_N| > 1$.

Item (3) is why the definition quantifies over all pairs of large indices. Consecutive pairs give a weaker requirement, which $\sqrt{n}$ satisfies while being unbounded. Item (2) shows that boundedness is not sufficient either. Item (1) is the case the rest of the section is about: a sequence of rationals whose limit exists in $\mathbb{R}$ and not in $\mathbb{Q}$.

One of the two implications between convergence and the Cauchy condition follows from the triangle inequality alone.

Proposition 2.3.3: Convergent Sequences Are Cauchy

Every convergent sequence of real numbers is a Cauchy sequence.

Proof:

Let $a_n \rightarrow a$ and let $\epsilon > 0$. Applying definition 2.1.2 with $\epsilon/2 > 0$ gives an $N \in \mathbb{N}$ with $|a_n - a| < \epsilon/2$ for all $n \geq N$. For $n, m \geq N$,

$$|a_n - a_m| = |(a_n - a) + (a - a_m)| \leq |a_n - a| + |a_m - a| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

using proposition 1.3.8(4) for the inequality and proposition 1.3.8(1) for $|a - a_m| = |a_m - a|$. The index N therefore satisfies the requirement of definition 2.3.1 for ϵ , as we sought to show.

The converse is the substantial direction, and it needs one preparation: a Cauchy sequence is bounded, which is what makes Bolzano-Weierstrass applicable to it. Lemma 1.1.6 is the same statement for sequences of rationals, proved there because the construction of $\mathbb{R}$ needed it before real ϵ existed.

Proposition 2.3.4: Cauchy Sequences in $\mathbb{R}$ Are Bounded

Every Cauchy sequence $(a_n)_{n \geq 1}$ of real numbers is bounded: there is an $M \geq 0$ with $|a_n| \leq M$ for all n .

Proof:

Apply definition 2.3.1 with $\epsilon = 1$ to get an $N \in \mathbb{N}$ with $|a_n - a_m| < 1$ for all $n, m \geq N$. In particular, for $n \geq N$,

$$|a_n| = |(a_n - a_N) + a_N| \leq |a_n - a_N| + |a_N| < 1 + |a_N|$$

by proposition 1.3.8(4). Let M be the largest of the finitely many numbers

$$|a_1|, |a_2|, \dots, |a_{N-1}|, 1 + |a_N|$$

Then $|a_n| \leq M$ for $n < N$ because $|a_n|$ is one of the listed numbers, and $|a_n| < 1 + |a_N| \leq M$ for $n \geq N$ by the displayed estimate. Since $M \geq |a_1| \geq 0$, this is the required bound, completing the proof.

The converse of proposition 2.3.4 fails, by example 2.3.2(2). What boundedness gives is a convergent subsequence, and the Cauchy condition is the extra information that forces the whole sequence to converge to the limit of that subsequence.

Theorem 2.3.5: The Cauchy Criterion

A sequence of real numbers converges if and only if it is a Cauchy sequence.

Proof:

One direction is proposition 2.3.3. For the other, let $(a_n)_{n \geq 1}$ be a Cauchy sequence. By proposition 2.3.4 it is bounded, which is the hypothesis of theorem 2.2.9, so it has a convergent subsequence $(a_{n_k})_{k \geq 1}$, say $a_{n_k} \rightarrow a$. The indices satisfy $n_k \geq k$ for every k , by induction on k : the base case is $n_1 \geq 1$, and $n_{k+1} > n_k$ with both integers gives $n_{k+1} \geq n_k + 1 \geq k + 1$.

Let $\epsilon > 0$. The Cauchy condition gives an $N \in \mathbb{N}$ with $|a_n - a_m| < \epsilon/2$ for all $n, m \geq N$, and convergence of the subsequence gives a $K \in \mathbb{N}$ with $|a_{n_k} - a| < \epsilon/2$ for all $k \geq K$. Fix one index

$k = \max\{K, N\}$. Then $k \geq K$, and $n_k \geq k \geq N$, so for every $n \geq N$,

$$|a_n - a| \leq |a_n - a_{n_k}| + |a_{n_k} - a| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

using proposition 1.3.8(4). Hence $a_n \rightarrow a$, and the sequence converges, as we sought to show.

Theorem 2.3.5 makes convergence provable without producing the limit, and that is what later sections use it for: a sequence given by a recursion, or by adding one more term at each stage, can be tested on its terms alone. It is not the only result that gives a limit without naming it. Theorem 2.2.2 does the same for a bounded monotone sequence, and for the increasing partial sums of a series of nonnegative terms that is the route section 2.4 takes.

The condition in definition 2.3.1 refers to $|a_n - a_m|$, the distance between two terms, and to no other feature of $\mathbb{R}$. Replacing that distance by an abstract one gives a Cauchy condition on any set equipped with a notion of distance, and such a set is called *Cauchy-complete* when every Cauchy sequence in it converges. The qualifier matters, because section 1.5★ has already fixed *complete* for an ordered field to mean the least upper bound property, and the two senses are not equivalent in general: it is exactly the Archimedean hypothesis that closes the gap, which is why it appears in theorem 2.3.6. Theorem 2.3.5 says that $\mathbb{R}$ is *Cauchy-complete*. That it is complete in the order sense is theorem 1.2.4. The general theory belongs to the companion volume *Metric and Function Spaces*, and no argument in this book uses it.

Five statements have now been proved about $\mathbb{R}$, each of them a way of saying that $\mathbb{R}$ has no gaps: the least upper bound property theorem 1.2.4, monotone convergence theorem 2.2.2, the nested interval theorem theorem 1.4.8, Bolzano-Weierstrass theorem 2.2.9, and the Cauchy criterion. Inside $\mathbb{R}$ each of them is a theorem, so an implication among them is true for the trivial reason that its conclusion has already been proved. The content appears one level up, where the five are treated as conditions an Archimedean ordered field may or may not satisfy: for such a field each one implies the other four, and which to install as the axiom is a matter of preference. The theorem below is therefore stated for an arbitrary Archimedean ordered field, in the sense of definition 1.1.1, and that hypothesis is not removable.

Fix an Archimedean ordered field F . Absolute value in F is given by the formula in definition 1.1.1, and the definitions of convergence, of the Cauchy condition, of boundedness and of monotonicity read verbatim in F , with ϵ ranging over the positive elements of F and with a positive integer n read as $n \cdot 1 \in F$. For $a \leq b$ in F write $[a, b] = \{x \in F \mid a \leq x \leq b\}$.

Theorem 2.3.6: Equivalent Forms of Completeness

Let F be an Archimedean ordered field. The following five statements are equivalent.

1. Every nonempty subset of F that is bounded above has a supremum in F .
2. Every bounded monotone sequence in F converges.
3. If $[a_n, b_n]$ for $n \geq 1$ are intervals with $a_n \leq b_n$ and $[a_{n+1}, b_{n+1}] \subseteq [a_n, b_n]$ for every n , then $\bigcap_{n \geq 1} [a_n, b_n]$ is nonempty.
4. Every bounded sequence in F has a convergent subsequence.
5. Every Cauchy sequence in F converges.

Proof:

The cycle $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)$ is proved. Two observations are used more than once.

Observation A (halving). For every $\ell > 0$ and every $\epsilon > 0$ in F there is a positive integer k with $\ell \cdot (2^k)^{-1} < \epsilon$, where $2 = 1 + 1$. Indeed $2^k \geq k + 1$ for every $k \geq 0$, by induction: the case $k = 0$ reads $1 \geq 1$, and $2^{k+1} = 2^k + 2^k \geq (k + 1) + 1$ because $2^k \geq 1$. Next, $\epsilon^{-1} > 0$, since $\epsilon^{-1} < 0$ would give $1 = \epsilon \cdot \epsilon^{-1} < 0$ by proposition 1.1.2(5), contradicting $1 > 0$ from proposition 1.1.2(6). So $\ell\epsilon^{-1} > 0$, and the Archimedean property gives a positive integer k with $\ell\epsilon^{-1} < k$. Then $2^k \geq k + 1 > \ell\epsilon^{-1} > 0$, so $(2^k)^{-1} < (\ell\epsilon^{-1})^{-1} = \epsilon\ell^{-1}$ by proposition 1.1.2(7), and multiplying by $\ell > 0$ (proposition 1.1.2(5)) gives $\ell(2^k)^{-1} < \epsilon$.

Observation B (width). If $u, v \in [a, b]$ then $|u - v| \leq b - a$. Indeed $a \leq u \leq b$ and $a \leq v \leq b$ give both $u - v \leq b - a$ and $v - u \leq b - a$, and $|u - v|$ is one of these two numbers.

Step 1: (1) implies (2). Let (x_n) be increasing and bounded. The set $E = \{x_n | n \geq 1\}$ is nonempty and bounded above, so by (1) it has a supremum s . Let $\epsilon > 0$. Since $s - \epsilon < s$ and s is the least upper bound, $s - \epsilon$ is not an upper bound of E , so there is an N with $x_N > s - \epsilon$. For $n \geq N$ monotonicity gives $s - \epsilon < x_N \leq x_n \leq s$, hence $|x_n - s| = s - x_n < \epsilon$. So $x_n \rightarrow s$. If instead (x_n) is decreasing and bounded, then $(-x_n)$ is increasing and bounded, so $-x_n \rightarrow t$ for some $t \in F$. The elements $x_n - (-t) = x_n + t$ and $(-x_n) - t = -(x_n + t)$ have the same absolute value, so $|x_n - (-t)| < \epsilon$ whenever $|(-x_n) - t| < \epsilon$, and (x_n) converges to $-t$.

Step 2: (2) implies (3). Let the intervals $[a_n, b_n]$ be nested. Since a_{n+1} and b_{n+1} both lie in $[a_n, b_n]$, the sequence (a_n) is increasing and (b_n) is decreasing. Moreover $a_n \leq b_m$ for all n and m : for $n \geq m$ nesting gives $a_n \leq b_n \leq b_m$, and for $n < m$ it gives $a_n \leq a_m \leq b_m$. In particular (a_n) is bounded above by b_1 and below by a_1 , so by (2) it converges, say $a_n \rightarrow a$. Fix m . If $a < a_m$, then putting $\epsilon = a_m - a > 0$, every $n \geq m$ has $a_n \geq a_m > a$ and hence $|a_n - a| = a_n - a \geq \epsilon$, contradicting convergence. If $a > b_m$, then putting $\epsilon = a - b_m > 0$, some n has $|a_n - a| < \epsilon$ and hence $a_n > a - \epsilon = b_m$, contradicting $a_n \leq b_m$. So $a_m \leq a \leq b_m$ for every m , and a lies in the intersection.

Step 3: (3) implies (4). Let (x_n) be bounded, say $|x_n| \leq M$ for all n , where $M > 0$ after replacing M by $M + 1$ if necessary. Then $x_n \in I_0 = [-M, M]$ for all n , by the definition of absolute value in definition 1.1.1. Intervals $I_k = [a_k, b_k]$ are built so that $a_k < b_k$, so that $I_{k+1} \subseteq I_k$, and so that $x_n \in I_k$ for infinitely many indices n . The set $\{n | x_n \in I_0\}$ is all of $\mathbb{N}$, so I_0 starts the construction. Given I_k , put $c = (a_k + b_k)(1 + 1)^{-1}$, so that $a_k < c < b_k$ by proposition 1.1.2(8). If $\{n | x_n \in [a_k, c]\}$ is infinite, set $I_{k+1} = [a_k, c]$, and otherwise set $I_{k+1} = [c, b_k]$. In the second case $\{n | x_n \in [c, b_k]\}$ is infinite, because $\{n | x_n \in I_k\}$ is the union of the two index sets and is infinite, while a union of two finite sets is finite. By construction $b_{k+1} - a_{k+1} = (b_k - a_k)(1 + 1)^{-1}$, so $b_k - a_k = (b_0 - a_0)(2^k)^{-1}$ for every k .

By (3) there is an $x \in \bigcap_{k \geq 0} I_k$. Define indices recursively: let n_1 be the least n with $x_n \in I_1$, and let n_{k+1} be the least $n > n_k$ with $x_n \in I_{k+1}$. Each exists because each index set is infinite, so the relevant set of positive integers is nonempty and has a least element. Both x_{n_k} and x lie in I_k , so Observation B gives $|x_{n_k} - x| \leq b_k - a_k = (b_0 - a_0)(2^k)^{-1}$. Let $\epsilon > 0$. Observation A gives a K with $(b_0 - a_0)(2^K)^{-1} < \epsilon$. For $k \geq K$ one has $2^k \geq 2^K > 0$, since each step from 2^j to $2^{j+1} = 2^j + 2^j$ adds a positive element, so $(2^k)^{-1} \leq (2^K)^{-1}$ by proposition 1.1.2(7) and therefore $|x_{n_k} - x| < \epsilon$. Hence $x_{n_k} \rightarrow x$.

Step 4: (4) implies (5). Let (x_n) be Cauchy. Taking $\epsilon = 1$ gives an N with $|x_n - x_N| < 1$ for $n \geq N$, so $|x_n| \leq |x_n - x_N| + |x_N| < 1 + |x_N|$ for those n by proposition 1.1.2(9), and (x_n) is

bounded by the largest of $|x_1|, \dots, |x_{N-1}|, 1 + |x_N|$. By (4) some subsequence converges, say $x_{n_k} \rightarrow x$, and $n_k \geq k$ for every k by induction on k , from $n_1 \geq 1$ and $n_{k+1} \geq n_k + 1 \geq k + 1$. Let $\epsilon > 0$, choose N_1 with $|x_n - x_m| < \epsilon(1+1)^{-1}$ for $n, m \geq N_1$, choose K with $|x_{n_k} - x| < \epsilon(1+1)^{-1}$ for $k \geq K$, and fix $k = \max\{K, N_1\}$, so that $n_k \geq k \geq N_1$. For $n \geq N_1$,

$$|x_n - x| \leq |x_n - x_{n_k}| + |x_{n_k} - x| < \epsilon$$

by proposition 1.1.2(9). Hence $x_n \rightarrow x$.

Step 5: (5) implies (1). Let $E \subseteq F$ be nonempty and bounded above, say by b_0 . Pick $x_0 \in E$ and put $a_0 = x_0 - 1$, so that $a_0 < x_0$ and a_0 is not an upper bound of E , while $a_0 < x_0 \leq b_0$. Build a_k and b_k recursively so that a_k is never an upper bound of E , b_k always is, and $a_k < b_k$: given the pair (a_k, b_k) , put $c = (a_k + b_k)(1+1)^{-1}$, which satisfies $a_k < c < b_k$ by proposition 1.1.2(8), and set $(a_{k+1}, b_{k+1}) = (a_k, c)$ if c is an upper bound of E and $(a_{k+1}, b_{k+1}) = (c, b_k)$ otherwise. Then (a_k) is increasing, (b_k) is decreasing, and $b_k - a_k = (b_0 - a_0)(2^k)^{-1}$.

The sequence (b_k) is Cauchy. For $m \geq k$ one has $a_k \leq a_m < b_m \leq b_k$, so b_k and b_m both lie in $[a_k, b_k]$ and Observation B gives $|b_k - b_m| \leq b_k - a_k = (b_0 - a_0)(2^k)^{-1}$. Given $\epsilon > 0$, Observation A gives a K with $(b_0 - a_0)(2^K)^{-1} < \epsilon$, and for $m \geq k \geq K$ the monotonicity of $(2^k)^{-1}$ recorded in Step 3 gives $|b_k - b_m| \leq (b_0 - a_0)(2^k)^{-1} \leq (b_0 - a_0)(2^K)^{-1} < \epsilon$. By (5) the sequence converges, say $b_k \rightarrow s$.

The sequence (a_k) converges to s as well. Let $\epsilon > 0$. Observation A and the monotonicity just used give a K_1 with $b_k - a_k < \epsilon(1+1)^{-1}$ for all $k \geq K_1$, and convergence gives a K_2 with $|b_k - s| < \epsilon(1+1)^{-1}$ for all $k \geq K_2$. For $k \geq \max\{K_1, K_2\}$,

$$|a_k - s| \leq |a_k - b_k| + |b_k - s| < \epsilon$$

by proposition 1.1.2(9).

It remains to identify s as $\sup E$. Suppose some $x \in E$ had $x > s$. Put $\epsilon = x - s > 0$ and choose k with $|b_k - s| < \epsilon$. Then $b_k < s + \epsilon = x$, contradicting that b_k is an upper bound of E , so s is an upper bound. Now suppose some upper bound u had $u < s$. Put $\epsilon = s - u > 0$ and choose k with $|a_k - s| < \epsilon$. Then $a_k > s - \epsilon = u$, and since a_k is not an upper bound of E there is an $x \in E$ with $x > a_k > u$, contradicting that u is an upper bound. So s is the least upper bound of E , which is (1) and closes the cycle, completing the proof.

The Archimedean hypothesis enters the proof at exactly two places, in Step 3 and in Step 5, and at both through Observation A, which is what makes an interval of a fixed width halve below a prescribed ϵ after finitely many steps. It is a hypothesis and not a convenience of this particular argument. An ordered field satisfying (1) is Archimedean by corollary 1.5.1, so a version of the theorem for ordered fields at large would have to produce the Archimedean property out of (5) alone. That cannot be done, because there are non-Archimedean ordered fields in which every Cauchy sequence converges, the formal Laurent series in a positive infinitesimal being one such field. Statement (5) therefore does not imply statement (1) once the hypothesis is dropped.

A single failure in $\mathbb{Q}$ now accounts for five. The field $\mathbb{Q}$ is an ordered field and is Archimedean, these being among the hypotheses on $\mathbb{Q}$ recorded in section 1.1, so theorem 2.3.6 applies to it. By example 2.3.2(1) there is a Cauchy sequence of rationals with no rational limit, so $\mathbb{Q}$ fails (5), and therefore fails the other four as well. It has bounded sets of rationals with no rational supremum, which is what example 1.2.3(3) exhibited by hand, and it also has bounded monotone sequences that do not converge, nested rational intervals with empty intersection, and bounded sequences with no

subsequence converging in $\mathbb{Q}$. These are four descriptions of the one defect, and theorem 2.3.6 is what identifies them.

Exercise 2.3.1

1. Let (a_n) and (b_n) be Cauchy sequences of real numbers. Prove directly from definition 2.3.1, without using theorem 2.3.5 or theorem 2.1.6, that $(a_n + b_n)$ and $(a_n b_n)$ are Cauchy sequences.
2. Let (a_n) be a Cauchy sequence of real numbers. Prove that $(|a_n|)$ is a Cauchy sequence, and show by an example that the converse fails: there is a sequence (a_n) such that $(|a_n|)$ is Cauchy and (a_n) is not.
3. Let $0 < r < 1$ be fixed and let $(a_n)_{n \geq 1}$ satisfy

$$|a_{n+1} - a_n| \leq r |a_n - a_{n-1}| \quad \text{for all } n \geq 2$$

Prove that (a_n) converges. (First bound $|a_{n+1} - a_n|$ by a multiple of r^{n-1} , then estimate $|a_m - a_n|$ for $m > n$ by summing the intermediate differences.)

4. For each of the following, either produce an example or prove that none exists.
 1. A Cauchy sequence of real numbers with a divergent subsequence.
 2. An unbounded sequence of real numbers with a Cauchy subsequence.
 3. A monotone sequence of real numbers that is not Cauchy.
5. Exhibit a bounded sequence of rational numbers no subsequence of which converges to a rational number, and prove that it has this property. (This is the failure of statement (4) of theorem 2.3.6 for $\mathbb{Q}$, produced directly, without passing through the cycle.)
6. For $n \geq 1$ let $J_n = \{x \in \mathbb{R} \mid 0 < x < 1/n\}$. Show that $J_{n+1} \subseteq J_n$ and $J_n \neq \emptyset$ for every n , and that $\bigcap_{n \geq 1} J_n = \emptyset$. Which hypothesis of theorem 1.4.8 does this family violate?

2.4 Series and the Comparison Tests

A series is rarely decided by computing its partial sums, since a closed form for s_N is the exception. The tests below decide convergence without one, and each of them works by measuring a series against another whose behaviour is already known. So the section first settles three reference series, the geometric series, the harmonic series and $\sum 1/n^p$, and then proves the comparison, ratio and root tests that reduce an arbitrary series to one of them.

The failure in $\mathbb{Q}$ sits at the criterion for nonnegative terms. Increasing bounded partial sums have a limit in $\mathbb{R}$ because of theorem 1.2.4, and over $\mathbb{Q}$ they need not: the partial sums of $\sum_{n \geq 0} 1/n!$ are rational and lie below 3, while proposition 2.4.19 shows that the number they approach is irrational. That example closes the section, alongside the decimal expansion of a real number, which is the geometric series applied to the digits produced by proposition 1.4.3.

Definition 2.4.1: [Real] Series and Partial Sums

Let $(a_n)_{n \geq 1}$ be a sequence of real numbers. For $N \geq 1$ the N -th *partial sum* is

$$s_N = \sum_{n=1}^N a_n$$

The *series* $\sum_{n \geq 1} a_n$ *converges* if the sequence $(s_N)_{N \geq 1}$ converges, and in that case the *sum* of the series is

$$\sum_{n=1}^{\infty} a_n := \lim_{N \rightarrow \infty} s_N$$

If (s_N) diverges, the series *diverges*. A series $\sum_{n \geq m} a_n$ indexed from an integer m other than 1 is treated identically, with $s_N = \sum_{n=m}^N a_n$ for $N \geq m$.

The symbol $\sum_{n \geq 1} a_n$ names the series, and the symbol $\sum_{n=1}^{\infty} a_n$ names its sum, which exists only when the series converges.

The sum of a convergent series is a limit and not an addition, so every statement about limits from sections 2.1 to 2.3 applies to it with no further work. Uniqueness of the sum is proposition 2.1.4 applied to (s_N) . Two series arise immediately: a *finite* sum is the series whose terms vanish from some index on, whose partial sums are eventually constant, and the series $\sum_{n \geq 1} 1$, whose partial sums are $s_N = N$ and which therefore diverges by proposition 2.1.5, since $(N)_{N \geq 1}$ is unbounded by theorem 1.4.1(2).

Every result below is stated for a series indexed from $n = 1$ and holds verbatim for a series indexed from any integer m , since the two differ only in the bookkeeping of indices: the partial sums of $\sum_{n \geq m} a_n$ are those of $\sum_{n \geq 1} a_{n+m-1}$.

Convergence of (s_N) was characterized in theorem 2.3.5 without reference to a limit, and translating that characterization into a statement about the terms a_n gives the criterion every test in this section is proved from.

Proposition 2.4.2: The Cauchy Criterion for Series

The series $\sum_{n \geq 1} a_n$ converges if and only if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$\left| \sum_{k=n}^m a_k \right| < \epsilon$$

for all integers $m \geq n \geq N$.

Proof:

For $m \geq n \geq 2$ the identity $\sum_{k=n}^m a_k = s_m - s_{n-1}$ holds, since s_m and s_{n-1} share the terms $a_1, \dots, a_{n-1}$, and for $n = 1$ the left side is s_m itself. Both directions are this identity together with theorem 2.3.5.

The condition is necessary. Suppose the series converges, so that (s_N) converges. By proposition 2.3.3, (s_N) is a Cauchy sequence: given $\epsilon > 0$ there is an $N_0 \in \mathbb{N}$ with $|s_p - s_q| < \epsilon$ for all

$p, q \geq N_0$. Put $N = N_0 + 1$. If $m \geq n \geq N$ then $m \geq N_0 + 1 \geq N_0$ and $n - 1 \geq N_0$, so

$$\left| \sum_{k=n}^m a_k \right| = |s_m - s_{n-1}| < \epsilon$$

which is the stated condition.

The condition is sufficient. Suppose the condition holds and let $\epsilon > 0$. Take N as in the condition and let $p > q \geq N$. Applying the condition with $n = q + 1$ and $m = p$, which satisfy $m \geq n \geq N + 1 \geq N$, gives

$$|s_p - s_q| = \left| \sum_{k=q+1}^p a_k \right| < \epsilon$$

For $p = q$ the difference is $0 < \epsilon$, and the case $q > p$ follows by symmetry. So (s_N) is a Cauchy sequence, and theorem 2.3.5 makes it convergent, completing the proof.

The criterion sees only the terms a_n with $n \geq N$, and N may be taken as large as one likes. Altering finitely many terms therefore changes whether a series converges not at all. Concretely, if $a_n = b_n$ for every $n > M$, then for $m \geq n > M$ the two sums $\sum_{k=n}^m a_k$ and $\sum_{k=n}^m b_k$ are the same number, so the condition of proposition 2.4.2 holds for one series exactly when it holds for the other. The two sums differ, by the amount $\sum_{n=1}^M (a_n - b_n)$.

Taking $m = n$ in the criterion leaves a single term, and the resulting necessary condition is the cheapest divergence test there is.

Proposition 2.4.3: The n -th Term Test

If $\sum_{n \geq 1} a_n$ converges, then $a_n \rightarrow 0$. Equivalently, if (a_n) does not converge to 0, then $\sum_{n \geq 1} a_n$ diverges.

Proof:

Let $\epsilon > 0$ and take N as in proposition 2.4.2. For $n \geq N$ the choice $m = n$ is admissible, and the sum $\sum_{k=n}^n a_k$ is the single term a_n , so $|a_n| < \epsilon$. As ϵ was arbitrary, $a_n \rightarrow 0$ by definition 2.1.2. The second sentence is the contrapositive of the first, completing the proof.

The converse is false, and example 2.4.8 below is the standard example: $1/n \rightarrow 0$ while $\sum 1/n$ diverges. The test therefore establishes divergence and never convergence.

For a series of nonnegative terms the partial sums increase, and an increasing sequence has exactly one obstruction to convergence. This is where completeness enters the theory of series, and it is the criterion the condensation and comparison tests are proved from.

Proposition 2.4.4: Nonnegative Terms and Bounded Partial Sums

Let $a_n \geq 0$ for every $n \geq 1$. Then $\sum_{n \geq 1} a_n$ converges if and only if the set $\{s_N | N \geq 1\}$ of partial sums is bounded above, and in that case

$$\sum_{n=1}^{\infty} a_n = \sup_{N \geq 1} s_N$$

Proof:

Since $s_{N+1} - s_N = a_{N+1} \geq 0$, the sequence (s_N) is increasing.

If the partial sums are bounded above, theorem 2.2.2 makes (s_N) convergent, so the series converges. Conversely, a convergent sequence is bounded by proposition 2.1.5, so the partial sums of a convergent series are bounded above.

Suppose now that the series converges and write $s = \lim_N s_N$. First, s is an upper bound for the partial sums: if $s_{N_0} > s$ for some N_0 , then $s_N \geq s_{N_0}$ for every $N \geq N_0$ because (s_N) is increasing, and proposition 2.1.8 applied to the constant sequence s_{N_0} gives $s \geq s_{N_0} > s$, which is impossible. Second, let $\epsilon > 0$. Since $s_N \rightarrow s$ there is an N with $|s_N - s| < \epsilon$, hence $s_N > s - \epsilon$. So s is an upper bound that is approached to within every $\epsilon > 0$ from inside the set of partial sums, and proposition 1.2.6 identifies $s = \sup_N s_N$, as we sought to show.

Over $\mathbb{Q}$ the first half of this proposition fails. The partial sums of $\sum_{n \geq 0} 1/n!$ are rational and bounded above by 3, as example 2.4.18 computes, and no rational number is their limit, by proposition 2.4.19. The limit exists in $\mathbb{R}$ because theorem 2.2.2 produces it from theorem 1.2.4, and an ordered field without that property gives nothing in its place.

Sums and scalar multiples of convergent series behave as the algebra of limits predicts, and the triangle inequality passes from finite sums to infinite ones.

Proposition 2.4.5: The Algebra of Series and the Triangle Inequality

Let $\sum_{n \geq 1} a_n$ and $\sum_{n \geq 1} b_n$ converge, with sums A and B , and let $c \in \mathbb{R}$.

1. The series $\sum_{n \geq 1} (a_n + b_n)$ converges with sum $A + B$, and $\sum_{n \geq 1} ca_n$ converges with sum cA .
2. For all integers $m \geq n \geq 1$,

$$\left| \sum_{k=n}^m a_k \right| \leq \sum_{k=n}^m |a_k|$$

3. If in addition $\sum_{n \geq 1} |a_n|$ converges, then

$$\left| \sum_{n=1}^{\infty} a_n \right| \leq \sum_{n=1}^{\infty} |a_n|$$

Proof:

1. Write A_N and B_N for the partial sums of the two series. Rearranging a finite sum gives $\sum_{n=1}^N (a_n + b_n) = A_N + B_N$ and $\sum_{n=1}^N ca_n = cA_N$. Since $A_N \rightarrow A$ and $B_N \rightarrow B$, theorem 2.1.6 gives $A_N + B_N \rightarrow A + B$ and $cA_N \rightarrow cA$, which are the two assertions.
2. Fix n and induct on $m \geq n$. For $m = n$ both sides equal $|a_n|$. Assuming the inequality for m , the triangle inequality of proposition 1.3.8(4) gives

$$\left| \sum_{k=n}^{m+1} a_k \right| \leq \left| \sum_{k=n}^m a_k \right| + |a_{m+1}| \leq \sum_{k=n}^m |a_k| + |a_{m+1}| = \sum_{k=n}^{m+1} |a_k|$$

which closes the induction.

3. Write s_N for the partial sums of $\sum a_n$ and t_N for those of $\sum |a_n|$, and let $t = \lim_N t_N$. Part (2) with $n = 1$ and $m = N$ gives $|s_N| \leq t_N$ for every N . The sequence $(|s_N|)$ converges to $|A|$: by proposition 1.3.8(5), $||s_N| - |A|| \leq |s_N - A|$, and $(s_N - A)$ is a null sequence, so the comparison test for null sequences in proposition 2.1.10 makes $(|s_N| - |A|)$ null. Now proposition 2.1.8 applied to $|s_N| \leq t_N$ gives $|A| \leq t$, as we sought to show.

The three reference series come next. The first is the one case where the partial sums collapse, the second is the geometric series that every test in this section compares against, and the third is the harmonic series, which separates $a_n \rightarrow 0$ from convergence.

Example 2.4.6: The Telescoping Series $\sum 1/(n(n+1))$

Suppose $a_n = b_n - b_{n+1}$ for a sequence (b_n) . Then

$$s_N = \sum_{n=1}^N (b_n - b_{n+1}) = b_1 - b_{N+1}$$

because every interior term $b_2, \dots, b_N$ appears once with each sign, a cancellation confirmed by induction on N : the case $N = 1$ reads $b_1 - b_2$, and passing from N to $N + 1$ adds $b_{N+1} - b_{N+2}$ to $b_1 - b_{N+1}$. So $\sum_{n \geq 1} a_n$ converges precisely when $(b_{N+1})_{N \geq 1}$ converges, and then its sum is $b_1 - \lim_N b_{N+1}$.

Take $b_n = 1/n$. Then

$$b_n - b_{n+1} = \frac{1}{n} - \frac{1}{n+1} = \frac{(n+1) - n}{n(n+1)} = \frac{1}{n(n+1)}$$

and $b_{N+1} = 1/(N+1) \rightarrow 0$ by example 2.1.3(1). Hence

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$$

This is the rare series whose sum is available directly from its partial sums.

Example 2.4.7: The Geometric Series

Let $x \in \mathbb{R}$. The series $\sum_{n \geq 0} x^n$ converges if and only if $|x| < 1$, and in that case

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

Suppose first that $|x| \geq 1$. An induction using proposition 1.3.8(3) gives $|x^n| = |x|^n$, and a second induction using proposition 1.1.2(5) gives $|x|^n \geq 1$ for every $n \geq 0$. So (x^n) does not converge to 0, and proposition 2.4.3 makes the series divergent.

Suppose now $|x| < 1$, so that $x \neq 1$. Multiplying out,

$$(1-x) \sum_{n=0}^N x^n = \sum_{n=0}^N x^n - \sum_{n=0}^N x^{n+1} = 1 - x^{N+1}$$

the middle step being the telescoping cancellation of example 2.4.6 applied to $b_n = x^n$. Dividing by $1-x \neq 0$,

$$s_N = \sum_{n=0}^N x^n = \frac{1-x^{N+1}}{1-x}$$

If $x = 0$ then $x^{N+1} = 0$ for every N . If $x \neq 0$ then $r = |x|$ satisfies $0 < r < 1$, so $r^{N+1} \rightarrow 0$ by proposition 2.1.10 with $j = 0$, and since $|x^{N+1}| = r^{N+1}$ the comparison test for null sequences in the same proposition gives $x^{N+1} \rightarrow 0$. In both cases theorem 2.1.6 gives $s_N \rightarrow (1-0)/(1-x) = 1/(1-x)$.

Example 2.4.8: The Harmonic Series

The series $\sum_{n \geq 1} 1/n$ diverges, although its terms tend to 0.

The terms tend to 0 by example 2.1.3(1). For divergence, fix $n \geq 1$ and estimate the block of terms from $n+1$ to $2n$. It contains exactly n terms, and each satisfies $1/k \geq 1/(2n)$ because $k \leq 2n$ and proposition 1.1.2(7) reverses the inequality on taking reciprocals of positive numbers. Hence

$$\sum_{k=n+1}^{2n} \frac{1}{k} \geq n \cdot \frac{1}{2n} = \frac{1}{2}$$

Take $\epsilon = 1/2$ in proposition 2.4.2. Whatever N is proposed, the choice $n = N$ and $m = 2N$ satisfies $m \geq n \geq N$ and produces a block sum of at least $1/2$, so the condition of the criterion fails and the series diverges.

The harmonic series diverges slowly, and the block estimate says how slowly: applying it at $n = 2^j$ gives $s_{2^{j+1}} - s_{2^j} \geq 1/2$, so $s_{2^k} \geq s_1 + k/2 = 1 + k/2$ and the partial sums gain at least $1/2$ each time the index doubles. What decided the series was therefore the sequence of terms at the indices $1, 2, 4, 8, \dots$, each counted once for every index in the block it heads. For a decreasing sequence of nonnegative terms that reduction is available in general.

Theorem 2.4.9: Cauchy's Condensation Test

Let $(a_n)_{n \geq 1}$ satisfy $a_1 \geq a_2 \geq a_3 \geq \cdots \geq 0$. Then $\sum_{n \geq 1} a_n$ converges if and only if

$$\sum_{k \geq 0} 2^k a_{2^k} = a_1 + 2a_2 + 4a_4 + 8a_8 + \cdots$$

converges.

Proof:

Write $s_N = \sum_{n=1}^N a_n$ and $t_k = \sum_{j=0}^k 2^j a_{2^j}$. All terms of both series are nonnegative, so both sequences of partial sums are increasing and proposition 2.4.4 applies to each: the statement to be proved is that (s_N) is bounded above if and only if (t_k) is.

Step 1: $s_{2^{k+1}-1} \leq t_k$ for every integer $k \geq 0$. Split the index range $1 \leq n \leq 2^{k+1} - 1$ into the blocks $2^j \leq n \leq 2^{j+1} - 1$ for $j = 0, 1, \dots, k$, which are disjoint and exhaust it. The j -th block contains $2^{j+1} - 2^j = 2^j$ indices, and each of its terms satisfies $a_n \leq a_{2^j}$ because $n \geq 2^j$ and (a_n) is decreasing. Hence

$$s_{2^{k+1}-1} = \sum_{j=0}^k \sum_{n=2^j}^{2^{j+1}-1} a_n \leq \sum_{j=0}^k 2^j a_{2^j} = t_k$$

Step 2: $t_k \leq 2s_{2^k}$ for every integer $k \geq 0$. Split the range $1 \leq n \leq 2^k$ into $\{1\}$ and the blocks $2^{j-1} + 1 \leq n \leq 2^j$ for $j = 1, \dots, k$. The j -th block contains $2^j - 2^{j-1} = 2^{j-1}$ indices, and each of its terms satisfies $a_n \geq a_{2^j}$ because $n \leq 2^j$. Hence

$$s_{2^k} = a_1 + \sum_{j=1}^k \sum_{n=2^{j-1}+1}^{2^j} a_n \geq a_1 + \sum_{j=1}^k 2^{j-1} a_{2^j} = a_1 + \frac{1}{2}(t_k - a_1) = \frac{a_1}{2} + \frac{t_k}{2}$$

Since $a_1 \geq 0$, this gives $s_{2^k} \geq t_k/2$, which is the claim.

Step 3: the two boundedness conditions agree. An induction gives $2^k \geq k + 1$ for every integer $k \geq 0$: the case $k = 0$ reads $1 \geq 1$, and $2^{k+1} = 2 \cdot 2^k \geq 2k + 2 \geq k + 2$ since $k \geq 0$.

Suppose $\sum_{k \geq 0} 2^k a_{2^k}$ converges, so that $t_k \leq M$ for all k and some M , by proposition 2.4.4. Given $N \geq 1$, the inequality just proved gives $2^{N+1} - 1 \geq (N + 2) - 1 > N$, so (s_N) increasing and Step 1 give

$$s_N \leq s_{2^{N+1}-1} \leq t_N \leq M$$

Hence (s_N) is bounded above and $\sum_{n \geq 1} a_n$ converges.

Conversely, suppose $\sum_{n \geq 1} a_n$ converges, so that $s_N \leq M'$ for all N and some M' . Step 2 gives $t_k \leq 2s_{2^k} \leq 2M'$ for every k , so (t_k) is bounded above and $\sum_{k \geq 0} 2^k a_{2^k}$ converges, completing the proof.

The test replaces a series by one whose terms are spaced geometrically, and that is what makes it decisive on the family the comparison test most often needs.

Example 2.4.10: The p -Series for Integer Exponents

Let $p \in \mathbb{Z}$. The series

$$\sum_{n \geq 1} \frac{1}{n^p}$$

converges if and only if $p \geq 2$.

Suppose first $p \leq 0$. Then $1/n^p = n^{-p}$ with $-p \geq 0$, and $n^{-p} \geq 1$ for every $n \geq 1$ by induction from proposition 1.1.2(5). So the terms do not converge to 0 and proposition 2.4.3 gives divergence.

Suppose $p \geq 1$. Then $a_n = 1/n^p$ is positive, and it is decreasing: if $1 \leq n < m$ then $n^p < m^p$ by induction from proposition 1.1.2(5), and proposition 1.1.2(7) inverts this to $1/m^p < 1/n^p$. So theorem 2.4.9 applies, and the condensed series has terms

$$2^k a_{2^k} = \frac{2^k}{(2^k)^p} = 2^k 2^{-kp} = (2^{1-p})^k$$

This is the geometric series of example 2.4.7 with ratio $q = 2^{1-p} > 0$, so it converges if and only if $q < 1$. For $p = 1$ the ratio is $q = 1$. For $p \geq 2$ it is $q = 1/2^{p-1} \leq 1/2 < 1$, since $2^{p-1} \geq 2$. So the condensed series converges exactly when $p \geq 2$, and theorem 2.4.9 transfers that verdict to $\sum_{n \geq 1} 1/n^p$.

The case $p = 1$ recovers example 2.4.8 by a second route, since the condensed series of $\sum 1/n$ is $\sum_{k \geq 0} 1$. The three examples above are the whole stock of series whose convergence this section decides directly. Everything else is decided by comparison, and the tests that make that comparison usable are stated next.

Each of the three tests measures $|a_n|$, with the sign of a_n discarded, so each concludes something formally stronger than convergence.

Definition 2.4.11: Absolute Convergence

The series $\sum_{n \geq 1} a_n$ *converges absolutely* if the series $\sum_{n \geq 1} |a_n|$ converges.

Since $|a_n| \geq 0$, absolute convergence is decided by proposition 2.4.4: it holds exactly when the sums $\sum_{n=1}^N |a_n|$ are bounded above. For a concrete pair, the series $\sum_{n \geq 1} (-1)^{n+1}/n^2$ converges absolutely, since $\sum_{n \geq 1} 1/n^2$ converges by example 2.4.10, while $\sum_{n \geq 1} (-1)^{n+1}/n$ does not, since $\sum_{n \geq 1} 1/n$ diverges by example 2.4.8. The second series does nevertheless converge, which no test in this section can show. Section 2.5 proves it, and studies the series that converge without converging absolutely.

Theorem 2.4.12: Absolute Convergence Implies Convergence

If $\sum_{n \geq 1} a_n$ converges absolutely, then it converges.

Proof:

Let $\epsilon > 0$. Applying proposition 2.4.2 to the convergent series $\sum_{n \geq 1} |a_n|$ produces an N such that

$$\sum_{k=n}^m |a_k| = \left| \sum_{k=n}^m a_k \right| < \epsilon$$

for all $m \geq n \geq N$, the first equality holding because a sum of nonnegative terms is nonnegative. For the same m and n , proposition 2.4.5(2) gives

$$\left| \sum_{k=n}^m a_k \right| \leq \sum_{k=n}^m |a_k| < \epsilon$$

So $\sum_{n \geq 1} a_n$ satisfies the condition of proposition 2.4.2 and converges, as we sought to show.

Absolute convergence is what the criterion of proposition 2.4.4 can certify, and theorem 2.4.12 is what carries that certificate back to the original series. The comparison test is the first application, and it is the statement from which the ratio and root tests are both derived.

Theorem 2.4.13: The Comparison Test

Let (a_n) , (c_n) and (d_n) be real sequences and let $N_0 \in \mathbb{N}$.

1. If $|a_n| \leq c_n$ for all $n \geq N_0$ and $\sum_{n \geq 1} c_n$ converges, then $\sum_{n \geq 1} a_n$ converges absolutely, hence converges.
2. If $a_n \geq d_n \geq 0$ for all $n \geq N_0$ and $\sum_{n \geq 1} d_n$ diverges, then $\sum_{n \geq 1} a_n$ diverges.

Proof:

1. Let $\epsilon > 0$. Since $\sum c_n$ converges, proposition 2.4.2 gives an N_1 with $|\sum_{k=n}^m c_k| < \epsilon$ for all $m \geq n \geq N_1$. Put $N = \max\{N_0, N_1\}$ and let $m \geq n \geq N$. Every index k in that range satisfies $k \geq N_0$, so $c_k \geq |a_k| \geq 0$, and summing these inequalities gives

$$\sum_{k=n}^m |a_k| \leq \sum_{k=n}^m c_k = \left| \sum_{k=n}^m c_k \right| < \epsilon$$

where the middle equality holds because the c_k in the range are nonnegative. The left side is its own absolute value for the same reason, so $\sum_{n \geq 1} |a_n|$ satisfies the condition of proposition 2.4.2 and converges. That is absolute convergence of $\sum a_n$, and theorem 2.4.12 gives convergence.

2. Suppose, for a contradiction, that $\sum_{n \geq 1} a_n$ converges. Let $\epsilon > 0$ and take N_1 from proposition 2.4.2 for this series, and put $N = \max\{N_0, N_1\}$. For $m \geq n \geq N$ every index k in the range satisfies $a_k \geq d_k \geq 0$, so both $\sum_{k=n}^m d_k$ and $\sum_{k=n}^m a_k$ are nonnegative and

$$\left| \sum_{k=n}^m d_k \right| = \sum_{k=n}^m d_k \leq \sum_{k=n}^m a_k = \left| \sum_{k=n}^m a_k \right| < \epsilon$$

So $\sum_{n \geq 1} d_n$ satisfies the condition of proposition 2.4.2 and converges, contradicting the hypothesis. Hence $\sum_{n \geq 1} a_n$ diverges, completing the proof.

Both clauses tolerate a finite number of exceptions, which matters in practice: an inequality between two sequences almost never holds at the first few indices. Two computations show the shape of an

application. The series $\sum_{n \geq 1} (n+1)/(n^3+2)$ converges, because $n+1 \leq 2n$ and $n^3+2 \geq n^3$ for $n \geq 1$ give

$$\frac{n+1}{n^3+2} \leq \frac{2n}{n^3} = \frac{2}{n^2}$$

while $\sum 2/n^2$ converges by example 2.4.10 and proposition 2.4.5(1). The series $\sum_{n \geq 1} 1/\sqrt{n}$ diverges, because $\sqrt{n} \geq 1$ for $n \geq 1$ gives $n = (\sqrt{n})^2 \geq \sqrt{n}$, hence $1/\sqrt{n} \geq 1/n > 0$ by proposition 1.1.2(7), and $\sum 1/n$ diverges by example 2.4.8. Here $\sqrt{n}$ is the nonnegative square root produced by theorem 1.3.12.

Applying the comparison test requires a candidate series to compare against, and the geometric series is almost always the candidate. The next two theorems automate that comparison: each extracts from (a_n) a number that decides whether $|a_n|$ is eventually dominated by β^n for some $\beta < 1$. They differ in how the number is extracted, and theorem 2.4.16 settles which extraction is sharper.

Theorem 2.4.14: The Ratio Test

Let $(a_n)_{n \geq 1}$ satisfy $a_n \neq 0$ for every n .

1. If $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$, then $\sum_{n \geq 1} a_n$ converges absolutely.
2. If there is an n_0 with $|a_{n+1}/a_n| \geq 1$ for all $n \geq n_0$, then $\sum_{n \geq 1} a_n$ diverges.

Proof:

1. Write α for the limit superior and choose a real β with $\alpha < \beta < 1$, which exists by proposition 1.1.2(8) applied to α and 1. Since $\beta > \alpha$, the eventual domination clause of theorem 2.2.12 gives an N with $|a_{n+1}/a_n| < \beta$ for all $n \geq N$, that is $|a_{n+1}| < \beta|a_n|$ for $n \geq N$ after multiplying by $|a_n| > 0$.

An induction on $p \geq 0$ now gives $|a_{N+p}| \leq \beta^p |a_N|$: the case $p = 0$ is an equality, and the inductive step multiplies the hypothesis by $\beta > 0$ and applies $|a_{N+p+1}| < \beta|a_{N+p}|$. Setting $C = |a_N|\beta^{-N} > 0$ and writing $n = N + p$, this reads

$$|a_n| \leq C\beta^n \quad (n \geq N)$$

Since $0 < \beta < 1$, the series $\sum_{n \geq 0} C\beta^n$ converges by example 2.4.7 and proposition 2.4.5(1). By theorem 2.4.13(1), with $N_0 = N$ and $c_n = C\beta^n$, the series $\sum_{n \geq 1} a_n$ converges absolutely.

2. The hypothesis gives $|a_{n+1}| \geq |a_n|$ for $n \geq n_0$, so an induction on $p \geq 0$ gives $|a_{n_0+p}| \geq |a_{n_0}|$. Because $a_{n_0} \neq 0$, the number $\delta = |a_{n_0}|$ is positive by proposition 1.3.8(1), and $|a_n| \geq \delta$ for all $n \geq n_0$. So no N has $|a_n - 0| < \delta$ for all $n \geq N$, and (a_n) does not converge to 0 by definition 2.1.2. By proposition 2.4.3 the series diverges, completing the proof.

Clause (2) is stated with a hypothesis on every large n , where clause (1) uses a limit superior, and the asymmetry is deliberate: a limit superior at least 1 is compatible with convergence, as $\sum (-1)^{n+1}/n^2$ shows, whose ratios have limit superior 1. The root test has no such asymmetry.

Theorem 2.4.15: The Root Test

Let $(a_n)_{n \geq 1}$ be a real sequence and put

$$\alpha = \limsup_{n \rightarrow \infty} |a_n|^{1/n} \in [0, +\infty]$$

where $|a_n|^{1/n}$ is the nonnegative n -th root of theorem 1.3.12.

1. If $\alpha < 1$, then $\sum_{n \geq 1} a_n$ converges absolutely.
2. If $\alpha > 1$, then $\sum_{n \geq 1} a_n$ diverges.
3. If $\alpha = 1$, both outcomes occur, so the test decides nothing.

Proof:

Throughout, the inequality $0 \leq u < v$ implies $u^n < v^n$ for every $n \geq 1$, by induction on n : multiplying $u < v$ by $u \geq 0$ gives $u^{n+1} \leq u \cdot v^n$ once $u^n \leq v^n$ is known, and multiplying $u < v$ by $v^n > 0$ gives $uv^n < v^{n+1}$, both steps by proposition 1.1.2(5).

1. Choose a real β with $\alpha < \beta < 1$, by proposition 1.1.2(8). The eventual domination clause of theorem 2.2.12 gives an N with $|a_n|^{1/n} < \beta$ for all $n \geq N$. Raising both sides to the n -th power by the inequality above, and using $(|a_n|^{1/n})^n = |a_n|$ from theorem 1.3.12, gives

$$|a_n| < \beta^n \quad (n \geq N)$$

The series $\sum_{n \geq 0} \beta^n$ converges by example 2.4.7 because $0 < \beta < 1$, so theorem 2.4.13(1) with $N_0 = N$ and $c_n = \beta^n$ gives absolute convergence.

2. Suppose $\alpha > 1$. Then $|a_n|^{1/n} > 1$ for infinitely many n . Otherwise there would be an N with $|a_n|^{1/n} \leq 1$ for every $n \geq N$, and the constant 1 would dominate the sequence from N on, forcing $\alpha \leq 1$ by theorem 2.2.12 and contradicting $\alpha > 1$. For each such n , raising $1 < |a_n|^{1/n}$ to the n -th power gives $|a_n| > 1$. So no N has $|a_n - 0| < 1$ for all $n \geq N$, hence (a_n) does not converge to 0 and proposition 2.4.3 gives divergence.
3. Take $a_n = 1$ for every n . Then $|a_n|^{1/n} = 1$ for every n , so $\alpha = 1$, and the series diverges by proposition 2.4.3. For a convergent example with $\alpha = 1$, take $a_n = 1/n^2$, which converges by example 2.4.10. Here $|a_n|^{1/n} = (n^{1/n})^{-2}$, and the evaluation $\alpha = 1$ rests on $n^{1/n} \rightarrow 1$. That limit is proved as proposition 2.6.1 in section 2.6, and this one step of the present proof is deferred to it, completing the proof.

Both tests detect the same thing, domination of $|a_n|$ by a geometric series, and both fail in the same way, by returning the value 1. They are not equally sharp. The next theorem compares the two numbers they extract, and its right-hand inequality says both that whenever the ratio test establishes convergence the root test does too, and that whenever the root test is inconclusive so is the ratio test. Its left-hand inequality is needed for neither of those, and exercise 2.4.1 (8) is where it is used.

Theorem 2.4.16: The Root Test Is at Least as Sharp as the Ratio Test

For every sequence $(c_n)_{n \geq 1}$ of positive numbers,

$$\liminf_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} \leq \liminf_{n \rightarrow \infty} c_n^{1/n} \leq \limsup_{n \rightarrow \infty} c_n^{1/n} \leq \limsup_{n \rightarrow \infty} \frac{c_{n+1}}{c_n}$$

Proof:

The middle inequality holds for every sequence, since definition 2.2.10 takes both as an infimum and a supremum of the same nonempty set E of subsequential limits, and theorem 2.2.12(2) places s_* and s^* in E . So only the outer two are at issue. Both use two preliminary facts. First, if $0 < s < 1$ then $s^n \rightarrow 0$ by proposition 2.1.10 with $j = 0$, so for every $D > 0$ there is an index beyond which $s^n < D$. Second, $0 \leq u < v$ implies $u^n < v^n$ for every $n \geq 1$, by the induction recorded in the proof of theorem 2.4.15, so an inequality between $c_n^{1/n}$ and a nonnegative number may be raised to the n -th power, where $(c_n^{1/n})^n = c_n$ by theorem 1.3.12.

Step 1: the right-hand inequality. Put $\alpha = \limsup_n c_{n+1}/c_n$. If $\alpha = +\infty$ there is nothing to prove, so assume α is a real number, necessarily $\alpha \geq 0$ because every ratio is positive. Let $\beta > \alpha$. Then $\beta > 0$. By the eventual domination clause of theorem 2.2.12 there is an N with $c_{n+1}/c_n < \beta$ for all $n \geq N$, hence $c_{n+1} < \beta c_n$ for $n \geq N$ after multiplying by $c_n > 0$. An induction on $p \geq 0$ gives $c_{N+p} \leq \beta^p c_N$, so with $C = c_N \beta^{-N} > 0$,

$$c_n \leq C\beta^n \quad (n \geq N) \quad (2.1)$$

Suppose, for a contradiction, that $\gamma = \limsup_n c_n^{1/n}$ satisfies $\gamma > \beta$. Choose a real β' with $\beta < \beta' < \gamma$, taking the midpoint of β and γ when γ is finite (proposition 1.1.2(8)) and $\beta' = \beta + 1$ when $\gamma = +\infty$. If $c_n^{1/n} \leq \beta'$ held for every n beyond some index, then β' would dominate the sequence eventually and theorem 2.2.12 would give $\gamma \leq \beta'$, against the choice $\beta' < \gamma$. So $c_n^{1/n} > \beta'$ for infinitely many n , and raising to the n -th power gives $c_n > (\beta')^n$ for those n . Combining with (2.1), for infinitely many $n \geq N$,

$$(\beta')^n < C\beta^n, \quad \text{that is} \quad \left(\frac{\beta}{\beta'}\right)^n > \frac{1}{C}$$

But $s = \beta/\beta'$ lies in $(0, 1)$, so by the preliminary computation $s^n < 1/C$ for all large n , and only finitely many n can satisfy $(\beta/\beta')^n > 1/C$. This contradiction gives $\gamma \leq \beta$. As $\beta > \alpha$ was arbitrary, $\gamma \leq \alpha + \epsilon$ for every $\epsilon > 0$, and proposition 1.4.7(1) gives $\gamma \leq \alpha$.

Step 2: the left-hand inequality. Put $\alpha' = \liminf_n c_{n+1}/c_n$ and $\gamma' = \liminf_n c_n^{1/n}$, and note $\gamma' \geq 0$ since every $c_n^{1/n}$ is positive. If $\alpha' = 0$ then $\alpha' \leq \gamma'$ holds because $\gamma' \geq 0$, so let β be any real with $0 < \beta < \alpha'$. The eventual domination clause of theorem 2.2.12, in its form for the limit inferior, gives an N with $c_{n+1}/c_n > \beta$ for all $n \geq N$, hence $c_{n+1} > \beta c_n$. An induction on $p \geq 0$ gives $c_{N+p} \geq \beta^p c_N$, so with $C = c_N \beta^{-N} > 0$,

$$c_n \geq C\beta^n \quad (n \geq N) \quad (2.2)$$

Suppose, for a contradiction, that $\gamma' < \beta$, and choose a real β' with $\gamma' < \beta' < \beta$ by proposition 1.1.2(8). If $c_n^{1/n} \geq \beta'$ held for every n beyond some index, then β' would bound the sequence

from below eventually and theorem 2.2.12 would give $\gamma' \geq \beta'$, against the choice $\beta' > \gamma'$. So $c_n^{1/n} < \beta'$ for infinitely many n , and raising to the n -th power gives $c_n < (\beta')^n$ for those n . Combining with (2.2), for infinitely many $n \geq N$,

$$C\beta^n < (\beta')^n, \quad \text{that is} \quad \left(\frac{\beta'}{\beta}\right)^n > C$$

Here $s = \beta'/\beta$ lies in $(0, 1)$, so $s^n < C$ for all large n and only finitely many n can satisfy $(\beta'/\beta)^n > C$. This contradiction gives $\gamma' \geq \beta$. As $\beta < \alpha'$ was arbitrary, $\gamma' \geq \alpha' - \epsilon$ for every $\epsilon \in (0, \alpha')$ when α' is finite, and proposition 1.4.7(1) gives $\gamma' \geq \alpha'$. When $\alpha' = +\infty$ the same conclusion $\gamma' \geq \beta$ for every real $\beta > 0$ makes γ' larger than every real number, so $\gamma' = +\infty = \alpha'$. This completes the proof.

Both consequences announced above follow from the right-hand inequality applied to $c_n = |a_n|$, which is positive because theorem 2.4.14 presupposes $a_n \neq 0$. If clause (1) of the ratio test applies, then $\limsup_n c_{n+1}/c_n < 1$, so $\limsup_n |a_n|^{1/n} < 1$ and theorem 2.4.15(1) applies as well. If the root test is inconclusive, so that $\limsup_n |a_n|^{1/n} = 1$, then $\limsup_n c_{n+1}/c_n \geq 1$ and clause (1) of the ratio test does not apply either. The reverse implications fail, and the following example is where.

Example 2.4.17: A Series the Root Test Decides and the Ratio Test Does Not

Define

$$a_n = \begin{cases} 2^{-n} & n \text{ odd} \\ 3^{-n} & n \text{ even} \end{cases}$$

so that the series is $\frac{1}{2} + \frac{1}{9} + \frac{1}{8} + \frac{1}{81} + \frac{1}{32} + \cdots$.

The root test decides it. By the uniqueness clause of theorem 1.3.12, $|a_n|^{1/n}$ equals $1/2$ for odd n and $1/3$ for even n , since $(1/2)^n = 2^{-n}$ and $(1/3)^n = 3^{-n}$. So the sequence $(|a_n|^{1/n})$ takes only the two values $1/2$ and $1/3$, is dominated by $1/2$ everywhere, and equals $1/2$ for infinitely many n , so the constant subsequence indexed by the odd n converges to $1/2$. By theorem 2.2.12 these two facts identify $\alpha = 1/2 < 1$, and theorem 2.4.15(1) gives absolute convergence.

The ratio test does not. For n odd,

$$\frac{a_{n+1}}{a_n} = \frac{3^{-(n+1)}}{2^{-n}} = \frac{2^n}{3^{n+1}} = \frac{1}{3} \left(\frac{2}{3}\right)^n \leq \frac{1}{3} \cdot \frac{2}{3} = \frac{2}{9}$$

using that $(2/3)^n \leq 2/3$ for $n \geq 1$, which is an induction from proposition 1.1.2(5) since $0 < 2/3 < 1$, while for n even,

$$\frac{a_{n+1}}{a_n} = \frac{2^{-(n+1)}}{3^{-n}} = \frac{3^n}{2^{n+1}} = \frac{1}{2} \left(\frac{3}{2}\right)^n \geq \frac{1}{2} \cdot \frac{9}{4} = \frac{9}{8}$$

using that $(3/2)^n \geq (3/2)^2 = 9/4$ for $n \geq 2$, again an induction from proposition 1.1.2(5) since $3/2 > 1$. Clause (2) of theorem 2.4.14 fails, since ratios below 1 occur beyond every index. Clause (1) fails as well: if the limit superior of the ratios were some $\alpha < 1$, then choosing β with $\alpha < \beta < 1$ and applying the eventual domination clause of theorem 2.2.12 would put every ratio below $\beta < 1$ from some index on, contradicting the bound $a_{n+1}/a_n \geq 9/8 > 1$ just proved at every even index. So the ratio test yields no verdict on a series the root test settles at once.

The series is not itself difficult, since $a_n \leq 2^{-n}$ for every n and theorem 2.4.13(1) settles it.

The tests now produce the number e , together with an error estimate sharp enough to prove it

irrational, and the decimal notation for a real number as a convergent series.

Example 2.4.18: The Series for e and Its Tail Estimate

The series $\sum_{n \geq 0} 1/n!$ converges. Its sum is written e , and it satisfies $2 < e < 3$. Writing $s_n = \sum_{k=0}^n 1/k!$ for its partial sums, every integer $n \geq 1$ satisfies

$$0 < e - s_n < \frac{1}{n!n} \quad (2.3)$$

Convergence and the bounds on e . An induction gives $n! \geq 2^{n-1}$ for every $n \geq 1$: the case $n = 1$ reads $1 \geq 1$, and $(n+1)! = (n+1) \cdot n! \geq 2 \cdot 2^{n-1} = 2^n$ since $n+1 \geq 2$. Hence $1/n! \leq (1/2)^{n-1}$ for $n \geq 1$ by proposition 1.1.2(7), and theorem 2.4.13(1) with the convergent geometric series $\sum_{n \geq 1} (1/2)^{n-1}$ of example 2.4.7 gives convergence. The same comparison bounds the sum. The inequality $1/n! \leq (1/2)^{n-1}$ is strict at $n = 3$, where $1/6 < 1/4$, so for every $N \geq 3$,

$$\sum_{n=1}^N \frac{1}{n!} \leq \sum_{n=1}^N \left(\frac{1}{2}\right)^{n-1} - \left(\frac{1}{4} - \frac{1}{6}\right) \leq 2 - \frac{1}{12}$$

the last step using that the geometric partial sums are bounded above by their limit 2, by proposition 2.4.4. Letting $N \rightarrow \infty$ and applying proposition 2.1.8,

$$e = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \leq 3 - \frac{1}{12} < 3$$

For the lower bound, $e \geq s_2 = 1 + 1 + 1/2 = 5/2 > 2$, since proposition 2.4.4 identifies e as the supremum of the partial sums and s_2 is one of them.

The tail estimate. Fix $n \geq 1$ and let $m > n$. For $k = n+1+j$ with $0 \leq j \leq m-n-1$, the factorial $k!$ contains the factors $n+2, n+3, \dots, k$ beyond $(n+1)!$, which are j factors each at least $n+2$, so $k! \geq (n+1)!(n+2)^j$ and hence $1/k! \leq (n+2)^{-j}/(n+1)!$ by proposition 1.1.2(7). Summing,

$$s_m - s_n = \sum_{k=n+1}^m \frac{1}{k!} \leq \frac{1}{(n+1)!} \sum_{j=0}^{m-n-1} \left(\frac{1}{n+2}\right)^j < \frac{1}{(n+1)!} \cdot \frac{1}{1 - \frac{1}{n+2}} = \frac{n+2}{(n+1)!(n+1)}$$

where the strict inequality holds because the finite geometric sum is smaller than the full geometric series of example 2.4.7, whose remaining terms are positive. Letting $m \rightarrow \infty$ and applying proposition 2.1.8,

$$e - s_n \leq \frac{n+2}{(n+1)!(n+1)}$$

Finally $n(n+2) = n^2 + 2n < n^2 + 2n + 1 = (n+1)^2$, so

$$\frac{n+2}{(n+1)!(n+1)} = \frac{1}{n!} \cdot \frac{n+2}{(n+1)^2} < \frac{1}{n!} \cdot \frac{1}{n}$$

which is the upper half of (2.3). The lower half holds because $e - s_n \geq s_{n+1} - s_n = 1/(n+1)! > 0$, again by proposition 2.1.8 and the fact that (s_m) is increasing.

Estimate (2.3) converges quickly enough to be used numerically, since s_{10} already approximates e

to within $1/(10! \cdot 10)$, a number below $3 \cdot 10^{-8}$. It also decides the arithmetic nature of e , because multiplying it by $n!$ produces a quantity confined between two consecutive integers.

Proposition 2.4.19: e Is Irrational

The number e is not rational.

Proof:

Suppose $e = p/q$ with p and q positive integers, which is possible for a positive rational number. Apply (2.3) with $n = q \geq 1$ and multiply through by $q! > 0$, which preserves the inequalities by proposition 1.1.2(5):

$$0 < q!(e - s_q) < \frac{q!}{q!q} = \frac{1}{q} \leq 1$$

Both quantities inside are integers. First, $q!e = q!p/q = p \cdot (q-1)!$, an integer because $q \geq 1$. Second,

$$q!s_q = \sum_{k=0}^q \frac{q!}{k!}$$

and each summand is an integer, since $k \leq q$ makes $q!/k!$ the product of the integers $k+1, k+2, \dots, q$, empty and equal to 1 when $k = q$. So $q!(e - s_q)$ is a difference of two integers, hence an integer, and it lies strictly between 0 and 1. No integer does, so the assumption is untenable and e is irrational, completing the proof.

This is the failure in $\mathbb{Q}$ promised at proposition 2.4.4, delivered in full: an increasing sequence of rational numbers, bounded above by 3, whose limit exists in $\mathbb{R}$ and is not itself rational.

The number e has a second description, as a limit, and the two are connected by the binomial theorem. The limit form is the one most applications of e use.

Theorem 2.4.20: e as a Limit

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$$

Proof:

Write $t_n = (1 + 1/n)^n$ and $s_n = \sum_{k=0}^n 1/k!$ as in example 2.4.18. The binomial theorem ([4, the binomial theorem]) expands

$$t_n = \sum_{k=0}^n \binom{n}{k} \frac{1}{n^k}$$

and the k -th summand rewrites as

$$\binom{n}{k} \frac{1}{n^k} = \frac{1}{k!} \cdot \frac{n(n-1) \cdots (n-k+1)}{n^k} = \frac{1}{k!} \prod_{j=0}^{k-1} \left(1 - \frac{j}{n}\right) = \frac{1}{k!} \prod_{j=1}^{k-1} \left(1 - \frac{j}{n}\right)$$

the last step dropping the factor $j = 0$, which is 1. For $0 \leq j \leq k-1 \leq n-1$ each factor lies in $(0, 1]$, so the product lies in $(0, 1]$ as well.

Step 1: $t_n \leq e$ for every n . Each summand is at most $1/k!$ by the previous paragraph, so $t_n \leq s_n$, and $s_n < e$ by (2.3).

Step 2: a lower bound depending on a fixed truncation. Fix an integer $m \geq 1$. For $n \geq m$ all $n + 1$ summands of t_n are nonnegative, so discarding those with $k > m$ gives

$$t_n \geq u_n^{(m)} := \sum_{k=0}^m \frac{1}{k!} \prod_{j=1}^{k-1} \left(1 - \frac{j}{n}\right)$$

Hold m fixed and let n grow. For each fixed j the sequence $(1 - j/n)_n$ converges to 1, since $j/n \rightarrow 0$ by example 2.1.3(1) and theorem 2.1.6. Each of the $m + 1$ summands is a product of at most m such factors times a constant, so theorem 2.1.6 applied finitely many times gives $u_n^{(m)} \rightarrow s_m$ as $n \rightarrow \infty$.

Step 3: the two bounds close. Let $\epsilon > 0$. Since $s_m \rightarrow e$, there is an m with $s_m > e - \epsilon/2$. With that m fixed, Step 2 provides an $N \geq m$ with $|u_n^{(m)} - s_m| < \epsilon/2$, hence $u_n^{(m)} > s_m - \epsilon/2$, for all $n \geq N$. For such n ,

$$e - \epsilon < s_m - \frac{\epsilon}{2} < u_n^{(m)} \leq t_n \leq e$$

so $|t_n - e| < \epsilon$. As $\epsilon > 0$ was arbitrary, $t_n \rightarrow e$ by definition 2.1.2, as we sought to show.

A decimal expansion is an infinite series, its convergence follows from the geometric series, and its digits are produced by proposition 1.4.3. Stating it as a theorem also settles the status of the identity $0.999\ldots = 1$.

Proposition 2.4.21: Decimal Expansions

Let $x \in \mathbb{R}$ with $x \geq 0$. Put $a_0 = \lfloor x \rfloor$ and, for $n \geq 1$,

$$a_n = \lfloor 10^n x \rfloor - 10 \lfloor 10^{n-1} x \rfloor$$

Then each a_n with $n \geq 1$ is an integer with $0 \leq a_n \leq 9$, and

$$x = a_0 + \sum_{n=1}^{\infty} \frac{a_n}{10^n}$$

Distinct digit sequences can represent the same real number: the series $\sum_{n \geq 1} 9 \cdot 10^{-n}$ converges to 1.

Proof:

Write $b_n = \lfloor 10^n x \rfloor$ for $n \geq 0$, so $a_0 = b_0$ and $a_n = b_n - 10b_{n-1}$ for $n \geq 1$. Each b_n is an integer and proposition 1.4.3 gives

$$b_n \leq 10^n x < b_n + 1 \tag{2.4}$$

Step 1: the digits lie in $\{0, 1, \dots, 9\}$. Multiplying (2.4) at index $n - 1$ by $10 > 0$ gives $10b_{n-1} \leq 10^n x < 10b_{n-1} + 10$. Combining the left half of this with the right half of (2.4) at index n gives $10b_{n-1} \leq 10^n x < b_n + 1$, hence $10b_{n-1} < b_n + 1$ and so $a_n = b_n - 10b_{n-1} > -1$. Since a_n is an integer, $a_n \geq 0$. Combining the left half of (2.4) at index n with the right half of the multiplied inequality gives $b_n \leq 10^n x < 10b_{n-1} + 10$, hence $a_n < 10$ and so $a_n \leq 9$.

Step 2: the partial sums telescope. Dividing $a_n = b_n - 10b_{n-1}$ by 10^n gives

$$\frac{a_n}{10^n} = \frac{b_n}{10^n} - \frac{b_{n-1}}{10^{n-1}}$$

so the interior terms cancel in the partial sums and

$$a_0 + \sum_{n=1}^N \frac{a_n}{10^n} = \frac{b_N}{10^N} \quad (N \geq 1)$$

This is an induction on N : for $N = 1$ the left side is $b_0 + (b_1/10 - b_0) = b_1/10$, and the step from N to $N + 1$ adds $b_{N+1}/10^{N+1} - b_N/10^N$ to $b_N/10^N$.

Step 3: the partial sums converge to x . Dividing (2.4) by $10^N > 0$ gives

$$0 \leq x - \frac{b_N}{10^N} < \frac{1}{10^N}$$

and $(1/10)^N \rightarrow 0$ by proposition 2.1.10 with $j = 0$ and $r = 1/10$. By theorem 2.1.7 the sequence $(x - b_N/10^N)$ converges to 0, so by theorem 2.1.6 the partial sums $b_N/10^N$ converge to x . That is the identity $x = a_0 + \sum_{n=1}^{\infty} a_n/10^n$.

Step 4: the nines. By example 2.4.7 and proposition 2.4.5(1),

$$\sum_{n=1}^{\infty} \frac{9}{10^n} = \frac{9}{10} \sum_{n=0}^{\infty} \left(\frac{1}{10}\right)^n = \frac{9}{10} \cdot \frac{1}{1 - \frac{1}{10}} = \frac{9}{10} \cdot \frac{10}{9} = 1$$

completing the proof.

Applied to $x = 1$ the recipe returns $a_0 = 1$ and $a_n = 0$ for $n \geq 1$, so the expansion it produces is the terminating one. Step 4 shows that the non-terminating sequence of nines represents the same number, which is the content of the identity $0.999\cdots = 1$: the two symbols are two convergent series with one sum, not two numbers. The proposition also gives a second proof that $\mathbb{Q}$ is dense in $\mathbb{R}$, established as theorem 1.4.4, since the truncation $a_0 + \sum_{n=1}^N a_n 10^{-n}$ is rational and Steps 2 and 3 together place it within 10^{-N} of x . The proposition asks for $x \geq 0$, and a negative x is reached by approximating $-x$ and negating.

Exercise 2.4.1

1. Show that $\sum_{n \geq 1} \frac{1}{n(n+2)}$ converges and compute its sum.
2. Compute $\sum_{n=1}^{\infty} \frac{2^n + 3^n}{6^n}$, justifying each step by a result of this section.
3. Show that $\sum_{n \geq 1} \frac{n}{2n+1}$ diverges.
4. Decide the convergence of each of the following, in each case naming the series used for comparison.

1. $\sum_{n \geq 1} \frac{n+1}{n^4+5}$

$$2. \sum_{n \geq 1} \frac{1}{2^n + n}$$

$$3. \sum_{n \geq 2} \frac{1}{n-1}$$

5. Use theorem 2.4.9 to give a second proof that $\sum_{n \geq 1} 1/n$ diverges, and identify where the hypothesis that the terms decrease is used.
6. Let $a_1 \geq a_2 \geq \dots \geq 0$ and suppose $\sum_{n \geq 1} a_n$ converges. Show that $na_{2n} \rightarrow 0$.
7. Apply theorem 2.4.14 to $\sum_{n \geq 1} 2^n/n!$ and to $\sum_{n \geq 1} n^2/2^n$, and in each case state the value of the limit superior of the ratios.
8. Let (c_n) be a sequence of positive numbers.
 1. Show that if $\lim_n c_{n+1}/c_n = L$ exists and is a real number, then $\lim_n c_n^{1/n} = L$.
 2. Deduce that $\lim_{n \rightarrow \infty} \frac{(n!)^{1/n}}{n} = \frac{1}{e}$.
9. Verify that $s_5 = 163/60$, and use (2.3) to deduce $163/60 < e < 1631/600$.
10. Show that $\frac{4}{10} + \sum_{n=2}^{\infty} \frac{9}{10^n} = \frac{1}{2}$, and determine which digit sequence the recipe of proposition 2.4.21 produces for $x = 1/2$.
11. Let $\sum_{n \geq 1} a_n$ converge absolutely and let (b_n) be a bounded sequence. Show that $\sum_{n \geq 1} a_n b_n$ converges absolutely, and show that boundedness of (b_n) cannot be dropped, by exhibiting an absolutely convergent $\sum_{n \geq 1} a_n$ and an unbounded (b_n) for which $\sum_{n \geq 1} a_n b_n$ diverges.

2.5 Conditional Convergence and Rearrangement

Every test proved in section 2.4 reaches its conclusion from a bound on $|a_n|$ alone, and so each of them establishes absolute convergence. A series whose terms change sign can converge for a different reason: not because its terms are small enough to be added with the signs discarded, but because consecutive terms cancel. This section proves the two tests that detect convergence of that second kind, names the series that converge for that reason and no other, and then measures the difference between the two kinds. An absolutely convergent series has the same sum in every order. A conditionally convergent one can be reordered so that its partial sums have any prescribed limit inferior and limit superior in $[-\infty, +\infty]$, subject only to the first not exceeding the second, and its Cauchy product with a convergent series can diverge.

Neither test below produces an expression for the sum whose existence it asserts. Each closes on the Cauchy criterion for series, proposition 2.4.2, which is theorem 2.3.5 read for the sequence of partial sums, so completeness is again what produces the number.

The tests of section 2.4 discard the signs of the terms at their first step, so none of them can see cancellation. What replaces them is an identity that exposes the cancellation directly: it trades the terms a_n for their partial sums and the terms b_n for their successive differences, so that a series

whose partial sums merely stay bounded can still be handled. The identity is the discrete analogue of integration by parts.

Lemma 2.5.1: Summation by Parts

Let $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ be sequences of real numbers, and put

$$A_N = \sum_{n=1}^N a_n \quad (N \geq 1), \quad A_0 = 0$$

Then for all integers $1 \leq p \leq q$,

$$\sum_{n=p}^q a_n b_n = \sum_{n=p}^{q-1} A_n (b_n - b_{n+1}) + A_q b_q - A_{p-1} b_p$$

where the first sum on the right is empty, and therefore 0, when $p = q$.

Proof:

For every $n \geq 1$, $a_n = A_n - A_{n-1}$, which for $n = 1$ reads $a_1 = A_1 - A_0 = A_1$. Hence

$$\sum_{n=p}^q a_n b_n = \sum_{n=p}^q (A_n - A_{n-1}) b_n = \sum_{n=p}^q A_n b_n - \sum_{n=p}^q A_{n-1} b_n$$

Re-indexing the second sum by $m = n - 1$ turns it into $\sum_{m=p-1}^{q-1} A_m b_{m+1}$. Splitting the first sum at its top term and the second at its bottom term,

$$\sum_{n=p}^q A_n b_n - \sum_{n=p-1}^{q-1} A_n b_{n+1} = A_q b_q + \sum_{n=p}^{q-1} A_n b_n - A_{p-1} b_p - \sum_{n=p}^{q-1} A_n b_{n+1}$$

Collecting the two sums over the common range $p \leq n \leq q - 1$ gives $\sum_{n=p}^{q-1} A_n (b_n - b_{n+1}) + A_q b_q - A_{p-1} b_p$, as we sought to show.

The right-hand side is small as soon as the numbers A_n are bounded and the increments $b_n - b_{n+1}$ have a small total over the range, and for a monotone (b_n) that total telescopes to $b_p - b_q$. This is what the next test uses.

Both the test and its corollary also use the following observation. A nonincreasing sequence with limit 0 has every term nonnegative: if $b_N < 0$ for some N , then $b_n \leq b_N$ for all $n \geq N$, and proposition 2.1.8 applied to the tail $(b_n)_{n \geq N}$ gives $0 = \lim_n b_n \leq b_N < 0$, which is false.

Theorem 2.5.2: Dirichlet's Test

Let $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ be sequences of real numbers such that

1. the partial sums $A_N = \sum_{n=1}^N a_n$ form a bounded sequence,
2. $b_1 \geq b_2 \geq b_3 \geq \dots$, and
3. $b_n \rightarrow 0$.

Then $\sum_{n \geq 1} a_n b_n$ converges.

Proof:

Choose $M > 0$ with $|A_n| \leq M$ for every $n \geq 0$, which is possible by (1) together with $A_0 = 0$. By the observation above, every b_n is nonnegative, and by (2) every difference $b_n - b_{n+1}$ is nonnegative as well.

Let $\epsilon > 0$. By (3) there is an N with $b_N \leq \epsilon/(2M)$. Fix integers $q \geq p \geq N$. Applying lemma 2.5.1 on the range $p \leq n \leq q$ and then the triangle inequality proposition 2.4.5(2) in its finite form,

$$\left| \sum_{n=p}^q a_n b_n \right| \leq \sum_{n=p}^{q-1} |A_n| (b_n - b_{n+1}) + |A_q| b_q + |A_{p-1}| b_p \leq M \left(\sum_{n=p}^{q-1} (b_n - b_{n+1}) + b_q + b_p \right)$$

where the nonnegativity of $b_n - b_{n+1}$, b_q and b_p is what allows the bound $|A_n| \leq M$ to be pulled out of each term. The remaining sum telescopes to $b_p - b_q$, so the right-hand side equals $M(b_p - b_q + b_q + b_p) = 2Mb_p$. Since $p \geq N$ and (b_n) is nonincreasing, $b_p \leq b_N \leq \epsilon/(2M)$, and therefore

$$\left| \sum_{n=p}^q a_n b_n \right| \leq 2Mb_p \leq \epsilon$$

for all $q \geq p \geq N$. This is the Cauchy criterion for series, proposition 2.4.2, so $\sum_{n \geq 1} a_n b_n$ converges, completing the proof.

Hypothesis (1) is much weaker than convergence of $\sum a_n$: it asks only that the partial sums stay in a bounded set, and permits them to oscillate forever. For a concrete case, let (a_n) repeat the block $1, 1, -2$ forever, so that $a_1 = a_2 = 1$, $a_3 = -2$, $a_4 = a_5 = 1$, $a_6 = -2$, and so on. The partial sums run $1, 2, 0, 1, 2, 0, \dots$, so $|A_N| \leq 2$ for every N , while $\sum_{n \geq 1} a_n$ itself does not converge, its partial sums taking the value 2 and the value 0 infinitely often. Taking $b_n = 1/n$, which decreases to 0, theorem 2.5.2 gives the convergence of

$$\sum_{n \geq 1} \frac{a_n}{n} = 1 + \frac{1}{2} - \frac{2}{3} + \frac{1}{4} + \frac{1}{5} - \frac{2}{6} + \dots$$

The same series with absolute values diverges: $|a_n| \geq 1$ for every n , so $|a_n|/n \geq 1/n$, and theorem 2.4.13 together with example 2.4.8 makes $\sum_n |a_n|/n$ divergent.

The case in which the block is $1, -1$ is the one that occurs most often, and it carries an estimate for the error committed by stopping at a finite stage. That estimate is what makes an alternating series usable for computation, and it is recorded here alongside the convergence statement.

Corollary 2.5.3: The Alternating Series Test

Let $b_1 \geq b_2 \geq b_3 \geq \cdots$ with $b_n \rightarrow 0$. Write $s_N = \sum_{n=1}^N (-1)^{n+1} b_n$. Then

1. $\sum_{n \geq 1} (-1)^{n+1} b_n$ converges. Call its sum s .
2. $s_{2k} \leq s \leq s_{2k-1}$ for every $k \geq 1$.
3. $|s - s_N| \leq b_{N+1}$ for every $N \geq 1$.

Proof:

1. Put $a_n = (-1)^{n+1}$. Pairing consecutive terms gives $A_{2m} = \sum_{j=1}^m (1 - 1) = 0$ and $A_{2m+1} = A_{2m} + 1 = 1$, so $|A_N| \leq 1$ for every N . Hypotheses (2) and (3) of theorem 2.5.2 are the hypotheses assumed here, so that theorem applies and $\sum_{n \geq 1} (-1)^{n+1} b_n$ converges.

2. As noted before theorem 2.5.2, every b_n is nonnegative. Three computations from the definition of s_N record the shape of the partial sums:

$$s_{2k+2} - s_{2k} = b_{2k+1} - b_{2k+2} \geq 0, \quad s_{2k+1} - s_{2k-1} = b_{2k+1} - b_{2k} \leq 0, \quad s_{2k+1} - s_{2k} = b_{2k+1} \geq 0$$

So $(s_{2k})_{k \geq 1}$ is nondecreasing, and the second computation, read for $k \geq 1$, says that $(s_{2k-1})_{k \geq 1}$ is nonincreasing. Both are subsequences of the convergent sequence (s_N) , so by proposition 2.2.7 both converge to s . Fix $k \geq 1$. For every $j \geq k$ the inequality $s_{2j} \geq s_{2k}$ holds, and letting $j \rightarrow \infty$ in it, proposition 2.1.8 gives $s \geq s_{2k}$. The same argument applied to the nonincreasing sequence $(s_{2j-1})_{j \geq k}$ gives $s \leq s_{2k-1}$.

3. Let $N \geq 1$. If $N = 2k$ is even, then part (2) read at $k + 1$ gives $s \leq s_{2(k+1)-1} = s_{2k+1}$, so with the third computation above,

$$0 \leq s - s_{2k} \leq s_{2k+1} - s_{2k} = b_{2k+1} = b_{N+1}$$

If $N = 2k - 1$ is odd with $k \geq 1$, then part (2) read at k gives $s_{2k} \leq s \leq s_{2k-1}$, while the definition of s_N gives $s_{2k} = s_{2k-1} + (-1)^{2k+1} b_{2k} = s_{2k-1} - b_{2k}$. Hence

$$0 \leq s_{2k-1} - s \leq s_{2k-1} - s_{2k} = b_{2k} = b_{N+1}$$

In each case $|s - s_N| \leq b_{N+1}$, as required.

Part (2) brackets the sum between two consecutive partial sums, and part (3) turns that bracketing into a bound on the truncation error: the error made by stopping after N terms is at most the first term omitted. The prototype is the alternating harmonic series $\sum_{n \geq 1} (-1)^{n+1}/n$, whose terms $1/n$ decrease to 0, so its sum s exists and satisfies $s_4 = 7/12 \leq s \leq 5/6 = s_3$.

That series converges while $\sum_{n \geq 1} 1/n$ diverges (example 2.4.8), so it converges for a reason that no test in section 2.4 could have detected. Series of this kind are the subject of the rest of the section.

Definition 2.5.4: Conditional Convergence and Rearrangement

Let $\sum_{n \geq 1} a_n$ be a series of real numbers.

1. The series *converges conditionally* if $\sum_{n \geq 1} a_n$ converges and $\sum_{n \geq 1} |a_n|$ diverges.
2. For a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$, the series $\sum_{n \geq 1} a_{\sigma(n)}$ is called a *rearrangement* of $\sum_{n \geq 1} a_n$.

The alternating harmonic series is conditionally convergent, by the two facts recorded above: it converges by corollary 2.5.3, and the series of absolute values is the harmonic series, which diverges by example 2.4.8. So is $\sum_{n \geq 1} a_n/n$ for the $1, 1, -2$ pattern computed after theorem 2.5.2. A convergent series is by definition either absolutely or conditionally convergent, and theorem 2.4.12 rules out the fourth possibility of a divergent series whose absolute values are summable, so every series of real numbers falls into exactly one of three classes: absolutely convergent, conditionally convergent, divergent.

A rearrangement changes nothing about which terms occur, only the order in which they are added, so a first guess is that it changes nothing about the sum. For absolutely convergent series that guess is correct.

Theorem 2.5.5: Absolutely Convergent Series Are Rearrangement Invariant

Let $\sum_{n \geq 1} a_n$ converge absolutely, with sum s . Then for every bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ the rearrangement $\sum_{n \geq 1} a_{\sigma(n)}$ converges absolutely, and its sum is s .

Proof:

Write $t_N = \sum_{n=1}^N |a_n|$ and $t = \lim_N t_N$, which exists by the hypothesis of absolute convergence, and write $s_N = \sum_{n=1}^N a_n$.

Step 1: the rearranged series converges absolutely. Fix $M \geq 1$ and put $T = \{\sigma(1), \dots, \sigma(M)\}$, a set of M distinct positive integers because σ is injective. Re-indexing the finite sum by the bijection σ from $\{1, \dots, M\}$ onto T ,

$$\sum_{n=1}^M |a_{\sigma(n)}| = \sum_{m \in T} |a_m| \leq \sum_{m=1}^{\max T} |a_m| = t_{\max T} \leq t$$

the middle inequality because $T \subseteq \{1, \dots, \max T\}$ and every $|a_m|$ is nonnegative, and the last because (t_N) is nondecreasing with limit t , so proposition 2.1.8 gives $t_N \leq t$ for every N . Thus the partial sums of $\sum_n |a_{\sigma(n)}|$ are bounded above, and its terms are nonnegative, so it converges by proposition 2.4.4. By theorem 2.4.12 the rearrangement converges.

Step 2: the sum is unchanged. Let $\epsilon > 0$. Since $t_N \rightarrow t$ and $s_N \rightarrow s$, choose N with $t - t_N \leq \epsilon$ and $|s - s_N| \leq \epsilon$. Because σ is a bijection, each of $1, \dots, N$ is $\sigma(i)$ for exactly one i , so

$$P = \max \{\sigma^{-1}(1), \dots, \sigma^{-1}(N)\}$$

is a well-defined positive integer, and $\{1, \dots, N\} \subseteq T$ whenever $M \geq P$, where $T = \{\sigma(1), \dots, \sigma(M)\}$ as above.

Fix $M \geq P$ and set $S = T \setminus \{1, \dots, N\}$, a finite set of integers all of which exceed N . Re-indexing

as in Step 1 and cancelling the terms indexed by $\{1, \dots, N\}$,

$$\sum_{n=1}^M a_{\sigma(n)} - s_N = \sum_{m \in T} a_m - \sum_{m=1}^N a_m = \sum_{m \in S} a_m$$

If S is empty this difference is 0. Otherwise the finite triangle inequality proposition 2.4.5(2) and the inclusion $S \subseteq \{N+1, \dots, \max S\}$ give

$$\left| \sum_{m \in S} a_m \right| \leq \sum_{m \in S} |a_m| \leq \sum_{m=N+1}^{\max S} |a_m| = t_{\max S} - t_N \leq t - t_N \leq \epsilon$$

Combining the two displays,

$$\left| \sum_{n=1}^M a_{\sigma(n)} - s \right| \leq \left| \sum_{n=1}^M a_{\sigma(n)} - s_N \right| + |s_N - s| \leq 2\epsilon$$

for every $M \geq P$. As $\epsilon > 0$ was arbitrary, the partial sums of the rearranged series converge to s , completing the proof.

The hypothesis was used twice, and in both places it was used in the same way: the tail $t - t_N$ of the series of absolute values controls every finite sum of terms with index beyond N , no matter which such terms are chosen and no matter how many. A conditionally convergent series has no such control, since the partial sums of $\sum |a_n|$ are unbounded past every stage. The next theorem measures how completely that control is lost: for a conditionally convergent series, the pair consisting of the limit inferior and the limit superior of the rearranged partial sums can be prescribed arbitrarily.

Theorem 2.5.6: Riemann's Rearrangement Theorem

Let $\sum_{n \geq 1} a_n$ converge conditionally, and let $\alpha, \beta \in [-\infty, +\infty]$ with $\alpha \leq \beta$. Then there is a bijection $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ whose partial sums $s'_N = \sum_{n=1}^N a_{\sigma(n)}$ satisfy

$$\liminf_{N \rightarrow \infty} s'_N = \alpha, \quad \limsup_{N \rightarrow \infty} s'_N = \beta$$

Proof:

Step 1: the positive and negative parts both diverge. For $n \geq 1$ put

$$p_n = \frac{|a_n| + a_n}{2}, \quad q_n = \frac{|a_n| - a_n}{2}$$

so that $p_n - q_n = a_n$ and $p_n + q_n = |a_n|$, and both are nonnegative because $-|a_n| \leq a_n \leq |a_n|$ by proposition 1.3.8. Suppose first that $\sum p_n$ and $\sum q_n$ both converge. Then $\sum |a_n| = \sum (p_n + q_n)$ converges by proposition 2.4.5, contradicting conditional convergence. Suppose next that exactly one of them converges, say $\sum p_n$. Since $\sum a_n$ converges by hypothesis, proposition 2.4.5 makes $\sum (p_n - a_n) = \sum q_n$ convergent, again a contradiction, and the same argument with the roles exchanged rules out the case that only $\sum q_n$ converges. So both diverge, and since their terms are nonnegative, proposition 2.4.4 says that the partial sums of each are unbounded above.

Step 2: the two extracted sequences. Let $P_1, P_2, P_3, \dots$ be the nonnegative terms of (a_n) , listed

in the order in which they occur, and let $Q_1, Q_2, Q_3, \dots$ be the absolute values of the negative terms, again in their original order. Both lists are infinite: p_n is nonzero only where $a_n > 0$, so the divergence of $\sum p_n$ forces infinitely many a_n to be positive, and likewise the divergence of $\sum q_n$ forces infinitely many a_n to be negative. The partial sums of $\sum_j P_j$ are exactly the partial sums of $\sum_n p_n$ with the repetitions caused by the zero terms deleted, so the two sets of values coincide and the two suprema agree, and the same comparison holds for $\sum_j Q_j$ against $\sum_n q_n$. Hence the partial sums of $\sum_j P_j$ and of $\sum_j Q_j$ are unbounded above, and this remains true for the tail $\sum_{j>J} P_j$ for any fixed J , since its partial sums differ from those of $\sum_j P_j$ by the constant $P_1 + \dots + P_J$. Finally, $\sum_n a_n$ converges, so $a_n \rightarrow 0$ by proposition 2.4.3, and (P_j) and (Q_j) are the subsequences of (a_n) and of $(-a_n)$ along the nonnegative and negative terms, so $P_j \rightarrow 0$ and $Q_j \rightarrow 0$ by proposition 2.2.7.

Step 3: two sequences of real targets. Define real numbers u_n and v_n for $n \geq 1$ by

$$u_n = \begin{cases} \alpha - 1/n, & \alpha \in \mathbb{R} \\ -n, & \alpha = -\infty \\ n, & \alpha = +\infty \end{cases} \quad v_n = \begin{cases} \beta + 1/n, & \beta \in \mathbb{R} \\ n, & \beta = +\infty \\ -n, & \beta = -\infty \end{cases}$$

and put $u_0 = u_1$,

$$\alpha_n = u_n, \quad \beta_n = \max \left\{ v_n, u_n + \frac{1}{n}, u_{n-1} + \frac{1}{n} \right\}$$

Two inequalities hold by construction, for every $n \geq 1$:

$$\alpha_n < \beta_n \quad (\text{because } \beta_n \geq u_n + \frac{1}{n}), \quad \alpha_n < \beta_{n+1} \quad (\text{because } \beta_{n+1} \geq u_n + \frac{1}{n+1})$$

The behaviour of (β_n) is recorded case by case, and each case is a direct computation from the displayed formulas. If $\beta \in \mathbb{R}$ and $\alpha \in \mathbb{R}$, then for $n \geq 2$ one has $u_n + 1/n = \alpha \leq \beta < v_n$ and $u_{n-1} + 1/n < \alpha \leq \beta < v_n$, so $\beta_n = v_n = \beta + 1/n$ and $\beta_n \rightarrow \beta$. If $\beta \in \mathbb{R}$ and $\alpha = -\infty$, then $u_n + 1/n = -n + 1/n$ and $u_{n-1} + 1/n = -(n-1) + 1/n$ are both eventually smaller than $v_n = \beta + 1/n$, namely as soon as $n > 1 - \beta$, so again $\beta_n = v_n$ for all large n and $\beta_n \rightarrow \beta$. If $\beta = +\infty$, then $\beta_n \geq v_n = n$. If $\beta = -\infty$, then $\alpha = -\infty$ as well, and for $n \geq 2$ the largest of the three numbers is $u_{n-1} + 1/n$, so $\beta_n = -n + 1 + 1/n$. Correspondingly $\alpha_n = \alpha - 1/n \rightarrow \alpha$ when $\alpha \in \mathbb{R}$, while $\alpha_n = -n$ when $\alpha = -\infty$ and $\alpha_n = n$ when $\alpha = +\infty$.

Step 4: the rearrangement. Define integers $0 = m_0 < m_1 < m_2 < \dots$ and $0 = k_0 < k_1 < k_2 < \dots$ recursively. Given m_{n-1} and k_{n-1} , let m_n be the smallest integer greater than m_{n-1} for which

$$x_n := P_1 + \dots + P_{m_1} - Q_1 - \dots - Q_{k_1} + \dots + P_{m_{n-1}+1} + \dots + P_{m_n} > \beta_n$$

and then let k_n be the smallest integer greater than k_{n-1} for which

$$y_n := x_n - Q_{k_{n-1}+1} - \dots - Q_{k_n} < \alpha_n$$

Both minima exist: by Step 2 the partial sums of the tails $\sum_{j>m_{n-1}} P_j$ and $\sum_{j>k_{n-1}} Q_j$ are unbounded above, so the running sum can be pushed above β_n by adding enough terms P_j and then below α_n by subtracting enough terms Q_j . Listing the terms in the order

$$P_1, \dots, P_{m_1}, -Q_1, \dots, -Q_{k_1}, P_{m_1+1}, \dots, P_{m_2}, -Q_{k_1+1}, \dots, -Q_{k_2}, \dots$$

defines a map $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ sending the position in this list to the original index of the term standing there. It is injective because each P_j and each Q_j is listed once, and surjective because $m_n \rightarrow \infty$ and $k_n \rightarrow \infty$, so every P_j and every Q_j eventually appears. Thus σ is a bijection, and

(s'_N) denotes the partial sums of the rearranged series. By construction $y_0 = 0$, and x_n and y_n are among the s'_N .

Step 5: the two estimates. Fix $n \geq 2$. The stopping rule gives $x_n > \beta_n$. For the reverse bound, note that the partial sum standing one place before x_n in the list is $x_n - P_{m_n}$, and that it is at most β_n : if $m_n > m_{n-1} + 1$ this is the minimality of m_n , and if $m_n = m_{n-1} + 1$ that partial sum is y_{n-1} , which satisfies $y_{n-1} < \alpha_{n-1} < \beta_n$ by Step 3. Hence

$$|x_n - \beta_n| = x_n - \beta_n \leq P_{m_n} \quad (n \geq 2)$$

Similarly $y_n < \alpha_n$ for every $n \geq 1$, and the partial sum one place before y_n is $y_n + Q_{k_n}$, which is at least α_n : if $k_n > k_{n-1} + 1$ this is the minimality of k_n , and if $k_n = k_{n-1} + 1$ that partial sum is $x_n > \beta_n > \alpha_n$. Hence

$$|y_n - \alpha_n| = \alpha_n - y_n \leq Q_{k_n} \quad (n \geq 1)$$

Step 6: the limits. Call the positions occupied by $P_{m_{n-1}+1}, \dots, P_{m_n}, -Q_{k_{n-1}+1}, \dots, -Q_{k_n}$ the n -th block. Every index $N \geq 1$ lies in exactly one block. Every P_j is nonnegative and every Q_j is nonnegative, so inside the P -part of the n -th block the partial sums are nondecreasing, running from y_{n-1} up to x_n , and inside the Q -part they are nonincreasing, running from x_n down to y_n . Hence

$$\min\{y_{n-1}, y_n\} \leq s'_N \leq x_n \quad \text{for every } N \text{ in the } n\text{-th block}$$

The inequality $\limsup_N s'_N \leq \limsup_n x_n$. If $\limsup_n x_n = +\infty$ there is nothing to prove, so let $t \in \mathbb{R}$ satisfy $t > \limsup_n x_n$. By theorem 2.2.12(1) applied to (x_n) there is an n_0 with $x_n < t$ for every $n > n_0$. The blocks $1, \dots, n_0$ occupy finitely many positions, so there is an N_0 with every $N \geq N_0$ lying in a block of index $n > n_0$, and for such N the display gives $s'_N \leq \sup_{n > n_0} x_n \leq t$. Comparing (s'_N) with the constant sequence t in proposition 2.2.13(1) gives $\limsup_N s'_N \leq t$. As t was an arbitrary real above $\limsup_n x_n$, the inequality follows.

The reverse inequality. This is immediate from definition 2.2.10 alone, with no tail formula: (x_n) is a subsequence of (s'_N) , so every subsequential limit of (x_n) is a subsequential limit of (s'_N) , whence the set E_x is contained in $E_{s'}$ and $\limsup_n x_n = \sup E_x \leq \sup E_{s'} = \limsup_N s'_N$.

The two arguments applied to (y_n) , with the inequalities reversed and theorem 2.2.12(1) replaced by its $\liminf$ counterpart, give $\liminf_N s'_N = \liminf_n y_n$. Therefore $\limsup_N s'_N = \limsup_n x_n$ and $\liminf_N s'_N = \liminf_n y_n$.

It remains to evaluate these two quantities, and Step 5 does it in each of the three cases for β . If $\beta \in \mathbb{R}$, then $\beta_n \rightarrow \beta$ by Step 3 and $x_n - \beta_n \rightarrow 0$ by theorem 2.1.7 applied to $0 \leq x_n - \beta_n \leq P_{m_n}$ for $n \geq 2$, so $x_n \rightarrow \beta$ by theorem 2.1.6, and $\limsup_n x_n = \beta$ by theorem 2.2.12. If $\beta = +\infty$, then $x_n > \beta_n \geq n$, so $\sup_{n > n_0} x_n = +\infty$ for every n_0 and $\limsup_n x_n = +\infty$. If $\beta = -\infty$, then (P_{m_n}) converges to 0 and is therefore bounded by some G (proposition 2.1.5), so $x_n \leq \beta_n + P_{m_n} \leq -n + 1 + 1/n + G$ for $n \geq 2$, and since $1/n \leq 1$ there, $\sup_{n > n_0} x_n \leq -n_0 + 1 + G$ for every $n_0 \geq 1$, whence $\limsup_n x_n = -\infty$. The three cases for α are identical with (y_n) , (α_n) and (Q_{k_n}) in place of (x_n) , (β_n) and (P_{m_n}) , using $y_n < \alpha_n = -n$ when $\alpha = -\infty$ and $y_n \geq \alpha_n - Q_{k_n} \geq n - G'$ when $\alpha = +\infty$, where G' bounds (Q_{k_n}) . In every case $\limsup_n x_n = \beta$ and $\liminf_n y_n = \alpha$, which completes the proof.

Taking $\alpha = \beta = s$ for a real number s makes the limit inferior and the limit superior of the rearranged

partial sums both equal to s , so by theorem 2.2.12 the rearrangement converges to s : every real number is the sum of some rearrangement of a given conditionally convergent series. Taking $\alpha = \beta = +\infty$ produces a rearrangement whose partial sums tend to $+\infty$, and taking $\alpha = -\infty$, $\beta = +\infty$ produces one whose partial sums are unbounded in both directions. Together with theorem 2.5.5, this makes the distinction between absolute and conditional convergence exact: a convergent series has a sum independent of the order of its terms if and only if it converges absolutely.

The proof reorders the terms by an algorithm, and does not exhibit the reordering in closed form. For the alternating harmonic series there is a reordering with a simple description whose sum can be computed exactly, and computed without any information about what the original sum is.

Example 2.5.7: The Alternating Harmonic Series Rearranged Two Positive, One Negative

Let $s = \sum_{n \geq 1} (-1)^{n+1}/n$, which exists by corollary 2.5.3, and let s_N be its partial sums. Consider the rearrangement that takes two positive terms, then one negative term, and repeats:

$$1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \frac{1}{9} + \frac{1}{11} - \frac{1}{6} + \cdots$$

This is a rearrangement in the sense of definition 2.5.4: the reciprocals of the odd numbers appear once each, in order, and the negatives of the reciprocals of the even numbers appear once each, in order.

Write $H_n = \sum_{j=1}^n 1/j$ and let t_N denote the partial sums of the rearranged series. After k groups the terms used are $1, 1/3, \dots, 1/(4k-1)$ and $-1/2, -1/4, \dots, -1/(2k)$, so

$$t_{3k} = \sum_{j=1}^{2k} \frac{1}{2j-1} - \sum_{j=1}^k \frac{1}{2j}$$

The first sum is H_{4k} with the even-denominator terms removed, and those terms total $\sum_{j=1}^{2k} 1/(2j) = \frac{1}{2} H_{2k}$, so it equals $H_{4k} - \frac{1}{2} H_{2k}$. The second sum is $\frac{1}{2} H_k$. Hence

$$t_{3k} = H_{4k} - \frac{1}{2} H_{2k} - \frac{1}{2} H_k = (H_{4k} - H_{2k}) + \frac{1}{2} (H_{2k} - H_k)$$

Now $s_{2n} = H_{2n} - H_n$ for every n , because $s_{2n} = \sum_{m=1}^{2n} 1/m - 2 \sum_{j=1}^n 1/(2j) = H_{2n} - H_n$. Applying this identity with $n = 2k$ to the first bracket of the previous display, and with $n = k$ to the second, gives

$$t_{3k} = s_{4k} + \frac{1}{2} s_{2k}$$

Both s_{4k} and s_{2k} converge to s by proposition 2.2.7, so $t_{3k} \rightarrow s + \frac{1}{2}s = \frac{3}{2}s$ by theorem 2.1.6. The two intermediate partial sums satisfy $t_{3k+1} = t_{3k} + 1/(4k+1)$ and $t_{3k+2} = t_{3k+1} + 1/(4k+3)$, and both correction terms converge to 0, so $t_{3k+1} \rightarrow \frac{3}{2}s$ and $t_{3k+2} \rightarrow \frac{3}{2}s$ as well. Every index $N \geq 1$ is of exactly one of the three forms $3k$, $3k+1$, $3k+2$, so given $\epsilon > 0$ each of the three subsequences lies within ϵ of $\frac{3}{2}s$ from some index onward, and beyond the largest of those three indices every t_N does. Hence $t_N \rightarrow \frac{3}{2}s$.

Finally $\frac{3}{2}s \neq s$, since corollary 2.5.3 gives $s \geq s_2 = 1/2 > 0$. The rearranged series therefore converges, and to a different number: reordering has multiplied the sum by $3/2$.

The rest of the section concerns two further interchanges, both of which are governed by absolute convergence in the same way theorem 2.5.5 is. The first is the interchange of two summations.

Theorem 2.5.8: Fubini's Theorem for Double Series

Let $a_{ij} \in \mathbb{R}$ be given for all integers $i, j \geq 1$.

1. For all $M, N \geq 1$, $\sum_{i=1}^M \sum_{j=1}^N a_{ij} = \sum_{j=1}^N \sum_{i=1}^M a_{ij}$.
2. Suppose that for each i the series $\sum_{j \geq 1} |a_{ij}|$ converges, with sum b_i , and that $\sum_{i \geq 1} b_i$ converges. Then every row sum $r_i = \sum_{j \geq 1} a_{ij}$ and every column sum $c_j = \sum_{i \geq 1} a_{ij}$ is defined, the series $\sum_{i \geq 1} r_i$ and $\sum_{j \geq 1} c_j$ both converge absolutely, and

$$\sum_{i \geq 1} \left(\sum_{j \geq 1} a_{ij} \right) = \sum_{j \geq 1} \left(\sum_{i \geq 1} a_{ij} \right)$$

Proof:

1. Fix N and induct on M . For $M = 1$ both sides equal $\sum_{j=1}^N a_{1j}$. Assume the identity for M . Splitting each finite sum $\sum_{i=1}^{M+1}$ into $\sum_{i=1}^M$ plus the term $i = M + 1$, and using that a finite sum of finite sums may be split termwise,

$$\sum_{j=1}^N \sum_{i=1}^{M+1} a_{ij} = \sum_{j=1}^N \left(\sum_{i=1}^M a_{ij} + a_{M+1,j} \right) = \sum_{j=1}^N \sum_{i=1}^M a_{ij} + \sum_{j=1}^N a_{M+1,j}$$

By the inductive hypothesis the first term equals $\sum_{i=1}^M \sum_{j=1}^N a_{ij}$, and adding the second term gives $\sum_{i=1}^{M+1} \sum_{j=1}^N a_{ij}$. This is the identity for $M + 1$.

2. Write $S = \sum_{i \geq 1} b_i$ and record once a bound used several times below. If $\sum_{i \geq 1} u_i$ and $\sum_{i \geq 1} |u_i|$ both converge, then applying the finite triangle inequality proposition 2.4.5(2) in the form $-\sum_{i=1}^M |u_i| \leq \sum_{i=1}^M u_i \leq \sum_{i=1}^M |u_i|$ and letting $M \rightarrow \infty$ in both inequalities using proposition 2.1.8 gives $|\sum_{i \geq 1} u_i| \leq \sum_{i \geq 1} |u_i|$.

The row sums are defined: $\sum_j |a_{ij}|$ converges by hypothesis, so $\sum_j a_{ij}$ converges by theorem 2.4.12, and the bound just recorded gives $|r_i| \leq b_i$. Since $\sum_i b_i$ converges, theorem 2.4.13 makes $\sum_i |r_i|$ convergent, so $\sum_i r_i$ converges absolutely. Write $R = \sum_{i \geq 1} r_i$.

The column sums are defined: for fixed j , the number $|a_{ij}|$ is one of the nonnegative terms whose sum is b_i , so $|a_{ij}| \leq b_i$ and therefore $\sum_{i=1}^M |a_{ij}| \leq \sum_{i=1}^M b_i \leq S$ for every M , the last inequality because the partial sums of the nonnegative convergent series $\sum_i b_i$ are bounded by its sum (proposition 2.1.8). Hence $\sum_i |a_{ij}|$ converges by proposition 2.4.4 and $\sum_i a_{ij}$ converges. For absolute convergence of $\sum_j c_j$, fix N . Then $|c_j| \leq \sum_{i \geq 1} |a_{ij}|$ by the recorded bound, and by theorem 2.1.6 applied to the N convergent sequences involved, together with part (1),

$$\sum_{j=1}^N \sum_{i \geq 1} |a_{ij}| = \lim_{M \rightarrow \infty} \sum_{j=1}^N \sum_{i=1}^M |a_{ij}| = \lim_{M \rightarrow \infty} \sum_{i=1}^M \sum_{j=1}^N |a_{ij}| \leq S$$

the last step because each term of the sequence is at most $\sum_{i=1}^M b_i \leq S$, so proposition 2.1.8 bounds the limit by S . Hence $\sum_{j=1}^N |c_j| \leq S$ for every N , and $\sum_j c_j$ converges absolutely by proposition 2.4.4.

It remains to prove $\sum_{j \geq 1} c_j = R$. The same interchange of a finite sum with a limit, now without absolute values, gives for each N

$$\sum_{j=1}^N c_j = \lim_{M \rightarrow \infty} \sum_{i=1}^M \sum_{j=1}^N a_{ij} = \sum_{i \geq 1} \left(\sum_{j=1}^N a_{ij} \right)$$

the last series converging because the limit exists. Subtracting $R = \sum_i r_i$ term by term, which proposition 2.4.5 permits since both series converge,

$$\sum_{j=1}^N c_j - R = \sum_{i \geq 1} \left(\sum_{j=1}^N a_{ij} - r_i \right) = - \sum_{i \geq 1} \rho_i^{(N)}, \quad \rho_i^{(N)} := \sum_{j > N} a_{ij}$$

where $\rho_i^{(N)}$ is the tail of a convergent series and satisfies $|\rho_i^{(N)}| \leq \sum_{j > N} |a_{ij}| \leq b_i$ by the recorded bound.

Let $\epsilon > 0$. Since $\sum_i b_i$ converges, choose I with $\sum_{i > I} b_i \leq \epsilon$. For each $i \leq I$ the series $\sum_j |a_{ij}|$ converges, so its tails tend to 0 and there is a J_i with $\sum_{j > J_i} |a_{ij}| \leq \epsilon/I$. Put $J = \max\{J_1, \dots, J_I\}$ and fix $N \geq J$. Then for each $i \leq I$, $|\rho_i^{(N)}| \leq \sum_{j > N} |a_{ij}| \leq \sum_{j > J_i} |a_{ij}| \leq \epsilon/I$, while for $i > I$ only $|\rho_i^{(N)}| \leq b_i$ is used. Applying the recorded bound to $\sum_i \rho_i^{(N)}$, whose absolute values are dominated by the summable b_i , and splitting the resulting sum at $i = I$,

$$\left| \sum_{j=1}^N c_j - R \right| \leq \sum_{i=1}^I |\rho_i^{(N)}| + \sum_{i > I} b_i \leq I \cdot \frac{\epsilon}{I} + \epsilon = 2\epsilon$$

Since $\epsilon > 0$ was arbitrary, $\sum_{j \geq 1} c_j = R$, which is the asserted equality of the two iterated sums, completing the proof.

Part (1) is the statement that a finite rectangular array may be added by rows or by columns, and part (2) says that the same holds for an infinite array as soon as the row sums of absolute values are themselves summable. Without that hypothesis the two iterated sums can differ, and exercise 2.5.1 (8) asks for an array where they do.

The second interchange multiplies two series. Expanding the product of two finite sums and collecting the terms $a_i b_j$ by the value of $i + j$ produces the following series, and the question is whether it converges to the product of the two sums when the sums are infinite.

Theorem 2.5.9: Mertens' Theorem on Cauchy Products

Let $\sum_{n \geq 1} a_n$ converge absolutely with sum A , and let $\sum_{n \geq 1} b_n$ converge with sum B . Define the *Cauchy product* of the two series to be $\sum_{n \geq 1} c_n$, where

$$c_n = \sum_{k=1}^n a_k b_{n+1-k}$$

so that c_n collects exactly the products $a_i b_j$ with $i + j = n + 1$. Then $\sum_{n \geq 1} c_n$ converges, with sum AB .

Proof:

Write $A_N = \sum_{n=1}^N a_n$, $B_N = \sum_{n=1}^N b_n$ and $C_N = \sum_{n=1}^N c_n$, and put $\gamma_N = B_N - B$, so that $\gamma_N \rightarrow 0$ by theorem 2.1.6.

Step 1: an expression for C_N . Define $d_{nk} = a_k b_{n+1-k}$ for $1 \leq k \leq n \leq N$ and $d_{nk} = 0$ for $1 \leq n < k \leq N$. Then $c_n = \sum_{k=1}^N d_{nk}$ for each $n \leq N$, so by theorem 2.5.8(1) applied to the square array (d_{nk}) ,

$$C_N = \sum_{n=1}^N \sum_{k=1}^N d_{nk} = \sum_{k=1}^N \sum_{n=1}^N d_{nk} = \sum_{k=1}^N a_k \sum_{n=k}^N b_{n+1-k} = \sum_{k=1}^N a_k B_{N+1-k}$$

where the inner sum was re-indexed by $j = n + 1 - k$, which runs over $1 \leq j \leq N + 1 - k$. Substituting $B_{N+1-k} = B + \gamma_{N+1-k}$ and separating the two parts,

$$C_N = BA_N + \delta_N, \quad \delta_N := \sum_{k=1}^N a_k \gamma_{N+1-k}$$

Step 2: $\delta_N \rightarrow 0$. Let $\tau = \sum_{n \geq 1} |a_n|$, finite by hypothesis, and let $\tau_N = \sum_{n=1}^N |a_n|$. The sequence (γ_m) converges, hence is bounded (proposition 2.1.5), say $|\gamma_m| \leq G$ for all m .

Let $\epsilon > 0$. Choose M with $|\gamma_m| \leq \epsilon$ for all $m > M$. Since $\tau_N \rightarrow \tau$, choose $N_0 > M$ with $\tau_N - \tau_{N-M} \leq \epsilon$ for all $N \geq N_0$, which is possible because $\tau_N - \tau_{N-M} \rightarrow \tau - \tau = 0$ by theorem 2.1.6. Fix $N \geq N_0$ and split the sum defining δ_N at $k = N - M$. For $1 \leq k \leq N - M$ the index $N + 1 - k$ exceeds M , so $|\gamma_{N+1-k}| \leq \epsilon$, while for $N - M < k \leq N$ only the crude bound G is used. The finite triangle inequality proposition 2.4.5(2) then gives

$$|\delta_N| \leq \epsilon \sum_{k=1}^{N-M} |a_k| + G \sum_{k=N-M+1}^N |a_k| \leq \epsilon \tau + G(\tau_N - \tau_{N-M}) \leq \epsilon(\tau + G)$$

As $\tau + G$ does not depend on ϵ and $\epsilon > 0$ was arbitrary, $\delta_N \rightarrow 0$.

By Step 1 and theorem 2.1.6, $C_N = BA_N + \delta_N \rightarrow BA + 0 = AB$, completing the proof.

Absolute convergence is assumed of one factor only, and the theorem is silent when neither factor converges absolutely. It has to be: two conditionally convergent series can have a Cauchy product whose terms do not even tend to 0.

Example 2.5.10: A Divergent Cauchy Product of Two Convergent Series

Let $a_n = b_n = (-1)^{n+1}/\sqrt{n}$ for $n \geq 1$, where $\sqrt{n}$ is the nonnegative square root given by theorem 1.3.12. Square roots preserve the order on $[0, \infty)$: if $0 \leq x < y$ then $x^2 < y^2$ by proposition 1.1.2, so $\sqrt{u} > \sqrt{v} \geq 0$ would force $u > v$, and therefore $0 \leq u \leq v$ implies $\sqrt{u} \leq \sqrt{v}$.

The sequence $1/\sqrt{n}$ decreases to 0, so $\sum_{n \geq 1} a_n$ converges by corollary 2.5.3. It does not converge absolutely: $n \leq n^2$ for $n \geq 1$ gives $\sqrt{n} \leq n$ by the order fact just recorded, hence $1/\sqrt{n} \geq 1/n$ by proposition 1.1.2, and theorem 2.4.13 with example 2.4.8 makes $\sum_n 1/\sqrt{n}$ divergent. So both factors are conditionally convergent.

Their Cauchy product has terms

$$c_n = \sum_{k=1}^n \frac{(-1)^{k+1}}{\sqrt{k}} \cdot \frac{(-1)^{n+2-k}}{\sqrt{n+1-k}} = (-1)^{n+1} \sum_{k=1}^n \frac{1}{\sqrt{k}\sqrt{n+1-k}}$$

since $(-1)^{k+1}(-1)^{n+2-k} = (-1)^{n+3} = (-1)^{n+1}$. For $1 \leq k \leq n$, the uniqueness clause of theorem 1.3.12 identifies $\sqrt{k}\sqrt{n+1-k}$ with $\sqrt{k(n+1-k)}$, because the left-hand side is nonnegative and its square is $k(n+1-k)$. Writing $m = n+1$, the algebraic identity

$$\frac{m^2}{4} - k(m-k) = \frac{m^2 - 4km + 4k^2}{4} = \frac{(m-2k)^2}{4} \geq 0$$

gives $k(n+1-k) \leq \left(\frac{n+1}{2}\right)^2$. By the order fact recorded at the start, $\sqrt{k(n+1-k)} \leq \frac{n+1}{2}$, so each of the n summands is at least $\frac{2}{n+1}$ and

$$|c_n| \geq n \cdot \frac{2}{n+1} = \frac{2n}{n+1} \geq 1$$

for every $n \geq 1$. The terms c_n therefore do not converge to 0, and $\sum_{n \geq 1} c_n$ diverges by proposition 2.4.3.

Exercise 2.5.1

1. Let $s = \sum_{n \geq 1} (-1)^{n+1}/n$. Using only corollary 2.5.3, prove the bounds $1/2 \leq s \leq 1$ and then the sharper bounds $7/12 \leq s \leq 5/6$. Show more generally that for a sequence $b_1 \geq b_2 \geq \dots$ with $b_n \rightarrow 0$, the sum s of $\sum_{n \geq 1} (-1)^{n+1}b_n$ satisfies $b_1 - b_2 \leq s \leq b_1$.
2. Let $p \geq 1$ be an integer and let $(a_n)_{n \geq 1}$ be periodic with period p , meaning $a_{n+p} = a_n$ for every n , and suppose $a_1 + a_2 + \dots + a_p = 0$. Prove that $\sum_{n \geq 1} a_n/n$ converges. Prove furthermore that if $a_n \neq 0$ for at least one n , then the convergence is conditional in the sense of definition 2.5.4.
3. Show that the monotonicity hypothesis (2) of theorem 2.5.2 cannot be dropped, even when the remaining hypotheses are strengthened to $b_n > 0$ for all n . Specifically, let $a_n = (-1)^{n+1}$ and let $b_n = 1/n$ for odd n and $b_n = 1/n^2$ for even n . Verify that $b_n \rightarrow 0$ and that (A_N) is bounded, and prove that $\sum_{n \geq 1} a_n b_n$ diverges.
4. With s as in exercise 2.5.1 (1), consider the rearrangement of $\sum_{n \geq 1} (-1)^{n+1}/n$ that takes one

positive term followed by two negative terms and repeats:

$$1 - \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{6} - \frac{1}{8} + \frac{1}{5} - \frac{1}{10} - \frac{1}{12} + \cdots$$

Show that it converges, and that its sum is $\frac{1}{2}s$. (Follow the pattern of example 2.5.7, computing the partial sum after k groups in terms of $H_n = \sum_{j=1}^n 1/j$.)

5. Prove *Abel's test*: if $\sum_{n \geq 1} a_n$ converges and (b_n) is monotone and bounded, then $\sum_{n \geq 1} a_n b_n$ converges. (Treat the nonincreasing and nondecreasing cases separately. For a nonincreasing (b_n) write $b_n = b + c_n$ with $b = \lim_n b_n$, so that (c_n) is nonincreasing with limit 0. For a nondecreasing (b_n) put $c_n = b - b_n$ instead. Either way (c_n) is nonnegative, nonincreasing and tends to 0.)
6. Let $|x| < 1$. Compute the Cauchy product of $\sum_{n \geq 1} x^{n-1}$ with itself, and use theorem 2.5.9 to conclude that $\sum_{n \geq 1} nx^{n-1}$ converges with sum $1/(1-x)^2$. Explain why theorem 2.5.9 applies.
7. Suppose $\sum_{n \geq 1} a_n$ converges and every rearrangement of it converges as well. Prove that $\sum_{n \geq 1} a_n$ converges absolutely. (The converse is theorem 2.5.5.)
8. Define $a_{ij} = 1$ if $i = j$, $a_{ij} = -1$ if $i = j + 1$, and $a_{ij} = 0$ otherwise. Compute every row sum $r_i = \sum_{j \geq 1} a_{ij}$ and every column sum $c_j = \sum_{i \geq 1} a_{ij}$, and deduce that $\sum_i \sum_j a_{ij} \neq \sum_j \sum_i a_{ij}$. Identify precisely which hypothesis of theorem 2.5.8(2) fails.

2.6 Power Series and the Radius of Convergence

The tests of section 2.4 decide one series at a time. A power series presents a whole family at once, one series for each real x , and the question is which members of the family converge. The answer is a dichotomy with a single exception: there is a number R computed from the coefficients alone, such that the series converges absolutely at every x within distance R of the centre and diverges at every x farther away. Only the two points at distance exactly R are left undecided, and example 2.6.4 exhibits all three of the behaviours that can occur there.

The same computation is what makes the exponential and the trigonometric functions available. Each is defined below by a power series whose radius is infinite, and the identity $\exp(x+y) = \exp(x)\exp(y)$, which is what makes $\exp$ an exponential, is the Cauchy product of section 2.5 evaluated with the binomial theorem.

The coefficient whose n -th root is wanted is usually a constant, a power of n , or a product of the two. Two limits cover both cases, and both of their proofs use one fact about powers: for $s, t \geq 0$ and an integer $n \geq 1$,

$$s < t \quad \Rightarrow \quad s^n < t^n \tag{2.5}$$

For $n = 1$ this is the hypothesis itself. If it holds for n , then $t > s \geq 0$ gives $t > 0$, so multiplying $s^n < t^n$ by $s \geq 0$ gives $s^{n+1} \leq s t^n$ and multiplying $s < t$ by $t^n > 0$ gives $s t^n < t^{n+1}$, both by proposition 1.1.2(5), whence $s^{n+1} < t^{n+1}$. Taking contrapositives, $s^n \leq t^n$ implies $s \leq t$ for $s, t \geq 0$. Together with theorem 1.3.12, which produces a unique nonnegative n -th root of every nonnegative real, (2.5) says that raising to the n -th power and extracting the nonnegative n -th root both preserve the order on the nonnegative reals.

Proposition 2.6.1: Two Special Limits

1. $\lim_{n \rightarrow \infty} n^{1/n} = 1$.
2. $\lim_{n \rightarrow \infty} c^{1/n} = 1$ for every real $c > 0$.

Proof:

1. For $n \geq 1$ put $x_n = n^{1/n} - 1$. If $n^{1/n} < 1$ then (2.5), together with the identity $(n^{1/n})^n = n$ of theorem 1.3.12, gives $n = (n^{1/n})^n < 1^n = 1$, which contradicts $n \geq 1$, so $x_n \geq 0$. Let $n \geq 2$. The binomial theorem [4, the binomial theorem] expands

$$n = (1 + x_n)^n = \sum_{k=0}^n \binom{n}{k} x_n^k \geq \binom{n}{2} x_n^2 = \frac{n(n-1)}{2} x_n^2$$

where the inequality discards terms that are all nonnegative, since $x_n \geq 0$. Dividing by the positive number $n(n-1)/2$ gives

$$0 \leq x_n^2 \leq \frac{2}{n-1}$$

Now let $\epsilon > 0$. By theorem 1.4.1(2) there is an $N \in \mathbb{N}$ with $N > 1 + 2/\epsilon^2$, and $N \geq 2$ because $1 + 2/\epsilon^2 > 1$. For $n \geq N$ one has $n-1 > 2/\epsilon^2$ and therefore $x_n^2 \leq 2/(n-1) < \epsilon^2$. Were $x_n \geq \epsilon$, then (2.5) with exponent 2 would give $x_n^2 \geq \epsilon^2$, so in fact $x_n < \epsilon$. Thus $|n^{1/n} - 1| = x_n < \epsilon$ for all $n \geq N$, which is definition 2.1.2 for the limit 1.

2. For $c = 1$ the sequence is constant, since $1^n = 1$ and n -th roots are unique by theorem 1.3.12. Let $c > 1$ and put $y_n = c^{1/n} - 1$. If $c^{1/n} \leq 1$ then the contrapositive form of (2.5) gives $c = (c^{1/n})^n \leq 1^n = 1$, contradicting $c > 1$, so $y_n > 0$. The binomial expansion of $(1 + y_n)^n$ has nonnegative terms, and discarding all but the first two of them gives $c = (1 + y_n)^n \geq 1 + ny_n$, so

$$0 < y_n \leq \frac{c-1}{n}$$

Given $\epsilon > 0$, choose $N \in \mathbb{N}$ with $N > (c-1)/\epsilon$ by theorem 1.4.1(2). For $n \geq N$ one has $1/n \leq 1/N$ by proposition 1.1.2(7), so $y_n \leq (c-1)/N < \epsilon$, the last inequality being $N > (c-1)/\epsilon$ multiplied by the positive number ϵ/N . Hence $c^{1/n} \rightarrow 1$.

Finally let $0 < c < 1$ and put $d = 1/c$, so that $d > 1$ by proposition 1.1.2(7). The number $1/d^{1/n}$ is positive and satisfies $(1/d^{1/n})^n = 1/(d^{1/n})^n = 1/d = c$, so uniqueness in theorem 1.3.12 identifies it with $c^{1/n}$. The previous case gives $d^{1/n} \rightarrow 1$, and since the limit 1 is not 0 the quotient rule of theorem 2.1.6 gives $c^{1/n} = 1/d^{1/n} \rightarrow 1$, as we sought to show.

The two limits together control every coefficient sequence built from a constant and a power of n . For $c > 0$ and an integer k , the number $c^{1/n} (n^{1/n})^k$ is positive and its n -th power is cn^k , so uniqueness in theorem 1.3.12 gives $(cn^k)^{1/n} = c^{1/n} (n^{1/n})^k$, and theorem 2.1.6 then sends the right-hand side to 1. Coefficients of that shape all produce the same radius below, whatever c and k are.

A power series is written down by naming its coefficients, and everything about its convergence is computable from them. The definition accordingly attaches a number to the coefficient sequence first, and the theorem afterwards says what that number does.

Definition 2.6.2: Power Series and its Radius of Convergence

Let $a \in \mathbb{R}$ and let $(c_n)_{n \geq 0}$ be real numbers. The *power series with coefficients (c_n) centred at a* assigns to each $x \in \mathbb{R}$ the series

$$\sum_{n \geq 0} c_n (x - a)^n$$

in which $(x - a)^0$ is read as 1 for every x , including $x = a$. Put

$$\alpha = \limsup_{n \rightarrow \infty} |c_n|^{1/n}$$

the limit superior being taken over $n \geq 1$, where the root is defined. Every term of that sequence is nonnegative, so $\alpha \in [0, +\infty]$. The *radius of convergence* of the power series is

$$R = \begin{cases} +\infty & \text{if } \alpha = 0 \\ 1/\alpha & \text{if } 0 < \alpha < +\infty \\ 0 & \text{if } \alpha = +\infty \end{cases}$$

Two conventions apply below. The coefficients are indexed from $n = 0$, where section 2.4 started at $n = 1$, and every statement of section 2.4 and section 2.5 about a series $\sum_{n \geq 1} a_n$ applies after the relabelling $b_n = a_{n-1}$, which leaves the sequence of partial sums unchanged apart from its indexing. That relabelling is left implicit below. The extended values 0 and $+\infty$ for α are the ones given by definition 2.2.4, and both of them occur.

Take $a = 0$ and $c_n = 1$ for every n , so that the power series is $\sum_{n \geq 0} x^n$. The sequence $|c_n|^{1/n}$ is constantly 1, and a convergent sequence has its limit for a limit superior: by proposition 2.2.7 every subsequence of a convergent sequence has the same limit, while by theorem 2.2.12 the limit superior is the limit of some subsequence. So $\alpha = 1$ and $R = 1$. The geometric series of example 2.4.7 converges for $|x| < 1$, and for $|x| \geq 1$ the contrapositive form of (2.5) gives $|x^n| = |x|^n \geq 1^n = 1$, so the terms do not tend to 0 and proposition 2.4.3 makes the series divergent. The set of convergence is exactly $(-1, 1)$, which is the pattern the next theorem asserts in general.

Theorem 2.6.3: Cauchy-Hadamard: Convergence Inside the Radius, Divergence Outside

Let $\sum_{n \geq 0} c_n (x - a)^n$ be a power series with radius of convergence R as in definition 2.6.2, and let $x \in \mathbb{R}$.

1. If $|x - a| < R$, the series converges absolutely.
2. If $|x - a| > R$, the series diverges.

Proof:

Write $\alpha = \limsup_{n \rightarrow \infty} |c_n|^{1/n}$, so that R is the value attached to α in definition 2.6.2.

1. Suppose $|x - a| < R$. Then $R > 0$, so $\alpha < +\infty$. The first task is to produce a real ρ with

$$|x - a| < \rho \quad \text{and} \quad \frac{1}{\rho} > \alpha$$

If $\alpha = 0$, take $\rho = |x - a| + 1$, which is positive, so that $1/\rho > 0 = \alpha$. If $0 < \alpha < +\infty$ then $R = 1/\alpha$ and $|x - a| < 1/\alpha$, so $\rho = (|x - a| + 1/\alpha)(1 + 1)^{-1}$ satisfies $|x - a| < \rho < 1/\alpha$ by proposition 1.1.2(8), and then $0 < \rho < 1/\alpha$ gives $1/\rho > \alpha$ by proposition 1.1.2(7).

Since $1/\rho > \alpha$, the eventual domination clause of theorem 2.2.12 gives an $N \in \mathbb{N}$ with $|c_n|^{1/n} < 1/\rho$ for all $n \geq N$. Raising this to the n -th power with (2.5), and using theorem 1.3.12 to write $|c_n| = (|c_n|^{1/n})^n$, gives $|c_n| < \rho^{-n}$, that is $|c_n|\rho^n < 1$, for all $n \geq N$.

Put $r = |x - a|/\rho$, so that $0 \leq r < 1$, and put

$$M = 1 + \sum_{n=0}^{N-1} |c_n|\rho^n$$

a finite sum of nonnegative numbers increased by 1. For $n < N$ the number $|c_n|\rho^n$ is one of the summands, so $|c_n|\rho^n \leq M$, and for $n \geq N$ the previous paragraph gives $|c_n|\rho^n < 1 \leq M$. Since $|x - a| = \rho r$, applying the multiplicativity of proposition 1.3.8(3) n times yields, for every $n \geq 0$,

$$|c_n(x - a)^n| = |c_n||x - a|^n = (|c_n|\rho^n)r^n \leq Mr^n$$

The geometric series $\sum_{n \geq 0} r^n$ converges because $0 \leq r < 1$ (example 2.4.7), so $\sum_{n \geq 0} Mr^n$ converges by the scalar rule of proposition 2.4.5, and theorem 2.4.13 then makes $\sum_{n \geq 0} |c_n(x - a)^n|$ convergent. That is absolute convergence in the sense of definition 2.4.11.

2. Suppose $|x - a| > R$. Then $|x - a| > 0$, so $\beta = 1/|x - a|$ is a positive real, and $\beta < \alpha$ in each of the three cases of definition 2.6.2: if $\alpha = 0$ then $R = +\infty$ and no x satisfies the hypothesis, so there is nothing to prove; if $0 < \alpha < +\infty$ then $|x - a| > R = 1/\alpha > 0$, which proposition 1.1.2(7) inverts to $\beta < \alpha$; if $\alpha = +\infty$ then $\beta < \alpha$ because β is a real number.

By theorem 2.2.12 the value α is the limit of a subsequence $(|c_{n_k}|^{1/n_k})_{k \geq 1}$, in the extended sense that this subsequence exceeds every bound when $\alpha = +\infty$. In either case there is a K with $|c_{n_k}|^{1/n_k} > \beta$ for all $k \geq K$: for finite α apply definition 2.1.2 with $\epsilon = \alpha - \beta > 0$, and for $\alpha = +\infty$ apply the unboundedness with the bound β .

Raising this to the n_k -th power with (2.5) gives $|c_{n_k}| > \beta^{n_k}$, and multiplying by the positive number $|x - a|^{n_k}$ (proposition 1.1.2(5)) gives

$$|c_{n_k}(x - a)^{n_k}| > \beta^{n_k}|x - a|^{n_k} = (\beta|x - a|)^{n_k} = 1$$

for every $k \geq K$. So infinitely many terms of the series have absolute value larger than 1. Were the terms to tend to 0, all but finitely many would have absolute value below 1, so the terms do not tend to 0, and proposition 2.4.3 makes the series divergent, completing the proof.

The theorem justifies the word radius. Write S for the set of $|x - a|$ with x a point at which the series converges. Part (1) gives $[0, R) \subseteq S$, part (2) gives $S \subseteq [0, R]$, and $0 \in S$ because at $x = a$ every term after the first vanishes and the partial sums are constantly c_0 . When $R = +\infty$ the first inclusion makes S unbounded. When R is finite, the second inclusion gives $\sup S \leq R$, and the reverse inequality holds as well: an upper bound b of S satisfies $b \geq 0$ because $0 \in S$, and $b < R$ would be contradicted by the number $(b + R)(1 + 1)^{-1}$, which lies in $[0, R) \subseteq S$ and exceeds b by proposition 1.1.2(8). So $\sup S = R$. When $0 < R < +\infty$ the set of points of convergence therefore contains the open interval $(a - R, a + R)$ and is contained in the closed interval $[a - R, a + R]$, and the two endpoints are the only points theorem 2.6.3 leaves undecided. The degenerate cases were settled above and leave nothing undecided: for $R = +\infty$ the series converges at every real, and for $R = 0$ it converges at $x = a$ alone.

Example 2.6.4: Five Radii, and Three Behaviours at an Endpoint

All five power series below are centred at $a = 0$.

1. $\sum_{n \geq 0} n! x^n$ has $R = 0$. Fix $x \neq 0$ and use theorem 1.4.1(2) to choose $N \in \mathbb{N}$ with $N > 2/|x|$, so that $j|x| > N|x| > 2$ for every integer $j > N$. Passing from n to $n + 1$ multiplies $|n! x^n|$ by $(n + 1)|x|$, which exceeds $2 > 1$ once $n \geq N$, so induction gives $|n! x^n| \geq |N! x^N|$ for every $n \geq N$. Since $|N! x^N| > 0$, the terms do not tend to 0, and proposition 2.4.3 gives divergence at every $x \neq 0$. If R were positive there would be an $x \neq 0$ with $|x| < R$, namely $x = R/2$ for finite R and $x = 1$ for $R = +\infty$, and part (1) of theorem 2.6.3 would make the series converge there. So $R = 0$. The formula of definition 2.6.2 is not evaluated directly here, because the root $(n!)^{1/n}$ is not among the limits available.
2. $\sum_{n \geq 0} x^n/n!$ has $R = +\infty$. At $x = 0$ every term after the first vanishes. For $x \neq 0$ every term is nonzero and the ratios are

$$\frac{|x^{n+1}/(n+1)!|}{|x^n/n!|} = \frac{|x|}{n+1}$$

which tend to 0: given $\epsilon > 0$, theorem 1.4.1(2) provides an N with $N > |x|/\epsilon$, and then $|x|/(n+1) < |x|/N < \epsilon$ for $n \geq N$. So theorem 2.4.14 gives absolute convergence at every $x \neq 0$. The series therefore converges at every real point, and a finite R would contradict part (2) of theorem 2.6.3 at any x with $|x| > R$, one of which exists by theorem 1.4.1(2).

3. $\sum_{n \geq 0} x^n$ has $R = 1$, since every $|c_n|^{1/n}$ equals 1, a constant sequence converges to its value, and a convergent sequence has its limit for a limit superior (proposition 2.2.7 and theorem 2.2.12). It diverges at both endpoints: the terms at $x = 1$ are constantly 1 and the terms at $x = -1$ have absolute value 1, so in neither case do they tend to 0.
4. $\sum_{n \geq 1} x^n/n$ has $R = 1$. Here $c_0 = 0$ and $c_n = 1/n$ for $n \geq 1$, and $1/n^{1/n}$ is positive with n -th power $1/n$, so $|c_n|^{1/n} = 1/n^{1/n}$ by theorem 1.3.12. By proposition 2.6.1(1) and the quotient rule of theorem 2.1.6 this tends to 1, and a convergent sequence has its limit for a limit superior (proposition 2.2.7 and theorem 2.2.12), so $\alpha = 1$ and $R = 1$. At $x = 1$ the series is the harmonic series and diverges (example 2.4.8). At $x = -1$ it is $\sum_{n \geq 1} (-1)^n/n$, which converges by corollary 2.5.3 because $1/n$ decreases to 0, while $\sum_{n \geq 1} 1/n$ diverges. So the convergence at $x = -1$ is conditional in the sense of definition 2.5.4.
5. $\sum_{n \geq 1} x^n/n^2$ has $R = 1$. The number $1/(n^{1/n})^2$ is positive with n -th power $1/n^2$, so it is $|c_n|^{1/n}$ by theorem 1.3.12, and it tends to 1 by proposition 2.6.1(1) and theorem 2.1.6. Hence

$\alpha = 1$ and $R = 1$. At $x = 1$ and at $x = -1$ the series of absolute values is $\sum_{n \geq 1} 1/n^2$, which converges by example 2.4.10, so both endpoints give absolute convergence.

Items (3), (4) and (5) have the same radius and differ at the boundary in all three possible ways: divergence at both endpoints, convergence at one and divergence at the other, and absolute convergence at both. So no statement phrased in terms of R alone can decide an endpoint, and each such point is settled by applying the tests of section 2.4 and section 2.5 to the single numerical series that the endpoint produces.

A power series of infinite radius defines a function on all of $\mathbb{R}$, and the exponential and trigonometric functions are introduced in exactly that way. The three series below are their definitions in this book, and each converges absolutely at every real point.

For exp that is item (2) of example 2.6.4. For the other two, fix $x \in \mathbb{R}$ and consider the cosine series, whose m -th coefficient is $(-1)^n/(2n)!$ when $m = 2n$ is even and 0 when m is odd. Writing $u_n = (-1)^n x^{2n}/(2n)!$ for the terms that can be nonzero, the same ratio computation as in item (2) gives

$$\frac{|u_{n+1}|}{|u_n|} = \frac{x^2}{(2n+1)(2n+2)} \leq \frac{x^2}{2n+1}$$

for $x \neq 0$, and the right-hand side falls below any prescribed $\epsilon > 0$ once $n \geq N$, where $N > x^2/\epsilon$ is chosen by theorem 1.4.1(2), since then $2n+1 > n \geq N$. The ratios are nonnegative, so they converge to 0, and $\sum_{n \geq 0} |u_n|$ converges by theorem 2.4.14. The partial sum $\sum_{m=0}^M |c_m x^m|$ omits the odd terms and so equals $\sum_{n: 2n \leq M} |u_n|$, which is at most $\sum_{n=0}^M |u_n|$, and this in turn is at most $\sum_{n \geq 0} |u_n|$, because the partial sums of a series of nonnegative terms increase and proposition 2.1.8, applied to the tail of that increasing sequence, bounds each of them by the limit. Bounded partial sums and nonnegative terms give convergence by proposition 2.4.4, so the cosine series converges absolutely at x , and theorem 2.4.12 makes it convergent. The case $x = 0$ leaves only the term $n = 0$, and the sine series is treated identically with $v_n = (-1)^n x^{2n+1}/(2n+1)!$, whose ratios are $x^2/((2n+2)(2n+3))$. Since all three series converge at every real point, a finite radius R would contradict part (2) of theorem 2.6.3 at any x with $|x| > R$, one of which exists by theorem 1.4.1(2). So all three radii are $+\infty$.

Definition 2.6.5: The Exponential, Cosine and Sine Functions

For $x \in \mathbb{R}$ set

$$\exp(x) = \sum_{n \geq 0} \frac{x^n}{n!}, \quad \cos x = \sum_{n \geq 0} \frac{(-1)^n x^{2n}}{(2n)!}, \quad \sin x = \sum_{n \geq 0} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$$

with the conventions $0! = 1$ and $x^0 = 1$ for every x . Each of the three series converges absolutely at every real x and has radius of convergence $+\infty$, so each defines a function $\mathbb{R} \rightarrow \mathbb{R}$.

Setting $x = 1$ in the first series gives the series of example 2.4.18 term by term, so $\exp(1) = e$. Every coefficient $1/n!$ lies in $\mathbb{Q}$ and so does the point $x = 1$, and yet the value does not, since proposition 2.4.19 shows $e \notin \mathbb{Q}$. The value of a power series is a limit of rationals whenever its coefficients and its argument are rational, and the limit of a sequence of rationals need not be rational, which is the failure that section 1.1 built $\mathbb{R}$ to repair. The values at $x = 0$ come straight from the series: only the term $n = 0$ is nonzero in the first two and every term of the third vanishes, so $\exp(0) = 1$, $\cos 0 = 1$

and $\sin 0 = 0$.

Nothing in definition 2.6.5 explains the name of $\exp$. What does explain it is the identity below, which converts addition of arguments into multiplication of values, and its proof is the Cauchy product of section 2.5 followed by one application of the binomial theorem.

Theorem 2.6.6: The Functional Equation of the Exponential

For all $x, y \in \mathbb{R}$,

$$\exp(x + y) = \exp(x) \exp(y)$$

Proof:

Fix $x, y \in \mathbb{R}$. By definition 2.6.5 the series $\sum_{n \geq 0} x^n/n!$ converges absolutely and the series $\sum_{n \geq 0} y^n/n!$ converges, which are the hypotheses of theorem 2.5.9. That theorem therefore applies: the Cauchy product of the two, whose n -th term is

$$d_n = \sum_{k=0}^n \frac{x^k}{k!} \cdot \frac{y^{n-k}}{(n-k)!}$$

converges, and $\sum_{n \geq 0} d_n = \exp(x) \exp(y)$.

For $0 \leq k \leq n$ the binomial coefficient is $\binom{n}{k} = n!/(k!(n-k)!)$, so $1/(k!(n-k)!) = \binom{n}{k}/n!$ and

$$d_n = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} = \frac{(x+y)^n}{n!}$$

the second equality being the binomial theorem [4, the binomial theorem] applied to $(x+y)^n$.

Hence $\sum_{n \geq 0} d_n$ is the series defining $\exp(x+y)$, and comparing the two evaluations of that one convergent series gives $\exp(x+y) = \exp(x) \exp(y)$, as we sought to show.

Two consequences follow at once. Taking $y = -x$ in theorem 2.6.6 gives $\exp(x) \exp(-x) = \exp(0) = 1$, so $\exp(x)$ is never 0 and $\exp(-x) = 1/\exp(x)$. And induction on n , with theorem 2.6.6 giving the step $\exp(n+1) = \exp(n) \exp(1)$, gives $\exp(n) = e^n$ for every $n \in \mathbb{N}$. The functional equation is what makes the alternative notation e^x for $\exp(x)$ consistent with integer powers of e .

Exercise 2.6.1

1. Compute the radius of convergence of $\sum_{n \geq 1} \frac{2^n}{n} x^n$, and decide the behaviour of the series at each of the two endpoints $x = 1/2$ and $x = -1/2$, saying in the convergent case whether the convergence is absolute or conditional.
2. Let $c_n = 2^n$ for even n and $c_n = 3^n$ for odd n . Show that the power series $\sum_{n \geq 0} c_n x^n$ has radius of convergence $1/3$. Show also that the sequence of ratios $|c_{n+1}/c_n|$ does not converge, so that no test phrased in terms of a limit of consecutive ratios reaches this radius.
3. Show that $\sum_{n \geq 1} x^n/n^n$ has radius of convergence $+\infty$, and that $\sum_{n \geq 1} n^k x^n$ has radius of convergence 1 for every fixed integer $k \geq 1$.
4. Prove that $\exp(x) \geq 1+x$ for every $x \geq 0$, and deduce that $\exp(x) > 0$ for every $x \in \mathbb{R}$. (For $x < 0$, use theorem 2.6.6.)

5. Working directly from definition 2.6.5, prove that $\cos(-x) = \cos x$ and $\sin(-x) = -\sin x$ for every $x \in \mathbb{R}$.
6. Let $c_m = 1$ when m is a perfect square and $c_m = 0$ otherwise. Show that $\sum_{m \geq 0} c_m x^m$ has radius of convergence 1, and that it diverges at $x = 1$ and at $x = -1$. (For the radius, verify both clauses of theorem 2.2.12 for the candidate value 1.)

Limits and Continuity

Chapter 2 spent completeness on sequences. This chapter spends it on functions, and almost every theorem below is one of its results read through a single translation.

That translation is the sequential criterion of Section 3.1: a function has limit L at a point exactly when it carries every sequence approaching that point to a sequence converging to L . Once it is proved, the algebra of limits, the squeeze and theorem 2.2.9 transfer with no further work, and the counterexamples that settle the delicate questions become sequences chosen by hand.

Stating the definition at all takes vocabulary this book has not yet needed. A limit at c says nothing about $f(c)$, and c need not lie in the domain, so what the definition does require is that c be approachable from the domain. Naming that condition means naming limit points, closure, and the closed subsets of $\mathbb{R}$. Those notions belong to a general theory the later volumes develop, and only the part the real line needs is built here.

The last two sections are where completeness is spent. On a closed bounded interval a continuous function attains every intermediate value, stays bounded, attains its bounds, and is uniformly continuous. All four fail over $\mathbb{Q}$, and the first fails visibly, since $x^2 - 2$ is negative at 0 and positive at 2 with no rational root between them. Every proof runs through theorem 2.2.9 or through theorem 1.2.4 directly.

Section 3.1 builds the point-set vocabulary, defines the functional limit, proves the sequential criterion, and constructs the Cantor set as the worked example of a closed set containing no interval. Section 3.2 defines continuity, establishes the algebra and the composition rule, and separates the standard pathologies: the Dirichlet function, continuous nowhere; Thomae's function, whose discontinuity set is exactly $\mathbb{Q}$; and the monotone functions, whose discontinuity sets can be no larger than that. Section 3.3 proves the intermediate value and extreme value theorems, with the counterexamples showing that neither half of the hypothesis can be dropped. Section 3.4 strengthens continuity by making the modulus independent of the point, shows that a closed bounded interval forces the strengthening at no cost, and records the three consequences the later chapters spend.

3.1 Limits of Functions

A sequence has a single limiting direction. Its index runs through $\mathbb{N}$, and the only question is the behaviour of the terms for large n . A function $f: E \rightarrow \mathbb{R}$ with $E \subseteq \mathbb{R}$ can be examined near any point c that has other points of E arbitrarily close to it, so before the limit can be defined at all, the points c at which the question makes sense have to be identified. This section fixes that vocabulary for subsets of the line, defines $\lim_{x \rightarrow c} f(x)$, and proves the characterization that turns a statement about convergent sequences into a statement about functions.

An interval is a subset of $\mathbb{R}$ with no gaps in it. A betweenness condition says this without naming endpoints, so that the unbounded and the degenerate cases fall under the same words, and the nine standard notations are fixed alongside it.

Definition 3.1.1: [Real] Interval

A set $I \subseteq \mathbb{R}$ is an *interval* if for all $x, y \in I$ and all $z \in \mathbb{R}$,

$$x < z < y \quad \text{implies} \quad z \in I$$

For $a, b \in \mathbb{R}$ the following nine sets carry the names shown, and each is an interval:

$$\begin{aligned} (a, b) &= \{x \in \mathbb{R} \mid a < x < b\}, & [a, b] &= \{x \in \mathbb{R} \mid a \leq x \leq b\}, \\ [a, b) &= \{x \in \mathbb{R} \mid a \leq x < b\}, & (a, b] &= \{x \in \mathbb{R} \mid a < x \leq b\}, \\ (a, \infty) &= \{x \in \mathbb{R} \mid x > a\}, & [a, \infty) &= \{x \in \mathbb{R} \mid x \geq a\}, \\ (-\infty, b) &= \{x \in \mathbb{R} \mid x < b\}, & (-\infty, b] &= \{x \in \mathbb{R} \mid x \leq b\}, \end{aligned}$$

together with $(-\infty, \infty) = \mathbb{R}$. An interval of the form $[a, b]$ with $a \leq b$ is called *closed and bounded*.

The symbols ∞ and $-\infty$ in these names record that an end is unbounded and are not elements of $\mathbb{R}$. They are the two points adjoined in definition 2.2.4. Two degenerate sets satisfy the definition, and later statements quantify over intervals without excluding them. The empty set is an interval, since the condition on x, y, z is never triggered, and so is a one-point set $\{a\} = [a, a]$.

Example 3.1.2: Intervals and Three Sets That Are Not Intervals

1. Each of the nine sets named in definition 3.1.1 is an interval. For $(a, b]$, suppose $x, y \in (a, b]$ and $x < z < y$. Then $a < x < z$ and $z < y \leq b$, so $a < z \leq b$ and $z \in (a, b]$. The other eight are identical, with the strict and weak inequalities changed to match.
2. $\mathbb{Q} \cap [0, 1]$ is not an interval. Both 0 and 1 lie in it, and corollary 1.4.5 gives a $t \in \mathbb{R} \setminus \mathbb{Q}$ with $0 < t < 1$. This t lies between two points of the set and is not in it.
3. $[0, 1] \cup [2, 3]$ is not an interval. It contains 0 and 3, and $3/2$ lies strictly between them, but $3/2 > 1$ and $3/2 < 2$, so $3/2$ belongs to neither piece.
4. $\mathbb{R} \setminus \{0\}$ is not an interval, since -1 and 1 lie in it and 0 does not.

Four definitions now separate the points that a set approaches from the points that it merely contains.

Three of them are stated with the absolute value of definition 1.3.6, so that $|y - x| < \delta$ says that y is within δ of x , as computed in example 1.3.7.

Definition 3.1.3: Adherent Point

Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is *adherent* to E if for every real $\delta > 0$ there is a $y \in E$ with $|y - x| < \delta$.

Definition 3.1.4: [Real] Closure

Let $E \subseteq \mathbb{R}$. The *closure* of E , written $\overline{E}$, is the set of all points of $\mathbb{R}$ adherent to E .

Definition 3.1.5: [Real] Limit Point

Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is a *limit point* of E if for every real $\delta > 0$ there is a $y \in E$ with

$$0 < |y - x| < \delta$$

The set of limit points of E is written E' .

Definition 3.1.6: [Real] Isolated Point

Let $E \subseteq \mathbb{R}$ and $x \in E$. Then x is an *isolated point* of E if there is a real $\delta > 0$ such that the only $y \in E$ with $|y - x| < \delta$ is $y = x$.

The single character of difference between definition 3.1.3 and definition 3.1.5 is the strict inequality $0 < |y - x|$, which forbids the witness y from being x itself. Every point of E is adherent to E for the trivial reason that it can serve as its own witness, so adherence records nothing new at points of E . Being a limit point does, because the witnesses must be other points. The terminology of chapter 2 is not reused here: a *subsequential limit* of a sequence is the notion of definition 2.2.6, and a limit point in the present sense is always a limit point of a set.

Example 3.1.7: Limit Points and Closures of Four Subsets of the Line

1. $E = (0, 1)$. Every $x \in [0, 1]$ is a limit point of E . Let $\delta > 0$ and put $r = \min(\delta, 1)$, so $0 < r \leq 1$. If $x < 1$, set $y = \min(x + r/2, (x + 1)/2)$. Both entries exceed x , the first because $r/2 > 0$ and the second by proposition 1.1.2(8), so $y > x \geq 0$. The same part (8) gives $(x + 1)/2 < 1$, and $y \leq (x + 1)/2$, so $y < 1$ and $y \in (0, 1)$. Since $y \leq x + r/2$, also $0 < y - x \leq r/2 < \delta$. If $x = 1$, set $y = \max(1 - r/2, 1/2)$, so $1/2 \leq y < 1$ and $y \in (0, 1)$, while $0 < 1 - y \leq r/2 < \delta$. No point outside $[0, 1]$ is adherent to E : if $x < 0$ then every $y \in (0, 1)$ has $|y - x| = y - x > -x$, and if $x > 1$ then every such y has $|y - x| = x - y > x - 1$. So $E' = [0, 1]$ and $\overline{E} = [0, 1]$, and E has no isolated points.
2. $E = \mathbb{Z}$. Distinct integers differ by at least 1, so for $m \in \mathbb{Z}$ the value $\delta = 1$ shows m is isolated in $\mathbb{Z}$ and shows that m is not a limit point of $\mathbb{Z}$. A point $x \in \mathbb{R} \setminus \mathbb{Z}$ is not adherent either: with $m = \lfloor x \rfloor$ from proposition 1.4.3 one has $m < x < m + 1$, and $\delta = \min(x - m, m + 1 - x) > 0$ leaves no integer within δ of x , since an integer $k \leq m$ has $x - k \geq x - m \geq \delta$ and an integer $k \geq m + 1$ has $k - x \geq m + 1 - x \geq \delta$. So $\mathbb{Z}' = \emptyset$ and $\overline{\mathbb{Z}} = \mathbb{Z}$.

3. $E = \mathbb{Q}$. Every $x \in \mathbb{R}$ is a limit point. Given $\delta > 0$, theorem 1.4.4 applied to $x < x + \delta$ produces a $p \in \mathbb{Q}$ with $x < p < x + \delta$, so $0 < |p - x| < \delta$. Hence $\mathbb{Q}' = \overline{\mathbb{Q}} = \mathbb{R}$, and $\mathbb{Q}$ has no isolated points. This is the failure that section 1.1 was built to repair: $\mathbb{Q}$ is a proper subset of the line and is adherent to all of it.
4. $E = \{0, 1, 2\}$. Every point of E is isolated, by the same computation as in (2) with $\delta = 1$. No $x \in \mathbb{R}$ is a limit point of E . If $x \notin E$, the number $\delta = \min\{|y - x| \mid y \in E\}$ is a minimum over a finite nonempty set of positive reals, so $\delta > 0$, and no $y \in E$ has $|y - x| < \delta$, so x is not even adherent to E . If $x \in E$, then $\delta = 1$ admits no $y \in E$ with $0 < |y - x| < 1$, since the only $y \in E$ within 1 of x is x itself. Hence $E' = \emptyset$ and $\overline{E} = E$.

The four computations above already show the pattern that proposition 3.1.8 records: the closure of E is E with its limit points added, and inside E the two possibilities for a point are to be isolated or to be a limit point. Part (3) disposes of every finite set at once, and part (5) is what example 3.1.14 uses to see that a union of finitely many closed sets is closed.

Proposition 3.1.8: Adherent Points, Limit Points and Isolated Points

Let $E, A, B \subseteq \mathbb{R}$ and let E' be the set of limit points of E .

1. $E \subseteq \overline{E}$ and $E' \subseteq \overline{E}$, and $\overline{E} = E \cup E'$.
2. Every point of E is either an isolated point of E or a limit point of E , and no point of E is both.
3. If $x \in E'$, then for every $\delta > 0$ the set $\{y \in E \mid |y - x| < \delta\}$ is infinite. In particular a finite subset of $\mathbb{R}$ has no limit points.
4. If $A \subseteq B$ then $\overline{A} \subseteq \overline{B}$.
5. $\overline{A \cup B} = \overline{A} \cup \overline{B}$.

Proof:

1. If $x \in E$ then for every $\delta > 0$ the point $y = x$ lies in E and satisfies $|y - x| = 0 < \delta$, using proposition 1.3.8(1). Hence $E \subseteq \overline{E}$. If $x \in E'$ then a y with $0 < |y - x| < \delta$ in particular has $|y - x| < \delta$, so $E' \subseteq \overline{E}$, and therefore $E \cup E' \subseteq \overline{E}$. For the reverse inclusion let $x \in \overline{E}$ with $x \notin E$, and let $\delta > 0$. There is a $y \in E$ with $|y - x| < \delta$, and $y \neq x$ because $y \in E$ while $x \notin E$, so $|y - x| > 0$ by proposition 1.3.8(1). Hence $x \in E'$.
2. Let $x \in E$ and suppose x is not isolated in E . Then for every $\delta > 0$ the statement defining isolation fails, so there is a $y \in E$ with $|y - x| < \delta$ and $y \neq x$. As in (1) this gives $0 < |y - x| < \delta$, so $x \in E'$. Conversely, if x is isolated with witness δ , then no $y \in E$ satisfies $0 < |y - x| < \delta$, because such a y would be a point of E within δ of x and different from x , so $x \notin E'$.
3. Let $x \in E'$, let $\delta > 0$, and put $F = \{y \in E \mid 0 < |y - x| < \delta\}$. Since $x \in E'$, the set F is nonempty. Suppose F were finite. A finite nonempty set of positive real numbers has a least element, so $r = \min\{|y - x| \mid y \in F\}$ exists and $r > 0$. Applying the definition of a limit point with r in place of δ produces a $y_0 \in E$ with $0 < |y_0 - x| < r$. Then $|y_0 - x| < r \leq \delta$, so $y_0 \in F$ and hence $r \leq |y_0 - x|$, contradicting $|y_0 - x| < r$. So F is

infinite, and so is the larger set $\{y \in E \mid |y - x| < \delta\}$. If E is finite and $x \in E'$, then taking $\delta = 1$ exhibits an infinite subset of E , which is impossible, so $E' = \emptyset$.

4. If $x \in \overline{A}$ and $\delta > 0$, a witness $y \in A$ with $|y - x| < \delta$ is also a point of B .
5. The inclusion $\overline{A \cup B} \subseteq \overline{A} \cup \overline{B}$ is (4) applied twice. For the reverse, let $x \in \overline{A \cup B}$ and suppose $x \notin \overline{A}$ and $x \notin \overline{B}$. Then there are $\delta_1 > 0$ with no $y \in A$ satisfying $|y - x| < \delta_1$, and $\delta_2 > 0$ with no $y \in B$ satisfying $|y - x| < \delta_2$. Put $\delta = \min(\delta_1, \delta_2) > 0$. Then no $y \in A \cup B$ satisfies $|y - x| < \delta$, contradicting $x \in \overline{A \cup B}$, and completing the proof.

The closure has a second description that will be used more often than the definition itself, because it converts a question about all $\delta > 0$ into a question about a single sequence. Section 3.4 extends a uniformly continuous function from a set to its closure, and it is proposition 3.1.9 that produces the point at which the extended function must be evaluated.

Proposition 3.1.9: The Closure Described by Sequences

Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$.

1. $x \in \overline{E}$ if and only if there is a sequence (x_n) with $x_n \in E$ for every n and $x_n \rightarrow x$.
2. x is a limit point of E if and only if there is a sequence (x_n) with $x_n \in E$ and $x_n \neq x$ for every n , and $x_n \rightarrow x$.

Proof:

1. Suppose $x \in \overline{E}$. For each $n \in \mathbb{N}$ the number $1/n$ is positive, so there is a point of E within $1/n$ of x . Choose one and call it x_n .^a Let $\epsilon > 0$. By theorem 1.4.1(3) there is an $N \in \mathbb{N}$ with $0 < 1/N < \epsilon$, and for $n \geq N$ one has $1/n \leq 1/N$ by proposition 1.1.2(7), so $|x_n - x| < 1/n \leq 1/N < \epsilon$. Hence $x_n \rightarrow x$. Conversely, suppose $x_n \in E$ and $x_n \rightarrow x$, and let $\delta > 0$. Definition 2.1.2 gives an N with $|x_N - x| < \delta$, and $x_N \in E$, so $x \in \overline{E}$.
2. The same argument with $0 < |x_n - x| < 1/n$ in place of $|x_n - x| < 1/n$ proves the forward direction, the choice being available because x is a limit point. For the converse, let $\delta > 0$ and take N with $|x_N - x| < \delta$. Since $x_N \neq x$, proposition 1.3.8(1) gives $|x_N - x| > 0$, so $0 < |x_N - x| < \delta$ with $x_N \in E$, as we sought to show.

^aChoosing one such x_n for every n at once is an appeal to countable choice.

A set that contains every point adherent to it is one on which limits of sequences cannot escape, and that is the property the extreme value theorem of section 3.3 will need of $[a, b]$.

Definition 3.1.10: [Real] Closed Set

A set $E \subseteq \mathbb{R}$ is *closed* if $\overline{E} = E$.

Since $E \subseteq \overline{E}$ always holds by proposition 3.1.8(1), a set is closed exactly when $\overline{E} \subseteq E$, that is, when every point adherent to E already lies in E .

Proposition 3.1.11: Closed Sets Contain Their Limit Points

A set $E \subseteq \mathbb{R}$ is closed if and only if every limit point of E belongs to E .

Proof:

By proposition 3.1.8(1), $\overline{E} = E \cup E'$. If $E' \subseteq E$ then $E \cup E' = E$, so $\overline{E} = E$ and E is closed. If E is closed then $E' \subseteq E \cup E' = \overline{E} = E$, completing the proof.

Proposition 3.1.12: Closed Sets Are Exactly the Sequentially Closed Sets

A set $E \subseteq \mathbb{R}$ is closed if and only if the following holds: whenever (x_n) is a sequence with $x_n \in E$ for every n and $x_n \rightarrow x$ for some $x \in \mathbb{R}$, the limit x lies in E .

Proof:

Suppose E is closed and let (x_n) be a sequence in E with $x_n \rightarrow x$. By proposition 3.1.9(1), $x \in \overline{E} = E$. Conversely, suppose the stated condition holds and let $x \in \overline{E}$. By proposition 3.1.9(1) there is a sequence (x_n) in E with $x_n \rightarrow x$, so the condition gives $x \in E$. Hence $\overline{E} \subseteq E$, and with proposition 3.1.8(1) this gives $\overline{E} = E$, as we sought to show.

Example 3.1.13: Closed and Non-Closed Subsets of the Line

1. $\emptyset$ is closed, since it has no adherent points at all: the definition asks for a $y \in \emptyset$, and there is none. And $\mathbb{R}$ is closed, since $\mathbb{R} \subseteq \mathbb{R}$ by definition.
2. For $a \leq b$ the interval $[a, b]$ is closed. Let (x_n) be a sequence in $[a, b]$ with $x_n \rightarrow x$. Then $a \leq x_n \leq b$ for every n , so $a \leq x \leq b$ by proposition 2.1.8(2), and proposition 3.1.12 applies. The same argument with only one of the two inequalities shows that $[a, \infty)$ and $(-\infty, b]$ are closed.
3. $(0, 1)$ is not closed, and neither is $[0, 1)$. By example 3.1.7(1) the closure of $(0, 1)$ is $[0, 1]$, which is not $(0, 1)$. The point 1 is adherent to $[0, 1)$ by the computation there, while $1 \notin [0, 1)$.
4. $\mathbb{Z}$ is closed and $\mathbb{Q}$ is not, by example 3.1.7(2) and (3).
5. Every finite subset of $\mathbb{R}$ is closed. It has no limit points by proposition 3.1.8(3), so proposition 3.1.11 applies.
6. A union $F_1 \cup \dots \cup F_m$ of finitely many closed sets is closed. For $m = 1$ this is the hypothesis, and if it holds for m then proposition 3.1.8(5) with $A = F_1 \cup \dots \cup F_m$ and $B = F_{m+1}$ gives

$$\overline{F_1 \cup \dots \cup F_{m+1}} = \overline{F_1 \cup \dots \cup F_m} \cup \overline{F_{m+1}} = (F_1 \cup \dots \cup F_m) \cup F_{m+1}$$

so the union of $m + 1$ of them is closed. No such statement holds for infinite unions: each $\{1/n\}$ is closed by (5), while $\bigcup_{n \in \mathbb{N}} \{1/n\}$ has 0 as an adherent point, since $|1/n - 0| < \delta$ for any n with $1/n < \delta$, which theorem 1.4.1(3) gives.

The closed sets met so far are unions of intervals, or so sparse that they have no limit points at

all. The following construction produces one that is neither. It is built from theorem 1.4.8 and the geometric series of example 2.4.7, and nothing else.

Example 3.1.14: The Cantor Set

For $A \subseteq \mathbb{R}$, $\lambda \in \mathbb{R}$ and $t \in \mathbb{R}$ write $\lambda A = \{\lambda a \mid a \in A\}$ and $t + A = \{t + a \mid a \in A\}$. Define sets $C_n \subseteq \mathbb{R}$ by

$$C_0 = [0, 1], \quad C_{n+1} = \frac{1}{3}C_n \cup \left(\frac{2}{3} + \frac{1}{3}C_n\right)$$

and put $C = \bigcap_{n \geq 0} C_n$, the *Cantor set*. Then:

1. $C_n \subseteq [0, 1]$ and $C_{n+1} \subseteq C_n$ for every n , and C_n is a union of 2^n closed intervals each of length 3^{-n} and each with both endpoints of the form $k/3^n$ for an integer k .
2. C is nonempty. More precisely, for every sequence $(\sigma_n)_{n \geq 1}$ with $\sigma_n \in \{0, 2\}$ for each n there is a point of C obtained from it, and 0, 1 and $1/4$ all lie in C .
3. C is closed.
4. If $I \subseteq C$ is an interval, then I has at most one point.
5. For every $\epsilon > 0$ there are finitely many closed intervals whose union contains C and the sum of whose lengths is less than ϵ .

Proof of (1). The map $x \mapsto x/3$ preserves the order by proposition 1.1.2(5), and so does $x \mapsto 2/3 + x/3$. Hence $\frac{1}{3}[\alpha, \beta] = [\alpha/3, \beta/3]$ and $\frac{2}{3} + \frac{1}{3}[\alpha, \beta] = [2/3 + \alpha/3, 2/3 + \beta/3]$ for $\alpha \leq \beta$, the inclusion in each direction being the observation that $x \mapsto x/3$ and $x \mapsto 2/3 + x/3$ are bijections of $\mathbb{R}$ with inverses $y \mapsto 3y$ and $y \mapsto 3y - 2$. Now argue by induction. Certainly $C_0 = [0, 1]$ is one closed interval of length 1, and $1 = 2^0$, $1 = 3^{-0}$. If $C_n \subseteq [0, 1]$, then $\frac{1}{3}C_n \subseteq [0, 1/3]$ and $\frac{2}{3} + \frac{1}{3}C_n \subseteq [2/3, 1]$, so $C_{n+1} \subseteq [0, 1]$. If C_n is the union of the 2^n closed intervals $[\alpha_i, \alpha_i + 3^{-n}]$ with each $\alpha_i = k_i/3^n$ for an integer k_i , then $\frac{1}{3}C_n$ is the union of the 2^n closed intervals $[\alpha_i/3, \alpha_i/3 + 3^{-n-1}]$ and $\frac{2}{3} + \frac{1}{3}C_n$ is the union of the 2^n closed intervals $[2/3 + \alpha_i/3, 2/3 + \alpha_i/3 + 3^{-n-1}]$, so C_{n+1} is a union of 2^{n+1} closed intervals of length 3^{-n-1} . The new left endpoints are $k_i/3^{n+1}$ and $(2 \cdot 3^n + k_i)/3^{n+1}$, and the right endpoints exceed them by $1/3^{n+1}$, so all are of the required form. For the nesting, first $C_1 = [0, 1/3] \cup [2/3, 1] \subseteq [0, 1] = C_0$. If $C_{n+1} \subseteq C_n$, then $\frac{1}{3}C_{n+1} \subseteq \frac{1}{3}C_n$ and $\frac{2}{3} + \frac{1}{3}C_{n+1} \subseteq \frac{2}{3} + \frac{1}{3}C_n$, whence $C_{n+2} \subseteq C_{n+1}$. Iterating the nesting gives $C_n \subseteq C_m$ whenever $m \leq n$.

Proof of (2). Write $T_0(x) = x/3$ and $T_2(x) = (x + 2)/3$, so that $C_{n+1} = T_0(C_n) \cup T_2(C_n)$ and, by the computation in (1), $T_0([\alpha, \beta]) = [\alpha/3, \beta/3]$ and $T_2([\alpha, \beta]) = [(\alpha + 2)/3, (\beta + 2)/3]$, each of length $(\beta - \alpha)/3$. Given $\sigma_1, \sigma_2, \dots$ in $\{0, 2\}$, set $J_0 = [0, 1]$ and

$$J_n = T_{\sigma_1} \circ T_{\sigma_2} \circ \dots \circ T_{\sigma_n}([0, 1])$$

Each J_n is a closed interval of length 3^{-n} , by induction on n from the length computation just made. Since $T_0([0, 1]) = [0, 1/3]$ and $T_2([0, 1]) = [2/3, 1]$ are both contained in $[0, 1]$, and each T_σ carries a subset into the image of any set containing it, applying $T_{\sigma_1} \circ \dots \circ T_{\sigma_n}$ to $T_{\sigma_{n+1}}([0, 1]) \subseteq [0, 1]$ gives $J_{n+1} \subseteq J_n$. Next, $J_n \subseteq C_n$. Prove instead the stronger statement that $T_{\tau_1} \circ \dots \circ T_{\tau_n}([0, 1]) \subseteq C_n$ for every $\tau_1, \dots, \tau_n$ in $\{0, 2\}$, by induction on n . For $n = 0$ this is $[0, 1] = C_0$. For the step, write $T_{\tau_1} \circ \dots \circ T_{\tau_{n+1}}([0, 1]) = T_{\tau_1}(K)$ where $K = T_{\tau_2} \circ \dots \circ T_{\tau_{n+1}}([0, 1])$ is built from a list of length n , so $K \subseteq C_n$ by the inductive hypothesis, and then $T_{\tau_1}(K) \subseteq T_{\tau_1}(C_n) \subseteq C_{n+1}$.

The lengths 3^{-n} form a null sequence by proposition 2.1.10(3) with $j = 0$ and $r = 1/3$, so for

every $\epsilon > 0$ there is an n with $3^{-n} < \epsilon$. The hypotheses of theorem 1.4.8(2) are therefore met by $(J_n)_{n \geq 0}$, reindexed to start at 1, and it gives a point x lying in every J_n . Then $x \in C_n$ for every n , so $x \in C$. Taking every $\sigma_n = 0$ gives $J_n = [0, 3^{-n}]$ and $x = 0$, and taking every $\sigma_n = 2$ gives $x = 1$. For $1/4$, argue instead that $1/4 \in C_n$ and $3/4 \in C_n$ for every n , by simultaneous induction: both lie in $C_0 = [0, 1]$, and if both lie in C_n then $1/4 = \frac{1}{3} \cdot \frac{3}{4} \in \frac{1}{3}C_n \subseteq C_{n+1}$ and $3/4 = \frac{2}{3} + \frac{1}{3} \cdot \frac{1}{4} \in \frac{2}{3} + \frac{1}{3}C_n \subseteq C_{n+1}$, the second because $2/3 + 1/12 = 8/12 + 1/12 = 9/12 = 3/4$. So $1/4 \in C$. Together with 0 and 1 this proves (2).

The point $1/4$ is not an endpoint of any of the intervals of any C_n : by (1) such an endpoint is $k/3^n$ for an integer k , and $1/4 = k/3^n$ would give $3^n = 4k$, which is impossible because 3^n is odd and $4k$ is even. It is the point that the argument above produces from the sequence with $\sigma_n = 0$ for odd n and $\sigma_n = 2$ for even n . Indeed $T_\sigma(x) = (x + \sigma)/3$ for both values of σ , so applying T_{σ_1} to a closed interval with left endpoint α returns one with left endpoint $(\alpha + \sigma_1)/3$, and induction on the length of the list makes the left endpoint of J_n equal to $\sum_{i=1}^n \sigma_i 3^{-i}$. That endpoint is within the length 3^{-n} of the point of C lying in every J_n , so that point is the limit of these endpoints. Dropping the term $k = 0$, whose value is 1, from the geometric series of example 2.4.7 with $x = 1/9$ evaluates the limit as

$$2 \sum_{k=1}^{\infty} 9^{-k} = 2 \left(\frac{1}{1 - 1/9} - 1 \right) = 2 \left(\frac{9}{8} - 1 \right) = \frac{1}{4}$$

so the sequence alternating 0 and 2 produces $1/4$.

Proof of (3). Each of the 2^n closed intervals composing C_n is closed by example 3.1.13(2), so C_n is closed by example 3.1.13(6). Now let (x_k) be a sequence in C with $x_k \rightarrow x$. Fix n . Every x_k lies in C and hence in C_n , so proposition 3.1.12 applied to the closed set C_n gives $x \in C_n$. As n was arbitrary, $x \in C$, and proposition 3.1.12 applied to C shows C is closed.

Proof of (4). First a claim: for every $n \geq 0$, if $a \leq b$ and $[a, b] \subseteq C_n$, then $b - a \leq 3^{-n}$. For $n = 0$ this says $[a, b] \subseteq [0, 1]$ forces $b - a \leq 1$, which holds since $0 \leq a \leq b \leq 1$. Assume the claim for n and let $[a, b] \subseteq C_{n+1}$ with $a \leq b$. If $a = b$ there is nothing to prove, so assume $a < b$. By (1), $C_{n+1} \subseteq [0, 1/3] \cup [2/3, 1]$, since $\frac{1}{3}C_n \subseteq [0, 1/3]$ and $\frac{2}{3} + \frac{1}{3}C_n \subseteq [2/3, 1]$. Suppose $b > 1/3$ and $a < 2/3$, and put $u = \max(a, 1/3)$ and $v = \min(b, 2/3)$. Each of the four comparisons $a < b$, $a < 2/3$, $1/3 < b$ and $1/3 < 2/3$ holds, so $u < v$. By proposition 1.1.2(8) the point $w = (u + v)/2$ satisfies $u < w < v$, hence $a \leq u < w < v \leq b$ and $w \in [a, b] \subseteq C_{n+1}$, while $1/3 \leq u < w < v \leq 2/3$ puts w outside $[0, 1/3] \cup [2/3, 1]$. That is a contradiction, so $b \leq 1/3$ or $a \geq 2/3$. In the first case every $x \in [a, b]$ satisfies $x \leq 1/3 < 2/3$, so $x \notin [2/3, 1]$ and therefore $x \notin \frac{2}{3} + \frac{1}{3}C_n$. Being in C_{n+1} , such an x lies in $\frac{1}{3}C_n$, that is $3x \in C_n$. Every $y \in [3a, 3b]$ is $3x$ for the point $x = y/3$ of $[a, b]$, so $[3a, 3b] \subseteq C_n$, and the inductive hypothesis gives $3b - 3a \leq 3^{-n}$, that is $b - a \leq 3^{-n-1}$. In the second case every $x \in [a, b]$ satisfies $x \geq 2/3 > 1/3$, so $x \notin \frac{1}{3}C_n$ and hence $3x - 2 \in C_n$. The same reasoning gives $[3a - 2, 3b - 2] \subseteq C_n$ and again $b - a \leq 3^{-n-1}$. This proves the claim. Now let $I \subseteq C$ be an interval with two distinct points $a < b$. Every z with $a < z < b$ lies in I by definition 3.1.1, and a, b lie in I , so $[a, b] \subseteq I \subseteq C_n$ for every n . The claim gives $0 < b - a \leq 3^{-n}$ for every n , and since $3^{-n} \rightarrow 0$ by proposition 2.1.10(3), proposition 2.1.8(2) forces $b - a \leq 0$, a contradiction.

Proof of (5). The sequence $(2/3)^n$ is null by proposition 2.1.10(3) with $j = 0$ and $r = 2/3$, so there is an n with $(2/3)^n < \epsilon$. By (1), $C \subseteq C_n$ and C_n is a union of 2^n closed intervals of length 3^{-n} , whose lengths sum to $2^n \cdot 3^{-n} = (2/3)^n < \epsilon$.

So C is a closed subset of $[0, 1]$ containing no interval of positive length, and yet it is not thin in the way a finite set is. Each of the 2^n intervals from which C_n is assembled is the interval J_n of some sequence (σ_k) , and the proof of (2) puts a point of C inside every such J_n , so C meets all 2^n of them.

It also contains points such as $1/4$ that are not endpoints of any of them. The vocabulary is now complete enough to define the limit.

Definition 3.1.15: Limit of a Function

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let c be a limit point of E , and let $L \in \mathbb{R}$. Then f has limit L at c , written

$$\lim_{x \rightarrow c} f(x) = L$$

if for every real $\epsilon > 0$ there is a real $\delta > 0$ such that

$$|f(x) - L| < \epsilon \quad \text{for every } x \in E \text{ with } 0 < |x - c| < \delta$$

3.1.16 Convention: The Deleted Limit Two features of definition 3.1.15 hold throughout the book and are not restated later.

1. The point c need not belong to E , and when it does, the value $f(c)$ plays no part. The condition $0 < |x - c|$ excludes $x = c$ from every requirement the definition makes. A limit of this kind is called *deleted*.
2. The point c must be a limit point of E . Without that hypothesis the definition is empty: if c is not a limit point, some $\delta > 0$ admits no $x \in E$ with $0 < |x - c| < \delta$, so the displayed condition holds for every $L \in \mathbb{R}$ at once and f would have every real number as a limit at c .

The two work together. Deleting c is what allows a limit to be taken at a point where f is undefined or takes an unrelated value, and requiring c to be a limit point is what stops that deletion from removing all the information at once.

Example 3.1.17: Four Functional Limits

1. $E = \mathbb{R}$, $f(x) = 3x - 1$, $c \in \mathbb{R}$. Then $\lim_{x \rightarrow c} f(x) = 3c - 1$. Every $c \in \mathbb{R}$ is a limit point of $\mathbb{R}$, since for $\delta > 0$ the point $y = c + \delta/2$ satisfies $0 < |y - c| = \delta/2 < \delta$. Given $\epsilon > 0$ put $\delta = \epsilon/3$. If $0 < |x - c| < \delta$, then $|f(x) - (3c - 1)| = |3(x - c)| = 3|x - c| < 3\delta = \epsilon$, using proposition 1.3.8(3).
2. $E = \mathbb{R} \setminus \{1\}$, $f(x) = (x^2 - 1)/(x - 1)$, $c = 1$. Here $c \notin E$. For $x \in E$ one has $x - 1 \neq 0$, so $f(x) = (x - 1)(x + 1)/(x - 1) = x + 1$. Also 1 is a limit point of E , since for $\delta > 0$ the point $x = 1 + \min(\delta, 1)/2$ lies in E and satisfies $0 < |x - 1| < \delta$. Given $\epsilon > 0$ put $\delta = \epsilon$. If $x \in E$ and $0 < |x - 1| < \delta$, then $|f(x) - 2| = |x - 1| < \epsilon$. So $\lim_{x \rightarrow 1} f(x) = 2$ although $f(1)$ is not defined.
3. $E = \mathbb{R}$, $g(x) = x + 1$ for $x \neq 1$ and $g(1) = 17$. Here 1 is a limit point of $\mathbb{R}$ by (1). The points at which definition 3.1.15 tests g are those with $0 < |x - 1| < \delta$, and $x = 1$ is not among them, so g agrees there with the f of (2) and the same choice $\delta = \epsilon$ works. Hence $\lim_{x \rightarrow 1} g(x) = 2$ while $g(1) = 17$.
4. $E = (0, 1) \cup \{2\}$ and $c = 2$. Every $x \in (0, 1)$ has $|x - 2| = 2 - x > 1$, so $\delta = 1/2$ admits no $x \in E$ with $0 < |x - 2| < \delta$. Hence 2 is not a limit point of E , and definition 3.1.15 assigns no limit at 2 to any function on E . This is the situation box 3.1.16(2) excludes.

Proposition 3.1.18: A Function Has at Most One Limit at a Point

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, and let c be a limit point of E . If $\lim_{x \rightarrow c} f(x) = L$ and $\lim_{x \rightarrow c} f(x) = L'$, then $L = L'$.

Proof:

Suppose $L \neq L'$ and put $\epsilon = |L - L'|/2$. Then $\epsilon > 0$ by proposition 1.3.8(1). The definition gives a $\delta > 0$ with $|f(x) - L| < \epsilon$ for every $x \in E$ with $0 < |x - c| < \delta$, and a $\delta' > 0$ with $|f(x) - L'| < \epsilon$ for every $x \in E$ with $0 < |x - c| < \delta'$. Since c is a limit point of E and $\min(\delta, \delta') > 0$, there is an $x_0 \in E$ with $0 < |x_0 - c| < \min(\delta, \delta')$, and this x_0 satisfies both estimates. By proposition 1.3.8(4),

$$|L - L'| \leq |L - f(x_0)| + |f(x_0) - L'| < \epsilon + \epsilon = |L - L'|$$

which is impossible. So $L = L'$, completing the proof.

The limit-point hypothesis is what made that proof possible: it produced the point x_0 at which the two estimates could be compared. Uniqueness is what entitles the notation $\lim_{x \rightarrow c} f(x)$ to name a single real number, and from here on the phrase “the limit of f at c exists” means that some $L \in \mathbb{R}$ satisfies definition 3.1.15.

Everything chapter 2 proved about convergent sequences now becomes available, through a single translation. Theorem 3.1.19 is the one result of this section that later sections use most often. The algebra of limits, the squeeze theorem and the counterexamples of section 3.2 are corollaries of it together with theorem 2.1.6.

Theorem 3.1.19: The Sequential Criterion for Functional Limits

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let c be a limit point of E , and let $L \in \mathbb{R}$.

1. $\lim_{x \rightarrow c} f(x) = L$ if and only if $f(x_n) \rightarrow L$ for every sequence (x_n) with $x_n \in E$ and $x_n \neq c$ for all n and $x_n \rightarrow c$.
2. If two such sequences (x_n) and (y_n) have $f(x_n) \rightarrow L_1$ and $f(y_n) \rightarrow L_2$ with $L_1 \neq L_2$, or if one such sequence has $(f(x_n))$ divergent, then f has no limit at c .

Proof:

1. Sequences of the stated kind exist, by proposition 3.1.9(2) applied to the limit point c . Suppose first that $\lim_{x \rightarrow c} f(x) = L$, and let (x_n) be such a sequence. Let $\epsilon > 0$ and take $\delta > 0$ from definition 3.1.15. Since $x_n \rightarrow c$, there is an N with $|x_n - c| < \delta$ for every $n \geq N$, and $x_n \neq c$ gives $|x_n - c| > 0$ by proposition 1.3.8(1). So $0 < |x_n - c| < \delta$ and hence $|f(x_n) - L| < \epsilon$ for every $n \geq N$, which is $f(x_n) \rightarrow L$.
For the converse, suppose $\lim_{x \rightarrow c} f(x) = L$ fails. Then there is an $\epsilon_0 > 0$ such that no $\delta > 0$ works, that is, such that for every $\delta > 0$ there is an $x \in E$ with $0 < |x - c| < \delta$ and $|f(x) - L| \geq \epsilon_0$. For each $n \in \mathbb{N}$ apply this with $\delta = 1/n$ and call the resulting point x_n . Then $x_n \in E$ and $x_n \neq c$, and $|x_n - c| < 1/n$ gives $x_n \rightarrow c$ by the argument used in proposition 3.1.9(1). But $|f(x_n) - L| \geq \epsilon_0$ for every n , so $(f(x_n))$ does not converge to L .

So the sequential condition fails as well, which is the contrapositive of the claim.

2. Suppose f had a limit L at c . In the first case part (1) gives $f(x_n) \rightarrow L$ and $f(y_n) \rightarrow L$, so $L = L_1$ and $L = L_2$ by proposition 2.1.4, contradicting $L_1 \neq L_2$. In the second case part (1) gives $f(x_n) \rightarrow L$, so $(f(x_n))$ converges, contrary to hypothesis. Hence no such L exists, completing the proof.

Part (2) is how a limit is shown not to exist: produce two sequences approaching c along which f does two different things. Parts (1) and (2) together say that the functional limit carries exactly as much information as the collection of all sequential limits along $E \setminus \{c\}$, which is why corollary 3.1.20 and corollary 3.1.21 need no new estimates.

Corollary 3.1.20: The Algebra of Functional Limits

Let $E \subseteq \mathbb{R}$, let c be a limit point of E , let $f, g: E \rightarrow \mathbb{R}$ with $\lim_{x \rightarrow c} f(x) = A$ and $\lim_{x \rightarrow c} g(x) = B$, and let $\lambda \in \mathbb{R}$.

1. $\lim_{x \rightarrow c} (f(x) + g(x)) = A + B$.
2. $\lim_{x \rightarrow c} \lambda f(x) = \lambda A$.
3. $\lim_{x \rightarrow c} f(x)g(x) = AB$.
4. If $B \neq 0$, then c is a limit point of $E^* = \{x \in E \mid g(x) \neq 0\}$, and the function $f/g: E^* \rightarrow \mathbb{R}$ satisfies $\lim_{x \rightarrow c} f(x)/g(x) = A/B$.

Proof:

1. Let (x_n) be a sequence in E with $x_n \neq c$ and $x_n \rightarrow c$. By theorem 3.1.19(1), $f(x_n) \rightarrow A$ and $g(x_n) \rightarrow B$, so $f(x_n) + g(x_n) \rightarrow A + B$ by theorem 2.1.6(1). Since (x_n) was arbitrary, theorem 3.1.19(1) applied to $f + g$ gives the claim.
2. The same argument with theorem 2.1.6(2).
3. The same argument with theorem 2.1.6(3).
4. Apply definition 3.1.15 to g with $\epsilon = |B|/2$, which is positive by proposition 1.3.8(1): there is a $\delta_0 > 0$ such that $|g(x) - B| < |B|/2$ for every $x \in E$ with $0 < |x - c| < \delta_0$. For such an x , proposition 1.3.8(1) and (5) give

$$|B| - |g(x)| \leq ||B| - |g(x)|| \leq |B - g(x)| = |g(x) - B| < \frac{|B|}{2}$$

so $|g(x)| > |B|/2 > 0$ and hence $g(x) \neq 0$, that is $x \in E^*$. To see that c is a limit point of E^* , let $\delta > 0$ and put $\delta' = \min(\delta, \delta_0) > 0$. Since c is a limit point of E there is an $x \in E$ with $0 < |x - c| < \delta'$, and the previous sentence puts this x in E^* while $0 < |x - c| < \delta$. Now let (x_n) be a sequence in E^* with $x_n \neq c$ and $x_n \rightarrow c$. It is in particular a sequence in E , so theorem 3.1.19(1) gives $f(x_n) \rightarrow A$ and $g(x_n) \rightarrow B$, and $g(x_n) \neq 0$ for every n because $x_n \in E^*$. Theorem 2.1.6(4) gives $f(x_n)/g(x_n) \rightarrow A/B$. Since (x_n) was arbitrary, theorem 3.1.19(1) applied to f/g on E^* completes the proof.

Corollary 3.1.21: The Squeeze Theorem for Functions

Let $E \subseteq \mathbb{R}$, let c be a limit point of E , and let $f, g, h: E \rightarrow \mathbb{R}$. Suppose there is an $r > 0$ with

$$f(x) \leq g(x) \leq h(x) \quad \text{for every } x \in E \text{ with } 0 < |x - c| < r$$

If $\lim_{x \rightarrow c} f(x) = L$ and $\lim_{x \rightarrow c} h(x) = L$, then $\lim_{x \rightarrow c} g(x) = L$.

Proof:

Let (x_n) be a sequence in E with $x_n \neq c$ for every n and $x_n \rightarrow c$. Since $r > 0$, there is an N_0 with $|x_n - c| < r$ for every $n \geq N_0$, and $|x_n - c| > 0$ because $x_n \neq c$. So $f(x_n) \leq g(x_n) \leq h(x_n)$ for every $n \geq N_0$. By theorem 3.1.19(1), $f(x_n) \rightarrow L$ and $h(x_n) \rightarrow L$, so theorem 2.1.7 gives $g(x_n) \rightarrow L$. Since (x_n) was arbitrary, theorem 3.1.19(1) gives $\lim_{x \rightarrow c} g(x) = L$, as we sought to show.

The limit at c is determined by the values of f arbitrarily close to c and by nothing else. Proposition 3.1.22 makes that precise, and one-sided limits are then instances of definition 3.1.15 applied to a restricted domain.

Proposition 3.1.22: The Limit Depends Only on the Values Near the Point

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let c be a limit point of E , let $r > 0$, and put $E_r = E \cap (c - r, c + r)$.

1. c is a limit point of E_r .
2. For $L \in \mathbb{R}$, $\lim_{x \rightarrow c} f(x) = L$ if and only if $\lim_{x \rightarrow c} (f|_{E_r})(x) = L$.
3. If $g: E \rightarrow \mathbb{R}$ satisfies $g(x) = f(x)$ for every $x \in E_r$ with $x \neq c$, then f has a limit at c exactly when g does, and the two limits are equal.

Proof:

1. Let $\delta > 0$ and put $\delta' = \min(\delta, r) > 0$. Since c is a limit point of E there is a $y \in E$ with $0 < |y - c| < \delta'$. Then $|y - c| < r$ gives $c - r < y < c + r$ by proposition 1.3.8(2), so $y \in E_r$, and $0 < |y - c| < \delta$.
2. Suppose $\lim_{x \rightarrow c} f(x) = L$ and let $\epsilon > 0$. The δ furnished by definition 3.1.15 for f also works for $f|_{E_r}$, since every $x \in E_r$ with $0 < |x - c| < \delta$ is a point of E with the same property. Conversely, suppose $\lim_{x \rightarrow c} (f|_{E_r})(x) = L$ and let $\epsilon > 0$. Take $\delta > 0$ from the definition for $f|_{E_r}$ and put $\delta'' = \min(\delta, r) > 0$. If $x \in E$ and $0 < |x - c| < \delta''$, then $|x - c| < r$ puts x in E_r , and $0 < |x - c| < \delta$ gives $|f(x) - L| < \epsilon$.
3. The functions $f|_{E_r}$ and $g|_{E_r}$ agree at every point of E_r other than c , and definition 3.1.15 never evaluates either at c , so the two satisfy exactly the same instances of the definition and have the same limits at c . Now apply (2) to f and then to g , completing the proof.

Proposition 3.1.23: Limits Along a Subset

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let $c \in \mathbb{R}$ and $L \in \mathbb{R}$.

1. Let $D \subseteq E$ and suppose c is a limit point of D . Then c is a limit point of E , and if $\lim_{x \rightarrow c} f(x) = L$ then $\lim_{x \rightarrow c} (f|_D)(x) = L$.
2. Let $D_1, D_2 \subseteq E$ with $E \setminus \{c\} \subseteq D_1 \cup D_2$, suppose c is a limit point of both D_1 and D_2 , and suppose $\lim_{x \rightarrow c} (f|_{D_1})(x) = L$ and $\lim_{x \rightarrow c} (f|_{D_2})(x) = L$. Then $\lim_{x \rightarrow c} f(x) = L$.

Proof:

1. For each $\delta > 0$, a point $y \in D$ with $0 < |y - c| < \delta$ is also a point of E , so c is a limit point of E . Given $\epsilon > 0$, take $\delta > 0$ from definition 3.1.15 for f . Every $x \in D$ with $0 < |x - c| < \delta$ lies in E , so $|f|_D(x) - L| = |f(x) - L| < \epsilon$.
2. By (1) applied to D_1 , the point c is a limit point of E , so $\lim_{x \rightarrow c} f(x)$ is a meaningful assertion. Given $\epsilon > 0$, take $\delta_1 > 0$ for $f|_{D_1}$ and $\delta_2 > 0$ for $f|_{D_2}$, and put $\delta = \min(\delta_1, \delta_2) > 0$. Let $x \in E$ with $0 < |x - c| < \delta$. Then $x \neq c$ by proposition 1.3.8(1), so $x \in D_1 \cup D_2$. If $x \in D_1$ then $0 < |x - c| < \delta_1$ gives $|f(x) - L| < \epsilon$, and if $x \in D_2$ the same holds with δ_2 , completing the proof.

Definition 3.1.24: One-Sided Limits

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let $c \in \mathbb{R}$. Put

$$E_c^+ = E \cap (c, \infty), \quad E_c^- = E \cap (-\infty, c)$$

If c is a limit point of E_c^+ , the *right-hand limit* of f at c is

$$\lim_{x \rightarrow c^+} f(x) := \lim_{x \rightarrow c} (f|_{E_c^+})(x)$$

whenever the right-hand side exists. If c is a limit point of E_c^- , the *left-hand limit* $\lim_{x \rightarrow c^-} f(x)$ is defined the same way with E_c^- in place of E_c^+ .

For $E = \mathbb{R} \setminus \{0\}$, $f(x) = |x|/x$ and $c = 0$, the two sets are $E_0^+ = (0, \infty)$ and $E_0^- = (-\infty, 0)$, and f takes the constant value 1 on the first and the constant value -1 on the second. The right-hand limit at 0 is then 1 and the left-hand limit is -1 , and example 3.1.30(1) carries out the estimates.

A one-sided limit is therefore the limit of a restriction, and it needs no theory of its own. What it does need is a direct description, so that a computation does not have to pass through the restricted domain each time.

Proposition 3.1.25: The Direct Description of a One-Sided Limit

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let $c \in \mathbb{R}$ and $L \in \mathbb{R}$.

1. c is a limit point of E_c^+ if and only if for every $\delta > 0$ there is an $x \in E$ with $c < x < c + \delta$.
2. Assume c is a limit point of E_c^+ . Then $\lim_{x \rightarrow c^+} f(x) = L$ if and only if for every $\epsilon > 0$ there is a $\delta > 0$ with $|f(x) - L| < \epsilon$ for every $x \in E$ satisfying $c < x < c + \delta$.
3. Assume c is a limit point of E_c^+ . Then $\lim_{x \rightarrow c^+} f(x) = L$ if and only if $f(x_n) \rightarrow L$ for every sequence (x_n) with $x_n \in E$ and $x_n > c$ for all n and $x_n \rightarrow c$.

The three statements hold with E_c^- , $c - \delta < x < c$ and $x_n < c$ in place of E_c^+ , $c < x < c + \delta$ and $x_n > c$.

Proof:

For $x \in E$ the two conditions

$$x \in E_c^+ \text{ and } 0 < |x - c| < \delta, \quad c < x < c + \delta$$

are equivalent. Indeed, if $x > c$ then $x - c > 0$, so $|x - c| = x - c$ by definition 1.3.6, and $0 < x - c < \delta$ is exactly $c < x < c + \delta$. Conversely $c < x < c + \delta$ gives $x > c$ and $0 < x - c < \delta$. Parts (1) and (2) are now definition 3.1.5 and definition 3.1.15 for the function $f|_{E_c^+}$, with the left-hand condition of the display replaced by the right-hand one. Part (3) is theorem 3.1.19(1) applied to $f|_{E_c^+}$, once one notes that a sequence in E_c^+ with no term equal to c is the same thing as a sequence in E with every term greater than c , the case $x_n = c$ being impossible inside E_c^+ . The left-handed statements follow by the same computation with $|x - c| = c - x$ for $x < c$, completing the proof.

Proposition 3.1.26: A Limit Exists Exactly When the One-Sided Limits Agree

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$, let c be a limit point of E and let $L \in \mathbb{R}$. Then c is a limit point of at least one of E_c^+ and E_c^- , and:

1. If c is a limit point of both, then $\lim_{x \rightarrow c} f(x) = L$ if and only if both one-sided limits at c exist and are equal to L .
2. If c is a limit point of E_c^+ and not of E_c^- , then $\lim_{x \rightarrow c} f(x) = L$ if and only if $\lim_{x \rightarrow c^+} f(x) = L$. Symmetrically with the two sides exchanged.

Proof:

Suppose c were a limit point of neither. Then there are $\delta_+ > 0$ admitting no $y \in E_c^+$ with $0 < |y - c| < \delta_+$, and $\delta_- > 0$ admitting no such $y \in E_c^-$. Put $\delta = \min(\delta_+, \delta_-) > 0$. Since c is a limit point of E there is a $y \in E$ with $0 < |y - c| < \delta$. This y satisfies $y \neq c$ by proposition 1.3.8(1), so $y > c$ or $y < c$, placing y in E_c^+ or in E_c^- and contradicting the choice of δ_+ or of δ_- .

1. If $\lim_{x \rightarrow c} f(x) = L$, then proposition 3.1.23(1) applied to $D = E_c^+$ and to $D = E_c^-$ gives both one-sided limits, equal to L . Conversely, every $x \in E$ with $x \neq c$ satisfies $x > c$ or

$x < c$, so $E \setminus \{c\} \subseteq E_c^+ \cup E_c^-$, and proposition 3.1.23(2) with $D_1 = E_c^+$ and $D_2 = E_c^-$ gives $\lim_{x \rightarrow c} f(x) = L$.

2. The forward implication is proposition 3.1.23(1) with $D = E_c^+$. For the converse, since c is not a limit point of E_c^- there is an $r > 0$ admitting no $y \in E_c^-$ with $0 < |y - c| < r$. Let $\epsilon > 0$ and take $\delta_+ > 0$ from proposition 3.1.25(2). Put $\delta = \min(\delta_+, r) > 0$ and let $x \in E$ with $0 < |x - c| < \delta$. Then $x \neq c$. If $x < c$ then $x \in E_c^-$ and $0 < |x - c| < r$, which is impossible, so $x > c$, and then $c < x < c + \delta_+$ gives $|f(x) - L| < \epsilon$. Hence $\lim_{x \rightarrow c} f(x) = L$, completing the proof.

Two ends of the definition remain. A domain may be unbounded above, and the question is then what f does at large x . The values of f may also grow beyond every bound. Both cases are recorded by relaxing one half of definition 3.1.15 while keeping the other.

Definition 3.1.27: Limits at Infinity

Let $E \subseteq \mathbb{R}$ be unbounded above, let $f: E \rightarrow \mathbb{R}$ and let $L \in \mathbb{R}$. Write $\lim_{x \rightarrow \infty} f(x) = L$ if for every real $\epsilon > 0$ there is an $M \in \mathbb{R}$ such that

$$|f(x) - L| < \epsilon \quad \text{for every } x \in E \text{ with } x > M$$

For E unbounded below, $\lim_{x \rightarrow -\infty} f(x) = L$ is defined the same way with $x < M$ in place of $x > M$.

These limits are unique as well. If $\lim_{x \rightarrow \infty} f(x)$ were both L and L' with $L \neq L'$, then taking $\epsilon = |L - L'|/2$ produces M and M' . Since E is unbounded above there is an $x_0 \in E$ with $x_0 > \max(M, M')$, and proposition 1.3.8(4) gives $|L - L'| \leq |L - f(x_0)| + |f(x_0) - L'| < |L - L'|$, which is impossible. The unboundedness of E here does the work that the limit-point hypothesis does in proposition 3.1.18.

Definition 3.1.28: Infinite Limits

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let c be a limit point of E . Write $\lim_{x \rightarrow c} f(x) = +\infty$ if for every $M \in \mathbb{R}$ there is a real $\delta > 0$ such that

$$f(x) > M \quad \text{for every } x \in E \text{ with } 0 < |x - c| < \delta$$

and $\lim_{x \rightarrow c} f(x) = -\infty$ if for every $M \in \mathbb{R}$ there is a $\delta > 0$ with $f(x) < M$ for every such x . The one-sided forms $\lim_{x \rightarrow c^\pm} f(x) = \pm\infty$ are the corresponding statements for the restrictions of definition 3.1.24. For E unbounded above, $\lim_{x \rightarrow \infty} f(x) = +\infty$ means that for every $M \in \mathbb{R}$ there is an $N \in \mathbb{R}$ with $f(x) > M$ for every $x \in E$ with $x > N$, and the remaining three combinations of signs are defined analogously.

The symbols $+\infty$ and $-\infty$ are those of definition 2.2.4 and are not real numbers, so “ $\lim_{x \rightarrow c} f(x) = +\infty$ ” is a statement about divergence and not the assertion that some limit exists. The two situations never overlap: if $\lim_{x \rightarrow c} f(x) = L$ for a real L , then taking $\epsilon = 1$ gives a δ with $|f(x) - L| < 1$, hence $f(x) < L + 1$ by proposition 1.3.8(2), for every $x \in E$ with $0 < |x - c| < \delta$, while $\lim_{x \rightarrow c} f(x) = +\infty$ with $M = L + 1$ gives a δ' with $f(x) > L + 1$ for every such x with δ' in place of δ . Since c is a limit point there is a point of E satisfying both restrictions, and it would give $f(x) < L + 1 < f(x)$.

Two facts about these limits are used later. The first is the algebra of corollary 3.1.20 with M in place of δ , and the second is the function-side form of theorem 2.2.2.

Proposition 3.1.29: Limits at Infinity: Algebra and Monotone Convergence

Let $E \subseteq \mathbb{R}$ be unbounded above and let $f, g: E \rightarrow \mathbb{R}$.

1. If $\lim_{x \rightarrow \infty} f(x) = L$ and $\lim_{x \rightarrow \infty} g(x) = L'$, then $\lim_{x \rightarrow \infty} (f + g)(x) = L + L'$ and $\lim_{x \rightarrow \infty} (f - g)(x) = L - L'$.
2. If f is increasing on E and bounded above, then $\lim_{x \rightarrow \infty} f(x) = \sup_{x \in E} f(x)$, which is finite. If f is increasing and not bounded above, then $\lim_{x \rightarrow \infty} f(x) = +\infty$.

Proof:

Part (1). Let $\epsilon > 0$. Take M_1 from definition 3.1.27 for f at $\epsilon/2$ and M_2 for g at $\epsilon/2$. For $x \in E$ with $x > \max(M_1, M_2)$, proposition 1.3.8(4) gives $|(f + g)(x) - (L + L')| \leq |f(x) - L| + |g(x) - L'| < \epsilon$. The difference is the same computation with $-g$, whose limit is $-L'$ because $|-g(x) + L'| = |g(x) - L'|$.

Part (2). Suppose f is bounded above and put $S = \sup_{x \in E} f(x)$, which exists by theorem 1.2.4 since the range is nonempty and bounded above. Let $\epsilon > 0$. By proposition 1.2.6 there is an $x_0 \in E$ with $f(x_0) > S - \epsilon$. For $x \in E$ with $x > x_0$ the increase of f gives $S - \epsilon < f(x_0) \leq f(x) \leq S$, so $|f(x) - S| < \epsilon$, and $M = x_0$ answers ϵ in definition 3.1.27. If instead f is not bounded above, then for every $M \in \mathbb{R}$ there is an $x_0 \in E$ with $f(x_0) > M$, and every $x > x_0$ in E has $f(x) \geq f(x_0) > M$, which is definition 3.1.28's condition at infinity.

Example 3.1.30: Three One-Sided and Infinite Limits

1. $E = \mathbb{R} \setminus \{0\}$ and $f(x) = |x|/x$, with $c = 0$. Every $\delta > 0$ admits $x = \min(\delta, 1)/2 \in E$ with $0 < x < \delta$ and admits $-x$, so 0 is a limit point of both E_0^+ and E_0^- . For $x > 0$ one has $|x| = x$ and $f(x) = 1$, so proposition 3.1.25(2) with any δ gives $\lim_{x \rightarrow 0^+} f(x) = 1$; for $x < 0$ one has $|x| = -x$ and $f(x) = -1$, giving $\lim_{x \rightarrow 0^-} f(x) = -1$. Since $1 \neq -1$, proposition 3.1.26(1) shows that f has no limit at 0.
2. $E = \mathbb{R} \setminus \{0\}$ and $f(x) = 1/x^2$, with $c = 0$. Then $\lim_{x \rightarrow 0} f(x) = +\infty$. Every $x \in E$ has $x \neq 0$, so $x^2 > 0$ by proposition 1.1.2(6), and applying proposition 1.1.2(7) to $0 < x^2 < x^2 + 1$ gives $1/x^2 > 0$. Let $M \in \mathbb{R}$. If $M \leq 0$, take $\delta = 1$, and the previous sentence gives $1/x^2 > 0 \geq M$ for every $x \in E$. If $M > 0$, then $1/M > 0$ by proposition 1.1.2(7), and theorem 1.3.12 with $n = 2$ gives a $\delta \geq 0$ with $\delta^2 = 1/M$. Here $\delta \neq 0$, since $\delta = 0$ would give $1/M = 0$. For $x \in E$ with $0 < |x - 0| < \delta$ one has $0 < |x| < \delta$. Two applications of proposition 1.1.2(5), first multiplying by $|x| > 0$ and then by $\delta > 0$, give $|x|^2 < |x|\delta < \delta^2$, and $|x|^2 = |x^2| = x^2$ by proposition 1.3.8(3) and definition 1.3.6. So $0 < x^2 < \delta^2 = 1/M$, and proposition 1.1.2(7) applied to this gives $1/x^2 > M$.
3. $E = (3, \infty)$ and $f(x) = (2x + 1)/(x - 3)$. Then E is unbounded above: given $y \in \mathbb{R}$, theorem 1.4.1(2) produces an $n \in \mathbb{N}$ with $n > \max(y, 3)$, and such an n lies in E and exceeds

y. Moreover $\lim_{x \rightarrow \infty} f(x) = 2$. For $x \in E$ one has $x - 3 > 0$, and

$$f(x) - 2 = \frac{2x + 1 - 2(x - 3)}{x - 3} = \frac{7}{x - 3}$$

Given $\epsilon > 0$ put $M = 3 + 7/\epsilon$. If $x \in E$ and $x > M$ then $x - 3 > 7/\epsilon > 0$, so $0 < 7/(x - 3) < \epsilon$ by proposition 1.1.2(7) and (5), and $|f(x) - 2| < \epsilon$.

The last two definitions inherit no algebra. Nothing like corollary 3.1.20 holds for them, and the difference of two functions tending to $+\infty$ can do anything. On $E = \mathbb{R} \setminus \{0\}$ with $c = 0$, the function $1/x^2$ tends to $+\infty$ by example 3.1.30(2), and so does $2/x^2$, since $2/x^2 > 1/x^2 > M$ wherever the estimate there holds. Yet $2/x^2 - 1/x^2 = 1/x^2$ tends to $+\infty$, the difference of $1/x^2$ with itself is the constant 0, and $1/x^2 - 2/x^2 = -1/x^2$ tends to $-\infty$, because $1/x^2 > -M$ gives $-1/x^2 < M$. What these definitions give instead is a vocabulary precise enough to say what a function does at a point where it has no limit.

Exercise 3.1.1

1. Let $E = \{1/n | n \in \mathbb{N}\}$. Determine the set of limit points of E , determine which points of E are isolated, and compute $\overline{E}$. Prove each of the three answers.
2. Let $E \subseteq \mathbb{R}$. Prove that $\overline{\overline{E}} = \overline{E}$, and conclude that $\overline{E}$ is closed for every E .
3. Let $A, B \subseteq \mathbb{R}$. Prove that $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$, and give sets A and B for which the inclusion is strict. Compare with proposition 3.1.8(5), where the corresponding statement for unions is an equality.
4. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = x^2$ and let $c \in \mathbb{R}$. Given $\epsilon > 0$, exhibit an explicit $\delta > 0$, depending on ϵ and c , that witnesses $\lim_{x \rightarrow c} f(x) = c^2$ in definition 3.1.15. Then show that no single $\delta > 0$ works for $\epsilon = 1$ at every $c \in \mathbb{R}$ simultaneously.
5. Let $E \subseteq \mathbb{R}$, let c be a limit point of E and let $f: E \rightarrow \mathbb{R}$ with $\lim_{x \rightarrow c} f(x) = L$. Prove that $\lim_{x \rightarrow c} |f(x)| = |L|$. Then give an E , a c and an f for which $\lim_{x \rightarrow c} |f(x)|$ exists and $\lim_{x \rightarrow c} f(x)$ does not.
6. Let $E \subseteq \mathbb{R}$ be nonempty and bounded above. Prove that $\sup E \in \overline{E}$, and deduce that a nonempty closed set that is bounded above contains its supremum.
7. Let $f: (0, \infty) \rightarrow \mathbb{R}$ be $f(x) = 1/x$. Prove directly from definition 3.1.27 that $\lim_{x \rightarrow \infty} f(x) = 0$, and use proposition 3.1.25(3) to show that there is no $L \in \mathbb{R}$ with $\lim_{x \rightarrow 0^+} f(x) = L$.
8. Let $E \subseteq \mathbb{R}$, let c be a limit point of E , and let $f, g: E \rightarrow \mathbb{R}$ satisfy $f(x) \leq g(x)$ for every $x \in E$. Suppose both limits at c exist. Prove that $\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x)$. Then give an example with $f(x) < g(x)$ for every $x \in E$ in which the two limits are equal.

3.2 Continuous Functions

Section 3.1 attached a limit to a point its domain approaches and read nothing from the function's value there. Continuity at a point is the condition that the value and the limit agree, and the definition below states it by dropping the restriction $x \neq c$ from the ϵ - δ sentence. Two things change

with it. The point c must now lie in the domain, since $f(c)$ occurs in the inequality, and c need no longer be a limit point of the domain, since the condition can be satisfied with no nearby domain points at all.

Definition 3.2.1: [Real] Continuity at a Point

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let $c \in E$. Then f is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that

$$|f(x) - f(c)| < \epsilon \quad \text{for every } x \in E \text{ with } |x - c| < \delta$$

If f is not continuous at c , then f is *discontinuous at c* , and c is a *discontinuity* of f .

Definition 3.2.2: Continuous Function

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$. Then f is *continuous* if it is continuous at every point of E . For $A \subseteq E$, the function f is *continuous on A* if the restriction of f to A is continuous in this sense.

The clause about a subset is not idle. The restriction of f to A is a function on the smaller domain A , so its continuity at $c \in A$ quantifies only over the points of A within δ of c , which is a weaker condition than continuity of f at c . The Dirichlet function of example 3.2.10 is continuous on $\mathbb{Q}$ and continuous at no point of $\mathbb{R}$.

A constant function $f(x) = \lambda$ on E is continuous at every $c \in E$, since $|f(x) - f(c)| = |\lambda - \lambda| = 0 < \epsilon$ for every $x \in E$, so any $\delta > 0$ answers any ϵ . For $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 3x - 4$ and any $c \in \mathbb{R}$,

$$|f(x) - f(c)| = |3x - 3c| = 3|x - c|$$

the last equality by proposition 1.3.8(3). Given $\epsilon > 0$, put $\delta = \epsilon/3$, which is positive by proposition 1.1.2(7). Then $|x - c| < \delta$ gives $|f(x) - f(c)| = 3|x - c| < 3\delta = \epsilon$, using proposition 1.1.2(5) to multiply the inequality by $3 > 0$. So f is continuous.

The definition of the functional limit required c to be a limit point of E , and box 3.1.16(2) showed why: at an isolated point every candidate value satisfies the ϵ - δ condition vacuously, so no limit is determined. Definition 3.2.1 carries no such hypothesis, and the vacuous case does no harm here, because $f(c)$ is already fixed by f and nothing has to be determined. The next proposition records both halves.

Proposition 3.2.3: Continuity Against the Functional Limit

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let $c \in E$.

1. If c is a limit point of E , then f is continuous at c if and only if $\lim_{x \rightarrow c} f(x)$ exists and equals $f(c)$.
2. If c is an isolated point of E , then f is continuous at c .

Proof:

1. Suppose f is continuous at c and let $\epsilon > 0$. Take $\delta > 0$ as in definition 3.2.1. Every $x \in E$ with $0 < |x - c| < \delta$ satisfies $|f(x) - f(c)| < \epsilon$. Since c is a limit point of E , this is the condition of definition 3.1.15 for the value $f(c)$, so $\lim_{x \rightarrow c} f(x) = f(c)$.

Conversely suppose $\lim_{x \rightarrow c} f(x) = f(c)$ and let $\epsilon > 0$. Definition 3.1.15 gives a $\delta > 0$ with $|f(x) - f(c)| < \epsilon$ for every $x \in E$ satisfying $0 < |x - c| < \delta$. The only point of E with $|x - c| < \delta$ not covered by that statement is $x = c$ itself, and there $|f(c) - f(c)| = 0 < \epsilon$ by proposition 1.3.8(1). So the same δ answers ϵ in definition 3.2.1.

2. By definition 3.1.6 the point c lies in E and is not a limit point of E . Negating definition 3.1.5, there is a $\delta_0 > 0$ such that no $x \in E$ satisfies $0 < |x - c| < \delta_0$. Let $\epsilon > 0$ and take $\delta = \delta_0$. Suppose $x \in E$ and $|x - c| < \delta_0$. The quantity $|x - c|$ is nonnegative by proposition 1.3.8(1) and cannot be positive by the previous sentence, so it is 0 and $x = c$, again by proposition 1.3.8(1). Hence $|f(x) - f(c)| = 0 < \epsilon$, and f is continuous at c , completing the proof.

Part (2) says that continuity detects nothing at an isolated point. A function on $\mathbb{Z}$, or on $\{1/n \mid n \in \mathbb{N}\}$, is continuous no matter how its values are assigned, since each point of either set is isolated. Continuity is a statement about a function together with the way its domain approaches each point, and where the domain offers no approach the statement is empty.

The characterization by sequences is the form in which continuity is applied below. It holds at every point of the domain, with no limit-point hypothesis, which theorem 3.1.19 cannot dispense with.

Proposition 3.2.4: The Sequential Characterization of Continuity

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let $c \in E$. Then f is continuous at c if and only if $f(x_n) \rightarrow f(c)$ for every sequence $(x_n)_{n \geq 1}$ of points of E with $x_n \rightarrow c$.

Proof:

Step 1: Continuity gives the sequential condition. Suppose f is continuous at c and let (x_n) be a sequence in E with $x_n \rightarrow c$. Let $\epsilon > 0$ and take $\delta > 0$ as in definition 3.2.1. By definition 2.1.2 applied to (x_n) with the positive real δ , there is an $N \in \mathbb{N}$ with $|x_n - c| < \delta$ for every $n \geq N$. Each such x_n lies in E , so $|f(x_n) - f(c)| < \epsilon$ for every $n \geq N$. Since $\epsilon > 0$ was arbitrary, $f(x_n) \rightarrow f(c)$.

Step 2: The sequential condition gives continuity. The argument is by contraposition. Suppose f is not continuous at c . Negating definition 3.2.1, there is an $\epsilon_0 > 0$ such that for every $\delta > 0$ some $x \in E$ satisfies $|x - c| < \delta$ and $|f(x) - f(c)| \geq \epsilon_0$. For each $n \in \mathbb{N}$ apply this with $\delta = 1/n$, positive by proposition 1.1.2(7), and let $x_n \in E$ be a point it produces.^a

The sequence (x_n) converges to c . Given $\epsilon > 0$, theorem 1.4.1(3) gives an $N \in \mathbb{N}$ with

$0 < 1/N < \epsilon$, and for $n \geq N$ the bound $|x_n - c| < 1/n \leq 1/N < \epsilon$ holds, the middle inequality by proposition 1.1.2(7). But $|f(x_n) - f(c)| \geq \epsilon_0$ for every n , so no index can force $|f(x_n) - f(c)| < \epsilon_0$, and $(f(x_n))$ does not converge to $f(c)$. So the sequential condition fails, completing the proof.

^aSelecting one such x_n for every n at once is an appeal to countable choice. Unlike the least-index choices used for subsequences in theorem 2.2.9, the sets drawn from here are sets of real numbers and carry no canonical element.

The results of chapter 2 about sequences now transfer to functions, and the first case is the algebra. The proof does not repeat the ϵ - δ estimates of theorem 2.1.6, which were carried out once there and enter here through proposition 3.2.4.

Theorem 3.2.5: The Algebra of Continuous Functions

Let $E \subseteq \mathbb{R}$, let $f, g: E \rightarrow \mathbb{R}$ be continuous at $c \in E$ and let $\lambda \in \mathbb{R}$. Then

1. $f + g$ is continuous at c ;
2. λf is continuous at c ;
3. fg is continuous at c ;
4. if $g(c) \neq 0$, then f/g , viewed as a function on $E^* = \{x \in E \mid g(x) \neq 0\}$, is continuous at c .

Consequently, if f and g are continuous on E , then so are $f + g$, λf and fg , and f/g is continuous on E^* .

Proof:

1. Let (x_n) be a sequence in E with $x_n \rightarrow c$. By proposition 3.2.4 applied to f and then to g , $f(x_n) \rightarrow f(c)$ and $g(x_n) \rightarrow g(c)$. By theorem 2.1.6(1), $(f + g)(x_n) = f(x_n) + g(x_n) \rightarrow f(c) + g(c) = (f + g)(c)$. Since (x_n) was an arbitrary sequence in E converging to c , proposition 3.2.4 makes $f + g$ continuous at c .
2. The same sequence argument with theorem 2.1.6(2) gives $\lambda f(x_n) \rightarrow \lambda f(c)$.
3. The same sequence argument with theorem 2.1.6(3) gives $f(x_n)g(x_n) \rightarrow f(c)g(c)$.
4. Since $g(c) \neq 0$, the point c lies in E^* , so f/g is defined at c and the statement has content. Let (x_n) be a sequence in E^* with $x_n \rightarrow c$. Every x_n lies in E , so $f(x_n) \rightarrow f(c)$ and $g(x_n) \rightarrow g(c)$ as above. The hypotheses of theorem 2.1.6(4) hold: the limit $g(c)$ is nonzero, and $g(x_n) \neq 0$ for every n because $x_n \in E^*$. So $f(x_n)/g(x_n) \rightarrow f(c)/g(c)$, and proposition 3.2.4, applied to the function f/g on the domain E^* , makes f/g continuous at c .

The final sentence is each part applied at every point of the relevant domain, completing the proof.

Taking $\lambda = -1$ in part (2) shows that f is continuous at c exactly when $-f$ is, since applying (2) twice returns f . That symmetry transfers every statement below about increasing functions to decreasing ones.

Example 3.2.6: Polynomials and Rational Functions

1. The identity function $f(x) = x$ on $\mathbb{R}$ is continuous: given $c \in \mathbb{R}$ and $\epsilon > 0$, the choice $\delta = \epsilon$ gives $|f(x) - f(c)| = |x - c| < \epsilon$ whenever $|x - c| < \delta$. Constant functions are continuous by the computation following definition 3.2.2. By induction on n , the function $x \mapsto x^n$ is continuous on $\mathbb{R}$ for every integer $n \geq 0$: the case $n = 0$ is the constant 1, and if $x \mapsto x^n$ is continuous then so is $x \mapsto x^{n+1} = x \cdot x^n$ by theorem 3.2.5(3). Applying theorem 3.2.5(2) to each $a_k x^k$ and then theorem 3.2.5(1) repeatedly, every polynomial

$$p(x) = a_0 + a_1 x + \cdots + a_n x^n$$

with real coefficients is continuous on $\mathbb{R}$. If p and q are polynomials, theorem 3.2.5(4) makes the rational function p/q continuous on $\{x \in \mathbb{R} \mid q(x) \neq 0\}$.

2. The general argument hides the estimate, so here is one case run in full. Fix $c \in \mathbb{R}$ and let $f(x) = x^2$. For any x ,

$$|f(x) - f(c)| = |x^2 - c^2| = |x - c| |x + c|$$

by proposition 1.3.8(3), since $x^2 - c^2 = (x - c)(x + c)$. Write $x + c = (x - c) + 2c$, so that $|x + c| \leq |x - c| + 2|c|$ by parts (3) and (4) of the same proposition. Given $\epsilon > 0$, put

$$\delta = \min \left\{ 1, \frac{\epsilon}{1 + 2|c|} \right\}$$

which is positive because $1 + 2|c| \geq 1 > 0$ by proposition 1.3.8(1). If $|x - c| < \delta$ then $|x - c| < 1$, so $|x + c| < 1 + 2|c|$, and therefore

$$|f(x) - f(c)| < \delta (1 + 2|c|) \leq \frac{\epsilon}{1 + 2|c|} (1 + 2|c|) = \epsilon$$

So $x \mapsto x^2$ is continuous at c . The δ depends on c as well as on ϵ , and it shrinks as $|c|$ grows. Section 3.4 takes up the question of when that dependence can be removed.

Composition is the operation the functional limit handles badly and continuity handles cleanly.

Theorem 3.2.7: Compositions of Continuous Functions

Let $E, F \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ satisfy $f(E) \subseteq F$ and let $g: F \rightarrow \mathbb{R}$. If f is continuous at $c \in E$ and g is continuous at $f(c)$, then $g \circ f: E \rightarrow \mathbb{R}$ is continuous at c . If f is continuous on E and g is continuous on F , then $g \circ f$ is continuous on E .

Proof:

Write $b = f(c)$, a point of F because $f(E) \subseteq F$. Let $\epsilon > 0$. Since g is continuous at b , definition 3.2.1 gives an $\eta > 0$ such that

$$|g(y) - g(b)| < \epsilon \quad \text{for every } y \in F \text{ with } |y - b| < \eta$$

Since f is continuous at c and $\eta > 0$, definition 3.2.1 gives a $\delta > 0$ such that $|f(x) - b| < \eta$ for every $x \in E$ with $|x - c| < \delta$. Fix such an x and put $y = f(x)$. Then $y \in F$ and $|y - b| < \eta$, so the displayed inequality applies and gives

$$|(g \circ f)(x) - (g \circ f)(c)| = |g(y) - g(b)| < \epsilon$$

So $g \circ f$ is continuous at c . The second sentence is the first applied at every $c \in E$, using that $f(c) \in F$ for each such c , completing the proof.

The step that carries the proof is the one where $y = f(x)$ is fed to the inequality for g without asking whether $y = b$. A functional limit forbids exactly that question, since $\lim_{y \rightarrow b} g(y)$ is computed with $y = b$ excluded, and the next example shows that the corresponding statement for limits is false.

Example 3.2.8: The Composition of Two Functional Limits

Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by

$$g(y) = \begin{cases} 1 & \text{if } y = 0 \\ 0 & \text{if } y \neq 0 \end{cases}$$

Then $\lim_{y \rightarrow 0} g(y) = 0$: given $\epsilon > 0$, every y with $0 < |y - 0| < 1$ satisfies $y \neq 0$ and hence $|g(y) - 0| = 0 < \epsilon$, so $\delta = 1$ answers every ϵ .

1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be the constant function $f(x) = 0$. Then $\lim_{x \rightarrow 0} f(x) = 0$ by the same argument, and $g(f(x)) = g(0) = 1$ for every x , so $\lim_{x \rightarrow 0} g(f(x)) = 1$. Both inner and outer limits are 0 while the composite has limit 1.
2. The failure does not depend on f being constant. Define $h: \mathbb{R} \rightarrow \mathbb{R}$ by $h(x) = 0$ for $x \in \mathbb{Q}$ and $h(x) = x$ for $x \notin \mathbb{Q}$. Then $\lim_{x \rightarrow 0} h(x) = 0$, since $|h(x) - 0| \leq |x|$ for every x , so $\delta = \epsilon$ answers ϵ . The composite $g \circ h$ has no limit at 0 at all. For each $n \in \mathbb{N}$, theorem 1.4.4 applied to $0 < 1/n$ gives a rational p_n with $0 < p_n < 1/n$, and corollary 1.4.5 applied to the same pair gives an irrational t_n with $0 < t_n < 1/n$. Both sequences converge to 0: given $\epsilon > 0$, theorem 1.4.1(3) produces N with $0 < 1/N < \epsilon$, and for $n \geq N$ both $|p_n| < 1/n \leq 1/N < \epsilon$ and $|t_n| < 1/n \leq 1/N < \epsilon$. Both sequences also avoid 0. But $g(h(p_n)) = g(0) = 1$ for every n , while $g(h(t_n)) = g(t_n) = 0$ for every n , since $t_n \neq 0$. Two sequences approaching 0 from

within $\mathbb{R} \setminus \{0\}$ produce composite values with different limits, so theorem 3.1.19 rules out any limit for $g \circ h$ at 0.

What breaks in both cases is that the inner function attains the value 0, and 0 is the one point deleted in the computation of the outer limit. Redefining $g(0) = 0$ makes g continuous at 0 and removes both pathologies, which is theorem 3.2.7 at work.

Example 3.2.9: The Absolute Value and the Pointwise Maximum

The function $x \mapsto |x|$ is continuous on $\mathbb{R}$. Indeed proposition 1.3.8(5) gives

$$||x| - |c|| \leq |x - c|$$

for all $x, c \in \mathbb{R}$, so $\delta = \epsilon$ answers every $\epsilon > 0$ at every c . Combining this with theorem 3.2.7, if $f: E \rightarrow \mathbb{R}$ is continuous then so is $|f|$, the function $x \mapsto |f(x)|$.

For $f, g: E \rightarrow \mathbb{R}$ write $\max(f, g)$ for the function $x \mapsto \max\{f(x), g(x)\}$ and $\min(f, g)$ for $x \mapsto \min\{f(x), g(x)\}$. These are continuous whenever f and g are, because of the identities

$$\max(f, g) = \frac{f + g + |f - g|}{2}, \quad \min(f, g) = \frac{f + g - |f - g|}{2}$$

which hold pointwise. At a point where $f(x) \geq g(x)$ the right-hand side of the first is $(f(x) + g(x) + f(x) - g(x))/2 = f(x)$, and at a point where $f(x) < g(x)$ it is $(f(x) + g(x) + g(x) - f(x))/2 = g(x)$. The second identity is checked the same way. The right-hand sides are built from f , g and $|f - g|$ by the operations of theorem 3.2.5, so both are continuous.

That is the whole catalogue available at this point: polynomials, rational functions, the absolute value, and everything obtained from them by the algebra and by composition. The exponential, cosine and sine functions were defined by their power series in definition 2.6.5, but nothing proved so far says that a power series defines a continuous function, and their continuity is established in Section 4.6. The n -th root function is absent for a different reason: its continuity follows from a theorem about inverses of continuous *strictly* monotone functions, which is theorem 3.3.12 in the next section.

The catalogue is short, and the three functions that follow show why a longer one would still leave open the question of which subsets of $\mathbb{R}$ occur as sets of discontinuities. Each treats rational and irrational arguments differently, and between them they realize $\mathbb{R}$, $\mathbb{R} \setminus \{0\}$ and $\mathbb{Q}$.

Example 3.2.10: The Dirichlet Function

The *Dirichlet function* $\mathbf{1}_{\mathbb{Q}}: \mathbb{R} \rightarrow \mathbb{R}$ is the indicator of the rationals,

$$\mathbf{1}_{\mathbb{Q}}(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

It is discontinuous at every point of $\mathbb{R}$.

Let $c \in \mathbb{R}$. For each $n \in \mathbb{N}$, theorem 1.4.4 applied to $c < c + 1/n$ gives a rational p_n with $c < p_n < c + 1/n$, and corollary 1.4.5 applied to the same pair gives an irrational t_n with $c < t_n < c + 1/n$. Both sequences converge to c : given $\epsilon > 0$, theorem 1.4.1(3) produces N with

$0 < 1/N < \epsilon$, and for $n \geq N$ both $|p_n - c| < 1/n \leq 1/N < \epsilon$ and $|t_n - c| < 1/n \leq 1/N < \epsilon$, the middle inequality by proposition 1.1.2(7).

Now $\mathbf{1}_{\mathbb{Q}}(p_n) = 1$ for every n and $\mathbf{1}_{\mathbb{Q}}(t_n) = 0$ for every n , so the two image sequences converge to 1 and to 0. At most one of these equals $\mathbf{1}_{\mathbb{Q}}(c)$, since $1 \neq 0$, so proposition 3.2.4 fails for one of the two sequences and $\mathbf{1}_{\mathbb{Q}}$ is discontinuous at c .

The restriction of $\mathbf{1}_{\mathbb{Q}}$ to $\mathbb{Q}$ is the constant 1 and its restriction to $\mathbb{R} \setminus \mathbb{Q}$ is the constant 0, so $\mathbf{1}_{\mathbb{Q}}$ is continuous on each of those two sets in the sense of definition 3.2.2 while being continuous at no point of $\mathbb{R}$. It is also bounded and not Riemann integrable, which example 5.2.3 proves once the integral exists.

Discontinuity everywhere is one extreme. The next function differs from $\mathbf{1}_{\mathbb{Q}}$ by a single factor of x and stops one point short of that extreme, being continuous at exactly one point and discontinuous at every other.

Example 3.2.11: The Dirichlet Function Multiplied by x

Define $u: \mathbb{R} \rightarrow \mathbb{R}$ by $u(x) = x \mathbf{1}_{\mathbb{Q}}(x)$, so that $u(x) = x$ for rational x and $u(x) = 0$ for irrational x . Then u is continuous at 0 and discontinuous at every other point.

Continuity at 0. For every $x \in \mathbb{R}$ the value $u(x)$ is either x or 0, so $|u(x)| \leq |x|$ by proposition 1.3.8(1). Since $u(0) = 0$, given $\epsilon > 0$ the choice $\delta = \epsilon$ gives $|u(x) - u(0)| = |u(x)| \leq |x| < \epsilon$ whenever $|x - 0| < \delta$.

Discontinuity elsewhere. Let $c \neq 0$ and take the sequences (p_n) of rationals and (t_n) of irrationals converging to c constructed in example 3.2.10. Then $u(p_n) = p_n \rightarrow c$ and $u(t_n) = 0$ for every n , so $u(t_n) \rightarrow 0$. Since $c \neq 0$, the two limits differ, so at most one equals $u(c)$ and proposition 3.2.4 fails for one of the two sequences. So u is discontinuous at c .

Both functions so far are discontinuous at all but at most one point, and neither settles whether a set whose complement is itself dense, such as $\mathbb{Q}$, can be exactly the set of discontinuities of a function. Thomae answered that in 1875 with the function below, whose value at a rational records how large a denominator that rational needs. The argument proves a single statement, that the limit of t is 0 at every real number, and both the continuity and the discontinuity follow from it.

Example 3.2.12: Thomae's Function

Every rational number has exactly one representation p/q with $p \in \mathbb{Z}$, q a positive integer and p/q in lowest terms, which is among the facts about $\mathbb{Z}$ and $\mathbb{Q}$ taken as given in section 1.1. Thomae's function $t: \mathbb{R} \rightarrow \mathbb{R}$ is

$$t(x) = \begin{cases} 1/q & \text{if } x \in \mathbb{Q} \text{ and } x = p/q \text{ in lowest terms with } q \geq 1 \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

In particular $t(0) = 1$, since $0 = 0/1$ in lowest terms, and $t(50/100) = t(1/2) = 1/2$, the reduction to lowest terms being what makes the value depend on the number alone, whatever fraction is written for it. Then t is continuous at every irrational number and discontinuous at every rational number.

Step 1: The rationals within 1 of c whose reduced denominator is at most N . Fix $c \in \mathbb{R}$ and let $\epsilon > 0$. By theorem 1.4.1(3) choose $N \in \mathbb{N}$ with $0 < 1/N < \epsilon$, and set

$$F = \{p/q \mid q \in \{1, \dots, N\}, p \in \mathbb{Z}, \lfloor q(c-1) \rfloor \leq p \leq \lfloor q(c+1) \rfloor\}$$

the integer parts being those of proposition 1.4.3. For each of the N values of q the integer p is confined between two fixed integers, so only finitely many p occur, and F is a finite set of rational numbers.

Step 2: F catches every rational near c with a small denominator. Let $x \in \mathbb{Q}$ satisfy $|x - c| \leq 1$ and write $x = p/q$ in lowest terms with $1 \leq q \leq N$. By proposition 1.3.8(2) the hypothesis reads $c - 1 \leq x \leq c + 1$, and multiplying through by $q > 0$ using proposition 1.1.2(5) gives $q(c-1) \leq p \leq q(c+1)$. Now proposition 1.4.3 gives $\lfloor q(c-1) \rfloor \leq q(c-1)$, so $\lfloor q(c-1) \rfloor \leq p$. And if $p > \lfloor q(c+1) \rfloor$ then $p \geq \lfloor q(c+1) \rfloor + 1 > q(c+1)$, again by proposition 1.4.3, contradicting $p \leq q(c+1)$. So $p \leq \lfloor q(c+1) \rfloor$, and hence $x \in F$.

Step 3: The choice of δ . The set $F \setminus \{c\}$ is finite, and $|c - r| > 0$ for each of its elements r by proposition 1.3.8(1). Let δ be the least of the number 1 and the numbers $|c - r|$ for $r \in F \setminus \{c\}$, a minimum over a finite set of positive numbers, so $\delta > 0$.

Step 4: The estimate. Let $x \in \mathbb{R}$ with $0 < |x - c| < \delta$. If $x \notin \mathbb{Q}$ then $|t(x) - 0| = 0 < \epsilon$. If $x \in \mathbb{Q}$, write $x = p/q$ in lowest terms with $q \geq 1$, and suppose $q \leq N$. Since $|x - c| < \delta \leq 1$, Step 2 puts x in F , and $|x - c| > 0$ gives $x \neq c$, so $x \in F \setminus \{c\}$ and therefore $|c - x| \geq \delta$ by the choice of δ . That contradicts $|x - c| < \delta$. So $q > N$, and proposition 1.1.2(7) gives

$$|t(x) - 0| = \frac{1}{q} < \frac{1}{N} < \epsilon$$

Since $\epsilon > 0$ was arbitrary and c is a limit point of $\mathbb{R}$, definition 3.1.15 gives $\lim_{x \rightarrow c} t(x) = 0$.

Step 5: Continuity and its failure. Every $c \in \mathbb{R}$ is a limit point of $\mathbb{R}$, so proposition 3.2.3(1) makes t continuous at c exactly when $t(c) = \lim_{x \rightarrow c} t(x) = 0$. If $c \notin \mathbb{Q}$ then $t(c) = 0$ and t is continuous at c . If $c \in \mathbb{Q}$, write $c = p/q$ in lowest terms, so that $t(c) = 1/q > 0$ by proposition 1.1.2(7) and t is discontinuous at c .

So the set of discontinuities of a function can be all of $\mathbb{R}$, all of $\mathbb{R}$ with one point removed, or exactly $\mathbb{Q}$. Determining which subsets of $\mathbb{R}$ occur in general is beyond what has been built here. For one class of functions the constraint is severe and provable now, and reaching it takes a way of separating failures of continuity into kinds, since the three examples above fail in visibly different ways.

Definition 3.2.13: Discontinuities of the First and Second Kind

Let $a < b$ be real numbers, let $f: (a, b) \rightarrow \mathbb{R}$ and let $c \in (a, b)$ be a discontinuity of f . The discontinuity is *simple*, or *of the first kind*, if both one-sided limits $\lim_{x \rightarrow c^-} f(x)$ and $\lim_{x \rightarrow c^+} f(x)$ exist in $\mathbb{R}$, and *of the second kind* otherwise. A simple discontinuity is *removable* if the two one-sided limits are equal, and a *jump* if they are not.

The two names for a simple discontinuity describe the two ways it arises. If the one-sided limits agree,

then $\lim_{x \rightarrow c} f(x)$ exists by proposition 3.1.26, and the only obstruction to continuity is the value $f(c)$, which can be redefined to that common limit. If they disagree, no value assigned at c can help.

Each of the three named functions is now classified. Thomae's function has a removable discontinuity at every rational: example 3.2.12 gives $\lim_{x \rightarrow c} t(x) = 0$, so proposition 3.1.26 makes both one-sided limits exist and equal 0, while $t(c) > 0$. The Dirichlet function has a discontinuity of the second kind at every $c \in (a, b)$, since neither one-sided limit exists. For the right-hand side, apply theorem 1.4.4 and corollary 1.4.5 inside the interval with endpoints c and $\min\{b, c + 1/n\}$ to get a rational p_n and an irrational t_n in (c, b) converging to c . Their images are the constant sequences 1 and 0, so theorem 3.1.19, applied to the restriction of $\mathbf{1}_{\mathbb{Q}}$ to (c, b) as proposition 3.1.25 permits, rules out a right-hand limit. The left-hand side is the same argument inside the interval with endpoints $\max\{a, c - 1/n\}$ and c . A jump is easiest to exhibit directly: let $f: (-1, 1) \rightarrow \mathbb{R}$ be $f(x) = 0$ for $x < 0$ and $f(x) = 1$ for $x \geq 0$. Then $\lim_{x \rightarrow 0^-} f(x) = 0$ and $\lim_{x \rightarrow 0^+} f(x) = 1$, each because f is constant on the relevant side, so f has a jump at 0. It is continuous at every other point of $(-1, 1)$, since every such point has a neighbourhood inside $(-1, 0)$ or inside $(0, 1)$ on which f is constant. The arithmetic of $\mathbb{Q}$ is not the only source of a second-kind discontinuity. Oscillation produces one too, in the function $x \mapsto \sin(1/x)$ extended by the value 0 at 0, treated in Section 4.6 once $\sin$ is available.

Monotone functions are the class for which the second kind is impossible, and the reason is that a supremum can be produced whenever a value is needed. The definition applies to a function on any subset of $\mathbb{R}$, while the results that follow are stated on an open interval, where both sides of every point carry domain points.

Definition 3.2.14: [Real] Monotone Function

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$. Then f is

1. *increasing* if $f(x) \leq f(y)$ whenever $x, y \in E$ satisfy $x \leq y$, and *strictly increasing* if $f(x) < f(y)$ whenever $x, y \in E$ satisfy $x < y$;
2. *decreasing* if $f(x) \geq f(y)$ whenever $x, y \in E$ satisfy $x \leq y$, and *strictly decreasing* if $f(x) > f(y)$ whenever $x, y \in E$ satisfy $x < y$.

It is *monotone* if it is increasing or decreasing, and *strictly monotone* if it is strictly increasing or strictly decreasing.

The affine function $f(x) = 2x + 5$ on $\mathbb{R}$ is strictly increasing, since $x < y$ gives $2x < 2y$ by proposition 1.1.2(5) and then $2x + 5 < 2y + 5$ by the third ordered-field axiom of definition 1.1.1. The same two steps with the factor -2 make $x \mapsto -2x + 5$ strictly decreasing, using proposition 1.1.2(5) in its second form. A constant function is both increasing and decreasing and neither strictly. The Dirichlet function is neither on any interval containing more than one point, since theorem 1.4.4 and corollary 1.4.5 place a rational and an irrational between any two of its points. Definition 2.2.1 attached the same words to sequences. The two definitions agree when a sequence is read as a function on $\mathbb{N}$, and the one above is what the rest of this chapter uses.

Proposition 3.2.15: One-Sided Limits of an Increasing Function

Let $a < b$ be real numbers and let $f: (a, b) \rightarrow \mathbb{R}$ be increasing. Then for every $c \in (a, b)$ both one-sided limits of f at c exist in $\mathbb{R}$, and

$$\sup\{f(x) \mid a < x < c\} = \lim_{x \rightarrow c^-} f(x) \leq f(c) \leq \lim_{x \rightarrow c^+} f(x) = \inf\{f(x) \mid c < x < b\}$$

Moreover, if $a < c < d < b$, then $\lim_{x \rightarrow c^+} f(x) \leq \lim_{x \rightarrow d^-} f(x)$.

Proof:

Step 1: The left-hand limit. Let $A = \{f(x) \mid a < x < c\}$. This set is nonempty, since $a < c$ gives a point of (a, c) by proposition 1.1.2(8), and it is bounded above by $f(c)$, since $a < x < c$ gives $f(x) \leq f(c)$ by definition 3.2.14. By theorem 1.2.4 the number $L = \sup A$ exists, and $L \leq f(c)$ because $f(c)$ is an upper bound of A and L is the least one.

Let $\epsilon > 0$. By proposition 1.2.6 there is a point of A exceeding $L - \epsilon$, that is, an $x_0 \in (a, c)$ with $L - \epsilon < f(x_0)$. Put $\delta = c - x_0 > 0$. If $x \in (a, c)$ satisfies $|x - c| < \delta$, then $c - x < \delta = c - x_0$ gives $x_0 < x < c$, so $f(x_0) \leq f(x)$ by definition 3.2.14 and $f(x) \leq L$ because L is an upper bound of A . Combining, $L - \epsilon < f(x) \leq L$, which gives $-\epsilon < f(x) - L \leq 0$ and so $|f(x) - L| < \epsilon$ by proposition 1.3.8(2). Since c is a limit point of (a, c) , this is the condition of definition 3.1.24, and $\lim_{x \rightarrow c^-} f(x) = L$.

Step 2: The right-hand limit. Let $B = \{f(x) \mid c < x < b\}$, nonempty by proposition 1.1.2(8) and bounded below by $f(c)$, since $c < x < b$ gives $f(c) \leq f(x)$. By corollary 1.2.5 the number $M = \inf B$ exists, and $f(c) \leq M$ because $f(c)$ is a lower bound and M is the greatest one. Let $\epsilon > 0$. By lemma 1.3.1 there is an $x_1 \in (c, b)$ with $M \leq f(x_1) < M + \epsilon$. Put $\delta = x_1 - c > 0$. If $x \in (c, b)$ satisfies $|x - c| < \delta$, then $c < x < x_1$, so $M \leq f(x) \leq f(x_1) < M + \epsilon$ and $|f(x) - M| < \epsilon$ by proposition 1.3.8(2). So $\lim_{x \rightarrow c^+} f(x) = M$.

Step 3: Comparison across two points. Let $a < c < d < b$. By proposition 1.1.2(8) there is a w with $c < w < d$. Then $w \in (c, b)$, so $f(w) \in B$ and $\lim_{x \rightarrow c^+} f(x) = \inf B \leq f(w)$. Also $w \in (a, d)$, so $f(w)$ belongs to the set whose supremum Step 1 identifies with $\lim_{x \rightarrow d^-} f(x)$, giving $f(w) \leq \lim_{x \rightarrow d^-} f(x)$. Chaining the two inequalities gives $\lim_{x \rightarrow c^+} f(x) \leq \lim_{x \rightarrow d^-} f(x)$, completing the proof.

Proposition 3.2.16: Limit of a Monotone Function at an Endpoint

Let $a < b$ be real numbers and let $g: (a, b] \rightarrow \mathbb{R}$ be decreasing. If g is bounded above, then

$$\lim_{t \rightarrow a^+} g(t) = \sup\{g(t) \mid a < t \leq b\}$$

and that supremum is finite. If g is not bounded above, then $\lim_{t \rightarrow a^+} g(t) = +\infty$. The increasing case, and the case of the right-hand endpoint, follow by applying this to $-g$ and to $t \mapsto g(a + b - t)$.

Proof:

Suppose g is bounded above and put $S = \sup\{g(t) \mid a < t \leq b\}$, which exists by theorem 1.2.4. Let $\epsilon > 0$. By proposition 1.2.6 there is a $t_0 \in (a, b]$ with $g(t_0) > S - \epsilon$. Put $\delta = t_0 - a > 0$. For $t \in (a, b]$ with $0 < t - a < \delta$ we have $t < t_0$, so the decrease of g gives $S - \epsilon < g(t_0) \leq g(t) \leq S$ and hence $|g(t) - S| < \epsilon$. That is the condition of definition 3.1.24 for the right-hand limit at a .

If instead g is not bounded above, let $M \in \mathbb{R}$ and take $t_0 \in (a, b]$ with $g(t_0) > M$. Every $t \in (a, t_0)$ has $g(t) \geq g(t_0) > M$, so $\delta = t_0 - a$ meets the one-sided condition of definition 3.1.28.

The hypothesis that g is defined on $(a, b]$ and not at a is what separates this from proposition 3.2.15, whose proof bounds the relevant set by the value of the function at the point in question.

Applying the proposition to $-f$, which is increasing exactly when f is decreasing, and using corollary 3.1.20 to move the sign outside each one-sided limit, gives the decreasing case with every inequality reversed. So a monotone function has both one-sided limits at every interior point of its interval, and by definition 3.2.13 it has no discontinuity of the second kind. The gap between the two one-sided limits is the whole of the failure, and the next theorem bounds how often that gap can be nonzero.

Theorem 3.2.17: The Discontinuity Set of a Monotone Function

Let $a < b$ be real numbers and let $f: (a, b) \rightarrow \mathbb{R}$ be monotone. Then the set of points of (a, b) at which f is discontinuous is finite or countable.

Proof:

Step 1: Reduction to the increasing case. If f is decreasing then $-f$ is increasing, and f is continuous at a point exactly when $-f$ is, by theorem 3.2.5(2) applied with $\lambda = -1$ in both directions. So the two functions have the same set of discontinuities and it is enough to treat an increasing f . Write D for that set.

Step 2: A discontinuity is a strict gap. Let $c \in (a, b)$ and write $L(c) = \lim_{x \rightarrow c^-} f(x)$ and $M(c) = \lim_{x \rightarrow c^+} f(x)$, which exist by proposition 3.2.15 and satisfy $L(c) \leq f(c) \leq M(c)$. Suppose $L(c) = M(c)$. Then both numbers equal $f(c)$, and c is a limit point of (a, c) and of (c, b) , so proposition 3.1.26 gives $\lim_{x \rightarrow c} f(x) = f(c)$. Since c is a limit point of (a, b) , proposition 3.2.3(1) makes f continuous at c . Contrapositively, $c \in D$ forces $L(c) < M(c)$.

Step 3: An injection into $\mathbb{N}$. By [4, countable sets and the countability of $\mathbb{Q}$] the set $\mathbb{Q}$ is countable, so there is a function $r: \mathbb{N} \rightarrow \mathbb{Q}$ whose range is all of $\mathbb{Q}$. Let $c \in D$. Theorem 1.4.4 applied to $L(c) < M(c)$ gives a rational strictly between them, and that rational equals $r(n)$ for some n , so

$$S(c) = \{n \in \mathbb{N} \mid L(c) < r(n) < M(c)\}$$

is a nonempty set of positive integers. Let $n(c)$ be its least element, which exists by the well-ordering of $\mathbb{N}$ recorded in section 1.1. No choice is made, since $n(c)$ is determined by c .

Let $c_1, c_2 \in D$ with $c_1 \neq c_2$, say $c_1 < c_2$. The last clause of proposition 3.2.15 gives $M(c_1) \leq L(c_2)$, so

$$r(n(c_1)) < M(c_1) \leq L(c_2) < r(n(c_2))$$

and in particular $r(n(c_1)) \neq r(n(c_2))$, which forces $n(c_1) \neq n(c_2)$. So $c \mapsto n(c)$ is an injection of D into $\mathbb{N}$.

Step 4: Conclusion. The injection identifies D with a subset of $\mathbb{N}$, and $\mathbb{N}$ is countable, being the range of the identity function on $\mathbb{N}$. A subset of a finite or countable set is finite or countable, by [4, countable sets], so D is finite or countable, completing the proof.

The bound is exact. Every nonempty finite or countable subset of an interval is realized as a set of discontinuities, by a function built from a convergent series of positive numbers whose n -th term is deposited at the n -th point of the set.

Example 3.2.18: A Monotone Function with Prescribed Discontinuities

Let $a < b$ be real and let $S \subseteq (a, b)$ be nonempty, finite or countable. Then there is an increasing $f: (a, b) \rightarrow \mathbb{R}$ whose set of discontinuities is exactly S .

The construction. By [4, countable sets] there is a function $\sigma: \mathbb{N} \rightarrow S$ whose range is all of S . For $n \in \mathbb{N}$ define $u_n: (a, b) \rightarrow \mathbb{R}$ by $u_n(x) = 1$ if $\sigma(n) < x$ and $u_n(x) = 0$ otherwise, and set

$$f(x) = \sum_{n=1}^{\infty} 2^{-n} u_n(x)$$

The terms are nonnegative, and the partial sums satisfy $\sum_{n=1}^M 2^{-n} u_n(x) \leq \sum_{n=1}^M 2^{-n} = 1 - 2^{-M} \leq 1$, the middle equality by summing the identity $2^{-n} = 2^{-(n-1)} - 2^{-n}$ from $n = 1$ to M and cancelling as in example 2.4.6. So the partial sums are bounded above, proposition 2.4.4 makes the series convergent, and f is defined on all of (a, b) .

f is increasing. Let $x \leq y$ in (a, b) and let $n \in \mathbb{N}$. If $u_n(x) = 1$ then $\sigma(n) < x \leq y$, so $u_n(y) = 1$. Hence $u_n(x) \leq u_n(y)$ for every n , so the partial sums of the two series compare term by term, and proposition 2.1.8(1) gives $f(x) \leq f(y)$.

f is discontinuous at every point of S . Let $s \in S$ and choose $m \in \mathbb{N}$ with $\sigma(m) = s$, which exists because σ has range S . Let $x \in (a, b)$ with $x > s$. For every n , if $u_n(s) = 1$ then $\sigma(n) < s < x$ and $u_n(x) = 1$, so $u_n(x) - u_n(s) \geq 0$. Also $u_m(x) - u_m(s) = 1 - 0 = 1$, since $\sigma(m) = s < x$ while $\sigma(m) = s$ is not less than s . By proposition 2.4.5(1), applied with the scalar -1 and then with the sum, the series $\sum_{n \geq 1} 2^{-n} (u_n(x) - u_n(s))$ converges with sum $f(x) - f(s)$, and each of its partial sums with $M \geq m$ is at least 2^{-m} , because every term is nonnegative and the m -th equals 2^{-m} . By proposition 2.1.8(2), $f(x) - f(s) \geq 2^{-m}$.

Now take $\epsilon_0 = 2^{-m} > 0$ and let $\delta > 0$ be arbitrary. Since $s < b$ and $s < s + \delta$, proposition 1.1.2(8) gives a point x strictly between s and $\min\{b, s + \delta\}$, so $x \in (a, b)$ and $|x - s| < \delta$ while $|f(x) - f(s)| \geq \epsilon_0$. So no δ answers ϵ_0 in definition 3.2.1, and f is discontinuous at s .

f is continuous at every other point. Let $c \in (a, b)$ with $c \notin S$ and let $\epsilon > 0$. By proposi-

tion 2.1.10(3), applied with $j = 0$ and $r = 1/2$, the sequence $(2^{-k})_{k \geq 1}$ is null, so there is an $N \in \mathbb{N}$ with $2^{-N} < \epsilon$. For each $n \in \{1, \dots, N\}$ the point $\sigma(n)$ lies in S and so differs from c , whence $|c - \sigma(n)| > 0$ by proposition 1.3.8(1). Let δ be the least of those N numbers, a minimum over a finite set of positive numbers, so $\delta > 0$.

Let $x \in (a, b)$ with $|x - c| < \delta$ and let $n \leq N$. If $\sigma(n) < c$ then $c - \sigma(n) \geq \delta$, so $\sigma(n) \leq c - \delta < x$ and $u_n(x) = 1 = u_n(c)$. If $\sigma(n) > c$ then $\sigma(n) - c \geq \delta$, so $\sigma(n) \geq c + \delta > x$ and $u_n(x) = 0 = u_n(c)$. The case $\sigma(n) = c$ does not arise. So the first N terms of the series for $f(x) - f(c)$ vanish, and for every $M > N$, using proposition 2.4.5(2) and $|u_n(x) - u_n(c)| \leq 1$,

$$\left| \sum_{n=1}^M 2^{-n} (u_n(x) - u_n(c)) \right| = \left| \sum_{n=N+1}^M 2^{-n} (u_n(x) - u_n(c)) \right| \leq \sum_{n=N+1}^M 2^{-n} = 2^{-N} - 2^{-M} \leq 2^{-N}$$

the last equality again by summing $2^{-n} = 2^{-(n-1)} - 2^{-n}$, now from $n = N+1$ to M . These partial sums converge to $f(x) - f(c)$ by proposition 2.4.5(1), and every one of them lies between -2^{-N} and 2^{-N} , so proposition 2.1.8(2) gives $-2^{-N} \leq f(x) - f(c) \leq 2^{-N}$ and hence $|f(x) - f(c)| \leq 2^{-N} < \epsilon$ by proposition 1.3.8(2). So f is continuous at c .

Taking $(a, b) = (0, 1)$ and $S = \mathbb{Q} \cap (0, 1)$ produces an increasing function on $(0, 1)$ whose discontinuities are exactly the rationals, the same set that example 3.2.12 produced by an entirely different construction.

Exercise 3.2.1

1. Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ be continuous at $c \in E$ and suppose $f(c) > 0$. Show that there is a $\delta > 0$ such that $f(x) > f(c)/2$ for every $x \in E$ with $|x - c| < \delta$. State and prove the corresponding statement when $f(c) < 0$.
2. Let I be an interval containing more than one point and let $f, g: I \rightarrow \mathbb{R}$ be continuous with $f(x) = g(x)$ for every $x \in I \cap \mathbb{Q}$. Show that $f = g$. Where does the argument use that I contains more than one point?
3. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfy $f(x + y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$. Show that if f is continuous at 0, then f is continuous at every point of $\mathbb{R}$.
4. Let t be Thomae's function of example 3.2.12. Show that t is monotone on no interval containing more than one point.
5. Give a continuous $f: \mathbb{R} \rightarrow \mathbb{R}$ and a discontinuous $g: \mathbb{R} \rightarrow \mathbb{R}$ with $g \circ f$ continuous at every point of $\mathbb{R}$, and a discontinuous f and a continuous g with the same property.
6. Let $f, g: \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Show that $\{x \in \mathbb{R} \mid f(x) = g(x)\}$ is a closed subset of $\mathbb{R}$, and deduce that a continuous function agreeing with a polynomial at every rational number agrees with it at every real number.
7. Let $a < b$ be real, let $f: (a, b) \rightarrow \mathbb{R}$ be increasing and for $c \in (a, b)$ write $J(c) = \lim_{x \rightarrow c^+} f(x) - \lim_{x \rightarrow c^-} f(x)$. Show that $J(c) \geq 0$, and that for any points $u < c_1 < c_2 < \dots < c_k < v$ of

(a, b) ,

$$\sum_{i=1}^k J(c_i) \leq f(v) - f(u)$$

8. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = \lfloor x \rfloor$, the integer part of proposition 1.4.3. Show that f is increasing and continuous at every $c \notin \mathbb{Z}$. Fix $m \in \mathbb{Z}$ and restrict f to $(m-1, m+1)$, which puts it in the setting of definition 3.2.13; show that the restriction has a jump discontinuity at m , computing both one-sided limits there.

3.3 Continuous Functions on a Closed Bounded Interval

Section 3.1 and section 3.2 placed no condition on the domain: a continuous function was $f: E \rightarrow \mathbb{R}$ for an arbitrary $E \subseteq \mathbb{R}$, and the algebra and composition rules hold at that generality. This section imposes conditions on E and records what each one gives. That E is an interval produces the intermediate value theorem. That E is a closed bounded interval produces the boundedness of f , the attainment of its bounds, and the description of its image. Adding strict monotonicity to the first of these produces a continuous inverse.

Every hypothesis below constrains the domain, so the first thing to record is that shrinking a domain costs nothing.

Proposition 3.3.1: Restriction of a Continuous Function

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ be continuous and let $D \subseteq E$. Then the restriction of f to D is continuous.

Proof:

Let $x \in D$ and $\epsilon > 0$. The continuity of f at x gives a $\delta > 0$ such that every $t \in E$ with $|t - x| < \delta$ satisfies $|f(t) - f(x)| < \epsilon$ (definition 3.2.1). The points $t \in D$ with $|t - x| < \delta$ are among those t , so the same δ answers ϵ for the restriction at x .

In particular a polynomial function, regarded as a function on any subset of $\mathbb{R}$, is continuous, by proposition 3.3.1 together with example 3.2.6.

The intermediate value theorem fails over $\mathbb{Q}$, and the failure is concrete. Let $E_0 = \mathbb{Q} \cap [0, 2]$ and let $g: E_0 \rightarrow \mathbb{R}$ be $g(q) = q^2 - 2$. It is continuous, being a polynomial restricted to E_0 , and $g(0) = -2 < 0 < 2 = g(2)$, yet $g(q) = 0$ has no solution in E_0 because $q^2 = 2$ has no rational solution (proposition 1.1.3). What E_0 lacks is the betweenness condition of definition 3.1.1. The number $\sqrt{2}$ given by theorem 1.3.12 satisfies $1 < \sqrt{2} < 2$. It is positive, since $(\sqrt{2})^2 = 2 \neq 0$ forces $\sqrt{2} \neq 0$ while $\sqrt{2} \geq 0$. If $\sqrt{2} = 1$ then $2 = 1$, and if $\sqrt{2} < 1$ then multiplying by the positive number $\sqrt{2}$ gives $2 = (\sqrt{2})^2 < \sqrt{2} < 1$ (proposition 1.1.2(5)), so $\sqrt{2} > 1$. If $\sqrt{2} = 2$ then $2 = 4$, and if $\sqrt{2} > 2$ then the same multiplication gives $2 = (\sqrt{2})^2 > 2\sqrt{2}$ while $2\sqrt{2} > 4$, so $\sqrt{2} < 2$. Hence 1 and 2 lie in E_0 while the number $\sqrt{2}$ between them does not.

Theorem 3.3.2: The Intermediate Value Theorem

1. Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. If $y \in \mathbb{R}$ satisfies $\min\{f(a), f(b)\} < y < \max\{f(a), f(b)\}$, then $f(c) = y$ for some $c \in (a, b)$.
2. Let $I \subseteq \mathbb{R}$ be an interval, let $f: I \rightarrow \mathbb{R}$ be continuous, and let $u, v \in I$ with $u < v$. If $y \in \mathbb{R}$ satisfies $\min\{f(u), f(v)\} < y < \max\{f(u), f(v)\}$, then $f(c) = y$ for some $c \in (u, v)$.

Proof:

1. *Reduction to a sign change at 0.* If $f(a) < y < f(b)$, put $g = f - y$. If $f(b) < y < f(a)$, put $g = y - f$. In both cases $g: [a, b] \rightarrow \mathbb{R}$ is continuous by theorem 3.2.5, applied to f and to the constant function with value y , which is a polynomial restricted to $[a, b]$. In the first case $g(a) = f(a) - y < 0$ and $g(b) = f(b) - y > 0$; in the second, $g(a) = y - f(a) < 0$ and $g(b) = y - f(b) > 0$. In both cases $g(c) = 0$ holds exactly when $f(c) = y$. So it suffices to produce a $c \in (a, b)$ with $g(c) = 0$, given a continuous $g: [a, b] \rightarrow \mathbb{R}$ with $g(a) < 0 < g(b)$.

The supremum. Let

$$S = \{x \in [a, b] \mid g(x) < 0\}$$

Then $a \in S$, so S is nonempty, and b is an upper bound of S . By theorem 1.2.4 the number $c = \sup S$ exists, and $a \leq c \leq b$, since $a \in S$ forces $a \leq c$ and b is an upper bound.

Step 1: $a < c$. Apply definition 3.2.1 at a with $\epsilon = -g(a) > 0$: there is a $\delta_1 > 0$ such that every $x \in [a, b]$ with $|x - a| < \delta_1$ satisfies $|g(x) - g(a)| < -g(a)$, hence $g(a) < g(x) - g(a) < -g(a)$ by proposition 1.3.8(2), hence $g(x) < 0$. Put $x_0 = \min\{b, a + \delta_1/2\}$. Then $a < x_0 \leq b$, because $a < b$ and $a < a + \delta_1/2$, and $|x_0 - a| = x_0 - a \leq \delta_1/2 < \delta_1$. So $g(x_0) < 0$, that is $x_0 \in S$, and $c \geq x_0 > a$.

Step 2: $c < b$. Apply definition 3.2.1 at b with $\epsilon = g(b) > 0$: there is a $\delta_2 > 0$ such that every $x \in [a, b]$ with $|x - b| < \delta_2$ satisfies $|g(x) - g(b)| < g(b)$, hence $g(x) > 0$ by proposition 1.3.8(2). Put $x_1 = \max\{a, b - \delta_2/2\}$, so that $a \leq x_1 < b$. Every $x \in [a, b]$ with $x > x_1$ satisfies $b - x < b - (b - \delta_2/2) = \delta_2/2 < \delta_2$, so $g(x) > 0$ and $x \notin S$. Every element of S is therefore at most x_1 , and $c \leq x_1 < b$.

Step 3: $g(c) \leq 0$. Suppose $g(c) > 0$. Apply definition 3.2.1 at c with $\epsilon = g(c)$: there is a $\delta > 0$ such that every $x \in [a, b]$ with $|x - c| < \delta$ satisfies $g(x) > 0$, again by proposition 1.3.8(2). Put $t = \max\{a, c - \delta/2\}$. Then $t < c$, since $a < c$ by Step 1 and $c - \delta/2 < c$. If $x \in S$ satisfied $x > t$, then $x \leq c$ and $c - x < c - (c - \delta/2) = \delta/2 < \delta$, so $|x - c| < \delta$ and $g(x) > 0$, contradicting $x \in S$. So t is an upper bound of S strictly smaller than c , which contradicts the definition of c as the least upper bound (definition 1.2.2).

Step 4: $g(c) \geq 0$. Suppose $g(c) < 0$. Apply definition 3.2.1 at c with $\epsilon = -g(c)$: there is a $\delta > 0$ such that every $x \in [a, b]$ with $|x - c| < \delta$ satisfies $g(x) < 0$. Put $x_2 = \min\{b, c + \delta/2\}$. By Step 2, $c < b$, so $c < x_2 \leq b$, and $|x_2 - c| = x_2 - c \leq \delta/2 < \delta$. Hence $g(x_2) < 0$, that is $x_2 \in S$, and $x_2 > c = \sup S$ contradicts that c is an upper bound.

Steps 3 and 4 give $g(c) = 0$, and Steps 1 and 2 give $c \in (a, b)$.

2. Since I is an interval and $u, v \in I$ with $u < v$, every z with $u < z < v$ lies in I (definition 3.1.1), so $[u, v] \subseteq I$. The restriction of a continuous function to a subset of its domain is continuous, so f restricted to $[u, v]$ is continuous. Part (1) applied to that restriction on $[u, v]$ produces the required $c \in (u, v)$, completing the proof.

The theorem asserts existence and nothing else. It does not say how many solutions $f(c) = y$ has, and it gives no procedure for finding one. The two hypotheses enter at separate places. Continuity is what the four applications of definition 3.2.1 in Steps 1 to 4 use, and completeness is what the single line producing $c = \sup S$ uses. Over $\mathbb{Q}$ the second is unavailable, and the function $g(q) = q^2 - 2$ on $\mathbb{Q} \cap [0, 2]$ is what its absence looks like.

One immediate application settles a question about polynomials that the algebra of section 3.2 cannot reach.

Corollary 3.3.3: Odd-Degree Polynomials Have Real Roots

Let $n \geq 1$ be an odd integer, let $c_0, \dots, c_n \in \mathbb{R}$ with $c_n \neq 0$, and let

$$p(x) = c_n x^n + c_{n-1} x^{n-1} + \dots + c_1 x + c_0$$

Then $p(c) = 0$ for some $c \in \mathbb{R}$.

Proof:

The polynomial $-p$ has the same roots as p and leading coefficient $-c_n$, so it may be assumed that $c_n > 0$.

Step 1: a radius beyond which the leading term dominates. Put $A = |c_0| + \dots + |c_{n-1}|$ and $M = 1 + A/c_n$, so that $M \geq 1$ and $c_n M = c_n + A$. For every integer k with $0 \leq k \leq n-1$,

$$M^k \leq M^{n-1}$$

To see it, note that $M \geq 1 > 0$ makes every power M^j positive (proposition 1.1.2(3) and (5)), and multiplying $1 \leq M$ by M^j gives $M^j \leq M^{j+1}$, with equality when $M = 1$ and strict inequality by proposition 1.1.2(5) when $M > 1$. These inequalities chain from k up to $n-1$. Hence, using $|c_k M^k| = |c_k| M^k$ (proposition 1.3.8(3)) and applying proposition 1.3.8(4) $n-1$ times,

$$\left| \sum_{k=0}^{n-1} c_k M^k \right| \leq \sum_{k=0}^{n-1} |c_k| M^k \leq A M^{n-1}$$

The same estimate holds with M replaced by $-M$, since $|(-M)^k| = M^k$ by proposition 1.3.8(3) applied k times.

Step 2: the two sign evaluations. From $c_n M^n = (c_n M) M^{n-1} = (c_n + A) M^{n-1}$ and $M^{n-1} > 0$,

$$p(M) = c_n M^n + \sum_{k=0}^{n-1} c_k M^k \geq c_n M^n - A M^{n-1} = c_n M^{n-1} > 0$$

where the first inequality uses $s \geq -|s|$ (proposition 1.3.8(1)) with $s = \sum_{k < n} c_k M^k$. Since n is odd, $(-M)^n = -M^n$, so the same computation gives

$$p(-M) = -c_n M^n + \sum_{k=0}^{n-1} c_k (-M)^k \leq -c_n M^n + AM^{n-1} = -c_n M^{n-1} < 0$$

Step 3: the intermediate value theorem. The function p is continuous on $\mathbb{R}$ (example 3.2.6), the set $\mathbb{R}$ is an interval, and $-M < M$ because $M \geq 1 > 0$. Since $\min\{p(-M), p(M)\} = p(-M) < 0 < p(M) = \max\{p(-M), p(M)\}$, the value 0 lies strictly between $p(-M)$ and $p(M)$, and theorem 3.3.2(2) produces a $c \in (-M, M)$ with $p(c) = 0$, as we sought to show.

The hypothesis that n be odd is what makes the two evaluations have opposite signs, and it cannot be dropped: $p(x) = x^2 + 1$ satisfies $p(x) \geq 1$ for every real x , since $x^2 > 0$ for $x \neq 0$ by proposition 1.1.2(6) and $0 \cdot 0 = 0$ by proposition 1.1.2(3).

Theorem 3.3.2(2) says that a continuous function on an interval attains every value between any two of its values. That conclusion is a property a function may have on its own, without reference to continuity, and it is given a name because the two conditions are not equivalent.

Definition 3.3.4: Intermediate Value Property

Let $I \subseteq \mathbb{R}$ be an interval. A function $f: I \rightarrow \mathbb{R}$ has the *intermediate value property* if for all $u, v \in I$ with $u < v$ and every $y \in \mathbb{R}$ with $\min\{f(u), f(v)\} < y < \max\{f(u), f(v)\}$ there is a $c \in (u, v)$ with $f(c) = y$.

Every continuous function on an interval has the property, which is exactly theorem 3.3.2(2). The polynomial $p(x) = x^3 - x$ on $\mathbb{R}$ has it for that reason. Not every function does: define $f: [-1, 1] \rightarrow \mathbb{R}$ by $f(x) = -1$ for $x < 0$ and $f(x) = 1$ for $x \geq 0$. Then $f(-1) = -1$ and $f(1) = 1$, and 0 lies strictly between them, but the only values f takes are -1 and 1 , so no c satisfies $f(c) = 0$.

The converse also fails: a function can have the intermediate value property at every pair of points and still be discontinuous. The standard example is $x \mapsto \sin(1/x)$ on $[-1, 1]$, extended by the value 0 at the origin, which oscillates between -1 and 1 on every interval around 0 and therefore omits no intermediate value, while failing to have a limit at 0. Constructing $\sin$ and establishing both facts is the work of section 4.6. The intermediate value property is therefore strictly weaker than continuity, and theorem 3.3.2 cannot be reversed.

Theorem 3.3.5 and theorem 3.3.7 replace “interval” by “closed bounded interval”, and both are false without the strengthening. They also fail over $\mathbb{Q}$, and again it is the absence of $\sqrt{2}$ that breaks them. Let $h: E_0 \rightarrow \mathbb{R}$ be $h(q) = 1/(2 - q^2)$, again with $E_0 = \mathbb{Q} \cap [0, 2]$. It is defined at every point of E_0 , since $q^2 = 2$ has no rational solution (proposition 1.1.3), and it is continuous by theorem 3.2.5 applied to the polynomials 1 and $2 - q^2$, the second of which does not vanish on E_0 . It is unbounded. Given $n \in \mathbb{N}$, theorem 1.4.4 produces a rational q with $\sqrt{2} - 1/n < q < \sqrt{2}$. Then $q > 0$ because $\sqrt{2} - 1/n \geq \sqrt{2} - 1 > 0$, and $q < 2$ because $\sqrt{2} < 2$, so $q \in E_0$. Factoring,

$$0 < 2 - q^2 = (\sqrt{2} - q)(\sqrt{2} + q) < 4/n$$

using $0 < \sqrt{2} - q < 1/n$ and $0 < \sqrt{2} + q < 4$, so $h(q) > n/4$ by proposition 1.1.2(7). Since $\{n/4 \mid n \in \mathbb{N}\}$ has no upper bound in $\mathbb{R}$ (theorem 1.4.1(2) applied to $4M$ for a proposed bound M), neither does the set of values of h .

Theorem 3.3.5: A Continuous Function on a Closed Bounded Interval Is Bounded

Let $a \leq b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. Then there is an $M \in \mathbb{R}$ with $|f(x)| \leq M$ for every $x \in [a, b]$.

Proof:

Suppose no such M exists. Then for every $n \in \mathbb{N}$ the number n is not such an M , so there is an $x_n \in [a, b]$ with $|f(x_n)| > n$.

Step 1: a convergent subsequence. The sequence $(x_n)_{n \geq 1}$ satisfies $a \leq x_n \leq b$ for every n , so it is bounded. By theorem 2.2.9(1) it has a convergent subsequence $(x_{n_k})_{k \geq 1}$, say $x_{n_k} \rightarrow x$. Applying proposition 2.1.8(2) to the constant bounds $a \leq x_{n_k} \leq b$ gives $a \leq x \leq b$, so $x \in [a, b]$ and $f(x)$ is defined.

Step 2: the contradiction. By proposition 3.2.4, applied to f at the point $x \in [a, b]$ and to the sequence (x_{n_k}) in $[a, b]$ converging to x , the sequence $(f(x_{n_k}))_{k \geq 1}$ converges to $f(x)$. A convergent sequence is bounded (proposition 2.1.5), so there is a $C \in \mathbb{R}$ with $|f(x_{n_k})| \leq C$ for every k . Because $n_1 < n_2 < \dots$ in $\mathbb{N}$, an induction gives $n_k \geq k$ for every k : indeed $n_1 \geq 1$, and $n_{k+1} > n_k \geq k$ forces $n_{k+1} \geq k+1$, since no element of $\mathbb{N}$ lies strictly between k and $k+1$. Choose $k_0 \in \mathbb{N}$ with $k_0 > C$, which theorem 1.4.1(2) provides. Then

$$C \geq |f(x_{n_{k_0}})| > n_{k_0} \geq k_0 > C$$

which is impossible. So some M works, completing the proof.

Both halves of “closed and bounded” are used, and the proof shows where. That the interval is bounded is what makes (x_n) a bounded sequence, so that theorem 2.2.9 applies. That it is closed is what puts the limit x back inside the domain, so that $f(x)$ exists and proposition 3.2.4 can be invoked. Dropping either one breaks the argument, and the next example shows that each failure is real.

Example 3.3.6: Unbounded Continuous Functions on $(0, 1]$ and on $[0, \infty)$

1. Let $f: (0, 1] \rightarrow \mathbb{R}$ be $f(x) = 1/x$. It is continuous by theorem 3.2.5, applied to the polynomials 1 and x , the second of which does not vanish on $(0, 1]$. It is unbounded: given $M \in \mathbb{R}$, theorem 1.4.1(2) gives an $n \in \mathbb{N}$ with $n > M$, and $1/n \in (0, 1]$ has $f(1/n) = n > M$. The domain is a bounded interval, and it is not closed, since 0 is a limit point of $(0, 1]$ that does not belong to it. That 0 is a limit point is theorem 1.4.1(3): every $\delta > 0$ admits an n with $1/n < \delta$, and $1/n$ is a point of $(0, 1]$ other than 0 within δ of 0. So $(0, 1]$ is not closed by proposition 3.1.11.
2. Let $f: [0, \infty) \rightarrow \mathbb{R}$ be $f(x) = x$, continuous as a polynomial restricted to $[0, \infty)$. Given $M \in \mathbb{R}$, theorem 1.4.1(2) gives an $n \in \mathbb{N}$ with $n > M$, and $f(n) = n > M$, so f is unbounded. The domain is a closed interval and it is not bounded.

With boundedness in hand, the supremum of the set of values exists. The substantial question is whether it is attained, and the answer uses theorem 2.2.9 a second time.

Theorem 3.3.7: The Extreme Value Theorem

Let $a \leq b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. Then there are $p, q \in [a, b]$ with

$$f(p) \leq f(x) \leq f(q) \quad \text{for every } x \in [a, b]$$

Consequently the set $f([a, b]) = \{f(x) | x \in [a, b]\}$ has both a supremum and an infimum, and both belong to it: $\sup f([a, b]) = f(q)$ and $\inf f([a, b]) = f(p)$.

Proof:

Write $E = f([a, b])$.

Step 1: the supremum exists. The set E is nonempty, since $a \in [a, b]$. By theorem 3.3.5 there is an M with $|f(x)| \leq M$ for every $x \in [a, b]$, hence $f(x) \leq M$ by proposition 1.3.8(1), so M is an upper bound of E . By theorem 1.2.4 the number $s = \sup E$ exists.

Step 2: a sequence approaching the supremum. Let $n \in \mathbb{N}$. By proposition 1.2.6, applied with $\epsilon = 1/n$, there is an element of E exceeding $s - 1/n$, that is, there is an $x_n \in [a, b]$ with $s - 1/n < f(x_n)$. Since s is an upper bound of E ,

$$s - \frac{1}{n} < f(x_n) \leq s$$

Step 3: passing to a subsequence. The sequence (x_n) lies in $[a, b]$ and is therefore bounded, so theorem 2.2.9(1) gives a convergent subsequence $x_{n_k} \rightarrow q$, and $q \in [a, b]$ by proposition 2.1.8(2) applied to $a \leq x_{n_k} \leq b$. By proposition 3.2.4, $f(x_{n_k}) \rightarrow f(q)$.

Step 4: identifying the limit. The sequence $(1/n)_{n \geq 1}$ converges to 0 by example 2.1.3(1), so its subsequence $(1/n_k)_{k \geq 1}$ converges to 0 by proposition 2.2.7(1). The constant sequence with value s converges to s by example 2.1.3(2), so $s - 1/n_k \rightarrow s$ by theorem 2.1.6(1) and (2). The estimate of Step 2 reads $s - 1/n_k < f(x_{n_k}) \leq s$ for every k , so theorem 2.1.7, applied with the lower sequence $(s - 1/n_k)_{k \geq 1}$ and the upper sequence constantly equal to s , gives $f(x_{n_k}) \rightarrow s$. A sequence has at most one limit (proposition 2.1.4), so $f(q) = s$. Since s is an upper bound of E , this gives $f(x) \leq f(q)$ for every $x \in [a, b]$.

Step 5: the minimum. The function $-f$ is continuous on $[a, b]$ by theorem 3.2.5, so Steps 1 to 4 applied to it produce a $p \in [a, b]$ with $-f(x) \leq -f(p)$ for every $x \in [a, b]$. Each such inequality gives $f(p) \leq f(x)$: if $-f(x) = -f(p)$ then $f(x) = f(p)$, and if $-f(x) < -f(p)$ then $f(p) < f(x)$ by proposition 1.1.2(6).

Step 6: the two extrema are the supremum and the infimum. The number $f(q)$ is an upper bound of E and belongs to E , so any upper bound u of E satisfies $f(q) \leq u$, and $f(q) = \sup E$ by definition 1.2.2. The number $f(p)$ is a lower bound of E and belongs to E , so any lower bound ℓ satisfies $\ell \leq f(p)$, and $f(p) = \inf E$, completing the proof.

The content is the last clause: the supremum of the set of values is one of the values. For an arbitrary bounded set of reals the supremum need not belong to the set (example 1.2.3(2)), and the theorem says that a continuous function on a closed bounded interval never produces such a set. Both hypotheses on the domain are again needed, and this time the failure is subtler than in example 3.3.6, since the functions of example 3.3.8 are bounded.

Example 3.3.8: Bounded Continuous Functions That Do Not Attain a Bound

1. Let $f: (0, 1) \rightarrow \mathbb{R}$ be $f(x) = x$, continuous as a polynomial restricted to $(0, 1)$, with values in $(0, 1)$ and therefore bounded. Its set of values is $(0, 1)$, whose supremum is 1. Indeed 1 is an upper bound, and no $u < 1$ is one. If $u < 1/2$, then $1/2 \in (0, 1)$ exceeds u . If $1/2 \leq u$, then proposition 1.1.2(8) gives a v with $u < v < 1$, and $v > u \geq 1/2 > 0$ puts v in $(0, 1)$. Since $1 \notin (0, 1)$, no point of the domain has $f(x) = 1$, so the supremum is not attained. The same argument with the order reversed gives $\inf f((0, 1)) = 0$. The number 0 is a lower bound, and no $\ell > 0$ is one: if $\ell > 1/2$ then $1/2 \in (0, 1)$ is smaller than ℓ , and if $\ell \leq 1/2$ then proposition 1.1.2(8) applied to $0 < \ell$ gives a v with $0 < v < \ell \leq 1/2 < 1$, so $v \in (0, 1)$ is smaller than ℓ . Again $0 \notin (0, 1)$. The domain is a bounded interval that is not closed.
2. Let $f: [0, \infty) \rightarrow \mathbb{R}$ be $f(x) = 1/(1 + x^2)$. For every real x the number x^2 is nonnegative (proposition 1.1.2(3) and (6)), so $1 + x^2 \geq 1 > 0$ and f is continuous by theorem 3.2.5. Its values satisfy $0 < f(x) \leq 1$: for $x = 0$ the value is 1, and for $x \neq 0$ the inequality $0 < 1 < 1 + x^2$ and proposition 1.1.2(7) give $0 < f(x) < 1$. So f is bounded and $f(0) = 1$ is a maximum. There is no minimum. The number 0 is a lower bound that no value attains, and it is the infimum: given $\epsilon > 0$, theorem 1.4.1(3) gives an $n \in \mathbb{N}$ with $1/n < \epsilon$, and multiplying $1 \leq n$ by $n > 0$ gives $n \leq n^2 < 1 + n^2$, so proposition 1.1.2(7) gives $f(n) = 1/(1 + n^2) < 1/n < \epsilon$. The domain is a closed interval that is not bounded.

Theorem 3.3.2 and theorem 3.3.7 combine into a single statement about the image, and it is this packaged form that later chapters quote.

Corollary 3.3.9: The Continuous Image of a Closed Bounded Interval

Let $a \leq b$, let $f: [a, b] \rightarrow \mathbb{R}$ be continuous, and put $m = \inf f([a, b])$ and $M = \sup f([a, b])$, both of which exist by theorem 3.3.7. Then

$$f([a, b]) = [m, M]$$

In particular the image is again a closed bounded interval, degenerating to the single point $\{m\}$ when $m = M$.

Proof:

By theorem 3.3.7 there are $p, q \in [a, b]$ with $f(p) = m$, $f(q) = M$ and $m \leq f(x) \leq M$ for every $x \in [a, b]$. The last inequality is the inclusion $f([a, b]) \subseteq [m, M]$.

For the reverse inclusion let $y \in [m, M]$. If $y = m$ then $y = f(p)$, and if $y = M$ then $y = f(q)$, so assume $m < y < M$. Then $f(p) \neq f(q)$, so $p \neq q$. Put $u = \min\{p, q\}$ and $v = \max\{p, q\}$, so that $u < v$ and both lie in $[a, b]$. The two values $f(u)$ and $f(v)$ are m and M in one order or the other,

so $\min\{f(u), f(v)\} = m < y < M = \max\{f(u), f(v)\}$. Since $[a, b]$ is an interval (definition 3.1.1), theorem 3.3.2(2) gives a $c \in (u, v)$ with $f(c) = y$, and $c \in [a, b]$. Hence $[m, M] \subseteq f([a, b])$, as we sought to show.

The corollary is the reason the two theorems are usually quoted together: it converts a continuous function on $[a, b]$ into a surjection onto a closed bounded interval whose endpoints are values of the function. One consequence is a statement about self-maps that requires no further hypothesis at all.

Theorem 3.3.10: Fixed Points of a Continuous Self-Map of $[0, 1]$

Let $f: [0, 1] \rightarrow [0, 1]$ be continuous. Then $f(c) = c$ for some $c \in [0, 1]$.

Proof:

Let $g: [0, 1] \rightarrow \mathbb{R}$ be $g(x) = f(x) - x$. It is continuous by theorem 3.2.5, applied to f and to the polynomial x restricted to $[0, 1]$. Since f takes values in $[0, 1]$, its values satisfy $0 \leq f(x) \leq 1$, so

$$g(0) = f(0) \geq 0 \quad \text{and} \quad g(1) = f(1) - 1 \leq 0$$

If $g(0) = 0$ then $f(0) = 0$ and $c = 0$ works. If $g(1) = 0$ then $f(1) = 1$ and $c = 1$ works. Otherwise $g(1) < 0 < g(0)$, so $\min\{g(0), g(1)\} = g(1) < 0 < g(0) = \max\{g(0), g(1)\}$, and theorem 3.3.2(1) gives a $c \in (0, 1)$ with $g(c) = 0$, that is $f(c) = c$. In every case a fixed point exists, completing the proof.

The hypothesis that f map $[0, 1]$ into itself is what forces the two sign conditions, and dropping it destroys the conclusion: $f(x) = x + 1$ is continuous on $[0, 1]$ and $f(c) = c$ would give $1 = 0$.

Theorem 3.3.12 concerns inverses. A function is invertible onto its image exactly when it is injective, and a strictly monotone function (definition 3.2.14) is injective. The question is whether the inverse of a continuous strictly monotone function on an interval is again continuous. It is, and the interval hypothesis carries most of the argument: the continuity of f is used only to see that the image is again an interval.

Proposition 3.3.11: Powers Are Strictly Increasing on the Nonnegative Reals

Let $n \geq 1$ be an integer. If $0 \leq s < t$ then $s^n < t^n$. So $x \mapsto x^n$ is strictly increasing on $[0, \infty)$ in the sense of definition 3.2.14.

Proof:

Induction on n . For $n = 1$ the claim is the hypothesis $s < t$. Suppose it holds for n , and let $0 \leq s < t$, so that $t > 0$. Every power t^j is then positive, by proposition 1.1.2(3) and (5) applied repeatedly to $0 < t$. If $s = 0$ then $s^{n+1} = 0 < t^{n+1}$. If $s > 0$, then multiplying $s^n < t^n$ by $s > 0$ gives $s^{n+1} < st^n$, and multiplying $s < t$ by $t^n > 0$ gives $st^n < t^{n+1}$, both by proposition 1.1.2(5). Transitivity of the order gives $s^{n+1} < t^{n+1}$, completing the proof.

Theorem 3.3.12: Continuous Strictly Monotone Functions Have Continuous Inverses

Let $I \subseteq \mathbb{R}$ be an interval and let $f: I \rightarrow \mathbb{R}$ be continuous and strictly monotone (definition 3.2.14). Put $J = f(I) = \{f(x) | x \in I\}$.

1. J is an interval.
2. The map $f: I \rightarrow J$ is a bijection.
3. The inverse $f^{-1}: J \rightarrow I$ is strictly monotone in the same direction as f , and it is continuous.

Proof:

Assume first that f is strictly increasing. The strictly decreasing case is reduced to this one at the end.

1. Let $y_1, y_2 \in J$ and let $z \in \mathbb{R}$ satisfy $y_1 < z < y_2$. Write $y_1 = f(u_1)$ and $y_2 = f(u_2)$ with $u_1, u_2 \in I$. Then $u_1 < u_2$: the case $u_1 = u_2$ would give $y_1 = y_2$, and the case $u_1 > u_2$ would give $f(u_1) > f(u_2)$ by strict increase, both contradicting $y_1 < y_2$. Now $\min\{f(u_1), f(u_2)\} = y_1 < z < y_2 = \max\{f(u_1), f(u_2)\}$, so theorem 3.3.2(2), applied to the interval I and the points $u_1 < u_2$, gives a $c \in (u_1, u_2)$ with $f(c) = z$. Hence $z \in J$. This is the betweenness condition of definition 3.1.1, so J is an interval.
2. The map $f: I \rightarrow J$ is surjective by the definition of J . It is injective: if $u \neq v$ in I , say $u < v$, then $f(u) < f(v)$ by strict increase, so $f(u) \neq f(v)$.
3. Write $g = f^{-1}: J \rightarrow I$, which exists by part (2).

Step 1: g is strictly increasing. Let $y_1 < y_2$ in J and put $u_i = g(y_i)$, so $f(u_i) = y_i$. The argument in part (1) gives $u_1 < u_2$, which is $g(y_1) < g(y_2)$.

Step 2: a one-sided approximation inside I . Fix $y_0 \in J$, put $u_0 = g(y_0) \in I$, and let $\epsilon > 0$. Claim: if I contains a point greater than u_0 , then it contains a point u_+ with $u_0 < u_+ \leq u_0 + \epsilon$. Indeed, let $w \in I$ with $w > u_0$. If $w \leq u_0 + \epsilon$, take $u_+ = w$. Otherwise $u_0 < u_0 + \epsilon < w$ with $u_0, w \in I$, so $u_0 + \epsilon \in I$ by definition 3.1.1, and $u_+ = u_0 + \epsilon$ serves. Symmetrically, if I contains a point smaller than u_0 , it contains a point u_- with $u_0 - \epsilon \leq u_- < u_0$.

Step 3: the choice of δ . Strict increase gives $f(u_-) < f(u_0) = y_0 < f(u_+)$ whenever the relevant point exists. Set

$$\delta = \begin{cases} \min\{f(u_+) - y_0, y_0 - f(u_-)\} & \text{if both } u_+ \text{ and } u_- \text{ exist} \\ f(u_+) - y_0 & \text{if only } u_+ \text{ exists} \\ y_0 - f(u_-) & \text{if only } u_- \text{ exists} \\ 1 & \text{if neither exists} \end{cases}$$

In each case $\delta > 0$.

Step 4: δ works. Let $y \in J$ with $|y - y_0| < \delta$ and put $u = g(y)$. If $u = u_0$ then $|g(y) - g(y_0)| = 0 < \epsilon$. Suppose $u > u_0$. Then I contains a point greater than u_0 , namely u , so u_+ exists and $\delta \leq f(u_+) - y_0$. From $|y - y_0| < \delta$ and proposition 1.3.8(2), $y < y_0 + \delta \leq f(u_+)$. If $u \geq u_+$ held, then $f(u) \geq f(u_+)$, since $u = u_+$ gives equality and $u > u_+$ gives strict inequality, so $y \geq f(u_+)$, which is false. Hence $u < u_+ \leq u_0 + \epsilon$, and $|g(y) - g(y_0)| = u - u_0 < \epsilon$. The case $u < u_0$ is the mirror image: u_- exists, $\delta \leq y_0 - f(u_-)$, $y > y_0 - \delta \geq f(u_-)$ forces $u > u_- \geq u_0 - \epsilon$, and $|g(y) - g(y_0)| = u_0 - u < \epsilon$. In all three cases $|g(y) - g(y_0)| < \epsilon$, which is the condition of definition 3.2.1 for g at y_0 .

The strictly decreasing case. Let f be continuous and strictly decreasing on I and put $h = -f$, which is continuous by theorem 3.2.5 and strictly increasing, since $u < v$ gives $f(u) > f(v)$ and hence $-f(u) < -f(v)$ by proposition 1.1.2(6). By the case already proved, $K = h(I)$ is an interval, $h: I \rightarrow K$ is a bijection, and $h^{-1}: K \rightarrow I$ is continuous and strictly increasing. Now $J = f(I) = \{-z \mid z \in K\}$. If $y_1 < z < y_2$ with $y_1, y_2 \in J$, then $-y_2 < -z < -y_1$ with $-y_1, -y_2 \in K$ (proposition 1.1.2(6)), so $-z \in K$ and $z \in J$, and J is an interval. The map $f: I \rightarrow J$ is a bijection, being surjective by the definition of J and injective because $u < v$ gives $f(u) > f(v)$. For $y \in J$ the element $x = h^{-1}(-y) \in I$ satisfies $-f(x) = -y$, so $f(x) = y$ and $f^{-1}(y) = h^{-1}(-y)$. The map $\nu(y) = -y$ is a polynomial, hence continuous on J , and it carries J into K , so $f^{-1} = h^{-1} \circ \nu$ is continuous by theorem 3.2.7. Finally $y_1 < y_2$ in J gives $-y_1 > -y_2$, hence $h^{-1}(-y_1) > h^{-1}(-y_2)$, that is $f^{-1}(y_1) > f^{-1}(y_2)$, so f^{-1} is strictly decreasing, completing the proof.

The interval hypothesis is used twice. Part (1) uses it through theorem 3.3.2(2), and its conclusion fails without it: the identity map on $[0, 1] \cup [2, 3]$ is continuous and strictly increasing, and its image is that same set, which is not an interval (example 3.1.2(3)). Step 2 of part (3) uses it again, and that step is where the continuity of the inverse would break. On a domain that is not an interval, the points above u_0 can be separated from u_0 by a gap, so no u_+ within ϵ of u_0 need exist. A strictly increasing continuous bijection from a set that is not an interval onto an interval can then have a discontinuous inverse, and exercise 3.3.1 (7) exhibits one. Continuity of f , by contrast, entered only in part (1), through theorem 3.3.2: parts (2) and (3) hold for any strictly monotone f on an interval, with no continuity assumed.

The immediate application extends the catalogue of continuous functions of section 3.2, which stopped at polynomials, rational functions and $|x|$.

Corollary 3.3.13: The n -th Root Function Is Continuous

Let $n \geq 1$ be an integer, and for $x \geq 0$ write $x^{1/n}$ for the unique $y \geq 0$ with $y^n = x$ furnished by theorem 1.3.12. The function $r: [0, \infty) \rightarrow \mathbb{R}$ given by $r(x) = x^{1/n}$ is a strictly increasing continuous bijection from $[0, \infty)$ onto $[0, \infty)$.

Proof:

Let $p: [0, \infty) \rightarrow \mathbb{R}$ be $p(x) = x^n$. It is continuous, being a polynomial restricted to $[0, \infty)$, and strictly increasing there by proposition 3.3.11. The set $[0, \infty)$ is an interval.

Its image is $[0, \infty)$. Every value $p(y) = y^n$ with $y \geq 0$ is nonnegative: for $y = 0$ it is 0, and

for $y > 0$ repeated use of proposition 1.1.2(3) and (5) gives $y^n > 0$. Conversely, given $x \geq 0$, theorem 1.3.12 produces exactly one $y \geq 0$ with $y^n = x$, namely $y = x^{1/n}$, so $x = p(x^{1/n})$ lies in the image.

By theorem 3.3.12, applied to p on the interval $[0, \infty)$, the map $p: [0, \infty) \rightarrow [0, \infty)$ is a bijection and its inverse is continuous and strictly increasing. That inverse sends x to the unique $y \geq 0$ with $y^n = x$, which is $x^{1/n}$ by the uniqueness clause of theorem 1.3.12. So $r = p^{-1}$ has the stated properties, as we sought to show.

Combined with theorem 3.2.5 and theorem 3.2.7, this shows that any function built from polynomials, quotients with nonvanishing denominators and n -th roots is continuous on the set where those operations are defined. The function $x \mapsto (1 + x^2)^{1/2}$ on $\mathbb{R}$ is continuous for this reason: it is r with $n = 2$ composed with the polynomial $1 + x^2$, whose values lie in $[1, \infty) \subseteq [0, \infty)$.

Exercise 3.3.1

1. Let $f: [0, 1] \rightarrow \mathbb{R}$ be continuous and suppose that $f(x) \in \mathbb{Q}$ for every $x \in [0, 1]$. Show that f is constant.
2. Let $f: [0, 2] \rightarrow \mathbb{R}$ be continuous with $f(0) = f(2)$. Show that there is a $c \in [0, 1]$ with $f(c) = f(c + 1)$.
3. Show that the polynomial $p(x) = x^5 - 3x + 1$ has at least three distinct real roots, and name an interval of length at most 2 containing each of them.
4. Let $a \leq b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous with $f(x) > 0$ for every $x \in [a, b]$. Show that there is an $\eta > 0$ with $f(x) \geq \eta$ for every $x \in [a, b]$. Then give a continuous $g: (0, 1] \rightarrow \mathbb{R}$ with $g(x) > 0$ for every x and $\inf g((0, 1]) = 0$, and say which hypothesis of theorem 3.3.7 the domain $(0, 1]$ fails.
5. Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous and injective. Show that f is strictly monotone. (Begin by showing that for $u < v < w$ in $[a, b]$ the value $f(v)$ lies strictly between $f(u)$ and $f(w)$, using theorem 3.3.2 on $[u, v]$ and on $[v, w]$ to contradict injectivity otherwise.)
6. Prove that $x \mapsto x^{1/2}$ is continuous on $[0, \infty)$ directly from definition 3.2.1, without appealing to theorem 3.3.12, by factoring $s - t = (s^{1/2} - t^{1/2})(s^{1/2} + t^{1/2})$. Treat $t > 0$ and $t = 0$ separately, and say in each case which δ answers a given ϵ . (The δ you produce at $t > 0$ depends on t . That dependence is the subject of section 3.4.)
7. Let $E = [0, 1) \cup [2, 3]$ and define $f: E \rightarrow \mathbb{R}$ by $f(x) = x$ for $x \in [0, 1)$ and $f(x) = x - 1$ for $x \in [2, 3]$. Show that f is continuous and strictly increasing, that $f(E) = [0, 2]$, and that the inverse map $[0, 2] \rightarrow E$ is not continuous at 1. Which hypothesis of theorem 3.3.12 does E fail?
8. Show that $\{x^2 \mid x \in (-1, 1)\} = [0, 1)$. Conclude that corollary 3.3.9 becomes false if the domain is a bounded interval that is not closed, even though the image is still an interval.

3.4 Uniform Continuity

Continuity at a point is a statement about that point. The δ answering a given ϵ in definition 3.2.1 is produced after the point has been named, and for most functions it has to be, since the same ϵ needs a smaller δ wherever the function moves faster. This section names the condition that holds when a single δ answers ϵ at every point of the domain at once, exhibits continuous functions that fail it, and shows that on a closed bounded interval continuity already implies it. Three consequences follow: such a function carries Cauchy sequences to Cauchy sequences, extends continuously to the closure of its domain, and carries bounded sets to bounded sets.

Definition 3.4.1: [Real] Uniform Continuity

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$. Then f is *uniformly continuous* on E if for every $\epsilon > 0$ there is a $\delta > 0$ such that

$$|f(x) - f(y)| < \epsilon \quad \text{for all } x, y \in E \text{ with } |x - y| < \delta$$

The condition constrains the pair (x, y) , no point of E is distinguished, and E itself is unrestricted. The function $x \mapsto 2x + 1$ on $\mathbb{R}$ satisfies it: given $\epsilon > 0$, the choice $\delta = \epsilon/2$ gives $|(2x + 1) - (2y + 1)| = 2|x - y| < \epsilon$ whenever $|x - y| < \delta$.

3.4.2 Note: The Order of the Quantifiers Written out, continuity of f on E says

$$\forall c \in E \quad \forall \epsilon > 0 \quad \exists \delta > 0 \quad \forall x \in E \quad (|x - c| < \delta \Rightarrow |f(x) - f(c)| < \epsilon)$$

and uniform continuity of f on E says

$$\forall \epsilon > 0 \quad \exists \delta > 0 \quad \forall c \in E \quad \forall x \in E \quad (|x - c| < \delta \Rightarrow |f(x) - f(c)| < \epsilon)$$

The two strings differ only in the position of $\exists \delta$, which has moved to the left of $\forall c$. In the first, δ may depend on c as well as on ϵ ; in the second it may depend on ϵ only.

One implication holds with no hypothesis on E : a uniformly continuous function is continuous. Given $c \in E$ and $\epsilon > 0$, the δ produced by definition 3.4.1 satisfies $|f(x) - f(c)| < \epsilon$ for every $x \in E$ with $|x - c| < \delta$, which is what definition 3.2.1 asks for at c . The converse is false, and example 3.4.3 gives three witnesses.

Negating definition 3.4.1, a function f fails to be uniformly continuous on E exactly when there is an $\epsilon_0 > 0$ such that for every $\delta > 0$ there are $x, y \in E$ with $|x - y| < \delta$ and $|f(x) - f(y)| \geq \epsilon_0$. Such an ϵ_0 is a tolerance that no single δ meets across all of E , and for each of the three functions below, uniform continuity is refuted by exhibiting one.

Example 3.4.3: $1/x$ on $(0, 1)$, x^2 on $\mathbb{R}$, and $1/(2 - q^2)$ on the Rationals in $[0, 2]$

1. Let $f(x) = 1/x$ on $E = (0, 1)$. The function $x \mapsto x$ is continuous and vanishes nowhere on E , so f is continuous on E by theorem 3.2.5. Take $\epsilon_0 = 1$ and let $\delta > 0$. By theorem 1.4.1(2) there is an $n \in \mathbb{N}$ with $n > \max\{2, 1/\delta\}$, and from $0 < 1/\delta < n$ and proposition 1.1.2(7) it follows that $0 < 1/n < \delta$. Put $x = 1/n$ and $y = 1/(2n)$, both of which lie in $(0, 1)$ because

$n > 2$. Then

$$|x - y| = \frac{1}{n} - \frac{1}{2n} = \frac{1}{2n} < \frac{1}{n} < \delta$$

while $|f(x) - f(y)| = |n - 2n| = n > 2 > \epsilon_0$. So no δ answers ϵ_0 , and f is not uniformly continuous on $(0, 1)$.

2. Let $g(x) = x^2$ on $\mathbb{R}$, which is continuous by example 3.2.6. Take $\epsilon_0 = 1$ and let $\delta > 0$. By theorem 1.4.1(2) choose $n \in \mathbb{N}$ with $n > 1/\delta$, and multiply this inequality by $\delta > 0$ using proposition 1.1.2(5) to get $\delta n > 1$. Put $x = n$ and $y = n + \delta/2$, so that $0 < x < y$ and $|x - y| = \delta/2 < \delta$. Then

$$|g(y) - g(x)| = y^2 - x^2 = (y - x)(y + x) = \frac{\delta}{2} \left(2n + \frac{\delta}{2} \right) = \delta n + \frac{\delta^2}{4} > \delta n > 1 = \epsilon_0$$

So g is not uniformly continuous on $\mathbb{R}$. Here the domain is all of $\mathbb{R}$ and the function is bounded on every bounded set, so the obstruction is neither of the ones that defeat (1) and (3).

3. Let $E_0 = \mathbb{Q} \cap [0, 2]$ and let $h: E_0 \rightarrow \mathbb{R}$ be $h(q) = 1/(2 - q^2)$. This is the function Section 3.3 sets against theorem 3.3.5, and it is defined, continuous and unbounded above on E_0 for the reasons given there. Throughout, $\sqrt{2}$ is the number given by theorem 1.3.12, and $1 < \sqrt{2} < 2$ is proved in Section 3.3.

For each $n \in \mathbb{N}$, theorem 1.4.4 produces a rational q_n with $\sqrt{2} - 1/n < q_n < \sqrt{2}$. Then $q_n > 0$, since $\sqrt{2} - 1/n \geq \sqrt{2} - 1 > 0$, and $q_n < 2$, so $q_n \in E_0$. From $|q_n - \sqrt{2}| < 1/n$, together with example 2.1.3(1) and definition 2.1.2, it follows that $q_n \rightarrow \sqrt{2}$, and then theorem 2.1.6(3) and (1) give $2 - q_n^2 \rightarrow 0$. Each $2 - q_n^2$ is positive, since $0 \leq q_n < \sqrt{2}$ forces $q_n^2 < (\sqrt{2})^2 = 2$ by proposition 3.3.11. Now let $M > 0$. Applying definition 2.1.2 to $(2 - q_n^2)$ with $\epsilon = 1/M$ produces an N' such that $0 < 2 - q_n^2 < 1/M$ for every $n \geq N'$, and inverting with proposition 1.1.2(7) gives $h(q_n) > M$ for every $n \geq N'$.

Now take $\epsilon_0 = 1$ and let $\delta > 0$. Since $q_n \rightarrow \sqrt{2}$, there is an $N \in \mathbb{N}$ with $|q_n - \sqrt{2}| < \delta/2$ for all $n \geq N$, so proposition 1.3.8(4) gives $|q_n - q_m| < \delta$ for all $n, m \geq N$. The value $h(q_N)$ is positive, so the previous paragraph applies to $M = h(q_N) + 1$ and returns an N' with $h(q_n) > h(q_N) + 1$ for every $n \geq N'$. Put $m = \max\{N, N'\}$. Then $|h(q_m) - h(q_N)| > 1 = \epsilon_0$ while $|q_m - q_N| < \delta$. So h is not uniformly continuous on E_0 .

Two of these three failures occur on a bounded domain, and in both the function is unbounded near a point the domain does not contain: 0 in item (1) and $\sqrt{2}$ in item (3). Item (3) is the failure over $\mathbb{Q}$. Its domain $E_0 = \mathbb{Q} \cap [0, 2]$ is bounded and contains both of its bounds, and it is nonetheless neither an interval in the sense of definition 3.1.1 nor a closed set in the sense of definition 3.1.10. Both failures occur at $\sqrt{2}$, which lies outside E_0 by proposition 1.1.3 and satisfies $1 < \sqrt{2} < 2$, as Section 3.3 proves. The betweenness condition fails for the pair $0, 2 \in E_0$, and the rationals q_n of item (3) lie in E_0 and converge to $\sqrt{2}$, so E_0 is not closed by proposition 3.1.12. The theorem below therefore cannot be proved from order and boundedness alone, and its proof goes through theorem 2.2.9.

Theorem 3.4.4: Continuity on a Closed Bounded Interval Is Uniform

Let $a < b$ be real numbers and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. Then f is uniformly continuous on $[a, b]$.

Proof:

Suppose f is not uniformly continuous on $[a, b]$. Negating definition 3.4.1, there is an $\epsilon_0 > 0$ such that every $\delta > 0$ admits $x, y \in [a, b]$ with $|x - y| < \delta$ and $|f(x) - f(y)| \geq \epsilon_0$. Applying this to $\delta = 1/n$ for each $n \in \mathbb{N}$ produces points $x_n, y_n \in [a, b]$ with

$$|x_n - y_n| < \frac{1}{n} \quad \text{and} \quad |f(x_n) - f(y_n)| \geq \epsilon_0 \quad (3.1)$$

Step 1: a convergent subsequence of (x_n) . Every term of (x_n) lies in $[a, b]$, so the sequence is bounded, and theorem 2.2.9(1) gives a subsequence $(x_{n_k})_{k \geq 1}$ converging to some $x \in \mathbb{R}$. Since $a \leq x_{n_k} \leq b$ for every k , two applications of proposition 2.1.8(2) give $a \leq x \leq b$. So $x \in [a, b]$ and f is continuous at x .

Step 2: the matching subsequence of (y_n) has the same limit. The sequence $(1/n)_{n \geq 1}$ converges to 0 by example 2.1.3(1), and $(1/n_k)_{k \geq 1}$ is a subsequence of it because $k \mapsto n_k$ is strictly increasing, so $1/n_k \rightarrow 0$ by proposition 2.2.7(1). Let $\epsilon > 0$ and take K with $1/n_k < \epsilon$ for every $k \geq K$. For such k the first half of (3.1) gives

$$|(y_{n_k} - x_{n_k}) - 0| = |y_{n_k} - x_{n_k}| < \frac{1}{n_k} < \epsilon$$

so $y_{n_k} - x_{n_k} \rightarrow 0$ by definition 2.1.2. Writing $y_{n_k} = x_{n_k} + (y_{n_k} - x_{n_k})$ and applying theorem 2.1.6(1) gives $y_{n_k} \rightarrow x$.

Step 3: the contradiction. Both (x_{n_k}) and (y_{n_k}) are sequences in $[a, b]$ converging to x , and f is continuous at x by Step 1, so proposition 3.2.4 gives $f(x_{n_k}) \rightarrow f(x)$ and $f(y_{n_k}) \rightarrow f(x)$. By theorem 2.1.6(2) with $c = -1$ and then theorem 2.1.6(1), the sequence $d_k = f(x_{n_k}) - f(y_{n_k})$ converges to $f(x) - f(x) = 0$. Applying definition 2.1.2 to (d_k) with $\epsilon = \epsilon_0$ produces a k with $|d_k| < \epsilon_0$, while the second half of (3.1) gives $|d_k| \geq \epsilon_0$ for every k . This contradiction shows that f is uniformly continuous on $[a, b]$, as we sought to show.

On a closed bounded interval the two conditions therefore coincide, and nothing is gained by checking the stronger one. This is one of the statements about a continuous function on $[a, b]$ that section 3.3 and this section prove from completeness, and it is the one whose failure over $\mathbb{Q}$ is visible in example 3.4.3(3). The proof applied continuity at the single point x , and everything else was theorem 2.2.9 and the algebra of limits.

The three results that follow hold for an arbitrary domain $E \subseteq \mathbb{R}$. The first is what the other two rest on.

Proposition 3.4.5: Uniform Continuity Carries Cauchy Sequences to Cauchy Sequences

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ be uniformly continuous, and let $(x_n)_{n \geq 1}$ be a Cauchy sequence with $x_n \in E$ for every n . Then $(f(x_n))_{n \geq 1}$ is a Cauchy sequence.

Proof:

Let $\epsilon > 0$. By definition 3.4.1 there is a $\delta > 0$ with $|f(x) - f(y)| < \epsilon$ for all $x, y \in E$ satisfying $|x - y| < \delta$. Since (x_n) is Cauchy, definition 2.3.1 applied to δ gives an $N \in \mathbb{N}$ with $|x_n - x_m| < \delta$ for all $n, m \geq N$. For such n and m both x_n and x_m lie in E , so the choice of δ gives $|f(x_n) - f(x_m)| < \epsilon$. As ϵ was arbitrary, $(f(x_n))$ satisfies definition 2.3.1, completing the proof.

Continuity alone does not give this. Take $f(x) = 1/x$ on $(0, 1)$ and $x_n = 1/(n + 1)$. Every x_n lies in $(0, 1)$, and (x_n) is a subsequence of $(1/n)_{n \geq 1}$, hence converges to 0 by example 2.1.3(1) and proposition 2.2.7(1), hence is Cauchy by proposition 2.3.3. But $f(x_n) = n + 1$, and no $M \geq 0$ bounds these values, since theorem 1.4.1(2) produces an $n \in \mathbb{N}$ with $n > M$. So $(f(x_n))$ is unbounded, and proposition 2.3.4 then says it is not Cauchy. This settles example 3.4.3(1) a second time, without producing an explicit pair of nearby points.

Once Cauchy sequences are preserved, a uniformly continuous function has a value forced on it at every point its domain approaches, because the images of the approaching sequences converge and the limit does not depend on which sequence is used. Proposition 3.1.9 is what turns a point of the closure into such a sequence.

Theorem 3.4.6: Uniformly Continuous Functions Extend to the Closure

Let $E \subseteq \mathbb{R}$ be nonempty and let $f: E \rightarrow \mathbb{R}$ be uniformly continuous. Then there is exactly one continuous function $g: \bar{E} \rightarrow \mathbb{R}$ with $g(x) = f(x)$ for every $x \in E$, and this g is uniformly continuous on $\bar{E}$.

Proof:

Step 1: a candidate value at each point of the closure. Let $a \in \bar{E}$, the closure of definition 3.1.4. By proposition 3.1.9 there is a sequence (x_n) with $x_n \in E$ for every n and $x_n \rightarrow a$. By proposition 2.3.3 the sequence (x_n) is Cauchy, so $(f(x_n))$ is Cauchy by proposition 3.4.5, and therefore converges by theorem 2.3.5.

Step 2: the value does not depend on the sequence. Let (x_n) and (y_n) be sequences in E with $x_n \rightarrow a$ and $y_n \rightarrow a$, and let L and L' be the limits of $(f(x_n))$ and $(f(y_n))$ produced in Step 1. Define $z_{2n-1} = x_n$ and $z_{2n} = y_n$ for $n \geq 1$, so that every z_m lies in E . Then $z_m \rightarrow a$. Indeed, given $\epsilon > 0$, pick N_1 with $|x_n - a| < \epsilon$ for $n \geq N_1$ and N_2 with $|y_n - a| < \epsilon$ for $n \geq N_2$, and set $N = 2 \max\{N_1, N_2\}$. If $m \geq N$ and $m = 2n - 1$, then $2n = m + 1 > N$, so $n > N/2 = \max\{N_1, N_2\} \geq N_1$. If $m \geq N$ and $m = 2n$, then $n = m/2 \geq N/2 \geq N_2$. In both cases $|z_m - a| < \epsilon$.

By Step 1 applied to (z_n) , the sequence $(f(z_n))$ converges, to M say. The index maps $n \mapsto 2n - 1$ and $n \mapsto 2n$ are strictly increasing, so $(f(x_n))$ and $(f(y_n))$ are subsequences of $(f(z_n))$, and proposition 2.2.7(1) gives $f(x_n) \rightarrow M$ and $f(y_n) \rightarrow M$. By proposition 2.1.4, $L = M = L'$. Setting $g(a) = L$ therefore defines a function $g: \bar{E} \rightarrow \mathbb{R}$.

Step 3: g extends f . Let $a \in E$. The constant sequence $x_n = a$ lies in E and converges to a

by example 2.1.3(2), so $a \in \overline{E}$ by proposition 3.1.9, and in particular $E \subseteq \overline{E}$. By Step 2 the value $g(a)$ may be computed from this sequence, and $f(x_n) = f(a)$ is constant, so $g(a) = f(a)$ by example 2.1.3(2) and proposition 2.1.4.

Step 4: g is uniformly continuous. Let $\epsilon > 0$. Apply definition 3.4.1 to f and the positive number $\epsilon/3$ to get a $\delta > 0$ with $|f(x) - f(y)| < \epsilon/3$ for all $x, y \in E$ with $|x - y| < \delta$. Let $a, b \in \overline{E}$ satisfy $|a - b| < \delta/3$, and choose sequences (x_n) and (y_n) in E with $x_n \rightarrow a$ and $y_n \rightarrow b$, so that $f(x_n) \rightarrow g(a)$ and $f(y_n) \rightarrow g(b)$ by Steps 1 and 2. Pick N with $|x_n - a| < \delta/3$ and $|y_n - b| < \delta/3$ for all $n \geq N$. For such n , two applications of proposition 1.3.8(4) give

$$|x_n - y_n| \leq |x_n - a| + |a - b| + |b - y_n| < \frac{\delta}{3} + \frac{\delta}{3} + \frac{\delta}{3} = \delta$$

so $|f(x_n) - f(y_n)| < \epsilon/3$, which proposition 1.3.8(2) turns into $-\epsilon/3 \leq f(x_n) - f(y_n) \leq \epsilon/3$ for every $n \geq N$. The sequence $f(x_n) - f(y_n)$ converges to $g(a) - g(b)$ by theorem 2.1.6(2) with $c = -1$ and then theorem 2.1.6(1), so proposition 2.1.8(2) applied twice gives $-\epsilon/3 \leq g(a) - g(b) \leq \epsilon/3$, and proposition 1.3.8(2) converts this to $|g(a) - g(b)| \leq \epsilon/3 < \epsilon$. Thus $\delta/3$ answers ϵ for g , so g is uniformly continuous on $\overline{E}$, and by box 3.4.2 it is continuous there.

Step 5: uniqueness. Let $h: \overline{E} \rightarrow \mathbb{R}$ be continuous with $h(x) = f(x)$ for every $x \in E$, and let $a \in \overline{E}$. Choose a sequence (x_n) in E with $x_n \rightarrow a$, using proposition 3.1.9. Every x_n lies in $E \subseteq \overline{E}$, which is the domain of h , and h is continuous at a , so proposition 3.2.4 gives $h(x_n) \rightarrow h(a)$. But $h(x_n) = f(x_n) \rightarrow g(a)$ by Steps 1 and 2, so $h(a) = g(a)$ by proposition 2.1.4. Since a was an arbitrary point of $\overline{E}$, the functions h and g are equal, completing the proof.

The extension is unique for a reason that has nothing to do with uniform continuity. By proposition 3.1.9 every point of $\overline{E}$ is a limit of points of E , and proposition 3.2.4 then forces any two continuous functions agreeing on E to agree on $\overline{E}$. What uniform continuity provides is existence, and continuity does not provide it: the function $x \mapsto 1/x$ on $(0, 1)$ takes the value $n + 1$ at $1/(n + 1)$, so it is unbounded on $(0, 1)$ by theorem 1.4.1(2), whereas every continuous function on $[0, 1]$ is bounded by theorem 3.3.5. No continuous function on $[0, 1]$ restricts to it. The closure of $(0, 1)$ is $[0, 1]$ by example 3.1.7(1), so the extension that fails here is exactly the one the theorem asserts.

That computation is an instance of the third consequence, which explains both failures on a bounded domain in example 3.4.3 at once. Recall that a subset $S \subseteq \mathbb{R}$ is bounded when it is bounded above and below in the sense of definition 1.2.1. This is the same as the existence of an $M \geq 0$ with $|t| \leq M$ for every $t \in S$: given such an M , proposition 1.3.8(2) makes $-M$ a lower bound and M an upper bound, and conversely if $\ell \leq t \leq u$ for every $t \in S$ then $M = \max\{|\ell|, |u|\}$ works, since proposition 1.3.8(1) gives $t \leq u \leq |u| \leq M$ and $-t \leq -\ell \leq |\ell| \leq M$, whence $|t| \leq M$ by proposition 1.3.8(2).

Proposition 3.4.7: Uniform Continuity Carries Bounded Sets to Bounded Sets

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$ be uniformly continuous. If $B \subseteq E$ is bounded, then $f(B)$ is bounded. In particular, a uniformly continuous function on a bounded domain is bounded.

Proof:

Suppose $f(B)$ is not bounded. Then for each $n \in \mathbb{N}$ there is a $b_n \in B$ with $|f(b_n)| > n$, since otherwise $|f(x)| \leq n$ would hold for every $x \in B$, making $-n$ a lower bound and n an upper bound for $f(B)$ by proposition 1.3.8(2). The sequence (b_n) lies in the bounded set B , so it is bounded, and theorem 2.2.9(1) gives a convergent subsequence $(b_{n_k})_{k \geq 1}$.

By proposition 2.3.3 the sequence (b_{n_k}) is Cauchy, and all of its terms lie in $B \subseteq E$, so

proposition 3.4.5 makes $(f(b_{n_k}))$ Cauchy and proposition 2.3.4 gives an $M \geq 0$ with $|f(b_{n_k})| \leq M$ for every k . Combining this with $|f(b_{n_k})| > n_k$ gives $n_k < M$ for every k .

This is impossible. The sequence $(1/n_k)_{k \geq 1}$ is a subsequence of $(1/n)_{n \geq 1}$, which converges to 0 by example 2.1.3(1), so $1/n_k \rightarrow 0$ by proposition 2.2.7(1). Applying definition 2.1.2 with $\epsilon = 1/(M+1)$, a positive number because $M \geq 0$, produces a k with $0 < 1/n_k < 1/(M+1)$, and proposition 1.1.2(7) inverts this to $M+1 < n_k$. Together with $n_k < M$ this gives $M+1 < M$, which is false. Hence $f(B)$ is bounded. Taking $B = E$ gives the final sentence, completing the proof.

Items (1) and (3) of example 3.4.3 are now settled in one line each, since both domains are bounded and both functions are unbounded on them. Item (2) is not settled this way, because $x \mapsto x^2$ is bounded on every bounded subset of $\mathbb{R}$. That item also shows the converse of proposition 3.4.7 to be false: a continuous function may carry bounded sets to bounded sets without being uniformly continuous.

A single quantitative condition implies uniform continuity outright, and it is the one that most concrete estimates produce.

Definition 3.4.8: [Real] Lipschitz

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$. Then f is *Lipschitz* on E if there is an $L \geq 0$ with

$$|f(x) - f(y)| \leq L|x - y| \quad \text{for all } x, y \in E$$

Any such L is called a *Lipschitz constant* for f on E .

The absolute value $x \mapsto |x|$ on $\mathbb{R}$ is Lipschitz with $L = 1$, which is exactly proposition 1.3.8(5). An affine function $x \mapsto cx + d$ on $\mathbb{R}$ is Lipschitz with $L = |c|$, since $|(cx + d) - (cy + d)| = |c||x - y|$ by proposition 1.3.8(3). And for $R > 0$ the function $x \mapsto x^2$ on $[0, R]$ is Lipschitz with $L = 2R$, since for $x, y \in [0, R]$,

$$|x^2 - y^2| = |x + y||x - y| \leq 2R|x - y|$$

where the equality is proposition 1.3.8(3) and the inequality uses $0 \leq x + y \leq 2R$ together with $|x - y| \geq 0$. Compare the last instance with example 3.4.3(2): squaring is Lipschitz on each bounded interval, with a constant that grows with the interval, and no single constant serves all of $\mathbb{R}$.

Proposition 3.4.9: Lipschitz Implies Uniformly Continuous

Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$ be Lipschitz on E . Then f is uniformly continuous on E .

Proof:

Let $L \geq 0$ be a Lipschitz constant for f and put $L' = L + 1$, so that $L' > 0$. For $x, y \in E$ the number $|x - y|$ is nonnegative and $L \leq L'$, so $L|x - y| \leq L'|x - y|$ and therefore

$$|f(x) - f(y)| \leq L'|x - y|$$

Let $\epsilon > 0$ and put $\delta = \epsilon/L' > 0$. If $x, y \in E$ and $|x - y| < \delta$, then multiplying by $L' > 0$ and using proposition 1.1.2(5) gives $L'|x - y| < L'\delta = \epsilon$, so $|f(x) - f(y)| < \epsilon$. Hence δ answers ϵ in definition 3.4.1, as we sought to show.

The implication is strict, and the square root on $[0, 1]$ separates the two conditions.

Example 3.4.10: The Square Root on $[0, 1]$

Let $s: [0, 1] \rightarrow \mathbb{R}$ be $s(x) = x^{1/2}$, which is defined because every $x \in [0, 1]$ satisfies $x \geq 0$ (theorem 1.3.12 with $n = 2$).

First, s is uniformly continuous on $[0, 1]$. By corollary 3.3.13 the function $x \mapsto x^{1/2}$ is continuous on $[0, \infty)$, so s is continuous by proposition 3.3.1. Now theorem 3.4.4 applies to s on $[0, 1]$.

Second, s is not Lipschitz on $[0, 1]$. Suppose $L \geq 0$ satisfied $|s(x) - s(y)| \leq L|x - y|$ for all $x, y \in [0, 1]$. Fix $n \in \mathbb{N}$ and take $y = 0$ and $x = 1/n^2$, which lies in $[0, 1]$ because $n \geq 1$ forces $n^2 \geq 1$. Since $(1/n)^2 = 1/n^2$ and $1/n \geq 0$, the uniqueness clause of theorem 1.3.12 gives $s(1/n^2) = 1/n$, and the same clause gives $s(0) = 0$. The assumed inequality reads

$$\frac{1}{n} \leq \frac{L}{n^2}$$

and multiplying by $n^2 > 0$ gives $n \leq L$. This holds for every $n \in \mathbb{N}$, contradicting theorem 1.4.1(2), which produces an $n \in \mathbb{N}$ with $n > L$. So no Lipschitz constant exists.

The failure is a statement about how large a δ a given ϵ permits. A Lipschitz function admits $\delta = \epsilon/L'$, which is proportional to ϵ , while the square root on $[0, 1]$ needs a δ that shrinks faster than that near 0. How much faster is the subject of exercise 3.4.1 (1), where $\delta = \epsilon^2$ is shown to answer ϵ on all of $[0, \infty)$, a domain that is not bounded and to which theorem 3.4.4 does not apply.

Exercise 3.4.1

1. Let $x, y \geq 0$. Prove that $|x^{1/2} - y^{1/2}| \leq |x - y|^{1/2}$, and deduce that $x \mapsto x^{1/2}$ is uniformly continuous on $[0, \infty)$, with $\delta = \epsilon^2$ answering ϵ .
2. Let $E, F \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ be uniformly continuous with $f(E) \subseteq F$, and let $g: F \rightarrow \mathbb{R}$ be uniformly continuous. Prove that $g \circ f$ is uniformly continuous on E .
3. Let $E \subseteq \mathbb{R}$ and let $f, g: E \rightarrow \mathbb{R}$ be uniformly continuous. Prove that $f + g$ is uniformly continuous on E . Then show that the product fg need not be, by taking $E = \mathbb{R}$ and $f = g$ the identity function.
4. Let $c > 0$. Prove that $x \mapsto 1/x$ is uniformly continuous on $[c, \infty)$ by exhibiting a Lipschitz constant, and prove that it is not uniformly continuous on $(0, \infty)$.
5. Let $f: (0, 1] \rightarrow \mathbb{R}$ be continuous. Prove that f is uniformly continuous on $(0, 1]$ if and only if there is a continuous $g: [0, 1] \rightarrow \mathbb{R}$ with $g(x) = f(x)$ for every $x \in (0, 1]$. Begin by showing that the closure of $(0, 1]$ is $[0, 1]$.
6. Let $E \subseteq \mathbb{R}$ and let $f: E \rightarrow \mathbb{R}$. Prove that f is uniformly continuous on E if and only if $f(x_n) - f(y_n) \rightarrow 0$ for every pair of sequences (x_n) and (y_n) in E with $x_n - y_n \rightarrow 0$.

Differentiation

The derivative measures how a function changes, and the definition is a limit of the kind Section 3.1 already built. What makes the chapter more than an application of that machinery is a single theorem. The mean value theorem converts a hypothesis about the derivative at every point into a statement about the function across an interval, and every use of the derivative as a tool passes through it.

The route to it runs through Chapter 3. A differentiable function on a closed bounded interval attains an interior extremum, the derivative vanishes there, and the extremum exists because theorem 3.3.7 says so. So the mean value theorem rests on completeness exactly as the theorems of the previous chapter do, and the chapter inherits their standing prohibition: no compactness, no connectedness.

The second half of the chapter is about interchanging two limits. A sequence of functions may converge at every point while the limit function loses every property the terms had, and the condition that prevents this is uniform convergence. It is the same strengthening Section 3.4 applied to a single function, made this time in the index where that one was made in the point. Applied to power series it turns Section 2.6' formal objects into functions that can be differentiated term by term, which is what makes $\exp$, $\cos$ and $\sin$ available as functions at last.

Section 4.1 defines the derivative, proves the rules, and gives Newton's approximation, the formulation that keeps the chain rule free of a division by zero. Section 4.2 proves Rolle's theorem and the mean value theorem, then two consequences that are not corollaries of each other: Darboux's theorem, that a derivative has the intermediate value property of definition 3.3.4 even when it is discontinuous, and L'Hôpital's rule. Section 4.3 approximates by polynomials and bounds the error. Section 4.4 defines uniform convergence for sequences and series of functions and proves the three theorems that make it useful, ending with term-by-term differentiation. Section 4.5 specializes to power series, where the coefficients turn out to be determined by the function. Section 4.6 establishes the exponential, the logarithm, the trigonometric functions, their addition formulas, and π . Section 4.7 closes by asking which smooth functions are sums of their own Taylor series, and answers with a growth condition on the derivatives.

4.1 The Derivative

A difference quotient records the average rate of change of a function across an interval of positive length, and the derivative at a point is the limit of those averages as the interval closes on the point. Two features of the setting are fixed here and used without comment afterwards. The domain is an interval with more than one point, which makes every point of the domain a limit point of it, so definition 3.1.15 applies at every point with no side condition left to check. And the difference quotient at c is undefined at c itself, which does no harm because the limit of definition 3.1.15 is deleted (box 3.1.16).

The first of these follows from definition 3.1.1, and is recorded as a lemma because three later sections spend it.

Lemma 4.1.1: Every Point of a Nondegenerate Interval Is a Limit Point of It

Let $I \subseteq \mathbb{R}$ be an interval with more than one point and let $c \in I$. Then c is a limit point of I , and of $I \setminus \{c\}$.

Proof:

Let $\delta > 0$. Since I has a point other than c , there is a $y \in I$ with $y \neq c$. Suppose first that $y > c$ and put $z = \min(y, c + \delta/2)$, so that $c < z \leq y$. If $z < y$, then $c < z < y$ with $c, y \in I$ gives $z \in I$ by definition 3.1.1. If $z = y$, then $z \in I$ already. In both cases $0 < |z - c| = z - c \leq \delta/2 < \delta$. If instead $y < c$, then $z = \max(y, c - \delta/2)$ satisfies $y \leq z < c$, the same two cases put $z \in I$, and $0 < |z - c| = c - z \leq \delta/2 < \delta$. So c is a limit point of I in the sense of definition 3.1.5. Each witness z produced here differs from c , so it lies in $I \setminus \{c\}$, and c is a limit point of that set as well.

Definition 4.1.2: [Real] Derivative

Let $I \subseteq \mathbb{R}$ be an interval with more than one point, let $f: I \rightarrow \mathbb{R}$ and let $c \in I$. The *difference quotient of f at c* is the function $q_{f,c}: I \setminus \{c\} \rightarrow \mathbb{R}$ given by

$$q_{f,c}(x) = \frac{f(x) - f(c)}{x - c}$$

Then f is *differentiable at c* if $\lim_{x \rightarrow c} q_{f,c}(x)$ exists in $\mathbb{R}$. That limit is written $f'(c)$ and called the *derivative of f at c* .

If f is differentiable at every point of I , then f is *differentiable*, and the function $f': I \rightarrow \mathbb{R}$ is its *derivative*. Put $f^{(0)} = f$, and for $k \geq 0$ put $f^{(k+1)} = (f^{(k)})'$ whenever the right-hand side exists. Then $f^{(k)}$ is the *k -th derivative of f* . The first three are also written f' , f'' and f''' .

The limit in the definition is taken over a domain that excludes c , so the number $f(c)$ enters only inside the numerator and never as a point at which anything is evaluated. This is what makes the definition usable: the quotient would be $0/0$ at $x = c$, and box 3.1.16 removes that point from every requirement the limit makes. The limit-point argument above is what guarantees there are still enough points left to determine the limit, and proposition 3.1.18 then makes $f'(c)$ a single real number.

Example 4.1.3: Four Derivatives from the Definition

1. Let I be any interval with more than one point, let $\lambda \in \mathbb{R}$ and let $f(x) = \lambda$ for every $x \in I$. For $c \in I$ and $x \in I \setminus \{c\}$, $q_{f,c}(x) = (\lambda - \lambda)/(x - c) = 0$. A constant function has that constant as its limit, directly from definition 3.1.15 with any $\delta > 0$ answering any ϵ . So $f'(c) = 0$ for every $c \in I$.
2. Let $f(x) = x$ on $\mathbb{R}$ and let $c \in \mathbb{R}$. For $x \neq c$ the quotient is $q_{f,c}(x) = (x - c)/(x - c) = 1$, so by the same remark $f'(c) = 1$.
3. Let $f(x) = x^2$ on $\mathbb{R}$ and let $c \in \mathbb{R}$. For $x \neq c$,

$$q_{f,c}(x) = \frac{x^2 - c^2}{x - c} = \frac{(x - c)(x + c)}{x - c} = x + c$$

The identity function has $\lim_{x \rightarrow c} x = c$, since $\delta = \epsilon$ answers ϵ in definition 3.1.15. The constant function with value c has limit c by the remark in item (1), so corollary 3.1.20(1) gives $\lim_{x \rightarrow c} (x + c) = 2c$, and $f'(c) = 2c$.

4. Let $f(x) = x^{1/2}$ on $(0, \infty)$ and let $c > 0$. For $x \in (0, \infty)$ with $x \neq c$ the two square roots differ, by the uniqueness clause of theorem 1.3.12, and $x - c = (x^{1/2})^2 - (c^{1/2})^2 = (x^{1/2} - c^{1/2})(x^{1/2} + c^{1/2})$, so

$$q_{f,c}(x) = \frac{x^{1/2} - c^{1/2}}{(x^{1/2} - c^{1/2})(x^{1/2} + c^{1/2})} = \frac{1}{x^{1/2} + c^{1/2}}$$

The root function is continuous on $[0, \infty)$ by corollary 3.3.13, so $\lim_{x \rightarrow c} x^{1/2} = c^{1/2}$ by proposition 3.2.3(1), and proposition 3.1.23(1) transfers this limit to the subset $(0, \infty) \setminus \{c\}$, of which c is a limit point. Adding the constant $c^{1/2}$ gives $\lim_{x \rightarrow c} (x^{1/2} + c^{1/2}) = 2c^{1/2}$, which is nonzero because $c > 0$. Now corollary 3.1.20(4), applied with numerator the constant function 1, gives $f'(c) = 1/(2c^{1/2})$.

Item (4) was computed by cancelling the factor $x^{1/2} - c^{1/2}$, and that cancellation is valid only because the factor is nonzero for $x \neq c$. The denominator $x - c$ of a difference quotient is nonzero for the same reason. Both facts fail once the denominator becomes a difference of values of some other function, which may vanish at points arbitrarily close to c . The next theorem restates differentiability with no division in it at all.

Theorem 4.1.4: Newton's Approximation

Let $I \subseteq \mathbb{R}$ be an interval with more than one point, let $f: I \rightarrow \mathbb{R}$, let $c \in I$ and let $L \in \mathbb{R}$. The following are equivalent.

1. f is differentiable at c with $f'(c) = L$.
2. For every $\epsilon > 0$ there is a $\delta > 0$ such that

$$|f(x) - f(c) - L(x - c)| \leq \epsilon|x - c| \quad \text{for every } x \in I \text{ with } |x - c| < \delta$$

Proof:

Throughout, c is a limit point of I and of $I \setminus \{c\}$, by the argument opening this section, so definition 4.1.2 applies to f at c .

Step 1: (1) implies (2). Let $\epsilon > 0$. By definition 4.1.2 and definition 3.1.15 applied to $q_{f,c}$ there is a $\delta > 0$ with $|q_{f,c}(x) - L| < \epsilon$ for every $x \in I \setminus \{c\}$ with $0 < |x - c| < \delta$. Fix such an x . Multiplying $q_{f,c}(x) - L$ by $x - c$ gives $f(x) - f(c) - L(x - c)$, so proposition 1.3.8(3) gives

$$|f(x) - f(c) - L(x - c)| = |q_{f,c}(x) - L| |x - c| \leq \epsilon |x - c|$$

The remaining point of I with $|x - c| < \delta$ is $x = c$, where both sides of the displayed inequality are 0. So the inequality holds for every $x \in I$ with $|x - c| < \delta$.

Step 2: (2) implies (1). Let $\epsilon > 0$ and apply (2) with $\epsilon/2$ in place of ϵ , which is positive by proposition 1.1.2(7). This gives a $\delta > 0$ with $|f(x) - f(c) - L(x - c)| \leq (\epsilon/2)|x - c|$ for every $x \in I$ with $|x - c| < \delta$. Let $x \in I \setminus \{c\}$ with $0 < |x - c| < \delta$. Then $|x - c| > 0$, and dividing the last inequality by it, using proposition 1.3.8(3) once more, gives

$$|q_{f,c}(x) - L| \leq \frac{\epsilon}{2} < \epsilon$$

Since $\epsilon > 0$ was arbitrary, $\lim_{x \rightarrow c} q_{f,c}(x) = L$, which is statement (1), completing the proof.

The content of (2) is that $x \mapsto f(c) + L(x - c)$ approximates f near c with an error small compared with $|x - c|$, and that no other slope does. Condition (2) is an inequality between quantities defined at every point of I , the point c included, so it may be applied where the increment $x - c$ has been replaced by the increment of some other function. It also carries an explicit bound on $|f(x) - f(c)|$, which is what the next proposition extracts.

Proposition 4.1.5: Differentiability Implies Continuity

Let $I \subseteq \mathbb{R}$ be an interval with more than one point, let $f: I \rightarrow \mathbb{R}$ and let $c \in I$. If f is differentiable at c , then f is continuous at c .

Proof:

Write $L = f'(c)$ and apply theorem 4.1.4(2) with $\epsilon = 1$: there is a $\delta_0 > 0$ with $|f(x) - f(c) - L(x - c)| \leq |x - c|$ for every $x \in I$ with $|x - c| < \delta_0$. For such an x , proposition 1.3.8(4) and (3) give

$$|f(x) - f(c)| \leq |f(x) - f(c) - L(x - c)| + |L| |x - c| \leq (1 + |L|)|x - c|$$

Now let $\epsilon > 0$ and put $\delta = \min(\delta_0, \epsilon/(1 + |L|))$, which is positive because $1 + |L| > 0$ by proposition 1.3.8(1). If $x \in I$ satisfies $|x - c| < \delta$, then multiplying that inequality by the positive number $1 + |L|$, as in proposition 1.1.2(5), gives

$$|f(x) - f(c)| \leq (1 + |L|)|x - c| < (1 + |L|)\delta \leq \epsilon$$

This is definition 3.2.1 for f at c , as we sought to show.

The converse fails, and it fails at one point of a function that is continuous on all of $\mathbb{R}$.

Example 4.1.6: The Absolute Value at the Origin

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = |x|$.

1. f is continuous at every $c \in \mathbb{R}$, by example 3.2.9.
2. f is not differentiable at 0. For $x \neq 0$, $q_{f,0}(x) = |x|/x$, which equals 1 when $x > 0$ and -1 when $x < 0$, the second case because $|x| = -x$ there. Take $x_n = 1/n$ and $y_n = -1/n$. Both are sequences in $\mathbb{R} \setminus \{0\}$, and both converge to 0, the first by example 2.1.3(1) and the second by theorem 2.1.6(2) with the scalar -1 . Meanwhile $q_{f,0}(x_n) = 1 \rightarrow 1$ and $q_{f,0}(y_n) = -1 \rightarrow -1$ by example 2.1.3(2). Since $1 \neq -1$, theorem 3.1.19(2) shows that $q_{f,0}$ has no limit at 0.
3. f is differentiable at every $c \neq 0$, with $f'(c) = 1$ for $c > 0$ and $f'(c) = -1$ for $c < 0$. Suppose $c > 0$ and put $r = c > 0$. Every x with $|x - c| < r$ satisfies $x > 0$, so for x in the set $(\mathbb{R} \setminus \{c\}) \cap (c - r, c + r)$ both $|x| = x$ and $|c| = c$, and

$$q_{f,c}(x) = \frac{|x| - |c|}{x - c} = \frac{x - c}{x - c} = 1$$

By proposition 3.1.22(2) the limit of $q_{f,c}$ at c is therefore the limit of the constant function 1, which is 1. The case $c < 0$ is the same with $r = -c$, where $|x| = -x$ and $|c| = -c$ give $q_{f,c}(x) = -1$.

So differentiability at a point is strictly stronger than continuity there, and the gap already opens at a single point of the absolute value. Every derivative computed so far came from evaluating one limit by hand. The four algebraic operations that build new functions from old each convert into an operation on the derivatives, which reduces most computations to the four formulas below.

Theorem 4.1.7: The Rules of Differentiation

Let $I \subseteq \mathbb{R}$ be an interval with more than one point, let $f, g: I \rightarrow \mathbb{R}$ be differentiable at $c \in I$, and let $\lambda \in \mathbb{R}$.

1. $f + g$ is differentiable at c , with $(f + g)'(c) = f'(c) + g'(c)$.
2. λf is differentiable at c , with $(\lambda f)'(c) = \lambda f'(c)$.
3. fg is differentiable at c , with $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$.
4. If $g(x) \neq 0$ for every $x \in I$, then f/g is differentiable at c , with

$$\left(\frac{f}{g}\right)'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}$$

5. For every integer $n \geq 1$ the function $p_n: \mathbb{R} \rightarrow \mathbb{R}$ given by $p_n(x) = x^n$ is differentiable, with $p'_n(x) = nx^{n-1}$, where x^0 is read as 1 for every $x \in \mathbb{R}$.

Proof:

Write $E = I \setminus \{c\}$, a set of which c is a limit point. Each of (1) to (4) is proved by rewriting the difference quotient of the new function on E in terms of $q_{f,c}$ and $q_{g,c}$ and applying corollary 3.1.20.

1. For $x \in E$,

$$q_{f+g,c}(x) = \frac{f(x) + g(x) - f(c) - g(c)}{x - c} = q_{f,c}(x) + q_{g,c}(x)$$

and corollary 3.1.20(1) gives the limit $f'(c) + g'(c)$.

2. For $x \in E$, $q_{\lambda f,c}(x) = \lambda q_{f,c}(x)$, and corollary 3.1.20(2) gives the limit $\lambda f'(c)$.

3. The identity to check is that for $x \in E$,

$$q_{fg,c}(x) = \frac{f(x)g(x) - f(c)g(c)}{x - c} = q_{f,c}(x)g(x) + f(c)q_{g,c}(x)$$

Over the common denominator $x - c$, the numerator of the right-hand side is

$$(f(x) - f(c))g(x) + f(c)(g(x) - g(c)) = f(x)g(x) - f(c)g(x) + f(c)g(x) - f(c)g(c)$$

and the two middle terms cancel, leaving $f(x)g(x) - f(c)g(c)$, which is the numerator on the left.

The function g is continuous at c by proposition 4.1.5, so $\lim_{x \rightarrow c} g(x) = g(c)$ by proposition 3.2.3(1), and proposition 3.1.23(1) transfers that limit to the restriction of g to E . Applying corollary 3.1.20(3) to the first summand and (2) to the second, then (1) to the sum, gives

$$\lim_{x \rightarrow c} q_{fg,c}(x) = f'(c)g(c) + f(c)g'(c)$$

4. The hypothesis makes f/g a function on I . For $x \in E$,

$$q_{f/g,c}(x) = \frac{1}{x - c} \left(\frac{f(x)}{g(x)} - \frac{f(c)}{g(c)} \right) = \frac{f(x)g(c) - f(c)g(x)}{(x - c)g(x)g(c)} = \frac{q_{f,c}(x)g(c) - f(c)q_{g,c}(x)}{g(x)g(c)}$$

the second equality by putting the two fractions over the denominator $g(x)g(c)$, and the third because

$$(f(x) - f(c))g(c) - f(c)(g(x) - g(c)) = f(x)g(c) - f(c)g(c) - f(c)g(x) + f(c)g(c)$$

whose two $f(c)g(c)$ terms cancel to leave $f(x)g(c) - f(c)g(x)$.

By the previous part, $\lim_{x \rightarrow c} g(x) = g(c)$ along E , so corollary 3.1.20(2) gives $\lim_{x \rightarrow c} g(x)g(c) = g(c)^2$, which is nonzero by proposition 1.1.2(3) since $g(c) \neq 0$. The numerator has limit $f'(c)g(c) - f(c)g'(c)$ by parts (1) and (2) of corollary 3.1.20. The denominator $g(x)g(c)$ is nonzero at every $x \in E$, again by proposition 1.1.2(3), so the set E^* appearing in corollary 3.1.20(4) is all of E , and that part gives

$$\lim_{x \rightarrow c} q_{f/g,c}(x) = \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}$$

5. Induction on n . For $n = 1$, example 4.1.3(2) gives $p'_1(x) = 1 = 1 \cdot x^0$. Suppose $p'_n(x) = nx^{n-1}$ for every $x \in \mathbb{R}$. Since $p_{n+1} = p_n p_1$, part (3) applies at every $x \in \mathbb{R}$ and gives

$$p'_{n+1}(x) = p'_n(x)p_1(x) + p_n(x)p'_1(x) = nx^{n-1} \cdot x + x^n \cdot 1 = nx^n + x^n = (n+1)x^n$$

where $nx^{n-1} \cdot x = nx^n$ holds for $n = 1$ as well, because $x^0 = 1$ there. This is the claim for $n + 1$, completing the proof.

Parts (1) and (2) say that differentiation at c is linear on the functions differentiable at c . Parts (1), (2) and (5) make every polynomial differentiable on $\mathbb{R}$, with the derivative computed term by term, and parts (3) and (4) then make every rational function differentiable at every point where its denominator is nonzero. What remains is composition, and it is the one rule the difference quotient does not deliver on its own. Writing

$$\frac{f(g(x)) - f(g(c))}{x - c} = \frac{f(g(x)) - f(g(c))}{g(x) - g(c)} \cdot \frac{g(x) - g(c)}{x - c}$$

requires $g(x) \neq g(c)$, and that can fail at points arbitrarily close to c . For a constant g the middle expression is undefined at every x , so the identity above says nothing at any point. Theorem 4.1.4 states differentiability without a division, and the proof below is carried out with it.

Theorem 4.1.8: The Chain Rule

Let $I, J \subseteq \mathbb{R}$ be intervals, each with more than one point, let $g: I \rightarrow \mathbb{R}$ satisfy $g(I) \subseteq J$, and let $f: J \rightarrow \mathbb{R}$. Let $c \in I$ and suppose g is differentiable at c and f is differentiable at $g(c)$. Then $f \circ g: I \rightarrow \mathbb{R}$ is differentiable at c , and

$$(f \circ g)'(c) = f'(g(c)) g'(c)$$

Proof:

Write $d = g(c) \in J$, $A = g'(c)$ and $B = f'(d)$. Let $\epsilon > 0$ and put

$$\eta = \frac{\epsilon}{2(|A| + 1)}, \quad \eta' = \min\left(1, \frac{\epsilon}{2(|B| + 1)}\right)$$

Both are positive, since $|A| + 1$ and $|B| + 1$ are positive by proposition 1.3.8(1).

Step 1: the approximation for f at d . By theorem 4.1.4(2) applied to f at d with tolerance η , there is a $\delta_1 > 0$ such that

$$|f(y) - f(d) - B(y - d)| \leq \eta|y - d| \quad \text{for every } y \in J \text{ with } |y - d| < \delta_1 \quad (4.1)$$

Step 2: the approximation for g at c , and a bound on the increment. By theorem 4.1.4(2) applied to g at c with tolerance η' , there is a $\delta_2 > 0$ such that

$$|g(x) - d - A(x - c)| \leq \eta'|x - c| \quad \text{for every } x \in I \text{ with } |x - c| < \delta_2 \quad (4.2)$$

For such an x , proposition 1.3.8(4) and (3) give

$$|g(x) - d| \leq |g(x) - d - A(x - c)| + |A||x - c| \leq (\eta' + |A|)|x - c| \leq (1 + |A|)|x - c|$$

the last step because $\eta' \leq 1$.

Step 3: the two estimates combined. Put

$$\delta = \min\left(\delta_2, \frac{\delta_1}{1 + |A|}\right) > 0$$

and let $x \in I$ with $|x - c| < \delta$. Then $y = g(x)$ lies in J , and Step 2 gives $|y - d| \leq (1 + |A|)|x - c| < (1 + |A|)\delta \leq \delta_1$, so (4.1) applies to this y and yields

$$|f(g(x)) - f(d) - B(g(x) - d)| \leq \eta|g(x) - d| \leq \eta(1 + |A|)|x - c| = \frac{\epsilon}{2}|x - c|$$

Separately, (4.2) and proposition 1.3.8(3) give

$$|B(g(x) - d) - BA(x - c)| = |B||g(x) - d - A(x - c)| \leq |B|\eta'|x - c| \leq \frac{\epsilon}{2}|x - c|$$

the last step because $\eta' \leq \epsilon/(2(|B| + 1))$ and $|B| \leq |B| + 1$, so that $|B|\eta' \leq \epsilon/2$.

Step 4: conclusion. Adding the two displays through proposition 1.3.8(4), and using $f(d) = f(g(c))$,

$$|f(g(x)) - f(g(c)) - BA(x - c)| \leq \frac{\epsilon}{2}|x - c| + \frac{\epsilon}{2}|x - c| = \epsilon|x - c|$$

for every $x \in I$ with $|x - c| < \delta$. Since $\epsilon > 0$ was arbitrary, theorem 4.1.4 in the direction (2) implies (1), applied to $f \circ g$ on I at c with $L = BA$, shows that $f \circ g$ is differentiable at c with derivative $BA = f'(g(c))g'(c)$, completing the proof.

The proof never divided by $g(x) - g(c)$, and the hypothesis $g(I) \subseteq J$ is what let the outer approximation be evaluated at the point $y = g(x)$. For a worked instance, take $g: \mathbb{R} \rightarrow \mathbb{R}$ with $g(x) = 1 + x^2$ and $f: (0, \infty) \rightarrow \mathbb{R}$ with $f(y) = y^{1/2}$. Every value of g satisfies $g(x) \geq 1 > 0$, so $g(\mathbb{R}) \subseteq (0, \infty)$, and g is differentiable with $g'(x) = 0 + 2x = 2x$ by theorem 4.1.7(1) and (5) together with example 4.1.3(1). By example 4.1.3(4), f is differentiable at every point of $(0, \infty)$ with $f'(y) = 1/(2y^{1/2})$. So $h(x) = (1 + x^2)^{1/2}$ is differentiable on $\mathbb{R}$ with

$$h'(x) = \frac{1}{2(1 + x^2)^{1/2}} \cdot 2x = \frac{x}{(1 + x^2)^{1/2}}$$

The same h is continuous by corollary 3.3.13 and theorem 3.2.7, and it is now differentiable as well.

The last rule of the section takes the inverse of a function. Theorem 3.3.12 produced a continuous inverse for a continuous strictly monotone function on an interval, and what remains is its derivative. The chain rule already forces the formula: if g is the inverse of f and both are differentiable, then $g(f(x)) = x$ gives $g'(f(x))f'(x) = 1$. That argument presupposes g differentiable, which is half of what has to be proved. The theorem below establishes the differentiability of g and the formula together, from the sequential criterion.

Theorem 4.1.9: The Derivative of an Inverse Function

Let $I \subseteq \mathbb{R}$ be an interval with more than one point and let $f: I \rightarrow \mathbb{R}$ be continuous and strictly monotone. Put $J = f(I)$, let $c \in I$ and put $d = f(c)$. Suppose f is differentiable at c with $f'(c) \neq 0$. Then J is an interval with more than one point, the map $f: I \rightarrow J$ is a bijection, and its inverse $g: J \rightarrow I$ is differentiable at d with

$$g'(d) = \frac{1}{f'(c)}$$

Proof:

Step 1: the setting. By theorem 3.3.12, J is an interval, $f: I \rightarrow J$ is a bijection, and $g = f^{-1}$ is continuous and strictly monotone. Since I has two distinct points u and v , and f is injective, $f(u)$ and $f(v)$ are two distinct points of J . So J is an interval with more than one point, and definition 4.1.2 applies to g at $d \in J$.

Step 2: transferring a sequence. Let (y_n) be any sequence in J with $y_n \neq d$ for every n and $y_n \rightarrow d$. Put $x_n = g(y_n) \in I$. Since g is injective and $g(d) = c$, the condition $y_n \neq d$ gives $x_n \neq c$. Since g is continuous at d , proposition 3.2.4 gives $x_n \rightarrow g(d) = c$. So (x_n) is a sequence in $I \setminus \{c\}$ with $x_n \neq c$ and $x_n \rightarrow c$.

Step 3: the difference quotients are reciprocal. For each n , $f(x_n) = y_n$ and $f(c) = d$, so

$$q_{f,c}(x_n) = \frac{f(x_n) - f(c)}{x_n - c} = \frac{y_n - d}{x_n - c}$$

This is nonzero, because $y_n - d \neq 0$ by the hypothesis on (y_n) and $x_n - c \neq 0$ by Step 2. Its reciprocal is

$$\frac{x_n - c}{y_n - d} = \frac{g(y_n) - g(d)}{y_n - d} = q_{g,d}(y_n)$$

Step 4: passing to the limit. By theorem 3.1.19(1) applied to $q_{f,c}$ on $I \setminus \{c\}$, whose limit at c is $f'(c)$, the sequence $(q_{f,c}(x_n))$ converges to $f'(c)$. Every term is nonzero by Step 3 and the limit $f'(c)$ is nonzero by hypothesis, so theorem 2.1.6(4), applied to the constant sequence 1 over $(q_{f,c}(x_n))$, gives

$$q_{g,d}(y_n) = \frac{1}{q_{f,c}(x_n)} \rightarrow \frac{1}{f'(c)}$$

Step 5: conclusion. The sequence (y_n) was an arbitrary sequence in J with $y_n \neq d$ and $y_n \rightarrow d$, and d is a limit point of $J \setminus \{d\}$ by the argument opening this section applied to the interval J . So theorem 3.1.19(1), applied to $q_{g,d}$ on $J \setminus \{d\}$, gives $\lim_{y \rightarrow d} q_{g,d}(y) = 1/f'(c)$. That is, g is differentiable at d with $g'(d) = 1/f'(c)$, as we sought to show.

The hypothesis $f'(c) \neq 0$ cannot be dropped, and its failure is visible in the simplest case. Let $f(x) = x^3$ on $[0, \infty)$, which is continuous and strictly increasing by proposition 3.3.11, with image $[0, \infty)$ and inverse $g(y) = y^{1/3}$ by corollary 3.3.13. The function $x \mapsto x^3$ on $\mathbb{R}$ has derivative $3x^2$ by theorem 4.1.7(5), and proposition 3.1.23(1) applied to its difference quotient at 0, restricted to the subset $[0, \infty) \setminus \{0\}$, gives $f'(0) = 3 \cdot 0^2 = 0$. Take $y_n = 1/n^3$, a sequence in $(0, \infty)$. It converges to 0 by proposition 2.1.10(2) applied with $b_n = 1/n$, which is null by example 2.1.3(1), since $n \leq n^3$ gives $0 < 1/n^3 \leq 1/n$ by proposition 1.1.2(7). Then $g(y_n) = 1/n$ by the uniqueness clause of theorem 1.3.12, so

$$q_{g,0}(y_n) = \frac{1/n - 0}{1/n^3 - 0} = n^2$$

The sequence (n^2) is unbounded, since theorem 1.4.1(2) produces for each $M \in \mathbb{R}$ an $n \in \mathbb{N}$ with $n > M$, and then $n^2 \geq n > M$ because $n \geq 1$. An unbounded sequence diverges by proposition 2.1.5, so theorem 3.1.19(2) shows $q_{g,0}$ has no limit at 0 and g is not differentiable there.

Where the hypothesis does hold, the theorem gives derivatives that the definition reaches only through an algebraic identity chosen for the occasion. Fix an integer $n \geq 1$ and let $f(x) = x^n$ on $[0, \infty)$, which is strictly increasing by proposition 3.3.11 and continuous by example 3.2.6 together with proposition 3.3.1, with image $[0, \infty)$, and whose inverse is $r(y) = y^{1/n}$ by corollary 3.3.13. Let $d > 0$ and put $c = d^{1/n} > 0$. Then $f'(c) = nc^{n-1}$, by theorem 4.1.7(5) together with proposition 3.1.23(1), which transfers the limit of the difference quotient on $\mathbb{R} \setminus \{c\}$ to the subset $[0, \infty) \setminus \{c\}$, and this value is nonzero because $c > 0$. So r is differentiable at d with

$$r'(d) = \frac{1}{n(d^{1/n})^{n-1}}$$

For $n = 2$ this reads $r'(d) = 1/(2d^{1/2})$, which is what example 4.1.3(4) obtained by rationalizing a difference of square roots. The general case needs no such identity.

Exercise 4.1.1

1. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = x^3$ and let $c \in \mathbb{R}$. Compute $f'(c)$ directly from definition 4.1.2, by factoring $x^3 - c^3 = (x - c)(x^2 + xc + c^2)$ and evaluating the resulting limit with corollary 3.1.20. Do not use theorem 4.1.7(5).
2. Let I be an interval with more than one point, let $f: I \rightarrow \mathbb{R}$, let $c \in I$, and suppose L_1 and L_2 both satisfy condition (2) of theorem 4.1.4 for f at c . Prove that $L_1 = L_2$, without appealing to proposition 3.1.18. Deduce that the number $f'(c)$ is the only slope for which the approximation of theorem 4.1.4 holds.
3. Define $f: \mathbb{R} \rightarrow \mathbb{R}$ by $f(x) = x^2$ for $x \in \mathbb{Q}$ and $f(x) = 0$ for $x \notin \mathbb{Q}$. Prove that f is differentiable at 0 with $f'(0) = 0$, and that f is differentiable at no other point.
4. Let I be an interval with more than one point and let $f, g, h: I \rightarrow \mathbb{R}$ be differentiable at $c \in I$. Prove that fgh is differentiable at c with

$$(fgh)'(c) = f'(c)g(c)h(c) + f(c)g'(c)h(c) + f(c)g(c)h'(c)$$

5. Let $p: \mathbb{R} \rightarrow \mathbb{R}$ be $p(x) = (x^2 + 1)^5$. Use theorem 4.1.8 to compute $p'(x)$, naming the two functions being composed and checking that the image of the inner one lies in the domain of the outer one.
6. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = x^3 + x$. Prove that f is strictly increasing on $\mathbb{R}$, using the factorization $x^3 - y^3 = (x - y)(x^2 + xy + y^2)$ together with $x^2 + xy + y^2 = (x + y/2)^2 + 3y^2/4$. Then let g be the inverse furnished by theorem 3.3.12 and compute $g'(2)$.
7. Let I be an interval with more than one point, let $f: I \rightarrow \mathbb{R}$ be differentiable at $c \in I$ and suppose $f'(c) > 0$. Using theorem 4.1.4 with $\epsilon = f'(c)/2$, prove that there is a $\delta > 0$ such that $f(x) > f(c)$ for every $x \in I$ with $c < x < c + \delta$, and $f(x) < f(c)$ for every $x \in I$ with $c - \delta < x < c$.
8. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be $f(x) = x|x|$. Prove that f is differentiable on $\mathbb{R}$ with $f'(x) = 2|x|$, and deduce from example 4.1.6 and theorem 4.1.7(2) that $f''(0)$ does not exist.

4.2 The Mean Value Theorem

A derivative is a limit taken at a single point, and on its own it constrains the function only near that point. The passage from a hypothesis at every point to a conclusion across an interval runs through one observation. At an interior maximum of a differentiable function the difference quotients formed to the left are nonnegative and those formed to the right are nonpositive, so the limit of the quotients is 0. Theorem 3.3.7 produces such a maximum on a closed bounded interval, and subtracting a straight line converts the resulting statement into the mean value theorem. Once that theorem is available, a bound on f' across an interval becomes a bound on the increments of f , and the sign, the vanishing and the boundedness of f' each give a different reading of the same identity.

Two results stand slightly apart from that pattern. Darboux's theorem says that f' takes every value between two of its values, which is the intermediate value property of definition 3.3.4 for a function that need not be continuous anywhere. L'Hôpital's rule compares two functions through their derivatives and rests on the two-function form of the mean value theorem.

Throughout, I denotes an interval containing more than one point, so that lemma 4.1.1 makes every $x \in I$ a limit point of $I \setminus \{x\}$. So definition 3.1.15 is available at every point of I , which is what makes definition 4.1.2 a statement about every point of such an interval, the endpoints included.

Definition 4.2.1: Local Extremum

Let $E \subseteq \mathbb{R}$, let $f: E \rightarrow \mathbb{R}$ and let $p \in E$. Then f has a *local maximum* at p if there is a $\delta > 0$ such that

$$f(x) \leq f(p) \quad \text{for every } x \in E \text{ with } |x - p| < \delta$$

and a *local minimum* at p if there is a $\delta > 0$ such that $f(x) \geq f(p)$ for every such x . A point at which f has a local maximum or a local minimum is a *local extremum* of f .

Three instances fix the notion. The function $f(x) = x^2$ on $\mathbb{R}$ has a local minimum at 0, and any δ serves: for $x \neq 0$, proposition 1.1.2(6) gives $x^2 > 0$, while $0^2 = 0$ by proposition 1.1.2(3), so $x^2 \geq 0 = f(0)$ for every real x . The identity function $f(x) = x$ on $[0, 1]$ has a local minimum at 0 and a local maximum at 1, again with any δ , because $0 \leq x \leq 1$ for every x in the domain. Its difference quotient at any $p \in [0, 1]$ is

$$\frac{f(x) - f(p)}{x - p} = \frac{x - p}{x - p} = 1$$

for every $x \in [0, 1]$ with $x \neq p$, so $f'(p) = 1$ at every point, the two extrema included. A constant function on $\mathbb{R}$ has a local maximum and a local minimum at every point, and its difference quotient is the constant 0, so its derivative vanishes everywhere.

The identity function on $[0, 1]$ is the reason the theorem below carries a hypothesis about where p sits. At an endpoint of the domain there is only one side from which to approach, and the sign obtained from that side is contradicted by nothing. The hypothesis that rules this out is that some open interval around p lies inside the domain.

Theorem 4.2.2: The Derivative Vanishes at an Interior Extremum

Let $I \subseteq \mathbb{R}$ be an interval, let $f: I \rightarrow \mathbb{R}$ and let $p \in I$ be a point for which there is an $\eta > 0$ with $(p - \eta, p + \eta) \subseteq I$. If f is differentiable at p and p is a local extremum of f , then $f'(p) = 0$.

Proof:

Suppose first that f has a local maximum at p , and take $\delta_0 > 0$ from definition 4.2.1, so that $f(x) \leq f(p)$ for every $x \in I$ with $|x - p| < \delta_0$. Put $\delta = \min(\delta_0, \eta) > 0$ and write

$$D(x) = \frac{f(x) - f(p)}{x - p} \quad (x \in I, x \neq p)$$

so that $f'(p) = \lim_{x \rightarrow p} D(x)$ by definition 4.1.2.

Step 1: the two sign estimates. Let $x \in I$ satisfy $0 < |x - p| < \delta$. Then $|x - p| < \delta_0$, so $f(x) - f(p) \leq 0$. Suppose $x > p$, so $x - p > 0$. If $D(x)$ were positive, then proposition 1.1.2(5) applied to $0 < D(x)$ and the positive factor $x - p$ would give $0 = 0 \cdot (x - p) < D(x)(x - p) = f(x) - f(p)$, using proposition 1.1.2(3) for the first equality. That contradicts $f(x) - f(p) \leq 0$, so $D(x) \leq 0$. Suppose instead $x < p$, so $x - p < 0$. If $D(x)$ were negative, then proposition 1.1.2(5) applied to $D(x) < 0$ and the negative factor $x - p$ would give $0 = 0 \cdot (x - p) < D(x)(x - p) = f(x) - f(p)$, again a contradiction, so $D(x) \geq 0$.

Step 2: passing to the limit from each side. For each $n \in \mathbb{N}$ put

$$x_n = p + \frac{\delta}{n+1}, \quad y_n = p - \frac{\delta}{n+1}$$

The sequence $(\delta/(n+1))$ is null by the argument recorded before definition 4.2.1, so $x_n \rightarrow p$ and $y_n \rightarrow p$ by example 2.1.3(2) and theorem 2.1.6(1). For every n ,

$$0 < |x_n - p| = \frac{\delta}{n+1} \leq \frac{\delta}{2} < \delta \leq \eta$$

so $x_n \in (p - \eta, p + \eta) \subseteq I$ and $x_n \neq p$, and the same bound holds for y_n . In particular (x_n) is a sequence in $I \setminus \{p\}$ converging to p , so proposition 3.1.9(2) makes p a limit point of $I \setminus \{p\}$ and theorem 3.1.19(1) applies to D , giving

$$D(x_n) \rightarrow f'(p), \quad D(y_n) \rightarrow f'(p)$$

Step 1 gives $D(x_n) \leq 0$ and $D(y_n) \geq 0$ for every n , so proposition 2.1.8(2) gives $f'(p) \leq 0$ from the first sequence and $f'(p) \geq 0$ from the second. Hence $f'(p) = 0$.

Step 3: the minimum case. If f has a local minimum at p , then $-f$ has a local maximum at p , since $f(x) \geq f(p)$ says $-f(x) \leq -f(p)$. By theorem 4.1.7 the function $-f$ is differentiable at p with $(-f)'(p) = -f'(p)$, so the case already proved gives $-f'(p) = 0$ and therefore $f'(p) = 0$, completing the proof.

The implication runs one way only. A vanishing derivative does not produce an extremum: the function $f(x) = x^3$ on $\mathbb{R}$ has $f'(0) = 0$, and the factorization of $v^3 - u^3$ displayed after corollary 4.2.7 below makes f strictly increasing, so 0 is neither a local maximum nor a local minimum. The theorem gives a finite list of candidates for the location of an extremum, and that is what the next proof needs.

Theorem 4.2.3: Rolle's Theorem

Let $I \subseteq \mathbb{R}$ be an interval, let $f: I \rightarrow \mathbb{R}$ be continuous, and let $u, v \in I$ with $u < v$ and $f(u) = f(v)$. If f is differentiable at every point of (u, v) , then there is a $c \in (u, v)$ with $f'(c) = 0$.

Proof:

Since $u, v \in I$ and I is an interval, every z with $u < z < v$ lies in I by definition 3.1.1, so $[u, v] \subseteq I$. The restriction of f to $[u, v]$ is continuous by proposition 3.3.1, so theorem 3.3.7 provides $p, q \in [u, v]$ with

$$f(p) \leq f(x) \leq f(q) \quad \text{for every } x \in [u, v]$$

Step 1: the case where both extrema sit at the ends. Suppose $p \in \{u, v\}$ and $q \in \{u, v\}$. Since $f(u) = f(v)$, this forces $f(p) = f(q)$, and the displayed inequalities then give $f(x) = f(p)$ for every $x \in [u, v]$. Take $c = (u + v)(1 + 1)^{-1}$, which lies in (u, v) by proposition 1.1.2(8), and put $\eta = \min(c - u, v - c) > 0$. Because $c \in (u, v)$, the function f is differentiable at c , so its difference quotient $D(x) = (f(x) - f(c))/(x - c)$ on $I \setminus \{c\}$ has a limit at c . At every point x of $(c - \eta, c + \eta)$ other than c we have $x \in [u, v]$, hence $f(x) = f(c)$ and $D(x) = 0$. So D agrees with the constant function 0 on that punctured interval, and proposition 3.1.22(3) gives $f'(c) = 0$, the constant function having limit 0 by definition 3.1.15 with any δ .

Step 2: the case where an extremum sits strictly inside. Otherwise at least one of p, q lies in (u, v) . Suppose first that $q \in (u, v)$ and put $\eta = \min(q - u, v - q) > 0$. Then $(q - \eta, q + \eta) \subseteq (u, v) \subseteq I$, and every $x \in I$ with $|x - q| < \eta$ lies in (u, v) and therefore satisfies $f(x) \leq f(q)$. So q is a local maximum of f in the sense of definition 4.2.1, and f is differentiable at q because $q \in (u, v)$. Theorem 4.2.2 gives $f'(q) = 0$, and $c = q$ works. If instead $p \in (u, v)$, the same three sentences with the inequality reversed exhibit p as a local minimum, and theorem 4.2.2 gives $f'(p) = 0$, completing the proof.

Rolle's theorem is the mean value theorem for a function whose two endpoint values agree. The general case comes from subtracting the straight line through the two endpoints, which changes the derivative by a constant and leaves the hypotheses of Rolle's theorem in place.

Theorem 4.2.4: The Mean Value Theorem

Let $I \subseteq \mathbb{R}$ be an interval, let $f: I \rightarrow \mathbb{R}$ be continuous, and let $u, v \in I$ with $u < v$. If f is differentiable at every point of (u, v) , then there is a $c \in (u, v)$ with

$$f(v) - f(u) = f'(c)(v - u)$$

Proof:

Put $m = (f(v) - f(u))/(v - u)$, which is defined because $v - u \neq 0$, and define $h: I \rightarrow \mathbb{R}$ by

$$h(x) = f(x) + k(x), \quad k(x) = -f(u) - m(x - u)$$

The function k is a polynomial function of x , hence continuous on I by example 3.2.6 together with proposition 3.3.1, so h is continuous on I by theorem 3.2.5(1).

Step 1: the two endpoint values agree. Directly, $h(u) = f(u) - f(u) - m \cdot 0 = 0$, and

$$h(v) = f(v) - f(u) - m(v - u) = f(v) - f(u) - (f(v) - f(u)) = 0$$

so $h(u) = h(v)$.

Step 2: the derivative of the correction term. Let $t \in (u, v)$. For $x \in I$ with $x \neq t$, the difference quotient of k at t is

$$\frac{k(x) - k(t)}{x - t} = \frac{-m(x - u) + m(t - u)}{x - t} = \frac{-m(x - t)}{x - t} = -m$$

a constant function of x , whose limit at t is $-m$ by definition 3.1.15 with any δ . So k is differentiable at t with $k'(t) = -m$, and theorem 4.1.7 makes h differentiable at t with $h'(t) = f'(t) - m$.

Step 3: Rolle. The hypotheses of theorem 4.2.3 now hold for h on I with the same u and v , so there is a $c \in (u, v)$ with $h'(c) = 0$, which says $f'(c) = m$. Multiplying by $v - u$ gives $f'(c)(v - u) = f(v) - f(u)$, as we sought to show.

The number m in the proof is the slope of the line joining the two endpoints of the graph, and the theorem says that some tangent line has that slope. The reading used below is the reverse one. An inequality satisfied by f' at every point of (u, v) transfers to an inequality about $f(v) - f(u)$, because the location of c inside (u, v) is never used and only the bound on f' enters. The three corollaries after the next example apply that reading to the bounds $f' \geq 0$, $f' = 0$ and $|f'| \leq L$.

A second version compares two functions at once. Applying the mean value theorem separately to f and to g produces two points, and nothing forces them to coincide, so a statement relating f' and g' at a single point needs its own proof.

Theorem 4.2.5: Cauchy's Mean Value Theorem

Let $I \subseteq \mathbb{R}$ be an interval, let $f, g: I \rightarrow \mathbb{R}$ be continuous, and let $u, v \in I$ with $u < v$. If f and g are differentiable at every point of (u, v) , then there is a $c \in (u, v)$ with

$$(f(v) - f(u))g'(c) = (g(v) - g(u))f'(c)$$

Proof:

Define $h: I \rightarrow \mathbb{R}$ by

$$h(x) = (f(v) - f(u))g(x) - (g(v) - g(u))f(x)$$

The two coefficients are fixed real numbers, so h is continuous on I by theorem 3.2.5(1) and (2), and differentiable at every point $t \in (u, v)$ with

$$h'(t) = (f(v) - f(u))g'(t) - (g(v) - g(u))f'(t)$$

by theorem 4.1.7.

Step 1: the two endpoint values agree. Expanding,

$$h(u) = f(v)g(u) - f(u)g(u) - g(v)f(u) + g(u)f(u) = f(v)g(u) - g(v)f(u)$$

and

$$h(v) = f(v)g(v) - f(u)g(v) - g(v)f(v) + g(u)f(v) = g(u)f(v) - f(u)g(v)$$

The two right-hand sides are the same number, so $h(u) = h(v)$.

Step 2: Rolle. By theorem 4.2.3 there is a $c \in (u, v)$ with $h'(c) = 0$, which is the displayed identity, completing the proof.

Taking $g(x) = x$ recovers theorem 4.2.4, since the difference quotient of the identity function is the constant 1, so $g' \equiv 1$, and $g(v) - g(u) = v - u$. The next example shows that the two theorems are different statements, and that neither hypothesis of Rolle's theorem can be dropped.

Example 4.2.6: Three Failures Around the Mean Value Theorem

1. *Differentiability at a single interior point.* Let $f(x) = |x|$ on $[-1, 1]$. It is continuous by example 3.2.9, and $f(-1) = 1 = f(1)$. Fix $x \in (0, 1)$ and put $\eta = \min(x, 1 - x) > 0$. For $t \in (0, 1)$ with $t \neq x$ the difference quotient of f at x is $(t - x)/(t - x) = 1$, so it agrees with the constant function 1 at every point of $(x - \eta, x + \eta)$ other than x , and proposition 3.1.22(3) gives $f'(x) = 1$. The same argument on $(-1, 0)$, where the quotient is $(-t + x)/(t - x) = -1$, gives $f'(x) = -1$ for $x \in (-1, 0)$. At 0 the function is not differentiable (example 4.1.6). So there is no $c \in (-1, 1)$ at which $f'(c) = 0$, and the conclusion of theorem 4.2.3 fails. The single point 0 is the whole obstruction.
2. *Continuity at a single endpoint.* Define $f: [0, 1] \rightarrow \mathbb{R}$ by $f(x) = x$ for $0 \leq x < 1$ and $f(1) = 0$. Then $f(0) = 0 = f(1)$, and for every $x \in (0, 1)$ the difference quotient computation of item (1) gives $f'(x) = 1$. But f is not continuous at 1. The set $E_1^- = [0, 1] \cap (-\infty, 1) = [0, 1)$ carries the restriction $f(x) = x$, and it has 1 as a limit point by the mirrored form of proposition 3.1.25(1), since $1 - \min(\delta, 1)/2$ lies in $[0, 1)$ and within δ of 1 for every $\delta > 0$. For $\epsilon > 0$ the choice $\delta = \epsilon$ makes $|f(x) - 1| = |x - 1| < \epsilon$ for every $x \in [0, 1]$ with $1 - \delta < x < 1$, so proposition 3.1.25(2) gives $\lim_{x \rightarrow 1^-} f(x) = 1$. Since 1 is not a limit point of $E_1^+ = \emptyset$, proposition 3.1.26(2) makes $\lim_{x \rightarrow 1} f(x) = 1 \neq 0 = f(1)$, and proposition 3.2.3(1) then denies continuity at 1. No $c \in (0, 1)$ has $f'(c) = 0$, and no c satisfies $f(1) - f(0) = f'(c)(1 - 0)$ either, since the left side is 0 and the right side is 1.
3. *Cauchy's theorem is not two applications of the mean value theorem.* Let $f(x) = x^2$ and

$g(x) = x^3$ on $[0, 1]$. The identity function on $[0, 1]$ has derivative 1 at every point, as computed after definition 4.2.1, so the product rule of theorem 4.1.7 gives

$$f'(t) = t \cdot 1 + 1 \cdot t = 2t, \quad g'(t) = f'(t)t + f(t) \cdot 1 = 2t^2 + t^2 = 3t^2$$

The mean value theorem for f on $[0, 1]$ asks for $1 = f(1) - f(0) = 2c$, whose only solution is $c = 1/2$. For g it asks for $1 = 3c^2$, that is $c^2 = 1/3$, whose only nonnegative solution is $c = \sqrt{1/3}$ by theorem 1.3.12. Cauchy's theorem asks for $1 \cdot 3c^2 = 1 \cdot 2c$, that is $c(3c - 2) = 0$, whose only root in $(0, 1)$ is $c = 2/3$ by proposition 1.1.2(3). The three numbers are distinct: multiplying by 6 turns $1/2 = 2/3$ into $3 = 4$, and squaring turns $2/3 = \sqrt{1/3}$ into $4/9 = 3/9$.

Corollary 4.2.7: Monotonicity From the Sign of the Derivative

Let $I \subseteq \mathbb{R}$ be an interval containing more than one point and let $f: I \rightarrow \mathbb{R}$ be differentiable at every point of I .

1. If $f'(x) \geq 0$ for every $x \in I$, then f is increasing.
2. If $f'(x) > 0$ for every $x \in I$, then f is strictly increasing.
3. If $f'(x) \leq 0$ for every $x \in I$, then f is decreasing, and if $f'(x) < 0$ for every $x \in I$, then f is strictly decreasing.
4. Conversely, if f is increasing then $f'(x) \geq 0$ for every $x \in I$, and if f is decreasing then $f'(x) \leq 0$ for every $x \in I$.

Proof:

By proposition 4.1.5 the function f is continuous at every point of I , hence continuous on I in the sense of definition 3.2.2.

Parts (1) to (3). Let $u, v \in I$ with $u < v$. Every point of (u, v) lies in I by definition 3.1.1, so f is differentiable at every point of (u, v) , and theorem 4.2.4 gives a $c \in (u, v)$ with $f(v) - f(u) = f'(c)(v - u)$. The factor $v - u$ is positive. If $f'(c) > 0$ then proposition 1.1.2(5) applied to $0 < f'(c)$ gives $0 = 0 \cdot (v - u) < f'(c)(v - u) = f(v) - f(u)$, so $f(u) < f(v)$. Under the weaker hypothesis $f'(c) \geq 0$ the same computation gives $f(u) \leq f(v)$, the case $f'(c) = 0$ producing equality. The two decreasing cases are the same computation with the inequalities reversed. Since $u < v$ were arbitrary and the case $u = v$ is an equality, this is definition 3.2.14 in each of the four statements.

Part (4). Suppose f is increasing and let $x \in I$. Write $D(t) = (f(t) - f(x))/(t - x)$ for $t \in I$ with $t \neq x$. If $t > x$ then $f(t) - f(x) \geq 0$ and $t - x > 0$. Were $D(t)$ negative, proposition 1.1.2(5) applied to $D(t) < 0$ and the positive factor $t - x$ would give $f(t) - f(x) = D(t)(t - x) < 0$, so $D(t) \geq 0$. If $t < x$ then $f(t) - f(x) \leq 0$ and $t - x < 0$, and the same argument with the negative factor again gives $D(t) \geq 0$. Choose $y \in I$ with $y \neq x$, which exists because I has more than one point, and put $t_n = x + (y - x)/(n + 1)$. As recorded before definition 4.2.1, each t_n lies in $I \setminus \{x\}$ and $t_n \rightarrow x$, so theorem 3.1.19(1) gives $D(t_n) \rightarrow f'(x)$. Every $D(t_n)$ is nonnegative, so proposition 2.1.8(2) gives $f'(x) \geq 0$. For a decreasing f , apply what was just proved to $-f$, which is increasing and has derivative $-f'$ by theorem 4.1.7, completing the proof.

Parts (2) and (4) do not combine into an equivalence, and the failure is one-directional. Strict monotonicity does not force $f' > 0$. For $f(x) = x^3$ on $\mathbb{R}$ and $u < v$,

$$v^3 - u^3 = (v - u) \left(\left(v + \frac{u}{2} \right)^2 + \frac{3}{4}u^2 \right)$$

as multiplying out confirms, and the bracket is a sum of two squares that vanishes only when $u = 0$ and $v = 0$, which $u < v$ excludes. So $v^3 > u^3$ and f is strictly increasing, while the product-rule computation of example 4.2.6(3), carried out on $\mathbb{R}$, gives $f'(x) = 3x^2$ and hence $f'(0) = 0$.

Corollary 4.2.8: A Vanishing Derivative Forces a Constant

Let $I \subseteq \mathbb{R}$ be an interval containing more than one point.

1. If $f: I \rightarrow \mathbb{R}$ is differentiable at every point of I with $f'(x) = 0$ for every $x \in I$, then f is constant.
2. If $f, g: I \rightarrow \mathbb{R}$ are differentiable at every point of I with $f'(x) = g'(x)$ for every $x \in I$, then $f - g$ is constant. If in addition $f(x_0) = g(x_0)$ for some $x_0 \in I$, then $f = g$.

Proof:

1. The hypothesis gives both $f'(x) \geq 0$ and $f'(x) \leq 0$ for every $x \in I$, so f is increasing by corollary 4.2.7(1) and decreasing by part (3) of the same corollary. For $u, v \in I$ with $u < v$ this yields $f(u) \leq f(v)$ and $f(u) \geq f(v)$, hence $f(u) = f(v)$. Any two distinct points of I stand in that relation once the smaller is named u , so f takes one value on I .
2. By theorem 4.1.7 the function $f - g$ is differentiable at every point of I with $(f - g)'(x) = f'(x) - g'(x) = 0$, so part (1) makes $f - g$ constant. If $f(x_0) = g(x_0)$ then that constant is $f(x_0) - g(x_0) = 0$, so $f(x) = g(x)$ for every $x \in I$, completing the proof.

Part (2) is what makes an antiderivative unique up to an additive constant. Part (1) is what Section 4.6 rests on: the identity $\sin^2 + \cos^2 = 1$ and the addition formulas for $\sin$ and $\cos$ are each proved there by exhibiting a function whose derivative vanishes on all of $\mathbb{R}$ and evaluating it at 0.

Section 3.4 defined the Lipschitz condition (definition 3.4.8) and proved that it implies uniform continuity, and it stopped there because it had no way to verify the condition for a given function. A bound on the derivative is that way.

Corollary 4.2.9: A Bounded Derivative Makes a Function Lipschitz

Let $I \subseteq \mathbb{R}$ be an interval, let $f: I \rightarrow \mathbb{R}$ be differentiable at every point of I , and let $L \geq 0$ satisfy $|f'(x)| \leq L$ for every $x \in I$. Then f is Lipschitz on I with Lipschitz constant L , and therefore uniformly continuous on I .

Proof:

Let $x, y \in I$. If $x = y$ then both sides of the Lipschitz inequality are 0 and there is nothing to check. Both $|f(x) - f(y)|$ and $|x - y|$ are unchanged when x and y are exchanged, by proposition 1.3.8(1), so assume $x < y$.

By proposition 4.1.5 the function f is continuous on I , and it is differentiable at every point of

(x, y) , which lies in I by definition 3.1.1. So theorem 4.2.4 gives a $c \in (x, y)$ with $f(y) - f(x) = f'(c)(y - x)$. Taking absolute values and using proposition 1.3.8(3),

$$|f(y) - f(x)| = |f'(c)| |y - x| \leq L|y - x|$$

the inequality being $|f'(c)| \leq L$ multiplied by the nonnegative number $|y - x|$. This is definition 3.4.8 with constant L , and proposition 3.4.9 then makes f uniformly continuous on I , as we sought to show.

A derivative can fail to be continuous. Example 4.6.17, which waits for Section 4.6 because it is built from $\sin$, gives a function differentiable at every point of $\mathbb{R}$ whose derivative is discontinuous at the origin. That the discontinuity is of the second kind is not special to this example, and it is proved below for every derivative. So the derivatives on an interval are not the continuous functions on it. They are also not all functions on it, and the reason is that the argument used for Rolle's theorem applies to every function of the form $f(t) - \lambda t$.

Theorem 4.2.10: Darboux's Theorem

Let $I \subseteq \mathbb{R}$ be an interval containing more than one point and let $f: I \rightarrow \mathbb{R}$ be differentiable at every point of I . Then $f': I \rightarrow \mathbb{R}$ has the intermediate value property of definition 3.3.4: for all $u, v \in I$ with $u < v$ and every $\lambda \in \mathbb{R}$ with $\min\{f'(u), f'(v)\} < \lambda < \max\{f'(u), f'(v)\}$, there is a $c \in (u, v)$ with $f'(c) = \lambda$.

Proof:

Fix $u < v$ in I and such a λ . As in the proof of theorem 4.2.3, $[u, v] \subseteq I$ by definition 3.1.1. Suppose first that $f'(u) < \lambda < f'(v)$, and define $g: I \rightarrow \mathbb{R}$ by $g(t) = f(t) - \lambda t$. The difference quotient of $t \mapsto \lambda t$ at any point is the constant λ , so that function has derivative λ everywhere and theorem 4.1.7 makes g differentiable at every point of I with $g'(t) = f'(t) - \lambda$. In particular $g'(u) < 0 < g'(v)$. By proposition 4.1.5 and proposition 3.3.1 the restriction of g to $[u, v]$ is continuous, so theorem 3.3.7 provides a $p \in [u, v]$ with $g(p) \leq g(t)$ for every $t \in [u, v]$.

Step 1: the minimum is not at u . The difference quotient $Q(t) = (g(t) - g(u))/(t - u)$, defined for $t \in I$ with $t \neq u$, has limit $g'(u)$ at u . Apply definition 3.1.15 with $\epsilon = -g'(u) > 0$: there is a $\delta > 0$ such that $|Q(t) - g'(u)| < -g'(u)$ for every $t \in I$ with $0 < |t - u| < \delta$, and proposition 1.3.8(2) turns this into $Q(t) < g'(u) + (-g'(u)) = 0$. Take $t_1 = u + \min(\delta, v - u)/2$, so that $u < t_1 < v$ and $0 < t_1 - u < \delta$, whence $Q(t_1) < 0$. Were $g(t_1) - g(u)$ nonnegative, proposition 1.1.2(5) applied to $Q(t_1) < 0$ and the positive factor $t_1 - u$ would give $g(t_1) - g(u) = Q(t_1)(t_1 - u) < 0$. So $g(t_1) < g(u)$, and $p \neq u$.

Step 2: the minimum is not at v . The difference quotient $R(t) = (g(t) - g(v))/(t - v)$ has limit $g'(v) > 0$ at v . Applying definition 3.1.15 with $\epsilon = g'(v)$ gives a $\delta' > 0$ such that $R(t) > g'(v) - g'(v) = 0$ for every $t \in I$ with $0 < |t - v| < \delta'$. Take $t_2 = v - \min(\delta', v - u)/2$, so that $u < t_2 < v$ and $0 < v - t_2 < \delta'$, whence $R(t_2) > 0$. Since $t_2 - v < 0$, proposition 1.1.2(5) applied to $0 < R(t_2)$ and the negative factor $t_2 - v$ gives $g(t_2) - g(v) = R(t_2)(t_2 - v) < 0$. So $g(t_2) < g(v)$, and $p \neq v$.

Step 3: the interior extremum. By Steps 1 and 2, $p \in (u, v)$. Put $\eta = \min(p - u, v - p) > 0$,

so that $(p - \eta, p + \eta) \subseteq (u, v) \subseteq I$ and every $x \in I$ with $|x - p| < \eta$ lies in $[u, v]$ and satisfies $g(p) \leq g(x)$. So p is a local minimum of g , and g is differentiable at p . By theorem 4.2.2, $g'(p) = 0$, which is $f'(p) = \lambda$. Take $c = p$.

Step 4: the reversed case. Suppose instead $f'(v) < \lambda < f'(u)$. By theorem 4.1.7 the function $-f$ is differentiable at every point of I with $(-f)' = -f'$, and proposition 1.1.2(6) turns the two inequalities into $(-f)'(u) < -\lambda < (-f)'(v)$. The case already proved gives a $c \in (u, v)$ with $(-f)'(c) = -\lambda$, that is $f'(c) = \lambda$. The two cases exhaust the hypothesis on λ , completing the proof.

One consequence is a restriction on how a derivative may fail to be continuous. Let $a < b$, let $f: (a, b) \rightarrow \mathbb{R}$ be differentiable at every point, and let $z \in (a, b)$ be a discontinuity of f' of the first kind in the sense of definition 3.2.13, so that $\alpha = \lim_{x \rightarrow z^-} f'(x)$ and $\beta = \lim_{x \rightarrow z^+} f'(x)$ both exist in $\mathbb{R}$. Suppose $\alpha \neq f'(z)$. Replacing f by $-f$ if necessary, which by theorem 4.1.7 replaces f' by $-f'$ and α by $-\alpha$, assume $\alpha < f'(z)$, and choose λ with $\alpha < \lambda < f'(z)$ using proposition 1.1.2(8). By proposition 3.1.25(2) with $\epsilon = \lambda - \alpha$ there is a $\delta > 0$ with $f'(x) < \lambda$ for every $x \in (a, b)$ satisfying $z - \delta < x < z$. Fix such an x and call it u . Applying theorem 4.2.10 on (a, b) to the pair $u < z$ produces a $c \in (u, z)$ with $f'(c) = \lambda$, while $z - \delta < u < c < z$ forces $f'(c) < \lambda$. So $\alpha = f'(z)$. The same four sentences with the right-hand limit, with the mirrored form of proposition 3.1.25(2), and with theorem 4.2.10 applied to a pair $z < v$ inside $(z, z + \delta)$ give $\beta = f'(z)$. Then proposition 3.1.26(1) gives $\lim_{x \rightarrow z} f'(x) = f'(z)$, so f' is continuous at z by proposition 3.2.3(1), contradicting the choice of z . Every discontinuity of a derivative is therefore of the second kind, and example 4.6.17 exhibits one.

A quotient $f(x)/g(x)$ whose numerator and denominator both tend to 0 is not determined by those two limits. Cauchy's mean value theorem rewrites such a quotient as a quotient of derivatives at a single unknown point, and that point can be confined to an arbitrarily short interval by shrinking the range of x . The following theorem is that argument.

Theorem 4.2.11: L'Hôpital's Rule

Let $a < b$ be real numbers and let $f, g: (a, b) \rightarrow \mathbb{R}$ be differentiable at every point of (a, b) , with $g'(x) \neq 0$ for every $x \in (a, b)$. Let $A \in \mathbb{R}$ and suppose

$$\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} = A$$

Suppose in addition that at least one of the following holds:

1. $\lim_{x \rightarrow a^+} f(x) = 0$ and $\lim_{x \rightarrow a^+} g(x) = 0$;
2. $\lim_{x \rightarrow a^+} g(x) = +\infty$.

Then there is a $c_0 \in (a, b]$ with $g(x) \neq 0$ for every $x \in (a, c_0)$, and the quotient f/g formed on (a, c_0) satisfies

$$\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = A$$

Every domain in the statement is a subinterval of (a, ∞) , so the set E_a^+ of definition 3.1.24 is in each case the domain itself, and each one-sided limit above is the limit of definition 3.1.15 taken over that

domain at the limit point a . The proof uses that reading throughout.

Proof:

By proposition 4.1.5, f and g are continuous on (a, b) .

Step 1: the increment quotient is a quotient of derivatives. Let $x, y \in (a, b)$ with $x < y$. If $g(x) = g(y)$, then theorem 4.2.3 applied to g on (a, b) with these two points gives a $t \in (x, y)$ with $g'(t) = 0$, which the hypothesis forbids. So $g(y) - g(x) \neq 0$. By theorem 4.2.5 applied to f and g on (a, b) with these two points there is a $t \in (x, y)$ with $(f(y) - f(x))g'(t) = (g(y) - g(x))f'(t)$, and dividing by the nonzero numbers $g'(t)$ and $g(y) - g(x)$ gives

$$\frac{f(y) - f(x)}{g(y) - g(x)} = \frac{f'(t)}{g'(t)} \quad \text{for some } t \in (x, y) \quad (4.3)$$

Step 2: a uniform estimate near a . Let $\epsilon > 0$. By the hypothesis on f'/g' and definition 3.1.15 there is a $\delta > 0$ with $|f'(t)/g'(t) - A| < \epsilon/2$ for every $t \in (a, b)$ with $t < a + \delta$. Put $c = \min(a + \delta, (a + b)(1 + 1)^{-1})$, so that $a < c < b$ by proposition 1.1.2(8) and every point of (a, c) lies in (a, b) and is below $a + \delta$. For $a < x < y < c$ the point t produced by (4.3) lies in $(x, y) \subseteq (a, c)$, so

$$A - \frac{\epsilon}{2} < \frac{f(y) - f(x)}{g(y) - g(x)} < A + \frac{\epsilon}{2} \quad (a < x < y < c) \quad (4.4)$$

by proposition 1.3.8(2). Multiplying numerator and denominator by -1 shows the middle term equals $(f(x) - f(y))/(g(x) - g(y))$, and both forms are used below.

Step 3: case (1), where g never vanishes. Assume f and g tend to 0 at a^+ . Each value $g'(x)$ is nonzero, hence positive or negative. Suppose g' took a negative value at one point of (a, b) and a positive value at another, and name the smaller of the two points u and the larger v . Then 0 lies strictly between $\min\{g'(u), g'(v)\}$ and $\max\{g'(u), g'(v)\}$, so theorem 4.2.10 would produce a $w \in (u, v)$ with $g'(w) = 0$, which the hypothesis forbids. So either $g'(x) > 0$ for every $x \in (a, b)$ or $g'(x) < 0$ for every $x \in (a, b)$, and corollary 4.2.7(2) or (3) makes g strictly increasing or strictly decreasing on (a, b) .

Suppose g is strictly increasing, and suppose some $x_0 \in (a, b)$ had $g(x_0) < 0$. Applying definition 3.1.15 to $\lim_{x \rightarrow a^+} g(x) = 0$ with $\epsilon = -g(x_0) > 0$ gives a $\delta_1 > 0$ with $|g(x)| < -g(x_0)$ for every $x \in (a, b)$ with $x < a + \delta_1$. The point $x_1 = a + \min(\delta_1, x_0 - a)/2$ satisfies $a < x_1 < x_0 < b$ and $x_1 < a + \delta_1$, so $g(x_0) < g(x_1)$ by proposition 1.3.8(2), while $x_1 < x_0$ and strict monotonicity give $g(x_1) < g(x_0)$. That is impossible, so $g(x) \geq 0$ for every $x \in (a, b)$. Now for any $y \in (a, b)$, choosing $x_0 = (a + y)(1 + 1)^{-1} \in (a, y)$ gives $g(y) > g(x_0) \geq 0$. If instead g is strictly decreasing, apply this argument to $-g$, which has derivative $-g' > 0$ by theorem 4.1.7 and is therefore strictly increasing by corollary 4.2.7(2), and which tends to 0 at a^+ by corollary 3.1.20(2), to get $-g(y) > 0$. Either way $g(y) \neq 0$ for every $y \in (a, b)$, so $c_0 = b$ serves.

Fix $y \in (a, c)$ and consider $H(x) = (f(y) - f(x))/(g(y) - g(x))$ on (a, y) , which is defined by Step 1. The points $x_n = a + (y - a)/(n + 1)$ lie in (a, y) and converge to a , by the argument recorded before definition 4.2.1, so proposition 3.1.9(2) makes a a limit point of (a, y) . By proposition 3.1.23(1) the restrictions of f and g to (a, y) still tend to 0 at a , and a constant function has itself as its limit by definition 3.1.15 with any δ , so corollary 3.1.20(1) and (2) give $\lim_{x \rightarrow a} (f(y) - f(x)) = f(y)$ and $\lim_{x \rightarrow a} (g(y) - g(x)) = g(y)$. Since $g(y) \neq 0$, corollary 3.1.20(4)

gives $\lim_{x \rightarrow a} H(x) = f(y)/g(y)$, and theorem 3.1.19(1) makes $H(x_n) \rightarrow f(y)/g(y)$. Each $H(x_n)$ lies between $A - \epsilon/2$ and $A + \epsilon/2$ by (4.4), so proposition 2.1.8(2) gives

$$A - \frac{\epsilon}{2} \leq \frac{f(y)}{g(y)} \leq A + \frac{\epsilon}{2}$$

and therefore $|f(y)/g(y) - A| \leq \epsilon/2 < \epsilon$ by proposition 1.3.8(2). This holds for every $y \in (a, c)$, so $c - a$ answers ϵ in definition 3.1.15 for f/g at a . As $\epsilon > 0$ was arbitrary, $\lim_{x \rightarrow a^+} f(x)/g(x) = A$.

Step 4: case (2), where g tends to $+\infty$. Assume $\lim_{x \rightarrow a^+} g(x) = +\infty$. Applying definition 3.1.28 with $M = 0$ gives a $\delta_2 > 0$ with $g(x) > 0$ for every $x \in (a, b)$ with $x < a + \delta_2$, so $c_0 = \min(a + \delta_2, b)$ has the property required in the statement. Fix $y \in (a, c)$, with c as in Step 2, and apply definition 3.1.28 with $M = \max\{g(y), 0\}$ to get a $\delta_3 > 0$ with $g(x) > g(y)$ and $g(x) > 0$ for every $x \in (a, b)$ with $x < a + \delta_3$. Put $c_1 = \min(a + \delta_3, y)$.

Let $x \in (a, c_1)$ and put $\mu(x) = (g(x) - g(y))/g(x)$, which is positive because $g(x) - g(y) > 0$ and $g(x) > 0$. Multiplying the two inequalities of (4.4), in the form using $(f(x) - f(y))/(g(x) - g(y))$, by the positive number $\mu(x)$ and adding $f(y)/g(x)$ gives

$$\left(A - \frac{\epsilon}{2}\right)\mu(x) + \frac{f(y)}{g(x)} < \frac{f(x)}{g(x)} < \left(A + \frac{\epsilon}{2}\right)\mu(x) + \frac{f(y)}{g(x)}$$

the first step by proposition 1.1.2(5) and the second by the order axiom of definition 1.1.1 that adding a fixed element preserves a strict inequality.

Write $\varphi(x) = 1/g(x)$ on (a, c_1) . Then φ tends to 0 at a : given $\eta > 0$, definition 3.1.28 with $M = 1/\eta$ gives a $\delta_4 > 0$ with $g(x) > 1/\eta > 0$ for every $x \in (a, c_1)$ with $x < a + \delta_4$, and proposition 1.1.2(7) inverts this to $0 < \varphi(x) < \eta$. Since $\mu(x) = 1 - g(y)\varphi(x)$, the two bounding expressions can be rewritten as

$$\left(A \mp \frac{\epsilon}{2}\right) + \left(f(y) - \left(A \mp \frac{\epsilon}{2}\right)g(y)\right)\varphi(x)$$

a constant plus a constant multiple of φ , so corollary 3.1.20(1) and (2) give them the limits $A - \epsilon/2$ and $A + \epsilon/2$ at a . Applying definition 3.1.15 to each with tolerance $\epsilon/2$ produces $\delta_5, \delta_6 > 0$ such that the lower expression exceeds $A - \epsilon$ for $x \in (a, c_1)$ with $x < a + \delta_5$, and the upper expression falls below $A + \epsilon$ for $x \in (a, c_1)$ with $x < a + \delta_6$.

Put $\delta_\epsilon = \min(\delta_5, \delta_6, c_1 - a) > 0$. For $x \in (a, a + \delta_\epsilon)$ the displayed chain gives $A - \epsilon < f(x)/g(x) < A + \epsilon$, hence $|f(x)/g(x) - A| < \epsilon$ by proposition 1.3.8(2). As $\epsilon > 0$ was arbitrary, this is definition 3.1.15 for f/g at a and gives $\lim_{x \rightarrow a^+} f(x)/g(x) = A$, completing the proof.

The mirror image of the theorem, at the right-hand endpoint, follows by reflecting the line. For $E \subseteq \mathbb{R}$ write $-E = \{-x | x \in E\}$, and for $h: E \rightarrow \mathbb{R}$ let $\tilde{h}: -E \rightarrow \mathbb{R}$ be $\tilde{h}(t) = h(-t)$. For $t \in -E$ and $c \in \mathbb{R}$,

$$|(-t) - c| = |-(t + c)| = |t - (-c)|$$

by proposition 1.3.8(1), while $|\tilde{h}(t) - L| = |h(-t) - L|$. So each instance of definition 3.1.15 for h at c is the corresponding instance for $\tilde{h}$ at $-c$ under the substitution $x = -t$, and the same holds for definition 3.1.28. Hence $-c$ is a limit point of $-E$ exactly when c is a limit point of E , and $\lim_{x \rightarrow c} h(x)$ and $\lim_{t \rightarrow -c} \tilde{h}(t)$ exist together and are equal, whether the common value is real or $\pm\infty$.

Applied to a difference quotient this identifies the reflected derivative. If $f: (a, b) \rightarrow \mathbb{R}$ is differentiable

at $-t_0$ for $t_0 \in (-b, -a)$, then for $t \in (-b, -a)$ with $t \neq t_0$,

$$\frac{\tilde{f}(t) - \tilde{f}(t_0)}{t - t_0} = \frac{f(-t) - f(-t_0)}{t - t_0} = - \frac{f(-t) - f(-t_0)}{(-t) - (-t_0)}$$

so $\tilde{f}$ is differentiable at t_0 with $\tilde{f}'(t_0) = -f'(-t_0)$, by corollary 3.1.20(2) for the factor -1 and the reflection just established for the substitution. Consequently $\tilde{f}'(t)/\tilde{g}'(t) = f'(-t)/g'(-t)$ and $\tilde{g}'(t) \neq 0$. A pair f, g on (a, b) whose data satisfy the hypotheses of theorem 4.2.11 at b^- therefore transports to the pair $\tilde{f}, \tilde{g}$ on $(-b, -a)$ satisfying them at $(-b)^+$, and the conclusion transports back to give $\lim_{x \rightarrow b^-} f(x)/g(x) = A$.

Both endpoint forms together give the rule at an interior point. Let $a < p < b$, let f and g be differentiable at every point of (a, p) and of (p, b) with g' nowhere zero on either, and suppose $\lim_{x \rightarrow p^-} f'(x)/g'(x) = \lim_{x \rightarrow p^+} f'(x)/g'(x) = A$ with hypothesis (1) or (2) holding on each side. Then both one-sided limits of f/g at p equal A , and proposition 3.1.26(1) gives $\lim_{x \rightarrow p} f(x)/g(x) = A$.

Hypothesis (1) cannot be dropped. Let $f(x) = x + 1$ and $g(x) = x + 2$ on $(0, 1)$. The difference quotient of each at any point is the constant 1, so $f'(x) = g'(x) = 1$ for every x , and g' never vanishes while $f'(x)/g'(x) = 1$ gives $\lim_{x \rightarrow 0^+} f'(x)/g'(x) = 1$. The quotient of the functions behaves differently. For $\epsilon > 0$ the choice $\delta = \epsilon$ gives $|f(x) - 1| = |x| < \epsilon$ and $|g(x) - 2| = |x| < \epsilon$ for every $x \in (0, 1)$ with $0 < x < \delta$, so $\lim_{x \rightarrow 0^+} f(x) = 1$ and $\lim_{x \rightarrow 0^+} g(x) = 2$, and corollary 3.1.20(4) gives $\lim_{x \rightarrow 0^+} f(x)/g(x) = 1/2$. Neither alternative of the theorem holds here, since f and g tend to 1 and 2 at 0^+ and g does not tend to $+\infty$, and the two limits differ.

Exercise 4.2.1

1. Let $f: [0, 1] \rightarrow \mathbb{R}$ be continuous and differentiable at every point of $(0, 1)$, with $f(0) = 0$ and $f'(x) \leq 1$ for every $x \in (0, 1)$. Show that $f(x) \leq x$ for every $x \in [0, 1]$.
2. Let $p(x) = x^3 - 3x + 1$. Show that p has at least three distinct real roots by evaluating p at $-2, 0, 1$ and 2 and applying theorem 3.3.2. Then show that p has at most three distinct real roots, by supposing it had four and applying theorem 4.2.3 between consecutive roots.
3. Let $f: [0, \infty) \rightarrow \mathbb{R}$ be differentiable at every point of $[0, \infty)$, with $f(0) = 0$, and suppose f' is increasing on $[0, \infty)$. Show that $x \mapsto f(x)/x$ is increasing on $(0, \infty)$. Use theorem 4.2.4 on $[0, x]$ to compare $f(x)$ with $xf'(x)$, and then differentiate the quotient.
4. Let $a < b$, let $f, g: [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable at every point of (a, b) , and suppose $g'(x) \neq 0$ for every $x \in (a, b)$. Show first that $g(a) \neq g(b)$, and then that there is a $c \in (a, b)$ with

$$\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}$$

5. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be differentiable at every point and suppose $f'(x) \in \{0, 1\}$ for every $x \in \mathbb{R}$. Show that f' is constant, and deduce that either f is constant or there is a $k \in \mathbb{R}$ with $f(x) = x + k$ for every $x \in \mathbb{R}$.
6. Define $h: \mathbb{R} \rightarrow \mathbb{R}$ by $h(x) = x$ for $x < 0$ and $h(x) = x + 1$ for $x \geq 0$. Show that no differentiable $F: \mathbb{R} \rightarrow \mathbb{R}$ has $F' = h$.
7. Let $f, g: (0, 1) \rightarrow \mathbb{R}$ be given by $f(x) = 1/x + 1$ and $g(x) = 1/x$. Verify every hypothesis of theorem 4.2.11 under its alternative (2), compute $\lim_{x \rightarrow 0^+} f'(x)/g'(x)$, and check the conclusion against a direct computation of $f(x)/g(x)$.

8. Let $a < b$, let $f: [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable at every point of (a, b) , and suppose $\lim_{x \rightarrow a^+} f'(x) = L$ for some $L \in \mathbb{R}$. Show that f is differentiable at a with $f'(a) = L$. Apply theorem 4.2.4 on $[a, x]$ and note where the intermediate point lies.

4.3 Taylor's Theorem

The mean value theorem controls a function across an interval from a single value of its derivative. When f has n derivatives at a point a , more local information is on hand, and this section converts it into a statement of the same shape: the polynomial built from $f(a), f'(a), \dots, f^{(n-1)}(a)$ approximates f near a , and the error is the next term of that polynomial with the derivative evaluated at an unknown point between a and x . The mean value theorem is the case $n = 1$. A bound on that one higher derivative becomes a bound on the approximation error, and section 4.7 uses exactly that to decide when a function is the sum of the infinite series these polynomials approach.

A polynomial is determined by its coefficients, and its coefficients are determined by its derivatives at any one point: differentiating $\sum_j b_j(x-a)^j$ k times and evaluating at a kills every term except the k -th, which leaves $k!b_k$. In the other direction, the derivatives of an arbitrary f at a name a polynomial, and that polynomial is the one whose derivatives at a match those of f as far as they reach.

Definition 4.3.1: Taylor Polynomial

Let $I \subseteq \mathbb{R}$ be an interval, let $a \in I$, let $n \geq 0$ be an integer, and let $f: I \rightarrow \mathbb{R}$ be a function for which $f^{(1)}(a), \dots, f^{(n)}(a)$ all exist. The *Taylor polynomial of degree n of f at a* is

$$P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k$$

written with the conventions $f^{(0)} = f$ and $0! = 1$. The point a is the *centre* of the expansion.

For $f^{(n)}(a)$ to exist, $f^{(n-1)}$ must already exist at every point of some interval around a inside I , since a derivative at a point is a limit of difference quotients of a function defined near that point (definition 4.1.2). The degree of P_n is at most n : it drops when $f^{(n)}(a) = 0$, and P_n is the zero polynomial when every $f^{(k)}(a)$ vanishes.

The derivatives of the powers $(x-a)^m$ carry both the matching property and the proof of the main theorem below. Fix a and write $q_m(x) = (x-a)^m$ for an integer $m \geq 0$. Then q_0 is constant and $q'_0 = 0$, while $q'_1 = 1$ straight from definition 4.1.2, the difference quotient of q_1 between two distinct points x and t being $((t-a) - (x-a))/(t-x) = 1$. For $m \geq 2$, writing $q_m = q_1 q_{m-1}$ and applying the product rule of theorem 4.1.7 with the inductive hypothesis $q'_{m-1} = (m-1)q_{m-2}$ gives

$$q'_m(x) = q_{m-1}(x) + (x-a)(m-1)(x-a)^{m-2} = m(x-a)^{m-1}$$

so $q'_m = m q_{m-1}$ for every $m \geq 1$. Iterating that identity k times,

$$q_m^{(k)}(x) = m(m-1) \cdots (m-k+1) (x-a)^{m-k} \quad (0 \leq k \leq m), \quad q_m^{(k)} = 0 \quad (k > m) \quad (4.5)$$

where the second half follows from the first at $k = m$, which reads $q_m^{(m)} = m!$, a constant whose derivative is 0. Evaluating at $x = a$ leaves $q_m^{(k)}(a) = 0$ for $0 \leq k < m$ and $q_m^{(m)}(a) = m!$.

Differentiation is additive and homogeneous by theorem 4.1.7, so (4.5) applies term by term to $P_n = \sum_{j=0}^n \frac{f^{(j)}(a)}{j!} q_j$. At $x = a$ every summand with $j \neq k$ contributes 0, which leaves

$$P_n^{(k)}(a) = \frac{f^{(k)}(a)}{k!} q_k^{(k)}(a) = f^{(k)}(a) \quad (0 \leq k \leq n) \quad (4.6)$$

The same computation shows the coefficients are forced: a polynomial $Q(x) = \sum_{j=0}^n b_j(x-a)^j$ has $Q^{(k)}(a) = k! b_k$, so requiring $Q^{(k)}(a) = f^{(k)}(a)$ for $0 \leq k \leq n$ leaves $b_k = f^{(k)}(a)/k!$ and $Q = P_n$.

Example 4.3.2: Taylor Polynomials of x^3 , $1/(1-x)$ and $|x|^3$

1. Let $f(x) = x^3$ on $\mathbb{R}$ and $a = 1$. By (4.5) with centre 0, $f'(x) = 3x^2$, $f''(x) = 6x$ and $f'''(x) = 6$, so $f(1) = 1$, $f'(1) = 3$, $f''(1) = 6$ and $f'''(1) = 6$. Hence

$$P_3(x) = 1 + 3(x-1) + \frac{6}{2}(x-1)^2 + \frac{6}{6}(x-1)^3 = 1 + 3(x-1) + 3(x-1)^2 + (x-1)^3$$

Putting $u = x - 1$ and multiplying out, $1 + 3u + 3u^2 + u^3 = (1+u)^3 = x^3$, so $P_3 = f$ as functions. The lower-degree polynomials do not recover f : $P_2(x) = 1 + 3(x-1) + 3(x-1)^2$ differs from f by $(x-1)^3$ at every $x \neq 1$.

2. Let $f(x) = 1/(1-x)$ on $I = (-\infty, 1)$ and $a = 0$. Write $h_k(x) = (1-x)^{-k}$ for $k \geq 1$. The quotient rule of theorem 4.1.7, together with the derivative -1 of $x \mapsto 1-x$, gives $h'_1(x) = 1/(1-x)^2 = h_2(x)$. If $h'_k = k h_{k+1}$ for some $k \geq 1$, then the product rule applied to $h_{k+1} = h_k h_1$ gives

$$h'_{k+1} = h'_k h_1 + h_k h'_1 = k h_{k+1} h_1 + h_k h_2 = (k+1) h_{k+2}$$

so $h'_k = k h_{k+1}$ for every $k \geq 1$. A second induction now gives $f^{(k)} = k! h_{k+1}$ for every $k \geq 0$: it holds at $k = 0$ because $f = h_1$, and if it holds at k then $f^{(k+1)} = (k! h_{k+1})' = k! (k+1) h_{k+2} = (k+1)! h_{k+2}$. Since $h_{k+1}(0) = 1$, this gives $f^{(k)}(0) = k!$ and

$$P_n(x) = \sum_{k=0}^n \frac{k!}{k!} x^k = \sum_{k=0}^n x^k$$

which is the partial sum of the geometric series of example 2.4.7 through the term x^n .

3. Let $f(x) = |x|^3$ on $\mathbb{R}$ and $a = 0$. For $x > 0$ the function is x^3 , so $f'(x) = 3x^2 = 3x|x|$, and for $x < 0$ it is $-x^3$, so $f'(x) = -3x^2 = 3x|x|$. At 0 the difference quotient is $|h|^3/h$, whose absolute value is $|h|^2 \rightarrow 0$, so $f'(0) = 0$ and the formula $f'(x) = 3x|x|$ holds at every point of $\mathbb{R}$. Differentiating again, $f''(x) = 6x = 6|x|$ for $x > 0$ and $f''(x) = -6x = 6|x|$ for $x < 0$, while at 0 the quotient $3h|h|/h = 3|h| \rightarrow 0$ gives $f''(0) = 0 = 6|0|$. But $f'''(0)$ does not exist. The quotient $6|h|/h$ equals 6 for $h > 0$ and -6 for $h < 0$, so it has no limit at 0 (theorem 3.1.19 applied to $h = 1/n$ and $h = -1/n$). So f has Taylor polynomials at 0 only up to degree 2, and $P_0 = P_1 = P_2 = 0$.

Matching derivatives at a is a statement about f at the one point a , and by itself it says nothing about how close P_n stays to f as x moves away. Example 4.3.2(3) is the extreme case: there P_2 is identically zero while f is positive at every $x \neq 0$. What is needed is a formula for the difference $f - P_{n-1}$,

and the mean value theorem already gives one when $n = 1$, in the form $f(x) - f(a) = f'(\xi)(x - a)$ for some ξ between a and x . The general statement replaces f' by $f^{(n)}$ and the factor $x - a$ by $(x - a)^n/n!$.

Theorem 4.3.3: Lagrange Form of the Remainder

Let $n \geq 1$ be an integer, let $I \subseteq \mathbb{R}$ be an interval, and let $f: I \rightarrow \mathbb{R}$ be a function such that $f^{(n-1)}$ exists and is continuous on I , and $f^{(n)}(t)$ exists at every interior point t of I . Let $a, x \in I$ with $x \neq a$, and let P_{n-1} be the Taylor polynomial of degree $n - 1$ of f at a . Then there is a point ξ strictly between a and x with

$$f(x) = P_{n-1}(x) + \frac{f^{(n)}(\xi)}{n!}(x - a)^n$$

Proof:

Step 1: an auxiliary function. Since $x \neq a$, the number

$$M = \frac{f(x) - P_{n-1}(x)}{(x - a)^n}$$

is defined, and the conclusion is exactly the assertion that $n!M = f^{(n)}(\xi)$ for some ξ strictly between a and x . Put

$$g(t) = f(t) - P_{n-1}(t) - M(t - a)^n \quad (t \in I)$$

Step 2: the values of g at the two ends. In the notation $q_m(t) = (t - a)^m$ used above, $g = f - P_{n-1} - Mq_n$, so $g^{(k)} = f^{(k)} - P_{n-1}^{(k)} - Mq_n^{(k)}$ for every k at which these derivatives exist, differentiation being additive and homogeneous by theorem 4.1.7. Now $P_{n-1}^{(k)}(a) = f^{(k)}(a)$ for $0 \leq k \leq n - 1$ by (4.6), and $q_n^{(k)}(a) = 0$ for $0 \leq k \leq n - 1$ by (4.5). Evaluating at a therefore gives

$$g^{(k)}(a) = 0 \quad (0 \leq k \leq n - 1)$$

The definition of M gives $g(x) = f(x) - P_{n-1}(x) - M(x - a)^n = 0$.

Step 3: the n -th derivative of g . The polynomial P_{n-1} is a linear combination of $q_0, \dots, q_{n-1}$, and (4.5) makes $q_j^{(n)} = 0$ for every $j \leq n - 1$, so $P_{n-1}^{(n)} = 0$ identically. The same display gives $q_n^{(n)} = n!$. At an interior point t of I the derivative $f^{(n)}(t)$ exists by hypothesis, so $g^{(n)}(t)$ exists as well and

$$g^{(n)}(t) = f^{(n)}(t) - n!M \quad (4.7)$$

Step 4: n applications of Rolle's theorem. Set $x_0 = x$. The claim is that for each j with $1 \leq j \leq n$ there is a point x_j strictly between a and x_{j-1} satisfying $g^{(j)}(x_j) = 0$, and it is proved by induction on j .

Suppose x_{j-1} has been produced. Then $g^{(j-1)}(a) = 0$ by Step 2, which applies because $j - 1 \leq n - 1$, and $g^{(j-1)}(x_{j-1}) = 0$ by Step 2 when $j = 1$ and by the previous stage of the induction when $j \geq 2$. Let J be the closed interval whose endpoints are a and x_{j-1} , both of which lie in I , so that $J \subseteq I$ by definition 3.1.1. If p and q are points of I with $p < q$, then $(p, q) \subseteq I$ by definition 3.1.1, so every point strictly between p and q is an interior point of I . In particular every interior point of J is an interior point of I . Two cases now discharge the hypotheses of theorem 4.2.3 for $g^{(j-1)}$ on J .

1. If $j \leq n-1$, then $f^{(j)}$ exists at every point of I , because $j \leq n-1$ and $f^{(n-1)}$ exists on I . Polynomials are differentiable everywhere by theorem 4.1.7, so $g^{(j)}$ exists at every point of I . Thus $g^{(j-1)}$ is differentiable on I , and continuous there by proposition 4.1.5.
2. If $j = n$, then $g^{(n-1)} = f^{(n-1)} - P_{n-1}^{(n-1)} - Mq_n^{(n-1)}$ is continuous on I , because $f^{(n-1)}$ is continuous by hypothesis and the other two terms are polynomials, which are continuous by proposition 4.1.5. And $g^{(n)}(t)$ exists at every interior point t of I by Step 3.

In either case $g^{(j-1)}$ is continuous on J , differentiable at every interior point of J , and takes the value 0 at both endpoints of J . Theorem 4.2.3 produces a point x_j in the interior of J , that is, strictly between a and x_{j-1} , with $g^{(j)}(x_j) = 0$.

Step 5: conclusion. Put $\xi = x_n$. Each x_j lies strictly between a and x_{j-1} , so an induction on j places every x_j strictly between a and $x_0 = x$. In particular ξ is an interior point of I , so (4.7) applies at ξ and gives $0 = g^{(n)}(\xi) = f^{(n)}(\xi) - n!M$, that is, $M = f^{(n)}(\xi)/n!$. Multiplying the defining equation of M by $(x-a)^n$ and substituting this value,

$$f(x) = P_{n-1}(x) + \frac{f^{(n)}(\xi)}{n!}(x-a)^n$$

as we sought to show.

At $n = 1$ the Taylor polynomial P_0 is the constant $f(a)$ and the theorem reads $f(x) = f(a) + f'(a)(x-a)$, which is theorem 4.2.4. The general case moves the unknown point onto a higher derivative and replaces the factor $x-a$ by $(x-a)^n/n!$.

Suppose $|f^{(n)}(t)| \leq M_n$ at every interior point t of I . Taking absolute values in the conclusion of the theorem and using the multiplicativity of the absolute value (proposition 1.3.8(3)),

$$|f(x) - P_{n-1}(x)| \leq \frac{M_n}{n!} |x-a|^n \quad (4.8)$$

for every $x \in I$. When $x \neq a$ the point ξ the theorem produces is an interior point of I , so $|f^{(n)}(\xi)| \leq M_n$, and at $x = a$ both sides of the display are 0. So a bound on the single derivative $f^{(n)}$ across an interval containing a bounds the error of the approximation of f by P_{n-1} at every point of that interval. Whether the right-hand side tends to 0 as n grows is a question about how fast M_n grows against $n!$, and that comparison is the subject of section 4.7.

For a polynomial the remainder vanishes outright. If Q is a polynomial of degree at most m , then applying (4.5) with centre 0 to each of its terms gives $Q^{(m+1)} = 0$ identically, so the theorem with $n = m+1$ reads $Q(x) = P_m(x)$ for every $x \neq a$ and every centre a , while $Q(a) = P_m(a)$ holds by (4.6) at $k = 0$. A polynomial of degree at most m is therefore its own Taylor polynomial of degree m about any point, which is what example 4.3.2(1) exhibited for x^3 at $a = 1$. Combined with the uniqueness of the matching coefficients recorded after (4.6), this makes P_m the only polynomial of degree at most m whose derivatives at a through order m agree with those of f .

The point ξ is given by no formula in general, and the theorem asserts only that one exists. For $f(x) = 1/(1-x)$ at $a = 0$, as in example 4.3.2(2), both sides can be computed and ξ comes out explicitly. Fix $n \geq 1$ and let $0 < x < 1$. The theorem provides ξ strictly between 0 and x with

$$\frac{1}{1-x} - \sum_{k=0}^{n-1} x^k = \frac{f^{(n)}(\xi)}{n!} x^n = \frac{x^n}{(1-\xi)^{n+1}}$$

using $f^{(n)} = n! h_{n+1}$ from that example. The finite geometric identity $(1-x) \sum_{k=0}^{n-1} x^k = 1-x^n$ computed in example 2.4.7 evaluates the left-hand side as $x^n/(1-x)$, so $(1-\xi)^{n+1} = 1-x$ and theorem 1.3.12 identifies ξ completely:

$$\xi = 1 - (1-x)^{1/(n+1)}$$

That this ξ does lie strictly between 0 and x can be checked against the formula. Write $v = (1-x)^{1/(n+1)}$, which is positive because $1-x$ is, and satisfies $v < 1$, since $v \geq 1$ would give $v^{n+1} \geq 1 > 1-x$ by proposition 3.3.11. Then $0 < v^n < 1$, so $1-x = v^{n+1} = v \cdot v^n < v$, and $\xi = 1-v$ satisfies $0 < \xi < x$. For a numerical instance, take $n = 4$ and $x = 1/2$: the left-hand side above is $2 - (1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8}) = \frac{1}{8}$, while $(1-\xi)^5 = \frac{1}{2}$ makes the right-hand side $(\frac{1}{2})^4 / \frac{1}{2} = \frac{1}{8}$ as well.

Exercise 4.3.1

1. Let $f(x) = 1/(1+x)$ on $(-1, \infty)$. Show that $f^{(k)}(x) = (-1)^k k! (1+x)^{-(k+1)}$ for every integer $k \geq 0$, write down its Taylor polynomial P_3 of degree 3 at $a = 0$, and use theorem 4.3.3 to prove that $|f(x) - P_3(x)| \leq x^4$ for every $x \geq 0$.
2. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable with $f(0) = 0$, $f'(0) = 1$ and $|f''(t)| \leq 2$ for every $t \in \mathbb{R}$. Prove that $|f(x) - x| \leq x^2$ for every $x \in \mathbb{R}$, and name the Taylor polynomial and the constant M_2 with which (4.8) delivers this bound.
3. Let $p(x) = x^4 - 2x^2 + 5$. Compute the Taylor polynomial of degree 4 of p at $a = 1$, and verify by direct expansion that it equals p .
4. Let $f(x) = x^3$ on $\mathbb{R}$, let $a = 0$ and let $n = 2$. Show that for each $x \neq 0$ the point ξ whose existence theorem 4.3.3 asserts is $\xi = x/3$, and that it is the only point strictly between 0 and x for which the conclusion holds.
5. Let I be an open interval, let $c \in I$, and let $f: I \rightarrow \mathbb{R}$ be twice differentiable with f'' continuous on I . Suppose $f'(c) = 0$ and $f''(c) > 0$. Prove that there is a $\delta > 0$ with $f(x) > f(c)$ for every x satisfying $0 < |x - c| < \delta$, so that c is a local minimum of f in the sense of definition 4.2.1.

4.4 Uniform Convergence

Chapter 2 produced functions as limits. A power series names a function only through the convergence of its partial sums, and definition 2.6.5 defines $\exp$, $\cos$ and $\sin$ in exactly that way. Convergence at every point of a set is a condition imposed at each point separately, and it carries almost nothing about the limit function as a whole: example 4.4.3(1) gives continuous functions on $[0, 1]$ whose pointwise limit is discontinuous at one point.

The strengthened condition asks a single index N to answer a given ϵ at every point of the domain at once. Under it continuity does pass to the limit, a limit at a point can be interchanged with the limit in the index, and, with the hypothesis carried by the derivatives, a sequence of functions can be differentiated term by term. The last of these is the theorem section 4.5 applies to power series.

Definition 4.4.1: Pointwise Convergence

Let $E \subseteq \mathbb{R}$, let $f_n: E \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$, and let $f: E \rightarrow \mathbb{R}$. The sequence $(f_n)_{n \geq 1}$ *converges pointwise* to f on E if $f_n(x) \rightarrow f(x)$ for every $x \in E$. Written out, the condition is that for every $x \in E$ and every $\epsilon > 0$ there is an $N \in \mathbb{N}$ with

$$|f_n(x) - f(x)| < \epsilon \quad \text{for every } n \geq N$$

Let $g_n: E \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$ and write $s_N = \sum_{n=1}^N g_n$ for the partial sums. The series $\sum_{n \geq 1} g_n$ *converges pointwise* on E if $(s_N)_{N \geq 1}$ does, and its *sum* is the pointwise limit.

The limit function is determined by the sequence, since $f(x)$ is the limit of the real sequence $(f_n(x))_{n \geq 1}$ and proposition 2.1.4 makes that limit unique. So the phrase “the pointwise limit of (f_n) ” names at most one function.

Definition 4.4.2: [Real] Uniform Convergence

Let $E \subseteq \mathbb{R}$, let $f_n: E \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$, and let $f: E \rightarrow \mathbb{R}$. The sequence $(f_n)_{n \geq 1}$ *converges uniformly* to f on E if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$|f_n(x) - f(x)| < \epsilon \quad \text{for every } n \geq N \text{ and every } x \in E$$

The series $\sum_{n \geq 1} g_n$ *converges uniformly* on E if its sequence of partial sums converges uniformly on E .

Written as quantifier strings, pointwise convergence of (f_n) to f on E says

$$\forall x \in E \quad \forall \epsilon > 0 \quad \exists N \in \mathbb{N} \quad \forall n \geq N \quad (|f_n(x) - f(x)| < \epsilon)$$

and uniform convergence says

$$\forall \epsilon > 0 \quad \exists N \in \mathbb{N} \quad \forall x \in E \quad \forall n \geq N \quad (|f_n(x) - f(x)| < \epsilon)$$

The two strings differ only in the position of $\exists N$, which has moved to the left of $\forall x$. In the first, N may depend on x as well as on ϵ ; in the second it may depend on ϵ only. That is the move box 3.4.2 recorded for the δ of definition 3.4.1, performed here on the index of a sequence.

One implication holds with no hypothesis on E : a uniformly convergent sequence converges pointwise to the same limit, since fixing $x \in E$ leaves the N produced for ϵ answering ϵ at that x . Negating definition 4.4.2 describes the gap between the two notions. The sequence (f_n) fails to converge uniformly to f on E exactly when there is an $\epsilon_0 > 0$ such that for every $N \in \mathbb{N}$ there are an $n \geq N$ and an $x \in E$ with $|f_n(x) - f(x)| \geq \epsilon_0$. Each item below refutes uniform convergence by producing such an ϵ_0 , and the three fail for three different reasons.

Example 4.4.3: x^n on $[0, 1]$, x/n on the Line, and $nx/(1 + n^2x^2)$ on $[0, 1]$

1. Let $f_n(x) = x^n$ on $E = [0, 1]$, each continuous by example 3.2.6 together with proposition 3.3.1. For $0 < x < 1$ the sequence (x^n) is null by proposition 2.1.10 with $j = 0$ and $r = x$, for $x = 0$ every term is 0, and for $x = 1$ every term is 1. So (f_n) converges pointwise to the function f with $f(x) = 0$ for $0 \leq x < 1$ and $f(1) = 1$.

This f is discontinuous at 1. The points $x_k = 1 - 1/k$ lie in $[0, 1]$ and satisfy $x_k \rightarrow 1$, while $f(x_k) = 0$ for every k , so $f(x_k) \rightarrow 0 \neq 1 = f(1)$ and proposition 3.2.4 denies continuity at 1.

The convergence is not uniform. Take $\epsilon_0 = 1/2$ and let $N \in \mathbb{N}$. By theorem 1.3.12 there is a unique $t \geq 0$ with $t^N = 1/2$. If $t = 0$ then $t^N = 0$, and if $t \geq 1$ then $t^N \geq 1^N = 1$ by proposition 3.3.11, so $0 < t < 1$ and hence $f(t) = 0$. Then $|f_N(t) - f(t)| = t^N = 1/2 \geq \epsilon_0$, so no N answers ϵ_0 .

2. Let $f_n(x) = x/n$ on $E = \mathbb{R}$. For fixed x the sequence $x \cdot (1/n)$ converges to 0 by example 2.1.3(1) and theorem 2.1.6, so (f_n) converges pointwise to the constant function 0. The convergence is not uniform on $\mathbb{R}$: take $\epsilon_0 = 1$, let $N \in \mathbb{N}$, and put $x = 2N$, so that $|f_N(x) - 0| = 2 \geq \epsilon_0$.

On a bounded set the same sequence does converge uniformly. Let $E_0 \subseteq \mathbb{R}$ and $M > 0$ satisfy $|x| \leq M$ for every $x \in E_0$, and let $\epsilon > 0$. By theorem 1.4.1(2) there is an $N \in \mathbb{N}$ with $N > M/\epsilon$. For $n \geq N$ and $x \in E_0$,

$$\left| \frac{x}{n} \right| = \frac{|x|}{n} \leq \frac{M}{n} \leq \frac{M}{N} < \epsilon$$

the first equality by proposition 1.3.8(3) and the last step by proposition 1.1.2(7). The obstruction in this item is therefore the size of the domain.

3. Let $f_n(x) = nx/(1 + n^2x^2)$ on $E = [0, 1]$. The denominator satisfies $1 + n^2x^2 \geq 1 > 0$, so each f_n is continuous by example 3.2.6. For $x = 0$ every term is 0. For $0 < x \leq 1$ the bounds $1 + n^2x^2 \geq n^2x^2 > 0$ and $nx > 0$ give

$$0 \leq \frac{nx}{1 + n^2x^2} \leq \frac{nx}{n^2x^2} = \frac{1}{x} \cdot \frac{1}{n}$$

and the right side converges to 0 by example 2.1.3(1) and theorem 2.1.6, so $f_n(x) \rightarrow 0$ by theorem 2.1.7. The pointwise limit is therefore the constant function 0, which is continuous.

The convergence is still not uniform. Take $\epsilon_0 = 1/2$, let $N \in \mathbb{N}$, and put $x = 1/N \in [0, 1]$. Then

$$|f_N(x) - 0| = \frac{N \cdot (1/N)}{1 + N^2 \cdot (1/N)^2} = \frac{1}{2} \geq \epsilon_0$$

so no N answers ϵ_0 . A continuous limit is therefore no guarantee of uniform convergence.

Item (1) shows that continuity is lost across a pointwise limit. Item (2) shows that uniform convergence is a property of the pair (formula, domain): one sequence of functions has it on every bounded set and fails it on $\mathbb{R}$. Item (3) refutes the converse of theorem 4.4.5 below.

Definition 4.4.2 names the limit function in its statement, so a test built directly on it needs f before it can be applied. For a sequence of real numbers theorem 2.3.5 removed that requirement, and the same removal works here once the index answering ϵ is again taken independent of the point.

Theorem 4.4.4: The Uniform Cauchy Criterion

Let $E \subseteq \mathbb{R}$ and let $f_n: E \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$. Then (f_n) converges uniformly on E to some $f: E \rightarrow \mathbb{R}$ if and only if for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ such that

$$|f_n(x) - f_m(x)| < \epsilon \quad \text{for all } n, m \geq N \text{ and every } x \in E$$

Proof:

The condition is necessary. Suppose $f_n \rightarrow f$ uniformly on E and let $\epsilon > 0$. Take $N \in \mathbb{N}$ with $|f_n(x) - f(x)| < \epsilon/2$ for every $n \geq N$ and every $x \in E$. For $n, m \geq N$ and $x \in E$, splitting the difference across $f(x)$ and applying proposition 1.3.8(4),

$$|f_n(x) - f_m(x)| \leq |f_n(x) - f(x)| + |f(x) - f_m(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

where $|f(x) - f_m(x)| = |f_m(x) - f(x)|$ by proposition 1.3.8(1).

The condition is sufficient. Suppose the condition holds and fix $x \in E$. Given $\epsilon > 0$, the N provided by the condition satisfies $|f_n(x) - f_m(x)| < \epsilon$ for all $n, m \geq N$, which is the requirement of definition 2.3.1 for the real sequence $(f_n(x))_{n \geq 1}$. By theorem 2.3.5 that sequence converges, and defining $f(x)$ to be its limit gives a function $f: E \rightarrow \mathbb{R}$ with $f_n(x) \rightarrow f(x)$ for every $x \in E$. Let $\epsilon > 0$ and take $N \in \mathbb{N}$ with $|f_n(x) - f_m(x)| < \epsilon/2$ for all $n, m \geq N$ and every $x \in E$. Fix $n \geq N$ and $x \in E$. For every $m \geq N$, proposition 1.3.8(2) turns the bound into

$$-\frac{\epsilon}{2} < f_n(x) - f_m(x) < \frac{\epsilon}{2}$$

The constant sequence with value $f_n(x)$ converges to $f_n(x)$, so theorem 2.1.6 makes the sequence $m \mapsto f_n(x) - f_m(x)$ converge to $f_n(x) - f(x)$. Applying proposition 2.1.8(2) to each of the two bounds, with $N_0 = N$, gives

$$-\frac{\epsilon}{2} \leq f_n(x) - f(x) \leq \frac{\epsilon}{2}$$

and proposition 1.3.8(2) reads this back as $|f_n(x) - f(x)| \leq \epsilon/2 < \epsilon$. The index N was chosen before x , so this holds for every $n \geq N$ and every $x \in E$, which is uniform convergence of (f_n) to f , as we sought to show.

The criterion tests a sequence of functions on its own terms. The two theorems that follow transfer a property from those terms to the limit, and the first transfers continuity.

Theorem 4.4.5: A Uniform Limit of Continuous Functions Is Continuous

Let $E \subseteq \mathbb{R}$ and let $f_n: E \rightarrow \mathbb{R}$ be continuous for each $n \in \mathbb{N}$. If (f_n) converges uniformly to f on E , then f is continuous.

Proof:

Let $c \in E$ and let $\epsilon > 0$. By definition 4.4.2 there is an $N \in \mathbb{N}$ with $|f_N(x) - f(x)| < \epsilon/3$ for every $x \in E$. The function f_N is continuous at c by definition 3.2.2, so definition 3.2.1 gives a $\delta > 0$ such that $|f_N(x) - f_N(c)| < \epsilon/3$ for every $x \in E$ with $|x - c| < \delta$.

Let $x \in E$ satisfy $|x - c| < \delta$. Writing

$$f(x) - f(c) = (f(x) - f_N(x)) + (f_N(x) - f_N(c)) + (f_N(c) - f(c))$$

and applying proposition 1.3.8(4) twice,

$$|f(x) - f(c)| \leq |f(x) - f_N(x)| + |f_N(x) - f_N(c)| + |f_N(c) - f(c)| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

the first and third bounds by the choice of N together with proposition 1.3.8(1). So f is continuous at c , and since $c \in E$ was arbitrary, f is continuous, completing the proof.

The contrapositive of theorem 4.4.5 refutes uniform convergence with no ϵ_0 to produce: continuous functions with a discontinuous pointwise limit cannot converge uniformly. That settles example 4.4.3(1) a second time, and example 4.4.3(3) shows the test works in one direction only, since there the limit is continuous and the convergence is not uniform.

The series form follows from the theorem applied to partial sums. If each $g_n: E \rightarrow \mathbb{R}$ is continuous and $\sum_{n \geq 1} g_n$ converges uniformly on E , then each partial sum $s_N = \sum_{n=1}^N g_n$ is continuous by theorem 3.2.5 and induction on N , and (s_N) converges uniformly by definition 4.4.2, so the sum of the series is continuous.

Theorem 4.4.5 transfers a statement made at points of E . The difference quotient $t \mapsto (f_n(t) - f_n(x))/(t - x)$ whose limit definition 4.1.2 takes is undefined at $t = x$, so its domain is $E \setminus \{x\}$ and the point at which its limit is wanted lies outside that domain. The transfer needed there asks only that the point be a limit point of the domain.

Theorem 4.4.6: Interchange of a Limit at a Point with a Limit in the Index

Let $E \subseteq \mathbb{R}$, let c be a limit point of E , and let $f_n: E \rightarrow \mathbb{R}$ converge uniformly to f on E . Suppose the limit

$$A_n = \lim_{x \rightarrow c} f_n(x)$$

exists in $\mathbb{R}$ for every $n \in \mathbb{N}$. Then $(A_n)_{n \geq 1}$ converges, the limit of f at c exists, and

$$\lim_{x \rightarrow c} f(x) = \lim_{n \rightarrow \infty} A_n$$

that is, $\lim_{x \rightarrow c} \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \lim_{x \rightarrow c} f_n(x)$.

Proof:

Step 1: the sequence (A_n) converges. Let $\epsilon > 0$. By theorem 4.4.4 there is an $N \in \mathbb{N}$ with $|f_n(y) - f_m(y)| < \epsilon/2$ for all $n, m \geq N$ and every $y \in E$. Since c is a limit point of E , proposition 3.1.9(2) produces a sequence $(y_k)_{k \geq 1}$ with $y_k \in E$, $y_k \neq c$ for every k , and $y_k \rightarrow c$. Fix $n, m \geq N$. By theorem 3.1.19(1) applied to f_n and to f_m along (y_k) , the sequences $(f_n(y_k))_k$ and $(f_m(y_k))_k$ converge to A_n and to A_m , so theorem 2.1.6 makes $f_n(y_k) - f_m(y_k) \rightarrow A_n - A_m$ as $k \rightarrow \infty$. Every k satisfies $-\epsilon/2 < f_n(y_k) - f_m(y_k) < \epsilon/2$ by proposition 1.3.8(2), so proposition 2.1.8(2) applied to each bound gives $-\epsilon/2 \leq A_n - A_m \leq \epsilon/2$, and proposition 1.3.8(2) reads this as $|A_n - A_m| \leq \epsilon/2 < \epsilon$. So (A_n) satisfies definition 2.3.1 and theorem 2.3.5 makes it converge. Write A for its limit.

Step 2: the limit of f at c is A . Let $\epsilon > 0$. Uniform convergence gives an $N_1 \in \mathbb{N}$ with

$|f_n(x) - f(x)| < \epsilon/3$ for every $n \geq N_1$ and every $x \in E$, and $A_n \rightarrow A$ gives an $N_2 \in \mathbb{N}$ with $|A_n - A| < \epsilon/3$ for every $n \geq N_2$. Fix $n = \max\{N_1, N_2\}$.

Since $\lim_{x \rightarrow c} f_n(x) = A_n$, definition 3.1.15 gives a $\delta > 0$ with $|f_n(x) - A_n| < \epsilon/3$ for every $x \in E$ with $0 < |x - c| < \delta$. For such an x , writing

$$f(x) - A = (f(x) - f_n(x)) + (f_n(x) - A_n) + (A_n - A)$$

and applying proposition 1.3.8(4) twice,

$$|f(x) - A| \leq |f(x) - f_n(x)| + |f_n(x) - A_n| + |A_n - A| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

the first bound by proposition 1.3.8(1) and the choice of N_1 . Since $\epsilon > 0$ was arbitrary and c is a limit point of E , definition 3.1.15 gives $\lim_{x \rightarrow c} f(x) = A$. Both iterated limits therefore exist and equal A , as we sought to show.

The point c is not required to lie in E , and that is what separates theorem 4.4.6 from theorem 4.4.5. When c does lie in E and each f_n is continuous at c , the numbers A_n may be taken to be $f_n(c)$ by proposition 3.2.3(1), and the conclusion becomes $\lim_{x \rightarrow c} f(x) = \lim_n f_n(c) = f(c)$, which is continuity of f at c . Theorem 4.4.8 uses the case $c \notin E$.

Testing a series against theorem 4.4.4 still requires estimating sums of the functions g_n . When each g_n is bounded on E by a number independent of x , those estimates reduce to a question about a single series of real numbers, which the tests of section 2.4 answer.

Theorem 4.4.7: The Weierstrass M-Test

Let $E \subseteq \mathbb{R}$ and let $g_n : E \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$. Suppose there are real numbers M_n with

$$|g_n(x)| \leq M_n \quad \text{for every } x \in E \text{ and every } n \in \mathbb{N}$$

and suppose $\sum_{n \geq 1} M_n$ converges. Then $\sum_{n \geq 1} g_n$ converges uniformly on E , and $\sum_{n \geq 1} g_n(x)$ converges absolutely for every $x \in E$.

Proof:

Write $s_N = \sum_{n=1}^N g_n$ for the partial sums and let $\epsilon > 0$. Since $\sum_{n \geq 1} M_n$ converges, proposition 2.4.2 gives an $N \in \mathbb{N}$ with

$$\left| \sum_{k=n}^m M_k \right| < \epsilon \quad \text{for all integers } m \geq n \geq N$$

Let $p, q \geq N$ and let $x \in E$. If $p = q$ then $|s_p(x) - s_q(x)| = 0 < \epsilon$. If $p > q$, then $s_p(x) - s_q(x) = \sum_{k=q+1}^p g_k(x)$ and the indices satisfy $p \geq q + 1 \geq N + 1 \geq N$, so

$$|s_p(x) - s_q(x)| = \left| \sum_{k=q+1}^p g_k(x) \right| \leq \sum_{k=q+1}^p |g_k(x)| \leq \sum_{k=q+1}^p M_k \leq \left| \sum_{k=q+1}^p M_k \right| < \epsilon$$

the first inequality by proposition 1.3.8(1) applied $p - q - 1$ times to the finite sum, whose convergence is not in question, the second by summing the hypothesis $|g_k(x)| \leq M_k$ over the

range, and the third by proposition 1.3.8(1). The case $q > p$ follows by exchanging the names, since $|s_p(x) - s_q(x)| = |s_q(x) - s_p(x)|$ by proposition 1.3.8(1).

The index N was chosen before x , so (s_N) satisfies the condition of theorem 4.4.4 on E and therefore converges uniformly on E , which by definition 4.4.2 is uniform convergence of the series. For a fixed $x \in E$, the hypothesis $|g_n(x)| \leq M_n$ holds for every $n \geq 1$ and $\sum_{n \geq 1} M_n$ converges, so theorem 2.4.13(1) with $a_n = g_n(x)$, $c_n = M_n$ and $N_0 = 1$ gives absolute convergence at x , completing the proof.

The test reaches the geometric series on every interval strictly inside $(-1, 1)$ and fails on $(-1, 1)$ itself, which is the pattern section 4.5 generalizes. Fix $0 < r < 1$, take $E = [-r, r]$, and put $g_n(x) = x^{n-1}$ for $n \geq 1$, so that $\sum_{n \geq 1} g_n$ is the series $\sum_{n \geq 0} x^n$ of example 2.4.7 with its terms reindexed. An induction using proposition 1.3.8(3) gives $|x^{n-1}| = |x|^{n-1}$, and a second induction using proposition 1.1.2(5) gives $|x|^{n-1} \leq r^{n-1}$ whenever $|x| \leq r$, so $M_n = r^{n-1}$ bounds g_n on E . The series $\sum_{n \geq 1} r^{n-1}$ is $\sum_{n \geq 0} r^n$, which converges by example 2.4.7 because $0 < r < 1$. So theorem 4.4.7 gives uniform convergence on $[-r, r]$.

On $E = (-1, 1)$ the convergence is not uniform, so no choice of bounds M_n can meet the hypothesis there. Write $f(x) = 1/(1-x)$ for the sum, which is what example 2.4.7 computes, and $s_N(x) = \sum_{n=0}^{N-1} x^n$ for the N -th partial sum of $\sum_{n \geq 1} g_n$. Take $\epsilon_0 = 1/2$ and let $N \in \mathbb{N}$. By theorem 1.3.12 there is a unique $t \geq 0$ with $t^N = 1/2$, and it satisfies $0 < t < 1$ for the reason given in example 4.4.3(1). For every integer $m \geq N$ the terms t^n with $N \leq n \leq m$ are positive, so

$$\sum_{n=N}^m t^n \geq t^N = \frac{1}{2}$$

and $\sum_{n=N}^m t^n = (\sum_{n=0}^m t^n) - s_N(t)$ converges to $f(t) - s_N(t)$ as $m \rightarrow \infty$, by example 2.4.7 and theorem 2.1.6. Now proposition 2.1.8(2) gives $f(t) - s_N(t) \geq 1/2 = \epsilon_0$, so the N -th partial sum misses f by at least ϵ_0 at the point $t \in (-1, 1)$, for every N .

Differentiation is the property that does not transfer under uniform convergence of the functions themselves. Let $f_n(x) = x/(1+n^2x^2)$ on $\mathbb{R}$. Expanding $(1-n|x|)^2 \geq 0$ gives $1+n^2x^2 \geq 2n|x|$, so for $x \neq 0$

$$|f_n(x)| = \frac{|x|}{1+n^2x^2} \leq \frac{|x|}{2n|x|} = \frac{1}{2n}$$

and $f_n(0) = 0$, so $|f_n(x)| \leq 1/(2n)$ for every $x \in \mathbb{R}$. Given $\epsilon > 0$, theorem 1.4.1(2) gives an $N \in \mathbb{N}$ with $N > 1/(2\epsilon)$, and then $|f_n(x) - 0| \leq 1/(2n) \leq 1/(2N) < \epsilon$ for all $n \geq N$ and all $x \in \mathbb{R}$. So $f_n \rightarrow 0$ uniformly on $\mathbb{R}$, and the limit function $f = 0$ has $f'(0) = 0$. The quotient rule of theorem 4.1.7 gives

$$f'_n(x) = \frac{(1+n^2x^2) - x \cdot 2n^2x}{(1+n^2x^2)^2} = \frac{1-n^2x^2}{(1+n^2x^2)^2}$$

so $f'_n(0) = 1$ for every n , and $\lim_n f'_n(0) = 1 \neq 0 = f'(0)$. The hypothesis in the next theorem is therefore placed on the derivatives, and the functions themselves are asked only to converge at one point.

Theorem 4.4.8: Term-by-Term Differentiation

Let $a < b$ and let $f_n: [a, b] \rightarrow \mathbb{R}$ be differentiable for each $n \in \mathbb{N}$. Suppose that

1. the sequence $(f_n(x_0))_{n \geq 1}$ converges for some $x_0 \in [a, b]$, and
2. the sequence $(f'_n)_{n \geq 1}$ converges uniformly on $[a, b]$.

Then (f_n) converges uniformly on $[a, b]$ to a function f , this f is differentiable, and

$$f'(x) = \lim_{n \rightarrow \infty} f'_n(x) \quad \text{for every } x \in [a, b]$$

Proof:

Step 1: a uniform Lipschitz estimate for the differences. The claim is that for every $\eta > 0$ there is an $N \in \mathbb{N}$ such that

$$|(f_n - f_m)(x) - (f_n - f_m)(t)| \leq \eta |x - t| \quad (4.9)$$

for all $n, m \geq N$ and all $x, t \in [a, b]$.

Let $\eta > 0$. Hypothesis (2) and theorem 4.4.4 give an $N \in \mathbb{N}$ with $|f'_n(s) - f'_m(s)| < \eta$ for all $n, m \geq N$ and every $s \in [a, b]$. Fix such n and m and put $\varphi = f_n - f_m$. By theorem 4.1.7 the function φ is differentiable at every point of $[a, b]$ with $\varphi' = f'_n - f'_m$, and by proposition 4.1.5 it is continuous on $[a, b]$.

Let $x, t \in [a, b]$. If $x = t$ both sides of (4.9) are 0. Otherwise put $u = \min\{x, t\}$ and $v = \max\{x, t\}$, so that $a \leq u < v \leq b$. The restriction of φ to $[u, v]$ is continuous, and for each $\xi \in (u, v)$ the point ξ is a limit point of $[u, v] \setminus \{\xi\}$, so the difference quotient of φ at ξ has the same limit along $[u, v] \setminus \{\xi\}$ as along $[a, b] \setminus \{\xi\}$ by proposition 3.1.23(1). That restriction is therefore differentiable on (u, v) with derivative $\varphi'(\xi)$, and theorem 4.2.4 gives a $\xi \in (u, v)$ with $\varphi(v) - \varphi(u) = \varphi'(\xi)(v - u)$. Taking absolute values and using proposition 1.3.8(1) and (3),

$$|\varphi(x) - \varphi(t)| = |\varphi(v) - \varphi(u)| = |\varphi'(\xi)| (v - u) \leq \eta |x - t|$$

since $v - u = |x - t| > 0$ and $|\varphi'(\xi)| = |f'_n(\xi) - f'_m(\xi)| < \eta$. That is (4.9).

Step 2: the sequence (f_n) converges uniformly on $[a, b]$. Let $\epsilon > 0$. Hypothesis (1) says $(f_n(x_0))$ converges, so proposition 2.3.3 gives an $N_1 \in \mathbb{N}$ with $|f_n(x_0) - f_m(x_0)| < \epsilon/2$ for all $n, m \geq N_1$. Apply Step 1 with $\eta = \epsilon/(2(b - a))$, which is positive because $b - a > 0$, to get an $N_2 \in \mathbb{N}$, and put $N = \max\{N_1, N_2\}$.

Let $n, m \geq N$, put $\varphi = f_n - f_m$, and let $x \in [a, b]$. Both x and x_0 lie in $[a, b]$, so $|x - x_0| \leq b - a$ and (4.9) gives

$$|\varphi(x) - \varphi(x_0)| \leq \frac{\epsilon |x - x_0|}{2(b - a)} \leq \frac{\epsilon}{2}$$

Splitting $\varphi(x)$ across x_0 and applying proposition 1.3.8(4),

$$|f_n(x) - f_m(x)| = |\varphi(x)| \leq |\varphi(x) - \varphi(x_0)| + |\varphi(x_0)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

the strict inequality coming from $|\varphi(x_0)| = |f_n(x_0) - f_m(x_0)| < \epsilon/2$. The index N was chosen before x , so theorem 4.4.4 produces a function $f: [a, b] \rightarrow \mathbb{R}$ with $f_n \rightarrow f$ uniformly on $[a, b]$.

Step 3: the difference quotients converge uniformly away from a point. Fix $x \in [a, b]$ and put $E_x = [a, b] \setminus \{x\}$. The point x is a limit point of E_x by lemma 4.1.1. Define $\varphi_n, \varphi: E_x \rightarrow \mathbb{R}$ by

$$\varphi_n(t) = \frac{f_n(t) - f_n(x)}{t - x}, \quad \varphi(t) = \frac{f(t) - f(x)}{t - x}$$

so that $\lim_{t \rightarrow x} \varphi_n(t) = f'_n(x)$ for every n , by definition 4.1.2.

Let $\eta > 0$ and take $N \in \mathbb{N}$ from Step 1 for the value $\eta/2$. For $n, m \geq N$ and $t \in E_x$, the numerator of $\varphi_n(t) - \varphi_m(t)$ is $(f_n - f_m)(t) - (f_n - f_m)(x)$, so dividing (4.9) by $|t - x| > 0$ and using proposition 1.3.8(3),

$$|\varphi_n(t) - \varphi_m(t)| = \frac{|(f_n - f_m)(t) - (f_n - f_m)(x)|}{|t - x|} \leq \frac{\eta}{2} < \eta$$

So (φ_n) satisfies the condition of theorem 4.4.4 on E_x and converges uniformly there to some function. That function is φ : for each fixed $t \in E_x$, Step 2 gives $f_n(t) \rightarrow f(t)$ and $f_n(x) \rightarrow f(x)$, so theorem 2.1.6 gives $\varphi_n(t) \rightarrow \varphi(t)$. A uniform limit is also a pointwise limit, so proposition 2.1.4 identifies the two.

Step 4: interchanging the two limits. Apply theorem 4.4.6 on the set E_x with $c = x$, which is a limit point of E_x by Step 3, to the sequence (φ_n) , which converges uniformly to φ on E_x by Step 3 and whose members have the limits $A_n = f'_n(x)$ at x . The theorem gives

$$\lim_{t \rightarrow x} \varphi(t) = \lim_{n \rightarrow \infty} f'_n(x)$$

and asserts that both sides exist. The left side is the limit at x of the difference quotient of f over $[a, b] \setminus \{x\}$, so definition 4.1.2 says that f is differentiable at x with $f'(x) = \lim_n f'_n(x)$. The point $x \in [a, b]$ was arbitrary, completing the proof.

Neither hypothesis may be weakened to a statement about (f_n) alone. The example before the theorem has (f_n) converging uniformly on $\mathbb{R}$ while $\lim_n f'_n(0) \neq f'(0)$, so uniform convergence of the functions does not deliver the conclusion once hypothesis (2) is removed. With hypothesis (1) deleted, a sequence satisfying (2) can fail to converge at every point of $[a, b]$, which exercise 4.4.1 (8) exhibits. The derivatives f'_n are nowhere assumed continuous, and the only property of them the proof uses is the one theorem 4.2.4 extracts.

For series the theorem reads as follows. Let $g_n: [a, b] \rightarrow \mathbb{R}$ be differentiable, suppose $\sum_{n \geq 1} g_n(x_0)$ converges for some $x_0 \in [a, b]$, and suppose $\sum_{n \geq 1} g'_n$ converges uniformly on $[a, b]$. The partial sums $s_N = \sum_{n=1}^N g_n$ are differentiable with $s'_N = \sum_{n=1}^N g'_n$ by theorem 4.1.7 and induction on N , so theorem 4.4.8 applies to (s_N) and gives uniform convergence of $\sum_{n \geq 1} g_n$ on $[a, b]$ together with

$$\left(\sum_{n=1}^{\infty} g_n \right)'(x) = \sum_{n=1}^{\infty} g'_n(x) \quad \text{for every } x \in [a, b]$$

This is the form section 4.5 applies to a power series, where theorem 4.4.7 discharges the hypothesis on the differentiated series.

Exercise 4.4.1

1. For $n \in \mathbb{N}$ let $f_n: [0, 1] \rightarrow \mathbb{R}$ be $f_n(x) = x^n/n$.
 1. Show that (f_n) converges uniformly on $[0, 1]$, and identify the limit f .
 2. Compute f'_n and show that (f'_n) converges pointwise on $[0, 1]$ and not uniformly.
 3. Show that $f'(1) \neq \lim_n f'_n(1)$.

Which hypothesis of theorem 4.4.8 fails here?

2. Show that $\sum_{n \geq 1} x^n/n^2$ converges uniformly on $[-1, 1]$ and that its sum is continuous on $[-1, 1]$.
3. Let $E \subseteq \mathbb{R}$ and suppose $f_n \rightarrow f$ and $g_n \rightarrow g$ uniformly on E .
 1. Show that $f_n + g_n \rightarrow f + g$ uniformly on E .
 2. Give functions $f_n, g_n: \mathbb{R} \rightarrow \mathbb{R}$ converging uniformly on $\mathbb{R}$ to f and g for which $(f_n g_n)$ does not converge uniformly to fg on $\mathbb{R}$.
4. Let $E \subseteq \mathbb{R}$, let each $f_n: E \rightarrow \mathbb{R}$ be bounded, and suppose $f_n \rightarrow f$ uniformly on E . Show that f is bounded and that there is a single $M \in \mathbb{R}$ with $|f_n(x)| \leq M$ for every $n \in \mathbb{N}$ and every $x \in E$. Then give bounded functions $f_n: (0, 1] \rightarrow \mathbb{R}$ converging pointwise to a bounded function for which no such M exists.
5. Let $E = (0, 1]$ and let $f_n(x) = nx/(1 + nx)$.
 1. Show that (f_n) converges pointwise on E to the constant function 1.
 2. Show that $\lim_{x \rightarrow 0} f_n(x) = 0$ for every $n \in \mathbb{N}$, the limit being taken over E .
 3. Deduce from theorem 4.4.6 that the convergence is not uniform on E .
6. Let $E \subseteq \mathbb{R}$, let each $f_n: E \rightarrow \mathbb{R}$ be uniformly continuous on E , and suppose $f_n \rightarrow f$ uniformly on E . Show that f is uniformly continuous on E .
7. Give functions $f_n: [0, 1] \rightarrow \mathbb{R}$ with the following two properties: for every $\epsilon > 0$ there is an $N \in \mathbb{N}$ with $|f_{n+1}(x) - f_n(x)| < \epsilon$ for every $n \geq N$ and every $x \in [0, 1]$, and $(f_n(x))_{n \geq 1}$ converges for no $x \in [0, 1]$. Conclude that a uniform bound on consecutive differences is weaker than the condition of theorem 4.4.4.
8. Give differentiable functions $f_n: [0, 1] \rightarrow \mathbb{R}$ for which (f'_n) converges uniformly on $[0, 1]$ while $(f_n(x))$ diverges for every $x \in [0, 1]$. Which hypothesis of theorem 4.4.8 does this delete, and what does the example establish about it?

Theorem 4.4.8 exchanges a uniform limit with differentiation. The same exchange with integration is theorem 5.2.8, which waits for Chapter 5 because the integral does not exist yet.

4.5 Power Series as Functions

Definition 2.6.2 attaches to a power series a radius R , and theorem 2.6.3 settles, at every point except the two endpoints, whether the series converges there. Convergence at each point of $(a - R, a + R)$ makes the sum of the series a function on that interval, and nothing so far says what kind of function it is. This section answers that. The sum has derivatives of every order, each obtained by differentiating the series term by term, and its coefficients are the Taylor coefficients of the sum at the centre, so a function has at most one power series expansion about a given point. The expansion can be recentred at any interior point, and the sum is determined by its values on any subset of the interval having a limit point inside. The last theorem concerns the endpoints, where theorem 2.6.3 is silent: at an endpoint where the series converges, its sum there is the limit of the function from inside the interval. Throughout this section $\sum_{n \geq 0} c_n(x - a)^n$ is a power series in the sense of definition 2.6.2 whose

radius of convergence R is positive, and I is the interval

$$I = (a - R, a + R)$$

on which theorem 2.6.3 gives convergence, read as $I = \mathbb{R}$ when $R = +\infty$. Its sum is the function

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n \quad (x \in I)$$

and the partial sums are written $s_N(x) = \sum_{n=0}^N c_n(x-a)^n$, each a polynomial in x .

Two facts about radii are used repeatedly below. The first computes the radius of a series whose coefficients have been multiplied by a factor whose n -th root tends to 1, which is what the coefficient n produced by term-by-term differentiation does. The second bounds the radius below using only the set of points at which the series is known to converge, which is all the information a construction that produces a series point by point makes available.

Lemma 4.5.1: Scaling an Upper Limit, and a Lower Bound for a Radius of Convergence

1. Let $(u_n)_{n \geq 1}$ be a sequence of nonnegative reals and let $(t_n)_{n \geq 1}$ be a sequence of positive reals with $t_n \rightarrow t$ for some real $t > 0$. Then

$$\limsup_{n \rightarrow \infty} (t_n u_n) = t \limsup_{n \rightarrow \infty} u_n$$

both sides being read as $+\infty$ when $\limsup_n u_n = +\infty$.

2. Let $\sum_{n \geq 0} c_n(x-a)^n$ be a power series with radius of convergence R , and let ρ be a real number with $\rho > 0$. If the series converges at every $x \in \mathbb{R}$ with $|x-a| < \rho$, then $R \geq \rho$.

Proof:

Part (1). Write $s = \limsup_n u_n$ and $\sigma = \limsup_n (t_n u_n)$. Both sequences have nonnegative terms, so comparing each with the constant sequence 0, whose limit and hence whose upper limit is 0 by theorem 2.2.12(5), gives $s \geq 0$ and $\sigma \geq 0$ by proposition 2.2.13(1). In particular both lie in $[0, +\infty]$.

Step 1: $\sigma \leq ts$. If $s = +\infty$ the right-hand side is $+\infty$ and there is nothing to prove, so assume s is real. Let ϵ be a real number with $0 < \epsilon \leq 1$. Since $t_n \rightarrow t$ there is an N_1 with $t_n < t + \epsilon$ for every $n \geq N_1$, and since $s + \epsilon > s = \limsup_n u_n$, theorem 2.2.12(1) provides an N_2 with $u_n < s + \epsilon$ for every $n \geq N_2$. For $n \geq \max\{N_1, N_2\}$ the two bounds multiply, because $u_n \geq 0$ and $t + \epsilon > 0$:

$$t_n u_n \leq (t + \epsilon) u_n \leq (t + \epsilon)(s + \epsilon)$$

Comparing $(t_n u_n)$ with the constant sequence $(t + \epsilon)(s + \epsilon)$ through proposition 2.2.13(1), and using theorem 2.2.12(5) for the upper limit of a constant sequence,

$$\sigma \leq (t + \epsilon)(s + \epsilon) = ts + \epsilon(t + s) + \epsilon^2 \leq ts + \epsilon(t + s + 1)$$

the last step because $\epsilon \leq 1$ gives $\epsilon^2 \leq \epsilon$. Now let $\delta > 0$ be arbitrary and apply this with $\epsilon = \min\{1, \delta/(t + s + 1)\}$, which is positive since $t + s + 1 > 0$. It gives $\sigma \leq ts + \delta$ for every $\delta > 0$, and σ is real because it is bounded above by a real number, so $\sigma \leq ts$ by proposition 1.4.7(1).

Step 2: $ts \leq \sigma$. Put $t'_n = 1/t_n$ and $u'_n = t_n u_n$. Each t'_n is positive, $t'_n \rightarrow 1/t$ by theorem 2.1.6 since $t \neq 0$, and $1/t > 0$. Each u'_n is nonnegative. Step 1, applied to these two sequences, gives

$$\limsup_{n \rightarrow \infty} (t'_n u'_n) \leq \frac{1}{t} \sigma$$

with the right side read as $+\infty$ when $\sigma = +\infty$. Since $t'_n u'_n = u_n$ for every n , the left side is s . If $\sigma = +\infty$ then $ts \leq \sigma$ holds outright. If σ is real then $s \leq \sigma/t$ is an inequality between real numbers, and multiplying by $t > 0$ gives $ts \leq \sigma$. Together with Step 1 this proves part (1).

Part (2). Suppose $R < \rho$. Then R is a real number, since ρ is, and $R \geq 0$. Put $x = a + (R + \rho)/2$, so that $|x - a| = (R + \rho)/2$. From $R < \rho$ it follows that $R < (R + \rho)/2 < \rho$. The left inequality and theorem 2.6.3(2) make the series diverge at x , while the right inequality and the hypothesis make it converge at x . A series cannot both converge and diverge, so $R \geq \rho$, completing the proof.

The first theorem upgrades the pointwise convergence of theorem 2.6.3 to uniform convergence, which is what theorem 4.4.5 and theorem 4.4.8 need. Uniformity cannot be expected on all of I : the geometric series $\sum_{n \geq 0} x^n$ has partial sums $s_N(x) = (1 - x^{N+1})/(1 - x)$ by example 2.4.7, and the difference $f(x) - s_N(x) = x^{N+1}/(1 - x)$ is unbounded on $(-1, 1)$ for every fixed N , so no single N makes the error small at every point of the interval. Retreating from the endpoints by any fixed positive amount removes the obstruction.

Theorem 4.5.2: Uniform Convergence of a Power Series on Closed Bounded Subintervals

Let $\sum_{n \geq 0} c_n(x - a)^n$ have radius of convergence $R > 0$, and let I , f and s_N be as above.

1. For every closed bounded interval $[\beta, \gamma] \subseteq I$, the sequence (s_N) converges to f uniformly on $[\beta, \gamma]$.
2. f is continuous at every point of I .

Proof:

Part (1). Put $r = \max\{|\beta - a|, |\gamma - a|\}$. If $x \in [\beta, \gamma]$ then $x - a \leq \gamma - a \leq |\gamma - a| \leq r$ and $-(x - a) \leq a - \beta \leq |\beta - a| \leq r$, the two middle inequalities by proposition 1.3.8(1), so $|x - a| \leq r$ by proposition 1.3.8(2). Both β and γ lie in I , so $|\beta - a| < R$ and $|\gamma - a| < R$, whence $r < R$. Set $M_0 = |c_0|$ and $M_n = |c_n| r^n$ for $n \geq 1$. For $x \in [\beta, \gamma]$ the reading $(x - a)^0 = 1$ of definition 2.6.2 gives $|c_0(x - a)^0| = M_0$, and for $n \geq 1$,

$$|c_n(x - a)^n| = |c_n| |x - a|^n \leq |c_n| r^n = M_n$$

the first equality by proposition 1.3.8(3) applied n times, and the inequality by proposition 3.3.11 when $|x - a| < r$ and by equality of the two sides when $|x - a| = r$. The series $\sum_{n \geq 0} M_n$ converges: since $r < R$, theorem 2.6.3(1) makes $\sum_{n \geq 0} c_n r^n$ converge absolutely, and the absolute values of its terms are $|c_0| = M_0$ and $|c_n| r^n = M_n$ for $n \geq 1$. Theorem 4.4.7 is stated over $\mathbb{N}$, which starts at 1, so apply it to the functions $x \mapsto c_n(x - a)^n$ on $[\beta, \gamma]$ for $n \geq 1$ against the bounds M_n , whose sum converges as a tail of a convergent series by proposition 2.4.5(1). It gives uniform convergence of $\sum_{n \geq 1} c_n(x - a)^n$ on $[\beta, \gamma]$, and adding the constant function c_0 changes no partial sum by more than 0, so the full series $\sum_{n \geq 0} c_n(x - a)^n$ converges uniformly on $[\beta, \gamma]$, which is uniform convergence of the sequence (s_N) of its partial sums. The limit is f ,

because at each $x \in [\beta, \gamma] \subseteq I$ the series converges to $f(x)$ by definition of f .

Part (2). Fix $x_0 \in I$ and choose a real r with $|x_0 - a| < r < R$, which is possible because $|x_0 - a| < R$: take r halfway between the two when R is finite, and $r = |x_0 - a| + 1$ when $R = +\infty$. Then $[a - r, a + r] \subseteq I$, and part (1) makes (s_N) converge to f uniformly on $[a - r, a + r]$. Each s_N is a polynomial, hence continuous by example 3.2.6, so theorem 4.4.5 makes the restriction $f|_{[a-r, a+r]}$ continuous, in particular at x_0 .

It remains to pass from the restriction to f itself. Put $\eta = r - |x_0 - a| > 0$. If $|x - x_0| < \eta$ then $|x - a| \leq |x - x_0| + |x_0 - a| < r$ by proposition 1.3.8(4), so $(x_0 - \eta, x_0 + \eta) \subseteq (a - r, a + r) \subseteq I$. The two sets $I \cap (x_0 - \eta, x_0 + \eta)$ and $[a - r, a + r] \cap (x_0 - \eta, x_0 + \eta)$ are therefore both equal to $(x_0 - \eta, x_0 + \eta)$, and f and $f|_{[a-r, a+r]}$ agree on it. Since I is an open interval, lemma 4.1.1 makes x_0 a limit point of I , and continuity at x_0 is the statement $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ by proposition 3.2.3(1). Applying proposition 3.1.22(2) once with $E = [a - r, a + r]$ and once with $E = I$, both with $c = x_0$ and the radius η , the limit of $f|_{[a-r, a+r]}$ at x_0 and the limit of f at x_0 exist together and are equal. The first is $f(x_0)$, so f is continuous at x_0 , as we sought to show.

Theorem 4.4.8 places its uniformity hypothesis on the sequence of derivatives and asks the functions themselves only to converge at one point. The derivatives of the partial sums s_N are the partial sums of the differentiated series, so theorem 4.5.2 gives that hypothesis on every closed bounded subinterval of I once the differentiated series is known to have radius R , which is the first part of the next theorem.

Theorem 4.5.3: A Power Series May Be Differentiated Term by Term

Let $\sum_{n \geq 0} c_n(x - a)^n$ have radius of convergence $R > 0$ and sum f on I .

1. The series $\sum_{n \geq 1} n c_n(x - a)^{n-1}$ has radius of convergence R .
2. f is differentiable at every point of I , and

$$f'(x) = \sum_{n=1}^{\infty} n c_n(x - a)^{n-1} \quad (x \in I)$$

3. f has derivatives of every order on I , and for every integer $k \geq 0$,

$$f^{(k)}(x) = \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) c_n(x - a)^{n-k} \quad (x \in I)$$

the product $n(n-1) \cdots (n-k+1)$ being read as 1 when $k = 0$.

Proof:

Step 1: an auxiliary series with radius R . Let $\alpha = \limsup_n |c_n|^{1/n}$, the upper limit over $n \geq 1$ that definition 2.6.2 uses to define R . Consider the power series $\sum_{n \geq 0} \tilde{c}_n(x - a)^n$ with $\tilde{c}_0 = 0$ and $\tilde{c}_n = n c_n$ for $n \geq 1$. For $n \geq 1$ the numbers $n^{1/n}$ and $|c_n|^{1/n}$ are nonnegative and their product raised to the n -th power is $n|c_n|$, so uniqueness in theorem 1.3.12 gives

$$|\tilde{c}_n|^{1/n} = (n|c_n|)^{1/n} = n^{1/n} |c_n|^{1/n}$$

Now $n^{1/n} \rightarrow 1$ by proposition 2.6.1(1), every $n^{1/n}$ is positive, every $|c_n|^{1/n}$ is nonnegative, and

$1 > 0$, so lemma 4.5.1(1) gives $\limsup_n |\tilde{c}_n|^{1/n} = \alpha$. The formula of definition 2.6.2 determines a radius from this upper limit alone, so $\sum_{n \geq 0} \tilde{c}_n(x-a)^n$ has radius R .

Step 2: the differentiated series has radius R . Re-indexing by $m = n - 1$ writes $\sum_{n \geq 1} nc_n(x-a)^{n-1}$ as the power series $\sum_{m \geq 0} d_m(x-a)^m$ with $d_m = (m+1)c_{m+1}$. The two have the same partial sums, so they converge at the same points and to the same values. Let R' be the radius of convergence of that power series. For $x \neq a$ the terms of the differentiated series and of the series of Step 1 are related by

$$\tilde{c}_n(x-a)^n = (x-a) \cdot nc_n(x-a)^{n-1} \quad (n \geq 1)$$

so by proposition 2.4.5(1), applied with the constant $x-a$ in one direction and with the constant $1/(x-a)$ in the other, one of the two series converges at x exactly when the other does. The term with $n = 0$ of the first series is 0 and contributes nothing to its partial sums, so the starting index plays no part in this.

Let ρ be any real number with $0 < \rho < R$. At every x with $|x-a| < \rho$ the series $\sum \tilde{c}_n(x-a)^n$ converges by theorem 2.6.3(1), hence so does the differentiated series when $x \neq a$. At $x = a$ every term of the differentiated series past the first vanishes, so it converges there too. Lemma 4.5.1(2) now gives $R' \geq \rho$. If R' were smaller than R there would be such a ρ with $R' < \rho < R$, namely the point halfway between them when R is finite and $R' + 1$ when $R = +\infty$, and $R' \geq \rho$ would fail. So $R' \geq R$.

Suppose $R' > R$. Then R is real, and there is a real ρ with $R < \rho < R'$, again the halfway point when R' is finite and $R + 1$ when $R' = +\infty$. At $x = a + \rho$, which differs from a because $\rho > 0$, the differentiated series converges by theorem 2.6.3(1) applied with radius R' , so $\sum \tilde{c}_n(x-a)^n$ converges there. But $|x-a| = \rho > R$, so theorem 2.6.3(2) applied with radius R makes that series diverge. This contradiction gives $R' \leq R$, and with the previous paragraph $R' = R$, which is part (1).

Step 3: differentiability. Write $g(x) = \sum_{n=1}^{\infty} nc_n(x-a)^{n-1}$ for $x \in I$, which is defined by Step 2 and theorem 2.6.3(1). Fix $x_0 \in I$ and choose a real r with $|x_0 - a| < r < R$ as in the proof of theorem 4.5.2, so that $[a-r, a+r] \subseteq I$.

Each partial sum s_N is a polynomial, and theorem 4.1.7 together with theorem 4.1.8 applied to the composition of $t \mapsto t^n$ with $x \mapsto x-a$ gives

$$s'_N(x) = \sum_{n=1}^N nc_n(x-a)^{n-1}$$

which is the N -th partial sum of the differentiated series. By theorem 4.5.2(1) applied to the original series, (s_N) converges to f uniformly on $[a-r, a+r]$, so in particular $(s_N(x))$ converges at every point of that interval. By Step 2 the differentiated series is a power series centred at a with radius R , so theorem 4.5.2(1) applied to it makes (s'_N) converge to g uniformly on $[a-r, a+r]$. These are the hypotheses of theorem 4.4.8 on the interval $[a-r, a+r]$: a sequence of differentiable functions, convergent at a point of the interval, whose derivatives converge uniformly on it. That theorem gives a function to which (s_N) converges, differentiable on $[a-r, a+r]$ with derivative g there. The limit of (s_N) is f on $[a-r, a+r]$, and limits are unique by proposition 2.1.4, so that function is $f|_{[a-r, a+r]}$, which is therefore differentiable with derivative g on $[a-r, a+r]$.

Put $\eta = r - |x_0 - a| > 0$, so that $(x_0 - \eta, x_0 + \eta) \subseteq (a-r, a+r)$ as in the proof of theorem 4.5.2(2). The difference quotient of f at x_0 and that of $f|_{[a-r, a+r]}$ at x_0 are the same function on $(x_0 - \eta, x_0 + \eta) \setminus \{x_0\}$, so proposition 3.1.22(2), applied once with $E = [a-r, a+r] \setminus \{x_0\}$ and

once with $E = I \setminus \{x_0\}$, makes the two limits at x_0 exist together and be equal. Hence f is differentiable at x_0 with $f'(x_0) = g(x_0)$. As x_0 was an arbitrary point of I , this is part (2).

Step 4: all orders. Argue by induction on k that $f^{(k)}$ exists on I and is the sum there of the power series

$$\sum_{n \geq k} n(n-1) \cdots (n-k+1) c_n (x-a)^{n-k}$$

which has radius of convergence R . For $k = 0$ this is the hypothesis of the theorem. Assume it for k . Re-indexed by $m = n - k$, the displayed series is the power series $\sum_{m \geq 0} b_m (x-a)^m$ with $b_m = (m+k)(m+k-1) \cdots (m+1) c_{m+k}$, of radius R , and its sum is $f^{(k)}$. Parts (1) and (2), already proved and applicable to any power series of positive radius, give that $f^{(k)}$ is differentiable on I with

$$f^{(k+1)}(x) = \sum_{m=1}^{\infty} m b_m (x-a)^{m-1}$$

and that this series has radius R . Undoing the re-indexing, the term with index $m \geq 1$ is $(n-k) \cdot n(n-1) \cdots (n-k+1) c_n (x-a)^{n-k-1}$ with $n = m+k \geq k+1$, and appending the factor $n-k$ to the product turns it into $n(n-1) \cdots (n-k)$, so the series is

$$\sum_{n \geq k+1} n(n-1) \cdots (n-k) c_n (x-a)^{n-(k+1)}$$

which is the assertion for $k+1$. The induction is complete, and part (3) follows, completing the proof.

Evaluating part (3) at the centre leaves a single term, because every other term of the series carries a positive power of $x-a$. What comes out is that the coefficients of a power series are recoverable from its sum.

Corollary 4.5.4: The Coefficients Are the Taylor Coefficients of the Sum

Let $\sum_{n \geq 0} c_n (x-a)^n$ have radius of convergence $R > 0$ and sum f on I . Then

$$c_k = \frac{f^{(k)}(a)}{k!} \quad (k \geq 0)$$

In particular, if two power series centred at a have positive radii of convergence and the same sum at every point of some interval $(a-\rho, a+\rho)$ with $\rho > 0$, their coefficients coincide.

Proof:

Fix $k \geq 0$ and put $x = a$ in theorem 4.5.3(3). The term with $n = k$ is $k(k-1) \cdots 1 \cdot c_k (a-a)^0 = k! c_k$, using the reading $(a-a)^0 = 1$ of definition 2.6.2, and every term with $n > k$ carries the factor $(a-a)^{n-k} = 0$. So the series has partial sums equal to $k! c_k$ from index k onward, and $f^{(k)}(a) = k! c_k$. Dividing by $k! \neq 0$ gives the displayed formula.

For the last statement, let $\sum_{n \geq 0} c_n (x-a)^n$ and $\sum_{n \geq 0} b_n (x-a)^n$ have positive radii R_1 and R_2 and equal sums on $(a-\rho, a+\rho)$. Shrinking ρ if necessary, assume $\rho \leq R_1$ and $\rho \leq R_2$, which changes no hypothesis. Write f and h for the two sums. They agree at every point of the open interval $(a-\rho, a+\rho)$, so at each point of that interval their difference quotients agree on a punctured neighbourhood, and $f' = h'$ there by proposition 3.1.22(2). An induction on k , each step repeating

that argument for $f^{(k)}$ and $h^{(k)}$, gives $f^{(k)} = h^{(k)}$ on $(a - \rho, a + \rho)$ for every k , and in particular at a . The displayed formula applied to each series gives $c_k = f^{(k)}(a)/k! = h^{(k)}(a)/k! = b_k$ for every k , as we sought to show.

The corollary settles the relationship between a power series and the Taylor polynomials of definition 4.3.1: the N -th partial sum s_N is precisely the Taylor polynomial of f of order N at a . So a function that is the sum of a power series about a has one candidate expansion there and no other, and theorem 4.3.3 measures how far the truncations of that one candidate sit from f . It says nothing yet about which functions are sums of power series, which is the question of section 4.7.

Example 4.5.5: Differentiating the Geometric Series

The series $\sum_{n \geq 0} x^n$ has $|c_n|^{1/n} = 1$ for every $n \geq 1$, and the upper limit of a constant sequence is that constant by theorem 2.2.12(5), so $\alpha = 1$ and its radius is $R = 1$ by definition 2.6.2. Its sum on $(-1, 1)$ is $f(x) = 1/(1 - x)$ by example 2.4.7.

Theorem 4.5.3(2) gives $f'(x) = \sum_{n \geq 1} nx^{n-1}$ on $(-1, 1)$, while theorem 4.1.7 applied to the quotient $1/(1 - x)$ gives $f'(x) = 1/(1 - x)^2$. Comparing the two,

$$\sum_{n=1}^{\infty} nx^{n-1} = \frac{1}{(1-x)^2} \quad (-1 < x < 1)$$

Differentiating k times, an induction using the quotient rule gives $f^{(k)}(x) = k!/(1 - x)^{k+1}$, and theorem 4.5.3(3) gives $f^{(k)}(x) = \sum_{n \geq k} n(n-1) \cdots (n-k+1)x^{n-k}$. At $x = 0$ the first expression is $k!$, so $f^{(k)}(0)/k! = 1$, which is the coefficient $c_k = 1$ that the series started with, as corollary 4.5.4 requires.

A power series about a carries information about f near every interior point, not only near a , because the derivatives $f^{(m)}(b)$ at an interior point b are computable from the coefficients. The next theorem shows that assembling them into a power series about b reproduces f near b . The proof rearranges a double array, and the hypothesis that makes the rearrangement legitimate is absolute convergence of the rows, which is why the reach of the new expansion is governed by $|x - b| + |b - a|$.

Theorem 4.5.6: Re-expansion of a Power Series About a New Centre

Let $\sum_{n \geq 0} c_n(x - a)^n$ have radius of convergence $R > 0$ and sum f on I , and let $b \in I$. Then for every $x \in \mathbb{R}$ with

$$|x - b| + |b - a| < R$$

the point x lies in I , the series $\sum_{m \geq 0} \frac{f^{(m)}(b)}{m!}(x - b)^m$ converges absolutely, and

$$f(x) = \sum_{m=0}^{\infty} \frac{f^{(m)}(b)}{m!}(x - b)^m$$

Proof:

Fix such an x and put $\sigma = |x - b| + |b - a|$, so $0 \leq \sigma < R$. By proposition 1.3.8(4), $|x - a| = |(x - b) + (b - a)| \leq \sigma < R$, so $x \in I$.

Step 1: the array. For integers $i, j \geq 1$ set $n = i - 1$ and $m = j - 1$ and define

$$u_{ij} = \begin{cases} \binom{n}{m} c_n (b-a)^{n-m} (x-b)^m & \text{if } m \leq n \\ 0 & \text{if } m > n \end{cases}$$

For fixed i only finitely many entries are nonzero, so $\sum_{j \geq 1} |u_{ij}|$ converges, and by proposition 1.3.8(3) its sum is

$$b_i = |c_n| \sum_{m=0}^n \binom{n}{m} |b-a|^{n-m} |x-b|^m = |c_n| \sigma^n$$

the second equality being the binomial theorem [4, the binomial theorem] applied to $(|x-b| + |b-a|)^n$. Since $\sigma < R$, theorem 2.6.3(1) makes $\sum_{n \geq 0} c_n \sigma^n$ converge absolutely, and the absolute values of its terms are exactly the numbers b_i . So $\sum_{i \geq 1} b_i$ converges, and the hypotheses of theorem 2.5.8(2) hold for the array (u_{ij}) .

Step 2: the row sums. For fixed i , the binomial theorem again gives

$$\sum_{j \geq 1} u_{ij} = c_n \sum_{m=0}^n \binom{n}{m} (b-a)^{n-m} (x-b)^m = c_n ((x-b) + (b-a))^n = c_n (x-a)^n$$

so $\sum_{i \geq 1} (\sum_{j \geq 1} u_{ij})$ is the original series evaluated at x , whose sum is $f(x)$ because $x \in I$.

Step 3: the column sums. Fix j , so $m = j - 1$ is fixed. Since $b \in I$, theorem 4.5.3(3) applies at the point b and gives

$$f^{(m)}(b) = \sum_{n=m}^{\infty} n(n-1) \cdots (n-m+1) c_n (b-a)^{n-m}$$

the series on the right being convergent. Cancelling $(n-m)!$ in $\binom{n}{m} = n!/(m!(n-m)!)$ gives $n(n-1) \cdots (n-m+1) = m! \binom{n}{m}$, so dividing the displayed identity by $m!$ and then multiplying by the constant $(x-b)^m$, both steps by proposition 2.4.5(1),

$$\sum_{i \geq 1} u_{ij} = \sum_{n=m}^{\infty} \binom{n}{m} c_n (b-a)^{n-m} (x-b)^m = \frac{f^{(m)}(b)}{m!} (x-b)^m$$

Step 4. Theorem 2.5.8(2) states that the series of column sums converges absolutely and that the two iterated sums agree. Reading Steps 2 and 3 into that identity gives

$$f(x) = \sum_{m=0}^{\infty} \frac{f^{(m)}(b)}{m!} (x-b)^m$$

with the series on the right absolutely convergent, as we sought to show.

When R is finite the condition $|x-b| + |b-a| < R$ says $|x-b| < R - |b-a|$, so the re-expanded series converges at every point of an interval of radius $R - |b-a|$ about b , and lemma 4.5.1(2) makes its radius of convergence at least $R - |b-a|$. That is the distance from b to the nearer endpoint of I , and it is the most the theorem promises. The bound is sometimes exactly the radius of the recentred series, as the next example shows, and sometimes far below it, as the exercises show.

Example 4.5.7: The Geometric Series Recentred at 1/2

Take $f(x) = 1/(1-x) = \sum_{n \geq 0} x^n$ on $I = (-1, 1)$, so $a = 0$ and $R = 1$, and put $b = 1/2$. From $f^{(m)}(x) = m!/(1-x)^{m+1}$ in example 4.5.5,

$$\frac{f^{(m)}(1/2)}{m!} = \frac{1}{(1/2)^{m+1}} = 2^{m+1}$$

so theorem 4.5.6 gives $f(x) = \sum_{m \geq 0} 2^{m+1}(x - 1/2)^m$ whenever $|x - 1/2| + 1/2 < 1$, that is, for $|x - 1/2| < 1/2$.

The same expansion comes out of a direct computation. For $|x - 1/2| < 1/2$ the number $y = 2(x - 1/2)$ satisfies $|y| < 1$, and

$$\frac{1}{1-x} = \frac{1}{\frac{1}{2} - (x - \frac{1}{2})} = \frac{2}{1-y} = 2 \sum_{m=0}^{\infty} y^m = \sum_{m=0}^{\infty} 2^{m+1} \left(x - \frac{1}{2}\right)^m$$

by example 2.4.7 and proposition 2.4.5(1). For $|x - 1/2| > 1/2$ the number y satisfies $|y| > 1$, so the terms $2y^m$ of the recentred series have absolute value at least 2 and do not tend to 0, and proposition 2.4.3 makes the series diverge. So here the radius of the recentred series is exactly $R - |b - a| = 1/2$.

The re-expansion turns information about f near one interior point into information about its coefficients there. If f vanishes on a set having a limit point x_0 in I , the expansion about x_0 turns out to have all its coefficients equal to 0, so f vanishes on a whole subinterval about x_0 , and a supremum argument then carries the vanishing out to all of I . A sequence of distinct zeros of f converging to an interior point already meets the hypothesis.

Theorem 4.5.8: The Identity Theorem

Let $\sum_{n \geq 0} c_n(x-a)^n$ have radius of convergence $R > 0$ and sum f on I .

1. If the set $E = \{x \in I \mid f(x) = 0\}$ has a limit point belonging to I , then $c_n = 0$ for every $n \geq 0$, and f vanishes at every point of I .
2. Let $\rho > 0$ be real and suppose the two power series $\sum_{n \geq 0} c_n(x-a)^n$ and $\sum_{n \geq 0} b_n(x-a)^n$ both converge at every point of $J = (a - \rho, a + \rho)$, with sums f and h there. If $\{x \in J \mid f(x) = h(x)\}$ has a limit point belonging to J , then $c_n = b_n$ for every $n \geq 0$.

Proof:

Step 1: f vanishes on a subinterval about any limit point of E lying in I . Let $x_0 \in I$ be a limit point of E and put $\rho_0 = R - |x_0 - a|$ when R is finite, and $\rho_0 = 1$ when $R = +\infty$. In both cases $\rho_0 > 0$, and $|x - x_0| < \rho_0$ implies $|x - x_0| + |x_0 - a| < R$. The claim is that f vanishes at every point of $(x_0 - \rho_0, x_0 + \rho_0)$.

First, $f(x_0) = 0$. For each $j \geq 1$, definition 3.1.5 provides an $x_j \in E$ with $0 < |x_j - x_0| < 1/j$, and $x_j \rightarrow x_0$ because $1/j \rightarrow 0$. Each x_j lies in I and $f(x_j) = 0$, and f is continuous at x_0 by theorem 4.5.2(2), so proposition 3.2.4 gives $f(x_0) = \lim_j f(x_j) = 0$.

By theorem 4.5.6 with $b = x_0$, writing $d_m = f^{(m)}(x_0)/m!$,

$$f(x) = \sum_{m=0}^{\infty} d_m(x-x_0)^m \quad (|x-x_0| < \rho_0)$$

Suppose some d_m is nonzero and let k be the least index with $d_k \neq 0$. Since $d_0 = f(x_0) = 0$, $k \geq 1$. Define

$$g(x) = \sum_{m=0}^{\infty} d_{k+m}(x-x_0)^m$$

for $|x-x_0| < \rho_0$. This converges: at $x = x_0$ every term past the first vanishes, and for $0 < |x-x_0| < \rho_0$ the series $\sum_{m \geq 0} d_{k+m}(x-x_0)^{k+m}$ is a tail of the convergent series above, hence converges, and multiplying by the constant $(x-x_0)^{-k}$ preserves convergence by proposition 2.4.5(1). So the power series defining g converges at every point of $(x_0 - \rho_0, x_0 + \rho_0)$, its radius is at least ρ_0 by lemma 4.5.1(2), and g is continuous on that interval by theorem 4.5.2(2).

Because $d_m = 0$ for $m < k$, another application of proposition 2.4.5(1), this time with the constant $(x-x_0)^k$, gives

$$f(x) = \sum_{m \geq k} d_m(x-x_0)^m = (x-x_0)^k g(x) \quad (|x-x_0| < \rho_0)$$

where the identity also holds at $x = x_0$, both sides being 0 there since $k \geq 1$. Now $g(x_0) = d_k \neq 0$. Applying continuity of g at x_0 with $\epsilon = |d_k|/2 > 0$ produces a δ with $0 < \delta \leq \rho_0$ such that $|g(x) - d_k| < |d_k|/2$ for every x with $|x-x_0| < \delta$, and then proposition 1.3.8(5) gives

$$|g(x)| \geq |d_k| - |g(x) - d_k| > \frac{|d_k|}{2} > 0$$

For $0 < |x-x_0| < \delta$ both factors of $(x-x_0)^k g(x)$ are nonzero, so $f(x) \neq 0$ by proposition 1.1.2(3), which makes a product vanish only when one of its factors does. Then no point of E satisfies $0 < |x-x_0| < \delta$, contradicting the assumption that x_0 is a limit point of E . So every d_m is 0, and the displayed expansion makes f vanish on $(x_0 - \rho_0, x_0 + \rho_0)$, which is the claim.

Step 2: spreading the vanishing across I . Let $x_0 \in I$ be a limit point of E , which the hypothesis provides, and let $y \in I$ with $y > x_0$. Put

$$S = \{ t \in [x_0, y] \mid f(x) = 0 \text{ for every } x \in [x_0, t] \}$$

Step 1 gives $f(x_0) = 0$, so $x_0 \in S$ and S is nonempty, and S is bounded above by y . Let $\tau = \sup S$, which exists by theorem 1.2.4 and satisfies $x_0 \leq \tau \leq y$.

First, f vanishes on $[x_0, \tau]$. For $x_0 \leq x < \tau$, proposition 1.2.6 applied with $\epsilon = \tau - x$ produces a $t \in S$ with $t > x$, and then $x \in [x_0, t]$ gives $f(x) = 0$. For the endpoint, if $\tau = x_0$ then $f(\tau) = 0$ already. If $\tau > x_0$ then the points $x_j = \tau - (\tau - x_0)/(j+1)$ for $j \geq 1$ lie in $[x_0, \tau)$ and converge to τ , so $f(\tau) = \lim_j f(x_j) = 0$ by theorem 4.5.2(2) and proposition 3.2.4. Hence $\tau \in S$.

Suppose $\tau < y$. Then τ is a limit point of E : if $\tau = x_0$ this is the hypothesis, and if $\tau > x_0$ then every punctured neighbourhood of τ contains points of $[x_0, \tau)$, all of which lie in E . Step 1 applied at τ gives a $\rho_1 > 0$ with f vanishing on $(\tau - \rho_1, \tau + \rho_1)$. Choose t strictly between τ and $\min\{y, \tau + \rho_1\}$, which exceeds τ . Every $x \in [x_0, t]$ satisfies either $x \leq \tau$, where $f(x) = 0$ because $\tau \in S$, or $\tau < x \leq t < \tau + \rho_1$, where $f(x) = 0$ by Step 1. So $t \in S$ with $t > \tau = \sup S$, a contradiction. Hence $\tau = y$ and $f(y) = 0$.

For $y \in I$ with $y < x_0$ the argument runs the same way with every inequality reversed: take $S = \{t \in [y, x_0] \mid f(x) = 0 \text{ for every } x \in [t, x_0]\}$ and $\tau = \inf S$, which exists by corollary 1.2.5, use lemma 1.3.1 in place of proposition 1.2.6 to see that f vanishes on $(\tau, x_0]$ and, by continuity at τ , at τ itself, and then use Step 1 at τ to contradict $\tau > y$. So f vanishes at every point of I .

Step 3: the coefficients. A function vanishing at every point of an interval has vanishing derivative there, since every difference quotient is 0, so an induction on k gives $f^{(k)}(x) = 0$ for every $k \geq 0$ and every $x \in I$. By corollary 4.5.4, $c_k = f^{(k)}(a)/k! = 0$. This proves part (1).

Part (2). By proposition 2.4.5(1) the series $\sum_{n \geq 0} (c_n - b_n)(x - a)^n$ converges at every point of J , with sum $f - h$ there. Its radius of convergence R'' is at least ρ by lemma 4.5.1(2), so $J \subseteq (a - R'', a + R'')$. The set where its sum vanishes contains $\{x \in J \mid f(x) = h(x)\}$, which has a limit point in J and hence in $(a - R'', a + R'')$. Part (1) applied to this series gives $c_n - b_n = 0$ for every n , completing the proof.

Two power series about the same centre whose sums agree along a sequence of distinct points converging to an interior point have the same coefficients, so an identity between power series that has been checked along such a sequence holds at every point of the common interval. Corollary 4.5.4 reaches the same conclusion from the values of f and of all its derivatives at the single point a , and the identity theorem reaches it from the values of f alone on a set that need not contain a at all.

Theorem 2.6.3 says nothing at the endpoints $a \pm R$, and all three behaviours occur there: among the series $\sum x^n$, $\sum x^n/n$ and $\sum x^n/n^2$ of example 2.6.4, all of radius 1, the first diverges at both endpoints, the second converges at $x = -1$ and diverges at $x = 1$, and the third converges at both. When the series does converge at an endpoint, its sum there is the limit of f from inside the interval.

Theorem 4.5.9: Abel's Limit Theorem

Let $\sum_{n \geq 0} c_n(x - a)^n$ have finite radius of convergence $R > 0$ and sum f on $I = (a - R, a + R)$.

1. If $\sum_{n \geq 0} c_n R^n$ converges, then

$$\lim_{x \rightarrow (a+R)^-} f(x) = \sum_{n=0}^{\infty} c_n R^n$$

2. If $\sum_{n \geq 0} c_n (-R)^n$ converges, then

$$\lim_{x \rightarrow (a-R)^+} f(x) = \sum_{n=0}^{\infty} c_n (-R)^n$$

Proof:

Step 1: the normalized statement. Let $(b_n)_{n \geq 0}$ be real numbers with $\sum_{n \geq 0} b_n$ convergent, write $s_N = \sum_{n=0}^N b_n$ and $s = \lim_N s_N$, and set $s_{-1} = 0$. The claim is that for every $\epsilon > 0$ there is a δ_0 with $0 < \delta_0 \leq 1$ such that

$$\left| \sum_{n=0}^{\infty} b_n u^n - s \right| < \epsilon \quad \text{whenever } 1 - \delta_0 < u < 1$$

The series there converges for every u with $|u| < 1$: the sequence $(s_N)_{N \geq 0}$ converges, hence is

bounded by proposition 2.1.5, say $|s_N| \leq M$ for every $N \geq 0$, and $|s_{-1}| = 0 \leq M$ as well, so

$$|b_n u^n| = |s_n - s_{n-1}| |u|^n \leq 2M |u|^n$$

and theorem 2.4.13(1) applies with the geometric series of example 2.4.7.

Fix u with $|u| < 1$. For $N \geq 1$, substituting $b_n = s_n - s_{n-1}$ and separating the two sums,

$$\sum_{n=0}^N b_n u^n = \sum_{n=0}^N s_n u^n - \sum_{n=1}^N s_{n-1} u^n = \sum_{n=0}^N s_n u^n - u \sum_{n=0}^{N-1} s_n u^n = (1-u) \sum_{n=0}^{N-1} s_n u^n + s_N u^N$$

the term $n = 0$ having been dropped from the second sum because $s_{-1} = 0$. The series $\sum_{n \geq 0} s_n u^n$ converges absolutely by theorem 2.4.13(1), compared with $M|u|^n$, and $s_N u^N \rightarrow 0$ because $|s_N u^N| \leq M|u|^N$ and $|u|^N \rightarrow 0$ by proposition 2.1.10(3) when $u \neq 0$, the case $u = 0$ being immediate. Letting $N \rightarrow \infty$ and using theorem 2.1.6,

$$\sum_{n=0}^{\infty} b_n u^n = (1-u) \sum_{n=0}^{\infty} s_n u^n$$

Also $(1-u) \sum_{n \geq 0} u^n = 1$ by example 2.4.7, so $s = (1-u) \sum_{n \geq 0} s_n u^n$, and subtracting the two identities through proposition 2.4.5(1),

$$\sum_{n=0}^{\infty} b_n u^n - s = (1-u) \sum_{n=0}^{\infty} (s_n - s) u^n$$

Now let $\epsilon > 0$. Since $s_n \rightarrow s$ there is an $N_1 \geq 1$ with $|s_n - s| < \epsilon/2$ for every $n \geq N_1$. Restrict to $0 \leq u < 1$ and split the last series at N_1 into the finite sum over $n < N_1$ and the tail over $n \geq N_1$, both convergent, and bound the two separately by proposition 1.3.8(4). The finite sum is handled by proposition 2.4.5(2) together with $u^n \leq 1$, and the tail by proposition 2.4.5(3), which applies because the series converges absolutely, together with $|s_n - s| < \epsilon/2$ on the tail and $\sum_{n \geq N_1} u^n \leq \sum_{n \geq 0} u^n = 1/(1-u)$. This gives

$$\left| \sum_{n=0}^{\infty} b_n u^n - s \right| \leq (1-u) \sum_{n=0}^{N_1-1} |s_n - s| + (1-u) \frac{\epsilon}{2} \frac{1}{1-u} = (1-u)M_1 + \frac{\epsilon}{2}$$

where $M_1 = \sum_{n=0}^{N_1-1} |s_n - s| \geq 0$. Put $\delta_0 = \min\{1, \epsilon/(2M_1 + 2)\}$, which is positive. If $1 - \delta_0 < u < 1$ then $u \geq 0$, since $\delta_0 \leq 1$, and $1 - u < \delta_0$, so

$$(1-u)M_1 \leq \delta_0 M_1 \leq \frac{\epsilon M_1}{2M_1 + 2} < \frac{\epsilon}{2}$$

and the two halves add to less than ϵ . This is the claim of Step 1.

Step 2: transferring to the endpoint. Let $\theta = 1$ for part (1) and $\theta = -1$ for part (2), put $b_n = c_n(\theta R)^n$, so that $\sum_{n \geq 0} b_n$ is the convergent series in the hypothesis, and write s for its sum. For $|u| < 1$ the point $x = a + \theta R u$ satisfies $|x - a| = R|u| < R$, so $x \in I$, and

$$f(x) = \sum_{n=0}^{\infty} c_n(\theta R u)^n = \sum_{n=0}^{\infty} b_n u^n$$

Let $\epsilon > 0$ and take δ_0 from Step 1. Put $\delta = R\delta_0 > 0$ and let $c = a + \theta R$, the endpoint in question. Let $x \in I$ satisfy $0 < |x - c| < \delta$, and put $u = \theta(x - a)/R$, so that $|u| < 1$ and $x = a + \theta Ru$. In part (1) membership of I forces $x < c$, so $a + R - \delta < x < a + R$, whence $R - \delta < x - a < R$ and $1 - \delta_0 < u < 1$. In part (2) membership of I forces $x > c$, so $a - R < x < a - R + \delta$, whence $-R < x - a < \delta - R$ and again $1 - \delta_0 < u < 1$. Step 1 therefore gives $|f(x) - s| < \epsilon$. In the notation of definition 3.1.24 with $E = I$, the set E_c^- is all of I in part (1) and E_c^+ is all of I in part (2), and c is a limit point of I in both cases. What has been shown is exactly that the corresponding one-sided limit of f at c exists and equals s , as we sought to show.

The theorem asserts an equality between two numbers that both exist under its hypothesis. It does not define f at the endpoint, and the hypothesis that the series converges there cannot be dropped. The implication runs one way only, and the geometric series shows it: $\sum_{n \geq 0} x^n$ has sum $f(x) = 1/(1 - x)$ on $(-1, 1)$, and f agrees there with a rational function continuous at -1 by example 3.2.6, so proposition 3.2.3(1) and definition 3.1.24 give $\lim_{x \rightarrow (-1)^+} f(x) = 1/2$, while $\sum_{n \geq 0} (-1)^n$ diverges by proposition 2.4.3. A limit from inside can exist while the series at the endpoint has no sum at all.

Exercise 4.5.1

1. Show that the power series $\sum_{n \geq 1} x^n/n$ has radius of convergence 1, and that its sum f satisfies $f'(x) = 1/(1 - x)$ at every $x \in (-1, 1)$.
2. Show that $\sum_{n \geq 1} n^2 x^n$ has radius of convergence 1, and that its sum on $(-1, 1)$ is $x(1 + x)/(1 - x)^3$. (Differentiate the geometric series twice, multiplying by x between the two differentiations to restore the exponents.)
3. Show that there is no power series $\sum_{n \geq 0} c_n x^n$ with radius of convergence greater than $1/2$ whose sum f satisfies $f(1/k) = 0$ for every odd integer $k \geq 3$ and $f(1/2) = 1/2$.
4. Let $\sum_{n \geq 0} c_n x^n$ have radius of convergence $R > 0$ and sum f on $(-R, R)$, and suppose $f(-x) = f(x)$ for every $x \in (-R, R)$. Show that $c_n = 0$ for every odd n . State and prove the corresponding statement for a sum satisfying $f(-x) = -f(x)$.
5. Recentre the geometric series at $b = -1/2$: compute $f^{(m)}(-1/2)/m!$ for $f(x) = 1/(1 - x)$, write down the series that theorem 4.5.6 produces, and determine the exact set of x at which that series converges. Compare what you find with the interval $|x + 1/2| < R - |b|$ on which the theorem guarantees the expansion is valid.
6. Show that $\sum_{n \geq 1} (-1)^{n+1} x^n/n$ has radius of convergence 1 and that its coefficient series converges at $x = 1$, and deduce that its sum f on $(-1, 1)$ satisfies $\lim_{x \rightarrow 1^-} f(x) = \sum_{n \geq 1} (-1)^{n+1}/n$.
7. Show that the hypothesis $t > 0$ in lemma 4.5.1(1) cannot be dropped: exhibit a sequence (t_n) of positive reals with $t_n \rightarrow 0$ and a sequence (u_n) of nonnegative reals for which $\limsup_n (t_n u_n)$ is nonzero, and say which of $\limsup_n u_n$ and $\limsup_n (t_n u_n)$ is finite.
8. Theorem 4.5.3 says that a power series and its differentiated series have the same radius, and says nothing about the endpoints. Exhibit a power series of radius 1 centred at 0 that converges at $x = 1$ while its differentiated series diverges at $x = 1$, and one whose differentiated series converges at $x = -1$ while the series obtained by differentiating twice diverges there.

4.6 The Elementary Functions

Definition 2.6.5 attached the names $\exp$, $\cos$ and $\sin$ to three power series of infinite radius, and theorem 2.6.6 proved one identity about the first of them. Neither statement uses a derivative. Theorem 4.5.3 now differentiates all three, and each derivative is again one of the three: the exponential is its own, and the cosine and sine are each other's up to a sign.

From those three identities the exponential is strictly increasing with range $(0, \infty)$, so it has an inverse, and that inverse is the logarithm. The two together define a^x for real exponents and identify it with the n -th roots of theorem 1.3.12. The cosine changes sign between 0 and 2π , and twice its first positive zero is π .

Theorem 4.6.1: The Exponential Is Its Own Derivative

The function $\exp: \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at every $x \in \mathbb{R}$, and $\exp'(x) = \exp(x)$.

Proof:

The series $\sum_{n \geq 0} x^n/n!$ is the power series centred at 0 with coefficients $c_n = 1/n!$ in the sense of definition 2.6.2, and definition 2.6.5 records that its radius of convergence is $+\infty$. Its interval of convergence is therefore all of $\mathbb{R}$, so theorem 4.5.3 applies at every $x \in \mathbb{R}$: the function $\exp$ is differentiable there and

$$\exp'(x) = \sum_{n \geq 1} n \frac{x^{n-1}}{n!}$$

For $n \geq 1$ the factorial factors as $n! = n \cdot (n-1)!$, so $n/n! = 1/(n-1)!$ and the n -th term of that series is $x^{n-1}/(n-1)!$. Writing $m = n-1$, which runs over $m \geq 0$ as n runs over $n \geq 1$ and leaves every partial sum unchanged apart from its indexing, gives

$$\exp'(x) = \sum_{m \geq 0} \frac{x^m}{m!} = \exp(x)$$

the last equality being definition 2.6.5, as we sought to show.

The equation $\exp' = \exp$ says that the exponential grows at a rate equal to its own value, and it determines every higher derivative at once: applying the theorem k times gives $\exp^{(k)} = \exp$ for every integer $k \geq 0$, by induction on k with the theorem itself as the step. A function whose derivatives are all equal to the function is the extreme case of the growth condition that section 4.7 measures.

The cosine and sine series are power series in which every second coefficient vanishes. Writing $\cos x = \sum_{m \geq 0} c_m x^m$ with $c_{2n} = (-1)^n/(2n)!$ and $c_{2n+1} = 0$ puts the cosine series into the shape of definition 2.6.2, and the same padding with zeros puts the sine series there with the roles of the even and odd indices exchanged. Both radii are $+\infty$ by definition 2.6.5, so theorem 4.5.3 again applies at every real point.

Theorem 4.6.2: The Derivatives of the Cosine and the Sine

The functions $\cos: \mathbb{R} \rightarrow \mathbb{R}$ and $\sin: \mathbb{R} \rightarrow \mathbb{R}$ are differentiable at every $x \in \mathbb{R}$, and

$$\cos'(x) = -\sin x, \quad \sin'(x) = \cos x$$

Proof:

The cosine. With the coefficients $c_{2n} = (-1)^n/(2n)!$ and $c_{2n+1} = 0$, theorem 4.5.3 gives $\cos'(x) = \sum_{m \geq 1} m c_m x^{m-1}$. The terms with m odd vanish, since $c_m = 0$ there. A term with $m = 2n$ and $n \geq 1$ equals

$$2n \frac{(-1)^n}{(2n)!} x^{2n-1} = \frac{(-1)^n}{(2n-1)!} x^{2n-1}$$

because $(2n)! = 2n \cdot (2n-1)!$. Writing $n = k+1$, so that k runs over $k \geq 0$, the exponent $2n-1$ becomes $2k+1$ and the factorial $(2n-1)!$ becomes $(2k+1)!$, while $(-1)^n = (-1)^{k+1} = -(-1)^k$. Hence

$$\cos'(x) = \sum_{k \geq 0} \frac{-(-1)^k}{(2k+1)!} x^{2k+1} = -\sin x$$

the last equality being definition 2.6.5 together with the scalar rule of proposition 2.4.5.

The sine. Write $\sin x = \sum_{m \geq 0} d_m x^m$ with $d_{2n+1} = (-1)^n/(2n+1)!$ and $d_{2n} = 0$. Then $\sin'(x) = \sum_{m \geq 1} m d_m x^{m-1}$, in which the terms with m even vanish, and a term with $m = 2n+1$ and $n \geq 0$ equals

$$(2n+1) \frac{(-1)^n}{(2n+1)!} x^{2n} = \frac{(-1)^n}{(2n)!} x^{2n}$$

because $(2n+1)! = (2n+1) \cdot (2n)!$. Summing over $n \geq 0$ gives definition 2.6.5's cosine series, so $\sin'(x) = \cos x$, completing the proof.

The catalogue of continuous functions in section 3.2 stopped at polynomials, rational functions and the absolute value, because no result available there made the sum of a power series continuous. Differentiability gives it at once.

Example 4.6.3: The Continuity of the Exponential, Cosine and Sine

Each of $\exp$, $\cos$ and $\sin$ is continuous at every point of $\mathbb{R}$. Each is differentiable at every point of $\mathbb{R}$ by theorem 4.6.1 and theorem 4.6.2, and proposition 4.1.5 turns differentiability at a point into continuity at that point. Being continuous at every point of $\mathbb{R}$ is continuity on $\mathbb{R}$ in the sense of definition 3.2.2.

The function $x \mapsto 1/x$ is continuous on $\mathbb{R} \setminus \{0\}$ by theorem 3.2.5, applied to the polynomials 1 and x , the second of which does not vanish there. So theorem 3.2.7 makes $x \mapsto \sin(1/x)$ and $x \mapsto \cos(1/x)$ continuous at every point of $\mathbb{R} \setminus \{0\}$. The same theorem applied to the polynomial x^2 makes $x \mapsto \exp(x^2)$ continuous on $\mathbb{R}$.

The exponents in the three series also fix how the cosine and the sine behave under $x \mapsto -x$. For every $n \geq 0$ the identities $(-1)^{2n} = 1$ and $(-1)^{2n+1} = -1$ give $(-x)^{2n} = x^{2n}$ and $(-x)^{2n+1} = -x^{2n+1}$, so the partial sums of the cosine series at $-x$ are exactly those at x , and the partial sums of the sine series at $-x$ are the negatives of those at x . Passing to the limit with theorem 2.1.6 gives, for every $x \in \mathbb{R}$,

$$\cos(-x) = \cos x, \quad \sin(-x) = -\sin x \quad (4.10)$$

The exponential is positive everywhere and its derivative is itself, so its derivative is positive everywhere, which is the hypothesis of corollary 4.2.7. A continuous strictly increasing function on an interval has a continuous inverse by theorem 3.3.12.

Theorem 4.6.4: The Exponential Is a Strictly Increasing Bijection onto $(0, \infty)$

1. $\exp(x) \geq 1 + x$ for every $x \geq 0$, and $\exp(x) > 0$ for every $x \in \mathbb{R}$.
2. $\exp$ is strictly increasing on $\mathbb{R}$.
3. $\exp(\mathbb{R}) = (0, \infty)$, the map $\exp: \mathbb{R} \rightarrow (0, \infty)$ is a bijection, and its inverse is strictly increasing and continuous.

Proof:

1. Let $x \geq 0$. Every term $x^n/n!$ of the exponential series is then nonnegative, so for $N \geq 1$ the partial sum $s_N = \sum_{n=0}^N x^n/n!$ is at least the sum of its first two terms, giving $s_N \geq 1 + x$. Since $s_N \rightarrow \exp(x)$ by definition 2.6.5, proposition 2.1.8 gives $\exp(x) \geq 1 + x \geq 1 > 0$. Taking $y = -x$ in theorem 2.6.6 and using $\exp(0) = 1$, which is the value of the series at 0, gives

$$\exp(x) \exp(-x) = 1 \quad (x \in \mathbb{R}) \quad (4.11)$$

Now let $x < 0$. Then $-x > 0$, so $\exp(-x) \geq 1 > 0$ by what was just proved, and $\exp(x) = 1/\exp(-x) > 0$ by (4.11) and proposition 1.1.2(7).

2. The set $\mathbb{R}$ is an interval, $\exp$ is differentiable at each of its points by theorem 4.6.1, and the same theorem with part (1) gives $\exp'(x) = \exp(x) > 0$ at every $x \in \mathbb{R}$. Those are the hypotheses of corollary 4.2.7, which makes $\exp$ strictly increasing on $\mathbb{R}$.
3. Put $J = \exp(\mathbb{R})$. Part (1) gives $J \subseteq (0, \infty)$. The function $\exp$ is continuous by example 4.6.3 and strictly monotone by part (2), and $\mathbb{R}$ is an interval, so theorem 3.3.12 applies: J is an interval, $\exp: \mathbb{R} \rightarrow J$ is a bijection, and its inverse is strictly increasing and continuous. It remains to show $(0, \infty) \subseteq J$. Let $y > 0$. By theorem 1.4.1(2) there is an $n \in \mathbb{N}$ with $n > y$, and then $\exp(n) \geq 1 + n > y$ by part (1). By theorem 1.4.1(3) there is an $m \in \mathbb{N}$ with $0 < 1/m < y$. Part (1) gives $\exp(m) \geq 1 + m > m > 0$, so $\exp(-m) = 1/\exp(m) < 1/m < y$ by (4.11) and proposition 1.1.2(7). The two numbers $\exp(-m)$ and $\exp(n)$ lie in J and satisfy $\exp(-m) < y < \exp(n)$, so the betweenness condition of definition 3.1.1 puts $y \in J$. Hence $J = (0, \infty)$, as we sought to show.

Definition 4.6.5: [Real] Logarithm

The *logarithm* is the inverse $\log: (0, \infty) \rightarrow \mathbb{R}$ of the bijection $\exp: \mathbb{R} \rightarrow (0, \infty)$ of theorem 4.6.4(3). So $\log y$ is the unique real number x with $\exp(x) = y$, and

$$\log(\exp x) = x \quad (x \in \mathbb{R}), \quad \exp(\log y) = y \quad (y > 0)$$

Three values come straight from the definition. The exponential series at $x = 0$ has only its constant term 1, so $\exp(0) = 1$ and the logarithm satisfies $\log 1 = 0$. The exponential series at $x = 1$ is the series

$\sum_{n \geq 0} 1/n!$ of example 2.4.18, whose sum is e , so $\exp(1) = e$ and $\log e = 1$. And $\exp(-1) = 1/\exp(1)$ by (4.11), so $\log(1/e) = -1$. By theorem 4.6.4(3) the logarithm is strictly increasing and continuous on $(0, \infty)$, so it is negative on $(0, 1)$ and positive on $(1, \infty)$.

The functional equation of the exponential becomes a statement about the logarithm. For $u, v > 0$ put $s = \log u$ and $t = \log v$, so that $\exp(s) = u$ and $\exp(t) = v$. Then $uv = \exp(s)\exp(t) = \exp(s+t)$ by theorem 2.6.6, and applying $\log$ to both ends gives $\log(uv) = s+t$. That is

$$\log(uv) = \log u + \log v \quad (u, v > 0) \quad (4.12)$$

Taking $v = 1/u$ in (4.12) and using $\log 1 = 0$ gives $\log(1/u) = -\log u$, and combining the two gives $\log(u/v) = \log u - \log v$.

Theorem 4.6.6: The Derivative of the Logarithm

The function $\log$ is differentiable at every $y \in (0, \infty)$, and

$$\log'(y) = \frac{1}{y}$$

Proof:

Fix $y > 0$ and put $x = \log y$, so that $\exp(x) = y$ by definition 4.6.5. The hypotheses of theorem 4.1.9 hold for $\exp$ at x : the domain $\mathbb{R}$ is an interval, $\exp$ is continuous on it by example 4.6.3 and strictly monotone by theorem 4.6.4(2), it is differentiable at x by theorem 4.6.1, and its derivative there is $\exp'(x) = \exp(x) = y \neq 0$ because $y > 0$. The inverse of $\exp$ on the image $(0, \infty)$ is $\log$, by definition 4.6.5. So theorem 4.1.9 makes $\log$ differentiable at $y = \exp(x)$ with

$$\log'(y) = \frac{1}{\exp'(x)} = \frac{1}{\exp(\log y)} = \frac{1}{y}$$

completing the proof.

Proposition 4.6.7: The Logarithm Is Unbounded

$$\lim_{y \rightarrow \infty} \log y = +\infty.$$

Proof:

Let $M \in \mathbb{R}$. By theorem 4.6.4 the function $\exp$ maps $\mathbb{R}$ onto $(0, \infty)$ and $\log$ is its inverse, so $\log$ is defined at $\exp(M) > 0$ and $\log(\exp(M)) = M$. The function $\log$ is increasing, being the inverse of an increasing bijection, so every $y > \exp(M)$ has $\log y > M$. Taking $N = \exp(M)$ meets the at-infinity condition of definition 3.1.28.

Theorem 1.3.12 produced $x^{1/n}$ for integers $n \geq 1$, and repeated multiplication produces x^n . Between them they leave most exponents undefined: the symbol $2^{\sqrt{2}}$ has had no meaning in this book. The pair $\exp$ and $\log$ closes that gap, because $a^n = \exp(n \log a)$ holds for the integer powers already defined and the right-hand side makes sense for every real exponent.

Definition 4.6.8: General Powers

Let $a > 0$ and $x \in \mathbb{R}$. The *power* a^x is the real number

$$a^x = \exp(x \log a)$$

For $a = e$ the definition reads $e^x = \exp(x \log e) = \exp(x)$, since $\log e = 1$, so the notation e^x used informally after theorem 2.6.6 is now the general power at the base e . For $a = 4$ and $x = 1/2$ it reads $4^{1/2} = \exp((\log 4)/2)$, and theorem 4.6.9(4) identifies that number with the square root 2 produced by theorem 1.3.12.

Theorem 4.6.9: The Laws of Real Exponents

Let $a, b > 0$ and let $x, y \in \mathbb{R}$.

1. $a^0 = 1$, $a^1 = a$, $a^{x+y} = a^x a^y$, and $a^{-x} = 1/a^x$.
2. $a^x > 0$ and $\log(a^x) = x \log a$.
3. $(a^x)^y = a^{xy}$ and $(ab)^x = a^x b^x$.
4. For every integer $n \geq 1$, the power a^n is the product of n copies of a , and $a^{1/n}$ is the n -th root of theorem 1.3.12.
5. $e^x = \exp(x)$, where $e = \exp(1)$.

Proof:

1. By definition 4.6.8, $a^0 = \exp(0) = 1$ and $a^1 = \exp(\log a) = a$, the last step being definition 4.6.5. For the third identity,

$$a^{x+y} = \exp((x+y) \log a) = \exp(x \log a + y \log a) = \exp(x \log a) \exp(y \log a) = a^x a^y$$

where the middle equality is distributivity and the next is theorem 2.6.6. Taking $y = -x$ in it gives $a^x a^{-x} = a^0 = 1$, so $a^{-x} = 1/a^x$.

2. a^x is a value of $\exp$, hence positive by theorem 4.6.4(1). Applying $\log$ to $a^x = \exp(x \log a)$ and using definition 4.6.5 gives $\log(a^x) = x \log a$.
3. Using definition 4.6.8 for the outer power and part (2) for the inner one,

$$(a^x)^y = \exp(y \log(a^x)) = \exp(yx \log a) = a^{xy}$$

For the second identity, (4.12) gives $\log(ab) = \log a + \log b$, so

$$(ab)^x = \exp(x(\log a + \log b)) = \exp(x \log a) \exp(x \log b) = a^x b^x$$

again by theorem 2.6.6.

4. The first claim is an induction on n . For $n = 1$ it is $a^1 = a$ from part (1). If a^n is the product of n copies of a , then $a^{n+1} = a^n a^1 = a^n a$ by part (1), which is the product of $n + 1$ copies.

For the second claim put $y = a^{1/n}$, a positive number by part (2). Part (3) with the exponents $1/n$ and n gives $(a^{1/n})^n = a^{(1/n) \cdot n} = a^1 = a$, and the first claim of this part, applied to the positive base y , says that $(a^{1/n})^n = y^n$ is the product of n copies of y . So y is a nonnegative real whose n -th power is a , and the uniqueness clause of theorem 1.3.12 identifies it with the n -th root of a .

5. By definition 4.6.5 the value $\log e$ is the unique x with $\exp(x) = e$, and $\exp(1) = e$, so $\log e = 1$. Hence $e^x = \exp(x \log e) = \exp(x)$, completing the proof.

Part (4) is what makes the notation unambiguous: the symbol $a^{1/n}$ has one meaning, and $4^{1/2} = 2$. Part (2) turns a power into a product under log, and it is the identity from which part (3) and theorem 4.6.10 are computed.

Theorem 4.6.10: The Derivative of a Real Power

Let $\alpha \in \mathbb{R}$ and let $a > 0$.

1. The function $x \mapsto x^\alpha$ on $(0, \infty)$ is differentiable at every $x > 0$, with derivative $\alpha x^{\alpha-1}$.
2. The function $x \mapsto a^x$ on $\mathbb{R}$ is differentiable at every $x \in \mathbb{R}$, with derivative $(\log a) a^x$.

Proof:

1. Write $q(x) = x^\alpha$. By definition 4.6.8 this is $q = \exp \circ u$ with $u(x) = \alpha \log x$ on the interval $(0, \infty)$. Theorem 4.6.6 and the constant-multiple rule of theorem 4.1.7 make u differentiable at every $x > 0$ with $u'(x) = \alpha/x$, and theorem 4.6.1 makes $\exp$ differentiable at $u(x)$ with derivative $\exp(u(x))$. So theorem 4.1.8 applies and gives

$$q'(x) = \exp(u(x)) u'(x) = x^\alpha \cdot \frac{\alpha}{x}$$

By theorem 4.6.9(1) and (4), $x^{-1} = 1/x^1 = 1/x$ and $x^{\alpha-1} = x^\alpha x^{-1}$, so the right-hand side is $\alpha x^{\alpha-1}$.

2. Write $r(x) = a^x$, which is $r = \exp \circ v$ with $v(x) = x \log a$ on the interval $\mathbb{R}$, and $v'(x) = \log a$ by the constant-multiple rule of theorem 4.1.7. As above theorem 4.1.8 gives $r'(x) = \exp(x \log a) \log a = (\log a) a^x$, completing the proof.

For an integer $\alpha = n \geq 1$ part (1) recovers the power rule that theorem 4.1.7 already gives on $(0, \infty)$, and for $\alpha = 1/n$ it recovers the derivative of the n -th root function, which theorem 4.1.9 established. The irrational exponents are what is new here. For those, definition 4.6.8 is the only route to a value of x^α at all, so the rules of section 4.1 have nothing to differentiate. Part (2) identifies e as the only base whose exponential is its own derivative: the values a^x are never 0 by theorem 4.6.9(2), so $(\log a) a^x = a^x$ for all x forces $\log a = 1$, and $\log$ is injective with $\log e = 1$.

How fast the exponential grows is settled by a single term of its series. For $x > 0$ every term is positive, so $\exp(x)$ exceeds each of them separately, and among them is $x^{n+1}/(n+1)!$ for every integer $n \geq 0$.

Proposition 4.6.11: The Exponential Outgrows Every Power

For every integer $n \geq 0$,

$$\lim_{x \rightarrow \infty} \frac{x^n}{\exp(x)} = 0$$

Proof:

Let $x > 0$. Every term $x^k/k!$ of the exponential series is then positive, so for $N \geq n+1$ the partial sum $s_N = \sum_{k=0}^N x^k/k!$ is at least its own term with $k = n+1$, giving $s_N \geq x^{n+1}/(n+1)!$. Since $s_N \rightarrow \exp(x)$, proposition 2.1.8 gives

$$\exp(x) \geq \frac{x^{n+1}}{(n+1)!} > 0$$

Inverting this with proposition 1.1.2(7) and multiplying by $x^n > 0$ using proposition 1.1.2(5),

$$0 < \frac{x^n}{\exp(x)} \leq \frac{(n+1)! x^n}{x^{n+1}} = \frac{(n+1)!}{x}$$

The function $x \mapsto x^n/\exp(x)$ is defined on all of $\mathbb{R}$, since $\exp$ never vanishes by theorem 4.6.4(1), and $\mathbb{R}$ is unbounded above, so definition 3.1.27 applies. Let $\epsilon > 0$ and put $M = (n+1)!/\epsilon$, a positive real. If $x > M$ then $x > 0$, so the display above holds, and $(n+1)!/x < (n+1)!/M = \epsilon$ by proposition 1.1.2(7). Hence $|x^n/\exp(x) - 0| < \epsilon$ for every $x > M$, which is the assertion, as we sought to show.

Written with $\exp(-x) = 1/\exp(x)$ from (4.11), the conclusion reads $x^n \exp(-x) \rightarrow 0$: no power of x , however large the exponent, keeps pace with $\exp(x)$ as x grows. The proof gives the rate as well, bounding $x^n/\exp(x)$ by $(n+1)!/x$ for every $x > 0$.

The cosine and sine were defined by series and differentiated above, and nothing so far connects their values at two different points. The two identities that do are both consequences of corollary 4.2.8, applied to functions built from $\cos$ and $\sin$ whose derivatives vanish identically. Throughout, $\cos^2 x$ abbreviates $(\cos x)^2$ and $\sin^2 x$ abbreviates $(\sin x)^2$.

Theorem 4.6.12: The Pythagorean Identity

For every $x \in \mathbb{R}$,

$$\cos^2 x + \sin^2 x = 1$$

In particular $|\cos x| \leq 1$ and $|\sin x| \leq 1$ for every $x \in \mathbb{R}$.

Proof:

Put $g(x) = \cos^2 x + \sin^2 x$. Both $\cos$ and $\sin$ are differentiable at every real point by theorem 4.6.2, so the product and sum rules of theorem 4.1.7 make g differentiable at every real point with

$$g'(x) = 2 \cos x \cos'(x) + 2 \sin x \sin'(x) = 2 \cos x (-\sin x) + 2 \sin x \cos x = 0$$

the second equality being theorem 4.6.2. Since $\mathbb{R}$ is an interval and g' vanishes at each of its points, corollary 4.2.8 makes g constant on $\mathbb{R}$.

The constant is $g(0)$. At $x = 0$ the cosine series has only its term with $n = 0$ nonzero, which is 1, and every term of the sine series vanishes, so $\cos 0 = 1$ and $\sin 0 = 0$ by definition 2.6.5. Hence $g(0) = 1 + 0 = 1$ and $g \equiv 1$.

For the bounds, $\cos^2 x = 1 - \sin^2 x \leq 1$ because $\sin^2 x \geq 0$. Multiplicativity of the absolute value, proposition 1.1.2(9), gives $|\cos x|^2 = |\cos^2 x| = \cos^2 x$, the second equality because a square is nonnegative. If $|\cos x| > 1$ held, then proposition 3.3.11 with $n = 2$ applied to $1 < |\cos x|$ would give $1 = 1^2 < |\cos x|^2 = \cos^2 x$, contradicting $\cos^2 x \leq 1$. So $|\cos x| \leq 1$, and the same argument with the roles of $\cos$ and $\sin$ exchanged gives $|\sin x| \leq 1$, as we sought to show.

The identity ties the two values at one point to each other, and it bounds both functions by 1 in absolute value at every point. The addition formulas tie values at different points together, and they are proved by the same device, applied to a function measuring the failure of the formula.

Theorem 4.6.13: The Addition Formulas

For all $x, y \in \mathbb{R}$,

$$\sin(x + y) = \sin x \cos y + \cos x \sin y, \quad \cos(x + y) = \cos x \cos y - \sin x \sin y$$

Proof:

Fix $y \in \mathbb{R}$ and define $h: \mathbb{R} \rightarrow \mathbb{R}$ by

$$h(x) = \sin(x + y) - \sin x \cos y - \cos x \sin y$$

in which $\cos y$ and $\sin y$ are constants.

Step 1: $h'' = -h$. The map $x \mapsto x + y$ is differentiable with derivative 1 by theorem 4.1.7, so theorem 4.1.8 with theorem 4.6.2 makes $x \mapsto \sin(x + y)$ differentiable with derivative $\cos(x + y)$, and $x \mapsto \cos(x + y)$ differentiable with derivative $-\sin(x + y)$. Adding the sum and constant-multiple rules of theorem 4.1.7,

$$h'(x) = \cos(x + y) - \cos x \cos y + \sin x \sin y$$

and differentiating once more by the same rules,

$$h''(x) = -\sin(x + y) + \sin x \cos y + \cos x \sin y = -h(x)$$

Step 2: $h(0) = h'(0) = 0$. At $x = 0$ the only nonzero term of the cosine series is its constant term 1, and every term of the sine series vanishes, so $\cos 0 = 1$ and $\sin 0 = 0$ by definition 2.6.5. Substituting them,

$$h(0) = \sin y - 0 \cdot \cos y - 1 \cdot \sin y = 0, \quad h'(0) = \cos y - 1 \cdot \cos y + 0 \cdot \sin y = 0$$

Step 3: h vanishes identically. Put $E(x) = h(x)^2 + h'(x)^2$. Both h and h' are differentiable at every real point by Step 1, so the product and sum rules of theorem 4.1.7 give

$$E'(x) = 2h(x)h'(x) + 2h'(x)h''(x) = 2h(x)h'(x) - 2h'(x)h(x) = 0$$

using $h'' = -h$. Since $\mathbb{R}$ is an interval, corollary 4.2.8 makes E constant, and Step 2 gives $E(0) = 0 + 0 = 0$. So $h(x)^2 + h'(x)^2 = 0$ for every x . Both summands are nonnegative, so each

is 0, and in particular $h(x)^2 = 0$, whence $h(x) = 0$. That is the first formula.

Step 4: the cosine formula. By Step 3 the function h is the constant function 0, whose derivative at every point is 0 directly from definition 4.1.2. So $h'(x) = 0$ for every x , and the expression for h' in Step 1 gives $\cos(x+y) = \cos x \cos y - \sin x \sin y$. Since $y \in \mathbb{R}$ was arbitrary and x ranged over $\mathbb{R}$ throughout, both formulas hold for all $x, y \in \mathbb{R}$, as we sought to show.

The formulas express the values at $x+y$ through the values at x and at y , and together with a point at which the cosine is negative they give periodicity. The alternating series estimate of corollary 2.5.3 produces such a point, and two further facts about the interval $[0, 2]$ come with it.

First, $\sin x > 0$ for $0 < x \leq 2$. Fix such an x and put $b_n = x^{2n-1}/(2n-1)!$ for $n \geq 1$, so that $\sin x = \sum_{n \geq 1} (-1)^{n+1} b_n$ is definition 2.6.5's sine series after the relabelling that replaces the index n by $n-1$. Each b_n is positive, and

$$\frac{b_{n+1}}{b_n} = \frac{x^2}{2n(2n+1)} \leq \frac{4}{6} = \frac{2}{3}$$

because $0 < x \leq 2$ gives $x^2 \leq 4$ by proposition 3.3.11, with equality when $x = 2$, and $2n(2n+1) \geq 6$ for $n \geq 1$. So $b_{n+1} \leq \frac{2}{3}b_n$, which makes (b_n) nonincreasing, and induction on n gives $b_n \leq (2/3)^{n-1}b_1 = \frac{3}{2}b_1(2/3)^n$. Here $(2/3)^n \rightarrow 0$ by proposition 2.1.10 with the exponent $j = 0$, so the right-hand side tends to 0 by the scalar rule of theorem 2.1.6, and $0 \leq b_n \leq \frac{3}{2}b_1(2/3)^n$ with theorem 2.1.7 gives $b_n \rightarrow 0$. The hypotheses of corollary 2.5.3 therefore hold, and its part (2) at $k = 1$ gives $s_2 \leq \sin x \leq s_1$, that is

$$x - \frac{x^3}{6} \leq \sin x \leq x$$

Since $0 < x \leq 2$ gives $x^2 \leq 4 < 6$, the left-hand side equals $x(6 - x^2)/6 > 0$, so $\sin x > 0$.

Second, $\cos 2 \leq -1/3$. For $N \geq 2$, splitting off the first two terms of the cosine series at $x = 2$ gives

$$\sum_{n=0}^N \frac{(-1)^n 2^{2n}}{(2n)!} = 1 - 2 + \sum_{n=2}^N \frac{(-1)^n 2^{2n}}{(2n)!}$$

The left-hand side tends to $\cos 2$ by definition 2.6.5, so the partial sums on the right tend to $\cos 2 + 1$ by theorem 2.1.6. Writing $T = \sum_{n \geq 2} (-1)^n 2^{2n}/(2n)!$ for that limit, $\cos 2 = -1 + T$. Put $c_k = 2^{2k+2}/(2k+2)!$ for $k \geq 1$, so that $T = \sum_{k \geq 1} (-1)^{k+1} c_k$ after the relabelling $n = k+1$, which turns $(-1)^n$ into $(-1)^{k+1}$. Each c_k is positive and

$$\frac{c_{k+1}}{c_k} = \frac{4}{(2k+3)(2k+4)} \leq \frac{4}{30}$$

for $k \geq 1$, so (c_k) is nonincreasing, and $c_k \leq (4/30)^{k-1}c_1 \rightarrow 0$ by the same three steps used for (b_n) : induction on k , then proposition 2.1.10 with the scalar rule of theorem 2.1.6, then theorem 2.1.7. Corollary 2.5.3(2) at $k = 1$ now gives $T \leq s_1 = c_1 = 2^4/4! = 2/3$, whence $\cos 2 \leq -1 + 2/3 = -1/3$.

Third, $\cos$ is strictly decreasing on $(0, 2)$. Every $x \in (0, 2)$ satisfies $0 < x \leq 2$, so $\sin x > 0$ there and $\cos'(x) = -\sin x < 0$ at every $x \in (0, 2)$ by theorem 4.6.2. Corollary 4.2.7 applied to the interval $(0, 2)$ gives the claim.

The cosine is continuous on $[0, 2]$, by example 4.6.3 and proposition 3.3.1, with $\cos 0 = 1$ and $\cos 2 \leq -1/3 < 0$. So $\min\{\cos 0, \cos 2\} < 0 < \max\{\cos 0, \cos 2\}$, and theorem 3.3.2(1) produces a $c \in (0, 2)$ with $\cos c = 0$. Strict decrease makes $\cos$ injective on $(0, 2)$, so c is the only zero there, and it gives $\cos x > \cos c = 0$ for $0 < x < c$, while $\cos 0 = 1 > 0$.

Definition 4.6.14: The Number π

The cosine has exactly one zero in the interval $(0, 2)$. Writing c for that zero, so that $\cos c = 0$ and $\cos x > 0$ for $0 \leq x < c$, define

$$\pi = 2c$$

So $0 < \pi < 4$, the value $\cos(\pi/2)$ is 0, and $\cos x > 0$ for $0 \leq x < \pi/2$.

The companion value is $\sin(\pi/2) = 1$. Indeed theorem 4.6.12 at $x = \pi/2$ gives $\sin^2(\pi/2) = 1 - 0 = 1$, and $\pi/2$ lies in $(0, 2)$ by definition 4.6.14, so $\sin(\pi/2) > 0$ because $\sin x > 0$ for $0 < x \leq 2$. Hence $\sin(\pi/2)$ is the positive number whose square is 1, which is 1. The estimate $\sin x \leq x$ for $0 < x \leq 2$, from the same alternating series bound, gives $1 \leq \pi/2$ when read at $x = \pi/2$, so

$$2 \leq \pi < 4 \quad (4.13)$$

Theorem 4.6.15: Periodicity of the Cosine and the Sine

Let $x \in \mathbb{R}$ and $k \in \mathbb{Z}$.

1. $\sin(x + \pi/2) = \cos x$ and $\cos(x + \pi/2) = -\sin x$.
2. $\sin(x + \pi) = -\sin x$ and $\cos(x + \pi) = -\cos x$, and more generally $\sin(x + k\pi) = (-1)^k \sin x$ and $\cos(x + k\pi) = (-1)^k \cos x$. Taking $k = 2$ shows that $\cos$ and $\sin$ are periodic with period 2π .
3. $\cos x = 0$ exactly when $x = \pi/2 + k\pi$ for some $k \in \mathbb{Z}$, and $\sin x = 0$ exactly when $x = k\pi$ for some $k \in \mathbb{Z}$.
4. $\cos x = 1$ exactly when $x = 2k\pi$ for some $k \in \mathbb{Z}$. If $p > 0$ satisfies $\cos(x + p) = \cos x$ for every $x \in \mathbb{R}$, or satisfies $\sin(x + p) = \sin x$ for every $x \in \mathbb{R}$, then $p \geq 2\pi$.

Proof:

1. Theorem 4.6.13 with $y = \pi/2$, together with $\cos(\pi/2) = 0$ from definition 4.6.14 and $\sin(\pi/2) = 1$ from the paragraph above, gives

$$\sin(x + \pi/2) = \sin x \cdot 0 + \cos x \cdot 1 = \cos x, \quad \cos(x + \pi/2) = \cos x \cdot 0 - \sin x \cdot 1 = -\sin x$$

2. Applying part (1) twice, $\sin(x + \pi) = \sin((x + \pi/2) + \pi/2) = \cos(x + \pi/2) = -\sin x$ and $\cos(x + \pi) = -\sin(x + \pi/2) = -\cos x$. For $k \geq 0$ the two general identities follow by induction on k : the case $k = 0$ is trivial, and the step is the case $k = 1$ just proved, applied at the point $x + k\pi$. For $k < 0$ the integer $-k$ is positive, so the case already proved, applied at the point $x + k\pi$ with $-k$ in place of k , gives $\sin x = \sin((x + k\pi) + (-k)\pi) = (-1)^{-k} \sin(x + k\pi)$. Now $(-1)^{-k} = (-1)^k$, since k and $-k$ have the same parity, and multiplying both sides by $(-1)^k$ gives $(-1)^k \sin x = \sin(x + k\pi)$. The same induction gives the two cosine identities.
3. By definition 4.6.14, $\cos > 0$ on $[0, \pi/2)$ and $\cos(\pi/2) = 0$. For $x \in (\pi/2, \pi]$ write $x = t + \pi/2$ with $t \in (0, \pi/2]$. Then $0 < t < 2$, since $\pi/2 < 2$, so $\sin t > 0$ because the sine is positive on $(0, 2]$, and part (1) gives $\cos x = -\sin t < 0$. Hence $\pi/2$ is the only zero of

$\cos$ in $[0, \pi)$.

Let $x \in \mathbb{R}$. Proposition 1.4.3 applied to x/π gives a unique $k \in \mathbb{Z}$ with $k \leq x/\pi < k+1$, and multiplying by $\pi > 0$ puts $x - k\pi$ in $[0, \pi)$. Part (2) gives $\cos x = (-1)^k \cos(x - k\pi)$, so $\cos x = 0$ holds exactly when $\cos(x - k\pi) = 0$, which by the previous paragraph holds exactly when $x - k\pi = \pi/2$. That is the first assertion. For the second, part (1) gives $\sin x = -\cos(x + \pi/2)$, so $\sin x = 0$ holds exactly when $\cos(x + \pi/2) = 0$, which by the first assertion holds exactly when $x + \pi/2 = \pi/2 + k\pi$, that is $x = k\pi$.

4. Suppose $\cos x = 1$. Then theorem 4.6.12 gives $\sin^2 x = 1 - 1 = 0$, so $\sin x = 0$ and $x = k\pi$ for some $k \in \mathbb{Z}$ by part (3). Part (2) then gives $\cos x = (-1)^k \cos 0 = (-1)^k$, and $(-1)^k = 1$ forces k to be even, say $k = 2m$, so $x = 2m\pi$. Conversely $\cos(2m\pi) = (-1)^{2m} \cos 0 = 1$. Let $p > 0$ satisfy $\cos(x + p) = \cos x$ for every x . Taking $x = 0$ gives $\cos p = \cos 0 = 1$, so $p = 2m\pi$ for some integer m by the previous paragraph, and $p > 0$ with $\pi > 0$ forces $m \geq 1$, whence $p \geq 2\pi$. Let $p > 0$ satisfy $\sin(x + p) = \sin x$ for every x . Taking $x = \pi/2$ gives $\sin(\pi/2 + p) = \sin(\pi/2) = 1$. If $\sin z = 1$ then theorem 4.6.12 gives $\cos^2 z = 0$, so $\cos z = 0$ and $z = \pi/2 + k\pi$ for some $k \in \mathbb{Z}$ by part (3), and part (2) gives $\sin z = (-1)^k \sin(\pi/2) = (-1)^k$, so k is even. Applying this to $z = \pi/2 + p$ gives $\pi/2 + p = \pi/2 + 2m\pi$ for an integer m , so $p = 2m\pi \geq 2\pi$ as before, completing the proof.

The number π arrives here as a zero of a power series, and no length, angle or circle enters its construction. Part (4) is what makes 2π the smallest period: no positive number below it is a period of either function. Part (3) locates every zero of both functions, and example 4.6.16 and example 4.6.17 use those zeros to produce oscillation on every interval around the origin.

Example 4.6.16: The Function $\sin(1/x)$ at the Origin

Define $f: [-1, 1] \rightarrow \mathbb{R}$ by $f(x) = \sin(1/x)$ for $x \neq 0$ and $f(0) = 0$. Then f is continuous at every $x \neq 0$, its restriction to $(-1, 1)$ has a discontinuity of the second kind at 0 in the sense of definition 3.2.13, and f has the intermediate value property of definition 3.3.4 on $[-1, 1]$.

Continuity away from 0. On $[-1, 1] \setminus \{0\}$ the function agrees with $x \mapsto \sin(1/x)$, which example 4.6.3 makes continuous at every point of $\mathbb{R} \setminus \{0\}$. For $x \neq 0$ the set $[-1, 1] \setminus \{0\}$ contains every point of $[-1, 1]$ within $|x|$ of x , so proposition 3.3.1 and proposition 3.1.22(1) together give continuity of f at x as a function on $[-1, 1]$.

Two sequences of oscillation points. For $k \in \mathbb{N}$ put

$$a_k = \frac{1}{\pi/2 + 2k\pi}, \quad b_k = \frac{1}{3\pi/2 + 2k\pi}$$

Both denominators are positive and the second exceeds the first, so $0 < b_k < a_k$ by proposition 1.1.2(7). By theorem 4.6.15(2), whose integer parameter is taken to be $2k$ here,

$$\sin\left(\frac{1}{a_k}\right) = \sin(\pi/2 + 2k\pi) = \sin(\pi/2) = 1, \quad \sin\left(\frac{1}{b_k}\right) = \sin(3\pi/2 + 2k\pi) = \sin(3\pi/2)$$

and $\sin(3\pi/2) = \sin(\pi/2 + \pi) = -\sin(\pi/2) = -1$ by theorem 4.6.15(2) with the parameter 1. So $f(a_k) = 1$ and $f(b_k) = -1$. Both sequences tend to 0: given $\epsilon > 0$, theorem 1.4.1(1) with $x = 2\pi > 0$ and $y = 1/\epsilon$ gives an $N \in \mathbb{N}$ with $2N\pi > 1/\epsilon$, and for $k \geq N$ both denominators exceed $2N\pi$, so $0 < b_k < a_k < \epsilon$ by proposition 1.1.2(7). Every one of these points lies in $(0, 1)$, since (4.13) gives $\pi/2 + 2k\pi \geq 1 + 4k > 1$ and hence $0 < b_k < a_k < 1$. Finally (4.10) gives $f(-a_k) = -1$ and $f(-b_k) = 1$, and those points lie in $(-1, 0)$.

The discontinuity at 0 is of the second kind. Suppose the right-hand limit $\lim_{x \rightarrow 0^+} f(x)$ existed and equalled $L \in \mathbb{R}$. By proposition 3.1.25 that is the limit at 0 of the restriction of f to $(0, 1]$. The point 0 is a limit point of $(0, 1]$ by definition 3.1.5, since every $\delta > 0$ admits an $n \in \mathbb{N}$ with $0 < 1/n < \min\{\delta, 1\}$ by theorem 1.4.1(3). So theorem 3.1.19 applied to the sequences (a_k) and (b_k) , both in $(0, 1)$ and both tending to 0, gives $f(a_k) \rightarrow L$ and $f(b_k) \rightarrow L$. The two sequences of values are constantly 1 and constantly -1 , so $L = 1$ and $L = -1$ by proposition 2.1.4, which is impossible. The same argument with $(-a_k)$ and $(-b_k)$, which lie in $(-1, 0)$ and tend to 0, rules out the left-hand limit. Neither one-sided limit exists in $\mathbb{R}$, so definition 3.2.13 classifies 0 as a discontinuity of the second kind.

The intermediate value property. Let $u < v$ in $[-1, 1]$ and let y lie strictly between $f(u)$ and $f(v)$. Every value of f lies in $[-1, 1]$, by theorem 4.6.12 and $f(0) = 0$, so $-1 < y < 1$.

If $0 \notin [u, v]$, then f is continuous on $[u, v]$ by the first paragraph and proposition 3.3.1, and theorem 3.3.2(1) gives a $c \in (u, v)$ with $f(c) = y$.

If $0 \in [u, v]$, then $u \leq 0 \leq v$. Suppose first that $v > 0$. Since $a_k \rightarrow 0$, there is a k with $a_k < v$, and for that k , $u \leq 0 < b_k < a_k < v$, so $[b_k, a_k] \subseteq (u, v)$, and f is continuous on $[b_k, a_k]$ because 0 is not in it. Since $\min\{f(b_k), f(a_k)\} = -1 < y < 1 = \max\{f(b_k), f(a_k)\}$, theorem 3.3.2(1) gives a $c \in (b_k, a_k) \subseteq (u, v)$ with $f(c) = y$. If instead $v = 0$, then $u < 0$, and a k with $a_k < -u$ gives $u < -a_k < -b_k < 0 = v$, so $[-a_k, -b_k] \subseteq (u, v)$, on which f is continuous with $f(-a_k) = -1$ and $f(-b_k) = 1$, and theorem 3.3.2(1) again produces the required c .

This function settles the question left open at definition 3.3.4. It takes every intermediate value between any two of its values and is discontinuous at 0, so the intermediate value property does not imply continuity and theorem 3.3.2 has no converse. In the next example the function with the intermediate value property is itself a derivative, which is the case theorem 4.2.10 covers.

Example 4.6.17: A Differentiable Function Whose Derivative Is Discontinuous

Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) = x^2 \sin(1/x)$ for $x \neq 0$ and $g(0) = 0$. Then g is differentiable at every point of $\mathbb{R}$, and g' is not continuous at 0.

Differentiability away from 0. On the interval $(0, \infty)$ the map $x \mapsto 1/x$ is differentiable with derivative $-1/x^2$ by theorem 4.1.7, so theorem 4.1.8 with theorem 4.6.2 makes $x \mapsto \sin(1/x)$ differentiable with derivative $-\cos(1/x)/x^2$, and the product rule of theorem 4.1.7 gives

$$g'(x) = 2x \sin(1/x) + x^2 \cdot \left(-\frac{\cos(1/x)}{x^2}\right) = 2x \sin(1/x) - \cos(1/x)$$

The same computation on the interval $(-\infty, 0)$ gives the same formula, so it holds at every $x \neq 0$. *Differentiability at 0.* For $x \neq 0$ the difference quotient is

$$\frac{g(x) - g(0)}{x - 0} = \frac{x^2 \sin(1/x)}{x} = x \sin(1/x)$$

and $|\sin(1/x)| \leq 1$ by theorem 4.6.12, so multiplicativity of the absolute value, proposition 1.1.2(9), gives $|x \sin(1/x)| = |x| |\sin(1/x)| \leq |x|$, that is $-|x| \leq x \sin(1/x) \leq |x|$. Both bounds tend to 0 as $x \rightarrow 0$, since $|\pm|x| - 0| = |x| < \epsilon$ whenever $0 < |x - 0| < \epsilon$, which is definition 3.1.15 with $\delta = \epsilon$. So corollary 3.1.21 gives $\lim_{x \rightarrow 0} x \sin(1/x) = 0$, and by definition 4.1.2 the function g is differentiable at 0 with $g'(0) = 0$.

The derivative is discontinuous at 0. Put $x_k = 1/(2k\pi)$ for $k \in \mathbb{N}$. Each x_k is nonzero, and $x_k \rightarrow 0$: given $\epsilon > 0$, theorem 1.4.1(1) with $x = 2\pi > 0$ and $y = 1/\epsilon$ gives an $N \in \mathbb{N}$ with $2N\pi > 1/\epsilon$, so $0 < x_k < \epsilon$ for $k \geq N$ by proposition 1.1.2(7). Theorem 4.6.15(3) gives $\sin(2k\pi) = 0$, since $2k\pi$ is

an integer multiple of π , and theorem 4.6.15(4) gives $\cos(2k\pi) = 1$. So

$$g'(x_k) = 2x_k \cdot 0 - 1 = -1$$

for every k . If g' were continuous at 0, then proposition 3.2.4 would give $g'(x_k) \rightarrow g'(0) = 0$, while the sequence $(g'(x_k))$ is constantly -1 and therefore converges to -1 . Two different limits for one sequence contradict proposition 2.1.4, so g' is not continuous at 0.

The derivative g' is therefore discontinuous at 0, while theorem 4.2.10 says that it still has the intermediate value property. That is the example section 4.2 deferred to this section, and it measures exactly how far theorem 4.2.10 improves on continuity: a derivative need not be continuous, and the intermediate value property is the weaker conclusion that remains available.

Exercise 4.6.1

1. Show that $\log x \leq x - 1$ for every $x > 0$, with equality exactly when $x = 1$.
2. Show that $\log x < 2\sqrt{x}$ for every $x > 1$, and deduce that

$$\lim_{x \rightarrow \infty} \frac{\log x}{x} = 0$$

(For the first part, apply the inequality of the previous question at $\sqrt{x}$ and use theorem 4.6.9(2).)

3. Compute $\lim_{x \rightarrow 0} \frac{\sin x}{x}$ directly from the definition of the derivative, and then compute $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$.
4. Show that $\sin x < x$ for every $x > 0$. (Treat $x \geq 2\pi$ separately from $0 < x < 2\pi$, and on the second range use theorem 4.6.15(4) to see that $1 - \cos x > 0$.)
5. Show that $\sin x > 0$ for every $x \in (0, \pi)$, that $\cos$ is strictly decreasing on $[0, \pi]$, and that $\cos \pi = -1$.
6. Define $\cosh x = (\exp(x) + \exp(-x))/2$ and $\sinh x = (\exp(x) - \exp(-x))/2$. Prove that $\cosh^2 x - \sinh^2 x = 1$, that $\sinh' = \cosh$ and $\cosh' = \sinh$, and that $\sinh: \mathbb{R} \rightarrow \mathbb{R}$ is a strictly increasing bijection.
7. Define $F: \mathbb{R} \rightarrow \mathbb{R}$ by $F(x) = \exp(-1/x^2)$ for $x \neq 0$ and $F(0) = 0$. Prove that F is differentiable at 0 with $F'(0) = 0$. (Bound $\exp(1/x^2)$ below by a single term of its series.)
8. Let $a > 0$ with $a \neq 1$. Show that $x \mapsto a^x$ is strictly increasing on $\mathbb{R}$ when $a > 1$ and strictly decreasing when $a < 1$, and that $\lim_{x \rightarrow \infty} a^x = +\infty$ when $a > 1$.

4.7 Analyticity and the Growth of Derivatives

Differentiability is a local, one-step condition, and iterating it asks nothing about how the successive derivatives behave. A power series expansion asks for much more: that the whole infinite list $f(a), f'(a), f''(a), \dots$ determine f near a . This section measures the gap between the two, and finds that a single growth condition on the derivatives closes it.

Call a function *smooth* on an interval if it has derivatives of every order there. Let f be smooth on an open interval. At a point a one may form the Taylor series

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (t-a)^k$$

and ask whether it agrees with f on some interval around a .

Definition 4.7.1: [Real] Analytic Function

Let $U \subseteq \mathbb{R}$ be open. A function $f: U \rightarrow \mathbb{R}$ is *analytic* if for each $a \in U$ there is an $R > 0$ and real coefficients $c_0, c_1, c_2, \dots$ such that

$$f(a+h) = \sum_{k=0}^{\infty} c_k h^k \quad \text{for all } |h| < R$$

the series converging for every such h .

Every analytic function is smooth by theorem 4.5.3, which differentiates a power series term by term at every point strictly inside its interval of convergence, and corollary 4.5.4 then forces $c_k = f^{(k)}(a)/k!$: the only candidate expansion is the Taylor series. The converse fails, and the standard counterexample is

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

Every derivative of this function at the origin vanishes. For $x \neq 0$ the product and chain rules give, by induction on k , a polynomial p_k with $f^{(k)}(x) = p_k(1/x)e^{-1/x^2}$: differentiating $p_k(1/x)e^{-1/x^2}$ produces $(2x^{-3}p_k(1/x) - x^{-2}p'_k(1/x))e^{-1/x^2}$, which has that shape again. Substituting $u = 1/x^2$, which tends to ∞ as $x \rightarrow 0$, turns proposition 4.6.11 into the statement that $p(1/x)e^{-1/x^2} \rightarrow 0$ as $x \rightarrow 0$ for every polynomial p . The values at the origin now follow by induction: $f(0) = 0$, and if $f^{(k)}(0) = 0$ then the difference quotient there is $f^{(k)}(h)/h = (1/h)p_k(1/h)e^{-1/h^2}$, an expression of that same shape, so it tends to 0 and $f^{(k+1)}(0) = 0$. The Taylor series at the origin is therefore

$$T_0 f = \sum_{k=0}^{\infty} 0 \cdot x^k = 0$$

which agrees with f at the single point $x = 0$ and nowhere else. The Taylor series exists, converges everywhere, and converges to the wrong function.

What goes wrong is visible in the remainder. Truncating the series after k terms splits f into a polynomial and an error,

$$f(t) = \underbrace{\sum_{j=0}^{k-1} \frac{f^{(j)}(a)}{j!} (t-a)^j}_{P_{k-1}(t)} + \underbrace{R_{k-1}(t)}_{\text{remainder}}$$

and the series converges to f at t precisely when $R_{k-1}(t) \rightarrow 0$. So the question is how large the remainder can be, and the following classical estimate answers it in terms of a single derivative.

Theorem 4.3.3 answers it: with I an open interval containing a and f smooth on it, the hypotheses of that theorem hold for every k , so for each $t \in I$ with $t \neq a$ there is a ξ_t strictly between a and t with

$$R_{k-1}(t) = \frac{f^{(k)}(\xi_t)}{k!}(t-a)^k$$

The remainder is therefore controlled by the size of the k -th derivative on the interval, and nothing else. Fix a radius $r > 0$, write $I = [a-r, a+r]$, and let f be smooth on an open interval containing I . Each $f^{(k)}$ is then differentiable there, hence continuous by proposition 4.1.5, so theorem 3.3.5 bounds it on the closed bounded interval I and

$$M_k = \sup_{t \in I} |f^{(k)}(t)|$$

is finite. For $t \in I$ the Lagrange form bounds the remainder by

$$|R_{k-1}(t)| = \frac{|f^{(k)}(\xi_t)|}{k!} |t-a|^k \leq \frac{M_k}{k!} r^k$$

This is the worst the error can be, and whether it tends to zero is a question about the sequence $M_k/k!$ alone. Theorem 2.4.15 compares that sequence's decay against a geometric one through the threshold quantity

$$\limsup_{k \rightarrow \infty} \sqrt[k]{\frac{M_k}{k!} r^k} = r \limsup_{k \rightarrow \infty} \sqrt[k]{\frac{M_k}{k!}}$$

which factors into the radius and an intrinsic property of f .

Definition 4.7.2: Derivative Growth Rate

Let f be smooth on an open interval containing $I = [a-r, a+r]$, and set $M_k = \sup_{t \in I} |f^{(k)}(t)|$. The *derivative growth rate* of f on I is

$$\alpha_r = \limsup_{k \rightarrow \infty} \sqrt[k]{\frac{M_k}{k!}}$$

and f has *bounded derivative growth rate* on I if $\alpha_r < \infty$. It has *locally bounded derivative growth rate* on an open set U if for each $x \in U$ there is a $\rho > 0$ with $[x-\rho, x+\rho] \subseteq U$ such that f is smooth on an open interval containing $[x-\rho, x+\rho]$ and its derivative growth rate α_ρ on that interval is finite.

For $n \geq 1$ every term of the series defining $\exp$ (definition 2.6.5) is positive at $x = n$, so $e^n = \exp(n) \geq n^n/n!$ and

$$\sqrt[n]{n!} \geq \frac{n}{e} \tag{4.14}$$

which sends $\sqrt[n]{n!} \rightarrow \infty$.

Example 4.7.3: Derivative Growth Rate of the Exponential

For $f = \exp$ on $I = [-r, r]$ every derivative is $\exp$ itself by theorem 4.6.1, and $\exp$ is increasing by

theorem 4.6.4(2), so $M_k = \sup_{t \in I} e^t = e^r$ for all k and

$$\alpha_r = \limsup_{k \rightarrow \infty} \sqrt[k]{\frac{e^r}{k!}} = \lim_{k \rightarrow \infty} \frac{(e^r)^{1/k}}{\sqrt[k]{k!}} = 0$$

since $(e^r)^{1/k} \rightarrow 1$ while $\sqrt[k]{k!} \rightarrow \infty$ by (4.14). The growth rate is 0 at every radius, so $\alpha_r r = 0 < 1$ however large r is, and the Taylor series of $\exp$ converges to it on every interval. The same computation with $M_k = 1$ handles $\sin$ and $\cos$: each derivative of either one is $\pm \sin$ or $\pm \cos$ by theorem 4.6.2, and theorem 4.6.12 gives $|\sin t| \leq 1$ and $|\cos t| \leq 1$ at every t .

The threshold quantity is $r\alpha_r$, and a bound on it converts directly into convergence.

Theorem 4.7.4: Taylor Convergence Under Bounded Growth

Let f be smooth on an open interval containing $I = [a - \sigma, a + \sigma]$, with derivative growth rate α on I . If $\alpha\sigma < 1$ then the partial sums of the Taylor series of f at a converge to f uniformly on I , in the sense of definition 4.4.2.

Proof:

Choose $\delta > 0$ small enough that $(\alpha + \delta)\sigma < 1$. By theorem 2.2.12(1) there is an N with

$$\sqrt[n]{\frac{M_n}{n!}} \leq \alpha + \delta \quad \text{for all } n \geq N$$

Let $0 < |h| \leq \sigma$, the case $h = 0$ being trivial because both sides of the estimate below are then 0. Applying theorem 4.3.3 at $t = a + h$, whose hypothesis $t \neq a$ is what this restriction discharges, gives a θ between a and $a + h$ with

$$\left| f(a + h) - \sum_{k=0}^{n-1} \frac{f^{(k)}(a)}{k!} h^k \right| = \frac{|f^{(n)}(\theta)|}{n!} |h|^n \leq \frac{M_n}{n!} \sigma^n = \left(\sqrt[n]{\frac{M_n}{n!}} \sigma \right)^n$$

so for $n \geq N$ this is at most $((\alpha + \delta)\sigma)^n$. The bound does not depend on h , and $(\alpha + \delta)\sigma < 1$ makes it tend to 0, completing the proof.

The converse direction starts from a function presented as a convergent power series and bounds its derivatives. Differentiating such a series n times multiplies its coefficients by factors of size roughly k^n , and the geometric decay of those coefficients absorbs the factors.

Theorem 4.7.5: Bounded Growth Rate of an Analytic Function

Suppose $f(a + h) = \sum_{k \geq 0} c_k h^k$ for all $|h| < R$, and let $0 < \sigma < R$. Then f has bounded derivative growth rate on $I = [a - \sigma, a + \sigma]$.

Proof:

Step 1: a binomial estimate. Fix λ with $0 < \lambda < 1$. The estimate is

$$\limsup_{n \rightarrow \infty} \sqrt[n]{\sum_{k=n}^{\infty} \binom{k}{n} \lambda^k} < \infty \quad (4.15)$$

The geometric series $\sum_{k \geq 0} x^k = 1/(1-x)$ has radius of convergence 1, so theorem 4.5.3, applied n times at each point strictly inside $(-1, 1)$, differentiates it term by term and gives

$$\sum_{k=n}^{\infty} k(k-1) \cdots (k-n+1) x^{k-n} = \frac{n!}{(1-x)^{n+1}} \quad \text{for } |x| < 1$$

Since $k(k-1) \cdots (k-n+1) = n! \binom{k}{n}$, dividing by $n!$, setting $x = \lambda$ and multiplying by λ^n evaluates the sum in (4.15) exactly:

$$\sum_{k=n}^{\infty} \binom{k}{n} \lambda^k = \frac{\lambda^n}{(1-\lambda)^{n+1}}$$

Its n -th root is $\lambda(1-\lambda)^{-1-1/n}$, and $(1-\lambda)^{-1/n} \rightarrow 1$ because $1-\lambda$ is a positive constant, so those roots converge to $\lambda/(1-\lambda)$. That limit is finite, proving (4.15).

Step 2: bounding the derivatives. The series converges at every h with $|h| < R$, so its radius of convergence is at least R , which gives $\limsup_k \sqrt[k]{|c_k|} \leq 1/R$. Since $\sigma < R$ there is a λ with $\sigma/R < \lambda < 1$, and then

$$\limsup_{k \rightarrow \infty} \sqrt[k]{|c_k| \sigma^k} = \sigma \limsup_{k \rightarrow \infty} \sqrt[k]{|c_k|} \leq \frac{\sigma}{R} < \lambda$$

so there is a K with $|c_k| \sigma^k \leq \lambda^k$ for all $k \geq K$. By theorem 4.5.3, applied n times at each point strictly inside $(-R, R)$ and evaluated at $|h| \leq \sigma < R$, differentiating the series term by term gives, for $n \geq K$,

$$\begin{aligned} |f^{(n)}(a+h)| &\leq \sum_{k=n}^{\infty} k(k-1) \cdots (k-n+1) |c_k| |h|^{k-n} \\ &\leq \frac{n!}{\sigma^n} \sum_{k=n}^{\infty} \binom{k}{n} |c_k| \sigma^k \leq \frac{n!}{\sigma^n} \sum_{k=n}^{\infty} \binom{k}{n} \lambda^k \end{aligned}$$

using $k(k-1) \cdots (k-n+1) = n! \binom{k}{n}$ and $|h|^{k-n} \leq \sigma^k / \sigma^n$. The last inequality calls on $|c_k| \sigma^k \leq \lambda^k$ only for $k \geq n$, which $n \geq K$ guarantees. Taking the supremum over $|h| \leq \sigma$ and then n -th roots, and discarding the finitely many $n < K$, which leave the lim sup unchanged,

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{\frac{M_n}{n!}} \leq \frac{1}{\sigma} \limsup_{n \rightarrow \infty} \sqrt[n]{\sum_{k=n}^{\infty} \binom{k}{n} \lambda^k}$$

which is finite by (4.15), as we sought to show.

The two directions now fit together, and the result is a characterization of analyticity that never mentions series at all.

Theorem 4.7.6: Analyticity Theorem

A smooth function on an open interval is analytic if and only if it has locally bounded derivative growth rate.

Proof:

Suppose first that $f: (a, b) \rightarrow \mathbb{R}$ is smooth with locally bounded derivative growth rate, and let $x \in (a, b)$. Then x lies in an interval on which the growth rate α is finite, and shrinking that interval can only shrink the suprema M_k , hence cannot increase α . So choose $\sigma > 0$ small enough that $I = [x - \sigma, x + \sigma]$ sits inside it and $\alpha\sigma < 1$. By theorem 4.7.4 the Taylor series of f at x converges to f on I , so f is analytic at x .

Conversely, suppose f is analytic and let $x \in (a, b)$. Some power series $\sum c_k h^k$ converges to $f(x + h)$ for all h in an interval $(-R, R)$ with $R > 0$. Choosing any σ with $0 < \sigma < R$, theorem 4.7.5 shows f has bounded derivative growth rate on $[x - \sigma, x + \sigma]$, completing the proof.

The characterization makes the difference between the two classes quantitative. An analytic function satisfies a bound of the form

$$|f^{(k)}(x)| \leq C \cdot \frac{k!}{R^k}$$

on a neighbourhood of each of its points: a finite growth rate α_r bounds $\sqrt[k]{M_k/k!}$ by some A for all large k , which is the displayed bound with $R = 1/A$, the constant C absorbing the finitely many remaining k . Smoothness imposes no such bound. The function e^{-1/x^2} is smooth on all of $\mathbb{R}$ and its Taylor series at the origin is the zero series, so it is not analytic there, and theorem 4.7.6 therefore denies it locally bounded derivative growth rate on every open interval about the origin.

The Riemann Integral

The integral of a positive function over an interval ought to be the area under its graph, and the difficulty is that area is exactly what has not been defined. What is available is the area of a rectangle. So the construction approximates the region from inside and outside by finite unions of rectangles, takes the best inner approximation and the best outer one, and calls the function integrable when the two agree.

The two approximations are a supremum and an infimum of sets of real numbers, so the definition rests on theorem 1.2.4, as every construction in this book has. What is new is that the two numbers it produces need not be equal. Both numbers come from one function, and the gap between them is a property of that function. The chapter is the study of when that gap closes.

Chapter 3 and Chapter 4 give the answer for the functions that matter most. A continuous function on a closed bounded interval is uniformly continuous by theorem 3.4.4, and uniform continuity is what lets one partition be chosen fine enough to work at every point of the interval at once. That is the whole proof that continuous functions are integrable, and it is the reason Section 3.4 was written the way it was.

The chapter closes the loop between the two halves of the book. Differentiation and integration are inverse to each other, which is the fundamental theorem, and once that is available the integral becomes a way of deciding questions about series: theorem 5.3.9 settles the convergence of $\sum 1/(n(\log n)^p)$, which none of the tests of Section 2.4 can reach.

Section 5.1 builds the integral from partitions and Darboux sums, proves the criterion that decides integrability, and shows that the definition agrees with the Riemann sums of elementary calculus. Section 5.2 identifies the functions the criterion admits, which include the continuous and the monotone ones, and exhibits a bounded function it excludes. Section 5.3 proves the fundamental theorem in both of its forms, derives integration by parts and substitution, and extends the integral to unbounded intervals. Section 5.4★ replaces the increment Δx by the increment of a fixed increasing function, which turns sums and integrals into two cases of one construction. It is enrichment, and

nothing later depends on it.

5.1 The Riemann Integral

A subdivision of $[a, b]$ into finitely many pieces turns the two-sided approximation into arithmetic. On each piece a bounded f has a supremum and an infimum, and multiplying each by the length of its piece produces one sum lying above the region under the graph and one lying below. Every statement in this section describes how those two sums respond when the subdivision changes: they move toward each other when the pieces are cut smaller, no sum built from below ever exceeds a sum built from above, and the size of the gap between them is what decides whether the two approximations name a single number.

The number produced this way is Darboux's. Riemann's own definition samples f at an arbitrary point of each piece and forms a single sum, with no suprema anywhere in it, and theorem 5.1.11 identifies the two. The section closes with the arithmetic every later computation uses: the integral is linear in f , it respects the order on functions, and it adds across a point of $[a, b]$.

Throughout the chapter a and b are real numbers with $a < b$, and $[a, b]$ is the closed bounded interval of definition 3.1.1. Every function integrated is bounded, and boundedness appears as a hypothesis of every statement below.

Definition 5.1.1: [Real] Partition

A *partition* of $[a, b]$ is a finite set $P = \{x_0, x_1, \dots, x_n\}$ of points of $[a, b]$ with

$$a = x_0 < x_1 < \dots < x_n = b$$

Its *subintervals* are the n closed bounded intervals $[x_{i-1}, x_i]$ for $1 \leq i \leq n$, and their lengths are written

$$\Delta x_i = x_i - x_{i-1} > 0$$

The *mesh* of P is $\|P\| = \max_{1 \leq i \leq n} \Delta x_i$.

Two facts about a partition are used constantly. The lengths of its subintervals add to the length of the whole interval,

$$\sum_{i=1}^n \Delta x_i = \sum_{i=1}^n (x_i - x_{i-1}) = x_n - x_0 = b - a \quad (5.1)$$

since the sum telescopes, and every subinterval is nonempty, since $x_{i-1} < x_i$. The set $\{a, b\}$ is a partition of $[a, b]$ with one subinterval, so partitions exist. For each $n \in \mathbb{N}$ the *uniform* partition P_n with $x_i = a + i(b-a)/n$ has $\Delta x_i = (b-a)/n$ for every i and mesh $(b-a)/n$. Given $t > 0$, applying theorem 1.4.1(3) to the positive number $t/(b-a)$ produces an n with $1/n < t/(b-a)$, that is $\|P_n\| < t$, so the uniform meshes fall below any prescribed positive number.

A bounded $f: [a, b] \rightarrow \mathbb{R}$ is bounded on each subinterval of a partition, since $f([x_{i-1}, x_i]) \subseteq f([a, b])$, so the supremum and the infimum of f over each subinterval exist by theorem 1.2.4 and corollary 1.2.5. These $2n$ numbers are the data from which the two sums are assembled.

Definition 5.1.2: Upper and Lower Darboux Sums

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded and let $P = \{x_0, \dots, x_n\}$ be a partition of $[a, b]$. For $1 \leq i \leq n$ write

$$m_i = \inf_{[x_{i-1}, x_i]} f, \quad M_i = \sup_{[x_{i-1}, x_i]} f$$

The *lower Darboux sum* and the *upper Darboux sum* of f over P are

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i, \quad U(P, f) = \sum_{i=1}^n M_i \Delta x_i$$

Since f is bounded there are real numbers m and M_0 with $m \leq f(x) \leq M_0$ for every $x \in [a, b]$. The number $M = |m| + |M_0| + 1$ is then positive and satisfies $|f(x)| \leq M$ for every x : indeed $f(x) \leq M_0 \leq |M_0| \leq M$ and $-M \leq -|m| \leq m \leq f(x)$ by proposition 1.3.8(1), so proposition 1.3.8(2) applies with $c = M$. Such an M is fixed whenever an argument below needs one.

The two sums are the areas of two families of rectangles, one family containing the region under the graph and one contained in it, and a single partition already produces a numerical bracket. Take $f(x) = x^2$ on $[0, 1]$ and $P = \{0, 1/2, 1\}$. For $0 \leq x \leq y$, multiplying $x \leq y$ first by $x \geq 0$ and then by $y \geq 0$ gives $x^2 \leq xy \leq y^2$, so f is increasing on $[0, 1]$ and attains its extreme values on a subinterval at the endpoints. Hence $m_1 = f(0) = 0$, $M_1 = f(1/2) = 1/4$, $m_2 = 1/4$ and $M_2 = f(1) = 1$, while $\Delta x_1 = \Delta x_2 = 1/2$. So

$$L(P, f) = 0 \cdot \frac{1}{2} + \frac{1}{4} \cdot \frac{1}{2} = \frac{1}{8}, \quad U(P, f) = \frac{1}{4} \cdot \frac{1}{2} + 1 \cdot \frac{1}{2} = \frac{5}{8}$$

and any number deserving the name “the integral of x^2 over $[0, 1]$ ” lies between $1/8$ and $5/8$. The bracket is wide because the partition is coarse. Cutting each subinterval into smaller pieces replaces M_i by suprema over smaller sets, which can only be smaller, and this is the operation the next definition names.

Definition 5.1.3: [Real] Refinement

Let P and P^* be partitions of $[a, b]$. Then P^* is a *refinement* of P if $P \subseteq P^*$. Given two partitions P and Q of $[a, b]$, their union $P \cup Q$ is a partition of $[a, b]$, called the *common refinement* of P and Q , and it refines each of them.

The union of two partitions is a finite subset of $[a, b]$ containing a and b , and listing its elements in increasing order exhibits it as a partition, so the common refinement is well defined. Refining subdivides some subintervals of P and keeps the others unchanged.

One observation about suprema over subsets is used at every step below. Let $\emptyset \neq A \subseteq B \subseteq [a, b]$ and let f be bounded on B . Applying proposition 1.3.2(1) to the nonempty sets $f(A) \subseteq f(B)$ gives $\sup_A f \leq \sup_B f$. For infima, $\inf_B f$ is a lower bound of $f(B)$ and therefore of the subset $f(A)$, so $\inf_B f \leq \inf_A f$ because $\inf_A f$ is the greatest lower bound of $f(A)$ (definition 1.2.2). Cutting a set down cannot raise its supremum and cannot lower its infimum.

Proposition 5.1.4: Refinement Moves the Darboux Sums Together

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, and write $m = \inf_{[a,b]} f$ and $M = \sup_{[a,b]} f$.

1. $L(P, f) \leq U(P, f)$ for every partition P of $[a, b]$.
2. If P^* refines P , then $L(P, f) \leq L(P^*, f)$ and $U(P^*, f) \leq U(P, f)$.
3. $m(b-a) \leq L(P, f)$ and $U(P, f) \leq M(b-a)$ for every partition P of $[a, b]$.

Proof:

1. Fix i and pick any point z of the nonempty set $[x_{i-1}, x_i]$. Then $m_i \leq f(z) \leq M_i$, so $m_i \leq M_i$. Multiplying by $\Delta x_i > 0$ gives $m_i \Delta x_i \leq M_i \Delta x_i$, and summing over $i = 1, \dots, n$ gives $L(P, f) \leq U(P, f)$.
2. It is enough to treat the case where P^* has exactly one point more than P , since a refinement P^* of P is obtained from P by adjoining the finitely many points of $P^* \setminus P$ one at a time, and each intermediate stage is again a partition refining the previous one. The general case then follows by induction on the number of adjoined points, the case of zero points being the equality $P^* = P$.

So suppose $P^* = P \cup \{y\}$ with $y \notin P$, say $x_{i-1} < y < x_i$. The subintervals of P^* are those of P with $[x_{i-1}, x_i]$ replaced by $[x_{i-1}, y]$ and $[y, x_i]$, and every other term of the two upper sums is identical. Writing $w_1 = \sup_{[x_{i-1}, y]} f$ and $w_2 = \sup_{[y, x_i]} f$, the subset observation above gives $w_1 \leq M_i$ and $w_2 \leq M_i$, so

$$w_1(y - x_{i-1}) + w_2(x_i - y) \leq M_i(y - x_{i-1}) + M_i(x_i - y) = M_i \Delta x_i$$

which is exactly $U(P^*, f) \leq U(P, f)$ after the common terms are cancelled. Writing $v_1 = \inf_{[x_{i-1}, y]} f$ and $v_2 = \inf_{[y, x_i]} f$, the same observation gives $m_i \leq v_1$ and $m_i \leq v_2$, so

$$m_i \Delta x_i = m_i(y - x_{i-1}) + m_i(x_i - y) \leq v_1(y - x_{i-1}) + v_2(x_i - y)$$

which is $L(P, f) \leq L(P^*, f)$.

3. Each $[x_{i-1}, x_i]$ is a nonempty subset of $[a, b]$, so the subset observation gives $m \leq m_i$ and $M_i \leq M$. Multiplying by $\Delta x_i > 0$ and summing,

$$m(b-a) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i = L(P, f)$$

using (5.1), and the same computation with $M_i \leq M$ gives $U(P, f) \leq M(b-a)$, completing the proof.

Part (2) is the reason the two sums are worth watching at all: cutting the subdivision finer can only raise the lower sum and only lower the upper one, so the bracket produced by a partition never widens under refinement. Part (1) compares the two sums attached to the *same* partition. What the theory needs is the comparison for two unrelated partitions, and the common refinement is what delivers it.

Theorem 5.1.5: Every Lower Sum Is Below Every Upper Sum

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded.

1. $L(P, f) \leq U(Q, f)$ for all partitions P and Q of $[a, b]$.
2. The set of all lower sums of f is nonempty and bounded above, the set of all upper sums is nonempty and bounded below, and

$$\sup_P L(P, f) \leq \inf_Q U(Q, f)$$

the supremum and infimum being taken over all partitions of $[a, b]$.

Proof:

1. Let $P^* = P \cup Q$, which by definition 5.1.3 is a partition refining both P and Q . Since P^* refines P , proposition 5.1.4(2) gives $L(P, f) \leq L(P^*, f)$. Since P^* refines Q , the same part gives $U(P^*, f) \leq U(Q, f)$. Combining these with proposition 5.1.4(1) applied to P^* ,

$$L(P, f) \leq L(P^*, f) \leq U(P^*, f) \leq U(Q, f)$$

2. Both sets are nonempty because $\{a, b\}$ is a partition of $[a, b]$. Fix a partition Q . By part (1) the number $U(Q, f)$ is an upper bound of the set of all lower sums, so that set is bounded above and its supremum $s = \sup_P L(P, f)$ exists by theorem 1.2.4, with $s \leq U(Q, f)$. Since Q was arbitrary, s is a lower bound of the set of all upper sums. That set is therefore bounded below, its infimum exists by corollary 1.2.5, and $s \leq \inf_Q U(Q, f)$ because the infimum is the greatest lower bound, completing the proof.

The two numbers produced in part (2) exist for every bounded function, whether or not the approximation converges, and they are what the definition of the integral compares.

Definition 5.1.6: Upper and Lower Integrals

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. The *lower integral* and the *upper integral* of f over $[a, b]$ are

$$\int_a^b f = \sup_P L(P, f), \quad \overline{\int}_a^b f = \inf_Q U(Q, f)$$

the supremum and infimum being taken over all partitions of $[a, b]$. Both exist and satisfy $\int_a^b f \leq \overline{\int}_a^b f$ by theorem 5.1.5(2).

For the function $f(x) = x^2$ on $[0, 1]$ computed above, the partition $\{0, 1/2, 1\}$ already gives $1/8 \leq \int_0^1 f$ and $\overline{\int}_0^1 f \leq 5/8$, since a lower integral is at least each lower sum and an upper integral at most each upper sum. Refining the partition tightens both bounds, and the question is whether the two numbers coincide.

Definition 5.1.7: [Real] Riemann Integrable

A bounded function $f: [a, b] \rightarrow \mathbb{R}$ is *Riemann integrable* on $[a, b]$ if

$$\int_a^b f = \overline{\int_a^b f}$$

The set of such functions is written $\mathcal{R}[a, b]$, and for $f \in \mathcal{R}[a, b]$ the common value is the *Riemann integral* of f over $[a, b]$, written $\int_a^b f$ or $\int_a^b f(x) dx$.

Unboundedness is excluded at the outset, since an unbounded f is unbounded on some subinterval of every partition P , where it then has no supremum or no infimum, so that one of $U(P, f)$ and $L(P, f)$ is undefined for every P . Deciding integrability from definition 5.1.7 means comparing a supremum over all partitions with an infimum over all partitions, which is a statement about two infinite families at once. The next theorem replaces it by a condition on a single partition at a time, and every existence theorem in section 5.2 is proved by producing such a partition.

Theorem 5.1.8: The Integrability Criterion

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded.

1. $f \in \mathcal{R}[a, b]$ if and only if for every $\epsilon > 0$ there is a partition P of $[a, b]$ with

$$U(P, f) - L(P, f) < \epsilon$$

2. If $f \in \mathcal{R}[a, b]$, then $L(P, f) \leq \int_a^b f \leq U(P, f)$ for every partition P of $[a, b]$, and $\int_a^b f$ is the only real number with that property.

Proof:

Step 1: the condition is sufficient. Suppose that for every $\epsilon > 0$ some partition P satisfies $U(P, f) - L(P, f) < \epsilon$. Fix such an ϵ and such a P . By definition 5.1.6 the lower integral is at least $L(P, f)$ and the upper integral is at most $U(P, f)$, so

$$\overline{\int_a^b f} - \int_a^b f \leq U(P, f) - L(P, f) < \epsilon$$

Also $\overline{\int_a^b f} - \int_a^b f \geq 0$ by definition 5.1.6. So the number $d = \overline{\int_a^b f} - \int_a^b f$ satisfies $0 \leq d \leq \epsilon$ for every $\epsilon > 0$, and $|d - 0| = d$ by definition 1.3.6, so proposition 1.4.7(2) gives $d = 0$. Hence $f \in \mathcal{R}[a, b]$.

Step 2: the condition is necessary. Suppose $f \in \mathcal{R}[a, b]$ and let $\epsilon > 0$. Applying proposition 1.2.6 to the set of lower sums, whose supremum is $\int_a^b f$, produces a partition P_1 with

$$L(P_1, f) > \int_a^b f - \frac{\epsilon}{2}$$

Applying lemma 1.3.1 to the set of upper sums, whose infimum is $\int_a^b f$, produces a partition P_2

with $U(P_2, f) < \overline{\int_a^b} f + \epsilon/2$. Let $P = P_1 \cup P_2$. By proposition 5.1.4(2), $L(P, f) \geq L(P_1, f)$ and $U(P, f) \leq U(P_2, f)$, so

$$U(P, f) - L(P, f) < \left(\overline{\int_a^b} f + \frac{\epsilon}{2} \right) - \left(\underline{\int_a^b} f - \frac{\epsilon}{2} \right) = \epsilon$$

the last equality because the two integrals are equal.

Step 3: the bracket and its uniqueness. Let $f \in \mathcal{R}[a, b]$ and let P be any partition. Then $L(P, f) \leq \underline{\int_a^b} f = \int_a^b f$ and $\int_a^b f = \overline{\int_a^b} f \leq U(P, f)$, again by definition 5.1.6. Now let A be any real number with $L(P, f) \leq A \leq U(P, f)$ for every partition P . The first inequality says that A is an upper bound of the set of lower sums, so $\underline{\int_a^b} f \leq A$ because that lower integral is the least upper bound. The second says that A is a lower bound of the set of upper sums, so $A \leq \overline{\int_a^b} f$. The two integrals agree, so $A = \int_a^b f$, completing the proof.

The criterion says that integrability is a statement about the *gap* $U(P, f) - L(P, f)$, and part (2) says that once the gap is small the integral is located by a single partition to within that gap. Both halves are used in the computation below, the first to prove that the integral exists and the second to identify its value.

Example 5.1.9: The Integrals of x and of a Step Function

1. Let $f(x) = x$ on $[0, 1]$ and let P_n be the uniform partition with $x_i = i/n$, so $\Delta x_i = 1/n$. The function is increasing, so $m_i = f(x_{i-1}) = (i-1)/n$ and $M_i = f(x_i) = i/n$. Hence

$$U(P_n, f) = \sum_{i=1}^n \frac{i}{n} \cdot \frac{1}{n} = \frac{1}{n^2} \sum_{i=1}^n i, \quad L(P_n, f) = \frac{1}{n^2} \sum_{i=1}^n (i-1)$$

The identity $\sum_{i=1}^n i = n(n+1)/2$ holds for $n = 1$, since both sides equal 1, and if it holds for n then

$$\sum_{i=1}^{n+1} i = \frac{n(n+1)}{2} + (n+1) = \frac{(n+1)(n+2)}{2}$$

so it holds for every n by induction. Therefore $U(P_n, f) = (n+1)/(2n)$, and $\sum_{i=1}^n (i-1) = \sum_{i=1}^n i - n = n(n-1)/2$ gives $L(P_n, f) = (n-1)/(2n)$. The gap is

$$U(P_n, f) - L(P_n, f) = \frac{2}{2n} = \frac{1}{n}$$

Given $\epsilon > 0$, theorem 1.4.1(3) produces an n with $1/n < \epsilon$, so theorem 5.1.8(1) gives $f \in \mathcal{R}[0, 1]$. For the value, theorem 5.1.8(2) gives

$$\frac{n-1}{2n} \leq \int_0^1 f \leq \frac{n+1}{2n}$$

for every n , that is $-1/(2n) \leq \int_0^1 f - 1/2 \leq 1/(2n)$. By proposition 1.3.8(2) this reads $|\int_0^1 f - 1/2| \leq 1/(2n)$, and given $\epsilon > 0$ an n with $1/n < \epsilon$ makes the right side smaller than ϵ . So proposition 1.4.7(2) gives $\int_0^1 x \, dx = 1/2$.

2. Let $g: [0, 2] \rightarrow \mathbb{R}$ be $g(x) = 0$ for $0 \leq x \leq 1$ and $g(x) = 1$ for $1 < x \leq 2$, a bounded function with a single discontinuity. Fix δ with $0 < \delta < 1$ and take $P = \{0, 1, 1 + \delta, 2\}$. On $[0, 1]$ the function is constantly 0, so $m_1 = M_1 = 0$. On $[1, 1 + \delta]$ it takes both the value 0 (at $x = 1$) and the value 1 (at every x in $(1, 1 + \delta]$), so $m_2 = 0$ and $M_2 = 1$. On $[1 + \delta, 2]$ it is constantly 1, so $m_3 = M_3 = 1$. Hence

$$U(P, g) = 0 \cdot 1 + 1 \cdot \delta + 1 \cdot (1 - \delta) = 1, \quad L(P, g) = 0 \cdot 1 + 0 \cdot \delta + 1 \cdot (1 - \delta) = 1 - \delta$$

so $U(P, g) - L(P, g) = \delta$. Since $\delta \in (0, 1)$ was arbitrary and theorem 1.4.1(3) makes $1/n$ smaller than any prescribed positive number, theorem 5.1.8(1) gives $g \in \mathcal{R}[0, 2]$. By theorem 5.1.8(2), $1 - \delta \leq \int_0^2 g \leq 1$ for every such δ , so $1 \leq \int_0^2 g + \delta$ for every $\delta > 0$ and proposition 1.4.7(1) gives $1 \leq \int_0^2 g$. Together with $\int_0^2 g \leq 1$ this gives $\int_0^2 g = 1$.

The step function of part (2) is the first sign that continuity is not required: a function with a jump can still have its upper and lower sums forced together, because the partition can confine the jump to one short subinterval whose contribution to the gap is at most the oscillation of g times the length of that subinterval. Section 5.2 turns this observation into three existence theorems.

Nothing so far samples f anywhere. Both Darboux sums are built from suprema and infima, and neither m_i nor M_i need be a value of f . Riemann's own construction picks an actual point of each subinterval and evaluates f there.

Definition 5.1.10: Tagged Partitions and Riemann Sums

A *tagged partition* of $[a, b]$ is a pair (P, ξ) consisting of a partition $P = \{x_0, \dots, x_n\}$ together with a choice of points $\xi = (\xi_1, \dots, \xi_n)$ with $\xi_i \in [x_{i-1}, x_i]$ for each i . The points ξ_i are the *tags*. For $f: [a, b] \rightarrow \mathbb{R}$ the *Riemann sum* of f over (P, ξ) is

$$R(P, \xi, f) = \sum_{i=1}^n f(\xi_i) \Delta x_i$$

Since $\xi_i \in [x_{i-1}, x_i]$, the value $f(\xi_i)$ lies between m_i and M_i , so multiplying by $\Delta x_i > 0$ and summing gives

$$L(P, f) \leq R(P, \xi, f) \leq U(P, f) \quad (5.2)$$

for every tagged partition. A Riemann sum therefore lies in the same bracket the Darboux sums produce. For $f(x) = x$ on $[0, 1]$, the uniform partition P_n with the midpoint tags $\xi_i = (2i - 1)/(2n)$ gives

$$R(P_n, \xi, f) = \sum_{i=1}^n \frac{2i-1}{2n} \cdot \frac{1}{n} = \frac{1}{2n^2} \left(2 \sum_{i=1}^n i - n \right) = \frac{1}{2n^2} (n(n+1) - n) = \frac{1}{2}$$

using the identity established in example 5.1.9. Every midpoint sum equals the integral computed there exactly, while the two Darboux sums approach it from either side.

The bracket in (5.2) is available for every choice of tags, and the next theorem converts it into a limit statement: once the mesh is small enough, every Riemann sum over every choice of tags is within ϵ of the integral. The condition characterizes integrability and the value of the integral at the same time.

Theorem 5.1.11: Riemann Sums Compute the Darboux Integral

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded and let $A \in \mathbb{R}$. Then $f \in \mathcal{R}[a, b]$ with $\int_a^b f = A$ if and only if for every $\epsilon > 0$ there is a $\delta > 0$ such that every tagged partition (P, ξ) of $[a, b]$ with $\|P\| < \delta$ satisfies

$$|R(P, \xi, f) - A| < \epsilon$$

Proof:

Fix $M > 0$ with $|f(x)| \leq M$ on $[a, b]$, as produced after definition 5.1.2.

Step 1: adjoining one point to a partition changes an upper sum by at most $2M$ times the mesh. Let Q be a partition of $[a, b]$ and let $y \notin Q$, say $u < y < v$ for consecutive points u, v of Q . As in the proof of proposition 5.1.4(2), the sums $U(Q, f)$ and $U(Q \cup \{y\}, f)$ differ only in the terms coming from $[u, v]$, so

$$U(Q, f) - U(Q \cup \{y\}, f) = \sup_{[u, v]} f \cdot (v - u) - \sup_{[u, y]} f \cdot (y - u) - \sup_{[y, v]} f \cdot (v - y)$$

Every value of f lies between $-M$ and M , so each of the three suprema lies between $-M$ and M as well, the first because M is an upper bound and the last two because $-M$ is a lower bound and a supremum is at least each element of a nonempty set. Bounding the first supremum above by M and the other two below by $-M$,

$$U(Q, f) - U(Q \cup \{y\}, f) \leq M(v - u) + M(y - u) + M(v - y) = 2M(v - u) \leq 2M\|Q\|$$

The same computation with infima, using $\inf_{[u, v]} f \geq -M$ and $\inf_{[u, y]} f, \inf_{[y, v]} f \leq M$, gives $L(Q \cup \{y\}, f) - L(Q, f) \leq 2M\|Q\|$. Adjoining a point never increases the mesh, since it replaces one subinterval by two shorter ones.

Step 2: integrability gives the ϵ - δ condition. Suppose $f \in \mathcal{R}[a, b]$ with $\int_a^b f = A$, and let $\epsilon > 0$. By theorem 5.1.8(1) there is a partition P_0 with $U(P_0, f) - L(P_0, f) < \epsilon/3$. Let k be the number of points of P_0 lying strictly between a and b , and put

$$\delta = \frac{\epsilon}{6M(k+1)} > 0$$

Let (P, ξ) be any tagged partition with $\|P\| < \delta$ and set $P^* = P \cup P_0$. Then P^* is obtained from P by adjoining at most k points, namely those points of P_0 strictly between a and b that are not already in P . Adjoining them one at a time and applying Step 1 at each stage, where the mesh is at most $\|P\| < \delta$,

$$U(P, f) - U(P^*, f) \leq 2Mk\delta = \frac{k\epsilon}{3(k+1)} < \frac{\epsilon}{3}$$

and likewise $L(P^*, f) - L(P, f) < \epsilon/3$. Since P^* refines P_0 , proposition 5.1.4(2) gives $U(P^*, f) - L(P^*, f) \leq U(P_0, f) - L(P_0, f) < \epsilon/3$. Adding the three estimates,

$$U(P, f) - L(P, f) = (U(P, f) - U(P^*, f)) + (U(P^*, f) - L(P^*, f)) + (L(P^*, f) - L(P, f)) < \epsilon$$

Both $R(P, \xi, f)$ and A lie in the interval from $L(P, f)$ to $U(P, f)$, the first by (5.2) and the second by theorem 5.1.8(2). Two numbers in an interval differ by at most its length, so $|R(P, \xi, f) - A| \leq U(P, f) - L(P, f) < \epsilon$.

Step 3: the ϵ - δ condition gives integrability. Suppose the condition holds. Let $\epsilon > 0$ and take $\delta > 0$ for the number $\epsilon/4$. Fix any partition P with $\|P\| < \delta$, say with subintervals $[x_{i-1}, x_i]$ for $1 \leq i \leq n$, and put $\eta = \epsilon/(4(b-a)) > 0$. By proposition 1.2.6 applied to the nonempty set $f([x_{i-1}, x_i])$, whose supremum is M_i , there is $s_i \in [x_{i-1}, x_i]$ with $f(s_i) > M_i - \eta$. By lemma 1.3.1 applied to the same set, there is $t_i \in [x_{i-1}, x_i]$ with $f(t_i) < m_i + \eta$. Writing s and t for the resulting tag choices and using (5.1),

$$U(P, f) - R(P, s, f) = \sum_{i=1}^n (M_i - f(s_i)) \Delta x_i < \eta \sum_{i=1}^n \Delta x_i = \eta(b-a) = \frac{\epsilon}{4}$$

and the same computation gives $R(P, t, f) - L(P, f) < \epsilon/4$. Both (P, s) and (P, t) have mesh less than δ , so the hypothesis gives $|R(P, s, f) - A| < \epsilon/4$ and $|R(P, t, f) - A| < \epsilon/4$, whence $R(P, s, f) - A < \epsilon/4$ and $A - R(P, t, f) < \epsilon/4$ by proposition 1.3.8(2). Adding the four estimates,

$$U(P, f) - L(P, f) = (U(P, f) - R(P, s, f)) + (R(P, s, f) - A) + (A - R(P, t, f)) + (R(P, t, f) - L(P, f)) < \epsilon$$

So $f \in \mathcal{R}[a, b]$ by theorem 5.1.8(1).

Step 4: the value is A . Keep the notation of Step 3 and let $\eta' > 0$ be given. Running Step 3 with $\epsilon = 4\eta'/5$ produces a partition P and tags s with $U(P, f) - L(P, f) < \epsilon$ and $|R(P, s, f) - A| < \epsilon/4$. Both $\int_a^b f$ and $R(P, s, f)$ lie between $L(P, f)$ and $U(P, f)$, by theorem 5.1.8(2) and (5.2), so they differ by at most $U(P, f) - L(P, f) < \epsilon$. Then proposition 1.3.8(4) gives

$$\left| \int_a^b f - A \right| \leq \left| \int_a^b f - R(P, s, f) \right| + |R(P, s, f) - A| < \epsilon + \frac{\epsilon}{4} = \frac{5\epsilon}{4} = \eta'$$

As $\eta' > 0$ was arbitrary, proposition 1.4.7(2) gives $\int_a^b f = A$, completing the proof.

The theorem makes each of the two constructions available wherever it is the convenient one. Existence proofs run on Darboux sums, because a supremum is available without knowing where f is large, while a concrete evaluation runs on Riemann sums, because those are sums of actual values of f . Nothing in the statement constrains how the tags are chosen: left endpoints, right endpoints and midpoints all converge to the same number, and so does any other rule.

One consequence of a small gap is recorded separately, because it is the form in which the criterion is applied when two different tag choices have to be compared.

Proposition 5.1.12: A Small Gap Controls Every Pair of Sample Points

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, let $\epsilon > 0$, and let $P = \{x_0, \dots, x_n\}$ be a partition with $U(P, f) - L(P, f) < \epsilon$. Then for any points $s_i, t_i \in [x_{i-1}, x_i]$, $1 \leq i \leq n$,

$$\sum_{i=1}^n |f(s_i) - f(t_i)| \Delta x_i < \epsilon$$

Proof:

Fix i . The set $[x_{i-1}, x_i]$ is nonempty and f is bounded on it, with supremum M_i and infimum m_i , so proposition 1.3.10(1) applied to f on $[x_{i-1}, x_i]$ gives $|f(s_i) - f(t_i)| \leq M_i - m_i$. Multiplying by $\Delta x_i > 0$ and summing over i ,

$$\sum_{i=1}^n |f(s_i) - f(t_i)| \Delta x_i \leq \sum_{i=1}^n (M_i - m_i) \Delta x_i = U(P, f) - L(P, f) < \epsilon$$

as we sought to show.

The proposition converts a bound on the single number $U(P, f) - L(P, f)$ into a bound on a whole family of sums, one for each way of sampling f twice in every subinterval, which is what makes it usable when the sample points are chosen by something other than the argument at hand. It is used only in the proof of theorem 5.4.7, which a reader may skip.

The rest of the section establishes the arithmetic of the integral. Linearity is the statement that $\mathcal{R}[a, b]$ is closed under sums and scalar multiples and that $\int_a^b$ respects both operations. The failure of the sum rule for suprema, recorded in proposition 1.3.4, shapes the proof: the upper sum of $f + g$ is at most the sum of the upper sums and can be strictly smaller, so the estimate available for upper sums and the one available for lower sums run in opposite directions, and the criterion is what closes the space between them.

Theorem 5.1.13: Linearity of the Integral

Let $f, g \in \mathcal{R}[a, b]$ and let $c \in \mathbb{R}$. Then $f + g$ and cf belong to $\mathcal{R}[a, b]$, and

$$\int_a^b (f + g) = \int_a^b f + \int_a^b g, \quad \int_a^b cf = c \int_a^b f$$

Proof:

Step 1: the sum is integrable. Both f and g are bounded, so $f + g$ is bounded, since $|f(x) + g(x)| \leq |f(x)| + |g(x)|$ by proposition 1.3.8(4). Let $\epsilon > 0$. By theorem 5.1.8(1) there are partitions P_1 and P_2 with $U(P_1, f) - L(P_1, f) < \epsilon/2$ and $U(P_2, g) - L(P_2, g) < \epsilon/2$. Put $P = P_1 \cup P_2$. By proposition 5.1.4(2) the same two estimates hold for P in place of P_1 and of P_2 , since refining raises a lower sum and lowers an upper sum.

On each subinterval $[x_{i-1}, x_i]$ of P , the functions f and g are bounded, so proposition 1.3.4 applies with $D = [x_{i-1}, x_i]$ and gives

$$\sup_{[x_{i-1}, x_i]} (f + g) \leq \sup_{[x_{i-1}, x_i]} f + \sup_{[x_{i-1}, x_i]} g, \quad \inf_{[x_{i-1}, x_i]} (f + g) \geq \inf_{[x_{i-1}, x_i]} f + \inf_{[x_{i-1}, x_i]} g$$

Multiplying each by $\Delta x_i > 0$ and summing over i ,

$$U(P, f + g) \leq U(P, f) + U(P, g), \quad L(P, f + g) \geq L(P, f) + L(P, g) \quad (5.3)$$

Subtracting,

$$U(P, f + g) - L(P, f + g) \leq (U(P, f) - L(P, f)) + (U(P, g) - L(P, g)) < \epsilon$$

so $f + g \in \mathcal{R}[a, b]$ by theorem 5.1.8(1).

Step 2: the value of the sum. Keep P and ϵ from Step 1. By theorem 5.1.8(2) applied to $f + g$ and then to (5.3),

$$\int_a^b (f + g) \leq U(P, f + g) \leq U(P, f) + U(P, g)$$

and $U(P, f) < L(P, f) + \epsilon/2 \leq \int_a^b f + \epsilon/2$, with the same estimate for g , so $\int_a^b (f + g) < \int_a^b f + \int_a^b g + \epsilon$. In the other direction,

$$\int_a^b (f + g) \geq L(P, f + g) \geq L(P, f) + L(P, g) > \left(U(P, f) - \frac{\epsilon}{2}\right) + \left(U(P, g) - \frac{\epsilon}{2}\right) \geq \int_a^b f + \int_a^b g - \epsilon$$

using $\int_a^b f \leq U(P, f)$ and $\int_a^b g \leq U(P, g)$. The two displays say that the difference between $\int_a^b (f + g)$ and $\int_a^b f + \int_a^b g$ lies strictly between $-\epsilon$ and ϵ , hence has absolute value at most ϵ by proposition 1.3.8(2). Since $\epsilon > 0$ was arbitrary, proposition 1.4.7(2) gives $\int_a^b (f + g) = \int_a^b f + \int_a^b g$.

Step 3: positive scalars. Let $c > 0$. On each subinterval, $(cf)([x_{i-1}, x_i]) = c \cdot f([x_{i-1}, x_i])$, so proposition 1.3.2(2) gives $\sup_{[x_{i-1}, x_i]}(cf) = c \sup_{[x_{i-1}, x_i]} f$. For the infimum, proposition 1.3.2(3) applied to a nonempty bounded set S says $\sup(-S) = -\inf S$, and applying it to $-S$ in place of S gives $\inf(-S) = -\sup S$. Hence

$$\inf(cS) = -\sup(-cS) = -\sup(c(-S)) = -c \sup(-S) = -c(-\inf S) = c \inf S$$

where the third equality is proposition 1.3.2(2) again. Applying this with $S = f([x_{i-1}, x_i])$ gives $\inf_{[x_{i-1}, x_i]}(cf) = c \inf_{[x_{i-1}, x_i]} f$. Multiplying by Δx_i and summing, $U(P, cf) = cU(P, f)$ and $L(P, cf) = cL(P, f)$ for every partition P . Given $\epsilon > 0$, choose P with $U(P, f) - L(P, f) < \epsilon/c$, which theorem 5.1.8(1) provides; then $U(P, cf) - L(P, cf) = c(U(P, f) - L(P, f)) < \epsilon$, so $cf \in \mathcal{R}[a, b]$. For the value, $L(P, cf) \leq \int_a^b cf \leq U(P, cf)$ reads $cL(P, f) \leq \int_a^b cf \leq cU(P, f)$, while $cL(P, f) \leq c \int_a^b f \leq cU(P, f)$, so the two numbers $\int_a^b cf$ and $c \int_a^b f$ lie in an interval of length $c(U(P, f) - L(P, f)) < \epsilon$ and differ by at most ϵ . As $\epsilon > 0$ was arbitrary, proposition 1.4.7(2) gives $\int_a^b cf = c \int_a^b f$.

Step 4: the remaining scalars. If $c = 0$ then cf is the constant function 0, every Darboux sum of which is 0 by definition 5.1.2, so $cf \in \mathcal{R}[a, b]$ with $\int_a^b cf = 0 = c \int_a^b f$. For $c = -1$, the identity $\inf(-S) = -\sup S$ and its companion $\sup(-S) = -\inf S$ from Step 3 give $U(P, -f) = -L(P, f)$ and $L(P, -f) = -U(P, f)$ for every P , so $U(P, -f) - L(P, -f) = U(P, f) - L(P, f)$ and $-f \in \mathcal{R}[a, b]$ by theorem 5.1.8(1). Moreover $L(P, -f) \leq \int_a^b (-f) \leq U(P, -f)$ reads $-U(P, f) \leq \int_a^b (-f) \leq -L(P, f)$, while $-U(P, f) \leq -\int_a^b f \leq -L(P, f)$ follows from $L(P, f) \leq \int_a^b f \leq U(P, f)$. So $\int_a^b (-f)$ and $-\int_a^b f$ both lie in an interval of length $U(P, f) - L(P, f)$, which theorem 5.1.8(1) makes smaller than any prescribed $\epsilon > 0$ for a suitable P , and proposition 1.4.7(2) gives $\int_a^b (-f) = -\int_a^b f$. Finally, for $c < 0$ write $cf = (-c)(-f)$ with $-c > 0$ and apply Step 3 to the integrable function $-f$, obtaining

$$\int_a^b cf = (-c) \int_a^b (-f) = (-c) \left(-\int_a^b f \right) = c \int_a^b f$$

completing the proof.

Linearity makes $\mathcal{R}[a, b]$ a real vector space of functions and $\int_a^b$ a linear functional on it. The remaining facts concern the order and the interval. The first two are immediate from the bracket in theorem 5.1.8(2), and the third is what allows an integral to be computed piece by piece and is what section 5.3 needs in order to differentiate $x \mapsto \int_a^x f$.

Proposition 5.1.14: Order Bounds and Additivity over a Subdivision Point

Let $f, g: [a, b] \rightarrow \mathbb{R}$ be bounded.

1. If $f, g \in \mathcal{R}[a, b]$ and $f(x) \leq g(x)$ for every $x \in [a, b]$, then $\int_a^b f \leq \int_a^b g$.
2. If $f \in \mathcal{R}[a, b]$ and $m \leq f(x) \leq M$ for every $x \in [a, b]$, then $m(b-a) \leq \int_a^b f \leq M(b-a)$.
3. Let $a < c < b$. Then $f \in \mathcal{R}[a, b]$ if and only if the restrictions of f to $[a, c]$ and to $[c, b]$ belong to $\mathcal{R}[a, c]$ and $\mathcal{R}[c, b]$ respectively, and in that case

$$\int_a^b f = \int_a^c f + \int_c^b f$$

Proof:

1. Fix a partition P and a subinterval $[x_{i-1}, x_i]$. For every x there, $f(x) \leq g(x) \leq \sup_{[x_{i-1}, x_i]} g$, so $\sup_{[x_{i-1}, x_i]} f$ is an upper bound of $f([x_{i-1}, x_i])$ and therefore $\sup_{[x_{i-1}, x_i]} f \leq \sup_{[x_{i-1}, x_i]} g$, the supremum of f being the least upper bound (definition 1.2.2). Multiplying by $\Delta x_i > 0$ and summing gives $U(P, f) \leq U(P, g)$ for every P . By theorem 5.1.8(2), $\int_a^b f \leq U(P, f) \leq U(P, g)$ for every P , so $\int_a^b f$ is a lower bound of the set of upper sums of g and hence $\int_a^b f \leq \overline{\int_a^b g} = \int_a^b g$, the infimum being the greatest lower bound.
2. Apply theorem 5.1.8(2) to the partition $P = \{a, b\}$, whose single subinterval is $[a, b]$ and whose two Darboux sums are $(\inf_{[a, b]} f)(b-a)$ and $(\sup_{[a, b]} f)(b-a)$. The number m is a lower bound of $f([a, b])$ and M an upper bound, so $m \leq \inf_{[a, b]} f$ and $\sup_{[a, b]} f \leq M$ (definition 1.2.2). Multiplying by $b-a > 0$,

$$m(b-a) \leq L(P, f) \leq \int_a^b f \leq U(P, f) \leq M(b-a)$$

3. Note first that f is bounded on $[a, c]$ and on $[c, b]$, both being subsets of $[a, b]$. A partition P of $[a, b]$ containing c splits as $P = P_1 \cup P_2$ with $P_1 = P \cap [a, c]$ a partition of $[a, c]$ and $P_2 = P \cap [c, b]$ a partition of $[c, b]$, and the subintervals of P are exactly those of P_1 together with those of P_2 . Hence

$$U(P, f) = U(P_1, f) + U(P_2, f), \quad L(P, f) = L(P_1, f) + L(P_2, f) \quad (5.4)$$

and conversely any partition P_1 of $[a, c]$ and P_2 of $[c, b]$ have union a partition P of $[a, b]$ containing c , for which (5.4) holds.

Necessity. Suppose $f \in \mathcal{R}[a, b]$ and let $\epsilon > 0$. By theorem 5.1.8(1) there is a partition P with $U(P, f) - L(P, f) < \epsilon$. Put $P^* = P \cup \{c\}$, a refinement of P , so proposition 5.1.4(2) gives $U(P^*, f) - L(P^*, f) \leq U(P, f) - L(P, f) < \epsilon$. Splitting P^* as above and using (5.4),

$$(U(P_1, f) - L(P_1, f)) + (U(P_2, f) - L(P_2, f)) < \epsilon$$

Both bracketed numbers are nonnegative by proposition 5.1.4(1), so each is less than ϵ . By theorem 5.1.8(1) applied on $[a, c]$ and on $[c, b]$, both restrictions are integrable.

Sufficiency and the value. Suppose both restrictions are integrable and let $\epsilon > 0$. Choose partitions P_1 of $[a, c]$ and P_2 of $[c, b]$ with $U(P_1, f) - L(P_1, f) < \epsilon/2$ and $U(P_2, f) - L(P_2, f) < \epsilon/2$, and let $P = P_1 \cup P_2$. By (5.4), $U(P, f) - L(P, f) < \epsilon$, so $f \in \mathcal{R}[a, b]$ by theorem 5.1.8(1). For the value, theorem 5.1.8(2) on the three intervals gives

$$\int_a^b f \leq U(P, f) = U(P_1, f) + U(P_2, f) < L(P_1, f) + L(P_2, f) + \epsilon \leq \int_a^c f + \int_c^b f + \epsilon$$

and, in the other direction,

$$\int_a^b f \geq L(P, f) = L(P_1, f) + L(P_2, f) > U(P_1, f) + U(P_2, f) - \epsilon \geq \int_a^c f + \int_c^b f - \epsilon$$

So the difference between $\int_a^b f$ and $\int_a^c f + \int_c^b f$ lies strictly between $-\epsilon$ and ϵ , hence has absolute value at most ϵ by proposition 1.3.8(2). As $\epsilon > 0$ was arbitrary, proposition 1.4.7(2) gives $\int_a^b f = \int_a^c f + \int_c^b f$, completing the proof.

Part (3) is an equivalence in both directions, and the direction that is easy to overlook is the first: integrability on the whole interval already forces it on each half. Neither half can be badly behaved while the whole is well behaved, because the gap over $[a, b]$ computed on a partition containing c is the sum of the two gaps and both are nonnegative. Part (2) with $M = \sup_{[a, b]} |f|$ gives the crude bound $|\int_a^b f| \leq M(b - a)$, which is the estimate every convergence argument in section 5.2 and section 5.3 begins from.

Exercise 5.1.1

1. Let $c \in \mathbb{R}$ and let $f: [a, b] \rightarrow \mathbb{R}$ be the constant function $f(x) = c$. Show that $L(P, f) = U(P, f) = c(b - a)$ for every partition P of $[a, b]$, and conclude that $f \in \mathcal{R}[a, b]$ with $\int_a^b f = c(b - a)$.
2. Let $p \in [a, b]$ and let $f: [a, b] \rightarrow \mathbb{R}$ be given by $f(p) = 1$ and $f(x) = 0$ for $x \neq p$. Show that $L(P, f) = 0$ for every partition P , that $U(P, f) \leq 2\|P\|$, and hence that $f \in \mathcal{R}[a, b]$ with $\int_a^b f = 0$.
3. Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded and let P be a partition of $[a, b]$. Write ω_i for the oscillation of f on $[x_{i-1}, x_i]$ in the sense of proposition 1.3.10. Show that $U(P, f) - L(P, f) = \sum_{i=1}^n \omega_i \Delta x_i$, and deduce that $f \in \mathcal{R}[a, b]$ if and only if for every $\epsilon > 0$ there is a partition with $\sum_{i=1}^n \omega_i \Delta x_i < \epsilon$.
4. Let $f \in \mathcal{R}[a, b]$ and let $g: [a, b] \rightarrow \mathbb{R}$ agree with f at every point of $[a, b]$ outside a finite set. Show that $g \in \mathcal{R}[a, b]$ and that $\int_a^b g = \int_a^b f$. (Consider $h = g - f$ and adapt exercise 5.1.1(2).)
5. Let $f(x) = x$ on $[a, b]$. Show that $U(P, f) - L(P, f) \leq \|P\|(b - a)$ for every partition P , so that $f \in \mathcal{R}[a, b]$. Then show that for every partition P , the Riemann sum with the midpoint tags $\xi_i = (x_{i-1} + x_i)/2$ equals $(b^2 - a^2)/2$ exactly, and use theorem 5.1.11 to conclude that $\int_a^b x dx = (b^2 - a^2)/2$.
6. Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded and let $S = \{A \in \mathbb{R} \mid L(P, f) \leq A \leq U(P, f) \text{ for every partition } P \text{ of } [a, b]\}$.

Show that S is exactly the set of A with $\int_a^b f \leq A \leq \overline{\int_a^b f}$, and conclude that f is Riemann integrable if and only if S has exactly one element.

7. Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded and let $a < c < b$. Show that

$$\overline{\int_a^b f} = \overline{\int_a^c f} + \overline{\int_c^b f}$$

without assuming that f is integrable. (For one inequality use the partitions $P_1 \cup P_2$; for the other, adjoin c to an arbitrary partition of $[a, b]$.)

8. Let $f: [0, 1] \rightarrow \mathbb{R}$ be given by $f(x) = 1/n$ if $x = 1/n$ for some $n \in \mathbb{N}$, and $f(x) = 0$ otherwise. Show that $0 \leq f(x) \leq x$ for every $x \in [0, 1]$, that f agrees with the zero function outside a finite subset of $[c, 1]$ for each $c \in (0, 1)$, and hence, using exercise 5.1.1 (4) and exercise 5.1.1 (7), that $f \in \mathcal{R}[0, 1]$ with $\int_0^1 f = 0$.

5.2 Classes of Integrable Functions

Theorem 5.1.8 decides integrability by exhibiting one partition on which the two Darboux sums are close, and every existence proof below produces such a partition the same way: bound the oscillation of f on each subinterval, multiply by the length of that subinterval, and add. Uniform continuity bounds every oscillation at once; monotonicity makes the oscillations telescope. A finite set of bad points can be confined to subintervals whose total length is small, and on those the oscillation may be bounded by twice a bound on $|f|$ at negligible cost.

Boundedness alone is not enough. The Dirichlet function of example 3.2.10 has upper and lower integrals differing by the whole length of the interval, which is the computation section 3.2 promised, while Thomae's function is integrable with infinitely many discontinuities. The section closes with the operations that carry $\mathcal{R}[a, b]$ into itself: products, absolute values, pointwise extrema, and uniform limits, the last being the statement section 4.4 deferred until the integral existed.

Theorem 5.2.1: Continuous Functions Are Integrable

Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. Then f is bounded on $[a, b]$ and $f \in \mathcal{R}[a, b]$.

Proof:

Step 1: f is bounded. By theorem 3.4.4 the function f is uniformly continuous on $[a, b]$. The set $[a, b]$ is bounded above by b and below by a , so the final clause of proposition 3.4.7 makes f bounded. The Darboux sums of definition 5.1.2 are therefore defined for every partition.

Step 2: a mesh small enough to control every subinterval. Let $\epsilon > 0$ and put

$$\eta = \frac{\epsilon}{2(b-a)} > 0$$

By definition 3.4.1 there is a $\delta > 0$ with $|f(s) - f(t)| < \eta$ for all $s, t \in [a, b]$ satisfying $|s - t| < \delta$. The number $\delta/(b-a)$ is positive, so theorem 1.4.1(3) produces an $n \in \mathbb{N}$ with $1/n < \delta/(b-a)$. Let P be the uniform partition $x_i = a + i(b-a)/n$ introduced after definition 5.1.1, whose

subintervals all have length $\Delta x_i = (b - a)/n < \delta$.

Step 3: the oscillation on each subinterval. Fix i and let $s, t \in [x_{i-1}, x_i]$. Then $|s - t| \leq \Delta x_i < \delta$, so $|f(s) - f(t)| < \eta$. Thus η is an upper bound of the set $\{|f(s) - f(t)| \mid s, t \in [x_{i-1}, x_i]\}$, whose supremum is $M_i - m_i$ by proposition 1.3.10(2). A supremum is at most any upper bound by definition 1.2.2, so $M_i - m_i \leq \eta$.

Step 4: the criterion. Multiplying the bound of Step 3 by $\Delta x_i > 0$ and summing over i ,

$$U(P, f) - L(P, f) = \sum_{i=1}^n (M_i - m_i) \Delta x_i \leq \eta \sum_{i=1}^n \Delta x_i = \eta(b - a) = \frac{\epsilon}{2} < \epsilon$$

using (5.1) for the third equality. As $\epsilon > 0$ was arbitrary, theorem 5.1.8(1) gives $f \in \mathcal{R}[a, b]$, completing the proof.

The proof spends uniform continuity exactly once, at Step 3, and it is the whole reason section 3.4 was written. Continuity at each point separately would give a δ depending on the point, and no single mesh would then serve all n subintervals at once. What replaces uniform continuity in the next theorem is a hypothesis under which the n oscillations, however large individually, add up to a fixed number.

Theorem 5.2.2: Monotone Functions Are Integrable

Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be monotone in the sense of definition 3.2.14. Then f is bounded on $[a, b]$ and $f \in \mathcal{R}[a, b]$.

Proof:

Step 1: reduction to the increasing case. Suppose f is decreasing. Then $-f$ is increasing, since $x \leq y$ gives $f(x) \geq f(y)$ and hence $-f(x) \leq -f(y)$. A function is bounded exactly when its negative is, and if $-f \in \mathcal{R}[a, b]$ then theorem 5.1.13 with $c = -1$ puts $f = (-1)(-f)$ in $\mathcal{R}[a, b]$. So it is enough to treat an increasing f .

Step 2: f is bounded. Every $x \in [a, b]$ satisfies $a \leq x \leq b$, so definition 3.2.14 gives $f(a) \leq f(x) \leq f(b)$. Hence $f(a)$ is a lower bound and $f(b)$ an upper bound of $f([a, b])$, and f is bounded.

Step 3: the two sums on a uniform partition. Let $n \in \mathbb{N}$ and let P be the uniform partition $x_i = a + i(b - a)/n$, with $\Delta x_i = (b - a)/n$. Fix i . Every $x \in [x_{i-1}, x_i]$ satisfies $x_{i-1} \leq x \leq x_i$, so $f(x_{i-1}) \leq f(x) \leq f(x_i)$ by definition 3.2.14. Both bounds are themselves values of f on $[x_{i-1}, x_i]$, so an upper bound belonging to the set is its least upper bound and a lower bound belonging to it is its greatest lower bound (definition 1.2.2), giving $m_i = f(x_{i-1})$ and $M_i = f(x_i)$. Therefore

$$U(P, f) - L(P, f) = \frac{b-a}{n} \sum_{i=1}^n (f(x_i) - f(x_{i-1})) = \frac{(b-a)(f(b) - f(a))}{n}$$

because the sum telescopes, every $f(x_i)$ with $0 < i < n$ occurring once with each sign.

Step 4: the criterion. Put $K = (b - a)(f(b) - f(a))$, which is nonnegative because $b - a > 0$ and $f(a) \leq f(b)$. Let $\epsilon > 0$. The number $\epsilon/(K + 1)$ is positive, so theorem 1.4.1(3) produces an $n \in \mathbb{N}$ with $1/n < \epsilon/(K + 1)$. Multiplying by $K \geq 0$ and using $K/(K + 1) < 1$ together with $\epsilon > 0$,

$$U(P, f) - L(P, f) = \frac{K}{n} \leq \frac{K\epsilon}{K + 1} < \epsilon$$

where the strict inequality holds because $K\epsilon/(K + 1) < \epsilon$ when $K > 0$ and both sides of the display vanish when $K = 0$. So theorem 5.1.8(1) gives $f \in \mathcal{R}[a, b]$, completing the proof.

Monotonicity is the only hypothesis, and it admits functions far from continuous. Example 3.2.18 produces an increasing function on $(0, 1)$ whose set of discontinuities is exactly the set of rational points of $(0, 1)$, and theorem 3.2.17 says that a finite or countable set is the most that can occur for a monotone function. The gap $U(P, f) - L(P, f)$ is insensitive to all of this, because Step 3 evaluates it from the two endpoint values alone.

Both theorems produce integrable functions. The criterion also excludes functions, and the first exclusion is the one example 3.2.10 recorded without proof.

Example 5.2.3: The Dirichlet Function Is Not Integrable

Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be given by $f(x) = \mathbf{1}_{\mathbb{Q}}(x)$, the Dirichlet function of example 3.2.10 read on $[a, b]$. Then f is bounded,

$$\int_a^b f = 0, \quad \overline{\int_a^b} f = b - a$$

and $f \notin \mathcal{R}[a, b]$.

Boundedness. Every value of f is 0 or 1, so 0 is a lower bound and 1 an upper bound of f on $[a, b]$.

The two sums on an arbitrary partition. Let $P = \{x_0, \dots, x_n\}$ be a partition of $[a, b]$ and fix i . Since $x_{i-1} < x_i$, theorem 1.4.4 produces a rational p with $x_{i-1} < p < x_i$ and corollary 1.4.5 produces an irrational s with $x_{i-1} < s < x_i$. Both lie in $[x_{i-1}, x_i]$, and $f(p) = 1$ while $f(s) = 0$. The number 1 is an upper bound of f on $[x_{i-1}, x_i]$ and is attained there, so $M_i = 1$ by definition 1.2.2, and the same argument with 0 gives $m_i = 0$. Therefore

$$L(P, f) = \sum_{i=1}^n 0 \cdot \Delta x_i = 0, \quad U(P, f) = \sum_{i=1}^n 1 \cdot \Delta x_i = b - a$$

the second by (5.1).

The two integrals. The set of lower sums of f is $\{0\}$ and the set of upper sums is $\{b - a\}$, since P was arbitrary. A one-element set is its own supremum and its own infimum, so definition 5.1.6 gives $\int_a^b f = 0$ and $\overline{\int_a^b} f = b - a$. These differ because $b - a > 0$, so definition 5.1.7 excludes f from $\mathcal{R}[a, b]$.

No refinement improves the bracket at all: the gap is the full length of the interval for every partition, because every subinterval, however short, carries both values of f . The obstruction is that f is

discontinuous at every point of $[a, b]$. Weakening the hypothesis of theorem 5.2.1 to allow finitely many discontinuities costs nothing, and the proof needs one observation about restricting a domain.

Continuity at a point is preserved by restriction. If $h: E \rightarrow \mathbb{R}$ is continuous at c and $A \subseteq E$ contains c , then the δ that definition 3.2.1 produces for ϵ at c on E serves at c on A as well, because every $x \in A$ with $|x - c| < \delta$ lies in E .

Theorem 5.2.4: Finitely Many Discontinuities Are Harmless

Let $a < b$, let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, and suppose there is a finite set $D \subseteq [a, b]$ such that f is continuous at every point of $[a, b] \setminus D$. Then $f \in \mathcal{R}[a, b]$.

Proof:

Fix $M > 0$ with $|f(x)| \leq M$ for every $x \in [a, b]$, as produced after definition 5.1.2. For any partition of any subinterval of $[a, b]$ the numbers $-M$ and M are a lower and an upper bound of f on each block, so $-M \leq m_i \leq M_i \leq M$ and

$$M_i - m_i \leq 2M \quad (5.5)$$

Step 1: one bad left endpoint. Let $u < v$ in $[a, b]$ and suppose f is continuous at every point of $(u, v]$. The claim is that f lies in $\mathcal{R}[u, v]$. Let $\epsilon > 0$ and put

$$\theta = \min \left\{ \frac{v - u}{2}, \frac{\epsilon}{8M} \right\} > 0, \quad w = u + \theta$$

Then $u < w < v$, since $\theta \leq (v - u)/2 < v - u$. Every point of $[w, v]$ lies in $(u, v]$, so f is continuous at each of them, and by the observation above its restriction to $[w, v]$ is continuous there. By theorem 5.2.1 that restriction lies in $\mathcal{R}[w, v]$, so theorem 5.1.8(1) produces a partition P_1 of $[w, v]$ with $U(P_1, f) - L(P_1, f) < \epsilon/2$.

Let $P = \{u\} \cup P_1$, a partition of $[u, v]$ whose first subinterval is $[u, w]$ and whose remaining subintervals are exactly those of P_1 . The two sums split accordingly, so (5.5) applied to $[u, w]$ gives

$$U(P, f) - L(P, f) = \left(\sup_{[u, w]} f - \inf_{[u, w]} f \right) (w - u) + (U(P_1, f) - L(P_1, f)) \leq 2M\theta + \frac{\epsilon}{2} \leq \frac{\epsilon}{4} + \frac{\epsilon}{2} < \epsilon$$

using $\theta \leq \epsilon/(8M)$. As $\epsilon > 0$ was arbitrary, theorem 5.1.8(1) gives $f \in \mathcal{R}[u, v]$.

Step 2: one bad right endpoint. Suppose instead that f is continuous at every point of $[u, v)$. Run Step 1 with $w = v - \theta$ and with $P = P_1 \cup \{v\}$ for a partition P_1 of $[u, w]$ produced by theorem 5.2.1 and theorem 5.1.8(1). The single new subinterval is $[w, v]$, of length θ , and the same two estimates give $U(P, f) - L(P, f) < \epsilon$. So $f \in \mathcal{R}[u, v]$ in this case as well.

Step 3: two bad endpoints. Suppose f is continuous at every point of (u, v) . Put $w = (u + v)/2$, so that $u < w < v$. Every point of $(u, w]$ lies in (u, v) , so Step 1 applies on $[u, w]$, and every point of $[w, v)$ lies in (u, v) , so Step 2 applies on $[w, v]$. Both restrictions are integrable, and proposition 5.1.14(3) gives $f \in \mathcal{R}[u, v]$.

Step 4: the general case. Let $T = \{a, b\} \cup D$, a finite subset of $[a, b]$ containing a and b , and list its elements in increasing order as $a = t_0 < t_1 < \cdots < t_p = b$, which is possible because T is finite and $a < b$ forces $p \geq 1$. Fix i with $1 \leq i \leq p$. Every point of D belongs to T , and t_{i-1} and t_i are consecutive in the increasing list of T , so no element of T and hence no point of D lies in the open interval (t_{i-1}, t_i) . So f is continuous at every point of (t_{i-1}, t_i) , and Step 3 applied on $[t_{i-1}, t_i]$ gives $f \in \mathcal{R}[t_{i-1}, t_i]$.

Finally, $f \in \mathcal{R}[t_0, t_i]$ for every i , by induction on i . The case $i = 1$ is the previous paragraph. If $f \in \mathcal{R}[t_0, t_i]$ with $i < p$, then $t_0 < t_i < t_{i+1}$ and f is integrable on $[t_0, t_i]$ and on $[t_i, t_{i+1}]$, so proposition 5.1.14(3) puts f in $\mathcal{R}[t_0, t_{i+1}]$. Taking $i = p$ gives $f \in \mathcal{R}[a, b]$, completing the proof.

Boundedness is a separate hypothesis here and cannot be dropped, since definition 5.1.7 asks for it outright. A function with a single discontinuity may fail to be bounded, and $x \mapsto 1/x$ on $(0, 1]$ extended by any value at 0 is such a function: it takes the value $n+1$ at $1/(n+1)$, and theorem 1.4.1(2) produces an n exceeding any proposed bound. What the theorem does reach is every bounded function continuous on the interior of the interval, and section 5.3 uses that case when it matches an improper integral against a Riemann integral.

Theorem 5.2.5: Bounded and Continuous on an Open Interval

Let $a < b$ and let $f: (a, b) \rightarrow \mathbb{R}$ be continuous and bounded.

1. Every $g: [a, b] \rightarrow \mathbb{R}$ with $g(x) = f(x)$ for all $x \in (a, b)$ belongs to $\mathcal{R}[a, b]$.
2. Any two such functions have the same integral over $[a, b]$.

Proof:

Step 1: such a g is bounded. Let $K \geq 0$ satisfy $|f(x)| \leq K$ for every $x \in (a, b)$, which exists because f is bounded. Then $|g(x)| \leq \max\{K, |g(a)|, |g(b)|\}$ for every $x \in [a, b]$, since x is either a , or b , or a point of (a, b) where g agrees with f .

Step 2: g is continuous at every point of (a, b) . Let $x_0 \in (a, b)$ and let $\epsilon > 0$. The numbers $x_0 - a$ and $b - x_0$ are positive, so $\rho = \min\{x_0 - a, b - x_0\}$ is positive, and every $x \in \mathbb{R}$ with $|x - x_0| < \rho$ satisfies $a < x < b$ by proposition 1.3.8(2). Continuity of f at x_0 gives a $\delta_0 > 0$ with $|f(x) - f(x_0)| < \epsilon$ for every $x \in (a, b)$ satisfying $|x - x_0| < \delta_0$. Put $\delta = \min\{\rho, \delta_0\} > 0$. If $x \in [a, b]$ and $|x - x_0| < \delta$, then $x \in (a, b)$ and $|g(x) - g(x_0)| = |f(x) - f(x_0)| < \epsilon$. So δ answers ϵ in definition 3.2.1.

Step 3: part (1). By Steps 1 and 2 the function g is bounded and continuous at every point of $[a, b] \setminus \{a, b\}$, and $\{a, b\}$ is finite, so theorem 5.2.4 gives $g \in \mathcal{R}[a, b]$.

Step 4: part (2). Let g_1 and g_2 both agree with f on (a, b) , both in $\mathcal{R}[a, b]$ by part (1), and let $K_0 > 0$ bound $|g_1|$ and $|g_2|$ on $[a, b]$, obtained by taking the larger of the two constants of Step 1. Let $\epsilon > 0$ and put

$$\theta = \min \left\{ \frac{b-a}{3}, \frac{\epsilon}{8K_0} \right\} > 0, \quad c = a + \theta, \quad d = b - \theta$$

Then $a < c < d < b$, since $2\theta \leq 2(b-a)/3 < b-a$. Fix $j \in \{1, 2\}$. Applying proposition 5.1.14(3) at c splits g_j into integrable restrictions on $[a, c]$ and $[c, b]$, and applying it again at d inside $[c, b]$ splits the second of these, so all four restrictions used below are integrable and

$$\int_a^b g_j = \int_a^c g_j + \int_c^d g_j + \int_d^b g_j$$

The functions g_1 and g_2 agree at every point of $[c, d]$, because $[c, d] \subseteq (a, b)$, so the middle terms are equal.

Each of g_1 and g_2 satisfies $-K_0 \leq g_j(x) \leq K_0$ on $[a, b]$, so proposition 5.1.14(2) gives $-K_0(c-a) \leq \int_a^c g_j \leq K_0(c-a)$ and the same bound over $[d, b]$ with $b-d$ in place of $c-a$. By proposition 1.3.8(2) and then (4), the difference of the two whole integrals satisfies

$$\left| \int_a^b g_1 - \int_a^b g_2 \right| \leq 2K_0(c-a) + 2K_0(b-d) = 4K_0\theta \leq \frac{\epsilon}{2} < \epsilon$$

As $\epsilon > 0$ was arbitrary, proposition 1.4.7(2) gives $\int_a^b g_1 = \int_a^b g_2$, completing the proof.

Part (2) says that the two endpoint values contribute nothing, so the symbol $\int_a^b f$ may be written for a bounded continuous f on (a, b) with no extension named. Boundedness is what separates the theorem from a statement about continuity alone. The function $x \mapsto 1/x$ is continuous on $(0, 1)$ by theorem 3.2.5 and example 3.2.6, and no function on $[0, 1]$ agreeing with it on $(0, 1)$ is bounded, so definition 5.1.7 reaches none of them.

Of the three existence theorems so far, only theorem 5.2.2 admits infinitely many discontinuities, and it pays for them with a hypothesis constraining the function at every pair of points. Thomae's function meets no such hypothesis. Inside any interval with more than one point, theorem 1.4.4 and corollary 1.4.5 produce a rational p , an irrational $s > p$ and a rational $q > s$, and the values $t(p) > 0$, $t(s) = 0$, $t(q) > 0$ leave the function neither increasing nor decreasing there. It is nonetheless integrable.

Example 5.2.6: Thomae's Function Is Integrable

Let $f: [0, 1] \rightarrow \mathbb{R}$ be given by $f(x) = t(x)$, where t is Thomae's function of example 3.2.12. Then $f \in \mathcal{R}[0, 1]$ and $\int_0^1 f = 0$, while f is discontinuous at every rational point of $[0, 1]$.

Boundedness and the lower sums. For irrational x the value is 0, and for $x = p/q$ in lowest terms with $q \geq 1$ the value is $1/q$, which lies in $(0, 1]$ by proposition 1.1.2(7). So $0 \leq f(x) \leq 1$ on $[0, 1]$ and f is bounded. Let $P = \{x_0, \dots, x_n\}$ be any partition of $[0, 1]$ and fix i . Since $x_{i-1} < x_i$, corollary 1.4.5 produces an irrational s with $x_{i-1} < s < x_i$, and $f(s) = 0$. So 0 is a lower bound of f on $[x_{i-1}, x_i]$ and is attained there, giving $m_i = 0$ by definition 1.2.2. Hence $L(P, f) = 0$ for every partition P .

The points where f is not small. Let $\epsilon > 0$. By theorem 1.4.1(3) there is an $N \in \mathbb{N}$ with $1/N < \epsilon/2$. Put

$$F = \{x \in [0, 1] \mid f(x) \geq 1/N\}$$

A point of F is rational, since an irrational point has $f(x) = 0 < 1/N$. Writing such a point as

p/q in lowest terms with $q \geq 1$, the condition $1/q \geq 1/N$ gives $q \leq N$ by proposition 1.1.2(7), and $0 \leq p/q \leq 1$ multiplied by $q > 0$ gives $0 \leq p \leq q$. For each of the N admissible values of q there are at most $q + 1 \leq N + 1$ admissible values of p , so F has at most $N(N + 1)$ elements and is finite. It is nonempty, since $1 = 1/1$ in lowest terms and $f(1) = 1 \geq 1/N$. Write r for the number of its elements, so $r \geq 1$.

A uniform partition. The number $\epsilon/(4r)$ is positive, so theorem 1.4.1(3) produces an $n \in \mathbb{N}$ with $1/n < \epsilon/(4r)$. Let P be the uniform partition of $[0, 1]$ with $x_i = i/n$ and $\Delta x_i = 1/n$. Call an index i marked if $[x_{i-1}, x_i]$ contains a point of F . A single point of $[0, 1]$ lies in at most two of the subintervals of P , since it lies in two exactly when it is one of the interior division points and in one otherwise, so there are at most $2r$ marked indices.

The upper sum. For a marked i the bound $M_i \leq 1$ holds because 1 is an upper bound of f on $[0, 1]$. For an unmarked i every $x \in [x_{i-1}, x_i]$ lies outside F and therefore satisfies $f(x) < 1/N$, so $1/N$ is an upper bound of f there and definition 1.2.2 gives $M_i \leq 1/N$. Splitting the upper sum into the two groups and using (5.1),

$$U(P, f) = \sum_{i=1}^n M_i \Delta x_i \leq 2r \cdot \frac{1}{n} + \frac{1}{N} \sum_{i=1}^n \Delta x_i = \frac{2r}{n} + \frac{1}{N} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

the last step using $2r/n < 2r \cdot \epsilon/(4r) = \epsilon/2$ and $1/N < \epsilon/2$.

Conclusion. Since $L(P, f) = 0$, the gap $U(P, f) - L(P, f)$ is smaller than ϵ , and $\epsilon > 0$ was arbitrary, so theorem 5.1.8(1) gives $f \in \mathcal{R}[0, 1]$. By theorem 5.1.8(2) the integral satisfies $0 = L(P, f) \leq \int_0^1 f \leq U(P, f) < \epsilon$, so $|\int_0^1 f - 0| \leq \epsilon$ for every $\epsilon > 0$ and proposition 1.4.7(2) gives $\int_0^1 f = 0$. The final assertion is example 3.2.12, which shows t discontinuous at every rational and continuous at every irrational.

The set of discontinuities is infinite, so theorem 5.2.4 does not reach this function. What the argument uses is that the points where f is at least $1/N$ form a finite set for each N , so the subintervals on which the upper sum is large have total length under the partitioner's control while every other subinterval contributes at most $1/N$ times its length. The Dirichlet function fails exactly this: its values are 0 and 1 with nothing between, and the points where it equals 1 are dense.

The remaining results ask which operations carry integrable functions to integrable functions. Theorem 5.1.13 settled sums and scalar multiples, and the products, absolute values and pointwise extrema below are what section 5.3 spends on integration by parts and on substitution.

Theorem 5.2.7: Products, Absolute Values and Pointwise Extrema

Let $a < b$ and let $f, g \in \mathcal{R}[a, b]$.

1. $|f| \in \mathcal{R}[a, b]$, and

$$\left| \int_a^b f \right| \leq \int_a^b |f|$$

2. $f^2 \in \mathcal{R}[a, b]$ and $fg \in \mathcal{R}[a, b]$.

3. $\max(f, g)$ and $\min(f, g)$ belong to $\mathcal{R}[a, b]$.

Proof:

For a bounded $h: [a, b] \rightarrow \mathbb{R}$ and a partition $P = \{x_0, \dots, x_n\}$ write

$$\omega_i(h) = \sup_{[x_{i-1}, x_i]} h - \inf_{[x_{i-1}, x_i]} h$$

for the oscillation of h on the i -th subinterval, in the sense of proposition 1.3.10. By definition 5.1.2, $U(P, h) - L(P, h) = \sum_{i=1}^n \omega_i(h) \Delta x_i$, and $\omega_i(f)$ is what the display of theorem 5.1.8(1) controls.

Step 1: the absolute value. The function $|f|$ is bounded, since $|f|$ takes the same values as f up to sign. Fix a partition P and an index i , and let $s, t \in [x_{i-1}, x_i]$. By proposition 1.3.8(5) and then proposition 1.3.10(1),

$$||f(s)| - |f(t)|| \leq |f(s) - f(t)| \leq \omega_i(f)$$

So $\omega_i(f)$ is an upper bound of the set whose supremum is $\omega_i(|f|)$ by proposition 1.3.10(2), and definition 1.2.2 gives $\omega_i(|f|) \leq \omega_i(f)$. Multiplying by $\Delta x_i > 0$ and summing,

$$U(P, |f|) - L(P, |f|) \leq U(P, f) - L(P, f)$$

for every P . Given $\epsilon > 0$, the partition theorem 5.1.8(1) produces for f therefore serves $|f|$, and $|f| \in \mathcal{R}[a, b]$.

Step 2: the inequality of part (1). By proposition 1.3.8(1), $-|f(x)| \leq f(x) \leq |f(x)|$ for every x . The function $-|f|$ lies in $\mathcal{R}[a, b]$ with integral $-\int_a^b |f|$ by theorem 5.1.13, so two applications of proposition 5.1.14(1) give

$$-\int_a^b |f| \leq \int_a^b f \leq \int_a^b |f|$$

Since $|f| \geq 0$ on $[a, b]$, proposition 5.1.14(2) with $m = 0$ makes $\int_a^b |f|$ nonnegative, so proposition 1.3.8(2) applies with that number as c and yields the asserted inequality.

Step 3: squares. Let $h \in \mathcal{R}[a, b]$ and fix $M > 0$ with $|h(x)| \leq M$ on $[a, b]$, as produced after definition 5.1.2. Then h^2 is bounded by M^2 , using proposition 1.3.8(3). Fix a partition P and an index i , and let $s, t \in [x_{i-1}, x_i]$. By proposition 1.3.8(3) applied to the factorization $h(s)^2 - h(t)^2 = (h(s) - h(t))(h(s) + h(t))$, then proposition 1.3.8(4) and (1), and then proposition 1.3.10(1),

$$|h(s)^2 - h(t)^2| = |h(s) - h(t)| |h(s) + h(t)| \leq 2M |h(s) - h(t)| \leq 2M \omega_i(h)$$

As in Step 1, proposition 1.3.10(2) and definition 1.2.2 give $\omega_i(h^2) \leq 2M\omega_i(h)$, and multiplying by $\Delta x_i > 0$ and summing gives

$$U(P, h^2) - L(P, h^2) \leq 2M(U(P, h) - L(P, h))$$

Let $\epsilon > 0$. The number $\epsilon/(2M)$ is positive, so theorem 5.1.8(1) produces a partition P with $U(P, h) - L(P, h) < \epsilon/(2M)$, and then $U(P, h^2) - L(P, h^2) < \epsilon$. So $h^2 \in \mathcal{R}[a, b]$.

Step 4: products. Taking $h = f$ in Step 3 gives $f^2 \in \mathcal{R}[a, b]$. By theorem 5.1.13 the functions $f + g$ and $f - g = f + (-1)g$ lie in $\mathcal{R}[a, b]$, so Step 3 applied to each of them puts $(f + g)^2$ and $(f - g)^2$ in $\mathcal{R}[a, b]$. Expanding both squares pointwise, $(f + g)^2 - (f - g)^2 = 4fg$, so

$$fg = \frac{1}{4} \left((f + g)^2 - (f - g)^2 \right)$$

and theorem 5.1.13 puts fg in $\mathcal{R}[a, b]$.

Step 5: pointwise extrema. The identities

$$\max(f, g) = \frac{f + g + |f - g|}{2}, \quad \min(f, g) = \frac{f + g - |f - g|}{2}$$

hold pointwise, as verified in example 3.2.9. The function $f - g$ lies in $\mathcal{R}[a, b]$ by theorem 5.1.13, hence so does $|f - g|$ by Step 1, and one more application of theorem 5.1.13 to the two right-hand sides completes the proof.

Every part is a statement about membership and only part (1) computes anything. The integral of a product is not the product of the integrals, and $f = g = 1$ on $[0, 2]$ already separates them, since $\int_0^2 fg = 2$ while $\int_0^2 f \cdot \int_0^2 g = 4$. What the theorem provides is that $\int_a^b fg$ exists at all, which is what section 5.3 needs before it may write $\int_a^b Fg$ down.

The last operation is a limit. Section 4.4 showed that a pointwise limit of continuous functions may be discontinuous and that uniform convergence repairs it, and the same pattern holds for the integral.

Theorem 5.2.8: Uniform Convergence and Integration

Let $a < b$, let $f_n \in \mathcal{R}[a, b]$ for every $n \in \mathbb{N}$, and suppose $(f_n)_{n \geq 1}$ converges uniformly on $[a, b]$ to $f: [a, b] \rightarrow \mathbb{R}$. Then $f \in \mathcal{R}[a, b]$ and

$$\int_a^b f = \lim_{n \rightarrow \infty} \int_a^b f_n$$

Proof:

Step 1: f is bounded. Applying definition 4.4.2 with $\epsilon = 1$ produces an $N_0 \in \mathbb{N}$ with $|f_{N_0}(x) - f(x)| < 1$ for every $x \in [a, b]$. Boundedness is part of definition 5.1.7, so there is an M_0 with $|f_{N_0}(x)| \leq M_0$ on $[a, b]$. By proposition 1.3.8(4) and (1),

$$|f(x)| \leq |f(x) - f_{N_0}(x)| + |f_{N_0}(x)| < 1 + M_0$$

for every $x \in [a, b]$, so f is bounded and its Darboux sums are defined.

Step 2: the two gaps compare. Let $\epsilon > 0$ and put

$$\eta = \frac{\epsilon}{4(b-a)} > 0$$

By definition 4.4.2 there is an $N \in \mathbb{N}$ with $|f_n(x) - f(x)| < \eta$ for every $n \geq N$ and every $x \in [a, b]$. Fix such an n and let $P = \{x_0, \dots, x_k\}$ be any partition. By proposition 1.3.8(2), every $x \in [x_{i-1}, x_i]$ satisfies $f(x) \leq f_n(x) + \eta$, and $f_n(x)$ is at most the supremum of f_n on that subinterval, so that supremum plus η is an upper bound of f there and definition 1.2.2 gives

$$\sup_{[x_{i-1}, x_i]} f \leq \sup_{[x_{i-1}, x_i]} f_n + \eta$$

The same argument with the inequality $f(x) \geq f_n(x) - \eta$ gives $\inf_{[x_{i-1}, x_i]} f \geq \inf_{[x_{i-1}, x_i]} f_n - \eta$. Subtracting, multiplying by $\Delta x_i > 0$, summing over i and using (5.1),

$$U(P, f) - L(P, f) \leq (U(P, f_n) - L(P, f_n)) + 2\eta(b-a) = (U(P, f_n) - L(P, f_n)) + \frac{\epsilon}{2}$$

Step 3: f is integrable. Keep ϵ, η and N from Step 2 and take $n = N$. Since $f_N \in \mathcal{R}[a, b]$, theorem 5.1.8(1) produces a partition P with $U(P, f_N) - L(P, f_N) < \epsilon/2$, and Step 2 then gives $U(P, f) - L(P, f) < \epsilon$. As $\epsilon > 0$ was arbitrary, theorem 5.1.8(1) gives $f \in \mathcal{R}[a, b]$.

Step 4: the integrals converge. Let $\epsilon > 0$ and take η and N as in Step 2. Let $n \geq N$. By theorem 5.1.13 the function $f_n - f$ lies in $\mathcal{R}[a, b]$ with $\int_a^b (f_n - f) = \int_a^b f_n - \int_a^b f$, and proposition 1.3.8(2) turns the choice of N into $-\eta \leq f_n(x) - f(x) \leq \eta$ for every $x \in [a, b]$. By proposition 5.1.14(2),

$$-\eta(b-a) \leq \int_a^b f_n - \int_a^b f \leq \eta(b-a)$$

so proposition 1.3.8(2) gives $|\int_a^b f_n - \int_a^b f| \leq \eta(b-a) = \epsilon/4 < \epsilon$. This holds for every $n \geq N$, so definition 2.1.2 gives $\int_a^b f_n \rightarrow \int_a^b f$, completing the proof.

Both conclusions are needed. The first says that the class $\mathcal{R}[a, b]$ is closed under uniform limits; the second says that the two operations commute: $\int_a^b \lim_n f_n = \lim_n \int_a^b f_n$. Applied to partial sums, the theorem integrates a uniformly convergent series one term at a time.

Corollary 5.2.9: Term-by-Term Integration of a Series

Let $a < b$, let $g_n \in \mathcal{R}[a, b]$ for every $n \in \mathbb{N}$, and suppose the series $\sum_{n \geq 1} g_n$ converges uniformly on $[a, b]$ with sum g . Then $g \in \mathcal{R}[a, b]$, the series $\sum_{n \geq 1} \int_a^b g_n$ converges, and

$$\int_a^b g = \sum_{n=1}^{\infty} \int_a^b g_n$$

Proof:

Write $s_N = \sum_{n=1}^N g_n$ for the partial sums. By theorem 5.1.13 and induction on N , each s_N lies in $\mathcal{R}[a, b]$ with

$$\int_a^b s_N = \sum_{n=1}^N \int_a^b g_n \quad (5.6)$$

the case $N = 1$ being trivial and the inductive step applying theorem 5.1.13 to $s_{N+1} = s_N + g_{N+1}$.

By definition 4.4.2 the uniform convergence of the series is the uniform convergence of $(s_N)_{N \geq 1}$ to g on $[a, b]$. So theorem 5.2.8 applies to that sequence and gives $g \in \mathcal{R}[a, b]$ together with $\int_a^b s_N \rightarrow \int_a^b g$. By (5.6) the left-hand side is the N -th partial sum of $\sum_{n \geq 1} \int_a^b g_n$, so that series converges with sum $\int_a^b g$, as we sought to show.

Uniform convergence cannot be relaxed to pointwise convergence, and the integrals may then converge to the wrong number. For $n \in \mathbb{N}$ write $c_n = 1/(n+1)$, which satisfies $0 < c_n < 1$, and let $f_n: [0, 1] \rightarrow \mathbb{R}$ be $f_n(x) = n+1$ for $0 < x < c_n$ and $f_n(x) = 0$ for every other $x \in [0, 1]$.

On the open interval $(0, c_n)$ the function f_n agrees with the constant $n+1$, which is continuous and bounded there, so theorem 5.2.5 puts the restriction of f_n to $[0, c_n]$ in $\mathcal{R}[0, c_n]$ and gives it the same integral as the constant function $n+1$ on $[0, c_n]$. That constant is continuous, hence integrable by theorem 5.2.1, and proposition 5.1.14(2) with $m = M = n+1$ evaluates its integral as $(n+1)c_n = 1$. On $[c_n, 1]$ the function f_n is constantly 0, so the same two results give $\int_{c_n}^1 f_n = 0$. By proposition 5.1.14(3), $f_n \in \mathcal{R}[0, 1]$ with $\int_0^1 f_n = 1$ for every n .

The sequence converges pointwise to the constant function 0. Indeed $f_n(0) = 0$ for every n , and for $0 < x \leq 1$ the number x is positive, so theorem 1.4.1(3) produces an $N \in \mathbb{N}$ with $1/N < x$, and every $n \geq N$ has $c_n < 1/n \leq 1/N < x$ and therefore $f_n(x) = 0$. The limit function has integral 0 while every $\int_0^1 f_n$ equals 1, so the conclusion of theorem 5.2.8 fails and the convergence is not uniform. The height $n+1$ grows at exactly the rate at which the length c_n of the interval carrying it shrinks, and the product of the two is what the integral sees.

Exercise 5.2.1

1. Let $f: [0, 1] \rightarrow \mathbb{R}$ be $f(x) = 1$ for $x \in \mathbb{Q}$ and $f(x) = -1$ for $x \notin \mathbb{Q}$. Show that $|f|$ and f^2 both belong to $\mathcal{R}[0, 1]$ while f does not. (For the last part express $\mathbf{1}_{\mathbb{Q}}$ in terms of f and use theorem 5.1.13 together with example 5.2.3.) Which implications in theorem 5.2.7 does this show to be one-directional?
2. Let $f: [a, b] \rightarrow \mathbb{R}$ be increasing and let P_n be the uniform partition of $[a, b]$ into n subintervals. Show that

$$U(P_n, f) - L(P_n, f) = \frac{(b-a)(f(b) - f(a))}{n}$$

and deduce that every tagged partition (P_n, ξ) of definition 5.1.10 satisfies $|\int_a^b f - R(P_n, \xi, f)| \leq (b-a)(f(b) - f(a))/n$.

3. Let $f: [0, 1] \rightarrow \mathbb{R}$ be given by $f(0) = 0$ and $f(x) = 1/n$ for $x \in (1/(n+1), 1/n]$ and $n \in \mathbb{N}$. Show that f is increasing, that it is continuous at 0 and at 1, and that its set of discontinuities is exactly $\{1/n | n \in \mathbb{N}, n \geq 2\}$. Conclude that theorem 5.2.2 reaches f and theorem 5.2.4 does

not.

4. Let $f, g \in \mathcal{R}[a, b]$ and suppose there is a $c > 0$ with $|g(x)| \geq c$ for every $x \in [a, b]$. Show that $1/g \in \mathcal{R}[a, b]$, and deduce that $f/g \in \mathcal{R}[a, b]$. (Bound the oscillation of $1/g$ on a subinterval by c^{-2} times the oscillation of g .)
5. Let $f \in \mathcal{R}[a, b]$, let $M > 0$ satisfy $|f(x)| \leq M$ on $[a, b]$, and let $\varphi: [-M, M] \rightarrow \mathbb{R}$ be Lipschitz in the sense of definition 3.4.8. Show that $\varphi \circ f \in \mathcal{R}[a, b]$, and obtain parts (1) and (2) of theorem 5.2.7 a second time from this.
6. Let $f: [0, 1] \rightarrow \mathbb{R}$ be Thomae's function of example 5.2.6 and let $\varphi: [0, 1] \rightarrow \mathbb{R}$ be $\varphi(0) = 0$ and $\varphi(y) = 1$ for $y > 0$. Show that φ has exactly one discontinuity, that $\varphi \circ f$ is the Dirichlet function of example 5.2.3 on $[0, 1]$, and hence that a composition of two integrable functions need not be integrable. Why does exercise 5.2.1 (5) not apply?
7. Let $f_n \in \mathcal{R}[a, b]$ converge uniformly on $[a, b]$ to f . Show that the functions $F_n(x) = \int_a^x f_n$ converge uniformly on $(a, b]$ to $F(x) = \int_a^x f$.
8. Let $f \in \mathcal{R}[a, b]$ with $f(x) \geq 0$ for every $x \in [a, b]$, let $M > 0$ satisfy $f(x) \leq M$ on $[a, b]$, and let $\sqrt{f}$ be the function $x \mapsto f(x)^{1/2}$ of theorem 1.3.12. Show that $\sqrt{f} \in \mathcal{R}[a, b]$. (The square root is not Lipschitz, by example 3.4.10, so exercise 5.2.1 (5) is unavailable. Given $\epsilon > 0$, use theorem 3.4.4 to obtain a δ for ϵ on $[0, M]$, take a partition with $U(P, f) - L(P, f) < \delta^2$, and treat separately the subintervals on which the oscillation of f is below δ and those on which it is not.)

5.3 The Fundamental Theorem of Calculus

Sections 5.1 and 5.2 decide which functions have an integral, and the few values they compute are computed one at a time from the definition. That definition compares a supremum and an infimum taken over every partition of $[a, b]$ at once, and it offers no method that carries from one integrand to the next. What closes the gap is the derivative, through two statements. Integrating a function and then differentiating the result returns the function; differentiating a function and then integrating the result returns the difference of its values at the two endpoints. Both are called the fundamental theorem of calculus, and between them they turn every derivative computed in chapter 4 into an integral whose value is known.

Integration by parts and substitution are the second form applied to a function assembled from the integrand, and the identity $\log x = \int_1^x dt/t$ is the first form applied to theorem 4.6.6. Dropping the standing restriction to a bounded interval and a bounded integrand gives the improper integral, whose convergence decides the convergence of series that no test of section 2.4 reaches.

Two pieces of notation are fixed first. The symbol $\int_a^b f$ has so far been written only for $a < b$, and the statements below produce endpoints in either order, so put

$$\int_a^a f = 0, \quad \int_b^a f = -\int_a^b f \quad (a < b, f \in \mathcal{R}[a, b]) \quad (5.7)$$

Every integral in this section whose endpoints are distinct and written in increasing order is the integral of definition 5.1.7, and (5.7) extends the symbol to the two remaining cases without altering any of them.

The second piece is an inequality used at four points below. Let $u < v$, let $f \in \mathcal{R}[u, v]$, and let m and M be the infimum and the supremum of f on $[u, v]$, which exist because an integrable function is bounded. Then $m \leq f(x) \leq M$ on $[u, v]$, so proposition 5.1.14(2) applied on that interval gives

$$m(v - u) \leq \int_u^v f \leq M(v - u) \quad (5.8)$$

This is the whole of what a single partition says, and it is enough to prove both forms of the fundamental theorem.

The third piece is needed wherever a function differentiated on a large domain is integrated on a small one.

Proposition 5.3.1: Differentiability Passes to a Restriction

Let $I \subseteq \mathbb{R}$ be an interval, let $g: I \rightarrow \mathbb{R}$ be differentiable at c , and let $J \subseteq I$ be an interval with more than one point containing c . Then $g|_J$ is differentiable at c and $(g|_J)'(c) = g'(c)$.

Proof:

The difference quotient of $g|_J$ at c is the restriction to $J \setminus \{c\}$ of the difference quotient of g at c , and c is a limit point of $J \setminus \{c\}$ by lemma 4.1.1. So proposition 3.1.23(1) carries the limit over, and the two derivatives are the same number.

This is not automatic: definition 4.1.2 takes the limit over the stated domain, so a function differentiable on $\mathbb{R}$ is not by definition differentiable on $[0, 1]$ until the restriction is taken.

The first form takes an integrable f and integrates it over a moving upper limit. The resulting function of that limit is better behaved than f : it is continuous whatever f does, and at each point where f is continuous it is differentiable with derivative f . Integration improves regularity by one degree, which reverses the effect of differentiation.

Theorem 5.3.2: The Fundamental Theorem of Calculus, First Form

Let $a < b$, let $f \in \mathcal{R}[a, b]$, and define $F: [a, b] \rightarrow \mathbb{R}$ by

$$F(x) = \int_a^x f$$

with $F(a) = 0$ as in (5.7).

1. F is Lipschitz on $[a, b]$, and hence continuous.
2. If f is continuous at $x_0 \in [a, b]$, then F is differentiable at x_0 and $F'(x_0) = f(x_0)$.
3. If f is continuous on $[a, b]$, then F is differentiable at every point of $[a, b]$ and $F' = f$.

Proof:

Step 1: an increment of F is an integral. Let $a \leq u < v \leq b$. If $u = a$, then $F(v) - F(u) = F(v) = \int_a^v f$ by the definition of F together with $F(a) = 0$. If $a < u$, then proposition 5.1.14 makes f integrable on $[a, u]$ and on $[u, v]$ and gives $\int_a^v f = \int_a^u f + \int_u^v f$, which rearranges to the

same conclusion. So

$$F(v) - F(u) = \int_u^v f \quad (a \leq u < v \leq b) \quad (5.9)$$

Step 2: part (1). The function f is bounded, so the set $\{|f(t)| \mid t \in [a, b]\}$ is nonempty and bounded above and has a supremum $K \geq 0$ by theorem 1.2.4. Fix $a \leq u < v \leq b$ and let m and M be the infimum and the supremum of f on $[u, v]$. Every $t \in [u, v]$ satisfies $-K \leq -|f(t)| \leq f(t) \leq |f(t)| \leq K$ by proposition 1.3.8(1), so $-K$ is a lower bound and K an upper bound for f on $[u, v]$, giving $-K \leq m$ and $M \leq K$. Multiplying by $v - u > 0$ preserves both inequalities by proposition 1.1.2(5), so combining them with (5.8) and (5.9) gives

$$-K(v - u) \leq F(v) - F(u) \leq K(v - u)$$

By proposition 1.3.8(2) this reads $|F(v) - F(u)| \leq K|v - u|$. The inequality is symmetric in u and v and holds trivially when $u = v$, so it holds for all $u, v \in [a, b]$, which is definition 3.4.8 with Lipschitz constant K . Then F is uniformly continuous by proposition 3.4.9 and continuous by box 3.4.2.

Step 3: part (2). Let $\epsilon > 0$. Continuity of f at x_0 gives a $\delta > 0$ with

$$|f(t) - f(x_0)| < \frac{\epsilon}{2} \quad \text{for every } t \in [a, b] \text{ with } |t - x_0| < \delta$$

Let $x \in [a, b]$ with $0 < |x - x_0| < \delta$, and write $u = \min(x, x_0)$ and $v = \max(x, x_0)$, so that $u < v$ and $v - u = |x - x_0|$. Every $t \in [u, v]$ lies between x and x_0 , so $|t - x_0| \leq |x - x_0| < \delta$, and therefore $f(x_0) - \epsilon/2 \leq f(t) \leq f(x_0) + \epsilon/2$ by proposition 1.3.8(2). The infimum m and the supremum M of f on $[u, v]$ then satisfy $f(x_0) - \epsilon/2 \leq m$ and $M \leq f(x_0) + \epsilon/2$, and feeding these into (5.8) and using (5.9),

$$\left(f(x_0) - \frac{\epsilon}{2}\right)(v - u) \leq F(v) - F(u) \leq \left(f(x_0) + \frac{\epsilon}{2}\right)(v - u)$$

which by proposition 1.3.8(2) is

$$|F(v) - F(u) - f(x_0)(v - u)| \leq \frac{\epsilon}{2}(v - u)$$

The quantity $F(x) - F(x_0) - f(x_0)(x - x_0)$ equals $F(v) - F(u) - f(x_0)(v - u)$ when $x > x_0$ and equals its negative when $x < x_0$, so the two have the same absolute value by proposition 1.3.8(1). Dividing by $|x - x_0| = v - u > 0$ and using proposition 1.3.8(3),

$$\left| \frac{F(x) - F(x_0)}{x - x_0} - f(x_0) \right| \leq \frac{\epsilon}{2} < \epsilon$$

The left-hand side is $|q_{F, x_0}(x) - f(x_0)|$ for the difference quotient of definition 4.1.2, whose domain $[a, b] \setminus \{x_0\}$ has x_0 as a limit point by lemma 4.1.1. So δ answers ϵ in definition 3.1.15, and $F'(x_0) = f(x_0)$.

Step 4: part (3). If f is continuous on $[a, b]$, then it is continuous at every $x_0 \in [a, b]$ by definition 3.2.2, and part (2) applies at each of them, completing the proof.

Part (1) asks least of f : it may be as discontinuous as integrability allows, and F is still Lipschitz with the bound on $|f|$ as its constant. Part (2) uses continuity at the single point x_0 and nothing more, and continuity somewhere is genuinely needed: exercise 5.3.1 (1) exhibits an integrable f with one jump whose F fails to be differentiable exactly there.

The second form runs the other way. It starts from a function known to be a derivative and evaluates its integral from two values of the antiderivative. The derivative F' is assumed integrable and is not assumed continuous, which matters because a derivative need not be continuous: example 4.6.17 produces one that is not. What theorem 4.2.10 adds is that such a derivative still has the intermediate value property, so the discontinuity is never a simple one.

Theorem 5.3.3: The Fundamental Theorem of Calculus, Second Form

Let $a < b$, let $F: [a, b] \rightarrow \mathbb{R}$ be differentiable at every point of $[a, b]$, and suppose $f = F'$ belongs to $\mathcal{R}[a, b]$. Then

$$\int_a^b f = F(b) - F(a)$$

Proof:

Let $\epsilon > 0$. Since $f \in \mathcal{R}[a, b]$, theorem 5.1.8 produces a partition $P = \{a = x_0 < x_1 < \cdots < x_n = b\}$ with

$$U(P, f) - L(P, f) < \epsilon \quad (5.10)$$

Step 1: the mean value theorem on each block. Fix i with $1 \leq i \leq n$. The function F is differentiable at every point of $[a, b]$, hence continuous at every point of $[a, b]$ by proposition 4.1.5, and so continuous on $[a, b]$ in the sense of definition 3.2.2. The points $x_{i-1} < x_i$ lie in the interval $[a, b]$, and every point of (x_{i-1}, x_i) lies in $[a, b]$ as well, so F is differentiable there. Theorem 4.2.4, applied to F on the interval $[a, b]$ with the two points x_{i-1} and x_i , gives a $t_i \in (x_{i-1}, x_i)$ with

$$F(x_i) - F(x_{i-1}) = F'(t_i)(x_i - x_{i-1}) = f(t_i) \Delta x_i$$

Step 2: the sum telescopes. Summing the previous display over $i = 1, \dots, n$, every value $F(x_i)$ with $1 \leq i \leq n - 1$ occurs once with each sign, so

$$\sum_{i=1}^n f(t_i) \Delta x_i = F(x_n) - F(x_0) = F(b) - F(a) \quad (5.11)$$

Step 3: both numbers lie between the two Darboux sums. For each i the point t_i lies in $[x_{i-1}, x_i]$, so $m_i \leq f(t_i) \leq M_i$, these being the infimum and the supremum of f on that block. Multiplying by $\Delta x_i > 0$ using proposition 1.1.2(5) and summing over i ,

$$L(P, f) \leq \sum_{i=1}^n f(t_i) \Delta x_i \leq U(P, f)$$

which by (5.11) places $F(b) - F(a)$ between $L(P, f)$ and $U(P, f)$. The number $\int_a^b f$ lies between the same two bounds, by definition 5.1.6 and definition 5.1.7 as in (5.8). Two numbers lying in

one interval differ in absolute value by at most its length, so (5.10) gives

$$\left| F(b) - F(a) - \int_a^b f \right| \leq U(P, f) - L(P, f) < \epsilon$$

Step 4: conclusion. The last display holds for every $\epsilon > 0$, with the partition depending on ϵ and the two numbers $F(b) - F(a)$ and $\int_a^b f$ fixed throughout. So proposition 1.4.7(2) gives $F(b) - F(a) = \int_a^b f$, as we sought to show.

Every antiderivative computed in chapter 4 now evaluates an integral. Theorem 4.1.7(5) and (2) make $x \mapsto x^3/3$ differentiable on $\mathbb{R}$ with derivative x^2 , hence on $[0, 1]$ by proposition 5.3.1, which is continuous and therefore integrable by theorem 5.2.1, so $\int_0^1 x^2 dx = 1/3 - 0 = 1/3$. Theorem 4.6.1 makes $\exp$ its own derivative, so $\int_0^1 \exp = \exp(1) - \exp(0) = e - 1$. Only the two endpoint values of F survive, and any two antiderivatives of f differ by a constant by corollary 4.2.8(2), so the answer does not depend on which antiderivative is used.

Each form needs what the other produces. Theorem 5.3.3 needs an antiderivative handed to it and produces nothing when none is known; theorem 5.3.2 manufactures one for every continuous f and says nothing about how to write it down. Used together they give the two rules for rewriting an integral. The first rewrites the integral of a product.

Theorem 5.3.4: Integration by Parts

Let $a < b$ and let $F, G: [a, b] \rightarrow \mathbb{R}$ be differentiable at every point of $[a, b]$, with $f = F'$ and $g = G'$ both in $\mathcal{R}[a, b]$. Then Fg and fG belong to $\mathcal{R}[a, b]$ and

$$\int_a^b Fg = F(b)G(b) - F(a)G(a) - \int_a^b fG$$

Proof:

Step 1: the products are integrable. By proposition 4.1.5 both F and G are continuous at every point of $[a, b]$, hence continuous on $[a, b]$, and theorem 5.2.1 puts both in $\mathcal{R}[a, b]$. Since f and g lie in $\mathcal{R}[a, b]$ by hypothesis, theorem 5.2.7 puts the products Fg and fG in $\mathcal{R}[a, b]$, and theorem 5.1.13 puts their sum there as well.

Step 2: the product rule gives an antiderivative. Put $H = FG$ on $[a, b]$. The interval $[a, b]$ has more than one point, and F and G are differentiable at each of its points, so theorem 4.1.7(3) makes H differentiable at every point of $[a, b]$ with

$$H' = F'G + FG' = fG + Fg$$

which lies in $\mathcal{R}[a, b]$ by Step 1.

Step 3: the second form of the fundamental theorem. The hypotheses of theorem 5.3.3 hold for H and H' , so

$$\int_a^b (fG + Fg) = H(b) - H(a) = F(b)G(b) - F(a)G(a)$$

By theorem 5.1.13 the left-hand side is $\int_a^b fG + \int_a^b Fg$, and subtracting $\int_a^b fG$ from both sides completes the proof.

The rule converts an integral into another integral together with two evaluations, and it is useful whenever the second integral is easier than the first. Take $F(t) = \log t$ and $G(t) = t$ on $[1, X]$ with $X > 1$. The derivatives are $f(t) = 1/t$ by theorem 4.6.6 and $g(t) = 1$ by theorem 4.1.7(5), both continuous on $[1, X]$ and therefore integrable by theorem 5.2.1. The product fG is the constant function 1, whose integral over $[1, X]$ is $X - 1$ by (5.8) with $m = M = 1$. So

$$\int_1^X \log t \, dt = X \log X - 1 \cdot \log 1 - (X - 1) = X \log X - X + 1$$

using $\log 1 = 0$ from definition 4.6.5. The rule has produced an antiderivative of the logarithm, which none of the differentiation rules of section 4.6 produces directly.

The second rule rewrites the variable of integration. Its statement is where (5.7) is needed, since a substitution may reverse the order of the endpoints.

Theorem 5.3.5: Substitution

Let $c < d$, let $\varphi: [c, d] \rightarrow \mathbb{R}$ be differentiable at every point of $[c, d]$ with $\varphi' \in \mathcal{R}[c, d]$, let $p < q$, and let $f: [p, q] \rightarrow \mathbb{R}$ be continuous with $\varphi([c, d]) \subseteq [p, q]$. Then $(f \circ \varphi)' \in \mathcal{R}[c, d]$ and

$$\int_c^d f(\varphi(t)) \varphi'(t) \, dt = \int_{\varphi(c)}^{\varphi(d)} f(x) \, dx$$

the right-hand side being read through (5.7) when $\varphi(d) \leq \varphi(c)$.

Proof:

Step 1: an antiderivative for f . The function f is continuous on $[p, q]$, so theorem 5.2.1 gives $f \in \mathcal{R}[p, q]$, and theorem 5.3.2(3) makes $F(y) = \int_p^y f$ differentiable at every $y \in [p, q]$ with $F'(y) = f(y)$.

Step 2: the chain rule. The sets $[c, d]$ and $[p, q]$ are intervals with more than one point, $\varphi([c, d]) \subseteq [p, q]$ by hypothesis, φ is differentiable at each $t \in [c, d]$, and F is differentiable at $\varphi(t) \in [p, q]$ by Step 1. So theorem 4.1.8 makes $F \circ \varphi$ differentiable at every $t \in [c, d]$, with

$$(F \circ \varphi)'(t) = F'(\varphi(t)) \varphi'(t) = f(\varphi(t)) \varphi'(t)$$

Step 3: the integrand is integrable. The map φ is continuous on $[c, d]$ by proposition 4.1.5, and f is continuous on $[p, q]$, so $f \circ \varphi$ is continuous on $[c, d]$ by theorem 3.2.7 and lies in $\mathcal{R}[c, d]$ by theorem 5.2.1. With $\varphi' \in \mathcal{R}[c, d]$ by hypothesis, theorem 5.2.7 puts the product $(f \circ \varphi) \varphi'$ in $\mathcal{R}[c, d]$.

Step 4: evaluation. Steps 2 and 3 are the hypotheses of theorem 5.3.3 for the function $F \circ \varphi$ on $[c, d]$, so

$$\int_c^d f(\varphi(t)) \varphi'(t) \, dt = F(\varphi(d)) - F(\varphi(c))$$

Write $\alpha = \varphi(c)$ and $\beta = \varphi(d)$, both in $[p, q]$. Suppose $\alpha < \beta$. If $p < \alpha$, then proposition 5.1.14 gives $\int_p^\beta f = \int_p^\alpha f + \int_\alpha^\beta f$, and if $p = \alpha$ then $F(\alpha) = 0$ by (5.7). Either way $F(\beta) - F(\alpha) = \int_\alpha^\beta f$. If $\alpha = \beta$, both sides are 0 by (5.7). If $\beta < \alpha$, the first case with the roles of α and β exchanged gives $F(\alpha) - F(\beta) = \int_\beta^\alpha f$, so $F(\beta) - F(\alpha) = -\int_\beta^\alpha f = \int_\alpha^\beta f$ by (5.7), completing the proof.

The hypothesis on φ says nothing about monotonicity, so the endpoints $\varphi(c)$ and $\varphi(d)$ may fall in either order, which is the case (5.7) was fixed for. For a computation, take $\varphi(t) = t^2$ on $[0, 1]$, whose derivative $2t$ is continuous and therefore integrable, and $f = \exp$ on $[0, 1]$, a closed bounded interval containing $\varphi([0, 1]) = [0, 1]$. The theorem gives

$$\int_0^1 2t \exp(t^2) dt = \int_0^1 \exp(x) dx = e - 1$$

the last value being the computation after theorem 5.3.3. Read from left to right, the rule recognizes an integrand as a composite times the derivative of the inner function. Read from right to left, it replaces a hard integrand by an easier one.

The identity recovering the logarithm as an integral is the first form applied to $1/t$. The logarithm itself was defined in definition 4.6.5 as the inverse of the exponential, so what follows is a theorem about that function.

Example 5.3.6: The Logarithm as an Integral

For every $x > 0$,

$$\log x = \int_1^x \frac{dt}{t}$$

where the right-hand side is read through (5.7) when $x < 1$.

The restriction of the logarithm. For an interval $J \subseteq (0, \infty)$ with more than one point, proposition 5.3.1 carries the derivative $\log'(c) = 1/c$ of theorem 4.6.6 over to $\log|_J$ at every $c \in J$.

The case $x > 1$. Put $J = [1, x]$ and let $h: J \rightarrow \mathbb{R}$ be $h(t) = 1/t$, which is continuous on J by example 3.2.6 since $t \neq 0$ there, and hence integrable by theorem 5.2.1. By theorem 5.3.2(3) the function $G(y) = \int_1^y h$ is differentiable at every $y \in J$ with $G'(y) = 1/y$. Put $H = G - \log|_J$ on J . By theorem 4.1.7(1) and (2) it is differentiable at every $y \in J$ with

$$H'(y) = \frac{1}{y} - \frac{1}{y} = 0$$

so H is constant by corollary 4.2.8(1). Its value at $y = 1$ is $G(1) - \log 1 = 0 - 0 = 0$, using $G(1) = 0$ from (5.7) and $\log 1 = 0$ from definition 4.6.5. So $H(x) = 0$, which is $\int_1^x dt/t = \log x$.

The case $x < 1$. Put $J = [x, 1]$ and run the same argument with $G(y) = \int_x^y h$, again differentiable with $G'(y) = 1/y$ by theorem 5.3.2(3). The function $H = G - \log|_J$ has $H' = 0$ and is therefore constant, and its value at $y = x$ is $G(x) - \log x = -\log x$. Taking $y = 1$ gives $G(1) - \log 1 = -\log x$, that is $\int_x^1 dt/t = -\log x$, and (5.7) turns this into $\int_1^x dt/t = \log x$.

The case $x = 1$. Both sides are 0.

The identity gives an antiderivative for $1/t$, which is what makes example 5.3.10 computable, and exercise 5.3.1 (4) uses it to evaluate an integral that theorem 5.3.5 reduces to $\int_1^2 dx/x$.

Everything so far requires a bounded function on a closed bounded interval, because definition 5.1.7 requires both. Two situations fall outside: an interval that is not bounded, and a function unbounded

near an endpoint of a bounded interval. Both are handled by integrating over a subinterval where the definition applies and then taking a limit.

Definition 5.3.7: Improper Integral

1. Let $a \in \mathbb{R}$ and let $f: [a, \infty) \rightarrow \mathbb{R}$ satisfy $f \in \mathcal{R}[a, X]$ for every $X > a$. The *improper integral* $\int_a^\infty f$ *converges* if the limit

$$\lim_{X \rightarrow \infty} \int_a^X f$$

exists in $\mathbb{R}$ in the sense of definition 3.1.27, applied to the function $X \mapsto \int_a^X f$ on (a, ∞) , and in that case $\int_a^\infty f$ denotes that limit. Otherwise the improper integral *diverges*.

2. Let $a < b$ and let $f: (a, b] \rightarrow \mathbb{R}$ satisfy $f \in \mathcal{R}[t, b]$ for every $t \in (a, b)$. The improper integral $\int_a^b f$ *converges* if the one-sided limit

$$\lim_{t \rightarrow a^+} \int_t^b f$$

exists in $\mathbb{R}$ in the sense of definition 3.1.24, applied to the function $t \mapsto \int_t^b f$ on $(a, b]$, and in that case $\int_a^b f$ denotes that limit. Otherwise the improper integral *diverges*.

The integrals $\int_{-\infty}^b f$ for $f: (-\infty, b] \rightarrow \mathbb{R}$ and $\int_a^b f$ for $f: [a, b) \rightarrow \mathbb{R}$ are defined by the limits $\lim_{X \rightarrow -\infty} \int_X^b f$ and $\lim_{t \rightarrow b^-} \int_a^t f$.

Form (2) writes $\int_a^b f$ for a symbol that definition 5.1.7 may already have defined, and the two agree whenever both apply. Suppose $f \in \mathcal{R}[a, b]$, and write K for the supremum of $|f|$ on $[a, b]$. For $t \in (a, b)$ the additivity of proposition 5.1.14 gives $\int_t^b f = \int_a^b f - \int_a^t f$, and (5.8) together with proposition 1.3.8(2) bounds $|\int_a^t f|$ by $K(t - a)$, as in Step 2 of the proof of theorem 5.3.2. Given $\epsilon > 0$, the choice $\delta = \epsilon/(K + 1)$ makes $|\int_t^b f - \int_a^b f| \leq K(t - a) < \epsilon$ for every $t \in (a, a + \delta)$, so the improper integral converges to the Riemann integral and the notation is unambiguous.

For a computation, take $f(t) = t^{-2}$ on $[1, \infty)$. It is continuous on each $[1, X]$ by example 3.2.6, and theorem 4.6.10(1) makes $t \mapsto -t^{-1}$ differentiable on $[1, X]$ with derivative t^{-2} , so theorem 5.3.3 gives $\int_1^X t^{-2} dt = 1 - 1/X$. Given $\epsilon > 0$, theorem 1.4.1(3) produces an $n \in \mathbb{N}$ with $1/n < \epsilon$, and every $X > n$ has $0 < 1/X < 1/n < \epsilon$ by proposition 1.1.2(7). So $\lim_{X \rightarrow \infty} (1 - 1/X) = 1$ and $\int_1^\infty t^{-2} dt = 1$.

Deciding convergence by evaluating the limit requires an antiderivative, and for most integrands none is available. The tests below decide convergence without one. They are the integral forms of proposition 2.4.4 and theorems 2.4.12 and 2.4.13, and they hold for the same reason: a nonnegative integrand makes $X \mapsto \int_a^X f$ increasing, and an increasing function has a limit exactly when it is bounded.

Theorem 5.3.8: Convergence Tests for Improper Integrals

Let $a \in \mathbb{R}$ and let $f, g: [a, \infty) \rightarrow \mathbb{R}$ satisfy $f, g \in \mathcal{R}[a, X]$ for every $X > a$.

1. Suppose $f(t) \geq 0$ for every $t \geq a$. Then $\int_a^\infty f$ converges if and only if the set $S = \{\int_a^X f \mid X > a\}$ is bounded above, and in that case $\int_a^\infty f = \sup S$.
2. Suppose $0 \leq f(t) \leq g(t)$ for every $t \geq a$. If $\int_a^\infty g$ converges, then $\int_a^\infty f$ converges and $\int_a^\infty f \leq \int_a^\infty g$. If $\int_a^\infty f$ diverges, then $\int_a^\infty g$ diverges.
3. If $\int_a^\infty |f|$ converges, then $\int_a^\infty f$ converges.
4. Let $b > a$. All three statements hold for the improper integrals of definition 5.3.7(2) of functions $f, g: (a, b] \rightarrow \mathbb{R}$ with $f, g \in \mathcal{R}[t, b]$ for every $t \in (a, b)$, with the set $\{\int_t^b f \mid t \in (a, b)\}$ in place of S .

Proof:

Write $\Phi(X) = \int_a^X f$ for $X > a$.

Step 1: Φ increases under the hypothesis of part (1). Let $a < X < Y$. By proposition 5.1.14 the function f is integrable on $[X, Y]$ and $\Phi(Y) - \Phi(X) = \int_X^Y f$. The infimum of f on $[X, Y]$ is at least 0, so the left-hand inequality of (5.8) gives $\int_X^Y f \geq 0$ and hence $\Phi(X) \leq \Phi(Y)$.

Step 2: part (1). Suppose first that S is bounded above. The set (a, ∞) is unbounded above and Φ is increasing on it, so proposition 3.1.29(2) gives $\lim_{X \rightarrow \infty} \Phi(X) = \sup S$, a real number, which is convergence with the stated value.

Suppose instead that $\int_a^\infty f$ converges, with value L . Taking $\epsilon = 1$ in definition 3.1.27 produces an $M \in \mathbb{R}$ with $\Phi(X) < L + 1$ for every $X > M$, by proposition 1.3.8(2). Put $X_0 = \max(M, a) + 1$, so that $X_0 > M$ and $X_0 > a$. Every $X \in (a, X_0]$ has $\Phi(X) \leq \Phi(X_0) < L + 1$ by Step 1, and every $X > X_0$ has $\Phi(X) < L + 1$ directly. So $L + 1$ is an upper bound for S .

Step 3: part (2). Both f and g are nonnegative, so part (1) applies to each. Let $X > a$. By proposition 5.1.14 the inequality $f \leq g$ on $[a, X]$ gives $\int_a^X f \leq \int_a^X g$. If $\int_a^\infty g$ converges, then part (1) applied to g makes $\int_a^\infty g$ an upper bound for every $\int_a^X g$, hence for every $\int_a^X f$, so S is bounded above by $\int_a^\infty g$. Part (1) applied to f now gives convergence of $\int_a^\infty f$ with value $\sup S \leq \int_a^\infty g$. The second assertion is the contrapositive of the first.

Step 4: part (3). Put

$$f^+ = \frac{|f| + f}{2}, \quad f^- = \frac{|f| - f}{2}$$

on $[a, \infty)$. By proposition 1.3.8(1), $-|f(t)| \leq f(t) \leq |f(t)|$, so $0 \leq f^+(t) \leq |f(t)|$ and $0 \leq f^-(t) \leq |f(t)|$ for every $t \geq a$, and $f = f^+ - f^-$ by subtraction. For each $X > a$, theorem 5.2.7 puts $|f|$ in $\mathcal{R}[a, X]$ and theorem 5.1.13 puts f^+ and f^- there. Part (2), applied to the pair $(f^+, |f|)$ and to the pair $(f^-, |f|)$, makes $\int_a^\infty f^+$ and $\int_a^\infty f^-$ convergent. By theorem 5.1.13,

$\int_a^X f = \int_a^X f^+ - \int_a^X f^-$ for every $X > a$, and proposition 3.1.29(1) gives

$$\lim_{X \rightarrow \infty} \int_a^X f = \int_a^\infty f^+ - \int_a^\infty f^-$$

a real number, so $\int_a^\infty f$ converges.

Step 5: the analogue of part (1) at an endpoint. Write $\Psi(t) = \int_t^b f$ for $t \in (a, b)$ and $\Psi(b) = 0$. For $a < t < t' \leq b$, proposition 5.1.14 gives $\Psi(t) - \Psi(t') = \int_t^{t'} f$, so under the hypothesis $f \geq 0$ the left-hand inequality of (5.8) makes that difference nonnegative and Ψ decreasing on $(a, b]$. If Ψ is bounded above, proposition 3.2.16 gives $\lim_{t \rightarrow a^+} \Psi(t) = \sup\{\Psi(t) \mid a < t \leq b\}$, a real number, which is convergence with that value. If instead the improper integral converges with value L , then $\epsilon = 1$ in definition 3.1.24 produces a $\delta > 0$ with $\Psi(t) < L + 1$ for every $t \in (a, a + \delta)$. Fix one such t_1 . Every $t \in [t_1, b]$ has $\Psi(t) \leq \Psi(t_1) < L + 1$ because Ψ decreases, so $L + 1$ bounds Ψ above.

Step 6: the analogues of parts (2) and (3) at an endpoint. Suppose $0 \leq f \leq g$ on $(a, b]$ with both integrable on each $[t, b]$, and suppose $\int_a^b g$ converges. For $t \in (a, b)$, proposition 5.1.14 gives $\int_t^b f \leq \int_t^b g$, and Step 5 applied to g makes $\int_a^b g$ an upper bound for the latter. So the set of numbers $\int_t^b f$ is bounded above by $\int_a^b g$, and Step 5 applied to f gives convergence with $\int_a^b f \leq \int_a^b g$. For the third statement, define f^+ and f^- on $(a, b]$ by the formulas of Step 4. The bounds and the integrability established there hold on each $[t, b]$ verbatim, and the endpoint analogue of part (2) just proved makes $\int_a^b f^+$ and $\int_a^b f^-$ convergent. By theorem 5.1.13, $\int_t^b f = \int_t^b f^+ - \int_t^b f^-$ for every $t \in (a, b)$. A one-sided limit is the limit of a restriction by proposition 3.1.25, so corollary 3.1.20 applied to those restrictions makes $\lim_{t \rightarrow a^+} \int_t^b f$ equal to $\int_a^b f^+ - \int_a^b f^-$, a real number, completing the proof.

Part (3) is the integral form of theorem 2.4.12, and it is what exercise 5.3.1 (8) runs on. The next theorem turns the parallel with series into a theorem of its own: for a decreasing nonnegative integrand, the integral and the series of its values at the integers converge together. That decides series whose terms sit too close to $1/n$ for the tests of section 2.4 to separate them.

Theorem 5.3.9: The Integral Test

Let $N_0 \in \mathbb{N}$ and let $f: [N_0, \infty) \rightarrow \mathbb{R}$ be nonnegative and decreasing. Then $f \in \mathcal{R}[N_0, X]$ for every $X > N_0$, and the series $\sum_{n \geq N_0} f(n)$ converges if and only if the improper integral $\int_{N_0}^\infty f$ converges. When they converge,

$$\int_{N_0}^\infty f \leq \sum_{n=N_0}^\infty f(n) \leq f(N_0) + \int_{N_0}^\infty f$$

Proof:

Step 1: integrability and the block estimate. Fix $X > N_0$. On $[N_0, X]$ the function f is decreasing, so $0 \leq f(t) \leq f(N_0)$ for every t there and f is bounded, and theorem 5.2.2 gives $f \in \mathcal{R}[N_0, X]$. Now let $n \geq N_0$ be an integer. On $[n, n+1]$ the decrease of f gives $f(n+1) \leq f(t) \leq f(n)$ for every t , and both bounds are values of f there, so the infimum is $f(n+1)$ and the supremum is $f(n)$. With $(n+1) - n = 1$, (5.8) reads

$$f(n+1) \leq \int_n^{n+1} f \leq f(n) \quad (5.12)$$

Step 2: summing the blocks. Let $N > N_0$ be an integer. Repeated use of the additivity in proposition 5.1.14, by induction on N , gives $\int_{N_0}^N f = \sum_{n=N_0}^{N-1} \int_n^{n+1} f$. Summing (5.12) over $n = N_0, \dots, N-1$ therefore gives

$$\sum_{n=N_0+1}^N f(n) \leq \int_{N_0}^N f \leq \sum_{n=N_0}^{N-1} f(n) \quad (5.13)$$

Step 3: the integral converges when the series does. Suppose $\sum_{n \geq N_0} f(n)$ converges, with sum S . Its terms are nonnegative, so proposition 2.4.4 identifies S as the supremum of the partial sums, and the right-hand half of (5.13) gives $\int_{N_0}^N f \leq S$ for every integer $N > N_0$. Let $X > N_0$ be real. By theorem 1.4.1(2) there is an $N \in \mathbb{N}$ with $N > X$, and such an N satisfies $N > N_0$. For $N_0 < X < N$, proposition 5.1.14 gives $\int_{N_0}^N f - \int_{N_0}^X f = \int_X^N f$, which is nonnegative by the left-hand inequality of (5.8), so $\int_{N_0}^X f \leq \int_{N_0}^N f \leq S$. Thus S bounds the set of theorem 5.3.8(1) above, and that part gives convergence of $\int_{N_0}^\infty f$ together with $\int_{N_0}^\infty f \leq S$.

Step 4: the series converges when the integral does. Suppose $\int_{N_0}^\infty f$ converges, with value I . By theorem 5.3.8(1) the number I is the supremum of the numbers $\int_{N_0}^X f$, so $\int_{N_0}^N f \leq I$ for every integer $N > N_0$, and the left-hand half of (5.13) bounds the partial sums of $\sum_{n \geq N_0+1} f(n)$ by I . Those terms are nonnegative, so proposition 2.4.4 makes $\sum_{n \geq N_0+1} f(n)$ convergent with sum at most I . Adding the single term $f(N_0)$ leaves convergence unaffected and gives

$$\sum_{n=N_0}^{\infty} f(n) \leq f(N_0) + I$$

Step 5: the two inequalities. When both converge, Step 3 gives $\int_{N_0}^\infty f \leq \sum_{n \geq N_0} f(n)$ and Step 4 gives $\sum_{n \geq N_0} f(n) \leq f(N_0) + \int_{N_0}^\infty f$, which is the displayed pair, completing the proof.

The two inequalities are as useful as the equivalence. They bound the sum of a convergent series by an integral with an error of at most the first term, which is how such a sum is estimated numerically. For a divergent series they say nothing, but the block estimate (5.12) inside the proof holds for every integer $n \geq N_0$ whether or not either side converges: exercise 5.3.1 (7) sums it for $f(t) = 1/t$ to reach the bounds $\log(N+1) \leq \sum_{n=1}^N 1/n \leq 1 + \log N$ on the harmonic partial sums. The hypothesis that f decreases is what makes (5.12) available, and it cannot be dropped: without it the values $f(n)$ say

nothing about f between consecutive integers.

The series below is the one the test was stated for. Its terms sit between $1/n$ and $1/n^{1+\epsilon}$ for every $\epsilon > 0$, so theorem 2.4.13 against example 2.4.10 decides nothing, and both theorem 2.4.14 and theorem 2.4.15 return the inconclusive value 1.

Example 5.3.10: The Series $\sum 1/(n(\log n)^p)$

Let $p \in \mathbb{R}$. The series

$$\sum_{n \geq 2} \frac{1}{n(\log n)^p}$$

converges if and only if $p > 1$. Its terms are defined because $\log n \geq \log 2 > \log 1 = 0$ for $n \geq 2$, the strict increase of log being theorem 4.6.4(3), so definition 4.6.8 applies to the base $\log n$.

The case $p \leq 0$. By example 2.4.18 the number e satisfies $e < 3$, so $\log 3 > \log e = 1$ by the strict increase of log and definition 4.6.5, and $\log n \geq 1$ for every $n \geq 3$. Write $\beta = -p \geq 0$. The map $u \mapsto u^\beta$ is increasing on $(0, \infty)$, by theorem 4.6.10(1) and corollary 4.2.7(1) applied to its derivative $\beta u^{\beta-1} \geq 0$, so $(\log n)^\beta \geq 1^\beta = 1$ for $n \geq 3$. Since $(\log n)^p = 1/(\log n)^\beta$ by theorem 4.6.9(1),

$$\frac{1}{n(\log n)^p} = \frac{(\log n)^\beta}{n} \geq \frac{1}{n} \quad (n \geq 3)$$

and theorem 2.4.13(2) against the divergent harmonic series of example 2.4.8 makes the series diverge.

The integrand for $p > 0$. Let $f(t) = 1/(t(\log t)^p)$ on $[2, \infty)$, which is defined and positive there. For $2 \leq s < t$ the strict increase of log gives $0 < \log s < \log t$, and $u \mapsto u^p$ is strictly increasing on $(0, \infty)$ by theorem 4.6.10(1) and corollary 4.2.7(2), so $0 < (\log s)^p < (\log t)^p$ and hence $0 < s(\log s)^p < t(\log t)^p$. Inverting with proposition 1.1.2(7) gives $f(t) < f(s)$, so f is decreasing. It is also continuous on each $[2, X]$: the restriction of log is continuous there by theorem 4.6.4(3) and proposition 3.3.1, the map $u \mapsto u^p$ is continuous on $(0, \infty)$ by theorem 4.6.10(1) and proposition 4.1.5, so $t \mapsto (\log t)^p$ is continuous by theorem 3.2.7, and theorem 3.2.5 handles the product with t and the reciprocal, whose denominator does not vanish.

An antiderivative. Fix $X > 2$. As in example 5.3.6, the restriction of log to $[2, X]$ is differentiable at every point with derivative $1/t$, by lemma 4.1.1 and proposition 3.1.23(1), and its image lies in $(0, \infty)$. For $p \neq 1$ put

$$\Lambda(t) = \frac{(\log t)^{1-p}}{1-p} \quad (2 \leq t \leq X)$$

Theorem 4.6.10(1) makes $u \mapsto u^{1-p}$ differentiable on $(0, \infty)$ with derivative $(1-p)u^{-p}$, so theorem 4.1.8 and theorem 4.1.7(2) make Λ differentiable on $[2, X]$ with

$$\Lambda'(t) = \frac{1}{1-p} \cdot (1-p)(\log t)^{-p} \cdot \frac{1}{t} = \frac{1}{t(\log t)^p} = f(t)$$

using theorem 4.6.9(1) for $(\log t)^{-p} = 1/(\log t)^p$. For $p = 1$ the same two theorems applied to $\Lambda(t) = \log(\log t)$ give $\Lambda'(t) = (1/\log t)(1/t) = f(t)$. In both cases f is continuous on $[2, X]$, hence integrable by theorem 5.2.1, and theorem 5.3.3 gives $\int_2^X f = \Lambda(X) - \Lambda(2)$.

A limit at infinity. Let $\beta > 0$. Then $\lim_{X \rightarrow \infty} (\log X)^\beta = +\infty$. Indeed, let $M > 0$ and put $u_0 = M^{1/\beta} > 0$, so that $u_0^\beta = M$ by theorem 4.6.9(3). The map $u \mapsto u^\beta$ is strictly increasing on $(0, \infty)$ as above, and proposition 4.6.7 produces an $N \in \mathbb{R}$ with $\log X > u_0$ for every $X > N$. For such X , $(\log X)^\beta > u_0^\beta = M$, which is the condition of definition 3.1.28 at infinity. Taking $M = 1/\epsilon$

and using proposition 1.1.2(7) gives $0 < (\log X)^{-\beta} < \epsilon$ for every such X , so $\lim_{X \rightarrow \infty} (\log X)^{-\beta} = 0$. The verdict for $p > 1$. Here $\beta = p - 1 > 0$ and $(\log X)^{1-p} = (\log X)^{-\beta}$ by theorem 4.6.9(1), so the previous paragraph and proposition 3.1.29(1) give $\lim_{X \rightarrow \infty} \Lambda(X) = 0$ and hence $\lim_{X \rightarrow \infty} \int_2^X f = -\Lambda(2) = (\log 2)^{1-p}/(p-1)$. The improper integral converges, so theorem 5.3.9 with $N_0 = 2$ makes the series converge.

The verdict for $0 < p \leq 1$. Suppose $0 < p < 1$ and let $M \in \mathbb{R}$. Applying the limit above with $\beta = 1 - p > 0$ to the number $(1-p)(M + \Lambda(2))$ produces an X with $(\log X)^{1-p} > (1-p)(M + \Lambda(2))$, and dividing by $1 - p > 0$ gives $\int_2^X f = \Lambda(X) - \Lambda(2) > M$. Suppose instead $p = 1$. Given $M \in \mathbb{R}$, proposition 4.6.7 gives an N_1 with $\log u > M$ for every $u > N_1$, and then an N_2 with $\log X > N_1$ for every $X > N_2$, so $\int_2^X f = \log(\log X) - \log(\log 2) > M - \log(\log 2)$ for such X . In both cases the numbers $\int_2^X f$ are unbounded above, so theorem 5.3.8(1) makes the improper integral diverge and theorem 5.3.9 makes the series diverge.

The exponent $p = 1$ is the boundary in both families at once: $\sum 1/n^p$ turns at $p = 1$, and $\sum 1/(n(\log n)^p)$ turns at $p = 1$ again, one logarithm further out. Replacing $\log n$ by $\log(\log n)$ produces a third family with the same boundary, and the process continues without terminating, so no series of this shape is a last divergent one.

Exercise 5.3.1

1. Define $f: [-1, 1] \rightarrow \mathbb{R}$ by $f(t) = 0$ for $t < 0$ and $f(t) = 1$ for $t \geq 0$. Show that $f \in \mathcal{R}[-1, 1]$, compute $F(x) = \int_{-1}^x f$ for every $x \in [-1, 1]$, and show that F fails to be differentiable at 0. Which hypothesis of theorem 5.3.2(2) fails at that point?
2. Let $a < b$ and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous and nonnegative with $\int_a^b f = 0$. Show that $f(x) = 0$ for every $x \in [a, b]$. Then show that continuity cannot be dropped, by exhibiting a nonnegative $f \in \mathcal{R}[a, b]$ with $\int_a^b f = 0$ that is not identically zero.
3. Show that $\int_0^x t \exp(t) dt = (x - 1) \exp(x) + 1$ for every $x > 0$.
4. Show that $\int_0^1 \frac{2t}{1+t^2} dt = \log 2$.
5. Let $p \in \mathbb{R}$ with $p > 0$. Show that the improper integral $\int_1^\infty t^{-p} dt$ converges if and only if $p > 1$, and that its value is $1/(p-1)$ in that case. Then deduce the verdict of example 2.4.10 for integer $p \geq 1$ from theorem 5.3.9, and say what the argument adds for non-integer exponents.
6. Show that the improper integral $\int_0^1 t^{-1/2} dt$ of definition 5.3.7(2) converges with value 2, and that $\int_0^1 t^{-1} dt$ diverges.
7. Apply (5.12) to $f(t) = 1/t$ on $[1, \infty)$ to prove that

$$\log(N+1) \leq \sum_{n=1}^N \frac{1}{n} \leq 1 + \log N$$

for every integer $N \geq 1$, and deduce that every $\gamma_N = \sum_{n=1}^N 1/n - \log N$ lies in $[0, 1]$.

8. Show that $\int_1^\infty \frac{\sin t}{t^2} dt$ converges. Identify the step at which theorem 5.3.8(3) is used, and say why theorem 5.3.8(2) alone does not settle the question.

5.4★ The Riemann-Stieltjes Integral

★ *Optional. This section is an aside; nothing later in the book depends on it.*

The weight attached to the subinterval $[x_{i-1}, x_i]$ in a Darboux sum is its length Δx_i , and the Darboux theory of section 5.1 rests on two properties of that weight. It is nonnegative, and the weights of the pieces into which a subinterval is cut add up to the weight of the subinterval. Any increasing function $\alpha: [a, b] \rightarrow \mathbb{R}$ produces weights with both properties, namely the increments $\alpha(x_i) - \alpha(x_{i-1})$, so the arguments behind the refinement inequalities and the integrability criterion go through with those in place of the lengths. The arguments that name the lengths themselves do not, and theorem 5.1.11 is the one that matters. For an integrator constant on either side of a single interior jump, exactly one subinterval of any partition carries the whole increase of α . So the sum $\sum_i f(\xi_i) \Delta \alpha_i$, formed as in definition 5.1.10 with the increments of α in place of the lengths, reduces to one value of f at a tag of that subinterval, and both endpoints of that subinterval are admissible tags. Shrinking the mesh does not bring those two values together when f is discontinuous at the jump point. Every statement below is therefore proved afresh from the increments of α , and both weights appear in those proofs, since uniform continuity of f and the integrability of α' are conditions on lengths.

One reading of the resulting integral is a distribution of mass along a rod occupying $[a, b]$, with $\alpha(x)$ the mass lying in $[a, x]$, so that a mass sitting at the point x shows up in the jump of α there. A rod of continuously varying density has a differentiable α , and theorem 5.4.7 identifies the integral against such an α with the Riemann integral of $f\alpha'$. A rod carrying beads at finitely many points has an α that increases only by jumps, and example 5.4.6 identifies the integral against that α with a finite sum. The moment of inertia of the rod is $\int_a^b x^2 d\alpha$ in both cases, computed by one formula. Nothing later in the book uses this section.

Fix the data once. Throughout, $a < b$ are real numbers, $\alpha: [a, b] \rightarrow \mathbb{R}$ is increasing in the sense of definition 3.2.14, and $f: [a, b] \rightarrow \mathbb{R}$ is bounded. Being increasing, α satisfies $\alpha(a) \leq \alpha(x) \leq \alpha(b)$ for every $x \in [a, b]$, so α is bounded as well. For a partition $P = \{x_0 < x_1 < \cdots < x_n\}$ of $[a, b]$ as in definition 5.1.1, write

$$\Delta \alpha_i = \alpha(x_i) - \alpha(x_{i-1})$$

Each $\Delta \alpha_i$ is nonnegative because α is increasing, and the sum $\sum_{i=1}^n \Delta \alpha_i$ telescopes: each $\alpha(x_i)$ with $0 < i < n$ occurs once with a plus sign and once with a minus sign, leaving $\alpha(b) - \alpha(a)$.

The two families of sums that the definition below compares are bounded, which is what makes the definition possible at all. Write $m = \inf_{[a, b]} f$ and $M = \sup_{[a, b]} f$, which exist by theorem 1.2.4 and corollary 1.2.5 because f is bounded, and let m_i and M_i denote the infimum and supremum of f on $[x_{i-1}, x_i]$, as in definition 5.1.2. Then $m \leq m_i \leq M_i \leq M$, and multiplying these by $\Delta \alpha_i \geq 0$ and summing over i gives

$$m(\alpha(b) - \alpha(a)) \leq \sum_{i=1}^n m_i \Delta \alpha_i \leq \sum_{i=1}^n M_i \Delta \alpha_i \leq M(\alpha(b) - \alpha(a)) \quad (5.14)$$

for every partition P . So the sums on the left of (5.14) form a set bounded above and the sums on the right form a set bounded below, and theorem 1.2.4 and corollary 1.2.5 produce the supremum and the infimum named next.

Definition 5.4.1: Riemann-Stieltjes Integral

Let $a < b$, let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing, and let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. For a partition $P = \{x_0 < x_1 < \cdots < x_n\}$ of $[a, b]$ set

$$L(P, f, \alpha) = \sum_{i=1}^n m_i \Delta\alpha_i, \quad U(P, f, \alpha) = \sum_{i=1}^n M_i \Delta\alpha_i$$

where m_i and M_i are the infimum and the supremum of f on $[x_{i-1}, x_i]$. The *lower* and *upper Riemann-Stieltjes integrals* of f with respect to α are

$$\int_a^b f d\alpha = \sup_P L(P, f, \alpha), \quad \overline{\int_a^b} f d\alpha = \inf_P U(P, f, \alpha)$$

the supremum and the infimum being taken over all partitions of $[a, b]$. When the two agree, f is *Riemann-Stieltjes integrable with respect to α* , written $f \in \mathcal{R}(\alpha)$, and the common value is written $\int_a^b f d\alpha$. The function α is called the *integrator*.

The integrator enters only through its increments, so two integrators differing by a constant give the same integral, and an integrator that is constant on a subinterval gives that subinterval no weight at all. A short computation shows how the weights redistribute. Take $[a, b] = [0, 1]$, the integrand $f(x) = x$, the integrator $\alpha(x) = x^2$, and the partition $P = \{0, 1/2, 1\}$. The increments are $\Delta\alpha_1 = 1/4 - 0 = 1/4$ and $\Delta\alpha_2 = 1 - 1/4 = 3/4$, while f is increasing, so $m_1 = 0$, $M_1 = 1/2$, $m_2 = 1/2$ and $M_2 = 1$. Hence

$$L(P, f, \alpha) = 0 \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{3}{4} = \frac{3}{8}, \quad U(P, f, \alpha) = \frac{1}{2} \cdot \frac{1}{4} + 1 \cdot \frac{3}{4} = \frac{7}{8}$$

The same f and the same partition against the integrator $\alpha(x) = x$ have increments $1/2$ and $1/2$, giving $L(P, f) = 1/4$ and $U(P, f) = 3/4$. The integrator x^2 grows slowly on the left half of the interval and quickly on the right half, and it is the right half where f is large, so both sums rise.

Setting $\alpha(x) = x$ returns the construction of section 5.1 unchanged, so the family just defined contains it as one member.

Proposition 5.4.2: The Riemann Integral as the Case $\alpha(x) = x$

Let $a < b$ and let $\alpha: [a, b] \rightarrow \mathbb{R}$ be given by $\alpha(x) = x$. Then α is increasing, and for every bounded $f: [a, b] \rightarrow \mathbb{R}$ and every partition P of $[a, b]$,

$$L(P, f, \alpha) = L(P, f), \quad U(P, f, \alpha) = U(P, f)$$

Consequently $f \in \mathcal{R}(\alpha)$ if and only if $f \in \mathcal{R}[a, b]$, and in that case the two integrals are equal.

Proof:

If $x \leq y$ then $\alpha(x) = x \leq y = \alpha(y)$, so α is increasing. For a partition $P = \{x_0 < \cdots < x_n\}$,

$$\Delta\alpha_i = \alpha(x_i) - \alpha(x_{i-1}) = x_i - x_{i-1} = \Delta x_i$$

for each i , and the numbers m_i and M_i do not involve α . So the two sums of definition 5.4.1 are

term by term the two sums of definition 5.1.2.

The set of all lower sums $L(P, f, \alpha)$ is therefore the same set of real numbers as the set of all lower Darboux sums $L(P, f)$, and likewise for the two sets of upper sums. A set has one supremum and one infimum, so the lower and upper Riemann-Stieltjes integrals of f with respect to α are exactly the lower and upper integrals of f in the sense of definition 5.1.6. So the two numbers compared in definition 5.4.1 are the two numbers compared in definition 5.1.7, which makes the two memberships equivalent and the two integrals equal whenever they exist, as we sought to show.

The theory needs refinement statements of its own. Those of section 5.1 are written with the increments Δx_i in them and do not transfer, and every proof below runs on the three facts collected next: refining a partition raises the lower sum and lowers the upper sum, any lower sum is below any upper sum, and the lower integral is below the upper one.

Proposition 5.4.3: Refinement and the Two Stieltjes Integrals

Let $a < b$, let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing, let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, and let P be a partition of $[a, b]$.

1. If P^* is a refinement of P , then $L(P, f, \alpha) \leq L(P^*, f, \alpha)$ and $U(P^*, f, \alpha) \leq U(P, f, \alpha)$.
2. $L(P, f, \alpha) \leq U(Q, f, \alpha)$ for all partitions P and Q of $[a, b]$.
3. $\int_a^b f d\alpha \leq \int_a^b f d\alpha$.

Proof:

1. Take first the case $P^* = P \cup \{y\}$ with $y \notin P$, say $x_{i-1} < y < x_i$. Write w_1 and w_2 for the suprema of f on $[x_{i-1}, y]$ and on $[y, x_i]$. Each of these two intervals is a subset of $[x_{i-1}, x_i]$, so the image of f on each is a subset of the image of f on $[x_{i-1}, x_i]$, and proposition 1.3.2(1) gives $w_1 \leq M_i$ and $w_2 \leq M_i$. The two increments $\alpha(y) - \alpha(x_{i-1})$ and $\alpha(x_i) - \alpha(y)$ are nonnegative and add to $\Delta\alpha_i$, so

$$w_1(\alpha(y) - \alpha(x_{i-1})) + w_2(\alpha(x_i) - \alpha(y)) \leq M_i(\alpha(y) - \alpha(x_{i-1})) + M_i(\alpha(x_i) - \alpha(y)) = M_i \Delta\alpha_i$$

Every other term of $U(P^*, f, \alpha)$ is a term of $U(P, f, \alpha)$, so $U(P^*, f, \alpha) \leq U(P, f, \alpha)$.

For the infima, let v_1 and v_2 be the infima of f on $[x_{i-1}, y]$ and on $[y, x_i]$. Applying proposition 1.3.2(3) to each image set turns each infimum into the negative of a supremum, and proposition 1.3.2(1) applied to those suprema gives $v_1 \geq m_i$ and $v_2 \geq m_i$. The same two lines with the inequalities reversed give $L(P, f, \alpha) \leq L(P^*, f, \alpha)$.

A general refinement P^* of P contains P together with finitely many further points. Adjoining those points one at a time produces a chain of partitions from P to P^* , each obtained from its predecessor by adding a single point, and the previous two paragraphs apply at each link of the chain. Composing the inequalities along the chain gives the general case.

2. Let $R = P \cup Q$, which is a refinement of P and of Q by definition 5.1.3. Since $m_i \leq M_i$ and $\Delta\alpha_i \geq 0$ for the partition R , comparing the two sums of definition 5.4.1 term by term gives $L(R, f, \alpha) \leq U(R, f, \alpha)$. Part (1) gives $L(P, f, \alpha) \leq L(R, f, \alpha)$ and $U(R, f, \alpha) \leq U(Q, f, \alpha)$, and the three inequalities chain to $L(P, f, \alpha) \leq U(Q, f, \alpha)$.
3. Fix a partition Q . By part (2), $U(Q, f, \alpha)$ is an upper bound of the set of all lower sums, so it is at least that set's least upper bound by definition 1.2.2, which is $\int_a^b f d\alpha$. As Q was arbitrary, $\int_a^b f d\alpha$ is a lower bound of the set of all upper sums, so it is at most that set's greatest lower bound by definition 1.2.2, completing the proof.

Deciding integrability from definition 5.4.1 means computing two numbers, each an extremum over all partitions, and then comparing them. The criterion below replaces that by the exhibition of a single partition on which the two sums are close, which is what every existence proof in this section produces.

Theorem 5.4.4: The Stieltjes Integrability Criterion

Let $a < b$, let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing and let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then $f \in \mathcal{R}(\alpha)$ if and only if for every $\epsilon > 0$ there is a partition P of $[a, b]$ with

$$U(P, f, \alpha) - L(P, f, \alpha) < \epsilon$$

Proof:

Suppose first that every $\epsilon > 0$ admits such a partition, let $\epsilon > 0$, and let P be a partition answering it. By definition 1.2.2 the lower integral is at least the lower sum $L(P, f, \alpha)$ and the upper integral is at most the upper sum $U(P, f, \alpha)$, and proposition 5.4.3(3) puts the lower integral below the upper one, so

$$L(P, f, \alpha) \leq \int_a^b f d\alpha \leq \overline{\int_a^b f d\alpha} \leq U(P, f, \alpha)$$

Adding the first and the last of these inequalities gives

$$\overline{\int_a^b f d\alpha} \leq \int_a^b f d\alpha + (U(P, f, \alpha) - L(P, f, \alpha)) < \int_a^b f d\alpha + \epsilon$$

This holds for every $\epsilon > 0$, so proposition 1.4.7(1) gives that the upper integral is at most the lower one. With proposition 5.4.3(3) the two are equal, which is $f \in \mathcal{R}(\alpha)$.

Conversely, let $f \in \mathcal{R}(\alpha)$, write A for the common value of the two integrals, and let $\epsilon > 0$. Since A is the least upper bound of the set of lower sums and $A - \epsilon/2 < A$, the number $A - \epsilon/2$ is not an upper bound of that set by definition 1.2.2, so there is a partition P_1 with $L(P_1, f, \alpha) > A - \epsilon/2$. Symmetrically, $A + \epsilon/2$ is not a lower bound of the set of upper sums, so there is a partition P_2 with $U(P_2, f, \alpha) < A + \epsilon/2$.

Let $P = P_1 \cup P_2$, a refinement of both. By proposition 5.4.3(1),

$$L(P, f, \alpha) \geq L(P_1, f, \alpha) > A - \frac{\epsilon}{2}, \quad U(P, f, \alpha) \leq U(P_2, f, \alpha) < A + \frac{\epsilon}{2}$$

Combining the two gives $U(P, f, \alpha) - L(P, f, \alpha) < (A + \epsilon/2) - (A - \epsilon/2) = \epsilon$, completing the proof.

The criterion is the exact analogue of theorem 5.1.8 and is used the same way. What it needs from the integrator is only that the increments are nonnegative, which is what the hypothesis that α be increasing provides. With increments of both signs the inequality $L(P, f, \alpha) \leq U(P, f, \alpha)$ can fail, and with it the two-sided approximation the whole construction rests on.

Continuity of the integrand is enough for integrability against every increasing integrator, however badly the integrator behaves. The reason is that uniform continuity controls the oscillation of f on every subinterval at once, while the increments of α contribute only their fixed total $\alpha(b) - \alpha(a)$.

Theorem 5.4.5: Continuous Integrands Are Integrable Against Every Increasing Integrator

Let $a < b$, let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing and let $f: [a, b] \rightarrow \mathbb{R}$ be continuous. Then $f \in \mathcal{R}(\alpha)$.

Proof:

By theorem 3.3.5 the function f is bounded on $[a, b]$, so the sums of definition 5.4.1 are defined. Write $K = \alpha(b) - \alpha(a)$, a nonnegative number, and let $\epsilon > 0$. Set $\eta = \epsilon/(K + 1)$, which is positive because $K + 1 > 0$.

Step 1: a partition fine enough for f . By theorem 3.4.4 the function f is uniformly continuous on $[a, b]$, so there is a $\delta > 0$ such that $|f(s) - f(t)| < \eta$ for all $s, t \in [a, b]$ with $|s - t| < \delta$. Since $b - a > 0$, the number $\delta/(b - a)$ is positive, and theorem 1.4.1(3) gives an $n \in \mathbb{N}$ with $1/n < \delta/(b - a)$, hence $(b - a)/n < \delta$. Let

$$x_i = a + i \frac{b - a}{n} \quad (i = 0, 1, \dots, n)$$

These points increase from $x_0 = a$ to $x_n = b$, so they form a partition P , and $\Delta x_i = (b - a)/n < \delta$ for every i .

Step 2: the oscillation on each subinterval. Fix i and let $s, t \in [x_{i-1}, x_i]$. Then $|s - t| \leq x_i - x_{i-1} < \delta$, so $|f(s) - f(t)| < \eta$. Thus η is an upper bound of the set $\{|f(s) - f(t)| \mid s, t \in [x_{i-1}, x_i]\}$, whose supremum is $M_i - m_i$ by proposition 1.3.10(2). By definition 1.2.2 a supremum is at most any upper bound, so $M_i - m_i \leq \eta$.

Step 3: the criterion. Each $\Delta \alpha_i$ is nonnegative, so multiplying the bound of Step 2 by $\Delta \alpha_i$ and summing gives

$$U(P, f, \alpha) - L(P, f, \alpha) = \sum_{i=1}^n (M_i - m_i) \Delta \alpha_i \leq \eta \sum_{i=1}^n \Delta \alpha_i = \eta K = \frac{K}{K + 1} \epsilon$$

Since $0 \leq K < K + 1$ and $\epsilon > 0$, the right-hand side is smaller than ϵ . So P answers ϵ in theorem 5.4.4, and as $\epsilon > 0$ was arbitrary that theorem gives $f \in \mathcal{R}(\alpha)$, completing the proof.

Two extremes of integrator sit at the ends of the family. The first is an integrator that is constant

except for finitely many jumps. Its increments are zero except on the subintervals containing a jump, so the sums of definition 5.4.1 have only finitely many nonzero terms no matter how fine the partition, and the integral collapses to a finite sum. The construction below is the finite case of the increasing function assembled in example 3.2.18, where a convergent series of positive numbers deposits its n -th term at the n -th point of a prescribed set.

Example 5.4.6: A Step Integrator and Its Integral

Let $a < s_1 < s_2 < \cdots < s_N < b$ and let $c_1, \dots, c_N$ be positive real numbers. For $1 \leq k \leq N$ define $u_k: [a, b] \rightarrow \mathbb{R}$ by $u_k(x) = 1$ if $s_k < x$ and $u_k(x) = 0$ otherwise, and set

$$\alpha(x) = \sum_{k=1}^N c_k u_k(x)$$

Then α is increasing, and for every continuous $f: [a, b] \rightarrow \mathbb{R}$,

$$\int_a^b f d\alpha = \sum_{k=1}^N c_k f(s_k)$$

α is increasing. Let $x \leq y$ in $[a, b]$ and let k be an index. If $u_k(x) = 1$ then $s_k < x \leq y$, so $u_k(y) = 1$. Hence $u_k(x) \leq u_k(y)$ in every case, and multiplying by $c_k > 0$ and summing over k gives $\alpha(x) \leq \alpha(y)$. In particular f belongs to $\mathcal{R}(\alpha)$ by theorem 5.4.5, so only the value of the integral is at issue.

The choice of a partition. Write $S = \sum_{k=1}^N c_k f(s_k)$ and $C = \sum_{k=1}^N c_k$, a positive number. Let $\epsilon > 0$. For each k the function f is continuous at s_k , so definition 3.2.1 gives a $\delta_k > 0$ with $|f(x) - f(s_k)| < \epsilon$ for all $x \in [a, b]$ with $|x - s_k| < \delta_k$. Let δ be one third of the least of the finitely many positive numbers $\delta_1, \dots, \delta_N$, $s_1 - a$, $b - s_N$, and, when $N \geq 2$, the gaps $s_{k+1} - s_k$ for $1 \leq k \leq N - 1$. Then $\delta > 0$, each of those numbers is at least 3δ , and

$$a < s_1 - \delta, \quad s_k + \delta < s_{k+1} - \delta \quad (1 \leq k \leq N - 1), \quad s_N + \delta < b$$

so the $2N + 2$ numbers $a, s_1 - \delta, s_1 + \delta, \dots, s_N - \delta, s_N + \delta, b$ increase, and they form a partition P of $[a, b]$.

The increments of α on P . For $x < y$ in $[a, b]$ the difference $u_k(y) - u_k(x)$ equals 1 when $x \leq s_k < y$ and equals 0 otherwise, so $\alpha(y) - \alpha(x)$ is the sum of those c_k whose index satisfies $x \leq s_k < y$. On the subinterval $[s_k - \delta, s_k + \delta]$ the indices j with $s_k - \delta \leq s_j < s_k + \delta$ are exactly $j = k$, since $s_j \leq s_{k-1} \leq s_k - 3\delta < s_k - \delta$ for $j < k$ and $s_j \geq s_{k+1} \geq s_k + 3\delta > s_k + \delta$ for $j > k$. So that subinterval carries increment c_k . On each of the remaining subintervals, namely $[a, s_1 - \delta]$, the intervals $[s_k + \delta, s_{k+1} - \delta]$, and $[s_N + \delta, b]$, no s_j lies in the half-open interval determined by its endpoints, so the increment is 0.

The two sums. Only the N subintervals around the s_k contribute, so writing $M^{(k)}$ and $m^{(k)}$ for the supremum and the infimum of f on $[s_k - \delta, s_k + \delta]$,

$$U(P, f, \alpha) = \sum_{k=1}^N M^{(k)} c_k, \quad L(P, f, \alpha) = \sum_{k=1}^N m^{(k)} c_k$$

For $x \in [s_k - \delta, s_k + \delta]$ one has $|x - s_k| \leq \delta < \delta_k$, hence $f(s_k) - \epsilon < f(x) < f(s_k) + \epsilon$ by proposition 1.3.8(2). So $f(s_k) + \epsilon$ is an upper bound and $f(s_k) - \epsilon$ a lower bound of the values of f there, and definition 1.2.2 gives $M^{(k)} \leq f(s_k) + \epsilon$ and $m^{(k)} \geq f(s_k) - \epsilon$. Multiplying by $c_k > 0$ and summing,

$$L(P, f, \alpha) \geq S - \epsilon C, \quad U(P, f, \alpha) \leq S + \epsilon C$$

Conclusion. The integral equals both the lower and the upper Stieltjes integral, so definition 1.2.2 places it between $L(P, f, \alpha)$ and $U(P, f, \alpha)$. With the two bounds of the previous paragraph,

$$S - \epsilon C \leq \int_a^b f d\alpha \leq S + \epsilon C$$

so $|\int_a^b f d\alpha - S| \leq \epsilon C$ by proposition 1.3.8(2). Given any $\theta > 0$, running the argument with $\epsilon = \theta/C$ gives $|\int_a^b f d\alpha - S| \leq \theta$, so proposition 1.4.7(2) gives $\int_a^b f d\alpha = S$, which is the asserted value of the integral.

A finite sum is therefore an integral, against an integrator built to have the right jumps. Taking $c_k = 1$ for every k and $f(x) = x^2$ turns $\int_a^b x^2 d\alpha$ into $s_1^2 + \cdots + s_N^2$, the moment of inertia of N unit beads placed at the points s_k . The integrator records where the mass sits, and one formula covers a mass spread along the rod and a mass concentrated at finitely many points.

The opposite extreme is an integrator with no jumps at all, differentiable at every point, whose derivative is itself integrable. Then the increment $\Delta\alpha_i$ is a derivative value times a length by the mean value theorem, and the whole Stieltjes integral reduces to an ordinary one.

Theorem 5.4.7: Reduction to a Riemann Integral

Let $a < b$, let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing and differentiable at every point of $[a, b]$, and suppose $\alpha' \in \mathcal{R}[a, b]$. Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then $f \in \mathcal{R}(\alpha)$ if and only if $f\alpha' \in \mathcal{R}[a, b]$, and in that case

$$\int_a^b f d\alpha = \int_a^b f(x) \alpha'(x) dx$$

Proof:

Write $M = \sup_{[a,b]} |f|$, which exists by theorem 1.2.4 and corollary 1.2.5 because f is bounded, and note $M \geq 0$. Boundedness is part of definition 5.1.7, so α' is bounded, and writing $M' = \sup_{[a,b]} |\alpha'|$ the estimate $|f(x)\alpha'(x)| = |f(x)||\alpha'(x)| \leq MM'$ of proposition 1.3.8(3) makes the product $f\alpha'$ bounded as well.

Step 1: an upper sum as a supremum over tags. Let $P = \{x_0 < \cdots < x_n\}$ be a partition of $[a, b]$, let $g: [a, b] \rightarrow \mathbb{R}$ be bounded and let $w_1, \dots, w_n$ be nonnegative reals. Then

$$\sup \left\{ \sum_{i=1}^n g(s_i) w_i \mid s_i \in [x_{i-1}, x_i] \text{ for each } i \right\} = \sum_{i=1}^n \left(\sup_{[x_{i-1}, x_i]} g \right) w_i \quad (5.15)$$

To see it, put $A_i = \{g(s)w_i \mid s \in [x_{i-1}, x_i]\}$, which is nonempty and bounded above. Then $A_i = w_i \cdot g([x_{i-1}, x_i])$, so proposition 1.3.2(2) with the nonnegative constant w_i gives $\sup A_i =$

$w_i \sup_{[x_{i-1}, x_i]} g$. An element of the set on the left of (5.15) is a sum $\sum_i g(s_i)w_i$ with the s_i chosen independently, which is exactly an element of $A_1 + \cdots + A_n$, so the two sets are equal. Proposition 1.3.2(4) gives $\sup(A_1 + A_2) = \sup A_1 + \sup A_2$, and since $A_1 + \cdots + A_{k+1} = (A_1 + \cdots + A_k) + A_{k+1}$, an induction on k extends the identity to any finite number of summands. Combining it with the values of the $\sup A_i$ proves (5.15).

Taking $g = f$ and $w_i = \Delta\alpha_i$ in (5.15) identifies $U(P, f, \alpha)$ with the supremum over tags of $\sum_i f(s_i)\Delta\alpha_i$, and taking $g = f\alpha'$ and $w_i = \Delta x_i$ identifies $U(P, f\alpha')$ with the supremum over tags of $\sum_i f(s_i)\alpha'(s_i)\Delta x_i$.

Step 2: the two upper sums on a fine partition. Let $\epsilon > 0$. Since $\alpha' \in \mathcal{R}[a, b]$, theorem 5.1.8 gives a partition P_0 with $U(P_0, \alpha') - L(P_0, \alpha') < \epsilon$. Let $P = \{x_0 < \cdots < x_n\}$ be any refinement of P_0 . By proposition 5.1.4, refining raises the lower sum and lowers the upper sum, so

$$U(P, \alpha') - L(P, \alpha') < \epsilon \quad (5.16)$$

The function α is differentiable at every point of $[a, b]$, hence continuous on $[a, b]$ by proposition 4.1.5. So theorem 4.2.4, applied with the interval $[a, b]$, the function α , and the pair $u = x_{i-1} < v = x_i$, produces a $t_i \in (x_{i-1}, x_i)$ with

$$\Delta\alpha_i = \alpha'(t_i) \Delta x_i \quad (5.17)$$

Let $s_i \in [x_{i-1}, x_i]$ be arbitrary and put $D = \sum_{i=1}^n f(s_i)(\alpha'(t_i) - \alpha'(s_i))\Delta x_i$. For each i write $u_i = f(s_i)(\alpha'(t_i) - \alpha'(s_i))$. Then $-|u_i| \leq u_i \leq |u_i|$ by proposition 1.3.8(1), and $|u_i| = |f(s_i)| |\alpha'(t_i) - \alpha'(s_i)| \leq M |\alpha'(t_i) - \alpha'(s_i)|$ by proposition 1.3.8(3) together with $|f(s_i)| \leq M$. Multiplying these by $\Delta x_i > 0$ and summing over i gives

$$-M \sum_{i=1}^n |\alpha'(t_i) - \alpha'(s_i)| \Delta x_i \leq D \leq M \sum_{i=1}^n |\alpha'(t_i) - \alpha'(s_i)| \Delta x_i$$

The hypothesis of proposition 5.1.12 for the function α' and the partition P is (5.16), and s_i and t_i both lie in $[x_{i-1}, x_i]$, so that proposition gives $\sum_i |\alpha'(s_i) - \alpha'(t_i)| \Delta x_i < \epsilon$. The two absolute values agree by proposition 1.3.8(1), so the sum in the display is smaller than ϵ , and multiplying by $M \geq 0$ gives $-M\epsilon \leq D \leq M\epsilon$.

Substituting (5.17) into $\sum_i f(s_i)\Delta\alpha_i$ and then adding and subtracting $\sum_i f(s_i)\alpha'(s_i)\Delta x_i$,

$$\sum_{i=1}^n f(s_i) \Delta\alpha_i = \sum_{i=1}^n f(s_i) \alpha'(t_i) \Delta x_i = \sum_{i=1}^n f(s_i) \alpha'(s_i) \Delta x_i + D$$

Each term of $\sum_i f(s_i)\alpha'(s_i)\Delta x_i$ is at most $(\sup_{[x_{i-1}, x_i]} f\alpha')\Delta x_i$, so that sum is at most $U(P, f\alpha')$. With $D \leq M\epsilon$ the display gives $\sum_i f(s_i)\Delta\alpha_i \leq U(P, f\alpha') + M\epsilon$ for every choice of tags. The right-hand side does not depend on the tags, so it is an upper bound of the set on the left of (5.15) with $g = f$ and $w_i = \Delta\alpha_i$, whose supremum is $U(P, f, \alpha)$. A supremum is at most any upper bound by definition 1.2.2, so $U(P, f, \alpha) \leq U(P, f\alpha') + M\epsilon$.

Reading the same display the other way, $\sum_i f(s_i)\alpha'(s_i)\Delta x_i = \sum_i f(s_i)\Delta\alpha_i - D$. Here each term

of $\sum_i f(s_i)\Delta\alpha_i$ is at most $M_i\Delta\alpha_i$, so that sum is at most $U(P, f, \alpha)$, and $-D \leq M\epsilon$. Hence $\sum_i f(s_i)\alpha'(s_i)\Delta x_i \leq U(P, f, \alpha) + M\epsilon$ for every choice of tags, and taking the supremum over tags through (5.15) with $g = f\alpha'$ and $w_i = \Delta x_i$ gives $U(P, f\alpha') \leq U(P, f, \alpha) + M\epsilon$.

Step 3: the two upper integrals. Let Q be any partition of $[a, b]$. Then $P_0 \cup Q$ is a refinement of P_0 , so Step 2 applies to it, and it is a refinement of Q , so proposition 5.4.3(1) and proposition 5.1.4 give $U(P_0 \cup Q, f, \alpha) \leq U(Q, f, \alpha)$ and $U(P_0 \cup Q, f\alpha') \leq U(Q, f\alpha')$. Chaining,

$$\overline{\int_a^b} f d\alpha \leq U(P_0 \cup Q, f, \alpha) \leq U(P_0 \cup Q, f\alpha') + M\epsilon \leq U(Q, f\alpha') + M\epsilon$$

the first inequality by definition 1.2.2. So $\overline{\int_a^b} f d\alpha - M\epsilon$ is a lower bound of the set of all upper Darboux sums of $f\alpha'$, and definition 1.2.2 places it at or below the greatest lower bound, giving

$$\overline{\int_a^b} f d\alpha \leq \overline{\int_a^b} f\alpha' + M\epsilon$$

The same chain with the two upper sums exchanged, using the second inequality of Step 2 and $U(P_0 \cup Q, f, \alpha) \leq U(Q, f, \alpha)$, gives $\overline{\int_a^b} f\alpha' \leq \overline{\int_a^b} f d\alpha + M\epsilon$.

Step 4: removing ϵ . The two inequalities of Step 3 say that the difference of the two upper integrals has absolute value at most $M\epsilon$, by proposition 1.3.8(2). Let $\theta > 0$. Running Steps 2 and 3 with $\epsilon = \theta/(M+1)$, which is positive, bounds that absolute value by $M\theta/(M+1) \leq \theta$. So proposition 1.4.7(2) gives

$$\overline{\int_a^b} f d\alpha = \overline{\int_a^b} f\alpha'$$

Step 5: the lower integrals. Let $h: [a, b] \rightarrow \mathbb{R}$ be bounded. On each subinterval, proposition 1.3.2(3) gives $\sup_{[x_{i-1}, x_i]}(-h) = -\inf_{[x_{i-1}, x_i]} h$, and multiplying by $\Delta\alpha_i$ and summing gives $U(P, -h, \alpha) = -L(P, h, \alpha)$ for every partition P . Now let B be the set whose elements are the numbers $-L(P, h, \alpha)$, which is bounded below because the lower sums are bounded above by (5.14). Then $-B$ is the set of lower sums of h , so proposition 1.3.2(3) applied to B reads $\sup_P L(P, h, \alpha) = -\inf_P (-L(P, h, \alpha))$, and the infimum on the right is $\overline{\int_a^b} (-h) d\alpha$. Rearranging,

$$\overline{\int_a^b} (-h) d\alpha = -\underline{\int_a^b} h d\alpha$$

and the identical computation with Δx_i in place of $\Delta\alpha_i$ gives the same relation between the upper and lower integrals of definition 5.1.6. The function $-f$ is bounded with the same M , and $(-f)\alpha' = -(f\alpha')$, so Steps 2 to 4 applied to $-f$ give $\overline{\int_a^b} (-f) d\alpha = \overline{\int_a^b} (-f)\alpha'$. By the two relations just proved the left side is $-\underline{\int_a^b} f d\alpha$ and the right side is $-\underline{\int_a^b} f\alpha'$, so multiplying by -1 gives

$$\underline{\int_a^b} f d\alpha = \underline{\int_a^b} f\alpha'$$

Step 6: conclusion. By definition 5.4.1, f lies in $\mathcal{R}(\alpha)$ exactly when its lower and upper Stieltjes

integrals agree. Steps 4 and 5 identify those two numbers with the lower and upper integrals of $f\alpha'$, which agree exactly when $f\alpha' \in \mathcal{R}[a, b]$ by definition 5.1.7. So the two memberships are equivalent, and when they hold the common value of the first pair is the common value of the second, which is the asserted equality of integrals, completing the proof.

The theorem says that a differentiable integrator acts through its derivative, so the symbol $d\alpha$ behaves as $\alpha'(x) dx$ whenever α' exists and is integrable. The computation opened earlier can now be finished: for $\alpha(x) = x^2$ and $f(x) = x$ on $[0, 1]$, the derivative $\alpha'(x) = 2x$ is continuous and hence lies in $\mathcal{R}[0, 1]$ by theorem 5.2.1, so

$$\int_0^1 x d\alpha = \int_0^1 2x^2 dx = \frac{2}{3}$$

by theorem 5.4.7 and theorem 5.3.3, the antiderivative being $2x^3/3$. The partition $\{0, 1/2, 1\}$ gave the bracket $3/8 \leq I \leq 7/8$ for the integral I , and $2/3$ lies inside it.

Two limitations mark where the section stops. The theorem asks α to be differentiable at every point of $[a, b]$, which no integrator with a jump satisfies, so theorem 5.4.7 and example 5.4.6 describe disjoint families of integrators. And an increasing α that both jumps and grows continuously falls under neither result. The general theorem covering all increasing integrators at once belongs to the Lebesgue theory of integration, which this book does not develop.

Exercise 5.4.1

1. Let $\alpha: [0, 2] \rightarrow \mathbb{R}$ be the step integrator of example 5.4.6 with jumps of size $c_1 = 2$ at $s_1 = 1$ and $c_2 = 3$ at $s_2 = 3/2$, that is, $\alpha(x) = 0$ for $0 \leq x \leq 1$, $\alpha(x) = 2$ for $1 < x \leq 3/2$, and $\alpha(x) = 5$ for $3/2 < x \leq 2$. Compute $\int_0^2 x^2 d\alpha$, and check that the value lies between $m(\alpha(2) - \alpha(0))$ and $M(\alpha(2) - \alpha(0))$, where m and M are the infimum and the supremum of x^2 on $[0, 2]$.
2. Let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing and let $f_1, f_2: [a, b] \rightarrow \mathbb{R}$ be bounded with $f_1, f_2 \in \mathcal{R}(\alpha)$. Show that $f_1 + f_2 \in \mathcal{R}(\alpha)$ and that

$$\int_a^b (f_1 + f_2) d\alpha = \int_a^b f_1 d\alpha + \int_a^b f_2 d\alpha$$

Follow the route of theorem 5.1.13, using proposition 1.3.4 to compare the sums for $f_1 + f_2$ with those for f_1 and f_2 on a common refinement, and theorem 5.4.4 in both directions.

3. Define $\alpha: [0, 2] \rightarrow \mathbb{R}$ by $\alpha(x) = 0$ for $0 \leq x \leq 1$ and $\alpha(x) = 1$ for $1 < x \leq 2$. Show that α is increasing and that $U(P, \alpha, \alpha) = 1$ and $L(P, \alpha, \alpha) = 0$ for every partition P of $[0, 2]$. Conclude that $\alpha \notin \mathcal{R}(\alpha)$, and hence that the continuity hypothesis in example 5.4.6 cannot be dropped.
4. Let $\alpha(x) = x^3$ on $[0, 1]$. Check that α satisfies the hypotheses of theorem 5.4.7, and compute $\int_0^1 x d\alpha$.
5. Let $\alpha: [a, b] \rightarrow \mathbb{R}$ be increasing, let $f: [a, b] \rightarrow \mathbb{R}$ be bounded with $f \in \mathcal{R}(\alpha)$, and let m and M be real numbers with $m \leq f(x) \leq M$ for all $x \in [a, b]$. Prove that

$$m(\alpha(b) - \alpha(a)) \leq \int_a^b f d\alpha \leq M(\alpha(b) - \alpha(a))$$

Show also that the two inequalities can both be equalities, by exhibiting a single f and α for which they are.

What Next?

The theorems of this book are all theorems about one set. Each proof used some combination of three things: that $\mathbb{R}$ is ordered, that it is complete, and that its points are separated by a distance. The natural continuations are the three ways of keeping some of that and discarding the rest.

Analysis on Metric and Function Spaces [1] keeps the distance and discards the order. That is a larger change than it sounds, and Section 3.3 is where the reader has already met the reason: the proofs there that a continuous function on a closed bounded interval is bounded and attains its bounds went through theorem 2.2.9, and every step used that a bounded sequence of reals has a convergent subsequence. In a space of functions that is false, and recovering it becomes a hypothesis with a name. The same book generalizes Section 4.4, where the reader has already seen that the uniform limit of continuous functions is continuous, into a statement that a certain space of functions is complete.

Multivariable Calculus [2] keeps everything and raises the dimension. The definition it starts from is theorem 4.1.4, which was stated here in the form that survives the move: differentiability as approximation by a linear map, with no division anywhere in it. The mean value theorem does not survive in the same form, and seeing what replaces it is most of the work of that book's opening chapters.

Real Analysis [5] keeps the line and changes the integral. The motivation is already on the page. Example 5.2.3 produces a bounded function on $[0, 1]$ that this book's integral cannot handle, and theorem 5.2.8 needs uniform convergence to exchange a limit with an integral, which is a strong hypothesis and often false in practice. Both defects have the same cure, and it is a different definition of the integral, not a repair of this one. That book also settles the question Section 5.2 leaves open, by characterizing exactly which bounded functions are Riemann integrable.

Two external books are worth naming. Rudin's *Principles of Mathematical Analysis* covers the same ground as this book and then continues through metric spaces, the Riemann-Stieltjes integral and Fourier series in the same volume; a reader who wants one book to carry them further along the

same line should use it. Folland's *Real Analysis: Modern Techniques and Their Applications* is the standard destination for the measure-theoretic direction, and begins at roughly the point *Real Analysis* [5] does.

A reader with no particular destination in mind should take the second of these three. Multivariable calculus is where the largest number of applications live, and it is the only one of the three that needs nothing from this book beyond what Chapter 4 already established.

A

The Construction by Dedekind

Cuts

★ *Optional. This chapter is an aside; nothing later in the book depends on it.*

The main text builds $\mathbb{R}$ in section 1.1 as a set of equivalence classes of Cauchy sequences of rationals. This appendix carries the older construction, due to Dedekind, which builds $\mathbb{R}$ by partitioning $\mathbb{Q}$. It is self-contained and nothing in the book depends on it.

Two reasons to have it. The cut construction reaches the least upper bound property almost immediately, by taking a union of lower sets, where theorem 1.2.4 needed a bisection. Seeing that contrast is worth the detour. And theorem 1.5.2 identifies the field built here with the field built in the main text, so the appendix is an alternative route to the same object, which is a concrete use of a theorem that otherwise reads as an abstraction.

A.1 Cuts and the Field They Form

This appendix assumes the main text's definition 1.1.1 for the ordered-field axioms and the Archimedean property, its proposition 1.1.2 for their consequences, and its proposition 1.1.3 for the irrationality of a square root of two. Only the construction itself is redone here.

Definition A.1.1: [Dedekind] Cut

A *cut* is a pair $A|B$ of subsets of $\mathbb{Q}$ such that

1. $A \neq \emptyset$ and $B \neq \emptyset$;
2. $A \cup B = \mathbb{Q}$ and $A \cap B = \emptyset$;
3. $a < b$ for every $a \in A$ and every $b \in B$;
4. A has no largest element, that is, for every $a \in A$ there is $a' \in A$ with $a < a'$.

The set A is the *lower set* of the cut.

Condition (2) makes $B = \mathbb{Q} \setminus A$, so a cut is determined by its lower set alone, and condition (3) is a condition on that lower set: given (1) and (2), condition (3) holds if and only if A is closed downward, meaning that $p < q$ and $q \in A$ force $p \in A$. Indeed, if (3) holds and $p < q \in A$ with $p \notin A$, then $p \in B$, so $q < p$ by (3), contradicting trichotomy. Conversely, if A is closed downward and $a \in A$, $b \in B$ satisfy $b \leq a$, then $b \in A$, contradicting (2). Both formulations are used below.

Example A.1.2: Rational Cuts, the Cut at the Square Root of Two, and Two Non-Examples

1. For $q \in \mathbb{Q}$ put $q^* = \{p \in \mathbb{Q} \mid p < q\}$. Then $q^*|(\mathbb{Q} \setminus q^*)$ is a cut. It is nonempty since $q - 1 \in q^*$, and its complement is nonempty since $q \notin q^*$. It is closed downward, since $p < p' < q$ gives $p < q$. And it has no largest element: if $p < q$ then $p < (p + q)(1 + 1)^{-1} < q$ by proposition 1.1.2(8), so $(p + q)(1 + 1)^{-1}$ is an element of q^* exceeding p . This cut is called the *rational cut* at q . A cut is determined by its lower set, so q^* is written both for the set displayed above and for the cut, and therefore for the real number, it determines. Context fixes which is meant. The two cases used constantly below are 0^* , the set of negative rationals, and 1^* , the set of rationals less than 1.
2. Let $A = \{q \in \mathbb{Q} \mid q \leq 0 \text{ or } q^2 < 2\}$ and $B = \mathbb{Q} \setminus A$. Then $A|B$ is a cut, and it is not a rational cut. The set A contains 0, and $2 \notin A$ because $2^2 = 4 > 2$, so both A and B are nonempty. For downward closure, let $p < q$ with $q \in A$. If $p \leq 0$ then $p \in A$ by definition. If $p > 0$ then $q > p > 0$, so $q \in A$ forces $q^2 < 2$, and proposition 1.1.2(5) applied twice gives $p^2 < pq < q^2 < 2$, so $p \in A$. For the absence of a largest element, let $q \in A$. If $q \leq 0$ then $1 \in A$ (since $1^2 = 1 < 2$) and $1 > q$. If $q > 0$, so that $q^2 < 2$, put

$$q' = \frac{2q + 2}{q + 2}$$

which is a rational number since $q + 2 > 0$. Two computations settle the matter:

$$q' - q = \frac{2q + 2 - q(q + 2)}{q + 2} = \frac{2 - q^2}{q + 2}, \quad (q')^2 - 2 = \frac{(2q + 2)^2 - 2(q + 2)^2}{(q + 2)^2} = \frac{2(q^2 - 2)}{(q + 2)^2}$$

Since $q^2 < 2$ and $q + 2 > 0$, the first is positive, so $q' > q$. Since $(q + 2)^2 > 0$, the second is negative, so $(q')^2 < 2$ and $q' \in A$. Hence A has no largest element and $A|B$ is a cut. It is not the rational cut at any q : suppose $A = q^*$. Then $q \notin A$, so $q > 0$, and $q^2 \neq 2$ by proposition 1.1.3, so $q^2 > 2$. Form $q' = (2q + 2)/(q + 2)$ again. It is positive, since $2q + 2 > 0$

and $q + 2 > 0$. The first display above gives $q' - q = (2 - q^2)/(q + 2) < 0$, so $q' < q$. The second gives $(q')^2 - 2 = 2(q^2 - 2)/(q + 2)^2 > 0$, so $q' \notin A$. But $q' < q$ means $q' \in q^* = A$, a contradiction.

3. The pair $\emptyset|\mathbb{Q}$ satisfies condition (2) of definition A.1.1, satisfies (3) and (4) vacuously, and fails condition (1). It is excluded deliberately: were it admitted, it would be a real number below every other, and the order constructed in definition A.1.4 would have a least element.
4. The pair $\{q \in \mathbb{Q} | q \leq 0\}|\{q \in \mathbb{Q} | q > 0\}$ satisfies (1), (2) and (3) and fails (4), since 0 is a largest element of the lower set. Without condition (4) each rational would give rise to two distinct cuts, q^* and $q^* \cup \{q\}$, and exercise A.1.1 (6) works out what then goes wrong with addition.

Part (2) is the cut that proposition 1.1.3 produces: A has no largest element, and B has no least element either (exercise A.1.1 (4)), because proposition 1.1.3 forces every element of B to satisfy $q^2 > 2$. Parts (1) and (2) together say that the cuts include copies of all the rational numbers and at least one object that is not among them.

Definition A.1.3: The Real Numbers

The set of all cuts of $\mathbb{Q}$ is denoted $\mathbb{R}$, and its elements are called *real numbers*.

Since a cut is determined by its lower set, the following abbreviation is used throughout: a real number is written α , its lower set is written A , and $p \in \alpha$ means $p \in A$. An operation on real numbers is then specified by saying which rationals belong to the lower set of the result, and part of verifying such a specification is checking that the set produced is the lower set of a cut.

Definition A.1.4: Order on the Real Numbers

For real numbers $\alpha = A|B$ and $\gamma = C|D$, write $\alpha < \gamma$ if $A \subseteq C$ and $A \neq C$, and $\alpha \leq \gamma$ if $A \subseteq C$.

The order compares two real numbers by comparing how much of $\mathbb{Q}$ lies below each. Write $\sigma = A|B$ for the cut of example A.1.2(2), so that $A = \{q \in \mathbb{Q} | q \leq 0 \text{ or } q^2 < 2\}$. Then $1^* < \sigma < 2^*$. For the left inequality, a rational $p < 1$ satisfies $p \leq 0$ or $0 < p < 1$, and in the second case $p^2 < 1 < 2$. Hence $1^* \subseteq A$, while $1 \in A$ and $1 \notin 1^*$. For the right inequality, a rational $q \in A$ with $q > 0$ has $q^2 < 2 < 4$, so $q < 2$ by proposition 1.1.2(5), and a rational $q \leq 0$ satisfies $q < 2$ as well. Hence $A \subseteq 2^*$, while $3/2 \in 2^*$ and $3/2 \notin A$ because $(3/2)^2 = 9/4 > 2$.

Each arithmetic operation on $\mathbb{R}$ has to be specified by saying which rationals lie below the result, and the three operations differ sharply in how much care that takes. For addition the answer is the direct one: the rationals below $\alpha + \beta$ are the sums of a rational below α and a rational below β . For negation the direct answer fails. The set of negatives of the rationals not below α has a largest element whenever that complement has a least one, which condition (4) of definition A.1.1 forbids, so the extreme element has to be discarded, and the definition below does it by requiring a strict gap $r > 0$. For multiplication the direct answer fails differently: a product of two rationals below α and β is large when both factors are large and negative, so the products of rationals below α and β are not bounded above at all. The definition therefore treats positive real numbers first and extends to the rest by the sign rules of proposition 1.1.2(4).

Theorem A.1.5: The Real Numbers Form an Ordered Field

For real numbers α and β define the following subsets of $\mathbb{Q}$.

1. $\alpha + \beta := \{r + s \mid r \in \alpha \text{ and } s \in \beta\}$.
2. $-\alpha := \{p \in \mathbb{Q} \mid \text{there is } r \in \mathbb{Q} \text{ with } r > 0 \text{ and } -p - r \notin \alpha\}$.
3. If $\alpha > 0^*$ and $\beta > 0^*$,

$$\alpha\beta := \{p \in \mathbb{Q} \mid p \leq rs \text{ for some } r \in \alpha \text{ and } s \in \beta \text{ with } r > 0 \text{ and } s > 0\}$$

Further, $\alpha \cdot 0^* := 0^*$ and $0^* \cdot \alpha := 0^*$; and for $\alpha, \beta \neq 0^*$ not both positive,

$$\alpha\beta := \begin{cases} (-\alpha)(-\beta) & \text{if } \alpha < 0^* \text{ and } \beta < 0^* \\ -((-\alpha)\beta) & \text{if } \alpha < 0^* < \beta \\ -(\alpha(-\beta)) & \text{if } \beta < 0^* < \alpha \end{cases}$$

4. If $\alpha > 0^*$,

$$\alpha^{-1} := \{p \in \mathbb{Q} \mid p \leq 0\} \cup \{p \in \mathbb{Q} \mid p > 0 \text{ and } bp < 1 \text{ for some } b \notin \alpha\}$$

and if $\alpha < 0^*$, $\alpha^{-1} := -((-\alpha)^{-1})$.

Each of these subsets is the lower set of a cut. With these operations and with the order of definition A.1.4, the set $\mathbb{R}$ is an ordered field whose zero element is 0^* and whose unit element is 1^* .

Proof:

Step 1: the order is transitive and satisfies trichotomy. Transitivity is transitivity of inclusion together with the observation that $A \subseteq C \subseteq E$ with $A \neq C$ and $C \neq E$ gives $A \neq E$: if $A = E$ then $C \subseteq E = A \subseteq C$, so $C = A$. For trichotomy, let $\alpha = A|B$ and $\gamma = C|D$ be cuts and suppose A is not contained in C . Take $p \in A$ with $p \notin C$. For any $q \in C$ we cannot have $p < q$, since C is closed downward and would then contain p . Nor is $q = p$ possible, since $p \notin C$. So $q < p$, and A closed downward gives $q \in A$. Hence $C \subseteq A$, and $C \neq A$ because $p \in A \setminus C$, so $\gamma < \alpha$. Thus at least one of $\alpha < \gamma$, $\alpha = \gamma$, $\gamma < \alpha$ holds, and at most one holds because $A \subseteq C$ and $C \subseteq A$ together give $A = C$.

Step 2: addition, and the additive axioms. First, $\alpha + \beta$ is the lower set of a cut. It is nonempty, being a set of sums of elements of two nonempty sets. It is not all of $\mathbb{Q}$: choose $r' \notin \alpha$ and $s' \notin \beta$. For $r \in \alpha$ and $s \in \beta$ we have $r < r'$ and $s < s'$ by definition A.1.1(3), hence $r + s < r' + s < r' + s'$, so $r' + s' \notin \alpha + \beta$. It is closed downward: if $q < r + s$ with $r \in \alpha$ and $s \in \beta$, then $q - s < r$, so $q - s \in \alpha$, and $q = (q - s) + s \in \alpha + \beta$. It has no largest element: given $r + s$, choose $t \in \alpha$ with $t > r$, and then $t + s > r + s$ lies in $\alpha + \beta$.

Commutativity and associativity are inherited from $\mathbb{Q}$, since $\alpha + \beta$ and $\beta + \alpha$ are the same set of sums, and $(\alpha + \beta) + \gamma$ and $\alpha + (\beta + \gamma)$ are both the set of all $r + s + t$ with $r \in \alpha$, $s \in \beta$, $t \in \gamma$.

For the identity, if $r \in \alpha$ and $s \in 0^*$ then $s < 0$, so $r + s < r$ and $r + s \in \alpha$. Hence $\alpha + 0^* \subseteq \alpha$. Conversely, given $p \in \alpha$ choose $r \in \alpha$ with $r > p$. Then $p - r < 0$, so $p - r \in 0^*$ and $p = r + (p - r) \in \alpha + 0^*$. Thus $\alpha + 0^* = \alpha$.

Next, $-\alpha$ is the lower set of a cut. It is nonempty: choose $s \notin \alpha$ and put $p = -s - 1$. Then $-p - 1 = s \notin \alpha$, so $p \in -\alpha$. It is not all of $\mathbb{Q}$: if $q \in \alpha$ and $-q \in -\alpha$, there would be $r > 0$ with $q - r \notin \alpha$, but $q - r < q$ and α is closed downward, so $q - r \in \alpha$. Hence $-q \notin -\alpha$. It is closed downward: if $q < p \in -\alpha$, take $r > 0$ with $-p - r \notin \alpha$. Then $-q - r > -p - r$, and since α is closed downward and does not contain $-p - r$, it does not contain $-q - r$ either, so $q \in -\alpha$. It has no largest element: with p and r as above, put $t = p + r/2$. Then $t > p$ and $-t - r/2 = -p - r \notin \alpha$, so $t \in -\alpha$.

Now $\alpha + (-\alpha) = 0^*$. For one inclusion, let $r \in \alpha$ and $p \in -\alpha$, and take $s > 0$ with $-p - s \notin \alpha$. Then $r < -p - s$ by definition A.1.1(3), so $r + p < -s < 0$ and $r + p \in 0^*$. For the other, let $v \in 0^*$ and put $w = -v/(1+1)$, which is a positive rational. The set S of integers n with $nw \in \alpha$ is nonempty and bounded above. Indeed, choose $a \in \alpha$ and $b \notin \alpha$. Since $\mathbb{Q}$ is Archimedean there is a positive integer m with $mw > -a$, so $-mw < a$ and hence $-mw \in \alpha$, giving $-m \in S$. There is likewise a positive integer k with $kw > b$, and every rational at least as large as b lies outside α because α is closed downward, so $nw \notin \alpha$ for every $n \geq k$ and S is bounded above by k . A nonempty set of integers bounded above has a largest element, so let n be the largest element of S : then $nw \in \alpha$ and $(n+1)w \notin \alpha$. Put $p = -(n+2)w$. Then $-p - w = (n+1)w \notin \alpha$, so $p \in -\alpha$, and $nw + p = nw - (n+2)w = -2w = v$. Hence $v \in \alpha + (-\alpha)$, and $\alpha + (-\alpha) = 0^*$. Two consequences of what has just been proved are used repeatedly below. The additive inverse is unique: if $\alpha + \eta = 0^*$ then

$$\eta = 0^* + \eta = ((-\alpha) + \alpha) + \eta = (-\alpha) + (\alpha + \eta) = (-\alpha) + 0^* = -\alpha$$

Consequently $-(-\alpha) = \alpha$, because $(-\alpha) + \alpha = 0^*$, and $-(\alpha + \beta) = (-\alpha) + (-\beta)$, because $(\alpha + \beta) + ((-\alpha) + (-\beta)) = (\alpha + (-\alpha)) + (\beta + (-\beta)) = 0^*$.

Step 3: the order is compatible with addition. Let $\beta < \gamma$. Every element of $\alpha + \beta$ is $r + s$ with $r \in \alpha$ and $s \in \beta \subseteq \gamma$, so $\alpha + \beta \leq \alpha + \gamma$. Suppose the two were equal. Using only the properties of addition proved in Step 2,

$$\beta = 0^* + \beta = ((-\alpha) + \alpha) + \beta = (-\alpha) + (\alpha + \beta) = (-\alpha) + (\alpha + \gamma) = \gamma$$

contradicting $\beta < \gamma$. So $\alpha + \beta < \alpha + \gamma$. In particular $\alpha > 0^*$ if and only if $-\alpha < 0^*$: adding $-\alpha$ to $0^* < \alpha$ gives $-\alpha < 0^*$, and adding α to $-\alpha < 0^*$ gives $0^* < \alpha$.

Step 4: multiplication of positive real numbers. Let $\alpha > 0^*$ and $\beta > 0^*$. Three observations about an arbitrary real number $\delta > 0^*$ are used throughout this step. Its lower set properly contains 0^* , so it contains some $q \geq 0$, and if $q = 0$ then, δ having no largest element, it contains a larger element, which is a positive rational. In either case δ contains a positive rational. Every rational outside δ is then positive, being larger than that one by definition A.1.1(3). And δ contains every nonpositive rational, since it is closed downward and contains a positive one.

The set $\alpha\beta$ is the lower set of a cut. It is nonempty, containing rs for positive $r \in \alpha$ and $s \in \beta$. It is not all of $\mathbb{Q}$: choose $r' \notin \alpha$ and $s' \notin \beta$, both positive by the previous paragraph. If $r's' \leq rs$ with $r \in \alpha$, $s \in \beta$, $r, s > 0$, then $r < r'$ and $s < s'$, so $rs < r's < r's'$ by proposition 1.1.2(5), a contradiction. Hence $r's' \notin \alpha\beta$. It is closed downward, since $q < p \leq rs$ gives $q \leq rs$. It has no largest element: given $p \leq rs$ as in the definition, choose $r' \in \alpha$ with $r' > r$. Then $r's > rs \geq p$ and $r's \in \alpha\beta$. Since $\alpha\beta$ contains the positive rational rs and contains every nonpositive rational, $\alpha\beta > 0^*$, which is the second order axiom of definition 1.1.1.

Commutativity holds because $rs = sr$ in $\mathbb{Q}$, so the condition defining $\alpha\beta$ and the condition defining $\beta\alpha$ are the same. For associativity, $p \in (\alpha\beta)\gamma$ means $p \leq ut$ for some positive $u \in \alpha\beta$ and positive $t \in \gamma$, and such a u satisfies $u \leq rs$ for positive $r \in \alpha$, $s \in \beta$. Then $ut \leq rst$, so $p \leq rst$. Conversely, if $p \leq rst$ with r, s, t positive elements of α, β, γ , then rs is a positive

element of $\alpha\beta$ and $p \leq (rs)t$, so $p \in (\alpha\beta)\gamma$. Hence $(\alpha\beta)\gamma$ is the set of p with $p \leq rst$ for some such r, s, t , and this description is symmetric in the three factors, so it also describes $\alpha(\beta\gamma)$.

For the unit, if $p \leq rs$ with $r \in \alpha$, $r > 0$ and $0 < s < 1$, then $rs < r$, so $p < r$ and $p \in \alpha$. Hence $\alpha 1^* \subseteq \alpha$. Conversely let $p \in \alpha$. If $p \leq 0$, choose positive $r \in \alpha$ and take $s = 1/2 \in 1^*$. Then $rs > 0 \geq p$. If $p > 0$, choose $r \in \alpha$ with $r > p$ and put $s = p/r$. Then $0 < s < 1$, so $s \in 1^*$, and $p = rs$. In both cases $p \in \alpha 1^*$, so $\alpha 1^* = \alpha$.

For distributivity among positive real numbers, let $\alpha, \beta, \gamma > 0^*$. Observe first that $rs \in \alpha\beta$ whenever $r \in \alpha$ is positive and $s \in \beta$ is arbitrary: for $s > 0$ this is the definition, and for $s \leq 0$ the product rs is nonpositive and $\alpha\beta$ contains every nonpositive rational. Now let $p \in \alpha(\beta + \gamma)$, so $p \leq rm$ with $r \in \alpha$ positive and $m \in \beta + \gamma$ positive. Write $m = s + t$ with $s \in \beta$ and $t \in \gamma$. Then $rm = rs + rt$ with $rs \in \alpha\beta$ and $rt \in \alpha\gamma$, so $rm \in \alpha\beta + \alpha\gamma$, and that set is closed downward, so p belongs to it. Conversely let $u \in \alpha\beta$ and $v \in \alpha\gamma$, say $u \leq r_1 s_1$ and $v \leq r_2 t_1$ with all of r_1, r_2, s_1, t_1 positive and lying in the respective real numbers. Let r be the larger of r_1 and r_2 , an element of α . Then $u + v \leq r_1 s_1 + r_2 t_1 \leq rs_1 + rt_1 = r(s_1 + t_1)$, and $s_1 + t_1$ is a positive element of $\beta + \gamma$, so $u + v \in \alpha(\beta + \gamma)$. Hence $\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma$.

Step 5: multiplicative inverses of positive real numbers. Let $\alpha > 0^*$. Every rational outside α is then positive. The set α^{-1} is the lower set of a cut. It is nonempty, containing 0. It is not all of $\mathbb{Q}$: choose a positive $a \in \alpha$ and suppose $1/a \in \alpha^{-1}$. Since $1/a > 0$ there would be $b \notin \alpha$ with $b/a < 1$, that is $b < a$, contradicting $a < b$. It is closed downward: if $q < p \in \alpha^{-1}$ and $q \leq 0$ then $q \in \alpha^{-1}$, while if $q > 0$ then $p > 0$, so there is $b \notin \alpha$ with $bp < 1$, and $b > 0$ gives $bq < bp < 1$. It has no largest element: if $p \leq 0$, choose $b \notin \alpha$ and note that $q = 1/(2b)$ is positive with $bq = 1/2 < 1$, so $q \in \alpha^{-1}$ and $q > p$; if $p > 0$ with $bp < 1$ for some $b \notin \alpha$, then $p < 1/b$, and by proposition 1.1.2(8) any rational q with $p < q < 1/b$ lies in α^{-1} and exceeds p . Since α^{-1} contains a positive rational and every nonpositive one, $\alpha^{-1} > 0^*$.

The identity $\alpha\alpha^{-1} = 1^*$ rests on the following approximation of α from outside. *Claim: for every rational $\lambda > 1$ there are $r \in \alpha$ with $r > 0$ and $b \notin \alpha$ with $b \leq \lambda r$.* To see it, fix a positive $a_0 \in \alpha$ and some $b_0 \notin \alpha$, and put $w = (\lambda - 1)a_0 > 0$. Since $\mathbb{Q}$ is Archimedean there is a positive integer n_0 with $n_0 w > b_0 - a_0$, so $a_0 + n_0 w > b_0$ and therefore $a_0 + n_0 w \notin \alpha$, because α is closed downward and omits b_0 . The set of positive integers n with $a_0 + nw \notin \alpha$ is thus nonempty. Let m be its least element. Put $r = a_0 + (m - 1)w$ and $b = a_0 + mw$. Then $b \notin \alpha$ by the choice of m , and $r \in \alpha$: if $m = 1$ then $r = a_0 \in \alpha$, and if $m > 1$ then $m - 1$ is a positive integer smaller than m , so $a_0 + (m - 1)w \in \alpha$ by the minimality of m . Also $r \geq a_0 > 0$, and

$$b = r + w = r + (\lambda - 1)a_0 \leq r + (\lambda - 1)r = \lambda r$$

which proves the claim.

Now let $p \in \alpha\alpha^{-1}$, say $p \leq rs$ with $r \in \alpha$, $s \in \alpha^{-1}$ both positive, and take $b \notin \alpha$ with $bs < 1$. Then $r < b$, so $rs < bs < 1$ and $p < 1$, giving $\alpha\alpha^{-1} \subseteq 1^*$. Conversely let $p < 1$. If $p \leq 0$ then $p \in \alpha\alpha^{-1}$, since that real number is positive and so contains every nonpositive rational. If $0 < p < 1$ then $1/p > 1$, so proposition 1.1.2(8) provides a rational λ with $1 < \lambda < 1/p$. Apply the claim to get $r \in \alpha$ positive and $b \notin \alpha$ with $b \leq \lambda r$. Then $pb \leq p\lambda r < r$ because $p\lambda < 1$ and $r > 0$, and dividing by the positive rational rb gives $p/r < 1/b$. Choose a rational s with $p/r < s < 1/b$, again by proposition 1.1.2(8). Then $s > 0$ and $bs < 1$, so $s \in \alpha^{-1}$, and $rs > p$, so $p \in \alpha\alpha^{-1}$. Hence $\alpha\alpha^{-1} = 1^*$.

Step 6: multiplication on all of $\mathbb{R}$. By the trichotomy of Step 1, a real number other than 0^* is either positive or negative, and in the second case it is $-(-\alpha)$ with $-\alpha$ positive by Step 3. So the four clauses of the statement define $\alpha\beta$ for every pair, and by Step 4 the result is again a real number.

Commutativity: for $\alpha, \beta > 0^*$ it is Step 4; if both are negative then $\alpha\beta = (-\alpha)(-\beta) =$

$(-\beta)(-\alpha) = \beta\alpha$; if $\alpha < 0^* < \beta$ then $\alpha\beta = -((-\alpha)\beta)$ and $\beta\alpha = -(\beta(-\alpha))$, which agree because $(-\alpha)\beta$ and $\beta(-\alpha)$ are products of positive real numbers and Step 4 gives commutativity there. The case $\beta < 0^* < \alpha$ is the same with the roles interchanged. And if either factor is 0^* both products are 0^* .

The sign rule $(-\alpha)\beta = -(\alpha\beta) = \alpha(-\beta)$ holds for all α, β . The second equality follows from the first by commutativity, so only the first needs checking, and $-0^* = 0^*$ handles the cases where either factor is 0^* . If $\alpha > 0^*$ and $\beta > 0^*$ then $-\alpha < 0^* < \beta$, so $(-\alpha)\beta = -((-\alpha)\beta) = -(\alpha\beta)$. If $\alpha > 0^* > \beta$ then $-\alpha < 0^*$ and $\beta < 0^*$, so $(-\alpha)\beta = (-(-\alpha))(-\beta) = \alpha(-\beta)$, while $\alpha\beta = -(\alpha(-\beta))$ by definition, so the two sides agree. If $\alpha < 0^* < \beta$ then $-\alpha > 0^*$ and $\alpha\beta = -((-\alpha)\beta)$ by definition, which is the assertion. If $\alpha < 0^*$ and $\beta < 0^*$ then $-\alpha > 0^* > \beta$, so $(-\alpha)\beta = -((-\alpha)(-\beta))$, while $\alpha\beta = (-\alpha)(-\beta)$ by definition, and again the two sides agree.

Associativity: if any of α, β, γ equals 0^* , both $(\alpha\beta)\gamma$ and $\alpha(\beta\gamma)$ equal 0^* . So let α, β, γ be nonzero and suppose the identity $(\alpha\beta)\gamma = \alpha(\beta\gamma)$ holds for the triple (α, β, γ) . It then holds for $(-\alpha, \beta, \gamma)$, since the sign rule turns the left side into $(-\alpha\beta)\gamma = -((\alpha\beta)\gamma)$ and the right side into $-(\alpha(\beta\gamma))$. It holds for $(\alpha, -\beta, \gamma)$, since the two sides become $(-\alpha\beta)\gamma = -((\alpha\beta)\gamma)$ and $\alpha(-\beta\gamma) = -(\alpha(\beta\gamma))$. And it holds for $(\alpha, \beta, -\gamma)$, since the two sides become $(-\alpha\beta)\gamma = -((\alpha\beta)\gamma)$ and $\alpha(-\beta\gamma) = -(\alpha(\beta\gamma))$. Every nonzero real number is either positive or the negative of a positive one, so every triple of nonzero real numbers is obtained from a triple of positive ones by negating some of its entries, and the identity holds for positive triples by Step 4.

Unit and inverses: $0^* \subseteq 1^*$ and $0 \in 1^* \setminus 0^*$, so $1^* > 0^*$. In particular $1^* \neq 0^*$. Hence $\alpha 1^* = \alpha$ for $\alpha > 0^*$ by Step 4, while for $\alpha < 0^*$ the sign rule gives $\alpha 1^* = -((-\alpha)1^*) = -(-\alpha) = \alpha$, and $0^* \cdot 1^* = 0^*$. For $\alpha > 0^*$, Step 5 gives $\alpha\alpha^{-1} = 1^*$. For $\alpha < 0^*$ we have $-\alpha > 0^*$, so $(-\alpha)^{-1} > 0^*$ and hence $\alpha^{-1} = -((-\alpha)^{-1}) < 0^*$. Both factors being negative, $\alpha\alpha^{-1} = (-\alpha)(-(\alpha^{-1})) = (-\alpha)(-\alpha)^{-1} = 1^*$.

Distributivity: let $\alpha, \beta, \gamma \in \mathbb{R}$. If $\alpha = 0^*$ both sides of $\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma$ are 0^* . If $\alpha < 0^*$ and the identity is known for $-\alpha > 0^*$, then by the sign rule and the identity $-(\mu + \nu) = (-\mu) + (-\nu)$ of Step 2,

$$\alpha(\beta + \gamma) = -((-\alpha)(\beta + \gamma)) = -((-\alpha)\beta + (-\alpha)\gamma) = \alpha\beta + \alpha\gamma$$

So assume $\alpha > 0^*$. If β or γ equals 0^* the identity follows from $\alpha 0^* = 0^*$ and Step 2. If $\beta > 0^*$ and $\gamma > 0^*$ it is Step 4. If $\beta < 0^*$ and $\gamma < 0^*$ then $\beta + \gamma < 0^*$ by Step 3, while $-\beta$ and $-\gamma$ are positive, and using $-(\beta + \gamma) = (-\beta) + (-\gamma)$,

$$\alpha(\beta + \gamma) = -(\alpha((-\beta) + (-\gamma))) = -(\alpha(-\beta) + \alpha(-\gamma)) = -((-\alpha\beta) + (-\alpha\gamma)) = \alpha\beta + \alpha\gamma$$

the last step being the Step 2 identity $-(\mu + \nu) = (-\mu) + (-\nu)$ together with $-(-\mu) = \mu$. There remains the case where β and γ have opposite signs, and by commutativity of addition it suffices to treat $\beta > 0^* > \gamma$. Put $\delta = \beta + \gamma$. If $\delta > 0^*$, then $\beta = \delta + (-\gamma)$ is a sum of two positive real numbers, so Step 4 gives $\alpha\beta = \alpha\delta + \alpha(-\gamma) = \alpha\delta - \alpha\gamma$, and adding $\alpha\gamma$ gives $\alpha\delta = \alpha\beta + \alpha\gamma$. If $\delta < 0^*$, then $-\gamma = \beta + (-\delta)$ is a sum of two positive real numbers, so $\alpha(-\gamma) = \alpha\beta + \alpha(-\delta)$, that is $-\alpha\gamma = \alpha\beta - \alpha\delta$, and rearranging again gives $\alpha\delta = \alpha\beta + \alpha\gamma$. If $\delta = 0^*$ then $\gamma = -\beta$, so $\alpha\beta + \alpha\gamma = \alpha\beta - \alpha\beta = 0^* = \alpha\delta$.

Steps 1 through 6 verify every clause of definition 1.1.1 for $\mathbb{R}$, completing the proof.

The theorem says that nothing was lost in passing from $\mathbb{Q}$ to $\mathbb{R}$: every identity and inequality that follows from the ordered field axioms, and therefore every consequence recorded in proposition 1.1.2, holds for real numbers. What was gained is not yet visible, since the least upper bound property plays no part above. It is proved in theorem 1.2.4, and it is the only thing about $\mathbb{R}$ assumed later in this book beyond the ordered field axioms.

One thing still has to be checked, namely that the copy of $\mathbb{Q}$ inside $\mathbb{R}$ given by the rational cuts is a sub-ordered-field carrying the original arithmetic of $\mathbb{Q}$. Without that, a rational number could not be treated as a real number without ambiguity.

Proposition A.1.6: The Rational Numbers Embed in the Real Numbers

The map $\mathbb{Q} \rightarrow \mathbb{R}$, $q \mapsto q^*$, is injective and satisfies, for all $q, r \in \mathbb{Q}$,

$$q^* + r^* = (q + r)^*, \quad q^* r^* = (qr)^*, \quad -(q^*) = (-q)^*$$

and $q < r$ if and only if $q^* < r^*$. Moreover $0 \mapsto 0^*$ and $1 \mapsto 1^*$.

Proof:

That each q^* is a cut is example A.1.2(1), and 0^* and 1^* are the images of 0 and 1 by definition.

Order. If $q < r$ then every $p < q$ satisfies $p < r$, so $q^* \subseteq r^*$, and $q \in r^* \setminus q^*$, so $q^* < r^*$. Conversely suppose $q^* < r^*$. If $r \leq q$ then every $p < r$ satisfies $p < q$, so $r^* \subseteq q^*$, that is $r^* \leq q^*$, and this contradicts $q^* < r^*$ by the trichotomy established in Step 1 of theorem A.1.5. Hence $q < r$ by trichotomy in $\mathbb{Q}$. In particular the map is injective, since $q \neq r$ forces $q < r$ or $r < q$ and hence $q^* \neq r^*$.

Addition. If $p < q$ and $p' < r$ then $p + p' < q + r$, so $q^* + r^* \subseteq (q + r)^*$. Conversely let $t < q + r$ and put $h = (q + r - t)(1 + 1)^{-1}$, a positive rational. Then $q - h < q$ and $r - h < r$, and $(q - h) + (r - h) = q + r - 2h = t$, so $t \in q^* + r^*$.

Negation. A rational p lies in $-(q^*)$ exactly when there is $s > 0$ with $-p - s \notin q^*$, that is with $-p - s \geq q$, that is with $p \leq -q - s$. Such an s exists if and only if $p < -q$: if $p < -q$ take $s = (-q - p)(1 + 1)^{-1} > 0$, for which $-q - s > p$. Conversely, such an s gives $p \leq -q - s < -q$. Hence $-(q^*) = \{p \mid p < -q\} = (-q)^*$.

Multiplication. Suppose first $q > 0$ and $r > 0$, so that $q^* > 0^*$ and $r^* > 0^*$ by the order statement already proved. If $p \leq st$ with $0 < s < q$ and $0 < t < r$, then $st < qt < qr$ by proposition 1.1.2(5), so $p < qr$ and $p \in (qr)^*$. Conversely let $p < qr$. If $p \leq 0$, take $s = q(1 + 1)^{-1}$ and $t = r(1 + 1)^{-1}$, which are positive elements of q^* and r^* with $st > 0 \geq p$. If $p > 0$, put $c = p(qr)^{-1}$, so $0 < c < 1$. By proposition 1.1.2(8) choose a rational μ with $c < \mu < 1$, and then, since $c\mu^{-1} < 1$, a rational ν with $c\mu^{-1} < \nu < 1$. Put $s = \mu q$ and $t = \nu r$. Then $0 < s < q$ and $0 < t < r$, and $\mu\nu > c$ gives $st = \mu\nu qr > cqr = p$, so $p \in q^* r^*$. Hence $q^* r^* = (qr)^*$ for positive q and r .

For the remaining sign cases, if $q = 0$ or $r = 0$ then both sides are 0^* . If $q < 0 < r$ then $q^* < 0^*$, so by the definition of multiplication and the negation rule just proved,

$$q^* r^* = -((-q^*)r^*) = -((-q)^* r^*) = -(((-q)r)^*) = (-((-q)r))^* = (qr)^*$$

where the last step is $-((-q)r) = qr$ from proposition 1.1.2(4). The case $r < 0 < q$ is identical with the roles interchanged. If $q < 0$ and $r < 0$ then $q^* r^* = (-q^*)(-r^*) = (-q)^*(-r)^* = ((-q)(-r))^* = (qr)^*$, again by proposition 1.1.2(4) in the last step, as we sought to show.

From now on a rational number q and the real number q^* are identified, and the star is retained only where the distinction between the two is at issue. Under this identification $\mathbb{Q}$ is a subfield of $\mathbb{R}$ on which the order of definition A.1.4 restricts to the order of $\mathbb{Q}$, and example A.1.2(2) exhibits an element of $\mathbb{R}$ that is not in $\mathbb{Q}$.

The construction is not finished until the cut field is shown to have the property the main text's does, and this is where the cut route pays off: the supremum is a union, and no sequence has to be built.

Proposition A.1.7: The Cut Field Is Complete

Every nonempty set of cuts that is bounded above has a least upper bound.

Proof:

Let $\mathcal{C}$ be a nonempty set of cuts, bounded above by the cut β with lower set B . Let A be the union of the lower sets of the members of $\mathcal{C}$, and let γ be the pair $A | (\mathbb{Q} \setminus A)$.

γ is a cut. A is nonempty because $\mathcal{C}$ is and each lower set is. It is not all of $\mathbb{Q}$: every member of $\mathcal{C}$ is at most β , so every lower set is contained in B , hence $A \subseteq B \neq \mathbb{Q}$. It is closed downward, being a union of downward-closed sets. And it has no largest element: given $p \in A$, choose a member of $\mathcal{C}$ whose lower set contains p . That lower set has no largest element, so it contains some $r > p$, and $r \in A$.

γ is an upper bound. Each member of $\mathcal{C}$ has its lower set contained in A , which is exactly the statement that it is at most γ in the order of definition A.1.4.

γ is least. Let δ be an upper bound of $\mathcal{C}$, with lower set D . Every lower set of a member of $\mathcal{C}$ is contained in D , so their union A is contained in D , which is $\gamma \leq \delta$, completing the proof.

So the cut field is a complete ordered field, and theorem 1.5.2 applies to it and to the field of section 1.1 together: the two are isomorphic by a unique order-preserving isomorphism, and the appendix has constructed the same $\mathbb{R}$ by a different route.

A.1.8 Remark: How the Two Constructions Compare Cuts are not the only route from $\mathbb{Q}$ to $\mathbb{R}$. A second construction, the one Terence Tao follows in *Analysis I*, calls a sequence $q_1, q_2, q_3, \dots$ of rationals *Cauchy* when for every rational $\epsilon > 0$ there is an index N with $-\epsilon < q_n - q_m < \epsilon$ for all $n, m \geq N$. It calls two Cauchy sequences $q_1, q_2, \dots$ and $q'_1, q'_2, \dots$ equivalent when for every rational $\epsilon > 0$ there is an index N with $-\epsilon < q_n - q'_n < \epsilon$ for all $n \geq N$. And it takes a real number to be an equivalence class. That construction uses the arithmetic of $\mathbb{Q}$ and a notion of small difference, and nothing about the order, which is why it is the one that generalizes to the completion of a metric space. The cut construction uses the order of $\mathbb{Q}$ and nothing else. The two produce the same theory for the purposes of this book: every property of $\mathbb{R}$ used below is derived either from the ordered field axioms verified in theorem A.1.5 or from the least upper bound property proved in theorem 1.2.4.

Exercise A.1.1

1. Decide which of the following subsets $A \subseteq \mathbb{Q}$ are lower sets of cuts, and justify each answer against definition A.1.1.
 1. $A = \{q \in \mathbb{Q} \mid q \leq 3\}$.
 2. $A = \{q \in \mathbb{Q} \mid q \leq 0 \text{ or } q^2 < 3\}$.
 3. $A = \{q \in \mathbb{Q} \mid q^2 < 2\}$.

2. Let α be a real number with $\alpha > 0^*$. Show that there is a rational $r > 0$ with $r^* \leq \alpha$. Deduce that there is no real number α satisfying $0^* < \alpha < r^*$ for every rational $r > 0$.
3. Let $\alpha = A|B$ be a cut. Show that $\alpha = q^*$ for some $q \in \mathbb{Q}$ if and only if B has a least element.
4. Let $\alpha = A|B$ be the cut of example A.1.2(2), so that B is the set of rationals $q > 0$ with $q^2 > 2$. Show that B has no least element, by running the computation of example A.1.2(2) with a $q \in B$. Combined with exercise A.1.1 (3), this gives a second proof that α is not a rational cut.
5. Let $\alpha = A|B$ be a cut and let $r \in \mathbb{Q}$. Show that $\{a + r \mid a \in A\}$ is the lower set of a cut, and that this cut is $\alpha + r^*$.
6. Suppose condition (4) of definition A.1.1 were dropped. Show that for each $q \in \mathbb{Q}$ the set $\{p \in \mathbb{Q} \mid p \leq q\}$ would then satisfy conditions (1), (2) and (3), so that each rational would give rise to two objects. Writing α for that set and defining addition as in theorem A.1.5(1), show that $\alpha + 0^* \neq \alpha$, and identify which of the two objects attached to q the sum $\alpha + 0^*$ is.

Index

- Abel's Limit Theorem, 192
- Absolute Convergence, 68
 - Implies Convergence, 68
- Absolute Value, 16
- Adherent Point, 101
- Algebra of Limits, 36
- Alternating Series Test, 80
- Analytic Function, 208
 - characterization by derivative growth, 212
- Archimedean Property, 2, 22
 - of the real numbers, 8
- Bolzano-Weierstrass Theorem, 49
- Boundedness on a Closed Bounded Interval, 132
- Cantor Set, 105
- Cauchy Condensation Test, 66
- Cauchy Criterion, 57
 - for Series, 62
 - Uniform, 175
- Cauchy Product, 88
- Cauchy Sequence
 - in $\mathbb{R}$, 55
 - of rationals, 4
- Cauchy-Hadamard Theorem, 93
- Chain Rule, 153
- Change of Variables, 243
- Closed Set, 103
- Closure
 - of a Subset of the Real Line, 101
- Common Refinement, 215
- Comparison Test, 69
- Complete Ordered Field
 - uniqueness, 30
- Completeness
 - Equivalent Forms, 58
- Conditional Convergence, 81
- Continuity
 - algebra of continuous functions, 118
 - at a point, 116
 - of a composition, 119
 - of the Exponential, Cosine and Sine, 196
 - sequential characterization, 117
- Continuous Function, 116
- Continuous Inverse Theorem, 136
- Convergence
 - of a real sequence, 34
 - Pointwise, 173
- Darboux Sum, 214
- Darboux's Theorem, 164
- Decimal Expansion, 76
- Density
 - of $\mathbb{Q}$ in $\mathbb{R}$, 24
 - of the irrationals in $\mathbb{R}$, 24
- Derivative, 148
 - Discontinuous, 206
 - of a restriction, 239
 - of an Inverse Function, 154
 - rules, 151
- Derivative Growth Rate, 209
 - of an analytic function, 210

- Differentiability
 - implies continuity, 150
- Dirichlet Function, 121
 - not Riemann integrable, 229
- Dirichlet's Test, 79
- Discontinuity
 - of $\sin(1/x)$, 205
 - of the first and second kind, 123
- Epsilon Principle, 25
- Euler's Number, 74
 - as a Limit, 75
- Exponential Function, 96
 - Derivative, 195
 - Functional Equation, 97
 - Growth, 200
 - Range, 197
- Extended Real Line, 44
- Extreme Value Theorem, 133
- Fixed Point Theorem, 136
- Fubini's Theorem
 - for double series, 86
- Fundamental Theorem of Calculus
 - First Form, 239
 - Second Form, 241
- Geometric Series, 65
- Harmonic Series, 66
- Identity Theorem for Power Series, 190
- Improper Integral, 245
 - Convergence Tests, 245
- Infimum, 9
 - approximation property, 12
- Integer Part, 23
- Integrability Criterion, 218
- Integrable Function, 217
 - closure properties, 233
 - continuous functions, 227
 - continuous on an open interval, 231
 - finitely many discontinuities, 230
 - monotone functions, 228
- Integral
 - additivity over subintervals, 225
 - linearity of, 223
- Integral Test, 247
- Integration by Parts, 242
- Integrator, 251
- Interchange of Limits, 177
- Interior Extremum, 157
- Intermediate Value Property, 132
- Intermediate Value Theorem, 129
- Interval, 100
- Irrationality of the Square Root of Two, 4
- Isolated Point, 101
- L'Hôpital's Rule, 165
- Least Upper Bound Property, 10
 - for the cut construction, 270
- Limit
 - and weak inequalities, 39
 - at Infinity, 113
 - algebra of, 114
 - Infinite, 113
 - of a Function, 107
 - of a real sequence, 34
 - One-Sided, 111
 - uniqueness of, 35
- Limit Inferior, 50
- Limit Point, 101
- Limit Superior, 50
 - characterization, 51
 - Scaling by a Convergent Factor, 183
- Lipschitz
 - from a bounded derivative, 163
- Lipschitz Function, 145
- Local Extremum, 157
- Logarithm, 197
 - Derivative, 198
 - growth, 198
- Lower Bound, 9
- Lower Integral, 217
- Mean Value Theorem, 159
 - Cauchy's Form, 160
- Mertens' Theorem, 88
- Mesh, 214
- Monotone Convergence Theorem, 43
- Monotone Function, 124
 - discontinuity set, 126
 - from the sign of the derivative, 162
 - limit at an endpoint, 125
 - one-sided limits, 124
- Monotone Sequence, 42
- Nested Interval Theorem, 26

- Newton's Approximation, 149
- Null Sequence, 40
 - comparison test, 40
- Ordered Field, 2
- Oscillation, 17
- Partial Sum, 61
- Partition
 - of an Interval, 214
- Pi, 203
- Power
 - Derivative, 200
 - Laws of Exponents, 199
 - Real Exponent, 198
- Power Series, 93
 - Coefficients Are Taylor Coefficients, 187
 - Re-expansion About a New Centre, 188
 - Term-by-Term Differentiation, 185
 - Uniform Convergence on Subintervals, 184
- Pythagorean Identity, 201
- Radius of Convergence, 93
- Ratio Test, 70
- Rational Cut, 270
- Real Numbers, 6, 265
 - Order on, 265
 - Ordered Field Structure, 265
- Rearrangement, 81
 - of an absolutely convergent series, 82
- Refinement
 - of a Partition, 215
- Riemann Integral, 217
- Riemann Sum, 220
 - equivalence with Darboux sums, 220
- Riemann's Rearrangement Theorem, 83
- Riemann-Stieltjes Integral, 251
 - existence for continuous integrands, 255
 - integrability criterion, 254
 - reduction to a Riemann integral, 257
- Rolle's Theorem, 159
- Root
 - n -th, 19
- Root Function, 138
- Root Test, 70
 - Compared with the Ratio Test, 71
- Sequence, 34
 - bounded, 36
- Sequential Criterion for Functional Limits, 108
- Series, 61
 - of Nonnegative Terms, 63
- Squeeze Theorem, 38
- Subsequence, 46
- Subsequential Limit, 46
- Substitution, 243
- Summation by Parts, 79
- Supremum, 9
 - algebra of, 13
 - Epsilon-Characterization, 11
 - of a function, 15
- Tagged Partition, 220
- Taylor Polynomial, 169
- Taylor's Theorem, 171
 - convergence under bounded derivative growth, 210
 - Lagrange remainder, 171
- Thomae's Function, 122
 - Riemann integral of, 232
- Triangle Inequality, 16
- Trigonometric Functions
 - Addition Formulas, 202
 - Cosine, 96
 - Derivatives, 195
 - Periodicity, 204
 - Sine, 96
- Uncountability of $\mathbb{R}$, 27
- Uniform Continuity, 140
 - Extension to the Closure, 143
 - on a Closed Bounded Interval, 141
- Uniform Convergence, 174
 - and Continuity, 176
 - and Differentiation, 179
 - and Integration, 235
- Upper Bound, 9
- Upper Integral, 217
- Weierstrass M-Test, 178

Bibliography

- [1] Nathanael Chwojko-Srawley. *Everything You Need To Know About Metric and Function Spaces*. Bootstrapped 2026-07-29 to metric-function-spaces/ from NateElemAnalysis Part II. Series home for metric completeness, the Banach contraction principle, Stone-Weierstrass, Arzela-Ascoli and Baire category; see DECISIONS 2026-07-29. 2026.
- [2] Nathanael Chwojko-Srawley. *Everything You Need To Know About Multivariable Calculus*. Bootstrapped 2026-07-29 to calculus/. Owns partition-of-unity stratum A (smooth partitions on subsets of Euclidean n-space, Spivak Thm 3-11). Split from the forms/Stokes material now in NateCalcOnManifolds; see DECISIONS 2026-07-29. 2026.
- [3] Nathanael Chwojko-Srawley. *Everything You Need To Know About Topology*.
- [4] Nathanael Chwojko-Srawley. *Preliminary Chapter*. 2023. URL: /Users/nathanaelchwojko-srawkey/website-nate/nate-website/src/assets/latex/notes/preliminary_chapter.
- [5] Nathanael Chwojko-Srawley. *Everything You Need To Know About Real Analysis*. 2026. URL: <https://nathanaelsrawley.com/assets/pdfs/books/EYNTKA-real-analysis.pdf>.
- [6] Nathanael Chwojko-Srawley. *Everything You Need To Know About Harmonic Analysis*. 1st ed. Vol. 1. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_harmonic_analysis.pdf.