

Riemann Roch Generalized

Contents

1	Poincaré-Hopf Formula	4
2	About Normalization	5
3	The Riemann-Roch Theorem	9
3.1	Dimension of $\Omega^1(C)$	12
3.2	Two Important Theorems	18
3.3	Riemann Inequality	20
4	Application of Riemann-Roch Theorem	24
4.1	Genus 0	25
4.2	Genus 1	25
4.3	Genus ≥ 2 : Canonical Map	30
4.4	Hyperelliptic Compact Riemann Surfaces	32
4.5	Genus 2 always hyperelliptic	36
4.6	Genus 3	36
4.7	Genus 4	38
4.8	Abel's Theorem and its Applications	39
4.9	Applications of Abel's Theorem	41
5	Riemann Roch in Modern Language	43
6	Generalized Riemann Roch: Vector Bundles	43
7	Terms for G-bundles	46
8	Some Hodge Theory	46
9	L-Genus and Todd Genus	48
9.1	Todd Class Fixes Commutative Diagram	48

9.2	Multiplicative sequences	49
9.3	Todd Genus	54
9.4	Pontryagin and Chern Numbers	58
10	Some Cobordism	59
10.1	Index and Virtual index	63
11	RR for algebraic manifolds (i.e. projective verities)	63

Consider the Gauss-Bonnet theorem ([this](#)):

$$\int_M K dA = \int_{\partial M} k_g ds = 2\pi\chi(M)$$

where M is a 2-dimensional compact Riemann manifold, K the Gaussian curvature, k_g the geodesic curvature, and $\chi(M)$ the Euler characteristic. This can be thought of as a special case of The Riemann-Roch Theorem!

There is also a more direct topological generalization of the Gauss-Bonnet theorem known as the Chern theorem, or Chern–Gauss–Bonnet theorem, to $2n$ -dimensional Riemann manifolds without boundary:

$$\chi(M) = \int_M e(\Omega)$$

where $\chi(M)$ is the Euler characteristic, Ω is the curvature form (or Levi-Civita connection), and $e(\Omega)$ is the Euler class:

$$e(\Omega) = \frac{1}{(2\pi)^n} \text{Pf}(\Omega)$$

where Pf is the *Pfaffian*; namely as Ω can always be represented as a skew-symmetric matrix (when the dimension of M is even, hence the $2n$) it can always be written as the square of a polynomial in matrix entries, namely if A is the skew-symmetric matrix:

$$\text{Pf}(A)^2 = \det(A)$$

for example:

$$A = \begin{pmatrix} 0 & a \\ -a & b \end{pmatrix} \quad \text{Pf}(A) = a$$

$$B = \begin{pmatrix} 0 & a & b & c \\ -a & 0 & d & e \\ -b & -d & 0 & f \\ -c & -e & -f & 0 \end{pmatrix} \quad \text{Pf}(B) = af - be + dc$$

Atiyah–Singer index theorem seems to generalize what I’m working on further.

Rapid fire concepts we know:

- We shall take varieties and schemes for granted

- We shall be working with algebraic curves, usually algebraic plane curves; hence curves given by $f(x, y)$ for coordinate functions x, y .
- Such curves are projective ([2]). If they are smooth, we naturally ask about their relation to one-dimension complex manifolds; these are called Riemann surfaces (as they are 2 real dimensional)
- Compact Riemann surfaces are classified up to homeomorphism by their genus, and $\chi(X) = 2 - 2g$ where χ is the Euler Characteristic.
- Via normalization, we have the following (This is in fact proven in [5])

Theorem 0.1: Characterizing Riemann Surfaces Using Normalization

Any compact Riemann surface $\tilde{C}$ can be obtained through the normalization of a certain plane algebraic curve C with at most ordinary double points, i.e. There exists a holomorphic mapping $\sigma : \tilde{C} \rightarrow \mathbb{P}^2(\mathbb{C})$ such that $\sigma(\tilde{C})$ is an algebraic curve possessing at most ordinary double points

- The geometry polynomial function (by FTC) makes them surjective and n -to-1 except at the ramification points.
- We shall take holomorphic and meromorphic differential forms for granted
- Key result about holomorphic differential forms:

Theorem 0.2: Stokes Theorem For Holomorphic Differentials

Let C be a Riemann surface, $\Omega \subseteq C$ an open set, $\bar{\Omega}$ compact and $\partial\Omega = \gamma$ a piecewise smooth curve, and ω a holomorphic differential defined on an open set containing $\bar{\Omega}$. Then:

$$\int_{\partial\Omega} \omega = 0$$

Proof :

use local coordinates and apply Cauchy's theorem to get

$$\int_{\partial\Omega_i} \omega = 0$$

on the components, which then gluing gives the desired result

- Residue is winding number

1 Poincaré-Hopf Formula

Let C be a compact Riemann surface. Then we know that for any nonconstant $f \in K(C)$ (the set of meromorphic functions on C), we get:

$$\sum_{p \in C} \nu_p(f) = 0 \quad (1)$$

that is, the number of zeros plus the number of poles (counting multiplicity) is zero. Phillip actually has this really cool generalization of the proof that I got in [3], he proves the following:

Theorem 1.1: Residue Theorem For Forms

Let C be a compact Riemann surface and $\omega \in K^1(C)$ a differential form (the K^1 is his notation for diff. form). Then:

$$\sum_{p \in C} \text{Res}_p(\omega) = 0$$

Proof :
exercise

Then you can deduce the usual equation (1) by setting $\omega = \frac{df}{f}$. Now, we can ask the following: for a trivial $\omega \in K^1(C)$, what is:

$$\sum_{p \in C} \nu_p(\omega)$$

A general differential form on a compact Riemann surface has more flexibility than functions, and the order at p is certainly well-defined the same way the residue of a differential form is well defined, so this is a meaningful question. As one-dimensional differential forms are co-vector fields, this is equivalent to a problem on vector-fields, and we know that one interpretation of the Euler-characteristic on surfaces is to count the zero-points (which are sink and source points) and the saddle points (notice how closely this naturally relates to curvature; this is why it comes up naturally in the Gauss-Bonnet Theorem) and so we get:

$$\sum_{p \in C} \nu_p(\omega) = -\chi(C)$$

- There is an interesting word on the schematic representation of curves, [5, p.28]
- Note that meromorphic functions on complex manifold need not be able to map to $\mathbb{CP}^1$, consider z/w from $\mathbb{C}^2$.
- The end of chapter I, section 10, has a very very simple proof on when one can embed a Riemann surface into $\mathbb{CP}^n$!

2 About Normalization

- He starts with a general plane curve, defines singular points to be when the three partials are zero (hence no linear term when translated to $[0 : 0 : 1]$, or in his notation $[1 : 0 : 0]$). We can then consider any curve $f(x, y) = F(1, x, y)$ when considering the canonical embedding $(x, y) \mapsto [1 : x : y]$. Then it must be double/triple/tuple if the lowest degree is $\geq 2, \geq 3, \geq k$. It is ordinary if all the tangent lines are distinct.
- We can now consider irreducible components. If it is over a UFD, then the GCD is well-defined and so we get that f, g have no common factors if $(f) + (g) = (1)$ so that if $(f) + (g) = (h)$ then $f = Fh$ and $g = Gh$ so that $fG = gF$. We can detect this using the resultant, and hence we can detect singular points over non-characteristic 0 fields using the discriminant. We can then proceed to show that if C is a curve, S the singular points, and $C^* = C \setminus S$ the complement of the singular points, then C^* is connected, not necessarily compact, S is finite.
- Next comes normalization which he defines as follows:

Definition 2.1: Normalization

Suppose C is an irreducible plane algebraic curve, S the set of its singular points. Then there exists a compact Riemann surface $\tilde{C}$ and a holomorphic mapping

$$\sigma : \tilde{C} \rightarrow \mathbb{CP}^2$$

such that

1. $\sigma(\tilde{C}) = C$
2. $\sigma^{-1}(S)$ is a finite set
3. $\sigma : \tilde{C} \setminus \sigma^{-1}(S) \rightarrow C \setminus S$ is injective

Then $(\tilde{C}, \sigma)$ is the normalization of C , and we write just $\tilde{C}$ if it is unambiguous in context.

- We get the following result equivalent to birational maps:

Lemma 2.2: Dense Injective, then Biholomorphic

Suppose $\tilde{C}, \tilde{C}'$ are Riemann surfaces and

$$h : \tilde{C} \rightarrow \tilde{C}'$$

is a surjective holomorphic mapping which is injective on a dense open subset of $\tilde{C}$. Then h is a biholomorphic mapping.

Proof :

Prove it is biholomorphic in every neighbourhood.

- Then

Theorem 2.3: Uniqueness of Normalization

The normalization of an algebraic curve C is unique up to isomorphism (it commies in the way you expect)

Proof :

use above lemma.

The goal now is to prove a normalization always exists.

- Next Griffith's defines $\mathbb{C}\{x\}$ which is not quite $\mathbb{C}[[x]]$ because it is the set of all *convergent* power series of the form $\sum_k a_k x^k$. Similarly for $\mathbb{C}\{x, y\}$ which is convergent $\sum_{i,j} a_{ij} x^i y^j$. Then a Weierstrass polynomial is a polynomial of the form. He denotes the later by $\mathcal{O}$.

$$w = y^d + a_1(x)y^{d-1} + \cdots + a_d(x)$$

where

$$a_j(x) \in \mathbb{C}\{x\} \quad a_j(0) = 0 \quad j = 1, \dots, d$$

Then he goes on to prove the *Weierstrass Preparation theorem* which say that under the appropriate generic conditions, any $f \in \mathcal{O}$ can be written as $f = u \cdot w$ where u is a unit (so $1/u \in \mathcal{O}$ and w is a Weierstrass polynomial.

- Proving this, we get the following great corollary:

Corollary 2.4: Ring Structure of Analytic Func

$\mathcal{O}$ is a UFD, hence any $f \in \mathcal{O}$ can be expressed up to unit u as:

$$f = u f_1 f_2 \cdots f_n$$

where each f_i is irreducible in $\mathcal{O}$.

Proof :

Note that proving $\mathbb{C}\{x\}$ is a UFD is easy because any $f(x) = x^m h(x)$ with $h(x)$ a unit with nonzero constant. For the rest checkout [5].

- An important realization is that being irreducible in $\mathbb{C}\{x\}[y]$ and $\mathbb{C}\{x, y\}$ is the same thing.
- Next, let's take an irreducible curve, center the point of interest at $(0, 0)$, represent it as

$$y^n + a_1(x)y^{n-1} + \cdots + a_d(x)$$

Then locally at 0, we may have multiple local irreducible components. Let's take the curve over $\mathbb{C}^2$. Then around 0, if there are multiple components, it will look like multiple disks that are glued together at a point.

- Using this, we can locally normalize, and then use that to globally normalize
- Next Griffith looks at intersection numbers and Bézout Theorem.

Ramification Divisors and the Riemann-Hurwitz Formula

Let $\nu_f(p)$ be the order of a point of a holomorphic mapping $f : C \rightarrow C'$ between Riemann surfaces. As such a map is always surjective if it is non-constant, we have $\nu_f(p) \geq 1$. It ramifies if $\nu_f(p) > 1$, and as ramified points are isolated we know that $\{p \in C \mid \nu_f(p) > 1\}$ is finite. Thus we have the well-defined:

Definition 2.5: Ramification Divisor

The *ramification divisor* of f is :

$$R = \sum_{p \in C} (\nu_f(p) - 1)p \in \text{Div}(C)$$

Recall that if $D = \sum_{p \in C} n_p \cdot p$ is a divisor, $\deg D = \sum_{p \in C} n_p$.

Theorem 2.6: Riemann-Hurwitz Formula

Let C, C' be compact Riemann surfaces where:

$$\text{genus}(C) = g \quad \text{and} \quad \text{genus}(C') = g'$$

Let $f : C \rightarrow C'$ be a nonconstant holomorphic mapping of degree n . Let R be the ramification divisor of f . Then:

$$\deg R = 2(g + n - ng' - 1)$$

Proof :

First, we shall use the fact that on any compact Riemann surface, there exists a nontrivial meromorphic differential. This could be intuitively seen by the correspondence between compact Riemann surfaces and curves, and on any curve we have a non-trivial meromorphic differential.

Hence, choose $\omega \in K^1(C')$, and let $f^*\omega$ be its pullback. Locally we have:

$$\omega = z^\nu$$

and

$$\omega = g(\omega)d\omega$$

Hence:

$$f^*\omega = \nu g(z^\nu) z^{\nu-1} dz$$

and so the order of each point is:

$$\nu_{f^*\omega}(p) = \nu_f(p)\nu_\omega(f(p)) + \nu_f(p) - 1$$

and so the divisor of $(f^*\omega)$ is :

$$\begin{aligned}
 (f^*\omega) &= \sum_p \nu_f(p) \nu_\omega(f(p)) p + R \\
 &= \sum_q \nu_\omega(q) \left(\sum_{f(p)=q} \nu_f(p) p \right) + R \\
 &= \sum_q \nu_\omega(q) f^{-1}q + R \\
 &= f^{-1} \left(\sum_q \nu_\omega(q) q \right) + R = f^{-1}((\omega)) + R
 \end{aligned}$$

Thus:

$$\deg(f^*\omega) = \deg f^{-1}((\omega)) + \deg R$$

Then from the Poincaré-Hopf formula:

$$\deg(f^*\omega) = 2g - 2 \quad \deg(\omega) = 2g' - 2$$

Thus directly computing we get:

$$\begin{aligned}
 \deg f^{-1}((\omega)) &= \sum_q \nu_\omega(q) \left(\sum_{f(p)=q} \nu_f(p) \right) \\
 &= \deg(\omega) \deg f \\
 &= (2g' - 2)n
 \end{aligned}$$

then substituting we get:

$$2g - 2 = (2g' - 2)n + \deg R$$

re-arranging gives our desired result.

From this we can immediately conclude:

Corollary 2.7: Holomorphic Map and Genus

Let $f : C \rightarrow C'$ be a holomorphic map between genus surfaces of genus g, g' . Then if $n \geq 1$:

$$g \geq g'$$

Proof :

Apply the formula.

Corollary 2.8: Compact Riemann Surface As Algebraic Equation

Let C be a compact Riemann surface and let

$$x, y : C \rightarrow \mathbb{CP}^1$$

be holomorphic mapping where $\deg x = d$. Then there exists a polynomial equation f such that $f(x, y) = 0$.

Proof :

[5, p.94]

Next, we cover the genus formula.

Theorem 2.9: (Arithmetic) Genus Formula

Let $C \subseteq \mathbb{CP}^2$ be an irreducible algebraic curve of degree d with δ singular points all being ordinary double points, and let $\sigma : \tilde{C} \rightarrow C$ be its normalization where $\text{genus}(\tilde{C}) = g$. Then:

$$g = \frac{(d-1)(d-2)}{2} - \delta$$

The term “arithmetic” is because it relies on the arithmetic information. There is also the *geometric genus formula* which relies on the dimension of (co)homology groups.

Proof :

[5, p.95].

3 The Riemann-Roch Theorem

Let C be a compact Riemann surface. Define the divisor to be the formal sum:

$$D = \sum_i^k n_i p_i \in \text{Div}(C)$$

A divisor is called *effective* if each coefficient is ≥ 0 . and we write $D \geq 0$. Any divisor can be expressed as the difference of two effective divisors. We now define the Riemann-Roch space:

$$\mathcal{L}(D) = \{f \in K(C) \mid (f) \geq -D\}$$

That is, $\mathcal{L}(D)$ contains all $f \in K(C)$ whether f has a pole of multiplicity at most n_i at p_i if $n_i > 0$ and a zero of multiplicity at least $|n_i|$ at p_i if $n_i < 0$. We can then define:

$$l(D) = \dim \mathcal{L}(D)$$

We then define a similar notion for divisors:

$$K^1(D) = \{\omega \in K^1(C) \mid (\omega) \geq D\}$$

where now, if $n_i < 0$ then ω can have a p_i as a pole of multiplicity at most $|n_i|$ and if $n_i > 0$ then ω must have a p_i as a zero of multiplicity at least n_i . Again, $K^1(D)$ is a vector space and :

$$\dim K^1(D) = i(D)$$

We shall usually be focusing on effect divisors; $D \geq 0$; in that case we have that $\mathbb{C} \subseteq \mathcal{L}(D)$. Furthermore, to be more align with the usual differential geometry notation, we shall write $\Omega^1(D)$ instead of $K^1(D)$. Then for $D \geq 0$, $\omega \in \Omega^1(D)$ means ω is holomorphic and has a zero of appropriate multiplicity at p_i for appropriate points given by D .

Definition 3.1: Linearly Equivalent

Let $D, E \in \text{Div}(C)$. Then D, E are *linearly equivalent* if there exists a $f \in K(C)$ such that

$$(f) = D - E$$

and we write $D \sim E$.

We can also write $D = (f) + E$ for some (f) , and hence we are really saying that $D \sim E$ if they are equal up to principal divisor. Hence, two divisor are linearly equivalent if they belong to the same coset in $\text{Div}(C)/\text{PDiv}(C)$.

Proposition 3.2: Dimension and Linear Equivalence

Let $D, E \in \text{Div}(C)$, $D \sim E$. Then:

$$\mathcal{L}(D) \cong \mathcal{L}(E) \quad \Omega^1(D) \cong \Omega^1(E) \quad \deg D = \deg E$$

Proof :

Say $D \sim E$. Then there exists an $f \in K(C)$ such that

$$(f) = D - E$$

Now pick $g \in \mathcal{L}(D)$ so that $(g) + D \geq 0$. Then:

$$(fg) + E = (f) + (g) + E = D - E + (g) + E = (g) + D \geq 0$$

Thus, there is a mapping:

$$\mathcal{L}(D) \rightarrow \mathcal{L}(D) \quad g \mapsto fg$$

This mapping is certainly linear, and it is bijective as the inverse is:

$$\mathcal{L}(E) \rightarrow \mathcal{L}(D) \quad g \mapsto \frac{g}{f}$$

Thus, $\mathcal{L}(D) \cong \mathcal{L}(E)$. Similarly, we certainly get that $\Omega^1(D) \cong \Omega^1(E)$. To show the degree's are the same, recall that for any $f \in K(C)$:

$$\sum_{p \in C} \nu_p(f) = 0$$

Then:

$$\sum \nu_p(f) = \deg(f) = \deg D - \deg E$$

hence $\deg E = \deg D$, as we sought to show.

Proposition 3.3: Brill-Noether Reciprocity

Let $\omega \in \Omega^1(C)$ and $(\omega) = D + E$. Then:

$$\mathcal{L}(D) \cong \Omega^1(E) \quad \mathcal{L}(E) \cong \Omega^1(D)$$

This can also be seen as a special case of Serre Duality (see [2])

Proof :

Let $f \in \mathcal{L}(D)$ so that $(f) \geq -D$. Then:

$$(f\omega) = (f) + (\omega) \geq D + D + E = E$$

Thus define:

$$\mathcal{L}(D) \rightarrow \Omega^1(E) \quad f \mapsto f\omega$$

which is certainly a vector-space homomorphism with inverse:

$$\Omega^1(E) \rightarrow \mathcal{L}(D) \quad \varphi \mapsto \frac{\varphi}{\omega}$$

showing they are isomorphic. By symmetry, we have the similar isomorphism by swapping E and D , completing the proof.

Definition 3.4: Laurent Principal Part

Let $\sum_{i=-n}^{\infty} a_i z^i$ be a Laurent series. Then its singular part

$$\sum_{i=-n}^{-1} a_i z^i$$

is the *Laurent principal part*, or just *principal part*. If $a_{-n} \neq 0$, then N is called the order of the Laurent principal part.

Now suppose C is a Riemann surface, D an effective divisor. Let $z_i, i = 1, 2, \dots, k$ be local coordinates near p_i and $z_i(p_i) = 0$. Then if for every p_i , there is given a Laurent principal part of order n_i :

$$\eta_i = \frac{a_{in_i}}{z_i^{n_i}} + \dots + \frac{a_{i1}}{z_i}$$

we can ask the following question: under what condition does there exist an $f \in K(C)$ such that in a neighbourhood of each p_i , f has η_i as a Laurent principal part? This is called the *Mittag-Leffler problem*, and answering it will allow us to deduce the Riemann-Roch Theorem.

First off, a meromorphic function on a compact Riemann surface C is up to a constant uniquely determined by its Laurent principal part, for if $f_1, f_2 \in K(C)$ that have the same Laurent principal part then $f_1 - f_2$ is holomorphic on C , and hence is constant. Thus, we may identify $\mathcal{L}(D)/\mathbb{C}$ as a subspace of $\mathbb{C}^d$, $d = \deg D$, as follows. Fix $D = \sum_i^k n_i p_i \geq 0$, and let z_i be a coordinate function around p_i where $z_i(p_i) = 0$. Then for $f \in \mathcal{L}(D)$, let η_i be the principal part at p_i . Then we know

that:

$$\eta_i = \frac{a_{in}}{z^{n_i}} + \cdots + \frac{a_{i2}}{z_i^2} + \frac{a_{i1}}{z_i}$$

where $a_{in_i}, \dots, a_{i1} \in \mathbb{C}$. Now let P_i be the n_i -dimensional vector space over $\mathbb{C}$ spanned by $\{1/z_i^{n_i}, \dots, 1/z_i^2, 1/z\}$. Then we have the injective vector-space homomorphism:

$$\frac{\mathcal{L}(D)}{\mathbb{C}} \rightarrow P_1 \oplus \cdots \oplus P_k \quad [f] \mapsto (\eta_1, \dots, \eta_k)$$

Then by identifying each P_i with $\mathbb{C}^{n_i}$, we get that this space is a subspace of $\mathbb{C}^d$. Note that surjectivity is not so obvious or even necessary because we don't know if these maps are compatible (ex. the gluing conditions can impose restrictions).

3.1 Dimension of $\Omega^1(C)$

The goal is to show the following:

Theorem 3.5: Dimension of Differential Forms

Let C be a compact Riemann surface of genus g . Then:

$$\dim \Omega^1(C) = g$$

Note that this is easy in the case where $g = 0$, this is the Riemann sphere and we showed that the only holomorphic differential forms on the Riemann sphere are constant. The proof will proceed in two parts:

Proposition 3.6: Dimension of Differential Form: I

Let C be a compact Riemann surface of genus g . Then:

$$\dim \Omega^1(C) \geq g$$

The key idea is that we can “embed” the space of homogeneous functions of the right degree (minus the singularities) into $\Omega^1(C)$ by essentially taking it's differential with a bit of normalization.

Proof :

First, by normalization, for a compact Riemann surface C , there exists a holomorphic mapping $\pi : C \rightarrow \mathbb{P}^2$ such that $C' = \pi(C)$ is an irreducible algebraic curve of degree d in $\mathbb{P}^2$ possessing δ ordinary double points, say $p_1, \dots, p_\delta$, and no other singular points. Then if $d = 1$ or $d = 2$, then by the genus formula $g = 0$. Thus we shall assume $d \geq 3$.

Now suppose C is mapped holomorphically onto $\mathbb{P}^2$. We may apply a transformation of coordinates so that C' is not tangent to the line at infinity L_∞ in $\mathbb{P}^2$, that L_∞ does not pass through the points $p_1, \dots, p_\delta$, and that C' does not pass through $[0 : 0 : 1]$ (so the origin, $(0, 0)$). Indeed, these conditions can be guaranteed due to the following. We can guarantee that C' and L_∞ intersect at d distinct points for we can always find a linear transformation so that C' intersects a line at d distinct points. Taking this line to be L_∞ , suppose C' and L_∞ are tangent. Then at the point of tangency, their intersection number would be 2, and so we get by Bézout that $(C' \cdot L_\infty) \geq d + 1$, a contradiction. We can then do a similar reasoning to guarantee L_∞ does not pass through $p_1, \dots, p_\delta$.

Finally, for the $[0 : 0 : 1]$ condition, take a point which is on L_∞ but not on C' , and apply a transformation so that the point becomes $[0 : 0 : 1]$. Then we got our desired coordinates.

We can now write out our affine equation for C' such that $f(x, y) = 0$. Let $\Gamma = p_1 + \dots + p_\delta$, and let S^{d-3} represent the set of homogeneous polynomials with complex coefficients of degree $d - 3$ in three variables. Let:

$$S^{d-3}(-\Gamma) = \{G(\zeta^0, \zeta^1, \zeta^2) \in S^{d-3} \mid G(p_i) = 0, \forall i = 1, 2, \dots, \delta\}$$

Note that by some counting arguments we get:

$$\dim S^{d-3} = \frac{(d-1)(d-2)}{2}$$

On the other hand, as the collection of polynomials that vanish at p_i is equivalent to the fact that the coefficients of this polynomial satisfy a linear equation, we get:

$$\dim S^{d-3}(-\Gamma) \geq \frac{(d-1)(d-2)}{2} - \delta \stackrel{!}{=} g$$

where $\stackrel{!}{=}$ comes from the genus formula, theorem 2.9. Thus, if we show there is an injective homomorphism from $S^{d-3}(-\Gamma)$ into $\Omega^1(C)$, we have that:

$$\dim \Gamma^1(C) \geq \dim S^{d-3}(-\Gamma) = g$$

To that end, for any $G \in S^{d-3}(-\Gamma)$. We need to find a form $\omega_G \in \Omega^1(C)$ for it to map to. What we will do is find a pullback of a form α along π . Let $g(x, y) = G(1, x, y)$, $U_0 = \{[1, x, y] \in \mathbb{P}^2\}$ and $U_1 = \{[u, 1, v] \in \mathbb{P}^2\}$. Define the following:

$$\alpha_g = \begin{cases} \frac{g(x, y) dx}{\frac{\partial f(x, y)}{\partial y}} \Big|_{C'}, & \text{on } C' \cap U_0, \text{ when } \frac{\partial f(x, y)}{\partial y} \Big|_{C'} \neq 0 \\ -\frac{g(x, y) dy}{\frac{\partial f(x, y)}{\partial x}} \Big|_{C'}, & \text{on } C' \cap U_0 \text{ when } \frac{\partial f(x, y)}{\partial y} \Big|_{C'} = 0 \\ -\frac{g(\frac{1}{u}, \frac{v}{u}) \frac{dv}{u^2}}{\frac{\partial f}{\partial y}(\frac{1}{u}, \frac{v}{u})} \Big|_{C'}, & \text{on } C' \cap U_1 \end{cases}$$

where $\frac{\partial f}{\partial y}(1/u, v/u)$ is the value of $\frac{\partial f}{\partial y}$ at $(1/u, v/u)$. As C' does not pass through the point $[0, 0, 1]$, α is defined over all of C' . It is in fact uniquely defined. Indeed, inside U_0 the affine equation is $f(x, y) = 0$ and thus on C' :

$$\frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy = 0$$

and thus:

$$\frac{dx}{\frac{\partial f(x, y)}{\partial y}} \Big|_{C'} = -\frac{dy}{\frac{\partial f(x, y)}{\partial x}} \Big|_{C'}$$

Thus, we have the linear homomorphism:

$$S^{d-3}(-\Gamma) \rightarrow K^1(C) \quad G \mapsto \omega_G = \pi^*(\alpha_g)$$

Note this map is injective since if $\omega_G = 0$, then C' is contained in the curve $g(x, h) = 0$, but this is impossible since $\deg G = d - 3$. We must prove that ω_G is indeed holomorphic, and hence it maps into $\Omega^1(C)$, which shall complete the proof.

To that end, we must show that there are no poles. If a point $p' \in C'$ on $\pi^{-1}(C' \cap U_0)$ can become a pole of ω_G , then:

$$\left. \frac{\partial f(x, y)}{\partial y} \right|_{p'} = \left. \frac{\partial f(x, y)}{\partial x} \right|_{p'} = 0$$

where $p' = \pi(p)$. Then p' is a double point of C' , so $p' \in \Gamma$, so

$$g(x, y)|_{p'} = 0$$

But p' is a second-order zero of F , and hence a first order zero of $\partial f / \partial x$, so:

$$\alpha|_{p'} = - \left. \frac{g(x, y) dy}{\frac{\partial f(x, y)}{\partial x}} \right|_{C'} \neq \infty$$

hence ω_G is holomorphic at p .

Now consider ω_G on $\pi^{-1}(C' \cap L_\infty)$. Then by definition of α :

$$\alpha = - \left. \frac{u^{d-3} g\left(\frac{1}{u}, \frac{v}{u}\right) du}{u^{d-1} \frac{\partial f}{\partial y}\left(\frac{1}{u}, \frac{v}{u}\right)} \right|_{C'}$$

to simplify notation, denote the denominator as $h(u, v)$. Then we must show $h(u, v) \neq 0$ on $C' \cap L_\infty$. To see this, note that the equation of C' on U_1 is

$$k(u, v) = u^d f\left(\frac{1}{u}, \frac{v}{u}\right) = 0$$

And notice:

$$\frac{\partial k(u, v)}{\partial v} = u^d \frac{\partial f}{\partial u}\left(\frac{1}{u}, \frac{v}{u}\right) \cdot \frac{1}{u} = h(u, v)$$

With this setup, suppose $p' \in C' \cap L_\infty$ such that

$$h(u, v)|_{p'} = 0$$

Then since p' is not a double point of C' , it must be that :

$$\left. \frac{\partial k(u, v)}{\partial u} \right|_{p'} \neq 0$$

Since $p' \in L_\infty$, the coordinates of p' must be of the form $[0, 1, v]$. Then we can write the tangent line to C' at p' as :

$$\left. \frac{\partial k(u, v)}{\partial u} \right|_{p'} (u - 0) = 0$$

But then the tangent line is simply L_∞ , which contradicts our choice of coordinates, hence no such point exists, and ω_G is holomorphic, as we sought to show.

Let us now show the other direction of $\dim \Omega^1(C) \leq g$. This direction shall require some differential geometry reasoning.

Let C be a compact Riemann surface of genus g with $\gamma_1, \gamma_2, \dots, \gamma_{2g-1}, \gamma_{2g}$ framing a $\mathbb{Z}$ -basis of the homology group $H_1(C, \mathbb{Z})$. Then if γ is a closed curve on C ,

$$\gamma = \sum_{i=1}^{2g} n_i \gamma_i + \partial\Omega$$

where Ω is a region on C . Let $[\gamma]$ represent the homology class containing γ .

Proposition 3.7: Natural Dual of Homology Group

Let γ be a closed differential one-form on C . Then the mapping:

$$\eta_\lambda : H_1(C, \mathbb{Z}) \rightarrow \mathbb{C} \quad [\gamma] \mapsto \int_\gamma \lambda$$

is well-defined

Proof :

The key is that γ is closed. If γ' is homologous to γ then $\gamma - \gamma' = \partial\Omega$ where Ω is a region on C . By Stokes theorem

$$\eta_\lambda(\gamma) - \eta_\lambda(\gamma') = \int_{\gamma - \gamma'} \lambda = \int_{\partial\Omega} \lambda = \iint_\Omega d\lambda = 0$$

Proposition 3.8: Exactness

Let λ be a closed differential one-form on C . Then if for all i ,

$$\int_{\gamma_i} \lambda = 0$$

then λ is exact.

Proof :

We must find f such that $df = \lambda$. Let $p_0 \in C$ be some fixed point. Then for any point $p \in C$, choose a path γ from p_0 to p and define:

$$f(p) = \int_\gamma \lambda$$

If this is well-defined, this is the function we need. Indeed, if γ' is another path from p_0 to p then $\gamma - \gamma'$ is a closed curve on C , and hence:

$$\gamma - \gamma' = \sum_i^{2g} n_i \gamma_i + \partial\Omega$$

where $ni \in \mathbb{Z}$ and Ω is a region on C . Then as:

$$\int_{\gamma_i} \lambda = 0 \quad \forall i = 1, \dots, 2g$$

and

$$\int_{\partial\Omega} \lambda = \iint_{\Omega} d\lambda = 0$$

we get:

$$\int_{\gamma} \lambda = \int_{\gamma'} \lambda$$

Thus, f is a well-defined smooth function on C , and by the FTA we get $df = \lambda$, as we sought to show.

Proposition 3.9: Cauchy Riemann For Differential Forms

Let C be a compact Riemann surface, and $\omega, \varphi \in \Omega^1(C)$. Then if:

$$\omega + \bar{\varphi} = df$$

then

$$\omega = \varphi = 0$$

Proof :

Let $\omega = h(z)dz$ and $\varphi = g(z)dz$. Then as C is one-dimensional:

$$\omega \wedge \varphi = 0$$

and

$$\frac{i}{2} \varphi \wedge \bar{\varphi} = |g(z)|^2 \frac{i}{2} dz \wedge d\bar{z} = |g(z)|^2 du \wedge dv$$

We shall show $\varphi = 0$. For the sake of contradiction suppose $\varphi \neq 0$. Then:

$$\frac{i}{2} \iint_C \varphi \wedge \bar{\varphi} = \iint_C |g(z)|^2 du \wedge dv > 0$$

But then:

$$\varphi \wedge \bar{\varphi} = \varphi \wedge \omega + \varphi \wedge \bar{\varphi} = \varphi \wedge (\omega + \bar{\varphi}) = \varphi \wedge df$$

If we show $\iint_C \varphi \wedge df = 0$, we are done. Indeed, notice that:

$$d(f\varphi) = df \wedge \varphi + f \wedge d\varphi$$

As $\varphi \in \Omega^1(C)$, g is holomorphic, and so:

$$\frac{\partial g}{\partial \bar{z}} = 0$$

Thus:

$$\begin{aligned} d\varphi &= d(gdz) = dg \wedge dz \\ &= \left(\frac{\partial g}{\partial z} dz + \frac{\partial g}{\partial \bar{z}} d\bar{z} \right) \wedge dz = 0 \end{aligned}$$

Hence:

$$\varphi \wedge df = -df \wedge \varphi = -d(f\varphi)$$

Note that $f\varphi$ is a differential one-form, so:

$$\iint_C d(f\varphi) = 0$$

But then:

$$\iint_C \varphi \wedge df = - \iint_C d(f\varphi) = 0$$

By a similar argument $\omega = 0$, completing the proof.

Proposition 3.10: Dimension of Differential Form: II

Let C be a compact Riemann surface of genus g . Then:

$$\dim \Omega^1(C) \leq g$$

Proof :

For the sake of contradiction suppose $\dim \Omega^1(C) \geq g + 1$ so that there are $g + 1$ elements $\omega_1, \dots, \omega_{g+1} \in \Omega^1(C)$ that are $\mathbb{C}$ -linearly independent. Take the equations:

$$\int_{\gamma_i} \sum_{j=1}^{g+1} (\lambda^j \omega_j + \eta^j \bar{\omega}_j) = 0 \quad i = 1, 2, \dots, 2g$$

where each γ_i is given by choosing a $\mathbb{Z}$ -basis for $H_1(C, \mathbb{Z})$. By treating $\lambda^1, \dots, \lambda^{g+1}, \eta^1, \dots, \eta^{g+1}$, then the above is $2g$ linear equations with $2g + 2$ unknowns. This system must then have a trivial solutions, suppose $\lambda_0^1, \dots, \lambda_0^{g+1}, \eta_0^1, \dots, \eta_0^{g+1}$ is a solution. Then choosing:

$$\omega = \sum_{j=1}^{g+1} \lambda_0^j \omega_j \in \Omega^1(C) \quad \varphi = \sum_{j=1}^{g+1} \overline{\eta_0^j} \omega_j \in \Omega^1(C)$$

then for all $i = 1, 2, \dots, 2g$:

$$k \int_{\gamma_i} (\omega + \bar{\varphi}) = \int_{\gamma_i} \sum_{j=1}^{g+1} (\lambda_0^j \bar{\omega}_j + \eta_0^j \omega_j) = 0$$

This implies it is exact, hence there exists a C^∞ function f such that

$$\omega + \bar{\varphi} = df$$

But then by proposition 3.9 $\omega = \varphi = 0$. But then as $\lambda_0^1, \dots, \lambda_0^{g+1}, \eta_0^1, \dots, \eta_0^{g+1}$ is a non-trivial solution so that they are not all 0, this contradicts the linear independence of $\omega_1, \dots, \omega_{g+1}$, as we sought to show.

Theorem 3.11: Dimension of Differential Forms

Let C be a compact Riemann surface of genus g . Then:

$$\dim \Omega^1(C) = g$$

Proof :

combine proposition 3.6 and proposition 3.10.

3.2 Two Important Theorems

Let $H_1(C, \mathbb{Z})$ be the collection of closed forms of a genus g Riemann surface. It will have a basis spanned by $2g$ elements. We have a natural dual definition:

Definition 3.12: 1st Cohomology Group

Let C be a compact Riemann surface. Then the first cohomology group of C is:

$$H^1(C, \mathbb{C}) = \text{Hom}_{\mathbb{C}}(H_1(C, \mathbb{Z}), \mathbb{C})$$

Then we saw in proposition 3.7 that we have a map:

$$\begin{aligned} \eta : \{\text{closed differential one-forms on } C\} &\rightarrow H^1(C, \mathbb{C}) \\ \lambda &\mapsto \eta_\lambda \end{aligned}$$

and furthermore this map is injective by proposition 3.8. Let $H_{\text{DR}}(C)$ be the equivalence classes of closed one forms mod exact one forms. Then by proposition 3.8, we have in fact that the above η is a map from $H_{\text{DR}}(C) \hookrightarrow H^1(C, \mathbb{C})$. Then, by the ‘‘Hodge’’ condition, we have that:

$$\begin{aligned} \Omega^1(C) \oplus \overline{\Omega^1(C)} &\hookrightarrow H_{\text{DR}}(C) \\ \omega \oplus \bar{\varphi} &\mapsto \omega + \bar{\varphi} \end{aligned}$$

is also injective. We thus get a triple of injective maps:

$$\Omega^1(C) \oplus \overline{\Omega^1(C)} \hookrightarrow H_{\text{DR}}(C) \hookrightarrow H^1(C, \mathbb{C})$$

Note that the vector spaces on both ends are $2g$, and hence all these spaces are isomorphic! The first isomorphism is a special case of the *Hodge Theorem*, and the second isomorphism is a special case of *DeRham’s Theorem*.

Definition 3.13: Period Vector

Let C be a compact Riemann surface of genus g , $\omega_1, \dots, \omega_g$ a basis for $\Omega^1(C)$, and $\gamma_1, \dots, \gamma_{2g}$ a $\mathbb{Z}$ -basis for $H_1(C, \mathbb{Z})$. Then the $2g$ g -dimensional vector:

$$\pi_i = \begin{pmatrix} \int_{\gamma_i} \omega_1 \\ \vdots \\ \int_{\gamma_i} \omega_g \end{pmatrix}$$

are called a *system of period vectors of C* . The matrix:

$$\Omega = (\pi_1, \dots, \pi_{2g})_{g \times 2g}$$

is called the *period matrix* of C

Proposition 3.14: Period Matrix Nonsingular

The $2g$ period vectors are $\mathbb{R}$ -linearly independent.

Proof :

For the sake of contradiction, suppose $\pi_1, \dots, \pi_{2g}$ are $\mathbb{R}$ -linearly dependent so that there exists $a_1, \dots, a_{2g} \in \mathbb{R}$ not all zero st

$$a_1 \pi_1 + \dots + a_{2g} \pi_{2g} = 0$$

We make take the complex conjugate of this (note conjugating an integral brings the conjugation to the inside) and get the matrix:

$$\Omega^* = \begin{pmatrix} \pi_1 & \dots & \pi_{2g} \\ \bar{\pi}_1 & \dots & \bar{\pi}_{2g} \end{pmatrix}_{2g \times 2g}$$

By our assumption, $\text{Rank}(\Omega^*) < 2g$, hence there is a vector in the kernel

$$\int_{\gamma_i} \left(\sum_{j=1}^g \lambda^j \omega_j + \sum_{j=1}^g \eta^j \bar{\omega}_j \right) = 0 \quad (i = 1, \dots, 2g)$$

Let $\omega = \sum_{j=1}^g \lambda^j \omega_j$ and $\varphi = \sum_{j=1}^g \eta^j \bar{\omega}_j$. Then:

$$\int_{\gamma_i} (\omega + \bar{\varphi}) = 0 \quad (i = 1, \dots, 2g)$$

But then we know $\omega = \varphi = 0$, contradicting the linear independence of $\omega_1, \dots, \omega_g$, as we sought to show.

3.3 Riemann Inequality

Theorem 3.15: Riemann Inequality

Let C be a compact Riemann surface of genus g , $D \in \text{Div}(C)$, and $\deg D = d$. Then;

$$l(D) \geq d - g + 1$$

see [5, p.113] for full details.

Proof :

Like before, let us link Riemann surfaces to curves to get the required context. Let

$$\pi : C \rightarrow C' \subseteq \mathbb{CP}^2$$

be the normalization map with δ ordinary double points $p_1, \dots, p_\delta$. Griffith's offers the following visualization:




Figure 1: normalizatoion

Let $\pi^{-1}(p_i) = p'_i + p''_i$. Then let:

$$\Delta = \sum_{i=1}^{\delta} (p'_i + p''_i) \in \text{Div}(C)$$

Then by a change of coordinates in $\mathbb{CP}^2$, without loss of generality we may assume neither $\pi(D)$ nor $\pi(\Delta)$ contain the point at infinity (like in the above figure). Now, let C' be given by the equation $F(\xi^0, \xi^1, \xi^2) = 0$, $\deg F = m$. Let $D = D' - D''$ where $D', D'' \geq 0$ and let

$$\deg D' = d' \quad \deg D'' = d''$$

Let S^n be the vector-space of complex homogeneous polynomials in three variables of degree n . Then:

$$\dim S^n = \frac{1}{2}(n+1)(n+2)$$

Now let's consider a polynomial $G(\xi^0, \xi^1, \xi^2) \in S^n$ satisfying:

1. $F \nmid G$
2. $G \cdot C \geq D' + \Delta$ where $G \cdot C = (\pi^*(G(\xi^0, \xi^1, \xi^2)))$ (that is, the divisor induced by the zeros of $\pi^*(G)$ on C). Note that $G \cdot C \in \text{Div}(C)$.

We shall elaborate why we would want such a polynomial, then show such a polynomial exists. First, for $p \in C$, let $p' = \pi(p) \in C'$. Then a neighbourhood of p gets mapped under π into a curve component of C' near p' ; denote it by B_p . Then after a possible coordinate change, we may take p' to be $[1 : 0 : 0]$. Label the equation after this coordinate change G_p . As the multiple points of C' are ordinary double points, the equation of B_p near p' is:

$$y = a_0 + a_1x^2 + \dots$$

Putting this into $G_p(1, x, y)$, we get a power series in terms of x . Now, let $n(p)$ denote the degree of the lowest term. We then get:

$$G \cdot C = \sum_{p \in C} n(p)p$$

the right hand side is certainly a finite sum. (I'll finish rest later)

Corollary 3.16: Comparing Degree and Genus

If $d \geq g$, then

$$l(D) \neq 0$$

Hence if $l(D) = 0$, then:

$$d \leq g - 1$$

Proof :

immediate

Proposition 3.17: Meromorphic Differentials and Riemann Inequality

$$i(D) \geq g - d - 1$$

Proof :

Let F, m, D', D'', d', d'' , and Δ be as in theorem 3.15. **finish here.**

Corollary 3.18: Comparing Degree and Genus – Meromorphic Differential

If $d \leq g - 2$, then $i(D) \neq 0$, hence if $i(D) = 0$ then

$$d \geq g - 1$$

Proof :
immediate

Finally this leads us to:

Theorem 3.19: Riemann-Roch Theorem

Let C be a compact Riemann surface of genus g and :

$$D = \sum_{i=1}^k n_i p_i \in \text{Div}(C)$$

Then:

$$l(D) = d - g + i(D) + 1$$

Proof :
see [5, p.118]

Let us first assume $D \geq 0$ and prove

$$l(D) \leq d - g + i(D) + 1$$

And then take cases and judiciously use the Riemann inequality and its corollary to get the results. Suppose $\eta = (\eta_1, \dots, \eta_k)$ where:

$$\eta_i = \frac{a^{in_i}}{z_i^{n_i}} + \dots + \frac{a_{i1}}{z_i}$$

is a Laurent principal part near p_i . The linear space consisting of all these η is isomorphic to $\mathbb{C}^d$. Then thinking of η as a vector in $\mathbb{C}^d$, consider:

$$\begin{aligned} \text{Res} : \mathbb{C}^d \times \Omega^1(C) &\rightarrow \mathbb{C} \\ (\eta, \omega) &\mapsto \sum_{i=1}^k \text{Res}_{p_i}(\eta_i \omega) \end{aligned}$$

This is certainly a bilinear mapping. Let us show that:

$$\text{Res}(\eta, \omega) = 0, \forall \eta \in \mathbb{C}^d \quad \text{if and only if} \quad \omega \in \Omega^1(D)$$

If $\omega \in \Omega^1(D)$, then $\eta_i \omega$ does not have p_i as a pole for all $i = 1, \dots, k$, hence:

$$\text{Res}_{p_i}(\eta_i \omega) = 0$$

and so the sum of residues is zero, giving the $\Leftarrow$ direction. For the $\Rightarrow$ direction, assume otherwise so that $\omega \in \Omega^1(C)$ such that $\text{Res}(\eta, \omega) = 0$ for all $\eta \in \mathbb{C}^d$ but $\omega \notin \Omega^1(D)$. Then near p_i , let

$$\omega = b_{i0} z_i^{m_i} + b_{i1} z_i^{m_i+1} + \dots$$

where $m_i \in \mathbb{Z}$, $b_{i0} \neq 0$. By assumption, $m_i < n_i$. Then there is a Laurent Principal part near p_i

$$\xi_i = (\xi_{i1}, \dots, \xi_{ik})$$

such that

$$\xi_{ii} = \frac{1}{z_i^{m_i+1}} \quad \xi_{ij} = 0 \quad (i \neq j)$$

As $m_i + 1 \leq n_i$, $\mathbb{C}^d$, by treating these as $\mathbb{C}^d$ we have $\xi_i \in \mathbb{C}^d$. However:

$$0 = \text{Res}(\xi_i, \omega) = \text{Res}_{p_i} \frac{b_{i0}}{z_i} = b_{i0} \neq 0$$

a contradiction. Thus, $m_i \geq n_i$, and so $\omega \in \Omega^1(D)$.

Next, recall the following fact (or see exercise ref:HERE). Let $B : V \times W \rightarrow \mathbb{C}$ be a bilinear map between two finite dimensional vector spaces, in particular $\dim V = n$ and $\dim W = m$. Then B is called non-degenerated if given $w \in W$, $B(v, w) = 0$ for all $v \in V$ implies $w = 0$. Then given $L \subseteq V$ satisfies $B(L \times W) = 0$, then:

$$\dim L \leq n - m$$

Then by the property we just found earlier about the residue map, we have a non-nondegenerate bilinear map:

$$\text{Res}_1 : \mathbb{C}^d \times \frac{\Omega^1(C)}{\Omega^1(D)} \rightarrow \mathbb{C}$$

Now recall we can interpret $\mathcal{L}(D)/\mathbb{C}$ as a subspace of $\mathbb{C}^d$, and by what we've just shown:

$$\text{Res}_1 \left(\frac{\mathcal{L}(D)}{\mathbb{C}} \times \frac{\Omega^1(C)}{\Omega^1(D)} \right) = 0$$

Then:

$$\dim \left(\frac{\mathcal{L}(D)}{\mathbb{C}} \right) \leq \dim \mathbb{C}^d - \dim \left(\frac{\Omega^1(C)}{\Omega^1(D)} \right)$$

That is:

$$l(D) - 1 \leq d - \dim \Omega^1(C) + i(D)$$

By theorem 3.5, $\dim \Omega^1(C) = g$, and so we get:

$$l(D) \leq d - g + i(D) + 1$$

This was the hardest part of the proof. We now cases cases on $l(D)$ and $i(D)$ to finish:

1. Consider $l(D) \neq 0$, $i(D) = 0$. As $l(D) \neq 0$, there exists an $f \in K(C)$ such that $(f) + D \geq 0$. Let $E = (f) + D$. Then $(f) = E - D$, so $E \sim D$. Then

$$l(D) = l(E) \quad i(D) = i(E) \quad \deg D = \deg E$$

Thus, we may assume $D \geq 0$. Thus we get our inequality, and by the Riemann inequality:

$$d - g + i(D) + 1 = d - g + 1 \stackrel{\text{R.}}{\leq} l(D) \leq d - g + 1$$

giving equality

2. Consider $l(D) \neq 0, i(D) \neq 0$. As $l(D) \neq 0$, we may do the same trick to assume $D \geq 0$. Since $i(D) \neq 0$, there exists a $\omega \in \Omega^1(C)$ such that

$$(\omega) \geq D$$

Let $E = (\omega) - D$ so that $E \geq 0$. Then by proposition 3.3 and proposition 3.17 we get:

$$i(D) = l(E) \leq \deg E - g + 1 + i(E)$$

By the Poincaré-Hopf formula, $\deg(\omega) = 2g - 2$ so

$$\deg E = \deg(\omega) - \deg(D) = 2g - 2 - d$$

As $i(E) = l(D)$, we get:

$$i(D) \leq 2g - 2 - d - g + 1 + l(D)$$

re-arranging and combining we get:

$$l(D) \geq d - g + i(D) + 1$$

giving us the other wise

3. Consider $l(D) = 0, i(D) \neq 0$. Since $i(D) \neq 0$, there exists a $\omega \in \Omega^1(C)$ such that $(\omega) = D + E$ where $E \geq 0$. Then by proposition 3.3

$$l(E) = i(D) \neq 0 \quad i(E) = l(D) = 0$$

Thus, by case (1) applied to E we get:

$$l(E) = \deg E - g + i(E) + 1$$

Note that

$$\deg E = \deg(\omega) - \deg D = 2g - 2 - d$$

Then substituting and rewriting $l(E), i(E)$ as $i(D)$ and $l(D)$, we get the equality.

4. Consider $l(D) = 0, i(D) = 0$. Then by corollary 3.16 and corollary 3.18

$$d = g - 1$$

which is the desired equation.

(then do the exercises for $C = \mathbb{P}^1$ direct prove, and the elliptic curve exercise)

4 Application of Riemann-Roch Theorem

It shall cover genus 0, 1, 2, 3, 4.

4.1 Genus 0

Proposition 4.1: Genus 0 then Rational

Let C be a compact Riemann surface of genus 0. Then

$$C \cong \mathbb{P}^1$$

Proof :

Let $p \in C$, $D = (p) \in \text{Div}(C)$. Then:

$$0 \leq i(D) \leq i(0) = \dim \Omega^1(C) = 0$$

Thus $i(D) = 0$. By the Riemann-Roch theorem:

$$l(D) = d - g + i(D) + 1 = 1 - 0 + 0 + 1 = 2$$

Thus, $\mathcal{L}(p)$ contains two linearly independent functions, one of which can be chosen to be $f_1 = 1$. The other function must be a meromorphic function (as C is a compact Riemann surface) and must have p as a pole of order 1; call this function f_2 . Then by definition $f_2^{-1}(\infty) = p$. Hence it follows that this is a holomorphic mapping:

$$f_2 : C \rightarrow \mathbb{P}^1$$

where $\deg f_2 = \#\{f_2^{-1}(\infty)\} = 1$ implying that f_2 is in fact biholomorphic (recall lemma 2.2). Thus:

$$C \cong \mathbb{P}^1$$

as we sought to show.

4.2 Genus 1

Proposition 4.2: Genus 1 Weierstrass Equation Representation

Let C be a compact Riemann surface of genus 1. Then C can be represented by a smooth algebraic curve of degree 3 in $\mathbb{P}^3$

Proof :

We shall construct a holomorphic injective map $f : C \hookrightarrow \mathbb{P}^2$ where f shall be a smooth algebraic curve of degree 3. First, for any nonzero $\omega \in \Omega^1(C)$, by Poincaré-Hopf formula we have:

$$\deg(\omega) = 2g - 2 = 2 - 2 = 0$$

Hence, if $\omega(p) = 0$, then it would have to have a pole, but it is holomorphic! Thus, for any $D > 0$, $i(D) = 0$. Thus, letting $D = 2p$, we get:

$$i(D) = 0$$

Thus by the Riemann-Roch theorem we have that:

$$l(2p) = d - g + i(2p) + 1 = 2 - 1 + 0 + 1 = 2$$

Hence $\mathcal{L}(D)$ has non-constant meromorphic functions which only have p as a singular point of order ≤ 2 . In fact, it *must* be of order 2, as by the Riemann-Roch theorem:

$$l(p) = 1 - 1 + 1 = 1$$

Hence $\mathcal{L}(p)$ only has constant functions; and so the order of any meromorphic function with single pole p is at least 2. Let $x \in \mathcal{L}(2p)$ be such a function. Then near p we have:

$$x = \frac{c_2}{t^2} + \frac{c_1}{t} + h_1(t)$$

for a local coordinate near p where $t(p) = 0$ and $c_2 \neq 0$. Let's start by showing that it must be that $c_1 = 0$. As $\dim \Omega^1(C) = g = 1$, take a $\omega \in \Omega^1(C)$ that is nonzero so that under t

$$\omega = g(t)dt \quad g(0) \neq 0$$

Then $x\omega$ has no poles other than p , so by the Residue Theorem (the sum of all residues is 0):

$$0 = \text{Res}_p(x\omega) = c_1$$

Thus, with a suitable change of coordinates we get:

$$x = \frac{1}{z^2} + h(z)$$

Note that this x has many similar properties to the Weierstrass function, as we shall see later.

Now, for any $\omega \in \Omega^1(C)$, let

$$y = \frac{dx}{\omega}$$

which is well defined as $\omega(p) \neq 0$ for all $p \in C$. Define:

$$\begin{aligned} f : C &\rightarrow \mathbb{P}^2 \\ q &\mapsto [1 : x(q) : y(q)] \quad (p \mapsto [0 : 0 : 1]) \end{aligned}$$

Note that our chosen point p was mapped to the point at infinity. Certainly f is holomorphic, we must just show it is injective and that $f(C)$ is a curve of degree 3.

Starting by showing $f(C)$ is a degree 3 curve in $\mathbb{P}^2$, consider $x : C \rightarrow \mathbb{P}^1$. Then by construction $x^{-1}(\infty) = 2p$. Thus:

$$\deg x = 2$$

Thus, the ramification can never exceed 1, and hence the ramification index at its ramification points never exceeds 1. In particular, the coefficients of every point for the ramification divisor of x must be 1. By the Riemann-Hurwitz formula;

$$\deg R = 2(\deg x + g - 1) = 2(2 + 1 - 1) = 4$$

As p ramified, there are 3 other Ramification points:

$$R = p_1 + p_2 + P_3 + p$$

each p_i being mutually distinct:

FIG HERE

let $x(p_i) = a_i$. Let us figure out what type of ramification we have. To be succinct with tracking roots and poles, for a meromorphic differential form $\alpha \in K^1(C)$, let $(\alpha)_0$ and $(\alpha)_\infty$ represented the zeros and poles (including multiplicity) respectively. As ω has no zeros or poles then since $y = dx/\omega$ we get:

$$(y)_0 = (dx)_0 \quad (y)_\infty = (x)_\infty$$

Hence, $(y)_0$ and $(y)_\infty$ can only contain the points of x with ramification index 1, that is p_1, p_2, P_3, p . Thus, each of these “zero” and “pole” divisors are shared between the two.

To calculate, suppose $z_i (i = 1, 2, 3)$ are local coordinates near p_i where $z_i(p_i) = 0$. Since the multiplicity of x at p_i is 2,

$$x - a_i = z_i^2 h_i(z_i)$$

where h_i is a holomorphic function and $h_i(0) \neq 0$. Thus:

$$\begin{aligned} dx &= 2z_i h_i(z_i) dz_i + z_i^2 h_i'(z_i) dz_i \\ &= z_i(2h_i(z_i) + z_i h_i'(z_i)) dz_i \end{aligned}$$

But then evaluating at $z_i = 0$:

$$(2h_i(z_i) + z_i h_i'(z_i))|_{z_i=0} = 2h_i(0) \neq 0$$

and so p_i is a simple zero of dx . For the point p , choosing a local coordinate function z near p as we did before we get:

$$x = \frac{1}{z^2} + h(z)$$

Neurally, dx has p as a pole of order 3, and so we found:

$$(y)_0 = p_1 + p_2 + p_3 \quad (y)_\infty = 3p \quad (y) = p_1 + p_2 + p_3 - 3p$$

We now know the roots/pole information our desired function must have. Let $g(x) \in K(\mathbb{P}^1)$ be given by:

$$g(x) = (x - a_1)(x - a_2)(x - a_3)$$

Considering g as a function on C (that is x^*g), then as near the points p_i and p in coordinates z_i and z :

$$x - a_i = z_i^2 h_i(z_i) \quad x = \frac{1}{z^2} h(z)$$

we get:

$$(g)_0 = 2p_1 + 2p_2 + 2p_3 \quad (g)_\infty = 6p$$

since each p_i takes p_i as a zero of order 2, and near p every $(x - a_i)$ takes p as pole of order 2.

Now, we see that the divisor of g is just the divisor of y^2 , and so:

$$\left(\frac{y^2}{g} \right) = 0$$

showing y^2/g is holomorphic on C , and as C is compact $y^2/g = a \in \mathbb{C}$ where $a \neq 0$. To choose a better constant replace y by $y/\sqrt{a}$. Then we get:

$$y^2 = g(x)$$

But now, we just showed that under f that C lies in the image of a curve of degree 3:

$$C' = \{[1 : x : y] \in \mathbb{P}^2 \mid y^2 - g(x) = 0\} \cup \{[0 : 0 : 1]\}$$

Giving us the degree. Let us now check that f is an injective mapping except at ∞ .

To that end, we shall need a bit more structure on C . We shall define a natural involution map on C . Let

$$\begin{aligned} j : C &\rightarrow C \\ q_1 &\mapsto q_2 \end{aligned}$$

where $q_1, q_2 = x^{-1}(p_r)$ are an arbitrary $p_r \in C'$ (where I chose subscript r for no particular reason, we're running out of letters). Then certainly $j^2 = \text{id}$. It is thus its own inverse, and so it is easy to verify it is an automorphism. Using this, we shall show some properties of j^* on $K(C)$, $K^1(C)$ and $\Omega^1(C)$. Certainly by definition of j :

$$j^*(x) = x \quad j^*(dx) = dx$$

Let's show $j^*y = -y$. As $y = dx/\omega$:

$$j^*y = \frac{j^*(dx)}{j^*(\omega)} = \frac{dx}{d^*(\omega)}$$

Hence, we show $j^*(\omega) = -\omega$. As $\dim \Omega^1(C) = g = 1$, we get $\Omega^1(C) \cong \mathbb{C}$. Then notice that j can be thought of as a linear transformation on $]C$ so that $j^2 = 1$ implies $(j^*)^2 = 1$, (hence it is diagonalizable) hence the eigenvalues of j^* are either 1 or -1 . Suppose j^* has eigenvalue 1 so that $j^*\omega = \omega$. Then there exists a holomorphic differential φ on $\mathbb{P}^1$ such that

$$\omega = \varphi \circ x = x^*\varphi$$

(exercise). But there is no nonzero holomorphic differential on $\mathbb{P}^1$, hence $j^*\omega = -\omega$.

With the properties of j established, consider $q_1, q_2 \in C$ where $f(q_1) = f(q_2)$. We want to show that $q_1 = q_2$. Naturally if either q_1, q_2 is p then by construction $f(q_1) = f(q_2) = \infty$ and so $q_1 = q_2 = p$; thus we assume neither q_1, q_2 is p . By definition:

$$\begin{aligned} f(q_1) &= [1 : x(q_1) : y(q_1)] \\ f(q_2) &= [2 : x(q_2) : y(q_2)] \end{aligned}$$

and as these are equal $x(q_1) = x(q_2)$ and $y(q_1) = y(q_2)$. Now if $q_1 \neq q_2$, as $x(q_1) = x(q_2)$ it must be that $j(q_1) = q_2$. Thus, we have by definition of the pullback:

$$j^*y(q_1) = y(q_2)$$

however as $j^*(y) = -y$

$$j^*y(q_1) = -y(q_1) = -y(q_2)$$

and so

$$y(q_2) = -y(q_2)$$

telling us $y(q_2) = 0$. But $(y)_0 = p_1 + p_2 + p_3$, so $q_2 \in \{p_1, p_2, p_3\}$. Thus:

$$q_1 = j(q_2) = q_2$$

but this contradicts $q_1 \neq q_2$, thus it must be that they are equal, showing f is injective.

Let's now show surjectivity. We shall use a connectedness argument. First, $y^2 - g(x) = 0$ is certainly an irreducible polynomial, hence it forms an irreducible subset, hence a connected subset. Next, C is compact, and as f is holomorphic $f(C)$ is closed. As f is holomorphic, $f(C)$ is also open (by the holomorphic open mapping theorem), and so $f(C) = C'$.

Finally, we prove C' is smooth. We can show this directly by computing. By everything shown so far, we know that the projective equation of C' can be written as:

$$F = \xi^0(\xi^2)^2 - (\xi^1 - a_1\xi^0)(\xi^1 - a_2\xi^0)(\xi^1 - a_3\xi^0) = 0$$

Thus:

$$\begin{aligned} \frac{\partial F}{\partial \xi^0} &= (\xi^2)^2 + a_1(\xi^1 - a_2\xi^0)(\xi^1 - a_3\xi^0) \\ &\quad + a_2(\xi^1 - a_1\xi^0)(\xi^1 - a_3\xi^0) \\ &\quad + a_3(\xi^1 - a_1\xi^0)(\xi^1 - a_2\xi^0) \end{aligned}$$

$$\begin{aligned} \frac{\partial F}{\partial \xi^1} &= -(\xi^1 - a_2\xi^0)(\xi^1 - a_3\xi^0) \\ &\quad - (\xi^1 - a_1\xi^0)(\xi^1 - a_3\xi^0) \\ &\quad - (\xi^1 - a_1\xi^0)(\xi^1 - a_2\xi^0) \end{aligned}$$

$$\frac{\partial F}{\partial \xi^2} = 2\xi^0\xi^2$$

If $\partial F / \partial \xi^i$ ($i = 1, 2, 3$) are all zero, by the Euler formula, $F = 0$. Now from

$$2\xi^0\xi^2 = 0$$

we get

$$\xi^0(\xi^2)^2 = 0,$$

and substituting this expression into $F = 0$, we also get

$$(\xi^1 - a_1\xi^0)(\xi^1 - a_2\xi^0)(\xi^1 - a_3\xi^0) = 0.$$

At least one of the three factors on the left side of this expression is therefore equal to 0. Having only one factor equal to 0 however, contradicts

$$\frac{\partial F}{\partial \xi^1} = 0.$$

Thus at least two of the factors are equal to 0. But then $\xi^0 = \xi^1 = 0$ because a_1, a_2, a_3 are mutually distinct. Substituting into

$$\frac{\partial F}{\partial \xi^0} = 0,$$

we get $\xi^2 = 0$, i.e., we get

$$\xi^0 = \xi^1 = \xi^2 = 0,$$

and this is impossible. Thus, C' is smooth. As f is biholomorphic, we have that C is biholomorphic to a smooth degree 3 curve, as we sought to show.

4.3 Genus ≥ 2 : Canonical Map

In higher dimensions, there are generally many kinds of mapping from a compact Riemann surface into an n -dimensional projective space. However all compact Riemann surface of genus $g \geq 2$, they exists as special holomorphic mapping from this compact Riemann surface into $\mathbb{P}^{g-1}$ that is called the *canonical map*.

Definition 4.3: Canonical Map

Let C be a compact Riemann surface of genus $g \geq 2$ and $\{\omega_1, \dots, \omega_g\}$ a basis for $\Omega^1(C)$. Then:

$$\begin{aligned} \varphi_K : C &\rightarrow \mathbb{P}^{g-1} \\ p &\mapsto [\omega_1(p) : \dots : \omega_g(p)] \end{aligned}$$

Note that this is well-defined. It is certainly not coordinate dependent, and if, for the sake of contradiction, there is a p where $\omega_i(p) = 0$ for all p (which implies p maps to $[0 : \dots : 0]$ which doesn't exist), then notice that $\Omega^1(p) = \Omega^1(C)$ so

$$i(p) = \dim \Omega^1(C) = g$$

and by the Riemann Roch-theorem

$$l(p) = 1 - g + g + 1 = 2$$

and so there exists nonconstant meromorphic function $f \in \mathcal{L}(p)$ where $f^{-1}(\infty) = p$. Thus

$$\deg f = 1$$

giving us that $f : C \rightarrow \mathbb{P}^1$ is an isomorphism, but that is not possible as $\mathbb{P}^1$ has genus 0.

Proposition 4.4: Non-degeneracy of Canonical Map

Let φ_K be the canonical map. Then φ_K is non-degenerate.

Proof :

Assume φ_K is degenerate so that there exists $\lambda_\alpha \in \mathbb{C}$ not all zero st for all $p \in C$

$$\sum_{\alpha}^g \lambda_\alpha \omega_\alpha(p) = 0$$

Then:

$$\sum_{\alpha}^g \lambda_\alpha \omega_\alpha$$

contradicting that $\{\omega_1, \dots, \omega_g\}$ is a basis of $\Omega^1(C)$, as we sought to show.

Proposition 4.5: Degree 2 Mapping Condition

Let C be a Riemann surface of genus $g \geq 2$ and let φ_K be the canonical mapping. Then if

$$\varphi_K(p) = \varphi_K(q)$$

for some distinct $p, q \in C$, then there exists a holomorphic mapping of degree 2 from C into $\mathbb{P}^1$

Proof :

If $\varphi_K(p) = \varphi_K(q)$, then $\omega_\alpha(p) = \lambda \omega_\alpha(q)$ for all $\alpha = 1, \dots, g$ for some $0 \neq \lambda \in \mathbb{C}$. Let $D = p + q$. Let us find $i(D)$. We shall start by showing that:

$$\Omega^1(D) = \Omega^1(p)$$

Indeed, for any $\omega \in \Omega^1(C)$, if

$$\omega = \sum_{\alpha=1}^g \mu_\alpha \omega_\alpha$$

for $\mu_\alpha \in \mathbb{C}$, $\alpha = 1, \dots, g$, then:

$$\begin{aligned} \omega(p) = 0 &\Leftrightarrow \sum_{\alpha=1}^g \mu_\alpha \omega_\alpha(p) = 0 \\ &\Leftrightarrow \sum_{\alpha=1}^g \lambda \mu_\alpha \omega_\alpha(q) = 0 \\ &\Leftrightarrow \omega(q) = 0 \end{aligned}$$

Thus, $\Omega^1(p) = \Omega^1(p + q) = \Omega^1(D)$. Hence, to calculate $i(D)$ it suffices to calculate $i(p)$.

To that end, consider :

$$\lambda_1 \omega_1(p) + \lambda_2 \omega_2(p) + \dots + \lambda_g \omega_g(p) = 0$$

for $\lambda_1, \dots, \lambda_g \in \mathbb{C}$. Note not all these could be zero or else we would have a point map to $[0 : \dots : 0]$. Hence, this linear equation *must* have $g - 1$ linearly independent solution vectors $(\lambda_1, \dots, \lambda_g) \in \mathbb{C}^g$, giving rise to $g - 1$ linearly independent elements in $\Omega^1(p)$. Thus:

$$i(D) = i(p) = g - 1$$

Applying the Riemann-Roch there to $D = p + q$, we get:

$$l(D) = 2 - g + (g - 1) + 1 = 2$$

Thus, there exist a nonconstant meromorphic function $f \in \mathcal{L}(p+q)$. Thus there is a holomorphic $f : C \rightarrow \mathbb{P}^1$ where $f^{-1}(\infty) = p + q$. Note that $f^{-1}(\infty)$ cannot be a single point p or q otherwise we get $C \cong \mathbb{P}^1$, contradicting the fact that the genus g of C is ≥ 2 . Thus, it must be that:

$$\deg f = 2$$

as we sought to show.

This gives a rough division of Riemann surface of genus $g > 1$:

Definition 4.6: Hyperelliptic and Nonhyperelliptic

Let C be a compact Riemann surface of genus $g \geq 2$. Then it is *hyperelliptic* if there exists a holomorphic mapping of degree 2 $f : C \rightarrow \mathbb{P}^1$. Otherwise, C is said to be *nonhyperelliptic*.

We can deal with the nonhyperelliptic example holistically and then see the hyperelliptic examples by examining different genera:

Definition 4.7: Canonical Curve

Let C be a nonhyperelliptic compact Riemann surface of genus $g \geq 2$ with canonical map $\varphi_K : C \rightarrow \mathbb{P}^{g-1}$. Then

$$\varphi_K(C) \subseteq \mathbb{P}^{g-1}$$

is called the *canonical curve*.

Note that we put $g \geq 2$, but we shall show shortly that all $g = 2$ compact Riemann surface are hyperelliptic, but this shall not necessarily be the case for $g \geq 3$.

Proposition 4.8: Canonical Curve Smooth

Let C be a nonhyperelliptic compact Riemann surface. Then the canonical map φ_K is injective and $\varphi_K(C)$ is smooth.

Proof :

Proposition 4.5 show's φ_K must be injective, it remains to show φ_K is smooth.

[5, p.136]

4.4 Hyperelliptic Compact Riemann Surfaces

Proposition 4.9: Algebraic Characterization of Hyperelliptic Riemann Surface

Let C be a hyperelliptic compact Riemann surface. Then it can be represented as the normalization of a plane algebraic curve of degree $2g + 2$ where g is the genus of C . By a suitable linear projection, this plane curve will be a 2-sheered covering of a straight line (i.e. $\mathbb{C}$).

Proof :

Let C be a hyperelliptic compact Riemann surface of genus g . We shall construct a holomorphic mapping $f : C \rightarrow \mathbb{P}^2$ such that $f(C)$ can be given by a plane algebraic curve of degree $2g + 2$.

As C is hyperelliptic, there exist a degree 2 holomorphic mapping $x : C \rightarrow \mathbb{P}^1$. Let R be the ramification divisor of x . By the Riemann-Hurwitz formula (theorem 2.6):

$$\deg R = 2(2 + g - 1) = 2g + 2$$

As x is degree 2, the coefficients of every point in R must be equal to 1, hence we must have:

$$R = p_1 + p_2 + \cdots + p_{2g+2}$$

where each p_i is distinct on C . Now suppose:

$$x^{-1}(\infty) = p + q \quad x(p_i) = a_i$$

so that x can be visualized as:

FIG HERE

Then as in proposition 4.2 we can define $j : C \rightarrow C$ $j(q_1) = q_2$. Just like before, $j \in \text{Aut}(C)$ and $j^2 = 1$. Define again $j^* : K(C) \rightarrow K(C)$ by $j^*(a) = a \circ j$, which like before can be extended to $K^1(C)$ and restricted down to $\Omega^1(C)$. Then as $\Omega^1(C) \cong \mathbb{C}^g$, this is a linear involution of $\mathbb{C}^g$ with eigenvalues 1 or -1 . If the eigenvalue is 1, then the eigenvector corresponding to 1 will be a holomorphic differential form, inducing a nontrivial differential form on $\mathbb{P}^1$ a contradiction. Thus it must be that :

$$j^* : \Omega^1(C) \rightarrow \Omega^1(C) \quad \omega \mapsto -\omega$$

With the setup complete, let us show that such a function does exist by the Riemann-Roch Theorem. Let

$$D = (g+1)p + (g+1)q \in \text{Div}(C)$$

Recall that for any $\omega \in K^1(C)$ by the Poincaré-Hopf formula $\deg(\omega) = 2g - 2 < \deg G$, and so it cannot be that $(\omega) - D \geq 0$, and so $i(D) = 0$. Then by Riemann-Roch:

$$l(D) = (2g+2) - g + 1 = g + 3$$

Thus

$$\mathcal{L}(D) \cong \mathbb{C}^{g+3}$$

As $x(p) = x(q) = \infty$, it must be that $j(D) = D$, so j^* can be seen as a linear transformation on $\mathcal{L}(D)$. Then a $(j^*)^2 = 1$ and j^* also has eigenvalues ± 1 , we can split $\mathcal{L}(D)$ into eigenspaces:

$$\mathcal{L}(D) = \mathcal{L}(D)^+ \oplus \mathcal{L}(D)^-$$

where $\mathcal{L}(D)^+$ and $\mathcal{L}(D)^-$ are the eigenspace for eigenvalue $+1$ and -1 respectively. Note that the meromorphic functions in $\mathcal{L}(D)^+$ can be factored as the composition of x and a meromorphic function on $\mathbb{P}^1$ with pole poles only at p, q , and the order of the pole at either point not exceeding $g+1$. Thus, as meromorphic functions on C , the functions $1, x, x^2, \dots, x^{g+1}$ form a basis for $\mathcal{L}(D)^+$. Thus:

$$\dim \mathbb{C}(D)^+ = g + 2$$

As $l(D) = g + 3$, it follows that

$$\dim \mathbb{C}(D)^- = 1$$

In particular, there exists a meromorphic function $y \in \mathcal{L}(D)$ such that

$$j^*(y) = -y$$

With this build-up, let us compare divisors to prove that there exists a function $g(x) = (x - a_1)(x - a_2) \cdots (x - a_{2g+2})$ such that

$$y^2 = cg(x)$$

where $0 \neq c \in \mathbb{C}$. Let's first find (y^2) , in particular it suffices to find (y) . since

$$j^*(y) = -y \quad j(p_i) = p_i \quad (i = 1, 2, \dots, 2g+2)$$

we get:

$$y(p_i) = -y(p_i)$$

and so

$$y(p_i) = 0$$

Thus

$$\deg(y)_0 \geq \deg\left(\sum_{i=1}^{2g+2} p_i\right) = 2g+2$$

But by the Residue Theorem

$$\deg(y)_0 = \deg(y)_\infty = 2g+2$$

Thus

$$(y)_0 = \sum_{i=1}^{2g+2} p_i \quad (y)_\infty = D$$

For $(g(x))$, by definition $(x)_\infty = p + q$ and $(x - a_i)_\infty = p + q$ for all $i = 1, 2, \dots, 2g+2$. Thus:

$$(g(x))_\infty = (2g+2)p + (2g+2)q = 2D$$

Giving the pole divisor. For the root divisor, suppose z_i are local coordinate near p_i for each i where $z_i(p_i) = 0$. Then:

$$x - a_i = z_i^2 h_i(z_i)$$

where $h_i(0) \neq 0$ is holomorphic. But then:

$$(g(x))_0 = \sum_{i=1}^{2g+2} 2p_i$$

and so $(y^2) = (g(x))$, giving that $y^2/g(x)$ is holomorphic on C , which as is compact makes $y^2/g(x) = c \neq 0$ constant. By replacing y with $y/\sqrt{c}$ we get

$$y^2 = g(x)$$

From this, we can define the map:

$$\begin{aligned} f : C &\rightarrow \mathbb{P}^2 \\ t &\mapsto [1 : x(t) : y(t)] \\ p, q &\mapsto [0 : 0 : 1] \end{aligned}$$

As C is connected, the solution to an algebraic equation is irreducible (hence connected), f is holomorphic (hence an open map) and C is compact (hence f is a closed map), it must be that f is surjective.

Finally, it is easy to see that each a_i are distinct, hence C is smooth in $\mathbb{C}^2 = \{[1 : x : y] \in \mathbb{P}^2\}$ but singular at $[0 : 0 : 1]$ (which distinguishes from the case of $g = 1$). A linear projection $(x, y) \rightarrow x$ gives the 2-sheeted covering, giving everything asked for by the proposition, completing the proof.

Proposition 4.10: Canonical Map – Hyperelliptic Case

Let C be a hyperelliptic curve and let x, y be as in proposition 4.9. Then

$$\omega_1 = \frac{dx}{y}, \omega_2 = \frac{(xdx)}{y}, \dots, \omega_g = \frac{(x^{g-1}dx)}{y}$$

forms a basis for $\Omega^1(C)$.

Proof :

[5, p.142]

We thus get a very simple expression for the canonical map:

$$\begin{aligned} \varphi_K : C &\rightarrow \mathbb{P}^{g+1} \\ t &\mapsto [1 : x(t) : \dots : x(t)^{g-1}] \end{aligned}$$

Hence, φ_K factors in the following way:

$$\begin{array}{ccc} & & \mathbb{P}^{g-1} \\ & \nearrow \varphi_k & \uparrow r \\ C & \xrightarrow{x} & \mathbb{P}^1 \end{array}$$

where

$$\begin{aligned} r : \mathbb{P}^1 &\rightarrow \mathbb{P}^{g-1} \\ x &\mapsto [1 : x : \dots : x^{g-1}] \end{aligned}$$

Note that the converse of proposition 4.9 is true: given any $2g+2$ mutually distinct complex numbers $a_1, a_2, \dots, a_{2g+2}$, and letting:

$$C' = \{y^2 = \prod_{i=1}^{2g+2} (x - a_i) = 0\} \cup [0 : 0 : 1] \subseteq \mathbb{P}^2$$

then the normalization of C' is a hyperelliptic compact Riemann surface of genus g , giving an equivalence. Combining this with the commutative diagram given above, we see that C is uniquely determined by the equivalence of vectors in $\mathbb{C}^{2g+2}$ with (pairwise) distinct coordinates:

$$\{(a_1, \dots, a_{2g+2})\}$$

where $(a_1, \dots, a_{2g+2}) \sim (b_1, \dots, b_{2g+2})$ if each coordinate is related by the same linear automorphisms of $\mathbb{P}^1$, that is there exists $a, b, c, d \in \mathbb{C}$ ($ad - bc \neq 0$) such that

$$\frac{aa_i + b}{ca_i + d} = b_i \quad (i = 1, 2, \dots, 2g+2)$$

(recall the classification of $\text{Aut}(\mathbb{P}^1)$). Naturally, if replaced by $\lambda a, \lambda b, \lambda c, \lambda d$ the result is unchanged, hence without loss of generality we may assume one of them is fixed.

4.5 Genus 2 always hyperelliptic

Proposition 4.11: Genus 2 is Hyperelliptic

Let C be a compact Riemann surface of genus 2. Then C is hyperelliptic.

Proof :

Let C be a compact Riemann surface of genus 2. Then by proposition 4.5 it suffices to show that φ_K is not injective. But if $\varphi_K : C \rightarrow \mathbb{P}^{g-1} = \mathbb{P}^1$ were injective, C would have genus 0, completing the proof.

Thus, we always get C in $\mathbb{P}^1$ as:

$$y^2 = \prod_{i=1}^6 (x - a_i)$$

for pairwise distinct $a_1, \dots, a_6$ and forms

$$\omega_1 = \frac{dx}{y} \quad \omega_2 = \frac{x dx}{y}$$

where:

$$\varphi: C \rightarrow \mathbb{P}^1 \quad t \mapsto [\omega_1 : \omega_2] = [1 : x(t)]$$

4.6 Genus 3

We shall see in (it says at [5, p.212]) that almost all genus 3 curves are canonical curves (it will be because any smooth fourth-degree plane algebraic curve (where $2g - 2 = 4$) has genus $g = 3$ and hence is canonical). Let us start by showing that the canonical curve of genus 3 is a smooth plane algebraic curve of degree 3.

let $H \subseteq \mathbb{P}^n$ be a hyperplane given by:

$$L(\xi) = \sum_{i=0}^n l_i \xi^i = 0$$

Let C be a smooth algebraic curve in $\mathbb{P}^n$. Let H not contain any irreducible branches of C . Then pick the elements of $\text{Div}(C)$ in the following way. Choose a generic linear function:

$$L'(\xi) = \sum_{i=0}^n l'_i \xi^i$$

where by generic we mean:

$$\{L' = 0\} \cap C \cap \{L = 0\} = \emptyset$$

Then we get:

$$\frac{L}{L'} \in K(\mathbb{P}^n) \quad \frac{L}{L'}|_C \in K(C)$$

Then the set of all the zeros of $(L/L')|_C$ is given a name

We give a name to this set:

Definition 4.12: Hyperplane Section of Riemann Surface

Let C, H be as above. Then:

$$H \cdot C = \left(\frac{L}{L'} \Big|_C \right)_0$$

is the *hyperplane section* of C .

Intuitively:

$$H \cdot C = C \cap \{L(\xi) = 0\}$$

Certainly this definition is independent of choice of L' .

Definition 4.13: Degree of Smooth Curve

Let C be a smooth curve in $\mathbb{P}^n$. Then $\deg(H \cdot C)$ is called the *degree* of C and is denoted $\deg C$.

This too is well-defined as it is independent of choice of hyperplane H since if there exists another H_1 :

$$\deg(H_1 \cdot C) = \deg \left(\frac{L_1}{L'} \Big|_C \right)_0 = \deg \left(\frac{L_1}{L'} \Big|_C \right)_\infty = \deg \left(\frac{L}{L'} \Big|_C \right)_\infty = \deg \left(\frac{L}{L'} \Big|_C \right)_0 = \deg(H \cdot C)$$

Note that when $n = 2$, then H is a straight line in $\mathbb{P}^2$ not containing any irreducible curve components of C , and so by Bézout's theorem we know the curve degree is just the same as the plane curve degree given earlier.

In this section, we shall identify any nonhyperelliptic compact Riemann surface of genus $g > 1$ with its image under the canonical map, that is the canonical curve in $\mathbb{P}^{g-1}$ which is a smooth surface. We can do this as C is nonhyperelliptic and hence the canonical map is injective.

Proposition 4.14: Degree and Genus

Let C be a canonical curve of genus g . Then:

$$\deg C = 2g - 2$$

Proof :

Let $\omega_1, \dots, \omega_g$ be a basis for $\Omega^1(C)$ so that

$$C = \{[1w_1(p), \dots, \omega_g(p)] \mid p \in C\}$$

Suppose:

$$H = \left\{ \sum_{i=0}^{g-1} l_i \xi^i = 0 \right\} \quad H' = \left\{ \sum_{i=0}^{g-1} l'_i \xi^i = 0 \right\}$$

Then by going through the definition we see that $H \cdot C = (\omega)_0$ where $\omega = \sum_{i=0}^{g-1} l_i \omega_i$. As $\omega \in \Omega^1(C)$, $(\omega)_0 = (\omega)$, and hence:

$$H \cdot C = (\omega)$$

Thus

$$\deg C = \deg(H \cdot C) = \deg(\omega) = 2g - g$$

as we sought to show.

Proposition 4.15: Genus 3 By Degree 4 Curve

Let C be a canonical curve of genus 3. Then it is a smooth plane algebraic curve of degree 4.

Proof :

It is smooth by proposition 4.5. As the canonical curve of genus 3 must be in $\mathbb{P}^2$, by the proceeding we have:

$$\deg(C) = 2g - 2 = 2 \cdot 3 - 2 = 4$$

as we sought to show.

Note that if F is a degree 4 algebraic curve, then by the genus formula:

$$g = \frac{(d-1)(d-2)}{2} = \frac{3 \cdot 2}{2} = 3$$

From this we can do the same classification trick as in section 4.4 to show that the converse is mostly true (having to take into account that some canonical embeddings are not injective). Thus, it is as to verify that

$$F(x, y, z) = x^4 + y^4 + z^4 + \lambda x^2 y^2 + \mu y^2 z^2 + \nu z^2 x^2$$

is a canonical curve.

4.7 Genus 4

Canonical genus 4 curves have the amazing property that they can be characterized by the intersection of quadric¹ and a cubic hypersurface in $\mathbb{P}^3$.

Just like before, suppose C is a smooth algebraic curve in $\mathbb{P}^n$ and $H = \{\xi^0 = 0\}$ is a hypersurface. Let $D = H \cdot C \in \text{Div}(C)$. Notice that $\bigoplus_{k=0}^{\infty} \mathcal{L}(kD)$ and $\mathbb{C}[\xi^0, \dots, \xi^n]$ are graded rings. Let $F(\xi) \in \mathbb{C}[\xi^0, \dots, \xi^n]$ be a homogeneous polynomial and define:

$$r(F(\xi)) = \frac{F(\xi)}{(\xi^0)^k} \Big|_C \in \mathcal{L}(k(D))$$

Then extended r linearly to all of $\mathbb{C}[\xi^0, \dots, \xi^n]$, we get ring homomorphism:

$$r : \mathbb{C}[\xi^0, \dots, \xi^n] \rightarrow \bigoplus_{k=0}^{\infty} \mathcal{L}(kD)$$

I claimed this map is surjective; this was shown in [2] (but a judicious application of Noether Theorem also works).

I'll finish typing this up later: [5, p.148]

¹I've not seen the word "quadric" before. It means a degree 2 hypersurface embedded in D dimensions. Quadratic seems perfectly fine, but it seems that quadric is just a little more specific

4.8 Abel's Theorem and its Applications

Let C be a compact Riemann surface of genus $g \geq 1$. Then $H_1(C, \mathbb{Z})$ is a free abelian group of rank $2g$. Choose a canonical basis $\gamma_1, \gamma_2, \dots, \gamma_{2g-1}, \gamma_{2g}$ for $H_1(C, \mathbb{Z})$ satisfying the intersection requirements:

$$\gamma_i \cdot \gamma_{g+1} = 1 \quad \gamma_{g+i} \cdot \gamma_i = -1$$

for all i , and 0 otherwise; that is it is the usual vertical and horizontal circles around the handles when visualized in $\mathbb{R}^3$. Recall the period vector:

$$\pi_j = \begin{pmatrix} \int_{\gamma_j} \omega_1 \\ \vdots \\ \int_{\gamma_j} \omega_g \end{pmatrix} \in \mathbb{C}^g$$

for $j = 1, \dots, 2g$. Recall that we proved that this vector is well-defined as the paths are closed and that the period matrix is nonsingular (proposition 3.14). Thus, with $\mathbb{Z}$ -coordinates, we get a lattice in $\mathbb{C}^g$:

$$\Lambda = \left\{ \sum_{j=1}^{2g} m_j \pi_j \mid m_j \in \mathbb{Z} \right\} \subseteq \mathbb{C}^g$$

We thus can define the following:

Definition 4.16: Jacobian Variety

Let C be a compact Riemann surface. Then the g -dimensional complex torus $\mathbb{C}^g / \Lambda$ is called the *Jacobian variety* of C and will be denoted $J(C)$.

This variety will be useful to find when a divisor is principal, give a measure of the lack thereof. Namely, given a $D \in \text{Div}(C)$, does there exist a $f \in K^*(C)$ such that

$$(f) = D$$

First off, as we are working over compact Riemann surfaces, we can look at divisor of degree 0. Let:

$$\begin{aligned} \deg : \text{Div}(C) &\rightarrow \mathbb{Z} \\ D &\mapsto \deg D \end{aligned}$$

Let $\text{Div}^0(C) = \ker(\deg)$. Then if $(f) = D$, it must be that $D \in \text{Div}^0(C)$. The converse is not always true, and Abel's theorem will outline exactly how to associate these two.

To that end, let us start by defining a map from $\text{Div}(C)$ to $J(C)$. Let $q \in C$ be any fixed point and let $\omega \in \Omega^1(C)$ be a holomorphic differential. Then for a point $p \in C$, we can define:

$$\int_q^p \omega \tag{2}$$

Since there is nontrivial winding, the difference of two paths from q to p is a homology class that can be written as

$$\sum_{j=1}^{2g} m_j \gamma_j$$

hence two values of equation (2) will differ by:

$$\sum_{j=1}^{2g} m_j \int_{\gamma_j} \omega$$

By letting ω run through the basis of $\Omega^1(C)$, we get the following:

Definition 4.17: Abel-Jacobi Map

Let C be a compact Riemann surface, $q \in C$. Then the *Abel-Jacobi map*

$$u : \text{Div}(C) \rightarrow J(C)$$

is given by:

$$u(D) = \begin{pmatrix} \sum_{i=1}^k n_i \int_q^{p_i} \omega_1 \\ \vdots \\ \sum_{i=1}^k n_i \int_q^{p_i} \omega_g \end{pmatrix}$$

where $D = \sum_{i=1}^k n_i p_i$.

Certainly this is a homomorphism for if $D = \sum_{i=1}^k p_i - \sum_{i=1}^k q_i \in \text{Div}^0(C)$ then by the rules of integration bounds we get the homomorphism rule. We shall see that $(f) = D$ for $D \in \text{Div}^0(C)$ if and only if

$$u(D) = 0$$

Or, slightly more generally:

Theorem 4.18: Abel-Jacobi Theorem

Let C be a compact Riemann surface. Then the sequence:

$$K^*(C) \xrightarrow{(-)} \text{Div}^0(C) \xrightarrow{u} J(C) \rightarrow 0$$

is exact

We shall not give full proof of this, but [5] proves it. From this we get:

Definition 4.19: Picard Variety

Let C be a compact Riemann surface and $(-) : K^*(C) \rightarrow \text{Div}^0(C)$. Then:

$$\text{Pic}(C) = \frac{\text{Div}^0(C)}{\text{im}((-)}$$

The Abel-Jacobi map then becomes an isomorphism.

(a lot is done here in proving it, rather interesting!!)

(the Riemann bilinear relations puts important constrictions on the period matrix that shows how it reflects some of its topological properties!) (this gives us:)

Theorem 4.20: Torelli Theorem

Let C, C' be compact Riemann surfaces. Then a necessary and sufficient conditions for $C \cong C'$ as biholomorphisms is that they have the same normalized period matrix under a suitable choice of canonical homology bases

Proof :

[5, p.174]

(then it goes onto prove Jacobi inversion theorem; that $u : \text{Div}^0(C) \rightarrow J(C)$ is surjective)

4.9 Applications of Abel's Theorem

Let C be a nonhyperelliptic compact Riemann surface of genus 3; without loss of generality we may identify C with its canonical curve in $\mathbb{P}^2$. The nC is a fourth-degree smooth plane algebraic curve. Let L be a straight line in $\mathbb{P}^2$. Then:

$$D = L \cdot C = p_1 + p_2 + p_3 + p_4 \in C^{(4)}$$

is a hyperplane section. Let $E \in \mathbb{C}^{(4)}$. Then $u(E) = u(D)$ if and only if there exists a line L' such that

$$E = L' \cdot C$$

hence, E must be a hyperplane. Indeed, if $E = L' \cdot C$, let L and L' be given by linear equations l, l' so that $L = \{l = 0\}$ and $L' = \{l' = 0\}$. Letting:

$$f = \frac{l'}{l} \Big|_C \in K(C)$$

then $(f) = E - D$, and so $0 = u(E) - u(D)$, hence $u(E) = u(D)$. Conversely, if $u(D) = u(E)$, then we need to show that:

$$E \equiv q_1 + q_2 + q_3 + q_4$$

is a hyperplane section. Suppose that L' is the unique line passing through q_1, q_2 . Then:

$$L' \cdot C = q_1 + q_2 + q' + q''$$

If $L' \cdot C \neq E$, then it must be that :

$$q_3 + q_4 - q' - q'' \neq 0$$

If $u(D) = u(E)$, then $u(E) = u(L' \cdot C)$ (as D and $L' \cdot C$ are both hyperplanes sections), and so:

$$u(q_3 + q_4 - q' - q'') = 0$$

Thus by the Abel-Jacobi Theorem there exists an f such that

$$(f) = q_3 + q_4 - q' - q''$$

Now, consider $f : C \rightarrow \mathbb{P}^1$ as a holomorphic mapping. Then f has either degree 1 or 2 (depending on whether one of $\{q_3, q_4\}$ is equal to one of $\{q', q''\}$). Implying that C is either Riemann sphere or a

hyperelliptic Riemann surface, but that contradicts the fact that C is nonhyperelliptic of genus 3. Thus:

$$E = L' \cdot C$$

Thus, given a hyperplane section D of a canonical curve of genus 3, $|D|$ can be identified with the set of hyperplanes of $\mathbb{P}^2$, that is:

$$|D| \cong \mathbb{P}^2$$

Giving us a full geometric picture!

Next, if $g = 1$, we can rather easily classify all such surfaces up to biholomorphism. We first have:

Theorem 4.21: Jacobi Variety in Genus 1

Let C be a compact Riemann surface of genus 1. Then it is biholomorphic to its Jacobian variety

$$C \cong J(C)$$

Given a normalized period matrix

$$\Pi = (1 \ z)1 \times x$$

Then $C \cong \mathbb{C}/\Lambda$, where Λ is the lattice spanned by 1 and z .

Proof :

In [5, p.193], this was a consequence in the discussion of the Jacobi inversion theorem in the case of $d = 1$, where immediately concludes:

$$u : C \rightarrow J(C)$$

is a biholomorphic mapping.

Theorem 4.22: Up To Projective Special Automorphism

Let $C = \mathbb{C}/\Lambda$ and $C' = \mathbb{C}/\Lambda'$ where:

$$\begin{aligned} \Lambda &= \{m + nz \mid m, n \in \mathbb{Z}\} & \text{Im} z > 0 \\ \Lambda' &= \{m + nz' \mid m, n \in \mathbb{Z}\} & \text{Im} z' > 0 \end{aligned}$$

then $C \cong C'$ if and only if and only if

$$z = \frac{\alpha z' + \beta}{\gamma z' + \delta}$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$ and

$$\alpha\delta - \beta\gamma = 1$$

Proof :

In [5, p.194], but can also be found in [4] or [1].

(then he goes onto finish off with some commentary about elliptic integrals, the “usual” representation a $p(x) = 4x^3 + ax + b$, uses Bézouts theorem to get the number of flex points of an n th degree curve (that is $3n(n-2)$ counting multiplicity), and he uses all of this to show the original connection to integrating ellipses. The last thing he does is show some algebraic properties of $x(u)$; just like was shown a Weierstrass equation section in a complex analysis course).

5 Riemann Roch in Modern Language

here

6 Generalized Riemann Roch: Vector Bundles

Let X be a complex irreducible algebraic variety (not necessarily smooth). Define:

1. X_{Zar} : X with the Zariski topology (abbreviated to Z -topology) and $\mathcal{O}_X$ the germs of algebraic functions
2. $X_{\mathbb{C}}$: X with the complex topology (abbreviated to $\mathbb{C}$ -topology) and $\mathcal{O}_X$ the germs of algebraic functions
3. X^{an} : X with the complex topology and $\mathcal{O}_X$ the germs of holomorphic functions
4. X_{smooth} : X with the complex topology and $\mathcal{O}_X$ the germs of C^∞ -functions

Then a vector bundle $p : E \rightarrow X$ can either be locally trivial in the $\mathbb{Z}$ -topology or the $\mathbb{C}$ -topology. As X is irreducible, it is connected, so the rank is a well-defined. If $E \rightarrow X$ has rank r , then:

1. (covering) there exists an open covering $\{U_\alpha \rightarrow X\}_\alpha$ along with isomorphisms:

$$\varphi_\alpha : p^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^r$$

We shall sometimes denote $p^{-1}(U_\alpha)$ as $E|_{U_\alpha}$.

2. (local trivialization) For every α , the following diagram commutes:

$$\begin{array}{ccc} p^{-1}(U_\alpha) & \xrightarrow{\sim} & U_\alpha \times \mathbb{C}^r \\ & \searrow p & \swarrow \pi_1 \\ & X & \end{array}$$

3. (gluing – cocycle condition)

$$\begin{array}{ccc}
p^{-1}(U_\alpha) & \xrightarrow{\phi_\alpha} & U_\alpha \amalg \mathbb{C}^r \\
\downarrow & & \downarrow \\
p^{-1}(U_\alpha \cap U_\beta) & \xrightarrow{\phi_\alpha} & (U_\alpha \cap U_\beta) \amalg \mathbb{C}^r \\
\parallel & & \downarrow g_\alpha^\beta \\
p^{-1}(U_\beta \cap U_\alpha) & \xrightarrow{\phi_\beta} & (U_\beta \cap U_\alpha) \amalg \mathbb{C}^r \\
\downarrow & & \downarrow \\
p^{-1}(U_\beta) & \xrightarrow{\phi_\beta} & U_\beta \amalg \mathbb{C}^r
\end{array}$$

where by commutativity $g_\alpha^\beta = \varphi_\beta \circ \varphi_\alpha^{-1}|_{p^{-1}(U_\alpha \cap U_\beta)}$.

Note then that

$$g_\alpha^\beta|_{U_\alpha \cap U_\beta} = \text{id} \quad \text{and} \quad g_\alpha^\beta|_{\mathbb{C}^r} \in \text{GL}_r(\Gamma(U_\alpha \cap U_\beta), \mathcal{O}_X)$$

On a triple overlap, we get:

$$g_\beta^\gamma \circ g_\alpha^\beta = g_\alpha^\gamma \quad \text{and} \quad (g_\beta^\alpha)^{-1} = g_\alpha^\beta$$

Hence, the collection $\{g_\alpha^\beta\}$ forms a 1-cocycle in $Z^1(\{U_\alpha \rightarrow X\}, \mathbb{GL})$ where $\mathbb{GL}$ or $\mathbb{GL}(X)$ is:

$$\Gamma(U, \mathbb{GL}_r(X)) = \text{GL}_r(\Gamma(U, \mathcal{O}_X))$$

where the right hand side is the group of invertible linear maps on the free module $\Gamma(U, \mathcal{O}_X)^r \cong \Gamma(U, \mathcal{O}_X^r)$. When $r = 1$, we often denote $\mathbb{GL}_1(X)$ as $\mathbb{G}_m$ or $\mathcal{O}_X^*$.

Let's talk about choice. Say $\{\psi_\alpha\}_\alpha$ is another trivialization. Without loss of generality we may refine each cover so that $\{\varphi_\alpha\}_\alpha$ and $\{\psi_\alpha\}_\alpha$ use the same cover. Then there exists an isomorphism: $\sigma_\alpha : U_\alpha \times \mathbb{C}^r \rightarrow U_\alpha \times \mathbb{C}^r$ where the following diagram commutes:

$$\begin{array}{ccc}
& & U_\alpha \times \mathbb{C}^r \\
& \nearrow \varphi_\alpha & \downarrow \sigma_\alpha \\
p^{-1}(U_\alpha) & & \\
& \searrow \psi_\alpha & \downarrow \\
& & U_\alpha \times \mathbb{C}^r
\end{array}$$

Then $\{\sigma_\alpha\}_\alpha$ form 0-cochains in $C^0(\{U_\alpha \rightarrow X\}, \mathbb{GL}_r)$. if $\{h_\alpha^\beta\}$ is the gluing information for the $\{\psi_\alpha\}$ then:

$$\phi_\beta = g_\alpha^\beta \circ \phi_\alpha \psi_\beta = h_\alpha^\beta \circ \psi_\alpha \psi_\beta = \sigma_\alpha \circ \phi_\alpha.$$

Thus: $\sigma_\beta \circ \phi_\beta = \psi_\beta = h_\alpha^\beta \circ \sigma_\alpha \circ \phi_\alpha$, so:

$$\phi_\beta = (\sigma_\beta^{-1} \circ h_\alpha^\beta \circ \sigma_\alpha) \circ \phi_\alpha,$$

$$g_\alpha^\beta = \sigma_\beta^{-1} \circ h_\alpha^\beta \circ \sigma_\alpha$$
$$Z^1/\sim = H^1(\{U_\alpha \rightarrow X\}, \mathbb{GL}_r)$$

Theorem 6.1: Vector Bundle Classification Through Cohomology

$$\begin{array}{ccccccc}
1 & \longrightarrow & G'(X) & \longrightarrow & G(X) & \longrightarrow & G''(X) \\
& & & & & & \downarrow \delta_0 \\
& & & & & & \check{H}^1(X, G'') \\
& & & & & & \uparrow \\
& & & & & & \check{H}^1(X, G) \\
& & & & & & \uparrow \\
& & & & & & \check{H}^1(X, G')
\end{array}$$
$$\delta_0(\sigma) = s_\alpha s_\beta^{-1} \text{ on } U_\alpha \cap U_\beta / \sim.$$
$$0 \longrightarrow \mathbb{G}_m \longrightarrow \mathbb{G}_{r+1} \longrightarrow \mathbb{PGL}_r \longrightarrow 0 \text{ is exact.}$$

45

7 Terms for G -bundles

1. Principal G -bundle generalization of $X \times G \rightarrow G$. Let $P \rightarrow X$ be a whose fibers $p^{-1}(x)$ have a G -actions $G \times p^{-1}(x) \rightarrow p^{-1}(x)$ that is free and transitive. Given the appropriate topology, G is homeomorphic to $p^{-1}(x)$. The canonical example is the frame-bundle of a manifold M isomorphic to the appropriate subgroup of $GL_r(\mathbb{R})$. One application of principal bundles is detecting when a manifold may have additional structures; commonly it would be that there exists a quotient M/G (solggeojk quotient per fiber) that has a complex, quaternion, spin, etc, structure.
2. Principal G -bundles also important for Čech cohomology, as it has the transition information.

8 Some Hodge Theory

Let X be a complex analytic manifold of complex dimension n . As a real manifold, it is a smooth manifold of dimension $2n$. Let TX be the tangent bundle and $T_{X,x}$ the fiber at x , i.e. the $\mathbb{C}$ -vector space of (complex) dimension n . Let $z_1, \dots, z_n$ be complex local coordinates at $x \in X$ on U ; these are associated to $2n$ real coordinates $x_1, y_1, \dots, x_n, y_n$ where $z_j = x_j + iy_j$. Given such $2n$ real coordinates, there is two natural coordinates that can be from them; z_j and $\bar{z}_j$ via:

$$z_j = x_j + iy_j \quad \bar{z}_j = x_j - iy_j$$

and conversely using these we can form:

$$x_j = \frac{1}{2}(z_j + \bar{z}_j) \quad y_j = \frac{1}{2i}(z_j - \bar{z}_j)$$

If we treat TX as an $\mathbb{R}$ -vector space, we can complexity to naturally combine the z_j and $\bar{z}_j$ coordinates. To see this, a $\mathbb{C}$ -basis at $T_{X,x}$ is

$$\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}$$

And $\mathbb{R}$ -basis at $T_{X,x}$ is

$$\frac{\partial}{\partial x_1}, \frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial x_n}, \frac{\partial}{\partial y_n}$$

which is real $2n$ -dimensional (and each pair $\partial/\partial x_j, \partial/\partial y_j$ satisfy the Cauchy-Riemann equation). We can create complex dimensional spaces by tensoring:

$$T_{X,x\mathbb{C}} = T_{X,x} \otimes_{\mathbb{R}} \mathbb{C}$$

Then $T_{X,x\mathbb{C}}$ has a basis

$$\frac{\partial}{\partial z_j}, \frac{\partial}{\partial \bar{z}_j} \quad 1 \leq j \leq n$$

Then for any $f \in C^\infty(U)$:

$$\frac{\partial f}{\partial z_j} = \frac{\partial f}{\partial x_j} - i \frac{\partial f}{\partial y_j} \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}_j} = \frac{\partial f}{\partial x_j} + i \frac{\partial f}{\partial y_j}$$

This is what is going on in coordinates, but we can also view the theory more abstractly. Let $(e_1, \dots, e_n)$ be a $\mathbb{C}$ -basis for V . Then so is $(ie_1, \dots, ie_n)$. We can have an $\mathbb{R}$ -basis for V via $(f_1, \dots, f_n, g_1, \dots, g_n)$ and write $e_j = f_j + ig_j$. Then the map $(e_1, \dots, e_n) \mapsto (ie_1, \dots, ie_n)$ is represented in the $\mathbb{R}$ -basis as :

$$((f_1, g_1), \dots, (f_n, g_n)) \xrightarrow{J} ((-g_1, f_1), \dots, (-g_n, f_n))$$

We can view J as an $\mathbb{R}$ -endomorphism of V with the property that $J^2 = -\text{id}$. Does this in fact characterize complex multiplication? If we have an $\mathbb{R}$ -space of dimension $2n$ and an $\mathbb{R}$ -endomorphism J of V such that $J^2 = -\text{id}$, then we can give V a complex structure (i.e. a $\mathbb{C}$ action via:

$$(a + bi) \cdot v = av + bJ(v)$$

For example, on $\mathbb{R}^2$, there is $J(a_1, a_2) = (-a_2, a_1)$ and $J(a_1, a_2) = (a_2, -a_1)$. It is an exercise in algebra to see that on a real $2n$ -dimensional vector space V , the $\mathbb{C}$ -vector space structures on V corresponding bijective to the homogeneous to $\text{GL}_{2n}(\mathbb{R})/\text{GL}_n(\mathbb{C})$ via:

$$[A] \mapsto AJA^{-1}$$

Definition 8.1: Almost Complex Structure

Let X be a real smooth manifold. Then it has an *almost complex structure* if there exists a bundle morphism $J : TX \rightarrow TX$ such that $J^2 = -\text{id}$

Proposition 8.2: Complex Analytic, then Almost Complex

Let $(X, \mathcal{O}_X)$ be a complex analytic manifold. Then it is almost complex

Proof :

exercise (it is just pushing definitions around)

Now, consider a complex n -dimensional complex bundle TX ; let $z_1, \dots, z_n$ be its coordinate. Then as mentioned earlier, we can complexify to get a basis:

$$\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}, \frac{\partial}{\partial \bar{z}_1}, \dots, \frac{\partial}{\partial \bar{z}_n}$$

and so, in this basis, if we write $f = (w_1, \dots, w_n)$, where $w_\alpha = u_\alpha + iv_\alpha$, we get

$$J_{\mathbb{R}}(f) = \begin{pmatrix} \left(\frac{\partial w_\alpha}{\partial z_j} \right) & \left(\frac{\partial w_\alpha}{\partial \bar{z}_j} \right) \\ \left(\frac{\partial \bar{w}_\alpha}{\partial z_j} \right) & \left(\frac{\partial \bar{w}_\alpha}{\partial \bar{z}_j} \right) \end{pmatrix}$$

And if f is holomorphic:

$$\frac{\partial w_\alpha}{\partial \bar{z}_j} = \frac{\partial \bar{w}_\alpha}{\partial z_j} = 0,$$

giving:

$$J_{\mathbb{R}}(f) = \begin{pmatrix} \left(\frac{\partial w_\alpha}{\partial z_j} \right) & 0 \\ 0 & \left(\frac{\partial \bar{w}_\alpha}{\partial \bar{z}_j} \right) \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & \bar{A} \end{pmatrix}.$$

Now, write:

$$J(f) = \left(\frac{\partial w_\alpha}{\partial z_j} \right)$$

and call it the *holomorphic Jacobian*. Then:

1. $J_{\mathbb{R}}(f) = \begin{pmatrix} J(f) & 0 \\ 0 & \overline{J(f)} \end{pmatrix}$, so, $\mathbb{R}\text{-rank } J_{\mathbb{R}}(f) = 2\mathbb{C}\text{-rank } J(f)$.
2. We have $\det(J_{\mathbb{R}}(f)) = |\det(J(f))|^2 \geq 0$, and $\det(J_{\mathbb{R}}(f)) > 0$ if f is a holomorphic isomorphism (in this case, $m = n$ the common dimension of X, Y).

Then we see holomorphic maps preserve the orientation of a complex manifold and each complex manifold posses an orientation.

There is a lot more about Hodge stuff, but I don't feel like this is super helpful right now

9 L-Genus and Todd Genus

Let B be a commutative ring, and $Z, \alpha_1, \dots, \alpha_n$ be some independent intermediates (label the collection of alpha's α), all of degree 1. Let σ_j be symmetric functions in α . Let

$$q_j = \sigma_j(\alpha)$$

Define the ring $\mathcal{B} = B[[Z; \alpha_1, \alpha_2, \dots, \alpha_n]]$. Take the ring $\mathcal{P} = B[[Z; q_1, \dots, q_n]]$. Then $\mathcal{P}$ contains units called *one-units* that is:

$$1 = \sum_{j \geq 1} b_j^j \quad b \in B$$

If $Q(z)$ is a one-unit, then

9.1 Todd Class Fixes Commutative Diagram

The diagram:

$$\begin{array}{ccc} H^*(X; \mathbb{Q}) & \xrightarrow{f_*} & H^*(X; \mathbb{Q}) \\ \text{ch} \uparrow & & \uparrow \text{ch} \\ K(X) & \xrightarrow{f_*} & K(Y) \end{array}$$

Is not commutative, but the diagram:

$$\begin{array}{ccc} H^*(X; \mathbb{Q}) & \xrightarrow{f_*} & H^*(X; \mathbb{Q}) \\ \text{td}(X)\text{ch} \uparrow & & \uparrow \text{td}(X)\text{ch} \\ K(X) & \xrightarrow{f_*} & K(Y) \end{array}$$

So the Chern character is in some way an insufficient tool to naturally represent homological information of vector bundles, but “fixing” it by the Todd class is.

9.2 Multiplicative sequences

As we saw with Chern and Pontryagin classes, their product translates to a cup product. This gives a natural algebraic relation that we know codify for future reference. Let B be a commutative ring, and define:

$$\mathcal{B} = B[z, p_1, p_2, \dots]$$

Define a grading given by the following:

$$p_{j_1} p_{j_2} \cdots p_{j_r} \text{ has weight } j_1 + j_2 + \cdots + j_r$$

Then let $\mathcal{B}_k$ be the weight k -groups; note $\mathcal{B}_0 = B$ and $\mathcal{B}_r \mathcal{B}_s = \mathcal{B}_{r+s}$. Let $\pi(k)$ be all partitions of k . Then $\mathcal{B}_k$ is a B -module of rank $\pi(k)$.

Definition 9.1: Multiplicative Sequence

Let $\{K_j\}$ be a sequence of polynomials in indeterminates p_i where $K_0 = 1$ and $K_j \in \mathcal{B}_j$. Then the sequence $\{K_j\}$ is called a *multiplicative sequence* if whenever a power series is the product of two power series:

$$1 + p_1 z + p_2 z^2 + \cdots = (1 + p'_1 z + p'_2 z^2 + \cdots)(1 + p''_1 z + p''_2 z^2 + \cdots)$$

with indeterminates z, p'_i, p''_i , then we get the identity:

$$\sum_{j=0}^{\infty} K_j(p_1, p_2, \dots, p_j) z^j = \left(\sum_{i=1}^{\infty} K_i(p'_1, p'_2, \dots, p'_i) z^i \right) \left(\sum_{j=1}^{\infty} K_j(p''_1, p''_2, \dots, p''_j) z^j \right)$$

that is:

$$K \left(\sum_{i=0}^{\infty} q_i z^i \right) = \sum_{i=0}^{\infty} K_i(q_1, \dots, q_i) z^i$$

where $q_i \in \mathcal{B}$ of weight i . In other words a sequence $\{K_j\}$ is called multiplicative if it induces an endomorphism on

$$B[p_1, p_2, \dots][[z]]$$

Given a multiplicative sequence $\{K_j\}$, we associate to it its *characteristic power series* as the image of $K(1+z)$:

$$K(1+z) = \sum_j K_j(1, 0, \dots, 0) z^j = \sum_j b_j z^j$$

Recall we can define Chern roots. Hence we will be considering:

$$1 + p_1 z + \cdots + p_n z^n = \prod_i^n (1 + \beta_i z)$$

where β_i is a symmetric polynomial in the p_i 's. By some elementary ring theory and theory of symmetric functions, we can interpret $\mathcal{B}$ as the ring of all symmetric functions in $\beta_1, \dots, \beta_n$.

We can characterize m -sequences via 1-unit power series:

Proposition 9.2: Correspondence 1-unit Power Series and m -sequences

There is a bijective correspondence between m -sequences and formal power series with constant term 1.

Proof :

First, let's show any m -sequence $\{K_j\}$ is completely determined by its characteristic power series

$$Q(z) = K(1+z)$$

Proof. By decomposing:

$$1 + p_1z + \cdots + p_nz^n = \prod_i^n (1 + \beta_i z)$$

We get:

$$\sum_{j=0}^m K_j(p_1, \dots, p_j) z^j + \sum_{j=m+1}^{\infty} K_j(p_1, \dots, p_m, 0, \dots, 0) z^j = \prod_{i=1}^m Q(\beta_i z). \quad (6_m)$$

Therefore any polynomial K_j with $j \leq m$ is determined as a symmetric polynomial in the β_i and hence as a polynomial in the p_i . This holds for arbitrary m , completing the proof. $\square$

Next, let

$$Q(z) = 1 + \sum_{i=1}^{\infty} b_i z^i \quad b_i \in B$$

be a formal power series. Then there is an associated m -sequence $\{K_j\}$ where $K(1+z) = Q(z)$.

Proof. Consider the usual decomposition:

$$1 + p_1z + \cdots + p_nz^n = \prod_i^n (1 + \beta_i z)$$

and consider the product $\prod_{i=1}^m Q(\beta_i z)$ where β_i are symmetric functions. Then the coefficients z^j are homogeneous of weight j . Thus, it is in $K_j^{(m)}(p_1, \dots, p_j)$. It follows that $K_j^{(m)}$ does not depend on m for $m \geq j$.

Now, define $K_j = K_j^{(m)}$ for $m \geq j$. I claim $\{K_j\}$ is the required m -sequence. Indeed, the earlier proof the multiplicative property is true if the indeterminates p'_i, p''_i are replaced by 0 for large values of i , and so $\{K_j\}$ is an m -sequence. Finally applying $m = 1$ in equation:

$$\sum_{j=0}^m K_j(p_1, \dots, p_j) z^j + \sum_{j=m+1}^{\infty} K_j(p_1, \dots, p_m, 0, \dots, 0) z^j = \prod_{i=1}^m Q(\beta_i z). \quad (6_m)$$

shows that $K(1+z) = Q(z)$, completing the proof. $\square$

Combining these, we get the correspondence

The simplest example would be $\{p_j\}$ which has the characteristic series $1+z$. The notation p_i is due to the connection to Pontryagin classes. We shall be working with Chern classes so the following will be a useful substitution. Let $c_0 = p_0 = 1$, $z = x^2$, and γ_i be such that $\beta_i = \gamma_i^2$. then:

$$\sum_{i=1}^{\infty} p_i (-z)^i \left(\sum_{j=0}^{\infty} c_j (-x)^j \right) \left(\sum_{i=0}^{\infty} c_i x^i \right)$$

Naturally, if $\{K_j(p_1, \dots, p_j)\}$ is an m -sequence with $Q(z)$ as the characteristic power series, then let $\{\tilde{K}_j(c_1, \dots, c_j)\}$ be an m -sequence where $\tilde{Q}(x) = Q(x^2)$. Then:

$$\begin{aligned} K_j(p_1, \dots, p_j) &= \tilde{K}_{2j}(c_1, \dots, c_{2j}) \\ &\text{and} \\ 0 &= \tilde{K}_{2j+1}(c_1, \dots, c_{2j+1}) \end{aligned}$$

Thus, the m -sequence in c_i with the characteristic power series $1+x^2$ is given by

$$1, 0, p_1, 0, p_2, \dots$$

From this relations, we get:

$$\begin{aligned} p_1 &= -2c_2 + c_1^2 \\ p_2 &= 2c_4 - 2c_3c_1 + c_2^2 \\ p_3 &= -2c_6 + 2c_5c_1 + 5c_4c_2 + 1 = 2c_4c_2 + c_3^2, \dots \end{aligned}$$

Let's now find a way to calculate the explicit calculation of the polynomials of m -sequences. Given $Q(z) = 1 + \sum_{i=1}^{\infty} b_i z^i$, consider the factorization:

$$1 + b_1 z + b_2 z^2 + \dots + b_m z^m = (1 + \beta'_1 z) \dots (1 + \beta'_m z)$$

Then define the sum:

$$\sum (\beta'_1)^{j_1} (\beta'_2)^{j_2} (\beta'_r)^{j_r} \quad \text{where} \quad j_1 \geq j_2 \geq \dots \geq j_r \geq 1, \quad \sum_{s=1}^r j_s = k \leq m$$

which denotes the symmetric function in the β'_i which is the sum of all pairwise distinct monomials obtained by permuting $(\beta'_1, \dots, \beta'_m)$ to the monomial $(\beta'_1)^{j_1} (\beta'_2)^{j_2} (\beta'_r)^{j_r}$. The number of monomial in the sum is $m!/h$, h being the number of permutations living the aforementioned monomial fixed. Now:

Lemma 9.3: Computation of m -sequence

Let $\{K_j(p_1, \dots, p_j)\}$ be the m -sequence corresponding to $Q(z) = 1 + \sum_{i=1}^{\infty} b_i z^i$. Then the coefficient of $p_{j_1} p_{j_2} \dots p_{j_r}$ of K_s where $j_1 \geq j_2 \geq \dots \geq j_r \geq 1$ and $\sum_{s=1}^r j_s = k$ is equal to:

$$\sum (j_1, j_2, \dots, j_r)$$

Proof :

it is computations that will be omitted.

We shall want to define the m -sequence associated to Pontryagin classes associated to $\mathbb{CP}^n$ seen as the classifying space for n -dimensional complex vector bundles. Recall that $\chi(\mathbb{CP}^1 \oplus \mathbb{CP}^2) = \chi(\mathbb{CP}^1)\chi(\mathbb{CP}^2)$. And that $\chi(\mathbb{CP}^n) = 1$. We shall want to capture these properties using the Pontryagin class. Define:

$$Q(z) = \frac{\sqrt{z}}{\tanh \sqrt{z}} = 1 + \sum_{k=1}^{\infty} (-1)^{k-1} \frac{2^{2k}}{(2k)!} B_k z^k$$

where the B_k are the Bernoulli numbers. Note that in this case $B = \mathbb{Q}$. Furthermore, we shall have $B_k > 0$ skip the values of $1/2$. Thus we get:

$$B_1 = \frac{1}{6}, B_2 = \frac{1}{30}, B_3 = \frac{1}{42}, B_4 = \frac{1}{30}, B_5 = \frac{5}{66}, B_6 = \frac{691}{2730}, B_7 = \frac{7}{6}, B_8 = \frac{3617}{510}, \dots$$

We shall denote the m -sequence of $Q(z)$ as $\{L_j(p_1, \dots, p_j)\}$. Computing we get:

$$\begin{aligned} L_1 &= \frac{1}{3}p_1 \\ L_2 &= \frac{1}{5 \cdot 7}(7p_2 - p_1^2) \\ L_3 &= \frac{1}{3^3 \cdot 5 \cdot 7}(62p_3 - 13p_2p_1 + 2p_1^3) \\ &\vdots \end{aligned}$$

We will want the following normalization: $L_k(p_1, \dots, p_k) = 1$. To that end, if we let $p_i = \binom{2k+1}{i}$, then we get:

$$1 + p_1 z + p_2 z^2 + \dots + p_k z^k = (1 + z)^{2k+1} \bmod z^{k+1} + 1$$

From this we get:

Proposition 9.4: Coefficient of L-genus Power Series

Let $Q(z) = \frac{\sqrt{z}}{\tanh \sqrt{z}}$. Then for every k , the coefficient J_k of z^k in $(Q(z))^{2k+1}$ is equal to 1, and $Q(z)$ is the *only* power series with rational coefficient with this property.

Proof :

Apply Cauchy's integral formula and compute (see [6, p.12] to see the exact computations)

The above is useful for Pontryagin classes. We also requires the equivalent for Chern classes. Let:

$$Q(x) = \frac{x}{1 - e^{-x}} = 1 + \frac{1}{2}x + \sum_{k=1}^{\infty} (-1)^{k-1} \frac{B_k}{(2k)!} x^{2k}$$

Then we get polynomials T_k called *Todd polynomials*. For future reference, we have the identity:

$$\frac{x}{1 - \exp(-x)} = \exp\left(\frac{1}{2}x\right) \frac{\frac{1}{2}x}{\sinh \frac{1}{2}x}$$

By using the translation from Pontryagin to Chern class lemma, we get:

$$T + k(c_1, \dots, c_k) = \sum_{\substack{r,s \\ r+2s=k}} \frac{1}{2^{4s}r!} \left(\frac{1}{2}c_1\right)^r A_s(p_1, \dots, p_s)$$

We can compute the first few values:

$$\begin{aligned} T_1 &= \frac{1}{2}c_1 \\ T_2 &= \frac{1}{12}(c_2 + c_1^2) \\ T_3 &= \frac{1}{24}c_2c_1 \\ T_4 &= \frac{1}{720}(-c_4 + c_3c_1 + 3c_2^2 + 4c_2c_1^2 - c_1^4) \\ T_5 &= \frac{1}{1440}(-c_4c_1 + c_3c_1^2 + 3c_2^2c_1 - c_2c_1^3) \\ &\vdots \end{aligned}$$

Again, by substituting $c_i = \binom{n+i}{i}$ defined by:

$$1 + c_1x + \dots + c_nx^n = (1+x)^{n+1} \bmod x^{n+1}$$

gives

$$T_n(c_1, \dots, c_n) = 1$$

Just like before, $Q(x)$ is the only power series with rational coefficients with the property that $Q(x)^{k+1} = 1$ (see [6, p.14], or just repeat the above results with the appropriate modifications).

We shall need a special case. Let $B = \mathbb{Q}[y]$. Then consider the m -sequence $T_j(y; c_1, \dots, c_j)$ with characteristic power series:

$$Q(y; x) = \frac{x(y+1)}{1 - e^{-x(y+1)}} - yx = \frac{x(y+1)}{e^{x(y+1)} - 1} + x$$

Lemma 9.5

For every n the coefficient of x^n in $(Q(y; x))^{n+1}$ is equal to $\sum_{i=0}^n (-1)^i y^i$, and $Q(y, x)$ is the only power series with coefficients in $\mathbb{Q}[y]$ which has this property.

Proof :

Take substitution

$$c_i = \binom{n+1}{i}$$

then:

$$T_n(y; c_1, \dots, c_n) = 1 - y + y^2 - \dots + (-1)^n y^n.$$

Then as $Q(\frac{1}{y}; yx) = Q(y; -x)$:

$$y^n T_n(\frac{1}{y}; c_1, \dots, c_n) = (-1)^n T_n(y; c_1, \dots, c_n)$$

as we sought to show

The polynomial $T_n(y; c_1, \dots, c_n)$ can be written in a unique way in the form

$$T_n(y; c_1, \dots, c_n) = \sum_{p=0}^n T_n^p(c_1, \dots, c_n) y^p.$$

The polynomials $T_n^p(c_1, \dots, c_n)$ satisfy

$$T_n^p(c_1, \dots, c_n) = (-1)^n T_n^{n-p}(c_1, \dots, c_n)$$

more on p.15

9.3 Todd Genus

Let X be a complex smooth manifold, and ξ a continuous q -vector bundle on X . Let $c_i \in H^{2i}(X, \mathbb{Z})$ be its Chern classes. Then the *total Todd class* of ξ is :

$$\text{td}(\xi) = \sum_{j=1}^{\infty} T_j(c_1, \dots, c_j)$$

where $\{T_j(c_1, \dots, c_j)\}$ is given by the characteristic series:

$$Q(x) = \frac{x}{1 - e^{-x}} = 1 = \frac{1}{2}x + \sum_{j=1}^{\infty} (-1)^{j-1} \frac{B_j}{(2j)!} x^{2j}$$

where B_j is the Bernoulli numbers. To write down the m -sequence, recall that the m -sequence of the characteristic power series $Q(z) = \frac{2\sqrt{z}}{\sinh(2\sqrt{z})}$ is has as the first few terms:

$$A_1 = \frac{-2}{3}, \quad A_2 = \frac{2}{45}(-4p_2 + 7p_1^2) \quad A_3 = \frac{-4}{3^2 \cdot 5 \cdot 7} (16p_3 - 44p_2p_1 + 31p_1^3)$$

Then note that :

$$\frac{x}{1 - e^{-x}} = \exp\left(\frac{1}{2}x\right) \frac{\frac{1}{2}x}{\sinh \frac{1}{2}x}$$

Then:

$$T_j(c_1, \dots, c_j) = \sum \frac{1}{2^{4s} r!} \left(\frac{1}{2}c_1\right)^r A_s(p_1, \dots, p_j)$$

where $r + 2s = j$ and p_i are the Pontryagin classes. Recall this gives us:

$$\begin{aligned} T_1(c_1) &= \frac{1}{2}c_1 \\ T_2(c_1, c_2) &= \frac{1}{12}(c_2 + c_1^2) \\ T_3(c_1, c_2, c_3) &= \frac{1}{24}c_2c_1 \\ T_4(c_1, c_2, c_3, c_4) &= \frac{1}{720}(-c_4 + c_3c_1 + 3c_2^2 + 4c_2c_1^2 - c_1^4) \\ T_5(c_1, c_2, c_3, c_4, c_5) &= \frac{1}{1440}(-c_4c_1 + c_3c_1^2 + 3c_2^2c_1 - c_2c_1^3) \\ &\vdots \end{aligned}$$

Then by our earlier work:

$$\mathrm{td}(\xi \oplus \xi') = \mathrm{td}(\xi)\mathrm{td}(\xi')$$

If $q = 1$ and $c_1(\xi) = d \in H^2(X, \mathbb{Z})$, then:

$$\mathrm{td}(\xi) = \frac{d}{1 - e^{-d}}$$

For future reference, we shall require the following factorization. Given the factorization:

$$\sum_{j=0}^q c_j x^j = \prod_{i=1}^q (1 + \gamma_i x)$$

Then we get:

$$\mathrm{td}(\xi) = \prod_{i=1}^q \frac{\gamma_i}{1 - e^{-\gamma_i}}$$

Similarly, we can define the *total Chern Character* (not class):

$$\chi(\xi) = \sum_{i=1}^q e^{\gamma_i}$$

Then we get:

$$\begin{aligned} \mathrm{ch}(\xi \oplus \xi') &= \mathrm{ch}(\xi) + \mathrm{ch}(\xi') \\ \mathrm{ch}(\xi \otimes \xi') &= \mathrm{ch}(\xi)\mathrm{ch}(\xi') \end{aligned}$$

If $q = 1$ and $c_1(\xi) = d \in H^2(X, \mathbb{Z})$, then we get $\mathrm{ch}(\xi) = e^d$. In general, $\chi(\xi) = q + \sum_{k=1}^{\infty} \mathrm{ch}_k(\xi)$ where

$$\chi_k(\xi) = \frac{s_k}{k!} H^{2k}(X, \mathbb{Z}) \otimes \mathbb{Q} \quad s_k = \sum_{i=1}^q \gamma_i^k$$

Theorem 9.6: Relating Todd Class and Chern Character

Let ξ be a continuous $\mathrm{GL}_q(\mathbb{C})$ -bundle over X . Then:

$$\sum_{r=0}^q (-1)^r \mathrm{ch} \left(\bigwedge^r \xi^* \right) = (\mathrm{td}(\xi))^{-1} c_q(\xi)$$

Proof :

Proof: If $\sum_{j=0}^q c_j(\xi)x^j = \prod_{i=1}^q (1 + \gamma_i x)$ by the properties of Chern classes (though, I did not actually prove this result in my vect. bundle notes. thm 4.4.3 in Hirzebruch, p.64)

$$\mathrm{ch} \bigwedge^r \xi^* = \sum e^{-(\gamma_{i_1} + \cdots + \gamma_{i_r})}$$

where the sum is over all combinations $i_1, \dots, i_r$ with $1 \leq i_1 < \cdots < i_r \leq q$. Therefore

$$\sum_{r=0}^q (-1)^r \mathrm{ch} \bigwedge^r \xi^* = \prod_{i=1}^q (1 - e^{-\gamma_i}) = (\gamma_1 \cdots \gamma_q) \prod_{i=1}^q \frac{1 - e^{-\gamma_i}}{\gamma_i} = (\mathrm{td}(\xi))^{-1} c_q(\xi).$$

Now, recall that if we are talking about the Chern class of a complex manifold M , we are talking about the Chern class of its tangent bundle, and the K -genus is defined for such structures. Let

$$Q(x) = \frac{x}{1 - e^{-x}} \quad \text{and} \quad Q(y; x) = \frac{x(y+1)}{1 - e^{-x(y+1)}} - xy$$

Then:

Definition 9.7: Todd Genus

Let M_n be an n -dimensional almost complex manifold. The *Todd genus* (or *T-genus*) is the number:

$$T_n(M_n) = T_n(c_1, \dots, c_n)[M_n]$$

As usual, the $[M_n]$ represents the tangent bundle of M_n . The reader can verify that the direct product of manifolds $X \times Y$ is multiplicative on the Todd class. This means that if

$$0 \rightarrow \xi' \rightarrow \xi \rightarrow \xi'' \rightarrow 0$$

Then:

$$(1 + c'_1 t + \cdots + c'_{q'} t^{q'}) (1 + c''_1 t + \cdots + c''_{q''} t^{q''}) = 1 + c_1 t + \cdots + c_q t^q$$

and so:

$$\mathrm{td}(\xi')(t) \cdot \mathrm{td}(\xi'')(t) = \mathrm{td}(\xi)(t)$$

In many ways, the Todd class is “dual” to the Chern class. For example ([6, p.93]):

$$T_{-1}(M)n = \sum_{p=0}^n (-1)^p T^p(M_n) = c_n[M_n]$$

and:

$$T_1(M)n = \sum_{p=0}^n T^p(M_n) = \tau(M_n)$$

where τ is the index of M_n . More generally:

Proposition 9.8: Properties of Todd Genus

Let B be a commutative ring where $Q \subseteq B$. Then

1. $\sum_{l=0}^n T_n^{(l)}(c_1, \dots, c_n) = c_n$, for all n .
2. $T_n^{(0)}(c_1, \dots, c_n) = \text{td}(c_1, \dots, c_n) (= T_n(c_1, \dots, c_n))$.
- 3.

$$\sum_{l=0}^n T_n^{(l)}(c_1, \dots, c_n) = \begin{cases} 0 & \text{if } n \text{ is odd} \\ L_{\frac{n}{2}}(p_1, \dots, p_{\frac{n}{2}}) & \text{if } n \text{ is even.} \end{cases}$$

The *total Todd class* of a vector bundle, ξ , is

$$\text{td}(\xi)(t) = \sum_{j=0}^{\infty} \text{td}_j(c_1, \dots, c_j) t^j = 1 + \frac{1}{2} c_1(\xi) + \frac{1}{12} (c_1^2(\xi) + c_2(\xi)) t^2 + \frac{1}{24} (c_1(\xi) c_2(\xi)) t^3 + \dots$$

Proof :
exercise

Now, the Chern character is a bit different. If :

$$0 \rightarrow \xi'' \rightarrow \xi \rightarrow \xi' \rightarrow 0$$

then:

$$\text{ch}(\xi)(t) = \text{ch}(\xi')(t) + \text{ch}(\xi'')(t)$$

And similarly Here's the LaTeX code for the equation:

$$\text{ch}(\xi \otimes \eta)(t) = \sum_{i,j} e^{(\gamma_j + \delta_j)t} = \sum_{i,j} e^{\gamma_j t} e^{\delta_j t} = \left(\sum_i e^{\gamma_j t} \right) \left(\sum_j e^{\delta_j t} \right) = \text{ch}(\xi)(t) \text{ch}(\eta)(t).$$

Hence, it defines a ring homomorphism $K(X) \rightarrow H^*(X, \mathbb{Q})$.

Definition 9.9: T-characteristic

Let X be an admissible complex manifold, ξ a $U(q)$ -vector bundle. Then

$$T(X, \xi)(t) = \chi(\xi)(t) \text{td}(\xi)(t)$$

Is called the *T-characteristic* of ξ over X (see [7, p.226]).

The T-characteristic satisfies the duality:

$$(-1)^n T_n^{(l)}(c_1, \dots, c_n) T_n^{(n-l)}(c_1, \dots, c_n)$$

9.4 Pontryagin and Chern Numbers

Let V^n be an oriented compact differentiable manifold where n is the dimension. As V is compact and orientable, it is a cycle, and by cohomology theory V^n is generated by one element. The value of the n -dimensional cohomology class x on the fundamental cycle V^n is denoted $x[V^n]$. if the coefficients of the cohomology group are in A , then:

$$x \in H^n(V^n, A) \quad x[V^n] \in A$$

We can extend this by tensoring by a group B so that :

$$x[V^n] \in A \otimes B \quad x \in H^n(V^n, A) \otimes B$$

For our purposes, let $n = 4k$ and let $p_i \in H^{4i}(V^{4k}, \mathbb{Z})$ be the Pontryagin classes of V^{4k} (i.e. of its tangent bundle). We get a natural grading by the indices i so that given $k = j_1 + j_2 + \dots + j_r$ we get the product $p_{j_1} \dots p_{j_r}$ and an integer (as $A = \mathbb{Z}$):

$$p_{j_1} \dots p_{j_r}[V^{4k}]$$

Let $\pi(k)$ be the number of distinct partitions of k .

Definition 9.10: Pontryagin Numbers and Chern Number

The *Pontryagin numbers* are the $\pi(k)$ numbers :

$$p_{j_1} \dots p_{j_r}[V^{4k}]$$

where $n = 4k$. If n is not divisible by 4, then the Pontryagin number is 0.

Similarly the *Chern numbers* of a complex manifold M (with an almost complex structure so that we are given a particular orientation) are all:

$$c_{j_1} \dots c_{j_r}[V^{2k}]$$

where $n = j_1 + j_2 + \dots + j_r$ and the j_i run through all partitions.

By some characteristic class theory, if M_n is an oriented almost complex manifold, $c_n[M_n] = \chi(M_n, TM_n)$. We shall focus on Pontryagin numbers as we shall need them to prove an important result in cohomology, and return to Chern numbers once we cover the Todd genus in section 9.3.

We can define an equivalence relation where two manifolds $M \sim_p N$ if they have the same dimension and are the same Pontryagin number. Note that if the dimension is not divisible by 4, all manifolds are in the same equivalence relation given by $\sim_p$ as they are all 0.

Now let M, N be smooth $n = 4k$ dimensional manifolds. Then by classical differential geometry we have $T(M \times N) = \pi_M^*(TM) \times \pi_N^*(TM)$ (consider for now the real tangent bundle, so it is a

$GL_n(R)$ -bundle). Then considering the Pontryagin classes of M, N and $M \times N$, by the properties of characteristic classes we get:

$$1 + p_1'' + p_2'' + \cdots = (1 + p_1 + p_2 + \cdots)(1 + p_1' + p_2' + \cdots)$$

Or, as a generating function:

$$\sum_{k=0}^{\infty} p_k'' z^k = \sum_{i=0}^{\infty} f^*(p_i) z^i \sum_{j=0}^{\infty} g^*(p_j') z^j \text{ mod torsion.}$$

In terms of the fundamental class:

$$(f^*(x)g^*(y))[M \times N] = x[M] \cdot y[N] \quad \forall x \in H^n(M, \mathbb{Z}) \otimes B, y \in H^m(N, \mathbb{Z}) \otimes B$$

This last property is simply the properties of m -sequences:

Lemma 9.11: Pontryagin Character and M-sequence

HERE

Proof :

Definition 9.12: K-genus

Let V^n be an oriented n -dimensional manifolds and $\{K_j(p_1, \dots, p_j)\}$ be an m -sequence where $K(V^{4k}) = K_k[V^{4k}]$ and 0 otherwise. Then:

$$\begin{cases} 0 & \dim_{\mathbb{R}} X \not\equiv 0 \pmod{4} \\ K_n(p_1, \dots, p_n)[X] & \dim_{\mathbb{R}} X = 4n \end{cases}$$

is called the K -genus of V^n .

If we are working with Chern classes, then we would have $n = 2k$.

By what we've discussed earlier, the K -genus is multiplicative.

Example 9.13: K-genus

look at [6, p.77].

1. L-genus defined earlier
2. A-genus

10 Some Cobordism

Pontryagin classes will be important characterizing cobordism, and we shall need a bit of cobordism in the proof of HRR. What we shall show that that cobordant manifolds are characterized by

their Pontryagin class. We start by creating the algebraic structure that are naturally induced by Pontryagin classes and then quote the result by Thom that shows it is isomorphic to the cobordant ring.

Definition 10.1: Algebraic Cobordism Ring

Let V^n, W^n be two oriented manifolds of the same dimension. Let $-V^n$ represent the reverse orientation. It is easy to see that:

$$K(V^n \sqcup W^n) = K(V^n) + K(W^n) \quad K(-V^n) = -K(V^n)$$

Thus, $K(-)$ is compatible with the equivalence relation $\sim_p$ and creates an additive group $\tilde{\Omega}^n$. Note that $\tilde{\Omega}^n = 0$ if $4 \nmid n$. Let $\tilde{\Omega} \oplus_n \tilde{\Omega}^n$ so that by the above equations we have the natural grading:

$$\tilde{\Omega}^n \tilde{\Omega}^m \subseteq \tilde{\Omega}^{nm}$$

And so $\tilde{\Omega}$ is a commutative graded (torsion-free) ring.

The ring was called “algebraic” as it was defined using the algebraic properties of Pontryagin classes, we shall soon define the “geometric” cobordant ring that is the equivalence classes of cobordant manifolds and show they are isomorphic. For now, let us characterize the algebraic structure of the algebraic cobordism ring: the goal is to show that

$$\tilde{\Omega} \otimes \mathbb{Q} \cong \mathbb{Q}[x_1, x_2, \dots]$$

To that end, we must find which elements of $\tilde{\Omega}$ can represent each x_i . There is no natural choice, hence we define “candidate sequences” of elements given by manifolds $V^4, V^8, V^{12}, \dots$ that can be chosen to form this basis. We shall show these can be given by complex projective spaces of even dimension. Let us start by recalling that the Pontryagin class of V^{4k} can be written in indeterminate z and decomposed into:

$$1 + p_1 z + \dots + p_k z^k = \prod_{i=1}^k (1 + \beta_i z)$$

Define:

$$s(V^k) = (\beta_1^k + \dots + \beta_k^k)[V^{4k}]$$

By some symmetry function theory, we can think of this as needing the appropriate homogeneity and dimension properties. We shall define a *basis sequence* a sequence where:

$$s(V^{4k}) \neq 0$$

Theorem 10.2: Basis Sequence Correspondence

Let $\{V^{4k}\}$ be a basis sequence and B a [commutative] ring containing $\mathbb{Q}$. Then the set of sequences $\{a_k\}_k$ where $a_k \in B$ are in bijective correspondence to the set of m -sequences $\{K_j(p_1, \dots, p_j)\}$ with coefficients in B where $K(V^{4k}) = a_k$.

Proof :

[6, p.79] (but it is simple)

Theorem 10.3: Complex Projective Forms Basis Sequence

Let $\mathbb{CP}^{2k}$ be the $2k$ -dimensional complex projective space. Then they form a basis sequence where

$$s(\mathbb{CP}^{2k+1}) = 2k + 1$$

Proof :

ibid; simple proof.

Let us now consider $\tilde{\Omega} \otimes \mathbb{Q}$, so now $[V^{4k}]$ can be written as $\frac{a}{b}[V^{4k}]$. Certainly the Pontryagin number, K -genus, and $s(V^{4k})$ are all still well-defined (where the K -genus is well-defined as we required B to contain $\mathbb{Q}$). Now, the Pontryagin number of an element of $\tilde{\Omega} \otimes \mathbb{Q}$ can be a non-integer. This new ring can still be represented using a basis sequence:

Theorem 10.4: Basis Sequence With Rational Numbers Correspondence

Let $\{V^{4k}\}$ be a basis sequence of oriented manifolds, and let $(j) = (j_1, \dots, j_r)$ be a partition of k . Let $V_{(j)} = V^{4j_1} \times \dots \times V^{4j_r}$. Then every $\alpha \in \tilde{\Omega} \otimes \mathbb{Q}$ can be uniquely represented by the sum:

$$\alpha = \sum_{(j) \in \pi(j)} r_{(j)} [V_{(j)}] \quad r_{(j)} \in \mathbb{Q}$$

where the sum goes over all partitions of k . Furthermore, given a system $\{a_{(j)}\}$ of rational numbers, there exists an element $\alpha \in \tilde{\Omega} \otimes \mathbb{Q}$ where

$$p_{(j)}[\alpha] = a_{(j)}$$

Proof :

[6, p.80] (This is some linear algebra (recall the proof in comm. alg. using the Vandermonde Determinant))

Theorem 10.5: Isomorphism of Algebraic Cobordism Ring

$$\tilde{\Omega} \otimes \mathbb{Q} \cong \mathbb{Q}[x_1, x_2, \dots]$$

Proof :

Let $\{V^{4k}\}$ be an arbitrary sequence of oriented manifolds. What's left to show is that:

1. given $\alpha = \sum_{(j)} r_{(j)} [V_{(j)}]$ then $s(\alpha) = r_k s[V^{4k}]$
2. If for all k , every element of $\alpha \in \tilde{\Omega}^{4k} \otimes \mathbb{Q}$ can be represented as a sum $\alpha = \sum_{(j)} [V_{(j)}]$,

then $\{V^{4k}\}$ is a basis sequence

[6, p.80]

There is a natural question of smallest integers can represent a system of Pontryagin numbers and it involves the μ that I skipped.

Lastly, the K -genus can be characterized as being any homomorphism from the algebraic cobordism ring to $\mathbb{Q}$:

Theorem 10.6: Characterizing Rational Homomorphisms

Let $\varphi : \tilde{\Omega} \otimes \mathbb{Q} \rightarrow \mathbb{Q}$ be a homomorphism. Then there exists a K such that $\varphi(-) = K(-)$.

Proof :

easy on p.81

Definition 10.7: Geometric Cobordism Ring

Two oriented smooth n -dimensional manifolds M^n, N^n are *cobordant* if there exist a manifold C such that

$$\partial C \cong (M^n \sqcup -N^n)$$

where $-N^n$ is the reverse orientation; if such a C exists it defines an equivalence relation $M \sim_c N$ which respects disjoint union, reverse orientation, and direct product; $\sqcup, -, \times$. Then Ω^n is the collection of such equivalence classes and:

$$\Omega = \bigoplus_n \Omega^n$$

is the *geometric cobordism ring*.

Certainly this ring is graded, and it is *anti-commutative*, namely:

$$\alpha \cdot \beta = (-1)^{|\alpha||\beta|} \beta \cdot \alpha \quad \alpha \in \Omega^n, \beta \in \Omega^m$$

He now goes onto to prove the isomorphism between this ring and the algebraic one, skipping for now.

An important theorem:

Theorem 10.8: Hirzebruch Signature Theorem

Let M be a real differentiable oriented manifold, and let $I(M)$ be its index. Then:

1. if $\dim_{\mathbb{R}} M \not\equiv 0 \pmod{4}$, $I(M) = 0$
2. if $\dim_{\mathbb{R}} M \equiv 0 \pmod{4}$, then:

$$I(M) = L_k(p_1, \dots, p_k)[M]$$

Proof :
[7, p.231].

10.1 Index and Virtual index

[6, p.84-90]

11 RR for algebraic manifolds (i.e. projective varieties)

here

Index

1st Cohomology Group, 18

Abel-Jacobi Map, 40

Abel-Jacobi Theorem, 40

Algebraic Cobordism Ring, 60

Almost Complex Structure, 47

Canonical Curve, 32

Canonical Map, 30

Chern Numbers, 58

Degree of Smooth Curve, 37

Geometric Cobordism Ring, 62

Hyperelliptic and Nonhyperelliptic, 32

Hyperplane Section of Riemann Surface, 37

Jacobian Variety, 39

K-Chern genus, 59

K-genus, 59

K-Pontryagin genus, 59

Laurent Principal Part, 11

Linearly Equivalent, 10

Multiplicative Sequence, 49

Normalization, 5

Period Vector, 18

Picard Variety, 40

Pontryagin Numbers, 58

Ramification Divisor, 7

T-characteristic, 57

Todd Genus, 56

References

- [1] Nathanael Chwojko-Srawley. *Everything You Need To Know About Elliptic Curves*. NA. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_elliptic_curves.pdf.
- [2] Nathanael Chwojko-Srawley. *Everything You Need To Know About Algebraic Geometry*. 1 vols. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_algGeo.pdf.
- [3] Nathanael Chwojko-Srawley. *Everything You Need To Know About Complex Analysis*. 1st ed., p. 233. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_complex_analysis.pdf.
- [4] Joseph H. Silverman. *Advanced Topics in the Arithmetic of Elliptic Curves*. Graduate Texts in Mathematics. Springer, 1994. ISBN: 978-0-387-94328-2 978-1-4612-0851-8. URL: <http://link.springer.com/10.1007/978-1-4612-0851-8>.
- [5] Phillip A. Griffiths. *Introduction to Algebraic Curves*. Vol. 78. Mathematical Monographs. American Mathematical Society.
- [6] Friedrich Hirzebruch. *Topological Methods in Algebraic Geometry*. 1st ed. Springer, 1995. URL: <https://link-springer-com.myaccess.library.utoronto.ca/book/10.1007/978-3-642-62018-8>.
- [7] Jean Gallier and Stephen Shatz. *Complex Algebraic Geometry*. Vol. 1. 2011.