

Clock Arithmetic and the Mysteries of Prime Numbers

Foreword

This article is written for my patient friends to whom I give math riddles that require knowledge of modular arithmetic, or as I often call it *clock arithmetic*. This article serves as an easy reference for the necessary concepts and requires no mathematical background. For this entire article, we will only care about which hour the hour-hand is on; no AM or PM, and no adding half an hour (or anything that isn't an increment of 60 minutes).

Note that any exercise can be skipped; they are just there for fun!

1 Clock Arithmetic: Addition and Multiplication

We all have (at least) two ideas of addition in our heads. The first one is the usual notion of adding quantities¹: If I have 4 bottles and add 10 bottles, I have 14 bottles. In other words

$$4 + 10 = 14$$

The other notion of addition we often use comes from when we are calculating what time it is. Consider a 12-hour clock. If it is 4 o'clock, in 5 hours it will be 9 o'clock. If instead we add 10 hours, it will be 2 o'clock. In other words, we just comfortably did

$$4 + 10 = 2$$

in our heads. This is certainly different than the previous equality, but it still makes sense! Using the same symbol “+” may cause confusion, so to distinguish the two types of addition, we shall write $+_{12}$ for “12 hour clock” addition, and keep $+$ for quantity addition. Thus, we can write:

$$4 + 10 = 14 \quad \text{and} \quad 4 +_{12} 10 = 2$$

Let us call the addition using $+$ *natural arithmetic* and the addition using $+_{12}$ *clock arithmetic*. As good mathematicians, we should see what similarities and differences these different ideas of additions have.


The first thing we should try to do is add various hours to each other and see what we get. Let us start by just picking two random numbers and playing around with them. If it is 3 o'clock, then if I add 5 it will be 8 o'clock. But if I add 17 hours to 3 o'clock, it will *still* be 8 o'clock. More generally, adding any multiple of 12 will not change the result of my addition:

$$3 +_{12} 5 = 8 \quad 3 +_{12} 17 = 8 \quad 3 +_{12} 29 = 8 \quad 3 +_{12} 41 = 8 \cdots$$

So adding 5, $(5 + 1 \cdot 12)$, $(5 + 2 \cdot 12)$, ... to 3 gives the same result. Similarly, if I have any number n

¹A mathematician would say *discrete quantities* to distinguish between *continuous* quantities, which would describe something like water where the amount we add is on a “continuous” spectrum of possible amounts rather than “discrete” indivisible chunks of amounts

between 1 and 12, if I add b , or I add $(b + k \cdot 12)$ for some $k = 1, 2, \dots$ I get the same result.



$$3 +_{12} 12 = 3$$

$$3 +_{12} 24 = 3$$

- Starting position ($n = 3$)
- Adding 12 hours
- Adding 24 hours

+12 hours (1 full rotation)

+24 hours (2 full rotations)

We would like to remember that adding $k \cdot 12$ to any given hour gives the same result. Mathematicians usually use the symbol $\equiv_{12}$ between two such numbers:

$$5 \equiv_{12} 17 \equiv_{12} 29 \equiv_{12} \dots$$

where we say that 5 is *equivalent* to 17, which is *equivalent* to 29, and so on. Put differently, two numbers $n + a, n + b$ are equivalent if they give you the same time on the clock-face after addition:

$$n +_{12} a = m \quad \text{and} \quad n +_{12} b = m$$

Therefore, in clock arithmetic we only need to work with the following numbers:

$$1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12$$

All these numbers are *not* equivalent to each other (ex. $3 \not\equiv_{12} 5$). Adding any two of these numbers using $+_{12}$ will give another number in the above list:

$$1 +_{12} 1 = 2 \quad 11 +_{12} 7 = 6 \quad 9 +_{12} 12 = 9$$

The last of these three examples is somehow different than the others; it is the only equation where one of the numbers on the left hand side of the addition $+_{12}$ also appears on the right hand side of the equality; namely 9 appears on both sides. In fact, if we try adding any other number to 9, we see that only adding 12 returns 9:

$9 +_{12} 1 = 10$	$9 +_{12} 5 = 2$	$9 +_{12} 9 = 6$
$9 +_{12} 2 = 11$	$9 +_{12} 6 = 3$	$9 +_{12} 10 = 7$
$9 +_{12} 3 = 12$	$9 +_{12} 7 = 4$	$9 +_{12} 11 = 8$
$9 +_{12} 4 = 1$	$9 +_{12} 8 = 5$	$9 +_{12} 12 = 9$

The natural next question to ask is if adding 12 to any number n , not just 9, will return n . Try doing this for any n between 1 and 12 (including 12), and you'll see that adding 12 to any number in clock arithmetic gives back the same number, and 12 is the *only* number that does this!

$$n +_{12} 12 = n$$

Notice that in natural arithmetic, this is exactly the behaviour of 0! Namely, for any natural number n :

$$n + 0 = n$$

With this parallel in mind, when we are doing clock arithmetic, we should think of 12 as being the same as 0. Mathematicians would even go so far as re-labeling 12 as 0 so that the 12 numbers of clock arithmetic are:

$$0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11$$

This gives us a good understanding of addition in clock arithmetic: any number n can be re-written as one of the above 12 numbers since n is equivalent to one of these, and then it becomes easy to perform arithmetic's.

What about multiplication? Is there a natural way to multiply hours with each other. In natural arithmetic, we know that multiplying n by some natural number m means we are adding n to itself m times. Furthermore, $n \cdot 0 = 0$ ². Inspired by these properties, let us define multiplication $n \cdot_{12} m$ to mean that we are starting at 12 o'clock (which we should think of as 0 o'clock), and we are moving forward n hours a total of m times.

This matches our intuition on repeated addition. One difference from natural arithmetic and clock arithmetic is that in clock arithmetic there are numbers that are equivalent to each other. We should verify that if we multiply by two equivalent numbers, we ought to get back equivalent results. More precisely, if $n \cdot_{12} m = a$ and $n \cdot_{12} m' = b$, then if $m \equiv_{12} m'$ we want that $a \equiv_{12} b$ (try this out with some numbers!)

Let us start by showing that multiplying by 0 and multiplying by any number equivalent to 0 gives the same result (ex. $n \cdot_{12} 0 = n \cdot_{12} 12 = 0$). Let us first test if this is the case with an example. Let $n = 3$. Certainly if we add no multiples of 3 we stay at 12 o'clock (again, this should be thought of as 0 o'clock). If we add 12 multiples of 3, then we are adding 3 o'clock to itself 12 times. This brings us 36 hours forward, which on a clock-face is 12 o'clock. But we already saw that we should think of 12 as being 0! Hence, we see that adding 3 to itself 12 times will return 0. In fact, if we start with any number n between 0 and 11, adding the number to itself 12 times will always bring us back to the 12 o'clock position! Therefore, 12 really *does* behave like 0 in clock arithmetic! We just showed that multiplying by 12 is like multiplying by 0. To go back to our example using 3:

$$3 \cdot_{12} 12 = 0$$

because 36 hours after 12 o'clock is again 12 o'clock. In our re-labeled system (where 12 is 0) this becomes:

$$3 \cdot_{12} 0 = 0$$

Exercise 1: Multiplying by equivalent numbers

The above shows that if $0 \equiv_{12} m$, then $n \cdot_{12} m = 0$. Show this is true more generally:

$$\text{Let } n \cdot_{12} m = a \text{ and } n \cdot_{12} m' = b. \text{ If } m \equiv_{12} m', \text{ then } a \equiv_{12} b$$

²For those who have contested this by asking "what if we let n be infinity", there is [this article](#) addressing exactly how to think about infinity and why we shouldn't conclude $0 \cdot \infty = 0$ without putting in more care

Exercise 2: Multiplying by other numbers

When multiplying in natural arithmetic we get that $a \cdot b = 0$ if and only if $a = 0$ or $b = 0$. In clock arithmetic, it is now possible that $a \cdot_{12} b = 0$ but $a, b \neq 0$!

Can you think of two non-zero numbers a, b where $a \cdot_{12} b = 0$?

Exercise 3: Combining Addition and Multiplication

Show that for any natural numbers a, b, c

$$a \cdot_{12} (b +_{12} c) = (a \cdot_{12} b) +_{12} (a \cdot_{12} c)$$

Mathematicians call this *distributivity*

Exercise 4: Subtraction

Subtraction works just the same as addition. Convince yourself that:

$$1 -_{12} 2 = 11$$

that is, if it is 1 o'clock, two hours ago it was 11 o'clock. Convince yourself that:


$$a -_{12} 12 = a$$

2 From Clock Arithmetic to Modular Arithmetic

So we just showed how we can think of clock arithmetic which has 12 numbers. A mathematician would then naturally ask the following question:

What if we replace 12 with any other natural number³?

Let us do exactly this and see what we get. First of all, instead of writing $+_{12}$, we shall now write $+_n$. Let us call the arithmetic for all $+_n$ *modular arithmetic*. With this vocabulary, clock arithmetic is a special case of modular arithmetic. To distinguish the different modular arithmetics, we would usually say we are working *modulo* n . Clock arithmetic would be modular arithmetic modulo 12. When working modulo n , we can imagine a clock face with n positions. For example, if $n = 5$, then we can think of having a clock with 5 hours:



On this clock, $3 +_5 3 = 1$ and $2 \cdot_5 4 = 3$. When working modulo n , we will always have n numbers:

$$0, 1, 2, \dots, n-1$$

Just as for clock arithmetic, we can treat n like 0.

Exercise 5: Zero in Modular Arithmetic

Let a, n be any natural numbers. For a moment, let us not label n as 0. Convince yourself that:

$$a \cdot_n n = n$$

This shows that in modular arithmetic mod n , we can *always* treat n as 0, and hence why we will always write 0 instead of n .

Exercise 6: Special Property when Modulo is Prime

Remember in exercise 2 that in modulo 12, we can find two natural numbers a, b where $a \cdot_{12} b = 0$, but $a, b \neq 0$.

Convince yourself that if n is a prime number (ex. $n = 2, 3, 5, 7, 11, \dots$) then

$$a \cdot_n b = 0 \quad \text{if and only if} \quad a = 0 \text{ or } b = 0$$

This shows that primes are somehow “special” in that they behave more like natural arithmetic. Mathematicians even have a different word for when n is a prime: modular arithmetic modulo n where n is a prime is called a *finite field*.

³A natural number is of the form $1, 2, 3, \dots, n, \dots$. We in fact *can* replace it with non-natural numbers, but it took history a long time to understand what this means; this would be introduced in a course in *abstract algebra* (see[3] if you want my exposition on abstract algebra)

Applications

So now that we have defined modular arithmetic, what can we do with it? The rest of this article will now go over some interesting problems that demonstrate the usefulness of modular arithmetic, as well as reveal some hidden patterns in prime numbers

3 Solving Equations

The principal reason mathematicians care about modular arithmetic is that it allows us to give a lot more information about equations we are trying to solve. For example, consider $x^2 + x + 1$. A natural question to ask is whether there is an integer n (numbers of the form $\pm 1, \pm 2, \pm 3, \dots$) that solves this equation. In other words, we are looking for an n such that

$$n^2 + n + 1 = 0$$

It may take you a little while to figure out if such a number exists. A natural starting point is to just convince yourself that no small integer can solve it. By plugging in $\pm 1, \pm 2, \pm 3$ we get:

x	Calculation	Result
1	$1^2 + 1 + 1$	3
-1	$(-1)^2 + (-1) + 1$	1
2	$2^2 + 2 + 1$	7
-2	$(-2)^2 + (-2) + 1$	3
3	$3^2 + 3 + 1$	13
-3	$(-3)^2 + (-3) + 1$	7
$\vdots$	$\vdots$	$\vdots$

Doing this for the first 100 integers and their negatives would show you that the equation never equals 0, however there is no guarantee that you will *never* find a big enough number n such that $n^2 + n + 1 = 0$. Using modular arithmetic, it is in fact straightforward to show that *no* solutions exist.

To show this, let's assume that there *does* exist an n such that $n^2 + n + 1 = 0$ ⁴. Then since $n^2 + n + 1 = 0$, certainly $n^2 + n + 1 \equiv_2 0$: if we let $n^2 + n + 1 = m$ then $m \equiv_2 0$ means that m is a multiple of 2, but 0 is certainly a multiple of 2 since $0 \cdot 2 = 0$. What we will show is that it is *impossible* to find an integer n such that $n^2 + n + 1 \equiv_2 0$, in other words this equation has no solutions modulo 2.

First of all, for any integer n :

$$n \equiv_2 0 \quad \text{or} \quad n \equiv_2 1$$

⁴This is a usual trick called *proof by contradiction*. This idea is to assume the statement we want to prove false is true, and show this creates a contradiction

For example, if $n = -177$, then:

$$-177 = 1 + 2 \cdot (-89) \equiv_2 1$$

(look at exercise 4 if the subtraction is confusing). Thus, every natural number is equivalent to either 1 or 0. Thus, armed with the possible values of n modulo 2, we get one of two possible equations:

$$(1 \cdot_2 1) +_2 1 +_2 1 = 0 \quad \text{or} \quad (0 \cdot_2 0) +_2 0 +_2 1 = 0$$

But hang on, if we compute the left hand side of the first equation we get

$$1 +_2 1 +_2 1 = 0 +_2 1 = 1$$

which is *not* equal to the right hand side (i.e. $1 \neq 0$ and $1 \not\equiv_2 0$). The second equation is even more obvious since $(0 \cdot_2 0) +_2 0 +_2 1 = 1$, which is again not equal to 0.

So we arrived at a contradiction, because $0 \neq 1$ ⁵. What this tells us is that we made a wrong assumption earlier in the proof; and indeed this started by assuming that there exists an integer n such that $n^2 + n + 1 = 0$. Thus, it is *impossible* that such an integer exists! Notice how easy this was to prove: we only had to check *two* possible numbers in modular arithmetic modulo 2 as compared to the infinite possibilities when trying to solve the equation with natural numbers!

This “reduction to finitely many cases” is one of the **key** properties of modular arithmetic that mathematicians abuse to no end.

4 Application: Patterns in Prime Numbers

Let us now show that modular arithmetic reveals hidden patterns in primes. Consider the equation $x^2 = p$ for some prime number p . Then certainly no natural number solves this equation, because no natural number can be the square-root of a prime⁶. However, this question gets more interesting if we instead consider it in *modular arithmetic* modulo n .

let us first be precise what it means for a number to be a square root. A number a is the square root of a number m if $a^2 = m$. For example, 4 is the square-root of 16 because $4^2 = 16$. If m is a prime, then there is no *natural number* a such that $a^2 = p$.

So in natural arithmetics, square-roots of prime do not exist. Let us see if we can find a square-root in modular arithmetic in some modulo n . Let us demonstrate this with an example. Consider $p = 3$. Let us see if in modulo n , we can find a number m such that $m \cdot_n m = p$ (i.e. if p has a square-root modulo n). The following is a table where n goes over the first few prime numbers:

n	1 ²	2 ²	3 ²	4 ²	5 ²	6 ²	7 ²	8 ²	9 ²	10 ²
2	1									
3	1	1								
4	1	0	1							
5	1	4	4	1						
6	1	4	*3*	4	1					
7	1	4	2	2	4	1				
8	1	4	1	0	1	4	1			
9	1	4	0	7	7	0	4	1		

⁵The suspicious reader may ask why is $0 \neq 1$. In fact, there is nothing wrong with the arithmetic system where $0 = 1$, however if we let $0 = 1$ we get that every number equals 0 (convince yourself), and then nothing interesting happens.

⁶This is because the square-root of a prime number is *irrational*; see [2] for a friendly introduction to this notion

10	1	4	9	6	5	6	9	4	1	
11	1	4	9	5	*3*	*3*	5	9	4	1

Let us focus on the cases where n is a prime number. Notice that $5 \cdot_{11} 5 = 3$, since $5 \cdot 5 = 25 = 3 + 2 \cdot 11$. Therefore, in modulo 11, we have that:

$$\sqrt{3} = 5 \text{ (in modulo 11)}$$

Precisely because $5 \cdot_{11} 5 = 3$.

Hence, for a mysterious reason, there *does* exist a square-root of 3 in modulo 11, and there *doesn't* appear to be a square-root of 3 modulo 5 or 7 (among many other primes). But why 11? Why was the prime 3 somehow linked to the prime 11? The reader can check that there are in fact other primes in which 3 also has a square-root; so how can we determine for which primes 3 has a square-root?

This is a rather fascinating result: some primes have square-roots in modulo p , while other don't, in other words primes somehow have *relations* between each other! To be precise, two primes are tied to each other depending on whether we can take the square root of one modulo the other! Euler in the 19th century had the exact same realization, and did thousands of such computations looking for patterns. It turns out to be a rather hard problem, and it wasn't until Gauss around 50 years later that gave a definitive and satisfactory explanation how such primes relate via square-rooting: the result is now called *quadratic reciprocity*. As it turns out, there is a reason this question couldn't be quickly solved by Euler; answering this question would require a lot more math than this article covers (all the details are in [3], and requires a good deal of build-up before being proven in chapter 22.4). However, I strongly encourage the reader play around with this; after a few hours or a few days, the reader will most likely stumble upon the same pattern that many other mathematicians who worked on the problem (including Euler) have discovered!

5 Final Mystery

The problem in the above section turned out to be one of the first cornerstones in number theory, as it somehow connects different prime numbers to each other via equations; that is, somehow prime numbers are not completely “random” but can have subtle connections that are not obvious at first glance!

Here is the final problem we shall present:

Are there infinitely, or finitely, many primes p such that

$$p \equiv_5 3$$

In other words, are there finitely many or infinitely many primes for which p o'clock in a 5 hour clock will be the same time as 3 o'clock.

The numbers 3 and 5 can be replaced with any numbers a, b where $\gcd(a, b) = 1$ (i.e. they have no common factors).

For example,

prime	2	3	5	7	11	13	17	19	23	29	31	37	41	43	47	53	59	...
$p \equiv_5 m$	2	3	0	2	1	3	2	4	3	4	1	2	1	3	2	3	4	...

Table 2: First 17 primes and their values modulo 5

This question was pondered by Dirichlet in 1837. Essentially, it is asking if there is a meaningful way in which prime numbers can be grouped together. A mathematician would say that this is asking how the primes numbers are *distributed*. Just like the above problem, I strongly encourage the reader to think about this problem for a couple of hours and just see what they can come up with. Is the same true if one of the numbers is replaced by a non-prime number? The result has been proven, and a full proof requires more math than what can be covered in an expository article like this (for completeness, a proof can be found in [1]).

Bibliography

- [1] Nathanael Chwojko-Srawley. *Everything You Need To Know About Class Field Theory*. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_class_field_theory.pdf.
- [2] Shay Fuchs. *Introduction to Proofs and Proof Strategies*. Cambridge University Press. URL: <https://www.cambridge.org/highereducation/books/introduction-to-proofs-and-proof-strategies/3E17FC9C3CA060246BF1B5476687AEB0#overview>.
- [3] Nathanael Chwojko-Srawley. *Everything You Need To Know About Undergraduate Algebra*. 1st ed. URL: https://nathanaelsrawley.com/assets/pdfs/notes/EYNTKA_algebra.pdf.