

To Do

Read Sections 5.1 - 5.3 (Hypothesis Testing)

Do End-of-Chapter Problems 1 – 8 for Midterm Test 2.

Assignment 3 due Friday November 11.

See detailed information posted on Learn regarding material covered by Midterm Test 2 (4:40 - 6:10 on Tuesday November 15).

Last Class

- (1) Testing $H_0: \mu = \mu_0$ when σ is unknown for $G(\mu, \sigma)$ model.**
- (2) Statistical significance versus practical significance.**
- (3) Relationship between tests of hypothesis and confidence intervals**

Testing $H_0: \mu = \mu_0$ when σ is unknown

(1) Test $H_0: \mu = \mu_0$ based on observed random sample y_1, y_2, \dots, y_n from a $G(\mu, \sigma)$ distribution.

(2) Use the test statistic $D = \frac{|\bar{Y} - \mu_0|}{S / \sqrt{n}}$

with observed value $d = \frac{|\bar{y} - \mu_0|}{s / \sqrt{n}}$

(3) Calculate

$$p\text{-value} = 2[1 - P(T \leq d)] \text{ where } T \sim t(n-1)$$

(4) Draw conclusion based on p -value.

Relationship Between Tests of Hypothesis and Confidence Intervals

The p -value for testing $H_0: \theta = \theta_0$ is greater than or equal to q

if and only if

the value $\theta = \theta_0$ is inside a $100(1-q)\%$ confidence interval for θ
(assuming we use the same pivotal quantity).

Today's Class

- (1) Testing $H_0: \sigma^2 = \sigma_0^2$ when μ is unknown.
- (2) Likelihood Ratio Test for testing $H_0: \theta = \theta_0$.

$H_0: \sigma^2 = \sigma_0^2$ when μ is unknown

Suppose Y_1, Y_2, \dots, Y_n is a random sample from a $G(\mu, \sigma)$ distribution.

Recall the pivotal quantity

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$$

which is used to construct confidence intervals for σ and σ^2 .

$H_0: \sigma^2 = \sigma_0^2$ when μ is unknown

To test $H_0: \sigma^2 = \sigma_0^2$ we use the test statistic

$$U = \frac{(n-1)S^2}{\sigma_0^2}$$

If $H_0: \sigma^2 = \sigma_0^2$ is true then

$$\frac{(n-1)S^2}{\sigma_0^2} \sim \chi^2(n-1)$$

$H_0: \sigma^2 = \sigma_0^2$ when μ is unknown

Large and small values of

$$U = \frac{(n-1)S^2}{\sigma_0^2}$$

provide evidence against H_0 . Why?

The Chi-squared distribution is not symmetric about its mean which makes the determination of “large” and “small” values slightly more challenging.

$H_0: \sigma^2 = \sigma_0^2$ when μ is unknown

The following calculation approximates the p -value:

- (1) Let $u = (n-1)s^2/\sigma_0^2$ denote the observed value of U from the data.**
- (2) If $P(U \leq u) < 0.5$ then $p\text{-value} = 2P(U \leq u)$
If $P(U \geq u) < 0.5$ then $p\text{-value} = 2P(U \geq u)$
where $U \sim \chi^2(n-1)$.**

Note: Only one of these values will be between 0 and 1 and this is the value you want.

Female Handspan Example

Suppose we want to test the hypothesis $H_0: \sigma^2 = 1$.

For these data $s^2 = 2.055$

Previously we determined a 95% confidence interval for σ^2 which was $[1.491, 3.014]$.

Since $\sigma^2 = 1$ is not in this interval we already know that the $p\text{-value} \leq 0.05$.

Female Handspan Example

Following the suggested steps

(1) Calculate observed value of U :

$$\begin{aligned} u &= (n-1)s^2/\sigma_0^2 \\ &= (63)(2.055)/1 = 129.464375/1 = 129.46 \end{aligned}$$

$$\begin{aligned} (2) \text{ } p\text{-value} &= 2P(U \geq u) \\ &= 2P(U \geq 129.46) \approx 0 \end{aligned}$$

where $U \sim \chi^2(63)$.

There is very strong evidence against $H_0: \sigma^2 = 1$ based on the observed data.

Likelihood Ratio Test Statistic

Recall that, an exact pivotal quantity does not always exist for constructing a confidence interval for a parameter θ .

In these cases we used an approximate Normal pivotal quantity or the approximate pivotal quantity based on the likelihood ratio statistic:

$$\Lambda = -2 \log \left[\frac{L(\theta)}{L(\tilde{\theta})} \right] \sim \chi^2(1) \text{ approximately}$$

Likelihood Ratio Statistic

$$\Lambda = -2 \log \left[\frac{L(\theta)}{L(\tilde{\theta})} \right] \sim \chi^2(1) \text{ approximately}$$

Recall that we used this result to justify the fact that a 15% likelihood interval is an approximate 95% confidence interval.

We can also use this result to construct an approximate hypothesis test of $H_0: \theta = \theta_0$.