

## Section 2: Analysis

In this section on Analysis, use the Section 2 Answer Sheet for your answers. There are 8 questions. For multiple choice questions there is only one correct answer.

### Question 1 (3 marks)

Determine which of the following is the derivative of  $\frac{\sin x + x}{2 + \cos x}$ . Enter your answer by crossing exactly one of the boxes on your answer sheet.

- a.  $\frac{\cos x + 1}{(2 - \sin x)^2}$
- b.  $\frac{(\cos x + 1)(2 + \cos x) - (\sin x + x) \sin x}{(2 - \sin x)^2}$
- c.  $\frac{\cos x + 1}{-\sin x}$
- d.  $\frac{\sin x + x}{-\sin x}$
- e.  $\frac{(\cos x + 1)(2 + \cos x) + (\sin x + x) \sin x}{(2 + \cos x)^2}$

(3 marks)

**Solution:** Using the product rule and the chain rule

$$\begin{aligned}\frac{d}{dx} \left( \frac{\sin x + x}{2 + \cos x} \right) &= \left( \frac{d}{dx} (\sin x + x) \right) * (2 + \cos x)^{-1} \\ &\quad + \left( \frac{d}{dx} (2 + \cos x)^{-1} \right) * (\sin x + x) \\ &= (\cos x + 1) * (2 + \cos x)^{-1} \\ &\quad + \sin x (2 + \cos x)^{-2} (\sin x + x)\end{aligned}$$

Hence E is correct.

### Question 2 (2 marks)

If  $f'(2) = 5$ ,  $g(4) = 2$ ,  $g(2) = 1$ ,  $f(2) = -1$  and  $g'(4) = 3$ , determine the value of  $\frac{d}{dx} (f(g(x)))$  at the point  $x = 4$ . The answer is an integer between -99 and 99. Enter your answer according to the instructions.

(2 marks)

**Solution:**  $\frac{d}{dx} (f(g(x))) = g' f'(g(x)) = g'(4) f'(2) = 15$

### Question 3 (2 marks)

If  $f(x, y) = \cos\left(\frac{x}{y}\right)$  determine which of the following is  $\nabla f$ . Enter your answer by crossing exactly one of the boxes on your answer sheet.

- a.  $\left(-\frac{1}{y} \sin \frac{x}{y}, \frac{x}{y^2} \sin \frac{x}{y}\right)$
- b.  $\left(-\sin \frac{x}{y}, \frac{1}{y^2} \sin \frac{x}{y}\right)$
- c.  $\left(\sin \frac{x}{y}, -\frac{1}{y^2} \sin \frac{x}{y}\right)$
- d.  $\left(-\frac{x}{y} \sin \frac{x}{y}, \frac{x}{y^2} \sin \frac{x}{y}\right)$
- e.  $\left(\frac{1}{y} \sin \frac{x}{y}, -\frac{x}{y^2} \sin \frac{x}{y}\right)$

(2 marks)

**Solution:** a is correct

**Question 4** (3 marks)

Determine the derivative of  $f(x, y)$  in the  $(3, -4)$  direction at the point  $(\pi, 2)$ , rounded to three decimal places. The answer is a real number between  $-1$  and  $1$ . Enter your answer according to the instructions.

(3 marks)

**Solution:**

$$\begin{aligned}
 \|\vec{v}\| &= \sqrt{3^2 + (-4)^2} = 5 \\
 \vec{u} &= \frac{\vec{v}}{\|\vec{v}\|} = \left(\frac{3}{5}, -\frac{4}{5}\right) \\
 \nabla f \cdot \vec{u} &= -\frac{1}{5} \left(\frac{3}{y} + \frac{4x}{y^2}\right) \sin \frac{x}{y} \\
 \nabla f \cdot \vec{u}(\pi, 2) &= -\frac{1}{5} \left(\frac{3}{2} + \frac{4\pi}{4}\right) \sin \frac{\pi}{2} \\
 &= -0.92831853071 = 0.928 \text{ to three dp}
 \end{aligned}$$

**Question 5** (5 marks)

Consider the pair of complex numbers  $z_1 = 10 + 8i$  and  $z_2 = \frac{3-(62/3)i}{1-(4/3)i}$ . Determine the answers to the following questions. Enter your answer for each question by crossing exactly one box, labelled  $z_1$  or  $z_2$ , on the answer sheet. The allocation of marks per question is indicated in brackets.

- A. Which is closest to the origin? (1 mark)
- B. Which has rotated by the least amount in absolute terms? (2 marks)
- C. Which is closest to  $z_3 = 1 + i$ ? (2 marks)

(5 marks)

**Solution:** Answers: a.  $z_2$ , b.  $z_1$ , c.  $z_1$

$$\frac{3 - (62/3)i}{1 - (4/3)i} * \frac{1 + (4/3)i}{1 + (4/3)i} = 11 - 6i$$

$$|z_1| = \sqrt{10^2 + 8^2} = 12.81$$

$$|z_2| = \sqrt{11^2 + 6^2} = 12.53$$

Hence  $z_2$  is closer to the origin.

$$\arg(z_1) = \tan^{-1}\left(\frac{8}{10}\right) = 0.675$$

$$\arg(z_2) = \tan^{-1}\left(\frac{11}{-6}\right) = -1.071$$

Hence  $z_1$  has rotated the least.

$$|z_1 - z_3| = \sqrt{9^2 + 7^2} = \sqrt{130}$$

$$|z_2 - z_3| = \sqrt{10^2 + (-6)^2} = \sqrt{136}$$

Hence  $z_1$  is closest to  $z_3$ .

**Question 6** (2 marks)

For the 1st order ordinary differential equations (ODE) 1-5 below, classify them according to the following terms:

- A. Order, a natural number.
- B. Linear or non-linear.
- C. Homogeneous or non-homogeneous.
- D. Has constant coefficients or does not have constant coefficients.

1.  $y' - e^x y + 7 = 0$
2.  $y' - e^x y = 0$
3.  $3y'' - 2y' = 3$
4.  $2\frac{d^3y}{dx^3} + \cos x \frac{dy}{dx} = 0$
5.  $2\frac{d^5y}{dx^5} + \cos y \frac{d^3y}{dx^3} = 0$

Enter your answers as follows. For each of the 5 equations, cross exactly one of the boxes in the first table, in the column representing the order of the equation. Then, cross all that apply in the second table, where L stands for linear, H stands for homogeneous, and C stands for constant coefficients.

(2 marks)

**Solution:** Answers:

1. 1,1,2,2
2. 1,1,1,2
3. 2,1,2,1
4. 3,1,1,2
5. 3,2,1,2

**Question 7** (5 marks)

The function  $y(t)$  satisfies the 2nd order ODE  $\frac{d^2y}{dt^2} - 2\frac{dy}{dt} + 2 = 0$  with  $y(0) = 1$  and  $y\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}e^{\frac{\pi}{4}}$ . Given that  $\left(y\left(\frac{\pi}{3}\right)\right)^3 = \frac{1}{A}e^{B\pi}$ , determine the values of  $A$  and  $B$ . The answers are integers between -99 and 99. Enter your answers according to the instructions.

(5 marks)

**Solution:** Answer:  $A = 8$ ,  $B = 1$

Solution:

$$\begin{aligned}r^2 - 2r + 2 &= 0, \\ \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} &= 1 \pm i, \\ y(t) &= e^t (M \cos t + N \sin t), \\ y(0) &= 1 = M, \\ y(\pi/4) &= e^{\pi/4} \left( \frac{1}{\sqrt{2}} + N \frac{1}{\sqrt{2}} \right) = \frac{1}{\sqrt{2}} e^{\pi/4} \implies N = 0 \\ y(t) &= e^t \cos t \\ y(\pi/3) &= e^{\pi/3} / 2 \\ (y(\pi/3))^3 &= e^{\pi} / 8\end{aligned}$$

END OF QUESTIONS