

Lecture 9 - Projective Alignment

Robust Model Fitting

$$\min_T \sum_i \|T p_i - p_i'\|^2$$

$$\begin{bmatrix} \frac{ax+by+c}{gx+hy+k} \\ \frac{dx+ey+f}{gx+hy+k} \\ 1 \end{bmatrix} = \begin{bmatrix} \hat{x}/\hat{w} \\ \hat{y}/\hat{w} \\ 1 \end{bmatrix} \sim \begin{bmatrix} \hat{x} \\ \hat{y} \\ \hat{w} \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & k \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$r_{x_i}(H) = ax_i + by_i + c - x_i' (gx_i + hy_i + k)$$

$$r_{y_i}(H) = dx_i + ey_i + f - y_i' (gx_i + hy_i + k)$$

$$A \vec{x} = \vec{b}$$

$$\begin{bmatrix} x_1 & y_1 & 1 & 0 & 0 & 0 & -x_1'x_1 & -x_1'y_1 & -x_1' \\ 0 & 0 & 0 & x_1 & y_1 & 1 & -y_1'x_1 & -y_1'y_1 & -y_1' \\ \vdots & & & & & & & & \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \\ e \\ f \\ g \\ h \\ k \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \end{bmatrix}$$

$$\min_T \|A \vec{x}\|^2$$

$$\vec{x} = \vec{0}$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{s.t. } \|\vec{x}\| = 1$$


subject to

$$= 1$$

$$\min \|A \vec{x}\|^2 = \boxed{\vec{x}^T A^T A \vec{x}}$$

Singular value Decomposition

$$A = U \Sigma V^T$$



orthogonal: $U^T U = I$

$$\vec{x}^T (V \Sigma U^T) (U \Sigma V^T) \vec{x}$$

$$\vec{x}^T V \Sigma \boxed{\Sigma V^T \vec{x}}$$


$$\begin{bmatrix} \sigma_1 & & \\ & \sigma_2 & \\ & & \ddots \\ & & & \sigma_n \end{bmatrix} \begin{bmatrix} - & v_1 & - \\ - & v_2 & - \\ & \vdots & \\ & & v_n \end{bmatrix} \begin{bmatrix} \\ \\ \\ \vec{x} \end{bmatrix}$$

TL; DM^{math}

$$\begin{bmatrix} \sigma_1 v_1 \vec{x} \\ \sigma_2 v_2 \vec{x} \\ \vdots \\ \sigma_q v_q \vec{x} \end{bmatrix}$$

1. Compute SVD of A
2. Find index of smallest σ_i
3. Take corresponding row of V^T
(col of V)
set that to h .

Robust Model Fitting



RANSAC: RANdom Sample Consensus

For $i = 1$ to K :

$d_i \leftarrow$ random sample of S points

$M_i \leftarrow$ fit model to d_i

$\text{inlier_count} \leftarrow \sum \mathbb{1}(|M(x_i) - y_i| < \delta)$

if $\text{inl_count} > \text{best_count}$

$\text{best_model} \leftarrow M_i$

$\text{best_inliers} \leftarrow \{(x_i, y_i) : |M_i(x_i) - y_i| < \delta\}$

$\text{best_count} \leftarrow \text{inlier_count}$

$M_{\text{final}} \leftarrow$ fit model to best_inliers

k : # iterations

s : minimal set size ($\sqrt{\# \text{DOF}}$)



δ : inlier Threshold

δ : noise model

s : line fitting: 2

translation: 1 pair

affine: 3 pairs



homography: 4 pairs

k : Suppose we want s inliers w/ probability $\geq P$

knowing that $r = \frac{\# \text{ inliers}}{\# \text{ total pts}}$ is the inlier ratio.

$$P(\text{choose all inliers}) = r^s$$

$$P(\text{at least one outlier}) = 1 - r^s$$

\uparrow bad thing

$$P(\text{all at least one outlier after } k \text{ trials}) = (1 - r^s)^k$$

\uparrow bad thing k times

What K do I need for $P(\text{success}) \geq P$

$$P \leq 1 - (1 - r^s)^K$$

$$P = 1 - (1 - r^s)^{\boxed{K}}$$

$$1 - P = (1 - r^s)^K$$

$$\frac{\log(1 - P)}{\log(1 - r^s)} \geq K \cancel{\log(1 - r^s)}$$