

Matrix Factorization

CS 130 - Mathematical Methods in Computer Science

Jan Michael C. Yap

Department of Computer Science
University of the Philippines Diliman

LU(P) Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$$

LU(P) Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$$

\Downarrow

$$L = \begin{pmatrix} l_{11} & 0 & \cdots & 0 \\ l_{21} & l_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ l_{n1} & a_{n2} & \cdots & l_{nn} \end{pmatrix}, U = \begin{pmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ 0 & u_{22} & \cdots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & u_{nn} \end{pmatrix},$$

LU(P) Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$$

\Downarrow

$$L = \begin{pmatrix} l_{11} & 0 & \cdots & 0 \\ l_{21} & l_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ l_{n1} & a_{n2} & \cdots & l_{nn} \end{pmatrix}, U = \begin{pmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ 0 & u_{22} & \cdots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & u_{nn} \end{pmatrix},$$

$$P = \begin{pmatrix} p_{11} & p_{12} & \cdots & p_{1n} \\ p_{21} & p_{22} & \cdots & p_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ p_{n1} & p_{n2} & \cdots & p_{nn} \end{pmatrix}$$

LU(P) Decomposition

- $PA = LU$

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**
 - If A can be reduced to REF (i.e. to U) **without interchanging any rows**, then $P = I$

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**
 - If A can be reduced to REF (i.e. to U) **without interchanging any rows**, then $P = I$
- Mainly used to **solve $Ax = b$**

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**
 - If A can be reduced to REF (i.e. to U) **without interchanging any rows**, then $P = I$
- Mainly used to **solve $Ax = b$**

$$Ax = b \Rightarrow PAx = Pb \Rightarrow LUx = Pb$$

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**
 - If A can be reduced to REF (i.e. to U) **without interchanging any rows**, then $P = I$
- Mainly used to **solve $Ax = b$**

$$Ax = b \Rightarrow PAx = Pb \Rightarrow LUx = Pb$$

- Introduce alternative vector z , then **solve $Lz = Pb$**

LU(P) Decomposition

- $PA = LU$
- Applicable to **square** matrices (mainly)
 - Can be applied to **general** (i.e. $m \times n$) matrices
- P is a **rearrangement of the rows of I**
 - If A can be reduced to REF (i.e. to U) **without interchanging any rows**, then $P = I$
- Mainly used to **solve $Ax = b$**

$$Ax = b \Rightarrow PAx = Pb \Rightarrow LUx = Pb$$

- Introduce alternative vector z , then **solve $Lz = Pb$**
- Once z is solved, **solve $Ux = z$**

QR Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{mn} \end{pmatrix}$$

QR Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{mn} \end{pmatrix}$$

\Downarrow

$$Q = \begin{pmatrix} q_{11} & q_{12} & \cdots & q_{1m} \\ q_{21} & q_{22} & \cdots & q_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ q_{n1} & q_{n2} & \cdots & q_{mm} \end{pmatrix}, R = \begin{pmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{mn} \end{pmatrix},$$

QR Decomposition

- $A_{m \times n} = Q_{m \times m} R_{m \times n}$

QR Decomposition

- $A_{m \times n} = Q_{m \times m} R_{m \times n}$
- $Q^{-1} = Q^T$, i.e. Q is an **orthogonal** matrix

QR Decomposition

- $A_{m \times n} = Q_{m \times m} R_{m \times n}$
- $Q^{-1} = Q^T$, i.e. Q is an **orthogonal** matrix
- Applicable to **general** matrices

QR Decomposition

- $A_{m \times n} = Q_{m \times m} R_{m \times n}$
- $Q^{-1} = Q^T$, i.e. Q is an **orthogonal** matrix
- Applicable to **general** matrices
- Mainly used to **solve** $Ax = b$

QR Decomposition

- $A_{m \times n} = Q_{m \times m} R_{m \times n}$
- $Q^{-1} = Q^T$, i.e. Q is an **orthogonal** matrix
- Applicable to **general** matrices
- Mainly used to **solve** $Ax = b$

$$Ax = b \Rightarrow QRx = b \Rightarrow Rx = Q^{-1}b \Rightarrow Rx = Q^T b$$

Spectral Decomposition

$$P^{-1}AP = D$$

Spectral Decomposition

$$P^{-1}AP = D \Rightarrow A = PDP^{-1}$$

Spectral Decomposition

$$P^{-1}AP = D \Rightarrow A = PDP^{-1}$$

- Also called **eigendecomposition**

Spectral Decomposition

$$P^{-1}AP = D \Rightarrow A = PDP^{-1}$$

- Also called **eigendecomposition**
- Applicable only to **diagonalizable (square)** matrices

Spectral Decomposition

$$P^{-1}AP = D \Rightarrow A = PDP^{-1}$$

- Also called **eigendecomposition**
- Applicable only to **diagonalizable (square)** matrices
- Can be used for **analysis/solution of linear ordinary differential equations**

Singular Value Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

Singular Value Decomposition

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

\Downarrow

$$U = \begin{pmatrix} u_{11} & u_{12} & \cdots & u_{1m} \\ u_{21} & u_{22} & \cdots & u_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ u_{m1} & u_{m2} & \cdots & u_{mm} \end{pmatrix}, V = \begin{pmatrix} v_{11} & v_{12} & \cdots & v_{1n} \\ v_{21} & v_{22} & \cdots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \cdots & v_{nn} \end{pmatrix},$$

$$\Sigma = \begin{pmatrix} \sigma_{11} & 0 & \cdots & 0 & \cdots & 0 \\ 0 & \sigma_{22} & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \sigma_{mm} & \cdots & 0 \end{pmatrix}$$

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$
- The (main) diagonal entries of Σ are composed of the **square roots of the eigenvalues of AA^T (or $A^T A$)**

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$
- The (main) diagonal entries of Σ are composed of the **square roots of the eigenvalues of AA^T (or $A^T A$)**
 - The entries are also called the **singular values of A** .

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$
- The (main) diagonal entries of Σ are composed of the **square roots of the eigenvalues of AA^T (or $A^T A$)**
 - The entries are also called the **singular values of A** .
- The columns of V are composed of the **eigenvectors of $A^T A$** .

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$
- The (main) diagonal entries of Σ are composed of the **square roots of the eigenvalues of AA^T (or $A^T A$)**
 - The entries are also called the **singular values of A** .
- The columns of V are composed of the **eigenvectors of $A^T A$** .
 - $V^T V = I$

Singular Value Decomposition

- $A_{m \times n} = U_{m \times m} \Sigma_{m \times n} V_{n \times n}^T$
- The columns of U are composed of the **eigenvectors of AA^T** .
 - $U^T U = I$
- The (main) diagonal entries of Σ are composed of the **square roots of the eigenvalues of AA^T (or $A^T A$)**
 - The entries are also called the **singular values of A** .
- The columns of V are composed of the **eigenvectors of $A^T A$** .
 - $V^T V = I$
- Applicable to **general** matrices