

# Eigenvectors and Eigenvalues

CS 130 - Mathematical Methods in Computer Science

Jan Michael C. Yap

Department of Computer Science  
University of the Philippines Diliman

- 1 Eigenvectors and Eigenvalues
- 2 Computing for the Eigenvalues and Eigenvectors
- 3 Some Notes
- 4 Diagonalization of a Matrix

# Topics

- 1 Eigenvectors and Eigenvalues
- 2 Computing for the Eigenvalues and Eigenvectors
- 3 Some Notes
- 4 Diagonalization of a Matrix

# What does a matrix do to a vector?

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

# What does a matrix do to a vector?

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

# What does a matrix do to a vector?

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

- A “transforms”  $x$  by rotating and scaling the vector.

# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 18 \\ -18 \end{pmatrix}$$



# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 18 \\ -18 \end{pmatrix} = 9 \begin{pmatrix} 2 \\ -2 \end{pmatrix} = 9x$$

- A **only scales**  $x$ !

# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 18 \\ -18 \end{pmatrix} = 9 \begin{pmatrix} 2 \\ -2 \end{pmatrix} = 9x$$

- A **only scales**  $x$ !
- A has the **same effect as a scalar**! (At least to  $x$ )

# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 18 \\ -18 \end{pmatrix} = 9 \begin{pmatrix} 2 \\ -2 \end{pmatrix} = 9x$$

- A **only scales**  $x$ !
- A has the **same effect as a scalar**! (At least to  $x$ )
- 9 is an **eigenvalue of A** and  $x$  is the **associated eigenvector** for the eigenvalue 9.

# Eigenvectors and Eigenvalues

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, x = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 18 \\ -18 \end{pmatrix} = 9 \begin{pmatrix} 2 \\ -2 \end{pmatrix} = 9x$$

- A **only scales**  $x$ !
- A has the **same effect as a scalar!** (At least to  $x$ )
- 9 is an **eigenvalue of A** and  $x$  is the **associated eigenvector for the eigenvalue 9**.
  - General notation:  $\lambda$  is an **eigenvalue of A** and  $x$  is the **associated eigenvector for the eigenvalue  $\lambda$** .

# Topics

- 1 Eigenvectors and Eigenvalues
- 2 Computing for the Eigenvalues and Eigenvectors
- 3 Some Notes
- 4 Diagonalization of a Matrix

# Some preliminaries

$$Ax = \lambda x$$

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax$$

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = \lambda Ix - Ax$$



## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = \lambda I x - Ax$$

$$\Rightarrow 0 = (\lambda I - A)x$$

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = \lambda I x - Ax$$

$$\Rightarrow 0 = (\lambda I - A)x$$

- We need  $|(\lambda I - A)| = 0$

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = \lambda I x - Ax$$

$$\Rightarrow 0 = (\lambda I - A)x$$

- We need  $|(\lambda I - A)| = 0$ 
  - DAFUQ?!!! Y U NO WANT NONZERO DETERMINANT???

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = \lambda I x - Ax$$

$$\Rightarrow 0 = (\lambda I - A)x$$

- We need  $|(\lambda I - A)| = 0$ 
  - DAFUQ?!!! Y U NO WANT NONZERO DETERMINANT???
- Characteristic polynomial:
$$|(\lambda I - A)| = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \cdots + a_1 \lambda + a_0$$

## Some preliminaries

$$Ax = \lambda x \Rightarrow 0 = \lambda x - Ax \Rightarrow 0 = (\lambda I - A)x$$

$$\Rightarrow 0 = (\lambda I - A)x$$

- We need  $|(\lambda I - A)| = 0$ 
  - DAFUQ?!!! Y U NO WANT NONZERO DETERMINANT???
- Characteristic polynomial:
$$|(\lambda I - A)| = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \cdots + a_1 \lambda + a_0$$
- Characteristic equation:  $|(\lambda I - A)| = 0$

# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$

# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$
- Get the **characteristic polynomial**, i.e.  $|(\lambda I - A)|$  and set up the **characteristic equation**

# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$
- Get the **characteristic polynomial**, i.e.  $|(\lambda I - A)|$  and set up the **characteristic equation**
- Solve for the **real roots of the characteristic equation to get the eigenvalues**.



# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$
- Get the **characteristic polynomial**, i.e.  $|(\lambda I - A)|$  and set up the **characteristic equation**
- Solve for the **real roots of the characteristic equation to get the eigenvalues**.
- For each eigenvalue, **solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$**

# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}$$

# Computing for the Eigenvalues and Eigenvectors

- Form  $\lambda I - A$

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}$$

$\Downarrow$

$$\lambda I - A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} - \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} = \begin{pmatrix} \lambda - 11 & -2 \\ 9 & \lambda \end{pmatrix}$$

# Computing for the Eigenvalues and Eigenvectors

- Get the **characteristic polynomial** and set up the **characteristic equation**

$$\lambda I - A = \begin{pmatrix} \lambda - 11 & -2 \\ 9 & \lambda \end{pmatrix}$$

# Computing for the Eigenvalues and Eigenvectors

- Get the **characteristic polynomial** and set up the **characteristic equation**

$$\lambda I - A = \begin{pmatrix} \lambda - 11 & -2 \\ 9 & \lambda \end{pmatrix} \Rightarrow |\lambda I - A| = \lambda^2 - 11\lambda + 18$$

# Computing for the Eigenvalues and Eigenvectors

- Get the **characteristic polynomial** and set up the **characteristic equation**

$$\lambda I - A = \begin{pmatrix} \lambda - 11 & -2 \\ 9 & \lambda \end{pmatrix} \Rightarrow |\lambda I - A| = \lambda^2 - 11\lambda + 18$$

$\Downarrow$

$$|\lambda I - A| = 0$$

# Computing for the Eigenvalues and Eigenvectors

- Get the **characteristic polynomial** and set up the **characteristic equation**

$$\lambda I - A = \begin{pmatrix} \lambda - 11 & -2 \\ 9 & \lambda \end{pmatrix} \Rightarrow |\lambda I - A| = \lambda^2 - 11\lambda + 18$$

$\Downarrow$

$$|\lambda I - A| = 0 \Rightarrow \lambda^2 - 11\lambda + 18 = 0$$

# Computing for the Eigenvalues and Eigenvectors

- Solve for the **real roots of the characteristic equation to get the eigenvalues.**

$$\lambda^2 - 11\lambda + 18 = 0$$



# Computing for the Eigenvalues and Eigenvectors

- Solve for the **real roots of the characteristic equation to get the eigenvalues.**

$$\lambda^2 - 11\lambda + 18 = 0 \Rightarrow (\lambda - 9)(\lambda - 2) = 0$$

# Computing for the Eigenvalues and Eigenvectors

- Solve for the **real roots of the characteristic equation to get the eigenvalues.**

$$\lambda^2 - 11\lambda + 18 = 0 \Rightarrow (\lambda - 9)(\lambda - 2) = 0$$

$\Downarrow$

$$\lambda = 9, \lambda = 2$$

# Computing for the Eigenvalues and Eigenvectors

- For each eigenvalue, solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$   
 $\lambda = 9$  :

$$(9I - A)x = 0$$

## Computing for the Eigenvalues and Eigenvectors

- For each eigenvalue, solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$   
 $\lambda = 9$  :

$$(9I - A)x = 0 \Rightarrow \left( \begin{pmatrix} 9 & 0 \\ 0 & 9 \end{pmatrix} - \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

# Computing for the Eigenvalues and Eigenvectors

- For each eigenvalue, solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$   
 $\lambda = 9$  :

$$(9I - A)x = 0 \Rightarrow \left( \begin{pmatrix} 9 & 0 \\ 0 & 9 \end{pmatrix} - \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$
$$\Rightarrow \begin{pmatrix} -2 & -2 \\ 9 & 9 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

# Computing for the Eigenvalues and Eigenvectors

- For each eigenvalue, solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$   
 $\lambda = 9$  :

$$(9I - A)x = 0 \Rightarrow \left( \begin{pmatrix} 9 & 0 \\ 0 & 9 \end{pmatrix} - \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \right) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} -2 & -2 \\ 9 & 9 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$\Downarrow$

$$x = \begin{pmatrix} c \\ -c \end{pmatrix}, c \in \mathbb{R}$$

# Computing for the Eigenvalues and Eigenvectors

- For each eigenvalue, solve for the associated eigenvector by solving the homogenous system  $(\lambda I - A)x = 0$

$$\lambda = 2 : x = \begin{pmatrix} -\frac{2c}{9} \\ c \end{pmatrix}, c \in \mathbb{R}$$

# Topics

- 1 Eigenvectors and Eigenvalues
- 2 Computing for the Eigenvalues and Eigenvectors
- 3 Some Notes**
- 4 Diagonalization of a Matrix



# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**
- If there are  **$k$  distinct eigenvalues** for a matrix, then the  $k$  eigenvectors associated with those eigenvalues are **linearly independent**.

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**
- If there are  **$k$  distinct eigenvalues** for a matrix, then the  $k$  eigenvectors associated with those eigenvalues are **linearly independent**.
- If  $A$  is **invertible**, and  $x$  is an eigenvector for  $A$  with associated eigenvalue  $\lambda$ , then  **$x$  is also an eigenvector for  $A^{-1}$  with associated eigenvalue  $\frac{1}{\lambda}$**

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**
- If there are  **$k$  distinct eigenvalues** for a matrix, then the  $k$  eigenvectors associated with those eigenvalues are **linearly independent**.
- If  $A$  is **invertible**, and  $x$  is an eigenvector for  $A$  with associated eigenvalue  $\lambda$ , then  **$x$  is also an eigenvector for  $A^{-1}$  with associated eigenvalue  $\frac{1}{\lambda}$**
- If  $A$  is **noninvertible**, then **an eigenvalue for  $A$  is zero**.

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**
- If there are  **$k$  distinct eigenvalues** for a matrix, then the  $k$  eigenvectors associated with those eigenvalues are **linearly independent**.
- If  $A$  is **invertible**, and  $x$  is an eigenvector for  $A$  with associated eigenvalue  $\lambda$ , then  **$x$  is also an eigenvector for  $A^{-1}$  with associated eigenvalue  $\frac{1}{\lambda}$**
- If  $A$  is **noninvertible**, then **an eigenvalue for  $A$  is zero**.
- The  **$n$  eigenvalues of a square matrix  $A$  of order  $n$  (not all necessarily distinct)** has the following properties:

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of  $A$  corresponding to  $\lambda$**
- If there are  **$k$  distinct eigenvalues** for a matrix, then the  $k$  eigenvectors associated with those eigenvalues are **linearly independent**.
- If  $A$  is **invertible**, and  $x$  is an eigenvector for  $A$  with associated eigenvalue  $\lambda$ , then  **$x$  is also an eigenvector for  $A^{-1}$  with associated eigenvalue  $\frac{1}{\lambda}$**
- If  $A$  is **noninvertible**, then **an eigenvalue for  $A$  is zero**.
- The  **$n$  eigenvalues of a square matrix  $A$  of order  $n$  (not all necessarily distinct)** has the following properties:
  - $\sum_{i=1}^n \lambda_i = \sum_{i=1}^n a_{ii} = \text{tr}(A)$  (the **trace of matrix  $A$** )

# Eigenvalue and Eigenvector Notes

- The set of all eigenvectors associated with an eigenvalue  $\lambda$  is called the **eigenspace of A corresponding to  $\lambda$**
- If there are **k distinct eigenvalues** for a matrix, then the k eigenvectors associated with those eigenvalues are **linearly independent**.
- If A is **invertible**, and  $x$  is an eigenvector for A with associated eigenvalue  $\lambda$ , then  **$x$  is also an eigenvector for  $A^{-1}$  with associated eigenvalue  $\frac{1}{\lambda}$**
- If A is **noninvertible**, then **an eigenvalue for A is zero**.
- The **n eigenvalues of a square matrix A of order n (not all necessarily distinct)** has the following properties:
  - $\sum_{i=1}^n \lambda_i = \sum_{i=1}^n a_{ii} = \text{tr}(A)$  (the **trace of matrix A**)
  - $\prod_{i=1}^n \lambda_i = \det(A)$

# Topics

- 1 Eigenvectors and Eigenvalues
- 2 Computing for the Eigenvalues and Eigenvectors
- 3 Some Notes
- 4 Diagonalization of a Matrix



# Diagonalization

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, P = \begin{pmatrix} -1 & 2 \\ 1 & -9 \end{pmatrix}, P^{-1} = \begin{pmatrix} -\frac{9}{7} & -\frac{2}{7} \\ -\frac{1}{7} & -\frac{1}{7} \end{pmatrix}$$

# Diagonalization

$$A = \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix}, P = \begin{pmatrix} -1 & 2 \\ 1 & -9 \end{pmatrix}, P^{-1} = \begin{pmatrix} -\frac{9}{7} & -\frac{2}{7} \\ -\frac{1}{7} & -\frac{1}{7} \end{pmatrix}$$

$$\Downarrow$$

$$D = P^{-1}AP = \begin{pmatrix} -\frac{9}{7} & -\frac{2}{7} \\ -\frac{1}{7} & -\frac{1}{7} \end{pmatrix} \begin{pmatrix} 11 & 2 \\ -9 & 0 \end{pmatrix} \begin{pmatrix} -1 & 2 \\ 1 & -9 \end{pmatrix} = \begin{pmatrix} 9 & 0 \\ 0 & 2 \end{pmatrix}$$

## Connection to eigenvalues and eigenvectors

- A square matrix  $A$  with order  $n$  is diagonalizable if the  $n$  eigenvalues are real and distinct

## Connection to eigenvalues and eigenvectors

- A square matrix  $A$  with order  $n$  is diagonalizable if the  $n$  eigenvalues are real and distinct
- A square matrix  $A$  with order  $n$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors

## Connection to eigenvalues and eigenvectors

- A square matrix  $A$  with order  $n$  is diagonalizable if the  $n$  eigenvalues are real and distinct
- A square matrix  $A$  with order  $n$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors
  - The columns of  $P$  (the matrix used for diagonalization) are the  $n$  linearly independent eigenvectors

**END OF LESSON 4**