

# Fourier Series

## CS 130 - Mathematical Methods in Computer Science

Jan Michael C. Yap

Department of Computer Science  
University of the Philippines Diliman

March 12, 2013

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Periodic functions

- A function  $f$  is a **periodic function with period  $P$** , if

$$f(x + P) = f(x), \forall x$$

# Periodic functions

- A function  $f$  is a **periodic function with period  $P$** , if

$$f(x + P) = f(x), \forall x$$

- We'll limit ourselves to talking about the functions  **$\sin(kx)$  and  $\cos(kx)$**  for some constant  $k$ .

# Periodic functions

- A function  $f$  is a **periodic function with period  $P$** , if

$$f(x + P) = f(x), \forall x$$

- We'll limit ourselves to talking about the functions  **$\sin(kx)$  and  $\cos(kx)$**  for some constant  $k$ .
  - Quick question: What is the period  $P$  for  $\sin(kx)$  and  $\cos(kx)$ ?

# Periodic functions

- A function  $f$  is a **periodic function with period  $P$** , if

$$f(x + P) = f(x), \forall x$$

- We'll limit ourselves to talking about the functions  **$\sin(kx)$  and  $\cos(kx)$**  for some constant  $k$ .
  - Quick question: What is the period  $P$  for  $\sin(kx)$  and  $\cos(kx)$ ?  **$P = \frac{2\pi}{k}$**



# Topics

## 1 Preliminaries

- Periodic Functions
- **Even and Odd Functions**
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Even and Odd functions

- A function  $f$  is **odd**, if

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is



# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is an **even** function.

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is an **even** function.
- The product of **2 even functions** is

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is an **even** function.
- The product of **2 even functions** is an **even** function.

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is an **even** function.
- The product of **2 even functions** is an **even** function.
- The product of **an odd and an even function** is

# Even and Odd functions

- A function  $f$  is **odd**, if  $f(-x) = -f(x)$ 
  - $\sin(kx)$  is an odd function
- A function  $f$  is **even**, if  $f(-x) = f(x)$ 
  - $\cos(kx)$  is an even function
- The product of **2 odd functions** is an **even** function.
- The product of **2 even functions** is an **even** function.
- The product of **an odd and an even function** is an **odd** function.

# Even and Odd functions

- If  $f$  is **even**, then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

# Even and Odd functions

- If  $f$  is **even**, then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

- If  $f$  is **odd**, then

$$\int_{-a}^a f(x) dx = 0$$

# Even and Odd functions

- If  $f$  is **even**, then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

- If  $f$  is **odd**, then

$$\int_{-a}^a f(x) dx = 0$$

- The integration facts are only guaranteed to hold on a **“symmetric” interval**,  $[-a, a]$



# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Orthogonal functions

- Two **non-zero** functions  $f$  and  $g$  are **orthogonal on the interval**  $[a, b]$ , if

$$\int_a^b f(x)g(x)dx = 0$$

# Orthogonal functions

- Two **non-zero** functions  $f$  and  $g$  are **orthogonal on the interval**  $[a, b]$ , if

$$\int_a^b f(x)g(x)dx = 0$$

- $f(x) = \sin x, g(x) = \cos x, x \in [0, \pi]$

# Orthogonal functions

- Two **non-zero** functions  $f$  and  $g$  are **orthogonal on the interval**  $[a, b]$ , if

$$\int_a^b f(x)g(x)dx = 0$$

- $f(x) = \sin x, g(x) = \cos x, x \in [0, \pi]$

$$\int_0^\pi \sin x \cos x dx$$

# Orthogonal functions

- Two **non-zero** functions  $f$  and  $g$  are **orthogonal on the interval**  $[a, b]$ , if

$$\int_a^b f(x)g(x)dx = 0$$

- $f(x) = \sin x, g(x) = \cos x, x \in [0, \pi]$

$$\int_0^\pi \sin x \cos x dx = \left[ \frac{\sin^2 x}{2} \right]_0^\pi$$

# Orthogonal functions

- Two **non-zero** functions  $f$  and  $g$  are **orthogonal on the interval**  $[a, b]$ , if

$$\int_a^b f(x)g(x)dx = 0$$

- $f(x) = \sin x, g(x) = \cos x, x \in [0, \pi]$

$$\int_0^\pi \sin x \cos x dx = \left[ \frac{\sin^2 x}{2} \right]_0^\pi = 0$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{n\pi x}{L}\right) \mid n = 0, 1, 2, \dots \right\}, x \in [-L, L]$



# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j = k = 0$

$$2 \int_0^L dx = 2L$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j = k \neq 0$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{n\pi x}{L}\right) \mid n = 0, 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = 2 \int_0^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx$$

- $j = k \neq 0$

$$2 \int_0^L \cos^2\left(\frac{k\pi x}{L}\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{n\pi x}{L}\right) \mid n = 0, 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = 2 \int_0^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx$$

- $j = k \neq 0$

$$2 \int_0^L \cos^2\left(\frac{k\pi x}{L}\right) dx = \int_0^L \left(1 + \cos\left(\frac{2k\pi x}{L}\right)\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j = k \neq 0$

$$\int_0^L \left(1 + \cos(\frac{2k\pi x}{L})\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j = k \neq 0$

$$\int_0^L \left(1 + \cos(\frac{2k\pi x}{L})\right) dx = \left[x + \frac{L}{2k\pi} \sin(\frac{2k\pi x}{L})\right]_0^L$$



# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j = k \neq 0$

$$\int_0^L \left(1 + \cos(\frac{2k\pi x}{L})\right) dx = \left[x + \frac{L}{2k\pi} \sin(\frac{2k\pi x}{L})\right]_0^L = L$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j \neq k$

$$\int_0^L \left( \cos(\frac{(j-k)\pi x}{L}) + \cos(\frac{(j+k)\pi x}{L}) \right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j \neq k$

$$= \left[ \frac{L}{(j-k)\pi} \sin(\frac{(j-k)\pi x}{L}) + \frac{L}{(j+k)\pi} \sin(\frac{(j+k)\pi x}{L}) \right]_0^L$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\cos(\frac{n\pi x}{L}) | n = 0, 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx = 2 \int_0^L \cos(\frac{j\pi x}{L}) \cos(\frac{k\pi x}{L}) dx$$

- $j \neq k$

$$= \left[ \frac{L}{(j-k)\pi} \sin(\frac{(j-k)\pi x}{L}) + \frac{L}{(j+k)\pi} \sin(\frac{(j+k)\pi x}{L}) \right]_0^L = 0$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{n\pi x}{L}\right) \mid n = 0, 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{n\pi x}{L}\right) \mid n = 0, 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = \begin{cases} 2L & \text{if } j = k = 0 \\ L & \text{if } j = k \neq 0 \\ 0 & \text{if } j \neq k \end{cases}$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \sin\left(\frac{n\pi x}{L}\right) \mid n = 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \sin\left(\frac{n\pi x}{L}\right) \mid n = 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$



# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x)f_k(x)dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\{\sin(\frac{n\pi x}{L}) | n = 1, 2, \dots\}, x \in [-L, L]$

$$\int_{-L}^L \sin(\frac{j\pi x}{L}) \sin(\frac{k\pi x}{L}) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

- Why did we not begin at  $n = 0$ ?

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{m\pi x}{L}\right) \mid m = 0, 1, 2, \dots \right\} \cup \left\{ \sin\left(\frac{n\pi x}{L}\right) \mid n = 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{m\pi x}{L}\right) \mid m = 0, 1, 2, \dots \right\} \cup \left\{ \sin\left(\frac{n\pi x}{L}\right) \mid n = 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = 0$$

# Mutually orthogonal set of functions

- A **set** of non-zero functions  $F = \{f_n(x)\}$  is **mutually orthogonal on the interval  $[a, b]$** , if for any  $f_j(x), f_k(x) \in F$

$$\int_a^b f_j(x) f_k(x) dx = \begin{cases} 0 & \text{if } j \neq k \\ c > 0 & \text{if } j = k \end{cases}$$

- $\left\{ \cos\left(\frac{m\pi x}{L}\right) \mid m = 0, 1, 2, \dots \right\} \cup \left\{ \sin\left(\frac{n\pi x}{L}\right) \mid n = 1, 2, \dots \right\}, x \in [-L, L]$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = 0$$

- NOTE: each element in the set union is of a **different index**, hence **j can never be equal to k** in the case where we get the integral of the **product of a sine and a cosine function**.

# Summary

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx$$

# Summary

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = \begin{cases} 2L & \text{if } j = k = 0 \\ L & \text{if } j = k \neq 0 \\ 0 & \text{if } j \neq k \end{cases}$$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx$$

# Summary

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = \begin{cases} 2L & \text{if } j = k = 0 \\ L & \text{if } j = k \neq 0 \\ 0 & \text{if } j \neq k \end{cases}$$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx$$

# Summary

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \cos\left(\frac{k\pi x}{L}\right) dx = \begin{cases} 2L & \text{if } j = k = 0 \\ L & \text{if } j = k \neq 0 \\ 0 & \text{if } j \neq k \end{cases}$$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

$$\int_{-L}^L \cos\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = 0$$



# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Main idea of Fourier series

- Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

where  $f$  is differentiable at  $x = a$

# Main idea of Fourier series

- Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

where  $f$  is differentiable at  $x = a$

- On some “symmetric” interval,  $f(x)$  can be (somewhat) periodic.

# Main idea of Fourier series

- Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

where  $f$  is differentiable at  $x = a$

- On some “symmetric” interval,  $f(x)$  can be (somewhat) periodic.
  - Concept of odd and even functions is relevant in this case.

# Main idea of Fourier series

- Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

where  $f$  is differentiable at  $x = a$

- On some “symmetric” interval,  $f(x)$  can be (somewhat) periodic.
  - Concept of odd and even functions is relevant in this case.
- Instead of derivatives, we express  $f(x)$  as a sum of (weighted) sine and cosine functions on that interval.

# Main idea of Fourier series

- Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

where  $f$  is differentiable at  $x = a$

- On some “symmetric” interval,  $f(x)$  can be (somewhat) periodic.
  - Concept of odd and even functions is relevant in this case.
- Instead of derivatives, we express  $f(x)$  as a sum of (weighted) sine and cosine functions on that interval.
  - Choice of sines and cosines, apart from their obvious periodicity, is because the functions form a mutually orthogonal set of functions, which limit the number of variables we need to solve.

# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Fourier sine series

- Fourier sine series of  $f(x)$  on the interval  $[-L, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$



# Fourier sine series

- Fourier sine series of  $f(x)$  on the interval  $[-L, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- Now, how to solve for the coefficients  $B_n$ ?

# Solving for the coefficients of the Fourier sine series

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

# Solving for the coefficients of the Fourier sine series

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\Rightarrow f(x) \sin\left(\frac{m\pi x}{L}\right) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right), m \in \mathbb{Z}^+$$

# Solving for the coefficients of the Fourier sine series

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\Rightarrow f(x) \sin\left(\frac{m\pi x}{L}\right) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right), m \in \mathbb{Z}^+$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \int_{-L}^L \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

# Solving for the coefficients of the Fourier sine series

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\Rightarrow f(x) \sin\left(\frac{m\pi x}{L}\right) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right), m \in \mathbb{Z}^+$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \int_{-L}^L \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_{-L}^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

# Solving for the coefficients of the Fourier sine series

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\Rightarrow f(x) \sin\left(\frac{m\pi x}{L}\right) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right), m \in \mathbb{Z}^+$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \int_{-L}^L \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_{-L}^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m L$$

# Solving for the coefficients of the Fourier sine series

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m L$$

# Solving for the coefficients of the Fourier sine series

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m L$$

$$\Rightarrow \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m$$



# Solving for the coefficients of the Fourier sine series

$$\Rightarrow \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m L$$

$$\Rightarrow \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m$$

$$B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = x, x \in [-L, L]$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = x, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L x \sin\left(\frac{n\pi x}{L}\right) dx$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = x, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L x \sin\left(\frac{n\pi x}{L}\right) dx = \frac{2}{L} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = x, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L x \sin\left(\frac{n\pi x}{L}\right) dx = \frac{2}{L} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{(-1)^{(n+1)} 2L}{n\pi}$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = x, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L x \sin\left(\frac{n\pi x}{L}\right) dx = \frac{2}{L} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{(-1)^{(n+1)} 2L}{n\pi}$$

$$x = \frac{2L}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{(n+1)}}{n} \sin\left(\frac{n\pi x}{L}\right)$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [-L, L]$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx$$



# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx = 0$$

# Examples

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [-L, L]$

$$B_n = \frac{1}{L} \int_{-L}^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx = 0$$

$$1 + x^2 = 0$$

# Fourier sine series on a “half” interval

- Fourier sine series of  $f(x)$  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

# Fourier sine series on a “half” interval

- **Fourier sine series of  $f(x)$**  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

# Fourier sine series on a “half” interval

- **Fourier sine series of  $f(x)$**  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

# Fourier sine series on a “half” interval

- Fourier sine series of  $f(x)$  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\int_{-L}^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} L & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

$$\Rightarrow \int_0^L \sin\left(\frac{j\pi x}{L}\right) \sin\left(\frac{k\pi x}{L}\right) dx = \begin{cases} \frac{L}{2} & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

# Fourier sine series on a “half” interval

- **Fourier sine series of  $f(x)$**  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

# Fourier sine series on a “half” interval

- **Fourier sine series of  $f(x)$**  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m \frac{L}{2}$$



# Fourier sine series on a “half” interval

- Fourier sine series of  $f(x)$  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m \frac{L}{2}$$

$$\Rightarrow \frac{2}{L} \int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m$$

# Fourier sine series on a “half” interval

- **Fourier sine series of  $f(x)$**  on the interval  $[0, L]$

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \sum_{n=1}^{\infty} B_n \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

$$\Rightarrow \int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m \frac{L}{2}$$

$$\Rightarrow \frac{2}{L} \int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = B_m$$

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [0, L]$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [0, L]$

$$B_n = \frac{2}{L} \int_0^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx$$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [0, L]$

$$B_n = \frac{2}{L} \int_0^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{2}{n^3\pi^3} ((2L^2 - n^2\pi^2(1 + L^2))(-1)^n - 2L^2 + n^2\pi^2)$$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = 1 + x^2, x \in [0, L]$

$$B_n = \frac{2}{L} \int_0^L (1 + x^2) \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{2}{n^3 \pi^3} ((2L^2 - n^2 \pi^2 (1 + L^2)) (-1)^n - 2L^2 + n^2 \pi^2)$$

$$1 + x^2 = \sum_{n=1}^{\infty} \frac{2}{n^3 \pi^3} ((2L^2 - n^2 \pi^2 (1 + L^2)) (-1)^n - 2L^2 + n^2 \pi^2) \sin\left(\frac{n\pi x}{L}\right)$$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$

$$B_n = \frac{2}{L} \int_0^{\frac{L}{2}} \frac{L}{2} \sin\left(\frac{n\pi x}{L}\right) dx + \frac{2}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) \sin\left(\frac{n\pi x}{L}\right) dx$$



# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $$f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$$

$$B_n = \frac{2}{L} \int_0^{\frac{L}{2}} \frac{L}{2} \sin\left(\frac{n\pi x}{L}\right) dx + \frac{2}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{L}{n\pi} \left( 1 + (-1)^{(n+1)} - \cos\left(\frac{n\pi}{2}\right) - \frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right) \right)$$

# Example

$$f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

- $$f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$$

$$B_n = \frac{2}{L} \int_0^{\frac{L}{2}} \frac{L}{2} \sin\left(\frac{n\pi x}{L}\right) dx + \frac{2}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) \sin\left(\frac{n\pi x}{L}\right) dx$$

$$B_n = \frac{L}{n\pi} \left( 1 + (-1)^{(n+1)} - \cos\left(\frac{n\pi}{2}\right) - \frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right) \right)$$

$$f(x) = \frac{L}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left( 1 + (-1)^{(n+1)} - \cos\left(\frac{n\pi}{2}\right) - \frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right) \right) \sin\left(\frac{n\pi x}{L}\right)$$

# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- **Fourier Cosine Series**
- Fourier Series

# Fourier cosine series

- Fourier cosine series of  $f(x)$  on the interval  $[-L, L]$

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- On the interval  $[0, L]$ :

$$A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x dx = 0$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x dx = 0$$

$$A_n = \frac{1}{L} \int_{-L}^L x \cos\left(\frac{n\pi x}{L}\right) dx$$



# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x dx = 0$$

$$A_n = \frac{1}{L} \int_{-L}^L x \cos\left(\frac{n\pi x}{L}\right) dx = 0$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x dx = 0$$

$$A_n = \frac{1}{L} \int_{-L}^L x \cos\left(\frac{n\pi x}{L}\right) dx = 0$$

$$x = 0$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

$$A_0 = \frac{1}{L} \int_0^L x dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

$$A_0 = \frac{1}{L} \int_0^L x dx = \frac{L}{2}$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

$$A_0 = \frac{1}{L} \int_0^L x dx = \frac{L}{2}$$

$$A_n = \frac{2}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

$$A_0 = \frac{1}{L} \int_0^L x dx = \frac{L}{2}$$

$$A_n = \frac{2}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) dx = \frac{2L}{n^2 \pi^2} (1 - (-1)^n)$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x, x \in [0, L]$

$$A_0 = \frac{1}{L} \int_0^L x dx = \frac{L}{2}$$

$$A_n = \frac{2}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) dx = \frac{2L}{n^2\pi^2} (1 - (-1)^n)$$

$$x = \frac{L}{2} + \frac{2L}{n^2\pi^2} \sum_{n=1}^{\infty} (1 - (-1)^n) \cos\left(\frac{n\pi x}{L}\right)$$



# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx = \frac{1}{L} \int_0^L x^2 dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx = \frac{1}{L} \int_0^L x^2 dx = \frac{L^2}{3}$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx = \frac{1}{L} \int_0^L x^2 dx = \frac{L^2}{3}$$

$$A_n = \frac{2}{L} \int_0^L x^2 \cos\left(\frac{n\pi x}{L}\right) dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx = \frac{1}{L} \int_0^L x^2 dx = \frac{L^2}{3}$$

$$A_n = \frac{2}{L} \int_0^L x^2 \cos\left(\frac{n\pi x}{L}\right) dx = \frac{4L^2(-1)^n}{n^2\pi^2}$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{2L} \int_{-L}^L f(x) dx & n = 0 \\ \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{1}{2L} \int_{-L}^L x^2 dx = \frac{1}{L} \int_0^L x^2 dx = \frac{L^2}{3}$$

$$A_n = \frac{2}{L} \int_0^L x^2 \cos\left(\frac{n\pi x}{L}\right) dx = \frac{4L^2(-1)^n}{n^2\pi^2}$$

$$x^2 = \frac{L^2}{3} + \sum_{n=1}^{\infty} \frac{4L^2(-1)^n}{n^2\pi^2} \cos\left(\frac{n\pi x}{L}\right)$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$



# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$

$$A_0 = \frac{1}{L} \int_0^{\frac{L}{2}} \frac{L}{2} dx + \frac{1}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) dx$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$

$$A_0 = \frac{1}{L} \int_0^{\frac{L}{2}} \frac{L}{2} dx + \frac{1}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) dx$$

$$A_0 = \frac{3L}{8}$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

- $f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$

$$A_n = \frac{2}{L} \int_0^{\frac{L}{2}} \frac{L}{2} \cos\left(\frac{n\pi x}{L}\right) dx + \frac{2}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) \cos\left(\frac{n\pi x}{L}\right) dx$$

$$A_n = \frac{2L}{n^2\pi^2} \left( (-1)^n - \cos\left(\frac{n\pi}{2}\right) + \frac{n\pi}{2} \sin\left(\frac{n\pi}{2}\right) \right)$$

# Examples

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), A_n = \begin{cases} \frac{1}{L} \int_0^L f(x) dx & n = 0 \\ \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx & n > 0 \end{cases}$$

$$\bullet f(x) = \begin{cases} \frac{L}{2} & x \in [0, \frac{L}{2}] \\ x - \frac{L}{2} & x \in [\frac{L}{2}, L] \end{cases}$$

$$A_n = \frac{2}{L} \int_0^{\frac{L}{2}} \frac{L}{2} \cos\left(\frac{n\pi x}{L}\right) dx + \frac{2}{L} \int_{\frac{L}{2}}^L \left(x - \frac{L}{2}\right) \cos\left(\frac{n\pi x}{L}\right) dx$$

$$A_n = \frac{2L}{n^2\pi^2} \left( (-1)^n - \cos\left(\frac{n\pi}{2}\right) + \frac{n\pi}{2} \sin\left(\frac{n\pi}{2}\right) \right)$$

$$f(x) = \frac{3L}{8} + \frac{2L}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \left( (-1)^n - \cos\left(\frac{n\pi}{2}\right) + \frac{n\pi}{2} \sin\left(\frac{n\pi}{2}\right) \right) \cos\left(\frac{n\pi x}{L}\right)$$

# Topics

## 1 Preliminaries

- Periodic Functions
- Even and Odd Functions
- Orthogonal Functions

## 2 Fourier Series

- Fourier Sine Series
- Fourier Cosine Series
- Fourier Series

# Fourier series

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x, x \in [-L, L]$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = 0$$

$$A_n = 0$$



# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = 0$$

$$A_n = 0$$

$$B_n = \frac{(-1)^{(n+1)} 2L}{n\pi}$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x, x \in [-L, L]$

$$A_0 = 0$$

$$A_n = 0$$

$$B_n = \frac{(-1)^{(n+1)} 2L}{n\pi}$$

$$x = \frac{2L}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{(n+1)}}{n} \sin\left(\frac{n\pi x}{L}\right)$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x^2, x \in [-L, L]$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{L^2}{3}$$

$$A_n = \frac{4L^2(-1)^n}{n^2\pi^2}$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{L^2}{3}$$

$$A_n = \frac{4L^2(-1)^n}{n^2\pi^2}$$

$$B_n = 0$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = x^2, x \in [-L, L]$

$$A_0 = \frac{L^2}{3}$$

$$A_n = \frac{4L^2(-1)^n}{n^2\pi^2}$$

$$B_n = 0$$

$$x^2 = \frac{L^2}{3} + \sum_{n=1}^{\infty} \frac{4L^2(-1)^n}{n^2\pi^2} \cos\left(\frac{n\pi x}{L}\right)$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = \begin{cases} L & x \in [-L, 0] \\ 2x & x \in [0, L] \end{cases}$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = \begin{cases} L & x \in [-L, 0] \\ 2x & x \in [0, L] \end{cases}$

$$A_0 = L$$

$$A_n = \frac{2L^2}{n^2\pi^2} (-1 + (-1)^n)$$



# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = \begin{cases} L & x \in [-L, 0] \\ 2x & x \in [0, L] \end{cases}$

$$A_0 = L$$

$$A_n = \frac{2L^2}{n^2\pi^2} (-1 + (-1)^n)$$

$$B_n = -\frac{L}{n\pi} (1 + (-1)^n)$$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = \begin{cases} L & x \in [-L, 0] \\ 2x & x \in [0, L] \end{cases}$

# Examples

- **Fourier series of  $f(x)$**  on the interval  $[-L, L]$  (or  $[0, L]$ )

$$f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

- $f(x) = \begin{cases} L & x \in [-L, 0] \\ 2x & x \in [0, L] \end{cases}$

$$x^2 = L + \frac{2L^2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} (-1 + (-1)^n) \cos\left(\frac{n\pi x}{L}\right)$$

$$- \frac{L}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} (1 + (-1)^n) \sin\left(\frac{n\pi x}{L}\right)$$

**END OF LESSON 11**