

Laplace Transforms

CS 130 - Mathematical Methods in Computer Science

Jan Michael C. Yap

Department of Computer Science
University of the Philippines Diliman

February 28, 2013

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Topics

1 Laplace Transforms

- Laplace Transforms and Solving ODEs
- Laplace Transform
- Useful Rules and Theorems

2 Inverse Laplace Transforms

3 Solving First and Second Order ODEs

- Examples
- General Solutions of ODEs and Laplace Transforms

4 Solving Systems of ODEs

Topics

1 Laplace Transforms

- Laplace Transforms and Solving ODEs
- Laplace Transform
- Useful Rules and Theorems

2 Inverse Laplace Transforms

3 Solving First and Second Order ODEs

- Examples
- General Solutions of ODEs and Laplace Transforms

4 Solving Systems of ODEs

Laplace Transforms and Solving ODEs

(Systems of) ODEs \Leftrightarrow (Inverse) Laplace Transform \Leftrightarrow Algebraic System

Laplace Transforms and Solving ODEs

(Systems of) ODEs \Leftrightarrow (Inverse) Laplace Transform \Leftrightarrow Algebraic System

- (System of) Linear ODEs with constant coefficients

Laplace Transforms and Solving ODEs

(Systems of) ODEs \Leftrightarrow (Inverse) Laplace Transform \Leftrightarrow Algebraic System

- (System of) Linear ODEs with constant coefficients
- Initial conditions are usually needed to solve the ODEs

Laplace Transforms and Solving ODEs

(Systems of) ODEs \Leftrightarrow (Inverse) Laplace Transform \Leftrightarrow Algebraic System

- (System of) Linear ODEs with constant coefficients
- Initial conditions are usually needed to solve the ODEs
 - Implicitly, we are looking for a particular solution when solving ODEs using Laplace Transforms

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Laplace Transform

- Laplace transform of a function $f(t)$, $L[f(t)]$ defined for $t > 0$:

$$\int_0^{\infty} e^{-st} f(t) dt$$

Laplace Transform

- Laplace transform of a function $f(t)$, $L[f(t)]$ defined for $t > 0$:

$$\int_0^{\infty} e^{-st} f(t) dt$$

- s is an arbitrary positive number.

Laplace Transform

- Laplace transform of a function $f(t)$, $L[f(t)]$ defined for $t > 0$:

$$\int_0^{\infty} e^{-st} f(t) dt$$

- s is an arbitrary positive number.
- Alternative notation for the Laplace transform of $f(t)$ is $F(s)$ or $X(s)$

Laplace Transforms of Some Functions

- $f(t) = k$

Laplace Transforms of Some Functions

- $f(t) = k$

$$\int_0^{\infty} k e^{-st} dt$$

Laplace Transforms of Some Functions

- $f(t) = k$

$$\int_0^{\infty} k e^{-st} dt = k \left[\frac{e^{-st}}{-s} \right]_0^{\infty}$$

Laplace Transforms of Some Functions

- $f(t) = k$

$$\int_0^{\infty} k e^{-st} dt = k \left[\frac{e^{-st}}{-s} \right]_0^{\infty} = k \left(-\frac{0}{s} - \left(-\frac{1}{s} \right) \right)$$

Laplace Transforms of Some Functions

- $f(t) = k$

$$\int_0^{\infty} k e^{-st} dt = k \left[\frac{e^{-st}}{-s} \right]_0^{\infty} = k \left(-\frac{0}{s} - \left(-\frac{1}{s} \right) \right)$$

$$L[k] = \frac{k}{s}$$

Laplace Transforms of Some Functions

- $f(t) = e^{-at}$

Laplace Transforms of Some Functions

- $f(t) = e^{-at}$

$$\int_0^{\infty} e^{-at} e^{-st} dt$$

Laplace Transforms of Some Functions

- $f(t) = e^{-at}$

$$\int_0^{\infty} e^{-at} e^{-st} dt = \int_0^{\infty} e^{-(s+a)t} dt$$

Laplace Transforms of Some Functions

- $f(t) = e^{-at}$

$$\int_0^{\infty} e^{-at} e^{-st} dt = \int_0^{\infty} e^{-(s+a)t} dt = \left[\frac{e^{-(s+a)t}}{-(s+a)} \right]_0^{\infty}$$

Laplace Transforms of Some Functions

- $f(t) = e^{-at}$

$$\int_0^{\infty} e^{-at} e^{-st} dt = \int_0^{\infty} e^{-(s+a)t} dt = \left[\frac{e^{-(s+a)t}}{-(s+a)} \right]_0^{\infty}$$

$$L[e^{-at}] = \frac{1}{s+a}$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\int_0^{\infty} e^{-st} \cos(at) dt$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\int_0^{\infty} e^{-st} \cos(at) dt = \left[\frac{e^{-st} \sin(at)}{a} \right]_0^{\infty} + \frac{s}{a} \int_0^{\infty} e^{-st} \sin(at) dt$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\begin{aligned}\int_0^{\infty} e^{-st} \cos(at) dt &= \left[\frac{e^{-st} \sin(at)}{a} \right]_0^{\infty} + \frac{s}{a} \int_0^{\infty} e^{-st} \sin(at) dt \\ &= \frac{s}{a} \left(\left[\frac{-e^{-st} \cos(at)}{a} \right]_0^{\infty} - \frac{s}{a} \int_0^{\infty} e^{-st} \cos(at) dt \right)\end{aligned}$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\int_0^{\infty} e^{-st} \cos(at) dt = \left[\frac{e^{-st} \sin(at)}{a} \right]_0^{\infty} + \frac{s}{a} \int_0^{\infty} e^{-st} \sin(at) dt$$

$$= \frac{s}{a} \left(\left[\frac{-e^{-st} \cos(at)}{a} \right]_0^{\infty} - \frac{s}{a} \int_0^{\infty} e^{-st} \cos(at) dt \right)$$

$$\Rightarrow L[\cos(at)] = \frac{s}{a^2} - \frac{s^2}{a^2} L[\cos(at)]$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\int_0^{\infty} e^{-st} \cos(at) dt = \left[\frac{e^{-st} \sin(at)}{a} \right]_0^{\infty} + \frac{s}{a} \int_0^{\infty} e^{-st} \sin(at) dt$$

$$= \frac{s}{a} \left(\left[\frac{-e^{-st} \cos(at)}{a} \right]_0^{\infty} - \frac{s}{a} \int_0^{\infty} e^{-st} \cos(at) dt \right)$$

$$\Rightarrow L[\cos(at)] = \frac{s}{a^2} - \frac{s^2}{a^2} L[\cos(at)] \Rightarrow \left(1 + \frac{s^2}{a^2} \right) L[\cos(at)] = \frac{s}{a^2}$$

Laplace Transforms of Some Functions

- $f(t) = \cos(at)$

$$\int_0^{\infty} e^{-st} \cos(at) dt = \left[\frac{e^{-st} \sin(at)}{a} \right]_0^{\infty} + \frac{s}{a} \int_0^{\infty} e^{-st} \sin(at) dt$$

$$= \frac{s}{a} \left(\left[\frac{-e^{-st} \cos(at)}{a} \right]_0^{\infty} - \frac{s}{a} \int_0^{\infty} e^{-st} \cos(at) dt \right)$$

$$\Rightarrow L[\cos(at)] = \frac{s}{a^2} - \frac{s^2}{a^2} L[\cos(at)] \Rightarrow \left(1 + \frac{s^2}{a^2} \right) L[\cos(at)] = \frac{s}{a^2}$$

$$L[\cos(at)] = \frac{s}{s^2 + a^2}$$

Laplace Transforms of Some Functions

- $L[\sin(at)] = \frac{a}{s^2+a^2}$
- $L[t^n] = \frac{n!}{s^{n+1}}$
- $L[te^{-at}] = \frac{1}{(s+a)^2}$
- $L[t \cos(at)] = \frac{(s^2-a^2)}{(s^2+a^2)^2}$
- $L[t \sin(at)] = \frac{2as}{(s^2+a^2)^2}$
- $L[\sin(at) - at \cos(at)] = \frac{2a^3}{(s^2+a^2)^2}$
- $L[\cosh(at)] = \frac{s}{s^2-a^2}$
- $L[\sinh(at)] = \frac{a}{s^2-a^2}$

Topics

1 Laplace Transforms

- Laplace Transforms and Solving ODEs
- Laplace Transform
- Useful Rules and Theorems

2 Inverse Laplace Transforms

3 Solving First and Second Order ODEs

- Examples
- General Solutions of ODEs and Laplace Transforms

4 Solving Systems of ODEs

Laplace Transform Rules and Theorems

- **Linearity:** $L[Af(t) + Bg(t)] = AL[f(t)] + BL[g(t)]$, A and B are constants

Laplace Transform Rules and Theorems

- **Linearity:** $L[Af(t) + Bg(t)] = AL[f(t)] + BL[g(t)]$, A and B are constants

$$L[2t^5 + 7 \cos(4t) - 1]$$

Laplace Transform Rules and Theorems

- **Linearity:** $L[Af(t) + Bg(t)] = AL[f(t)] + BL[g(t)]$, A and B are constants

$$L[2t^5 + 7 \cos(4t) - 1] = 2L[t^5] + 7L[\cos(4t)] - L[1]$$

Laplace Transform Rules and Theorems

- **Linearity:** $L[Af(t) + Bg(t)] = AL[f(t)] + BL[g(t)]$, A and B are constants

$$L[2t^5 + 7 \cos(4t) - 1] = 2L[t^5] + 7L[\cos(4t)] - L[1]$$

$$= 2 \left(\frac{5!}{s^6} \right) + 7 \frac{s}{s^2 + 16} - \frac{1}{s}$$

Laplace Transform Rules and Theorems

- **First Shifting Theorem:** $L[e^{-at}f(t)] = F(s + a)$

Laplace Transform Rules and Theorems

- **First Shifting Theorem:** $L[e^{-at}f(t)] = F(s + a)$

$$L[e^{-2t} \sin(3t)]$$

Laplace Transform Rules and Theorems

- **First Shifting Theorem:** $L[e^{-at}f(t)] = F(s + a)$

$$L[e^{-2t} \sin(3t)] \Rightarrow F(s) = L[\sin(3t)] = \frac{3}{s^2 + 9}$$

Laplace Transform Rules and Theorems

- **First Shifting Theorem:** $L[e^{-at}f(t)] = F(s + a)$

$$L[e^{-2t} \sin(3t)] \Rightarrow F(s) = L[\sin(3t)] = \frac{3}{s^2 + 9}$$

$$\Rightarrow F(s + a) = \frac{3}{(s + 2)^2 + 9} = L[e^{-2t} \sin(3t)]$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t}$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0 \Rightarrow f'(t) = 3t^2 e^{-t} - t^3 e^{-t}$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0 \Rightarrow f'(t) = 3t^2 e^{-t} - t^3 e^{-t}$$

$$L[f'(t)] = 3\left(\frac{2}{(s+1)^3}\right) - \frac{6}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0 \Rightarrow f'(t) = 3t^2 e^{-t} - t^3 e^{-t}$$

$$L[f'(t)] = 3 \left(\frac{2}{(s+1)^3} \right) - \frac{6}{(s+1)^4} = \frac{6s}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0 \Rightarrow f'(t) = 3t^2 e^{-t} - t^3 e^{-t}$$

$$L[f'(t)] = 3 \left(\frac{2}{(s+1)^3} \right) - \frac{6}{(s+1)^4} = \frac{6s}{(s+1)^4}$$

$$sL[f(t)] - f(0) = s \left(\frac{6}{(s+1)^4} \right) - 0$$

Laplace Transform Rules and Theorems

- Laplace transform of a first derivative:

$$L[f'(t)] = sL[f(t)] - f(0) \text{ OR } L\left[\frac{dx}{dt}\right] = sX(s) - x(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = 0 \Rightarrow f'(t) = 3t^2 e^{-t} - t^3 e^{-t}$$

$$L[f'(t)] = 3 \left(\frac{2}{(s+1)^3} \right) - \frac{6}{(s+1)^4} = \frac{6s}{(s+1)^4}$$

$$sL[f(t)] - f(0) = s \left(\frac{6}{(s+1)^4} \right) - 0 = \frac{6s}{(s+1)^4} = L[f'(t)]$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = f'(0) = 0$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = f'(0) = 0$$

$$f'(t) = 3t^2 e^{-t} - t^3 e^{-t} \Rightarrow f''(t) = 6te^{-t} - 6t^2 e^{-t} + t^3 e^{-t}$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = f'(0) = 0$$

$$f'(t) = 3t^2 e^{-t} - t^3 e^{-t} \Rightarrow f''(t) = 6te^{-t} - 6t^2 e^{-t} + t^3 e^{-t}$$

$$L[f''(t)] = \frac{6s^2}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = f'(0) = 0$$

$$f'(t) = 3t^2 e^{-t} - t^3 e^{-t} \Rightarrow f''(t) = 6te^{-t} - 6t^2 e^{-t} + t^3 e^{-t}$$

$$L[f''(t)] = \frac{6s^2}{(s+1)^4}$$

$$s^2 L[f(t)] - sf(0) - f'(0) = s^2 \left(\frac{6}{(s+1)^4} \right) - s(0) - 0$$

Laplace Transform Rules and Theorems

- Laplace transform of a second derivative:

$$L[f''(t)] = s^2 L[f(t)] - sf(0) - f'(0) \text{ OR}$$

$$L\left[\frac{d^2x}{dt^2}\right] = s^2 X(s) - sx(0) - x'(0)$$

$$f(t) = t^3 e^{-t} \Rightarrow f(0) = f'(0) = 0$$

$$f'(t) = 3t^2 e^{-t} - t^3 e^{-t} \Rightarrow f''(t) = 6te^{-t} - 6t^2 e^{-t} + t^3 e^{-t}$$

$$L[f''(t)] = \frac{6s^2}{(s+1)^4}$$

$$s^2 L[f(t)] - sf(0) - f'(0) = s^2 \left(\frac{6}{(s+1)^4} \right) - s(0) - 0 = \frac{6s^2}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)]$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)] = \frac{s^2 - 49}{(s^2 + 49)^2}$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)] = \frac{s^2 - 49}{(s^2 + 49)^2}$$

$$\Rightarrow L[\cos(7t)] = \frac{s}{s^2 + 49}$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)] = \frac{s^2 - 49}{(s^2 + 49)^2}$$

$$\Rightarrow L[\cos(7t)] = \frac{s}{s^2 + 49}$$

$$-\frac{dL[\cos(7t)]}{ds} = -\left(\frac{1}{s^2 + 49} - \frac{2s^2}{(s^2 + 49)^2}\right)$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)] = \frac{s^2 - 49}{(s^2 + 49)^2}$$

$$\Rightarrow L[\cos(7t)] = \frac{s}{s^2 + 49}$$

$$-\frac{dL[\cos(7t)]}{ds} = -\left(\frac{1}{s^2 + 49} - \frac{2s^2}{(s^2 + 49)^2}\right) = -\left(\frac{49 - s^2}{(s^2 + 49)^2}\right)$$

Laplace Transform Rules and Theorems

- **Multiplication by t:** $L[tf(t)] = -\frac{dL[f(t)]}{ds}$

$$L[t \cos(7t)] = \frac{s^2 - 49}{(s^2 + 49)^2}$$

$$\Rightarrow L[\cos(7t)] = \frac{s}{s^2 + 49}$$

$$\begin{aligned} -\frac{dL[\cos(7t)]}{ds} &= -\left(\frac{1}{s^2 + 49} - \frac{2s^2}{(s^2 + 49)^2}\right) = -\left(\frac{49 - s^2}{(s^2 + 49)^2}\right) \\ &= \frac{s^2 - 49}{(s^2 + 49)^2} = L[t \cos(7t)] \end{aligned}$$

Laplace Transform Rules and Theorems

- **Initial Value Theorem:** $\lim_{t \rightarrow 0} f(t) = \lim_{s \rightarrow \infty} sL[f(t)]$

Laplace Transform Rules and Theorems

- **Initial Value Theorem:** $\lim_{t \rightarrow 0} f(t) = \lim_{s \rightarrow \infty} sL[f(t)]$

$$f(t) = te^{-t} \Rightarrow L[f(t)] = \frac{6}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- **Initial Value Theorem:** $\lim_{t \rightarrow 0} f(t) = \lim_{s \rightarrow \infty} sL[f(t)]$

$$f(t) = te^{-t} \Rightarrow L[f(t)] = \frac{6}{(s+1)^4}$$

$$\lim_{t \rightarrow 0} t^3 e^{-t} = 0$$

Laplace Transform Rules and Theorems

- **Initial Value Theorem:** $\lim_{t \rightarrow 0} f(t) = \lim_{s \rightarrow \infty} sL[f(t)]$

$$f(t) = te^{-t} \Rightarrow L[f(t)] = \frac{6}{(s+1)^4}$$

$$\lim_{t \rightarrow 0} t^3 e^{-t} = 0$$

$$\lim_{s \rightarrow \infty} \frac{6s}{(s+1)^4} = 0$$

Laplace Transform Rules and Theorems

- **Final Value Theorem:** $\lim_{t \rightarrow \infty} f(t) = \lim_{s \rightarrow 0} sL[f(t)]$

Laplace Transform Rules and Theorems

- **Final Value Theorem:** $\lim_{t \rightarrow \infty} f(t) = \lim_{s \rightarrow 0} sL[f(t)]$

$$f(t) = te^{-t} \Rightarrow L[f(t)] = \frac{6}{(s+1)^4}$$

Laplace Transform Rules and Theorems

- **Final Value Theorem:** $\lim_{t \rightarrow \infty} f(t) = \lim_{s \rightarrow 0} sL[f(t)]$

$$f(t) = te^{-t} \Rightarrow L[f(t)] = \frac{6}{(s+1)^4}$$

$$\lim_{t \rightarrow \infty} t^3 e^{-t} = 0$$

$$\lim_{s \rightarrow 0} \frac{6s}{(s+1)^4} = 0$$

Some Examples/Exercises

- $L[2 \sin(3t) - 3 \cos(2t)]$

Some Examples/Exercises

- $L[2 \sin(3t) - 3 \cos(2t)]$
- $L[(t + 1)^3]$

Some Examples/Exercises

- $L[2 \sin(3t) - 3 \cos(2t)]$
- $L[(t + 1)^3]$
- $L[t(e^t + e^{-2t})]$

Some Examples/Exercises

- $L[2 \sin(3t) - 3 \cos(2t)]$
- $L[(t + 1)^3]$
- $L[t(e^t + e^{-2t})]$
- Given $\frac{dx}{dt} + x = e^t$, and $x(0) = 0$, what is $L[x(t)]$?

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Inverse Laplace Transform

- Given $F(s)$, we can obtain the **inverse Laplace transform** of $F(s)$ denoted as $L^{-1}[F(s)]$

Inverse Laplace Transform

- Given $F(s)$, we can obtain the **inverse Laplace transform** of $F(s)$ denoted as $L^{-1}[F(s)]$
 - It is implicitly assumed (at least for our examples later) that $L^{-1}[F(s)]$ hold for $t > 0$.

Inverse Laplace Transform

- Given $F(s)$, we can obtain the **inverse Laplace transform** of $F(s)$ denoted as $L^{-1}[F(s)]$
 - It is implicitly assumed (at least for our examples later) that $L^{-1}[F(s)]$ hold for $t > 0$.
- Main idea: **Convert $F(s)$ into something familiar looking and just “back substitute” into the original function prior to applying Laplace transform**

Inverse Laplace Transform

- Given $F(s)$, we can obtain the **inverse Laplace transform** of $F(s)$ denoted as $L^{-1}[F(s)]$
 - It is implicitly assumed (at least for our examples later) that $L^{-1}[F(s)]$ hold for $t > 0$.
- Main idea: **Convert $F(s)$ into something familiar looking and just “back substitute” into the original function prior to applying Laplace transform**
- Useful tip: Inverse Laplace transforms are also **linear**, i.e.

$$L^{-1}[AF(s) + BG(s)] = AL^{-1}[F(s)] + BL^{-1}[G(s)]$$

Inverse Laplace Transform

- Given $F(s)$, we can obtain the **inverse Laplace transform** of $F(s)$ denoted as $L^{-1}[F(s)]$
 - It is implicitly assumed (at least for our examples later) that $L^{-1}[F(s)]$ hold for $t > 0$.
- Main idea: **Convert $F(s)$ into something familiar looking and just “back substitute” into the original function prior to applying Laplace transform**
- Useful tip: Inverse Laplace transforms are also **linear**, i.e.

$$L^{-1}[AF(s) + BG(s)] = AL^{-1}[F(s)] + BL^{-1}[G(s)]$$

- (Another) Useful tip: **Partial fraction decomposition** can help make some functions be more familiar

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

$$L^{-1} \left[\frac{3}{s^3} + \frac{4}{s-2} \right]$$

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

$$L^{-1} \left[\frac{3}{s^3} + \frac{4}{s-2} \right] \Rightarrow 3L^{-1} \left[\frac{1}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

$$L^{-1} \left[\frac{3}{s^3} + \frac{4}{s-2} \right] \Rightarrow 3L^{-1} \left[\frac{1}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

$$\Rightarrow 3L^{-1} \left[\frac{1}{2} \frac{2}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

$$L^{-1} \left[\frac{3}{s^3} + \frac{4}{s-2} \right] \Rightarrow 3L^{-1} \left[\frac{1}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

$$\Rightarrow 3L^{-1} \left[\frac{1}{2} \frac{2}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right] \Rightarrow \frac{3}{2}L^{-1} \left[\frac{2}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

Examples

- $F(s) = \frac{3}{s^3} + \frac{4}{s-2}$

$$L^{-1} \left[\frac{3}{s^3} + \frac{4}{s-2} \right] \Rightarrow 3L^{-1} \left[\frac{1}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

$$\Rightarrow 3L^{-1} \left[\frac{1}{2} \frac{2}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right] \Rightarrow \frac{3}{2}L^{-1} \left[\frac{2}{s^3} \right] + 4L^{-1} \left[\frac{1}{s-2} \right]$$

$$L^{-1}[F(s)] = f(t) = \frac{3}{2}t^2 + 4e^{2t}$$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

$$\frac{2s+3}{s^2+3s} \Rightarrow \frac{2s+3}{s(s+3)}$$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

$$\frac{2s+3}{s^2+3s} \Rightarrow \frac{2s+3}{s(s+3)} \Rightarrow \frac{A}{s} + \frac{B}{s+3}$$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

$$\frac{2s+3}{s^2+3s} \Rightarrow \frac{2s+3}{s(s+3)} \Rightarrow \frac{A}{s} + \frac{B}{s+3}$$

$$\Rightarrow \frac{1}{s} + \frac{1}{s+3}$$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

$$\frac{2s+3}{s^2+3s} \Rightarrow \frac{2s+3}{s(s+3)} \Rightarrow \frac{A}{s} + \frac{B}{s+3}$$

$$\Rightarrow \frac{1}{s} + \frac{1}{s+3} \Rightarrow L^{-1} \left[\frac{1}{s} \right] + L^{-1} \left[\frac{1}{s+3} \right]$$

Examples

- $F(s) = \frac{2s+3}{s^2+3s}$

$$\frac{2s+3}{s^2+3s} \Rightarrow \frac{2s+3}{s(s+3)} \Rightarrow \frac{A}{s} + \frac{B}{s+3}$$

$$\Rightarrow \frac{1}{s} + \frac{1}{s+3} \Rightarrow L^{-1} \left[\frac{1}{s} \right] + L^{-1} \left[\frac{1}{s+3} \right]$$

$$L^{-1}[F(s)] = f(t) = 1 + e^{-3t}$$

Examples

- $F(s) = \frac{1}{s^2+9}$

Examples

- $F(s) = \frac{1}{s^2+9}$

$$\frac{1}{s^2 + 9}$$

Examples

- $F(s) = \frac{1}{s^2+9}$

$$\frac{1}{s^2+9} \Rightarrow \frac{1}{3} \frac{3}{s^2+9}$$

Examples

- $F(s) = \frac{1}{s^2+9}$

$$\frac{1}{s^2+9} \Rightarrow \frac{1}{3} \frac{3}{s^2+9} \Rightarrow \frac{1}{3} L^{-1} \left[\frac{3}{s^2+9} \right]$$

Examples

- $F(s) = \frac{1}{s^2+9}$

$$\frac{1}{s^2+9} \Rightarrow \frac{1}{3} \frac{3}{s^2+9} \Rightarrow \frac{1}{3} L^{-1} \left[\frac{3}{s^2+9} \right]$$

$$L^{-1}[F(s)] = f(t) = \frac{1}{3} \sin(3t)$$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

$$\frac{1}{(s+2)^5} = G(s+2)$$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

$$\frac{1}{(s+2)^5} = G(s+2) \Rightarrow G(s) = \frac{1}{s^5} = \frac{1}{24} \frac{24}{s^5}$$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

$$\frac{1}{(s+2)^5} = G(s+2) \Rightarrow G(s) = \frac{1}{s^5} = \frac{1}{24} \frac{24}{s^5} \Rightarrow \frac{1}{24} L^{-1} \left[\frac{24}{s^5} \right] = \frac{1}{24} t^4$$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

$$\frac{1}{(s+2)^5} = G(s+2) \Rightarrow G(s) = \frac{1}{s^5} = \frac{1}{24} \frac{24}{s^5} \Rightarrow \frac{1}{24} L^{-1} \left[\frac{24}{s^5} \right] = \frac{1}{24} t^4$$

By the first shifting theorem, $L^{-1}[G(s+2)] = L^{-1}[G(s)]e^{-2t}$

Examples

- $F(s) = \frac{1}{(s+2)^5}$

$$\frac{1}{(s+2)^5} = G(s+2) \Rightarrow G(s) = \frac{1}{s^5} = \frac{1}{24} \frac{24}{s^5} \Rightarrow \frac{1}{24} L^{-1} \left[\frac{24}{s^5} \right] = \frac{1}{24} t^4$$

By the first shifting theorem, $L^{-1}[G(s+2)] = L^{-1}[G(s)]e^{-2t}$

$$L^{-1}[F(s)] = f(t) = \frac{1}{24} t^4 e^{-2t}$$

Examples

- $F(s) = \frac{s}{s^2+4s+13}$

Examples

- $F(s) = \frac{s}{s^2+4s+13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9}$$

Examples

- $F(s) = \frac{s}{s^2+4s+13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9} \Rightarrow \frac{s}{(s + 2)^2 + 9}$$

Examples

- $F(s) = \frac{s}{s^2 + 4s + 13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9} \Rightarrow \frac{s}{(s + 2)^2 + 9}$$

$$\Rightarrow \frac{(s + 2) - 2}{(s + 2)^2 + 9} = G(s + 2)$$

Examples

- $F(s) = \frac{s}{s^2 + 4s + 13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9} \Rightarrow \frac{s}{(s + 2)^2 + 9}$$

$$\Rightarrow \frac{(s + 2) - 2}{(s + 2)^2 + 9} = G(s + 2) \Rightarrow G(s) = \frac{s - 2}{s^2 + 9}$$

Examples

- $F(s) = \frac{s}{s^2+4s+13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9} \Rightarrow \frac{s}{(s + 2)^2 + 9}$$

$$\Rightarrow \frac{(s + 2) - 2}{(s + 2)^2 + 9} = G(s + 2) \Rightarrow G(s) = \frac{s - 2}{s^2 + 9}$$

$$L^{-1}[G(s)] = \cos(3t) - \frac{2}{3} \sin(3t)$$

Examples

- $F(s) = \frac{s}{s^2 + 4s + 13}$

$$\frac{s}{s^2 + 4s + 13} \Rightarrow \frac{s}{s^2 + 4s + 4 + 9} \Rightarrow \frac{s}{(s + 2)^2 + 9}$$

$$\Rightarrow \frac{(s + 2) - 2}{(s + 2)^2 + 9} = G(s + 2) \Rightarrow G(s) = \frac{s - 2}{s^2 + 9}$$

$$L^{-1}[G(s)] = \cos(3t) - \frac{2}{3} \sin(3t)$$

$$L^{-1}[F(s)] = f(t) = e^{-2t} \left(\cos(3t) - \frac{2}{3} \sin(3t) \right)$$

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Reminder

(Systems of) ODEs \Leftrightarrow (Inverse) Laplace Transform \Leftrightarrow Algebraic System

- (System of) Linear ODEs with constant coefficients
- Additionally, initial conditions are also needed to solve the ODEs
 - Implicitly, we are looking for a particular solution when solving ODEs using Laplace Transforms

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L \left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x \right] = L[0]$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L \left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x \right] = L[0]$$

$$\Rightarrow s^2X(s) - sx(0) - x'(0) + 4(sX(s) - x(0)) + 13X(s) = 0$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L\left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x\right] = L[0]$$

$$\Rightarrow s^2X(s) - sx(0) - x'(0) + 4(sX(s) - x(0)) + 13X(s) = 0$$

$$\Rightarrow s^2X(s) - 3s + 4sX(s) - 12 + 13X(s) = 0$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L\left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x\right] = L[0]$$

$$\Rightarrow s^2X(s) - sx(0) - x'(0) + 4(sX(s) - x(0)) + 13X(s) = 0$$

$$\Rightarrow s^2X(s) - 3s + 4sX(s) - 12 + 13X(s) = 0$$

$$\Rightarrow s^2X(s) + 4sX(s) + 13X(s) = 3s + 12$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L\left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x\right] = L[0]$$

$$\Rightarrow s^2X(s) - sx(0) - x'(0) + 4(sX(s) - x(0)) + 13X(s) = 0$$

$$\Rightarrow s^2X(s) - 3s + 4sX(s) - 12 + 13X(s) = 0$$

$$\Rightarrow s^2X(s) + 4sX(s) + 13X(s) = 3s + 12$$

$$\Rightarrow (s^2 + 4s + 13)X(s) = 3s$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$L\left[\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x\right] = L[0]$$

$$\Rightarrow s^2X(s) - sx(0) - x'(0) + 4(sX(s) - x(0)) + 13X(s) = 0$$

$$\Rightarrow s^2X(s) - 3s + 4sX(s) - 12 + 13X(s) = 0$$

$$\Rightarrow s^2X(s) + 4sX(s) + 13X(s) = 3s + 12$$

$$\Rightarrow (s^2 + 4s + 13)X(s) = 3s \Rightarrow X(s) = \frac{3s + 12}{s^2 + 4s + 13}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13} \Rightarrow X(s) = \frac{3s + 12}{(s + 2)^2 + 9}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13} \Rightarrow X(s) = \frac{3s + 12}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = \frac{3s + 6 + 6}{(s + 2)^2 + 9} \Rightarrow X(s) = \frac{3(s + 2) + 6}{(s + 2)^2 + 9}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13} \Rightarrow X(s) = \frac{3s + 12}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = \frac{3s + 6 + 6}{(s + 2)^2 + 9} \Rightarrow X(s) = \frac{3(s + 2) + 6}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = 3\frac{(s + 2)}{(s + 2)^2 + 9} + 2\frac{3}{(s + 2)^2 + 9}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13} \Rightarrow X(s) = \frac{3s + 12}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = \frac{3s + 6 + 6}{(s + 2)^2 + 9} \Rightarrow X(s) = \frac{3(s + 2) + 6}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = 3\frac{(s + 2)}{(s + 2)^2 + 9} + 2\frac{3}{(s + 2)^2 + 9}$$

$$\Rightarrow L^{-1}[X(s)] = 3L^{-1}\left[\frac{(s + 2)}{(s + 2)^2 + 9}\right] + 2L^{-1}\left[\frac{3}{(s + 2)^2 + 9}\right]$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 13x = 0$, where $x = 3$ and $\frac{dx}{dt} = 0$ when $t = 0$

$$X(s) = \frac{3s + 12}{s^2 + 4s + 13} \Rightarrow X(s) = \frac{3s + 12}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = \frac{3s + 6 + 6}{(s + 2)^2 + 9} \Rightarrow X(s) = \frac{3(s + 2) + 6}{(s + 2)^2 + 9}$$

$$\Rightarrow X(s) = 3\frac{(s + 2)}{(s + 2)^2 + 9} + 2\frac{3}{(s + 2)^2 + 9}$$

$$\Rightarrow L^{-1}[X(s)] = 3L^{-1}\left[\frac{(s + 2)}{(s + 2)^2 + 9}\right] + 2L^{-1}\left[\frac{3}{(s + 2)^2 + 9}\right]$$

$$x(t) = 3e^{-2t} \cos(3t) + 2e^{-2t} \sin(3t)$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$s^2X(s) - s - 4 + 6(sX(s) - 1) + 9X(s) = \frac{50}{s^2 + 1}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$s^2X(s) - s - 4 + 6(sX(s) - 1) + 9X(s) = \frac{50}{s^2 + 1}$$

$$\Rightarrow s^2X(s) + 6sX(s) + 9X(s) - s - 10 = \frac{50}{s^2 + 1}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$s^2X(s) - s - 4 + 6(sX(s) - 1) + 9X(s) = \frac{50}{s^2 + 1}$$

$$\Rightarrow s^2X(s) + 6sX(s) + 9X(s) - s - 10 = \frac{50}{s^2 + 1}$$

$$\Rightarrow (s^2 + 6s + 9)X(s) = \frac{50}{s^2 + 1} + s + 10$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$s^2X(s) - s - 4 + 6(sX(s) - 1) + 9X(s) = \frac{50}{s^2 + 1}$$

$$\Rightarrow s^2X(s) + 6sX(s) + 9X(s) - s - 10 = \frac{50}{s^2 + 1}$$

$$\Rightarrow (s^2 + 6s + 9)X(s) = \frac{50}{s^2 + 1} + s + 10$$

$$\Rightarrow X(s) = \frac{50}{(s^2 + 1)(s + 3)^2} + \frac{s + 10}{(s + 3)^2}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$\Rightarrow X(s) = \frac{50}{(s^2 + 1)(s + 3)^2} + \frac{s + 10}{(s + 3)^2}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$\Rightarrow X(s) = \frac{50}{(s^2 + 1)(s + 3)^2} + \frac{s + 10}{(s + 3)^2}$$

$$X(s) = \frac{5}{(s + 3)^2} + \frac{3}{s + 3} + \frac{-3s + 4}{s^2 + 1} + \frac{1}{s + 3} + \frac{7}{(s + 3)^2}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$\Rightarrow X(s) = \frac{50}{(s^2 + 1)(s + 3)^2} + \frac{s + 10}{(s + 3)^2}$$

$$X(s) = \frac{5}{(s + 3)^2} + \frac{3}{s + 3} + \frac{-3s + 4}{s^2 + 1} + \frac{1}{s + 3} + \frac{7}{(s + 3)^2}$$

$$X(s) = 12\frac{1}{(s + 3)^2} + 4\frac{1}{s + 3} - 3\frac{s}{s^2 + 1} + 4\frac{1}{s^2 + 1}$$

Examples

- $\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 50 \sin t$, where
 $x = 1$ and $\frac{dx}{dt} = 4$ when $t = 0$

$$\Rightarrow X(s) = \frac{50}{(s^2 + 1)(s + 3)^2} + \frac{s + 10}{(s + 3)^2}$$

$$X(s) = \frac{5}{(s + 3)^2} + \frac{3}{s + 3} + \frac{-3s + 4}{s^2 + 1} + \frac{1}{s + 3} + \frac{7}{(s + 3)^2}$$

$$X(s) = 12\frac{1}{(s + 3)^2} + 4\frac{1}{s + 3} - 3\frac{s}{s^2 + 1} + 4\frac{1}{s^2 + 1}$$

$$x(t) = 12te^{-3t} + 4e^{-3t} - 3\cos t + 4\sin t$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow s^2X(s) - s + 4 + 4(sX(s) - 1) - 3X(s) = \frac{4}{s - 1}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow s^2X(s) - s + 4 + 4(sX(s) - 1) - 3X(s) = \frac{4}{s-1}$$

$$(s^2 + 4s - 3)X(s) = \frac{4}{s-1} + s + 2$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow s^2X(s) - s + 4 + 4(sX(s) - 1) - 3X(s) = \frac{4}{s-1}$$

$$(s^2 + 4s - 3)X(s) = \frac{4}{s-1} + s + 2$$

$$\Rightarrow X(s) = \frac{4}{(s-1)(s^2 + 4s - 3)} + \frac{s+2}{s^2 + 4s - 3}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow s^2X(s) - s + 4 + 4(sX(s) - 1) - 3X(s) = \frac{4}{s-1}$$

$$(s^2 + 4s - 3)X(s) = \frac{4}{s-1} + s + 2$$

$$\Rightarrow X(s) = \frac{4}{(s-1)(s^2 + 4s - 3)} + \frac{s+2}{s^2 + 4s - 3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-2s-10}{s^2 + 4s - 3} + \frac{s+2}{s^2 + 4s - 3}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow s^2X(s) - s + 4 + 4(sX(s) - 1) - 3X(s) = \frac{4}{s-1}$$

$$(s^2 + 4s - 3)X(s) = \frac{4}{s-1} + s + 2$$

$$\Rightarrow X(s) = \frac{4}{(s-1)(s^2 + 4s - 3)} + \frac{s+2}{s^2 + 4s - 3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-2s-10}{s^2 + 4s - 3} + \frac{s+2}{s^2 + 4s - 3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2 + 4s - 3}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2+4s-3}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2+4s-3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-2-6}{(s+2)^2-7}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2+4s-3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-2-6}{(s+2)^2-7}$$

$$\Rightarrow X(s) = \frac{2}{s-1} - \frac{s+2}{(s+2)^2-7} - \frac{6}{(s+2)^2-7}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2+4s-3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-2-6}{(s+2)^2-7}$$

$$\Rightarrow X(s) = \frac{2}{s-1} - \frac{s+2}{(s+2)^2-7} - \frac{6}{(s+2)^2-7}$$

$$\Rightarrow X(s) = 2\frac{1}{s-1} + (-1)\frac{s+2}{(s+2)^2-7} + \left(-\frac{6}{\sqrt{7}}\right)\frac{\sqrt{7}}{(s+2)^2-7}$$

Examples

- $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} - 3x = 4e^t$, where $x = 1$ and $\frac{dx}{dt} = -2$ when $t = 0$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-8}{s^2+4s-3}$$

$$\Rightarrow X(s) = \frac{2}{s-1} + \frac{-s-2-6}{(s+2)^2-7}$$

$$\Rightarrow X(s) = \frac{2}{s-1} - \frac{s+2}{(s+2)^2-7} - \frac{6}{(s+2)^2-7}$$

$$\Rightarrow X(s) = 2\frac{1}{s-1} + (-1)\frac{s+2}{(s+2)^2-7} + \left(-\frac{6}{\sqrt{7}}\right)\frac{\sqrt{7}}{(s+2)^2-7}$$

$$x(t) = 2e^t - e^{-2t} \cosh(\sqrt{7}t) - \frac{6}{\sqrt{7}}e^{-2t} \sinh(\sqrt{7}t)$$

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

General Solution of an ODE using (Inverse) Laplace Transforms

- In the event that no boundary conditions are given, we can still solve an ODE using Laplace transforms **by setting $x(0) = A$ and $x'(0) = B$** , where A and B are arbitrary constants.

General Solution of an ODE using (Inverse) Laplace Transforms

- In the event that no boundary conditions are given, we can still solve an ODE using Laplace transforms **by setting $x(0) = A$ and $x'(0) = B$** , where A and B are arbitrary constants.
- In actuality, we could solve the previous examples by **solving first for the general solution, then plugging the boundary conditions** to get the particular solution.
 - Obviously, that would take a little bit more work (which we usually **DO NOT** like), 'no?

Example

- $\frac{d^2x}{dt^2} + 4x = 0$

Example

- $\frac{d^2x}{dt^2} + 4x = 0$

$$s^2X(s) - As - B + 4X(s) = 0$$

Example

- $\frac{d^2x}{dt^2} + 4x = 0$

$$s^2X(s) - As - B + 4X(s) = 0 \Rightarrow X(s) = \frac{As + B}{s^2 + 4}$$

Example

- $\frac{d^2x}{dt^2} + 4x = 0$

$$s^2X(s) - As - B + 4X(s) = 0 \Rightarrow X(s) = \frac{As + B}{s^2 + 4}$$

$$\Rightarrow X(s) = A \frac{s}{s^2 + 4} + B \frac{1}{s^2 + 4}$$

Example

- $\frac{d^2x}{dt^2} + 4x = 0$

$$s^2X(s) - As - B + 4X(s) = 0 \Rightarrow X(s) = \frac{As + B}{s^2 + 4}$$

$$\Rightarrow X(s) = A \frac{s}{s^2 + 4} + B \frac{1}{s^2 + 4}$$

$$x(t) = A \cos(2t) + B \sin(2t)$$

Topics

- 1 Laplace Transforms
 - Laplace Transforms and Solving ODEs
 - Laplace Transform
 - Useful Rules and Theorems
- 2 Inverse Laplace Transforms
- 3 Solving First and Second Order ODEs
 - Examples
 - General Solutions of ODEs and Laplace Transforms
- 4 Solving Systems of ODEs

Solving Systems of ODEs using Laplace Transforms

- Firstly, take the Laplace Transform of each equations.

Solving Systems of ODEs using Laplace Transforms

- Firstly, take the Laplace Transform of each equations.
- Solve the resulting linear system algebraically.
 - One dependent variable gets solved

Solving Systems of ODEs using Laplace Transforms

- Firstly, take the **Laplace Transform of each equations.**
- Solve the resulting **linear system algebraically.**
 - **One dependent variable gets solved**
- Apply **inverse Laplace transform** on the solution for one variable.

Solving Systems of ODEs using Laplace Transforms

- Firstly, take the **Laplace Transform of each equations.**
- Solve the resulting **linear system algebraically.**
 - **One dependent variable gets solved**
- Apply **inverse Laplace transform** on the solution for one variable.
- **“Back substitute”** to solve for the other variable(s) using the obtained solution for one dependent variable.

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Firstly, take the Laplace Transform of each equations.

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Firstly, take the **Laplace Transform of each equations.**

$$sY(s) - 2 + 2X(s) = \frac{1}{s-1}$$

$$sX(s) - 1 - 2Y(s) = \frac{1}{s} + \frac{1}{s^2}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Solve the resulting **linear system algebraically**.

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Solve the resulting **linear system algebraically**.

$$sY(s) + 2X(s) = \frac{1}{s-1} + 2$$

$$sX(s) - 2Y(s) = \frac{1}{s} + \frac{1}{s^2} + 1$$

Examples

$$\begin{aligned}\frac{dy}{dt} + 2x &= e^t \\ \frac{dx}{dt} - 2y &= 1 + t \\ x(0) &= 1 \text{ and } y(0) = 2\end{aligned}$$

- Solve the resulting **linear system algebraically**.

$$sY(s) + 2X(s) = \frac{1}{s-1} + 2$$

$$sX(s) - 2Y(s) = \frac{1}{s} + \frac{1}{s^2} + 1$$

$$\Rightarrow (4 + s^2)X(s) = \frac{2}{s-1} + 4 + 1 + \frac{1}{s} + s$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Solve the resulting **linear system algebraically**.

$$(4 + s^2)X(s) = \frac{2}{s-1} + 4 + 1 + \frac{1}{s} + s$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Solve the resulting **linear system algebraically**.

$$(4 + s^2)X(s) = \frac{2}{s-1} + 4 + 1 + \frac{1}{s} + s$$

$$\Rightarrow X(s) = \frac{2}{(s-1)(4+s^2)} + \frac{5}{4+s^2} + \frac{1}{s(4+s^2)} + \frac{s}{4+s^2}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2}{(s-1)(4+s^2)} + \frac{5}{4+s^2} + \frac{1}{s(4+s^2)} + \frac{s}{4+s^2}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2}{(s-1)(4+s^2)} + \frac{5}{4+s^2} + \frac{1}{s(4+s^2)} + \frac{s}{4+s^2}$$

$$\Rightarrow X(s) = \frac{2}{5} \frac{1}{s-1} + \frac{7}{20} \frac{s}{s^2+4} + \frac{23}{5} \frac{1}{s^2+4} + \frac{1}{4} \frac{1}{s}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2}{5} \frac{1}{s-1} + \frac{7}{20} \frac{s}{s^2+4} + \frac{23}{5} \frac{1}{s^2+4} + \frac{11}{4} \frac{1}{s}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2}{5} \frac{1}{s-1} + \frac{7}{20} \frac{s}{s^2+4} + \frac{23}{5} \frac{1}{s^2+4} + \frac{11}{4} \frac{1}{s}$$

$$x(t) = \frac{2}{5} e^t + \frac{7}{20} \cos(2t) + \frac{23}{10} \sin(2t) + \frac{1}{4}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$x(t) = \frac{2}{5}e^t + \frac{7}{20}\cos(2t) + \frac{23}{10}\sin(2t) + \frac{1}{4}$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$x(t) = \frac{2}{5}e^t + \frac{7}{20}\cos(2t) + \frac{23}{10}\sin(2t) + \frac{1}{4}$$

$$y = \frac{1}{2} \left(\frac{dx}{dt} - 1 - t \right)$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y = \frac{1}{2} \left(\frac{dx}{dt} - 1 - t \right)$$

Examples

$$\frac{dy}{dt} + 2x = e^t$$

$$\frac{dx}{dt} - 2y = 1 + t$$

$$x(0) = 1 \text{ and } y(0) = 2$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y = \frac{1}{2} \left(\frac{dx}{dt} - 1 - t \right)$$

$$y(t) = \frac{1}{5}e^t + \frac{23}{10}\cos(2t) - \frac{7}{20}\sin(2t) - \frac{1}{2} - \frac{t}{2}$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Firstly, take the **Laplace Transform of each equations.**

$$sX(s) - 3 - Y(s) = \frac{1}{s^2 + 1}$$

$$sY(s) - 4 + X(s) = \frac{s}{s^2 + 1}$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Solve the resulting **linear system algebraically**.

$$sX(s) - Y(s) = \frac{1}{s^2 + 1} + 3$$

$$sY(s) + X(s) = \frac{s}{s^2 + 1} + 4$$

Examples

$$\begin{aligned}\frac{dx}{dt} - y &= \sin t \\ \frac{dy}{dt} + x &= \cos t \\ x(0) &= 3 \text{ and } y(0) = 4\end{aligned}$$

- Solve the resulting **linear system algebraically**.

$$sX(s) - Y(s) = \frac{1}{s^2 + 1} + 3$$

$$sY(s) + X(s) = \frac{s}{s^2 + 1} + 4$$

$$\Rightarrow s^2 X(s) + X(s) = \frac{2s}{s^2 + 1} + 3s + 4$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Solve the resulting **linear system algebraically**.

$$s^2 X(s) + X(s) = \frac{2}{s^2 + 1} + 3s + 4$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Solve the resulting **linear system algebraically**.

$$s^2 X(s) + X(s) = \frac{2}{s^2 + 1} + 3s + 4$$

$$\Rightarrow X(s) = \frac{2s}{(s^2 + 1)^2} + \frac{3s}{s^2 + 1} + \frac{4}{s^2 + 1}$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2s}{(s^2 + 1)^2} + \frac{3s}{s^2 + 1} + \frac{4}{s^2 + 1}$$

Examples

$$\frac{dx}{dt} - y = \sin t$$

$$\frac{dy}{dt} + x = \cos t$$

$$x(0) = 3 \text{ and } y(0) = 4$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$X(s) = \frac{2s}{(s^2 + 1)^2} + \frac{3s}{s^2 + 1} + \frac{4}{s^2 + 1}$$

$$x(t) = t \sin t + 3 \cos t + 4 \sin t$$

Examples

$$\begin{aligned}\frac{dx}{dt} - y &= \sin t \\ \frac{dy}{dt} + x &= \cos t \\ x(0) &= 3 \text{ and } y(0) = 4\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$x(t) = t \sin t + 3 \cos t + 4 \sin t$$

Examples

$$\begin{aligned}\frac{dx}{dt} - y &= \sin t \\ \frac{dy}{dt} + x &= \cos t \\ x(0) &= 3 \text{ and } y(0) = 4\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$x(t) = t \sin t + 3 \cos t + 4 \sin t$$

$$y = \frac{dy}{dt} - \sin t$$

Examples

$$\begin{aligned}\frac{dx}{dt} - y &= \sin t \\ \frac{dy}{dt} + x &= \cos t \\ x(0) &= 3 \text{ and } y(0) = 4\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$x(t) = t \sin t + 3 \cos t + 4 \sin t$$

$$y = \frac{dy}{dt} - \sin t$$

$$y(t) = t \cos t - 4 \sin t + 4 \cos t$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Firstly, take the **Laplace Transform of each equations.**

$$sX(s) - 3 + 4sY(s) + 6Y(s) = 0$$

$$5sX(s) - 15 + 2sY(s) + 6X(s) = 0$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$sX(s) + (4s + 6)Y(s) = 3$$

$$(5s + 6)X(s) + 2sY(s) = 15$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$sX(s) + (4s + 6)Y(s) = 3$$

$$(5s + 6)X(s) + 2sY(s) = 15$$

$$\Rightarrow \left(\left(-\frac{5s+6}{s} \right) (4s+6) + 2s \right) Y(s) = -3\frac{5s+6}{s} + 15$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$\left(\left(\frac{-5s - 6}{s} \right) (4s + 6) + 2s \right) Y(s) = -3 \frac{5s + 6}{s} + 15$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$\left(\left(\frac{-5s-6}{s} \right) (4s+6) + 2s \right) Y(s) = -3 \frac{5s+6}{s} + 15$$

$$\Rightarrow \left(\frac{-18s^2 - 30s - 36}{s} \right) Y(s) = \frac{-18}{s}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$\left(\frac{-18s^2 - 54s - 36}{s} \right) Y(s) = \frac{-18}{s}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Solve the resulting **linear system algebraically**.

$$\left(\frac{-18s^2 - 54s - 36}{s} \right) Y(s) = \frac{-18}{s}$$

$$Y(s) = \frac{18}{18s^2 + 54s + 36} = \frac{1}{s^2 + 3s + 2}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$Y(s) = \frac{1}{s^2 + 3s + 2}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$Y(s) = \frac{1}{s^2 + 3s + 2} = \frac{-1}{s + 2} + \frac{1}{s + 1}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- Apply **inverse Laplace transform** on the solution for one variable.

$$Y(s) = \frac{1}{s^2 + 3s + 2} = \frac{-1}{s + 2} + \frac{1}{s + 1}$$

$$y(t) = -e^{-2t} + e^{-t}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

$$x = \frac{1}{6} \left(5\frac{dx}{dt} + 2\frac{dy}{dt} \right)$$

Examples

$$\frac{dx}{dt} + 4\frac{dy}{dt} + 6y = 0$$

$$5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x = 0$$

$$x(0) = 3 \text{ and } y(0) = 0$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

Examples

$$\begin{aligned}\frac{dx}{dt} + 4\frac{dy}{dt} + 6y &= 0 \\ 5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x &= 0 \\ x(0) = 3 \text{ and } y(0) &= 0\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

$$\frac{dx}{dt} = -4\frac{dy}{dt} - 6y$$

Examples

$$\begin{aligned}\frac{dx}{dt} + 4\frac{dy}{dt} + 6y &= 0 \\ 5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x &= 0 \\ x(0) = 3 \text{ and } y(0) &= 0\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

$$\frac{dx}{dt} = -4\frac{dy}{dt} - 6y = -4(2e^{-2t} - e^{-t}) - 6(-e^{-2t} + e^{-t})$$

Examples

$$\begin{aligned}\frac{dx}{dt} + 4\frac{dy}{dt} + 6y &= 0 \\ 5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x &= 0 \\ x(0) = 3 \text{ and } y(0) &= 0\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

$$\frac{dx}{dt} = -4\frac{dy}{dt} - 6y = -4(2e^{-2t} - e^{-t}) - 6(-e^{-2t} + e^{-t}) = -2e^{-2t} - 2e^{-t}$$

Examples

$$\begin{aligned}\frac{dx}{dt} + 4\frac{dy}{dt} + 6y &= 0 \\ 5\frac{dx}{dt} + 2\frac{dy}{dt} + 6x &= 0 \\ x(0) &= 3 \text{ and } y(0) = 0\end{aligned}$$

- “Back substitute” to solve for the other variable(s) using the obtained solution for one dependent variable.

$$y(t) = -e^{-2t} + e^{-t}$$

$$\frac{dx}{dt} = -4\frac{dy}{dt} - 6y = -4(2e^{-2t} - e^{-t}) - 6(-e^{-2t} + e^{-t}) = -2e^{-2t} - 2e^{-t}$$

$$x(t) = e^{-2t} + 2e^{-t}$$

END OF LESSON 10