

Linear Transformations

CS 130 - Mathematical Methods in Computer Science

Jan Michael C. Yap

Department of Computer Science
University of the Philippines Diliman

November 22, 2012

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes

- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range

- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Scaling and projecting a vector

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

Scaling and projecting a vector

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f \left(\left(\begin{pmatrix} 2 \\ -1 \\ -5 \end{pmatrix} \right) \right) = \begin{pmatrix} 4 \\ -2 \end{pmatrix}$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix}\right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix}\right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f\left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} d \\ e \\ f \end{pmatrix}\right)$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix}\right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f\left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} d \\ e \\ f \end{pmatrix}\right) = f\left(\begin{pmatrix} ka + d \\ kb + e \\ kc + f \end{pmatrix}\right)$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix}\right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f\left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} d \\ e \\ f \end{pmatrix}\right) = f\left(\begin{pmatrix} ka + d \\ kb + e \\ kc + f \end{pmatrix}\right) = \begin{pmatrix} 2(ka + d) \\ 2(kb + e) \end{pmatrix}$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \\ c \end{pmatrix}\right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$\begin{aligned} f\left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} d \\ e \\ f \end{pmatrix}\right) &= f\left(\begin{pmatrix} ka + d \\ kb + e \\ kc + f \end{pmatrix}\right) = \begin{pmatrix} 2(ka + d) \\ 2(kb + e) \end{pmatrix} \\ &= \begin{pmatrix} 2ka + 2d \\ 2kb + 2e \end{pmatrix} \end{aligned}$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f \left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) + f \left(\begin{pmatrix} d \\ e \\ f \end{pmatrix} \right)$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f \left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) + f \left(\begin{pmatrix} d \\ e \\ f \end{pmatrix} \right) = \begin{pmatrix} 2ka \\ 2kb \end{pmatrix} + \begin{pmatrix} 2d \\ 2e \end{pmatrix}$$

Linear transformation

- Let V and W be vector spaces with vectors $u, v \in V$, a scalar c , and a function $L : V \rightarrow W$. L is a **linear transformation** if

$$L(cu + v) = cL(u) + L(v)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$f \left(k \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) + f \left(\begin{pmatrix} d \\ e \\ f \end{pmatrix} \right) = \begin{pmatrix} 2ka \\ 2kb \end{pmatrix} + \begin{pmatrix} 2d \\ 2e \end{pmatrix} = \begin{pmatrix} 2ka + 2d \\ 2kb + 2e \end{pmatrix}$$

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Some Linear Transformations

- Projection

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^{(n-1)}, L \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{pmatrix} \right) = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \end{pmatrix}$$

Some Linear Transformations

- Projection

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^{(n-1)}, L \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{pmatrix} \right) = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \end{pmatrix}$$

- Dilation

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^n, L \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = r \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, r > 1$$

Some Linear Transformations

- Contraction

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^n, L \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = r \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, 0 < r < 1$$

Some Linear Transformations

- Contraction

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^n, L \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = r \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, 0 < r < 1$$

- Rotation in \mathbb{R}^2

$$L : \mathbb{R}^n \rightarrow \mathbb{R}^n, L \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

Some Linear Transformations

- Reflection about the main axis in \mathbb{R}^2

$$L_1 : \mathbb{R}^n \rightarrow \mathbb{R}^n, L_1 \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

Some Linear Transformations

- Reflection about the main axis in \mathbb{R}^2

$$L_1 : \mathbb{R}^n \rightarrow \mathbb{R}^n, L_1 \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$L_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n, L_2 \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Notes on Linear Transformations

- $L : V \rightarrow V$ is called a **linear operator**

Notes on Linear Transformations

- $L : V \rightarrow V$ is called a **linear operator**
- $L(u), u \in V$ is called the **image of u (under the linear transformation L)**

Notes on Linear Transformations

- $L : V \rightarrow V$ is called a **linear operator**
- $L(u), u \in V$ is called the **image of u (under the linear transformation L)**
- $L(0_V) = 0_W$

Notes on Linear Transformations

- $L : V \rightarrow V$ is called a **linear operator**
- $L(u), u \in V$ is called the **image of u (under the linear transformation L)**
- $L(0_V) = 0_W$
- $L(u - v) = L(u) - L(v)$

Notes on Linear Transformations

- $L : V \rightarrow V$ is called a **linear operator**
- $L(u), u \in V$ is called the **image of u (under the linear transformation L)**
- $L(0_V) = 0_W$
- $L(u - v) = L(u) - L(v)$
- If $B = \{b_1, b_2, \dots, b_n\}$ is a basis for the n -dimensional vector space V and $u \in V$, then **$L(u) = \sum_{i=1}^n c_i L(b_i)$**

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$\begin{pmatrix} ? \\ ? \\ ? \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$\begin{pmatrix} ? \\ ? \\ ? \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} 0 \\ 0 \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$\begin{pmatrix} ? \\ ? \\ ? \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{pmatrix} 0 \\ 0 \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\ker(f) = \left\{ \begin{pmatrix} 0 \\ 0 \\ c \end{pmatrix} \mid c \in \mathbb{R} \right\}$$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n, f \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = r \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, r > 1$$

Kernel

- The **kernel of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the subset of vectors in V such that, for a vector u in that subset, $L(u) = 0_W$

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n, f \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = r \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, r > 1$$

$$\ker(f) = \left\{ \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \right\}$$

Some Notes on the Kernel

- $\ker(L)$ of $L : V \rightarrow W$ is a **subspace** of V

Some Notes on the Kernel

- $\ker(L)$ of $L : V \rightarrow W$ is a **subspace** of V
- $\ker(L)$ is **never empty** (Y?)

Some Notes on the Kernel

- $\ker(L)$ of $L : V \rightarrow W$ is a **subspace** of V
- $\ker(L)$ is **never empty** (Y?)
- If $\ker(L)$ contains only 0_V , then L is **one-to-one**

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Range

- The **range** of a linear transformation $L : V \rightarrow W$, denoted $\text{ker}(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

Range

- The **range of a linear transformation** $L : V \rightarrow W$, denoted $\text{ker}(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

Range

- The **range of a linear transformation** $L : V \rightarrow W$, denoted $\text{ker}(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$\text{range}(f) = \mathbb{R}^2$$

Range

- The **range** of a linear transformation $L : V \rightarrow W$, denoted $\text{ker}(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \end{pmatrix} \right) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

Range

- The **range** of a linear transformation $L : V \rightarrow W$, denoted $\text{ker}(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \end{pmatrix} \right) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix}$$

Range

- The **range** of a linear transformation $L : V \rightarrow W$, denoted $\ker(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \end{pmatrix}\right) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & -1 & c \\ -1 & 1 & d \end{pmatrix}$$

Range

- The **range** of a linear transformation $L : V \rightarrow W$, denoted $\ker(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \end{pmatrix}\right) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & -1 & c \\ -1 & 1 & d \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 1 & -1 & c \\ 0 & 0 & c + d \end{pmatrix}$$

Range

- The **range of a linear transformation** $L : V \rightarrow W$, denoted $\ker(L)$, is the **subset of vectors in W containing all images of the vectors in V under L .**

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f\left(\begin{pmatrix} a \\ b \end{pmatrix}\right) = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & -1 & c \\ -1 & 1 & d \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 1 & -1 & c \\ 0 & 0 & c + d \end{pmatrix}$$

$$\text{range}(f) = \left\{ \begin{pmatrix} c \\ -c \end{pmatrix} \mid c \in \mathbb{R} \right\}$$

Some Notes on the Range

- $\text{range}(L)$ of $L : V \rightarrow W$ is a **subspace** of W

Some Notes on the Range

- $\text{range}(L)$ of $L : V \rightarrow W$ is a **subspace** of W
- $\text{range}(L)$ is **also never empty** ($Z?$)

Some Notes on the Range

- $\text{range}(L)$ of $L : V \rightarrow W$ is a **subspace** of W
- $\text{range}(L)$ is **also never empty** ($Z?$)
- If $\text{range}(L) = W$, then L is **onto**

Some Notes on the Range

- $\text{range}(L)$ of $L : V \rightarrow W$ is a **subspace** of W
- $\text{range}(L)$ is **also never empty** ($Z?$)
- If $\text{range}(L) = W$, then L is **onto**
- Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m, L(u) = A_{m \times n}u$, then $\text{range}(L)$ is the **column space of A** and $\ker(L)$ is the **null space of A**

Some Notes on the Range

- $\text{range}(L)$ of $L : V \rightarrow W$ is a **subspace** of W
- $\text{range}(L)$ is **also never empty** (Z ?)
- If $\text{range}(L) = W$, then L is **onto**
- Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $L(u) = A_{m \times n}u$, then $\text{range}(L)$ is the **column space of A** and $\text{ker}(L)$ is the **null space of A**
 - $\dim(\text{range}(L)) + \dim(\text{ker}(L)) = n = \dim(V)$

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Matrix of a Linear Transformation

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

Matrix of a Linear Transformation

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}$$

Matrix of a Linear Transformation

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}$$

$$\Downarrow$$

$$A \begin{pmatrix} a \\ b \\ c \end{pmatrix}$$

Matrix of a Linear Transformation

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}$$

$$\Downarrow$$

$$A \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix}$$

Matrix of a Linear Transformation

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2, f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right) = \begin{pmatrix} 2a \\ 2b \end{pmatrix}$$

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}$$

$$\Downarrow$$

$$A \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 2a \\ 2b \end{pmatrix} = f \left(\begin{pmatrix} a \\ b \\ c \end{pmatrix} \right)$$

Matrix of a Linear Transformation

- Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then there exists an $m \times n$ matrix A such that $L(x) = Ax, x \in \mathbb{R}^n$

Matrix of a Linear Transformation

- Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then there exists an $m \times n$ matrix A such that $L(x) = Ax, x \in \mathbb{R}^n$
 - $A = (L(e_1) \ L(e_2) \cdots L(e_n))$ where e_i s are the vectors of the standard/natural basis for \mathbb{R}^n

Matrix of a Linear Transformation

- Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then there exists an $m \times n$ matrix A such that $L(x) = Ax, x \in \mathbb{R}^n$
 - $A = (L(e_1) \ L(e_2) \ \cdots \ L(e_n))$ where e_i s are the vectors of the standard/natural basis for \mathbb{R}^n

$$A = \left(f \left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) \ f \left(\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right) \ f \left(\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right) \right)$$

Topics

- 1 Linear Transformations
 - Linear Transformations
 - Some Linear Transformations
 - Some Notes
- 2 Kernel and Range of a Linear Transformation
 - Kernel
 - Range
- 3 Matrix of a Linear Transformation
 - Matrix of a Linear Transformation
 - General Change of Basis

Change of basis redux

- Transition matrix from basis S to basis T : $P_{S \rightarrow T}$

Change of basis redux

- Transition matrix from basis S to basis T : $P_{S \rightarrow T}$
 - Form the augmented matrix $T|S$
 - Transform augmented matrix to RREF
 - The square matrix on the right hand side of the RREF matrix is $P_{S \rightarrow T}$

Change of basis redux

- Transition matrix from basis S to basis T : $P_{S \rightarrow T}$
 - Form the augmented matrix $T|S$
 - Transform augmented matrix to RREF
 - The square matrix on the right hand side of the RREF matrix is $P_{S \rightarrow T}$
- Extend this to work with linearly transformed vectors across different vector spaces

Generalizing the change of basis process

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $S = \{v_1, v_2, \dots, v_n\}$ and $T = \{w_1, w_2, \dots, w_m\}$ be bases for \mathbb{R}^n and \mathbb{R}^m respectively, then the **matrix of the linear transformation L w.r.t. S and T** , denoted as $P_{S \rightarrow T}^L$, is obtained as follows:

Generalizing the change of basis process

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $S = \{v_1, v_2, \dots, v_n\}$ and $T = \{w_1, w_2, \dots, w_m\}$ be bases for \mathbb{R}^n and \mathbb{R}^m respectively, then the **matrix of the linear transformation L w.r.t. S and T** , denoted as $P_{S \rightarrow T}^L$, is obtained as follows:

- Form the **$m \times (m + n)$ augmented matrix $T|L(S)$** , where $L(S)$ is the image of the vectors in S under L

Generalizing the change of basis process

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $S = \{v_1, v_2, \dots, v_n\}$ and $T = \{w_1, w_2, \dots, w_m\}$ be bases for \mathbb{R}^n and \mathbb{R}^m respectively, then the **matrix of the linear transformation L w.r.t. S and T** , denoted as $P_{S \rightarrow T}^L$, is obtained as follows:

- Form the **$m \times (m + n)$ augmented matrix $T|L(S)$** , where $L(S)$ is the image of the vectors in S under L
- ???

Generalizing the change of basis process

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $S = \{v_1, v_2, \dots, v_n\}$ and $T = \{w_1, w_2, \dots, w_m\}$ be bases for \mathbb{R}^n and \mathbb{R}^m respectively, then the **matrix of the linear transformation L w.r.t. S and T**, denoted as $P_{S \rightarrow T}^L$, is obtained as follows:

- Form the **$m \times (m + n)$ augmented matrix $T|L(S)$** , where $L(S)$ is the image of the vectors in S under L
- ???
- !!!

Generalizing the change of basis process

- Form the $m \times (m + n)$ augmented matrix $T|L(S)$, where $L(S)$ is the image of the vectors in S under L

$$S = \left\{ \begin{pmatrix} 0 \\ 5 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}, T = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}$$

Generalizing the change of basis process

- Form the $m \times (m + n)$ augmented matrix $T|L(S)$, where $L(S)$ is the image of the vectors in S under L

$$S = \left\{ \begin{pmatrix} 0 \\ 5 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}, T = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}$$

$$\Downarrow$$

$$T|L(S) = \begin{pmatrix} 2 & -1 & 0 & -2 & 2 \\ 1 & 1 & 10 & 6 & -2 \end{pmatrix}$$

Generalizing the change of basis process

- ???

$$\begin{pmatrix} 2 & -1 & 0 & -2 & 2 \\ 1 & 1 & 10 & 6 & -2 \end{pmatrix}$$

Generalizing the change of basis process

- ???

$$\begin{pmatrix} 2 & -1 & 0 & -2 & 2 \\ 1 & 1 & 10 & 6 & -2 \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & 0 & \frac{10}{3} & \frac{4}{3} & 0 \\ 0 & 1 & \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

Generalizing the change of basis process

• !!!

$$\begin{pmatrix} 1 & 0 & \frac{10}{3} & \frac{4}{3} & 0 \\ 0 & 1 & \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

Generalizing the change of basis process

• !!!

$$\begin{pmatrix} 1 & 0 & \frac{10}{3} & \frac{4}{3} & 0 \\ 0 & 1 & \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix} \Rightarrow P_{S \rightarrow T}^L = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

General change of basis

$$S = \left\{ \begin{pmatrix} 0 \\ 5 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}, T = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\},$$

$$x = \begin{pmatrix} 8 \\ 9 \\ 10 \end{pmatrix}, f(x) = \begin{pmatrix} 16 \\ 18 \end{pmatrix}$$

General change of basis

$$S = \left\{ \begin{pmatrix} 0 \\ 5 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}, T = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\},$$

$$x = \begin{pmatrix} 8 \\ 9 \\ 10 \end{pmatrix}, f(x) = \begin{pmatrix} 16 \\ 18 \end{pmatrix}$$

\Downarrow

$$[x]_S = \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}, [f(x)]_T = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \end{pmatrix}$$

General change of basis

$$[x]_S = \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}, [f(x)]_T = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \\ \frac{3}{3} \end{pmatrix}, P_{S \rightarrow T}^L = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

General change of basis

$$[x]_S = \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}, [f(x)]_T = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \\ \frac{3}{3} \end{pmatrix}, P_{S \rightarrow T}^L = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

$$\Downarrow$$

$$P_{S \rightarrow T}^L [x]_S$$

General change of basis

$$[x]_S = \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}, [f(x)]_T = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \\ \frac{3}{3} \end{pmatrix}, P_{S \rightarrow T}^L = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

$$\Downarrow$$

$$P_{S \rightarrow T}^L [x]_S = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix} \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}$$

General change of basis

$$[x]_S = \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix}, [f(x)]_T = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \\ \frac{14}{3} \end{pmatrix}, P_{S \rightarrow T}^L = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix}$$

$$\Downarrow$$

$$P_{S \rightarrow T}^L [x]_S = \begin{pmatrix} \frac{10}{3} & \frac{4}{3} & 0 \\ \frac{20}{3} & \frac{14}{3} & -2 \end{pmatrix} \begin{pmatrix} \frac{47}{9} \\ -\frac{41}{9} \\ \frac{31}{9} \end{pmatrix} = \begin{pmatrix} \frac{34}{3} \\ \frac{20}{3} \\ \frac{14}{3} \end{pmatrix} = [f(x)]_T$$

END OF LESSON 5