1

AI1103-Assignment 1

revanth badavathu CS20BTECH11007

Download all python codes from

https://github.com/cs20btech11007/assignment1/blob/main/assignment1/assignment1.py

and latex-tikz codes from

https://github.com/cs20btech11007/assignment1/blob/main/assignment1/assignment1.tex

PROBLEM 1.12

A coin is biased so that the head is 3 times as likely to occur as tail. If the coin is tossed twice, find the probability distribution of number of tails.

sol.

let X be random variable $X \in \{0, 1, 2\}$ denotes outcomes of tail in a experiment. using binomial distribution and now finding

using binomial distribution and now finding probability distribution of number of tails in the events.

$$Pr(X = k) = {}^{n}C_{k}p^{k}(1-p)^{n-k}$$
 (0.0.1)

Here,

Symbol	Meaning
k	no. of tails in two throws of a coin
n	no. of throws $= 2$
p	Pr of getting tail in single throw= $\frac{1}{4}$

TABLE 0: This table gives the meaning of each symbol used in the formula

To find the probability of getting tails in 2 throws of a single coin:

$$\Pr(X = 0) = {}^{2}C_{0} \times \left(\frac{1}{4}\right)^{0} \times \left(\frac{3}{4}\right)^{2} = \frac{9}{16} \qquad (0.0.2)$$

$$\Pr(X = 1) = {}^{2}C_{1} \times \left(\frac{1}{4}\right)^{1} \times \left(\frac{3}{4}\right)^{1} = \frac{6}{16} \qquad (0.0.3)$$

$$\Pr(X=2) = {}^{2}C_{2} \times \left(\frac{1}{4}\right)^{2} \times \left(\frac{3}{4}\right)^{0} = \frac{1}{16} \qquad (0.0.4)$$

(0.0.5)