

CS 4110

# Programming Languages & Logics

---

## Lecture 16 Fixed-Point Combinators



# Termination in the $\lambda$ -calculus

We have encoded lots of useful programming functionality that produces values.

Does every closed  $\lambda$ -term eventually terminate under CBN evaluation?

$$\forall \text{ closed term } e. \exists e'. e \rightarrow^* e' \wedge e' \not\rightarrow ?$$

# Termination in the $\lambda$ -calculus

We have encoded lots of useful programming functionality that produces values.

Does every closed  $\lambda$ -term eventually terminate under CBN evaluation?

$$\forall \text{ closed term } e. \exists e'. e \rightarrow^* e' \wedge e' \not\rightarrow ?$$

No!

$$\Omega \triangleq (\lambda x. x x) (\lambda x. x x)$$

# Termination in the $\lambda$ -calculus

We have encoded lots of useful programming functionality that produces values.

Does every closed  $\lambda$ -term eventually terminate under CBN evaluation?

$$\forall \text{ closed term } e. \exists e'. e \rightarrow^* e' \wedge e' \not\rightarrow ?$$

No!

$$\begin{aligned}\Omega &\triangleq (\lambda x. x x) (\lambda x. x x) \\ &\rightarrow (x x) \{(\lambda x. x x)/x\} \\ &= (\lambda x. x x) (\lambda x. x x) \\ &= \Omega\end{aligned}$$

# Recursive Functions

---

How would we write recursive functions, like factorial?

# Recursive Functions

---

How would we write recursive functions, like factorial?

We'd like to write it like this...

$$\text{FACT} \triangleq \lambda n. \text{IF } (\text{ISZERO } n) \text{ 1 } (\text{TIMES } n \text{ (FACT (PRED } n)))$$

# Recursive Functions

How would we write recursive functions, like factorial?

We'd like to write it like this...

$$\text{FACT} \triangleq \lambda n. \text{IF } (\text{ISZERO } n) \text{ 1 } (\text{TIMES } n \text{ (FACT (PRED } n)))$$

In slightly more readable notation this is...

$$\text{FACT} \triangleq \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times \text{FACT } (n - 1)$$

...but this is an equation, not a definition!

# Recursion removal trick

---

We can perform a “trick” to define a function FACT that satisfies the recursive equation on the previous slide.



# Recursion removal trick

We can perform a “trick” to define a function FACT that satisfies the recursive equation on the previous slide.

Define a new function FACT' that takes a function  $f$  as an argument. Then, for “recursive” calls, it uses  $f f$ :

$$\text{FACT}' \triangleq \lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))$$

# Recursion removal trick

We can perform a “trick” to define a function FACT that satisfies the recursive equation on the previous slide.

Define a new function FACT' that takes a function  $f$  as an argument. Then, for “recursive” calls, it uses  $f f$ :

$$\text{FACT}' \triangleq \lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))$$

Then define FACT as FACT' applied to itself:

$$\text{FACT} \triangleq \text{FACT}' \text{ FACT}'$$

# Example

---

Let's try evaluating FACT on 3...

FACT 3

# Example

---

Let's try evaluating FACT on 3...

$$\text{FACT } 3 = (\text{FACT}' \text{ FACT}') 3$$

# Example

---

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\ &= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((ff) (n - 1)))) \text{FACT}') 3\end{aligned}$$

# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\ &= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))) \text{ FACT}') 3 \\ &\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3\end{aligned}$$

# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\ &= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((ff) (n - 1)))) \text{FACT}' 3 \\ &\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3 \\ &\rightarrow \text{if } 3 = 0 \text{ then } 1 \text{ else } 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1))\end{aligned}$$

# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\&= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))) \text{ FACT}') 3 \\&\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3 \\&\rightarrow \text{if } 3 = 0 \text{ then } 1 \text{ else } 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&\rightarrow 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1))\end{aligned}$$



# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\&= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))) \text{ FACT}') 3 \\&\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3 \\&\rightarrow \text{if } 3 = 0 \text{ then } 1 \text{ else } 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&\rightarrow 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&= 3 \times (\text{FACT } (3 - 1))\end{aligned}$$

# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\&= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((f f) (n - 1))) \text{FACT}') 3 \\&\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3 \\&\rightarrow \text{if } 3 = 0 \text{ then } 1 \text{ else } 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&\rightarrow 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&= 3 \times (\text{FACT } (3 - 1)) \\&\rightarrow \dots \\&\rightarrow 3 \times 2 \times 1 \times 1\end{aligned}$$

# Example

Let's try evaluating FACT on 3...

$$\begin{aligned}\text{FACT } 3 &= (\text{FACT}' \text{ FACT}') 3 \\&= ((\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((ff) (n - 1)))) \text{FACT}' 3 \\&\rightarrow (\lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\text{FACT}' \text{ FACT}') (n - 1))) 3 \\&\rightarrow \text{if } 3 = 0 \text{ then } 1 \text{ else } 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&\rightarrow 3 \times ((\text{FACT}' \text{ FACT}') (3 - 1)) \\&= 3 \times (\text{FACT } (3 - 1)) \\&\rightarrow \dots \\&\rightarrow 3 \times 2 \times 1 \times 1 \\&\rightarrow^* 6\end{aligned}$$

# Fixed point combinators

---

Our “trick” requires following human-readable instructions.  
Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

# Fixed point combinators

---

Our “trick” requires following human-readable instructions.  
Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

There is another way: fixed points!

# Fixed point combinators

Our “trick” requires following human-readable instructions.  
Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

There is another way: fixed points!

Consider factorial again. It is a fixed point of the following:

$$G \triangleq \lambda f. \lambda n. \textbf{if } n = 0 \textbf{ then } 1 \textbf{ else } n \times (f(n - 1))$$

# Fixed point combinators

Our “trick” requires following human-readable instructions. Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

There is another way: fixed points!

Consider factorial again. It is a fixed point of the following:

$$G \triangleq \lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))$$

Recall that if  $g$  is a fixed point of  $G$ , then  $G g = g$ . To see that any fixed point  $g$  is a real factorial function, try evaluating it:

$$g \ 5 = (G g) \ 5$$

# Fixed point combinators

Our “trick” requires following human-readable instructions.  
Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

There is another way: fixed points!

Consider factorial again. It is a fixed point of the following:

$$G \triangleq \lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))$$

Recall that if  $g$  is a fixed point of  $G$ , then  $G g = g$ . To see that any fixed point  $g$  is a real factorial function, try evaluating it:

$$\begin{aligned} g\ 5 &= (G\ g)\ 5 \\ &\rightarrow^* 5 \times (g\ 4) \end{aligned}$$



# Fixed point combinators

Our “trick” requires following human-readable instructions. Write a different function  $f'$  that takes itself as an argument and uses self-application for recursive calls, and then define  $f$  as  $f' f'$ .

There is another way: fixed points!

Consider factorial again. It is a fixed point of the following:

$$G \triangleq \lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))$$

Recall that if  $g$  is a fixed point of  $G$ , then  $G g = g$ . To see that any fixed point  $g$  is a real factorial function, try evaluating it:

$$\begin{aligned} g\ 5 &= (G\ g)\ 5 \\ &\rightarrow^* 5 \times (g\ 4) \\ &= 5 \times ((G\ g)\ 4) \end{aligned}$$

# Fixed point combinators

---

How can we generate the fixed point of  $G$ ?

In denotational semantics, finding fixed points took a lot of math. In the  $\lambda$ -calculus, we just need a suitable combinator...

# Y Combinator

---

The (infamous) Y combinator is defined as

$$Y \triangleq \lambda f. (\lambda x. f (x x)) (\lambda x. f (x x))$$

We say that Y is a *fixed point combinator* because Y  $f$  is a fixed point of  $f$  (for any  $\lambda$ -term  $f$ ).

# Y Combinator

The (infamous) Y combinator is defined as

$$Y \triangleq \lambda f. (\lambda x. f (x x)) (\lambda x. f (x x))$$

We say that Y is a *fixed point combinator* because Y  $f$  is a fixed point of  $f$  (for any  $\lambda$ -term  $f$ ).

What happens when we evaluate Y  $G$  under CBV?

# Z Combinator

---

To avoid this issue, we'll use a slight variant of the Y combinator, called Z, which is easier to use under CBV.

# Z Combinator

---

To avoid this issue, we'll use a slight variant of the Y combinator, called Z, which is easier to use under CBV.

$$Z \triangleq \lambda f. (\lambda x. f (\lambda y. x x y)) (\lambda x. f (\lambda y. x x y))$$

# Example

---

Let's see  $Z$  in action, on our function  $G$ .

FACT

# Example

---

Let's see  $Z$  in action, on our function  $G$ .

$$\begin{aligned} & \text{FACT} \\ = & Z G \end{aligned}$$



# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f(\lambda y. x\ x\ y))) (\lambda x. f(\lambda y. x\ x\ y)))\ G \end{aligned}$$

# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f (\lambda y. x\ x\ y))) (\lambda x. f (\lambda y. x\ x\ y)))\ G \\ \rightarrow & (\lambda x. G (\lambda y. x\ x\ y)) (\lambda x. G (\lambda y. x\ x\ y)) \end{aligned}$$

# Example

Let's see Z in action, on our function G.

FACT

= Z G

=  $(\lambda f. (\lambda x. f (\lambda y. x x y)) (\lambda x. f (\lambda y. x x y))) G$

→  $(\lambda x. G (\lambda y. x x y)) (\lambda x. G (\lambda y. x x y))$

→  $G (\lambda y. (\lambda x. G (\lambda y. x x y)) (\lambda x. G (\lambda y. x x y))) y$

# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f(\lambda y. x\ x\ y)) (\lambda x. f(\lambda y. x\ x\ y)))\ G \\ \rightarrow & (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y)) \\ \rightarrow & G(\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ = & (\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))) \\ & (\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \end{aligned}$$

# Example

Let's see Z in action, on our function G.

```
FACT
= Z G
= (λf. (λx. f (λy. x x y)) (λx. f (λy. x x y))) G
→ (λx. G (λy. x x y)) (λx. G (λy. x x y))
→ G (λy. (λx. G (λy. x x y)) (λx. G (λy. x x y)) y)
= (λf. λn. if n = 0 then 1 else n × (f (n - 1)))
    (λy. (λx. G (λy. x x y)) (λx. G (λy. x x y)) y)
→ λn. if n = 0 then 1
    else n × ((λy. (λx. G (λy. x x y)) (λx. G (λy. x x y)) y) (n - 1))
```

# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f(\lambda y. x\ x\ y)) (\lambda x. f(\lambda y. x\ x\ y)))\ G \\ \rightarrow & (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y)) \\ \rightarrow & G(\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ = & (\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))) \\ & (\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ \rightarrow & \lambda n. \text{if } n = 0 \text{ then } 1 \\ & \text{else } n \times ((\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) (n - 1)) \\ =_{\beta} & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (\lambda y. (Z\ G)\ y) (n - 1) \end{aligned}$$

# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f(\lambda y. x\ x\ y)) (\lambda x. f(\lambda y. x\ x\ y)))\ G \\ \rightarrow & (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y)) \\ \rightarrow & G(\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ = & (\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))) \\ & (\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ \rightarrow & \lambda n. \text{if } n = 0 \text{ then } 1 \\ & \text{else } n \times ((\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) (n - 1)) \\ =_{\beta} & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (\lambda y. (Z\ G)\ y) (n - 1) \\ =_{\beta} & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((Z\ G)\ (n - 1)) \end{aligned}$$

# Example

Let's see Z in action, on our function G.

$$\begin{aligned} & \text{FACT} \\ = & Z\ G \\ = & (\lambda f. (\lambda x. f(\lambda y. x\ x\ y)) (\lambda x. f(\lambda y. x\ x\ y)))\ G \\ \rightarrow & (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y)) \\ \rightarrow & G(\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ = & (\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))) \\ & (\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) \\ \rightarrow & \lambda n. \text{if } n = 0 \text{ then } 1 \\ & \text{else } n \times ((\lambda y. (\lambda x. G(\lambda y. x\ x\ y)) (\lambda x. G(\lambda y. x\ x\ y))\ y) (n - 1)) \\ =_{\beta} & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (\lambda y. (Z\ G)\ y) (n - 1) \\ =_{\beta} & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((Z\ G)\ (n - 1)) \\ = & \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (\text{FACT}\ (n - 1)) \end{aligned}$$



## Other fixed point combinators

There are many (indeed infinitely many) fixed-point combinators. Here's a cute one:

$$Y_k \triangleq (\text{LLLLLLLLLLLLLLLLLLLLLLLLLLLLLLLL})$$

where

$$L \triangleq \lambda abcdefghijklmnopqrstuvwxyzr. \\ (r(\textit{thisisafixedpointcombinator}))$$

# Turing's Fixed Point Combinator

---

To gain some more intuition for fixed point combinators, let's derive a combinator  $\Theta$  originally discovered by Turing.

# Turing's Fixed Point Combinator

---

To gain some more intuition for fixed point combinators, let's derive a combinator  $\Theta$  originally discovered by Turing.

We know that  $\Theta f$  is a fixed point of  $f$ , so we have

$$\Theta f = f(\Theta f).$$

# Turing's Fixed Point Combinator

---

To gain some more intuition for fixed point combinators, let's derive a combinator  $\Theta$  originally discovered by Turing.

We know that  $\Theta f$  is a fixed point of  $f$ , so we have

$$\Theta f = f(\Theta f).$$

We can write the following recursive equation:

$$\Theta = \lambda f. f(\Theta f)$$

# Turing's Fixed Point Combinator

To gain some more intuition for fixed point combinators, let's derive a combinator  $\Theta$  originally discovered by Turing.

We know that  $\Theta f$  is a fixed point of  $f$ , so we have

$$\Theta f = f(\Theta f).$$

We can write the following recursive equation:

$$\Theta = \lambda f. f(\Theta f)$$

Now use the recursion removal trick:

$$\begin{aligned}\Theta' &\triangleq \lambda t. \lambda f. f(t t f) \\ \Theta &\triangleq \Theta' \Theta'\end{aligned}$$

# $\theta$ Example

---

$$\text{FACT} = \Theta G$$

## $\theta$ Example

---

$$\begin{aligned}\text{FACT} &= \Theta G \\ &= ((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f))) G\end{aligned}$$

## $\theta$ Example

$$\begin{aligned}\text{FACT} &= \Theta G \\ &= ((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f))) G \\ &\rightarrow (\lambda f. f((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) f)) G\end{aligned}$$



## $\theta$ Example

$$\begin{aligned}\text{FACT} &= \Theta G \\ &= ((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f))) G \\ &\rightarrow (\lambda f. f((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) f)) G \\ &\rightarrow G((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) G)\end{aligned}$$

## $\theta$ Example

$$\begin{aligned}\text{FACT} &= \Theta G \\ &= ((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f))) G \\ &\rightarrow (\lambda f. f((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) f)) G \\ &\rightarrow G((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) G) \\ &= G(\Theta G)\end{aligned}$$

## $\theta$ Example

$$\begin{aligned}\text{FACT} &= \Theta G \\&= ((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f))) G \\&\rightarrow (\lambda f. f((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) f)) G \\&\rightarrow G((\lambda t. \lambda f. f(t t f)) (\lambda t. \lambda f. f(t t f)) G) \\&= G(\Theta G) \\&= (\lambda f. \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (f(n - 1))) (\Theta G) \\&\rightarrow \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times ((\Theta G) (n - 1)) \\&= \lambda n. \text{if } n = 0 \text{ then } 1 \text{ else } n \times (\text{FACT } (n - 1))\end{aligned}$$