

1vr 20 feb 2023 10:29Quantum Mechanics I

A Modern Physics Review

Clark Saben

February 20, 2023

1.1 Representations of Spin Half

(a) Write \hat{S}_z in the S_x basis, both in Dirac Notation and as a matrix. Hint: However you do this, you will need to know how \hat{S}_z acts on the eigenstates of S_x .

Find \hat{S}_z in the S_x basis:

$$\hat{1}_x \hat{S}_z \hat{1}_x = (|+x\rangle\langle+x| + |-x\rangle\langle-x|) \hat{S}_z (|+x\rangle\langle+x| + |-x\rangle\langle-x|)$$

Need \hat{S}_z (hint):

$$\begin{aligned}\hat{1}_z |\pm x\rangle &= | +z\rangle\langle+z|\pm x\rangle + | -z\rangle\langle-z|\pm x\rangle \\ &= \frac{1}{\sqrt{2}} (| +z\rangle|\pm| -z\rangle) \\ \implies \hat{S}_z |\pm x\rangle &= \frac{\hbar}{2\sqrt{2}} (\langle+z|\pm \langle-z|)\end{aligned}$$

So,

$$\begin{aligned}\hat{1}_x \hat{S}_z \hat{1}_x &= (|+x\rangle\langle+x| + |-x\rangle\langle-x|) \left(\frac{\hbar}{2} |-x\rangle\langle+x| + \frac{\hbar}{2} |+x\rangle\langle-x| \right) \\ &= \frac{\hbar}{2} (|+x\rangle\langle-x| + |-x\rangle\langle+x|) \\ \implies \mathbb{S}_z^{(S_x)} &= \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\end{aligned}$$

(b) Verify the equations (in the S_x basis),
using your matrix representation for \hat{S}_z in the S_x basis.

$$\hat{S}_z|\pm z\rangle = \pm \frac{\hbar}{2}|\pm z\rangle$$

$$\begin{aligned}\mathbb{S}_z^{(S_x)}\vec{v}_{+z}^{(S_x)} &= \frac{\hbar}{2\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ &= \frac{\hbar}{2\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ &= \frac{\hbar}{2} \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} \\ &= \frac{\hbar}{2} \vec{v}_{+z}^{(S_x)}\end{aligned}$$

$$\hat{1}_x|-z\rangle = |+x\rangle\langle+x|-z\rangle + |-x\rangle\langle-x|-z\rangle$$

where

$$\langle+x|-z\rangle = \frac{1}{\sqrt{2}}$$

and

$$\langle-x|-z\rangle = \frac{-1}{\sqrt{2}}$$

$$\begin{aligned}\Rightarrow \mathbb{S}_z^{(S_x)}\vec{v}_{-z}^{(S_x)} &= \frac{\hbar}{2\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ &= \frac{\hbar}{2\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix} \\ &= \frac{-\hbar}{2} \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} \end{pmatrix} \\ &= \frac{-\hbar}{2} \vec{v}_{-z}^{(S_x)}\end{aligned}$$

(c) Write the projection operator \hat{P}_{+z} in the S_x basis, both in Dirac Notation and as a matrix.

Find $\hat{P}_{+z}^{(S_x)}$ and \hat{P}_{+z} in the S_x basis:

$$\begin{aligned}\hat{P}_{+z}^{(S_x)} &= \hat{1}_x \hat{P}_{+z} \hat{1}_x \\ &= (|+x\rangle\langle+x| + |-x\rangle\langle-x|) \left(\frac{1}{\sqrt{2}}|+z\rangle\langle+x| + \frac{1}{\sqrt{2}}|+z\rangle\langle-x| \right) \\ &= \frac{1}{2} (|+x\rangle\langle+x| + |+x\rangle\langle-x| + |-x\rangle\langle+x| + |-x\rangle\langle-x|)\end{aligned}$$

$$\mathbb{P}_{+z}^{(S_x)} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

(d) Find eigenvalues and eigenvectors of your matrix representation of \hat{P}_{+z} in the S_x basis, thus verifying that $\hat{P}_{+z}|+z\rangle = |+z\rangle$.

Diagonalize $\mathbb{P}_{+z}^{(S_x)}$ and verify $\mathbb{P}_{+z}^{S_x} \vec{v}_{+z}^{S_x} = \vec{v}_{+z}^{S_x}$.

$$\begin{aligned}\lambda_1 \lambda_2 &= 0 ; \lambda_1 + \lambda_2 = 1 \\ \implies \lambda_1 &= 1 \text{ and } \lambda_2 = 0\end{aligned}$$

$$\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$$

(e) Repeat parts (c) and (d) for \hat{P}_{-z} .