
P1239 JNJ

99526687SCC1001 · Ac-225 DLL3 antibody + Zr-89 PET · Phase 1

Janssen · v1.0 08 June 2026 · 19-page elements document



JNJ-99526687 Protocol Elements Document 99526687SCC1001

PROTOCOL ELEMENTS DOCUMENT

TITLE PAGE

Sponsor Confidentiality Statement:	The information provided herein is considered trade secrets, commercial or financial, or privileged or confidential information that we customarily hold close and treat as confidential. The information is being provided under the assurance that the recipient will maintain the confidentiality of the information under the applicable statutes, regulations, rules, case law, contractual provisions, protective orders, or as otherwise necessary to protect the information from disclosure. The recipient agrees and acknowledges that release of information considered trade secrets, commercial or financial, or privileged or confidential will result in competitive harm and, as such, the information provided herein is exempt from disclosure under the Freedom of Information Act ("FOIA"), or other similar law or regulation. Particularly, the information shall not be utilized in any AI solutions, including GenAI, without Company's prior written approval.
Full Title:	**A Phase 1 Study of JNJ-99526687, an Actinium-225-Labeled Antibody Targeting Delta-Like Ligand 3 (DLL3), for Small Cell Lung Cancer**
Sponsor Protocol Identifier:	99526687SCC1001
EDMS number; Version Number:	EDMS-RIM-1986262; Version 1.0
Version Date:	08 June 2026
Sponsor's Investigational Product Codes:	JNJ-99526687 JNJ-99526726
Trial Phase:	Phase 1
Sponsor Name and Address:	Janssen Research & Development LLC, 920 Route 202 South, Raritan, 08869 New Jersey, USA

What this study is, in one breath

Give a zirconium-89 DLL3 antibody PET so you can see the tumor, then 7-10 days later give the same antibody labeled with actinium-225 to treat recurrent extensive-stage small-cell lung cancer.

Alpha-antibody therapy plus companion PET. Not tarlatamab. Not PSMA. Not FDG.

> **If the protocol feels like a wall of words**
>
> A protocol is a recipe: who can join, what is injected, how much, when you scan or treat, and how you decide if a picture or outcome was right. This sheet is the same idea in normal English. It does not replace the signed site copy.

1. Two products, one DLL3 antibody

Janssen 99526687SCC1001 is a first-in-human Phase 1 protocol-elements document (v1.0, 08 June 2026) for extensive-stage small-cell lung cancer. Same ~100 kDa bivalent spFv-Fc targeting DLL3, two labels. Not tarlatamab (IMDELLTRA), not a T-cell engager, not PSMA, not FDG.

IMP	Role	Starting numbers
JNJ-99526726	Companion PET · Zr-89 · DFO chelator	2 mCi IV. Mass starts 3 mg, SET may go to 20 mg. Serial PET 0-4 h, 24+/-6 h, 42-78 h, 120-216 h.
JNJ-99526687	Therapy · Ac-225 · macropa chelator JNJ-95981860	Start 0.75 uCi/kg on 3 mg, IV 7-10 days after the imaging dose. BOIN on uCi.

Both products mix labeled plus unlabeled spFv-Fc to hit the assigned mass. If blood-pool uptake is much greater than tumor, SET may repeat Zr-89 PET at a revised mass before allowing therapy.

2. Who

Adults with histologically confirmed recurrent ES-SCLC after platinum-based chemotherapy and a checkpoint inhibitor. ECOG 0-1. About 100 people: ~60 image/mass optimization plus ~40 dose expansion. Additional DLL3 tumor types may be added later.

- Part 1: measurable or evaluable disease. Part 2: at least one RECIST 1.1 lesion.
- eGFR >= 50 mL/min (CKD-EPI, adjusted BSA). Hb >= 9, ANC >= 1.5 (or 1.0 if benign ethnic neutropenia), platelets >= 100.
- Exclude ILD/pneumonitis in the prior 12 months, NYHA III/IV, QTcF >470 ms, untreated active CNS disease.

3. Design

Open-label BOIN. n=8-12 at 3 mg / 2 mCi Zr-89 until imaging mass is optimized, then radioactivity is escalated (schema: 50 then 100 then X uCi Ac-225). Stagger >= 1 week between the first patient on a new dose and the next. SET (investigators + sponsor, chaired by the sponsor Study Responsible Physician) owns mass and uCi decisions. No randomisation in this elements document.

- CT C/A/P/brain (or MRI) every 6 +/- 1 weeks for 24 weeks, then every 12 +/- 1 weeks. RECIST 1.1.
- Ac-225 SPECT after therapy for selected patients: post-treatment dosimetry vs EBRT-equivalent organ thresholds.
- Subsequent Ac-225 doses no sooner than 8 weeks, and only if Table 4 retreatment gates are met 7+/-1 days before.

4. DLT window is 42 days after first Ac-225

Bucket	Counts as DLT (related)
Heme	G4 ANC >5 days or febrile neutropenia; G3 TCP with bleed or G4 TCP; any Grade 5.
Lung	Grade >=2 pneumonitis / ILD.
Other	>= G3 non-heme with listed exceptions (short-lived fatigue/GI, isolated ALP/GGT, Hy's law still counts).

Alpha antibody, not tarlatamab

DLL3 is the same target as IMDELLTRA, but this construct carries actinium-225. Pulmonary DLT starts at Grade 2. Do not copy VECTRA 12 MBq PSMA numbers or BET-121 lutetium cycles.

What to know cold

Q. Target?
A. DLL3 on a ~100 kDa bivalent spFv-Fc. Not a T-cell engager.

Q. Imaging?
A. JNJ-99526726, Zr-89 DFO, 2 mCi IV. Mass 3 mg start, may go to 20 mg.

Q. Therapy?
A. JNJ-99526687, Ac-225 macropa. Start 0.75 uCi/kg on 3 mg, 7-10 d after PET.

Q. Who / n?
A. Recurrent ES-SCLC after platinum + ICI. ECOG 0-1. ~100 (60 + 40).

Q. DLT lung?
A. Grade >=2 pneumonitis/ILD. 42-day window after first Ac-225.

How to use this with Board Studio

Open the matching protocol module and press play. The left pane is the signed page. Use this PDF when legalese stalls you. If anything conflicts, the signed protocol wins.

Disclaimer. Educational summary for board-style study. Not medical advice, not a recruitment brochure, not a substitute for the signed protocol, and not affiliated with the sponsor, ABR, ABNM, ACGME, SNMMI, NRC, or FDA. Confirm numbers against the current site copy before any clinical use.