lab14

Catherina Santoso (PID: A18552334)

Background

The authors report on differential analysis of lung fibroblasts in response to loss of the developmental transcription factor HOXA1. Their results and others indicate that HOXA1 is required for lung fibroblast and HeLa cell cycle progression. In particular their analysis show that "loss of HOXA1 results in significant expression level changes in thousands of individual transcripts, along with isoform switching events in key regulators of the cell cycle". For our session we have used their Sailfish gene-level estimated counts and hence are restricted to protein-coding genes only.

Data Import

We have 2 key input files: counts and metadata

```
library(DESeq2)
```

```
Warning: package 'DESeq2' was built under R version 4.5.2

Warning: package 'S4Vectors' was built under R version 4.5.2

Warning: package 'IRanges' was built under R version 4.5.2

Warning: package 'GenomicRanges' was built under R version 4.5.2

Warning: package 'Biobase' was built under R version 4.5.2
```

```r
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"

# Import metadata and take a peek
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```
              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```r
# Import countdata
countData = read.csv(countFile, row.names=1)
head(countData)
```

```
                length SRR493366 SRR493367 SRR493368 SRR493369 SRR493370
ENSG00000186092    918         0         0         0         0         0
ENSG00000279928    718         0         0         0         0         0
ENSG00000279457   1982        23        28        29        29        28
ENSG00000278566    939         0         0         0         0         0
ENSG00000273547    939         0         0         0         0         0
ENSG00000187634   3214       124       123       205       207       212
                SRR493371
ENSG00000186092         0
ENSG00000279928         0
ENSG00000279457        46
ENSG00000278566         0
ENSG00000273547         0
ENSG00000187634       258
```

> Q1. Complete the code below to remove the troublesome first column from countData

We need to remove

```r
# Note we need to remove the odd first $length col
countData <- as.matrix(countData[,-1])
head(countData)
```

```
                 SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000186092         0         0         0         0         0         0
ENSG00000279928         0         0         0         0         0         0
ENSG00000279457        23        28        29        29        28        46
ENSG00000278566         0         0         0         0         0         0
ENSG00000273547         0         0         0         0         0         0
ENSG00000187634       124       123       205       207       212       258
```

```
rownames(colData) == colnames(countData)
```

```
[1] TRUE TRUE TRUE TRUE TRUE TRUE
```

> Q2.Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns).

Tip: What will rowSums() of countData return and how could you use it in this context?

```
# Filter count data where you have 0 read count across all samples.
countData = countData[rowSums(countData) > 0, ]
head(countData)
```

```
                 SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000279457        23        28        29        29        28        46
ENSG00000187634       124       123       205       207       212       258
ENSG00000188976      1637      1831      2383      1226      1326      1504
ENSG00000187961       120       153       180       236       255       357
ENSG00000187583        24        48        65        44        48        64
ENSG00000187642         4         9        16        14        16        16
```

Check and Tidy

```
colData
```

```
              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

Setup for DESeq

Run DESeq

```
dds = DESeqDataSetFromMatrix(countData = countData,
                        colData = colData,
                        design = ~condition)
```

```
Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in
design formula are characters, converting to factors
```

```
dds = DESeq(dds)
```

```
estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing
```

```
dds
```

```
class: DESeqDataSet
dim: 15975 6
metadata(1): version
assays(4): counts mu H cooks
rownames(15975): ENSG00000279457 ENSG00000187634 ... ENSG00000276345
  ENSG00000271254
rowData names(22): baseMean baseVar ... deviance maxCooks
colnames(6): SRR493366 SRR493367 ... SRR493370 SRR493371
colData names(2): condition sizeFactor
```

```r
res = results(dds)
res
```

```
log2 fold change (MLE): condition hoxa1 kd vs control sirna
Wald test p-value: condition hoxa1 kd vs control sirna
DataFrame with 15975 rows and 6 columns
                 baseMean log2FoldChange     lfcSE       stat      pvalue
                <numeric>      <numeric> <numeric>  <numeric>   <numeric>
ENSG00000279457   29.9136      0.1792571 0.3248216   0.551863 5.81042e-01
ENSG00000187634  183.2296      0.4264571 0.1402658   3.040350 2.36304e-03
ENSG00000188976 1651.1881     -0.6927205 0.0548465 -12.630158 1.43990e-36
ENSG00000187961  209.6379      0.7297556 0.1318599   5.534326 3.12428e-08
ENSG00000187583   47.2551      0.0405765 0.2718928   0.149237 8.81366e-01
...                   ...            ...       ...        ...         ...
ENSG00000273748  35.30265       0.674387  0.303666   2.220817 2.63633e-02
ENSG00000278817   2.42302      -0.388988  1.130394  -0.344117 7.30758e-01
ENSG00000278384   1.10180       0.332991  1.660261   0.200565 8.41039e-01
ENSG00000276345  73.64496      -0.356181  0.207716  -1.714752 8.63908e-02
ENSG00000271254 181.59590      -0.609667  0.141320  -4.314071 1.60276e-05
                      padj
                 <numeric>
ENSG00000279457 6.86555e-01
ENSG00000187634 5.15718e-03
ENSG00000188976 1.76549e-35
ENSG00000187961 1.13413e-07
ENSG00000187583 9.19031e-01
...                     ...
ENSG00000273748 4.79091e-02
ENSG00000278817 8.09772e-01
ENSG00000278384 8.92654e-01
ENSG00000276345 1.39762e-01
ENSG00000271254 4.53648e-05
```

Get results >Q. Call the summary() function on your results to get a sense of how many genes are up or down-regulated at the default 0.1 p-value cutoff.

```r
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
```

```
LFC > 0 (up)       : 4349, 27%
LFC < 0 (down)     : 4396, 28%
outliers [1]       : 0, 0%
low counts [2]     : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```

Volcano plot

```r
library(ggplot2)

ggplot(res) +
  aes(x = log2FoldChange,
      y = -log(padj)) +
  geom_point()
```

```
Warning: Removed 1237 rows containing missing values or values outside the scale range
(`geom_point()`).
```



Q. Improve this plot by completing the below code, which adds color, axis labels and cutoff lines:

```r
# Make a color vector for all genes
mycols <- rep("gray", nrow(res) )

# Color blue the genes with fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "blue"

# Color gray those with adjusted p-value more than 0.01
mycols[ res$padj > 0.01 ] <- "gray"

ggplot(res) +
  aes(x = log2FoldChange,
      y = -log10(padj)) +
  geom_point(color = mycols) +
  xlab("Log2(FoldChange)") +
  ylab("-Log(P-value)") +
  geom_vline(xintercept = c(-2,2)) +
  geom_hline(yintercept = -log10(0.01))
```

```
Warning: Removed 1237 rows containing missing values or values outside the scale range
(`geom_point()`).
```



Q. Use the mapIDs() function multiple times to add SYMBOL, ENTREZID and GENENAME annotation to our results by completing the code below.

Since we mapped and counted against the Ensembl annotation, our results only have information about Ensembl gene IDs. However, our pathway analysis downstream will use KEGG pathways, and genes in KEGG pathways are annotated with Entrez gene IDs. So lets add them as we did the last day.

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

```
 [1] "ACCNUM"       "ALIAS"        "ENSEMBL"      "ENSEMBLPROT"  "ENSEMBLTRANS"
 [6] "ENTREZID"     "ENZYME"       "EVIDENCE"     "EVIDENCEALL"  "GENENAME"
[11] "GENETYPE"     "GO"           "GOALL"        "IPI"          "MAP"
[16] "OMIM"         "ONTOLOGY"     "ONTOLOGYALL"  "PATH"         "PFAM"
[21] "PMID"         "PROSITE"      "REFSEQ"       "SYMBOL"       "UCSCKG"
[26] "UNIPROT"
```

```
res$symbol = mapIds(org.Hs.eg.db,
                    keys= row.names(res),
                    keytype="ENSEMBL",
                    column="SYMBOL",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```
res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```r
res$name =   mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="GENENAME",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```r
head(res, 10)
```

```
log2 fold change (MLE): condition hoxa1 kd vs control sirna
Wald test p-value: condition hoxa1 kd vs control sirna
DataFrame with 10 rows and 9 columns
                   baseMean log2FoldChange     lfcSE       stat      pvalue
                  <numeric>      <numeric> <numeric>  <numeric>   <numeric>
ENSG00000279457   29.913579      0.1792571 0.3248216   0.551863 5.81042e-01
ENSG00000187634  183.229650      0.4264571 0.1402658   3.040350 2.36304e-03
ENSG00000188976 1651.188076     -0.6927205 0.0548465 -12.630158 1.43990e-36
ENSG00000187961  209.637938      0.7297556 0.1318599   5.534326 3.12428e-08
ENSG00000187583   47.255123      0.0405765 0.2718928   0.149237 8.81366e-01
ENSG00000187642   11.979750      0.5428105 0.5215598   1.040744 2.97994e-01
ENSG00000188290  108.922128      2.0570638 0.1969053  10.446970 1.51282e-25
ENSG00000187608  350.716868      0.2573837 0.1027266   2.505522 1.22271e-02
ENSG00000188157 9128.439422      0.3899088 0.0467163   8.346304 7.04321e-17
ENSG00000237330    0.158192      0.7859552 4.0804729   0.192614 8.47261e-01
                      padj      symbol      entrez                   name
                  <numeric> <character> <character>            <character>
ENSG00000279457 6.86555e-01          NA          NA                     NA
ENSG00000187634 5.15718e-03      SAMD11      148398 sterile alpha motif ..
ENSG00000188976 1.76549e-35       NOC2L       26155 NOC2 like nucleolar ..
ENSG00000187961 1.13413e-07      KLHL17      339451 kelch like family me..
ENSG00000187583 9.19031e-01     PLEKHN1       84069 pleckstrin homology ..
ENSG00000187642 4.03379e-01       PERM1       84808 PPARGC1 and ESRR ind..
ENSG00000188290 1.30538e-24        HES4       57801 hes family bHLH tran..
ENSG00000187608 2.37452e-02       ISG15        9636 ISG15 ubiquitin like..
ENSG00000188157 4.21963e-16        AGRN      375790                   agrin
ENSG00000237330          NA      RNF223      401934 ring finger protein ..
```

Q. Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

```r
res = res[order(res$pvalue),]
write.csv(res, file="deseq_results.csv")
```

Section 2 Pathway Analysis

```r
library(pathview)
```

```
##############################################################################
Pathview is an open source software package distributed under GNU General
Public License version 3 (GPLv3). Details of GPLv3 is available at
http://www.gnu.org/licenses/gpl-3.0.html. Particullary, users are required to
formally cite the original Pathview paper (not just mention it) in publications
or products. For details, do citation("pathview") within R.

The pathview downloads and uses KEGG data. Non-academic uses may require a KEGG
license agreement (details at http://www.kegg.jp/kegg/legal.html).
##############################################################################
```

```r
library(gage)
```

```r
library(gageData)

data(kegg.sets.hs)
data(sigmet.idx.hs)

# Focus on signaling and metabolic pathways only
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

# Examine the first 3 pathways
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
[1] "10"   "1544" "1548" "1549" "1553" "7498" "9"

$`hsa00983 Drug metabolism - other enzymes`
 [1] "10"     "1066"   "10720"  "10941"  "151531" "1548"   "1549"   "1551"
 [9] "1553"   "1576"   "1577"   "1806"   "1807"   "1890"   "221223" "2990"
```

```
[17] "3251"   "3614"   "3615"   "3704"   "51733"  "54490"  "54575"  "54576"
[25] "54577"  "54578"  "54579"  "54600"  "54657"  "54658"  "54659"  "54963"
[33] "574537" "64816"  "7083"   "7084"   "7172"   "7363"   "7364"   "7365"
[41] "7366"   "7367"   "7371"   "7372"   "7378"   "7498"   "79799"  "83549"
[49] "8824"   "8833"   "9"      "978"

$`hsa00230 Purine metabolism`
  [1] "100"    "10201"  "10606"  "10621"  "10622"  "10623"  "107"    "10714"
  [9] "108"    "10846"  "109"    "111"    "11128"  "11164"  "112"    "113"
 [17] "114"    "115"    "122481" "122622" "124583" "132"    "158"    "159"
 [25] "1633"   "171568" "1716"   "196883" "203"    "204"    "205"    "221823"
 [33] "2272"   "22978"  "23649"  "246721" "25885"  "2618"   "26289"  "270"
 [41] "271"    "27115"  "272"    "2766"   "2977"   "2982"   "2983"   "2984"
 [49] "2986"   "2987"   "29922"  "3000"   "30833"  "30834"  "318"    "3251"
 [57] "353"    "3614"   "3615"   "3704"   "377841" "471"    "4830"   "4831"
 [65] "4832"   "4833"   "4860"   "4881"   "4882"   "4907"   "50484"  "50940"
 [73] "51082"  "51251"  "51292"  "5136"   "5137"   "5138"   "5139"   "5140"
 [81] "5141"   "5142"   "5143"   "5144"   "5145"   "5146"   "5147"   "5148"
 [89] "5149"   "5150"   "5151"   "5152"   "5153"   "5158"   "5167"   "5169"
 [97] "51728"  "5198"   "5236"   "5313"   "5315"   "53343"  "54107"  "5422"
[105] "5424"   "5425"   "5426"   "5427"   "5430"   "5431"   "5432"   "5433"
[113] "5434"   "5435"   "5436"   "5437"   "5438"   "5439"   "5440"   "5441"
[121] "5471"   "548644" "55276"  "5557"   "5558"   "55703"  "55811"  "55821"
[129] "5631"   "5634"   "56655"  "56953"  "56985"  "57804"  "58497"  "6240"
[137] "6241"   "64425"  "646625" "654364" "661"    "7498"   "8382"   "84172"
[145] "84265"  "84284"  "84618"  "8622"   "8654"   "87178"  "8833"   "9060"
[153] "9061"   "93034"  "953"    "9533"   "954"    "955"    "956"    "957"
[161] "9583"   "9615"
```

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
     1266     54855      1465      2034      2150      6659
-2.422719  3.201955 -2.313738 -1.888019  3.344508  2.392288
```

```
# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)
```

```
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

```r
# Look at the first few down (less) pathways
head(keggres$less)
```

```
                                        p.geomean stat.mean        p.val
hsa04110 Cell cycle                   8.995727e-06 -4.378644 8.995727e-06
hsa03030 DNA replication              9.424076e-05 -3.951803 9.424076e-05
hsa03013 RNA transport                1.375901e-03 -3.028500 1.375901e-03
hsa03440 Homologous recombination     3.066756e-03 -2.852899 3.066756e-03
hsa04114 Oocyte meiosis               3.784520e-03 -2.698128 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 8.961413e-03 -2.405398 8.961413e-03
                                            q.val set.size         exp1
hsa04110 Cell cycle                   0.001448312      121 8.995727e-06
hsa03030 DNA replication              0.007586381       36 9.424076e-05
hsa03013 RNA transport                0.073840037      144 1.375901e-03
hsa03440 Homologous recombination     0.121861535       28 3.066756e-03
hsa04114 Oocyte meiosis               0.121861535      102 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 0.212222694       53 8.961413e-03
```

```r
pathview(gene.data=foldchanges, pathway.id="hsa04110.png")
```

```
Info: Downloading xml files for hsa04110.png, 1/1 pathways..

Warning in download.file(xml.url, xml.target, quiet = T): cannot open URL
'https://rest.kegg.jp/get/hsa04110.png/kgml': HTTP status was '400 Bad Request'

Warning: Download of hsa04110.png xml file failed!
This pathway may not exist!

Info: Downloading png files for hsa04110.png, 1/1 pathways..

Warning: Download of hsa04110.png png file failed!
This pathway may not exist!

Warning: Failed to download KEGG xml/png files, hsa04110.png skipped!
```




```
# A different PDF based output of the same data
pathview(gene.data=foldchanges, pathway.id="hsa04110", kegg.native=FALSE)
```

```
'select()' returned 1:1 mapping between keys and columns

Warning: reconcile groups sharing member nodes!

     [,1] [,2]
[1,] "9"  "300"
[2,] "9"  "306"

Info: Working in directory /Users/ninasantoso/Desktop/BIMM143/lab14

Info: Writing image file hsa04110.pathview.pdf
```

```r
## Focus on top 5 upregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$greater)[1:5]

# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```r
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```


HEMATOPOIETIC CELL LINEAGE
Lymphoid Related Dendritic cell
γδ T cell
CD8 T cell
CD4 T cell
Regulatory T cell
NKT cell
Thymus
Pro T cell (DN2)
DN3
DN4
Intermediate single-positive cell (ISP)
Double-positive cell (DP)
NK cell Precursor
NK cell
Lymphoid stem cell, Double-negative cell (DN1)
Pro B Cell
Pre B I cell
Pre B II cell
Immature B cell
B Cell
Hematopoietic stem cell
CFU-Mast
Mast cell
CFU-Bas
Myeloblast
Basophilic Myelocyte
Basophil
CFU-E0
Eosinophilic Myelocyte
Eosinophil
CFU-M/DC
Myeloid Related Dendritic Cell
Monoblast
Promonocyte
Monocyte
Macrophage
Myeloid Stem Cell
CFU-GEMM
CFU-GM
CFU-G
Neutrophilic Myelocyte
Neutrophil
Bone marrow
BFU-E
CFU-E
Proerythroblast
Erythrocyte
BFU-MK
CFU-MK
Mega-karyocyte
Platelets
Data on KEGG graph
Rendered by Pathview


LYSOSOME
Exocytosis
Phagocytosis
bacterium
exosome
Phagosome
phagosome
endolysosome
phagolysosome
MVB
Endocytosis
Endocytosis
clathrin coat
AP-2
clathrin
early endosome
M6P
M6P receptor
late endosome
mannose
mature lysosomal hydrase
multivesicular body (MVB)
cytosol
pH~ 7.2
ATP
ADP
ATPeV
H+
pH~ 5.0
acid hydrolase
lysosomal membrane protein
lysosome
AP-3
BLOC1
VAMP8
SNAP29
Stx17
AP-1
recycling endosome
Receptor recycling
Receptor-dependent transport
clathrin coated vesicle
clathrin
CLN5
CLN3
Retromer
Rab7
PLEKHM
autolysosome
degradation and recycling
Transport of synthesized lysosomal enzymes
AP-1
AP-3
GGAs
AP-4
AP-5
SPG11
SPG15
MCOLN1
M6P receptor
MPR
sortilin
LIMP2
Activation of lysosomal sulfatase precursor
FGE
lysosomal hydrase precursor
mannose
from ER
CLN8
CLN6
LYSET
GNPT
NAGPA
cis Golgi network
Golgi body
trans Golgi network
transport vesicle
Snare interactions in vesicular transport
autophagosome
LMP
membrane repair & proton homeostasis
Autophagy
Regulation of autophagy
mitochondria
phagophore
plasma membrane
-1
0
1
Lysosomal membrane proteins
transporters
ion channels, transporters
MCOLN
LAPTM
NRAMP
CLN7
CLC-7
TPC
TMEM175
lipid
NPC1
ABCA3/5
WS12C
ORPs
sugar
GLUT6
nucleoside
SLC29A3
amino acid
PAT
PHT
CTNS
sialic acid
sialin
peptide
ABCB9
pH~ 7.2
ATP
ADP
ATPeV
H+
pH~ 5.0
acidification regulators
DMXL
ROGDI
WDR7
NCOA7
major lysosomal membrane proteins
LAMP
LIMP
fusion
mkb/Vps
SNAREs
VAMP7
VAMP8
GTPase
Rab7
ARL8
tethering factors
HOPS
BORC
motor adaptors
Lysosomal acid hydrolases
proteases
cathepsins
saposin
LGMN
TPP1
glycosidases
NPC2
GLA
GLB
GAA
GBA
IDUA
NAGA
NAGLU
GALC
GUSB
FUCA1
HEXA/B
MANB
LAMAN
HEU1
HYAL1
sulfatases
ARS
GALNS
GNS
IDS
SGSH
lipases
LIPA
LYPLA2
nuclease
DNase II
phosphatase
ACP2
ACP5
sphingomyelinase
SMPD1
ceramidase
ASAH1
aspartylglucosaminidase
AGA
Other lysosomal enzymes and activators
saposin
GM2A
CLN1
lysosome
signalling complexes
mTORC1
RagA/B
RagC/D
ragulator
mEAK7
transcription factors
MITF
TFEB
TFE3
sensors
SLC38A9
LYCHOS
mTOR signaling pathway
enzymes
FIG4
PIKfyve
LAL/LIP
HEXMA1
Glycosaminoglycan degradation
Other glycan degradation
immunity
TLR2
TLR7/8
TLR3
LITAF
Toll-like receptor signaling pathway
Data on KEGG graph
Rendered by Pathview


-1
0
1
NOTCH SIGNALING PATHWAY
Fringe
Delta
Serrate
Notch
ADAM17
S2
γ-Secretase complex
PSE2
PSEN
NCSTN
APH-1
S3
+u
Deltex
Itch
Numb
Dvl
NICD
(Notch intracellular domain)
Co-activator
MAML
HATs
SKIP
CSL
DNA
Hes1/5
Hey
PreTα
NRARP
Co-repressor
Hairless
SMRT
CtBP
Gro/TLE
CIR
SHARP
HDAC
ATXN1/L
Hes1
NRARP
Ras/MAPK
MAPK signaling pathway
Gene expression
Data on KEGG graph
Rendered by Pathview


STEROID HORMONE BIOSYNTHESIS
-1
0
1
Steroid biosynthesis
Cholesterol
Cholesterol sulfate
20α-Hydroxy-cholesterol
22β-Hydroxy-cholesterol
20α,22β-Dihydroxy-cholesterol
21-Hydroxy-pregnenolone
11-Deoxy-corticosterone
Corticosterone
11-Dehydro-corticosterone
21-Hydroxy-5β-pregnane-3,11,20-trione
Tetrahydro-corticosterone
11β,21-Dihydroxy-5β-pregnane-3,20-dione
3α,21-Dihydroxy-5β-pregnane-11,20-dione
3α,20α,21-Trihydroxy-5β-pregnane-11-one
11β,21-Dihydroxy-3,20-oxo-5β-pregnan-18-al
3α,11β,21-Trihydroxy-20-oxo-5β-pregnan-18-al
18-Hydroxy-corticosterone
Aldosterone
Aldosterone hemiacetal
5α-Dihydro-deoxycorticosterone
Allotetrahydro-deoxycorticosterone
4-Methylpentanal
Pregnenolone
7α-Hydroxy-pregnenolone
20α-Hydroxy-progesterone
Progesterone
11β-Hydroxy-progesterone
Pregnenolone-sulfate
3α-Hydroxy-5β-pregnane-20-one
Pregnanediol
5β-Pregnane-3,20-dione
5α-Pregnan-20α-ol-3-one
5α-Pregnan-3,20-dione
3α-Hydroxy-5α-pregnan-20-one
5α-Pregnane-3α,20α-diol
C21-Sterolds
17α,20α-Dihydroxy-cholesterol
17α-Hydroxy-pregnenolone
17α,20α-Dihydroxy-pregn-4-en-3-one
17α-Hydroxy-progesterone
21-Deoxycortisol
5α-Pregnan-17α-ol-3,20-dione
5α-Pregnane-3α,17α-diol-20-one
11β,17α,21-Trihydroxy-5β-pregnane-3,20-dione
Urocortisol
Cortol
17α,21-Dihydroxy-pregnenolone
11-Deoxycortisol
Cortisol
17α,21-Dihydroxy-5β-pregnane-3,11,20-trione
Cortisone
Urocortisone
Cortolone
11β,17α,21-Trihydroxy-pregnenolone
Dehydro-epiandrosterone
Dehydroepiandro-steron sulfate
7α-Hydroxydehydro-epiandrosterone
16α-Hydroxyandrost-4-ene-3,17-dione
16α-Hydroxydehydro-epiandrosterone
11β-Hydroxyandrost-4-ene-3,17-dione
11β-Hydroxy-testosterone
Adrenosterone
Estrone 3-sulfate
5β-Androstane-3,17-dione
Etiocholan-3α-ol-17-one
Etiocholan-3α-ol-17-one 3-glucuronide
3-Oxo-13,17-secoandrost-4-ene-17,13α-lactone
19-Hydroxyandrost-4-ene-3,17-dione
19-Oxoandrost-4-ene-3,17-dione
Androst-4-ene-3,17-dione
Estrone glucuronide
Estradiol-17α
2-Methoxyestrone-3-sulfate
2-Hydroxyestrone
2-Methoxyestrone
2-Methoxyestrone-3-glucuronide
Estrone
16-α-Hydroxyestrone
Androsterone
5α-Androstane-3,17-dione
Androsterone-glucuronide
7α-Hydroxy-androstenedione
C18-Sterolds
7α-Hydroxy-testosterone
5α-Dihydro-testosterone
Androstan-3alpha,17beta-diol
Testosterone
3β,17β-Dihydroxy-androst-5-ene
19-Hydroxy-testosterone
19-Oxo-testosterone
Estradiol-17β
16-Glucuronide-estriol
Estriol
2-Methoxy-estradiol-17β
2-Hydroxy-estradiol-17β
2-Methoxy-estradiol-17β-3-glucuronide
6β-Hydroxy-estradiol-17β
2-Methoxy-estradiol-17β-3-sulfate
Estradiol-17β-3-glucuronide
C19-Sterolds
5β-Dihydro-testosterone
Testosterone glucuronide
Estradiol-17β-3-sulfate
Data on KEGG graph
Rendered by Pathview



Q. Can you do the same procedure as above to plot the pathview figures for the top 5 down-regulated pathways?

```
## Focus on top 5 downregulated pathways here for demo purposes only
keggrespathways_less <- rownames(keggres$less)[1:5]

# Extract the 8 character long IDs part of each string
keggresids_less = substr(keggrespathways_less, start=1, stop=8)
keggresids_less
```

```
[1] "hsa04110" "hsa03030" "hsa03013" "hsa03440" "hsa04114"
```

```r
pathview(gene.data = foldchanges, pathway.id = keggresids_less, species = "hsa")
```




DNA REPLICATION
-1
0
1
Replication complex (Bacteria)
Lagging strand
5'
3'
RNase H / Pol I
Removal of RNA primer
Gap-filling
Lig
DNA ligase
Joining of
Okazaki fragment
Pol III core
Clamp
β
τ
γδ
complex
Primase
DnaG
Primer
DnaB
Leading strand
Pol III core
DNA polymerase III holoenzyme
SSB
Helicase
DNA polymerase III holoenzyme
θ
ε
α
γ & τ
Pol III core
Clamp
ψ
δ'
γδ complex
β
χ
δ
Helicase
DnaB
Primase
DnaG
SSB
RNaseH
RNaseHI
RNaseHII
RNaseHIII
DNA polymerase I
DpoI
DNA ligase
Lig
Replication complex (Archaea)
Lagging strand
RNase H or Dna2
Lig
FEN 1
DNA ligase
DNA polymerase
Pol D/B
Clamp
Clamp loader
RFC
Primase
Leading strand
RFC
Pol B
MCM
RPA/SSB
Helicase
DNA polymerase B
PolB
DNA polymerase D
PolD1
PolD2
Helicase
MCM
Primase
Pri1
Pri2
RPA/SSB
RPA
Clamp
PCNA
Clamp loader
RfcS
RfcL
RNaseH
RNaseHI
RNaseHIII
Helicase
Dna2
Fen1
DNA ligase
Lig
Replication complex (Eukaryotes)
Lagging strand
RNase H or Dna2
Lig I
FEN 1
DNA ligase I
DNA polymerase δ
complex
δ
PCNA
RFC
DNA polymerase α-primase
complex
α-Prim
Leading strand
PCNA
RFC
ε
DNA polymerase ε
complex
MCM
2-7
RPA
Helicase
DNA polymerase α-primase complex
α1
α2
Pri1
Pri2
DNA polymerase δ complex
δ1
δ2
δ3
δ4
DNA polymerase ε complex
ε1
ε2
ε3
ε4
MCM complex (helicase)
Mcm2
Mcm3
Mcm4
Mcm5
Mcm6
Mcm7
RPA
RPA1
RPA2/4
RPA3
Clamp
PCNA
Clamp loader
RFC1
RFC2/4
RFC3/5
RNaseHI
RNaseHI
RNaseHIII
RNaseHIA
RNaseHIB
RNaseHIC
Helicase
Dna2
Fen1
DNA ligase
Lig1
Data on KEGG graph
Rendered by Pathview


NUCLEOCYTOPLASMIC TRANSPORT
-1
0
1
Import
Importin
NLS
NPC
Cytoplasmic fibrils
Cytoplasm
Cytoplasmic ring
Central channel
Lumenal ring
Spoke complex
Lumen
Nucleoplasmic ring
Nucleus
Nuclear basket
NLS
Ran GTP
Importin
Export
Exportin
NES
Pi
Ran GDP
Rae1
DDX 19
Nup 98
Nup358 complex
Nup 214
Nup62 complex
Nup107-160 complex
ELYS
Nup 153
Tpr
Exportin
Ran GTP
NES
mRNA Export
EJC
PYM
AUG
PABP
AAAAA
Upf1
Upf2
Tap
Ref/ Aly
Cytoplasm
Lumen
NPC
Nucleus
mRNA surveilance pathway
SRm160
Pinin
EJC
p15
Upf3
Tap
TREX
Ref/Aly
CBC
m7G
AAAAA
Nuclear Pore complex (NPC)
Cytoplasmic fibrils
ALADIN
hCG1
Gle1
DDX19
Rae1
Nup98
Nup214
Nup88
Nup358 complex
RanBP2
RanGAP
UBC9
SUMO
Cytoplasmic ring / Nucleoplasmic ring (Symmetrical nups)
Nup160
Nup85
Sec13
Nup107
Nup133
Nup96
Nup145
Seh1
Nup43
Nup37
ELYS
Central channel
Nup62
Nup58/45
Nup54
Spoke complex
Nup205
Nup188
Nup155
Nup93
Nup53
Nup59
Lumenal ring
NDC1
gp210
pom121
pom152
pom34
pom33
Nuclear basket
Tpr
Nup50
Nup153
Senp2
Nup2
Nup1
Nup60
Nuclear transport complex
Importin
IPOA
IPOB
Adaptor proteins
SPN1
Exportin
XPO
Ran
eEF1A
PHAX
CBC
NMD3
Exon-junction complex (EJC)
EJC inner core
Y14
MAGOH
MLN51
EIF4A3
EJC outer shell
ACIN1
SAP18
RNPS1
Pinin
Ref/Aly
Transiently interacting factors
Upf1
Upf2
Upf3
Tap
p15
UAP56
SRm160
PYM
Transcription-export (TREX) complex
THO subcomplex
THOC1
THOC2
THOC5
THOC6
THOC7
TEX1
Data on KEGG graph
Rendered by Pathview


HOMOLOGOUS RECOMBINATION
Prokaryotic type
Eukaryotic type
Double strand break
5' to 3' resection
(Escherichia coli)
RecJ
SSB
RecBCD
RecB
RecC
RecD
Filament formation
RecF
RecO
RecR
RecA
RecFOR
RecA filament
Strand invasion
DNA synthesis
DpoI
DpoIII
Holliday junction intermediate
Branch migration and resolution of Holliday junction
RuvA
RuvB
RuvC
or
RecG
RecFOR pathway
Replication restart
PriA
PriB
PriC
DnaT
RecBC pathway
Rad51B
Rad51C
Rad51D
XRCC2
XRCC3
Rad51 paralogs
Recognition
-1
0
1
ATM
+p
(Saccharomyces cerevisiae)
(Mammals)
MRX complex
MRN complex
Rad50
Mre11
XRS2
Nbs1
CtIP
BARD1
BRCA1
BRIP1
TOPBP1
RPA
BRCA2-DSS1
Rad51
NBA1
BRE
BRCC36
Abraxas
RAP80
(S. cerevisiae)
(Common)
Rad55
Rad57
Rad52
PALB2
DSS1
BRCA2
SYCP3
Rad51 filament
Rad59
Rad54
D-loop
DNA synthesis
pol δ
Second end capture
Strand displacement
Flap removal and annealing
Ligation
BLM
TOP3
Mus81
Eme1
Crossover
Non-crossover
DSBR
Double-strand break repair
SDSA
Synthesis-dependent strand annealing
BIR
Break-induced replication
Data on KEGG graph
Rendered by Pathview



Section 3 Gene Ontology

```
data(go.sets.hs)
data(go.subs.hs)

# Focus on Biological Process subset of GO
gobpsets = go.sets.hs[go.subs.hs$BP]

gobpres = gage(foldchanges, gsets=gobpsets)

lapply(gobpres, head)
```

```
$greater
                                                    p.geomean stat.mean        p.val
```

```
GO:0007156 homophilic cell adhesion       8.519724e-05  3.824205 8.519724e-05
GO:0002009 morphogenesis of an epithelium 1.396681e-04  3.653886 1.396681e-04
GO:0048729 tissue morphogenesis           1.432451e-04  3.643242 1.432451e-04
GO:0007610 behavior                       1.925222e-04  3.565432 1.925222e-04
GO:0060562 epithelial tube morphogenesis  5.932837e-04  3.261376 5.932837e-04
GO:0035295 tube development               5.953254e-04  3.253665 5.953254e-04
                                              q.val set.size         exp1
GO:0007156 homophilic cell adhesion       0.1951953      113 8.519724e-05
GO:0002009 morphogenesis of an epithelium 0.1951953      339 1.396681e-04
GO:0048729 tissue morphogenesis           0.1951953      424 1.432451e-04
GO:0007610 behavior                       0.1967577      426 1.925222e-04
GO:0060562 epithelial tube morphogenesis  0.3565320      257 5.932837e-04
GO:0035295 tube development               0.3565320      391 5.953254e-04

$less
                                            p.geomean stat.mean        p.val
GO:0048285 organelle fission             1.536227e-15 -8.063910 1.536227e-15
GO:0000280 nuclear division              4.286961e-15 -7.939217 4.286961e-15
GO:0007067 mitosis                       4.286961e-15 -7.939217 4.286961e-15
GO:0000087 M phase of mitotic cell cycle 1.169934e-14 -7.797496 1.169934e-14
GO:0007059 chromosome segregation        2.028624e-11 -6.878340 2.028624e-11
GO:0000236 mitotic prometaphase          1.729553e-10 -6.695966 1.729553e-10
                                                q.val set.size         exp1
GO:0048285 organelle fission             5.841698e-12      376 1.536227e-15
GO:0000280 nuclear division              5.841698e-12      352 4.286961e-15
GO:0007067 mitosis                       5.841698e-12      352 4.286961e-15
GO:0000087 M phase of mitotic cell cycle 1.195672e-11      362 1.169934e-14
GO:0007059 chromosome segregation        1.658603e-08      142 2.028624e-11
GO:0000236 mitotic prometaphase          1.178402e-07       84 1.729553e-10

$stats
                                          stat.mean     exp1
GO:0007156 homophilic cell adhesion        3.824205 3.824205
GO:0002009 morphogenesis of an epithelium  3.653886 3.653886
GO:0048729 tissue morphogenesis            3.643242 3.643242
GO:0007610 behavior                        3.565432 3.565432
GO:0060562 epithelial tube morphogenesis   3.261376 3.261376
GO:0035295 tube development                3.253665 3.253665
```

#Section 4 Reactome Analysis

```r
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

[1] "Total number of significant genes: 8149"

```r
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quote=
```

> Q8: What pathway has the most significant "Entities p-value"? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

The cell cycle, mitotic has the most significant "Entities p-value" at 2.1E-5. The most significant pathways listed do match my previous KEGG results. Factors such as filtering and mapping contribute to differences between the two methods.