

Recovering the Metric - An Overview of Lie Algebra's

Chaskin Saroff

Oswego State University
Department of Mathematics

16 September 2014



Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

① $[x, x] = 0$ for all $x \in \mathfrak{g}$,

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in V$. and

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in V$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in V$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in V$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④ $[x, y] = -[y, x]$ for all $x, y \in V$.

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in V$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④ $[x, y] = -[y, x]$ for all $x, y \in V$.

The product $[\cdot, \cdot]$ is known as a *Lie bracket* on V .

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[,]$ is alternating then

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ , \]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ , \]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

1 $\langle x, y \rangle = \langle y, x \rangle$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

- 1 $\langle x, y \rangle = \langle y, x \rangle$
- 2 If $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$
then $x = \vec{0}$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

- 1 $\langle x, y \rangle = \langle y, x \rangle$
- 2 If $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$
then $x = \vec{0}$
- 3 $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$
and $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

- 1 $\langle x, y \rangle = \langle y, x \rangle$
- 2 If $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$
then $x = \vec{0}$
- 3 $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$
and $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$
- 4 $\langle x, x \rangle \geq 0$
and $\langle x, x \rangle = 0 \implies x = \vec{0}$

An *inner product* on the vector \mathfrak{g} is a *symmetric, non-degenerate, bilinear and positive definite* function on \mathfrak{g} . That is, an inner product on \mathfrak{g} is a function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ that satisfies the following properties: For $x, y, z \in \mathfrak{g}$

- 1 $\langle x, y \rangle = \langle y, x \rangle$
- 2 If $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$
then $x = \vec{0}$
- 3 $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$
and $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$
- 4 $\langle x, x \rangle \geq 0$
and $\langle x, x \rangle = 0 \implies x = \vec{0}$

A First Example

Let V be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

A First Example

Let V be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

A First Example

Let V be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

A First Example

Let V be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

A First Example

Let V be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ = (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \end{aligned}$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} & (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ &= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \\ &= 0. \end{aligned}$$

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z .

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0.

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

The basis vectors x, y , and z can be represented by the following matrices

$$x = \begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{and} \quad z = \begin{pmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

The basis vectors x, y , and z can be represented by the following matrices

$$x = \begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{and} \quad z = \begin{pmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Then

$$\begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

The basis vectors x, y , and z can be represented by the following matrices

$$x = \begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{and} \quad z = \begin{pmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Then

$$\begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \mathbf{1} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & \mathbf{1} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

- \mathfrak{h}_3 is *not* a subalgebra of $\mathfrak{gl}(3, \mathbb{R})$ because it doesn't have the same bracket.

\mathfrak{h}_{2k+1} : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

\mathfrak{h}_{2k+1} : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider \mathbb{R}^{2k+1} , $k \geq 1$, with basis vectors x_1, x_2, \dots, x_k , y_1, y_2, \dots, y_k , and z .

\mathfrak{h}_{2k+1} : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider \mathbb{R}^{2k+1} , $k \geq 1$, with basis vectors $x_1, x_2, \dots, x_k, y_1, y_2, \dots, y_k$, and z . Define a Lie bracket on these generators *via*

$$[x_i, x_j] = 0, \quad [y_i, y_j] = 0, \quad [x_i, y_j] = \delta_{ij}, \quad \text{and} \quad [z, \cdot] = 0.$$

\mathfrak{h}_{2k+1} : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider \mathbb{R}^{2k+1} , $k \geq 1$, with basis vectors x_1, x_2, \dots, x_k , y_1, y_2, \dots, y_k , and z . Define a Lie bracket on these generators *via*

$$[x_i, x_j] = 0, \quad [y_i, y_j] = 0, \quad [x_i, y_j] = \delta_{ij}, \quad \text{and} \quad [z, \cdot] = 0.$$

A general element of this space can be represented by a matrix of the form

$$X = \begin{pmatrix} 1 & x_1 & x_2 & \cdots & z \\ 0 & 1 & 0 & \cdots & y_1 \\ 0 & 0 & 1 & \cdots & y_2 \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}$$

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- 1 If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V$, $a, b \in \mathbb{R}$.

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n :

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n : Let $x, y \in \mathbb{R}^n$; then $x \cdot y = x^T y$.

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V, a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n : Let $x, y \in \mathbb{R}^n$; then $x \cdot y = x^T y$.

Example The *Minkowski inner product* on \mathbb{R}^n :

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n : Let $x, y \in \mathbb{R}^n$; then $x \cdot y = x^T y$.

Example The *Minkowski inner product* on \mathbb{R}^n : Let $x = (t, x_1, x_2, x_3)$ and $y = (s, y_1, y_2, y_3)$ be two vectors in \mathbb{R}^4 .

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V, a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n : Let $x, y \in \mathbb{R}^n$; then $x \cdot y = x^T y$.

Example The *Minkowski inner product* on \mathbb{R}^n : Let $x = (t, x_1, x_2, x_3)$ and $y = (s, y_1, y_2, y_3)$ be two vectors in \mathbb{R}^4 . Then the Minkowski inner product of x and y is given by

$$\langle x, y \rangle = -(ts) + x_1 y_1 + x_2 y_2 + x_3 y_3.$$

Inner Products

An *inner product* on a vector space V is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in V$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in V$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in V, a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^n : Let $x, y \in \mathbb{R}^n$; then $x \cdot y = x^T y$.

Example The *Minkowski inner product* on \mathbb{R}^n : Let $x = (t, x_1, x_2, x_3)$ and $y = (s, y_1, y_2, y_3)$ be two vectors in \mathbb{R}^4 . Then the Minkowski inner product of x and y is given by

$$\langle x, y \rangle = -(ts) + x_1 y_1 + x_2 y_2 + x_3 y_3.$$

This is the inner product that gives rise to Einstein's Special Relativity!

Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra $(\mathfrak{g}, [,])$ on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra $(\mathfrak{g}, [,])$ on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing $x \in \mathfrak{g}$, $\mathrm{ad}_x = [x, \]$ becomes a function from \mathfrak{g} to itself.

Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra $(\mathfrak{g}, [,])$ on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing $x \in \mathfrak{g}$, $\mathrm{ad}_x = [x, \]$ becomes a function from \mathfrak{g} to itself. One can then define a map $j(x) : \mathfrak{g} \rightarrow \mathfrak{g}$ by the formula

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

for all $y, z \in \mathfrak{g}$.

Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra $(\mathfrak{g}, [,])$ on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing $x \in \mathfrak{g}$, $\mathrm{ad}_x = [x, \]$ becomes a function from \mathfrak{g} to itself. One can then define a map $j(x) : \mathfrak{g} \rightarrow \mathfrak{g}$ by the formula

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

for all $y, z \in \mathfrak{g}$.

The map $j(x)$ is called the *adjoint* to ad_x with respect to the inner product \langle , \rangle . (Confusing, I know.)

j -maps

The equation that we used to define $j(x)$,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

j -maps

The equation that we used to define $j(x)$,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

j -maps

The equation that we used to define $j(x)$,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j(x) = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

j -maps

The equation that we used to define $j(x)$,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j(x) = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

j -maps

The equation that we used to define $j(x)$,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j(x) = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

This makes the computation of many complicated geometric objects—such as curvatures—into “simple” calculations in linear algebra!

Lie Algebra Seminar

There is a “Group Study Seminar” on Lie Algebras that meets every week!

Day: Thursdays

Time: 5:30 – 6:30 pm

Room: Math Commons

We'll start with a “review” of vector spaces, then spend a lot of time building examples. Eventually we'll prove some theorems, and study a little geometry (time permitting).