

Encoding the Bracket - An Overview of Lie Algebras

Alexander Jansing, Chaskin Saroff

Oswego State University
Department of Mathematics

2 May 2015



A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\ , \] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

① $[x, x] = 0$ for all $x \in \mathfrak{g}$,

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- 3 $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- 3 $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④ $[x, y] = -[y, x]$ for all $x, y \in \mathfrak{g}$.

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④ $[x, y] = -[y, x]$ for all $x, y \in \mathfrak{g}$.

The product $[\cdot, \cdot]$ is known as a *Lie bracket* on \mathfrak{g} .

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[,]$ is alternating then

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ , \]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[,]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

Conversely, if $[,]$ is skew-symmetric, then $[x, x] + [x, x] = 0$ implies that $2[x, x] = 0$. Now we see that this implies $[x, x] = 0$ so long as our field is not of characteristic 2, for in those spaces $2 = 0$ and we can deduce nothing about $[x, x]$.

The *center* of a Lie Algebra, \mathfrak{g} is

The *center* of a Lie Algebra, \mathfrak{g} is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

The *center* of a Lie Algebra, \mathfrak{g} is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

A vector, z , of \mathfrak{g} is said to be in the center of \mathfrak{g} if

$$[x, z] = \mathbf{0} \ \forall x \in \mathfrak{g}.$$

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ = (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \end{aligned}$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} & (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ &= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \\ &= 0. \end{aligned}$$

Consider a 3-dimensional vector space with basis vectors x, y, z .

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0.

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .




Figure: The Heisenberg Algebra, \mathfrak{h}_3

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ④ If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^3 :

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3$$

The equation that we used to define j_z ,

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

The equation that we used to define j_z ,

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

The equation that we used to define j_z ,

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j_z = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

The equation that we used to define j_z ,

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j_z = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

The equation that we used to define j_z ,

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define $j_z = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$.

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

This makes the computation of many complicated geometric objects—such as curvatures—into “simple” calculations in linear algebra!

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form.

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

$$[e_i, e_i] = \mathbf{0}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$?

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$?

$$[e_i, e_j] = -[e_j, e_i]$$

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Recall from before the adjoint map, $\text{ad}_{e_1} : \mathfrak{g} \rightarrow \mathfrak{g}$ to itself. Can anyone see a matrix representation of ad_{e_1} ?

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Recall from before the adjoint map, $\text{ad}_{e_1} : \mathfrak{g} \rightarrow \mathfrak{g}$ to itself. Can anyone see a matrix representation of ad_{e_1} ?

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket “map”?

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket “map”?
Notice that we can retrieve information about the bases.

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

So it will be enough find a way to write all Lie Brackets as a linear combination of basis brackets.

We can write $[x, y]$ as a linear combination their component vectors's brackets.
Let's look at an example of a 2 dimensional space

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\text{So } [x, y] = [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2]$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2]\end{aligned}$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2]\end{aligned}$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2]\end{aligned}$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\&= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\&= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\&= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\&= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22}\end{aligned}$$

We can write $[x, y]$ as a linear combination their component vectors's brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\ &= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22}\end{aligned}$$

This means that the Lie Bracket is fully described by the matrix, L

As before, we'll investigate the 2 dimensional example.

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$

That is $[x, y] = x^T L y$

What does Abelian \mathbb{R}^3 look like in this form?

What does Abelian \mathbb{R}^3 look like in this form?
Recall that a Lie Algebra is Abelian if

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of Abelian \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = e_2$$

$$[e_2, e_3] = e_1$$

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

What is the center of \mathbb{R}^3 with the cross product?

How could the three dimensional Heisenberg Algebra be represented?

How could the three dimensional Heisenberg Algebra be represented?
 \mathfrak{h}_3 is two-step nilpotent

How could the three dimensional Heisenberg Algebra be represented?

\mathfrak{h}_3 is two-step nilpotent

$$[[x, y], z] \quad \forall x, y, z \in \mathfrak{h}_3$$

How could the three dimensional Heisenberg Algebra be represented?

\mathfrak{h}_3 is two-step nilpotent

$[[x, y], z] \quad \forall x, y, z \in \mathfrak{h}_3$

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

How could the three dimensional Heisenberg Algebra be represented?

\mathfrak{h}_3 is two-step nilpotent

$[[x, y], z] \quad \forall x, y, z \in \mathfrak{h}_3$

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ?

How could the three dimensional Heisenberg Algebra be represented?

\mathfrak{h}_3 is two-step nilpotent

$[[x, y], z] \quad \forall x, y, z \in \mathfrak{h}_3$

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ?

The center is the span of $\{e_3\}$.

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Recall that $\langle e_i, e_j \rangle = \langle e_j, e_i \rangle$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And $\mathbf{0}$ cannot be a basis vector because it is linearly dependent with all vectors.

So $e_{ij} > 0$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And $\mathbf{0}$ cannot be a basis vector because it is linearly dependent with all vectors.

So $e_{ij} > 0$

The Dot Product can be represented in this way.

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

$$E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Recall that for any $z \in \mathfrak{z}$, the map $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ is defined by the identity

$$\langle y, j_z(x) \rangle_{\mathfrak{v}} = \langle z, [x, y] \rangle_{\mathfrak{z}}. \quad (7)$$

We wish to find a matrix J_z that represents the map j_z . Since j_z maps \mathfrak{v} to itself, J_z should be a matrix in $\mathbb{R}^{n \times n}$. Rewriting (7) in matrix form, we get

$$\begin{aligned} \langle y, j_z(x) \rangle_{\mathfrak{v}} &= \langle z, [x, y] \rangle_{\mathfrak{z}} \\ y^T E(J_z x) &= z^t (x^T L y) \\ y^T (E J_z) x &= z^t (-y^T L x) \\ &= -z^t (y^T L x) \\ &= -y^T (z^t L) x \\ &= y^T (z^t (-L)) x \\ &= y^T (z^t L^T) x. \end{aligned}$$

The equation $y^T(EJ_z)x = y^T(z^{\mathfrak{t}}L^T)x$ must hold for arbitrary vectors x and y in \mathfrak{v} , implying that

$$EJ_z = z^{\mathfrak{t}}L^T.$$

Since $\det(E) \neq 0$, it is invertible and

$$J_z = E^{-1}(z^{\mathfrak{t}}L^T). \quad (8)$$

This gives us a method to compute the j -map for any z using only matrix computations. Furthermore, if $z = z_k$ is a basis vector of \mathfrak{z} , then $z_k^{\mathfrak{t}}L^T$ is just $(L^k)^T$. Thus, for basis vectors z_k , the map j_{z_k} is represented by the matrix

$$J_{z_k} = E^{-1}(L^k)^T.$$

Since the j -maps are linear, knowing how they act on a basis of \mathfrak{z} is good enough. Indeed, we should be able to define a stack of j -maps by

$$J = E^{-1}L^T. \tag{9}$$

The j -map j_z *should* then be $J_z = z^\dagger J$.

Using the programming language, sage, we created a program that would compute the j -maps of an arbitrary Lie algebra.




Figure: The L-Stack




Figure: The L-Stack