Encoding the Bracket - An Overview of Lie Algebras

Alexander Jansing, Chaskin Saroff

Oswego State University Department of Mathematics

1 May 2015





Definition

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g \times \mathfrak g \to \mathfrak g$ that satisfies three properties:

Definition

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g\times\mathfrak g\to\mathfrak g$ that satisfies three properties:

Definition

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g\times\mathfrak g\to\mathfrak g$ that satisfies three properties:

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g\times\mathfrak g\to\mathfrak g$ that satisfies three properties:

- **②** [[x,y],z] + [[y,z],x] + [[z,x],y] = 0 for all $x,y,z \in \mathfrak{g}$. and
- **3** [x + y, z] = [x, z] + [y, z] and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g\times\mathfrak g\to\mathfrak g$ that satisfies three properties:

- ② [[x,y],z] + [[y,z],x] + [[z,x],y] = 0 for all $x,y,z \in \mathfrak{g}$. and
- **3** [x + y, z] = [x, z] + [y, z] and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

A $\mathit{Lie\ Algebra}$ is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g\times\mathfrak g\to\mathfrak g$ that satisfies three properties:

- ② [[x,y],z] + [[y,z],x] + [[z,x],y] = 0 for all $x,y,z \in \mathfrak{g}$. and
- **3** [x + y, z] = [x, z] + [y, z] and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called skew symmetry:

A Lie Algebra is a (finite-dimensional) vector space $\mathfrak g$ together with a bilinear multiplication $[\ ,\]:\mathfrak g \times \mathfrak g \to \mathfrak g$ that satisfies three properties:

- ② [[x,y],z] + [[y,z],x] + [[z,x],y] = 0 for all $x,y,z \in \mathfrak{g}$. and
- **3** [x + y, z] = [x, z] + [y, z] and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called skew symmetry:

The product [,] is known as a Lie bracket on g.



skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ ,\]$ is alternating then

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ ,\]$ is alternating then

$$0 = [x+y, x+y] \tag{1}$$

$$= [x + y, x] + [x + y, y]$$
 (2)

$$= [x, x] + [y, x] + [x + y, y]$$
 (3)

$$= [x, x] + [y, x] + [x, y] + [y, y]$$
 (4)

$$0 = [x, y] + [y, x]$$
 (5)

$$\implies [x, y] = -[y, x] \tag{6}$$

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if $[\ ,\]$ is alternating then

$$0 = [x+y, x+y] \tag{1}$$

$$= [x + y, x] + [x + y, y]$$
 (2)

$$= [x, x] + [y, x] + [x + y, y]$$
(3)

$$= [x, x] + [y, x] + [x, y] + [y, y]$$
 (4)

$$0 = [x, y] + [y, x]$$
 (5)

$$\implies [x,y] = -[y,x] \tag{6}$$

Conversely, if $[\ ,\]$ is skew-symmetric, then [x,x]+[x,x]=0 implies that 2[x,x]=0. Now we see that this implies [x,x]=0 so long as our field is not of characteristic 2, for in those spaces 2=0 and we can deduce nothing about [x,x].

The Center

The *center* of a Lie Algebra, $\mathfrak g$ is

The center of a Lie Algebra, g is

$$\mathfrak{z}=\{z\in\mathfrak{g}\mid [z,x]=\mathbf{0}\ \forall x\in\mathfrak{g}\}.$$

The center of a Lie Algebra, g is

$$\mathfrak{z} = \{ z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g} \}.$$

A vector, z, of $\mathfrak g$ is said to be in the center of $\mathfrak g$ if

$$[x,z] = \mathbf{0} \ \forall x \in \mathfrak{g}.$$

Let ${\mathfrak g}$ be any vector space, and define

$$[x,y]=0 \ \text{ for all } x,y\in \mathfrak{g}.$$

Let ${\mathfrak g}$ be any vector space, and define

$$[x, y] = 0$$
 for all $x, y \in \mathfrak{g}$.

Clearly $[\;,\;]$ is alternating:

$$[x,x]=0$$
 for all $x\in\mathfrak{g}$.

Let ${\mathfrak g}$ be any vector space, and define

$$[x, y] = 0$$
 for all $x, y \in \mathfrak{g}$.

Clearly [,] is alternating:

$$[x,x]=0$$
 for all $x\in\mathfrak{g}$.

And the Jacobi Identity is trivial:

Let $\mathfrak g$ be any vector space, and define

$$[x, y] = 0$$
 for all $x, y \in \mathfrak{g}$.

Clearly [,] is alternating:

$$[x,x]=0$$
 for all $x\in\mathfrak{g}$.

And the Jacobi Identity is trivial:

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = [0, z] + [0, x] + [0, y]$$

= 0 + 0 + 0 = 0.

Let $\mathfrak g$ be any vector space, and define

$$[x, y] = 0$$
 for all $x, y \in \mathfrak{g}$.

Clearly [,] is alternating:

$$[x,x]=0$$
 for all $x\in\mathfrak{g}$.

And the Jacobi Identity is trivial:

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = [0, z] + [0, x] + [0, y]$$

= 0 + 0 + 0 = 0.

This Algebra is called Abelian or One-Step Nilpotent



Let $\mathfrak g$ be any vector space, and define

$$[x, y] = 0$$
 for all $x, y \in \mathfrak{g}$.

Clearly [,] is alternating:

$$[x,x]=0$$
 for all $x\in\mathfrak{g}$.

And the Jacobi Identity is trivial:

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = [0, z] + [0, x] + [0, y]$$

= 0 + 0 + 0 = 0.

This Algebra is called Abelian or One-Step Nilpotent



Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The cross product of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The cross product of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x$$
 for all $x, y \in \mathbb{R}^3$.

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x$$
 for all $x, y \in \mathbb{R}^3$.



Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The cross product of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x$$
 for all $x, y \in \mathbb{R}^3$.

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The cross product of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x$$
 for all $x, y \in \mathbb{R}^3$.

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

= $(x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z$



Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The cross product of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x$$
 for all $x, y \in \mathbb{R}^3$.

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

$$= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z$$

$$= 0.$$



\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x, y, z.

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x,y,z. Define a Lie bracket on this space by

$$[x,y]=z,$$

with all other brackets equal to 0.

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x,y,z. Define a Lie bracket on this space by

$$[x,y]=z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

\mathfrak{h}_3 : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors x,y,z. Define a Lie bracket on this space by

$$[x,y]=z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .




Figure: The Heisenberg Algebra, \$\opena_3\$

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle : \mathfrak g \times \mathfrak g \to \mathbb R.$

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

- If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

- If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

- If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;

Example The *dot product* on \mathbb{R}^3 :

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

- If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

Inner Products

An inner product on a vector space $\mathfrak g$ is a non-degenerate, symmetric, bilinear function $\langle \ , \ \rangle: \mathfrak g \times \mathfrak g \to \mathbb R.$

- If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

$$\langle x,y\rangle=x_1y_1+x_2y_2+x_3y_3$$

j-maps

The equation that we used to define j_z ,

$$\langle [x,y],z\rangle = \langle y,j_zx\rangle$$

could be re-written to look like

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \operatorname{\mathsf{ad}}_{\mathsf{x}} y, z \rangle = \langle y, \operatorname{\mathsf{ad}}_{\mathsf{x}}^{\dagger} z \rangle.$$

$$\langle [x,y],z\rangle = \langle y,j_zx\rangle$$

could be re-written to look like

$$\langle \mathsf{ad}_x y, z \rangle = \langle y, \mathsf{ad}_x^\dagger z \rangle.$$

Now we could define $j_z = \operatorname{\mathsf{ad}}_x^\dagger : \mathfrak{g} \to \mathfrak{g}$.

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

could be re-written to look like

$$\langle \mathsf{ad}_x y, z \rangle = \langle y, \mathsf{ad}_x^{\dagger} z \rangle.$$

Now we could define $j_z = \operatorname{ad}_x^{\dagger} : \mathfrak{g} \to \mathfrak{g}$.

The j-maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

$$\langle [x,y],z\rangle = \langle y,j_zx\rangle$$

could be re-written to look like

$$\langle \mathsf{ad}_x y, z \rangle = \langle y, \mathsf{ad}_x^{\dagger} z \rangle.$$

Now we could define $j_z = \operatorname{ad}_x^{\dagger} : \mathfrak{g} \to \mathfrak{g}$.

The j-maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

This makes the computation of many complicated geometric objects—such as curvatures—into "simple" calculations in linear algebra!

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form.

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

$$[e_i,e_i]=\boldsymbol{0}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$?

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$? $[e_i, e_j] = -[e_j, e_i]$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Recall from before the adjoint map, $ad_{e_1}: \mathfrak{g} \to \mathfrak{g}$ to itself. Can anyone see a matrix representation of ad_{e_1} ?

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \ldots, e_n Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} 0 & L_{12} & \cdots & L_{1n} \\ -L_{12} & 0 & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & 0 \end{pmatrix}$$

Recall from before the adjoint map, $ad_{e_1}: \mathfrak{g} \to \mathfrak{g}$ to itself. Can anyone see a matrix representation of ad_{e_1} ?

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding to as the Lie Bracket "map"?

$$[e_1,e_2] = L_{12}$$

$$[e_1,e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & \boxed{L_{12}} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & \boxed{L_{12}} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

So it will be enough find a way to write all Lie Brackets as a linear combination of basis brackets.

We can write [x,y] as a linear combination their component vectors's brackets. Let's look at an example of a 2 dimensional space Let $x,y\in\mathfrak{g}$ with

We can write [x, y] as a linear combination their component vectors's brackets. Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x=w_1e_1+w_2e_2$$

Let
$$x, y \in \mathfrak{g}$$
 with

$$x=w_1e_1+w_2e_2$$

$$y=v_1e_1+v_2e_2$$

Let
$$x, y \in \mathfrak{g}$$
 with

$$x=w_1e_1+w_2e_2$$

$$y = v_1 e_1 + v_2 e_2$$

So
$$[x, y] = [w_1e_1 + w_2e_2, v_1e_1 + v_2e_2]$$

Let
$$x,y\in\mathfrak{g}$$
 with

$$x=w_1e_1+w_2e_2$$

$$y=v_1e_1+v_2e_2$$

So
$$[x, y] = [w_1e_1 + w_2e_2, v_1e_1 + v_2e_2]$$

= $[w_1e_1, v_1e_1 + v_2e_2] + [w_2e_2, v_1e_1 + v_2e_2]$

Let
$$x, y \in \mathfrak{g}$$
 with $x = w_1 e_1 + w_2 e_2$
 $y = v_1 e_1 + v_2 e_2$

So
$$[x, y] = [w_1e_1 + w_2e_2, v_1e_1 + v_2e_2]$$

 $= [w_1e_1, v_1e_1 + v_2e_2] + [w_2e_2, v_1e_1 + v_2e_2]$
 $= [w_1e_1, v_1e_1] + [w_1e_1, v_2e_2] + [w_2e_2, v_1e_1] + [w_2e_2, v_2e_2]$

Let
$$x, y \in \mathfrak{g}$$
 with $x = w_1e_1 + w_2e_2$ $y = v_1e_1 + v_2e_2$

So
$$[x, y] = [w_1e_1 + w_2e_2, v_1e_1 + v_2e_2]$$

 $= [w_1e_1, v_1e_1 + v_2e_2] + [w_2e_2, v_1e_1 + v_2e_2]$
 $= [w_1e_1, v_1e_1] + [w_1e_1, v_2e_2] + [w_2e_2, v_1e_1] + [w_2e_2, v_2e_2]$
 $= w_1v_1[e_1, e_1] + w_1v_2[e_1, e_2] + w_2v_1[e_2, e_1] + w_2v_2[e_2, e_2]$

Let
$$x, y \in \mathfrak{g}$$
 with $x = w_1e_1 + w_2e_2$
 $y = v_1e_1 + v_2e_2$

So
$$[x, y] = [w_1e_1 + w_2e_2, v_1e_1 + v_2e_2]$$

 $= [w_1e_1, v_1e_1 + v_2e_2] + [w_2e_2, v_1e_1 + v_2e_2]$
 $= [w_1e_1, v_1e_1] + [w_1e_1, v_2e_2] + [w_2e_2, v_1e_1] + [w_2e_2, v_2e_2]$
 $= w_1v_1[e_1, e_1] + w_1v_2[e_1, e_2] + w_2v_1[e_2, e_1] + w_2v_2[e_2, e_2]$
 $= w_1v_1L_{11} + w_1v_2L_{12} + w_2v_1L_{21} + w_2v_2L_{22}$

We can write [x,y] as a linear combination their component vectors's brackets. Let's look at an example of a 2 dimensional space

Let
$$x, y \in \mathfrak{g}$$
 with $x = w_1 e_1 + w_2 e_2$ $y = v_1 e_1 + v_2 e_2$ $y = v_1 e_1 + v_2 e_2$

$$So [x, y] = [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2]$$

$$= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2]$$

$$= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2]$$

$$= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2]$$

 $= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22}$

This means that the Lie Bracket is fully described by the matrix, L

As before, we'll investigate the 2 dimensional example.

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$\mathbf{x}^{\mathsf{T}} L = \begin{pmatrix} w_1 w_2 \end{pmatrix} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix}$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$
$$x^T L = \begin{pmatrix} w_1 w_2 \end{pmatrix} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix}$$
$$(x^T L) y = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = \begin{pmatrix} w_1 w_2 \end{pmatrix} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix}$$

$$(x^T L) y = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= \begin{pmatrix} w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{12} + w_2 v_2 L_{22} \end{pmatrix}$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = \begin{pmatrix} w_1 w_2 \end{pmatrix} \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix}$$

$$(x^T L) y = \begin{pmatrix} w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= \begin{pmatrix} w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{12} + w_2 v_2 L_{22} \end{pmatrix}$$

That is $[x, y] = x^T L y$

What does Abelian \mathbb{R}^3 look like in this form?

What does Abelian \mathbb{R}^3 look like in this form? Recall that a Lie Algebra is Abelian if

What does Abelian \mathbb{R}^3 look like in this form? Recall that a Lie Algebra is Abelian if $[x,y]=\mathbf{0} \ \forall x,y\in\mathbb{R}^3$

What does Abelian \mathbb{R}^3 look like in this form? Recall that a Lie Algebra is Abelian if

$$[x,y] = \mathbf{0} \ \forall x,y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What does Abelian \mathbb{R}^3 look like in this form? Recall that a Lie Algebra is Abelian if $[x,y] = \mathbf{0} \ \forall x,y \in \mathbb{R}^3$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of Abelian \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1,e_2]=e_3$$

$$[e_1,e_3]=e_2$$

$$\left[e_2,e_3\right]=e_1$$

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1,e_2]=e_3$$

$$[e_1, e_3] = e_2$$

 $[e_2, e_3] = e_1$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_3 & -e_1 & \mathbf{0} \end{pmatrix}$$

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1,e_2]=e_3$$

$$[e_1,e_3]=e_2$$

$$[e_2,e_3]=e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

What is the center of \mathbb{R}^3 with the cross product?

β₃'s Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

\$\mathbf{h}_3\end{array}\text{s Lie Bracket}

How could the three dimensional Heisenberg Algebra be represented? \mathfrak{h}_3 is two-step nilpotent

η₃'s Lie Bracket

How could the three dimensional Heisenberg Algebra be represented? \mathfrak{h}_3 is two-step nilpotent $[[x,y],z] \ \forall x,y,z \in \mathfrak{h}_3$

\$\textit{h}_3's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented? \mathfrak{h}_3 is two-step nilpotent

$$[[x,y],z] \ \forall x,y,z \in \mathfrak{h}_3$$

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

\$\textit{h}_3's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented? \mathfrak{h}_3 is two-step nilpotent

$$\left[\left[x,y\right],z\right] \ \forall x,y,z\in\mathfrak{h}_{3}$$

$$\begin{pmatrix} \mathbf{0} & e_3 & \mathbf{0} \\ -e_3 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ?

\$\textit{h}_3's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented? \mathfrak{h}_3 is two-step nilpotent $[[x,y],z] \ \forall x,y,z \in \mathfrak{h}_3$

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ? The center is the span of $\{e_3\}$.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Recall that $\langle e_i, e_j \rangle = \langle e_j, e_i \rangle$

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And ${\bf 0}$ canot be a basis vector because it is linearly dependent with all vectors.

So $e_{ii} > 0$

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And ${\bf 0}$ canot be a basis vector because it is linearly dependent with all vectors.

So
$$e_{ii} > 0$$

Dot Product on \mathbb{R}^3

Dot Product on \mathbb{R}^3

$$\langle e_i, e_j \rangle = 0$$

Dot Product on \mathbb{R}^3

$$\langle e_i, e_i \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

$$E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Recall that for any $z \in \mathfrak{z}$, the map $j_z : \mathfrak{v} \to \mathfrak{v}$ is defined by the identity

$$\langle y, j_z(x) \rangle_{\mathfrak{v}} = \langle z, [x, y] \rangle_{\mathfrak{z}}.$$
 (7)

We wish to find a matrix J_z that represents the map j_z . Since j_z maps v to itself, J_z should be a matrix in $\mathbb{R}^{n \times n}$. Rewriting (7) in matrix form, we get

$$\langle y, j_z(x) \rangle_v = \langle z, [x, y] \rangle_{\delta}$$

$$y^T E(J_z x) = z^{t}(x^T L y)$$

$$y^T (EJ_z) x = z^{t}(-y^T L x)$$

$$= -z^{t}(y^T L x)$$

$$= -y^T(z^{t} L) x$$

$$= y^T(z^{t}(-L)) x$$

$$= y^T(z^{t} L^T) x.$$

The equation $y^T(EJ_z)x = y^T(z^tL^T)x$ must hold for arbitrary vectors x and y in v, implying that

$$EJ_z=z^{\mathrm{t}}L^{\mathrm{T}}.$$

Since $det(E) \neq 0$, it is invertible and

$$J_z = E^{-1}(z^{\mathrm{t}}L^{\mathrm{T}}). \tag{8}$$

This gives us a method to compute the j-map for any z using only matrix computations. Furthermore, if $z=z_k$ is a basis vector of \mathfrak{z} , then $z_k^{\mathtt{t}} L^{\mathrm{T}}$ is just $(L^k)^{\mathrm{T}}$. Thus, for basis vectors z_k , the map j_{z_k} is represented by the matrix

$$J_{z_k} = E^{-1}(L^k)^{\mathrm{T}}.$$

Since the j-maps are linear, knowing how they act on a basis of \mathfrak{z} is good enough. Indeed, we should be able to define a stack of j-maps by

$$J = E^{-1}L^{\mathrm{T}}. (9)$$

The *j*-map j_z should then be $J_z = z^t J$.

What we did

Using the programming language, sage, we created a program that would compute the j-maps of an arbitrary Lie algebra.

TikZ

Acknowledgements




Figure : The L-Stack




Figure : The L-Stack