

# Recovering the Metric - An Overview of Lie Algebra's

Chaskin Saroff

Oswego State University  
Department of Mathematics

16 September 2014



# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[ , ] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[ , ] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- 1  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- 2  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in V$ . and

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in V$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in V$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[ , ] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in V$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④  $[x, y] = -[y, x]$  for all  $x, y \in V$ .

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in V$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y]$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 is known as *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④  $[x, y] = -[y, x]$  for all  $x, y \in V$ .

The product  $[\cdot, \cdot]$  is known as a *Lie bracket* on  $V$ .



# skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if  $[ , ]$  is alternating then

## skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if  $[ , ]$  is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

## skew-symmetry

By bilinearity, every alternating product is also skew-symmetric, regardless of the characteristic of the underlying field. Indeed, if  $[ , ]$  is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

Conversely, if  $[ , ]$  is skew-symmetric, then  $[x, x] + [x, x] = 0$  implies that  $2[x, x] = 0$ . Now we see that this implies  $[x, x] = 0$  so long as our field is not of characteristic 2, for in those spaces  $2 = 0$  and we can deduce nothing about  $[x, x]$ .

# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

1  $\langle x, y \rangle = \langle y, x \rangle$

# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

- ①  $\langle x, y \rangle = \langle y, x \rangle$
- ② If  $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$   
then  $x = \vec{0}$

# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

- ①  $\langle x, y \rangle = \langle y, x \rangle$
- ② If  $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$   
then  $x = \vec{0}$
- ③  $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$   
and  $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$

# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

- ①  $\langle x, y \rangle = \langle y, x \rangle$
- ② If  $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$   
then  $x = \vec{0}$
- ③  $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$   
and  $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$
- ④  $\langle x, x \rangle \geq 0$   
and  $\langle x, x \rangle = 0 \implies x = \vec{0}$



# Inner Product

An *inner product* on the vector  $\mathfrak{g}$  is a *symmetric, non-degenerate, bilinear and positive definite* function on  $\mathfrak{g}$ . That is, an inner product on  $\mathfrak{g}$  is a function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  that satisfies the following properties: For  $x, y, z \in \mathfrak{g}$

- ①  $\langle x, y \rangle = \langle y, x \rangle$
- ② If  $\langle x, y \rangle = 0 \quad \forall y \in \mathfrak{g}$   
then  $x = \vec{0}$
- ③  $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$   
and  $\alpha \langle x, y \rangle = \langle \alpha x, y \rangle = \langle x, \alpha y \rangle$
- ④  $\langle x, x \rangle \geq 0$   
and  $\langle x, x \rangle = 0 \implies x = \vec{0}$

# The Center

The *center* of a Lie Algebra,  $\mathfrak{g}$  is

# The Center

The *center* of a Lie Algebra,  $\mathfrak{g}$  is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \vec{0} \ \forall x \in \mathfrak{g}\}.$$

# The Center

The *center* of a Lie Algebra,  $\mathfrak{g}$  is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \vec{0} \ \forall x \in \mathfrak{g}\}.$$

A vector,  $z$ , of  $\mathfrak{g}$  is said to be in the center of  $\mathfrak{g}$  if

$$[x, z] = \vec{0} \ \forall x \in \mathfrak{g}.$$

# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$



# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*

# One-Step Nilpotence

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in V.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in V.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ = (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \end{aligned}$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} & (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ &= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \\ &= 0. \end{aligned}$$



# $\mathfrak{h}_3$ : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors  $x, y, z$ .

## $\mathfrak{h}_3$ : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0.

## $\mathfrak{h}_3$ : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by  $\mathfrak{h}_3$ .

## $\mathfrak{h}_3$ : The Heisenberg Algebra

Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by  $\mathfrak{h}_3$ .




Figure : The Heisenberg Algebra,  $\mathfrak{h}$

# $\mathfrak{h}_{2k+1}$ : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

## $\mathfrak{h}_{2k+1}$ : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider  $\mathbb{R}^{2k+1}$ ,  $k \geq 1$ , with basis vectors  $x_1, x_2, \dots, x_k$ ,  $y_1, y_2, \dots, y_k$ , and  $z$ .

## $\mathfrak{h}_{2k+1}$ : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider  $\mathbb{R}^{2k+1}$ ,  $k \geq 1$ , with basis vectors  $x_1, x_2, \dots, x_k, y_1, y_2, \dots, y_k$ , and  $z$ . Define a Lie bracket on these generators *via*

$$[x_i, x_j] = 0, \quad [y_i, y_j] = 0, \quad [x_i, y_j] = \delta_{ij}, \quad \text{and} \quad [z, \cdot] = 0.$$

## $\mathfrak{h}_{2k+1}$ : Higher Dimensional Heisenberg Algebras

Higher odd-dimensional analogues of the Heisenberg algebra can also be defined.

Consider  $\mathbb{R}^{2k+1}$ ,  $k \geq 1$ , with basis vectors  $x_1, x_2, \dots, x_k$ ,  $y_1, y_2, \dots, y_k$ , and  $z$ . Define a Lie bracket on these generators *via*

$$[x_i, x_j] = 0, \quad [y_i, y_j] = 0, \quad [x_i, y_j] = \delta_{ij}, \quad \text{and} \quad [z, \cdot] = 0.$$

A general element of this space can be represented by a matrix of the form

$$X = \begin{pmatrix} 1 & x_1 & x_2 & \cdots & z \\ 0 & 1 & 0 & \cdots & y_1 \\ 0 & 0 & 1 & \cdots & y_2 \\ \vdots & & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}$$



# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- 1 If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in V$ ;

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in V$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in V$ ,  $a, b \in \mathbb{R}$ .

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in V$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in V$ ,  $a, b \in \mathbb{R}$ .

**Example** The *dot product* on  $\mathbb{R}^3$ :

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in V$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in V$ ,  $a, b \in \mathbb{R}$ .

**Example** The *dot product* on  $\mathbb{R}^3$ : Let  $x, y \in \mathbb{R}^3$ ; then  $x \cdot y = x^T y$ . The dot product is given by

# Inner Products

An *inner product* on a vector space  $V$  is a non-degenerate, symmetric, bilinear function  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in V$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in V$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in V$ ,  $a, b \in \mathbb{R}$ .

**Example** The *dot product* on  $\mathbb{R}^3$ : Let  $x, y \in \mathbb{R}^3$ ; then  $x \cdot y = x^T y$ . The dot product is given by

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3$$

# Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.



# Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra  $(\mathfrak{g}, [ , ])$  on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

# Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra  $(\mathfrak{g}, [ , ])$  on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing  $x \in \mathfrak{g}$ ,  $\mathrm{ad}_x = [x, \ ]$  becomes a function from  $\mathfrak{g}$  to itself.

# Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra  $(\mathfrak{g}, [ , ])$  on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing  $x \in \mathfrak{g}$ ,  $\mathrm{ad}_x = [x, \ ]$  becomes a function from  $\mathfrak{g}$  to itself. One can then define a map  $j(x) : \mathfrak{g} \rightarrow \mathfrak{g}$  by the formula

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

for all  $y, z \in \mathfrak{g}$ .

# Adjoint of Adjoint

If you put an inner product on a Lie algebra (rather than just a vector space), then you will have two different types of products on the space.

The *adjoint representation* of a Lie algebra  $(\mathfrak{g}, [ , ])$  on itself is the function defined by

$$\mathrm{ad}_x y = [x, y].$$

Fixing  $x \in \mathfrak{g}$ ,  $\mathrm{ad}_x = [x, \ ]$  becomes a function from  $\mathfrak{g}$  to itself. One can then define a map  $j(x) : \mathfrak{g} \rightarrow \mathfrak{g}$  by the formula

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

for all  $y, z \in \mathfrak{g}$ .

The map  $j(x)$  is called the *adjoint* to  $\mathrm{ad}_x$  with respect to the inner product  $\langle , \rangle$ . (Confusing, I know.)

# $j$ -maps

The equation that we used to define  $j(x)$ ,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

# $j$ -maps

The equation that we used to define  $j(x)$ ,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

# $j$ -maps

The equation that we used to define  $j(x)$ ,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \operatorname{ad}_x y, z \rangle = \langle y, \operatorname{ad}_x^\dagger z \rangle.$$

Now we could define  $j(x) = \operatorname{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$ .

# $j$ -maps

The equation that we used to define  $j(x)$ ,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define  $j(x) = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$ .

The  $j$ -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra  $\mathfrak{g}$ .



# $j$ -maps

The equation that we used to define  $j(x)$ ,

$$\langle [x, y], z \rangle = \langle y, j(x)z \rangle$$

could be re-written to look like

$$\langle \text{ad}_x y, z \rangle = \langle y, \text{ad}_x^\dagger z \rangle.$$

Now we could define  $j(x) = \text{ad}_x^\dagger : \mathfrak{g} \rightarrow \mathfrak{g}$ .

The  $j$ -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra  $\mathfrak{g}$ .

This makes the computation of many complicated geometric objects—such as curvatures—into “simple” calculations in linear algebra!

# Acknowledgements