

# Matrix Computations in Lie Algebras

Alexander Jansing, Chaskin Saroff

Oswego State University  
Department of Mathematics

5 May 2015



# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[ , ] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in \mathfrak{g}$ . and

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in \mathfrak{g}$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in \mathfrak{g}$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in \mathfrak{g}$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④  $[x, y] = -[y, x]$  for all  $x, y \in \mathfrak{g}$ .

# Definition

A *Lie Algebra* is a (finite-dimensional) vector space  $\mathfrak{g}$  together with a bilinear multiplication  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$  that satisfies three properties:

- ①  $[x, x] = 0$  for all  $x \in \mathfrak{g}$ ,
- ②  $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$  for all  $x, y, z \in \mathfrak{g}$ . and
- ③  $[x + y, z] = [x, z] + [y, z]$  and  $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ④  $[x, y] = -[y, x]$  for all  $x, y \in \mathfrak{g}$ .

The product  $[\cdot, \cdot]$  is known as a *Lie bracket* on  $\mathfrak{g}$ .



# skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if  $[ , ]$  is alternating then

## skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if  $[\cdot, \cdot]$  is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

## skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if  $[\cdot, \cdot]$  is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

Conversely, if  $[\cdot, \cdot]$  is skew-symmetric, then  $[x, x] + [x, x] = 0$  implies that  $2[x, x] = 0$ . Now we see that this implies  $[x, x] = 0$  so long as our field is not of characteristic 2, for in those spaces  $2 = 0$  and we can deduce nothing about  $[x, x]$ .

# The Center

The *center* of a Lie Algebra,  $\mathfrak{g}$  is

# The Center

The *center* of a Lie Algebra,  $\mathfrak{g}$  is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

# The Center

The *center* of a Lie Algebra,  $\mathfrak{z}$  is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

Thus  $z$ , is in the center of  $\mathfrak{g}$  if and only if

$$[z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}.$$

The *non-center*,  $\mathfrak{v}$ , is given by  $\mathfrak{v} = \mathfrak{g} - \mathfrak{z}$

# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly  $[\ , \ ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$



# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly  $[ , ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly  $[\ , \ ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly  $[\ , \ ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*.

# A First Example: One-Step Nilpotent Lie Algebras

Let  $\mathfrak{g}$  be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly  $[\ , \ ]$  is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*.

# Two-Step Nilpotent

A Lie Algebra,  $\mathfrak{g}$  is called *two-step nilpotent* if and only if

$$[[x, y], z] = 0 \quad \forall x, y, z \in \mathfrak{g} \text{ and } z \neq 0$$

# Two-Step Nilpotent

A Lie Algebra,  $\mathfrak{g}$  is called *two-step nilpotent* if and only if

$$[[x, y], z] = \mathbf{0} \quad \forall x, y, z \in \mathfrak{g} \text{ and } z \neq \mathbf{0}$$

- 1 Applying the bracket twice to any arbitrary set of three vectors in the Algebra, always returns  $\mathbf{0}$ .

# Two-Step Nilpotent

A Lie Algebra,  $\mathfrak{g}$  is called *two-step nilpotent* if and only if

$$[[x, y], z] = \mathbf{0} \quad \forall x, y, z \in \mathfrak{g} \text{ and } z \neq \mathbf{0}$$

- ① Applying the bracket twice to any arbitrary set of three vectors in the Algebra, always returns  $\mathbf{0}$ .
- ② Applying the bracket once to any arbitrary set of two vectors does not always return  $\mathbf{0}$ .

$\mathbb{R}^3$  as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$



# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ = (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \end{aligned}$$

# $\mathbb{R}^3$ as a non-Abelian Lie Algebra

Let  $x = (x_1, x_2, x_3)$  and  $y = (y_1, y_2, y_3)$ . The *cross product* of  $x$  with  $y$  is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} & (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ &= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \\ &= 0. \end{aligned}$$

## $\mathfrak{h}_3$ : The Heisenberg Algebra

We can construct another example from  $\mathbb{R}^3$  by defining a different Lie Bracket.  
Consider a 3-dimensional vector space with basis vectors  $x, y, z$ .

## $\mathfrak{h}_3$ : The Heisenberg Algebra

We can construct another example from  $\mathbb{R}^3$  by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0.

## $\mathfrak{h}_3$ : The Heisenberg Algebra

We can construct another example from  $\mathbb{R}^3$  by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by  $\mathfrak{h}_3$ .



## $\mathfrak{h}_3$ : The Heisenberg Algebra

We can construct another example from  $\mathbb{R}^3$  by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by  $\mathfrak{h}_3$ .




Figure : The Heisenberg Algebra,  $\mathfrak{h}_3$

## $\mathfrak{h}_3$ : The Heisenberg Algebra

We can construct another example from  $\mathbb{R}^3$  by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors  $x, y, z$ . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by  $\mathfrak{h}_3$ .




Figure : The Heisenberg Algebra,  $\mathfrak{h}_3$

Can anyone tell me if the Heisenberg Algebra is one-step nilpotent, two-step nilpotent, neither, or both?

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in \mathfrak{g}$ ,  $a, b \in \mathbb{R}$ .
- ④  $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in \mathfrak{g}$ ,  $a, b \in \mathbb{R}$ .
- ④  $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in \mathfrak{g}$ ,  $a, b \in \mathbb{R}$ .
- ④  $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

**Example** The *dot product* on  $\mathbb{R}^3$ :



# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in \mathfrak{g}$ ,  $a, b \in \mathbb{R}$ .
- ④  $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

**Example** The *dot product* on  $\mathbb{R}^3$ : Let  $x, y \in \mathbb{R}^3$ ; then  $x \cdot y = x^T y$ . The dot product is given by

# Inner Products

An *inner product* on a vector space  $\mathfrak{g}$  is a non-degenerate, symmetric, bilinear, positive definite function  $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ .

- ① If  $\langle x, y \rangle = 0$  for all  $y \in \mathfrak{g}$ , then  $x$  must be 0;
- ②  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \mathfrak{g}$ ;
- ③  $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$  for all  $x, y, z \in \mathfrak{g}$ ,  $a, b \in \mathbb{R}$ .
- ④  $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

**Example** The *dot product* on  $\mathbb{R}^3$ : Let  $x, y \in \mathbb{R}^3$ ; then  $x \cdot y = x^T y$ . The dot product is given by

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3$$

# $j$ -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map,  $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ .

# $j$ -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map,  $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ .  $j_z$  is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

# $j$ -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map,  $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ .  $j_z$  is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

The  $j$ -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra  $\mathfrak{g}$ .

# $j$ -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map,  $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ .  $j_z$  is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

The  $j$ -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra  $\mathfrak{g}$ .

This makes the computation of many seemingly complicated objects “simple” calculations in linear algebra!

# The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form.

# The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$



# The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

# The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

We will denote the  $k$ th entry of  $L_{ij}$  as  $L_{ij}^k$ .

# The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

We will denote the  $k$ th entry of  $L_{ij}$  as  $L_{ij}^k$ .

$$[e_i, e_i] = \mathbf{0}$$

# The Lie Bracket as a Matrix with Vector entries

Recall that the Lie Bracket is a matrix of vectors.

# The Lie Bracket as a Matrix with Vector entries

Recall that the Lie Bracket is a matrix of vectors.




Figure : The Lie Bracket as a Matrix,  $L$

# The Lie Bracket as a Matrix with Vector entries

Recall that the Lie Bracket is a matrix of vectors.




Figure : The Lie Bracket as a Matrix,  $L$

# The Lie Bracket as a Matrix with Vector entries

Recall that the Lie Bracket is a matrix of vectors.





Figure : The Lie Bracket as a Matrix,  $L$

# The Lie Bracket as a vector of matrices

It can also be represented by a “stack” of matrices



A vector of matrices.

$$L_{ij} = \begin{pmatrix} L_{ij}^1 \\ L_{ij}^2 \\ \vdots \\ L_{ij}^m \end{pmatrix}.$$



$$L_{ij} = \begin{pmatrix} L_{ij}^1 \\ L_{ij}^2 \\ \vdots \\ L_{ij}^m \end{pmatrix}.$$

These  $L_{ij}$  can then be arranged in an  $n \times n$  matrix

$$L = (L_{ij}) = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}.$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?  
 Notice that we can retrieve information about the bases.

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

So it will be enough find a way to write all Lie Brackets as a linear combination of basis brackets.

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with



# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\text{So } [x, y] = [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2]$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2]\end{aligned}$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2]\end{aligned}$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2]\end{aligned}$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\ &= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22}\end{aligned}$$

# Lie Bracket as Linear Combination of Bases

We can write  $[x, y]$  as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let  $x, y \in \mathfrak{g}$  with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\ &= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22}\end{aligned}$$

This means that the Lie Bracket is fully described by the matrix,  $L$



# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about  $[e_i, e_i]$ ?

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about  $[e_i, e_i]$ ?

$$[e_i, e_i] = \mathbf{0}$$

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given  $[e_i, e_j]$ , what do we know about  $[e_j, e_i]$ ?

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given  $[e_i, e_j]$ , what do we know about  $[e_j, e_i]$ ?

$$[e_i, e_j] = -[e_j, e_i]$$



# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Consider the map  $[e_1, ] : \mathfrak{g} \rightarrow \mathfrak{g}$ .

Can anyone see the matrix representation of  $[e_1, ]$ ?

# The Lie Bracket as a Matrix with Vector entries

Let's take another look at  $L$

Consider the Lie Bracket, with bases  $e_1, e_2, \dots, e_n$

Formally,  $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \boxed{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Consider the map  $[e_1, ] : \mathfrak{g} \rightarrow \mathfrak{g}$ .

Can anyone see the matrix representation of  $[e_1, ]$ ?

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example.

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22})$$

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{12} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$



# Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$

That is  $[x, y] = x^T L y$

# Example Abelian $\mathbb{R}^3$

What does Abelian  $\mathbb{R}^3$  look like in this form?

## Example Abelian $\mathbb{R}^3$

What does Abelian  $\mathbb{R}^3$  look like in this form?  
 Recall that a Lie Algebra is Abelian if

# Example Abelian $\mathbb{R}^3$

What does Abelian  $\mathbb{R}^3$  look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

Example Abelian  $\mathbb{R}^3$ 

What does Abelian  $\mathbb{R}^3$  look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

# Example Abelian $\mathbb{R}^3$

What does Abelian  $\mathbb{R}^3$  look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of Abelian  $\mathbb{R}^3$

# Examples of Cross Product over $\mathbb{R}^3$

What does the the cross product on  $\mathbb{R}^3$  look like in this form?

# Examples of Cross Product over $\mathbb{R}^3$

What does the the cross product on  $\mathbb{R}^3$  look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$



Examples of Cross Product over  $\mathbb{R}^3$ 

What does the the cross product on  $\mathbb{R}^3$  look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} 0 & e_3 & -e_2 \\ -e_3 & 0 & e_1 \\ e_2 & -e_1 & 0 \end{pmatrix}$$

# Examples of Cross Product over $\mathbb{R}^3$

What does the the cross product on  $\mathbb{R}^3$  look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

What is the center of  $\mathbb{R}^3$  with the cross product?

# $\mathfrak{h}_3$ 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

# $\mathfrak{h}_3$ 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$  for bases  $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

# $\mathfrak{h}_3$ 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$  for bases  $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

# $\mathfrak{h}_3$ 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$  for bases  $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of  $\mathfrak{h}_3$ ?

# $\mathfrak{h}_3$ 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$  for bases  $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of  $\mathfrak{h}_3$ ?

The center is the span of  $\{e_3\}$ .

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.



# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Recall that  $\langle e_i, e_j \rangle = \langle e_j, e_i \rangle$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite:  $\langle x, x \rangle \geq 0$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And  $\mathbf{0}$  cannot be a basis vector because it is linearly dependent with all vectors.

So  $e_{ii} > 0$



# Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite:  $\langle x, x \rangle \geq 0$  and  $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And  $\mathbf{0}$  cannot be a basis vector because it is linearly dependent with all vectors.

So  $e_{ii} > 0$

# Dot Product on $\mathbb{R}^3$

The Dot Product can be represented in this way.

# Dot Product on $\mathbb{R}^3$

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = \delta_{ij}$$

# Dot Product on $\mathbb{R}^3$

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

Dot Product on  $\mathbb{R}^3$ 

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

$$E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

# Matrix representation of the $j$ -maps

Let  $\mathfrak{n}$  be a 2-step nilpotent Lie Algebra with  $\mathfrak{v}$  as the non-center and  $\mathfrak{z}$  as the center.

Recall that the Lie bracket on a 2-step nilpotent Lie algebra is a bilinear map  $[ , ] : \mathfrak{v} \times \mathfrak{v} \rightarrow \mathfrak{z}$ .

So for every  $z \in \mathfrak{z}$ , one can define a linear transformation  $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$  by the identity

$$\langle y, j_z(x) \rangle_{\mathfrak{v}} = \langle z, [x, y] \rangle_{\mathfrak{z}}.$$

We can then use the same methods as before to construct matrix representations of  $\langle , \rangle_{\mathfrak{v}}$  and  $[ , ]$  to find a matrix representation for  $j_z$ .

# Matrix representation of the $j$ -maps

By linearity, we will know how to construct any  $j$ -map, if we know the  $j$ -maps corresponding to basis vectors of  $\mathfrak{z}$ .

Suppose  $\mathfrak{z} = \text{span}\{z_1, z_2, \dots, z_m\}$ .

Then for any  $z_k$ , the  $j$ -map  $j_{z_k} : \mathfrak{v} \rightarrow \mathfrak{v}$  is given by

$$\begin{aligned}\langle y, j_{z_k}(x) \rangle_{\mathfrak{v}} &= \langle z_k, [x, y] \rangle_{\mathfrak{z}} \\ y^T E(J_{z_k} x) &= z_k^T (x^T L y) \\ y^T (E J_{z_k}) x &= y^T (L^k)^T x\end{aligned}$$

taking advantage of some clever stack-matrix manipulations.

# Matrix representation of the $j$ -maps

Since  $y^T(EJ_{z_k})x = y^T(L^k)^Tx$  for arbitrary  $x$  and  $y$  in  $\mathfrak{v}$ , we deduce that  $EJ_{z_k} = (L^k)^T$ .

Since  $\det(E) \neq 0$ , we may solve for  $J_{z_k}$  to obtain

$$J_{z_k} = E^{-1}(L^k)^T \in \mathbb{R}^{n \times n}.$$

If  $z = \zeta_1 z_1 + \zeta_2 z_2 + \cdots + \zeta_m z_m$ , where  $\zeta_i$  are coefficients of the linear combination of  $z_i$ , then the map  $j_z$  is represented by the matrix

$$J_z = \zeta_1 J_{z_1} + \zeta_2 J_{z_2} + \cdots + \zeta_m J_{z_m}.$$



# The matrix $J$ as a stack

The  $j$ -maps of a 2-step nilpotent Lie algebra can be described by a stack of matrices of the same type as  $L$ .

Indeed, if we let  $J^k = J_{z_k}$ ,  
then

$$J = \begin{pmatrix} J^1 \\ J^2 \\ \vdots \\ J^m \end{pmatrix}$$

is an  $(m \times 1)$  stack of  $(n \times n)$  matrices.

Then for  $z = (\zeta_1, \zeta_2, \dots, \zeta_m)^T \in \mathfrak{z}$ , the map  $j_z$  is represented by

$$J_z = z^T J.$$

The equation  $y^T(EJ_z)x = y^T(z^t L^T)x$  must hold for arbitrary vectors  $x$  and  $y$  in  $\mathfrak{v}$ , implying that

$$EJ_z = z^t L^T.$$

Since  $\det(E) \neq 0$ , it is invertible and

$$J_z = E^{-1}(z^t L^T). \quad (7)$$

This gives us a method to compute the  $j$ -map for any  $z$  using only matrix computations. Furthermore, if  $z = z_k$  is a basis vector of  $\mathfrak{z}$ , then  $z_k^t L^T$  is just  $(L^k)^T$ . Thus, for basis vectors  $z_k$ , the map  $j_{z_k}$  is represented by the matrix

$$J_{z_k} = E^{-1}(L^k)^T.$$

Since the  $j$ -maps are linear, knowing how they act on a basis of  $\mathfrak{z}$  is good enough. Indeed, we should be able to define a stack of  $j$ -maps by

$$J = E^{-1}L^T. \quad (8)$$

The  $j$ -map  $j_z$  *should* then be  $J_z = z^t J$ .

# What we did

Using the programming language, sage, we created a program that would compute the  $j$ -maps of an arbitrary Lie algebra.

# Acknowledgements

Justin Ryan for introducing us to Lie Algebras and providing us with Tex Templates, papers and illustrations.

Jonathan Mckibbin for his illustration of the Heisenberg Algebra.