

Encoding the Bracket - An Overview of Lie Algebras

Alexander Jansing, Chaskin Saroff

Oswego State University
Department of Mathematics

2 May 2015



Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

① $[x, x] = 0$ for all $x \in \mathfrak{g}$,

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[,] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\ , \] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- ① $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- ② $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- ③ $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- ① $[x, y] = -[y, x]$ for all $x, y \in \mathfrak{g}$.

Definition

A *Lie Algebra* is a (finite-dimensional) vector space \mathfrak{g} together with a bilinear multiplication $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ that satisfies three properties:

- 1 $[x, x] = 0$ for all $x \in \mathfrak{g}$,
- 2 $[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$ for all $x, y, z \in \mathfrak{g}$. and
- 3 $[x + y, z] = [x, z] + [y, z]$ and $[\alpha x, y] = [x, \alpha y] = \alpha [x, y] \quad \forall x, y, z \in \mathfrak{g} \text{ and } \alpha \in \mathbb{R}$

Property 1 is known as *alternating*, property 2 is known as the *Jacobi Identity* and property 3 describes *bilinearity*.

Property 1 and 3 imply another property called *skew symmetry*:

- 1 $[x, y] = -[y, x]$ for all $x, y \in \mathfrak{g}$.

The product $[\cdot, \cdot]$ is known as a *Lie bracket* on \mathfrak{g} .

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if $[,]$ is alternating then

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if $[,]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x + y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

skew-symmetry

By bilinearity, every alternating product is also skew-symmetric. Indeed, if $[\cdot, \cdot]$ is alternating then

$$0 = [x + y, x + y] \quad (1)$$

$$= [x + y, x] + [x + y, y] \quad (2)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (3)$$

$$= [x, x] + [y, x] + [x, y] + [y, y] \quad (4)$$

$$0 = [x, y] + [y, x] \quad (5)$$

$$\implies [x, y] = -[y, x] \quad (6)$$

Conversely, if $[\cdot, \cdot]$ is skew-symmetric, then $[x, x] + [x, x] = 0$ implies that $2[x, x] = 0$. Now we see that this implies $[x, x] = 0$ so long as our field is not of characteristic 2, for in those spaces $2 = 0$ and we can deduce nothing about $[x, x]$.

The Center

The *center* of a Lie Algebra, \mathfrak{g} is

The Center

The *center* of a Lie Algebra, \mathfrak{g} is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

The Center

The *center* of a Lie Algebra, \mathfrak{g} is

$$\mathfrak{z} = \{z \in \mathfrak{g} \mid [z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}\}.$$

Thus z , is in the center of \mathfrak{g} if and only if

$$[z, x] = \mathbf{0} \ \forall x \in \mathfrak{g}.$$

The *non-center*, \mathfrak{v} , is given by $\mathfrak{v} = \mathfrak{g} - \mathfrak{z}$

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[,]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[\ , \]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[\cdot, \cdot]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*.

A First Example: One-Step Nilpotent Lie Algebras

Let \mathfrak{g} be any vector space, and define

$$[x, y] = 0 \text{ for all } x, y \in \mathfrak{g}.$$

Clearly $[\cdot, \cdot]$ is alternating:

$$[x, x] = 0 \text{ for all } x \in \mathfrak{g}.$$

And the Jacobi Identity is trivial:

$$\begin{aligned} [[x, y], z] + [[y, z], x] + [[z, x], y] &= [0, z] + [0, x] + [0, y] \\ &= 0 + 0 + 0 = 0. \end{aligned}$$

This Algebra is called *Abelian* or *One-Step Nilpotent*.

Two-Step Nilpotent

A Lie Algebra, \mathfrak{g} is called *two-step nilpotent* if and only if

$$[[x, y], z] = \mathbf{0} \quad \forall x, y, z \in \mathfrak{g} \text{ and } \mathfrak{z} \neq \mathfrak{g}$$

Two-Step Nilpotent

A Lie Algebra, \mathfrak{g} is called *two-step nilpotent* if and only if

$$[[x, y], z] = \mathbf{0} \quad \forall x, y, z \in \mathfrak{g} \text{ and } z \neq \mathbf{0}$$

- ① Applying the bracket twice to any arbitrary set of three vectors in the Algebra, always returns $\mathbf{0}$.

Two-Step Nilpotent

A Lie Algebra, \mathfrak{g} is called *two-step nilpotent* if and only if

$$[[x, y], z] = \mathbf{0} \quad \forall x, y, z \in \mathfrak{g} \text{ and } \mathfrak{z} \neq \mathfrak{g}$$

- 1 Applying the bracket twice to any arbitrary set of three vectors in the Algebra, always returns $\mathbf{0}$.
- 2 Applying the bracket once to any arbitrary set of two vectors does not always return $\mathbf{0}$.

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2 y_3 - x_3 y_2, x_3 y_1 - x_1 y_3, x_1 y_2 - x_2 y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$(x \times y) \times z + (y \times z) \times x + (z \times x) \times y$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ = (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \end{aligned}$$

\mathbb{R}^3 as a non-Abelian Lie Algebra

Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$. The *cross product* of x with y is defined by

$$x \times y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1).$$

The cross product is skew symmetric:

$$x \times y = -y \times x \quad \text{for all } x, y \in \mathbb{R}^3.$$

Using some identities from Calc III, we can show that the cross product satisfies the Jacobi Identity:

$$\begin{aligned} (x \times y) \times z + (y \times z) \times x + (z \times x) \times y \\ &= (x \cdot z)y - (y \cdot z)x + (y \cdot x)z - (z \cdot x)y + (z \cdot y)x - (x \cdot y)z \\ &= 0. \end{aligned}$$

\mathfrak{h}_3 : The Heisenberg Algebra

We can construct another example from \mathbb{R}^3 by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors x, y, z .

\mathfrak{h}_3 : The Heisenberg Algebra

We can construct another example from \mathbb{R}^3 by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0.

\mathfrak{h}_3 : The Heisenberg Algebra

We can construct another example from \mathbb{R}^3 by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .

\mathfrak{h}_3 : The Heisenberg Algebra

We can construct another example from \mathbb{R}^3 by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .




Figure: The Heisenberg Algebra, \mathfrak{h}_3

\mathfrak{h}_3 : The Heisenberg Algebra

We can construct another example from \mathbb{R}^3 by defining a different Lie Bracket. Consider a 3-dimensional vector space with basis vectors x, y, z . Define a Lie bracket on this space by

$$[x, y] = z,$$

with all other brackets equal to 0. This is called the *Heisenberg Algebra*, and is denoted by \mathfrak{h}_3 .




Figure: The Heisenberg Algebra, \mathfrak{h}_3

Can anyone tell me if the Heisenberg Algebra is one-step nilpotent, two-step nilpotent, neither, or both?

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- 1 If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.
- ④ $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.
- ④ $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.
- ④ $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

Example The *dot product* on \mathbb{R}^3 :

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.
- ④ $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

Inner Products

An *inner product* on a vector space \mathfrak{g} is a non-degenerate, symmetric, bilinear, positive definite function $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$.

- ① If $\langle x, y \rangle = 0$ for all $y \in \mathfrak{g}$, then x must be 0;
- ② $\langle x, y \rangle = \langle y, x \rangle$ for all $x, y \in \mathfrak{g}$;
- ③ $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$ for all $x, y, z \in \mathfrak{g}$, $a, b \in \mathbb{R}$.
- ④ $\langle x, x \rangle \geq 0 \ \forall x \in \mathfrak{g}$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

A Lie Algebra with an inner product is called a *Metric Lie Algebra*

Example The *dot product* on \mathbb{R}^3 : Let $x, y \in \mathbb{R}^3$; then $x \cdot y = x^T y$. The dot product is given by

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + x_3 y_3$$

j -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map, $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$.

j -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map, $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$. j_z is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

j -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map, $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$. j_z is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

j -maps

In order to separate the interaction between the center and the non-center of a two-step nilpotent lie algebra, we can define a map, $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$. j_z is defined by the equation

$$\langle [x, y], z \rangle = \langle y, j_z x \rangle$$

The j -maps are extremely useful because they simultaneously encode information about the algebraic structure and geometric structure of the Lie algebra \mathfrak{g} .

This makes the computation of many seemingly complicated objects “simple” calculations in linear algebra!

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form.

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

The Lie Bracket as a Matrix with Vector entries

In order to do math with Lie Algebras on a computer, we would like to represent the Lie Bracket in a matrix form. Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

$$[e_i, e_i] = \mathbf{0}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?
Notice that we can retrieve information about the bases.

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

It's nice to see that we can *encode* the bracket into a matrix representation, but how can we use that encoding as the Lie Bracket “map”?

Notice that we can retrieve information about the bases.

$$[e_1, e_2] = L_{12}$$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

So it will be enough find a way to write all Lie Brackets as a linear combination of basis brackets.

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\text{So } [x, y] = [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2]$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2]\end{aligned}$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2]\end{aligned}$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned}\text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2]\end{aligned}$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned} \text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\ &= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22} \end{aligned}$$

Lie Bracket as Linear Combination of Bases

We can write $[x, y]$ as a linear combination of their component vectors' brackets.

Let's look at an example of a 2 dimensional space

Let $x, y \in \mathfrak{g}$ with

$$x = w_1 e_1 + w_2 e_2$$

$$y = v_1 e_1 + v_2 e_2$$

$$\begin{aligned} \text{So } [x, y] &= [w_1 e_1 + w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1 + v_2 e_2] + [w_2 e_2, v_1 e_1 + v_2 e_2] \\ &= [w_1 e_1, v_1 e_1] + [w_1 e_1, v_2 e_2] + [w_2 e_2, v_1 e_1] + [w_2 e_2, v_2 e_2] \\ &= w_1 v_1 [e_1, e_1] + w_1 v_2 [e_1, e_2] + w_2 v_1 [e_2, e_1] + w_2 v_2 [e_2, e_2] \\ &= w_1 v_1 L_{11} + w_1 v_2 L_{12} + w_2 v_1 L_{21} + w_2 v_2 L_{22} \end{aligned}$$

This means that the Lie Bracket is fully described by the matrix, L

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} L_{11} & L_{12} & \cdots & L_{1n} \\ L_{21} & L_{22} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & L_{nn} \end{pmatrix}$$

What is special about $[e_i, e_i]$?

$$[e_i, e_i] = \mathbf{0}$$

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$?

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ L_{21} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ L_{n1} & L_{n2} & \cdots & \mathbf{0} \end{pmatrix}$$

Given $[e_i, e_j]$, what do we know about $[e_j, e_i]$?

$$[e_i, e_j] = -[e_j, e_i]$$

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \mathbf{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & \mathbf{0} & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & \mathbf{0} \end{pmatrix}$$

Consider the map $[e_1, \cdot] : \mathfrak{g} \rightarrow \mathfrak{g}$.

Can anyone see the matrix representation of $[e_1, \cdot]$?

The Lie Bracket as a Matrix with Vector entries

Let's take another look at L

Consider the Lie Bracket, with bases e_1, e_2, \dots, e_n

Formally, $L_{ij} = [e_i, e_j]$

$$L = \begin{pmatrix} \boxed{0} & L_{12} & \cdots & L_{1n} \\ -L_{12} & 0 & \cdots & L_{2n} \\ \vdots & & \ddots & \vdots \\ -L_{1n} & -L_{2n} & \cdots & 0 \end{pmatrix}$$

Consider the map $[e_1,] : \mathfrak{g} \rightarrow \mathfrak{g}$.

Can anyone see the matrix representation of $[e_1,]$?

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example.

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$

Generalizing the Lie Bracket

As before, we'll investigate the 2 dimensional example. Let

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix},$$

with

$$x = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}, \quad y = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$x^T L = (w_1 \ w_2) \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22})$$

$$(x^T L)y = (w_1 L_{11} + w_2 L_{12}, w_1 L_{21} + w_2 L_{22}) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$$

$$= (w_1 v_1 L_{11} + w_2 v_1 L_{12} + w_1 v_2 L_{21} + w_2 v_2 L_{22})$$

That is $[x, y] = x^T L y$

Example Abelian \mathbb{R}^3

What does Abelian \mathbb{R}^3 look like in this form?

Example Abelian \mathbb{R}^3

What does Abelian \mathbb{R}^3 look like in this form?
Recall that a Lie Algebra is Abelian if

Example Abelian \mathbb{R}^3

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

Example Abelian \mathbb{R}^3

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Example Abelian \mathbb{R}^3

What does Abelian \mathbb{R}^3 look like in this form?

Recall that a Lie Algebra is Abelian if

$$[x, y] = \mathbf{0} \quad \forall x, y \in \mathbb{R}^3$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of Abelian \mathbb{R}^3

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

Examples of Cross Product over \mathbb{R}^3

What does the the cross product on \mathbb{R}^3 look like in this form?

$$[e_1, e_2] = e_3$$

$$[e_1, e_3] = -e_2$$

$$[e_2, e_3] = e_1$$

$$\begin{pmatrix} \mathbf{0} & e_3 & -e_2 \\ -e_3 & \mathbf{0} & e_1 \\ e_2 & -e_1 & \mathbf{0} \end{pmatrix}$$

What is the center of \mathbb{R}^3 with the cross product?

\mathfrak{h}_3 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

\mathfrak{h}_3 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$ for bases $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

\mathfrak{h}_3 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$ for bases $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

\mathfrak{h}_3 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$ for bases $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ?

\mathfrak{h}_3 's Lie Bracket

How could the three dimensional Heisenberg Algebra be represented?

$[e_1, e_2] = e_3$ for bases $e_1, e_2, e_3 \in \mathfrak{h}_3$

And all other brackets zero

$$\begin{pmatrix} 0 & e_3 & 0 \\ -e_3 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

What is the center of \mathfrak{h}_3 ?

The center is the span of $\{e_3\}$.

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Recall that $\langle e_i, e_j \rangle = \langle e_j, e_i \rangle$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{21} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{n1} & E_{n2} & \cdots & E_{nn} \end{pmatrix}$$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And $\mathbf{0}$ cannot be a basis vector because it is linearly dependent with all vectors.

So $e_{ij} > 0$

Inner Product as a Matrix

The Inner Product can be represented by a matrix very similarly to the Lie Bracket.

$$\langle e_i, e_j \rangle = E_{ij}$$

$$E = \begin{pmatrix} E_{11} & E_{12} & \cdots & E_{1n} \\ E_{12} & E_{22} & \cdots & E_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ E_{1n} & E_{2n} & \cdots & E_{nn} \end{pmatrix}$$

Recall also that the inner product has additional properties.

Postive Definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0 \implies x = \mathbf{0}$

And $\mathbf{0}$ cannot be a basis vector because it is linearly dependent with all vectors.

So $e_{ii} > 0$

Dot Product on \mathbb{R}^3

The Dot Product can be represented in this way.

Dot Product on \mathbb{R}^3

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

Dot Product on \mathbb{R}^3

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

Dot Product on \mathbb{R}^3

The Dot Product can be represented in this way.

$$\langle e_i, e_j \rangle = 0$$

$$\langle e_i, e_i \rangle = 1$$

$$E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

j -map as a matrix

Recall that for any $z \in \mathfrak{z}$, the map $j_z : \mathfrak{v} \rightarrow \mathfrak{v}$ is defined by the identity

$$\langle y, j_z(x) \rangle_{\mathfrak{v}} = \langle z, [x, y] \rangle_{\mathfrak{z}}. \quad (7)$$

We wish to find a matrix J_z that represents the map j_z . Since j_z maps \mathfrak{v} to itself, J_z should be a matrix in $\mathbb{R}^{n \times n}$. Rewriting (7) in matrix form, we get

$$\begin{aligned} \langle y, j_z(x) \rangle_{\mathfrak{v}} &= \langle z, [x, y] \rangle_{\mathfrak{z}} \\ y^T E(J_z x) &= z^t (x^T L y) \\ y^T (E J_z) x &= z^t (-y^T L x) \\ &= -z^t (y^T L x) \\ &= -y^T (z^t L) x \\ &= y^T (z^t (-L)) x \\ &= y^T (z^t L^T) x. \end{aligned}$$

The equation $y^T(EJ_z)x = y^T(z^{\mathfrak{t}}L^T)x$ must hold for arbitrary vectors x and y in \mathfrak{v} , implying that

$$EJ_z = z^{\mathfrak{t}}L^T.$$

Since $\det(E) \neq 0$, it is invertible and

$$J_z = E^{-1}(z^{\mathfrak{t}}L^T). \quad (8)$$

This gives us a method to compute the j -map for any z using only matrix computations. Furthermore, if $z = z_k$ is a basis vector of \mathfrak{z} , then $z_k^{\mathfrak{t}}L^T$ is just $(L^k)^T$. Thus, for basis vectors z_k , the map j_{z_k} is represented by the matrix

$$J_{z_k} = E^{-1}(L^k)^T.$$

Since the j -maps are linear, knowing how they act on a basis of \mathfrak{z} is good enough. Indeed, we should be able to define a stack of j -maps by

$$J = E^{-1}L^T. \quad (9)$$

The j -map j_z *should* then be $J_z = z^\dagger J$.

What we did

Using the programming language, sage, we created a program that would compute the j -maps of an arbitrary Lie algebra.

TikZ

Acknowledgements




Figure: The L-Stack




Figure: The L-Stack