

Axiomatic Potentialism

Chris Scambler

All Souls College,
Oxford University

chris.scambler@all-souls.ox.ac.uk

January 19, 2022

Overview

- 1 Background
- 2 Warm Up: Height Potentialism
- 3 Height and Width Potentialism

Table of Contents

1 Background

2 Warm Up: Height Potentialism

3 Height and Width Potentialism

Potentialism

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.
- Ideas like this have deep roots in set theory, e.g. Zermelo and even Cantor

Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.
- Ideas like this have deep roots in set theory, e.g. Zermelo and even Cantor
- Still deeper roots in mathematics in general.

- The recent literature has seen two branches of study here:

- The recent literature has seen two branches of study here:
 - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.

- The recent literature has seen two branches of study here:
 - 1 Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
 - 2 Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.

- The recent literature has seen two branches of study here:
 - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
 - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.

- The recent literature has seen two branches of study here:
 - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
 - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.
- The model-theoretic side has seen thorough examination in the recent literature

- The recent literature has seen two branches of study here:
 - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
 - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.
- The model-theoretic side has seen thorough examination in the recent literature
- Here we will be focused on axiomatic potentialism, and on relations between potentialist axiom systems and their first order counterparts.

Table of Contents

1 Background

2 Warm Up: Height Potentialism

3 Height and Width Potentialism

Motivation

Motivation

- Imagine one has the ability to take things and make a set containing them.

Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is able to do this arbitrarily many times.

Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is able to do this arbitrarily many times.
- Axiomatize this conception and work out its non-modal counterpart.

The Language \mathcal{L}_0

- object variables x, y, z

The Language \mathcal{L}_0

- object variables x, y, z
- plural variables X, Y, Z

The Language \mathcal{L}_0

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$

The Language \mathcal{L}_0

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$
- \square

The Language \mathcal{L}_0

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$
- \square
- \in

Axioms for the theory L

Logical Axioms

Axioms for the theory L

Logical Axioms

- 1 Free FO logic

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box \forall X \Diamond \exists x[Set(x, X)]$

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box \forall X \Diamond \exists x[Set(x, X)]$
- 3 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF (or free S5)
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$
- 5 A modal translation of replacement

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

$$UI = \forall x [\forall y \varphi y \rightarrow \varphi x];$$

Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

$UI = \forall x[\forall y \varphi y \rightarrow \varphi x]$;

Instantiation requires assumption of existence.

Relative Consistency

- In fact ZFC interprets L.

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$
- $t(\Box \varphi)(T) := \forall S \supseteq T [t(\varphi)(S)]$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$
- $t(\Box \varphi)(T) := \forall S \supseteq T [t(\varphi)(S)]$

Theorem

$L \vdash \varphi$ implies $ZFC \vdash t(\varphi)(T)$

Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$. (!)
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := t(x) \in t(X) \subseteq T$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$
- $t(\Box \varphi)(T) := \forall S \supseteq T [t(\varphi)(S)]$

Theorem

$L \vdash \varphi$ implies $ZFC \vdash t(\varphi)(T)$

Note: convention on variables T, S .

Converse Translation

Mirroring theorem

Converse Translation

Mirroring theorem

For φ in \mathcal{L}_\in , let φ^\diamond be the result of prefixing all atomic formulas and existential quantifiers by \diamond (and universal quantifiers by \Box .) Then we have

Converse Translation

Mirroring theorem

For φ in \mathcal{L}_\in , let φ^\diamond be the result of prefixing all atomic formulas and existential quantifiers by \diamond (and universal quantifiers by \Box .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Converse Translation

Mirroring theorem

For φ in \mathcal{L}_\in , let φ^\diamond be the result of prefixing all atomic formulas and existential quantifiers by \diamond (and universal quantifiers by \Box .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$.

Converse Translation

Mirroring theorem

For φ in \mathcal{L}_\in , let φ^\diamond be the result of prefixing all atomic formulas and existential quantifiers by \diamond (and universal quantifiers by \Box .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$.

Linnebo Interpretation Theorem

$L \vdash ZFC^\diamond$.

Converse Translation

Mirroring theorem

For φ in \mathcal{L}_\in , let φ^\diamond be the result of prefixing all atomic formulas and existential quantifiers by \diamond (and universal quantifiers by \Box .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$.

Linnebo Interpretation Theorem

$L \vdash ZFC^\diamond$.

Proof: use mirroring.

Axioms for the theory L

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$
- 5 A modal translation of replacement

Mini-conclusion

Mini-conclusion

Equivalence

We have an exact proof-theoretic equivalence, $L \equiv ZFC$.

Mini-conclusion

Equivalence

We have an exact proof-theoretic equivalence, $L \equiv ZFC$.

Bi-interpretation

In fact, $ZFC \vdash t(\phi^\diamond)(T) \leftrightarrow \phi$;

Mini-conclusion

Equivalence

We have an exact proof-theoretic equivalence, $L \equiv ZFC$.

Bi-interpretation

In fact, $ZFC \vdash t(\phi^\diamond)(T) \leftrightarrow \phi$; and
 $L \vdash Univ(T) \rightarrow (t(\phi)(T)^\diamond \leftrightarrow \phi)$

Mini-conclusion

Equivalence

We have an exact proof-theoretic equivalence, $L \equiv ZFC$.

Bi-interpretation

In fact, $ZFC \vdash t(\phi^\diamond)(T) \leftrightarrow \phi$; and
 $L \vdash Univ(T) \rightarrow (t(\phi)(T)^\diamond \leftrightarrow \phi)$

This is essentially what Button calls ‘near-synonymy’.

Table of Contents

- 1 Background
- 2 Warm Up: Height Potentialism
- 3 Height and Width Potentialism**

Motivation

Motivation

- Imagine one has the ability to take things and make a set containing them.

Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;

Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;
- Or, equivalently, to take some things and add an enumerating function.

Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;
- Or, equivalently, to take some things and add an enumerating function.
- Axiomatize this conception and work out its non-modal counterpart.

The Language \mathcal{L}_1

- object variables x, y, z

The Language \mathcal{L}_1

- object variables x, y, z
- plural variables X, Y, Z

The Language \mathcal{L}_1

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$

The Language \mathcal{L}_1

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$
- $\Box_{\uparrow}, \Box_{\leftarrow}, \Box$

The Language \mathcal{L}_1

- object variables x, y, z
- plural variables X, Y, Z
- $\wedge, \neg, \forall, =$
- $\Box_{\uparrow}, \Box_{\leftarrow}, \Box$
- \in

Axioms for the theory M

Logical Axioms

Axioms for the theory M

Logical Axioms

- 1 Free FO logic

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box\uparrow\varphi$, same for $\Box\leftarrow$.
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box\forall X\Diamond_{\uparrow}\exists x[\text{Set}(x, X)]$

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box\forall X\Diamond_{\uparrow}\exists x[\text{Set}(x, X)]$
- 3 $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box\forall X\Diamond_{\uparrow}\exists x[\text{Set}(x, X)]$
- 3 $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box\forall X\Diamond_{\uparrow}\exists x[\text{Set}(x, X)]$
- 3 $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$
- 5 $\Box\forall X\Diamond_{\leftarrow}\exists f[f : \mathbb{N} \twoheadrightarrow X]$

Axioms for the theory M

Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal (or free S5)
- 3 $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$, same for \Box_{\leftarrow} .
- 4 $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$, rigidity, Choice, Comp

Set-theoretic axioms

- 1 Extensionality, \in -rigidity, foundation
- 2 $\Box\forall X\Diamond_{\uparrow}\exists x[\text{Set}(x, X)]$
- 3 $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4 $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$
- 5 $\Box\forall X\Diamond_{\leftarrow}\exists f[f : \mathbb{N} \twoheadrightarrow X]$
- 6 A modal translation of replacement

Some basic facts

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^{\Diamond\uparrow}$.

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^{\diamond\uparrow}$.
- M interprets ZFC^- under the translation $\varphi \mapsto \varphi^{\diamond}$.

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^{\diamond\uparrow}$.
- M interprets ZFC^- under the translation $\varphi \mapsto \varphi^{\diamond}$.
- M proves $\neg \text{Pow}^{\diamond}$.

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^{\diamond\uparrow}$.
- M interprets ZFC^- under the translation $\varphi \mapsto \varphi^{\diamond}$.
- M proves $\neg Pow^{\diamond}$.
- M proves $V = HC^{\diamond}$ and hence SOA^{\diamond} .

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^\diamond_\uparrow$.
- M interprets ZFC^- under the translation $\varphi \mapsto \varphi^\diamond$.
- M proves $\neg \text{Pow}^\diamond$.
- M proves $V = \text{HC}^\diamond$ and hence SOA^\diamond .
- M proves the universal possibility of forcing.

Some basic facts

- M interprets ZFC under the translation $\varphi \mapsto \varphi^\diamond$.
- M interprets ZFC^- under the translation $\varphi \mapsto \varphi^\diamond$.
- M proves $\neg \text{Pow}^\diamond$.
- M proves $V = \text{HC}^\diamond$ and hence SOA^\diamond .
- M proves the universal possibility of forcing.
- e.g., it proves it is possible for the continuum to exist and have a cardinality at least as great as any \aleph number whose existence is provable in ZFC.

Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond_{\leftarrow} \exists y [y \subseteq x \wedge y = z] \wedge \Box_{\uparrow} \neg \exists y [y = z])$$

Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond_{\leftarrow} \exists y [y \subseteq x \wedge y = z] \wedge \Box_{\uparrow} \neg \exists y [y = z])$$

abbreviate the formula in parentheses by $\Psi(x, z)$.

Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by $\Psi(x, z)$.

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by $\Psi(x, z)$.

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

By height potentialism/rigidity,

$$\Diamond \exists x \Psi(x, z) \wedge \Diamond \uparrow \exists w \forall y [y \in w \leftrightarrow y \in z]$$

Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by $\Psi(x, z)$.

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

By height potentialism/rigidity,

$$\Diamond \exists x \Psi(x, z) \wedge \Diamond \uparrow \exists w \forall y [y \in w \leftrightarrow y \in z]$$

But then the rigidity/extensionality imply $w = z$ after all, so we have a contradiction.

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee z even exists (indeed this is what we are trying to establish.)

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee z even exists (indeed this is what we are trying to establish.)
Natural solution: ‘restrict’ comp to closed form:

$$\Box \forall z \Box \forall Z \exists X \forall y [Xy \leftrightarrow \varphi(y, z, Z)]$$

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee z even exists (indeed this is what we are trying to establish.)
Natural solution: ‘restrict’ comp to closed form:

$$\Box \forall z \Box \forall Z \exists X \forall y [Xy \leftrightarrow \varphi(y, z, Z)]$$

Amounts to ‘restricting’ ourselves to parameters that exist at the world of evaluation.

Relative Consistency

Key Point

It turns out that M exhibits roughly the same relation to second order arithmetic together with the Π_1^1 -*Perfect Set Property* as L did to ZFC.

Background on PSP

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.
- E.g. every closed set has the PSP.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.
- E.g. every closed set has the PSP.
- If the reals can be well-ordered (AC), not all sets have the PSP.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.
- E.g. every closed set has the PSP.
- If the reals can be well-ordered (AC), not all sets have the PSP.
- The Γ -PSP is the assertion that every set of reals defined by a formula of type Γ has the PSP.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.
- E.g. every closed set has the PSP.
- If the reals can be well-ordered (AC), not all sets have the PSP.
- The Γ -PSP is the assertion that every set of reals defined by a formula of type Γ has the PSP.
- The Γ -PSP implies no formula in Γ defines a well-order of the reals.

Background on PSP

- A set of reals is *perfect* iff it is closed and has no isolated points. There are only 2^{\aleph_0} such sets.
- A set of reals is said to *have the perfect set property* iff it is countable, or has a perfect subset.
- E.g. every closed set has the PSP.
- If the reals can be well-ordered (AC), not all sets have the PSP.
- The Γ -PSP is the assertion that every set of reals defined by a formula of type Γ has the PSP.
- The Γ -PSP implies no formula in Γ defines a well-order of the reals.
- Γ -definable sets are thus ‘regular’ or ‘well behaved’.

Relative Consistency

Intuitive idea

Relative Consistency

Intuitive idea

(From now on, I will ignore the difference between SOA and $\text{ZFC}^- + \text{V} = \text{HC}$. Replacement is formulated as collection.)

Relative Consistency

Intuitive idea

(From now on, I will ignore the difference between SOA and $ZFC^- + V = HC$. Replacement is formulated as collection.)

- We will use the fact that $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$, and in fact T proves that $L[r]$ is a model of ZFC for every real r .

Relative Consistency

Intuitive idea

(From now on, I will ignore the difference between SOA and $ZFC^- + V = HC$. Replacement is formulated as collection.)

- We will use the fact that $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$, and in fact T proves that $L[r]$ is a model of ZFC for every real r .
- Our translation will be doubly parameterized, once by a real and once by a transitive set. (These are the 'worlds' in the sense of the interp)

Relative Consistency

Intuitive idea

(From now on, I will ignore the difference between SOA and $ZFC^- + V = HC$. Replacement is formulated as collection.)

- We will use the fact that $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$, and in fact T proves that $L[r]$ is a model of ZFC for every real r .
- Our translation will be doubly parameterized, once by a real and once by a transitive set. (These are the 'worlds' in the sense of the interp)
- Our interpretation for \Diamond_\uparrow will involve holding r fixed and climbing transitive sets in $L[r]$;

Relative Consistency

Intuitive idea

(From now on, I will ignore the difference between SOA and $ZFC^- + V = HC$. Replacement is formulated as collection.)

- We will use the fact that $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$, and in fact T proves that $L[r]$ is a model of ZFC for every real r .
- Our translation will be doubly parameterized, once by a real and once by a transitive set. (These are the 'worlds' in the sense of the interp)
- Our interpretation for \diamond_{\uparrow} will involve holding r fixed and climbing transitive sets in $L[r]$;
- while our interpretation for \diamond_{\leftarrow} will involve allowing new reals to be added but not extending the height of the transitive set parameter.

Relative consistency

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\epsilon, (\varphi, T, r) \mapsto \psi(T, r)$$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_{\uparrow} \varphi)(T, r) := \forall S \supseteq T[S \in L[r] \rightarrow t(\varphi)(S, r)]$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_{\uparrow} \varphi)(T, r) := \forall S \supseteq T[S \in L[r] \rightarrow t(\varphi)(S, r)]$
- $t(\Box_{\leftarrow} \varphi)(T, r) := \forall s[r \in L[s] \rightarrow \forall S \in L[s][T \subseteq S \wedge \text{rank}(S) = \text{rank}(T) \rightarrow t(\varphi)(S, s)]]$

Relative consistency

Let $M \models SOA + \Pi_1^1PSP$.

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign singular/plural variables even/odd numbered variables $t(x), t(X)$.
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := t(x) \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_{\uparrow} \varphi)(T, r) := \forall S \supseteq T[S \in L[r] \rightarrow t(\varphi)(S, r)]$
- $t(\Box_{\leftarrow} \varphi)(T, r) := \forall s[r \in L[s] \rightarrow \forall S \in L[s][T \subseteq S \wedge rank(S) = rank(T) \rightarrow t(\varphi)(S, s)]]$

Theorem

$M \vdash \varphi$ implies $T \vdash \forall r, T \in L[r][t(\varphi)(X, r)]$

Converse Translation

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Solovay: If only countably many reals are constructible from any r , then the Π_1^1 -PSP holds.

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Solovay: If only countably many reals are constructible from any r , then the Π_1^1 -PSP holds.

By mirroring, suffices to establish that (only countably many reals are constructible from r) $^\diamond$ holds.

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Solovay: If only countably many reals are constructible from any r , then the Π_1^1 -PSP holds.

By mirroring, suffices to establish that (only countably many reals are constructible from r) $^\diamond$ holds.

Since we have $ZFC^{\diamond\uparrow}$, we get that (there is a set of all reals constructible from r) $^{\diamond\uparrow}$.

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Solovay: If only countably many reals are constructible from any r , then the Π_1^1 -PSP holds.

By mirroring, suffices to establish that (only countably many reals are constructible from r) $^\diamond$ holds.

Since we have $ZFC^{\diamond\uparrow}$, we get that (there is a set of all reals constructible from r) $^{\diamond\uparrow}$.

Some absoluteness lemmas imply (there is a set of all reals constructible from r) $^\diamond$.

Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

Proof (sketch)

Solovay: If only countably many reals are constructible from any r , then the Π_1^1 -PSP holds.

By mirroring, suffices to establish that (only countably many reals are constructible from r) $^\diamond$ holds.

Since we have $ZFC^{\diamond\uparrow}$, we get that (there is a set of all reals constructible from r) $^{\diamond\uparrow}$.

Some absoluteness lemmas imply (there is a set of all reals constructible from r) $^\diamond$.

Since (all sets are countable) $^\diamond$, the result follows.

Interpretations

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$.

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$.
The latter is in fact equiconsistent with ZFC. $M \equiv L \equiv ZFC$.

Interpretations

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$. The latter is in fact equiconsistent with ZFC. $M \equiv L \equiv ZFC$.

Bi-interpretation

As before, $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(X, r) \leftrightarrow \phi$.

Interpretations

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$.
The latter is in fact equiconsistent with ZFC. $M \equiv L \equiv ZFC$.

Bi-interpretation

As before, $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(X, r) \leftrightarrow \phi$.

We also have a partial converse:

for the language without \diamondleftarrow , $M \vdash Univ(T) \rightarrow (t(\phi)(T)^\diamond \leftrightarrow \phi)$

Interpretations

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$. The latter is in fact equiconsistent with ZFC. $M \equiv L \equiv ZFC$.

Bi-interpretation

As before, $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(X, r) \leftrightarrow \phi$.

We also have a partial converse:

for the language without \diamondleftarrow , $M \vdash Univ(T) \rightarrow (t(\phi)(T)^\diamond \leftrightarrow \phi)$

Thus we have at least ‘near’-near synonymy in this case.

Interpretations

Equivalence

We thus have an exact proof-theoretic equivalence, $M \equiv SOA + \Pi_1^1PSP$. The latter is in fact equiconsistent with ZFC. $M \equiv L \equiv ZFC$.

Bi-interpretation

As before, $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(X, r) \leftrightarrow \phi$.

We also have a partial converse:

for the language without \diamond_{\leftarrow} , $M \vdash Univ(T) \rightarrow (t(\phi)(T)^\diamond \leftrightarrow \phi)$

Thus we have at least ‘near’-near synonymy in this case.

The exact details of \diamond_{\leftarrow} are incidental anyway; if we formulate the theory with just \diamond and \diamond_{\uparrow} , we will get near synonymy.

Concluding remarks

Concluding remarks

- The 'purely quantificational picture' corresponding to the height + width potentialist view enshrined in M seems to be second order arithmetic + Π_1^1 PSP.

Concluding remarks

- The 'purely quantificational picture' corresponding to the height + width potentialist view enshrined in M seems to be second order arithmetic + Π_1^1 PSP.
- What happens if we strengthen the regularity properties?

Concluding remarks

- The 'purely quantificational picture' corresponding to the height + width potentialist view enshrined in M seems to be second order arithmetic + Π_1^1 PSP.
- What happens if we strengthen the regularity properties?
- You get more large cardinals in inner models.

Concluding remarks

- The 'purely quantificational picture' corresponding to the height + width potentialist view enshrined in M seems to be second order arithmetic + Π_1^1 PSP.
- What happens if we strengthen the regularity properties?
- You get more large cardinals in inner models.
- In the modal theory, this corresponds to the \Diamond_{\uparrow} -possibility of large cardinals.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.
- Such theories are strong in terms of consistency strength, and can interpret substantial portions of the traditional large cardinal hierarchy.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.
- Such theories are strong in terms of consistency strength, and can interpret substantial portions of the traditional large cardinal hierarchy.
- But they are also ontologically parsimonious, in only admitting countable sets, and offer hope of certain benefits in applied mathematics – e.g. the possibility of universal lebesgue measurability.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.
- Such theories are strong in terms of consistency strength, and can interpret substantial portions of the traditional large cardinal hierarchy.
- But they are also ontologically parsimonious, in only admitting countable sets, and offer hope of certain benefits in applied mathematics – e.g. the possibility of universal lebesgue measurability.
- Traditionally, the stronger regularity properties are believed to hold only in inner models, while the large cardinals are in the ‘real’ universe of sets.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.
- Such theories are strong in terms of consistency strength, and can interpret substantial portions of the traditional large cardinal hierarchy.
- But they are also ontologically parsimonious, in only admitting countable sets, and offer hope of certain benefits in applied mathematics – e.g. the possibility of universal lebesgue measurability.
- Traditionally, the stronger regularity properties are believed to hold only in inner models, while the large cardinals are in the ‘real’ universe of sets.
- But I am not convinced that there are decisive reasons for taking one of these, rather than the other, to hold only in inner models.

Concluding remarks

- These results have led me to consider second order arithmetic + regularity properties as a possible ‘foundation for mathematics’.
- Such theories are strong in terms of consistency strength, and can interpret substantial portions of the traditional large cardinal hierarchy.
- But they are also ontologically parsimonious, in only admitting countable sets, and offer hope of certain benefits in applied mathematics – e.g. the possibility of universal lebesgue measurability.
- Traditionally, the stronger regularity properties are believed to hold only in inner models, while the large cardinals are in the ‘real’ universe of sets.
- But I am not convinced that there are decisive reasons for taking one of these, rather than the other, to hold only in inner models.
- This corresponds, as we have effectively seen in this talk, to not thinking there are decisive reasons to favor pure height over height-and-width potentialism.

Thanks!