

# Axiomatic Potentialism

Chris Scambler

All Souls College,  
Oxford University

*chris.scambler@all-souls.ox.ac.uk*

November 24, 2021

# Overview

- 1 Background
- 2 Warm Up: Height Potentialism
- 3 Height and Width Potentialism
- 4 Concluding Remarks

# Table of Contents

1 Background

2 Warm Up: Height Potentialism

3 Height and Width Potentialism

4 Concluding Remarks

# Potentialism

## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct

## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble

## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.



## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.
- Ideas like this have deep roots in set theory, e.g. Zermelo and even Cantor

## Potentialism

is the idea that a mathematical object (e.g. a set) is the sort of thing that may *merely possibly* exist.

- E.g. a geometric object as a figure one can construct
- A set as a certain sort of data structure one could assemble
- Or perhaps a structure that is instantiated given enough objects.
- Ideas like this have deep roots in set theory, e.g. Zermelo and even Cantor
- Still deeper roots in mathematics in general.

- The recent literature has seen two branches of study here:

- The recent literature has seen two branches of study here:
  - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.

- The recent literature has seen two branches of study here:
  - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
  - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.

# Potentialism

- The recent literature has seen two branches of study here:
  - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
  - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.

# Potentialism

- The recent literature has seen two branches of study here:
  - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
  - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.
- E.g. Hamkins and Linnebo showed in MT potentialism with the structures initial segments of  $V$  that  $S5$  at a world  $V_\kappa$  is equivalent to  $\Sigma_3$  correctness of  $\kappa$ .

# Potentialism

- The recent literature has seen two branches of study here:
  - ① Model-theoretic: study Kripke models whose worlds are structures with the accessibility relation (some refinement of) the substructure relation.
  - ② Axiomatic: Develop axiom systems designed to characterize this or that form of potentialism directly, without appeal to models.
- In each case interesting questions arise concerning the relation between assertions in the modal framework and in first order set theory.
- E.g. Hamkins and Linnebo showed in MT potentialism with the structures initial segments of  $V$  that  $S5$  at a world  $V_\kappa$  is equivalent to  $\Sigma_3$  correctness of  $\kappa$ .
- Here we will be focused on axiomatic potentialism, and on relations between potentialist axiom systems and their first order counterparts.



# Table of Contents

- 1 Background
- 2 Warm Up: Height Potentialism**
- 3 Height and Width Potentialism
- 4 Concluding Remarks

# Motivation

# Motivation

- Imagine one has the ability to take things and make a set containing them.

# Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is able to do this arbitrarily many times.

# Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is able to do this arbitrarily many times.
- Axiomatize this conception and work out its non-modal counterpart.

## The Language $\mathcal{L}_0$

- object variables  $x, y, z$

## The Language $\mathcal{L}_0$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$

## The Language $\mathcal{L}_0$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$



## The Language $\mathcal{L}_0$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$
- $\square$

## The Language $\mathcal{L}_0$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$
- $\square$
- $\in$

# Axioms for the theory L

## Logical Axioms

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation



# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box \forall X \Diamond \exists x[Set(x, X)]$

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box \forall X \Diamond \exists x[Set(x, X)]$
- 3  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$
- 5 A modal translation of replacement

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$



# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

$$UI = \forall x [\forall y \varphi y \rightarrow \varphi x];$$

# Inconsistency?

Standard modal model theory validates the rule

$$\frac{\varphi \rightarrow \Box\psi}{\varphi \rightarrow \Box\forall x\psi}$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \forall y \Box \neg \text{Set}(y, X) \quad (1)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \neg \text{Set}(y, X) \quad (2)$$

$$(Xx \leftrightarrow x \notin x) \rightarrow \Box \forall y \neg \text{Set}(y, X) \quad (3)$$

Hence the need for free logic.

$UI = \forall x [\forall y \varphi y \rightarrow \varphi x]$ ;

Instantiation requires assumption of existence.

# Relative Consistency

- In fact ZFC interprets L.

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := x \in t(X)$



# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := x \in t(X)$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := x \in t(X)$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := x \in t(X)$
- $t(\forall x\varphi)(T) := \forall x \in T[t(\varphi)(T)]$
- $t(\forall X\varphi)(T) := \forall x \subseteq T[t(\varphi)(T)]$
- $t(\Box\varphi)(t) := \forall S \supseteq T[Tran(S) \rightarrow t(\varphi)(S)]$

# Relative Consistency

- In fact ZFC interprets L.

$$t : \mathcal{L}_0 \times V \rightarrow \mathcal{L}_\infty, (\varphi, T) \mapsto \psi(T)$$

- assign plural variables odd numbered variables  $t(X)$ .
- membership claims = id, commutes with propositional connectives
- $t(Xx)(T) := x \in t(X)$
- $t(\forall x \varphi)(T) := \forall x \in T [t(\varphi)(T)]$
- $t(\forall X \varphi)(T) := \forall x \subseteq T [t(\varphi)(T)]$
- $t(\Box \varphi)(t) := \forall S \supseteq T [Tran(S) \rightarrow t(\varphi)(S)]$

## Theorem

$L \vdash \varphi$  implies  $ZFC \vdash t(\varphi)(\emptyset)$

# Converse Translation

## Mirroring theorem

# Converse Translation

## Mirroring theorem

For  $\varphi$  in  $\mathcal{L}_\in$ , let  $\varphi^\diamond$  be the result of prefixing all universal quantifiers by a  $\Box$  (and existential quantifiers by  $\Diamond$ .) Then we have

# Converse Translation

## Mirroring theorem

For  $\varphi$  in  $\mathcal{L}_\in$ , let  $\varphi^\diamond$  be the result of prefixing all universal quantifiers by a  $\Box$  (and existential quantifiers by  $\Diamond$ .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$



# Converse Translation

## Mirroring theorem

For  $\varphi$  in  $\mathcal{L}_\in$ , let  $\varphi^\diamond$  be the result of prefixing all universal quantifiers by a  $\Box$  (and existential quantifiers by  $\Diamond$ .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$ .

# Converse Translation

## Mirroring theorem

For  $\varphi$  in  $\mathcal{L}_\in$ , let  $\varphi^\diamond$  be the result of prefixing all universal quantifiers by a  $\Box$  (and existential quantifiers by  $\Diamond$ .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$ .

## Linnebo Interpretation Theorem

$L \vdash ZFC^\diamond$ .

# Converse Translation

## Mirroring theorem

For  $\varphi$  in  $\mathcal{L}_\in$ , let  $\varphi^\diamond$  be the result of prefixing all universal quantifiers by a  $\Box$  (and existential quantifiers by  $\Diamond$ .) Then we have

$$\Gamma \vdash_{FOL} \varphi \Leftrightarrow \Gamma^\diamond \vdash_L \varphi^\diamond$$

Note on replacement $^\diamond$ .

## Linnebo Interpretation Theorem

$L \vdash ZFC^\diamond$ .

Proof: use mirroring.

# Axioms for the theory L

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF
- 3  $\forall z[Xz \leftrightarrow Yz] \rightarrow X = Y$ , rigidity, Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box \forall X \Diamond \exists x[\text{Set}(x, X)]$
- 3  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4  $\Diamond \exists X \Box \forall x[Xx \leftrightarrow x \subseteq y]$
- 5 A modal translation of replacement

# Mini-conclusion

# Mini-conclusion

## Equivalence

We have an exact proof-theoretic equivalence,  $L \equiv ZFC$ .

# Mini-conclusion

## Equivalence

We have an exact proof-theoretic equivalence,  $L \equiv ZFC$ .

## Bi-interpretation

In fact,  $ZFC \vdash t(\phi^\diamond)(\emptyset) \leftrightarrow \phi$ , and conversely.

# Table of Contents

- 1 Background
- 2 Warm Up: Height Potentialism
- 3 Height and Width Potentialism**
- 4 Concluding Remarks



# Motivation

# Motivation

- Imagine one has the ability to take things and make a set containing them.

# Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;

# Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;
- Or, equivalently, to take some things and add an enumerating function.

# Motivation

- Imagine one has the ability to take things and make a set containing them.
- Imagine one is also able to take a partial order and add a filter meeting all its (current) dense sets;
- Or, equivalently, to take some things and add an enumerating function.
- Axiomatize this conception and work out its non-modal counterpart.

## The Language $\mathcal{L}_1$

- object variables  $x, y, z$

## The Language $\mathcal{L}_1$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$

## The Language $\mathcal{L}_1$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$



## The Language $\mathcal{L}_1$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$
- $\Box_{\uparrow}, \Box_{\leftarrow}, \Box$

## The Language $\mathcal{L}_1$

- object variables  $x, y, z$
- plural variables  $X, Y, Z$
- $\wedge, \neg, \forall, =$
- $\Box_{\uparrow}, \Box_{\leftarrow}, \Box$
- $\in$

# Axioms for the theory M

## Logical Axioms

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

# Axioms for the theory M

## Logical Axioms

- ① Free FO logic
- ② S4.2 modal logic + CBF for each modal
- ③  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- ④ Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms



# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box\forall X\Diamond_{\uparrow}\exists x[Set(x, X)]$

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box\forall X\Diamond_{\uparrow}\exists x[Set(x, X)]$
- 3  $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow Nx]$

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box\forall X\Diamond_{\uparrow}\exists x[Set(x, X)]$
- 3  $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4  $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box\forall X\Diamond_{\uparrow}\exists x[Set(x, X)]$
- 3  $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow \mathbb{N}x]$
- 4  $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$
- 5  $\Box\forall\mathbb{P}, X[D(\mathbb{P}, X) \rightarrow \Diamond_{\leftarrow}\exists g[Fmeets(g, xx)]]$

# Axioms for the theory M

## Logical Axioms

- 1 Free FO logic
- 2 S4.2 modal logic + CBF for each modal
- 3  $\Box\varphi \rightarrow \Box_{\uparrow}\varphi$ , same for  $\Box_{\leftarrow}$ .
- 4 Ext for  $X$ ,  $\Diamond Xx \rightarrow \Box Xx$ ,  $\Diamond\exists x[Xx \wedge x = y] \rightarrow \exists x[Xx \wedge x = y]$ ,  
Choice, Comp

## Set-theoretic axioms

- 1 Extensionality,  $\in$ -rigidity, foundation
- 2  $\Box\forall X\Diamond_{\uparrow}\exists x[Set(x, X)]$
- 3  $\Diamond_{\uparrow}\exists X\Box\forall x[Xx \leftrightarrow Nx]$
- 4  $\Diamond_{\uparrow}\exists X\Box_{\uparrow}\forall x[Xx \leftrightarrow x \subseteq y]$
- 5  $\Box\forall\mathbb{P}, X[D(\mathbb{P}, X) \rightarrow \Diamond_{\leftarrow}\exists g[Fmeets(g, xx)]]$
- 6 A modal translation of replacement

# Some basic facts

# Some basic facts

- $M$  interprets ZFC under the translation  $\varphi \mapsto \varphi^{\diamond\uparrow}$ .



# Some basic facts

- $M$  interprets ZFC under the translation  $\varphi \mapsto \varphi^{\diamond\uparrow}$ .
- $M$  interprets  $\text{ZFC}^-$  under the translation  $\varphi \mapsto \varphi^{\diamond}$ .

# Some basic facts

- $M$  interprets ZFC under the translation  $\varphi \mapsto \varphi^{\diamond\uparrow}$ .
- $M$  interprets  $\text{ZFC}^-$  under the translation  $\varphi \mapsto \varphi^{\diamond}$ .
- $M$  proves  $\neg \text{Pow}^{\diamond}$ .

# Some basic facts

- $M$  interprets ZFC under the translation  $\varphi \mapsto \varphi^\diamond \uparrow$ .
- $M$  interprets  $ZFC^-$  under the translation  $\varphi \mapsto \varphi^\diamond$ .
- $M$  proves  $\neg Pow^\diamond$ .
- $M$  proves  $V = HC^\diamond$  and hence  $SOA^\diamond$ .

# Some basic facts

- $M$  interprets ZFC under the translation  $\varphi \mapsto \varphi^{\diamond\uparrow}$ .
- $M$  interprets  $\text{ZFC}^-$  under the translation  $\varphi \mapsto \varphi^{\diamond}$ .
- $M$  proves  $\neg \text{Pow}^{\diamond}$ .
- $M$  proves  $V = \text{HC}^{\diamond}$  and hence  $\text{SOA}^{\diamond}$ .
- $M$  proves it is possible for the continuum to exist and have a cardinality at least as great as any  $\aleph$  number whose existence is provable in ZFC.

# Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond_{\leftarrow} \exists y [y \subseteq x \wedge y = z] \wedge \Box_{\uparrow} \neg \exists y [y = z])$$

# Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by  $\Psi(x, z)$ .

# Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by  $\Psi(x, z)$ .

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

# Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by  $\Psi(x, z)$ .

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

By height potentialism/rigidity,

$$\Diamond \exists x \Psi(x, z) \wedge \Diamond \uparrow \exists w \forall y [y \in w \leftrightarrow y \in z]$$



# Inconsistency?

The axioms imply

$$\Diamond \exists x (\Diamond \leftarrow \exists y [y \subseteq x \wedge y = z] \wedge \Box \uparrow \neg \exists y [y = z])$$

abbreviate the formula in parentheses by  $\Psi(x, z)$ .

By comprehension,

$$\Diamond \exists x \Psi(x, z) \wedge \exists X \forall y [Xy \leftrightarrow y \in z]$$

By height potentialism/rigidity,

$$\Diamond \exists x \Psi(x, z) \wedge \Diamond \uparrow \exists w \forall y [y \in w \leftrightarrow y \in z]$$

But then the rigidity/extensionality imply  $w = z$  after all, so we have a contradiction.

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee  $z$  even exists (indeed this is what we are trying to establish.)

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee  $z$  even exists (indeed this is what we are trying to establish.)  
Natural solution: restrict comp to closed form:

$$\Box \forall z \Box \forall Z \exists X \forall y [Xy \leftrightarrow \varphi(y, z, Z)]$$

The argument just sketched uses comprehension with arbitrary parameters:

$$\exists X \forall y [Xy \leftrightarrow y \in z]$$

And in the crucial application, it applies when we have no *a priori* guarantee  $z$  even exists (indeed this is what we are trying to establish.)  
Natural solution: restrict comp to closed form:

$$\Box \forall z \Box \forall Z \exists X \forall y [Xy \leftrightarrow \varphi(y, z, Z)]$$

Amounts to restricting ourselves to parameters that exist at the world of evaluation.

# Relative Consistency

Intuitive idea

# Relative Consistency

## Intuitive idea

(From now on, I will ignore the difference between  $\text{SOA}$  and  $\text{ZFC}^- + \text{V} = \text{HC}$ . Replacement is formulated as collection.)

# Relative Consistency

## Intuitive idea

(From now on, I will ignore the difference between SOA and  $ZFC^- + V = HC$ . Replacement is formulated as collection.)

- We will use the fact that  $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$ , and in fact  $T$  proves that  $L[r]$  is a model of ZFC for every real  $r$ .



# Relative Consistency

## Intuitive idea

(From now on, I will ignore the difference between SOA and  $ZFC^- + V = HC$ . Replacement is formulated as collection.)

- We will use the fact that  $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$ , and in fact  $T$  proves that  $L[r]$  is a model of ZFC for every real  $r$ .
- Our translation will be doubly parameterized, once by a real and once by a transitive set.

# Relative Consistency

## Intuitive idea

(From now on, I will ignore the difference between SOA and  $ZFC^- + V = HC$ . Replacement is formulated as collection.)

- We will use the fact that  $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$ , and in fact  $T$  proves that  $L[r]$  is a model of ZFC for every real  $r$ .
- Our translation will be doubly parameterized, once by a real and once by a transitive set.
- Our interpretation for  $\Diamond_\uparrow$  will involve holding  $r$  fixed and climbing transitive sets in  $L[r]$ ;

# Relative Consistency

## Intuitive idea

(From now on, I will ignore the difference between SOA and  $ZFC^- + V = HC$ . Replacement is formulated as collection.)

- We will use the fact that  $T = SOA + \Pi_1^1\text{-PSP} \equiv ZFC$ , and in fact  $T$  proves that  $L[r]$  is a model of ZFC for every real  $r$ .
- Our translation will be doubly parameterized, once by a real and once by a transitive set.
- Our interpretation for  $\diamond_{\uparrow}$  will involve holding  $r$  fixed and climbing transitive sets in  $L[r]$ ;
- while our interpretation for  $\diamond_{\leftarrow}$  will involve allowing new reals to be added but not extending the height of the transitive set parameter.

# Relative consistency

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives



# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$
- $t(\forall x\varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X\varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_\uparrow\varphi)(T, r) := \forall S \supseteq T[Tran(S) \wedge S \in L[r] \rightarrow t(\varphi)(S, r)]$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$
- $t(\forall x\varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X\varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_{\uparrow}\varphi)(T, r) := \forall S \supseteq T[Tran(S) \wedge S \in L[r] \rightarrow t(\varphi)(S, r)]$
- $t(\Box_{\leftarrow}\varphi)(T, r) := \forall s[r \in L[s] \rightarrow \forall S[S \leq T \wedge S \in L[s] \rightarrow t(\varphi)(S, s)]]$

# Relative consistency

Let  $M \models SOA + \Pi_1^1PSP$ .

$t : \mathcal{L}_1 \times M \times \mathbb{R}^M \rightarrow \mathcal{L}_\infty, (\varphi, T, r) \mapsto \psi(T, r)$

- assign plural variables odd numbered variables  $t(X)$ .
- Membership = id, commutes with propositional connectives
- $t(Xx)(T, r) := x \in t(X)$
- $t(\forall x \varphi)(T, r) := \forall x \in T[t(\varphi)(T, r)]$
- $t(\forall X \varphi)(T, r) := \forall x \subseteq T[x \in L[r] \rightarrow t(\varphi)(T, r)]$
- $t(\Box_{\uparrow} \varphi)(T, r) := \forall S \supseteq T[Tran(S) \wedge S \in L[r] \rightarrow t(\varphi)(S, r)]$
- $t(\Box_{\leftarrow} \varphi)(T, r) := \forall s[r \in L[s] \rightarrow \forall S[S \leq T \wedge S \in L[s] \rightarrow t(\varphi)(S, s)]]$

## Theorem

$M \vdash \varphi$  implies  $T \vdash t(\varphi)(\emptyset, 0)$

# Converse Translation

# Converse Translation

Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$



# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas.

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas. One can also show by forcing that “ $\diamond \mathbb{R}^{L[r]}$  is countable.”

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas. One can also show by forcing that “ $\diamond \mathbb{R}^{L[r]}$  is countable.” This yields the  $\diamond$ -translation of “ $\mathbb{R}^{L[r]}$  exists for every  $r$ , and is countable.”

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas. One can also show by forcing that “ $\diamond \mathbb{R}^{L[r]}$  is countable.” This yields the  $\diamond$ -translation of “ $\mathbb{R}^{L[r]}$  exists for every  $r$ , and is countable.” By a result of Solovay, if only countably many reals are constructible from  $r$ , then the  $\Pi_1^1[r]$  PSP holds.

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas. One can also show by forcing that “ $\diamond \mathbb{R}^{L[r]}$  is countable.” This yields the  $\diamond$ -translation of “ $\mathbb{R}^{L[r]}$  exists for every  $r$ , and is countable.” By a result of Solovay, if only countably many reals are constructible from  $r$ , then the  $\Pi_1^1[r]$  PSP holds. Hence by mirroring  $\Pi_1^1[r] PSP^\diamond$ .

# Converse Translation

## Theorem

$$M \vdash \Pi_1^1 PSP^\diamond$$

## Proof (sketch)

Given any possible real  $r$ , one can show  $\diamond \exists x [x = \mathbb{R}^{L[r]}]$  using the  $\diamond_\uparrow$  translation of ZFC and some absoluteness lemmas. One can also show by forcing that “ $\diamond \mathbb{R}^{L[r]}$  is countable.” This yields the  $\diamond$ -translation of “ $\mathbb{R}^{L[r]}$  exists for every  $r$ , and is countable.” By a result of Solovay, if only countably many reals are constructible from  $r$ , then the  $\Pi_1^1[r]$  PSP holds. Hence by mirroring  $\Pi_1^1[r] PSP^\diamond$ . But  $r$  is arbitrary.

# Table of Contents

- 1 Background
- 2 Warm Up: Height Potentialism
- 3 Height and Width Potentialism
- 4 Concluding Remarks**



# Conclusions

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .  
The latter is in fact equiconsistent with ZFC.  $M \equiv L \equiv ZFC$ .

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .  
The latter is in fact equiconsistent with ZFC.  $M \equiv L \equiv ZFC$ .

## Bi-interpretation

However,  $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(\emptyset, 0) \leftrightarrow \phi$ , and conversely.

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .  
The latter is in fact equiconsistent with ZFC.  $M \equiv L \equiv ZFC$ .

## Bi-interpretation

However,  $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(\emptyset, 0) \leftrightarrow \phi$ , and conversely.

## Philosophical Interest?

There seems to be some relation between height + width potentialism and topological regularity.

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .  
The latter is in fact equiconsistent with ZFC.  $M \equiv L \equiv ZFC$ .

## Bi-interpretation

However,  $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(\emptyset, 0) \leftrightarrow \phi$ , and conversely.

## Philosophical Interest?

There seems to be some relation between height + width potentialism and topological regularity.

The 'first order picture' corresponding to the height + width potentialist view may be second order arithmetic + topological regularity.

# Conclusions

## Equivalence

We have an exact proof-theoretic equivalence,  $M \equiv SOA + \Pi_1^1PSP$ .  
The latter is in fact equiconsistent with ZFC.  $M \equiv L \equiv ZFC$ .

## Bi-interpretation

However,  $SOA + \Pi_1^1PSP \vdash t(\phi^\diamond)(\emptyset, 0) \leftrightarrow \phi$ , and conversely.

## Philosophical Interest?

There seems to be some relation between height + width potentialism and topological regularity.

The 'first order picture' corresponding to the height + width potentialist view may be second order arithmetic + topological regularity.

What happens with stronger regularity properties? How does PD/AD affect things?

# Thanks!