

Monic arrows in Ω -set .. An arrow $f : A \rightarrow B$ in Ω -set is monic iff it satisfies

$$f(a_0, b) \wedge f(a_1, b) \leq \llbracket a_0 = a_1 \rrbracket_A \quad (1)$$

Proof. Let $g : C \rightarrow A$ in Ω -set.

We first prove that (1) implies

$$g(c, a_0) = \bigsqcup_{b \in B} g \circ f(c, b) \wedge f(a_0, b) \quad (2)$$

for any c, a_0 . To do this we show each of

$$g(c, a_0) \leq \bigsqcup_{b \in B} g \circ f(c, b) \wedge f(a_0, b) \quad (3)$$

and

$$g(c, a_0) \geq \bigsqcup_{b \in B} g \circ f(c, b) \wedge f(a_0, b) \quad (4)$$

For (3):

$$g(c, a_0) = g(c, a_0) \wedge \llbracket a_0 = a_0 \rrbracket \quad (5)$$

$$= \bigsqcup_{b \in B} g(c, a_0) \wedge f(a_0, b) \quad (6)$$

$$= \bigsqcup_{b \in B} (\bigsqcup_{a \in A} g(c, a) \wedge f(a, b)) \wedge f(a_0, b) \wedge g(c, a_0) \quad (7)$$

$$= \bigsqcup_{b \in B} f \circ g(c, b) \wedge f(a_0, b) \wedge g(c, a_0) \quad (8)$$

which latter is what we want.

For (4), let $b \in B$. We have:

$$(\bigsqcup_{a \in A} g(c, a) \wedge f(a, b)) \wedge f(a_0, b) = \bigsqcup_{a \in A} g(c, a) \wedge f(a, b) \wedge f(a_0, b) \quad (9)$$

$$\leq \bigsqcup_{a \in A} g(c, a) \wedge \llbracket a = a_0 \rrbracket_A \quad (10)$$

$$\leq g(c, a_0) \quad (11)$$

where the transition from (9) to (10) uses (1), and the transition from (10) to (11) uses (v) p 277. (Is equality true?)

From here it is easy. Suppose $g \circ f(c, b) = h \circ f(c, b)$ for all c and b . Then for any a , $\bigsqcup_{b \in B} g \circ f(c, b) \wedge f(a, b) = \bigsqcup_{b \in B} h \circ f(c, b) \wedge f(a, b)$. Hence $g(c, a) = f(c, a)$ by (2). \square

Ω axiom in Ω -set .. Show that with $\Omega(p, q) := p \Leftrightarrow q$ and $\top(0, p) = p$ we get the Ω axiom for

$$\chi_f = \llbracket Ed \rrbracket \wedge \llbracket s_f(d) = p \rrbracket_{\Omega} \quad (12)$$

Proof. Let $f : \mathbf{A} \rightarrow \mathbf{D}$ be monic. First, it will be handy to have some identities. Note that

$$\top \circ ! (a, 0) = \bigsqcup_{x \in 1} \llbracket Ea \rrbracket \wedge p = \llbracket Ea \rrbracket \wedge p \quad (13)$$

Second, since $s_f(d) = \bigsqcup_{a \in A} f(a, d)$ (12) implies

$$\chi_f(d, p) = \bigsqcup_{a \in A} f(a, d) \Leftrightarrow p \quad (14)$$

which then implies

$$\chi_f \circ f(a, p) = \bigsqcup_{d \in D} f(a, d) \wedge \bigsqcup_{a' \in A} f(a', d) \Leftrightarrow p \quad (15)$$

putting (13), (15) together, it suffices show

$$\llbracket Ea \rrbracket \wedge p = \bigsqcup_{d \in D} f(a, d) \wedge \bigsqcup_{a' \in A} f(a', d) \Leftrightarrow p \quad (16)$$

to show the square commutes.

It is immediate that

$$\bigsqcup_{d \in D} f(a, d) \wedge \bigsqcup_{a' \in A} f(a', d) \Leftrightarrow p = \bigsqcup_{d \in D} f(a, d) \wedge f(a, d) \Leftrightarrow p \quad (17)$$

by algebra (lol) we get

$$\bigsqcup_{d \in D} f(a, d) \wedge f(a, d) \Leftrightarrow p = \bigsqcup_{d \in D} f(a, d) \wedge p \quad (18)$$

but putting (17) and (18) together implies (16) by vii on p 277.

□