

Homework problem set 1

Please hand in your solutions to these exercises in digital form (typed, or scanned from a neatly hand-written version) through Moodle no later than **Friday June 16, 20:00h**.

Problem 1: Purity

Purity of a quantum state is defined as $\text{Tr}\rho^2$. Consider a d -dimensional quantum state $\rho \in \mathbb{C}^{d \times d}$.

- What is the maximal value of purity, and what states achieve this value?
- What is the minimal value of purity, what state achieves this value?
- Any qubit density matrix can be represented by the Bloch vector \vec{r} , satisfying $|\vec{r}| \leq 1$. For any quantum state $\tau \in \mathbb{C}^{2 \times 2}$ we have that $\tau = \frac{1}{2}(\mathbb{1} + \vec{r} \cdot \vec{\sigma})$, where $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)^T$ is the vector of Pauli matrices. How does purity of τ relate to \vec{r} ?

Problem 2: Minimum-error measurement of qubits

Alice is given one of the states $|\theta\rangle := \cos\left(\frac{\theta}{2}\right)|0\rangle + \sin\left(\frac{\theta}{2}\right)|1\rangle$, $|\theta\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle - \sin\left(\frac{\theta}{2}\right)|1\rangle$, each with probability $p := \frac{1}{2}$. Design a measurement $\{\Pi_+, \Pi_-\}$, where $\Pi_+, \Pi_- \succeq 0$ and $\Pi_+ + \Pi_- = \mathbb{1}$, such that the probability of error is minimal. The probability of error is defined as $p_e := p\text{Tr}(\Pi_-|\theta\rangle\langle\theta|) + (1-p)\text{Tr}(\Pi_+|-\theta\rangle\langle-\theta|)$.

Problem 3: Parity measurements (Exercise 1.5.1)

Use a projective measurement to measure the parity, in the Hadamard basis, of the state $|00\rangle\langle 00|$. Compute the probabilities of obtaining measurement outcomes "even" and "odd", and the resulting post-measurement states. What would the post-measurement states have

been if you had first measured the qubits individually in the Hadamard basis, and then taken the parity?

Problem 4: Partial trace (Exercise 1.6.1)

Verify that the state $\rho_A = \sum_x (\mathbb{1} \otimes \langle u_x |) \rho_{AB} (\mathbb{1} \otimes |u_x\rangle)$ does not depend on the choice of basis $\{|u_x\rangle\}$. [Hint: first argue that if two density matrices ρ, σ satisfy $\langle \phi | \rho | \phi \rangle = \langle \phi | \sigma | \phi \rangle$ for all unit vectors $|\phi\rangle$ then $\rho = \sigma$. Then compute $\langle \phi | \rho_A | \phi \rangle$, and use the POVM condition $\sum_x M_x = \mathbb{1}$ to check that you can get an expression independent of $\{|u_x\rangle\}$. Conclude that ρ_A itself does not depend on $\{|u_x\rangle\}$.]

Problem 5: Robustness of GHZ and W states (Problem 6.1)

- Now we generalize to the N -qubit case. As you might expect, $|GHZ_N\rangle = \frac{1}{\sqrt{2}}(|0\rangle^{\otimes N} + |1\rangle^{\otimes N})$.
What is the overlap $\text{Tr}\left(|GHZ_{N-1}\rangle\langle GHZ_{N-1}| \text{tr}_N(|GHZ_N\rangle\langle GHZ_N|)\right)$ in the limit $N \rightarrow \infty$?
- $|W_N\rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^N \underbrace{|0 \cdots 0 1 0 \cdots 0\rangle}_{\text{1 at the } i\text{-th place}}$ is an equal superposition of all N -bit strings with exactly one 1 and $N-1$ 0's

What is the overlap $\text{Tr}\left(|W_{N-1}\rangle\langle W_{N-1}| \text{tr}_N(|W_N\rangle\langle W_N|)\right)$ in the limit $N \rightarrow \infty$?

Problem 6: Relation between min-entropy and ignorance

Let K be a classical (key) register, and let E be Eve's quantum register. Prove the following statement for arbitrary classical-quantum states ρ_{KE} : Eve is ignorant about K if and only if $H_{\min}(K|E) = \log|K|$.