

Finite-Center Accountability and the Positive Route to Binary Rank

Live review note of June 20, 2026

Abstract

Two finite-center obstruction notes show that quotient-visible operational, sector, modular, recovery, continuum, or entropy data do not by themselves reconstruct hidden finite central rank or the predicate $q = 2$. This short note states the matching positive criterion. For a supplied finite center $\mathcal{Z}_F \simeq \ell^\infty(F)$, let $V_F = \mathcal{Z}_F/\mathcal{C}1$ be its contrast space. A declared family of probes, effects, outcome statistics, sector labels, recovery tasks, modular records, finite-alphabet laws, or direct measurements generates a readout map $R : V_F \rightarrow W$. The supplied data determine exactly the quotient $V_F/\ker R$. Hence, in a comparison class where the hidden finite center may vary and q is not supplied as an external label, full finite central cardinality is recoverable from the readout data only when R is faithful on the physical finite-center contrast space and an admissibility or maximality premise says this represented center is the whole physical finite center. Under those premises $q = \text{rank } R + 1$. A binary conclusion $q = 2$ additionally needs one accounted contrast plus a theorem or measurement identifying that contrast as the full finite center rather than a quotient of a larger hidden extension.

This is deliberately demotion-ready. In finite statistical language it is a Blackwell/sufficiency statement. In operator-algebraic language it is a Petz-style relative-to-the-supplied-channel boundary. The possible value is only the translation into black-hole/AQFT reconstruction: recovery, entropy, sector, modular, or locally covariant records force q or $q = 2$ only after finite-center accountability has been supplied.

Status. This is live non-frozen review metadata. It does not modify either frozen Team 1 paper, Lean archive, public mirror manifest, or SHA-256 package. Its purpose is external classification: known theorem, false statement, missing hypothesis, too broad physically, or useful expository corollary.

1 Question

Positive reconstruction theorems reconstruct the object supplied to them: a represented von Neumann algebra, a locally covariant functor, a sector category, a correctable code

algebra, a channel tested by specified preparations and effects, or a Type II algebra with trace and normalization data. The finite-center obstruction is the converse question. If the displayed data have already passed through a quotient or coarse readout, what extra premise is needed before hidden finite central rank, and in particular the binary predicate $q = 2$, becomes reconstructible?

The word “reconstructible” is used throughout in the fiber sense: a target must be a function of the displayed readout data on the stated comparison class. There is no obstruction if q , the full finite center, or a complete atomic measurement is simply supplied as part of the input. The obstruction is only to recovering that target from records that have already forgotten some finite central directions.

2 Finite Accountability

Definition 1 (finite-center contrast and readout). Let F be a finite set with $q = |F|$, and let

$$\mathcal{Z}_F = \ell^\infty(F), \quad V_F = \mathcal{Z}_F / \mathbb{C}1.$$

Thus $\dim V_F = q - 1$. A finite-center readout is a linear map

$$R : V_F \longrightarrow W$$

generated by the declared observation family: normal probes, effects, outcome statistics, sector records, recovery tasks, modular records, a finite-alphabet law, or a direct measurement channel. Its null directions are

$$N_R = \ker R.$$

Theorem 1 (finite accountability quotient). *Let D_R be any data record whose dependence on finite-center contrasts factors through the readout map $R : V_F \rightarrow W$. Work on a comparison class in which the hidden finite center may be refined or extended inside directions not seen by R , and in which q is not separately supplied as a label. Then:*

1. D_R factors through V_F / N_R .
2. No invariant of the finite center that varies inside a coset of N_R is reconstructible from D_R .
3. The full finite central cardinality q is reconstructible from D_R on that comparison class only after adding both $N_R = 0$ and an admissibility or maximality premise saying that the represented finite center is the whole physical finite center, not a quotient, null sector, gauge reduction, excluded sector, or hidden extension.
4. Under those added premises,

$$q = \dim(V_F / N_R) + 1 = \text{rank } R + 1.$$

Proof. If two contrasts $v, v' \in V_F$ have the same image under R , then $v - v' \in N_R$. Every record built from R assigns the same value to v and v' , so it factors through V_F/N_R . Consequently any target that changes within a coset of N_R is not a function of the supplied data.

The finite commutative algebra $\ell^\infty(F)$ has one atomic coordinate for each element of F . After constants are quotiented out, its contrast space has dimension $q - 1$. If $N_R = 0$, the readout is faithful on that contrast space, so it determines $\dim V_F$. If the admissibility premise also says the represented finite center is physically exhaustive, this dimension is the physical finite-center contrast dimension, and $q = \dim V_F + 1$. If $N_R \neq 0$, or if the represented center is only a quotient or admissible subcenter of a larger hidden center, refinements inside the null or hidden directions change the unaccounted finite cardinality while preserving all R -visible data. $\square$

Remark 1 (supplied labels). The theorem does not deny that a laboratory, model definition, or microscopic postulate may hand over q directly. In that case q is an input label, not a reconstruction from quotient-visible readout data. The Team 1 boundary concerns black-hole/AQFT arguments where entropy, recovery, sector, modular, or locally covariant records are used to infer a finite alphabet that has not already been supplied.

Corollary 1 (binary rank requires two inputs). *A conclusion $q = 2$ from finite-center readout data requires both:*

$$\dim(V_F/N_R) = 1$$

and a maximality, admissibility, direct-measurement, or finite-alphabet theorem identifying that accounted one-dimensional contrast as the full physical finite center. One accounted contrast alone proves only that the declared experiment sees a binary quotient.

3 Relation to Standard Sufficiency

The theorem should be demoted if a standard source already states this exact translation. Blackwell comparison of experiments makes sufficiency relative to a declared experiment or decision problem [1, 2]. Petz sufficiency makes recovery and equality of relative entropy relative to a declared subalgebra or channel [3, 4]. Von Neumann algebra ideal theory says ultraweakly closed two-sided ideals are central-projection summands [5, 6]. None of these facts is new.

The possible finite-center contribution is the reconstruction boundary: black-hole/AQFT records determine a finite central cardinality only when their hypotheses include faithful finite-center accountability. Without that premise, quotient-visible entropy, recovery, sector, modular, or locally covariant records reconstruct the supplied quotient, not hidden cardinality.

4 Claim Boundary for Reviewers

Claim	Status
R determines $V_F/\ker R$.	Elementary finite-dimensional linear algebra.
R determines full q when $\ker R = 0$ and the represented center is physically exhaustive.	Positive accountability criterion.
One accounted central contrast implies $q = 2$.	False without maximality, admissibility, direct measurement, or a finite-alphabet theorem.
Blackwell/Petz sufficiency may already contain the criterion.	Reviewer should identify the exact theorem if so; Team 1 should demote.
AQFT/OAQEC hypotheses may already supply maximality or faithful support.	Reviewer should name the hypothesis and decide whether it is present in the Team 1 statement.

5 Reviewer Question

Is Theorem 1 merely finite Blackwell/Petz sufficiency under a different name, or does the black-hole/AQFT translation isolate a useful claim boundary? If it is standard, the desired response is the citation and the exact theorem name. If it is false or too broad, the desired response is a counterexample or missing physical hypothesis.

References

- [1] D. Blackwell, Comparison of experiments, in *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, University of California Press, 1951, 93–102.
- [2] D. Blackwell, Equivalent comparisons of experiments, *Annals of Mathematical Statistics* **24** (1953), 265–272.
- [3] D. Petz, Sufficient subalgebras and the relative entropy of states of a von Neumann algebra, *Communications in Mathematical Physics* **105** (1986), 123–131.
- [4] D. Petz, Sufficiency of channels over von Neumann algebras, *Quarterly Journal of Mathematics* **39** (1988), 97–108.
- [5] J. Lurie, *Math 261y: von Neumann Algebras*, Lecture 9, Proposition 1.
- [6] C. Houdayer, *An Invitation to von Neumann Algebras*, lecture notes, Exercise 3.7.
- [7] R. Brunetti, K. Fredenhagen, and R. Verch, The generally covariant locality principle: a new paradigm for local quantum field theory, *Communications in Mathematical Physics* **237** (2003), 31–68.

- [8] C. J. Fewster and R. Verch, Dynamical locality and covariance: what makes a physical theory the same in all spacetimes?, *Annales Henri Poincare* **13** (2012), 1613–1674.
- [9] D. L. Jafferis, A. Lewkowycz, J. Maldacena, and S. J. Suh, Relative entropy equals bulk relative entropy, *Journal of High Energy Physics* **2016**, 004 (2016).