

# Herbal Medicine in the Management of Liver Diseases

Luk Bahadur Chetry <sup>1\*</sup>, Moromi Engtipi <sup>2</sup>

<sup>1</sup> Department of Zoology, Jagannath Barooah University, Jorhat-785001, Assam, India

<sup>2</sup> Department of Life Science, Assam University, Diphu Campus, Diphu- 782462, Assam, India

<sup>1\*</sup> Corresponding author: [luk.chetry@jbu.ac.in](mailto:luk.chetry@jbu.ac.in)

## Abstract

*Liver diseases remain a global health challenge, with conventional drugs often limited in efficacy and sometimes caused hepatotoxicity. Herbal medicines offer promising alternatives due to their safety, potency and cost-effectiveness. Phytoconstituents such as flavonoids, polyphenols, glycosides and essential oils have demonstrated the hepatoprotective effects by reducing oxidative stress, enhancing antioxidant defenses and regulating liver enzyme activity. The major medicinal plants with reported hepatoprotective potential include C. sinensis, A. paniculata, S. marianum, Z. officinale, C. pepo, C. reticulata and P. crispum. The therapeutic efficacy of herbal formulations often arises from synergistic interactions among multiple constituents. Advanced techniques like high-performance liquid chromatography and microdialysis aid in characterizing these compounds, while understanding their pharmacokinetics is crucial for optimizing dosage and safety. Plant-based biomolecules thus represent a promising source for the development of effective hepatoprotective drugs.*

**Keywords:** Liver, herbal medicine, liver diseases, medicinal plants, hepatoprotective drugs.

## 1. Introduction

Currently, phytochemicals from natural resources are being adopted as alternative treatment options to treat several acute to chronic diseases. Despite considerable advancements in modern medicine, a shortage of proper and potent hepatoprotective agents has remained a chronic concern (Hasan et al., 2018). Many conventional medications have been removed from the market due to reports of drug-induced liver injury, and serious cases of liver dysfunction can require liver transplantation or lead to death (Au et al., 2011). Modern drugs provide little relief for liver disorders are primarily treated using plant-based formulations. Herbal drugs have grown in importance and popularity in recent years due to their potency, purity and cost-effectiveness. Therefore, an attempt has been made to address herbal medicine for the treatment of liver diseases, which are widely used for the therapeutic purpose. A number of chemical constituents, such as phenols, organic acids, lignans, monoterpenes, alkaloids, flavonoids, glycosides, coumarins, lipids, and xanthenes, have been identified in hepatoprotective plants (Bhawna and Kumar, 2010). A wide range of plants are thought to have hepatoprotective efficacy, suggesting that the advancement of plant-based hepatoprotective drugs on the global market should be prioritized (Abu et al., 2017).

Across the world, liver disorders, such as jaundice, cirrhosis, fatty liver, hepatitis, drug-induced hepatitis, alcohol-induced hepatitis, non-alcoholic steatohepatitis and hepatocellular carcinoma, create conditions in which the functions and structure of the liver become impaired, and liver failure is the fifth most common cause of death globally, behind only heart failure, stroke, chest disease and cancer, making liver disease a public health issue (Azab and Albasha, 2018). The worldwide incidence of chronic liver disease is 18.5% and the prevalence of cirrhosis is 4.5%–9.5%, with 2 million people dying every year (Islam and

Alam, 2019). According to the World Health Organization (WHO), deaths due to liver disease in India reached 268,580 in 2020, accounting for 3.17% of total deaths. Chronic liver diseases, including cirrhosis, were the eighth leading cause of death in 2022, a significant increase from previous years (Swaroop et al., 2024). Hepatic disorder is associated with the disruption of the liver structures, tissues and function, which may be caused by biological origins, including bacteria, viruses, parasites and autoimmune diseases, induced by various conventional drugs acetaminophen, nimesulide and hepatotoxins, which are primarily metabolized by the enzymes cytochrome P450 and glutathione S-acyltransferases mixed oxidases; and the excessive consumption of alcohol (Kumar et al., 2011; Madrigal-santill'an et al., 2014; Mishra et al., 2014). The present study discusses major medicinal plants with significant hepatoprotective properties and their contributions to the management of liver diseases.

## 2. Methodology

Different types of electronic bibliographic databases such as Web of Science Master Journal List, SpringerLink, ScienceDirect, Google Scholar, PubMed, Scopus, Wiley Online Library, Taylor & Francis, ResearchGate, and NISCAIR Online Periodicals Repository were extensively searched to obtain scientific information on “hepatoprotective traditional medicinal plants” by applying specific keywords. The keywords used included “plants against hepatoprotection,” “herbal medicine for liver problems,” “phytotherapy for liver disease,” and “plants and hepatoprotective effects.” The research articles were downloaded from various scientific journals and thoroughly examined for their relevance, authenticity, data quality and validity.

## 3. Medicinal Plants for Hepatic Disorders

### 3.1 *Zingiber officinale*

*Z. officinale* (Zingiberaceae) is a widely used medicinal plant around the world, commonly incorporated into foods as a spice. Traditionally, it has been an important ingredient in Chinese, Ayurvedic, and Unani herbal medicines for the treatment of various ailments such as catarrh, rheumatism, nervous disorders, gingivitis, toothache, asthma, stroke, constipation, and diabetes (Awang, 1992; Tapsell et al., 2006; Wang and Wang, 2005). The pharmacological effects of ginger have been reported to include antiplatelet, antioxidant, antitumor, anti-rhinoviral, anti-hepatotoxic, anti-arthritis, and anti-inflammatory activities (Lantz et al., 2007; Patel et al., 2000). The antioxidant activity of gingerol and other constituents of ginger has also been confirmed (Aeschbach et al., 1994). It has been reported that different doses of ginger extract cause alterations in biochemical parameters, free radicals, and antioxidant enzymes induced by bromobenzene in a rat model, thereby improving bromobenzene-induced liver toxicity (El-Sharakly et al., 2009). Curcumin, another active component present in ginger, has been found to act as an antioxidant and anti-inflammatory agent, inducing haem-oxygenase-1 and protected endothelial cells against oxidative stress (Motterlini et al., 2000). Furthermore, ginger root extract has shown hepatoprotective effects against aspartame-induced hepatotoxicity by decreasing liver function biomarkers (ALT, AST, ALP,  $\gamma$ -GT), serum total protein, albumin, and total bilirubin levels, while increasing antioxidant enzyme levels and reducing malondialdehyde concentrations (Hozayen and Abou Seif, 2014).

### 3.2 *Cucurbita pepo* L.

Pumpkin is frequently reported as a functional food or medicine (Caili et al., 2006). Pumpkin seeds are a rich source of unsaturated fatty acids, antioxidants, and fiber, and are known to have anti-atherogenic, hepatoprotective (Makni et al., 2008), and antidiabetic activities (Al-Zuhair et al., 2000). Phytochemical analyses have revealed that pumpkin is rich in unsaturated fatty acids particularly linoleic and oleic acids as well as tocopherols with very high oxidative stability (Stevenson et al., 2007), carotenoids, and gamma-aminobutyric acid (GABA) in the fruit (Liu et al., 2001; Gossell-Williams et al., 2008). Pumpkin seed oil contains nutrients such as essential fatty acids (omega-6 and omega-9), phytosterols, and antioxidants including carotenoids, vitamin A, and vitamin E (Murkovic et al., 1996). Moreover, pumpkin oil may play an important role in protecting against alcohol and Tramadol-induced hepatotoxicity and oxidative stress (Abou Seif, 2014a; Ekpono et al., 2024).

### 3.3 *Citrus reticulata*

*C. reticulata*, also known as mandarin, is a small citrus tree with fruits resembling other orange varieties. Chemically, the plant is rich in vitamin C, flavonoids, organic acids, volatile oils, and rutin (Wang et al., 2008; Luo et al., 2008). In humans, rutin exhibits more potent antioxidant properties compared to quercetin, acacetin, morin, hispidulin, hesperidin and naringin (Chow et al., 2005; Zhou et al., 2016), which are mostly isolated from citrus peel. Hesperidin is another flavanone glycoside found abundantly in citrus fruits that showed an antioxidant, anti-inflammatory, anticarcinogenic activities in vitro study (Farombi et al., 2008; Emim et al., 1994; Tanaka et al., 2000). Both hesperidin and rutin plays an important role in the protection against doxorubicin-induced hepatotoxicity, by alleviating the activities of liver enzymes (ALT, AST, and ALP and GGT) in addition to the amelioration in the levels of total bilirubin, albumin and sialic acid (Hozayen et al., 2014; Elhemiel et al., 2025).

### 3.4 *Petroselinum crispum* oil

The plant Parsley (Apiaceae) is a native herb of the central Mediterranean region (southern Italy, Algeria, and Tunisia) (Lopez et al., 1999). Various scientific studies have demonstrated its antibacterial and antioxidant activities (Wong and Kitts, 2006; Zhang et al., 2006; Kolarovic et al., 2010). Phytochemical analyses have shown that parsley contains several flavonoids, such as apiin and luteolin, while its essential oil is rich in apiol and myristicin (Alobaidi, 2024; Farzaei et al., 2013; Fusani et al., 2025). Parsley oil plays an important role in improving liver function by modulating enzyme activity, enhancing antioxidation and anti-lipid peroxidation, promoting detoxification, and protecting against glutathione depletion in cases of alcohol-induced hepatotoxicity and oxidative stress (Abou Seif, 2014b).

### 3.5 *Andrographis paniculata*

*A. paniculata* (Acanthaceae) is one of the most commonly used plants in the traditional systems of Unani and Ayurvedic medicine, and it is widely known as the king of bitters (Jarukamjorn and Nemoto, 2008). *A. paniculata* has been reported to possess antibacterial (Mishra et al., 2013), antimalarial (Dua et al., 2004), antiviral (Wiert et al., 2005), cardioprotective, antioxidant, anti-inflammatory (Sheeja et al., 2006), antidiabetic, and antitumor activities (Zhao et al., 2008). Treatment with whole-plant extracts of *A. paniculata* has been shown to reduce lipid peroxidation levels and enhance antioxidant enzyme activity, likely due to the presence of various flavonoids, phenols, and glycosides in the plant (Subramaniam et al., 2015).

### 3.6 *Silybum marianum*

Silymarin, derived from the seeds of *S. marianum*, is a member of the sunflower family and is commonly called milk thistle. The plant has been used for centuries as a natural remedy for liver and biliary tract diseases (Abou Seif, 2016). The active biomolecules of milk thistle are flavonolignans, including silybin, silydianin, and silychristin, collectively known as silymarin (Shaarawy et al., 2009). The pharmacological activity of silymarin has been reported to protect liver cells from a wide variety of toxins, including acetaminophen, ethanol, CCl<sub>4</sub> and D-galactosamine (Rasool et al., 2014). Its hepatoprotective efficacy is attributed to several mechanisms, including antioxidation, anti-lipid peroxidation, enhanced detoxification, and protection against glutathione depletion (Pradhan and Girish, 2006). *Silybum* seeds also contain betaine and essential fatty acids, which may contribute to silymarins' anti-inflammatory effect (Saller et al., 2001).

### 3.7 *Camellia sinensis*

Green tea leaves produce organic compounds that may be involved in the defense of plants against invading pathogens, and these metabolites are known as polyphenols (Friedman, 2007), which include catechin, epicatechin, epigallocatechin, tannins and caffeine (Wang and Goodman, 1999). Green tea displays antioxidants and free radical scavenger properties (Crespy and Williamson, 2004). The green tea extract and its main catechin polyphenols have medicinal value for the prevention of and therapeutics in several diseases (Ostrowska and Skrzydlewska, 2006). The green tea exerts improvement in liver function by preventing the production of reactive oxygen species (ROS) and enhancing the antioxidant defense system capacity. Thus, green tea extract has protective effects against ethanol toxicity (Bharali and Chetry, 2013; Lodhi et al., 2014).

## 4. Discussion and Conclusion

Although the number of patients with liver diseases has been increasing rapidly, treatment outcomes are still considered unsatisfactory. Herbal medicine has become a major contributor to the management of liver problems. The growing number of preclinical and traditional studies on various herbal medicines indicates a promising future for drug development from herbal-based products. The success of treating liver diseases with herbal medicines depends on understanding each chemical constituent and their interactions. Currently, a handful of herbal drugs, such as silymarin, rutin, epicatechin, and epigallocatechin, have been extensively studied. These plant-based biomolecules, along with other constituents mentioned in this study, have demonstrated their significance and potential as major treatment modalities for liver diseases.

Unlike conventional drugs, which are composed of known chemical constituents that can be accurately quantified, herbal drugs consist of complex mixtures of ingredients. Due to this complexity, studies face major challenges, with the greatest difficulty being the purification of herbal medicines and the identification and quantification of their individual components. Currently, advanced techniques such as high-performance liquid chromatography, protein precipitation, and microdialysis are being used to separate these chemical constituents.

However, studying the clinical effects of individual chemical constituents in isolation is of limited use for several reasons such as the neutralization of harmful chemicals in a mixture by other compounds, as well as the synergistic or inhibitory interactions among constituents that together create an effective in vitro therapeutic combination. Relying solely on

information about the pharmacodynamics of herbal medicines in liver diseases provides insufficient detail for developing drugs with similar effects. Factors such as metabolism, absorption, distribution, and the intrinsic concentration of the drug must be accurately understood to determine the appropriate dosage, duration of treatment, and safety margin.

#### Acknowledgements:

The authors have nothing to report.

#### Conflict of interests:

The authors declare that there are no conflicts of interest related to this article.

#### Data availability statement:

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

## References

- 1) Abou Seif, H. S. (2014b). Ameliorative effect of parsley oil (*Petroselinum crispum*) against alcohol-induced hepatotoxicity and oxidative stress. *Medical Research Journal*, 13(2), 100–107.
- 2) Abou Seif, H. S. (2014a). Ameliorative effect of pumpkin oil (*Cucurbita pepo* L.) against alcohol-induced hepatotoxicity and oxidative stress in albino rats. *Beni-Suef University Journal of Basic and Applied Sciences*, 3(3), 178–185.
- 3) Abou Seif, H. S. (2016). Physiological changes due to hepatotoxicity and the protective role of some medicinal plants. *Beni-Suef University Journal of Basic and Applied Sciences*, 5(2), 134–146.
- 4) Abu, M., Nyeem, B., Alam, K., & Hossain, A. (2017). A review of evidence-based hepatoprotective selected medicinal plants used in Bangladeshi traditional medicine. *Bangladeshi Traditional Medicine*, 2(1), 13–17.
- 5) Aeschbach, R., Loliger, J., Scott, B. C., Murcia, A., Butler, J., Halliwell, B., & et al. (1994). Antioxidant actions of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. *Food and Chemical Toxicology*, 32(1), 31–36.
- 6) Alobaidi, S. (2024). Renal health benefits and therapeutic effects of parsley (*Petroselinum crispum*): A review. *Frontiers in Medicine*, 11, 1494740.
- 7) Al-Zuhair, H., Abdel-Fattah, A. A., & El-Sayed, M. I. (2000). Pumpkin-seed oil modulates the effect of felodipine and captopril in spontaneously hypertensive rats. *Pharmacological Research*, 41(5), 555–563.
- 8) Au, J. S., Navarro, V. J., & Rossi, S. (2011). Drug-induced liver injury – Its pathophysiology and evolving diagnostic tools. *Alimentary Pharmacology & Therapeutics*, 34(1), 11–20.
- 9) Awang, D. V. C. (1992). Ginger. *Canadian Pharmaceutical Journal*, 125(6), 309–311.
- 10) Azab, A. E., & Albasha, M. O. (2018). Hepatoprotective effect of some medicinal plants and herbs against hepatic disorders induced by hepatotoxic agents. *Journal of Biotechnology & Bioengineering*, 2(1), 8–23.
- 11) Bharali, M. K., & Chetry, L. B. (2013). Hepatoprotective role of green tea extract in acetaminophen induced liver injury in mice. *South Asian Journal of Experimental Biology*, 3(2), 84–91.
- 12) Bhawna, S., & Kumar, S. U. (2010). Hepatoprotective activity of some indigenous plants. *International Journal of PharmTech Research*, 2(1), 568–572.



- 13) Caili, F., Huan, S., & Quanhong, L. (2006). A review on pharmacological activities and utilization technologies of pumpkin. *Plant Foods for Human Nutrition*, 61(2), 73–80.
- 14) Chow, J. M., Shen, S. C., Huan, S. K., Lin, H. Y., & Chen, Y. C. (2005). Quercetin, but not rutin and quercitrin, prevents H<sub>2</sub>O<sub>2</sub>-induced apoptosis via antioxidant activity and heme oxygenase-1 gene expression in macrophages. *Biochemical Pharmacology*, 69(12), 1839–1851.
- 15) Crespy, V., & Williamson, G. (2004). A review of the health effects of green tea catechins in vivo animal models. *Journal of Nutrition*, 134(12), 3431S–3440S.
- 16) Dua, V. K., Ojha, V. P., Roy, R., Joshi, B. C., Valecha, N., Devi, C. U., et al. (2004). Antimalarial activity of some xanthenes isolated from the roots of *Andrographis paniculata*. *Journal of Ethnopharmacology*, 95(2–3), 247–251.
- 17) Ekpono, E. U., Eze, E. D., Adam, A. M., Ibiam, U. A., Obasi, O. U., Ifie, J. E., Ekpono, E. U., Alum, E. U., Noreen, S., Awuchi, C. G., & Aja, P. M. (2024). Ameliorative potential of pumpkin seed oil (*Cucurbita pepo* L.) against tramadol-induced oxidative stress. *Dose-Response*, 22(1), 15593258241226913.
- 18) Elhemiely, A. A., El-Fayoumi, S. H., Gadelmawla, M. H. A., Mahran, N. A., & Gad, A. M. (2025). Hesperidin reduces hepatic injury induced by doxorubicin in rat model through its antioxidative and anti-inflammatory effects, focusing on SIRT-1/NRF-2 pathways. *Journal of Biochemical and Molecular Toxicology*, 39(9), e70465.
- 19) El-Sharaky, A. S., Newairy, A. A., Kamel, M. A., & Eweda, S. M. (2009). Protective effect of ginger extract against bromobenzene-induced hepatotoxicity in male rats. *Food and Chemical Toxicology*, 47(7), 1584–1590.
- 20) Emim, J. A., Oliveira, A. B., & Lapa, A. J. (1994). Pharmacological evaluation of the anti-inflammatory activity of a citrus bioflavonoid, hesperidin, and the isoflavonoids, dauricin and claussequinone, in rats and mice. *Journal of Pharmacy and Pharmacology*, 46(2), 118–122.
- 21) Farombi, E. O., Shrotriya, S., Na, H. K., Kim, S. H., & Surh, Y. J. (2008). Curcumin attenuates dimethylnitrosamine-induced liver injury in rats through Nrf2-mediated induction of heme oxygenase-1. *Food and Chemical Toxicology*, 46(4), 1279–1287.
- 22) Farzaei, M. H., Abbasabadi, Z., Ardekani, M. R., Rahimi, R., & Farzaei, F. (2013). Parsley: A review of ethnopharmacology, phytochemistry and biological activities. *Journal of Traditional Chinese Medicine*, 33(6), 815–826.
- 23) Friedman, M. (2007). Overview of antibacterial, antitoxin, antiviral and antifungal activities of tea flavonoids and teas. *Molecular Nutrition & Food Research*, 51(1), 116–134.
- 24) Fusani, P., Biazzi, E., Mella, M., Doria, F., & Tava, A. (2025). The flavonoids and oil composition of parsley [*Petroselinum crispum* (Mill.) Nyman] fruits. *Phytochemistry*, 140, 107288.
- 25) Gossett-Williams, M., Lyttle, K., Clarke, T., Gardner, M., & Simon, O. (2008). Supplementation with pumpkin seed oil improves plasma lipid profile and cardiovascular outcomes of female non-ovariectomized and ovariectomized Sprague-Dawley rats. *Phytotherapy Research*, 22(7), 873–877.
- 26) Hasan, M., Ali, M. T., Khan, R., Palit, P., Islam, A., Seidel, V., Akter, R., & Nahar, L. (2018). Hepatoprotective, antihyperglycemic and antidiabetic effects of *Dendrophthoe pentandra* leaf extract in rats. *Clinical Phytoscience*, 4, 29.
- 27) Hozayen, W. G., & Abou Seif, H. S. (2014). Chemopreventive effects of *Zingiber officinale* extract against aspartame-induced hepatotoxicity and oxidative stress

- in rat model. *Journal of International Academic Research for Multidisciplinary*, 2(8), 215–230.
- 28) Hozayen, W. G., Abou Seif, H. S., & Amin, S. (2014). Protective effects of rutin and/or hesperidin against doxorubicin-induced hepatotoxicity. *International Journal of Clinical Nutrition*, 2(1), 11–17.
  - 29) Islam, R., Alam, M. J., Shanta, S. B., Rahman, M. H., Mahmud, S., & Khan, A. S. (2019). Evaluation of liver protective activity of *Moringa oleifera* bark extract in paracetamol-induced hepatotoxicity in rats. *Journal of Pharmaceutical Research International*, 25(4), 1–9.
  - 30) Jarukamjorn, K., & Nemoto, N. (2008). Pharmacological aspects of *Andrographis paniculata* on health and its major diterpenoid constituent andrographolide. *Journal of Health Science*, 54(4), 370–381.
  - 31) Kolarovic, J., Popovic, M., Zlinska, J., Trivic, S., & Vojnovic, M. (2010). Antioxidant activities of celery and parsley juices in rats treated with doxorubicin. *Molecules*, 15(8), 6193–6204.
  - 32) Kumar, C. H., Ramesh, A., Kumar, S. J., & Ishaq, B. M. (2011). A review on hepatoprotective activity of medicinal plants. *International Journal of Pharmaceutical Sciences and Research*, 2(3), 501–515.
  - 33) Lantz, R. C., Chen, G. J., Sarihan, M., Sólyom, A. M., Jolad, S. D., & Timmermann, B. N. (2007). The effect of extracts from ginger rhizome on inflammatory mediator production. *Phytomedicine*, 14(2–3), 123–128.
  - 34) Liu, J., Lin, H., & McIntosh, H. (2001). Genus *Phyllanthus* for chronic hepatitis B virus infection: A systematic review. *Journal of Viral Hepatitis*, 8(5), 358–366.
  - 35) Lodhi, P., Tandan, N., Singh, N., Kumar, D., & Kumar, M. (2014). *Camellia sinensis* (L.) Kuntze extract ameliorates chronic ethanol-induced hepatotoxicity in albino rats. *Evidence-Based Complementary and Alternative Medicine*, 2014, 1–7.
  - 36) Lopez, M. G., Sanchez-Mendoza, I. R., & Ochoa-Alejo, N. (1999). Comparative study of volatile components and fatty acids of plants and in vitro cultures of parsley (*Petroselinum crispum* (Mill.) Nym. ex-Hill). *Journal of Agricultural and Food Chemistry*, 47(8), 3292–3296.
  - 37) Luo, H. N., Jiang, N. B., King, S. M., & Chen, C. Y. (2008). Inhibition of cell growth and VEGF expression in ovarian cancer cells by flavonoids. *Nutrition and Cancer*, 60(6), 800–809.
  - 38) Madrigal-Santillán, E., Madrigal-Bujaidar, E., Alvarez-González, I., Sumaya-Martínez, M. T., Gutiérrez-Salinas, J., Bautista, M., Morales-González, A., Gonzalez-Rubio, M. G., Aguilar-Faisal, J. L., & Morales-González, J. A. (2014). Review of natural products with hepatoprotective effects. *International Journal of Molecular Sciences*, 20, 14787–14804.
  - 39) Makni, M., Fetoui, H., Gargouri, N. K., Garoui, M., Jaber, H., Makni, J., Boudawara, T., & Zeghal, N. (2008). Hypolipidemic and hepatoprotective effects of flax and pumpkin seed mixture rich in  $\omega$ -3 and  $\omega$ -6 fatty acids in hypercholesterolemic rats. *Food and Chemical Toxicology*, 46(12), 3714–3720.
  - 40) Mishra, P. K., Singh, R. K., Gupta, A., Chaturvedi, A., Pandey, R., Tiwari, S. P., & Mohapatra, T. M. (2013). Antibacterial activity of *Andrographis paniculata* (Burm. f.) Wall ex Nees leaves against clinical pathogens. *Journal of Pharmacy Research*, 7(5), 459–462.
  - 41) Mishra, S., Aeri, V., & Katare, D. P. (2014). Hepatoprotective medication for liver injury. *World Journal of Pharmaceutical and Pharmaceutical Sciences*, 3(6), 891–932.

- 42) Motterlini, R., Foresti, R., Bassi, R., & Green, C. J. (2000). Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cells against oxidative stress. *Free Radical Biology and Medicine*, 28(8), 1303–1312.
- 43) Murkovic, M., Hillebrand, A., Winkler, J., & Pfannhauser, W. (1996). Variability of vitamin E content in pumpkin seeds (*Cucurbita pepo* L.). *European Food Research and Technology*, 202(4), 275–278.
- 44) Ostrowska, J., & Skrzydlewska, E. (2006). The comparison of effect of catechins and green tea extract on oxidative modifications of LDL in vitro. *Advances in Medical Sciences*, 5(2), 298–303.
- 45) Patel, K., Krishna, K., Sokoloski, E., & Ito, Y. (2000). Preparative separation of curcuminoids from crude curcumin and turmeric powder by pH-zone refining countercurrent chromatography. *Journal of Liquid Chromatography & Related Technologies*, 23(14), 2209–2218.
- 46) Pradhan, S. C., & Girish, C. (2006). Hepatoprotective herbal drug, silymarins from experimental pharmacology to clinical medicine. *Indian Journal of Medical Research*, 124(5), 491–504.
- 47) Rasool, M., Iqbal, J., Malik, A., Ramzan, H. S., Qureshi, M. S., Asif, M., et al. (2014). Hepatoprotective effects of *Silybum marianum* (silymarin) and *Glycyrrhiza glabra* (glycyrrhizin) in combination: A possible synergy. *Evidence-Based Complementary and Alternative Medicine*, 2014, 1–10.
- 48) Saller, R., Meier, R., & Brignoli, R. (2001). The use of silymarin in the treatment of liver diseases. *Drugs*, 61(14), 2035–2063.
- 49) Shaarawy, S. M., Tohamy, A. A., Elgendy, S. M., Abd Elmageed, Z. Y., Bahnasy, A., Mohamed, M. S., et al. (2009). Protective effects of garlic and silymarin on NDEA-induced hepatotoxicity in rats. *International Journal of Biological Sciences*, 5(6), 549–557.
- 50) Sheeja, K., Shihab, P. K., & Kuttan, G. (2006). Antioxidant and anti-inflammatory activities of the plant *Andrographis paniculata* Nees. *Immunopharmacology and Immunotoxicology*, 28(1), 129–140.
- 51) Stevenson, D. G., Eller, F. J., Wang, L., Jane, J. L., Wang, T., & Inglett, G. E. (2007). Oil and tocopherol content and composition of pumpkin seed oil in 12 cultivars. *Journal of Agricultural and Food Chemistry*, 55(10), 4005–4013.
- 52) Subramaniam, S., Khan, H. B. H., Elumalai, N., & Lakshmi, S. Y. S. (2015). Hepatoprotective effect of ethanolic extract of whole plant of *Andrographis paniculata* against CCl<sub>4</sub>-induced hepatotoxicity in rats. *Comparative Clinical Pathology*, 24(1), 1–7.
- 53) Swaroop, S., Vaishnav, M., Arora, U., Biswas, S., Aggarwal, A., Sarkar, S., Khanna, P., Elhence, A., Kumar, R., Goel, A., & Shalimar. (2024). Etiological spectrum of cirrhosis in India: A systematic review and meta-analysis. *Journal of Clinical and Experimental Hepatology*, 14(2), 101291.
- 54) Tanaka, T., Kohno, H., Tsukio, Y., Honjo, S., Tanino, M., & Miyake, M. (2000). Citrus limonoids obacunone and limonin inhibit azoxymethane-induced colon carcinogenesis in rats. *BioFactors*, 13(1–4), 213–218.
- 55) Tapsell, L. C., Hemphill, I., Cobiac, L., Patch, C. S., Sullivan, D. R., Fenech, M., Roodenrys, S., Keogh, J. B., Clifton, P. M., Williams, P. G., Fazio, V. A., & Inge, K. E. (2006). Health benefits of herbs and spices: The past, the present, the future. *Medical Journal of Australia*, 185(4 Suppl), S4–S24.
- 56) Wang, W. H., & Wang, Z. M. (2005). Studies of commonly used traditional medicine—Ginger. *Zhongguo Zhong Yao Za Zhi*, 30(20), 1569–1573.



- 57) Wang, W., & Goodman, M. T. (1999). Antioxidant property of dietary phenolic agents in a human LDL-oxidation ex vivo model: Interaction of protein binding activity. *Nutrition Research*, 19(2), 191–202.
- 58) Wang, Y. C., Chuang, Y. C., & Hsu, H. W. (2008). The flavonoid, carotenoid and pectin content in peels of citrus cultivated in Taiwan. *Food Chemistry*, 106(1), 277–284.
- 59) Wiart, C., Kumar, K., Yusof, M. Y., Hamimah, H., Fauzi, Z. M., & Sulaiman, M. (2005). Antiviral properties of ent-labdan diterpenes of *Andrographis paniculata* Nees, inhibitors of herpes simplex virus type 1. *Phytotherapy Research*, 19(12), 1069–1070.
- 60) Wong, P. Y. Y., & Kitts, D. D. (2006). Studies on the dual antioxidant and antibacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food Chemistry*, 97(3), 505–515.
- 61) Zhang, H., Chen, F., Wang, X., & Yao, H. Y. (2006). Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents. *Food Research International*, 39(8), 833–839.
- 62) Zhao, F., He, E. Q., Wang, L., & Liu, K. (2008). Anti-tumor activities of andrographolide, a diterpene from *Andrographis paniculata*, by inducing apoptosis and inhibiting VEGF level. *Journal of Asian Natural Products Research*, 10(5–6), 467–473.
- 63) Zhou, Y. F., Guo, B., Ye, M. J., Liao, R. F., & Li, S. L. (2016). Protective effect of rutin against H<sub>2</sub>O<sub>2</sub>-induced oxidative stress and apoptosis in human lens epithelial cells. *Current Eye Research*, 41(7), 933–942.