

Indium Oxide-Catalyzed Synthesis of Diaryl Disulfides from Aryl Halides

Prasanta Gogoi¹, Rajmul Ahmed², Imamul Hoque³

¹Department of Chemistry, Pandit Deendayal Upadhyaya Adarsha Mahavidyalaya,
Amjonga, Assam, Goalpara, PIN-783124

²Mahatma Gandhi Central University, Raghunathpur, Motihari, East Champaran,
Bihar, PIN- 845401

³Girijananda Chowdhury University, Hathkhowapara, Azara, Guwahati, PIN-781017
*corresponding author: prasantanits.11@gmail.com

Abstract

An efficient and general method for the synthesis of Aryldisulfides using indium (III) oxide as a catalyst is reported. Reaction of various organyl halides afforded the desired disulfide in good yields.

1. Introduction


Disulfide bonds are ubiquitous in many naturally occurring substances. For example, diallyl disulfide, ajoene, S-allylmercaptocysteine, and allicin are found in garlic.¹⁻⁴ These compounds have shown promising anti-infective properties.⁵ Disulfides are also used in the synthesis of various organosulfur compounds.⁶ Industrially, disulfides are employed as vulcanizing agents for rubbers and elastomers to increase their tensile strength.⁷ Due to their importance, various strategies have been reported in the literature for the synthesis of disulfides. Among them, the oxidation of thiols to disulfides using various reagents and catalysts under controlled conditions is the most common.⁸ Other methods include the conversion of alkyl halides to disulfides using Na₂S/S and the reductive coupling of sulfonyl chlorides.⁹⁻¹⁰ However, these methods have several disadvantages. They often produce the desired disulfides along with solid waste by-products. Other drawbacks include the use of toxic reagents and metal oxidants, low yields, long reaction times, and high reaction temperatures.¹¹

Thus, there exists a need for an efficient and simple method for the synthesis of disulfides without the formation of by-products. In this view, we have employed indium(III) oxide as a catalyst for the conversion of aryl halides into their corresponding disulfides. The method

affords disulfides in good yields. The reaction is straightforward and involves simple work-up steps (Scheme 1).

Indium(III) oxide has been chosen as a catalyst for the following reasons:

- It is an easily recyclable and reproducible catalyst.¹²
- It is active, stable, and non-toxic.¹³
- Indium(III) compounds are mild and water-tolerant.¹⁴
- Moreover, their application in organic transformations is still limited.¹⁵



where Ar = m-(NO₂)₂C₆H₃, 2-C₅H₃N, o-NO₂C₆H₄, m-ClC₆H₄NO₂
o-C₆H₄COOH, m-C₆H₅CH₂, p-C₆H₅, o-NO₂C₆H₄

Scheme 1. Synthesis of symmetrical disulfides.

2. Results and Discussions

In a general procedure, chlorobenzene (in ethanol) was refluxed with a solution of sodium disulfide (in ethanol) in the presence of varying amounts of catalyst. The effect of catalyst loading on the reaction system was examined. It was observed that when 3 mol% of the catalyst was used for the conversion of chlorobenzene to diphenyl disulfide, the yield of the desired product was high (78%). The results are summarized in Table 1.

Table 1: Effect of amounts of Indium (III) oxide on the synthesis of dipenyl disulfide

Entry	Substrate	Disulfide	Catalyst	Amount (mol %)	Yield ^a (%)
1	8a	8a	In ₂ O ₃	1.0	25
2	8a	8a	In ₂ O ₃	2.5	64
3	8a	8a	In ₂ O ₃	3.0	78
4	8a	8a	In ₂ O ₃	5.0	70
8	8a	8a	No catalyst	-	35

Reaction conditions: A mixture of sulfur powder (0.16 g, 5 mmol), sodium sulfide (0.39 g, 5 mmol) and ethanol (10 ml) was stirred at room temperature for 15 min. The mixture was added to a solution containing substrate (6 mmol) and ethanol (10 ml) and 3 mol % of indium (III) oxide. The resulting mixture was refluxed for 48 min.

^a Isolated yields. The potential of various solvents were also examined in the synthesis of diphenyl disulfide. Ethanol was found to be effective for the reaction system (Table 2).



Table 2. Effect of various solvents on the synthesis of diphenyl disulfide















Entry	Substrate	Disulfide	Solvent	Yield ^a (%)
1	8a	8a	DMF	50
2	8a	8a	C ₆ H ₆	39
3	8a	8a	CH ₂ Cl ₂	55
4	8a	8a	CH ₃ Cl	60
5	8a	8a	C ₂ H ₅ OH	78

^aIsolated yield

The protocol was extended for the conversion of several aromatic halides to their corresponding disulfides (Table 3). All were converted into corresponding disulfides giving good yields of the disulfides. Disulfides are the only products in the reaction condition.

Table 3. Synthesis of disulfides using In₂O₃ as catalyst

Entry	Substrate (1)	Product (2)	Reaction Time (min)	Melting Point (°C)	Yield ^a (%)
1			30	193	85

2			40	176	89
3			45	203	70
4			45	288	83
5			10	233	90
6			47	68	70
7			60	57	60
8			48	59	78

Reaction conditions: A mixture of sulfur powder (0.16 g, 5 mmol), sodium sulfide (0.39 g, 5 mmol) and ethanol (10 ml) was stirred at room temperature for 15 min. The mixture was added to a solution containing substrate (6 mmol) and ethanol (10 ml) and 3 mol % of indium (III) oxide. The resulting mixture was refluxed.

^a Isolated yield.

Factors such as the presence of substituents in aryl chlorides influence the course of the reaction. For example, aryl chlorides containing nitro groups react rapidly to form the desired

disulfides (entries 1, 2, and 5), while the presence of a chloro group in aryl chlorides decreases their reactivity toward disulfide formation (entry 3). On the other hand, aryl chlorides without substituents react more slowly (entries 6, 7, and 8).

It was also observed that the catalyst could be reused up to three times after washing with ethyl acetate or water, followed by drying under vacuum.

3. Conclusions

In conclusion, we have developed a new and efficient method for the synthesis of disulfides using indium(III) oxide as a catalyst. By applying this methodology, a variety of disulfides were synthesized in good yields.

References

1. Cavallito, C. J., & Bailey, J. H. (1944). Allicin, the antibacterial principle of *Allium sativum*. I. Isolation, physical properties and antibacterial action. *Journal of the American Chemical Society*, 66(11), 1950-1951.
2. Ankri, S., & Mirelman, D. (1999). Antimicrobial properties of allicin from garlic. *Microbes and infection*, 1(2), 125-129.
3. Block, E. (1985). The chemistry of garlic and onions. *Scientific american*, 252(3), 114-121.
4. Ankri, S., & Mirelman, D. (1999). Antimicrobial properties of allicin from garlic. *Microbes and infection*, 1(2), 125-129.
5. Heldreth, B., & Turos, E. (2005). Microbiological properties and modes of action of organosulfur-based anti-infectives. *Current Medicinal Chemistry-Anti-Infective Agents*, 4(4), 295-315.
6. Arisawa, M., & Yamaguchi, M. (2020). Rhodium-catalyzed synthesis of organosulfur compounds involving SS bond cleavage of disulfides and sulfur. *Molecules*, 25(16), 3595.
7. Hernández, M., Grande, A. M., Dierkes, W., Bijleveld, J., Van Der Zwaag, S., & García, S. J. (2016). Turning vulcanized natural rubber into a self-healing polymer: Effect of the disulfide/polysulfide ratio. *ACS sustainable chemistry & engineering*, 4(10), 5776-5784.
8. (a) Ong, C. L., Titinchi, S., Juan, J. C., & Khaligh, N. G. (2021). An Overview of recent advances in the synthesis of organic unsymmetrical disulfides. *Helvetica Chimica Acta*, 104(8), e2100053. (b) Kim, K., & Chae, J. (2025). Synthetic applications of symmetrical disulfides and their synthesis from non-thiolic substrates. *Synthesis*. (c) Burmistrova, D. A., Smolyaninov, I. V., & Berberova, N. T. (2023). Modern Trends in the Synthesis of Disulfides: From Metal-Containing Catalysts to Nonmaterial Reagents. *Russian Journal of Coordination Chemistry*, 49(Suppl 2), S159-S195. (d) Ruano, J. L. G., Parra, A., & Alemán, J. (2008). Efficient synthesis of disulfides by air oxidation of thiols under sonication. *Green Chemistry*, 10(6), 706-711.
9. (a) Brandsma, L., & Wijers, H. E. (1963). A simple laboratory scale preparative method for dialkyl sulphides, selenides, tellurides, thiols and selenols. *Recueil des Travaux Chimiques des Pays-Bas*, 82(1), 68-74. (b) Shiao, M. J., Lai, L. L., Ku, W. S., Lin, P. Y., & Hwu, J. R. (1993). Chlorotrimethylsilane in combination with sodium sulfide as the equivalent of

- sodium trimethylsilanethiolate in organic reactions. *The Journal of Organic Chemistry*, 58(17), 4742-4744.
10. (a) Narayana, C., Padmanabhan, S., & Kabalka, G. W. (1991). Reductive dimerization of sulfonyl derivatives to disulfides and deoxygenation of sulfoxides to sulfides using the boron triiodide-n, n-diethylaniline complex. *Synlett*, 1991(02), 125-126. (b) Guo, H., Wang, J., & Zhang, Y. (1997). Novel reduction of sulfonyl halides to disulfides with SmI₂/Thf/Hmpa system. *Synthetic communications*, 27(1), 85-88.
 11. Guo, H., Wang, J., & Zhang, Y. (1997). Novel reduction of sulfonyl halides to disulfides with SmI₂/Thf/Hmpa system. *Synthetic communications*, 27(1), 85-88.
 12. Gándara, F., Gomez-Lor, B., Gutiérrez-Puebla, E., Iglesias, M., Monge, M. A., Proserpio, D. M., & Snejko, N. (2008). An indium layered MOF as recyclable Lewis acid catalyst. *Chemistry of Materials*, 20(1), 72-76.
 13. Niederberger, M., Garnweitner, G., Buha, J., Polleux, J., Ba, J., & Pinna, N. (2006). Nonaqueous synthesis of metal oxide nanoparticles: Review and indium oxide as case study for the dependence of particle morphology on precursors and solvents. *Journal of Sol-Gel Science and Technology*, 40(2), 259-266.
 14. Serban, M., Halasz, I., & Datta, R. (1999). New water-tolerant supported molten indium catalyst for the selective catalytic reduction of nitric oxide by ethanol. *Catalysis letters*, 63(3), 217-225.
 15. Ranu, B. C. (2000). Indium metal and its halides in organic synthesis. *European Journal of Organic Chemistry*, 2000(13), 2347-2356.