

MA-106 Linear Algebra

M.K. Keshari




Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai - 76


27th January, 2015
D1 - Lecture 10

Orthogonal or perpendicular vectors

The standard basis for \mathbb{R}^2



are vectors of length 1 and perpendicular to each other.
Similarly the standard basis of \mathbb{R}^3



are vectors of length 1 and perpendicular to each other.

Orthogonal vectors

Recall : the length of a vector $v = (x_1, x_2)^T \in \mathbb{R}^2$ is given by

$$\|v\| = \sqrt{x_1^2 + x_2^2} = \sqrt{v^T v}.$$

By Pythagores theorem, $v, w \in \mathbb{R}^2$ are perpendicular to each other if and only if


$$\begin{aligned}\|v\|^2 + \|w\|^2 &= \|v - w\|^2 \\ v^T v + w^T w &= (v - w)^T (v - w) \\ &= (v^T - w^T)(v - w) \\ &= v^T v - w^T v - v^T w + w^T w \\ v^T v + w^T w &= v^T v - 2v^T w + w^T w \quad (\text{since } w^T v = v^T w)\end{aligned}$$

Therefore, v and w are perpendicular to each other if and only if

$$v^T w = 0.$$

We will generalise this to \mathbb{R}^3 and then to \mathbb{R}^n .

length of a vector in \mathbb{R}^3 and \mathbb{R}^n



Apply Pythagores theorem to the blue triangle to get

$$\|(2, 3, 2)\| = \sqrt{\|(2, 3, 0)\|^2 + \|(0, 0, 2)\|^2} = \sqrt{2^2 + 3^2 + 2^2}.$$

Generalize using induction to define the length of a vector $v = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ as

$$\begin{aligned}\|v\| &= \sqrt{\|(x_1, \dots, x_{n-1}, 0)\|^2 + \|(0, 0, \dots, x_n)\|^2} \\ &= \sqrt{x_1^2 + \dots + x_{n-1}^2 + x_n^2} = \sqrt{v^T v}.\end{aligned}$$

Inner product and Orthogonal set

Define the **inner product** (dot product) of two vectors $x, y \in \mathbb{R}^n$ as $x \cdot y := x^T y$. Note that

- $x^T y = y^T x = x_1 y_1 + \cdots + x_n y_n$.
- (Bilinearity) $(x + y)^T z = x^T z + y^T z$.
 $(c x)^T y = c (x^T y) = x^T (c y)$.
- $x^T x = \|x\|^2 \geq 0$ and $x^T x = 0$ if and only if $x = 0$.

Definition

As done in \mathbb{R}^2 , two vectors $v, w \in \mathbb{R}^n$ are orthogonal or perpendicular if and only if $v^T w = 0$ (consider Span of v, w).

A set of vectors $\{v_1, v_2, \dots\} \subseteq \mathbb{R}^n$ is an **orthogonal set** if $v_i \neq 0$ for all i and $v_i^T v_j = 0$ for all $i \neq j$.

An orthogonal set $\{v_1, v_2, \dots\} \subseteq \mathbb{R}^n$ with $\|v_i\| = 1$ for all i is called an **orthonormal set**.

Matrices with columns as orthogonal set

①

$$I = \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}.$$

②

$$A = \begin{pmatrix} 2 & 1 & -2 \\ 2 & -2 & 1 \\ 1 & 2 & 2 \end{pmatrix}.$$

③ Use 1 and -1 only.

$$B = \begin{pmatrix} 1 & -1 & -1 & 1 \\ 1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \\ -1 & 1 & -1 & 1 \end{pmatrix}.$$

- ④ We will show that an orthogonal set is always linearly independent. Hence above matrices are invertible.
- ⑤ From A and B , we can get matrices with orthonormal columns by replacing each column vector v by $v/\|v\|$.

Theorem

Every orthogonal set in \mathbb{R}^n is a linearly independent set.

Proof.

Let $\{v_1, \dots, v_k\}$ be an orthogonal set in \mathbb{R}^n , i.e. $v_i \neq 0$ and $v_i^T v_j = 0$ for $i \neq j$.

Assume for some $a_1, \dots, a_k \in \mathbb{R}$,

$$\begin{aligned} a_1 v_1 + a_2 v_2 + \dots + a_k v_k &= 0 \\ \implies (a_1 v_1 + a_2 v_2 + \dots + a_k v_k)^T v_1 &= 0 v_1 = 0 \\ \implies (a_1 v_1^T + a_2 v_2^T + \dots + a_k v_k^T) v_1 &= 0 \\ \implies a_1 v_1^T v_1 + a_2 v_2^T v_1 + \dots + a_k v_k^T v_1 &= 0 \\ &\implies a_1 v_1^T v_1 = 0 \\ &\implies a_1 = 0 \end{aligned}$$

Similarly, we can show that $a_2 = \dots = a_n = 0$.

Hence $\{v_1, \dots, v_k\}$ is linearly independent. □

1. 140020003 AGRIM GUPTA
2. 140020019 BANERJEE TANAY
3. 140020027 SAUMIL AGARWAL
4. 140020041 KANDHARKAR SWAPNIL SANJAYRAO
5. 140020062 SATYAM AGNIHOTRI
6. 140020074 GURMEET SINGH BEDI
7. 140020095 OJASWA GARG
8. 140020111 ALLIKANTI TEJASWINI RAVINDER
9. 140020124 PUSHPENDRA KUMAR
10. 140050003 MAHAJAN HARSHAL RAJESH
11. 140050009 UTKARSH GAUTAM
12. 140050011 HIMANSHU PAYAL
13. 140050040 KANAPARTHI VEERA VENKATA
14. 140050046 RAPOLU VISHWANADH
15. 140050050 UPPARA RAGHUVeer
16. 140050070 SHAIK ABDUL BASITH
17. 140050083 K M. MUGILVANNAN

Orthogonal Subspaces

Definition

Let V and W be subspaces of \mathbb{R}^n . We say V and W are orthogonal to each other (notation: $V \perp W$) if every vector in V is orthogonal to every vector in W .

Example

$V = \text{Span of } \{v_1, \dots, v_r\}$, $W = \text{Span } \{w_1, \dots, w_t\}$. If $v_i^T w_j = 0$ for all i, j , then $V \perp W$.

Proof. If $v = a_1 v_1 + a_2 v_2 \in V$, $w = b_1 w_1 + b_2 w_2 \in W$, then

$$\begin{aligned} v^T w &= (a_1 v_1^T + a_2 v_2^T)(b_1 w_1 + b_2 w_2) \\ &= a_1 b_1 v_1^T w_1 + a_1 b_2 v_1^T w_2 + a_2 b_1 v_2^T w_1 + a_2 b_2 v_2^T w_2 \\ &= 0 \end{aligned}$$

using bilinearity.

Example

Let V be yz -plane (screen wall) and W be xz -plane (side wall). Are V and W orthogonal to each other (as subspaces of \mathbb{R}^3)?

No. The vector $e_3 = (0, 0, 1) \neq 0$ lies in V and W both .

Remark. If V and W are orthogonal subspaces of \mathbb{R}^n , then $V \cap W = (0)$. It is a necessary condition.

Is it sufficient also? No.

Example

$V = \text{Span of } \begin{pmatrix} 1 \\ 0 \end{pmatrix}, W = \text{Span of } \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$

Then $V \cap W = (0)$, but V and W are not orthogonal, since

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 1.$$

Example. $V = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \right\}, W = \text{Span} \left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \\ 0 \end{pmatrix} \right\}.$

Then V is orthogonal to W .

Proof. It is enough to see that both generators of V are orthogonal to the given generator of W .

Question. Can we enlarge W to W' such that $V \perp W'$ and $\dim V + \dim W' = \dim \mathbb{R}^4 = 4$.

If $A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix}$, then $V = \text{row space of } A$ and $W \subset N(A)$.

By Rank-Nullity Theorem: $\text{rank}(A) + \text{nullity}(A) = 4$.

Since $\text{rank}(A) = 2$, we get $\text{Nullity}(A) = 2$.

Therefore if $W' = N(A)$, then $V \perp W'$ and $\dim V + \dim W' = \dim \mathbb{R}^4$.

By inspection, $W' = \text{Span}\{(1, -1, 1, 0)^T, (0, 0, 0, 1)^T\}$.

Theorem (Fundamental Theorem of Orthogonality - 1)

Let A be a $m \times n$ matrix.

1. The row space of A , $C(A^T)$ is orthogonal to $N(A)$.
2. The column space is orthogonal to the left nullspace, $N(A^T)$.

Example. Assume $\text{rank}(A) = 1$. Then A has 1 pivot. Therefore the column space has dimension 1 and row space has dimension 1.

$$\text{For } A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{pmatrix}, C(A^T) = \text{Span} \begin{pmatrix} 1 \\ 2 \end{pmatrix}, N(A) = \text{Span} \begin{pmatrix} -2 \\ 1 \end{pmatrix},$$

$$C(A) = \text{Span} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, N(A^T) = \text{plane } y_1 + 2y_2 + 3y_3 = 0.$$

$$\dim(C(A^T)) + \dim(N(A)) = 1 + 1 = 2,$$

$$\dim(C(A)) + \dim(N(A^T)) = 1 + 2 = 3.$$

Observe: $N(A)$ = set of all vectors orthogonal to row space.

$N(A^T)$ = set of all vectors orthogonal to column space.