

MA-106 Linear Algebra

M.K. Keshari



Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai - 76

19th January, 2015
D1 - Lecture 7

Recall: We have seen how to find all solutions of $Ax = b$.

$$x_{\text{Complete}} = x_{\text{Particular}} + x_{\text{NullSpace}}$$

We defined Column space $C(A)$ of A .

$b \in C(A)$ if and only if $Ax = b$ is consistent.

If $b \in C(A)$, then to write b as a linear combination of columns of A , we need to solve $Ax = b$.

We defined Vector space $(V, +, \cdot)$

It is closed under linear combinations with following properties

$+$ is commutative, associative, additive identity 0 , additive inverse

scalar multiplication \cdot is associative, $1 \cdot v = v$, distributive.

Examples of Vector spaces

- ① $M = \{2 \times 2 \text{ matrices with real entries}\}$ is a vector space.
- ② $V = C([0, 1], \mathbb{R})$ is a vector space.
In V , vectors are continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.
The zero vector is the zero function.
- ③ $P = \{\text{polynomial functions} : \mathbb{R} \rightarrow \mathbb{R}\}$ is a vector space.
A typical element of P : $a_0 + a_1x + \dots + a_nx^n : \mathbb{R} \rightarrow \mathbb{R}$
with $a_i \in \mathbb{R}$. The zero vector is the zero polynomial.
- ④ Let M , V and P be vector spaces considered above.
Let $W = \{(m, v, p) : m \in M, v \in V, p \in P\}$.
Define addition and scalar multiplication componentwise.
Then W is a vector space. What is the zero vector in W ?
- ⑤ $T = \{2 \times 2 \text{ matrices with entries in } W\}$ is a vector space
with componentwise addition and scalar multiplication.

Subspaces

A subspace of a vector space is a non-empty subset which itself satisfies the requirements of a vector space:

Linear combinations stay in the subspace.

Thus, if V is a vector space, and W is a non-empty subset, then W is a subspace of V if:

$$\bar{x}, \bar{y} \text{ in } W, \quad u, v \text{ in } \mathbb{R}, \Rightarrow u \cdot \bar{x} + v \cdot \bar{y} \text{ are in } W.$$

- 1 $\{(x_1, x_2) : x_1 \geq 0, x_2 \geq 0\}$ is not a subspace of \mathbb{R}^2 . Why?
- 2 The set of 2×2 symmetric matrices is a subspace of M .
So is the set of 2×2 lower triangular matrices.
- 3 The set of invertible 2×2 matrices is not a subspace of M . Why?

Examples: Subspaces of \mathbb{R}^n

- $\{0\}$: The zero subspace.
- \mathbb{R}^n itself.
- A line not passing through the origin is not a subspace of \mathbb{R}^2 . Why? e.g., $x - y = 1$.
- A line L passing through the origin is a subspace of \mathbb{R}^2 . Why? e.g., $x - y = 0$.
- Let A be an $m \times n$ matrix.
The null space of A , $N(A)$, is a subspace of \mathbb{R}^n .
The column space of A , $C(A)$, is a subspace of \mathbb{R}^m .
Recall: They are both closed under linear combinations.


Note: A subspace must contain the 0 vector!

Examples: Subspaces of \mathbb{R}^2

What are the subspaces of \mathbb{R}^2 ?

- $V = \{(0 \ 0)^T\}$.
- $V = \mathbb{R}^2$.

Suppose V
contains a
non-zero vector,
say
 $a = (-1 \ 1)^T$.



Hence V must contain the entire line $L : x + y = 0$, i.e., all multiples of \bar{a} .

- Thus if V is non-zero and not \mathbb{R}^2 , then it is a line passing through the origin. Why?

Examples: Subspaces of \mathbb{R}^2

Let V be a subspace of \mathbb{R}^2 containing $v_1 = (-1 \ 1)^T$. Then V must contain the entire line $L : x + y = 0$.

If $V \neq L$, it contains a vector v_2 , which is not a multiple of v_1 , say $v_2 = (0 \ 1)^T$.

Observe: $A = (v_1 \ v_2) = \begin{pmatrix} -1 & 0 \\ 1 & 1 \end{pmatrix}$ has two pivots,

$\Rightarrow A$ is invertible.

\Rightarrow for any v in \mathbb{R}^2 , $Ax = v$ is solvable,

$\Rightarrow v$ is in $C(A)$,

$\Rightarrow v$ can be written as a linear combination of v_1 and v_2 .

$\Rightarrow v$ is in V , i.e., $V = \mathbb{R}^2$

To summarise: A subspace of \mathbb{R}^2 , which is non-zero, and not a line passing through the origin, is \mathbb{R}^2 .

Linear Span: Definition

Given a collection S of vectors v_1, v_2, \dots, v_n in a vector space V , the *linear span* of S , denoted $\text{Span}(S)$ or $\text{Span}\{v_1, \dots, v_n\}$, is the set of all linear combinations of v_1, v_2, \dots, v_n , i.e.,

$$\text{Span}(S) = \{v = a_1 \cdot v_1 + \dots + a_n \cdot v_n, \text{ for scalars } a_1, \dots, a_n\}.$$

Remark: $\text{Span}(S)$ is a subspace of V . Why?

- $\text{Span}\{v_1, \dots, v_n\} = C(A)$ for $A = (v_1 \ \dots \ v_n)$.

Thus $v \in \text{Span}\{v_1, \dots, v_n\} \Leftrightarrow Ax = v$ is consistent.

- Let $v_1, \dots, v_n \in \mathbb{R}^n$ and $A = (v_1 \ \dots \ v_n)$. Then A is invertible $\Leftrightarrow A$ has n pivots

$$\Leftrightarrow Ax = v \text{ is consistent for every } v \text{ in } \mathbb{R}^n$$

$$\Leftrightarrow \text{Span}\{v_1, \dots, v_n\} = \mathbb{R}^n.$$

Linear Span: Examples

Examples:

- 1 $\text{Span}\{0\} = \{0\}$.
- 2 If $v \neq 0$ is a vector, $\text{Span}\{v\} = \{a \cdot v, \text{ for scalars } a\}$.
Geometrically: $\text{Span}\{v\}$ = the line in the direction of v passing through the origin.
- 3 $\text{Span}\left\{\begin{pmatrix} -1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}\right\} = \mathbb{R}^2$.
- 4 The columns e_1, \dots, e_n of the $n \times n$ identity matrix I span \mathbb{R}^n .
- 5 If A is $m \times n$, then $\text{Span}\{A_1, \dots, A_n\} = C(A)$.
- 6 If v_1, \dots, v_k are all the special solutions of A , then $\text{Span}\{v_1, \dots, v_k\} = N(A)$.

Remark: All of the above are subspaces.

1. 140020001 SHANTANU PATEL
2. 140020010 AVINASH CHATTANATHAN IYER
3. 140020020 VIDWANS NIRAJ ASHUTOSH
4. 140020028 ANKIT KUMAR GAUTAM
5. 140020039 GOUTHAM RAMAKRISHNAN
6. 140020091 KANCHAN KHETAN
7. 140020104 GAURAV JAIN
8. 140020120 NIPUN GUPTA
9. 140050012 AAKASH PRALIYA
10. 140050027 RAJAT CHATURVEDI
11. 140050074 KOPPISETTI NIKHILESWAR
12. 140050006 VAIDYA KAPIL EKNATH
13. 140020014 OMKAR RAVINDRA KAJROLKAR
14. 140020036 DRAVID ASHWIN NATRAJ
15. 140050062 POOSA NIHAL
16. 140050079 VUPPULA MANITEJA REDDY

Linear Span: Examples

Example: Is v in $\text{Span}\{v_1, v_2, v_3, v_4\}$, where:

$$v = \begin{pmatrix} 1 \\ 0 \\ 4 \end{pmatrix}, v_1 = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, v_2 = \begin{pmatrix} 2 \\ 4 \\ 6 \end{pmatrix}, v_3 = \begin{pmatrix} 3 \\ 8 \\ 7 \end{pmatrix}, v_4 = \begin{pmatrix} 5 \\ 12 \\ 13 \end{pmatrix}?$$

$$\text{Let } A = (v_1 \ v_2 \ v_3 \ v_4) = \begin{pmatrix} 1 & 2 & 3 & 5 \\ 2 & 4 & 8 & 12 \\ 3 & 6 & 7 & 13 \end{pmatrix}. \text{ Then}$$

$$(A|b) \rightarrow \left(\begin{array}{cccc|c} 1 & 2 & 3 & 5 & b_1 \\ 0 & 0 & 2 & 2 & b_2 - 2b_1 \\ 0 & 0 & 0 & 0 & b_3 + b_2 - 5b_1 \end{array} \right).$$

Linear Span: Examples

$$\text{Recall } A = \begin{pmatrix} 1 & 2 & 3 & 5 \\ 2 & 4 & 8 & 12 \\ 3 & 6 & 7 & 13 \end{pmatrix}$$

$$Ax = b \text{ is solvable} \Leftrightarrow 5b_1 - b_2 - b_3 = 0.$$

$\Rightarrow v = (1, 0, 4)^T$ is not in $\text{Span}\{v_1, v_2, v_3, v_4\}$, and

$w = (1 \ 0 \ 5)^T = 4v_1 + (-1)v_3$ is in it.

Observe: $v_2 = 2v_1$ and $v_4 = 2v_1 + v_3$.

Hence v_2, v_4 are in $\text{Span}\{v_1, v_3\}$.

Therefore, $\text{Span}\{v_1, v_3\} = \text{Span}\{v_1, v_2, v_3, v_4\} = C(A) =$ the plane $P : (5x - y - z = 0)$.

Q: Is the span of two vectors always a plane?

Linear Independence: Definition

Q: Is the span of two vectors always a plane? **A:** Not always.

If v is a multiple of $w \neq 0$, then $\text{Span}\{v, w\} = \text{Span}\{w\}$, which is a line through the origin.

Q: If v and w are not on the same line through the origin?

A: Yes. v, w are examples of the following:

The vectors v_1, v_2, \dots, v_n in a vector space V , are *linearly independent* if

$$a_1v_1 + a_2v_2 + \dots + a_nv_n = 0 \Rightarrow a_1 = 0, a_2 = 0, \dots, a_n = 0.$$

Observe: If $A = (v_1 \ v_2 \ \dots \ v_n)$, then v_1, v_2, \dots, v_n are linearly independent $\Leftrightarrow Ax = x_1v_1 + x_2v_2 + \dots + x_nv_n = 0$ has only the trivial solution $\Leftrightarrow N(A) = 0$.

The line in \mathbb{R}^2 not passing through origin is not a vector space.
e.g. If L is the line $x - y = 1$. Then $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ are on L , but their sum $\begin{pmatrix} 3 \\ 1 \end{pmatrix}$ is not on L . Hence L is not closed under linear combination. So L is not a vector space.