


PH 103 : Electricity and Magnetism

Tutorial Sheet 9 : Faraday's law

1. A rectangular metallic wire of mass m , width w , length L and resistance R falls through a magnetic field $\vec{B} = B_0\hat{i}$ in a specified region under the influence of gravity as shown in the figure. The loop first enters the region of magnetic field from the top.
 - (a) Draw a qualitative graph of (i) current in the loop vs time and (ii) velocity of the loop vs time, as the loop moves through the magnetic field.
 - (b) What is the terminal velocity of the loop when it is at the position shown in the figure?
 - (c) What will be the terminal velocity of the loop if the magnetic field is nonuniform, $\vec{B} = B_0(z)\hat{i}$



2. A rectangular coil PQRS (length $2L$ and width $2w$) is rotating with a uniform angular speed ω about the z -axis in a uniform magnetic field B directed along the y -direction. Determine the induced emf in the coil using only the concept of motional emf. Check your answer using the concept of change in flux.
3. A Faraday disk generator is a metal disk of radius R rotated about its axis with a constant angular velocity ω in a uniform magnetic field parallel to the axis of rotation. Calculate the emf between the axis and the rim. Verify your results both by Faraday's law and motional emf.
4. A circular loop of radius R rotates about its diameter with the angular velocity ω . A magnetic dipole carrying magnetic moment \vec{m} is placed at the centre of the loop. The axis of the dipole coincides with the rotating diameter of the loop. Find the emf induced between the north pole and a point on the equator of the loop. [Ans. $\mu_0 m \omega / 8\pi R$].
5. A long cylindrical shell of radius R carries a uniform surface charge density σ . If the cylinder is rotated about its axis with an angular velocity $\omega = \omega_0 e^{-t/\tau}$, find the electric field inside and outside the shell. [Ans. $E_{in} = \frac{\sigma \omega_0 R}{2\tau} r e^{-t/\tau} \hat{\phi}$, $E_{out} = \frac{\sigma \omega_0 R^3}{2\tau r} e^{-t/\tau} \hat{\phi} + \frac{\sigma R}{\epsilon_0 r} \hat{r}$]
6. The current through a long solenoid (radius a) varies as $I(t) = I_0 t$, ($I_0 > 0$). A circular metallic wire of radius b ($b > a$) and resistance R is kept concentric with the solenoid. Find the current through the wire. [Ans. $-\frac{\mu_0 N \pi a^2}{R} I_0$]