

MA-106 Linear Algebra

M.K. Keshari



Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai - 76

16th February, 2015
D1 - Lecture 18

Recall:

- The problem of solving linear system of 1-st order ODE with constant coefficients $\frac{du}{dt} = Au$ reduces to solving the eigenvalue problem $Ax = \lambda x$.
- The eigenvalues of A are roots of characteristic polynomial $\det(A - \lambda I)$ and the eigenspace associated to eigenvalue λ is $N(A - \lambda I)$.
- $\lambda = 0$ is an eigenvalue of $A \Leftrightarrow A$ is singular.
So A is non-singular $\Leftrightarrow 0$ is not an eigenvalue of A .
- If A is a diagonal matrix, then eigenvalues of A are $a_{11}, a_{22}, \dots, a_{nn}$ and the associated eigenvectors are e_1, \dots, e_n .
- Eigenvectors need not form a basis of \mathbb{R}^n . **Ex:** $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.
- **Def.** Let A and B be square matrices such that $S^{-1}AS = B$ for an invertible matrix S . Then A and B are called **similar**.

- Recall: Let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be linear map $L(x, y) = (x + y, x + 2y)$. Let $\mathcal{B} = \{(1, 0)^T, (0, 1)^T\}$ and $\mathcal{B}' = \{(1, 1)^T, (1, -1)^T\}$ be two bases of \mathbb{R}^2 . Let $A = [L]_{\mathcal{B}}^{\mathcal{B}}$ and $B = [L]_{\mathcal{B}'}^{\mathcal{B}'}$. Then A and B are similar. Consider

$$\mathbb{R}_{\mathcal{B}'}^2 \xrightarrow{id} \mathbb{R}_{\mathcal{B}}^2 \xrightarrow{L} \mathbb{R}_{\mathcal{B}}^2 \xrightarrow{id} \mathbb{R}_{\mathcal{B}'}^2$$

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, S = [id]_{\mathcal{B}'}^{\mathcal{B}} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, B = S^{-1}AS = \begin{pmatrix} \frac{5}{2} & -\frac{1}{2} \\ -1 & 1 \end{pmatrix}.$$

We can find B directly also

$$L(1, 1) = (2, 3) = \frac{5}{2}(1, 1) + \frac{-1}{2}(1, -1)$$

$$L(1, -1) = (0, -1) = \frac{-1}{2}(1, 1) + \frac{1}{2}(1, -1)$$

- If A and B are similar, then they have same characteristic polynomial, $\det(A - \lambda I) = \det(B - \lambda I)$, hence same eigenvalues.
- Def.** A square matrix A is called **diagonalizable** if A is similar to a diagonal matrix Λ , i.e., $S^{-1}AS = \Lambda$ for some S .
In this case, the eigenvalues of A are the diagonal entries of Λ .

1. 140020024 PRANSHU MAHENDRA JAIN
2. 140020025 AMIYA MAITREYA
3. 140020042 PRANAY AGARWAL
4. 140020045 MOHIT SINGHAL
5. 140020054 PRANAY LADIWALA
6. 140020089 SATYENDRA KUMAR
7. 140020105 ANUP PATTNAIK
8. 140050028 VARRE ADITYA VARDHAN
9. 140050031 C VISHWESH
10. 140050038 GUDIPATI KRISHNA CHAITANYA
11. 140050047 Y PUSHYARAG
12. 140050057 YOGENDRA VINAY
13. 140050065 ARUKONDA GOUTHAM SURYA
14. 140050071 DARNASI RAHUL KIRAN
15. 140050056 DANTAM MOHAN SAITEJA
16. 140050060 GANGAM ROHITH REDDY
17. 140050068 THANNEERU RANA PRATHAP

Diagonalization of a Matrix

Q: What is the advantage of a basis of \mathbb{R}^n consisting of eigenvectors?

Theorem:

Eigenvectors diagonalize a matrix

Assume an $n \times n$ matrix A has a basis consisting of eigenvectors $\{x_1, \dots, x_n\}$ with eigenvalues $\lambda_1, \dots, \lambda_n$.

Consider the invertible matrix $S = [x_1 \ \dots \ x_n]$ with columns x_i .

Then $S^{-1}AS = \text{diagonal matrix } \Lambda = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$.

Proof. (2×2 case). $Ax_i = \lambda_i x_i$. Hence

$$AS = A \begin{bmatrix} x_1 & x_2 \end{bmatrix} = \begin{bmatrix} Ax_1 & Ax_2 \end{bmatrix} = \begin{bmatrix} \lambda_1 x_1 & \lambda_2 x_2 \end{bmatrix} = S\Lambda.$$

Therefore $S^{-1}AS = \Lambda$, i.e., A is similar to a diagonal matrix.

Caution: $\Lambda S = \begin{bmatrix} \lambda_1 & \\ & \lambda_2 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} \lambda_1 a & \lambda_1 b \\ \lambda_2 c & \lambda_2 d \end{bmatrix} \neq S\Lambda.$

Diagonalization: Example

Ex: $A = \begin{bmatrix} 1 & 5 & 6 \\ 0 & 2 & -4 \\ 0 & 0 & 3 \end{bmatrix}$ is triangular.

$$\det(A - \lambda I) = (1 - \lambda)(2 - \lambda)(3 - \lambda).$$

Eigenvalues: $\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 3$.

Note: If A is triangular, its eigenvalues are sitting on the diagonal

$$\text{Eigenvectors: } x_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, x_2 = \begin{bmatrix} 5 \\ 1 \\ 0 \end{bmatrix}, x_3 = \begin{bmatrix} -7 \\ -4 \\ 1 \end{bmatrix}.$$

Further, $\{x_1, x_2, x_3\}$ is a basis of \mathbb{R}^3 .

Hence $S = [x_1 \ x_2 \ x_3]$ is invertible, and

$$AS = [Ax_1 \ Ax_2 \ Ax_3] = [x_1 \ 2x_2 \ 3x_3] = S\Lambda, \text{ where } \Lambda = \begin{bmatrix} 1 & & \\ & 2 & \\ & & 3 \end{bmatrix}.$$

Thus $S^{-1}AS = \Lambda$, i.e., A is diagonalizable.

Diagonalization and Change of Basis

With A as before, $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $Tx = Ax$ is linear. If $\mathcal{S} = \{e_1, e_2, e_3\}$ is the standard basis of \mathbb{R}^3 , then

$$[T]_{\mathcal{S}}^{\mathcal{S}} = \begin{bmatrix} [Te_1]_{\mathcal{S}} & [Te_2]_{\mathcal{S}} & [Te_3]_{\mathcal{S}} \end{bmatrix} = A.$$

Recall that for $x_1 = (1, 0, 0)^T$, $x_2 = (5, 1, 0)^T$ and $x_3 = (-7, -4, 1)^T$,
$$Tx_1 = x_1, \quad Tx_2 = 2x_2, \quad Tx_3 = 3x_3.$$

Furthermore, $\mathcal{B} = \{x_1, x_2, x_3\}$ is a basis of \mathbb{R}^3 . **Q:** What is $[T]_{\mathcal{B}}^{\mathcal{B}}$?

$$[T]_{\mathcal{B}}^{\mathcal{B}} = \begin{bmatrix} [Tx_1]_{\mathcal{B}} & [Tx_2]_{\mathcal{B}} & [Tx_3]_{\mathcal{B}} \end{bmatrix} = \Lambda = \begin{bmatrix} 1 & & \\ & 2 & \\ & & 3 \end{bmatrix}.$$

Consider $\mathbb{R}_{\mathcal{B}}^3 \xrightarrow{id} \mathbb{R}_{\mathcal{S}}^3 \xrightarrow{T} \mathbb{R}_{\mathcal{S}}^3 \xrightarrow{id} \mathbb{R}_{\mathcal{B}}^3$

Change of basis formula:

$$[id]_{\mathcal{S}}^{\mathcal{B}} [T]_{\mathcal{S}}^{\mathcal{S}} [id]_{\mathcal{B}}^{\mathcal{S}} = [T]_{\mathcal{B}}^{\mathcal{B}}, \text{ i.e., } [id]_{\mathcal{S}}^{\mathcal{B}} A [id]_{\mathcal{B}}^{\mathcal{S}} = \Lambda.$$

Observe: $[id]_{\mathcal{B}}^{\mathcal{S}} = \begin{bmatrix} [id(x_1)]_{\mathcal{S}} & [id(x_2)]_{\mathcal{S}} & [id(x_3)]_{\mathcal{S}} \end{bmatrix} := S.$

i.e., the change of basis formula gives: $S^{-1}AS = \Lambda.$

Thus diagonalization of a matrix is the same as finding a basis w.r.t. which the matrix is diagonal.

When is A Diagonalizable

- If x_1, \dots, x_r are eigenvectors of A associated to **distinct** eigenvalues $\lambda_1, \dots, \lambda_r$, then x_1, \dots, x_r are linearly independent.

Proof. Suppose x_1, \dots, x_r are linearly dependent. Choose a linear relation involving minimum number of x_i 's, say

$$(1) \quad a_1 x_1 + \dots + a_t x_t = 0. \quad (1 < t \leq r, t \text{ is minimal, } a_i \neq 0)$$

Apply A to get $a_1 \lambda_1 x_1 + \dots + a_t \lambda_t x_t = 0 \quad (2)$

$$\lambda_1 \cdot (1) - (2) \text{ gives } a_2(\lambda_1 - \lambda_2)x_2 + \dots + a_t(\lambda_1 - \lambda_t)x_t = 0,$$

which contradicts the minimality of t . □

- If A has n distinct eigenvalues, then A is diagonalizable.

Proof. If x_1, \dots, x_n are eigenvectors associated to distinct eigenvalues $\lambda_1, \dots, \lambda_n$, then $\{x_1, \dots, x_n\}$ is a linearly independent set.

Then $S = [x_1 \ \dots \ x_n]$ is invertible, and $S^{-1}AS = \Lambda$ as earlier.

Hence A is diagonalizable. □

When is A Diagonalizable

- A is diagonalizable $\Leftrightarrow A$ has n linearly independent eigenvectors.

Proof. We have seen (\Leftarrow). Let's prove (\Rightarrow).

Assume $S = [x_1 \ \dots \ x_n]$ is an invertible matrix such that

$$S^{-1}AS = \Lambda = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}.$$

Then $AS = S\Lambda$, i.e. $[Ax_1 \ \dots \ Ax_n] = [\lambda_1 x_1 \ \dots \ \lambda_n x_n]$.

Therefore x_1, \dots, x_n are eigenvectors of A . They are linearly independent since S is invertible. □

The columns of the diagonalizing matrix S are eigenvectors of A

S need not be unique, e.g., replace x_1 by $2x_1$ etc.

Diagonalizability: Non-examples

Ex: $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ has repeated eigenvalues 1, 1.

The eigenvectors of A are $\begin{bmatrix} y \\ 0 \end{bmatrix}$. Therefore A does not have a basis consisting of eigenvectors, so A is not diagonalizable.

Ex: Similarly, for any $a \in \mathbb{R}$, the matrix $\begin{bmatrix} a & 1 \\ 0 & a \end{bmatrix}$ has repeated eigenvalues a, a and is not diagonalizable.

Eigenvalues of AB and $A + B$

- If λ is an eigenvalue of A , μ is an eigenvalue of B , is $\lambda\mu$ an eigenvalue of AB ?

Proof. False Proof. $ABx = A(\mu x) = \mu(Ax) = \lambda\mu x.$

This is false since A and B may not have same eigenvector x .

- **Ex:** $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$, $AB = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$

The eigenvalues of A and B are 0,0 and that of AB are 1,0.

- Eigenvalues of $A + B$ are NOT $\lambda + \mu$.

In above example, $A + B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ has eigenvalues 1, -1.

- If A and B have **same eigenvectors** associated to λ and μ , then $\lambda\mu$ and $\lambda + \mu$ are eigenvalues of AB and $A + B$ respectively.

Q: When do A and B have the same eigenvectors?

Simultaneous Diagonalizability

- Assume A and B are diagonalizable. Then A and B have same eigenvector matrix S if and only if $AB = BA$.
- Proof.** (\Rightarrow) Assume $S^{-1}AS = \Lambda_1$ and $S^{-1}BS = \Lambda_2$, where Λ_1 and Λ_2 are diagonal matrices.

$$\text{Then } AB = (S\Lambda_1S^{-1})(S\Lambda_2S^{-1}) = S(\Lambda_1\Lambda_2)S^{-1}$$
$$\text{and } BA = S(\Lambda_2\Lambda_1)S^{-1}.$$

Since $\Lambda_1\Lambda_2 = \Lambda_2\Lambda_1$, we get $AB = BA$.

- (Part of \Leftarrow) Assume $AB = BA$.

If $Ax = \lambda x$, then $ABx = B(Ax) = B(\lambda x) = \lambda Bx$.

If $Bx = 0$, then x is an eigenvector of B , associated to $\mu = 0$.

If $Bx \neq 0$, then x and Bx both are eigenvectors of A , associated to λ .

Special case: Assume all the eigenspaces of A are one dimensional.

Then $Bx = \mu x$ for some scalar $\mu \Rightarrow x$ is an eigenvector of B .

We will not prove the general case. □