

# CS 207: Discrete Structures

## Graph theory

Bipartite subgraphs, graph isomorphism

Lecture 26

Sept 22 2015

## Topic 3: Graph theory

### Topics covered

- ▶ Eulerian graphs and a characterization
- ▶ Characterization of bipartite graphs using odd cycles.
- ▶ Subgraphs and degree sum formula.
- ▶ Cliques and independent sets.

### Today

- ▶ Finding a large bipartite subgraph of a given graph
- ▶ Graph representation as a matrix.
- ▶ Comparing graphs: isomorphism

# Bipartite graphs

## Definition

A graph is called **bipartite**, if the vertices of the graph can be partitioned into  $V = X \cup Y$ ,  $X \cap Y = \emptyset$  s.t.,  $\forall e = (u, v) \in E$ ,

- ▶ either  $u \in X$  and  $v \in Y$
- ▶ or  $v \in X$  and  $u \in Y$

Example:  $m$  jobs and  $n$  people,  $k$  courses and  $\ell$  students.

- ▶ How can we check if a graph is bipartite?
- ▶ Can we characterize bipartite graphs?

## Some basic stuff that we have already seen

Degree-Sum Formula (also called Handshake Lemma!)

For any graph  $G$  with vertex set  $V$  and edge set  $E$ :

$$\sum_{v \in V} d(v) = 2|E|$$

## Some basic stuff that we have already seen


### Degree-Sum Formula (also called Handshake Lemma!)

For any graph  $G$  with vertex set  $V$  and edge set  $E$ :

$$\sum_{v \in V} d(v) = 2|E|$$

### Subgraphs of a graph $G$

A subgraph  $H$  of a graph  $G$  is a graph  $H$  such that  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$  (and the assignment of endpoints to edges in  $H$  is same as in  $G$ ).



# Bipartite subgraphs of graphs

- ▶ Does a graph  $G$  always have a bipartite subgraph?
- ▶ How large is it?

# Bipartite subgraphs of graphs

- ▶ Does a graph  $G$  always have a bipartite subgraph?
- ▶ How large is it?

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

# Bipartite subgraphs of graphs

- ▶ Does a graph  $G$  always have a bipartite subgraph?
- ▶ How large is it?

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle)




# Bipartite subgraphs of graphs

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle) Let  $H$  be the bipartite subgraph of  $G$  s.t, # edges crossing bipartition is max possible.




# Bipartite subgraphs of graphs

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle) Let  $H$  be the bipartite subgraph of  $G$  s.t. # edges crossing bipartition is max possible.

- Either,  $\forall v \in V(G), d_H(v) \geq \frac{d_G(v)}{2}$ , then  $|E(H)| \geq \frac{|E(G)|}{2}$ .  
 $d_H(v)$  denotes degree of  $v$  in  $H$  and  $d_G(v)$  its degree in  $G$ .




# Bipartite subgraphs of graphs

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle) Let  $H$  be the bipartite subgraph of  $G$  s.t. # edges crossing bipartition is max possible.

- ▶ Either,  $\forall v \in V(G), d_H(v) \geq \frac{d_G(v)}{2}$ , then  $|E(H)| \geq \frac{|E(G)|}{2}$ .
- ▶ Or,  $\exists v \in V(G)$  such that  $d_H(v) \leq \frac{d_G(v)}{2}$ .




# Bipartite subgraphs of graphs

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle) Let  $H$  be the bipartite subgraph of  $G$  s.t. # edges crossing bipartition is max possible.

- ▶ Either,  $\forall v \in V(G), d_H(v) \geq \frac{d_G(v)}{2}$ , then  $|E(H)| \geq \frac{|E(G)|}{2}$ .
- ▶ Or,  $\exists v \in V(G)$  such that  $d_H(v) < \frac{d_G(v)}{2}$ . Then switching  $v$  to other partition contradicts choice of  $H$ .




# Bipartite subgraphs of graphs

## Theorem

Every simple graph  $G$  has a bipartite subgraph with at least  $|E|/2$  edges.

Proof: (by extremality principle) Let  $H$  be the bipartite subgraph of  $G$  s.t, # edges crossing bipartition is max possible.

- ▶ Either,  $\forall v \in V(G), d_H(v) \geq \frac{d_G(v)}{2}$ , then  $|E(H)| \geq \frac{|E(G)|}{2}$ .
- ▶ Or,  $\exists v \in V(G)$  such that  $d_H(v) \leq \frac{d_G(v)}{2}$ . Then switching  $v$  to other partition contradicts choice of  $H$ .  $\square$



- ▶ Can we make this into an algorithm to produce such  $H$ ?

# Representing and comparing graphs


We start with simple graphs...



To represent it, we need to name the vertices...

# Representing and comparing graphs

We start with simple graphs...




To represent it, we need to name the vertices...

► As an adjacency list:

$v_1$	$v_2, v_4$
$v_2$	$v_1, v_3$
$v_3$	$v_2, v_4$
$v_4$	$v_1, v_3$

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- As an adjacency matrix:

$$\begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} \begin{pmatrix} & v_1 & v_2 & v_3 & v_4 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$



# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- ▶ As an adjacency matrix:

$$\begin{array}{c} \begin{array}{cc} & \begin{array}{cccc} v_1 & v_2 & v_3 & v_4 \end{array} \\ \begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} & \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \end{array} \end{array}$$

- ▶ But we want to study properties that are independent of the naming, e.g., connectivity.
- ▶ Are two given graphs the “same”, wrt these properties?

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- As an adjacency matrix:

$$\begin{array}{c} v_1 \quad v_2 \quad v_3 \quad v_4 \\ \begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \end{array}$$

- But we want to study properties that are independent of the naming, e.g., connectivity.
- Are two given graphs the “same”, wrt these properties?

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- As an adjacency matrix:

$$\begin{array}{c} \begin{array}{cccc} & v_1 & v_2 & v_3 & v_4 \\ \begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} & \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & a & b & c & d \\ \begin{array}{c} a \\ b \\ c \\ d \end{array} & \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix} \end{array} \end{array}$$

- But we want to study properties that are independent of the naming, e.g., connectivity.
- Are two given graphs the “same”, wrt these properties?

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- As an adjacency matrix:

$$\begin{array}{c} \begin{array}{cccc} & v_1 & v_2 & v_3 & v_4 \\ \begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} & \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & v_1 & v_3 & v_2 & v_4 \\ \begin{array}{c} v_1 \\ v_3 \\ v_2 \\ v_4 \end{array} & \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & a & b & c & d \\ \begin{array}{c} a \\ b \\ c \\ d \end{array} & \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix} \end{array} \end{array}$$

- But we want to study properties that are independent of the naming, e.g., connectivity.
- Are two given graphs the “same”, wrt these properties?

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...


- As an adjacency matrix:

$$\begin{array}{c} \begin{array}{cccc} & v_1 & v_2 & v_3 & v_4 \\ \begin{array}{c} v_1 \\ v_2 \\ v_3 \\ v_4 \end{array} & \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & v_1 & v_3 & v_2 & v_4 \\ \begin{array}{c} v_1 \\ v_3 \\ v_2 \\ v_4 \end{array} & \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & a & b & c & d \\ \begin{array}{c} a \\ b \\ c \\ d \end{array} & \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix} \end{array} \end{array}$$

- **Reordering of vertices** is same as applying a **permutation** to rows and columns of  $A(G)$ .
- So, it seems two graphs are “same” if by reordering and renaming the vertices we get the same graph/matrix.

# Representing and comparing graphs

We start with simple graphs...



To represent it, we need to name the vertices...

- ▶ As an adjacency matrix:

$$\begin{array}{c} \begin{array}{cccc} & v_1 & v_2 & v_3 & v_4 \\ v_1 & \begin{pmatrix} 0 & 1 & 0 & 1 \end{pmatrix} \\ v_2 & \begin{pmatrix} 1 & 0 & 1 & 0 \end{pmatrix} \\ v_3 & \begin{pmatrix} 0 & 1 & 0 & 1 \end{pmatrix} \\ v_4 & \begin{pmatrix} 1 & 0 & 1 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & v_1 & v_3 & v_2 & v_4 \\ v_1 & \begin{pmatrix} 0 & 0 & 1 & 1 \end{pmatrix} \\ v_3 & \begin{pmatrix} 0 & 0 & 1 & 1 \end{pmatrix} \\ v_2 & \begin{pmatrix} 1 & 1 & 0 & 0 \end{pmatrix} \\ v_4 & \begin{pmatrix} 1 & 1 & 0 & 0 \end{pmatrix} \end{array} & \begin{array}{c} \begin{array}{cccc} & a & b & c & d \\ a & \begin{pmatrix} 0 & 0 & 1 & 1 \end{pmatrix} \\ b & \begin{pmatrix} 0 & 0 & 1 & 1 \end{pmatrix} \\ c & \begin{pmatrix} 1 & 1 & 0 & 0 \end{pmatrix} \\ d & \begin{pmatrix} 1 & 1 & 0 & 0 \end{pmatrix} \end{array} \end{array}$$

- ▶ **Reordering of vertices** is same as applying a **permutation** to rows and columns of  $A(G)$ .
- ▶ So, it seems two graphs are “same” if by reordering and renaming the vertices we get the same graph/matrix.
- ▶ How do we formalize this?

# Isomorphism

## Definition

An **isomorphism** from simple graph  $G$  to  $H$  is a **bijection**  $f : V(G) \rightarrow V(H)$  such that  $uv \in E(G)$  iff  $f(u)f(v) \in E(H)$ .

# Isomorphism

## Definition

An **isomorphism** from simple graph  $G$  to  $H$  is a **bijection**  $f : V(G) \rightarrow V(H)$  such that  $uv \in E(G)$  iff  $f(u)f(v) \in E(H)$ .

- ▶ Thus, it is a bijection that “preserves” the edge relation.
- ▶ Can be checked using adjacency matrix by reordering/renaming.
- ▶ What are the properties of this function/relation:  
 $R = \{(G, H) \mid \exists \text{ an isomorphism from } G \text{ to } H\}$ .



# Isomorphism

## Definition

An **isomorphism** from simple graph  $G$  to  $H$  is a **bijection**  $f : V(G) \rightarrow V(H)$  such that  $uv \in E(G)$  iff  $f(u)f(v) \in E(H)$ .

- ▶ Thus, it is a bijection that “preserves” the edge relation.
- ▶ Can be checked using adjacency matrix by reordering/renameing.
- ▶ What are the properties of this function/relation:  
 $R = \{(G, H) \mid \exists \text{ an isomorphism from } G \text{ to } H\}$ .

## Proposition

The isomorphism relation is an equivalence relation.

# Isomorphism

## Definition

An **isomorphism** from simple graph  $G$  to  $H$  is a **bijection**  $f : V(G) \rightarrow V(H)$  such that  $uv \in E(G)$  iff  $f(u)f(v) \in E(H)$ .

- ▶ Thus, it is a bijection that “preserves” the edge relation.
- ▶ Can be checked using adjacency matrix by reordering/renaming.
- ▶ What are the properties of this function/relation:  
 $R = \{(G, H) \mid \exists \text{ an isomorphism from } G \text{ to } H\}$ .

## Proposition

The isomorphism relation is an equivalence relation.

- ▶ The equivalence classes are called isomorphism classes.

# Isomorphism

## Definition

An **isomorphism** from simple graph  $G$  to  $H$  is a **bijection**  $f : V(G) \rightarrow V(H)$  such that  $uv \in E(G)$  iff  $f(u)f(v) \in E(H)$ .




- ▶ Thus, it is a bijection that “preserves” the edge relation.
- ▶ Can be checked using adjacency matrix by reordering/renaming.
- ▶ What are the properties of this function/relation:  
 $R = \{(G, H) \mid \exists \text{ an isomorphism from } G \text{ to } H\}$ .

## Proposition




The isomorphism relation is an equivalence relation.

- ▶ The equivalence classes are called isomorphism classes.
- ▶ When we talked about an “unlabeled” graph till now, we actually meant the isomorphism class of that graph!

# Graph isomorphism




# Graph isomorphism




- ▶ To show that two graphs are isomorphic, you have to
  1. give names to vertices
  2. specify a bijection
  3. check that it preserves the adjacency relation
- ▶ To show that two graphs are **non-isomorphic**, find a structural property that is different.

## Is checking graph isomorphism easy?




- Qn: Which of these graphs are isomorphic?

## Is checking graph isomorphism easy?



- ▶ Qn: Which of these graphs are isomorphic?
- ▶ A: All of them!


## Is checking graph isomorphism easy?



- ▶ Qn: Which of these graphs are isomorphic?
- ▶ A: All of them!
- ▶ This graph is called the **Petersen graph** and has some very interesting properties.




## Is checking graph isomorphism easy?




- ▶ Qn: Which of these graphs are isomorphic?
- ▶ A: All of them!
- ▶ This graph is called the **Petersen graph** and has some very interesting properties.
  - ▶ vertices are 2-element subsets of 5-element set and edges are pairs of disjoint 2-element subsets.

## Is checking graph isomorphism easy?



- ▶ Qn: Which of these graphs are isomorphic?
- ▶ A: All of them!
- ▶ This graph is called the **Petersen graph** and has some very interesting properties.
  - ▶ vertices are 2-element subsets of 5-element set and edges are pairs of disjoint 2-element subsets.
  - ▶ 2 vertices that do not share an edge, have exactly 1 common nbr.


## Is checking graph isomorphism easy?



- ▶ Qn: Which of these graphs are isomorphic?
- ▶ A: All of them!
- ▶ This graph is called the **Petersen graph** and has some very interesting properties.
  - ▶ vertices are 2-element subsets of 5-element set and edges are pairs of disjoint 2-element subsets.
  - ▶ 2 vertices that do not share an edge, have exactly 1 common nbr.

**Further reading:** Graph and sub-graph isomorphism problems.

## Some special graphs and notations



- ▶ Complete graphs  $K_n$
- ▶ Complete bipartite graphs  $K_{i,j}$
- ▶ Paths  $P_n$
- ▶ Cycles  $C_n$

## Some special graphs and notations




Figure: A whole graph zoo!

## Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:

1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

- Are  $C_5$  and  $P_5 \cup \{e\}$  isomorphic?

## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:

1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.



# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.


## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:


1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.
3.  $G, H$  have the same # vertices of degree  $k$ ,  $\forall k \in \mathbb{N}$ .

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.



$G$



$H$

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:

1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.
3.  $G, H$  have the same # vertices of degree  $k$ ,  $\forall k \in \mathbb{N}$ .

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:

1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.
3.  $G, H$  have the same # vertices of degree  $k$ ,  $\forall k \in \mathbb{N}$ .
4.  $G$  has  $k$  paths/cycles of length  $r$  iff  $H$  has  $k$  paths/cycles of length  $r$ .

# Properties of isomorphic graphs

Intuitively, if two graphs are isomorphic then all structural properties, i.e., properties that do not depend on the naming of vertices are preserved.

## Theorem

If  $G$  is isomorphic to  $H$ , then the following properties are preserved:

1.  $G, H$  have same # vertices.
2.  $G, H$  have same # edges.
3.  $G, H$  have the same # vertices of degree  $k$ ,  $\forall k \in \mathbb{N}$ .
4.  $G$  has  $k$  paths/cycles of length  $r$  iff  $H$  has  $k$  paths/cycles of length  $r$ .
5.  $G$  is bipartite iff  $H$  is bipartite.
6.  $G$  contains  $K_n$  as a subgraph iff  $H$  does.
7. ...